

ESTIMATION OF LOW RANK DENSITY MATRICES: BOUNDS IN SCHATTEN NORMS AND OTHER DISTANCES

BY DONG XIA ^{*} AND VLADIMIR KOLTCHINSKII [†]

*School of Mathematics
Georgia Institute of Technology*

Let \mathcal{S}_m be the set of all $m \times m$ density matrices (Hermitian positively semi-definite matrices of unit trace). Consider a problem of estimation of an unknown density matrix $\rho \in \mathcal{S}_m$ based on outcomes of n measurements of observables $X_1, \dots, X_n \in \mathbb{H}_m$ (\mathbb{H}_m being the space of $m \times m$ Hermitian matrices) for a quantum system identically prepared n times in state ρ . Outcomes Y_1, \dots, Y_n of such measurements could be described by a trace regression model in which $\mathbb{E}_\rho(Y_j|X_j) = \text{tr}(\rho X_j)$, $j = 1, \dots, n$. The design variables X_1, \dots, X_n are often sampled at random from the uniform distribution in an orthonormal basis $\{E_1, \dots, E_{m^2}\}$ of \mathbb{H}_m (such as Pauli basis). The goal is to estimate the unknown density matrix ρ based on the data $(X_1, Y_1), \dots, (X_n, Y_n)$. Let

$$\hat{Z} := \frac{m^2}{n} \sum_{j=1}^n Y_j X_j$$

and let $\check{\rho}$ be the projection of \hat{Z} onto the convex set \mathcal{S}_m of density matrices. It is shown that for estimator $\check{\rho}$ the minimax lower bounds in classes of low rank density matrices (established earlier) are attained up logarithmic factors for all Schatten p -norm distances, $p \in [1, \infty]$ and for Bures version of quantum Hellinger distance. Moreover, for a slightly modified version of estimator $\check{\rho}$ the same property holds also for quantum relative entropy (Kullback-Leibler) distance between density matrices.

1. Introduction. Let \mathbb{M}_m be the set of all $m \times m$ matrices with entries in \mathbb{C} . For $A \in \mathbb{M}_m$, let A^* denote its conjugate transpose and let $\text{tr}(A)$ denote the trace of A . The complex linear space \mathbb{M}_m of dimension m^2 will be equipped with the Hilbert-Schmidt inner product $\langle A, B \rangle = \text{tr}(AB^*)$, $A, B \in \mathbb{M}_m$. In what follows, the sign \otimes denotes the tensor product of vectors or matrices (linear transformations). For instance, for $u, v \in \mathbb{C}^m$, $u \otimes v$ is a linear transformation from \mathbb{C}^m into itself defined as follows: $(u \otimes v)x = u\langle x, v \rangle$, $x \in \mathbb{C}^m$.

Let

$$\mathbb{H}_m := \{A \in \mathbb{M}_m : A = A^*\}$$

^{*}Supported in part by NSF Grants DMS-1207808 and CCF-1523768

[†]Supported in part by NSF Grants DMS-1509739 and CCF-1523768

be the set of all Hermitian matrices. Clearly, \mathbb{H}_m is a linear space of dimension m^2 over the field of real numbers. For $A \in \mathbb{H}_m$, the notation $A \succcurlyeq 0$ means that A is positively semi-definite. A density matrix is a positively semi-definite Hermitian matrix of unit trace. The set of all $m \times m$ density matrices will be denoted by

$$\mathcal{S}_m := \{S \in \mathbb{H}_m : S \succcurlyeq 0, \text{tr}(S) = 1\}.$$

Density matrices are used in quantum mechanics to characterize the states of quantum systems. More generally, the states are represented by self-adjoint positively semidefinite operators of unit trace acting in an infinite-dimensional Hilbert space. In this case, density matrices of a large dimension m could be used to approximate the states of the system.

The goal of *quantum state tomography* is to estimate the density matrix for a system prepared in an unknown state based on specially designed measurements. Let $X \in \mathbb{H}_m$ be a Hermitian matrix (*an observable*) with spectral representation $X = \sum_{j=1}^{m'} \lambda_j P_j$, where $m' \leq m$, $\lambda_j \in \mathbb{R}, j = 1, \dots, m'$ being the distinct eigenvalues of X and $P_j, j = 1, \dots, m'$ being the corresponding eigenprojections. For a system prepared in state $\rho \in \mathcal{S}_m$, possible outcomes of a measurement of observable X are the eigenvalues $\lambda_j, j = 1, \dots, m'$ and they occur with probabilities $p_j := \text{tr}(\rho P_j), j = 1, \dots, m'$. If Y is a random variable representing such an outcome, then

$$\mathbb{E}_\rho Y = \text{tr}(\rho X) = \langle \rho, X \rangle.$$

In a simple model of quantum state tomography considered in this paper, an observable X is sampled at random from some probability distribution Π in \mathbb{H}_m , $\mathbb{E}_\rho(Y|X) = \langle \rho, X \rangle$ and $Y = \langle \rho, X \rangle + \xi$ with noise ξ such that $\mathbb{E}_\rho(\xi|X) = 0$. Given a sample X_1, \dots, X_n of n i.i.d. copies of X , n measurements of observables X_1, \dots, X_n are performed for a system identically prepared n times in the same unknown state $\rho \in \mathcal{S}_m$ resulting in outcomes Y_1, \dots, Y_n . This leads to the following *trace regression model*

$$(1.1) \quad Y_j = \langle \rho, X_j \rangle + \xi_j, j = 1, \dots, n$$

with design variables $X_j, j = 1, \dots, n$, response variables $Y_j, j = 1, \dots, n$ and noise $\xi_j, j = 1, \dots, n$ satisfying the assumption $\mathbb{E}_\rho(\xi_j|X_j) = 0, j = 1, \dots, n$ and $\mathbb{E}_\rho(Y_j|X_j) = \langle \rho, X_j \rangle$. The goal is to estimate the target density matrix ρ based on the data $(X_1, Y_1), \dots, (X_n, Y_n)$, with the estimation error being measured by one of the statistically meaningful distances between density matrices such as the Schatten p -norm distances for $p \in [1, \infty]$ or quantum versions of Hellinger and Kullback-Leibler distances.

This version of the problem of quantum state tomography has been intensively studied in the recent years. The noiseless case (quantum compressed sensing) was considered in [12] and [11]. In these papers, sharp bounds on the number n of measurements needed to recover a density matrix of rank r were obtained based on a subtle argument (so called “golfing scheme”) utilizing matrix Bernstein type inequalities. These developments were related to an earlier work on low rank matrix completion [7]. In the noisy case, trace

regression problems have been studied by many authors (see, e.g., [15] and references therein). The main focus was on nuclear norm penalized least squares estimator (matrix LASSO) and related methods such as matrix Dantzig selector (see [6], [19], [25], [14]). In [21], sharp bounds for matrix LASSO and matrix Dantzig selector, in particular, for Pauli measurements in quantum state tomography were obtained. Most of the results in these papers included upper bounds on the estimation error in Hilbert–Schmidt (Frobenius) norm as well as low rank oracle inequalities ([19], [15], [18]). In [19], an upper bound on the operator norm error of a nuclear norm penalized modified least squares estimator was also proved. This result was further developed in [22]. In [16], upper bounds and low rank oracle inequalities for von Neumann entropy penalized least squares estimators were studied (including the bounds on the error in Bures distance and quantum relative entropy distance). A rank penalized estimator of density matrix was studied in [1]. The minimax lower bounds on the Frobenius norm error for matrix completion problems in classes of matrices of rank r were obtained in [19] (the operator norm version could be found in [22]). In [23], a method of deriving lower bounds for unitary invariant matrix norms (including Schatten p -norms) was developed and, among other matrix estimation problems, such bounds were obtained for matrix completion. Minimax lower bounds on the nuclear norm error in density matrix estimation were obtained in [9], where it was also shown that these bounds are attained (up to logarithmic factors) for the matrix versions of LASSO and Dantzig selector. In our recent paper [20], we derived minimax lower bounds in classes of low rank density matrices for the whole range of Schatten p -norm distances as well as for Bures (quantum Hellinger) and quantum relative entropy distance. We also showed that these minimax bounds are attained (up to logarithmic factors) for von Neumann entropy penalized least squares estimators introduced in [16] simultaneously for Bures, relative entropy and Schatten p -norm distances for $p \in [1, 2]$.

The current paper could be viewed as a continuation of [20]. Our main goal is to study a minimal distance estimator $\check{\rho}$ of ρ (initially proposed in [17]) defined as the projection of a simple unbiased estimator

$$\hat{Z} = \frac{m^2}{n} \sum_{j=1}^n Y_j X_j$$

onto the convex set of density matrices \mathcal{S}_m . We show that the minimax error rates established in [20] for the classes of low rank density matrices are attained for this estimator up to logarithmic factors *in the whole range of Schatten p -norm distances* for $p \in [1, \infty]$ as well as for Bures and relative entropy distance. The proof of these results relies on simple properties of projections of Hermitian matrices onto the convex set \mathcal{S}_m of density matrices (see theorems 7 and 8) that might be of independent interest.

Throughout the paper, $\langle \cdot, \cdot \rangle$ denotes either Hilbert–Schmidt inner product (defined above), or (with a little abuse of notation) the canonical inner product of \mathbb{C}^m . The corresponding norm in \mathbb{C}^m is denoted by $|\cdot|$. For $A, B \geq 0$, the notation $A \lesssim B$ means that $A \leq CB$ for a numerical constant $C > 0$, $A \gtrsim B$ means that $B \lesssim A$ and $A \asymp B$ means that $B \lesssim A \lesssim B$. If needed, these signs might be provided with subscripts

indicating that the constant is allowed to depend on parameters. Say, $A \lesssim_\gamma B$ would mean that $A \leq CB$ with C depending on γ .

2. Preliminaries.

2.1. *Distances between density matrices.* The Schatten p -norm of a matrix $A \in \mathbb{H}_m$ is defined as

$$\|A\|_p := \left(\sum_{j=1}^m |\lambda_j(A)|^p \right)^{1/p}, p \in [1, +\infty],$$

where $\lambda_1(A) \geq \dots \geq \lambda_m(A)$ are the eigenvalues of A arranged in a non-increasing order. For $p = 1$, the norm $\|A\|_1$ is called the nuclear or the trace norm; for $p = 2$, $\|A\|_2$ is the Hilbert–Schmidt (generated by the Hilbert–Schmidt inner product) or Frobenius norm; for $p = +\infty$, $\|A\|_\infty = \max_{1 \leq j \leq m} |\lambda_j(A)|$ is called the operator or the spectral norm. Note that, for all $A \in \mathbb{H}_m$, $[1, \infty] \ni p \mapsto \|A\|_p$ is a non-increasing function. The following *interpolation inequality* is well known and can be easily deduced from a similar result for ℓ_p -norms. Let $1 \leq p < q < r \leq \infty$ and let $\mu \in [0, 1]$ be such that $\frac{\mu}{p} + \frac{1-\mu}{r} = \frac{1}{q}$, then

$$(2.1) \quad \|A\|_q \leq \|A\|_p^\mu \|A\|_r^{1-\mu}, A \in \mathbb{H}_m.$$

In addition to the distances generated by the Schatten p -norms, the following two distances (extending well known distances between probability distributions used in the classical statistics) are of importance in quantum statistics: Bures distance and Kullback–Leibler divergence. *The Bures distance* is a quantum version of Hellinger distance and it is defined as follows:

$$H^2(S_1, S_2) := 2 - 2\text{tr}\sqrt{S_1^{1/2}S_2S_1^{1/2}}, S_1, S_2 \in \mathcal{S}_m.$$

The quantity $\text{tr}\sqrt{S_1^{1/2}S_2S_1^{1/2}}$ is called *the fidelity* of states S_1, S_2 (a quantum version of Hellinger affinity). Note that $0 \leq H^2(S_1, S_2) \leq 2$ and that $H(S_1, S_2)$ defines a metric in the space \mathcal{S}_m . The non-commutative *Kullback–Leibler divergence*, or *relative entropy distance* is defined as

$$K(S_1\|S_2) := \text{tr}(S_1 \log S_1 - S_1 \log S_2), S_1, S_2 \in \mathcal{S}_m.$$

If S_2 is a density matrix of rank strictly smaller than m , $\log S_2$ is not well defined and $K(S_1\|S_2) := +\infty$. Clearly, $K(S_1\|S_2)$ is not a metric (in particular, it is not symmetric). It is well known that $K(S_1\|S_2)$ is the supremum of classical Kullback–Leibler divergences between the distributions of outcomes of all possible measurements (represented by positive operator valued measures (POVM)) for the system prepared in states S_1 and S_2 . Similar property holds also for the Bures (Hellinger) distance and for the nuclear norm distance $\|S_1 - S_2\|_1$ which is the supremum of classical total variation distances between

the distributions of outcomes of all measurements (see [26], [13]). These observations easily imply the following inequalities:

$$(2.2) \quad \frac{1}{4} \|S_1 - S_2\|_1^2 \leq H^2(S_1, S_2) \leq K(S_1 \| S_2) \wedge \|S_1 - S_2\|_1$$

(see also [16]).

2.2. Sampling from an orthonormal basis. Uniform sampling from an orthonormal basis is a model of design distribution in trace regression (1.1) that has been frequently used in the literature on quantum compressed sensing (see, [12], [11]). Let $\mathcal{E} := \{E_1, \dots, E_{m^2}\}$ be an orthonormal basis of the space \mathbb{H}_m of Hermitian matrices. Let

$$U := \max_{1 \leq j \leq m^2} \|E_j\|_\infty.$$

Clearly, $U \leq 1$ and

$$1 = \max_{1 \leq j \leq m^2} \|E_j\|_2 \leq m^{1/2} U,$$

implying that $U \geq m^{-1/2}$. In what follows, it will be assumed that Π is a uniform distribution on the basis \mathcal{E} . As a result, the response variables $Y_j, j = 1, \dots, n$ of trace regression model (1.1) could be viewed as noisy measurements of n randomly picked Fourier coefficients of the target density matrix ρ in basis \mathcal{E} . This model includes, in particular, the so called *Pauli measurements*, an important approach to quantum state tomography (see, e.g., [12], [11]).

Example: Pauli bases and Pauli measurements. The space of observables for a single qubit system is the space \mathbb{H}_2 of 2×2 Hermitian matrices. Let

$$\sigma_0 := \begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix}, \quad \sigma_1 := \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix}, \quad \sigma_2 := \begin{pmatrix} 0 & i \\ -i & 0 \end{pmatrix}, \quad \sigma_3 := \begin{pmatrix} 1 & 0 \\ 0 & -1 \end{pmatrix}.$$

The matrices $\sigma_1, \sigma_2, \sigma_3$ (often denoted $\sigma_x, \sigma_y, \sigma_z$) are called *Pauli matrices*. The matrices $W_i = \frac{1}{\sqrt{2}} \sigma_i, i = 0, 1, 2, 3$ form an orthonormal basis of the space \mathbb{H}_2 (the Pauli basis). For a system consisting of k qubits, the space of observables is \mathbb{H}_m , where $m = 2^k$. The Pauli basis of this space is defined by tensorizing the Pauli basis of \mathbb{H}_2 : it consists of $m^2 = 4^k$ tensor products $W_{i_1} \otimes \dots \otimes W_{i_k}, (i_1, \dots, i_k) \in \{0, 1, 2, 3\}^k$. Let $E_1 = W_0 \otimes \dots \otimes W_0$ and let E_2, \dots, E_{m^2} be the rest of the matrices of the Pauli basis of \mathbb{H}_m . It is straightforward to check that $E_1 = \frac{1}{\sqrt{m}} I_m$, where I_m denotes $m \times m$ identity matrix (thus, $\frac{1}{\sqrt{m}}$ is the only eigenvalue of E_1). Matrices E_2, \dots, E_{m^2} have eigenvalues $\pm \frac{1}{\sqrt{m}}$. Therefore, $\|E_j\|_\infty = m^{-1/2}$, implying that, for the Pauli basis, $U = m^{-1/2}$. The fact that the matrices of this basis have the smallest possible operator norms has been used in quantum compressed sensing (see [12], [11], [21]). Matrices E_j have the following spectral representations: $E_j = \frac{1}{\sqrt{m}} P_j^+ - \frac{1}{\sqrt{m}} P_j^-$ with eigenprojections $P_j^+, P_j^-, j = 1, \dots, m^2$ (for $E_1, P_1^- = 0$). A measurement of E_j for a k qubit system prepared in state ρ results in a random outcome

τ_j with two possible values $\pm \frac{1}{\sqrt{m}}$ taken with probabilities $\langle \rho, P_j^\pm \rangle$. For random variable τ_j , $\mathbb{E}_\rho \tau_j = \langle \rho, E_j \rangle$. The density matrix ρ admits the following representation in the Pauli basis:

$$\rho = \sum_{j=1}^{m^2} \frac{\alpha_j}{\sqrt{m}} E_j$$

with $\alpha_1 = 1$ and with some $\alpha_j \in \mathbb{R}, j = 2, \dots, m^2$. This implies that $\mathbb{E}_\rho \tau_j = \frac{\alpha_j}{\sqrt{m}}$,

$$\mathbb{P}_\rho \left\{ \tau_j = \pm \frac{1}{\sqrt{m}} \right\} = \frac{1 \pm \alpha_j}{2}$$

and $\text{Var}_\rho(\tau_j) = \frac{1-\alpha_j^2}{m}$. Note that, for $j = 1$, $\alpha_1 = 1$, $\mathbb{P}_\rho \left\{ \tau_1 = \frac{1}{\sqrt{m}} \right\} = 1$ and $\text{Var}_\rho(\tau_1) = 0$. For $j = 2, \dots, m^2$, $|\alpha_j| < 1$ and $\text{Var}_\rho(\tau_j) > 0$.

Let ν be picked at random from the set $\{1, \dots, m^2\}$ (with the uniform distribution) and let $X = E_\nu, Y = \tau_\nu$ (which corresponds to random sampling from the Pauli basis with a subsequent measurement of observable X resulting in the outcome Y). Then $\mathbb{E}_\rho(Y|X) = \langle \rho, X \rangle$ and $\text{Var}_\rho(Y|X) = \frac{1-\alpha_\nu^2}{m}$. Moreover, we have

$$\mathbb{P} \left\{ \text{Var}_\rho(Y|X) \leq \frac{1}{2m} \right\} = \mathbb{P} \left\{ \alpha_\nu^2 \geq \frac{1}{2} \right\} \leq 2\mathbb{E}\alpha_\nu^2 = \frac{2}{m} \sum_{j=1}^{m^2} \frac{\alpha_j^2}{m} = \frac{2\|\rho\|_2^2}{m}.$$

Since, for $\rho \in \mathcal{S}_m$, $\|\rho\|_2 \leq 1$, this means that, for $m > 2$ with probability at least $1 - \frac{2}{m}$, $\text{Var}_\rho(Y|X) > \frac{1}{2m}$. In other words, the number of $j = 1, \dots, m^2$ such that $\text{Var}_\rho(\tau_j) > \frac{1}{2m}$ is at least $m^2 - 2m$ implying that, for the most of the values of j , $\text{Var}_\rho(\tau_j) \asymp \frac{1}{m}$.

The variance could be further reduced by repeating the measurement of the observable X K times (for a system identically prepared in state ρ) and averaging the outcomes of the resulting K measurements. In this case, the response variable becomes $Y = \langle \rho, X \rangle + \xi$, where $\mathbb{E}_\rho(\xi|X) = 0$ and $\mathbb{E}_\rho(\xi^2|X) = \text{Var}_\rho(Y|X) = \frac{1-\alpha_\nu^2}{Km}$.

2.3. Minimax lower bounds. In [20], the problem of density matrix estimation was studied in the case of trace regression model (1.1) with i.i.d. random design variables X_1, \dots, X_n sampled from the uniform distribution in an orthonormal basis $\mathcal{E} = \{E_1, \dots, E_{m^2}\}$ in two different settings: trace regression with Gaussian noise and trace regression with a bounded response. In both cases, minimax lower bounds on the estimation error of the unknown target density matrix ρ of rank at most r were obtained for the Schatten p -norm distances ($p \in [1, +\infty]$) as well as for the Bures version of quantum Hellinger distance and for the quantum Kullback-Leibler (relative entropy) distance. These results of [20] are stated below.

Denote by $\mathcal{S}_{r,m}$ the set of all density matrices of rank at most r ($1 \leq r \leq m$).

ASSUMPTION 1 (Trace regression with Gaussian noise). *Let (X, Y) be a random couple with X being a random matrix sampled from the uniform distribution Π in an*

orthonormal basis $\mathcal{E} = \{E_1, \dots, E_{m^2}\} \subset \mathbb{H}_m$. Suppose that, for some density matrix $\rho \in \mathcal{S}_m$, $Y = \langle \rho, X \rangle + \xi$, where ξ is a mean zero normal random variable with variance σ_ξ^2 independent of X . Let $(X_1, Y_1), \dots, (X_n, Y_n)$ be n i.i.d. copies of (X, Y) .

In this model, the level of the noise ξ is characterized by its variance which should be involved in the error bound (this could be viewed as a normal approximation of the noise in the case when repeated measurements are performed for each observable X_j with averaging of the outcomes).

THEOREM 1. *Suppose Assumption 1 holds. For all $p \in [1, +\infty]$, there exist constants $c, c' > 0$ such that, the following bounds hold:¹*

$$(2.3) \quad \inf_{\hat{\rho}} \sup_{\rho \in \mathcal{S}_{r,m}} \mathbb{P}_\rho \left\{ \|\hat{\rho} - \rho\|_p \geq c \left(r^{1/p} \frac{\sigma_\xi m^{\frac{3}{2}}}{\sqrt{n}} \wedge \left(\frac{\sigma_\xi m^{3/2}}{\sqrt{n}} \right)^{1-\frac{1}{p}} \wedge 1 \right) \right\} \geq c',$$

$$(2.4) \quad \inf_{\hat{\rho}} \sup_{\rho \in \mathcal{S}_{r,m}} \mathbb{P}_\rho \left\{ H^2(\hat{\rho}, \rho) \geq c \left(r \frac{\sigma_\xi m^{\frac{3}{2}}}{\sqrt{n}} \wedge 1 \right) \right\} \geq c',$$

and

$$(2.5) \quad \inf_{\hat{\rho}} \sup_{\rho \in \mathcal{S}_{r,m}} \mathbb{P}_\rho \left\{ K(\rho \|\hat{\rho}) \geq c \left(r \frac{\sigma_\xi m^{\frac{3}{2}}}{\sqrt{n}} \wedge 1 \right) \right\} \geq c',$$

where $\inf_{\hat{\rho}}$ denotes the infimum over all estimators $\hat{\rho}$ in \mathcal{S}_m based on the data $(X_1, Y_1), \dots, (X_n, Y_n)$ satisfying the Gaussian trace regression model with noise variance σ_ξ^2 .

The trace regression model with a bounded response is characterized by the size U of the range of response variable Y , which usually coincides with the bound on the operator norms of the basis matrices E_j . It includes, in particular, Pauli measurements discussed above (for which $U = m^{-1/2}$).

ASSUMPTION 2 (Trace regression with a bounded response). *Let (X, Y) be a random couple with X being a random matrix sampled from the uniform distribution Π in an orthonormal basis $\mathcal{E} = \{E_1, \dots, E_{m^2}\} \subset \mathbb{H}_m$ with $U := \max_{1 \leq j \leq m^2} \|E_j\|_\infty$ and Y being a random variable with values in the interval $[-U, U]$. Suppose that, for some density matrix $\rho \in \mathcal{S}_m$, $\mathbb{E}(Y|X) = \langle \rho, X \rangle$ a.s. Let $(X_1, Y_1), \dots, (X_n, Y_n)$ be n i.i.d. copies of (X, Y) .*

Let $\mathcal{P}_{r,m}(U)$ denote the class of all distributions P of (X, Y) such that Assumption 2 holds for some $U > 0$ and $\mathbb{E}(Y|X) = \langle \rho_P, X \rangle$ for some $\rho_P \in \mathcal{S}_{r,m}$. For a given $P \in \mathcal{P}_{r,m}(U)$, \mathbb{P}_P denotes the corresponding probability measure such that $(X_1, Y_1), \dots, (X_n, Y_n)$ are i.i.d. copies of (X, Y) .

¹Here \mathbb{P}_ρ denotes a probability measure such that Assumption 1 is satisfied with density matrix ρ .

THEOREM 2. *Suppose Assumption 2 is satisfied and, for some constant $\gamma \in (0, 1)$,*

$$(2.6) \quad \left| \text{tr}(E_k) \right| \leq (1 - \gamma)Um, \quad k = 1, \dots, m^2.$$

Then, for all $p \in [1, +\infty]$, there exist constants $c_\gamma, c'_\gamma > 0$ such that the following bounds hold:

$$(2.7) \quad \inf_{\hat{\rho}} \sup_{P \in \mathcal{P}_{r,m}(U)} \mathbb{P}_P \left\{ \|\hat{\rho} - \rho_P\|_p \geq c_\gamma \left(r^{1/p} \frac{Um^{\frac{3}{2}}}{\sqrt{n}} \wedge \left(\frac{Um^{3/2}}{\sqrt{n}} \right)^{1-\frac{1}{p}} \wedge 1 \right) \right\} \geq c'_\gamma,$$

$$(2.8) \quad \inf_{\hat{\rho}} \sup_{P \in \mathcal{P}_{r,m}(U)} \mathbb{P}_P \left\{ H^2(\hat{\rho}, \rho_P) \geq c_\gamma \left(r \frac{Um^{\frac{3}{2}}}{\sqrt{n}} \wedge 1 \right) \right\} \geq c'_\gamma,$$

and

$$(2.9) \quad \inf_{\hat{\rho}} \sup_{P \in \mathcal{P}_{r,m}(U)} \mathbb{P}_P \left\{ K(\rho_P \|\hat{\rho}) \geq c_\gamma \left(r \frac{Um^{\frac{3}{2}}}{\sqrt{n}} \wedge 1 \right) \right\} \geq c'_\gamma,$$

where $\inf_{\hat{\rho}}$ denotes the infimum over all estimators $\hat{\rho}$ in \mathcal{S}_m based on the i.i.d. data $(X_1, Y_1), \dots, (X_n, Y_n)$ sampled from P .

As it was pointed out in [20] (see Remark 12), if γ in condition (2.6) is small enough (say, $\gamma < 1 - \frac{1}{\sqrt{2}}$), then, in a given orthonormal basis \mathcal{E} , there exists *at most one* matrix E_j such that $\text{tr}(E_j) > (1 - \gamma)Um$. In the case of Pauli basis, such a matrix indeed exists and it is $E_1 = W_0 \otimes \dots \otimes W_0$. Thus, Theorem 2 does not apply directly to the Pauli measurement model. However, the following result does hold (see [20], Theorem 10).

THEOREM 3. *Let $\{E_1, \dots, E_{m^2}\}$ be the Pauli basis in the space \mathbb{H}_m of $m \times m$ Hermitian matrices and let X_1, \dots, X_n be i.i.d. random variables sampled from the uniform distribution in $\{E_1, \dots, E_{m^2}\}$. Let Y_1, \dots, Y_n be outcomes of measurements of observables X_1, \dots, X_n for the system being identically prepared n times in state ρ . The corresponding probability measure will be denoted by \mathbb{P}_ρ . Then, for all $p \in [1, +\infty]$, there exist constants $c, c' > 0$ such that the following bounds hold:*

$$(2.10) \quad \inf_{\hat{\rho}} \sup_{\rho \in \mathcal{S}_{r,m}} \mathbb{P}_\rho \left\{ \|\hat{\rho} - \rho\|_p \geq c \left(r^{1/p} \frac{m}{\sqrt{n}} \wedge \left(\frac{m}{\sqrt{n}} \right)^{1-\frac{1}{p}} \wedge 1 \right) \right\} \geq c',$$

$$(2.11) \quad \inf_{\hat{\rho}} \sup_{\rho \in \mathcal{S}_{r,m}} \mathbb{P}_\rho \left\{ H^2(\hat{\rho}, \rho) \geq c \left(r \frac{m}{\sqrt{n}} \wedge 1 \right) \right\} \geq c',$$

and

$$(2.12) \quad \inf_{\hat{\rho}} \sup_{\rho \in \mathcal{S}_{r,m}} \mathbb{P}_\rho \left\{ K(\rho \|\hat{\rho}) \geq c \left(r \frac{m}{\sqrt{n}} \wedge 1 \right) \right\} \geq c',$$

where $\inf_{\hat{\rho}}$ denotes the infimum over all estimators $\hat{\rho}$ in \mathcal{S}_m based on the data $(X_1, Y_1), \dots, (X_n, Y_n)$.

It was also shown in [20] that, in the case of Schatten p -norm distances for $p \in [1, 2]$, Bures distance and Kullback-Leibler distance, the minimax lower bounds of theorems 1, 2 and 3 are attained up to logarithmic factors in m and n for a penalized least squares estimator with von Neumann entropy penalty introduced in [16]. In the current paper, our main goal is to show that the minimax optimal rates are attained up to logarithmic factors for a very simple minimal distance estimator (that does not require any penalization) *in the whole range* of Schatten p -norms, $p \in [1, \infty]$, as well as for Bures and Kullback-Leibler distances.

3. Main Results. For the model of uniform sampling from an orthonormal basis $\mathcal{E} = \{E_1, \dots, E_{m^2}\}$, the following simple estimator of unknown state $\rho \in \mathcal{S}_m$ is unbiased:

$$\hat{Z} := \frac{m^2}{n} \sum_{j=1}^n Y_j X_j.$$

Indeed,

$$\begin{aligned} \mathbb{E}_\rho \hat{Z} &= m^2 \mathbb{E}_\rho(YX) = m^2 \mathbb{E}(\mathbb{E}_\rho(Y|X)X) = m^2 \mathbb{E} \text{tr}(\rho X) X \\ &= m^2 \mathbb{E} \langle \rho, X \rangle X = m^2 \frac{1}{m^2} \sum_{j=1}^{m^2} \langle \rho, E_j \rangle E_j = \rho. \end{aligned}$$

Clearly, \hat{Z} is not necessarily a density matrix.

We will now define the minimal distance estimator $\check{\rho}$ as the projection of \hat{Z} onto the convex set \mathcal{S}_m of all density matrices. More precisely, for an arbitrary $Z \in \mathbb{H}_m$, define

$$(3.1) \quad \pi_{\mathcal{S}_m}(Z) := \operatorname{argmin}_{S \in \mathcal{S}_m} \|Z - S\|_2^2.$$

Clearly, $\pi_{\mathcal{S}_m}(Z)$ is the closest density matrix to Z with respect to the Hilbert–Schmidt norm distance (that is, the projection of Z onto \mathcal{S}_m ; such a closest density matrix exists in view of compactness of \mathcal{S}_m and it is unique in view of strict convexity of $S \mapsto \|Z - S\|_2^2$). Let

$$\check{\rho} := \pi_{\mathcal{S}_m}(\hat{Z}).$$

REMARK 1. *This definition is equivalent to the following*

$$(3.2) \quad \begin{aligned} \check{\rho} &:= \operatorname{argmin}_{S \in \mathcal{S}_m} \left[-\frac{2}{n} \sum_{j=1}^n Y_j \langle S, X_j \rangle + m^{-2} \|S\|_2^2 \right] = \\ &\operatorname{argmin}_{S \in \mathcal{S}_m} \left[\frac{1}{n} \sum_{j=1}^n Y_j^2 - \frac{2}{n} \sum_{j=1}^n Y_j \langle S, X_j \rangle + m^{-2} \|S\|_2^2 \right] \end{aligned}$$

that was considered in [17] (in [19], similar estimators involving nuclear norm penalty were studied). Note that replacing the term $m^{-2} \|S\|_2^2$ in the right hand side of (3.2) by its unbiased estimator $n^{-1} \sum_{j=1}^n \langle S, X_j \rangle^2$ yields the usual least squares estimator

$$(3.3) \quad \hat{\rho} := \operatorname{argmin}_{S \in \mathcal{S}_m} \left[n^{-1} \sum_{j=1}^n (Y_j - \langle S, X_j \rangle)^2 \right].$$

Note that we also have

$$(3.4) \quad \hat{\rho} := \operatorname{argmin}_{S \in \mathcal{S}_m} \left[n^{-1} \sum_{j=1}^n (Y_j - \langle S, X_j \rangle)^2 + \varepsilon \|S\|_1 \right]$$

since, for $S \in \mathcal{S}_m$, $\|S\|_1 = \operatorname{tr}(S) = 1$. Thus, $\hat{\rho}$ coincides with the nuclear norm penalized least squares estimator (also called the matrix LASSO estimator) for any value of the regularization parameter ε .

We will show that the upper bounds on the error rates in Schatten p -norm distances for $p \in [1, \infty]$ and in Bures distance that match the minimax lower bounds of theorems 1, 2 and 3 up to logarithmic factors hold for the estimator $\check{\rho}$. We will then introduce a simple modification of this estimator for which a matching upper bound holds also for Kullback-Leibler distance.

First, we consider the case of Gaussian trace regression model (Assumption 1). We need an additional assumption that $\sigma_\xi \geq \frac{U}{m^{1/2}}$ (the variance of the noise is not too small).

THEOREM 4. *Suppose Assumption 1 holds and $\sigma_\xi \geq \frac{U}{m^{1/2}}$. For all $p \in [1, +\infty]$, there exists a constant $C > 0$ such that, for all $A \geq 1$ the following bounds hold:*

$$(3.5) \quad \sup_{\rho \in \mathcal{S}_{r,m}} \mathbb{P}_\rho \left\{ \|\check{\rho} - \rho\|_p \geq C \left(r^{1/p} \frac{\sigma_\xi m^{\frac{3}{2}} \sqrt{A \log(2m)}}{\sqrt{n}} \wedge \left(\frac{\sigma_\xi m^{3/2} \sqrt{A \log(2m)}}{\sqrt{n}} \right)^{1-\frac{1}{p}} \wedge 1 \right) \right\} \leq (2m)^{-A}$$

and

$$(3.6) \quad \sup_{\rho \in \mathcal{S}_{r,m}} \mathbb{P}_\rho \left\{ H^2(\check{\rho}, \rho) \geq c \left(r \frac{\sigma_\xi m^{\frac{3}{2}} \sqrt{A \log(2m)}}{\sqrt{n}} \wedge 1 \right) \right\} \leq (2m)^{-A}.$$

If $\sigma_\xi < \frac{U}{m^{1/2}}$, the bounds still hold with σ_ξ replaced by $\frac{U}{m^{1/2}}$.

Similarly, in the case of trace regression with a bounded response, the following result holds.

THEOREM 5. *Suppose Assumption 2 is satisfied. Then, for all $p \in [1, +\infty]$, there exists a constant $C > 0$ such that, for all $A \geq 1$, the following bounds hold:*

$$(3.7) \quad \sup_{P \in \mathcal{P}_{r,m}(U)} \mathbb{P}_P \left\{ \|\check{\rho} - \rho_P\|_p \geq C \left(r^{1/p} \frac{U m^{\frac{3}{2}} \sqrt{A \log(2m)}}{\sqrt{n}} \wedge \left(\frac{U m^{3/2} \sqrt{A \log(2m)}}{\sqrt{n}} \right)^{1-\frac{1}{p}} \wedge 1 \right) \right\} \leq (2m)^{-A}$$

and

$$(3.8) \quad \sup_{P \in \mathcal{P}_{r,m}(U)} \mathbb{P}_P \left\{ H^2(\check{\rho}, \rho_P) \geq C \left(r \frac{U m^{\frac{3}{2}} \sqrt{A \log(2m)}}{\sqrt{n}} \wedge 1 \right) \right\} \leq (2m)^{-A}.$$

For completeness, we state also the upper bounds in the case of Pauli measurements (that immediately follow from Theorem 5).

THEOREM 6. *Suppose the assumptions of Theorem 3 hold. Then, for all $p \in [1, +\infty]$, there exists a constant C such that, for all $A \geq 1$, the following bounds hold:*

$$(3.9) \quad \sup_{\rho \in \mathcal{S}_{r,m}} \mathbb{P}_\rho \left\{ \|\check{\rho} - \rho\|_p \geq c \left(r^{1/p} \frac{m \sqrt{A \log(2m)}}{\sqrt{n}} \wedge \left(\frac{m \sqrt{A \log(2m)}}{\sqrt{n}} \right)^{1-\frac{1}{p}} \wedge 1 \right) \right\} \leq (2m)^{-A}$$

and

$$(3.10) \quad \sup_{\rho \in \mathcal{S}_{r,m}} \mathbb{P}_\rho \left\{ H^2(\check{\rho}, \rho) \geq c \left(r \frac{m}{\sqrt{n}} \wedge 1 \right) \right\} \leq (2m)^{-A}.$$

The proof of these results relies on the following fact that might be of independent interest and that essentially shows that $\pi_{\mathcal{S}_m}(Z)$ is the closest density matrix to Z not only in the Hilbert–Schmidt norm distance, but also in the operator norm distance.

THEOREM 7. *For all $Z \in \mathbb{H}_m$,*

$$\|Z - \pi_{\mathcal{S}_m}(Z)\|_\infty = \inf_{S \in \mathcal{S}_m} \|Z - S\|_\infty.$$

The proof of this theorem will be given in Section 4. Here we use it to establish the next result that is the main ingredient of the proofs of theorems 4, 5 and 6.

THEOREM 8. *Let $p \in [1, +\infty]$. For all $Z \in \mathbb{H}_m$ and all $S \in \mathcal{S}_{r,m}$,*

$$\|\pi_{\mathcal{S}_m}(Z) - S\|_p \leq \min \left(2^{3/p+1} r^{1/p} \|Z - S\|_\infty, 2 \|Z - S\|_\infty^{1-1/p} \right).$$

The proof relies on Theorem 7 and on a simple lemma stated below.

LEMMA 1. *Let $S, S' \in \mathcal{S}_m$ and $\text{rank}(S) = r$. Then, for all $p \in [1, \infty]$,*

$$\|S' - S\|_p \leq \min \left((8r)^{1/p} \|S' - S\|_\infty, 2^{1/p} \|S' - S\|_\infty^{1-1/p} \right).$$

PROOF. Let $S = \sum_{j=1}^r \lambda_j (\phi_j \otimes \phi_j)$ be the spectral decomposition of S with eigenvalues λ_j and eigenvectors ϕ_j . Let $L := \text{supp}(S)$ be the linear span of vectors $\phi_1, \dots, \phi_r \in \mathbb{C}^m$. Denote by P_L, P_{L^\perp} the orthogonal projection operators onto subspace L and its orthogonal complement L^\perp , respectively. We will need the following projection operators $\mathcal{P}_L, \mathcal{P}_L^\perp : \mathbb{H}_m \mapsto \mathbb{H}_m$:

$$\mathcal{P}_L^\perp(A) = P_{L^\perp} A P_{L^\perp}, \quad \mathcal{P}_L(A) = A - P_{L^\perp} A P_{L^\perp}, \quad A \in \mathbb{H}_m.$$

The following bounds are obvious:

$$\begin{aligned} \|S\|_1 = 1 &= \|S'\|_1 = \|S' - S + S\|_1 = \|\mathcal{P}_L(S' - S) + \mathcal{P}_L^\perp(S' - S) + S\|_1 \\ &\geq \|\mathcal{P}_L^\perp(S' - S) + S\|_1 - \|\mathcal{P}_L(S' - S)\|_1. \end{aligned}$$

Since $S = P_L S P_L$, we can use the pinching inequality for unitary invariant norm $\|\cdot\|_1$ (see [4], p. 97) to get:

$$\begin{aligned} \|\mathcal{P}_L^\perp(S' - S) + S\|_1 &= \|P_L S P_L + P_{L^\perp}(S' - S)P_{L^\perp}\|_1 \\ &= \|P_L S P_L\|_1 + \|P_{L^\perp}(S' - S)P_{L^\perp}\|_1 = \|S\|_1 + \|\mathcal{P}_L^\perp(S' - S)\|_1. \end{aligned}$$

Therefore,

$$\|S\|_1 \geq \|S\|_1 + \|\mathcal{P}_L^\perp(S' - S)\|_1 - \|\mathcal{P}_L(S' - S)\|_1,$$

implying that

$$\|\mathcal{P}_L^\perp(S' - S)\|_1 \leq \|\mathcal{P}_L(S' - S)\|_1.$$

It follows from the last bound that

$$\|S' - S\|_1 = \|\mathcal{P}_L(S' - S) + \mathcal{P}_L^\perp(S' - S)\|_1 \leq 2\|\mathcal{P}_L(S' - S)\|_1.$$

Since $\dim(L) = r$, the matrix $\mathcal{P}_L(S' - S)$ is of rank at most $2r$. This implies that

$$\begin{aligned} \|\mathcal{P}_L(S' - S)\|_1 &\leq 2r\|\mathcal{P}_L(S' - S)\|_\infty \\ &\leq 2r(\|(S' - S)P_L\|_\infty + \|P_L(S' - S)P_{L^\perp}\|_\infty) \leq 4r\|S' - S\|_\infty. \end{aligned}$$

Therefore, $\|S' - S\|_1 \leq 8r\|S' - S\|_\infty$, and since also $\|S' - S\|_1 \leq 2$, $S, S' \in \mathcal{S}_m$, we conclude that

$$\|S' - S\|_1 \leq \min(8r\|S' - S\|_\infty, 2).$$

Together with interpolation inequality this yields that for all $p \in [1, \infty]$

$$\|S' - S\|_p \leq \|S' - S\|_1^{1/p} \|S' - S\|_\infty^{1-1/p} \leq \min\left((8r)^{1/p} \|S' - S\|_\infty, 2^{1/p} \|S' - S\|_\infty^{1-1/p}\right).$$

□

PROOF. We now prove Theorem 8. It immediately follows from Theorem 7 that, for all $S \in \mathcal{S}_m$,

$$\|\pi_{\mathcal{S}_m}(Z) - S\|_\infty \leq \|\pi_{\mathcal{S}_m}(Z) - Z\|_\infty + \|Z - S\|_\infty \leq 2\|Z - S\|_\infty.$$

If $S \in \mathcal{S}_m$ is a density matrix of rank r , the last bound could be combined with the bound of Lemma 1 to get that for all $p \in [1, +\infty]$

$$\|\pi_{\mathcal{S}_m}(Z) - S\|_p \leq \min\left(2^{3/p+1} r^{1/p} \|Z - S\|_\infty, 2\|Z - S\|_\infty^{1-1/p}\right).$$

□

PROOF. We now turn to the proof of theorems 4, 5 and 6. To this end, we use the bound of Theorem 8 with $Z = \hat{Z}$ and $S = \rho \in \mathcal{S}_{r,m}$ that yields:

$$(3.11) \quad \|\check{\rho} - \rho\|_p \leq \min\left(2^{3/p+1}r^{1/p}\|\hat{Z} - \rho\|_\infty, 2\|\hat{Z} - \rho\|_\infty^{1-1/p}\right).$$

The control of

$$\|\hat{Z} - \rho\|_\infty = \left\| \frac{m^2}{n} \sum_{j=1}^n Y_j X_j - \rho \right\|_\infty$$

is based on a standard application of matrix Bernstein type inequalities. We give a detailed argument for completeness. Note that $\|\check{\rho} - \rho\|_p$ in the left-hand side of bound (3.11) is upper bounded by 2, so, if Bernstein bound on $\|\hat{Z} - \rho\|_\infty$ is larger than 1 (or even $\gtrsim 1$), it could be replaced by the trivial bound equal to 1. In the case of Theorem 5, we use the following version of Bernstein inequality for i.i.d. bounded random matrices (see, e.g., [28]).

LEMMA 2. *Let V, V_1, \dots, V_n be i.i.d. random matrices in \mathbb{H}_m with $\mathbb{E}V = 0$. Suppose that, for some constant $U > 0$, $\|V\|_\infty \leq U$ a.s. Let $\sigma^2 := \|\mathbb{E}V^2\|_\infty$. Then, for all $t > 0$ with probability at least $1 - e^{-t}$,*

$$\left\| \frac{V_1 + \dots + V_n}{n} \right\|_\infty \leq 2 \left[\sigma \sqrt{\frac{t + \log(2m)}{n}} \vee U \frac{t + \log(2m)}{n} \right].$$

For $V = YX - \mathbb{E}(YX)$, we get, under Assumption 2, that

$$\sigma^2 = \|\mathbb{E}V^2\|_\infty \leq \|\mathbb{E}(Y^2 X^2)\|_\infty \leq U^2 \|\mathbb{E}X^2\|_\infty.$$

It is also well known that, under the same assumption, $\|\mathbb{E}X^2\|_\infty = m^{-1}$. [Indeed, if $\{e_j, j = 1, \dots, m\}$ is an orthonormal basis of \mathbb{C}^m , then

$$\begin{aligned} \|\mathbb{E}X^2\|_\infty &= \sup_{v \in \mathbb{C}^m, |v| \leq 1} \mathbb{E}\langle X^2 v, v \rangle = \sup_{v \in \mathbb{C}^m, |v| \leq 1} \mathbb{E}|Xv|^2 = \sup_{v \in \mathbb{C}^m, |v| \leq 1} \mathbb{E} \sum_{j=1}^m |\langle Xv, e_j \rangle|^2 \\ &= \sup_{v \in \mathbb{C}^m, |v| \leq 1} \mathbb{E} \sum_{j=1}^m |\langle X, v \otimes e_j \rangle|^2 = \sup_{v \in \mathbb{C}^m, |v| \leq 1} \sum_{j=1}^m m^{-2} \sum_{k=1}^m |\langle E_k, v \otimes e_j \rangle|^2 = \sup_{v \in \mathbb{C}^m, |v| \leq 1} m^{-2} \sum_{j=1}^m \|v \otimes e_j\|_2^2 = \\ &\quad \sup_{v \in \mathbb{C}^m, |v| \leq 1} m^{-2} \sum_{j=1}^m |v|^2 |e_j|^2 = m^{-1}. \end{aligned}$$

We use the bound of Lemma 2 with $t = A \log(2m)$, $A \geq 1$ to get that with probability at least $1 - (2m)^{-A}$,

$$\left\| \frac{m^2}{n} \sum_{j=1}^n Y_j X_j - \rho \right\|_\infty \leq C \left[U m^{3/2} \sqrt{\frac{A \log(2m)}{n}} \vee \frac{U^2 m^2 A \log(2m)}{n} \right]$$

with some absolute constant $C \geq 1$. If

$$\frac{U^2 m^2 A \log(2m)}{n} \geq U m^{3/2} \sqrt{\frac{A \log(2m)}{n}},$$

then $U m^{1/2} \sqrt{\frac{A \log(2m)}{n}} \geq 1$ implying that $U m^{3/2} \sqrt{\frac{A \log(2m)}{n}} \geq 1$. Thus, when the bound on $\|\hat{Z} - \rho\|_\infty$ is substituted in bound (3.11), it is enough to keep only the first term $U m^{3/2} \sqrt{\frac{A \log(2m)}{n}}$, the second term could be dropped. This implies that with some constant $C' > 0$ (that does not depend on $\rho \in \mathcal{S}_{r,m}$) the inequality

$$\|\check{\rho} - \rho\|_p \leq C' \left(r^{1/p} \frac{U m^{3/2} \sqrt{A \log(2m)}}{\sqrt{n}} \wedge \left(\frac{U m^{3/2} \sqrt{A \log(2m)}}{\sqrt{n}} \right)^{1-\frac{1}{p}} \wedge 1 \right)$$

holds with probability at least $1 - (2m)^{-A}$, implying the first bound of Theorem 5. The second bound immediately follows from the inequality $H^2(\check{\rho}, \rho) \leq \|\check{\rho} - \rho\|_1$ (see (2.2)). Theorem 6 is an immediate consequence of Theorem 5.

The proof of Theorem 4 is very similar. In this case, Assumption 1 holds and it is natural to split $\hat{Z} - \rho$ into two parts

$$(3.12) \quad \hat{Z} - \rho = \frac{m^2}{n} \sum_{j=1}^n \langle \rho, X_j \rangle X_j - \rho + \frac{m^2}{n} \sum_{j=1}^n \xi_j X_j.$$

and to bound $\|\hat{Z} - \rho\|_\infty$ by triangle inequality. For the first part, an application of matrix Bernstein inequality of Lemma 2 yields the bound

$$(3.13) \quad \left\| \frac{m^2}{n} \sum_{j=1}^n \langle \rho, X_j \rangle X_j - \rho \right\|_\infty \leq C \left[U m \sqrt{\frac{A \log(2m)}{n}} \sqrt{\frac{U^2 m^2 A \log(2m)}{n}} \right]$$

that holds for some absolute constant $C \geq 1$ with probability at least $1 - (2m)^{-A}$. Indeed, in this case $V = \langle \rho, X \rangle X - \mathbb{E} \langle \rho, X \rangle X$ and

$$\sigma^2 \leq \|\mathbb{E} \langle \rho, X \rangle^2 X^2\|_\infty \leq U^2 \mathbb{E} \langle \rho, X \rangle^2 = \frac{U^2 \|\rho\|_2^2}{m^2} \leq \frac{U^2}{m^2},$$

$$\|\langle \rho, X \rangle X\|_\infty \leq \|\rho\|_1 \|X\|_\infty^2 \leq \|X\|_\infty^2 \leq U^2,$$

and Lemma 2 implies (3.13). As before, if $\frac{U^2 m^2 A \log(2m)}{n} \geq U m \sqrt{\frac{A \log(2m)}{n}}$, then $U m \sqrt{\frac{A \log(2m)}{n}} \geq 1$. Thus, the second term $\frac{U^2 m^2 A \log(2m)}{n}$ could be dropped when the bound on $\|\hat{Z} - \rho\|_\infty$ (for which the right hand side of (3.13) is a part) is substituted in (3.11).

As to the second part of representation (3.12) that involves normal random variables ξ_j , it is bounded using another version of matrix Bernstein inequality for not necessarily bounded random matrices (see [16], [15], [18]).

LEMMA 3. *Let V, V_1, \dots, V_n be i.i.d. random matrices in \mathbb{H}_m with $\mathbb{E}V = 0$. Suppose that, for some $\alpha \geq 1$, $U^{(\alpha)} := 2\| \|V\|_\infty \|_{\psi_\alpha} < +\infty$.² Let $\sigma^2 := \|\mathbb{E}V^2\|_\infty$. Then, for all $t > 0$ with probability at least $1 - e^{-t}$,*

$$\left\| \frac{V_1 + \dots + V_n}{n} \right\|_\infty \leq C \left[\sigma \sqrt{\frac{t + \log(2m)}{n}} \vee U^{(\alpha)} \log^{1/\alpha} \left(\frac{U^{(\alpha)}}{\sigma} \right) \frac{t + \log(2m)}{n} \right].$$

We apply the bound of Lemma 3 in the case when $V := \xi X$, $\alpha = 2$ and $t = A \log(2m)$ for $A \geq 1$. By an easy computation,

$$\sigma^2 = \sigma_\xi^2 \|\mathbb{E}X^2\|_\infty = \frac{\sigma_\xi^2}{m}$$

and

$$U^{(2)} = 2\|\xi\|X\|_\infty\|_{\psi_2} \leq 2U\|\xi\|_{\psi_2} \leq 4\sigma_\xi U.$$

This yields the following bound

$$(3.14) \quad \left\| \frac{m^2}{n} \sum_{j=1}^n \xi_j X_j \right\|_\infty \leq C \left[\sigma_\xi m^{3/2} \sqrt{\frac{A \log(2m)}{n}} \vee \sigma_\xi U \frac{m^2 A \log(2m) \log^{1/2}(4U\sqrt{m})}{n} \right]$$

that holds with probability at least $1 - (2m)^{-A}$ and with some absolute constant $C \geq 1$. If the second term in the maximum in the right hand side of (3.14) is dominant, then $Um^{1/2} \sqrt{\frac{A \log(2m)}{n}} \log^{1/2}(4U\sqrt{m}) \geq 1$. Under the condition that $\sigma_\xi \geq Um^{-1/2}$, this implies that also $\sigma_\xi m^{3/2} \sqrt{\frac{A \log(2m)}{n}} \gtrsim 1$. Thus, when the bound in the right hand side of (3.14) (used to control $\|\hat{Z} - \rho\|_\infty$) is substituted in (3.11), it is enough to keep only the first term in the maximum. Finally, under the assumption $\sigma_\xi \geq Um^{-1/2}$, the first term of bound (3.14) dominates the first term of (3.13), so, only this term is needed to control $\|\hat{Z} - \rho\|_\infty$ in bound (3.11). These considerations imply the bound

$$\|\check{\rho} - \rho\|_p \leq C' \left(r^{1/p} \frac{\sigma_\xi m^{\frac{3}{2}} \sqrt{A \log(2m)}}{\sqrt{n}} \wedge \left(\frac{\sigma_\xi m^{3/2} \sqrt{A \log(2m)}}{\sqrt{n}} \right)^{1 - \frac{1}{p}} \wedge 1 \right)$$

that holds with some constant $C' > 0$ (that does not depend on $\rho \in \mathcal{S}_{r,m}$) and with probability at least $1 - (2m)^{-A}$. The first bound of Theorem 4 now follows for all $p \in [1, \infty]$ (which also implies the second bound in view of (2.2)). \square

²Here $\|\cdot\|_{\psi_\alpha}$ denotes the ψ_α Orlicz norm in the space of random variables defined as follows:

$$\|\eta\|_{\psi_\alpha} := \inf \left\{ c > 0 : \mathbb{E} \exp \left\{ \frac{|\eta|^\alpha}{c^\alpha} \right\} \leq 2 \right\}.$$

It turns out that for a slightly modified version of estimator $\check{\rho}$, minimax lower bounds are also attained (up to logarithmic factors) in the case of Kullback-Leibler distance. For $S \in \mathcal{S}_m$ and $\delta \in [0, 1]$, define $S_\delta = (1 - \delta)S + \delta \frac{I_m}{m}$. Clearly, $S_\delta \in \mathcal{S}_m$. Let $\mathcal{S}_{m,\delta} := \{S_\delta : S \in \mathcal{S}_m\}$. Define $\pi_{\mathcal{S}_{m,\delta}}(Z)$ the projection of $Z \in \mathbb{H}_m$ onto the convex set $\mathcal{S}_{m,\delta}$:

$$\pi_{\mathcal{S}_{m,\delta}}(Z) := \operatorname{argmin}_{S \in \mathcal{S}_{m,\delta}} \|Z - S\|_2^2.$$

Let

$$\check{\rho}_\delta := \pi_{\mathcal{S}_{m,\delta}}(\hat{Z})$$

with $\check{\rho}_0 = \check{\rho}$. We will prove the following versions of theorems 4, 5 and 6 for the estimator $\check{\rho}_\delta$.

THEOREM 9. *Suppose Assumption 1 holds, $\sigma_\xi \geq \frac{U}{m^{1/2}}$ and*

$$\delta \leq \frac{\sigma_\xi m^{\frac{3}{2}} \sqrt{\log(2m)}}{\sqrt{n}} \wedge 1.$$

Then bounds (3.5) and (3.6) hold for estimator $\check{\rho}_\delta$. Moreover, for $A \geq 1$, define

$$\lambda := \frac{r \sigma_\xi m^{5/2} \sqrt{\frac{A \log(2m)}{n}} \wedge m}{\delta}.$$

Then, for some constant $c > 0$,

$$(3.15) \quad \sup_{\rho \in \mathcal{S}_{r,m}} \mathbb{P}_\rho \left\{ K(\rho \| \check{\rho}_\delta) \geq c \left(r \frac{\sigma_\xi m^{\frac{3}{2}} \sqrt{A \log(2m)}}{\sqrt{n}} \wedge 1 \right) \log(1 + c\lambda) \right\} \leq (2m)^{-A}.$$

If $\sigma_\xi < \frac{U}{m^{1/2}}$, the bounds still hold with σ_ξ replaced by $\frac{U}{m^{1/2}}$.

THEOREM 10. *Suppose Assumption 2 is satisfied and*

$$\delta \leq \frac{U m^{\frac{3}{2}} \sqrt{\log(2m)}}{\sqrt{n}} \wedge 1.$$

Then (3.7) and (3.8) hold for estimator $\check{\rho}_\delta$. Moreover, for $A \geq 1$, define

$$\lambda := \frac{r U m^{5/2} \sqrt{\frac{A \log(2m)}{n}} \wedge m}{\delta}.$$

Then, for some constant $c > 0$,

$$(3.16) \quad \sup_{\rho \in \mathcal{S}_{r,m}} \mathbb{P}_\rho \left\{ K(\rho \| \check{\rho}_\delta) \geq c \left(r \frac{U m^{\frac{3}{2}} \sqrt{A \log(2m)}}{\sqrt{n}} \wedge 1 \right) \log(1 + c\lambda) \right\} \leq (2m)^{-A}.$$

THEOREM 11. *Suppose the assumptions of Theorem 3 hold and*

$$\delta \leq \frac{m\sqrt{\log(2m)}}{\sqrt{n}} \wedge 1.$$

Then (3.9) and (3.10) hold for estimator $\check{\rho}_\delta$. Moreover, for $A \geq 1$, define

$$\lambda := \frac{rm^2\sqrt{\frac{A\log(2m)}{n}} \wedge m}{\delta}.$$

Then, for some constant $c > 0$,

$$(3.17) \quad \sup_{\rho \in \mathcal{S}_{r,m}} \mathbb{P}_\rho \left\{ K(\rho \| \check{\rho}_\delta) \geq c \left(r \frac{m\sqrt{A\log(2m)}}{\sqrt{n}} \wedge 1 \right) \log(1 + c\lambda) \right\} \leq (2m)^{-A}.$$

REMARK 2. *If, under the assumptions of Theorem 10, we choose*

$$\delta = \frac{Um^{\frac{3}{2}}\sqrt{\log(2m)}}{\sqrt{n}} \wedge 1,$$

then the logarithmic factor in bound (3.16) satisfies the inequality

$$\log(1 + c\lambda) \leq \log(1 + crm\sqrt{A}),$$

so it is of the order $\log m$. Under the assumptions of Theorem 9, this would require the choice of δ

$$\delta = \frac{\sigma_\xi m^{\frac{3}{2}}\sqrt{\log(2m)}}{\sqrt{n}} \wedge 1,$$

so δ would depend on an unknown parameter σ_ξ . Replacing σ_ξ in the definition of δ by the lower bound $Um^{-1/2}$ would result in a logarithmic factor $\lesssim \log\left(1 + crm\sqrt{A}\frac{\sigma_\xi}{Um^{-1/2}}\right)$.

PROOF. We start with the following modification of Theorem 8.

LEMMA 4. *Let $p \in [1, \infty]$. For all $Z \in \mathbb{H}_m$ and all $S \in \mathcal{S}_{r,m}$, the following bound holds:*

$$\|\pi_{\mathcal{S}_{m,\delta}}(Z) - S\|_p \leq \min\left(2^{3/p+1}r^{1/p}\left(\|Z - S\|_\infty + 2\delta\right), 2(1-\delta)^{1/p}\left(\|Z - S\|_\infty + 2\delta\right)^{1-1/p}\right) + 2\delta.$$

PROOF. The following formula is straightforward: for $\delta \in [0, 1)$,

$$\pi_{\mathcal{S}_{m,\delta}}(Z) = (1 - \delta)\pi_{\mathcal{S}_m}\left(\frac{Z}{1 - \delta} - \frac{\delta}{1 - \delta}\frac{I_m}{m}\right) + \delta\frac{I_m}{m}.$$

Indeed, $\pi_{\mathcal{S}_{m,\delta}}(Z)$ coincides with $(1 - \delta)S' + \delta\frac{I_m}{m}$, where

$$\begin{aligned} S' &:= \operatorname{argmin}_{S \in \mathcal{S}_m} \left\| Z - (1 - \delta)S - \delta\frac{I_m}{m} \right\|_2^2 \\ &= \operatorname{argmin}_{S \in \mathcal{S}_m} \left\| \frac{Z}{1 - \delta} - \frac{\delta}{1 - \delta}\frac{I_m}{m} - S \right\|_2^2 = \pi_{\mathcal{S}_m}\left(\frac{Z}{1 - \delta} - \frac{\delta}{1 - \delta}\frac{I_m}{m}\right), \end{aligned}$$

implying the claim.

Let $S \in \mathcal{S}_{r,m}$. Then, for $p \in [1, \infty]$,

$$\begin{aligned} (3.18) \quad \|\pi_{\mathcal{S}_{m,\delta}}(Z) - S\|_p &\leq \|\pi_{\mathcal{S}_{m,\delta}}(Z) - S_\delta\|_p + \|S_\delta - S\|_p \\ &\leq (1 - \delta) \left\| \pi_{\mathcal{S}_m}\left(\frac{Z}{1 - \delta} - \frac{\delta}{1 - \delta}\frac{I_m}{m}\right) - S \right\|_p + 2\delta. \end{aligned}$$

To control the first term in the right hand side, we use the bound of Theorem 8, which requires bounding $\left\| \frac{Z}{1 - \delta} - \frac{\delta}{1 - \delta}\frac{I_m}{m} - S \right\|_\infty$. We have

$$\begin{aligned} (3.19) \quad \left\| \frac{Z}{1 - \delta} - \frac{\delta}{1 - \delta}\frac{I_m}{m} - S \right\|_\infty &= \frac{1}{1 - \delta} \|Z - S_\delta\|_\infty \\ &\leq \frac{1}{1 - \delta} \|Z - S\|_\infty + \frac{1}{1 - \delta} \|S - S_\delta\|_\infty \leq \frac{1}{1 - \delta} \|Z - S\|_\infty + \frac{2\delta}{1 - \delta}. \end{aligned}$$

Using bounds (3.18), (3.19) along with the bound of Theorem 8, we get the bound of the lemma. \square

We will use the bound of Lemma 4 to control $\|\check{\rho}_\delta - \rho\|_p$ for $\rho \in \mathcal{S}_{r,m}$. To this end, we need to bound $\|\hat{Z} - \rho\|_\infty$ using matrix Bernstein inequalities exactly as it was done in the proof of theorems 4, 5 and 6 (under assumptions of these theorems). Denote by $\bar{\Delta}$ such an upper bound on $\|\hat{Z} - \rho\|_\infty$ that holds with probability at least $1 - (2m)^{-A}$. Recall that $\bar{\Delta} \asymp \sigma_\xi m^{3/2} \sqrt{\frac{A \log(2m)}{n}}$ under the conditions of Theorem 4 and $\bar{\Delta} \asymp Um^{3/2} \sqrt{\frac{A \log(2m)}{n}}$ under the conditions of Theorem 5 (it is the same under the conditions of Theorem 6 with $U = m^{-1/2}$). Setting $\Delta = \bar{\Delta} \wedge 1$, we get from the bound of Lemma 4 that

$$\|\check{\rho}_\delta - \rho\|_p \leq \min\left(2^{3/p+1}r^{1/p}(\Delta + 2\delta), 2(1 - \delta)^{1/p}(\Delta + 2\delta)^{1-1/p}\right) + 2\delta$$

that holds with the same probability at least $1 - (2m)^{-A}$. Recall that we replace $\bar{\Delta}$ by Δ since the left hand side $\|\check{\rho}_\delta - \rho\|_p \leq 2$; for the same reason, we can and do drop the

“exponential parts” of matrix Bernstein bounds leaving in the definition of Δ only the “Gaussian parts”. For $\delta \lesssim \Delta$, we get

$$\|\check{\rho}_\delta - \rho\|_p \lesssim \min(r^{1/p} \Delta, \Delta^{1-1/p}).$$

Exactly as in the proof of theorems 4, 5 and 6, this implies that bounds (3.5), (3.6), (3.7), (3.8), (3.9) and (3.10) hold for estimator $\check{\rho}_\delta$.

The bound on the Kullback-Leibler divergence $K(\rho\|\check{\rho}_\delta)$ is an immediate consequence of the bound on $\|\check{\rho}_\delta - \rho\|_1$ and the next lemma that follows from Corollary 1 in [3].

LEMMA 5. *Let $S_1, S_2 \in \mathcal{S}_m$ be density matrices and let $\beta := \lambda_{\min}(S_2)$ be the smallest eigenvalue of S_2 . Suppose that $\beta > 0$. Then*

$$K(S_1\|S_2) \leq \|S_1 - S_2\|_1 \log\left(1 + \frac{\|S_1 - S_2\|_1}{2\beta}\right).$$

We apply Lemma 5 to $S_1 = \rho, S_2 = \check{\rho}_\delta$, observing that $\check{\rho}_\delta \in \mathcal{S}_{m,\delta}$ and $\lambda_{\min}(\check{\rho}_\delta) \geq \delta/m$. We then use the bound on $\|\check{\rho}_\delta - \rho\|_1$ to complete the proof of the bound on $K(\rho\|\check{\rho}_\delta)$. \square

We conclude this section with a simple result concerning the least squares estimator $\hat{\rho}$ defined by (3.3). It shows that the estimators $\hat{\rho}$ and $\check{\rho}$ are close in the Hilbert-Schmidt norm. As a result, the bounds of the previous theorems could be applied to estimator $\hat{\rho}$ as well (at least, under some additional assumptions).

THEOREM 12. *Under the assumption that i.i.d. design variables X_1, \dots, X_n are sampled from the uniform distribution Π in an orthonormal basis $\mathcal{E} = \{E_1, \dots, E_{m^2}\}$, the following bound holds with some constant $C > 0$ for all $A \geq 1$ with probability at least $1 - (2m^2)^{-A}$:*

$$\|\check{\rho} - \hat{\rho}\|_2 \leq Cm \sqrt{\frac{A \log(2m)}{n}}.$$

PROOF. Note that the gradient (and subgradient) of convex function $S \mapsto \|S - Z\|_2^2$ is equal to $2(S - Z)$. By a necessary condition of minimum in convex minimization problem (3.1), for $\check{\rho} = \pi_{\mathcal{S}_m}(\hat{Z})$, $\hat{Z} - \check{\rho}$ should belong to the normal cone $N_{\mathcal{S}_m}(\check{\rho})$ of the convex set \mathcal{S}_m at point $\check{\rho}$ (see [2], Proposition 5, Chapter 4, Section 1). Since both $\check{\rho}, \hat{\rho} \in \mathcal{S}_m$, this implies that

$$(3.20) \quad \langle \check{\rho} - \hat{Z}, \check{\rho} - \hat{\rho} \rangle \leq 0.$$

Similar analysis of convex optimization problem (3.3) shows that

$$\left\langle \frac{m^2}{n} \sum_{j=1}^n (\langle \hat{\rho}, X_j \rangle - Y_j) X_j, \check{\rho} - \hat{\rho} \right\rangle \geq 0,$$

which could be rewritten as follows:

$$(3.21) \quad \left\langle \frac{m^2}{n} \sum_{j=1}^n \langle \hat{\rho}, X_j \rangle X_j - \hat{Z}, \check{\rho} - \hat{\rho} \right\rangle \geq 0.$$

Subtracting (3.21) from (3.20) yields

$$\left\langle \check{\rho} - \frac{m^2}{n} \sum_{j=1}^n \langle \hat{\rho}, X_j \rangle X_j, \check{\rho} - \hat{\rho} \right\rangle \leq 0,$$

implying that

$$(3.22) \quad \|\check{\rho} - \hat{\rho}\|_2^2 = \langle \check{\rho} - \hat{\rho}, \check{\rho} - \hat{\rho} \rangle \leq \left\langle \frac{m^2}{n} \sum_{j=1}^n \langle \hat{\rho}, X_j \rangle X_j - \hat{\rho}, \check{\rho} - \hat{\rho} \right\rangle.$$

We will now write ³

$$\begin{aligned} \frac{m^2}{n} \sum_{j=1}^n \langle \hat{\rho}, X_j \rangle X_j - \hat{\rho} &= \frac{m^2}{n} \sum_{j=1}^n \left(\langle \hat{\rho}, X_j \rangle X_j - \mathbb{E} \langle \hat{\rho}, X \rangle X \right) \\ &= m^2 \left[\frac{1}{n} \sum_{j=1}^n (X_j \otimes X_j - \mathbb{E}(X \otimes X)) \right] \hat{\rho}. \end{aligned}$$

It follows from (3.22) that

$$\|\check{\rho} - \hat{\rho}\|_2^2 \leq m^2 \left\| \frac{1}{n} \sum_{j=1}^n X_j \otimes X_j - \mathbb{E}(X \otimes X) \right\|_{\text{op}} \|\hat{\rho}\|_2 \|\check{\rho} - \hat{\rho}\|_2.$$

Since $\|\hat{\rho}\|_2 \leq 1$, we get

$$(3.23) \quad \|\check{\rho} - \hat{\rho}\|_2 \leq m^2 \left\| \frac{1}{n} \sum_{j=1}^n X_j \otimes X_j - \mathbb{E}(X \otimes X) \right\|_{\text{op}}.$$

It remains to control the operator norm in the right hand side for which we can again use matrix Bernstein inequality of Lemma 2 applying it to $V = X \otimes X - \mathbb{E}(X \otimes X)$. In this case,

$$\begin{aligned} \sigma^2 &= \|\mathbb{E}V^2\|_{\text{op}} \leq \|\mathbb{E}(X \otimes X)^2\|_{\text{op}} = \sup_{\|U\|_2 \leq 1} \mathbb{E} \langle (X \otimes X)^2 U, U \rangle = \sup_{\|U\|_2 \leq 1} \mathbb{E} \langle (X \otimes X) U, (X \otimes X) U \rangle \\ &= \sup_{\|U\|_2 \leq 1} \mathbb{E} |\langle U, X \rangle|^2 \|X\|_2^2 \leq \sup_{\|U\|_2 \leq 1} \mathbb{E} |\langle U, X \rangle|^2 = \sup_{\|U\|_2 \leq 1} \frac{\|U\|_2^2}{m^2} = \frac{1}{m^2} \end{aligned}$$

³Here we view the tensor product $A \otimes B$ of operators $A, B \in \mathbb{M}_m$ as an operator acting from the space \mathbb{M}_m of $m \times m$ matrices equipped with Hilbert-Schmidt inner product $\langle \cdot, \cdot \rangle$ into itself as follows: $(A \otimes B)C = A \langle C, B \rangle$. Let $\|\cdot\|_{\text{op}}$ denote the operator norm of linear operators from \mathbb{M}_m into itself, which corresponds to the $\|\cdot\|_{\infty}$ in the case of $m \times m$ matrices.

and

$$\|V\|_{\text{op}} \leq \|X \otimes X\|_{\text{op}} + \mathbb{E}\|X \otimes X\|_{\text{op}} = \|X\|_2^2 + \mathbb{E}\|X\|_2^2 \leq 2.$$

Bound (3.23) along with the bound of Lemma 2 with $t = A \log(2m^2)$, $A \geq 1$ yield the following inequality

$$\|\check{\rho} - \hat{\rho}\|_2 \lesssim m \sqrt{\frac{A \log(2m)}{n}} \sqrt{m^2 \frac{A \log(2m)}{n}}$$

that holds with probability at least $1 - (2m^2)^{-A}$. Since $\|\check{\rho} - \hat{\rho}\|_2 \leq 2$, the second term $m^2 \frac{A \log(2m)}{n}$ in the right hand side could be dropped (if this term is dominant, the bound is $\gtrsim 1$). This completes the proof of the theorem. \square

Since $\|\check{\rho} - \hat{\rho}\|_\infty \leq \|\check{\rho} - \hat{\rho}\|_2$, the bound of Theorem 12 also holds for $\|\check{\rho} - \hat{\rho}\|_\infty$. Combining this with the bound of Theorem 5 for $p = \infty$, it is easy to conclude that under conditions of this theorem

$$\|\hat{\rho} - \rho\|_\infty \lesssim U m^{3/2} \sqrt{\frac{A \log(2m)}{n}}$$

and that the last bound holds (with a proper choice of constant in relationship \lesssim) with probability at least $1 - (2m)^{-A}$. In view of Lemma 1, this immediately implies that all the bounds of Theorem 5 also hold for the least squares estimator $\hat{\rho}$. In a special case of Pauli measurements, this means that Theorem 6 holds for the estimator $\hat{\rho}$. Concerning Theorem 9, the same conclusion is true under the additional assumption that $\sigma_\xi \geq m^{-1/2}$. Moreover, if $\hat{\rho}_\delta$ is the following modification of estimator $\hat{\rho}$

$$(3.24) \quad \hat{\rho}_\delta := \operatorname{argmin}_{S \in \mathcal{S}_{m,\delta}} \left[n^{-1} \sum_{j=1}^n (Y_j - \langle S, X_j \rangle)^2 \right],$$

then the statements of theorems 9, 10 and 11 hold for the estimator $\hat{\rho}_\delta$ (in the case of Theorem 9, under the additional assumption that $\sigma_\xi \geq m^{-1/2}$).

4. Proof of Theorem 7.

Recall that

$$\pi_{\mathcal{S}_m}(Z) := \operatorname{argmin}_{S \in \mathcal{S}_m} \|Z - S\|_2^2, Z \in \mathbb{H}_m$$

defines the projection of Z onto \mathcal{S}_m . The mapping $\mathbb{H}_m \ni Z \mapsto \pi_{\mathcal{S}_m}(Z) \in \mathcal{S}_m$ possesses a couple of simple properties stated in the next proposition. Denote by \mathcal{S}_m^d the set of all diagonal density matrices.

PROPOSITION 1. 1. For all $m \times m$ unitary matrices U ,

$$\pi_{\mathcal{S}_m}(U^{-1}ZU) = U^{-1}\pi_{\mathcal{S}_m}(Z)U, Z \in \mathbb{H}_m.$$

2. If $D \in \mathbb{H}_m$ is a diagonal matrix, then $\pi_{\mathcal{S}_m}(D) \in \mathcal{S}_m^d$.

PROOF. To prove the first claim, note that, by the unitary invariance of the Hilbert–Schmidt norm,

$$\|U^{-1}ZU - S\|_2^2 = \|U^{-1}(Z - USU^{-1})U\|_2^2 = \|Z - USU^{-1}\|_2^2.$$

In addition, the mapping $S \mapsto USU^{-1}$ is a bijection from the set \mathcal{S}_m onto itself. This immediately implies that

$$\pi_{\mathcal{S}_m}(U^{-1}ZU) = \operatorname{argmin}_{S \in \mathcal{S}_m} \|Z - USU^{-1}\|_2^2 = U^{-1}\pi_{\mathcal{S}_m}(Z)U.$$

For an $m \times m$ matrix $A = (a_{ij})_{i,j=1}^m \in \mathbb{H}_m$, let A^d be the diagonal matrix with diagonal entries $a_{ii}, i = 1, \dots, m$. It is easy to see that if A is a density matrix, then A^d is also a density matrix. Moreover, it is also obvious that, for a diagonal matrix D ,

$$\|D - A^d\|_2^2 \leq \|D - A\|_2^2, A \in \mathcal{S}_m,$$

with a strict inequality if A is not diagonal. These observations immediately imply the second claim. \square

We will now state and prove a vector version of Theorem 7 in which the role of the set of density matrices \mathcal{S}_m is played by the simplex

$$\Delta_m := \left\{ u = (u_1, \dots, u_m) \in \mathbb{R}^m : u_j \geq 0, \sum_{j=1}^m u_j = 1 \right\}$$

in \mathbb{R}^m (this is equivalent to considering the set of diagonal density matrices). We will then show that the matrix version of the problem reduces to the vector case.

Define

$$\pi_{\Delta_m}(z) := \operatorname{argmin}_{u \in \Delta_m} \|z - u\|_{\ell_2^m}^2, z \in \mathbb{R}^m.$$

Since the function $u \mapsto \|z - u\|_{\ell_2^m}^2$ is strictly convex and Δ_m is a compact convex set, such a minimizer exists and is unique. In other words, $\pi_{\Delta_m}(z)$ is the projection of the point $z \in \mathbb{R}^m$ onto simplex Δ_m (the closest point to z in the set Δ_m with respect to the Euclidean ℓ_2^m -distance). The next lemma shows that the same point also minimizes the ℓ_∞^m -distance from z to the simplex Δ_m .

LEMMA 6. For all $z \in \mathbb{R}^m$,

$$\|z - \pi_{\Delta_m}(z)\|_{\ell_\infty^m} = \min_{v \in \Delta_m} \|z - v\|_{\ell_\infty^m}.$$

PROOF. Without loss of generality, assume that $z = (z_1, \dots, z_m) \in \mathbb{R}^m$ is a point with $z_1 \geq \dots \geq z_m$. Denote

$$\bar{z}_j := \frac{z_1 + \dots + z_j}{j}, j = 1, \dots, m.$$

Clearly, $\bar{z}_1 = z_1$ and $\bar{z}_j \geq z_j, j = 1, \dots, m$. Let

$$k := \max \left\{ j \leq m : \bar{z}_j \leq z_j + \frac{1}{j} \right\}.$$

Note that if $k > 1$, then, for all $j < k$, $\bar{z}_j \leq z_j + \frac{1}{j}$. Indeed,

$$\bar{z}_j = \frac{k\bar{z}_k - \sum_{i=j+1}^k z_i}{j} \leq \frac{kz_k + 1 - (k-j)z_k}{j} = \frac{jz_k + 1}{j} = z_k + \frac{1}{j} \leq z_j + \frac{1}{j}.$$

On the other hand, if $k < m$, then $\bar{z}_k > z_{k+1} + \frac{1}{k}$. Indeed, if $\bar{z}_k \leq z_{k+1} + \frac{1}{k}$, then

$$\bar{z}_{k+1} = \frac{k\bar{z}_k + z_{k+1}}{k+1} \leq \frac{kz_{k+1} + 1 + z_{k+1}}{k+1} = z_{k+1} + \frac{1}{k+1},$$

which would contradict the definition of k .

Let $\lambda = (\lambda_1, \dots, \lambda_m)$, where $\lambda_j = z_j - \bar{z}_k + \frac{1}{k}$ for $j = 1, \dots, k$ and $\lambda_j = 0$ for $j = k+1, \dots, m$. Since $\bar{z}_k \leq z_k + \frac{1}{k} \leq z_j + \frac{1}{k}$ for all $j \leq k$, we have $\lambda_j \geq 0, j = 1, \dots, m$ and

$$\sum_{j=1}^m \lambda_j = \sum_{j=1}^k \left(z_j - \bar{z}_k + \frac{1}{k} \right) = \sum_{j=1}^k z_j - k\bar{z}_k + 1 = 1.$$

Thus, $\lambda \in \Delta_m$. It turns out that $\pi_{\Delta_m}(z) = \lambda$.⁴ To prove this it is enough to show that $z - \lambda \in N_{\Delta_m}(\lambda)$, where

$$N_{\Delta_m}(\lambda) := \{u \in \mathbb{R}^m : \langle u, v - \lambda \rangle \leq 0, v \in \Delta_m\}$$

is the normal cone of the convex set Δ_m at point λ (see, e.g., [2], Proposition 5, Chapter 4, Section 1). Let $t := \bar{z}_k - \frac{1}{k}$. Clearly, we have $z_{k+1} < t \leq z_k$ if $k < m$ and $t \leq z_m$ if $k = m$. For $k = m$, $z - \lambda = (t, \dots, t)$ and

$$\langle z - \lambda, v - \lambda \rangle = \sum_{i=1}^m t(v_i - \lambda_i) = t \left(\sum_{i=1}^m v_i - \sum_{i=1}^m \lambda_i \right) = 0$$

since $v, \lambda \in \Delta_m$. For $k < m$, note that

$$z - \lambda = (t, \dots, t, z_{k+1}, \dots, z_m)$$

⁴The computation of the projection onto a simplex occurs in many applications and has been studied before: see, e.g., [24] and [27]. See also [8], where an explicit expression for the projection was derived. For completeness, we provide our version of the proof below.

and, for $v \in \Delta_m$,

$$\langle z - \lambda, v - \lambda \rangle = \sum_{i=1}^k t(v_i - \lambda_i) + \sum_{i=k+1}^m z_i v_i.$$

Using the facts that $\sum_{i=1}^m v_i = 1$ and $\sum_{i=1}^k \lambda_i = 1$, we get

$$\begin{aligned} \langle z - \lambda, v - \lambda \rangle &= t \left(\sum_{i=1}^k v_i - \sum_{i=1}^k \lambda_i \right) + \sum_{i=k+1}^m z_i v_i \\ &= -t \sum_{i=k+1}^m v_i + \sum_{i=k+1}^m z_i v_i = \sum_{i=k+1}^m (z_i - t) v_i \leq 0, \end{aligned}$$

where we also used that, for all $i = k+1, \dots, m$, $z_i - t \leq z_{k+1} - t \leq 0$ and $v_i \geq 0$. Thus, $z - \lambda \in N_{\Delta_m}(\lambda)$ and, by the uniqueness of the minimum, $\lambda = \pi_{\Delta_m}(z)$.

Note that

$$\|z - \lambda\|_{\ell_\infty^m} = \max(|t|, |z_{k+1}|, \dots, |z_m|).$$

For any $v \in \Delta_m$,

$$t = \bar{z}_k - \frac{1}{k} = \frac{1}{k} \sum_{i=1}^k z_i - \frac{1}{k} \sum_{i=1}^m v_i \leq \frac{1}{k} \sum_{i=1}^k z_i - \frac{1}{k} \sum_{i=1}^k v_i = \frac{1}{k} \sum_{i=1}^k (z_i - v_i) \leq \|z - v\|_{\ell_\infty^m}.$$

On the other hand,

$$z_m \geq z_m - v_m \geq -\|z - v\|_{\ell_\infty^m}.$$

Since

$$t = \bar{z}_k - \frac{1}{k} \geq z_{k+1} \geq \dots \geq z_m,$$

we conclude that, for all $v \in \Delta_m$,

$$\|z - \lambda\|_{\ell_\infty^m} \leq \|z - v\|_{\ell_\infty^m}.$$

□

We now turn to the proof of Theorem 7.

PROOF. Any matrix $Z \in \mathbb{H}_m$ admits spectral representation $Z = U^{-1}DU$, where D is the diagonal matrix with real entries d_1, \dots, d_m on the diagonal and U is a unitary $m \times m$ matrix. Let $d = (d_1, \dots, d_m) \in \mathbb{R}^m$. Given $v = (v_1, \dots, v_m) \in \Delta_m$, the diagonal matrix V with entries v_1, \dots, v_m is a density matrix. This defines a bijection $\Delta_m \ni v \mapsto V = J(v)$ between the simplex Δ_m and the set \mathcal{S}_m^d of all diagonal $m \times m$ density matrices. Moreover, J is an isometry of Δ_m and $\mathcal{S}_m^d : \|J(v) - J(u)\|_2^2 = \|u - v\|_{\ell_2^m}^2, u, v \in \Delta_m$.

We will now prove the following lemma.

LEMMA 7. *Let $Z = U^{-1}DU$ with a unitary $m \times m$ matrix U and diagonal matrix D with $d = (d_1, \dots, d_m) \in \mathbb{R}^m$ being the vector of its diagonal entries. Then*

$$\pi_{\mathcal{S}_m}(Z) = U^{-1}J(\pi_{\Delta_m}(d))U.$$

PROOF. This is an immediate consequence of Proposition 1 and the following simple fact:

$$\begin{aligned} \operatorname{argmin}_{A \in \mathcal{S}_m^d} \|D - A\|_2^2 &= J\left(\operatorname{argmin}_{v \in \Delta_m} \|J(d) - J(v)\|_2^2\right) \\ J\left(\operatorname{argmin}_{v \in \Delta_m} \|d - v\|_{\ell_2^m}^2\right) &= J(\pi_{\Delta_m}(d)). \end{aligned}$$

□

To complete the proof of Theorem 7, observe that, In view of lemmas 6, 7,

$$\begin{aligned} \|Z - \pi_{\mathcal{S}_m}(Z)\|_\infty &= \|U^{-1}(J(d) - J(\pi_{\Delta_m}(d)))U\|_\infty \\ &= \|J(d) - J(\pi_{\Delta_m}(d))\|_\infty = \|d - \pi_{\Delta_m}(d)\|_{\ell_\infty^m} = \inf_{v \in \Delta_m} \|d - v\|_{\ell_\infty^m}. \end{aligned}$$

Without loss of generality, assume that $d_1 \geq \dots \geq d_m$. Let $S \in \mathcal{S}_m$ be a density matrix with eigenvalues $v_1 \geq \dots \geq v_m$. Clearly, $v = (v_1, \dots, v_m) \in \Delta_m$. Therefore,

$$\|Z - \pi_{\mathcal{S}_m}(Z)\|_\infty \leq \|d - v\|_\infty \leq \|Z - S\|_\infty,$$

where to get the last bound we used Weyl's perturbation inequality (see [4], Corollary III.2.6).

□

5. Comments on computational aspects of the problem. An advantage of minimal distance estimator $\check{\rho} = \pi_{\mathcal{S}_m}(\hat{Z})$ is the simplicity of its computational implementation. The computation of the matrix $\hat{Z} = \frac{m^2}{n} \sum_{i=1}^n Y_i X_i$ requires $O(nm^2)$ operations. It is followed by an eigen-decomposition of Z that requires $O(m^3)$ operations (see [10]); there exist efficient software packages designed for this kind of tasks, for instance, LINPACK and PROPACK, etc.). As it is shown in the previous section, the problem of computing $\pi_{\mathcal{S}_m}(\hat{Z})$ then reduces to projecting of the vector of eigenvalues of Z arranged in a non-increasing order onto the simplex Δ_m . The last problem has been studied in the literature (see [24], [27], [8]) and it has an explicit solution of computational complexity proportional to m (see the proof of Lemma 6). Thus, the computational implementation of the minimal distance estimator $\check{\rho}$ requires $O((n+m)m^2)$ operations.

The matrix version of LASSO estimator for density matrices is equivalent to solving the following optimization problem

$$(5.1) \quad \hat{\rho} := \operatorname{argmin}_{S \in \mathcal{S}_m} \frac{1}{n} \sum_{i=1}^n \left(Y_i - \langle S, X_i \rangle \right)^2$$

that results in the least squares estimator. Clearly, there is no explicit solution for this optimization problem and it is usually solved by iterative algorithms. For example, a well know iterative singular value thresholding (SVT) algorithm was proposed in [5], and also implemented in quantum compressed sensing in [9]. The main idea is that (5.1) is equivalent to the following optimization problem: for any $\tau > 0$,

$$\hat{\rho} := \arg \min_{S \in \mathcal{S}_m, Z \in \mathbb{H}_m, S=Z} \frac{m^2}{n} \sum_{i=1}^n \left(Y_i - \langle Z, X_i \rangle \right)^2 + \tau \|S - Z\|_2^2.$$

The proposed algorithm updates Z and S alternatively, with the only constraint for S being that $S \in \mathcal{S}_m$. Therefore, the main ingredient of SVT is the following iterative updating rule (with initial $Z_0 = 0$): for $k = 1, 2, \dots$,

$$(5.2) \quad \begin{cases} S_k = \pi_{\mathcal{S}_m}(Z_{k-1}) \\ Z_k = S_k + \delta_k \left(\hat{Z} - \frac{m^2}{n} \sum_{i=1}^n \langle S_k, X_i \rangle X_i \right) \end{cases}$$

with certain pre-determined step sizes $\delta_k > 0$. The algorithm terminates at some step $k = N$ and outputs $S_N \in \mathcal{S}_m$ when $\|S_N - S_{N-1}\|_2 \leq \epsilon$ for some numerical threshold $\epsilon > 0$. It is clear that the minimal distance estimator $\check{\rho}$ can be produced by the above algorithm with one iteration and the initialization $Z_0 = \hat{Z}, \delta_1 = 0$. When the number of qubits k is not small (for instance, about 20) and the dimension m is very large, the iterative algorithm (5.2) is much more computationally expensive than the algorithm for the minimal distance estimator (since every iteration requires the eigen-decomposition of a high dimensional matrix).

References.

- [1] P. Alquier, C. Butucea, M. Hebiri, K. Meziani, and T. Morimae. Rank penalized estimation of a quantum system. *Physical Reviews A*, 88:032113, 2013.
- [2] J.-P. Aubin and I. Ekeland. *Applied Nonlinear Analysis*. Courier Corporation, 2006.
- [3] K. Audenaert and J. Eisert. Continuity bounds on the quantum relative entropy - ii. *Journal of Mathematical Physics*, 52(112201), 2011.
- [4] R. Bhatia. *Matrix Analysis*. Springer, 1997.
- [5] J.-F. Cai, E. J. Candès, and Z. Shen. A singular value thresholding algorithm for matrix completion. *SIAM Journal on Optimization*, 20(4):1956–1982, 2010.
- [6] E. J. Candès and Y. Plan. Tight oracle inequalities for low-rank matrix recovery from a minimal number of noisy random measurements. *IEEE Transactions on Information Theory*, 57(4):2342–2359, 2011.
- [7] E. J. Candès and T. Tao. The power of convex relaxation: Near-optimal matrix completion. *IEEE Transactions on Information Theory*, 56(5):2053–2080, 2010.
- [8] Y. Chen and X. Ye. Projection onto a simplex. *arXiv preprint arXiv:1101.6081*, 2011.
- [9] S. T. Flammia, D. Gross, Y.-K. Liu, and J. Eisert. Quantum tomography via compressed sensing: error bounds, sample complexity and efficient estimators. *New Journal of Physics*, 14(9):095022, 2012.
- [10] G. H. Golub and C. F. Van Loan. *Matrix computations*, volume 3. JHU Press, 2012.
- [11] D. Gross. Recovering low-rank matrices from few coefficients in any basis. *IEEE Transactions on Information Theory*, 57(3):1548–1566, 2011.

- [12] D. Gross, Y.-K. Liu, S. T. Flammia, S. Becker, and J. Eisert. Quantum state tomography via compressed sensing. *Physical Review Letters*, 105(15):150401, 2010.
- [13] H. Klauck, A. Nayak, A. Ta-Shma, and D. Zuckerman. Interaction in quantum communication. *IEEE Transactions on Information Theory*, 53(6):1970–1982, 2007.
- [14] O. Klopp. Noisy low-rank matrix completion with general sampling distribution. *Bernoulli*, 20(1):282–303, 2014.
- [15] V. Koltchinskii. *Oracle Inequalities in Empirical Risk Minimization and Sparse Recovery Problems: École d’Été de Probabilités de Saint-Flour XXXVIII-2008*. Springer, 2011.
- [16] V. Koltchinskii. von Neumann entropy penalization and low-rank matrix estimation. *The Annals of Statistics*, 39(6):2936–2973, 2011.
- [17] V. Koltchinskii. A remark on low rank matrix recovery and noncommutative Bernstein type inequalities. In *From Probability to Statistics and Back: High-Dimensional Models and Processes—A Festschrift in Honor of Jon A. Wellner*, pages 213–226. Institute of Mathematical Statistics, 2013.
- [18] V. Koltchinskii. Sharp oracle inequalities in low rank estimation. In *Empirical Inference*, pages 217–230. Springer, 2013.
- [19] V. Koltchinskii, K. Lounici, and A. B. Tsybakov. Nuclear-norm penalization and optimal rates for noisy low-rank matrix completion. *The Annals of Statistics*, 39(5):2302–2329, 2011.
- [20] V. Koltchinskii and D. Xia. Optimal estimation of low rank density matrices. *Journal of Machine Learning Research*, 16(Sep):1757–1792, 2015.
- [21] Y.-K. Liu. Universal low-rank matrix recovery from Pauli measurements. In *Advances in Neural Information Processing Systems*, pages 1638–1646, 2011.
- [22] K. Lounici. Optimal spectral norm rates for noisy low-rank matrix completion. arxiv:1110.5346, 2011.
- [23] Z. Ma and Y. Wu. Volume ratio, sparsity, and minimaxity under unitarily invariant norms. *IEEE Transactions on Information Theory*, 61(12):6939–6956, 2015.
- [24] C. Michelot. A finite algorithm for finding the projection of a point onto the canonical simplex of \mathbb{R}^n . *Journal of Optimization Theory and Applications*, 50(1):195–200, 1986.
- [25] S. Negahban and M. J. Wainwright. Estimation of (near) low-rank matrices with noise and high-dimensional scaling. *The Annals of Statistics*, 39(2):1069–1097, 2011.
- [26] M. Nielsen and I. Chuang. *Quantum Computation and Quantum Information*. Cambridge University Press, 2000.
- [27] S. Shalev-Shwartz and Y. Singer. Efficient learning of label ranking by soft projections onto polyhedra. *Journal of Machine Learning Research*, 7:1567–1599, 2006.
- [28] J. A. Tropp. User-friendly tail bounds for sums of random matrices. *Foundations of Computational Mathematics*, 12(4):389–434, 2012.

SCHOOL OF MATHEMATICS, GEORGIA INSTITUTE OF TECHNOLOGY, ATLANTA, GA, 30332, USA.
E-MAIL: vlad@math.gatech.edu; dxia7@math.gatech.edu