

A NOTE ON DICKSON POLYNOMIALS OF THE THIRD KIND AND LEGENDRE FUNCTIONS

NERANGA FERNANDO AND SOLOMON MANUKURE

ABSTRACT. In this paper, we show that the Dickson polynomials of the third kind satisfy a nonhomogeneous second order linear ordinary differential equation whose general solution contains Legendre functions.

1. INTRODUCTION

Let n be a non-negative integer. It is well known that the two elementary symmetric polynomials $x_1 + x_2$ and x_1x_2 form a \mathbb{Z} -basis of the ring of symmetric polynomials in $\mathbb{Z}[x_1, x_2]$. In [5], it was shown that there exists a polynomial $D_n(x, y) \in \mathbb{Z}[x, y]$ such that

$$x_1^n + x_2^n = D_n(x_1 + x_2, x_1x_2).$$

The explicit form of $D_n(x, y)$ is given by the Waring's formula [4, Theorem 1.1]

$$D_n(x, y) = \sum_{i=0}^{\lfloor \frac{n}{2} \rfloor} \frac{n}{n-i} \binom{n-i}{i} (-y)^i x^{n-2i}.$$

The polynomial $D_n(x, a) \in R[x]$ is the n -th Dickson polynomial of the first kind, where R is a commutative ring with identity and $a \in R$ is a parameter. In [4], it is shown that the Dickson polynomials of the first kind, $D_n(x, a)$, satisfy the homogeneous second order ordinary differential equation

$$(x^2 - 4a)D_n''(x, a) + xD_n'(x, a) - n^2 D_n(x, a) = 0.$$

The n -th Dickson polynomial of the second kind $E_n(x, a)$ is defined by

$$E_n(x, a) = \sum_{i=0}^{\lfloor \frac{n}{2} \rfloor} \binom{n-i}{i} (-a)^i x^{n-2i},$$

where $a \in R$ is a parameter. In [4], it is also shown that the Dickson polynomials of the second kind, $E_n(x, a)$, satisfy the homogeneous second order ordinary differential equation

$$(x^2 - 4a)E_n''(x, a) + 3xE_n'(x, a) - n(n+2)E_n(x, a) = 0.$$

Dickson polynomials have been widely studied over finite fields for their permutation behaviour. Let \mathbb{F}_q denote the finite field with q elements, where q is a prime power. A permutation polynomial (PP) of \mathbb{F}_q is a polynomial $f \in \mathbb{F}_q[x]$ such that the mapping $x \mapsto f(x)$ is a permutation of \mathbb{F}_q . Permutation polynomials over finite fields have

2010 *Mathematics Subject Classification.* 34A30, 11T06.

Key words and phrases. Dickson polynomial, Differential equation, Associated Legendre function, Hypergeometric function.

important applications in coding theory, cryptography, finite geometry, combinatorics and computer science, among other fields. The reader may find an excellent presentation of Dickson polynomials in [4]. Even though most studies on PPs have been over finite fields, several authors have studied PPs over finite commutative rings. We refer the reader to [3], [7] and references therein for further details on PPs over finite commutative rings.

In [8], Stoll considered Dickson-type polynomials f_n over \mathbb{R} , which depend on two real parameters a and B . The explicit expression for f_n is given by

$$(1.1) \quad f_n(x) = \sum_{i=0}^{\lfloor \frac{n}{2} \rfloor} \frac{n + (B-2)i}{n-i} \binom{n-i}{i} (-a)^i x^{n-2i}.$$

We point out that (1.1) had also appeared earlier in [2].

Again in [8], Stoll showed that the Dickson-type polynomials f_n with $a \neq 0, B \in \mathbb{R}$ satisfy the second order homogeneous ordinary differential equation (See [8, Lemma 17])

$$(1.2) \quad (A_4 x^4 + a A_2 x^2 + a^2 A_0) f_n'' + (B_3 x^3 + a B_1 x) f_n' - (C_2 x^2 + a C_0) f_n = 0,$$

where $A_4, A_2, A_0, B_3, B_1, C_2, C_0 \in \mathbb{R}$.

For $a \in \mathbb{F}_q$, the n -th Dickson polynomial of the $(k+1)$ -th kind $D_{n,k}(x, a)$ as defined by Wang and Yucas in [10] is

$$(1.3) \quad D_{n,k}(x, a) = \sum_{i=0}^{\lfloor \frac{n}{2} \rfloor} \frac{n - ki}{n-i} \binom{n-i}{i} (-a)^i x^{n-2i}.$$

Note that all the coefficients of $D_{n,k}(x, a)$ are integers and $D_{n,k}(x, a)$ also satisfies (1.2) with $A_4, A_2, A_0, B_3, B_1, C_2, C_0 \in \mathbb{Z}$.

The associated Legendre equation [12] is given by

$$(x^2 - 1) \frac{d^2 u}{dx^2} + 2x \frac{du}{dx} - \left[l(l+1) + \frac{m^2}{x^2 - 1} \right] u = 0,$$

where l and m are complex numbers. Its solutions are m -th derivatives of Legendre polynomials, which are a system of complete and orthogonal polynomials, multiplied by the adjustment factor $(1 - x^2)^{\frac{m}{2}}$. These are the associated Legendre functions. The complex numbers l and m are called the degree and order of the associated Legendre functions, respectively. The Legendre polynomials are the associated Legendre functions of order $m = 0$. We refer the reader to [1], [6], [9], and [12] for further details about Legendre functions.

Dickson polynomials have connections with special functions such as Legendre functions and hypergeometric functions. As stated previously, Dickson polynomials of the first and second kind satisfy some linear differential equations. It is well known that the solutions to these differential equations can be represented by Legendre functions. So far, there has been no work on how Dickson polynomials of the $(k+1)$ -th kind are related to Legendre functions. In this note, our main goal is to explore the relationship between the Dickson polynomials of the third kind and associated Legendre

functions. It turns out that the Dickson polynomials of the third kind satisfy a non-homogeneous second order linear ordinary differential equation whose solution contains Legendre functions. First, we show that the particular solution to the resulting differential equation involves the Dickson polynomials of the first kind and subsequently show that the general solution to the associated homogeneous equation involves the Associated Legendre functions which can be expressed in terms of the gamma function and hypergeometric functions.

The paper is organized as follows. In Section 2, we show that the Dickson polynomials of the third kind satisfy the non-homogeneous second-order linear ordinary differential equation given in Lemma 1.1. In Section 3, we find its general solution by proving Theorem 1.2.

By (1.3), the n -th Dickson polynomial of the third kind $D_{n,2}(x, a)$ is given by

$$(1.4) \quad D_{n,2}(x, a) = \sum_{i=0}^{\lfloor \frac{n}{2} \rfloor} \frac{n-2i}{n-i} \binom{n-i}{i} (-a)^i x^{n-2i},$$

where $a \in R$.

Throughout the paper, we denote the n -th Dickson polynomial of the third kind $D_{n,2}(x, a)$ by $F_n(x, a)$. Also, we assume that $R = \mathbb{C}$.

Lemma 1.1. *Let $a \neq 0$, $x = u + au^{-1}$ with $u \neq 0$ and $u^2 \neq a$. Then we have*

$$(1.5) \quad (x^2 - 4a) F_n''(x, a) + 3x F_n'(x, a) - n^2 F_n(x, a) = 2n D_n(x, a),$$

where $D_n(x, a)$ is the n -th Dickson polynomial of the first kind.

In Section 3, we solve (1.5) and prove the following theorem.

Theorem 1.2. *Let $F_c(x, a)$ and $F_p(x, a)$ be the general solution to the associated homogeneous equation of (1.5) and the particular solution of (1.5), respectively. Then*

$$F_c(x, a) = \frac{1}{\sqrt[4]{x^2 - 4a}} \left[A P_{\sqrt{n^2+1}-\frac{1}{2}}^{(\frac{1}{2})} \left(\frac{x}{2\sqrt{a}} \right) + B Q_{\sqrt{n^2+1}-\frac{1}{2}}^{(\frac{1}{2})} \left(\frac{x}{2\sqrt{a}} \right) \right],$$

where A and B are constants, and $P_{\sqrt{n^2+1}-\frac{1}{2}}^{(\frac{1}{2})}$ and $Q_{\sqrt{n^2+1}-\frac{1}{2}}^{(\frac{1}{2})}$ are the associated Legendre functions of the first and second kinds, respectively. Also,

$$F_p(x, a) = \sum_{k=0}^n b_k x^k,$$

where

$$\begin{cases} b_{k+2} = -\left(\frac{n^2 - k(k+2)}{4a(k+2)(k+1)} \right) b_k, & n \text{ is odd, } k \text{ is even or } n \text{ is even, } k \text{ is odd,} \\ b_{k+2} = \frac{1}{4a(k+2)(k+1)} \left\{ [k(k+2) - n^2] b_k - 2n (\text{coefficient of } x^{n-2i} \text{ in } D_n(x, a)) \right\}, \\ \text{otherwise,} \end{cases}$$

where $i = \frac{n-k}{2}$ and $D_n(x, a)$ is the n -th Dickson polynomial of the first kind.

In particular, we have

$$F_n(x, a) = F_c(x, a) + F_p(x, a).$$

1.1. Definitions. The Dickson polynomial $D_n(x, a)$ of the first kind of degree n in the indeterminate x and with parameter $a \in R$ is given as

$$D_n(x, a) = \sum_{i=0}^{\lfloor \frac{n}{2} \rfloor} \frac{n}{n-i} \binom{n-i}{i} (-a)^i x^{n-2i}.$$

Here $\lfloor \frac{n}{2} \rfloor$ denotes the largest integer $\leq n/2$. Let u_1 and u_2 be indeterminates. Then Waring's formula (see [4, Theorem 1.1]) yields

$$u_1^n + u_2^n = \sum_{i=0}^{\lfloor \frac{n}{2} \rfloor} \frac{n}{n-i} \binom{n-i}{i} (-u_1 u_2)^i (u_1 + u_2)^{n-2i}$$

and thus

$$u_1^n + u_2^n = D_n(u_1 + u_2, u_1 u_2).$$

If we let $u_1 = u$, $u_2 = \frac{a}{u}$ and $x = u + \frac{a}{u}$, then the functional equation of the Dickson polynomial of the first kind is given by

$$(1.6) \quad D_n\left(u + \frac{a}{u}, a\right) = u^n + \left(\frac{a}{u}\right)^n.$$

For $a \neq 0$, Let $x = u + au^{-1}$ with $u \neq 0$ and $u^2 \neq a$. Then the functional equation of the Dickson polynomial of the third kind, $F_n(x, a)$, is given by

$$(1.7) \quad F_n(x, a) = \left(u + \frac{a}{u}\right) \frac{u^n - \left(\frac{a}{u}\right)^n}{u - \frac{a}{u}},$$

see [10, Eq. 4.1].

For m a real number, the Leibniz rule for the product of two functions $u(z)$ and $v(z)$ is given by

$$(1.8) \quad D_z^m uv = \sum_{\alpha=0}^{\infty} \binom{m}{\alpha} D_z^{m-\alpha} u D_z^{\alpha} v,$$

where

$$\binom{m}{\alpha} = \frac{\Gamma(m+1)}{\Gamma(m-\alpha+1)\Gamma(\alpha+1)}.$$

2. PROOF OF LEMMA 1.1

From (1.7), we have

$$F_n(u + au^{-1}, a) \left(u - \frac{a}{u}\right) = \left(u + \frac{a}{u}\right) \left[u^n - \left(\frac{a}{u}\right)^n\right].$$

By differentiating both sides with respect to u , we get

$$\begin{aligned} & F'_n(u + au^{-1}, a) \left(u - \frac{a}{u}\right) \left(1 - \frac{a}{u^2}\right) + F_n(u + au^{-1}, a) \left(1 + \frac{a}{u^2}\right) \\ &= \left(u + \frac{a}{u}\right) \left[nu^{n-1} + \frac{na^n}{u^{n+1}}\right] + \left[u^n - \left(\frac{a}{u}\right)^n\right] \left(1 - \frac{a}{u^2}\right). \end{aligned}$$

Multiplying by u and differentiating both sides with respect to u again yields

$$\begin{aligned} & F''_n(u + au^{-1}, a) \left(u - \frac{a}{u}\right)^2 \left(1 - \frac{a}{u^2}\right) + 2 F'_n(u + au^{-1}, a) \left(u - \frac{a}{u}\right) \left(1 + \frac{a}{u^2}\right) \\ &+ F'_n(u + au^{-1}, a) \left(u + \frac{a}{u}\right) \left(1 - \frac{a}{u^2}\right) + F_n(u + au^{-1}, a) \left(1 - \frac{a}{u^2}\right) \\ &= n \left(u + \frac{a}{u}\right) \left[nu^{n-1} - \frac{na^n}{u^{n+1}}\right] + n \left[u^n + \left(\frac{a}{u}\right)^n\right] \left(1 - \frac{a}{u^2}\right) \\ &+ \left[u^n - \left(\frac{a}{u}\right)^n\right] \left(1 + \frac{a}{u^2}\right) + \left(u - \frac{a}{u}\right) \left[nu^{n-1} + \frac{na^n}{u^{n+1}}\right]. \end{aligned}$$

Multiplying by u gives

$$\begin{aligned} & \left(u - \frac{a}{u}\right)^3 F''_n(u + au^{-1}, a) + 3 \left[u^2 - \left(\frac{a}{u}\right)^2\right] F'_n(u + au^{-1}, a) \\ &+ \left(u - \frac{a}{u}\right) F_n(u + au^{-1}, a) \\ &= (n^2 + 1) \left(u - \frac{a}{u}\right) F_n(u + au^{-1}, a) + 2n \left(u - \frac{a}{u}\right) \left[u^n + \left(\frac{a}{u}\right)^n\right], \end{aligned}$$

which simplifies to

$$\begin{aligned} (2.1) \quad & \left(u - \frac{a}{u}\right)^2 F''_n(u + au^{-1}, a) + 3 \left(u + \frac{a}{u}\right) F'_n(u + au^{-1}, a) - n^2 F_n(u + au^{-1}, a) \\ &= 2n \left[u^n + \left(\frac{a}{u}\right)^n\right]. \end{aligned}$$

Note that, $D_n(x, a) = u^n + \left(\frac{a}{u}\right)^n$, where $D_n(x, a)$ is the n -th Dickson polynomial of the first kind (See (1.6)). Also note that, $\left(u - \frac{a}{u}\right)^2 = x^2 - 4a$.

From 2.1 we have,

$$(x^2 - 4a) F''_n(x, a) + 3x F'_n(x, a) - n^2 F_n(x, a) = 2n D_n(x, a).$$

Remark 2.1. Let $a \neq 0$ and $x = u + au^{-1}$ with $u \neq 0$. When $u^2 = a$, the functional equation of $F_n(a, x)$ is given by

$$(2.2) \quad F_n(x, a) = 2(\pm\sqrt{a})^n n;$$

see [10, Remark 2.4]. In this case, we have $F'_n(x, a) = 0$.

3. PROOF OF THEOREM 1.2

In Section 2, we showed that Dickson polynomials of the third kind satisfy the second order non-homogeneous differential equation

$$(3.1) \quad (x^2 - 4a) F''_n(x, a) + 3x F'_n(x, a) - n^2 F_n(x, a) = 2n D_n(x, a).$$

We first find the particular solution to (3.1), $F_p(x, a)$.

Let $F_p = \sum_{k=0}^n b_k x^k$ be the trial solution of the particular solution to (3.1). Then we have

$$F'_p = \sum_{k=1}^n k b_k x^{k-1} \quad \text{and} \quad F''_p = \sum_{k=2}^n k(k-1) b_k x^{k-2}.$$

Then

$$(3.2) \quad \begin{aligned} & (x^2 - 4a) F''_n(x, a) + 3x F'_n(x, a) - n^2 F_n(x, a) \\ &= (x^2 - 4a) \sum_{k=2}^n k(k-1) b_k x^{k-2} + 3x \sum_{k=1}^n k b_k x^{k-1} - n^2 \sum_{k=0}^n b_k x^k \\ &= \sum_{k=0}^{n-2} [k(k-1) b_k + 3k b_k - n^2 b_k - 4a(k+2)(k+1) b_{k+2}] x^k - b_{n-1} x^{n-1} + 2n b_n x^n \end{aligned}$$

Let $b_{n+1} = b_{n+2} = 0$. Now (3.2) can be written as

$$\begin{aligned} & (x^2 - 4a) F''_n(x, a) + 3x F'_n(x, a) - n^2 F_n(x, a) \\ &= \sum_{k=0}^n [k(k-1) b_k + 3k b_k - n^2 b_k - 4a(k+2)(k+1) b_{k+2}] x^k \end{aligned}$$

From (3.1) we have

$$(3.3) \quad \begin{aligned} & \sum_{k=0}^n [k(k-1) b_k + 3k b_k - n^2 b_k - 4a(k+2)(k+1) b_{k+2}] x^k \\ &= 2n \sum_{i=0}^{\lfloor \frac{n}{2} \rfloor} \frac{n}{n-i} \binom{n-i}{i} (-a)^i x^{n-2i}. \end{aligned}$$

Consider the following cases.

- (i) n is odd and k is even,
- (ii) n is even and k is odd,
- (iii) n is odd and k is odd, and
- (iv) n is even and k is even.

In cases (i) and (ii), by comparing the coefficients in (3.3) we have

$$k(k-1)b_k + 3kb_k - n^2b_k - 4a(k+2)(k+1)b_{k+2} = 0$$

which implies

$$b_{k+2} = -\left(\frac{n^2 - k(k+2)}{4a(k+2)(k+1)}\right)b_k.$$

Now consider cases (iii) and (iv). By comparing the coefficients in (3.3) we have

$$b_{k+2} = \frac{1}{4a(k+2)(k+1)} \left\{ [k(k+2) - n^2]b_k - 2n(\text{coefficient of } x^{n-2i} \text{ in } D_n(x, a)) \right\},$$

where $i = \frac{n-k}{2}$.

Now let's consider the associated homogeneous equation, i.e.

$$(3.4) \quad (x^2 - 4a) F_n''(x, a) + 3x F_n'(x, a) - n^2 F_n(x, a) = 0.$$

Denote $F_n(x, a)$ by $F(x)$ and let $X = \frac{x}{2\sqrt{a}}$. It then follows that

$$2\sqrt{a} \frac{d}{dX} \left[(X^2 - 1)^{\frac{3}{2}} F'(2\sqrt{a}X) \right] - n^2 (X^2 - 1)^{\frac{1}{2}} F(2\sqrt{a}X) = 0.$$

Letting $P(X) = F(2\sqrt{a}X)$ gives

$$(3.5) \quad \frac{d}{dX} \left[(X^2 - 1)^{\frac{3}{2}} P'(X) \right] - n^2 (X^2 - 1)^{\frac{1}{2}} P(X) = 0.$$

Now consider the Legendre equation:

$$(1 - x^2)P_l''(x) - 2x P_l'(x) + l(l+1)P_l(x) = 0$$

or equivalently

$$(3.6) \quad (x^2 - 1)P_l''(x) + 2x P_l'(x) - l(l+1)P_l(x) = 0,$$

where $l \in \mathbb{R}$ with $l \geq 0$.

Differentiating $m \in \mathbb{R}$ times using Leibniz rule (see (1.8)) gives the following.

$$(3.7) \quad \begin{aligned} D_x^m[(x^2 - 1)P_l''(x)] &= \sum_{\alpha=0}^{\infty} \frac{\Gamma(m+1)}{\Gamma(m-\alpha+1)\Gamma(\alpha+1)} D_x^\alpha(x^2 - 1) D^{m-\alpha} P_l''(x) \\ &= (x^2 - 1)P_l^{(m+2)}(x) + 2mx P_l^{(m+1)}(x) + m(m-1)P_l^{(m)}(x). \end{aligned}$$

Similarly,

$$(3.8) \quad D_x^m[2x P_l'(x)] = 2x P_l^{(m+1)}(x) + 2m P_l^{(m)}(x).$$

(3.6), (3.7), and (3.8) yield

$$(3.9) \quad (x^2 - 1)P_l^{(m+2)}(x) + 2(m+1)x P_l^{(m+1)}(x) - (l-m)(l+m+1)P_l^{(m)}(x) = 0.$$

Now let $y(x) = P_l^{(m)}$. Then (3.9) becomes

$$(3.10) \quad (x^2 - 1)\frac{d^2y}{dx^2} + 2(m+1)x \frac{dy}{dx} - (l-m)(l+m+1)y = 0.$$

By multiplying (3.10) by $(x^2 - 1)^m$, we get

$$(3.11) \quad \frac{d}{dx} \left[(x^2 - 1)^{m+1} \frac{dy}{dx} \right] - (l-m)(l+m+1) (x^2 - 1)^m y = 0.$$

Now let

$$(3.12) \quad y(x) = (x^2 - 1)^{-\frac{m}{2}} u(x).$$

Then

$$\frac{dy}{dx} = (x^2 - 1)^{-\frac{m}{2}} \frac{du}{dx} - mx(x^2 - 1)^{-\frac{m}{2}-1} u.$$

Thus (3.11) becomes

$$\begin{aligned} & (x^2 - 1)^{\frac{m}{2}+1} \frac{d^2u}{dx^2} + \left(\frac{m}{2} + 1\right)(2x)(x^2 - 1)^{\frac{m}{2}} \frac{du}{dx} - mx(x^2 - 1)^{\frac{m}{2}} \frac{du}{dx} - m(x^2 - 1)^{\frac{m}{2}} u \\ & - mx \frac{m}{2} (2x) (x^2 - 1)^{\frac{m}{2}-1} u - (l-m)(l+m+1) (x^2 - 1)^{\frac{m}{2}} u = 0. \end{aligned}$$

Dividing by $(x^2 - 1)^{\frac{m}{2}}$, we obtain

$$(3.13) \quad (x^2 - 1) \frac{d^2u}{dx^2} + 2x \frac{du}{dx} - \left[l(l+1) + \frac{m^2}{x^2 - 1} \right] u = 0,$$

which is the associated Legendre differential equation, with $0 \leq m \leq l$. Its general solution is given by (see [11, Ch. 5])

$$u_g = c_1 P_l^{(m)}(x) + c_2 Q_l^{(m)}(x),$$

where c_1 and c_2 are arbitrary constants, and $P_l^{(m)}(x)$ and $Q_l^{(m)}(x)$ are the associated Legendre functions of the first and second kinds, respectively. Thus, according to (3.12)

$$y_g(x) = (x^2 - 1)^{-\frac{m}{2}} u_g(x)$$

is the general solution to (3.11).

Now consider the equation

$$(3.14) \quad \frac{d}{dx} \left[(x^2 - 1)^{\frac{3}{2}} \frac{dy}{dx} \right] - n^2 (x^2 - 1)^{\frac{1}{2}} y = 0.$$

This is in the form of (3.11) with $m = \frac{1}{2}$ and $(l-m)(l+m+1) = n^2$. Since $l \in \mathbb{R}$ with $l \geq 0$, we have $l = \sqrt{n^2 + 1} - \frac{1}{2}$. Then the solution to (3.14) is

$$y = (x^2 - 1)^{-\frac{1}{4}} \left[c_1 P_{\sqrt{n^2+1}-\frac{1}{2}}^{(\frac{1}{2})}(x) + c_2 Q_{\sqrt{n^2+1}-\frac{1}{2}}^{(\frac{1}{2})}(x) \right].$$

So the solution to (3.5) is

$$P(X) = (X^2 - 1)^{-\frac{1}{4}} \left[c_1 P_{\sqrt{n^2+1}-\frac{1}{2}}^{(\frac{1}{2})}(X) + c_2 Q_{\sqrt{n^2+1}-\frac{1}{2}}^{(\frac{1}{2})}(X) \right].$$

Since $X = \frac{x}{2\sqrt{a}}$ and $P(X) = F(2\sqrt{a}X)$, we have

$$F(x) = (x^2 - 4a)^{-\frac{1}{4}} \left[\frac{c_1}{(4a)^{-\frac{1}{4}}} P_{\sqrt{n^2+1}-\frac{1}{2}}^{(\frac{1}{2})}\left(\frac{x}{2\sqrt{a}}\right) + \frac{c_2}{(4a)^{-\frac{1}{4}}} Q_{\sqrt{n^2+1}-\frac{1}{2}}^{(\frac{1}{2})}\left(\frac{x}{2\sqrt{a}}\right) \right].$$

If we denote the general solution to (3.4) by $F_c(x, a)$, then we have

$$F_c(x, a) = \frac{1}{\sqrt[4]{x^2 - 4a}} \left[A P_{\sqrt{n^2+1}-\frac{1}{2}}^{(\frac{1}{2})}\left(\frac{x}{2\sqrt{a}}\right) + B Q_{\sqrt{n^2+1}-\frac{1}{2}}^{(\frac{1}{2})}\left(\frac{x}{2\sqrt{a}}\right) \right]$$

where A and B are arbitrary constants. Hence the general solution to the differential equation

$$(x^2 - 4a) F_n''(x, a) + 3x F_n'(x, a) - n^2 F_n(x, a) = 2n D_n(x, a),$$

is

$$(3.15) \quad F_n(x, a) = \frac{1}{\sqrt[4]{x^2 - 4a}} \left[A P_{\sqrt{n^2+1}-\frac{1}{2}}^{(\frac{1}{2})}\left(\frac{x}{2\sqrt{a}}\right) + B Q_{\sqrt{n^2+1}-\frac{1}{2}}^{(\frac{1}{2})}\left(\frac{x}{2\sqrt{a}}\right) \right] + \sum_{k=0}^n b_k x^k,$$

where b_k is determined as explained in Theorem 1.2.

In terms of hypergeometric functions, the Legendre functions can be written as (see [11, Ch. 5])

$$(3.16) \quad P_{\sqrt{n^2+1}-\frac{1}{2}}^{(\frac{1}{2})}(z) = \frac{1}{\Gamma(\frac{1}{2})} \left[\frac{1+z}{1-z} \right]^{\frac{1}{4}} {}_2F_1\left(-\sqrt{n^2+1} + \frac{1}{2}, \sqrt{n^2+1} + \frac{1}{2}; \frac{1}{2}; \frac{1-z}{2}\right),$$

for $|1 - z| < 2$, and

$$(3.17) \quad \begin{aligned} & Q_{\sqrt{n^2+1}-\frac{1}{2}}^{(\frac{1}{2})}(z) \\ &= \frac{\sqrt{\pi}}{2\sqrt{n^2+1}+\frac{1}{2}} \frac{i(z^2-1)^{\frac{1}{4}}}{z\sqrt{n^2+1}+1} {}_2F_1\left(\frac{\sqrt{n^2+1}+1}{2}, \frac{\sqrt{n^2+1}+2}{2}; \sqrt{n^2+1}+1; \frac{1}{z^2}\right) \end{aligned}$$

for $|z| > 1$,

where Γ denotes the gamma function and ${}_2F_1(a, b; c; z)$ is the hypergeometric function defined by the power series

$${}_2F_1(a, b; c; z) = \sum_{n=0}^{\infty} \frac{(a)_n (b)_n}{(c)_n} \frac{z^n}{n!},$$

for $|z| < 1$.

Here $(\cdot)_n$ denotes the rising factorial defined by

$$(a)_n = \begin{cases} 1 & , \quad n = 0 \\ a(a+1) \cdots (a+n-1) & , \quad n > 0. \end{cases}$$

4. CONCLUDING REMARKS

In this note we have shown that there is a relationship between Dickson polynomials of the third kind and Legendre functions through the existence of a second order linear differential equation. Currently we do not know if such a differential equation exists for the Dickson polynomials of the $(k+1)$ -th kind. The nature of our computations clearly indicates that finding such a differential equation, if it exists, may be an arduous exercise.

REFERENCES

- [1] G. B. Arfken, H. J. Weber, *Mathematical methods for physicists*. Fifth edition. Harcourt/Academic Press, Burlington, MA, 2001. xiv+1112 pp.
- [2] A. Dujella, I. Gusić, *Decomposition of a recursive family of polynomials*, Monatsh. Math. **152** (2007), no. 2, 97 – 104.
- [3] D. Görcsös, G. Horváth, and A. Mészáros, *Permutation polynomials over finite rings*, Finite Fields Appl. **49** (2018), 198 – 211.
- [4] R. Lidl, G. L. Mullen, G. Turnwald, *Dickson polynomials*, Longman Scientific and Technical, Essex, United Kingdom, 1993.
- [5] I. G. Macdonald, *Symmetric Functions and Orthogonal Polynomials*, American Mathematical Society, Providence, RI, 1998.
- [6] F. W. J. Olver, D. W. Lozier, R. F. Boisvert, and C. W. Clark, *NIST Handbook of Mathematical Functions* National Institute of Standards and Technology and Cambridge University Press, Gaithersburg and Cambridge, 2010.
- [7] R. L. Rivest, *Permutation polynomials modulo 2^w* , Finite Fields Appl. **7** (2001), no. 2, 287 – 292.
- [8] T. Stoll, *Complete decomposition of Dickson-type polynomials and related Diophantine equations*, Journal of Number Theory **128** (2008), 1157 – 1181.
- [9] N. Virchenko, I. Fedotova, *Generalized associated Legendre functions and their applications*. With a foreword by Semyon Yakubovich. World Scientific Publishing Co., Inc., River Edge, NJ, 2001. xx+195 pp.
- [10] Q. Wang, J. L. Yucas, *Dickson polynomials over finite fields*, Finite Fields Appl. **18** (2012), 814 – 831.
- [11] Z. X. Wang, D. R. Guo, *Special Functions*, World Scientific Publishing Co. Pte. Ltd., 1989.
- [12] G. M. Wysin, *Associated Legendre Functions and Dipole Transition Matrix Elements*, online lecture notes.

DEPARTMENT OF MATHEMATICS, NORTHEASTERN UNIVERSITY, BOSTON, MA 02115
E-mail address: w.fernando@northeastern.edu

DEPARTMENT OF MATHEMATICS, THE UNIVERSITY OF TEXAS AT AUSTIN, AUSTIN, TX 78712
E-mail address: smanukure@math.utexas.edu