Nonexistence of a few binary orthogonal arrays *

Peter Boyvalenkov

Institute of Mathematics and Informatics, Bulgarian Academy of Sciences, 8 G.Bonchev Street, 1113, Sofia, BULGARIA and Faculty of Mathematics and Natural Sciences, South-Western University, Blagoevgrad, Bulgaria. email: peter@math.bas.bg Tanya Marinova, Maya Stoyanova Faculty of Mathematics and Informatics, Sofia University, 5 James Bourchier Blvd., 1164 Sofia, BULGARIA email: tanya.marinova@fmi.uni-sofia.bg email: stoyanova@fmi.uni-sofia.bg

Abstract

We develop and apply combinatorial algorithms for investigation of the feasible distance distributions of binary orthogonal arrays with respect to a point of the ambient binary Hamming space utilizing constraints imposed from the relations between the distance distributions of connected arrays. This turns out to be strong enough and we prove the nonexistence of binary orthogonal arrays of parameters (length, cardinality, strength) = $(9, 6.2^4 = 96, 4)$, $(10, 6.2^5, 5)$, $(10, 7.2^4 = 112, 4)$, $(11, 7.2^5, 5)$, $(11, 7.2^4, 4)$ and $(12, 7.2^5, 5)$, resolving the first cases where the existence was undecided so far. For the existing arrays our approach allows substantial reduction of the number of feasible distance distributions which could be helpful for classification results (uniqueness, for example).

Keywords.Binary Hamming space orthogonal arrays Krawtchouk polynomials distance distributions nonexistence Subclass.MSC 05B15 94B25 94B65

Subclass. MSC 05D15 94D25 94D05

^{*}The research of the first author was supported, in part, by a Bulgarian NSF contract I01/0003. The research of the second and third authors was supported, in part, by the Science Foundation of Sofia University under contract 144/2015 and its continuation "Discrete, Algebraic and Combinatorial Structures" in 2016.

1 Introduction

Orthogonal arrays have many connections to other combinatorial designs and have applications in coding theory, the statistical design of experiments, cryptography, various types of software testing and quality control. We refer to the book [4] as excellent exposition of the theory and practice of orthogonal arrays. In fact, there are enormous material about orthogonal arrays in internet.

An orthogonal array (OA) of strength τ and index λ in H(n, 2) (or binary orthogonal array, BOA), consists of the rows of an $M \times n$ matrix C with the property that every $M \times \tau$ submatrix of C contains all ordered τ -tuples of $H(\tau, 2)$, each one exactly $\lambda = M/2^{\tau}$ times as rows.

Let $C \subset H(n,2)$ be an (n, M, τ) BOA. The distance distribution of C with respect to $c \in H(n,2)$ if the (n + 1)-tuple

$$w = w(c) = (w_0(c), w_1(c), \dots, w_n(c)),$$

where $w_i(c) = |\{x \in C | d(x,c) = i\}|, i = 0, ..., n$. All feasible distance distributions of BOA of parameters (n, M, τ) can be computed effectively for relatively small n and τ as shown in [1]. Indeed, every distance distribution of C satisfies the system

$$\sum_{i=0}^{n} w_i(c) \left(1 - \frac{2i}{n} \right)^k = b_k |C|, \quad k = 0, 1, \dots, \tau,$$
(1)

where $b_k = \frac{1}{2^n} \sum_{d=0}^n {n \choose d} \left(1 - \frac{2d}{n}\right)^k$ and, in particular, $b_k = 0$ for k odd. The number b_k is in fact the first coefficient in the expansion of the poly-

The number b_k is in fact the first coefficient in the expansion of the polynomial t^k in terms of (binary) Krawtchouk polynomials. The Krawtchouk polynomials are zonal spherical functions for H(n, 2) (see [3, 6, 7]) and can be the defined by the three-term recurrence relation

$$(n-k)Q_{k+1}^{(n)}(t) = ntQ_k^{(n)}(t) - kQ_{k-1}^{(n)}(t)$$
 for $1 \le k \le n-1$,

with initial conditions $Q_0^{(n)}(t) = 1$ and $Q_1^{(n)}(t) = t$.

Let n, M and $\tau \leq n$ be fixed. We denote by $P(n, M, \tau)$ the set of all possible distance distributions of a (n, M, τ) BOA with respect to internal point c (in the beginning – all admissible solutions of the system (1) with $w_0(c) \geq 1$) and by $Q(n, M, \tau)$ the set of all possible distance distributions of a (n, M, τ) BOA with respect to external point (in the beginning – all admissible solutions of the system (1) with $w_0(c) = 0$). Denote also $W(n, M, \tau) = P(n, M, \tau) \cup Q(n, M, \tau)$. In this paper we describe an algorithm which works on the sets $P(n, M, \tau)$, $Q(n, M, \tau)$ and $W(n, M, \tau)$ utilizing connections between related BOAs. During the implementation of our algorithm these sets are changed¹ by ruling out some distance distributions.

In Section 2 we prove several assertions which connect the distance distributions of arrays under consideration and their relatives. This imposes significant constraints on the targeted BOAs and therefore allows us to collect rules for removing distance distributions from the sets $P(n, M, \tau)$, $Q(n, M, \tau)$ and $W(n, M, \tau)$. The logic of our algorithm is described in Section 3. The new nonexistence results are described in Section 4.

Algorithms for dealing with distance distributions were proposed earlier in [1] and [2] but in these papers the set $P(n, M, \tau)$ was only examined. Moreover, two seemingly crucial observations (Theorem 1 together with Corollary 2 and Theorem 13 together with Corollary 13) are new. Also, all complete versions (for the set $W(n, M, \tau)$) of the remaining assertions from the next section are new.

2 Relations between distance distributions of (n, M, τ) BOA and its derived BOAs

We start with a simple observation.

Theorem 1. If the distance distribution $w = (w_0, w_1, \ldots, w_n)$ belongs to the set $W(n, M, \tau)$, then the distance distribution $\overline{w} = (w_n, w_{n-1}, \ldots, w_0)$ also belongs to $W(n, M, \tau)$.

Proof. Let $C \,\subset\, H(n,2)$ be a BOA of parameters (n,M,τ) and \overline{C} is the array which is obtained from C by the permutation $(0 \to 1, 1 \to 0)$ in the whole C. Since the distances inside C are preserved by this transformation, \overline{C} is again (n, M, τ) BOA. On the other hand, distance i from external for C point to a point of C correspond to distance n - i to the transformed point of \overline{C} . This means that if $w = (w_0, w_1, \ldots, w_n)$ is the distance distribution of C with respect to some point $c \in H(n, 2)$ (internal or external for C), then the distance distribution of \overline{C} with respect to the same point (which can become either internal or external for \overline{C} , depending on whether $w_n > 0$ or $w_n = 0$) is $\overline{w} = (w_n, w_{n-1}, \ldots, w_0)$.

Corollary 2. The distance distribution $w = (w_0, w_1, \ldots, w_n) \in W(n, M, \tau)$ is ruled out if $\overline{w} = (w_n, w_{n-1}, \ldots, w_0) \notin W(n, M, \tau)$.

¹However, we prefer to keep the initial notation in order to avoid tedious notation.

Corollary 2 is important in all stages of our algorithm since it requires the non-symmetric distance distributions to be paired off and infeasibility of one element of the pair immediately implies the infeasibility for the other.

We proceed with analyzing relations between the BOA C and BOAs C' of parameters $(n - 1, M, \tau)$ which are obtained from C by deletion of one of its columns. Of course, the set $W(n - 1, M, \tau)$ of possible distance distributions of C' is sieved by Corollary 2 as well.

It is convenient to fix the removing of the first column of C. Let the distance distribution of C with respect to $c = \mathbf{0} = (0, 0, \dots, 0) \in H(n, 2)$ be $w = (w_0, w_1, \dots, w_n) \in W(n, M, \tau)$ and the distance distribution of C with respect to $c' = (0, 0, \dots, 0) \in H(n-1, 2)$ be $w' = (w'_0, w'_1, \dots, w'_{n-1}) \in W(n-1, M, \tau)$.

For every $i \in \{0, 1, ..., n\}$ the matrix which consists of the rows of C of weight i is called *i*-block. It follows from the above notations that the cardinality of the *i*-block is w_i . Next we denote by x_i (y_i , respectively) the number of the ones (zeros, respectively) in the intersection of the first column of C and the rows of the *i*-block.

Theorem 3. The numbers x_i and y_i , i = 0, 1, ..., n, satisfy the following system of linear equations

$$x_{i} + y_{i} = w_{i}, \ i = 1, 2, \dots, n-1$$

$$x_{i+1} + y_{i} = w'_{i}, \ i = 0, 1, \dots, n-1$$

$$y_{0} = w_{0}$$

$$x_{n} = w_{n}$$

$$x_{i}, y_{i} \in \mathbb{Z}, \ x_{i} > 0, \ y_{i} > 0, \ i = 0, 1, \dots, n$$
(2)

Proof. The equalities $x_i + y_i = w_i$, i = 1, ..., n - 1, $x_n = w_n$ and $y_0 = w_0$ follow directly from the definition of the numbers x_i and y_i . The relations $x_{i+1} + y_i = w'_i$, i = 0, 1, ..., n - 1, connecting w and w', follow from the fact that the rows of C', which are at distance i from c', are obtained in two ways: from the y_i rows of C at distance i from c and first coordinate 0, and from the x_{i+1} rows of C at distance i + 1 from c and first coordinate 1.

Corollary 4. The distance distribution $w = (w_0, w_1, \ldots, w_n) \in W(n, M, \tau)$ is ruled out if no system (2) obtained when w' runs $W(n - 1, M, \tau)$ has a solution.

Remark 5. Theorem 3 was firstly proved and used in 2013 by Boyvalenkov-Kulina [1] for $w \in P(n, M, \tau)$. Corollary 4 rules out some distance distributions w but it mainly serves to produce feasible pairs (w, w') which will be investigated further.

Our next step is based on the following property of BOAs: if we take the rows of C with first coordinate 0 (1, respectively) and remove that first coordinate then we obtain a BOA C_0 (C_1 , respectively) of parameters $(n-1, M/2, \tau - 1)$ (see Figure 1). At this stage the BOAs C_0 and C_1 have the same sets of admissible distance distributions – all these which have passed the sieves of Corollaries 2 and 4 for the set $W(n-1, M/2, \tau - 1)$.

Figure 1. $w' = (w'_0, w'_1, \dots, w'_{n-1})$ $C' = (n - 1, M, \tau)$ $0 \qquad y = (y_0, y_1, \dots, y_{n-1})$ $C_0 = (n - 1, M/2, \tau - 1)$ $C_1 = (n - 1, M/2, \tau - 1)$ $C_1 = (n - 1, M/2, \tau - 1)$ $W = (w_0, w_1, \dots, w_n)$ $C = (n, M, \tau)$

We continue with relations between the BOAs C, C', C_0 and C_1 using the numbers x_i and y_i , i = 0, 1, ..., n.

Theorem 6. The distance distribution of the $(n-1, M/2, \tau-1)$ BOA C_0 with respect to c' is $y = (y_0, y_1, \ldots, y_{n-1})$, i.e. $y \in W(n-1, M/2, \tau-1)$.

Proof. The number y_i is equal to the number of the points of C_0 at distance i from the point c'.

Remark 7. We have two possibilities in Theorem 6 – if $y_0 \ge 1$, then $c' \in C_0$ and therefore $y \in P(n-1, M/2, \tau-1)$ (this is Theorem 1a) in [2]), or $y_0 = 0$ when $c' \notin C_0$ and therefore $y \in Q(n-1, M/2, \tau-1)$.

Corollary 8. The pair (w, w') is ruled out if $y \notin W(n-1, M/2, \tau-1)$ or if $\overline{y} = (y_{n-1}, y_{n-2}, \ldots, y_0) \notin W(n-1, M/2, \tau-1)$.

Proof. Follows from Theorem 6 and Corollary 2 for $\overline{C_0}$.

Theorem 9. The distance distribution of the $(n-1, M/2, \tau - 1)$ BOA C_1 with respect to c' is $x = (x_1, x_2, \ldots, x_n)$, i.e. $x \in W(n-1, M/2, \tau - 1)$.

Proof. The number x_i is equal to the number of the points of C_1 at distance i-1 from the point c'.

Remark 10. Similarly to above, we have two possibilities in Theorem 9 – if $x_1 \ge 1$, then $c' \in C_1$ and therefore $x \in P(n-1, M/2, \tau-1)$ (this is Theorem 2a) in [2]), or $x_1 = 0$ when $c' \notin C_1$ and therefore $x \in Q(n-1, M/2, \tau-1)$.

Corollary 11. The pair (w, w') is ruled out if $x \notin W(n-1, M/2, \tau-1)$ or if $\overline{x} = (x_n, x_{n-1}, \ldots, x_1) \notin W(n-1, M/2, \tau-1)$.

Proof. Follows from Theorem 9 and Corollary 2 for $\overline{C_1}$.

In our next step we consider the effect of the permutation $(0 \rightarrow 1, 1 \rightarrow 0)$ in the first column of C. This transformation does not change the distances from C and thus we obtain a BOA $C^{1,0}$ of parameters (n, M, τ) again.

Theorem 12. If the distance distribution of C with respect to $c = \mathbf{0} \in H(n,2)$ is $w = (w_0, w_1, \ldots, w_{n-1}, w_n) = (y_0, x_1 + y_1, \ldots, x_{n-1} + y_{n-1}, x_n)$, then the distance distribution of $C^{1,0}$ with respect to c is $\widehat{w} = (x_1, x_2 + y_0, \ldots, x_n + y_{n-2}, y_{n-1})$, i.e. $\widehat{w} \in W(n, M, \tau)$.

Proof. There are x_i points in $C^{1,0}$ (coming from C_1) at distance i-1 from c. Analogously, there are y_i points in $C^{1,0}$ (coming from C_0) at distance i+1 from c. This means that the number of the points of $C^{1,0}$ at distance 0 from c is x_1 , the number of the points of $C^{1,0}$ at distance $i, 1 \leq i \leq n-1$, from c is $y_{i-1} + x_{i+1}$, and, finally, the number of the points of $C^{1,0}$ at distance n from c is y_{n-1} . Therefore the distance distribution of $C^{1,0}$ with respect to c is $\hat{w} = (x_1, x_2 + y_0, \dots, x_n + y_{n-2}, y_{n-1})$.

Corollary 13. The pair (w, w') is ruled out if $\widehat{w} \notin W(n, M, \tau)$ or if $\overline{\widehat{w}} \notin W(n, M, \tau)$.

Corollary 14. The distance distribution w is ruled out if all possible pairs $(w, w'), w' \in W(n-1, M, \tau)$, are ruled out.

Otherwise, we proceed with the remaining pairs as follows. Let

$$(x_0^{(j)} = 0, x_1^{(j)}, \dots, x_n^{(j)}; y_0^{(j)}, y_1^{(j)}, \dots, y_{n-1}^{(j)}, y_n^{(j)} = 0), \quad j = 1, \dots, s,$$
(3)

are all solutions of Theorem 3 when w' runs $W(n-1, M, \tau)$ which have passed the sieves of Corollaries 8, 11 and 13. We now free the cutting and thus consider all possible n cuts of columns of C. These cuts produce pairs (w, w') (where w is fixed) and corresponding solutions (3). Let the solutions (3) appear with multiplicities k_1, k_2, \ldots, k_s , respectively.

Theorem 15. The nonnegative integers k_1, k_2, \ldots, k_s satisfy the equations

$$k_{1} + k_{2} + \cdots + k_{s} = n$$

$$k_{1}x_{1}^{(1)} + k_{2}x_{1}^{(2)} + \cdots + k_{s}x_{1}^{(s)} = w_{1}$$

$$k_{1}x_{2}^{(1)} + k_{2}x_{2}^{(2)} + \cdots + k_{s}x_{2}^{(s)} = 2w_{2} . \qquad (4)$$

$$\vdots$$

$$k_{1}x_{n}^{(1)} + k_{2}x_{n}^{(2)} + \cdots + k_{s}x_{n}^{(s)} = nw_{n}$$

Proof. This follows for counting in two ways the number of the ones in the *i*-block of *C*. For fixed $i \in \{1, 2, ..., n\}$, this number is obviously iw_i , and, on the other hand, it is equal to the sum $k_1 x_i^{(1)} + k_2 x_i^{(2)} + \cdots + k_s x_i^{(s)}$. \Box

Corollary 16. The distance distribution w is ruled out if the system (4) does not have solutions.

Corollary 17. Let $j \in \{1, 2, ..., s\}$ be such that all solutions of the system (4) have $k_j = 0$. Then the pair (w, w'), which corresponds to j, is ruled out.

3 Our algorithm

We organize the results from the previous section to work together as follows.

All BOAs (in fact, their current sets of feasible distance distributions P, Q and W) of interest for the targeted BOA $C = (n, M, \tau)$ are collected in a table starting with first row

$$(\tau, M, \tau) (\tau + 1, M, \tau) (\tau + 2, M, \tau) \dots C = (n, M, \tau).$$

The next row consist of the derived BOAs

$$(\tau - 1, M/2, \tau - 1) (\tau, M/2, \tau - 1) (\tau + 1, M/2, \tau - 1) \dots (n - 1, M/2, \tau - 1)$$

and so on until it makes sense. We apply Corollaries 4, 14 and 16 in every row separately from left to right to reduce the sets P, Q and W. Of course, this process is fueled with information from the columns (starting from the bottom end) according to Corollaries 8, 11, 13 and 17. Every nonsymmetric distance distribution w which is ruled out, forces its mirror image \overline{w} to be ruled out according to Corollary 2. The algorithm stops when no new rulings out are possible. An entry at the right end, showing that some of the sets P, Q and W is empty², means nonexistence of the corresponding BOA. Otherwise, we collect the reduced sets for further analysis and classification results (in some cases, possibly, uniqueness).

Here is the pseudocode of the module of our algorithm which deals with the sets $W(n, M, \tau)$, $W(n - 1, M, \tau)$ and $W(n - 1, M/2, \tau - 1)$.

Algorithm 1

1:	procedure NDDA($W(n, M, \tau)$, $W(n-1, M, \tau)$, $W(n-1, M/2, \tau-1)$)
2:	filteredW = empty set
3:	for $w \in W(n, M, \tau)$ do
4:	all X = empty set
5:	for $w' \in W(n-1, M, \tau)$ do
6:	x, y = solve system (2) for integer nonnegative solutions
7:	if $x, \bar{x} \in W(n-1, M/2, \tau-1)$ and $y, \bar{y} \in W(n-1, M/2, \tau-1)$
	and $\widehat{w}, \overline{\widehat{w}} \in W(n, M, \tau)$ and $\widehat{w}, \overline{\widehat{w}} \notin filteredW$ then
8:	add x to $all X$
9:	if all is empty then
10:	add w to $filteredW$
11:	else
12:	if system (4) has no integer nonnegative solutions then
13:	add w to $filteredW$
14:	if <i>filteredW</i> is nonempty then
15:	return NDDA $(W(n, M, \tau) \setminus filteredW, W(n-1, M, \tau), W(n-1, M, \tau))$
	$1, M/2, \tau - 1))$
16:	else
17:	${f return} \ W(n,M, au)$

We believe that the above description is enough for smooth reproduction of our algorithm. Anyway, we are ready to supply the interested reader with all our programs and databases [9].

 $^{^2\}mathrm{In}$ fact, in all cases where we arrived at an empty set, the other two also became empty at the same step.

4 New nonexistence results

4.1 Nonexistence of (9,96,4) BOA and consequences

We apply the algorithm from the previous section on the table below targeting the (9, 96, 4) BOA.

(4, 96, 4)	(5, 96, 4)		(9, 96, 4)
(3, 48, 3)	(4, 48, 3)	•••	(8, 48, 3)
(2, 24, 2)	(3, 24, 2)	•••	(7, 24, 2)

The frame of the implementation is showed in the next table. In every entry we first show the number of distance distributions in the beginning and then (after the arrow) the number of the remaining distance distributions in the end of the implementation. The numbers in the brackets show how many distance distributions were left possible after [2].

P(n, 96, 4)	$1 \rightarrow 1$	$6 \rightarrow 6$	$12 \rightarrow 12$	$20(10) \rightarrow 10$	$34(12) \rightarrow 9$	$37(10) \rightarrow 0$
Q(n, 96, 4)	$0 \rightarrow 0$	$1 \rightarrow 1$	$4 \rightarrow 4$	$12 \rightarrow 6$	$41 \rightarrow 11$	$97 \rightarrow 0$
W(n, 96, 4)	$1 \rightarrow 1$	$7 \rightarrow 7$	$16 \rightarrow 16$	$32 \rightarrow 16$	$75 \rightarrow 20$	$134 \rightarrow 0$
P(n, 48, 3)	$1 \rightarrow 1$	$6 \rightarrow 6$	$13 \rightarrow 13$	$31(25) \rightarrow 25$	$53(41) \rightarrow 38$	$96(65) \rightarrow 62$
Q(n, 48, 3)	$0 \rightarrow 0$	$1 \rightarrow 1$	$4 \rightarrow 4$	$13 \rightarrow 9$	$41 \rightarrow 30$	$110 \rightarrow 85$
W(n, 48, 3)	$1 \rightarrow 1$	$7 \rightarrow 7$	$17 \rightarrow 17$	$44 \rightarrow 34$	$94 \rightarrow 68$	$206 \to 147$
P(n, 24, 2)	$1 \rightarrow 1$	$6 \rightarrow 6$	$13 \rightarrow 13$	$30 \rightarrow 28$	$49 \rightarrow 47$	$74 \rightarrow 69$
Q(n, 24, 2)	$0 \rightarrow 0$	$1 \rightarrow 1$	$5 \rightarrow 5$	$19 \rightarrow 17$	$54 \rightarrow 52$	$130 \rightarrow 125$
W(n, 24, 2)	$1 \rightarrow 1$	$7 \rightarrow 7$	$18 \rightarrow 18$	$49 \rightarrow 45$	$103 \rightarrow 99$	$204 \rightarrow 194$

Theorem 18. There exist no binary orthogonal arrays of parameters (9, 96, 4).

Proof. The zero entries in the right upper cells of the last table imply that there exists no binary orthogonal array of parameters (9, 96, 4). \Box

The implementation of the algorithm for Theorem 18 created database which is available at [9]. Note that intermediate results are also included.

In 1966 Seiden and Zemach [8] (see also [4, Theorem 2.24]) proved that BOAs of parameters $(n, N, \tau = 2k)$ and $(n + 1, 2N, \tau + 1 = 2k + 1)$ coexist. Therefore we have the following nonexistence result as well.

Corollary 19. There exist no binary orthogonal arrays of parameters (10, 192, 5).

The last Corollary follows also from the implementation of our algorithm with (10, 192, 5) as target. This is illustrated in the next table.

P(n, 192, 5)	$1 \rightarrow 1$	$6 \rightarrow 6$	$12 \rightarrow 12$	$21(8) \rightarrow 8$	$35(6) \rightarrow 4$	$35(4) \rightarrow 0$
Q(n, 192, 5)	$0 \rightarrow 0$	$1 \rightarrow 1$	$4 \rightarrow 4$	$12 \rightarrow 4$	$32 \rightarrow 4$	$85 \rightarrow 0$
W(n, 192, 5)	$1 \rightarrow 1$	$7 \rightarrow 7$	$16 \rightarrow 16$	$33 \rightarrow 12$	$67 \rightarrow 8$	$120 \rightarrow 0$

The nonexistence results of Theorem 18 and Corollary 19 give improvements in two entries of Table 12.1 from [4]. We have $7 \le L(n,\tau) \le 8$ instead of $6 \le L(n,\tau) \le 8$ for the pairs $(n,\tau) = (9,4)$ and (10,5).

4.2 Nonexistence of (10, 112, 4) BOA and consequences

Here we work in the table with C = (11, 112, 4) as target.

(4, 112, 4)	(5, 112, 4)	• • •	(11, 112, 4)
(3,56,3)	(4, 56, 3)	•••	(10, 56, 3)
(2, 28, 2)	(3,28,2)	•••	(9, 28, 2)

The results are shown below. Again, in every entry we first show the number of distance distributions in the beginning and then (after the arrow) the number of the remaining distance distributions in the end of the implementation. The numbers in the brackets show how many distance distributions were left possible after the implementation of the algorithm from [2].

P(n, 112, 4)	$1 \rightarrow 1$	$7 \rightarrow 7$	$15(13) \rightarrow 13$	$31(20) \rightarrow 12$	$58(25) \rightarrow 16$
Q(n, 112, 4)	$0 \rightarrow 0$	$1 \rightarrow 1$	$5 \rightarrow 3$	$17 \rightarrow 6$	$59 \rightarrow 18$
W(n, 112, 4)	$1 \rightarrow 1$	$8 \rightarrow 8$	$20 \rightarrow 16$	$48 \rightarrow 18$	$117 \rightarrow 34$
P(n, 56, 3)	$1 \rightarrow 1$	$7 \rightarrow 7$	$17(16) \rightarrow 16$	$49(40) \rightarrow 40$	$95(71) \rightarrow 68$
Q(n,56,3)	$0 \rightarrow 0$	$1 \rightarrow 1$	$4 \rightarrow 3$	$15 \rightarrow 14$	$59 \rightarrow 44$
W(n, 56, 3)	$1 \rightarrow 1$	$8 \rightarrow 8$	$21 \rightarrow 19$	$64 \rightarrow 54$	$154 \rightarrow 112$
P(n, 28, 2)	$1 \rightarrow 1$	$7 \rightarrow 7$	$17 \rightarrow 17$	$46(43) \rightarrow 43$	$87(82) \rightarrow 82$
Q(n, 28, 2)	$0 \rightarrow 0$	$1 \rightarrow 1$	$5 \rightarrow 4$	$23 \rightarrow 22$	$79 \rightarrow 76$
W(n, 28, 2)	$1 \rightarrow 1$	$8 \rightarrow 8$	$22 \rightarrow 21$	$69 \rightarrow 65$	$166 \rightarrow 158$
		72(28) -	→ 9 88(31)	$1) \rightarrow 0$	
		$158 \rightarrow 2$	24 373	$\rightarrow 0$	
		$230 \rightarrow 3$	33 461	$\rightarrow 0$	
	19	99(137) -	→ 135 311(205	$5) \rightarrow 193$	
		$181 \rightarrow 12$	29 451 -	$\rightarrow 313$	
		$380 \rightarrow 20$	64 762 -	$\rightarrow 506$	
	14	15(137) - 100	→ 137 208(196	$(5) \rightarrow 196$	
		$205 \rightarrow 19$		$\rightarrow 450$	
		$350 \rightarrow 33$	32 677 -	$\rightarrow 646$	

The zero entries in the upper right corner imply the nonexistence of BOAs of parameters (10, 112, 4).

Theorem 20. There exist no binary orthogonal arrays of parameters (10, 112, 4) and (11, 112, 4).

Proof. The zero entry in the right upper cell of the last table means that there exists no binary orthogonal array of parameters (10, 112, 4). This immediately implies the nonexistence of BOAs of parameters (11, 224, 4).

The data from the implementation of the algorithm for Theorem 20 is available at [9] with intermediate results included.

As above, we use the coexistence of (n, N, 2k) and (n + 1, 2N, 2k + 1)BOAs to obtain further nonexistence results.

Corollary 21. There exist no binary orthogonal arrays of parameters (11, 224, 5) and (12, 224, 5).

The last Corollary follows also from the implementation of our algorithm with (11, 224, 5) as target. This is illustrated in the next table, where the first two columns are missed.

P(n, 224, 5) $15(11) \rightarrow 11$ $32(19) \rightarrow 4$ $63(15) \rightarrow 5$ $74(11) \rightarrow 2$ $108(6) \to 0$ $16 \rightarrow 2$ $141 \rightarrow 4$ Q(n, 224, 5) $4 \rightarrow 2$ $47 \rightarrow 4$ $337 \rightarrow 0$ $19 \rightarrow 13$ $48 \rightarrow 6$ $110 \rightarrow 9$ $215 \to 6$ $445 \rightarrow 0$ W(n, 224, 5)

The nonexistence results of Theorem 20 and Corollary 21 give improvements in four entries of Table 12.1 from [4]. We have $L(n, \tau) = 8$ instead of $7 \le L(n, \tau) \le 8$ for the pairs $(n, \tau) = (10, 4)$, (11, 4), (11, 5) and (12, 5).

4.3 Other nonexistence results

Our algorithm gives other nonexistence results which however are superseded by the result of Khalyavin [5] from 2010. We list these in the next assertion.

Theorem 22. ([5] and our algorithm) There exist no binary orthogonal arrays of parameters $(10, 7.2^6 = 448, 6)$, $(11, 7.2^7 = 896, 7)$, $(12, 10.2^8 = 2560, 8)$, $(13, 10.2^9 = 5120, 9)$, $(12, 11.2^8 = 2816, 8)$, $(13, 11.2^9 = 5632, 9)$ and $(15, 13.2^{10} = 13312, 10)$.

5 Updated table for $L(n, \tau)$

We present an updated version of the situation with the possible values of function $L(n,\tau)$ – the minimum possible index λ of an $(n, M = \lambda 2^{\tau}, \tau)$ binary orthogonal array. Our table covers the range $4 \leq n \leq 16$ and $4 \leq \tau \leq 10$ (see Table 1).

All calculations in this paper were performed by programs in Maple. All results (in particular all possible distance distributions in the beginning) can be seen at [9]. All programs are available upon request.

References

- P. Boyvalenkov, H. Kulina, Investigation of binary orthogonal arrays via their distance distributions, *Problems of Information Transmission*, 49(4), 320-330 (2013).
- [2] P. Boyvalenkov, H. Kulina, T. Marinova, M. Stoyanova, Nonexistence of binary orthogonal arrays via their distance distributions, *Problems* of Information Transmission, 51(4), 326-334 (2015).
- [3] P. Delsarte, V. I. Levenshtein, Association schemes and coding theory, *IEEE Trans. Inform. Theory*, 44, 2477-2504 (1998).
- [4] A. Hedayat, N. J. A. Sloane, J. Stufken, Orthogonal Arrays: Theory and Applications, Springer-Verlag, New York, (1999).
- [5] A. V. Khalyavin, Estimates of the capacity of orthogonal arrays of large strength, *Moscow Univ. Math. Bull.*, 65, 130-131 (2010).
- [6] V. I. Levenshtein, Krawtchouk polynomials and universal bounds for codes and designs in Hamming spaces, *IEEE Trans. Infor. Theory*, 41, 1303-1321 (1995).
- [7] V. I. Levenshtein, Universal bounds for codes and designs, Handbook of Coding Theory, V. S. Pless and W. C. Huffman, Eds., Elsevier, Amsterdam, Ch. 6, 499-648 (1998).
- [8] E. Seiden, R. Zemach, On orthogonal arrays, Ann. Math. Statist., 37, 1355-1370 (1996).
- [9] https://drive.google.com/folderview?id=0B0r6JtxqB_RmLXRVQTBnMldmcGM&usp =sharing

n / $ au$	4	5	6	7	8	9	10
4	1						
5	1	1					
6	2	1	1				
7	sz4	2	1	1			
8	4^c	sz4	2	1	1		
9	^{bms} 7-8	4^c	4	2	1	1	
10	bms8	bms7-8	$^{kh}8$	4	2	1	1
11	bms8	bms8	8^c	$^{kh}8$	4	2	1
12	$^{bkms}8$	bms8	12-16	8^c	$^{kh}8$	4	2
13	8	$^{bkms}8$	16	12-16	$^{kh}16$	$^{kh}8$	4
14	8	8	16	16	16^c	$^{kh}16$	$^{kh}8$
15	8^{nr}	8	16^{rh}	16	26-32	16^c	$^{kh}16$
16	10-16	8^{nr}	21-32	16^{rh}	39-64	26-32	$^{kh}32$

Table 1: (see Table 12.1 in [4]) Minimum possible index λ of binary orthogonal array of length $n, 4 \leq n \leq 16$, and strength $\tau, 4 \leq \tau \leq 10$.

Key:

bkms Boyvalenkov, Kulina, Marinova, Stoyanova in [2]

c Cyclic code

kh Khalyavin (2010) (and see also Section 6 and [2, Section 4.3] and)

nr Nordstrom-Robinson (1967) code

rh Rao-Hamming construction

sz Seiden and Zemach (1966) bound

bms Theorem 18 and Corollary 19, or Theorem 20 and Corollary 21