

Global-Local Mixtures

Anindya Bhadra ^{*}
Purdue University

Jyotishka Datta [†]
University of Arkansas

Nicholas G. Polson [‡] and Brandon Willard [§]
The University of Chicago, Booth School of Business

January 19, 2022

Abstract

Global-local mixtures are derived from the Cauchy-Schlömilch and Liouville integral transformation identities. We characterize well-known normal-scale mixture distributions including the Laplace or lasso, logit and quantile as well as new global-local mixtures. We also apply our methodology to convolutions that commonly arise in Bayesian inference. Finally, we conclude with a conjecture concerning bridge and uniform correlation mixtures.

Keywords: Bayes regularization; Cauchy; Convolution; Global-local mixture; Lasso; Logistic; Quantile; Stable law.

1 Introduction

Many statistical problems involve regularization penalties derived from global-local mixture distributions (Polson & Scott, 2011; Hans, 2011; Bhadra et al., 2016a). A global-local mixture density, denoted by $p(x_1, \dots, x_p)$, takes the form

$$p(x_1, \dots, x_p) = \int_0^\infty \prod_{i=1}^p p(x_i | \tau) p(\tau) d\tau,$$

where $p(x_i | \tau) = \int_0^\infty p(x_i | \lambda_i, \tau) p(\lambda_i | \tau) d\lambda_i$ is a local mixture and $p(x_1, \dots, x_p)$ is a global mixture over $\tau \sim p(\tau)$. There is great interest in analytically calculating $p(x_i | \tau)$, and the associated regularization penalty $\phi(x_i, \tau) = -\log p(x_i | \tau)$. Convolution mixtures of the form $p(x_i | \tau) = \int p(x_i - \lambda_i) p(\lambda_i) d\lambda_i$ are also of interest. We show

^{*}email: bhadra@purdue.edu

[†]email: jd298@stat.duke.edu

[‡]email: ngp@chicagobooth.edu

[§]email: brandonwillard@gmail.com

how the Cauchy-Schlömilch and Liouville transformations can be used to derive closed-form global-local mixtures. We start by stating two key integral identities: the Cauchy-Schlömilch transformation

$$\int_0^\infty f \left\{ (ax - bx^{-1})^2 \right\} dx = \frac{1}{2a} \int_0^\infty f(y^2) dy, \quad (1)$$

and the Liouville transformation

$$\int_0^\infty f \left(ax + \frac{b}{x} \right) x^{-1/2} dx = a^{-1/2} \int_0^\infty f \left\{ 2(ab)^{1/2} + y \right\} y^{-1/2} dy, \quad a, b > 0. \quad (2)$$

See [Boros et al. \(2006\)](#), [Baker \(2008\)](#) and [Jones \(2014\)](#) for further discussion. Identity (1) follows from the simple transformation $t = b/(ax)$ as

$$I = \int_0^\infty f \left\{ (ax - b/x)^2 \right\} dx = \int_0^\infty f \left\{ (at - b/t)^2 \right\} \frac{b}{at^2} dt.$$

Adding the two terms in the last equality yields $2I = \int_0^\infty f \left\{ (at - b/t)^2 \right\} \{1 + b/(at^2)\} dt$ and transforming $y = b/t - at$ gives $dy = -a\{1 + b/(at^2)\} dt$, yielding $I = (2a)^{-1} \int_0^\infty f(y^2) dy$, as required. A useful generalization of the Cauchy-Schlömilch transformation is

$$\int_0^\infty f \left[\{x - s(x)\}^2 \right] dx = \int_0^\infty f(y^2) dy, \quad (3)$$

where $s(x) = s^{-1}(x)$ is a self-inverse function such as $s(x) = b/x$ or $s(x) = -a^{-1} \log\{1 - \exp(ax)\}$. The proof for the Liouville transformation identity follows in a similar manner, and is omitted for the sake of brevity. These identities can be used to construct new global-local mixture distributions. Let $f(x) = 2g\{t(x)\}$ and let $t(x)$ be of the form $x - s(x)$, where $s : \mathbb{R}^+ \rightarrow \mathbb{R}^+$ is a self-inverse, onto and monotone decreasing function. Together with the Cauchy-Schlömilch transformation, we have a rather surprising way to represent the resulting $g\{t(x)\}$ as a global-local scale mixture.

[Jones \(2014\)](#) shows that only a few choices of $t(x)$ leads to fully tractable formulae for its inverse $t^{-1} = \Pi$ and the integral $\Pi(y) = \int_{-\infty}^y \pi(\omega) d\omega$. Two special choices are the t -distribution with 2 degrees of freedom and the logistic.

$$\begin{aligned} \Pi_T(y) &= (1/2)\{y + (4b + y^2)^{1/2}\}, \Pi_T^{-1}(x) = t_T(x) = x - b/x, \quad b > 0, \\ \Pi_L(y) &= a^{-1} \log(1 + e^{ay}), \Pi_L^{-1}(x) = t_L(x) = a^{-1} \log(e^{ax} - 1), \quad a > 0. \end{aligned}$$

Now, the integral identity in (1) shows that if $f(x)$, $x \geq 0$ is a density function, so is $g(x) = 2af(|ax - b/x|)$, $x > 0$. The functions f and g are called mother and daughter density functions, respectively.

Apart from simplifying proofs involving global-local mixtures, the Cauchy-Schlömilch and Liouville transformations can generate new distributions via scale transformations. These transformations can take the form $f(x) = 2g\{t(x)\}$ for certain $f(x)$ under suitable conditions. For example, given a density $f(x)$ we can create a new global-local scale family, $f(ax - b/x)$, by effectively reallocating its probability mass. A particularly

useful tool for generating univariate and multivariate random variables is Khintchine's theorem. Khintchine's theorem states that any random variable X with a unimodal, univariate distribution and a mode at zero can be written as a product $X = ZU$, where $U \sim \mathcal{U}(0, 1)$ and Z has the density function $f_Z(z) = -zf'_X(z)$, $z \in \mathbb{R}$. [Bryson & Johnson \(1982\)](#), and subsequently [Jones \(2002\)](#), discuss how Khintchine's theorem allows one to construct both univariate and multivariate densities, even with special dependence structure. [Jones \(2014\)](#) develops an extended Khintchine's theorem that further allows one to generate random variables with unimodal densities of the form $2g\{t(x)\}$.

2 Global-local Scale Mixtures

2.1 Lasso as a normal scale mixture

The lasso penalty arises as a Laplace global-local mixture ([Andrews & Mallows, 1974](#)). A simple transformation proof follows using Cauchy-Schlömilch with $f(x) = e^{-x}$. Starting with the normal integral identity, $\int_0^\infty f(y^2)dy = \int_0^\infty e^{-y^2}dy = \pi^{1/2}/2$, we obtain

$$\int_0^\infty e^{-(ax)^2 - (b/x)^2} dx = \int_0^\infty \exp \left\{ -ab \left(\frac{a}{b}x^2 + \frac{b}{a}x^{-2} \right) \right\} dx = \frac{\pi^{1/2}}{2a} e^{-2ab} \quad a, b \in \mathbb{R}.$$

Substituting $t = (a/b)^{1/2}x$ and $c = ab$ yields the Laplace or Lasso penalty as

$$\int_0^\infty e^{-c(t-t^{-1})^2} dt = \frac{1}{2}(\pi/c)^{1/2} \Rightarrow \int_0^\infty e^{-c(t^2+t^{-2})} dt = \frac{1}{2}(\pi/c)^{1/2} e^{-2c}.$$

The Laplace density can be viewed as a transformed normal, via $y = t - t^{-1}$.

Proposition 1. *The usual identity for the lasso also follows from [Lévy \(1940\)](#) as*

$$\int_0^\infty \frac{a}{(2\pi)^{1/2}t^{3/2}} e^{-a^2/(2t)} e^{-\lambda t} dt = e^{-a(2\lambda)^{1/2}}. \quad (4)$$

For $a = 1$, and $\theta = (2\lambda)^{1/2}$, this can be written as

$$E[\exp\{-\theta^2/(2G)\}] = \exp(-\theta), \quad G \sim \mathcal{G}(1/2, 1/2) \quad (5)$$

Proof. First substitute $t^{-1} = x^2$, which makes the left hand side in (4) equal to

$$\int_0^\infty \frac{a}{(2\pi)^{1/2}t^{3/2}} e^{-a^2/(2t)} e^{-\lambda t} dt = \left(\frac{2}{\pi}\right)^{1/2} a e^{-a(2\lambda)^{1/2}} \int_0^\infty e^{-(2^{-1/2}ax - \lambda x^{-1})^2} dx = e^{-a(2\lambda)^{1/2}}.$$

The last step follows from Cauchy-Schlömilch formula. The second relationship (5) follows by fixing $a = 1$, $\theta = (2\lambda)^{1/2}$ and substituting $t = x^{-1}$

$$\int_0^\infty \frac{a}{(2\pi)^{1/2}t^{3/2}} e^{-a^2/(2t)} e^{-\lambda t} dt = \frac{1}{(2\pi)^{1/2}} \int_0^\infty e^{-\theta^2/(2x)} x^{-1/2} e^{-x/2} dx.$$

The left hand side can be identified as $E\left\{e^{-\theta^2/(2G)}\right\}$ for $G \sim \mathcal{G}(1/2, 1/2)$. \square

2.2 Logit and quantile as global-local mixtures

Logistic modeling can be viewed within the global-local mixture framework via the Pólya-Gamma distribution (Polson et al., 2013). This leads to efficient Markov chain Monte Carlo algorithms for inference.

Proposition 2. *The two key marginal distributions for the hyperbolic generalized inverse Gaussian (Barndorff-Nielsen et al., 1982) and Pólya-Gamma mixtures are*

$$\frac{\alpha^2 - \kappa^2}{2\alpha} e^{-\alpha|x-\mu| + \kappa(x-\mu)} = \int_0^\infty \phi(x | \mu + \kappa\lambda, \lambda) p_{\text{GIG}} \left\{ \lambda | 1, 0, (\alpha^2 - \kappa^2)^{1/2} \right\} d\lambda, \quad \alpha \geq \kappa \geq 0, \quad (6)$$

$$\frac{1}{B(\alpha, \kappa)} \frac{e^{\alpha(x-\mu)}}{(1 + e^{x-\mu})^{\alpha+\kappa}} = \int_0^\infty \phi(x | \mu + \kappa\lambda, \lambda) p_{\text{Polya}}(\lambda | \alpha, \kappa) d\lambda, \quad (7)$$

where $\phi(\mu + \kappa\lambda, \lambda)$ denotes the normal density function with mean $(\mu + \kappa\lambda)$ and variance λ . The functions p_{GIG} and p_{Polya} are the corresponding local mixture densities for the generalized inverse Gaussian and the Pólya-Gamma, respectively. The logit and quantile identities can be derived using Cauchy-Schlömilch identity.

Proof. Let $f(x) = e^{-x^2/2}$, $a = \alpha$ and $b = |x - \phi|$ in (1). Then,

$$(2/\pi)^{1/2} \int_0^\infty \exp \left\{ -\frac{1}{2} \left(ay - \frac{|x-\mu|}{y} \right)^2 \right\} dy = \frac{1}{\alpha} (2\pi)^{-1/2} \int_0^\infty e^{-\frac{1}{2}y^2} dy = \frac{1}{\alpha}.$$

Let $v = y^2$. Rearranging the constant terms yields

$$\frac{1}{\alpha} e^{-\alpha|x-\mu|} = \frac{1}{(2\pi v)^{1/2}} \int_0^\infty \exp \left[- \left\{ \frac{(x-\mu)^2}{2v} + \frac{\alpha^2}{2} v \right\} \right] dv.$$

Multiplying by $2^{-1}(\alpha^2 - \kappa^2)e^{\kappa(x-\mu)}$ and completing the square yields

$$\frac{\alpha^2 - \kappa^2}{2\alpha} \exp \{ -\alpha|x-\mu| + \kappa(x-\mu) \} = \int_0^\infty \phi(x | \mu + \kappa v, v) \frac{\alpha^2 - \kappa^2}{2} \exp \left(-\frac{\alpha^2 - \kappa^2}{2} v \right) dv.$$

The mixing distribution is exponential with rate parameter $(\alpha^2 - \kappa^2)/2$, a special case of the generalized inverse Gaussian distribution introduced by Etienne Halphen circa 1941 (Seshadri, 2004). The density with parameters $(\lambda, \delta, \gamma)$ has the form

$$p_{\text{GIG}}(x | \lambda, \delta, \gamma) = \frac{(\gamma/\delta)^\lambda}{2K_\lambda(\delta\gamma)} x^{\lambda-1} \exp \left\{ -\frac{1}{2}(\delta^2 x^{-1} + \gamma^2 x) \right\}, \quad x, \lambda, \delta > 0, \quad p \in \mathfrak{R},$$

where K_λ is the modified Bessel function of the second kind. The Liouville formula can be used to show that the above is a valid probability density function. When δ or γ is zero, the normalizing constant takes the limiting values given by $K_\lambda(u) \asymp \Gamma(|\lambda|) 2^{|\lambda|-1} u^{|\lambda|}$ for $\lambda > 0$. If $\delta = 0$, the generalized inverse Gaussian is identical to a gamma distribution:

$$p_{\text{GIG}}(x | \lambda, \delta = 0, \gamma) = \frac{\alpha^\lambda}{\Gamma(\lambda)} x^{\lambda-1} \exp(-\alpha x), \quad x > 0, \quad \alpha = \gamma^2/2.$$

We now present a simple proof for the Pólya-Gamma mixture in (7). First, write κ for $a - b/2$:

$$\frac{(e^\psi)^a}{(1 + e^\psi)^b} = 2^{-b} e^{\kappa\omega} \int_0^\infty e^{-\omega\psi^2/2} p(\omega) d\omega, \quad (8)$$

where $\omega \sim \text{PG}(b, 0)$, a Pólya-Gamma random variable with density

$$p(\omega | b, 0) = \frac{2^{b-1}}{\Gamma(b)} \sum_{n=0}^\infty (-1)^n \frac{\Gamma(n+b)}{\Gamma(n+1)} \frac{2n+b}{(2\pi)^{1/2} \omega^{3/2}} \exp \left\{ -\frac{(2n+b)^2}{8\omega} \right\}.$$

The logit function corresponds to $a = 0, b = 1$ in (8). Cauchy-Schlömilch identity yields

$$\frac{1}{1 + e^\psi} = \frac{1}{2} e^{-\psi/2} \int_0^\infty e^{-(\psi^2\omega)/2} p(\omega) d\omega, \quad p(\omega) = \sum_{n=0}^\infty (-1)^n \frac{2n+1}{(2\pi\omega^3)^{1/2}} e^{-(2n+1)^2/(8\omega)}. \quad (9)$$

To show (9), write the right-hand side interchanging the integral and summation:

$$I = \frac{1}{2} e^{-\psi/2} \sum_{n=0}^\infty (-1)^n \frac{2n+1}{(2\pi)^{1/2}} \int_0^\infty \exp \left[-\left\{ \frac{\psi^2}{2} \omega + \frac{(2n+1)^2}{8\omega} \right\} \right] \frac{1}{\omega^{3/2}} d\omega.$$

Using the change of variable $\omega = t^{-2}$ gives

$$I = \sum_{n=0}^\infty (-1)^n e^{-(n+1)\psi} \frac{2n+1}{(2\pi)^{1/2}} \left(\int_0^\infty \exp \left[-\frac{1}{2} \left\{ \frac{(2n+1)t}{2} - \frac{\psi}{t} \right\}^2 \right] dt \right).$$

Applying the Cauchy-Schlömilch identity to the inner integral yields

$$\int_0^\infty \exp \left[-\frac{1}{2} \left\{ \frac{(2n+1)t}{2} - \frac{\psi}{t} \right\}^2 \right] dt = \int_0^\infty \frac{e^{-y^2/2}}{2n+1} dy = \frac{(2\pi)^{1/2}}{2n+1},$$

which implies $I = \sum_{n=0}^\infty (-1)^n \exp\{-(n+1)\psi\} = \{1 + \exp(\psi)\}^{-1}$. \square

Remark 3. When $\alpha = \kappa$, we have the limiting result $(\alpha^2 - \kappa^2)^{-1} p_{\text{GIG}}\{1, 0, (\alpha^2 - \kappa^2)^{1/2}\} = 1$, or equivalently in terms of densities, with a marginal improper uniform prior, $p(\lambda) = 1$,

$$\int_0^\infty \phi(b | -a\lambda, c\lambda) d\lambda = a^{-1} \exp\{-2 \max(ab/c, 0)\}. \quad (10)$$

This pseudo-likelihood represents support vector machines as a global-local mixture. The identity for quantile regression, which is a limiting case of the above identities by applying Fatou-Lebesgue theorem, is the following:

$$c^{-1} \exp\{2c^{-1} \rho_q(b)\} = \int_0^\infty \phi(b | \lambda - 2\tau\lambda, c\lambda) e^{-2\tau(1-\tau)\lambda} d\lambda, \quad c, \tau > 0,$$

where $\rho_q(b) = |b|/2 + (q - 1/2)b$ is the check-loss function (Polson & Scott, 2013).

Polson & Scott (2011) derive this as a direct consequence of the lasso identity

$$\int_0^\infty p/(2\pi\lambda)^{1/2} \exp \left\{ - \left(p^2\lambda + q^2\lambda^{-1} \right) / 2 \right\} d\lambda = e^{-|pq|}.$$

Applying the Liouville identity yields

$$\int_0^\infty f \left(ax + \frac{b}{x} \right) x^{-1/2} dx = a^{-1/2} \int_0^\infty f \left\{ 2(ab)^{1/2} + y \right\} y^{-1/2} dy, \quad a, b > 0.$$

Setting $f(x) = e^{-x}$, $a = p^2/2$, and $b = q^2/2$ we get

$$\begin{aligned} \int_0^\infty \frac{e^{-(p^2\lambda + q^2\lambda^{-1})/2}}{\lambda^{1/2}} d\lambda &= \frac{2^{1/2}}{p} \int_0^\infty e^{-|pq|+y} y^{-1/2} dy \\ &= \frac{2^{1/2}e^{-|pq|}}{p} \int_0^\infty e^{-y} y^{-1/2} dy = \frac{(2\pi)^{1/2}e^{-|pq|}}{p}. \end{aligned}$$

Hans (2011) shows that the elastic-net regression can be recast as a global-local mixture with a mixing density belonging to the orthant-normal family of distributions. The orthant-normal prior on a single regression coefficient, β , given hyper-parameters λ_1 and λ_2 , has a density function with the following form:

$$p(\beta \mid \lambda_1, \lambda_2) = \begin{cases} \phi(\beta \mid \frac{\lambda_1}{2\lambda_2}, \frac{\sigma^2}{\lambda_2}) / 2\Phi \left(-\frac{\lambda_1}{2\sigma\lambda_2^{1/2}} \right), & \beta < 0, \\ \phi(\beta \mid \frac{-\lambda_1}{2\lambda_2}, \frac{\sigma^2}{\lambda_2}) / 2\Phi \left(-\frac{\lambda_1}{2\sigma\lambda_2^{1/2}} \right), & \beta \geq 0. \end{cases} \quad (11)$$

3 Convolution mixtures

Another interesting area of application is convolution mixtures and marginal densities for location-scale mixture problems. We show that the Cauchy convolution (Pillai & Meng, 2016) and inverse-gamma convolution can be derived similarly (Polson & Scott, 2012). Bhadra et al. (2016b) shows that the regularly varying tails of half-Cauchy priors work well for low-dimensional functions of normal vector mean, where flat priors give poorly calibrated inference.

Lemma 4. Let $X_i \sim \mathcal{C}(0, 1)$ ($i = 1, 2$) be Cauchy distributed random variates, then $Z = w_1X_1 + w_2X_2 \sim \mathcal{C}(0, w_1 + w_2)$. where $w_1, w_2 > 0$.

Lemma 5. Let $X_i \sim \mathcal{IG}(\alpha t_i, \alpha t_i^2)$ ($i = 1, 2$), then $Z = X_1 + X_2 \sim \mathcal{IG}\{\alpha(t_1 + t_2), \alpha(t_1^2 + t_2^2)\}$, where $\alpha, t_1, t_2 \geq 0$, and $\mathcal{IG}(\alpha t, \alpha t^2)$ is an inverse-Gaussian random variable with density

$$f(x) = \frac{t\alpha^{1/2}e^t}{(2\pi)^{1/2}x^{3/2}} \exp \left(-\frac{\alpha t^2}{2x} - \frac{x}{2\alpha} \right), \quad x \geq 0.$$

Both of these results follow from straightforward applications of the Cauchy-Schlömilch transformation. We give a proof for the Cauchy convolution identity below.

Proof. Exploiting symmetry and the Lagrange identity $(a^2 + b^2)(c^2 + d^2) = (ac + bd)^2 + (ad - bc)^2$, leads to the convolution density

$$\begin{aligned} f_Z(z) &= 2 \int_0^\infty \frac{1}{\pi w_1(1 + x^2/w_1^2)} \frac{1}{\pi w_2\{1 + (z - x)^2/w_2^2\}} dx \\ &= \frac{2}{\pi^2 w_1 w_2} \int_0^\infty \frac{1}{\{1 + w_1^{-1} w_2^{-1} x(z - x)\}^2 + \{w_2^{-1} z - (w_1^{-1} + w_2^{-1})x\}^2} dx. \end{aligned}$$

Transforming x to $x + w_2^{-1} z (w_1^{-1} + w_2^{-1})^{-1}$ and letting $a = 1 + z^2(w_1 + w_2)^{-2}$, $b = (w_1 w_2)^{-1}$, $c = z(w_2 - w_1)\{(w_1 + w_2)w_1 w_2\}^{-1}$, $d = z(w_2 - w_1)\{(w_1 + w_2)w_1 w_2\}^{-1}$ gives

$$\begin{aligned} f_Z(z) &= \frac{2}{\pi^2 w_1 w_2} \int_0^\infty \left[\left\{ 1 + \frac{z^2}{(w_1 + w_2)^2} - \frac{x^2}{w_1 w_2} + xz \frac{w_2 - w_1}{(w_1 + w_2)w_1 w_2} \right\}^2 + x^2 \left(\frac{w_1 + w_2}{w_1 w_2} \right)^2 \right]^{-1} dx \\ &= \frac{2}{\pi^2 w_1 w_2} \int_0^\infty \frac{dx}{(a - bx^2 + cx)^2 + x^2 d^2} = \frac{2}{\pi^2 w_1 w_2} \int_0^\infty \frac{dx/x^2}{(a/x - bx + c)^2 + d^2}. \end{aligned}$$

If we let $y = x^{-1}$ and apply the Cauchy-Schlömilch transformation, we arrive at

$$f_Z(z) = \frac{2}{\pi w_1 w_2} \int_0^\infty \frac{dy}{2a(y^2 + d^2)} = \frac{1}{\pi w_1 w_2} \frac{1}{ad} = \frac{1}{\pi(w_1 + w_2)} \frac{1}{1 + z^2/(w_1 + w_2)^2}.$$

A simple induction argument proves that the sum of any number of independent Cauchy random variates is also another Cauchy. \square

One can also use the characteristic function of $X \sim \mathcal{C}(\mu, \sigma)$, $\psi_X(t) = \exp(it\mu - |t|\sigma^2)$, and the relation $\psi_{X+Y}(t) = \psi_X(t)\psi_Y(t)$ to derive the result in just one step. For $X = \sum_{i=1}^p \omega_i C_i$ and $C_i \sim \mathcal{C}(0, 1)$, when $\sum_{i=1}^p \omega_i = 1$ we have $\phi_X(t) = \exp(-\sum_{i=1}^p \omega_i |t|) = \exp(-|t|) = \phi_C(t)$, where $C \sim \mathcal{C}(0, 1)$.

The most general result in this category is due to [Pillai & Meng \(2016\)](#), who they showed the following: Let (X_1, \dots, X_m) and (Y_1, \dots, Y_m) be independent and identically distributed $\mathcal{N}(0, \Sigma)$ for an arbitrary positive definite matrix Σ , then $Z = \sum_{j=1}^m w_j X_j / Y_j \sim \mathcal{C}(0, 1)$, as long as (w_1, \dots, w_m) is independent of (X, Y) , $w_j \geq 0$ ($j = 1, \dots, m$) and $\sum_{j=1}^m w_j = 1$.

4 Discussion

The Cauchy-Schlömilch and Liouville transformations not only guarantee an simple normalizing constant for $f(\cdot)$, it also establishes the wide class of unimodal densities as global-local scale mixtures. Global-local scale mixtures that are conditionally Gaussian hold a special place in statistical modeling and can be rapidly fit using an expectation-maximization algorithm, as pointed out by [Polson & Scott \(2013\)](#). [Palmer et al. \(2011\)](#) provides a similar tool for modeling multivariate dependence by writing general non-Gaussian multivariate densities as multivariate Gaussian scale mixtures.

We end our paper with conjectures that two other remarkable identities arise as corollaries of such transformation identities. The first one is a recent result by [Zhang et al.](#)

(2014) that proves a uniform correlation mixture of a bivariate Gaussian density with unit variance is a function of the maximum norm:

$$\int_{-1}^1 \frac{1}{4\pi(1-\rho^2)^{1/2}} \exp \left\{ -\frac{x_1^2 + x_2^2 - 2\rho x_1 x_2}{2(1-\rho^2)} \right\} d\rho = \frac{1}{2} \{1 - \Phi(\|x\|_\infty)\} , \quad (12)$$

where $\Phi(\cdot)$ is the standard normal distribution function and $\|x\|_\infty = \max\{x_1, x_2\}$. The bivariate density on the right side of (12) was introduced by Bryson & Johnson (1982) as uniform mixtures of a chi random variate with 3 degrees of freedom, but the representation as a uniform correlation mixture is a new find. We make a few remarks connected to the Erdelyi's integral identity, which is key to the proof of the uniform correlation mixture of (12).

Lemma 6. *Erdelyi's identity, defined by*

$$\int_{1/2}^{\infty} \frac{e^{-x^2 z}}{4\pi z(2z-1)^{1/2}} dz = \frac{1}{2} \{1 - \Phi(x)\} , \quad x \geq 0, \quad (13)$$

follows from the Laplace transformation $(1+u)^{-1} = \int_0^\infty \exp\{-v(1+u)\} dv$.

Proof. Apply the transform $u = 2z - 1$ to the left hand side of (13), denoted by I , to obtain

$$I = \int_0^\infty \frac{e^{-x^2/\{2(1+u)\}}}{4\pi u^{1/2}(1+u)} du .$$

Using the Laplace transformation $(1+u)^{-1} = \int_0^\infty e^{-v(1+u)} dv$ yields

$$\begin{aligned} I &= \int_0^\infty \frac{e^{-x^2/\{2(1+u)\}}}{4\pi u^{1/2}} \int_0^\infty e^{-v(1+u)} dv du = \int_{v=0}^\infty \int_{u=0}^\infty \frac{e^{-(x^2/2+v)(1+u)}}{4\pi u^{1/2}} dv du \\ &= \int_{v=0}^\infty \frac{1}{4\pi} e^{-(x^2/2+v)} \int_{u=0}^\infty u^{-1/2} e^{-(x^2/2+v)u} du dv = \int_{v=0}^\infty \frac{e^{-(x^2+2v)/2}}{2(2\pi)^{1/2}} \frac{1}{(x^2+2v)^{1/2}} dv , \end{aligned}$$

and letting $z^2 = x^2 + 2v$ we get

$$I = \frac{1}{2} \int_{z=|x|}^\infty \frac{1}{(2\pi)^{1/2}} e^{-z^2/2} dz = \frac{1}{2} \{1 - \Phi(|x|)\} .$$

□

The second candidate is the symmetric stable distribution, defined by its characteristic function $\phi(t) = \exp(-|t|^\alpha)$, $0 < \alpha \leq 2$. It admits a normal scale mixture representation with mixing density as $f(v) = 2^{-1} s_{\alpha/2}(v/2)$, $v > 0$, where $s_{\alpha/2}$ is the positive stable density with index $\alpha/2$ (Gneiting, 1997). The exponential power density arising as a dual of the symmetric stable density also has a normal scale mixture representation with important application in Bayesian bridge regression (Polson et al., 2014).

$$e^{-|x|^\alpha} = \int_0^\infty e^{-x\eta} g(\eta) d\eta, \quad g(\eta) = \sum_{j=1}^\infty (-1)^j \frac{\eta^{-j\alpha-1}}{j! \Gamma(-\alpha j)} ,$$

Polson et al. (2014) derive this as a limiting result of the scale-mixture of beta representation for k -montone densities and utilizing the complete monotonicity of exponential power density. Regularization, in this case, is an outcome of a normal scale mixture with respect to an α -stable random variable. We conjecture that these two results follow from the Cauchy-Schlömilch formula (1). Other potential applications include using Liouville formula to recognize and generate global-local mixtures, and to calculate higher-order closed-form moments $E(X^n)$ for random variables X that admit a global-local representation.

References

- ANDREWS, D. & MALLOWS, C. (1974). Scale mixtures of normal distributions. *Journal of the Royal Statistical Society. Series B: Statistical Methodology* **36**, 99–102.
- BAKER, R. (2008). Probabilistic applications of the Schlömilch transformation. *Communications in Statistics – Theory and Methods* **37**, 2162–2176.
- BARNDORFF-NIELSEN, O., KENT, J. & SØRENSEN, M. (1982). Normal variance-mean mixtures and z distributions. *International Statistical Review* **50**, 145–159.
- BHADRA, A., DATTA, J., POLSON, N. G. & WILLARD, B. (2016a). The Horseshoe+ Estimator of Ultra-Sparse Signals. *Bayesian Analysis (to appear)* .
- BHADRA, A., DATTA, J., POLSON, N. G. & WILLARD, B. T. (2016b). Default Bayesian Analysis with Global-Local Shrinkage Priors. *Biometrika (to appear)* .
- BOROS, G., MOLL, V. H. & FONCANNON, J. (2006). Irresistible integrals: symbolics, analysis and experiments in the evaluation of integrals. *The Mathematical Intelligencer* **28**, 65–68.
- BRYSON, M. C. & JOHNSON, M. E. (1982). Constructing and simulating multivariate distributions using Khintchine’s theorem. *Journal of Statistical Computation and Simulation* **16**, 129–137.
- GNEITING, T. (1997). Normal scale mixtures and dual probability densities. *Journal of Statistical Computation and Simulation* **59**, 375–384.
- HANS, C. (2011). Comment on Article by Polson and Scott. *Bayesian Analysis* **6**, 37–41.
- JONES, M. (2002). On Khintchine’s theorem and its place in random variate generation. *The American Statistician* **16**, 304–307.
- JONES, M. C. (2014). Generating distributions by transformation of scale. *Statist. Sinica* **24**, 749–772.
- LÉVY, P. (1940). Sur certains processus stochastiques homogènes. *Compositio mathematica* **7**, 283–339.

- PALMER, J. A., KREUTZ-DELGADO, K. & MAKEIG, S. (2011). AMICA: An adaptive mixture of independent component analyzers with shared components. Tech. rep., San Diego, CA: Technical report, Swartz Center for Computational Neuroscience.
- PILLAI, N. S. & MENG, X.-L. (2016). An unexpected encounter with Cauchy and Lévy. *Annals of Statistics (to appear)*.
- POLSON, N. G. & SCOTT, J. G. (2012). On the Half-Cauchy Prior for a Global Scale Parameter. *Bayesian Analysis* **7**, 887–902.
- POLSON, N. G. & SCOTT, J. G. (2013). Data augmentation for non-Gaussian regression models using variance-mean mixtures. *Biometrika* **100**, 459–471.
- POLSON, N. G., SCOTT, J. G. & WINDLE, J. (2013). Bayesian inference for logistic models using Pólya–Gamma latent variables. *Journal of the American Statistical Association* **108**, 1339–1349.
- POLSON, N. G., SCOTT, J. G. & WINDLE, J. (2014). The Bayesian bridge. *Journal of the Royal Statistical Society. Series B: Statistical Methodology* **76**, 713–733.
- POLSON, N. G. & SCOTT, S. L. (2011). Data augmentation for support vector machines. *Bayesian Analysis* **6**, 1–23.
- SESHADRI, V. (2004). Halphen’s laws. In *Encyclopedia of Statistical Sciences*. Hoboken, New Jersey: John Wiley and Sons, Inc.
- ZHANG, K., BROWN, L. D., GEORGE, E. & ZHAO, L. (2014). Uniform Correlation Mixture of Bivariate Normal Distributions and Hypercubically Contoured Densities That Are Marginally Normal. *The American Statistician* **68**, 183–187.