

Nonzero-sum stochastic differential games with impulse controls: a verification theorem with applications

René Aïd^{*} Matteo Basei[†] Giorgia Callegaro[‡]
 Luciano Campi[§] Tiziano Vargiolu[‡]

March 22, 2022

Abstract

We consider a general nonzero-sum impulse game with two players. The main mathematical contribution of the paper is a verification theorem which provides, under some regularity conditions, a suitable system of quasi-variational inequalities for the value functions and the optimal strategies of the two players. As an application, we study an impulse game with a one-dimensional state variable, following a real-valued scaled Brownian motion, and two players with linear and symmetric running payoffs. Thanks to the verification theorem, we find and fully characterize a Nash equilibrium by providing explicit expressions for the value functions and the optimal strategies of the players. Finally, we prove some asymptotic results with respect to the intervention costs for the one-dimensional symmetric game.

Keywords: stochastic differential game, impulse control, Nash equilibrium, quasi-variational inequality.

1 Introduction

In this article, we study a general two-player nonzero-sum stochastic differential game with impulse controls. In few words, after setting the general framework, we focus on the notion of Nash equilibrium and identify the corresponding system of quasi-variational inequalities (QVIs). As an application, we consider an impulse game with a one-dimensional state variable and fully solve the system of QVIs, obtaining explicit expressions for the value functions as well as the optimal strategies at equilibrium.

More specifically, we consider a game where two players can affect a continuous-time stochastic process X by discrete-time interventions which consist in shifting X to a new state. When none of the players intervenes, we assume X to diffuse according to a standard stochastic differential equation. Each intervention corresponds to a cost for the intervening player and a gain for the opponent. The strategy of player $i \in \{1, 2\}$ is determined by a couple $\varphi_i = (A_i, \xi_i)$, where A_i is a fixed subset of \mathbb{R}^d and ξ_i is a continuous function: namely, player i intervenes if and only if the process X exits from A_i and, when this happens, she shifts the process from state x to state $\xi_i(x)$. Once the strategies $\varphi_i = (A_i, \xi_i)$, $i \in \{1, 2\}$, and a starting point x have been chosen, a couple of impulse controls $u_i(x; \varphi_1, \varphi_2) = \{(\tau_{i,k}, \delta_{i,k})\}_{1 \leq k \leq M_i}$ is uniquely defined: $\tau_{i,k}$ is the k -th

^{*}Economics Department (LEDa), University Paris Dauphine, and Finance For Energy Market Research Centre (FiME).

[†]Department of Mathematics, University of Padova, and Laboratoire de Probabilités et Modèles Aléatoires (LPMA), University Paris Diderot.

[‡]Department of Mathematics, University of Padova.

[§]Department of Statistics, London School of Economics and Political Science. E-mail: L.Campi@lse.ac.uk

intervention time of player i and $\delta_{i,k}$ is the corresponding impulse. Each player aims at maximizing her payoff, defined as follows: for every x belonging to some fixed subset $S \subseteq \mathbb{R}^n$ and every couple of strategies (φ_1, φ_2) , we set

$$J^i(x; \varphi_1, \varphi_2) := \mathbb{E}_x \left[\int_0^{\tau_S} e^{-\rho_i s} f_i(X_s) ds + \sum_{1 \leq k \leq M_i : \tau_{i,k} < \tau_S} e^{-\rho_i \tau_{i,k}} \phi_i(X_{(\tau_{i,k})^-}, \delta_{i,k}) \right. \\ \left. + \sum_{1 \leq k \leq M_j : \tau_{j,k} < \tau_S} e^{-\rho_i \tau_{j,k}} \psi_i(X_{(\tau_{j,k})^-}, \delta_{j,k}) + e^{-\rho_i \tau_S} h_i(X_{(\tau_S)^-}) \mathbb{1}_{\{\tau_S < +\infty\}} \right], \quad (1.1)$$

where $i, j \in \{1, 2\}$, $i \neq j$ and τ_S is the exit time of X from S . The couple $(\varphi_1^*, \varphi_2^*)$ is a Nash equilibrium if, for every couple of strategies φ_1, φ_2 , we have

$$J^1(x; \varphi_1^*, \varphi_2^*) \geq J^1(x; \varphi_1, \varphi_2^*), \quad \text{and} \quad J^2(x; \varphi_1^*, \varphi_2^*) \geq J^2(x; \varphi_1^*, \varphi_2). \quad (1.2)$$

The game just described is connected to the following system of QVIs, where $i, j \in \{1, 2\}$ with $i \neq j$ and $\mathcal{M}_i, \mathcal{H}_i$ are suitable intervention operators defined in Section 3.1:

$$\begin{aligned} V_i &= h_i, & \text{in } \partial S, \\ \mathcal{M}_j V_j - V_j &\leq 0, & \text{in } S, \\ \mathcal{H}_i V_i - V_i &= 0, & \text{in } \{\mathcal{M}_j V_j - V_j = 0\}, \\ \max \{ \mathcal{A}V_i - \rho_i V_i + f_i, \mathcal{M}_i V_i - V_i \} &= 0, & \text{in } \{\mathcal{M}_j V_j - V_j < 0\}. \end{aligned} \quad (1.3)$$

The first contribution of our paper is the Verification Theorem 3.3: if two functions V_i , with $i \in \{1, 2\}$, are a solution to (1.3), have polynomial growth and satisfy the regularity condition

$$V_i \in C^2(\mathcal{D}_j \setminus \partial \mathcal{D}_i) \cap C^1(\mathcal{D}_j) \cap C(S), \quad (1.4)$$

where $j \in \{1, 2\}$ with $j \neq i$ and $\mathcal{D}_j = \{\mathcal{M}_j V_j - V_j < 0\}$, then they coincide with the value functions of the game and a characterization of the Nash strategy is possible.

We stress that, even though stochastic differential games have been widely studied in the last decades, the case of nonzero-sum impulse games did not deserve so much attention so far and it has never been considered, to the best of our knowledge, from a QVI perspective. Indeed, related former works only address zero-sum stopping games [12], the corresponding nonzero-sum problems [2] (with only two, very recent, explicit examples in [8] and [11]) and zero-sum impulse games [9]. We notice that the QVI system proposed in [9] for zero-sum impulse games can be obtained as a particular case of our framework. In the field of stochastic differential games, it is also worth mentioning [10], which provides a link between two-player nonzero-sum games of optimal stopping and two-player nonzero-sum games of singular control. Only the two papers [6, 7] deal with some nonzero-sum stochastic differential games with impulse controls and finite horizon using an approach based on backward stochastic differential equations and the maximum principle. Notice that in those two papers the sequence of stopping times along which impulses can be applied is given, hence the players can choose only the size of the impulses. In [6, 7] some examples are provided, where one Nash equilibrium is characterized in terms of solutions to suitable ordinary differential equations.

Our second contribution to this stream of research is a first example of a solvable impulse game. Using the Verification Theorem 3.3 described above and solving explicitly the system of QVIs (1.3) we are able to fully characterize one Nash equilibrium.

More in detail, we consider a two-player impulse game with a one-dimensional state variable X , modelled by a real-valued (scaled) Brownian motion. The two players have symmetric linear running payoffs and they can intervene on X by shifting it from its current state, say x , to some other state $x + \delta$, with $\delta \in \mathbb{R}$. When a player intervenes, she faces a penalty while her opponent faces a gain, both consisting in a fixed and in a variable part, which is assumed proportional to the size of the impulse. Hence, the players objective functions are

$$J^1(x; \varphi_1, \varphi_2) := \mathbb{E}_x \left[\int_0^\infty e^{-\rho s} (X_s - s_1) ds - \sum_{k \geq 1} e^{-\rho \tau_{1,k}} (c + \lambda |\delta_{1,k}|) + \sum_{k \geq 1} e^{-\rho \tau_{2,k}} (\tilde{c} + \tilde{\lambda} |\delta_{2,k}|) \right], \\ J^2(x; \varphi_1, \varphi_2) := \mathbb{E}_x \left[\int_0^\infty e^{-\rho s} (s_2 - X_s) ds - \sum_{k \geq 1} e^{-\rho \tau_{2,k}} (c + \lambda |\delta_{2,k}|) + \sum_{k \geq 1} e^{-\rho \tau_{1,k}} (\tilde{c} + \tilde{\lambda} |\delta_{1,k}|) \right],$$

where $\{(\tau_{i,k}, \delta_{i,k})\}_{k \geq 1}$ denotes the impulse controls of player i associated to the strategies φ_1, φ_2 .

Some preliminary heuristics on the QVIs in (1.3) leads us to consider a pair of candidates for the functions V_i . Then, a careful application of the verification theorem shows that such candidates actually coincide with the value functions of the game. In particular, a practical characterization of the Nash equilibria is possible: player 1 (resp. player 2) intervenes when the state X is smaller than \bar{x}_1 (resp. greater than \bar{x}_2) and moves the process to x_1^* (resp. x_2^*), for suitable \bar{x}_i, x_i^* . We provide explicit expressions for the value functions and for the parameters \bar{x}_i, x_i^* . Finally, we study the behaviour of the intervention region in some limit cases. In particular, we remark that in the case where $c = \tilde{c}$ and $\lambda = \tilde{\lambda}$, the game does not have an admissible Nash equilibrium. Moreover, such an example can be given different economic interpretations: it can be viewed as a competition between central banks of two countries controlling the exchange rate between their currencies or as an interaction model between an energy producer and a large consumer (see Section 4.1 for details).

The outline of the paper is the following. In Section 2, we rigorously formulate the general impulse game and give the notions of admissible strategies and of Nash equilibrium. Section 3 provides the associated system of QVIs and the corresponding verification theorem. In Section 4 we consider and fully compute a Nash equilibrium in a one-dimensional impulse game. Finally, Section 5 concludes.

2 Nonzero-sum stochastic impulse games

In this section we introduce a general class of two-player nonzero-sum stochastic differential games with impulse controls.

Let $(\Omega, \mathcal{F}, \{\mathcal{F}_t\}_{t \geq 0}, \mathbb{P})$ be a filtered probability space whose filtration satisfies the usual conditions of right-continuity and \mathbb{P} -completeness, and let $\{W_t\}_{t \geq 0}$ be a k -dimensional $\{\mathcal{F}_t\}_{t \geq 0}$ -adapted Brownian motion. For every $t \geq 0$ and $\xi \in L^p(\mathcal{F}_t)$ for some $p > 1$, we denote by $Y^{t,\xi} = \{Y_s^{t,\xi}\}_{s \geq t}$ a solution to the problem

$$dY_s^{t,\xi} = b(Y_s^{t,\xi})ds + \sigma(Y_s^{t,\xi})dW_s, \quad s \geq t, \quad (2.1)$$

with initial condition $Y_t^{t,\xi} = \xi$, where $b : \mathbb{R}^d \rightarrow \mathbb{R}^d$ and $\sigma : \mathbb{R}^d \rightarrow \mathbb{R}^{d \times k}$ are given functions. Throughout the whole paper, we assume that the coefficients b and σ are globally Lipschitz continuous, i.e. there exists a constant $K > 0$ such that for all $y_1, y_2 \in \mathbb{R}^d$ we have

$$\|b(y_1) - b(y_2)\| + \|\sigma(y_1) - \sigma(y_2)\| \leq K\|y_1 - y_2\|,$$

so that (2.1) admits a unique strong solution satisfying classical a-priori estimates (see, e.g., [13, Sect. 2.5] or [14, Thm. 2.2] among others).

We consider two players, that will be indexed by $i \in \{1, 2\}$. Let S be an open subset of \mathbb{R}^d and let Z_i be a fixed subset of \mathbb{R}^{l_i} , with $l_i \in \mathbb{N}$. Equation (2.1) models the underlying process when none of the players intervenes. If player i intervenes with some impulse $\delta \in Z_i$, the process is shifted from its current state x to a new state $\Gamma^i(x, \delta)$, where $\Gamma^i : \mathbb{R}^d \times Z_i \rightarrow S$ is a given continuous function. Each intervention corresponds to a cost for the intervening player and to a gain for the opponent, both depending on the state x and the impulse δ .

The action of the players is modelled via discrete-time controls: an impulse control for player i is a sequence

$$u_i = \{(\tau_{i,k}, \delta_{i,k})\}_{1 \leq k \leq M_i}, \quad (2.2)$$

where $M_i \in \mathbb{N} \cup \{\infty\}$ denotes the number of interventions of player i , $\{\tau_{i,k}\}_k$ is a non-decreasing sequence of stopping times (the intervention times) and $\{\delta_{i,k}\}_k$ are Z_i -valued $\mathcal{F}_{\tau_{i,k}}$ -measurable random variables (the corresponding impulses).

For the sake of tractability, we assume that the behaviour of the players, modelled by impulse controls, is driven by strategies, which are defined as follows.

Definition 2.1. *A strategy for player $i \in \{1, 2\}$ is a pair $\varphi_i = (A_i, \xi_i)$, where A_i is a fixed open subset of \mathbb{R}^d and ξ_i is a continuous function from \mathbb{R}^d to Z_i .*

Strategies determine the action of the players in the following sense. Let $x \in S$ be an initial value for the state variable. Once some strategies $\varphi_i = (A_i, \xi_i)$, $i \in \{1, 2\}$, have been chosen, a pair of impulse controls, which we denote by $u_i(x; \varphi_1, \varphi_2)$, is uniquely defined by the following procedure:

- player i intervenes if and only if the process exits from A_i ,
in which case the impulse is given by $\xi_i(y)$, where y is the state;
 - if both players want to act, player 1 has the priority;
 - the game ends when the process exits from S .
- (2.3)

In the following definition we provide a rigorous formalization of the controls associated to a pair of strategies and the corresponding controlled process, which we denote by $X^{x; \varphi_1, \varphi_2}$. Moreover O denotes a generic subset of \mathbb{R}^d and, finally, we adopt the conventions $\inf \emptyset = \infty$ and $[\infty, \infty[= \emptyset$.

Definition 2.2. Let $x \in S$ and let $\varphi_i = (A_i, \xi_i)$ be a strategy for player $i \in \{1, 2\}$. Let $\tilde{\tau}_0 = 0, x_0 = x, \tilde{X}^0 = Y^{\tilde{\tau}_0, x_0}, \alpha_0^S = \infty$. For every $k \in \mathbb{N}, k \geq 1$, we define, by induction,

$$\begin{aligned}
\alpha_k^O &= \inf\{s > \tilde{\tau}_{k-1} : \tilde{X}_s^{k-1} \notin O\}, & [\text{exit time from } O \subseteq \mathbb{R}^d] \\
\tilde{\tau}_k &= (\alpha_k^{A_1} \wedge \alpha_k^{A_2} \wedge \alpha_k^S) \mathbb{1}_{\{\tilde{\tau}_{k-1} < \alpha_{k-1}^S\}} + \tilde{\tau}_{k-1} \mathbb{1}_{\{\tilde{\tau}_{k-1} = \alpha_{k-1}^S\}}, & [\text{intervention time}] \\
m_k &= \mathbb{1}_{\{\alpha_k^{A_1} \leq \alpha_k^{A_2}\}} + 2 \mathbb{1}_{\{\alpha_k^{A_2} < \alpha_k^{A_1}\}}, & [\text{index of the player interv. at } \tilde{\tau}_k] \\
\tilde{\delta}_k &= \xi_{m_k}(\tilde{X}_{\tilde{\tau}_k}^{k-1}) \mathbb{1}_{\{\tilde{\tau}_k < \infty\}}, & [\text{impulse}] \\
x_k &= \Gamma^{m_k}(\tilde{X}_{\tilde{\tau}_k}^{k-1}, \tilde{\delta}_k) \mathbb{1}_{\{\tilde{\tau}_k < \infty\}}, & [\text{starting point for the next step}] \\
\tilde{X}^k &= \tilde{X}^{k-1} \mathbb{1}_{[0, \tilde{\tau}_k[} + Y^{\tilde{\tau}_k, x_k} \mathbb{1}_{[\tilde{\tau}_k, \infty[}. & [\text{contr. process up to the } k\text{-th interv.}]
\end{aligned}$$

Let $\bar{k} \in \mathbb{N} \cup \{\infty\}$ be the index of the last significant intervention, and let $M_i \in \mathbb{N} \cup \{\infty\}$ be the number of interventions of player i :

$$\begin{aligned}
\bar{k} &:= \sup\{k \in \mathbb{N} : \mathbb{P}(\tilde{\tau}_k = \alpha_k^S) < 1 \text{ and } \mathbb{P}(\tilde{\tau}_k = \infty) < 1\}, \\
M_i &:= \sum_{1 \leq k \leq \bar{k}} \mathbb{1}_{\{m_k = i\}}(k).
\end{aligned}$$

For $i \in \{1, 2\}$ and $1 \leq k \leq M_i$, let $\eta(i, k) = \min\{l \in \mathbb{N} : \sum_{1 \leq h \leq l} \mathbb{1}_{\{m_h = i\}} = k\}$ (index of the k -th intervention of player i) and let

$$\tau_{i,k} := \tilde{\tau}_{\eta(i,k)}, \quad \delta_{i,k} := \tilde{\delta}_{\eta(i,k)}. \quad (2.4)$$

Finally, the controls $u_i(x; \varphi_1, \varphi_2)$, $i \in \{1, 2\}$, the controlled process $X^{x; \varphi_1, \varphi_2}$ and the exit time from S are defined by

$$\begin{aligned}
u_i(x; \varphi_1, \varphi_2) &:= \{(\tau_{i,k}, \delta_{i,k})\}_{1 \leq k \leq M_i}, \\
X^{x; \varphi_1, \varphi_2} &:= \tilde{X}^{\bar{k}}, \\
\tau_S^{x; \varphi_1, \varphi_2} &= \inf\{s > 0 : X_s^{x; \varphi_1, \varphi_2} \notin S\}.
\end{aligned}$$

To shorten the notations, we will simply write X and τ_S . Notice that player 1 has the priority in the case of simultaneous interventions (i.e., if $\alpha_k^{A_1} = \alpha_k^{A_2}$). In the following lemma we give a rigorous formulation to the properties outlined in (2.3).

Lemma 2.3. Let $x \in S$ and let $\varphi_i = (A_i, \xi_i)$ be a strategy for player $i \in \{1, 2\}$.

- The process X admits the following representation (with the convention $[\infty, \infty[= \emptyset$):

$$X_s = \sum_{k=0}^{\bar{k}-1} Y_s^{\tilde{\tau}_k, x_k} \mathbb{1}_{[\tilde{\tau}_k, \tilde{\tau}_{k+1}[}(s) + Y_s^{\tilde{\tau}_{\bar{k}}, x_{\bar{k}}} \mathbb{1}_{[\tilde{\tau}_{\bar{k}}, \infty[}(s). \quad (2.5)$$

- The process X is right-continuous. More precisely, X is continuous and satisfies Equation (2.1) in $[0, \infty[\setminus \{\tau_{i,k} : \tau_{i,k} < \infty\}$, whereas X is discontinuous in $\{\tau_{i,k} : \tau_{i,k} < \infty\}$, where we have

$$X_{\tau_{i,k}} = \Gamma^i(X_{(\tau_{i,k})^-}, \delta_{i,k}), \quad \delta_{i,k} = \xi_i(X_{(\tau_{i,k})^-}), \quad X_{(\tau_{i,k})^-} \in \partial A_i. \quad (2.6)$$

- The process X never exits from the set $A_1 \cap A_2$.

Proof. We just prove the first property in (2.6), the other ones being immediate. Let $i \in \{1, 2\}$, $1 \leq k \leq M_i$ with $\tau_{i,k} < \infty$ and set $\sigma = \eta(i, k)$, with η as in Definition 2.2. By (2.4), (2.5) and Definition 2.2, we have

$$\begin{aligned} X_{\tau_{i,k}} &= X_{\tilde{\tau}_\sigma} = Y_{\tilde{\tau}_\sigma}^{\tilde{\tau}_\sigma, x_\sigma} = x_\sigma = \Gamma^i(\tilde{X}_{\tilde{\tau}_\sigma}^{\sigma-1}, \tilde{\delta}_\sigma) \\ &= \Gamma^i(\tilde{X}_{(\tilde{\tau}_\sigma)^-}^{\sigma-1}, \tilde{\delta}_\sigma) = \Gamma^i(X_{(\tilde{\tau}_\sigma)^-}, \tilde{\delta}_\sigma) = \Gamma^i(X_{(\tau_{i,k})^-}, \delta_{i,k}), \end{aligned}$$

where in the fifth equality we have used the continuity of the process $\tilde{X}^{\sigma-1}$ in $[\tilde{\tau}_{\sigma-1}, \infty[$ and in the next-to-last equality we exploited the fact that $\tilde{X}^{\sigma-1} \equiv X$ in $[0, \tilde{\tau}_\sigma[$. \square

Each player aims at maximizing her payoff, consisting of four discounted terms: a running payoff, the costs due to her interventions, the gains due to her opponent's interventions and a terminal payoff. More precisely, for each $i \in \{1, 2\}$ we consider $\rho_i > 0$ (the discount rate) and continuous functions $f_i : \mathbb{R}^d \rightarrow \mathbb{R}$ (the running payoffs), $h_i : \mathbb{R}^d \rightarrow \mathbb{R}$ (the terminal payoffs) and $\phi_i : \mathbb{R}^d \times Z_i \rightarrow \mathbb{R}$, $\psi_i : \mathbb{R}^d \times Z_j \rightarrow \mathbb{R}$ (the interventions' costs and gains), where $j \in \{1, 2\}$ with $j \neq i$. The payoff of player i is defined as follows.

Definition 2.4. Let $x \in S$, let (φ_1, φ_2) be a pair of strategies and let τ_S be defined as in Definition 2.2. For each $i \in \{1, 2\}$, provided that the right-hand side exists and is finite, we set

$$\begin{aligned} J^i(x; \varphi_1, \varphi_2) &:= \mathbb{E}_x \left[\int_0^{\tau_S} e^{-\rho_i s} f_i(X_s) ds + \sum_{1 \leq k \leq M_i : \tau_{i,k} < \tau_S} e^{-\rho_i \tau_{i,k}} \phi_i(X_{(\tau_{i,k})^-}, \delta_{i,k}) \right. \\ &\quad \left. + \sum_{1 \leq k \leq M_j : \tau_{j,k} < \tau_S} e^{-\rho_i \tau_{j,k}} \psi_i(X_{(\tau_{j,k})^-}, \delta_{j,k}) + e^{-\rho_i \tau_S} h_i(X_{(\tau_S)^-}) \mathbb{1}_{\{\tau_S < +\infty\}} \right], \quad (2.7) \end{aligned}$$

where $j \in \{1, 2\}$ with $j \neq i$ and $\{(\tau_{i,k}, \delta_{i,k})\}_{1 \leq k \leq M_i}$ is the impulse control of player i associated to the strategies φ_1, φ_2 .

As usual in control theory, the subscript in the expectation denotes conditioning with respect to the available information (hence, it recalls the starting point). Notice that in the summations above we do not consider stopping times which equal τ_S (since the game ends in τ_S , any intervention is meaningless).

In order for J^i in (2.7) to be well defined, we now introduce the set of admissible strategies in $x \in S$.

Definition 2.5. Let $x \in S$ and $\varphi_i = (A_i, \xi_i)$ be a strategy for player $i \in \{1, 2\}$. We use the notations of Definition 2.2 and we say that the pair (φ_1, φ_2) is x -admissible if:

1. for $i \in \{1, 2\}$, the following random variables are in $L^1(\Omega)$:

$$\begin{aligned} &\int_0^{\tau_S} e^{-\rho_i s} |f_i|(X_s) ds, \quad e^{-\rho_i \tau_S} |h_i|(X_{(\tau_S)^-}), \\ &\sum_{\tau_{i,k} < \tau_S} e^{-\rho_i \tau_{i,k}} |\phi_i|(X_{(\tau_{i,k})^-}, \delta_{i,k}), \quad \sum_{\tau_{i,k} < \tau_S} e^{-\rho_i \tau_{i,k}} |\psi_i|(X_{(\tau_{i,k})^-}, \delta_{i,k}); \end{aligned} \quad (2.8)$$

2. for each $p \in \mathbb{N}$, the random variable $\|X\|_\infty = \sup_{t \geq 0} |X_t|$ is in $L^p(\Omega)$:

$$\mathbb{E}_x[\|X\|_\infty^p] < \infty; \quad (2.9)$$

3. if $\tau_{i,k} = \tau_{i,k+1}$ for some $i \in \{1, 2\}$ and $1 \leq k \leq M_i$, then $\tau_{i,k} = \tau_{i,k+1} = \tau_S$;
4. if there exists $\lim_{k \rightarrow +\infty} \tau_{i,k} =: \eta$ for some $i \in \{1, 2\}$, then $\eta = \tau_S$.

We denote by \mathcal{A}_x the set of the x -admissible pairs.

Thanks to the first condition in Definition 2.5, the payoffs $J^i(x; \varphi_1, \varphi_2)$ are well-defined. The second condition will be used in the proof of the Verification Theorem 3.3. As for the third and the fourth conditions, they prevent each player to exercise twice at the same time and to accumulate the interventions before τ_S .

We conclude the section with the definition of Nash equilibrium and the corresponding value functions in our setting.

Definition 2.6. *Given $x \in S$, we say that $(\varphi_1^*, \varphi_2^*) \in \mathcal{A}_x$ is a Nash equilibrium of the game if*

$$\begin{aligned} J^1(x; \varphi_1^*, \varphi_2^*) &\geq J^1(x; \varphi_1, \varphi_2^*), & \forall \varphi_1 \text{ s.t. } (\varphi_1, \varphi_2^*) \in \mathcal{A}_x, \\ J^2(x; \varphi_1^*, \varphi_2^*) &\geq J^2(x; \varphi_1^*, \varphi_2), & \forall \varphi_2 \text{ s.t. } (\varphi_1^*, \varphi_2) \in \mathcal{A}_x. \end{aligned}$$

Finally, the value functions of the game are defined as follows: if $x \in S$ and a Nash equilibrium $(\varphi_1^*, \varphi_2^*) \in \mathcal{A}_x$ exists, we set for $i \in \{1, 2\}$

$$V_i(x) := J^i(x; \varphi_1^*, \varphi_2^*).$$

3 A verification theorem

In this section we define a suitable differential problem for the value functions of nonzero-sum impulse games (see Section 3.1) and prove a verification theorem for such games (see Section 3.2).

3.1 The quasi-variational inequality problem

We now introduce the differential problem that should be satisfied by the value functions of our games: this will be key for stating the verification theorem in the next section.

Let us consider an impulse game as in Section 2. Assume that the corresponding value functions V_1, V_2 are defined for each $x \in S$ and that for $i \in \{1, 2\}$ there exists a (unique) function δ_i from S to Z_i such that

$$\{\delta_i(x)\} = \arg \max_{\delta \in Z_i} \{V_i(\Gamma^i(x, \delta)) + \phi_i(x, \delta)\}, \quad (3.1)$$

for each $x \in S$. We define the four intervention operators by

$$\begin{aligned} \mathcal{M}_i V_i(x) &= V_i(\Gamma^i(x, \delta_i(x))) + \phi_i(x, \delta_i(x)), \\ \mathcal{H}_i V_i(x) &= V_i(\Gamma^j(x, \delta_j(x))) + \psi_i(x, \delta_j(x)), \end{aligned} \quad (3.2)$$

for $x \in S$ and $i, j \in \{1, 2\}$, with $i \neq j$. Notice that $\mathcal{M}_i V_i(\cdot) = \max_{\delta} \{V_i(\Gamma^i(\cdot, \delta)) + \phi_i(\cdot, \delta)\}$.

The functions in (3.1) and (3.2) have an immediate and intuitive interpretation. Let x be the current state of the process; if player i (resp. player j) intervenes with impulse δ , the present value of the game for player i can be written as $V_i(\Gamma^i(x, \delta)) + \phi_i(x, \delta)$ (resp. $V_i(\Gamma^j(x, \delta)) + \psi_i(x, \delta)$): we have considered the value in the new state and the intervention cost (resp. gain). Hence, $\delta_i(x)$ in (3.1) is the impulse that player i would use in case she wants to intervene.

Similarly, $\mathcal{M}_i V_i(x)$ (resp. $\mathcal{H}_i V_i(x)$) represents the value of the game for player i when player i (resp. player $j \neq i$) takes the best immediate action and behaves optimally afterwards. Notice that it is not always optimal to intervene, so $\mathcal{M}_i V_i(x) \leq V_i(x)$, for each $x \in S$, and that player i should intervene (with impulse $\delta_i(x)$) only if $\mathcal{M}_i V_i(x) = V_i(x)$. This gives a heuristic formulation of Nash equilibria, provided that an explicit expression for V_i is available. The verification theorem will give a rigorous proof to this heuristic argument. We now characterize the value functions V_i .

Assume $V_1, V_2 \in C^2(S)$ (weaker conditions will be given later) and define

$$\mathcal{A}V_i = b \cdot \nabla V_i + \frac{1}{2} \text{tr}(\sigma \sigma^t D^2 V_i),$$

where b, σ are as in (2.1), σ^t denotes the transpose of σ and $\nabla V_i, D^2 V_i$ are the gradient and the Hessian matrix of V_i , respectively. We are interested in the following quasi-variational inequalities

(QVIs) for V_1, V_2 , where $i, j \in \{1, 2\}$ and $i \neq j$:

$$V_i = h_i, \quad \text{in } \partial S, \quad (3.3a)$$

$$\mathcal{M}_j V_j - V_j \leq 0, \quad \text{in } S, \quad (3.3b)$$

$$\mathcal{H}_i V_i - V_i = 0, \quad \text{in } \{\mathcal{M}_j V_j - V_j = 0\}, \quad (3.3c)$$

$$\max \{\mathcal{A}V_i - \rho_i V_i + f_i, \mathcal{M}_i V_i - V_i\} = 0, \quad \text{in } \{\mathcal{M}_j V_j - V_j < 0\}. \quad (3.3d)$$

Notice that there is a small abuse of notation in (3.3a), as V_i is not defined in ∂S , so that (3.3a) means $\lim_{y \rightarrow x} V_i(y) = h_i(x)$, for each $x \in \partial S$.

We now provide some intuition behind conditions (3.3a)-(3.3d). First of all, the terminal condition is obvious. Moreover, as we already noticed, (3.3b) is a standard condition in impulse control theory. For (3.3c), if player j intervenes (i.e., $\mathcal{M}_j V_j - V_j = 0$), by the definition of Nash equilibrium we expect that player i does not lose anything: this is equivalent to $\mathcal{H}_i V_i - V_i = 0$, otherwise it would be in her interest to deviate. On the contrary, if player j does not intervene (hence $\mathcal{M}_j V_j - V_j < 0$), then the problem for player i becomes a classical one-player impulse control one, V_i satisfies $\max \{\mathcal{A}V_i - \rho_i V_i + f_i, \mathcal{M}_i V_i - V_i\} = 0$. In short, the latter condition says that $\mathcal{A}V_i - \rho_i V_i + f_i \leq 0$, with equality in case of non-intervention (i.e., $\mathcal{M}_i V_i - V_i < 0$).

Remark 3.1. We notice that $\mathcal{A}V_i$ only appears in $\{\mathcal{M}_j V_j - V_j < 0\}$, so that V_i needs to be of class C^2 only in such a region (indeed, this assumption can be slightly relaxed, as we will see). This represents a difference to the one-player case, where the value function is usually required to be twice differentiable almost everywhere in S , see [15, Thm. 6.2].

The zero-sum case. A verification theorem will be provided in the next section. Here, as a preliminary check, we show that we are indeed generalizing the system of QVIs provided in [9], where the zero-sum case is considered. We show that, if we assume

$$\begin{aligned} f &:= f_1 = -f_2, & \phi &:= \phi_1 = -\phi_2, & \psi &:= \psi_1 = -\phi_2, \\ h &:= h_1 = -h_2, & Z &:= Z_1 = Z_2, & \Gamma &:= \Gamma^1 = \Gamma^2, \end{aligned} \quad (3.4)$$

so that $V := V_1 = -V_2$, then the problem in (3.3) reduces to the one considered in [9]. To shorten the equations, we assume $\rho_1 = \rho_2 = 0$ (this makes sense since in [9] a finite-horizon problem is considered). First of all, we define

$$\begin{aligned} \widetilde{\mathcal{M}}V(x) &:= \sup_{\delta \in Z} \{V(\Gamma(x, \delta)) + \phi(x, \delta)\}, \\ \widetilde{\mathcal{H}}V(x) &:= \inf_{\delta \in Z} \{V(\Gamma(x, \delta)) + \psi(x, \delta)\}, \end{aligned}$$

for each $x \in S$. It is easy to see that, under the conditions in (3.4), we have

$$\mathcal{M}_1 V_1 = \widetilde{\mathcal{M}}V, \quad \mathcal{M}_2 V_2 = -\widetilde{\mathcal{H}}V, \quad \mathcal{H}_1 V_1 = \widetilde{\mathcal{H}}V, \quad \mathcal{H}_2 V_2 = -\widetilde{\mathcal{M}}V,$$

so that problem (3.3) becomes

$$V = h, \quad \text{in } \partial S, \quad (3.5a)$$

$$\widetilde{\mathcal{M}}V \leq V \leq \widetilde{\mathcal{H}}V, \quad \text{in } S, \quad (3.5b)$$

$$\mathcal{A}V + f \leq 0, \quad \text{in } \{V = \widetilde{\mathcal{M}}V\}, \quad (3.5c)$$

$$\mathcal{A}V + f = 0, \quad \text{in } \{\widetilde{\mathcal{M}}V < V < \widetilde{\mathcal{H}}V\}, \quad (3.5d)$$

$$\mathcal{A}V + f \geq 0, \quad \text{in } \{V = \widetilde{\mathcal{H}}V\}. \quad (3.5e)$$

Simple computations, reported below, show that problem (3.5) is equivalent to

$$V = h, \quad \text{in } \partial S, \quad (3.6a)$$

$$\widetilde{\mathcal{M}}V - V \leq 0, \quad \text{in } S, \quad (3.6b)$$

$$\min\{\max\{\mathcal{A}V + f, \widetilde{\mathcal{M}}V - V\}, \widetilde{\mathcal{H}}V - V\} = 0, \quad \text{in } S, \quad (3.6c)$$

which is exactly the problem studied in [9], as anticipated. We conclude this section by proving the equivalence of (3.5) and (3.6).

Lemma 3.2. *Problems (3.5) and (3.6) are equivalent.*

Proof. Step 1. We prove that (3.5) implies (3.6). The only property to be proved is (3.6c). We consider three cases.

First, assume $V = \widetilde{\mathcal{M}}V$. Since $\mathcal{A}V + f \leq 0$ and $\widetilde{\mathcal{M}}V - V = 0$, we have $\max\{\mathcal{A}V + f, \widetilde{\mathcal{M}}V - V\} = 0$, which implies (3.6c) since $\widetilde{\mathcal{H}}V - V \geq 0$. Then, assume $\widetilde{\mathcal{M}}V < V < \widetilde{\mathcal{H}}V$. Since $\mathcal{A}V + f = 0$ and $\widetilde{\mathcal{M}}V - V < 0$, we have $\max\{\mathcal{A}V + f, \widetilde{\mathcal{M}}V - V\} = 0$, which implies (3.6c) since $\widetilde{\mathcal{H}}V - V > 0$. Finally, assume $V = \widetilde{\mathcal{H}}V$. Since $\mathcal{A}V + f \geq 0$ and $\widetilde{\mathcal{M}}V - V \leq 0$, we have $\max\{\mathcal{A}V + f, \widetilde{\mathcal{M}}V - V\} \geq 0$, which implies (3.6c) since $\widetilde{\mathcal{H}}V - V = 0$.

Step 2. We show that (3.6) implies (3.5). The only properties to be proved are (3.5c), (3.5d) and (3.5e). We assume $\widetilde{\mathcal{M}}V < \widetilde{\mathcal{H}}V$ (the case $\widetilde{\mathcal{M}}V = \widetilde{\mathcal{H}}V$ being immediate) and consider three cases.

First, assume $V = \widetilde{\mathcal{M}}V$. Since $\widetilde{\mathcal{H}}V - V > 0$, from (3.6c) it follows that $\max\{\mathcal{A}V + f, 0\} = 0$, which implies $\mathcal{A}V + f \leq 0$. Then, assume $\widetilde{\mathcal{M}}V < V < \widetilde{\mathcal{H}}V$. Since $\min\{\max\{\alpha, \beta\}, \gamma\} \in \{\alpha, \beta, \gamma\}$ for every $\alpha, \beta, \gamma \in \mathbb{R}$, and since $\widetilde{\mathcal{M}}V - V < 0 < \widetilde{\mathcal{H}}V - V$, from (3.6c) it follows that $\mathcal{A}V + f = 0$. Finally, assume $V = \widetilde{\mathcal{H}}V$. From (3.6c) it follows that $\max\{\mathcal{A}V + f, \widetilde{\mathcal{M}}V - V\} \geq 0$, which implies $\mathcal{A}V + f \geq 0$ since $\widetilde{\mathcal{M}}V - V < 0$. \square

3.2 Statement and proof

We provide here the main mathematical contribution of this paper, which is a verification theorem for the problems formalized in Section 2.

Theorem 3.3 (Verification theorem). *Let all the notations and working assumptions in Section 2 be in force and let V_i be a function from S to \mathbb{R} , with $i \in \{1, 2\}$. Assume that (3.1) holds and set $\mathcal{D}_i := \{\mathcal{M}_i V_i - V_i < 0\}$, with $\mathcal{M}_i V_i$ as in (3.2). Moreover, for $i \in \{1, 2\}$ assume that:*

- V_i is a solution to (3.3a)-(3.3d);
- $V_i \in C^2(\mathcal{D}_i \setminus \partial\mathcal{D}_i) \cap C^1(\mathcal{D}_i) \cap C(S)$ and it has polynomial growth;
- $\partial\mathcal{D}_i$ is a Lipschitz surface¹, and V_i has locally bounded derivatives up to the second order in any neighbourhood of $\partial\mathcal{D}_i$.

Finally, let $x \in S$ and assume that $(\varphi_1^*, \varphi_2^*) \in \mathcal{A}_x$, with

$$\varphi_i^* = (\mathcal{D}_i, \delta_i),$$

where $i \in \{1, 2\}$, the set \mathcal{D}_i is as above and the function δ_i is as in (3.1). Then,

$$(\varphi_1^*, \varphi_2^*) \text{ is a Nash equilibrium and } V_i(x) = J^i(x; \varphi_1^*, \varphi_2^*) \text{ for } i \in \{1, 2\}.$$

Remark 3.4. Practically, the Nash strategy is characterized as follows: player i intervenes only if the controlled process exits from the region $\{\mathcal{M}_i V_i - V_i < 0\}$ (equivalently, only if $\mathcal{M}_i V_i(x) = V_i(x)$, where x is the current state). When this happens, her impulse is $\delta_i(x)$.

Remark 3.5. We notice that, for the (candidate) optimal strategies in the theorem above, the properties in Lemma 2.3 imply what follows (the notation is heavy, but it will be crucial to understand the proof of the theorem):

$$(\mathcal{M}_1 V_1 - V_1)(X_s^{x; \varphi_1^*, \varphi_2^*}) < 0, \tag{3.7a}$$

$$(\mathcal{M}_2 V_2 - V_2)(X_s^{x; \varphi_1^*, \varphi_2^*}) < 0, \tag{3.7b}$$

$$\delta_{1,k}^{x; \varphi_1^*, \varphi_2^*} = \delta_1 \left(X_{\left(\tau_{1,k}^{x; \varphi_1^*, \varphi_2^*} \right)^-}^{x; \varphi_1^*, \varphi_2^*} \right), \tag{3.7c}$$

$$\delta_{2,k}^{x; \varphi_1^*, \varphi_2^*} = \delta_2 \left(X_{\left(\tau_{2,k}^{x; \varphi_1^*, \varphi_2^*} \right)^-}^{x; \varphi_1^*, \varphi_2^*} \right), \tag{3.7d}$$

¹I.e. it is locally the graph of a Lipschitz function.

$$(\mathcal{M}_1 V_1 - V_1) \left(X_{\left(\tau_{1,k}^{x;\varphi_1^*,\varphi_2^*} \right)^-} \right) = 0, \quad (3.7e)$$

$$(\mathcal{M}_2 V_2 - V_2) \left(X_{\left(\tau_{2,k}^{x;\varphi_1,\varphi_2^*} \right)^-} \right) = 0, \quad (3.7f)$$

for every strategies φ_1, φ_2 such that $(\varphi_1, \varphi_2^*), (\varphi_1^*, \varphi_2) \in \mathcal{A}_x$, every $s \geq 0$ and every $\tau_{i,k}^{x;\varphi_1,\varphi_2^*}, \tau_{i,k}^{x;\varphi_1^*,\varphi_2} < \infty$.

Proof. By Definition 2.6, we have to prove that

$$V_i(x) = J^i(x; \varphi_1^*, \varphi_2^*), \quad V_1(x) \geq J^1(x; \varphi_1, \varphi_2^*), \quad V_2(x) \geq J^2(x; \varphi_1^*, \varphi_2),$$

for every $i \in \{1, 2\}$ and (φ_1, φ_2) strategies such that $(\varphi_1, \varphi_2^*) \in \mathcal{A}_x$ and $(\varphi_1^*, \varphi_2) \in \mathcal{A}_x$. We show the results for V_1 and J^1 , the arguments for V_2 and J^2 being symmetric.

Step 1: $V_1(x) \geq J^1(x; \varphi_1, \varphi_2^*)$. Let φ_1 be a strategy for player 1 such that $(\varphi_1, \varphi_2^*) \in \mathcal{A}_x$. Here we will use the following shortened notation:

$$X = X^{x;\varphi_1,\varphi_2^*}, \quad \tau_{i,k} = \tau_{i,k}^{x;\varphi_1,\varphi_2^*}, \quad \delta_{i,k} = \delta_{i,k}^{x;\varphi_1,\varphi_2^*}.$$

Thanks to the regularity assumptions and by standard approximation arguments, it is not restrictive to assume $V_1 \in C^2(\mathcal{D}_2) \cap C(S)$ (see [15, Thm. 3.1]). For each $r > 0$ and $n \in \mathbb{N}$, we set

$$\tau_{r,n} = \tau_S \wedge \tau_r \wedge n,$$

where $\tau_r = \inf\{s > 0 : X_s \notin B(0, r)\}$ is the exit time from the ball with radius r . We apply Itô's formula to the process $e^{-\rho_1 t} V_1(X_t)$ over the interval $[0, \tau_{r,n}]$ and take the conditional expectations (the initial point and the stochastic integral are integrable, so that the other terms are integrable too by equality): we get

$$\begin{aligned} V_1(x) = \mathbb{E}_x \left[- \int_0^{\tau_{r,n}} e^{-\rho_1 s} (\mathcal{A}V_1 - \rho_1 V_1)(X_s) ds - \sum_{\tau_{1,k} < \tau_{r,n}} e^{-\rho_1 \tau_{1,k}} \left(V_1(X_{\tau_{1,k}}) - V_1(X_{(\tau_{1,k})^-}) \right) \right. \\ \left. - \sum_{\tau_{2,k} < \tau_{r,n}} e^{-\rho_1 \tau_{2,k}} \left(V_1(X_{\tau_{2,k}}) - V_1(X_{(\tau_{2,k})^-}) \right) + e^{-\rho_1 \tau_{r,n}} V_1(X_{\tau_{r,n}}) \right]. \end{aligned} \quad (3.8)$$

We now estimate each term in the right-hand side of (3.8). As for the first term, since $(\mathcal{M}_2 V_2 - V_2)(X_s) < 0$ by (3.7b), from (3.3d) it follows that

$$(\mathcal{A}V_1 - \rho_1 V_1)(X_s) \leq -f_1(X_s), \quad (3.9)$$

for all $s \in [0, \tau_S]$. Let us now consider the second term: by (3.3b) and the definition of $\mathcal{M}_1 V_1$ in (3.2), for every stopping time $\tau_{1,k} < \tau_S$ we have

$$\begin{aligned} V_1(X_{(\tau_{1,k})^-}) &\geq \mathcal{M}_1 V_1(X_{(\tau_{1,k})^-}) \\ &= \sup_{\delta \in Z_1} \{ V_1(\Gamma^1(X_{(\tau_{1,k})^-}, \delta)) + \phi_1(X_{(\tau_{1,k})^-}, \delta) \} \\ &\geq V_1(\Gamma^1(X_{(\tau_{1,k})^-}, \delta_{1,k})) + \phi_1(X_{(\tau_{1,k})^-}, \delta_{1,k}) \\ &= V_1(X_{\tau_{1,k}}) + \phi_1(X_{(\tau_{1,k})^-}, \delta_{1,k}). \end{aligned} \quad (3.10)$$

As for the third term, let us consider any stopping time $\tau_{2,k} < \tau_S$. By (3.7f) we have $(\mathcal{M}_2 V_2 - V_2)(X_{(\tau_{2,k})^-}) = 0$; hence, the condition in (3.3c), the definition of $\mathcal{H}_1 V_1$ in (3.2) and the expression of $\delta_{2,k}$ in (3.7d) imply that

$$\begin{aligned} V_1(X_{(\tau_{2,k})^-}) &= \mathcal{H}_1 V_1(X_{(\tau_{2,k})^-}) \\ &= V_1(\Gamma^2(X_{(\tau_{2,k})^-}, \delta_2(X_{(\tau_{2,k})^-}))) + \psi_1(X_{(\tau_{2,k})^-}, \delta_2(X_{(\tau_{2,k})^-})) \\ &= V_1(\Gamma^2(X_{(\tau_{2,k})^-}, \delta_{2,k})) + \psi_1(X_{(\tau_{2,k})^-}, \delta_{2,k}) \\ &= V_1(X_{\tau_{2,k}}) + \psi_1(X_{(\tau_{2,k})^-}, \delta_{2,k}). \end{aligned} \quad (3.11)$$

By (3.8) and the estimates in (3.9)-(3.11) it follows that

$$V_1(x) \geq \mathbb{E}_x \left[\int_0^{\tau_{r,n}} e^{-\rho_1 s} f_1(X_s) ds + \sum_{\tau_{1,k} < \tau_{r,n}} e^{-\rho_1 \tau_{1,k}} \phi_1(X_{(\tau_{1,k})^-}, \delta_{1,k}) \right. \\ \left. + \sum_{\tau_{2,k} < \tau_{r,n}} e^{-\rho_1 \tau_{2,k}} \psi_1(X_{(\tau_{2,k})^-}, \delta_{2,k}) + e^{-\rho_1 \tau_{r,n}} V_1(X_{\tau_{r,n}}) \right].$$

Thanks to the conditions in (2.8), (2.9) and the polynomial growth of V_1 , we can use the dominated convergence theorem and pass to the limit, first as $r \rightarrow \infty$ and then as $n \rightarrow \infty$. In particular, for the fourth term we notice that

$$V_1(X_{\tau_{r,n}}) \leq C(1 + |X_{\tau_{r,n}}|^p) \leq C(1 + \|X\|_\infty^p) \in L^1(\Omega), \quad (3.12)$$

for suitable constants $C > 0$ and $p \in \mathbb{N}$; the corresponding limit immediately follows by the continuity of V_1 in the case $\tau_S < \infty$ and by (3.12) itself in the case $\tau_S = \infty$ (as a direct consequence of (2.9), we have $\|X\|_\infty^p < \infty$ a.s.). Hence, we finally get

$$V_1(x) \geq \mathbb{E}_x \left[\int_0^{\tau_S} e^{-\rho_1 s} f_1(X_s) ds + \sum_{\tau_{1,k} < \tau_S} e^{-\rho_1 \tau_{1,k}} \phi_1(X_{(\tau_{1,k})^-}, \delta_{1,k}) \right. \\ \left. + \sum_{\tau_{2,k} < \tau_S} e^{-\rho_1 \tau_{2,k}} \psi_1(X_{(\tau_{2,k})^-}, \delta_{2,k}) + e^{-\rho_1 \tau_S} h_1(X_{(\tau_S)^-}) \mathbb{1}_{\{\tau_S < +\infty\}} \right] = J^1(x; \varphi_1, \varphi_2^*).$$

Step 2: $V_1(x) = J^1(x; \varphi_1^*, \varphi_2^*)$. We argue as in Step 1, but here all the inequalities are equalities by the properties of φ_1^* . \square

As already noticed in Remark 3.1, we stress that, unlike one-player impulse control problems, in our verification theorem the candidates are not required to be twice differentiable everywhere. For example, consider the case of player 1: as in the proof we always consider pairs of strategies in the form (φ_1, φ_2^*) , by (3.7b) the controlled process never exits from $\mathcal{D}_2 = \{\mathcal{M}_2 V_2 - V_2 < 0\}$, which is then the only region where the function V_1 needs to be (almost everywhere) twice differentiable in order to apply Itô's formula.

We conclude this section with some considerations on how the theorem above will typically be used. First, when solving the system of QVIs, one deals with functions which are defined only piecewise, as it will be clear in the next section. Then, the regularity assumptions in the verification theorem will give us suitable pasting conditions, leading to a system of algebraic equations. If the regularity conditions are too strong, the system has more equations than parameters, making the application of the theorem more difficult. Hence, a crucial point when stating a verification theorem is to set regularity conditions giving a solvable system of equations. In Section 4 we show that, in an example of one-dimensional impulse game, the regularity conditions actually lead to a well-posed algebraic system having a unique solution.

4 A solvable one-dimensional impulse game

In this section we provide an application of the Verification Theorem 3.3 to an impulse game with a one-dimensional state variable modelled by a (scaled) Brownian motion, that can be shifted due to the interventions of two players with linear payoffs. We find a Nash equilibrium for such a game and provide explicit expressions for the value functions and for the optimal strategies at equilibrium.

4.1 Formulation of the problem

We consider a one-dimensional real process X and two players with opposite goals: player 1 prefers a high value for the process X , whereas the goal of player 2 is to force X to take a low value. More precisely, if x denotes the current value of the process, we assume that the running payoffs of the two players are given by

$$f_1(x) = x - s_1, \quad f_2(x) = s_2 - x, \quad s_1 < s_2,$$

where s_1, s_2 are fixed (possibly negative) constants.

We assume that each player can intervene and shift X from state x to state $x + \delta$, with $\delta \in \mathbb{R}$. Moreover, when none of the players intervenes, we assume that X follows a (scaled) Brownian motion. Hence, if x denotes the initial state and $u_i = \{(\tau_{i,k}, \delta_{i,k})\}_{k \geq 1}$ collects the intervention times and the corresponding impulses of player $i \in \{1, 2\}$, we have

$$X_s = X_s^{x; u_1, u_2} = x + \sigma W_s + \sum_{k: \tau_{1,k} \leq s} \delta_{1,k} + \sum_{k: \tau_{2,k} \leq s} \delta_{2,k}, \quad s \geq 0,$$

where W is a standard one-dimensional Brownian motion and $\sigma > 0$ is a fixed parameter.

As player 2 aims at lowering the level, we can assume that her impulses are negative: $\delta_{2,k} < 0$, for every $k \in \mathbb{N}$. Similarly, we assume $\delta_{1,k} > 0$, for every $k \in \mathbb{N}$. Affecting the process has a cost for the intervening player and we also assume that there is a corresponding gain for the opponent. In our model both intervention penalties and gains consist in a fixed cost and in a variable cost, assumed to be proportional to the absolute value of the impulse: if ϕ_i denotes the intervention penalty for player i and ψ_j denotes the corresponding gain for player j , we assume

$$\phi_i(\delta) = -c - \lambda|\delta|, \quad \psi_j(\delta) = \tilde{c} + \tilde{\lambda}|\delta|,$$

where $\delta \in \mathbb{R}$ is the impulse corresponding to the intervention of player i and $c, \tilde{c}, \lambda, \tilde{\lambda}$ are fixed constants such that

$$c \geq \tilde{c} \geq 0, \quad \lambda \geq \tilde{\lambda} \geq 0, \quad (c, \lambda) \neq (\tilde{c}, \tilde{\lambda}).$$

The order conditions have this justification: if we had $c < \tilde{c}$ or $\lambda < \tilde{\lambda}$, then, for a suitable impulse δ , the two players could realize a mutual gain by an (almost) instantaneous double intervention; by iterating this infinitely often in a finite interval, the two value functions would diverge (this phenomenon is analogous to the one already present in [11] for stopping games). The condition $(c, \lambda) \neq (\tilde{c}, \tilde{\lambda})$ will be explained in Remark 4.5 and Section 4.4. Finally, we assume

$$1 - \lambda\rho > 0,$$

where ρ is the discount rate, the same one for both the players.

This problem clearly belongs to the class described in Section 2, with

$$d = 1, \quad S = \mathbb{R}, \quad \Gamma^i(x, \delta) = x + \delta, \quad \rho_i = \rho, \quad Z_1 =]0, \infty[, \quad Z_2 =]-\infty, 0],$$

and with f_i, ϕ_i, ψ_i as above. In short, if $\varphi_i = (A_i, \xi_i)$ denotes the strategy of player i , the objective functions are

$$\begin{aligned} J^1(x; \varphi_1, \varphi_2) &:= \mathbb{E}_x \left[\int_0^\infty e^{-\rho s} (X_s - s_1) ds - \sum_{k \geq 1} e^{-\rho \tau_{1,k}} (c + \lambda|\delta_{1,k}|) + \sum_{k \geq 1} e^{-\rho \tau_{2,k}} (\tilde{c} + \tilde{\lambda}|\delta_{2,k}|) \right], \\ J^2(x; \varphi_1, \varphi_2) &:= \mathbb{E}_x \left[\int_0^\infty e^{-\rho s} (s_2 - X_s) ds - \sum_{k \geq 1} e^{-\rho \tau_{2,k}} (c + \lambda|\delta_{2,k}|) + \sum_{k \geq 1} e^{-\rho \tau_{1,k}} (\tilde{c} + \tilde{\lambda}|\delta_{1,k}|) \right], \end{aligned}$$

where $\{(\tau_{i,k}, \delta_{i,k})\}_{k \geq 1}$ denotes the impulse controls of player i associated to the strategies φ_1, φ_2 .

As already outlined, the players have different goals: we are going to investigate if a Nash equilibrium for such a problem exists. Indeed, since $s_1 < s_2$ both the players gain in the interval $[s_1, s_2]$, it seems that there is room for a Nash configuration. If a Nash equilibrium exists, we denote by $V_1(x), V_2(x)$ the corresponding value of the game with initial state $x \in \mathbb{R}$.

Interpretations. We provide two possible interpretations of the game just described.

First, let X denote the exchange rate between two currencies. The central banks of the corresponding countries (the players) have different targets for the rate: player 1 prefers a high value for X , while the goal of player 2 is yielding a low value. To have a tractable model, it is reasonable to assume that the payoffs of the two players are given, respectively, by $X - s_1$ and $s_2 - X$, where s_1, s_2 are fixed constants with $s_2 > s_1$, which leads to the one-dimensional game defined in this section. This interpretation (detailed in [1, Sect. 3.3]) corresponds to a two-player version of the model introduced and studied in, e.g., [4] and [5].

The second possible interpretation is related to energy markets. Let the process X model the forward price of energy. Player 1 is an energy producer and s_1 is the unitary production cost, so that, in a simplified model, her payoff is $X - s_1$. On the other hand, player 2 runs a company, which buys energy at price X and sells her products at some price s_2 , with $s_2 > s_1$, so that her payoff is $s_2 - X$. Since player 2 consumes a high volume of energy, she can be considered as a big market player; so, both the players affect the price of energy, and we get the nonzero-sum impulse game studied in this section (both the players have enough impact on the economy to obtain a financial compensation from the state to mitigate sudden adverse movements of the forward price).

4.2 Looking for candidates for the value functions

Our goal is to use the Verification Theorem 3.3. We start by looking for a solution to the problem in (3.3), in order to get a couple of candidates \tilde{V}_1, \tilde{V}_2 for the value functions V_1, V_2 .

First, consider the two equations in the QVI problem (3.3), that is

$$\begin{aligned} \mathcal{H}_i \tilde{V}_i - \tilde{V}_i &= 0, & \text{in } \{\mathcal{M}_j \tilde{V}_j - \tilde{V}_j = 0\}, \\ \max \{\mathcal{A} \tilde{V}_i - \rho \tilde{V}_i + f_i, \mathcal{M}_i \tilde{V}_i - \tilde{V}_i\} &= 0, & \text{in } \{\mathcal{M}_j \tilde{V}_j - \tilde{V}_j < 0\}, \end{aligned}$$

for $i, j \in \{1, 2\}$, with $i \neq j$; this suggests the following representation for \tilde{V}_i :

$$\tilde{V}_i(x) = \begin{cases} \mathcal{M}_i \tilde{V}_i(x), & \text{in } \{\mathcal{M}_i \tilde{V}_i - \tilde{V}_i = 0\}, \\ \varphi_i(x), & \text{in } \{\mathcal{M}_i \tilde{V}_i - \tilde{V}_i < 0, \mathcal{M}_j \tilde{V}_j - \tilde{V}_j < 0\}, \\ \mathcal{H}_i \tilde{V}_i(x), & \text{in } \{\mathcal{M}_j \tilde{V}_j - \tilde{V}_j = 0\}, \end{cases} \quad (4.1)$$

for $i \in \{1, 2\}$ and $x \in \mathbb{R}$, where φ_i is a solution to

$$\mathcal{A} \varphi_i - \rho \varphi_i + f_i = \frac{\sigma^2}{2} \varphi_i'' - \rho \varphi_i + f_i = 0.$$

Notice that an explicit formula for φ_i is available: for each $x \in \mathbb{R}$, we have

$$\begin{aligned} \varphi_1(x) &= \varphi_1^{A_{11}, A_{12}}(x) = A_{11} e^{\theta x} + A_{12} e^{-\theta x} + (x - s_1)/\rho, \\ \varphi_2(x) &= \varphi_2^{A_{21}, A_{22}}(x) = A_{21} e^{\theta x} + A_{22} e^{-\theta x} + (s_2 - x)/\rho, \end{aligned} \quad (4.2)$$

where A_{ij} are real parameters and the parameter θ is defined by

$$\theta = \sqrt{\frac{2\rho}{\sigma^2}}.$$

In order to go on, we need to guess an expression for the intervention regions. As the goal of player 1 is to keep a high value for the process, it is reasonable to assume that her intervention region is in the form $] - \infty, \bar{x}_1]$, for some threshold \bar{x}_1 . For a similar reason, we expect the intervention region of player 2 to be in the form $[\bar{x}_2, +\infty[$, for some other threshold \bar{x}_2 . Since $s_1 < s_2$, we guess that $\bar{x}_1 < \bar{x}_2$; as a consequence, the real line is heuristically partitioned into three intervals:

$$\begin{aligned}] - \infty, \bar{x}_1] &= \{\mathcal{M}_1 \tilde{V}_1 - \tilde{V}_1 = 0\}, \text{ where player 1 intervenes,} \\]\bar{x}_1, \bar{x}_2[&= \{\mathcal{M}_1 \tilde{V}_1 - \tilde{V}_1 < 0\} \cap \{\mathcal{M}_2 \tilde{V}_2 - \tilde{V}_2 < 0\}, \text{ where no one intervenes,} \\ [\bar{x}_2, +\infty[&= \{\mathcal{M}_2 \tilde{V}_2 - \tilde{V}_2 = 0\}, \text{ where player 2 intervenes.} \end{aligned}$$

By the representation (4.1), this leads to the following expressions for \tilde{V}_1 and \tilde{V}_2 :

$$\tilde{V}_1(x) = \begin{cases} \mathcal{M}_1 \tilde{V}_1(x), & \text{if } x \in] - \infty, \bar{x}_1], \\ \varphi_1(x), & \text{if } x \in]\bar{x}_1, \bar{x}_2[, \\ \mathcal{H}_1 \tilde{V}_1(x), & \text{if } x \in [\bar{x}_2, +\infty[, \end{cases} \quad \tilde{V}_2(x) = \begin{cases} \mathcal{H}_2 \tilde{V}_2(x), & \text{if } x \in] - \infty, \bar{x}_1], \\ \varphi_2(x), & \text{if } x \in]\bar{x}_1, \bar{x}_2[, \\ \mathcal{M}_2 \tilde{V}_2(x), & \text{if } x \in [\bar{x}_2, +\infty[. \end{cases}$$

Let us now investigate the form of $\mathcal{M}_i \tilde{V}_i$ and $\mathcal{H}_i \tilde{V}_i$. Recall that the impulses of player 1 (resp. player 2) are positive (resp. negative); then, we have

$$\begin{aligned} \mathcal{M}_1 \tilde{V}_1(x) &= \sup_{\delta \geq 0} \{\tilde{V}_1(x + \delta) - c - \lambda \delta\} = \sup_{y \geq x} \{\tilde{V}_1(y) - c - \lambda(y - x)\}, \\ \mathcal{M}_2 \tilde{V}_2(x) &= \sup_{\delta \leq 0} \{\tilde{V}_2(x + \delta) - c - \lambda(-\delta)\} = \sup_{y \leq x} \{\tilde{V}_2(y) - c - \lambda(x - y)\}. \end{aligned}$$

It is reasonable to assume that the maximum point of the function $y \mapsto \tilde{V}_1(y) - \lambda y$ (resp. $y \mapsto \tilde{V}_2(y) + \lambda y$) exists, is unique and belongs to the common continuation region $]\bar{x}_1, \bar{x}_2[$, where we have $\tilde{V}_1 = \varphi_1$ (resp. $\tilde{V}_2 = \varphi_2$). As a consequence, if we denote by x_i^* , $i \in \{1, 2\}$, such maximum points, that is

$$\begin{aligned}\varphi_1(x_1^*) &= \max_{y \in]\bar{x}_1, \bar{x}_2[} \{\varphi_1(y) - \lambda y\}, & \text{i.e.} \quad \varphi_1'(x_1^*) &= \lambda, \quad \varphi_1''(x_1^*) \leq 0, \quad \bar{x}_1 < x_1^* < \bar{x}_2, \\ \varphi_2(x_2^*) &= \max_{y \in]\bar{x}_1, \bar{x}_2[} \{\varphi_2(y) + \lambda y\}, & \text{i.e.} \quad \varphi_2'(x_2^*) &= -\lambda, \quad \varphi_2''(x_2^*) \leq 0, \quad \bar{x}_1 < x_2^* < \bar{x}_2,\end{aligned}$$

the functions $\mathcal{M}_i \tilde{V}_i$, $\mathcal{H}_i \tilde{V}_i$ have the following (heuristic, at the moment) expression:

$$\begin{aligned}\mathcal{M}_1 \tilde{V}_1(x) &= \varphi_1(x_1^*) - c - \lambda(x_1^* - x), & \mathcal{M}_2 \tilde{V}_2(x) &= \varphi_2(x_2^*) - c - \lambda(x - x_2^*), \\ \mathcal{H}_1 \tilde{V}_1(x) &= \varphi_1(x_2^*) + \tilde{c} + \tilde{\lambda}(x - x_2^*), & \mathcal{H}_2 \tilde{V}_2(x) &= \varphi_2(x_1^*) + \tilde{c} + \tilde{\lambda}(x_1^* - x).\end{aligned}$$

As for the parameters involved in \tilde{V}_1, \tilde{V}_2 , they must be chosen so as to satisfy the regularity assumptions in the verification theorem, which here write

$$\begin{aligned}\tilde{V}_1 &\in C^2(]-\infty, \bar{x}_1[\cup]\bar{x}_1, \bar{x}_2[) \cap C^1(]-\infty, \bar{x}_2[) \cap C(\mathbb{R}), \\ \tilde{V}_2 &\in C^2(]\bar{x}_1, \bar{x}_2[\cup]\bar{x}_2, +\infty[) \cap C^1(]\bar{x}_1, +\infty[) \cap C(\mathbb{R}).\end{aligned}$$

Since \tilde{V}_1 and \tilde{V}_2 are, by definition, smooth in $]-\infty, \bar{x}_1[\cup]\bar{x}_1, \bar{x}_2[\cup]\bar{x}_2, +\infty[$, we have to set the parameters so that \tilde{V}_i is continuous at \bar{x}_1, \bar{x}_2 and differentiable at \bar{x}_i (we underline that \tilde{V}_1 and \tilde{V}_2 might not be differentiable at, respectively, \bar{x}_2 and \bar{x}_1).

Finally, to summarize all the previous arguments, our candidates for the value functions are defined as follows.

Definition 4.1. *For every $x \in \mathbb{R}$, we set*

$$\begin{aligned}\tilde{V}_1(x) &= \begin{cases} \varphi_1(x_1^*) - c - \lambda(x_1^* - x), & \text{if } x \in]-\infty, \bar{x}_1[, \\ \varphi_1(x), & \text{if } x \in]\bar{x}_1, \bar{x}_2[, \\ \varphi_1(x_2^*) + \tilde{c} + \tilde{\lambda}(x - x_2^*), & \text{if } x \in [\bar{x}_2, +\infty[, \end{cases} \\ \tilde{V}_2(x) &= \begin{cases} \varphi_2(x_1^*) + \tilde{c} + \tilde{\lambda}(x_1^* - x), & \text{if } x \in]-\infty, \bar{x}_1[, \\ \varphi_2(x), & \text{if } x \in]\bar{x}_1, \bar{x}_2[, \\ \varphi_2(x_2^*) - c - \lambda(x - x_2^*), & \text{if } x \in [\bar{x}_2, +\infty[, \end{cases}\end{aligned}\tag{4.3}$$

where $\varphi_1 = \varphi_1^{A_{11}, A_{12}}$, $\varphi_2 = \varphi_2^{A_{21}, A_{22}}$ and the eight parameters involved

$$(A_{11}, A_{12}, A_{21}, A_{22}, \bar{x}_1, \bar{x}_2, x_1^*, x_2^*)$$

satisfy the order conditions

$$\bar{x}_1 < x_1^* < \bar{x}_2, \quad \bar{x}_1 < x_2^* < \bar{x}_2,\tag{4.4}$$

and the following conditions:

$$\begin{cases} \varphi_1'(x_1^*) = \lambda & \text{and } \varphi_1''(x_1^*) \leq 0, & (\text{optimality of } x_1^*) \\ \varphi_1'(\bar{x}_1) = \lambda, & (C^1\text{-pasting in } \bar{x}_1) \\ \varphi_1(\bar{x}_1) = \varphi_1(x_1^*) - c - \lambda(x_1^* - \bar{x}_1), & (C^0\text{-pasting in } \bar{x}_1) \\ \varphi_1(\bar{x}_2) = \varphi_1(x_2^*) + \tilde{c} + \tilde{\lambda}(\bar{x}_2 - x_2^*), & (C^0\text{-pasting in } \bar{x}_2) \end{cases}\tag{4.5}$$

$$\begin{cases} \varphi_2'(x_2^*) = -\lambda & \text{and } \varphi_2''(x_2^*) \leq 0, & (\text{optimality of } x_2^*) \\ \varphi_2'(\bar{x}_2) = -\lambda, & (C^1\text{-pasting in } \bar{x}_2) \\ \varphi_2(\bar{x}_1) = \varphi_2(x_1^*) + \tilde{c} + \tilde{\lambda}(x_1^* - \bar{x}_1), & (C^0\text{-pasting in } \bar{x}_1) \\ \varphi_2(\bar{x}_2) = \varphi_2(x_2^*) - c - \lambda(\bar{x}_2 - x_2^*). & (C^0\text{-pasting in } \bar{x}_2) \end{cases}\tag{4.6}$$

In order to have a well-posed definition, we need to show that the conditions in (4.4)-(4.5)-(4.6) actually admit a unique solution.

Proposition 4.2. *There exists a unique 8-uple $(A_{11}, A_{12}, A_{21}, A_{22}, \bar{x}_1, \bar{x}_2, x_1^*, x_2^*)$ which satisfies the conditions in (4.4)-(4.5)-(4.6). Moreover, there exists $\tilde{x} \in]x_2^*, \bar{x}_2[$ such that $\varphi_2'' < 0$ in $]\bar{x}_1, \tilde{x}[$ and $\varphi_2'' > 0$ in $]\tilde{x}, \bar{x}_2[$.*

Proof. We just focus on the existence of solutions: uniqueness will immediately follow from Proposition 4.7 in the next section (different solutions would imply different expressions for the value functions).

First, we reduce the number of equations. Notice that the running costs f_i are symmetric with respect to $\tilde{s} := (s_1 + s_2)/2$; we guess that the couples (\bar{x}_1, \bar{x}_2) , (x_1^*, x_2^*) and (φ_1, φ_2) have the same property, that is

$$\bar{x}_1 = 2\tilde{s} - \bar{x}_2, \quad x_1^* = 2\tilde{s} - x_2^*, \quad A_{11} = A_{22}e^{-2\theta\tilde{s}}, \quad A_{12} = A_{21}e^{2\theta\tilde{s}}. \quad (4.7)$$

Under condition (4.7), the systems in (4.5) and (4.6) are independent and equivalent: namely, the 4-uple $(A_{11}, A_{12}, \bar{x}_1, x_1^*)$ solves (4.5) if and only if $(A_{21}, A_{22}, \bar{x}_2, x_2^*)$, defined by (4.7), is a solution to (4.6). Hence, we just need to solve one of the two systems of equations: we decide to focus on (4.6), which, by the change of variable

$$\bar{y} = e^{\theta(\bar{x}_2 - \tilde{s})}, \quad y^* = e^{\theta(x_2^* - \tilde{s})}, \quad A_1 = 2\theta A_{21}e^{\theta\tilde{s}}, \quad A_2 = 2\theta A_{22}e^{-\theta\tilde{s}} \quad (4.8)$$

and by plugging (4.6a)-(4.6b) into (4.6c)-(4.6d), reads (we set $\eta = (1 - \lambda\rho)/\rho$, notice that $\eta > 0$)

$$\begin{cases} A_1(y^*)^2 - 2\eta y^* - A_2 = 0, & \text{with } y^* > 0 \text{ and } A_1 y^* - \eta \leq 0, \\ A_1 \bar{y}^2 - 2\eta \bar{y} - A_2 = 0, & \text{with } \bar{y} > 0, \\ (A_1 + A_2)^2(\bar{y} - y^*) + 2A_2[\theta(c - \tilde{c}) + (\lambda - \tilde{\lambda})\log(\bar{y}/y^*)] = 0, \\ A_1(\bar{y} - y^*) + \theta c - \eta \log(\bar{y}/y^*) = 0. \end{cases} \quad (4.9)$$

Also recall the order condition (4.4), which now reads

$$1 < \bar{y}y^* < \bar{y}^2. \quad (4.10)$$

We now prove that there exists a unique solution (A_1, A_2, \bar{y}, y^*) to (4.9)-(4.10). Given a fixed pair $(A_1, A_2) \in \mathcal{A}$, where

$$\mathcal{A} = \{(A_1, A_2) : A_1 > 0, \quad A_2 < 0, \quad \eta^2 + A_1 A_2 > 0\},$$

there exists a unique solution to (4.9a)-(4.9b), given by

$$\bar{y}(A_1, A_2) = \frac{\eta + \sqrt{\eta^2 + A_1 A_2}}{A_1}, \quad y^*(A_1, A_2) = \frac{\eta - \sqrt{\eta^2 + A_1 A_2}}{A_1}. \quad (4.11)$$

To conclude, we just need to prove that there exists a unique pair (A_1, A_2) such that

$$(A_1, A_2) \in \mathcal{A} \text{ and } (A_1, A_2, \bar{y}(A_1, A_2), y^*(A_1, A_2)) \text{ is a solution to (4.9c)-(4.9d)-(4.10),} \quad (4.12)$$

that is, by the expressions in (4.11), such that

$$\begin{cases} (A_1 + A_2)^2 \sqrt{\eta^2 + A_1 A_2} + A_1 A_2 \left[\theta(c - \tilde{c}) + (\lambda - \tilde{\lambda}) \log \left(\frac{\eta + \sqrt{\eta^2 + A_1 A_2}}{\eta - \sqrt{\eta^2 + A_1 A_2}} \right) \right] = 0, \\ 2\sqrt{\eta^2 + A_1 A_2} + \theta c - \eta \log \left(\frac{\eta + \sqrt{\eta^2 + A_1 A_2}}{\eta - \sqrt{\eta^2 + A_1 A_2}} \right) = 0, \\ A_1 + A_2 < 0, \\ A_1 > 0, \quad A_2 < 0, \quad \eta^2 + A_1 A_2 > 0. \end{cases} \quad (4.13)$$

For $x \in (0, \eta)$, define the function

$$F(x) = 2x + \theta c - \eta \log \left(\frac{\eta + x}{\eta - x} \right). \quad (4.14)$$

Since $F(0^+) = \theta c > 0$, $F(\eta^-) = -\infty$ and $F' < 0$, there exists a unique $\xi \in (0, \eta)$ such that $F(\xi) = 0$, so that we can rewrite (4.13b) as $\sqrt{\eta^2 + A_1 A_2} = \xi$. Consequently, (4.13) is equivalent to

$$\begin{cases} A_1 A_2 = -M, \\ A_1 + A_2 = -2N, \\ A_1 > 0, \quad A_2 < 0, \end{cases} \quad \text{with } M = \eta^2 - \xi^2 \text{ and } N = \sqrt{\frac{(\eta^2 - \xi^2)[\theta\eta(c - \tilde{c}) + (\lambda - \tilde{\lambda})(2\xi + \theta c)]}{4\eta\xi}}, \quad (4.15)$$

which trivially has a unique solution (notice that $N^2 + M > 0$), namely

$$A_1 = -N + \sqrt{N^2 + M}, \quad A_2 = -N - \sqrt{N^2 + M}. \quad (4.16)$$

Finally, it is immediate to see that $\varphi_2'' < 0$ in $] -\infty, \tilde{x}[$ and $\varphi_2'' > 0$ in $]\tilde{x}, +\infty[$, for a suitable $\tilde{x} \in \mathbb{R}$. By the change of variable, $\varphi_2''(\bar{x}_2) > 0$ (resp. $\varphi_2''(x_2^*) < 0$) if and only if $A_1 \bar{y} - \eta > 0$ (resp. $A_1 y^* - \eta < 0$), which is trivially true; hence, $\tilde{x} \in]x_2^*, \bar{x}_2[$. \square

Remark 4.3. From the proof of Proposition 4.2, we see that the systems (4.5)-(4.6) have more than one solution, but only one satisfies the order condition (4.4). In particular, we notice that the other solution of (4.6) is, in the new variables of the proofs, $\tilde{A}_1 = -A_2$, $\tilde{A}_2 = -A_1$, $\tilde{y}^* = -\bar{y}$, $\tilde{\bar{y}} = -y^*$.

Remark 4.4. By combining (4.7), (4.8), (4.11) and (4.16), we can get (semi-)explicit formulas for the 8-uple $(A_{11}, A_{12}, A_{21}, A_{22}, \bar{x}_1, \bar{x}_2, x_1^*, x_2^*)$ which solves (4.4)-(4.5)-(4.6): namely,

$$\begin{aligned} \bar{x}_i &= \tilde{s} + \frac{(-1)^i}{\theta} \log \left[\sqrt{\frac{\eta + \xi}{\eta - \xi}} \left(\sqrt{\Gamma + 1} + \sqrt{\Gamma} \right) \right], \\ x_i^* &= \tilde{s} + \frac{(-1)^i}{\theta} \log \left[\sqrt{\frac{\eta - \xi}{\eta + \xi}} \left(\sqrt{\Gamma + 1} + \sqrt{\Gamma} \right) \right], \\ A_{ij} &= e^{(-1)^j \theta \tilde{s}} \frac{\sqrt{\eta^2 - \xi^2}}{2\theta} \left((-1)^{i+j+1} \sqrt{\Gamma + 1} - \sqrt{\Gamma} \right), \end{aligned} \quad (4.17)$$

for $i, j \in \{1, 2\}$, where $\xi = \xi(c, \theta, \eta) \in (0, \eta)$ is the unique zero of the function F in (4.14) and the coefficients are defined by

$$\tilde{s} = \frac{s_1 + s_2}{2}, \quad \theta = \sqrt{\frac{2\rho}{\sigma^2}}, \quad \eta = \frac{1 - \lambda\rho}{\rho}, \quad \Gamma = \frac{\theta(c - \tilde{c})}{4\xi} + \frac{\theta c(\lambda - \tilde{\lambda})}{4\eta\xi} + \frac{\lambda - \tilde{\lambda}}{2\eta}. \quad (4.18)$$

Also, notice that (4.7) implies that $\tilde{V}_1(x) = \tilde{V}_2(2\tilde{s} - x)$, for $x \in \mathbb{R}$, i.e. the functions \tilde{V}_i are symmetric with respect to \tilde{s} .

Remark 4.5. From (4.17), we notice that $\tilde{x}_i = x_j^*$ when $\Gamma = 0$, which happens if and only if $(c, \lambda) = (\tilde{c}, \tilde{\lambda})$. This situation gives rise to a degenerate solution, where players intervene infinitely often in each instant. We analyze this in more detail in Section 4.4, where we study the case where $\lambda = \tilde{\lambda}$ and $c \rightarrow \tilde{c}^+$.

4.3 Application of the verification theorem

We now apply the Verification Theorem 3.3 and prove that the candidates \tilde{V}_1, \tilde{V}_2 in Definition 4.1 actually coincide with the value functions V_1, V_2 of the problem described in Section 4.1. We refer the reader to Section 3.1 for the definition of the functions $\delta_1, \delta_2, \mathcal{M}_1, \mathcal{M}_2$ used in the following lemma.

Lemma 4.6. *Let \tilde{V}_1, \tilde{V}_2 be as in Definition 4.1. For every $x \in \mathbb{R}$ we have*

$$\delta_1(x) = \begin{cases} x_1^* - x, & \text{in }] -\infty, x_1^*], \\ 0, & \text{in }]x_1^*, +\infty[, \end{cases} \quad \delta_2(x) = \begin{cases} 0, & \text{in }] -\infty, x_2^*[, \\ x_2^* - x, & \text{in }]x_2^*, +\infty[. \end{cases} \quad (4.19)$$

Moreover, we have

$$\begin{aligned} \mathcal{M}_1 \tilde{V}_1 - \tilde{V}_1 &\leq 0, & \{\mathcal{M}_1 \tilde{V}_1 - \tilde{V}_1 < 0\} &=]\bar{x}_1, +\infty[, & \{\mathcal{M}_1 \tilde{V}_1 - \tilde{V}_1 = 0\} &=]-\infty, \bar{x}_1], \\ \mathcal{M}_2 \tilde{V}_2 - \tilde{V}_2 &\leq 0, & \{\mathcal{M}_2 \tilde{V}_2 - \tilde{V}_2 < 0\} &=]-\infty, \bar{x}_2[, & \{\mathcal{M}_2 \tilde{V}_2 - \tilde{V}_2 = 0\} &= [\bar{x}_2, +\infty[. \end{aligned} \quad (4.20)$$

Proof. We give the proof only for δ_2 and $\mathcal{M}_2 \tilde{V}_2$, the arguments for δ_1 and $\mathcal{M}_1 \tilde{V}_1$ being the same. For every $x \in \mathbb{R}$, we have

$$\mathcal{M}_2 \tilde{V}_2(x) = \max_{\delta_2 \leq 0} \{ \tilde{V}_2(x + \delta_2) - c - \lambda(-\delta_2) \} = \max_{y \leq x} \{ \tilde{V}_2(y) - c - \lambda(x - y) \} = \max_{y \leq x} \{ \Gamma_2(y) \} - c - \lambda x, \quad (4.21)$$

where for each $y \in \mathbb{R}$ we have set

$$\Gamma_2(y) = \tilde{V}_2(y) + \lambda y.$$

By the definition of \tilde{V}_2 , we have $\Gamma'_2(x_2^*) = \Gamma'_2(\bar{x}_2) = 0$. Moreover, we notice that:

- $\Gamma'_2 = \lambda - \tilde{\lambda} \geq 0$ in $] - \infty, \bar{x}_1[$, by the definition of \tilde{V}_2 ;
- $\Gamma'_2 > 0$ in $] \bar{x}_1, x_2^*[$, as $\Gamma'_2(x_2^*) = 0$ and Γ'_2 is decreasing in $] \bar{x}_1, x_2^*[$ (since, by Proposition 4.2, we have $\Gamma''_2 = \varphi''_2 < 0$ in $] \bar{x}_1, x_2^*[$);
- $\Gamma'_2 < 0$ in $] x_2^*, \bar{x}_2[$, as $\Gamma'_2(x_2^*) = \Gamma'_2(\bar{x}_2) = 0$ and, in the interval $] x_2^*, \bar{x}_2[$, Γ'_2 is first decreasing and then increasing (since, by Proposition 4.2, $\Gamma''_2 = \varphi''_2$ is negative in $] x_2^*, \tilde{x}[$ and positive in $] \tilde{x}, \bar{x}_2[$);
- $\Gamma'_2 = 0$ in $] \bar{x}_2, +\infty[$, by the definition of \tilde{V}_2 .

As a consequence, the function Γ_2 has a unique global maximum point in x_2^* , so that

$$\max_{y \leq x} \Gamma_2(y) = \begin{cases} \Gamma_2(x), & \text{in }] - \infty, x_2^*], \\ \Gamma_2(x_2^*), & \text{in }] x_2^*, +\infty[; \end{cases}$$

therefore, by the computations in (4.21), we have

$$\mathcal{M}_2 \tilde{V}_2(x) = \begin{cases} \tilde{V}_2(x) - c, & \text{in }] - \infty, x_2^*], \\ \varphi_2(x_2^*) - c - \lambda(x - x_2^*), & \text{in }] x_2^*, +\infty[, \end{cases}$$

as $\tilde{V}_2(x_2^*) = \varphi_2(x_2^*)$, since $x_2^* \in] \bar{x}_1, \bar{x}_2[$. By the definition of \tilde{V}_2 , this can be written as

$$\mathcal{M}_2 \tilde{V}_2(x) = \begin{cases} \tilde{V}_2(x) - \xi_2(x), & \text{in }] - \infty, \bar{x}_2[, \\ \tilde{V}_2(x), & \text{in } [\bar{x}_2, +\infty[, \end{cases}$$

where, for each $x \in] - \infty, \bar{x}_2[$, we have set

$$\xi_2(x) = \begin{cases} c, & \text{in }] - \infty, x_2^*[, \\ \varphi_2(x) - \varphi_2(x_2^*) + c + \lambda(x - x_2^*), & \text{in } [x_2^*, \bar{x}_2[. \end{cases}$$

Let us prove that $\xi_2 > 0$. Recall by (4.6) that $\varphi_2(\bar{x}_2) = \varphi_2(x_2^*) - c - \lambda(\bar{x}_2 - x_2^*)$. Then, if $x \in [x_2^*, \bar{x}_2[$ we have that

$$\varphi_2(x) - \varphi_2(x_2^*) + c + \lambda(x - x_2^*) = \varphi_2(x) - \varphi_2(\bar{x}_2) - \lambda(\bar{x}_2 - x) = \Gamma_2(x) - \Gamma_2(\bar{x}_2) > 0,$$

as Γ_2 is decreasing in $[x_2^*, \bar{x}_2[$. Hence, ξ_2 is strictly positive, so that (4.20) holds. Finally, by the previous arguments it is clear that

$$\arg \max_{\delta_2 \leq 0} \{ \tilde{V}_2(x + \delta_2) - c - \lambda|\delta_2| \} = \begin{cases} \{0\}, & \text{in }] - \infty, x_2^*[, \\ \{x_2^* - x\}, & \text{in }] x_2^*, +\infty[, \end{cases}$$

which implies (4.19). □

Proposition 4.7. *A Nash equilibrium for the problem in Section 4.1 exists and is given by the strategies (A_1^*, ξ_1^*) , (A_2^*, ξ_2^*) defined by*

$$\begin{aligned} A_1^* &=]\bar{x}_1, +\infty[, & \xi_1^*(y) &= x_1^* - y, \\ A_2^* &=]-\infty, \bar{x}_2[, & \xi_2^*(y) &= x_2^* - y, \end{aligned}$$

with $y \in \mathbb{R}$ and x_i^*, \bar{x}_i ($i \in \{1, 2\}$) as in Definition 4.1. Moreover, the functions \tilde{V}_1, \tilde{V}_2 in Definition 4.1 coincide with the value functions V_1, V_2 :

$$V_1 \equiv \tilde{V}_1 \quad \text{and} \quad V_2 \equiv \tilde{V}_2.$$

Remark 4.8. Recall the practical characterization of the strategy: if x is the current state of the process, player 1 (resp. player 2) intervenes when $x \leq \bar{x}_1$ (resp. $x \geq \bar{x}_2$) and moves the process to the new state x_1^* (resp. x_2^*).

Proof. We have to check that the candidates \tilde{V}_1, \tilde{V}_2 satisfy all the assumptions of Theorem 3.3. We prove the claim for \tilde{V}_2 , the arguments for \tilde{V}_1 being the same. For the reader's convenience, we briefly report the conditions we have to check:

- (i) $\tilde{V}_2 \in C^2(]\bar{x}_1, +\infty[\setminus \{\bar{x}_2\}) \cap C^1(]\bar{x}_1, +\infty[) \cap C(\mathbb{R})$ and has polynomial growth;
- (ii) $\mathcal{M}_2 \tilde{V}_2 - \tilde{V}_2 \leq 0$;
- (iii) in $\{\mathcal{M}_1 \tilde{V}_1 - \tilde{V}_1 = 0\}$ we have $\tilde{V}_2 = \mathcal{H}_2 \tilde{V}_2$;
- (iv) in $\{\mathcal{M}_1 \tilde{V}_1 - \tilde{V}_1 < 0\}$ we have $\max\{\mathcal{A} \tilde{V}_2 - \rho \tilde{V}_2 + f_2, \mathcal{M}_2 \tilde{V}_2 - \tilde{V}_2\} = 0$;
- (v) the optimal strategies are x -admissible (see Definition 2.5) for every $x \in \mathbb{R}$.

Condition (i) and (ii). The first condition holds by the definition of \tilde{V}_2 , whereas the second condition has been proved in (4.20).

Condition (iii). Let $x \in \{\mathcal{M}_1 \tilde{V}_1 - \tilde{V}_1 = 0\} =]-\infty, \bar{x}_1]$. By the definition of $\mathcal{H}_2 \tilde{V}_2$ in (3.2), by (4.19) and by the definition of \tilde{V}_2 we have

$$\mathcal{H}_2 \tilde{V}_2(x) = \tilde{V}_2(x + \delta_1(x)) + \tilde{c} + \tilde{\lambda} |\delta_1(x)| = \tilde{V}_2(x_1^*) + \tilde{c} + \tilde{\lambda}(x_1^* - x) = \tilde{V}_2(x),$$

where we have used that $\tilde{V}_2(x_1^*) = \varphi_2(x_1^*)$, since $x_1^* \in]\bar{x}_1, \bar{x}_2[$.

Condition (iv). We have to prove that

$$\max\{\mathcal{A} \tilde{V}_2 - \rho \tilde{V}_2 + f_2, \mathcal{M}_2 \tilde{V}_2 - \tilde{V}_2\} = 0, \quad \text{in } \{\mathcal{M}_1 \tilde{V}_1 - \tilde{V}_1 < 0\} =]\bar{x}_1, +\infty[.$$

In $]\bar{x}_1, \bar{x}_2[$ the claim is true, as $\mathcal{M}_2 \tilde{V}_2 - \tilde{V}_2 < 0$ by (4.20) and $\mathcal{A} \tilde{V}_2 - \rho \tilde{V}_2 + f_2 = 0$ by definition (in $]\bar{x}_1, \bar{x}_2[$ we have $\tilde{V}_2 = \varphi_2$, which is a solution to the ODE). In $[\bar{x}_2, +\infty[$ we already know by (4.20) that $\mathcal{M}_2 \tilde{V}_2 - \tilde{V}_2 = 0$. Then, to conclude we have to check that

$$\mathcal{A} \tilde{V}_2(x) - \rho \tilde{V}_2(x) + f_2(x) \leq 0, \quad \forall x \in [\bar{x}_2, +\infty[.$$

As $\tilde{V}_2(x) = \varphi_2(x_2^*) - c - \lambda(x - x_2^*)$ by the definition of $\tilde{V}_2(x)$, the inequality can be written as

$$-\rho(\varphi_2(x_2^*) - c - \lambda(x - x_2^*)) + f_2(x) \leq 0, \quad \forall x \in [\bar{x}_2, +\infty[.$$

Since $\varphi_2(\bar{x}_2) = \varphi_2(x_2^*) - c - \lambda(\bar{x}_2 - x_2^*)$ by (4.6), we can rewrite the claim as

$$-\rho(\varphi_2(\bar{x}_2) - \lambda(x - \bar{x}_2)) + f_2(x) \leq 0, \quad \forall x \in [\bar{x}_2, +\infty[.$$

The function $x \mapsto \lambda \rho x + f_2(x) = (\lambda \rho - 1)x + s_2$ is decreasing, hence it is enough to prove the claim in $x = \bar{x}_2$:

$$-\rho \varphi_2(\bar{x}_2) + f_2(\bar{x}_2) \leq 0.$$

Since $\mathcal{A} \varphi_2(\bar{x}_2) - \rho \varphi_2(\bar{x}_2) + f_2(\bar{x}_2) = 0$, we can rewrite as

$$-\frac{\sigma^2}{2} \varphi_2''(\bar{x}_2) \leq 0,$$

which is true since $\varphi_2''(\bar{x}_2) \geq 0$ by Proposition 4.2.

Condition (v). Let x be the initial state of the process. By construction the controlled process never exists from $]\bar{x}_1, \bar{x}_2[\cup \{x\}$, so that condition (2.9) holds. It is easy to check that all the other conditions of Definition 2.5 are satisfied. The only non-trivial proof is the integrability of the intervention costs: let us prove that for $i \in \{1, 2\}$ we have (the result for $\tilde{c}, \tilde{\lambda}$ immediately follows, as $\tilde{\lambda} < \lambda$ and $\tilde{c} < c$)

$$\mathbb{E}_x \left[\sum_{k \geq 1} e^{-\rho \tau_{i,k}^*} (c + \lambda |\delta_{i,k}^*|) \right] < \infty, \quad (4.22)$$

where $\{\tau_{i,k}^*, \delta_{i,k}^*\}_k$ are the controls corresponding to the optimal strategies.

To start, let us assume that the initial state is either x_1^* or x_2^* . Since player i shifts the process to x_i^* when the state \bar{x}_i is hit, the idea is to write $\tau_{i,k}^*$ as a sum of independent exit times. First of all, we re-label the indexes and write $\{\tau_{i,k}^*\}_{i,k}$ as $\{\sigma_j\}_j$, with $\sigma_j < \sigma_{j+1}$ for every $j \in \mathbb{N}$ (this is possible: see Definition 2.2). Denote by μ_i the exit time of the process $x_i^* + \sigma W$ from $]\bar{x}_1, \bar{x}_2[$, where W is a real Brownian motion; then, each time σ_k can be written as $\sigma_k = \sum_{j=1}^k \zeta_j$, where the ζ_k are independent variables which are distributed either as μ_1 or as μ_2 . We can now estimate (4.22). As $\delta_{i,k}^* \in \{\bar{x}_2 - x_2^*, x_1^* - \bar{x}_1\}$, we have

$$\mathbb{E}_x \left[\sum_{i \in \{1,2\}} \sum_{k \geq 1} e^{-\rho \tau_{i,k}^*} (c + \lambda |\delta_{i,k}^*|) \right] \leq (c + \lambda \max\{\bar{x}_2 - x_2^*, x_1^* - \bar{x}_1\}) \mathbb{E}_x \left[\sum_{i \in \{1,2\}} \sum_{k \geq 1} e^{-\rho \tau_{i,k}^*} \right].$$

By the definition of $\{\sigma_j\}_j$ and the decomposition of σ_j ,

$$\mathbb{E}_x \left[\sum_{i \in \{1,2\}} \sum_{k \geq 1} e^{-\rho \tau_{i,k}^*} \right] = \mathbb{E}_x \left[\sum_{j \geq 1} e^{-\rho \sigma_j} \right] = \mathbb{E}_x \left[\sum_{j \geq 1} e^{-\rho \sum_{l=1}^j \zeta_l} \right] = \mathbb{E}_x \left[\sum_{j \geq 1} \prod_{l=1, \dots, j} e^{-\rho \zeta_l} \right].$$

By the Fubini-Tonelli theorem and the independence of the variables ζ_j , we get

$$\mathbb{E}_x \left[\sum_{j \geq 1} \prod_{l=1, \dots, j} e^{-\rho \zeta_l} \right] = \sum_{j \geq 1} \prod_{l=1, \dots, j} \mathbb{E}_x [e^{-\rho \zeta_l}] \leq \sum_{j \geq 1} (\mathbb{E}_x [e^{-\rho \min\{\mu_1, \mu_2\}}])^j,$$

which is a converging geometric series, since $\mu_1, \mu_2 \geq 0$. To sum up, we have shown

$$\mathbb{E}_x \left[\sum_{i \in \{1,2\}} \sum_{k \geq 1} e^{-\rho \tau_{i,k}^*} (\max\{c, \tilde{c}\} + \lambda |\delta_{i,k}^*|) \right] < \infty,$$

which clearly implies (4.22). The general case with initial state $x \in \mathbb{R}$ can be treated similarly: we have $\sigma_j = \eta + \sum_{l=1}^j \zeta_l$, where η is the exit time of $x + \sigma W$ from $[\bar{x}_1, \bar{x}_2]$, and the argument can be easily adapted. \square

4.4 Further comments and some limit properties

In order to understand the qualitative behaviour of the Nash equilibrium described in the previous section, we here study some asymptotic properties of the corresponding continuation regions and value functions. First, we recall some formulas from the previous sections, for reader's convenience. The value functions of the game are

$$V_2(x) = \begin{cases} \varphi_2^{A_{21}, A_{22}}(x_1^*) + \tilde{c} + \tilde{\lambda}(x_1^* - x), & \text{if } x \in]-\infty, \bar{x}_1], \\ \varphi_2^{A_{21}, A_{22}}(x), & \text{if } x \in]\bar{x}_1, \bar{x}_2[, \\ \varphi_2^{A_{21}, A_{22}}(x_2^*) - c - \lambda(x - x_2^*), & \text{if } x \in [\bar{x}_2, +\infty[, \end{cases} \quad V_1(x) = V_2(2\tilde{s} - x), \quad (4.23)$$

where the function $\varphi_2^{A_{21}, A_{22}}$ is defined in (4.2) and the parameters \bar{x}_i, x_i^*, A_{ij} are defined in (4.17). In particular, we recall the symmetry relations:

$$\bar{x}_1 = 2\tilde{s} - \bar{x}_2, \quad x_1^* = 2\tilde{s} - x_2^*,$$

with \tilde{s} as in (4.18). Also, recall that player 1 (resp. player 2) intervenes if the state is smaller than \bar{x}_1 (resp. greater than \bar{x}_2) and moves the process to x_1^* (resp. x_2^*).

Finally we remark that the parameter $\xi = \xi(c, \theta, \eta)$, defined as the unique zero of the function F in (4.14), satisfies the following properties: for given parameters θ and η , the function $c \mapsto \xi(c, \theta, \eta) =: \xi(c)$ belongs to $C^\infty(]0, \infty[)$ and we have

$$\xi'(c) = \frac{\theta \eta^2 - \xi^2(c)}{2 \xi^2(c)}, \quad \xi''(c) = -\theta \eta^2 \frac{\xi'(c)}{\xi^3(c)} = -\frac{\theta^2 \eta^2}{2} \frac{\eta^2 - \xi^2(c)}{\xi^5(c)}; \quad (4.24)$$

in particular, the following limits hold:

$$\lim_{c \rightarrow 0^+} \xi(c) = \lim_{c \rightarrow 0^+} \frac{c}{\xi(c)} = \lim_{c \rightarrow 0^+} c \xi'(c) = \lim_{c \rightarrow +\infty} c(\eta - \xi(c)) = 0, \quad \lim_{c \rightarrow +\infty} \xi(c) = \eta. \quad (4.25)$$

We now focus on the properties of the continuation region $]\bar{x}_1, \bar{x}_2[$ and the target states x_i^* with respect to the parameter c . All the other parameters are assumed to be fixed. To underline the dependence on this parameter, we will write $\bar{x}_i = \bar{x}_i(c)$, $x_i^* = x_i^*(c)$, $A_{ij} = A_{ij}(c)$ and $V_2 = V_2^c$, for $i, j \in \{1, 2\}$.

Limits as $c \rightarrow 0^+$. Since we are going to consider the limit $c \rightarrow 0^+$ and since by assumption we need $c > \tilde{c}$, with \tilde{c} fixed, we assume $\tilde{c} = 0$. If the fixed intervention cost vanishes, that is $c \rightarrow 0^+$, we expect that the players continuously intervene to keep the process in a state which satisfies both of them (namely \tilde{s} , for symmetry reasons): in other words, we expect the continuation region $]\bar{x}_1(c), \bar{x}_2(c)[$ to collapse to the singleton $\{\tilde{s}\}$, as $c \rightarrow 0^+$. Practically, if the initially state is x , either player 1 (if $x < \tilde{s}$) or player 2 (if $x > \tilde{s}$) shifts the process to \tilde{s} ; from then on, we constantly have $X_s \equiv \tilde{s}$. As a consequence, we guess that the value function for player 2 is

$$\begin{aligned} V_2^{0+}(x) &= \mathbb{E}_x \left[\int_0^\infty e^{-\rho s} (s_2 - \tilde{s}) ds - \lambda(x - \tilde{s}) \mathbb{1}_{\{x < \tilde{s}\}} + \tilde{\lambda}(\tilde{s} - x) \mathbb{1}_{\{x > \tilde{s}\}} \right] \\ &= \frac{s_2 - \tilde{s}}{\rho} - (\tilde{\lambda} \mathbb{1}_{\{x > \tilde{s}\}} + \lambda \mathbb{1}_{\{x < \tilde{s}\}})(x - \tilde{s}). \end{aligned}$$

We now rigorously prove these heuristic arguments by considering the explicit expression for the intervention region provided in (4.17). Actually, the limit situation is not as straightforward as it may appear: the parameters $\lambda, \tilde{\lambda}$ play an important role.

Proposition 4.9. *Assume $\tilde{c} = 0$ and $\lambda = \tilde{\lambda}$. Then we have, for $i \in \{1, 2\}$ and $x \in \mathbb{R}$*

$$\bar{x}_i(0^+) = x_i^*(0^+) = \tilde{s}, \quad V_1^{0+}(x) = \frac{s_2 - \tilde{s}}{\rho} - \lambda(x - \tilde{s}), \quad V_2^{0+}(x) = \frac{s_2 - \tilde{s}}{\rho} - \lambda(x - \tilde{s}).$$

Proof. By (4.17) and (4.25) it follows that $\bar{x}_2(c) \rightarrow \tilde{s}$ as $c \rightarrow 0^+$. The same result holds for \bar{x}_1 by symmetry and hence also for x_i^* , since $x_i^* \in]\bar{x}_1, \bar{x}_2[$. Moreover, again by (4.17) and (4.25), we get

$$A_{21}(0^+) = e^{-\theta \tilde{s}} \frac{\eta}{2\theta}, \quad A_{22}(0^+) = -e^{\theta \tilde{s}} \frac{\eta}{2\theta};$$

hence, by the first part of the proof, for each $x \in \mathbb{R}$ we have (recall that $\lambda = \tilde{\lambda}$)

$$V_2^{0+}(x) = \varphi_2^{A_{21}(0^+), A_{22}(0^+)}(\tilde{s}) - \lambda(x - \tilde{s}) = \frac{s_2 - \tilde{s}}{\rho} - \lambda(x - \tilde{s}).$$

The corresponding result for V_1 follows by symmetry. \square

Limits as $c \rightarrow +\infty$. If the intervention cost increases, the players rarely intervene. In the limit case $c \rightarrow +\infty$, they never intervene and we expect $]\bar{x}_1(c), \bar{x}_2(c)[$ to coincide with \mathbb{R} . Correspondingly, the state variable diffuses without being affected by the players, that is $X_s = x + \sigma W_s$, for each $s \geq 0$. As a consequence, we guess that the value function for player 2 is

$$V_2^{+\infty}(x) = \mathbb{E}_x \left[\int_0^\infty e^{-\rho s} (s_2 - x - \sigma W_s) ds \right] = \frac{s_2 - x}{\rho}.$$

Moreover, the intervening player clearly compensates the cost $c + \lambda|x_i^* - x|$ by moving the process to a state where her payoff is bigger than the opponent's one. In the case $c \rightarrow +\infty$, the intervening player has to compensate diverging costs, so that we guess that $x_i^*(c)$ diverges too. We now rigorously prove our guesses.

Proposition 4.10. *The following limits hold:*

$$\begin{aligned}\bar{x}_2(+\infty) &= x_1^*(+\infty) = +\infty, & \bar{x}_1(+\infty) &= x_2^*(+\infty) = -\infty, \\ V_1^{+\infty}(x) &= \frac{x - s_1}{\rho}, & V_2^{+\infty}(x) &= \frac{s_2 - x}{\rho}.\end{aligned}$$

Proof. By (4.17) and (4.25) it easily follows that $\bar{x}_2(+\infty) = +\infty$ and $x_2^*(+\infty) = -\infty$. By symmetry, corresponding results hold for \bar{x}_1, x_1^* . Moreover, by (4.17) and (4.25) we get

$$A_{21}(+\infty) = A_{21}(+\infty) = 0;$$

hence, by the first part of this proof, for each $x \in \mathbb{R}$ we have

$$V_2^{+\infty}(x) = \varphi_2^{A_{21}(+\infty), A_{22}(+\infty)}(x) = \frac{s_2 - x}{\rho}.$$

The corresponding result for V_1 follows by symmetry. \square

Monotonicity of \bar{x}_i, x_i^* . If the intervention cost c increases, we expect the common continuation region $]\bar{x}_1(c), \bar{x}_2(c)[$ to enlarge, since the players are less willing to intervene. Proposition 4.11 makes this guess rigorous.

Proposition 4.11. *The function $c \mapsto \bar{x}_2(c)$, with $c \in]\tilde{c}, +\infty[$, is increasing and the function $c \mapsto \bar{x}_1(c)$ is decreasing.*

Proof. Let us prove that $c \mapsto \bar{x}_2(c)$, with $c \in]\tilde{c}, +\infty[$, is increasing. By (4.17) it suffices to check that

$$c \mapsto \frac{\eta + \xi(c)}{\eta - \xi(c)} \left(\frac{\theta(\lambda - \tilde{\lambda} + \eta)}{4\eta} \frac{c}{\xi(c)} - \frac{\theta\tilde{c}}{4} \frac{1}{\xi(c)} + \frac{\lambda - \tilde{\lambda}}{2\eta} \right)$$

is an increasing function. Since $\xi' > 0$, a sufficient condition is that

$$c \mapsto \frac{c}{\xi(c)}, \quad c > \tilde{c},$$

is increasing, that is

$$H(c) = \xi(c) - c\xi'(c) \geq 0, \quad \forall c > \tilde{c},$$

which is true since by (4.25) we have $H(\tilde{c}^+) = 0$ (consider separately the cases $\tilde{c} = 0$ and $\tilde{c} > 0$) and $H'(c) > 0$. The result for $\bar{x}_1(c)$ follows by symmetry. \square

As for the monotonicity of x_i^* , it is not easy to make a guess. The formulas in (4.17) does not allow easy estimates; however, a monotonicity result can be proved in the case $\tilde{c} = 0$. We will see later by some numerical simulations that in the general case the function x_i^* is not monotone.

Proposition 4.12. *Assume $\tilde{c} = 0$. Then, the function $c \mapsto x_2^*(c)$, with $c \in]0, +\infty[$, is decreasing and the function $c \mapsto x_1^*(c)$ is increasing. Moreover, we have $x_2^* < \tilde{s} < x_1^*$ for each $c > 0$.*

Proof. Let us prove that $c \mapsto x_2^*(c)$, with $c \in]0, +\infty[$, is decreasing. By (4.17) it suffices to prove that

$$c \mapsto \frac{\theta(\lambda - \tilde{\lambda} + \eta)}{4\eta} \frac{c(\eta - \xi(c))}{\xi(c)(\eta + \xi(c))} + \frac{\lambda - \tilde{\lambda}}{2\eta} \frac{\eta - \xi(c)}{\eta + \xi(c)}$$

is a decreasing function. Since $\xi' > 0$, a sufficient condition is that

$$c \mapsto \frac{c(\eta - \xi(c))}{\xi(c)(\eta + \xi(c))}$$

is decreasing, that is

$$K(c) = \xi(c) - \frac{\theta}{2} \frac{c(\eta^2 - \xi^2(c))}{\xi^2(c)} - \theta\eta \frac{c}{\xi(c)} \leq 0, \quad \forall c > 0,$$

which is true since by (4.25) we have $K(0^+) = 0$ and $K'(c) < 0$. The results for $x_1^*(c)$ follows by symmetry. Finally, we get the inequalities by $(x_2^*)' < 0 < (x_1^*)'$ and $x_1^*(0^+) = x_2^*(0^+) = \tilde{s}$. \square

Limits as $c \rightarrow \tilde{c}^+$. We conclude this section with the behaviour as $c \rightarrow \tilde{c}^+$ in the case $\lambda = \tilde{\lambda}$, i.e. when each intervention practically becomes a transfer of money from the intervening player to the opponent. It is not easy to guess what happens in this case and the result is quite surprising: the limiting strategies are not admissible.

Proposition 4.13. *Assume $\lambda = \tilde{\lambda}$. For $i, j \in \{1, 2\}$ with $i \neq j$, the following limits hold:*

$$\bar{x}_i(\tilde{c}^+) = x_j^*(\tilde{c}^+) = \tilde{s} + \frac{(-1)^i}{2\theta} \log \left(\frac{\eta + \xi(\tilde{c})}{\eta - \xi(\tilde{c})} \right).$$

Proof. The result immediately follows by (4.17) and (4.25). \square

Essentially, the limit situation is as follows. Let $i, j \in \{1, 2\}$ with $i \neq j$; as soon as the process reaches \bar{x}_i , player i moves the process to $x_i^* = \bar{x}_j$, which is the boundary of the intervention region of player j , who moves the process back to \bar{x}_i , thus causing another intervention by player i and so on. We get a infinite sequence of simultaneous interventions, meaning that these strategies are not admissible.

Numerical simulations. Here, we present the results (obtained with Wolfram Mathematica) of some numerical simulations on the game we have described. We focus on player 2 and consider the following two sets of parameters:

Problem 1: $\rho = 0.02$, $\sigma = 0.15$, $s_1 = -3$, $s_2 = 3$, $\tilde{c} = 0$, $\lambda = \tilde{\lambda} = 15$.

Problem 2: $\rho = 0.02$, $\sigma = 0.15$, $s_1 = -3$, $s_2 = 3$, $\tilde{c} = 50$, $\lambda = \tilde{\lambda} = 0$.

Figure 4.1 represents the value function $x \mapsto V_2^c(x)$ for Problem 1 and $c = 100$ (the dashed lines correspond to the three components of the function). Similarly, in Figure 4.2 we plot the function $x \mapsto V_2^c(x)$ for Problem 2 and $c = 100$. In both the cases, we notice the C^1 -pasting in \bar{x}_1 , whereas, as noticed in Section 2, the functions are not differentiable in \bar{x}_1 . Also, when λ is non-zero, the function is unbounded.

Figure 4.3 (for Problem 1, with $c \in]\tilde{c}, \infty[=]0, \infty[$) and Figure 4.4 (for Problem 2, with $c \in]\tilde{c}, \infty[=]50, \infty[$) show the continuation region and the target states: namely, we plot $c \mapsto \bar{x}_1(c)$ (solid blue line), $\bar{x}_2(c)$ (solid green line), $x_1^*(c)$ (dashed blue line) and $x_2^*(c)$ (dashed green line). As proved above, the continuation region enlarges as c grows and diverges as $c \rightarrow \infty$. Consider the limit case $c \rightarrow \tilde{c}^+$: if $\tilde{c} = 0$, the four parameters converge to the same state \tilde{s} ; conversely, in the case $\tilde{c} > 0$, we see that x_1^* (resp. x_2^*) converges to \bar{x}_2 (resp. \bar{x}_1), which corresponds to an inadmissible game. Also, we notice that x_1^* (resp. x_2^*) is decreasing (resp. increasing) when $\tilde{c} = 0$, whereas such functions are not monotone in the $\tilde{c} > 0$ case.

We finally consider Problem 1 and the evolution of $x \mapsto V_2^c(x)$ as c grows: Figure 4.5 corresponds to $c = 0$, Figure 4.6 to $c = 250$, Figure 4.7 to $c = 500$ and Figure 4.8 to $c = 750$. The value function is a straight line in the limit case $c \rightarrow 0^+$, then a bell-shaped curve appears; as c grows, the local maximum moves to the left and the right side of the bell resembles more and more a straight line with slope $1/\rho$, which is actually the limit as $c \rightarrow +\infty$.




Figure 4.1: $x \mapsto V_2^c(x)$ for Pr. 1 and $c = 100$




Figure 4.2: $x \mapsto V_2^c(x)$ for Pr. 2 and $c = 100$




Figure 4.3: $c \mapsto \bar{x}_i(c), x_i^*(c)$ for Problem 1




Figure 4.4: $c \mapsto \bar{x}_i(c), x_i^*(c)$ for Problem 2




Figure 4.5: Problem 1, $c = 0$




Figure 4.6: Problem 1, $c = 250$




Figure 4.7: Problem 1, $c = 500$




Figure 4.8: Problem 1, $c = 750$

5 Conclusions

In this paper, we have considered a general two-player nonzero-sum impulse game, whose state variable follows a diffusive dynamics driven by a multi-dimensional Brownian motion. After setting the problem, we have provided a verification theorem giving sufficient conditions in order for the solutions of a suitable system of quasi-variational inequalities to coincide with the value functions of the two players. To the best of our knowledge this result is new to the literature on impulse games and it constitutes the major mathematical contribution of the present paper. As an application, we have provided a solvable one-dimensional impulse game where two players with linear running payoffs can shift a real-valued Brownian motion in order to maximize their objective functions. We have found a Nash equilibrium and explicitly characterized the corresponding optimal strategies. Finally, we have also studied some asymptotic properties of such a Nash equilibrium.

References

- [1] M. Basei, *Topics in stochastic control and differential game theory, with application to mathematical finance*, Ph.D. Thesis in Mathematics, University of Padua (2016).
- [2] A. Bensoussan, A. Friedman, *Nonzero-sum stochastic differential games with stopping times and free boundary problems*, Trans. Amer. Math. Society 231 (1977), no. 2, 275–327.
- [3] A. Bensoussan, J. L. Lions, *Contrôle impulsif et inéquations quasi variationnelles*, Vol. 1, Dunod, Paris, 1982.
- [4] G. Bertola, W. J. Runggaldier, K. Yasuda, *On Classical and Restricted Impulse Stochastic Control for the Exchange Rate*, Appl. Math. Optim. 74 (2016), no. 2, 423–454.
- [5] A. Cadenillas, F. Zapatero, *Classical and impulse stochastic control of the exchange rate using interest rates and reserves*, Math. Finance 10 (2000), no.2, 141–156.
- [6] D. Chang, H. Wang, Z. Wu, *Maximum principle for non-zero sum differential games of BSDEs involving impulse controls*, 32nd Chinese Control Conference (CCC 2013), IEEE, 1564–1569.
- [7] D. Chang, Z. Wu, *Stochastic maximum principle for non-zero sum differential games of FB-SDEs with impulse controls and its application to finance*, J. Ind. Manag. Optim. 11 (2015), no. 1, 27–40.
- [8] N. Chen, M. Dai, X. Wan, *A Nonzero-Sum Game Approach to Convertible Bonds: Tax Benefit, Bankruptcy Cost, and Early/Late Calls*, Math. Finance 23 (2013), no. 1, 57–93.
- [9] A. Cosso, *Stochastic differential games involving impulse controls and double-obstacle quasi-variational inequalities*, SIAM J. Control Optim. 51 (2013), no. 3, 2102–2131.
- [10] T. De Angelis, G. Ferrari, *Stochastic nonzero-sum games: a new connection between singular control and optimal stopping*, preprint (2016), arXiv:1601.05709v2 [math.OC].
- [11] T. De Angelis, G. Ferrari, J. Moriarty, *Nash equilibria of threshold type for two-player nonzero-sum games of stopping*, Ann. Appl. Probab., forthcoming.
- [12] A. Friedman, *Stochastic games and variational inequalities*, Arch. Rational Mech. Anal. 51 (1973), no. 5, 321–346.
- [13] N. V. Krylov, *Controlled diffusion processes*, Vol. 14, Springer Science & Business Media, 2008.
- [14] N. Touzi, *Optimal stochastic control, stochastic target problems, and backward SDE*, Vol. 29, Springer Science & Business Media, 2012.
- [15] B. K. Øksendal, A. Sulem, *Applied stochastic control of jump diffusions, Second Edition*, Springer-Verlag, Berlin-Heidelberg, 2007.