

Global Existence and Regularity Results for Large Cross Diffusion Models on Planar Domains.

DUNG LE¹

Abstract

The global existence of classical solutions to cross diffusion systems of more than 2 equations given on a planar domain is established. The results can apply to generalized Shigesada-Kawasaki-Teramoto (SKT) and food pyramid models whose diffusion and reaction can have polynomial growth of any order.

1 Introduction

We consider in this paper the following system

$$\begin{cases} u_t = \operatorname{div}(A(u)Du) + f(u, Du) & (x, t) \in Q = \Omega \times (0, T), \\ u(x, 0) = U_0(x) & x \in \Omega \\ u = 0 \text{ on } \partial\Omega \times (0, T). \end{cases} \quad (1.1)$$

Here, Ω is a bounded domain with smooth boundary $\partial\Omega$ in \mathbb{R}^2 ; $u : \Omega \rightarrow \mathbb{R}^m$ and $f : \mathbb{R}^m \times \mathbb{R}^{2m} \rightarrow \mathbb{R}^m$ are vector valued functions. $A(u)$ is a full matrix $m \times m$. Thus, the above is a system of m equations.

The system (1.1) arises in many mathematical biology and ecology applications. In the last few decades, papers concerning such strongly coupled parabolic systems usually assumed that the solutions under consideration were bounded, a very hard property to check as maximum principles had been unavailable for systems in general. Most of global existence results for cross diffusion systems relied on the following local existence result of Amann.

Theorem 1.1 ([2, 3]) *Suppose $\Omega \subset \mathbb{R}^n$, $n \geq 2$, with $\partial\Omega$ being smooth. Assume that (1.1) is normally elliptic. Let $p_0 \in (n, \infty)$ and U_0 be in $W^{1,p_0}(\Omega)$. Then there exists a maximal time $T_0 \in (0, \infty]$ such that the system (1.1) has a unique classical solution in $(0, T_0)$ with*

$$u \in C([0, T_0), W^{1,p_0}(\Omega)) \cap C^{1,2}((0, T_0) \times \bar{\Omega})$$

Moreover, if $T_0 < \infty$ then

$$\lim_{t \rightarrow T_0^-} \|u(\cdot, t)\|_{W^{1,p_0}(\Omega)} = \infty. \quad (1.2)$$

¹Department of Mathematics, University of Texas at San Antonio, One UTSA Circle, San Antonio, TX 78249. Email: Dung.Le@utsa.edu

Mathematics Subject Classifications: 35J70, 35B65, 42B37.

Key words: Cross diffusion systems, Hölder regularity, global existence.

Equivalently, the classical solution u will exist globally if its $W^{1,p_0}(\Omega)$ norm does not blow up in finite time. This requires the existence of a continuous function \mathcal{C} on $(0, \infty)$ such that

$$\|u(\cdot, t)\|_{W^{1,p_0}(\Omega)} \leq \mathcal{C}(t) \quad \forall t \in (0, T_0). \quad (1.3)$$

We refer the readers to [2] for the definition of normal ellipticity (roughly speaking, it means that the real parts of the eigenvalues of $A(u)$ are positive). The checking of (1.2) is the most difficult one as known techniques for the regularity of solutions to scalar equations could not be extended to systems and counterexamples were available.

In our recent work [12], see also [10, 11], we considered (1.1) on a domain in \mathbb{R}^n ($n \geq 2$) and were able to relax the condition (1.3) by

$$\|u(\cdot, t)\|_{W^{1,n}(\Omega)} \leq \mathcal{C}(t) \quad \forall t \in (0, T_0). \quad (1.4)$$

Obviously, (1.4) does not imply that $|u|$ is bounded so that (1.1) is not regularly elliptic, i.e. eigenvalues of $A(u)$ can still be unbounded. In [12], we only assume that (1.1) is uniformly elliptic, meaning the the eigenvalues of $A(u)$ are comparable.

In fact, we prove in [12] that the global existence result in Amann's theorems holds with a much weaker version of (1.3) and (1.4). Namely, one needs only to control the BMO norm of u and shows that u is VMO. The proof makes use several techniques from Harmonic Analysis and a generalized weighted Gagliardo-Nirenberg inequality involving BMO norms. Some mild structural conditions on the cross diffusion matrix $A(u)$ of (1.1) are imposed and easily verified, for examples, if $A(u)$ has a polynomial growth of order less than 5.

In this work, for planar domains ($n = 2$) we will present a new and simpler proof of the described result in [12]. Tools from Harmonic Analysis will not be needed here. More importantly, the structural conditions on $A(u)$ will be much weaker. As an example, we will allow $A(u)$ has a polynomial growth of *any* order. The new result is given in Theorem 2.1.

The checking of (1.4) for general n is by no means an easy task. Here, for $n = 2$ we will give two examples in applications where this can be done under very general assumptions.

In particular, we will consider a class of generalized Shigesada-Kawasaki-Teramoto (SKT) models consisting of more than 2 equations. Namely, we will establish the global existence of classical solutions to the following system

$$u_t = \Delta(\mathcal{P}(u)) + f(u, Du), \quad (1.5)$$

where $\mathcal{P}(u), f(u, Du)$ are vector valued functions. The above system is a special case of (1.1) with $A(u) = \frac{\partial \mathcal{P}(u)}{\partial u}$. The (SKT) models, consisting of only 2 equations, were introduced in [15] using quadratic growth \mathcal{P} 's and Lotka-Volterra reaction $f(u)$. The global existence for the (SKT) model for 2 species on planar domains was studied in [17]. In this work, we will consider (1.5) with $\mathcal{P}(u), f(u, Du)$ have polynomial growth of any order in u . More importantly, the number of species/equations can be arbitrary.

The second example is the food pyramid model which assumes that the first k equations of (1.1) do not involve with the j -unknown u_j if $j > k$. This model has been studied in literature (e.g., [1]) under the assumption that $A(u)$ is a constant diagonal matrix, with no cross diffusion.

2 Preliminaries and Main Results

Throughout this paper Ω is a bounded domain with smooth boundary in \mathbb{R}^2 . For any smooth (vector valued) function u defined on $\Omega \times (0, T)$, $T > 0$, its temporal and spatial derivatives are denoted by u_t, Du respectively. If A is a C^1 function in u then we also abbreviate $\frac{\partial A}{\partial u}$ by A_u .

As usual, $W^{1,p}(\Omega, \mathbb{R}^m)$, $p \geq 1$, will denote the standard Sobolev spaces whose elements are vector valued functions $u : \Omega \rightarrow \mathbb{R}^m$ with finite norm

$$\|u\|_{W^{1,p}(\Omega, \mathbb{R}^m)} = \|u\|_{L^p(\Omega)} + \|Du\|_{L^p(\Omega)}.$$

We assume the following structural conditions.

A) $A(u)$ is C^1 in u . Moreover, there are positive constants λ_0, C and a scalar C^1 function $\lambda(u)$ such that $\lambda(u) \geq \lambda_0$ for all $u \in \mathbb{R}^m$. Furthermore, for any $\zeta \in \mathbb{R}^{nm}$

$$\lambda(u)|\zeta|^2 \leq \langle A(u)\zeta, \zeta \rangle \text{ and } |A(u)| \leq C\lambda(u). \quad (2.1)$$

We also assume $|A_u| \leq C|\lambda_u|$ and

$$|\lambda_u(u)| \leq C\lambda(u). \quad (2.2)$$

Our first main theorem under this general assumption weakens the condition (1.2) of Amann's result in Theorem 1.1 by the condition (2.5) where we use much weaker norms.

Theorem 2.1 *Suppose $\Omega \subset \mathbb{R}^2$ with $\partial\Omega$ being smooth. Assume A) and U_0 be in $W^{1,p_0}(\Omega)$ for some $p_0 > 2$. Let $T_0 \in (0, \infty]$ be the maximal existence time for a unique classical solution $u \in C^{1,2}((0, T_0) \times \bar{\Omega})$ of*

$$\begin{cases} u_t = \operatorname{div}(A(u)Du) + \hat{f}(u, Du) & (x, t) \in Q = \Omega \times (0, T), \\ u(x, 0) = U_0(x) & x \in \Omega \\ \text{Boundary conditions for } u \text{ on } \partial\Omega \times (0, T). \end{cases} \quad (2.3)$$

For such u , assume that there are a constant C and a C^1 function $f(u)$ such that $|f_u(u)| \leq C\lambda(u)$ and $\hat{f}(u, Du)$ satisfies the following growth conditions

$$|\hat{f}(u, Du)| \leq C\lambda^{\frac{1}{2}}(u)|Du| + f(u), \quad (2.4)$$

Then if there exists a continuous function g on $(0, \infty)$ such that

$$\int_{\Omega} \lambda(u)|Du(x, t)|^2 dx \leq g(t) \quad \forall t \in (0, T_0), \quad (2.5)$$

then $T_0 = \infty$.

The proof of this theorem is fairly technical and we will postpone it to the end of the paper (see Section 4). Of course, the assumption (2.5) is a bit stronger than (1.2) but the structural conditions on (1.1) are much weaker than those in [12]. In fact, we will show that the crucial hypothesis (2.5) for global existence can be verified in many applications including the general (SKT) and pyramid systems.

For the sake of simplicity, we will consider first the case where the reaction term in (1.1) does not depend on Du . We then assume the following.

F) We assume that there are positive constants ε_0, C and nonnegative C^1 functions $P, F : \mathbb{R}^m \rightarrow \mathbb{R}^+$ satisfying $F(0) = P(0) = 0$ and

$$|F_u(u)| \leq C\lambda^{\frac{1}{2}}(u), \quad (2.6)$$

$$|P_u(u)| \leq C\lambda(u) \quad (2.7)$$

for all $u \in \mathbb{R}^m$ such that

$$|f(u)||u| \leq \varepsilon_0 F^2(u) + C, \quad (2.8)$$

$$\frac{\lambda^{\frac{1}{2}}(u)|f(u)|}{P(u) + 1} \leq C(F(u) + 1). \quad (2.9)$$

More generally, we can replace $f(u)$ by a function \hat{f} depending on u, Du and satisfying a linear growth in Du . Namely, we will assume the following.

F') There exist a constant C and a function $f(u)$ satisfying F) such that

$$|\hat{f}(u, Du)| \leq C\lambda^{\frac{1}{2}}(u)|Du| + f(u), \quad (2.10)$$

$$|f_u(u)| \leq C\lambda(u). \quad (2.11)$$

Let us discuss some applications where the conditions A) and F) can be verified. In many models, $A(u), f(u)$ have polynomial growths in u . That is, there are nonnegative numbers k, K such that $\lambda(u) \sim (1 + |u|)^k$ and $|f(u)| \sim (1 + |u|)^K$. Here and in the sequel, we will write $a \sim b$ if there are two generic positive constants C_1, C_2 such that $C_1 b \leq a \leq C_2 b$.

Obviously, (2.2) in A) holds for any $k \geq 0$. Concerning F), we can take $F(u) = |u|^{\frac{k+2}{2}}$ and $P(u) = |u|^{k+1}$. It is clear that (2.6) and (2.7) hold for such choice of F, P . We also have $|f(u)||u| \leq (1 + |u|)^{K+1} \sim (F(u) + 1)^2$ if $K = k + 1$. Thus, (2.8) is satisfied with ε_0 being the coefficient of the highest power of u in $f(u)$.

On the other hand, if $K \leq k + 2$ then (2.9) is satisfied because

$$\frac{\lambda^{\frac{1}{2}}(u)|f(u)|}{P(u) + 1} \leq C \frac{(1 + |u|)^{\frac{k}{2} + K}}{(1 + |u|)^{k+1}} \sim (1 + |u|)^{K - \frac{k}{2} - 1} \leq C(1 + |u|)^{\frac{k}{2} + 1} = C(F(u) + 1).$$

Hence, it is clear that the main assumptions in F) and (2.11) in F') are verified if $K = k + 1$.

We then prove the following next main result which shows that if the **excess**

$$\mathcal{A}(u) = \langle A(u)Du, DA(u)u_t - A(u)_t Du \rangle \quad (2.12)$$

of a classical solution u to (1.1) does not blow up in time then (2.5) holds to give the global existence of the classical solution u .

Theorem 2.2 *Assume A) and F'). Let u be a classical solution to (1.1) and T_0 be its maximal existence time. For any $t_0 \in (0, T_0)$ and $\varepsilon > 0$ assume that there are continuous*

function C_ε, C_A on $(0, \infty)$ which may also depend on $\|u(\cdot, t_0)\|_{W^{1,2}(\Omega)}$ and $\|u_t(\cdot, t_0)\|_{L^2(\Omega)}$ such that

$$\int_{t_0}^s \int_{\Omega} \mathcal{A}(u) \, dx \, dt \leq \int_{t_0}^s \int_{\Omega} [\varepsilon \lambda(u) |u_t|^2 + C_\varepsilon(t) |A(u) Du|^2] \, dx \, dt + C_A(s) \quad (2.13)$$

for all $s \in (t_0, T_0)$.

If either ε_0 or $d(\Omega)$ is sufficiently small then $T_0 = \infty$.

The next two theorems show that (2.13) holds for the generalized (SKT) and *pyramid* models.

Theorem 2.3 *Assume A) and F'). Suppose that there is a C^2 function $\mathcal{P} : \mathbb{R}^m \rightarrow \mathbb{R}^m$ such that $A(u) = \mathcal{P}_u$.*

If either ε_0 or $d(\Omega)$ is sufficiently small then $T_0 = \infty$.

Theorem 2.4 *For $k_0 = 1, \dots, m-1$ we suppose that the subsystems of the first k_0 equations of (1.1) do not depend on the unknowns u_i if $i > k_0$ and they satisfy A) and F'). Furthermore, there is a constant C such that for $u = (u_1, \dots, u_{k_0})^T$*

$$\left| \frac{\partial}{\partial u_{k_0}} a_{k_0 j}(u) \right| \leq C \lambda^{\frac{1}{2}}(u) \quad \forall j < k_0, k_0 = 2, \dots, m. \quad (2.14)$$

If either ε_0 or $d(\Omega)$ is sufficiently small then $T_0 = \infty$.

We remark that if $A(u)$ has polynomial growth in u and $\lambda(u) \sim (1 + |u|)^k$ then (2.14) holds if $k \in [0, 2]$. In fact, the reaction terms of the subsystem for the first k_0 preys (u_i , $i \leq k_0$) can depend on the predators u_i ($i > k_0$). We are interested in the effect of cross diffusion in this paper and assume very weak feeding rate of the predators in the reaction terms of the systems. Strong feeding rates of predators for cross diffusion systems will be reported in our forthcoming works.

3 Proof

In this section, we will consider a classical solution to (1.1) that exists in its maximal time interval $(0, T_0)$. We also fix a t_0 in $(0, T_0)$. We will frequently use functions \mathcal{C} satisfying the following property.

C) \mathcal{C} is continuous on $(0, \infty)$ and depends on $\|u(\cdot, t_0)\|_{W^{1,2}(\Omega)}$, $\|(u(\cdot, t_0))_t\|_{L^2(\Omega)}$ and $d(\Omega)$.

In the proof, when there is no ambiguity C, C_i will denote universal constants that can change from line to line in our argument. Furthermore, $C(\cdot \cdot \cdot)$ is used to denote quantities which are bounded in terms of their parameters. The same convention applies to functions $\mathcal{C}, \mathcal{C}_i$ satisfying the property C).

Lemma 3.1 *Assume (2.6) and (2.8) of F). If either ε_0 or $d(\Omega)$ is sufficiently small then there is a function \mathcal{C} satisfying C) such that*

$$\sup_{t \in [t_0, T]} \int_{\Omega} |u|^2 dx + \iint_{\Omega \times [t_0, T]} \lambda(u) |Du|^2 dz \leq \mathcal{C}(T) \quad \forall T \in (t_0, T_0). \quad (3.1)$$

Proof: Testing the system of u with u and integrating over $\Omega \times [t_0, T]$ by parts, we easily obtain for $Q = \Omega \times [t_0, T]$

$$\sup_{t \in [t_0, T]} \int_{\Omega} |u|^2 dx + \iint_Q \lambda(u) |Du|^2 dz \leq \iint_Q \langle f(u), u \rangle dz + \|u(\cdot, t_0)\|_{L^2(\Omega)}^2. \quad (3.2)$$

By (2.8) of F) we can find a constant C such that

$$\int_{\Omega} \langle f(u), u \rangle dx \leq \varepsilon_0 \int_{\Omega} F^2(u) dx + C|\Omega|.$$

Here, $|\Omega|$ is the Lebesgue measure of Ω .

Because $F(0) = 0$ and the boundary condition of u , we have $F(u) = 0$ on the boundary $\partial\Omega$. Using the Poincaré inequality and (2.6), we have

$$\int_{\Omega} F^2(u) dx \leq Cd^2(\Omega) \int_{\Omega} |DF(u)|^2 dx \leq Cd^2(\Omega) \int_{\Omega} \lambda(u) |Du|^2 dx.$$

Hence,

$$\int_{\Omega} \langle f(u), u \rangle dx \leq C\varepsilon_0 d^2(\Omega) \int_{\Omega} \lambda(u) |Du|^2 dx + C|\Omega|.$$

Thus, if either ε_0 or $d(\Omega)$ is sufficiently small then (3.2) and the above yield a constant C_1 such that

$$\sup_{t \in [t_0, T]} \int_{\Omega} |u|^2 dx + C_1 \iint_Q \lambda(u) |Du|^2 dz \leq C|\Omega|T + \|u(\cdot, t_0)\|_{L^2(\Omega)}^2.$$

The last quantity defines a function $\mathcal{C}(T)$ satisfying C). The proof is then complete. ■

To proceed, we need the following elementary fact. By the Ladyzhenskayas inequality ($n = 2$) we have

$$\left(\int_{\Omega} |U|^4 dx \right)^{\frac{1}{2}} \leq \left(\int_{\Omega} |U|^2 dx \right)^{\frac{1}{2}} \left(\int_{\Omega} |DU|^2 dx \right)^{\frac{1}{2}}, \quad (3.3)$$

if $U = 0$ on the boundary $\partial\Omega$. The Poincaré inequality applies to the first factor on the right gives

$$\left(\int_{\Omega} |U|^4 dx \right)^{\frac{1}{2}} \leq \int_{\Omega} |DU|^2 dx.$$

Hence, if U, V vanish on the boundary $\partial\Omega$ then

$$\int_{\Omega} |U|^2 |V|^2 dx \leq \left(\int_{\Omega} |U|^4 dx \right)^{\frac{1}{2}} \left(\int_{\Omega} |V|^4 dx \right)^{\frac{1}{2}} \leq C \int_{\Omega} |DU|^2 dx \int_{\Omega} |DV|^2 dx. \quad (3.4)$$

We also note that $\lambda(u)$ is the smallest eigenvalue of $(A + A^T)/2$ and $\Lambda(u)$ is the smallest eigenvalue of $A^T A$. Thus, if $\mu(u)$ is the eigenvalue of A with smallest real part then $\lambda(u) = \Re(\mu(u))$ and $\Lambda(u) = |\mu(u)|^2$. Therefore,

$$|A(u)\zeta|^2 = \langle A^T(u)A(u)\zeta, \zeta \rangle \geq \Lambda(u)|\zeta|^2 \geq \lambda^2(u)|\zeta|^2. \quad (3.5)$$

Proof: (*Proof of Theorem 2.2*) For any t_0 in $(0, T_0)$, our main goal is to show that there is a function \mathcal{C} satisfying C) such that

$$\int_{\Omega} |A(u)Du(x, t)|^2 dx \leq \mathcal{C}(t) \quad \forall t \in (t_0, T_0). \quad (3.6)$$

Once this is established, by (3.5) and the fact that $\lambda(u)$ is bounded from below by $\lambda_0 > 0$, the integral of $\lambda(u)|Du|^2$ over Ω does not blow up in finite time. The condition (2.5) of Theorem 2.1 follows and our theorem is then proved.

For any $T \in (t_0, T_0)$, test the system for u by $A(u)u_t$ (i.e. multiplying the i^{th} equation of (1.1) by $\sum_j a_{ij}(u_j)_t$, integrating over $\Omega \times [t_0, T]$, summing the results) and integrate by parts to get

$$\iint_{\Omega \times (t_0, T)} (\langle A(u)u_t, u_t \rangle + \langle A(u)Du, D(A(u)u_t) \rangle) dz = \iint_{\Omega \times (t_0, T)} \langle f(u), A(u)u_t \rangle dz. \quad (3.7)$$

We note that

$$\langle A(u)Du, D(A(u)u_t) \rangle = \langle A(u)Du, DAu_t \rangle + \langle A(u)Du, AD(u_t) \rangle,$$

$$\frac{1}{2} \frac{\partial}{\partial t} \|ADu\|^2 = \langle A(u)Du, (A(u)Du)_t \rangle = \langle A(u)Du, A(u)_t Du \rangle + \langle A(u)Du, ADu_t \rangle.$$

Hence, by the definition (2.12) of \mathcal{A}

$$\langle A(u)Du, D(A(u)u_t) \rangle - \frac{1}{2} \frac{\partial}{\partial t} |ADu|^2 = \langle A(u)Du, DAu_t - A(u)_t Du \rangle = \mathcal{A}(u).$$

Thus, we rewrite (3.7) as

$$\iint_{\Omega \times (t_0, T)} (\langle A(u)u_t, u_t \rangle + \frac{1}{2} \frac{\partial}{\partial t} |ADu|^2) dz = \iint_{\Omega \times (t_0, T)} (\langle f(u), A(u)u_t \rangle + \mathcal{A}(u)) dz.$$

The ellipticity of $A(u)$ and integrating in t then give for any $T \in (t_0, T_0)$

$$\begin{aligned} \iint_{\Omega \times (t_0, T)} \lambda(u)|u_t|^2 dz + \frac{1}{2} \int_{\Omega} |A(u(x, T))Du(x, T)|^2 dx \leq \\ \frac{1}{2} \int_{\Omega} |A(u(x, t_0))Du(x, t_0)|^2 dx + \iint_{\Omega \times (t_0, T)} (C|f(u)|\lambda(u)|u_t| + \mathcal{A}(u)) dz. \end{aligned}$$

Using Young's inequality to find a constant $C(\varepsilon)$ such that for any $\varepsilon > 0$

$$|f(u)|\lambda(u)|u_t| \leq \varepsilon \lambda(u)|u_t|^2 + C(\varepsilon)\lambda(u)|f(u)|^2.$$

For sufficiently small and fixed ε we then have

$$\begin{aligned} & \iint_{\Omega \times (t_0, T)} \lambda(u) |u_t|^2 dz + \int_{\Omega} |A(u(x, T)) Du(x, T)|^2 dx \leq \\ & \int_{\Omega} |A(u(x, t_0)) Du(x, t_0)|^2 dx + C \iint_{\Omega \times (t_0, T)} (\lambda(u) |f(u)|^2 + \mathcal{A}(u)) dz. \end{aligned} \quad (3.8)$$

Now, let $U = P(u)$ be the function in A) and $V = \frac{\lambda^{\frac{1}{2}}(u) |f(u)|}{P(u)+1}$. Then (2.9) gives $|V| \leq CF(u)$ ($F(u)$ was also defined in A)). We observe that

$$\begin{aligned} & \int_{\Omega} \lambda(u) |f(u)|^2 dx \leq C \int_{\Omega} (U^2 + 1)(F(u)^2 + 1) dx \\ & = C \int_{\Omega} P(u)^2 F(u)^2 dx + C \int_{\Omega} P(u)^2 dx + C \int_{\Omega} F(u)^2 dx + C(d(\Omega)) \\ & \leq C \int_{\Omega} |DP(u)|^2 dx \int_{\Omega} |DF(u)|^2 dx + C \int_{\Omega} (|DP(u)|^2 + |DF(u)|^2) dx + C(d(\Omega)), \end{aligned}$$

where we used (3.4) and then Poincaré's inequality for $P(u), F(u)$ in the last estimate. By (2.7) and (3.5), we have

$$|DP(u)|^2 \leq |P_u(u)|^2 |Du|^2 \leq C \lambda^2(u) |Du|^2 \leq C \langle A^T(u) A(u) Du, Du \rangle = C |A(u) Du|^2.$$

Since $|DF(u)|^2 \leq C \lambda(u) |Du|^2$ by (2.6), we can use the above estimates in (3.8) to get

$$\begin{aligned} & \iint_{\Omega \times (t_0, T)} \lambda(u) |u_t|^2 dz + \int_{\Omega} |AD(u(x, T))|^2 dx \leq \\ & \int_{\Omega} |AD(u(x, t_0))|^2 dx + C \int_{t_0}^T \left[\int_{\Omega} \lambda(u) |Du|^2 dx + 1 \right] \int_{\Omega} |A(u) Du|^2 dx dt \quad (3.9) \\ & + C \int_{t_0}^T \int_{\Omega} \lambda(u) |Du|^2 dx dt + C(d(\Omega))(T - t_0) + C \int_{t_0}^T \int_{\Omega} \mathcal{A}(u) dx dt. \end{aligned}$$

By (3.1) and the assumption (2.13) on $\mathcal{A}(u)$, There is a continuous function $C_{\mathcal{A}}(t)$ satisfying C) such that

$$\int_{t_0}^T \int_{\Omega} \mathcal{A}(u) dx dt \leq \int_{t_0}^T \int_{\Omega} [\varepsilon \lambda(u) |u_t|^2 + C_{\varepsilon}(t) |A(u) Du|^2] dx dt + C_{\mathcal{A}}(T).$$

For a sufficiently small and fixed ε we derive from the above and (3.9) that

$$\begin{aligned} & \int_{\Omega} |DA(u(x, T))|^2 dx \leq \int_{\Omega} |DA(u(x, t_0))|^2 dx + C_{\mathcal{A}}(T) \\ & + C \int_{t_0}^T \left[\int_{\Omega} (\lambda(u) |Du|^2 + 1) dx + C_{\varepsilon}(t) \right] \int_{\Omega} |AD(u)|^2 dx dt. \end{aligned} \quad (3.10)$$

We now set

$$y(t) = \int_{\Omega} |A(u) Du(x, t)|^2 dx, \quad \alpha(t) = \int_{\Omega} |A(u) Du(x, t_0)|^2 dx + C_{\mathcal{A}}(t),$$

and

$$\beta(t) = \int_{\Omega} (\lambda(u) |Du(x, t)|^2 + 1) dx + C_{\varepsilon}(t).$$

We obtain from (3.10)

$$y(t) \leq \alpha(t) + C \int_{t_0}^t \beta(s)y(s)ds \quad \forall t \in (t_0, T_0).$$

The integral form of Gronwall's inequality gives

$$y(t) \leq \alpha(t) + C \int_{t_0}^t \alpha(s)\beta(s) \exp\left(\int_s^t \beta(\tau)d\tau\right) ds.$$

Clearly, there are functions $\mathcal{C}_1, \mathcal{C}_2$ satisfying C) such that $\alpha(t)$ is bounded by $\mathcal{C}_1(t)$ and by Lemma 3.1

$$\int_s^t \beta(\tau)d\tau \leq \iint_{\Omega \times [t_0, t]} \lambda(u)|Du|^2 dz + Ct + \iint_{\Omega \times [t_0, t]} C_\varepsilon(t) dz \leq \mathcal{C}_2(t) \quad \forall t \in (t_0, T_0).$$

We conclude that there is a function \mathcal{C}_3 satisfying C) such that

$$y(t) = \int_{\Omega} |A(u)Du(x, t)|^2 dx \leq \mathcal{C}_3(t) \quad \forall t \in (t_0, T_0).$$

This gives the desired estimate (3.6) for Du . The proof is complete. ■

Remark 3.2 If we assume F') and replace $f(u)$ by $\hat{f}(u, Du)$ satisfying

$$|\hat{f}(u, Du)| \leq C\lambda^{\frac{1}{2}}(u)|Du| + f(u)$$

then the result continues to hold. Firstly, by Young's inequality

$$\langle \hat{f}(u, Du), u \rangle \leq \varepsilon\lambda(u)|Du|^2 + C(\varepsilon)|u|^2 + |f(u)||u|.$$

For sufficiently small ε , the argument in the proof of Lemma 3.1 will lead to (the last inequality in the proof with an extra term)

$$\sup_{t \in [t_0, T]} \int_{\Omega} |u|^2 dx \leq \mathcal{C}(T) + C(\varepsilon) \iint_{\Omega \times [0, T]} |u|^2 dz,$$

for some function \mathcal{C} satisfying C). A simple use of Gronwall's inequality gives

$$\sup_{t \in [t_0, T]} \int_{\Omega} |u|^2 dx \leq C(T, \|u(\cdot, t_0)\|_{L^2(\Omega)}, d(\Omega)),$$

and the assertion of the lemma still holds.

Next,

$$\begin{aligned} |\hat{f}(u, Du)|\lambda(u)|u_t| &\leq C\lambda(u)\lambda(u)|Du||u_t| + C|f(u)|\lambda(u)|u_t| \\ &\leq \varepsilon\lambda(u)|u_t|^2 + C(\varepsilon)\lambda^2(u)|Du|^2 + C|f(u)|\lambda(u)|u_t|. \end{aligned}$$

As $f(u)$ satisfies F), for small ε in the above the proof of Theorem 2.2 can continue.

We now consider the excess \mathcal{A} . If $A(u) = [a_{ij}(u)]$ then calculations give

$$\begin{aligned} A(u)Du &= \left[\sum_j a_{ij}(u)Du_j \right], \\ DA(u)u_t &= \left[\sum_k \frac{\partial}{\partial u_k} a_{ij}(u)Du_k \right] u_t = \left[\sum_{k,j} \frac{\partial}{\partial u_k} a_{ij}(u)Du_k(u_j)_t \right], \\ A(u)_t Du &= \left[\sum_k \frac{\partial}{\partial u_k} a_{ij}(u)(u_k)_t \right] Du = \left[\sum_{k,j} \frac{\partial}{\partial u_k} a_{ij}(u)(u_k)_t Du_j \right]. \end{aligned}$$

Thus,

$$DA(u)u_t - A(u)_t Du = \left[\sum_{k,j} \frac{\partial}{\partial u_k} a_{ij}(u)[Du_k(u_j)_t - (u_k)_t Du_j] \right].$$

In general, we have

$$\begin{aligned} \mathcal{A}(u) &= \langle A(u)Du, DA(u)u_t - A(u)_t Du \rangle \\ &= \sum_i \sum_l a_{il}(u)Du_l \sum_{k,j} \frac{\partial}{\partial u_k} a_{ij}(u)[Du_k(u_j)_t - (u_k)_t Du_j] \\ &= \sum_i \sum_{l,k,j} a_{il}(u) \frac{\partial}{\partial u_k} a_{ij}(u)[Du_k(u_j)_t - (u_k)_t Du_j] Du_l. \end{aligned}$$

Applying Theorem 2.2 to special structures of $A(u)$ we have the proof of the last two theorems.

Proof: (*Proof of Theorem 2.3*) If $\frac{\partial}{\partial u_k} a_{ij}(u) = \frac{\partial}{\partial u_i} a_{ik}(u)$ for any i and $k \neq j$ then it is clear that $\mathcal{A}(u) = 0$. This is the case if there are functions \mathcal{P}_i 's such that $a_{ij}(u) = \partial_{u_j} \mathcal{P}_i(u)$. The condition (2.13) is satisfied and our theorem follows from Theorem 2.2. ■

Proof: (*Proof of Theorem 2.4*) We assume that the i -th equation of the system (1.1) does not depend on the unknowns u_k if $k > i$. This means $a_{ik} = 0$ and $\frac{\partial}{\partial u_k} a_{ij}(u) = \frac{\partial}{\partial u_k} f_i(u) = 0$ if $k > i$.

Therefore, for the subsystem of the first k_0 equations ($k_0 \geq 1$), we have, writing $u = (u_1, \dots, u_{k_0})^T$

$$\begin{aligned} A_{k_0}(u)Du &= \left[\sum_{j \leq k_0} a_{ij}(u)Du_j \right], \\ DA_{k_0}(u)u_t &= \left[\sum_{j,k \leq k_0} \frac{\partial}{\partial u_k} a_{ij}(u)Du_k \right] u_t = \left[\sum_{k,j \leq k_0} \frac{\partial}{\partial u_k} a_{ij}(u)Du_k(u_j)_t \right], \\ A_{k_0}(u)_t Du &= \left[\sum_{k \leq k_0} \frac{\partial}{\partial u_k} a_{ij}(u)(u_k)_t \right] Du = \left[\sum_{k,j \leq k_0} \frac{\partial}{\partial u_k} a_{ij}(u)(u_k)_t Du_j \right]. \end{aligned}$$

Accordingly, the excess for the subsystem is

$$\begin{aligned} \mathcal{A}_{k_0}(u) &= \sum_{i,l,k,j \leq k_0} a_{il}(u) \frac{\partial}{\partial u_k} a_{ij}(u)[Du_k(u_j)_t - (u_k)_t Du_j] Du_l \\ &= I_{1,k_0} + I_{2,k_0} + I_{3,k_0}. \end{aligned}$$

Here,

$$I_{1,k_0} = \sum_{j < k_0} a_{k_0 k_0}(u) \frac{\partial}{\partial u_{k_0}} a_{k_0 j}(u) [Du_{k_0}(u_j)_t - (u_{k_0})_t Du_j] Du_{k_0},$$

$$I_{2,k_0} = \sum_{j, l < k_0} a_{k_0 l}(u) \frac{\partial}{\partial u_{k_0}} a_{k_0 j}(u) [Du_{k_0}(u_j)_t - (u_{k_0})_t Du_j] Du_l,$$

and

$$I_{3,k_0} = \sum_{i, j, k, l < k_0} a_{il}(u) \frac{\partial}{\partial u_k} a_{ij}(u) [Du_k(u_j)_t - (u_k)_t Du_j] Du_l.$$

It is clear that $\mathcal{A}_1 = 0$ so that u_1 exists globally and its derivatives do not blow up in finite time. We then argue by induction. Consider the induction hypothesis:

For some $k_0 \geq 1$ there is a continuous function \mathcal{C} satisfying C) such that

$$\mathbf{(K0)} \quad \|(u_i(\cdot, t))_t\|_{L^\infty(\Omega)}, \|u_i(\cdot, t)\|_{C^1(\Omega)} \leq \mathcal{C}_{k_0}(t) \quad \forall t \in (0, T_0), i = 1, \dots, k_0 - 1. \quad (3.11)$$

As we assume there is a constant C such that $|A(u)| \leq C\lambda(u)$ and (see (2.14))

$$\left| \frac{\partial}{\partial u_{k_0}} a_{k_0 j}(u) \right| \leq C\lambda^{\frac{1}{2}}(u) \quad \forall j < k_0, k_0 = 2, \dots, m.$$

Then by Young's inequality and our induction assumption (3.11)

$$I_{1,k_0}, I_{2,k_0} \leq \varepsilon\lambda(u)|u_t|^2 + C(\varepsilon)\mathcal{C}(t)\lambda^2(u)|Du|^2 + C(\varepsilon)\mathcal{C}(t) \quad \forall t \in (t_0, T_0)$$

for some continuous function \mathcal{C} depending on \mathcal{C}_{k_0} . Similarly, since I_{3,k_0} does not depend on u_{k_0} , (3.11) also shows that

$$\iint_{\Omega \times (t_0, t)} I_{3,k_0} dz \leq \mathcal{C}(t) \quad \forall t \in (t_0, T_0).$$

By (3.5), $\lambda^2(u)|Du|^2 \leq |A(u)Du|^2$. Therefore, we can conclude from the above estimates that there are continuous functions $\mathcal{C}, C_{\mathcal{A}}$ satisfying C) such that

$$\int_{t_0}^s \int_{\Omega} \mathcal{A}(u) dx dt \leq \int_{t_0}^s \int_{\Omega} [\varepsilon\lambda(u)|u_t|^2 + C(\varepsilon)\mathcal{C}(t)|A(u)Du|^2] dx dt + C_{\mathcal{A}}(s)$$

for all $s \in (t_0, T_0)$. Hence, Theorem 2.2 can be applied to the subsystem of the first k_0 equations and we see that $u = (u_1, \dots, u_{k_0})^T$ does not blow up in finite time. Therefore, (3.11) holds again for $k_0 + 1$ and our proof is complete by induction. ■

4 Proof of Theorem 2.1

In the sequel, we will denote $\Phi(u) = \frac{|\lambda_u(u)|^2}{\lambda(u)}$. Before going to the proof, we need some estimates for the integral of $\Phi(u)|Du|^{4p}$.

Lemma 4.1 *Assume (2.2) in A). For any $p \geq 1$ and any nonnegative function $\psi \in C_0^1(B_R)$ there is a constant C_1 such that*

$$\begin{aligned} \int_{B_R} \Phi(u) |Du|^{4p} \psi^4 dx &\leq C \int_{\Omega} \lambda^2(u) |Du|^{4p} \psi^4 dx \leq C_1 \int_{B_R} \lambda(u) |Du|^{2p} \psi^2 dx \times \\ &\int_{B_R} (\lambda(u) |Du|^{2p-2} |D^2u|^2 \psi^2 + \Phi(u) |Du|^{2p+2} \psi^2 + \lambda(u) |D\psi|^2 |Du|^{2p}) dx. \end{aligned} \quad (4.1)$$

Proof: By Ladyzhenskaya's inequality (3.3) with $U = \lambda^{\frac{1}{2}}(u) \psi |Du|^{p-1} Du$ we have

$$\begin{aligned} \int_{B_R} \lambda^2(u) |Du|^{4p} \psi^4 dx &= \int_{B_R} |U|^4 dx \leq C \int_{B_R} |U|^2 dx \int_{B_R} |DU|^2 dx \\ &\leq C \int_{B_R} \lambda(u) |Du|^{2p} \psi^2 dx \int_{B_R} |D(\lambda^{\frac{1}{2}}(u) \psi |Du|^{p-1} Du)|^2 dx. \end{aligned}$$

It is clear that there is a constant C_2 such that

$$\begin{aligned} |D(\lambda^{\frac{1}{2}}(u) \psi |Du|^{p-1} Du)|^2 &\leq \\ &C_2 \left[\lambda(u) |Du|^{2p-2} |D^2u|^2 \psi^2 + \frac{|\lambda_u(u)|^2}{\lambda(u)} |Du|^{2p+2} \psi^2 + \lambda(u) |D\psi|^2 |Du|^{2p} \right]. \end{aligned}$$

Because $\lambda(u)$ is bounded from below, (2.2) gives $\Phi(u) \leq C\lambda^2(u)$ for some constant C . Applying the above inequality in the previous estimate, we obtain the lemma. ■

Since u are C^2 in x , we can differentiate (1.1) with respect to x to get

$$(Du)_t = \operatorname{div}((A(u)D^2u + A_u(u)DuDu) + Df(u, Du)). \quad (4.2)$$

Furthermore, by [4, Lemma 2.1], if A is a matrix satisfying $\lambda_0|\zeta|^2 \leq \langle A\zeta, \zeta \rangle$ and $|A\zeta| \leq \Lambda_0|\zeta|$ then for any α and $\delta_\alpha \in (0, 1)$ are numbers such that $\frac{\alpha}{2+\alpha} = \delta_\alpha \frac{\lambda_0}{\Lambda_0}$ then there is a positive constant $\hat{\lambda}$ depending on $\lambda_0, \Lambda_0, \delta_\alpha$ such that

$$\langle AD\zeta, D(\zeta|\zeta|^\alpha) \rangle \geq \hat{\lambda}|\zeta|^\alpha |D\zeta|^2. \quad (4.3)$$

We also recall the following elementary iteration result (e.g., see [8, Lemma 6.1, p.192]).

Lemma 4.2 *Let f, g, h be bounded nonnegative functions in the interval $[\rho, R]$ with g, h being increasing. Assume that for $\rho \leq s < t \leq R$ we have*

$$f(s) \leq [(t-s)^{-\alpha} g(t) + h(t)] + \varepsilon f(t)$$

with $C \geq 0$, $\alpha > 0$ and $0 \leq \varepsilon < 1$. Then

$$f(\rho) \leq c(\alpha, \varepsilon)[(R-\rho)^{-\alpha} g(R) + h(R)].$$

The constant $c(\alpha, \varepsilon)$ can be taken to be $(1-\nu)^{-\alpha}(1-\nu^{-\alpha}\nu_0)^{-1}$ for any ν satisfying $\nu \in (0, 1)$ and $\nu^{-\alpha}\nu_0 < 1$.

Proof: Let $B_R = B_R(x_0)$ be a ball in Ω . We consider only the interior case because the boundary case, when the center x_0 of B_R is on the boundary $\partial\Omega$, is similar. For any s, t such that $0 \leq s < t \leq R$ let ψ be a cutoff function for two balls B_s, B_t centered at x_0 . That is, $\psi \equiv 1$ in B_s and $\psi \equiv 0$ outside B_t with $|D\psi| \leq 1/(t-s)$. In the sequel, we will fix two reals t_0, T such that $0 < t_0 < T < T_0$.

We divide the proof in three steps.

Step 1: (Local energy estimates) Clearly, by uniform ellipticity of $A(u)$, we can find a constant C_0 such that $|A(u)\zeta| \leq C_0\lambda(u)|\zeta|$. Thus, for any $p > 1$ there is $\delta_p \in (0, 1)$ such that $\alpha = 2p - 2$ satisfies

$$\frac{\alpha}{2 + \alpha} = \frac{2p - 2}{2p} = \delta_p C_0^{-1} = \delta_p \frac{\lambda(u)}{C_0 \lambda(u)}. \quad (4.4)$$

Testing (4.2) with $|Du|^{2p-2} Du \psi^2$. By the above, there is a positive constant $C(p)$ (see (4.3)) such that for $Q = \Omega \times [t_0, T_0]$

$$\begin{aligned} & \sup_{\tau \in (t_0, T)} \int_{\Omega} |Du|^{2p} \psi^2 dx + C(p) \iint_Q \lambda(u) |Du|^{2p-2} |D^2 u|^2 \psi^2 dz \leq \\ & \iint_Q |A(u)| |D^2 u| |Du|^{2p-1} \psi |D\psi| dz - \iint_Q A_u(u) Du Du D(|Du|^{2p-2} Du \psi^2) dz \\ & + \iint_Q D\hat{f}(u, Du) |Du|^{2p-2} Du \psi^2 dz + \int_{\Omega} |Du(x, t_0)|^{2p} \psi^2 dx. \end{aligned}$$

For simplicity, we will assume in the sequel that $\hat{f} \equiv 0$. The presence of \hat{f} will be discussed later in Remark 4.3. For any given positive ε we use Young's inequality to find a constant $C(\varepsilon)$ such that

$$\begin{aligned} |A(u)| |D^2 u| |Du|^{2p-1} \psi |D\psi| & \leq \varepsilon \lambda(u) |Du|^{2p-2} |D^2 u|^2 \psi^2 + C(\varepsilon) \lambda(u) |Du|^{2p} |D\psi|^2, \\ |A_u(u) Du Du D(|Du|^{2p-2} Du \psi^2)| & \leq |A_u(u)| |Du|^{2p} |D^2 u| \psi^2 + |A_u(u)| |Du|^{2p+1} \psi |D\psi| \\ & \leq \varepsilon \lambda(u) |Du|^{2p-2} |D^2 u|^2 + C(\varepsilon) \frac{|A_u|^2}{\lambda(u)} |Du|^{2p+2} \psi^2 + C(\varepsilon) \lambda(u) |Du|^{2p} |D\psi|^2. \end{aligned}$$

Therefore, taking ε small and the above two inequalities in the previous one, we easily deduce

$$\begin{aligned} & \sup_{\tau \in (t_0, T)} \int_{B_s} |Du|^{2p} dx + \iint_{Q_s} \lambda(u) |Du|^{2p-2} |D^2 u|^2 dz \leq \\ & C_1 \iint_{Q_t} \Phi(u) |Du|^{2p+2} \psi^2 dz + \frac{C_1}{(t-s)^2} \iint_{Q_t} \lambda(u) |Du|^{2p} dz + C(t_0). \end{aligned} \quad (4.5)$$

Here, we used the definition of ψ and $\Phi(u)$ and the fact that $|A_u| \sim \lambda_u$, and denoted

$$Q_t = B_t \times (t_0, T), \quad C(t_0) = \int_{\Omega} |Du(x, t_0)|^{2p} dx.$$

We now set

$$\mathcal{A}_p(t) = \sup_{\tau \in (t_0, T)} \int_{B_t} |Du|^{2p} dx, \quad \mathcal{H}_p(t) = \iint_{Q_t} \lambda(u) |Du|^{2p-2} |D^2 u|^2 dz,$$

$$\mathcal{B}_p(t) = \iint_{Q_t} \Phi(u) |Du|^{2p+2} dz, \quad \mathcal{G}_p(t) = \iint_{Q_t} \lambda(u) |Du|^{2p} dz + C(t_0).$$

So that, for any $p \geq 1$ and satisfies (4.4) and t is small, (4.5) can be rewritten as

$$\mathcal{A}_p(s) + \mathcal{H}_p(s) \leq C_1 \mathcal{B}_p(t) + \frac{C_1}{(t-s)^2} \mathcal{G}_p(t) \quad \forall s, t \text{ such that } 0 < s < t \text{ and } B_t \subset \Omega. \quad (4.6)$$

Step 2: (Estimates for the integral of $|Du|^4$ over $\Omega \times (0, T)$) By (2.5) and the uniform continuity of integrals (see Remark 4.4 following the proof) give for any given $\varepsilon_0 > 0$ a constant $R(\varepsilon_0, T)$ such that

$$\int_{B_R} \lambda(u) |Du(x, \tau)|^2 dx \leq \varepsilon_0 \quad \forall R < R(\varepsilon_0, T), \tau \in [t_0, T]. \quad (4.7)$$

Therefore, let $p = 1$ and ψ be the cutoff function for B_s, B_t in (4.1) and use the definition of $\psi, \mathcal{B}_p, \mathcal{H}_p$ and \mathcal{C}_p to have by integrating in $[t_0, T]$

$$\mathcal{B}_1(s) \leq \sup_{\tau \in [t_0, T]} \int_{B_R} \lambda(u) |Du(x, \tau)|^2 dx \left(\mathcal{H}_1(t) + \mathcal{B}_1(t) + \frac{1}{(t-s)^2} \mathcal{G}_1(t) \right)$$

for all s, t such that $0 < s < t < R(\varepsilon_0, T)$. We now choose ε_0 sufficiently small in (4.7) to have a number $\mu_0 \in (0, 1)$ such that

$$C_1 \mathcal{B}_1(s) \leq \frac{\mu_0}{2} \left(\mathcal{H}_1(t) + \mathcal{B}_1(t) + \frac{1}{(t-s)^2} \mathcal{G}_1(t) \right). \quad (4.8)$$

For $p = 1$, (4.6) gives

$$\mathcal{H}_1(s) \leq C_1 \mathcal{B}_1(t) + \frac{C_1}{(t-s)^2} \mathcal{G}_1(t), \quad 0 < s < t < R(\varepsilon_0, T).$$

Let $t_1 = (s+t)/2$ and use (4.8) with s being t_1 and the above with t being t_1 to obtain

$$\mathcal{H}_1(s) \leq \frac{\mu_0}{2} [\mathcal{H}_1(t) + \mathcal{B}_1(t)] + \frac{C_2}{(t-s)^2} \mathcal{G}_1(t). \quad (4.9)$$

Obviously, we can assume $C_1 \geq 1$ so that we can add (4.8) and (4.9) to have

$$\mathcal{H}_1(s) + \mathcal{B}_1(s) \leq \mu_0 [\mathcal{H}_1(t) + \mathcal{B}_1(t)] + \frac{C_3}{(t-s)^2} \mathcal{G}_1(t), \quad 0 < s < t < R(\varepsilon_0, T).$$

Since $\mu_0 \in (0, 1)$, we can use Lemma 4.2 with $f(t) = \mathcal{H}_1(t) + \mathcal{B}_1(t)$, $h(t) = 0$, $g(t) = \mathcal{G}_1(t)$ and $\alpha = 2$ to obtain a constant C_4 depending on μ_0, C_3 such that

$$\mathcal{H}_1(s) + \mathcal{B}_1(s) \leq \frac{C_4}{(t-s)^2} \mathcal{G}_1(t), \quad 0 < s < t < R(\varepsilon_0, T).$$

For $R = R(\varepsilon_0, T)/4$, the above with $s = R$, $t = 2R$ gives

$$\mathcal{H}_1(R) + \mathcal{B}_1(R) \leq \frac{C_4}{R^2} \iint_{Q_{2R}} \lambda(u) |Du|^2 dz + C(t_0). \quad (4.10)$$

Hence, by (4.8) and the estimate for the integral of $\lambda(u)|Du|^2$ over Q , we have

$$\iint_{Q_R} \Phi(u)|Du|^4 dz \leq C(T, R(\varepsilon_0), \|u(\cdot, t_0)\|_{C^1(\Omega)}) \quad \forall R < R(\varepsilon_0, T). \quad (4.11)$$

We also note that (4.10), (4.11) and the second inequality of (4.1) gives

$$\iint_{Q_R} \lambda^2(u)|Du|^4 dz \leq C(T, R(\varepsilon_0), \|u(\cdot, t_0)\|_{C^1(\Omega)}) \quad \forall R < R(\varepsilon_0, T). \quad (4.12)$$

Finite covering of Ω by balls $B_{R(\varepsilon_0, T)/2}$ yields

$$\iint_Q \lambda^2(u)|Du|^4 dz \leq C(T, R(\varepsilon_0, T), \Omega, \|u(\cdot, t_0)\|_{C^1(\Omega)}) \quad \forall T \in (0, T_0). \quad (4.13)$$

Step 3: (Estimates for the integral of $|Du|^{2p}$ over Ω) For any $p > 1$ we have by Hölder's inequality

$$\iint_{Q_t} \Phi(u)|Du|^{2p+2}\psi^2 dz \leq \left(\iint_{Q_t} \Phi^2(u)|Du|^4 dz \right)^{\frac{1}{2}} \left(\iint_{Q_t} |Du|^{4p}\psi^4 dz \right)^{\frac{1}{2}}. \quad (4.14)$$

Using Ladyzhenskaya's inequality (3.3) with $U = |Du|^{p-1}Du\psi$ and integrating the result over (t_0, T) , we have

$$\iint_{Q_t} |Du|^{4p}\psi^4 dz \leq C \sup_{\tau \in (t_0, T)} \int_{B_t} |Du|^{2p}\psi^2 dx \iint_{Q_t} |DU|^2 dz$$

Since $\lambda(u)$ is bounded from below by λ_0 , there is a constant $C(\lambda_0)$ such that

$$|DU|^2 \leq C(\lambda_0)[\lambda(u)|Du|^{2p-2}|D^2u|^2\psi^2 + \lambda(u)|Du|^{2p}|D\psi|^2].$$

Therefore,

$$\iint_{Q_t} |Du|^{4p}\psi^4 dz \leq C\mathcal{A}_p(t)[\mathcal{H}_p(t) + \frac{1}{(t-s)^2}\mathcal{G}_p(t)] \leq C[\mathcal{A}_p(t) + \mathcal{H}_p(t) + \frac{1}{(t-s)^2}\mathcal{G}_p(t)]^2.$$

Here, Cauchy's inequality was used in the last inequality. Using the above estimate in (4.14) and the fact that $\Phi^2(u) \leq C\lambda^2(u)$, we derive

$$\iint_{Q_t} \Phi(u)|Du|^{2p+2}\psi^2 dz \leq C \left(\iint_{Q_t} \lambda^2(u)|Du|^4\psi^4 dz \right)^{\frac{1}{2}} [\mathcal{A}_p(t) + \mathcal{H}_p(t) + \frac{1}{(t-s)^2}\mathcal{G}_p(t)].$$

By (4.13) and the continuity of integrals, the first factor on the right can be as small as we please if t is small. Hence, for any given $\mu_1 \in (0, 1)$, and the definition of \mathcal{B}_p , if $R \leq R(\mu_1, T)$ for some small $R(\mu_1, T)$ then the above gives

$$C_1 \iint_{Q_t} \Phi(u)|Du|^{2p+2}\psi^2 dz \leq \mu_1 [\mathcal{A}_p(t) + \mathcal{H}_p(t) + \frac{1}{(t-s)^2}\mathcal{G}_p(t)] \quad \text{for some } \mu_1 \in (0, 1).$$

If $p > 1$ and satisfies (4.4), we then have from (4.5) and the above inequality the following.

$$\mathcal{A}_p(s) + \mathcal{H}_p(s) \leq \mu_1(\mathcal{A}_p(t) + \mathcal{H}_p(t)) + \frac{1}{(t-s)^2} \mathcal{G}_p(t), \quad 0 < s < t < R(\mu_1, T).$$

For $f(t) = \mathcal{A}_p(t) + \mathcal{H}_p(t)$, $h(t) = 0$, $g(t) = \mathcal{G}_p(t)$ and $\alpha = 2$ we can use Lemma 4.2, as $\mu_1 \in (0, 1)$, to obtain

$$\mathcal{F}(\rho) \leq \frac{C_5(\mu_1)}{(R-\rho)^2} \mathcal{G}_p(R), \quad 0 < \rho < R < R(\mu_1, T).$$

We can assume that $2p < 4$. Because $\lambda(u)$ is bounded from below, by (4.12) and a simple use of Hölder's inequality, we can see that $\mathcal{G}_p(R)$ is bounded, using (4.12). Hence, the above yields

$$\sup_{t \in (t_0, T)} \int_{B_\rho} |Du|^{2p} dx + \iint_{Q_\rho} \lambda(u) |Du|^{2p-2} |D^2u|^2 dz \leq C(\rho, R, \|u(\cdot, t_0)\|_{C^1(\Omega)}) \quad (4.15)$$

if $0 < \rho < R$ and R is sufficiently small and some $p > 1$. Finite covering of Ω with balls $B_{R/2}$ yields

$$\sup_{t \in (t_0, T)} \int_{\Omega} |Du|^{2p} dx + \iint_Q \lambda(u) |Du|^{2p-2} |D^2u|^2 dz \leq C(\Omega, R, \|u(\cdot, t_0)\|_{C^1(\Omega)}). \quad (4.16)$$

Since $2p > 2$, Sobolev's imbedding theorem shows that u is Hölder continuous in x . From the system for u and the above, with $p = 1$, we see that u_t is in $L^2(Q)$. It is now standard to show that u is Hölder in (x, t) and Du is Hölder continuous. We now can refer to Amann's results to see that u exists globally. ■

Remark 4.3 If we replace $f(u)$ by a function \hat{f} depending on u, Du and satisfying a linear growth in Du then the proof can go on with minor modification. Namely, there exist a constant C and a function $f(u)$ satisfying F) such that

$$|\hat{f}(u, Du)| \leq C\lambda^{\frac{1}{2}}(u)|Du| + f(u).$$

We can assume that $|D\hat{f}(u, Du)| \leq C|D(\lambda^{\frac{1}{2}}(u)|Du)| + |f_u(u)||Du|$ so that

$$|D\hat{f}(u, Du)| \leq C\lambda^{\frac{1}{2}}|D^2u| + C\Phi^{\frac{1}{2}}(u)|Du|^2 + |f_u(u)||Du|.$$

Therefore, in **Step 1**, the extra term $|D\hat{f}(u, Du)||Du|^{2p-1}\psi^2$ can be handled by using the following estimates, which are the results of a simple use of Young's inequality.

$$\begin{aligned} |D\hat{f}(u, Du)||Du|^{2p-1} &\leq C[\lambda^{\frac{1}{2}}|D^2u| + C\Phi^{\frac{1}{2}}(u)|Du|^2 + |f_u(u)||Du|]|Du|^{2p-1} \\ &\leq \varepsilon\lambda|Du|^{2p-2}|D^2u|^2 + C(\varepsilon)\lambda|Du|^{2p} + \\ &\quad C\Phi(u)|Du|^{2p+2} + C|Du|^{2p} + C|f_u||Du|^{2p} \end{aligned} .$$

We can then assume that $|f_u| \leq C\lambda(u)$ for some constant C and see that the proof can continue to obtain the energy estimate (4.5). The result then follows.

Remark 4.4 The existence of $R = R(\varepsilon_0, T)$ in (4.7) is an easy consequence of a simple application of Hahn-Saks-Vitali's theorem and an argument by contradiction. Indeed, if there is no such uniform R for (4.7) to hold then there is a sequence $\{s_n\}$ in $[t_0, T]$ such that the integrals of $f_n := \lambda(u(\cdot, s_n))|Du(\cdot, s_n)|^2$ over $B_{\frac{1}{n}} \times \{t_n\}$ is greater than ε_0 . We can assume that $\{s_n\}$ converges in $[t_0, T]$. Since $\lambda(u)|Du|^2$ is continuous in $t \in [t_0, T]$, Hahn-Saks-Vitali's theorem, e.g. see [6], applies to the sequence f_n and shows the uniform continuity in R of the integral of $\lambda(u)|Du|^2$ over $B_R \times \{t_n\}$ and gives a contradiction.

5 Further Discussion

To compare Theorem 2.1 with some results in our earlier work [12], where we considered general dimension $n \geq 2$, let us recall the following results in [12]. There, we assumed the following structural conditions on the system (1.1)

A.1) (Uniform ellipticity) There are positive constants C, λ_0 and a smooth function $\lambda(u)$ such that $\lambda(u) \geq \lambda_0$ and

$$\lambda(u)|\xi|^2 \leq \langle A(u)\xi, \xi \rangle \leq C\lambda(u)|\xi|^2 \quad \forall u \in \mathbb{R}^m, \xi \in \mathbb{R}^{nm}.$$

A.2) Assume that $A \in C^1(\mathbb{R}^m)$. Let Φ_0, Φ be defined as

$$\Phi_0(u) = \lambda^{\frac{1}{2}}(u) \text{ and } \Phi(u) = \frac{|A_u(u)|}{\lambda^{\frac{1}{2}}(u)} \quad u \in \mathbb{R}^m.$$

Assume that the quantities

$$k_1 := \sup_{u \in \mathbb{R}^m} \frac{|\Phi_u|}{\Phi}, \quad k_2 := \sup_{u \in \mathbb{R}^m} \frac{\Phi}{\Phi_0} \quad (5.1)$$

are finite.

A.3) (Weights) If $u \in BMO(\Omega)$ then $\Phi^{\frac{2}{3}}(u)$ belongs to the $A_{\frac{4}{3}}$ class and the quantity $[\Phi(u)^{\frac{2}{3}}]_{\frac{4}{3}}$ can be controlled by the norm $\|u\|_{BMO(\Omega)}$.

As we discussed in [12], if $A(u)$ has a polynomial growth in u then A.2) is easily satisfied. In the general case, our assumption (2.2) in this paper is clearly much weaker than (5.1). Concerning the verification of A.3), a crucial factor for the validity of a generalized weighted Gagliardo-Nirenberg inequality involving BMO norms, extending a result in [14], we used a connection between BMO functions and A_γ weights (see [9, 13]) to see that if $\lambda(u) \sim (1 + |u|)^k$ for $0 \leq k < 5$ then A.3) holds. In this paper, we don't need such weighted Gagliardo-Nirenberg inequality so that we can allow k to be any nonnegative number.

We also assumed in [12] that the ellipticity constants $\lambda(u), \Lambda(u)$ of the matrix $A(u)$ were not too far apart.

R) (The ratio condition) There is $\delta \in [0, 1)$ such that

$$\frac{n-2}{n} = \delta \sup_{u \in \mathbb{R}^m} \frac{\lambda(u)}{\Lambda(u)}. \quad (5.2)$$

One should note that there are examples in [4] of blow up solutions to (1.1) if the condition R) is violated. In this paper, when $n = 2$, (5.2) is clearly not needed.

We assumed in [12] the following growth conditions on the nonlinearity f .

F) There are positive constants C, b such that for any vector valued functions $u \in C^1(\Omega, \mathbb{R}^m)$ and $p \in C^1(\Omega, \mathbb{R}^{mn})$

$$|f(u, p)| \leq C|p| + C|u|^b + C. \quad (5.3)$$

$$|Df(u, p)| \leq C|Dp| + C|u|^{b-1}|Du| + C. \quad (5.4)$$

Under the above structural conditions, we proved the following global existence result of classical solutions and improved Theorem 1.1.

Theorem 5.1 (Theorem 2.5 in [12]) *Assume A.1)-A.3), R) and F). Let $p_0 \in (n, \infty)$ and U_0 be in $W^{1,p_0}(\Omega)$. Suppose that $T_0 \in (0, \infty]$ is the maximal existence time for a classical solution*

$$u \in C([0, T_0), W^{1,p_0}(\Omega)) \cap C^{1,2}((0, T_0) \times \bar{\Omega})$$

for the system (1.1). Suppose that there is a function C in $C^0((0, T_0])$ such that

$$\|u(\cdot, t)\|_{BMO(\Omega)} \leq C(t) \quad \forall t \in (0, T_0).$$

Moreover, for any $\varepsilon > 0$ and $(x, t) \in Q$, there exists $R = R(\varepsilon) > 0$ such that

$$\|u(\cdot, t)\|_{BMO(B_R(x))} < \varepsilon \quad \forall t \in (0, T_0). \quad (5.5)$$

Then $T_0 = \infty$.

By Poincaré's inequality, the following consequence follows easily.

Corollary 5.2 (Corollary 2.6 in [12]) *In addition to the assumptions of Theorem 1.1, we assume R). Then there exists a maximal time $T_0 \in (0, \infty]$ such that the system (1.1) has a unique classical solution in $(0, T_0)$ with*

$$u \in C([0, T_0), W^{1,p_0}(\Omega)) \cap C^{1,2}((0, T_0) \times \bar{\Omega})$$

Moreover, if $T_0 < \infty$ then

$$\lim_{t \rightarrow T_0^-} \|u(\cdot, t)\|_{W^{1,n}(\Omega)} = \infty. \quad (5.6)$$

As we discussed earlier, our structural conditions for (1.1) in this paper, when $n = 2$, is much more general than those of [12] to obtain the same conclusion of the above corollary in Theorem 2.1. Although (2.5) is slightly stronger than (5.6) but we have proved that it could be verified in many applications. In fact, it is possible to assume (5.6) and obtain Theorem 2.1 under a bit stronger assumption than that of A) in this paper.

References

- [1] N.D. Alikakos An application of the invariance principle to reaction-diffusion equations, *J. Differential Equations*, 33 (1979), pp. 201-225.
- [2] H. Amann. Dynamic theory of quasilinear parabolic equations II. Reaction-diffusion systems, *Differential Integral Equations*, Vol. 3, no. 1 (1990), pp. 13-75.
- [3] H. Amann. Dynamic theory of quasilinear parabolic systems III. Global existence, *Math Z.* 202 (1989), pp. 219-250.
- [4] S. Ahmad and D. Le. Global and Blow Up Solutions to Cross Diffusion Systems. Nonlinear Analysis Series A: TMA. In press.
- [5] A. Friedman. *Partial Differential Equations*. New York, 1969.
- [6] N. Dunford and J.T. Schwartz. *Linear operators, Part I*. Interscience (1958)
- [7] B. Franchi, C. Perez and R. L. Wheeden. Self-Improving Properties of John Nirenberg and Poincaré Inequalities on Spaces of Homogeneous Type. *J. Functional Analysis*, 153, 108–146, 1998.
- [8] E. Giusti. *Direct Methods in the Calculus of Variations*. World Scientific, 2003.
- [9] R. L. Johnson and C. J. Neugebauer. Properties of BMO functions whose reciprocals are also BMO. *Z. Anal. Anwendungen*, 12(1):3-11, 1993.
- [10] D. Le. Regularity of BMO weak solutions to nonlinear parabolic systems via homotopy. *Trans. Amer. Math. Soc.* 365 (2013), no. 5, 2723–2753.
- [11] D. Le. Global existence results for near triangular nonlinear parabolic systems. *Adv. Nonlinear Studies.* 13 (2013), no. 4, 933-944.
- [12] D. Le. Global Existence and Regularity Results for Strongly Coupled Nonregular Parabolic Systems via Iterative Methods. *submitted*.
- [13] J. Orobitg and C. Pérez. A_p weights for nondoubling measures in \mathbb{R}^n and applications. *newblock Transactions of the American mathematical society*, 354 (2002), 2013-2033.
- [14] P. Strzelecki. Gagliardo Nirenberg inequalities with a BMO term. *Bull. London Math. Soc.* Vol. 38, pp. 294-300, 2006.
- [15] N. Shigesada, K. Kawasaki and E. Teramoto. *Spatial segregation of interacting species*. *J. Theor. Biol.*, 79(1979), 83– 99.
- [16] E. M. Stein. *Harmonic Analysis, Real Variable Methods, Orthogonality and Oscillatory Integrals*. Princeton Univ. Press, Princeton, NJ, 1993.
- [17] A. Yagi. *Global solution to some quasilinear parabolic systems in population dynamics*. *Nonlin. Anal.* 21 (1993), 603-630.