

ANGLE CRITERIA FOR UNIFORM CONVERGENCE OF AVERAGED PROJECTIONS AND CYCLIC OR RANDOM PRODUCTS OF PROJECTIONS

IZHAR OPPENHEIM

ABSTRACT. We apply a new notion of angle between projections to deduce criteria for uniform convergence results of the alternating projections method under several different settings: averaged projections, cyclic products, quasi-periodic products and random products.

Mathematics Subject Classification (2010). Primary 46B28, Secondary 41A65, 46A32.

1. INTRODUCTION

Given a set of orthogonal projections $\{P_1, \dots, P_n\}$ in a Hilbert space \mathcal{H} there is a well-known iterative method called the alternating projections method for approximating the orthogonal projection of some $v \in \mathcal{H}$ on $\text{Im}(P_1) \cap \dots \cap \text{Im}(P_n)$. There are several variations of this method, but perhaps its simplest form is to consider cyclic products of P_1, \dots, P_n , i.e., to look at the sequence $(P_n \dots P_1)^i$. In [9], Halperin showed that $(P_n \dots P_1)^i$ converges to the projection on $\text{Im}(P_1) \cap \dots \cap \text{Im}(P_n)$ in the strong operator topology, i.e., he showed that for every $v \in \mathcal{H}$

$$\lim_{i \rightarrow \infty} \|(P_n \dots P_1)^i v - P_{1, \dots, n} v\| = 0,$$

where $P_{1, \dots, n}$ is the orthogonal projection on $\text{Im}(P_1) \cap \dots \cap \text{Im}(P_n)$ (when $n = 2$ this result is due to von Neumann [22]).

Extending Halperin's result, one can consider non-cyclic sequences of the projections $\{P_1, \dots, P_n\}$. This can be described as follows - take $\tau \in \{1, \dots, n\}^{\mathbb{N}}$ and consider the sequence $\dots P_{\tau(i)} \dots P_{\tau(1)}$. Sakai [21] showed that if τ is quasi-periodic (see definition below) then the result of Halperin holds. Amemiya and Andô [1] showed that if for every $1 \leq k \leq n$, the set $\{j : \tau(j) = k\}$ is infinite, then $\dots P_{\tau(i)} \dots P_{\tau(1)}$ converges to $P_{1, \dots, n}$ in the weak-operator topology. This result was generalized by several authors, who gave sufficient conditions for stronger forms of convergence in the Hilbert or Banach settings (see [5], [7], [8], [4], [6]). On the other hand, it was shown by Kopecká and Müller in [10] (improving on a similar result of Paszkiewicz [16]) that in general even in the case of 3 orthogonal projections, $\dots P_{\tau(i)} \dots P_{\tau(1)}$ need not converge in the strong operator norm (this result was later strengthened in [11]).

Another of the method approximating the orthogonal projection on $\text{Im}(P_1) \cap \dots \cap \text{Im}(P_n)$ is averaged projections. In this methods, one fixes positive constants $\alpha_k > 0, k = 1, \dots, n$ such that $\sum_k \alpha_k = 1$ and considers the convergence of $(\sum_k \alpha_k P_k)^i$ as i tends to infinity. In [12], Lapidus showed that this method converges to a projection on $\text{Im}(P_1) \cap \dots \cap \text{Im}(P_n)$ in the strong operator topology. This result was extended to various types Banach spaces (such as uniformly convex spaces) in [20] and [3] (under the assumption that all the projections are of norm 1).

In this paper, we will give sufficient criteria for uniform convergence in the case of averaged projections and in the cases of cyclic, quasi-periodic and random products for projections on Banach spaces - see Theorem 2.6, Proposition 3.6, Theorem 3.8 and Theorem 3.9 below. Our criteria uses a new notion of angles between projections that we originally introduced in [14] (and later developed in [13]) in order to study representations of groups on Banach spaces. The basic idea behind all our criteria is that if the angle between any two projections is large enough (with respect to other parameters, such the norms of the projections) then uniform convergence is achieved.

Since various authors used conditions regarding angles in the Hilbert setting to deduce similar uniform convergence results (see for instance [2], [18] and [19]), we found it necessary to point out the new ideas in this paper:

- Unlike previous work, we generalize the notion of angles to a notion of angle between projections in Banach setting. Two things are new here - the notion of angles between projections (as opposed to angle between subspaces) and the generalization of this notion of an angle to Banach spaces. Let us remark that while there are other definitions of angles between subspaces in Banach spaces (see for instance [15][section 3]), as far as we know, these definitions are not applicable to ensure convergence of the alternating projections or the average projections methods.
- Our work does not require the projections to be of norm 1 (although the norm should be sufficiently close to 1).
- Our criteria refer only to angles between couples of projections (i.e., we ask that for each i, j , the angle between P_i and P_j will be large), while previous articles defined angles between the n -tuple of subspaces. This point was important in applying our results in concrete examples. Namely, in [14] and [13] we had concrete examples of projection (related to examples of groups) such that the angles between projection were effectively bounded for several classes of Banach spaces (e.g., in L^p spaces).

Acknowledgements. I would like to thank Piotr Nowak for pointing out a mistake in an early draft of this paper and Simeon Reich for his many useful suggestions.

2. ANGLE BETWEEN PROJECTIONS - DEFINITION AND UNIFORM CONVERGENCE OF AVERAGED PROJECTIONS

Let X be a Banach space. Recall that a projection P is a bounded operator $P \in \mathcal{B}(X)$ such $P^2 = P$. Note that $\|P\| \geq 1$ if $P \neq 0$.

Definition 2.1 (Angle between projections). *Let X be a Banach space and let P_1, P_2 be projections. Assume that there is a projection $P_{1,2}$ on $\text{Im}(P_1) \cap \text{Im}(P_2)$ such that $P_{1,2}P_1 = P_{1,2}$ and $P_{1,2}P_2 = P_{1,2}$ and define*

$$\cos(\angle(P_1, P_2)) = \max \{ \|P_1(P_2 - P_{1,2})\|, \|P_2(P_1 - P_{1,2})\| \}.$$

A few remarks regarding this definition are in order:

Remark 2.2. *In the above definition, we are actually defining the “cosine” of the angle. This is a little misleading, because it could be that in some cases $\cos(\angle(P_1, P_2)) > 1$.*

Remark 2.3. *We note that in the case where X is a Hilbert space and P_1, P_2 are orthogonal projections on V_1, V_2 , the orthogonal projection $P_{1,2}$ on $V_1 \cap V_2$ will always fulfill $P_{1,2}P_1 = P_{1,2}$ and $P_{1,2}P_2 = P_{1,2}$. Also, in this case, $\cos(\angle(P_1, P_2))$ will be equal to the Friedrichs angle between V_1 and V_2 defined by*

$$\cos(\angle(V_1, V_2)) = \sup \{ |\langle u, v \rangle| : \|u\| \leq 1, \|v\| \leq 1, u \in V_1 \cap (V_1 \cap V_2)^\perp, v \in V_2 \}.$$

Remark 2.4. The conditions $P_{1,2}P_1 = P_{1,2}$ and $P_{1,2}P_2 = P_{1,2}$ in the above definition can be understood as follows - in order to define the Friedrichs angle between two subspaces in a Hilbert space, we needed to quotient out their intersection. In Banach spaces, taking a quotient is a non linear operation in the sense that there is no canonical linear projection on the quotient. The condition $P_{1,2}P_1 = P_{1,2}$ replaces the notion of quotient by providing a linear projection $P_1 - P_{1,2}$ on the subspace of vectors fixed by P_1 but not fixed by $P_{1,2}$ (except for the 0-vector which is of course fixed by both). The condition $P_{1,2}P_2 = P_{1,2}$ serves the same purpose making $P_2 - P_{1,2}$ into a similar projection.

Remark 2.5. The above definition of angle might seem hard to compute or even bound. However, the reader can find a multitude of examples in [14] and [13] in which such bounds are computed in a large class of Banach spaces. The main theme of these bounds is that one can use bounds of the Friedrichs angle between two orthogonal projections in a Hilbert space in order to give bounds for the cosine of the angle between “similar” projections in Banach spaces, given that the Banach spaces are “close enough” to some Hilbert space.

Next, we’ll prove an “angle criterion” for uniform convergence of the method of averaged projections. A basic version of this result has already appeared in [14]. Namely, in [14][Theorem 3.12], we gave an “angle criterion” for uniform convergence of $(\frac{P_1 + \dots + P_n}{n})^i$ to a projection on $\text{Im}(P_1) \cap \dots \cap \text{Im}(P_n)$, given that the angle between every couple P_i and P_j is small enough and that the norm of all the P_i is sufficiently close to 1. Below, we’ll deal with the more general case of the convergence of any convex combination of P_1, \dots, P_n . We note that the angle condition we derive below is stronger than [14][Theorem 3.12] even in the simplified case of $(\frac{P_1 + \dots + P_n}{n})^i$, since it allows interplay between different angles, i.e., not all the angles have to be large (a few small angles can be compensated by many large angles).

Theorem 2.6. Let X be a Banach space and let P_1, \dots, P_n be projections in X ($n \geq 2$). Assume that for every $1 \leq j_1 < j_2 \leq n$, there is a projection P_{j_1, j_2} on $\text{Im}(P_{j_1}) \cap \text{Im}(P_{j_2})$, such that $P_{j_1, j_2}P_{j_1} = P_{j_1, j_2}$, $P_{j_1, j_2}P_{j_2} = P_{j_1, j_2}$. Let $\alpha_1, \dots, \alpha_n$ be positive constants such that $\sum_k \alpha_k = 1$. Denote $T = \sum_k \alpha_k P_k$. Assume there is a constant

$$\beta < \min \left\{ \frac{1}{1 - \alpha_1}, \dots, \frac{1}{1 - \alpha_n} \right\},$$

such that

$$\max\{\|P_1\|, \dots, \|P_n\|\} \leq \beta.$$

Assume further that for every $1 \leq j_1, j_2 \leq n$, $\cos(\angle(P_{j_1}, P_{j_2})) < 1$ and that

$$r = \min_{1 \leq j \leq n} \left((1 - \alpha_j)\beta + \sum_{k=1, k \neq j}^n \left(\frac{(\cos(\angle(P_j, P_k)) + \beta + \beta^2) \cos(\angle(P_j, P_k))}{1 - \cos(\angle(P_j, P_k))} 2\alpha_k \right) \right) < 1.$$

Then there is a projection T^∞ on $\bigcap_{j=1}^n \text{Im}(P_j)$, such that for every i ,

$$\|T^i - T^\infty\| \leq Cr^i$$

with $C = \frac{1+\beta}{1-r}$.

Remark 2.7. Note that in the theorem above the first condition

$$\beta < \min \left\{ \frac{1}{1 - \alpha_1}, \dots, \frac{1}{1 - \alpha_n} \right\}$$

is necessary for the fulfilment of the second condition $r < 1$.

In order to prove the above theorem, we will need the following lemma:

Lemma 2.8. *Let P_1, P_2 be projections in a Banach space X , such that there is a projection $P_{1,2}$ on $\text{Im}(P_1) \cap \text{Im}(P_2)$ with $P_{1,2}P_1 = P_{1,2}, P_{1,2}P_2 = P_{1,2}$. Assume that $\cos(\angle(P_1, P_2)) < 1$, and that β is a constant such that $\|P_1\| \leq \beta$ and $\|P_2\| \leq \beta$. Then for any operator $S \in B(X)$, we have that*

$$\|(P_1P_2 - P_2P_1)S\| \leq \frac{(\beta + \beta^2 + \cos(\angle(P_1, P_2))) \cos(\angle(P_1, P_2))}{1 - \cos(\angle(P_1, P_2))} (\|(I - P_1)S\| + \|(I - P_2)S\|)$$

Proof. Let $S \in B(X)$ be some operator. We start by observing that

$$\begin{aligned} \|(P_1P_2 - P_2P_1)S\| &\leq \\ \|(P_1P_2 - P_{1,2})S\| + \|(P_2P_1 - P_{1,2})S\| &= \|P_1(P_2 - P_{1,2})^2S\| + \|P_2(P_1 - P_{1,2})^2S\| \leq \\ \cos(\angle(P_1, P_2)) (\|(P_1 - P_{1,2})S\| + \|(P_2 - P_{1,2})S\|). \end{aligned}$$

In order to complete the proof, we will show that

$$\|(P_1 - P_{1,2})S\| + \|(P_2 - P_{1,2})S\| \leq \frac{\beta + \beta^2 + \cos(\angle(P_1, P_2))}{1 - \cos(\angle(P_1, P_2))} (\|(I - P_1)S\| + \|(I - P_2)S\|).$$

First, note that by the definition of $\cos(\angle(P_1, P_2))$, we have that

$$\|P_{1,2}\| \leq \|P_1P_2\| + \cos(\angle(P_1, P_2)) \leq \beta^2 + \cos(\angle(P_1, P_2)).$$

Observe that

$$\begin{aligned} \|(P_1 - P_{1,2})S\| &\leq \|(P_1 - P_{1,2})P_2S\| + \|(P_1 - P_{1,2})(I - P_2)S\| = \\ \|(P_1P_2 - P_{1,2})S\| + \|(P_1 - P_{1,2})(I - P_2)S\| &= \\ \|P_1(P_2 - P_{1,2})^2S\| + \|(P_1 - P_{1,2})(I - P_2)S\| &\leq \\ \cos(\angle(P_1, P_2)) \|(P_2 - P_{1,2})S\| + (\|P_1\| + \|P_{1,2}\|) \|(I - P_2)S\| &\leq \\ \cos(\angle(P_1, P_2)) \|(P_2 - P_{1,2})S\| + (\beta + \beta^2 + \cos(\angle(P_1, P_2))) \|(I - P_2)S\|, \end{aligned}$$

where the last inequality is due to the bound on $\|P_{1,2}\|$ noted above.

By similar computations

$$\begin{aligned} \|(P_2 - P_{1,2})S\| &\leq \\ \cos(\angle(P_1, P_2)) \|(P_1 - P_{1,2})S\| + (\beta + \beta^2 + \cos(\angle(P_1, P_2))) \|(I - P_1)S\|. \end{aligned}$$

Summing these two inequalities yields

$$\begin{aligned} \|(P_1 - P_{1,2})S\| + \|(P_2 - P_{1,2})S\| &\leq \\ \cos(\angle(P_1, P_2)) (\|(P_1 - P_{1,2})S\| + \|(P_2 - P_{1,2})S\|) + \\ (\beta + \beta^2 + \cos(\angle(P_1, P_2))) (\|(I - P_1)S\| + \|(I - P_2)S\|). \end{aligned}$$

Therefore

$$\|(P_1 - P_{1,2})S\| + \|(P_2 - P_{1,2})S\| \leq \frac{\beta + \beta^2 + \cos(\angle(P_1, P_2))}{1 - \cos(\angle(P_1, P_2))} (\|(I - P_1)S\| + \|(I - P_2)S\|),$$

as needed. \square

Next, we can prove Theorem 2.6:

Proof. We start by defining an energy function $E : B(X) \rightarrow \mathbb{R}_{\geq 0}$ as

$$E(S) = \sum_{j=1}^n \alpha_j \|(I - P_j)S\|.$$

Step 1: We will show that for $T = \sum_k \alpha_k P_k$, we have for every $S \in B(X)$ that $E(TS) \leq rE(S)$, where r is the constant defined as in Theorem 2.6. Fix some $S \in B(X)$, then

$$\begin{aligned}
E(TS) &= \sum_{j=1}^n \alpha_j \left\| (I - P_j) \left(\sum_{k=1}^n \alpha_k P_k \right) S \right\| \\
&= \sum_{j=1}^n \alpha_j \left\| \left(\sum_{k=1, k \neq j}^n \alpha_k (P_k - P_j P_k) \right) S \right\| \\
&= \sum_{j=1}^n \alpha_j \left\| \left(\sum_{k=1, k \neq j}^n \alpha_k (P_k - P_k P_j) + \alpha_k (P_k P_j - P_j P_k) \right) S \right\| \\
&= \sum_{j=1}^n \alpha_j \left\| \left(\sum_{k=1, k \neq j}^n \alpha_k P_k (I - P_j) + \alpha_k (P_k P_j - P_j P_k) \right) S \right\| \\
&\leq \left(\sum_{j=1}^n \alpha_j \sum_{k=1, k \neq j}^n \alpha_k \|P_k\| \|(I - P_j)S\| \right) \\
&\quad + \left(\sum_{k,j=1, k \neq j}^n \alpha_j \alpha_k \|(P_k P_j - P_j P_k)S\| \right).
\end{aligned}$$

We'll deal with each of the summands above separately. First, we notice that

$$\begin{aligned}
(1) \quad & \sum_{j=1}^n \alpha_j \sum_{k=1, k \neq j}^n \alpha_k \|P_k\| \|(I - P_j)S\| = \sum_{j=1}^n \alpha_j \|(I - P_j)S\| \sum_{k=1, k \neq j}^n \alpha_k \beta = \\
& \sum_{j=1}^n (1 - \alpha_j) \beta \alpha_j \|(I - P_j)S\|.
\end{aligned}$$

Second, we apply Lemma 2.8 to the second summand

$$\begin{aligned}
(2) \quad & \sum_{k,j=1, k \neq j}^n \alpha_j \alpha_k \|(P_k P_j - P_j P_k)S\| \leq \\
& \sum_{k,j=1, k \neq j}^n \alpha_j \alpha_k \frac{(\beta + \beta^2 + \cos(\angle(P_j, P_k))) \cos(\angle(P_j, P_k))}{1 - \cos(\angle(P_j, P_k))} (\|(I - P_j)S\| + \|(I - P_k)S\|) = \\
& \sum_{j=1}^n \alpha_j \|(I - P_j)S\| \sum_{k=1, k \neq j}^n \frac{(\beta + \beta^2 + \cos(\angle(P_j, P_k))) \cos(\angle(P_j, P_k))}{1 - \cos(\angle(P_j, P_k))} 2\alpha_k.
\end{aligned}$$

Combining (1) and (2) yields

$$\begin{aligned}
& E(TS) \\
& \leq \\
& \sum_{j=1}^n \alpha_j \|(I - P_j)S\| \left((1 - \alpha_j) \beta + \sum_{k=1, k \neq j}^n \frac{(\beta + \beta^2 + \cos(\angle(P_j, P_k))) \cos(\angle(P_j, P_k))}{1 - \cos(\angle(P_j, P_k))} 2\alpha_k \right) \\
& \leq rE(S).
\end{aligned}$$

Step 2: We'll show that T^i is a Cauchy sequence and therefore converges. Indeed, let $i \geq 1$, then

$$\|T^{i+1} - T^i\| = \left\| \left(\sum_{j=1}^n \alpha_j P_j - I \right) T^i \right\| \leq \sum_{j=1}^n \alpha_j \|(I - P_j)T^i\| = E(T^i) \leq r^i E(I),$$

where the last inequality is due to step 1. Recall that we assumed that $r < 1$ and note that $E(I)$ is a constant and therefore T^i is indeed a Cauchy sequence. Note that

$$E(I) = \sum_{j=1}^n \alpha_j \|I - P_j\| \leq \sum_{j=1}^n \alpha_j (1 + \|P_j\|) \leq 1 + \beta.$$

Denote by T^∞ the limit of T^i , then for every i we have that

$$\|T^\infty - T^i\| \leq \sum_{k=i}^{\infty} E(I) r^k = \frac{E(I)}{1-r} r^i \leq \frac{1+\beta}{1-r} r^i.$$

Step 3: We'll show that T^∞ is a projection on $\text{Im}(P_1) \cap \dots \cap \text{Im}(P_n)$. T^∞ is the limit of T^i and therefore $(T^\infty)^2 = T^\infty$, i.e., T^∞ is a projection. We are left to show that $\text{Im}(P_1) \cap \dots \cap \text{Im}(P_n) = \text{Im}(T^\infty)$. For each $v \in \text{Im}(P_1) \cap \dots \cap \text{Im}(P_n)$, we have that $P_j v = v$ for every $1 \leq j \leq n$ and therefore $Tv = v$ which implies that $T^\infty v = v$, i.e., $\text{Im}(P_1) \cap \dots \cap \text{Im}(P_n) \subseteq \text{Im}(T^\infty)$. On the other hand, $TT^\infty = T^\infty$ and therefore, by step 1, we have that $E(T^\infty) = E(TT^\infty) \leq rE(T^\infty)$, which implies that $E(T^\infty) = 0$. This yields $(I - P_j)T^\infty = 0$ for every j and therefore $\text{Im}(T^\infty) \subseteq \text{Im}(P_1) \cap \dots \cap \text{Im}(P_n)$. \square

As a corollary we can deduce a criterion for uniform convergence for the case $T = \frac{P_1 + \dots + P_n}{n}$ which is similar to the criterion of [14][Theorem 3.12] (the constants differ a little):

Corollary 2.9. *Let X be a Banach space and let P_1, \dots, P_n be projections in X ($n \geq 2$). Assume that for every $1 \leq j_1 < j_2 \leq n$, there is a projection P_{j_1, j_2} on $\text{Im}(P_{j_1}) \cap \text{Im}(P_{j_2})$, such that $P_{j_1, j_2} P_{j_1} = P_{j_1, j_2}$, $P_{j_1, j_2} P_{j_2} = P_{j_1, j_2}$.*

Denote $T = \frac{P_1 + \dots + P_n}{n}$ and assume there is a constant $0 \leq \beta < 1 + \frac{1}{n-1}$ such that $\max\{\|P_1\|, \dots, \|P_n\|\} \leq \beta$ and that $\cos(\angle(P_{j_1}, P_{j_2})) < 1$ for every $1 \leq j_1, j_2 \leq n$. Then there is $\gamma = \gamma(\beta, n) > 0$ such that if

$$\forall 1 \leq j_1, j_2 \leq n, \cos(\angle(P_{j_1}, P_{j_2})) \leq \gamma,$$

then there is a projection T^∞ on $\text{Im}(P_1) \cap \dots \cap \text{Im}(P_n)$ and constants $C = C > 0$, $0 < r < 1$ that depend only on β and γ such that $\|T^\infty - T^i\| \leq Cr^i$.

Proof. Fix $\beta < 1 + \frac{1}{n-1}$ and define a function

$$f(x) = \frac{n-1}{n} \beta + 2 \frac{n-1}{n} \frac{(x + \beta + \beta^2)x}{1-x}.$$

$f(x)$ is clearly continuous and strictly monotone increasing in the interval $[0, 1)$ and $\lim_{x \rightarrow 1} f(x) = \infty$. Also, note that $f(0) < 1$, by the choice of β and therefore there is $\gamma' > 0$ such that $f(\gamma') = 1$. Take γ to be any number such that $0 < \gamma < \gamma'$ and assume that $\max_{j_1, j_2} \cos(\angle(P_{j_1}, P_{j_2})) \leq \gamma$.

By this choice of γ , we have that

$$r = \min_{1 \leq j \leq n} \left(\frac{n-1}{n} \beta + 2 \frac{n-1}{n} \frac{(\cos(\angle(P_j, P_k)) + \beta + \beta^2) \cos(\angle(P_j, P_k))}{1 - \cos(\angle(P_j, P_k))} \right) \leq f(\gamma) < 1.$$

Therefore we can apply Theorem 2.6 with $\alpha_1 = \dots = \alpha_n = \frac{1}{n}$ and deduce that there is $C = \frac{1+\beta}{1-r}$ such that $\|T^\infty - T^i\| \leq Cr^i$, with T^∞ a projection on $\text{Im}(P_1) \cap \dots \cap \text{Im}(P_n)$. \square

Remark 2.10. The constants C, r, γ in the above corollary can be computed explicitly: note that finding γ' such that $f(\gamma') = 1$ boils down to solving a quadratic equation and therefore γ can be taken as $\frac{\gamma'}{2}$ and r, C can be taken as $r = f(\frac{\gamma'}{2}), C = \frac{1+\beta}{1-r}$.

Last, we note that T^i converges to a “canonical” projection with respect to P_1, \dots, P_n if such a projection exists.

Proposition 2.11. Let X be a Banach space and let P_1, \dots, P_n be projections in X ($n \geq 2$). Let $\alpha_1, \dots, \alpha_n$ be positive constants such that $\sum_k \alpha_k = 1$. Denote $T = \sum_k \alpha_k P_k$. Assume that T^i converges in the operator norm to T^∞ which is a projection on $\bigcap_{j=1}^n \text{Im}(P_j)$. If there is a projection $P_{1,2,\dots,n}$ on $\bigcap_{j=1}^n \text{Im}(P_j)$ such that for every j , $P_{1,2,\dots,n} P_j = P_{1,2,\dots,n}$, then $T^\infty = P_{1,2,\dots,n}$.

Proof. Note that for every i , we have that $P_{1,\dots,n} T^i = P_{1,\dots,n}$ and therefore $T^\infty = P_{1,\dots,n} T^\infty = P_{1,\dots,n}$. \square

3. CONVERGENCE OF RANDOM PRODUCTS OF PROJECTIONS

Below, we'll give “angle criteria” for the convergence of cyclic, quasi-periodic and random products for projections. Due to the messy nature of the computations, we keep these criteria non-effective, i.e., we do not specify the exact bounds on the angle to ensure uniform convergence, but only show these bounds exist. The interested reader can derive effective versions of all our criteria below by substituting our use of γ, r and C below by explicit constants such as the ones given in Remark 2.10 and then working out all the computations explicitly with these constants.

Throughout, let P_1, \dots, P_n be projections in a Banach space X . Below we will always assume that for every $1 \leq j_1 < j_2 \leq n$, there is a projection P_{j_1, j_2} on $\text{Im}(P_{j_1}) \cap \text{Im}(P_{j_2})$, such that $P_{j_1, j_2} P_{j_1} = P_{j_1, j_2}, P_{j_1, j_2} P_{j_2} = P_{j_1, j_2}$. and therefore $\cos(\angle(P_i, P_j))$ is defined.

Theorem 3.1. Let P_1, \dots, P_n be as above. Assume that there is a constant $\beta < 1 + \frac{1}{n-1}$ such that

$$\max\{\|P_1\|, \dots, \|P_n\|\} \leq \beta.$$

There is a constant $\gamma = \gamma(\beta, n)$ such that if

$$\max\{\cos(\angle(P_{j_1}, P_{j_2})) : 1 \leq j_1 < j_2 \leq n\} \leq \gamma,$$

then there is a projection T^∞ on $\bigcap_{j=1}^n \text{Im}(P_j)$ and constants $C > 0, 0 < r < 1$ that depend only on γ, β such that for every integer $m \geq n$ and every surjective map $\sigma : \{1, \dots, m\} \rightarrow \{1, \dots, n\}$ the following holds for every i :

$$\|P_{\sigma(m)} \dots P_{\sigma(1)} - T^\infty\| \leq \beta^m C r^i + 2\gamma \beta^{m-2} ((2 + \beta)^i - 1).$$

Proof. Fix m, σ as above. Denote $T = \frac{P_1 + \dots + P_n}{n}$ and let $\gamma = \gamma(\beta, n)$ be as in Corollary 2.9, then by this corollary there are $C > 0, 0 < r < 1$ such that for every i ,

$$\|T^\infty - T^i\| \leq C r^i,$$

where T^∞ is a projection on $\bigcap_{j=1}^n \text{Im}(P_j)$. Therefore for every i the following holds:

$$\begin{aligned} \|P_{\sigma(m)} \dots P_{\sigma(1)} - T^\infty\| &= \|P_{\sigma(m)} \dots P_{\sigma(1)} (I - T^\infty)\| \leq \\ &\|P_{\sigma(m)} \dots P_{\sigma(1)} (I - T^i)\| + \|P_{\sigma(m)} \dots P_{\sigma(1)} (T^i - T^\infty)\| \leq \\ &\|P_{\sigma(m)} \dots P_{\sigma(1)} (I - T^i)\| + \|P_{\sigma(m)} \dots P_{\sigma(1)}\| \|T^i - T^\infty\| \leq \\ &\|P_{\sigma(m)} \dots P_{\sigma(1)} (I - T^i)\| + \beta^m C r^i. \end{aligned}$$

By the above inequality, it is left to prove that

$$\|P_{\sigma(m)} \dots P_{\sigma(1)} (I - T^i)\| \leq 2\gamma \beta^{m-1} (2 + \beta)^i.$$

Observe that

$$I - T^i = I - (I - (I - T))^i = (I - T) \sum_{k=1}^i \binom{i}{k} (-1)^k (I - T)^{k-1}.$$

Therefore

$$\begin{aligned} \|P_{\sigma(m)} \dots P_{\sigma(1)}(I - T^i)\| &\leq \|P_{\sigma(m)} \dots P_{\sigma(1)}(I - T)\| \left\| \sum_{k=1}^i \binom{i}{k} (-1)^k (I - T)^{k-1} \right\| \leq \\ \|P_{\sigma(m)} \dots P_{\sigma(1)}(I - T)\| &\left(\sum_{k=1}^i \binom{i}{k} (1 + \beta)^{k-1} \right) = \frac{(2 + \beta)^i - 1}{1 + \beta} \|P_{\sigma(m)} \dots P_{\sigma(1)}(I - T)\|. \end{aligned}$$

In order to finish the proof, we must show that

$$\|P_{\sigma(m)} \dots P_{\sigma(1)}(I - T)\| \leq 2\gamma\beta^{m-2}(m-1)(1 + \beta).$$

Note that

$$\|P_{\sigma(m)} \dots P_{\sigma(1)}(I - T)\| \leq \frac{\|P_{\sigma(m)} \dots P_{\sigma(1)}(I - P_1)\| + \dots + \|P_{\sigma(m)} \dots P_{\sigma(1)}(I - P_n)\|}{n}$$

and therefore it is enough to show that for every j ,

$$\|P_{\sigma(m)} \dots P_{\sigma(1)}(I - P_j)\| \leq 2\gamma\beta^{m-2}(m-1)(1 + \beta).$$

Let $1 \leq l \leq m$ be the smallest integer such that $\sigma(l) = j$. We will finish the proof by showing by induction that

$$\|P_{\sigma(m)} \dots P_{\sigma(1)}(I - P_j)\| \leq 2\gamma\beta^{m-2}(l-1)(1 + \beta).$$

For $l = 1$, we have that $\sigma(1) = j$ and therefore

$$P_{\sigma(m)} \dots P_{\sigma(1)}(I - P_j) = P_{\sigma(m)} \dots P_{\sigma(2)} P_j (I - P_j) = 0$$

and we are done. Next, let $l > 1$ and assume that the above inequality holds for $l - 1$. Define $\sigma' : \{1, \dots, m\} \rightarrow \{1, \dots, n\}$ as

$$\sigma'(k) = \begin{cases} \sigma(k) & k > l \text{ or } k < l - 1 \\ \sigma(l) & k = l - 1 \\ \sigma(l - 1) & k = l \end{cases}.$$

Note that

$$\begin{aligned} &\|P_{\sigma(m)} \dots P_{\sigma(1)}(I - P_j) - P_{\sigma'(m)} \dots P_{\sigma'(1)}(I - P_j)\| = \\ &\|P_{\sigma(m)} \dots P_{\sigma(l+1)}(P_{\sigma(l)} P_{\sigma(l-1)} - P_{\sigma(l-1)} P_{\sigma(l)}) P_{\sigma(l-2)} \dots P_{\sigma(1)}(I - P_j)\| \leq \\ &\|P_{\sigma(m)} \dots P_{\sigma(l+1)}\| \|P_{\sigma(l)} P_{\sigma(l-1)} - P_{\sigma(l-1)} P_{\sigma(l)}\| \|P_{\sigma(l-2)} \dots P_{\sigma(1)}\| \|I - P_j\| \leq \\ &\beta^{n-2}(1 + \beta) \|P_{\sigma(l)} P_{\sigma(l-1)} - P_{\sigma(l-1)} P_{\sigma(l)}\| \leq \\ &\beta^{n-2}(1 + \beta) \|P_{\sigma(l)} P_{\sigma(l-1)} - P_{\sigma(l-1), \sigma(l)}\| + \beta^{n-2}(1 + \beta) \|P_{\sigma(l-1)} P_{\sigma(l)} - P_{\sigma(l-1), \sigma(l)}\| \leq \\ &2\gamma\beta^{m-2}(1 + \beta), \end{aligned}$$

where the last inequality is due to the assumption that $\cos(\angle(P_{\sigma(l-1)}, P_{\sigma(l)})) \leq \gamma$.

By definition $\sigma'(l - 1) = \sigma(l) = j$ and therefore by the induction assumption, we have that

$$\|P_{\sigma'(m)} \dots P_{\sigma'(1)}(I - P_j)\| \leq 2\gamma\beta^{n-2}(l-2)(1 + \beta).$$

Combining this inequality with the previous inequality, we get that

$$\begin{aligned} &\|P_{\sigma(m)} \dots P_{\sigma(1)}(I - P_j)\| \leq \\ &\|P_{\sigma(m)} \dots P_{\sigma(1)}(I - P_j) - P_{\sigma'(m)} \dots P_{\sigma'(1)}(I - P_j)\| + \|P_{\sigma'(m)} \dots P_{\sigma'(1)}(I - P_j)\| \leq \\ &2\gamma\beta^{m-2}(1 + \beta) + 2\gamma\beta^{m-2}(l-2)(1 + \beta) = 2\gamma\beta^{m-2}(l-1)(1 + \beta), \end{aligned}$$

as needed. \square

Corollary 3.2. *Let P_1, \dots, P_n be as above. For every integer $m \geq n$ and constants $0 \leq q < 1$, $0 \leq \beta < 1 + \frac{1}{n-1}$, there is a constant $\gamma = \gamma(\beta, n, m, q) > 0$ such that if*

$$\max\{\|P_1\|, \dots, \|P_n\|\} \leq \beta \text{ and } \max\{\cos(\angle(P_{j_1}, P_{j_2})) : 1 \leq j_1 < j_2 \leq n\} \leq \gamma,$$

then for every surjective map $\sigma : \{1, \dots, m\} \rightarrow \{1, \dots, n\}$, we have that

$$\|P_{\sigma(m)} \dots P_{\sigma(1)} - T^\infty\| \leq q,$$

where T^∞ is the projection on $\bigcap_{j=1}^n \text{Im}(P_j)$ given in the above theorem.

Proof. Fix m, q, β as above and fix a surjective map $\sigma : \{1, \dots, m\} \rightarrow \{1, \dots, n\}$. Let γ_1 be the bound on the cosine of the angles as in the above theorem and let r, C be the constants corresponding to γ_1 .

Therefore, if

$$\max\{\|P_1\|, \dots, \|P_n\|\} \leq \beta \text{ and } \max\{\cos(\angle(P_{j_1}, P_{j_2})) : 1 \leq j_1 < j_2 \leq n\} \leq \gamma_1,$$

then by the above theorem we have for every i that

$$\|P_{\sigma(m)} \dots P_{\sigma(1)} - T^\infty\| \leq \beta^m C r^{i-1} + 2\gamma_1 \beta^{m-2} ((2 + \beta)^i - 1).$$

Fix i_0 large enough such that $\beta^m C r^{i_0-1} \leq \frac{q}{2}$. Denote

$$\gamma_2 = \frac{1}{2\beta^{m-2}((2 + \beta)^{i_0} - 1)} \frac{q}{2}.$$

Then for $\gamma = \min\{\gamma_1, \gamma_2\}$, we have that

$$\|P_{\sigma(m)} \dots P_{\sigma(1)} - T^\infty\| \leq \beta^m C r^{i_0-1} + 2\gamma_2 \beta^{m-2} ((2 + \beta)^{i_0} - 1) \leq \frac{q}{2} + \frac{q}{2} = q.$$

□

The above corollary allows us to deduce uniform convergence criteria in the cases of quasi-periodic products (see definition below) and random products under an additional assumption which we call weak consistency:

Definition 3.3 (Consistency). *We shall say that the projections P_1, \dots, P_n are consistent if for every $\{i_1, \dots, i_k\} \subseteq \{1, \dots, n\}$, there is a projection P_{i_1, \dots, i_k} on $\text{Im}(P_{i_1}) \cap \text{Im}(P_{i_2}) \cap \dots \cap \text{Im}(P_{i_k})$, such that*

$$\forall 1 \leq j \leq k, P_{i_1, \dots, i_k} P_{i_j} = P_{i_1, \dots, i_k}.$$

We shall say that the projections P_1, \dots, P_n are weakly consistent if there is a projection $P_{1, \dots, n}$ on $\bigcap_j \text{Im}(P_j)$ such that

$$\forall 1 \leq j \leq n, P_{1, \dots, n} P_j = P_{1, \dots, n}.$$

Example 3.4. *In the case where $X = H$ is a Hilbert space and P_1, \dots, P_n are orthogonal projections on V_1, \dots, V_n correspondingly, we have that P_1, \dots, P_n are consistent by taking P_{i_1, \dots, i_k} to be the orthogonal projection on $V_{i_1} \cap \dots \cap V_{i_k}$. Other examples of consistent projections are given in [13].*

Remark 3.5. *We note that by Proposition 2.11, if P_1, \dots, P_n are weakly consistent and $T^i = (\frac{P_1 + \dots + P_n}{n})^i$ converges in the operator norm to a projection T^∞ on $\bigcap_j \text{Im}(P_j)$, then $T^\infty = P_{1, \dots, n}$, i.e.,*

$$\forall 1 \leq j \leq n, T^\infty P_j = T^\infty.$$

Next, we'll use Corollary 3.2 together with the assumption of weak consistency to deduce criteria for uniform convergence. We'll start with proving such a criterion for the convergence of $(P_n \dots P_1)^i$. We remark that although this case is well studied and various angle considerations were applied to it (see for instance [2], [17], [18], [19]), this result is new even in the setting of orthogonal projections in Hilbert spaces.

Proposition 3.6. *Let P_1, \dots, P_n be as above and let $0 \leq \beta < 1 + \frac{1}{n-1}$, $0 < q < 1$ be constants. Then there is $\gamma = \gamma(\beta, q)$ such that if*

- (1) $\max\{\|P_1\|, \dots, \|P_n\|\} \leq \beta$ and $\max\{\cos(\angle(P_{j_1}, P_{j_2})) : 1 \leq j_1 < j_2 \leq n\} \leq \gamma$.
- (2) P_1, \dots, P_n are weakly consistent.

Then $\|(P_n \dots P_1)^i - P_{1, \dots, n}\| \leq q^i$.

Proof. Fix β, q as above. Take $\gamma = \gamma(\beta, n, m, q) > 0$ given by Corollary 3.2. Note that by weak consistency, we have that

$$P_n \dots P_1 - P_{1, \dots, n} = P_n \dots P_1 (I - P_{1, \dots, n}) = (I - P_{1, \dots, n}) P_n \dots P_1 (I - P_{1, \dots, n}).$$

Therefore

$$\|(P_n \dots P_1)^i - P_{1, \dots, n}\| = \|((P_n \dots P_1)(I - P_{1, \dots, n}))^i\| \leq q^i,$$

where the last inequality is due to Corollary 3.2. \square

Next, we'll analyse the case of quasi-periodic products. Recall the following definition:

Definition 3.7. *A map $\tau \in \{1, \dots, n\}^{\mathbb{N}}$ will be called quasi-periodic if there is some fixed integer $m \geq n$ (which will be called the quasi-period of τ) such that for every $i \in \mathbb{N}$,*

$$\{\tau(i), \tau(i+1), \dots, \tau(i+m-1)\} = \{1, \dots, n\}.$$

For τ quasi-periodic, the infinite product $\dots P_{\tau(i)} \dots P_{\tau(1)}$ will be called a quasi-periodic product.

Theorem 3.8. *Let P_1, \dots, P_n be as above and let $0 \leq \beta < 1 + \frac{1}{n-1}$. Then for every $m \geq n$, there is $\gamma = \gamma(\beta, n, m)$ such that if*

- (1) $\max\{\|P_1\|, \dots, \|P_n\|\} \leq \beta$ and $\max\{\cos(\angle(P_{j_1}, P_{j_2})) : 1 \leq j_1 < j_2 \leq n\} \leq \gamma$.
- (2) P_1, \dots, P_n are weakly consistent.

Then every quasi-periodic product of P_1, \dots, P_n with a quasi-period m converges uniformly to $P_{1, \dots, n}$ as the rate of convergence is exponential.

Proof. Fix β and m and let $\tau \in \{1, \dots, n\}^{\mathbb{N}}$ be some quasi-periodic map with a quasi-period m . Fix some $q < 1$ and let $\gamma = \gamma(\beta, n, m, q)$ to be the constant given by Corollary 3.2. Then by Corollary 3.2 and by then quasi-periodicity of τ , if

$$\max\{\|P_1\|, \dots, \|P_n\|\} \leq \beta \text{ and } \max\{\cos(\angle(P_{j_1}, P_{j_2})) : 1 \leq j_1 < j_2 \leq n\} \leq \gamma,$$

then for every i , $\|P_{\tau(i+m-1)} \dots P_{\tau(i)} - P_{1, \dots, n}\| \leq q$.

From the assumption of weak consistency, we have that for every $i_1 < i_2$ the following holds

$$P_{\tau(i_2)} \dots P_{\tau(i_1)} (I - P_{1, \dots, n}) = (I - P_{1, \dots, n}) P_{\tau(i_2)} \dots P_{\tau(i_1)} (I - P_{1, \dots, n}).$$

Therefore, for every i we have that

$$\begin{aligned} \|P_{\tau(i)} \dots P_{\tau(1)} - P_{1, \dots, n}\| &= \\ \|P_{\tau(i)} \dots P_{\tau(\lfloor \frac{i}{m} \rfloor m + 1)} \prod_{l=1}^{\lfloor \frac{i}{m} \rfloor} (P_{\tau(lm)} \dots P_{\tau((l-1)m+1)} - P_{1, \dots, n})\| &\leq \\ \|P_{\tau(i)} \dots P_{\tau(\lfloor \frac{i}{m} \rfloor m + 1)}\| \prod_{l=1}^{\lfloor \frac{i}{m} \rfloor} \|P_{\tau(lm)} \dots P_{\tau((l-1)m+1)} - P_{1, \dots, n}\| &\leq \beta^{m-1} q^{\lfloor \frac{i}{m} \rfloor}. \end{aligned}$$

Therefore $\|P_{\tau(i)} \dots P_{\tau(1)} - P_{1, \dots, n}\|$ tends to 0 exponentially fast as i tends to infinity. \square

The proof of the analogous theorem when $\tau \in \{1, \dots, n\}^{\mathbb{N}}$ is chosen according to some product probability is similar, but requires a little more work:

Theorem 3.9. *Let P_1, \dots, P_n be as above and let $0 \leq \beta < 1 + \frac{1}{n-1}$. Also, let μ be a probability measure on $\{1, \dots, n\}$ such that for each j , $\mu(j) > 0$. Then there is $\gamma = \gamma(\beta, n, \mu)$ such that if the following holds:*

- (1) $\max\{\|P_1\|, \dots, \|P_n\|\} \leq \beta$ and $\max\{\cos(\angle(P_{j_1}, P_{j_2})) : 1 \leq j_1 < j_2 \leq n\} \leq \gamma$.
- (2) P_1, \dots, P_n are weakly consistent.

Then for the space $\{1, \dots, n\}^{\mathbb{N}}$ with the infinite product measure $\bar{\mu} = \prod_{i=1}^{\infty} \mu$, we have that for $\bar{\mu}$ -almost every $\tau \in \{1, \dots, n\}^{\mathbb{N}}$, the infinite product $\dots P_{\tau(i)} \dots P_{\tau(1)}$ converges uniformly to $P_{1, \dots, n}$ and the rate of convergence is exponential.

Proof. Fix β and μ as above. For $k \in \mathbb{N}$, $\tau \in \{1, \dots, n\}^{\mathbb{N}}$, define the sets

$$A(\tau, k) = \{1 \leq l \leq k : \{\tau(n(l-1)+1), \tau(n(l-1)+2), \dots, \tau(ln)\} = \{1, \dots, n\}\},$$

$$B(\tau, k) = \{1, \dots, k\} \setminus A(\tau, k),$$

$$A(\tau) = \bigcup_k A(\tau, k).$$

In a random n -tuple of numbers chosen according to the probability $\mu \times \mu \times \dots \times \mu$, the probability that each number $1, \dots, n$ will appear in the n -tuple exactly once is $\frac{1}{n! \mu(1) \mu(2) \dots \mu(n)}$. Therefore, by the law of large numbers, for $\bar{\mu}$ -almost every $\tau \in \{1, \dots, n\}^{\mathbb{N}}$ we have that

$$\lim_{k \rightarrow \infty} \frac{|A(\tau, k)|}{k} = \frac{1}{n! \mu(1) \mu(2) \dots \mu(n)}.$$

Fix $0 < \lambda < \frac{1}{n! \mu(1) \mu(2) \dots \mu(n)}$. Then, as a result of the law of large numbers mentioned above, for $\bar{\mu}$ -almost every $\tau \in \{1, \dots, n\}^{\mathbb{N}}$ there is $k_{\tau} \in \mathbb{N}$ such that for every $k \geq k_{\tau}$ we have that $\frac{|A(\tau, k)|}{k} \geq \lambda$, or equivalently, for every $k \geq k_{\tau}$ we have that $\frac{|B(\tau, k)|}{k} \leq 1 - \lambda$.

Fix $q > 0$ such that $\beta^{n(1-\lambda)} q^{\lambda} < 1$ and let $\gamma = \gamma(\beta, n, q)$ given by Corollary 3.2. Then by Corollary 3.2 above, if

$$\max\{\cos(\angle(P_{j_1}, P_{j_2})) : 1 \leq j_1 < j_2 \leq n\} \leq \gamma,$$

then for every $l \in A(\tau, k)$ we have that

$$\|P_{\tau(ln)} \dots P_{\tau((l-1)n+1)}(I - P_{1, \dots, n})\| = \|P_{\tau(ln)} \dots P_{\tau((l-1)n+1)} - P_{1, \dots, n}\| \leq q.$$

As in the previous proof, from the assumption of weak consistency, we have that for every $i_1 < i_2$ and every τ the following holds

$$P_{\tau(i_2)} \dots P_{\tau(i_1)}(I - P_{1, \dots, n}) = (I - P_{1, \dots, n}) P_{\tau(i_2)} \dots P_{\tau(i_1)}(I - P_{1, \dots, n}).$$

For $l \in \mathbb{N}$, define operators $L_l = L_l(\tau)$ as follows

$$L_l = \begin{cases} P_{\tau(ln)} \dots P_{\tau((l-1)n+1)}(I - P_{1, \dots, n}) & l \in A(\tau) \\ P_{\tau(ln)} \dots P_{\tau((l-1)n+1)} & l \notin A(\tau) \end{cases}.$$

Then for every i , we have (as a result of weak consistency) that

$$P_{\tau(i)} \dots P_{\tau(1)}(I - P_{1, \dots, n}) = P_{\tau(i)} \dots P_{\tau(\lfloor \frac{i}{n} \rfloor n + 1)} \left(\prod_{l=1}^{\lfloor \frac{i}{n} \rfloor} L_l \right) (I - P_{1, \dots, n}).$$

Note that

$$\begin{aligned} \forall l \in A(\tau), \|L_l\| &\leq q, \\ \forall l \notin A(\tau), \|L_l\| &\leq \beta^n. \end{aligned}$$

Therefore for every i , we have that

$$\begin{aligned} & \|P_{\tau(i)} \dots P_{\tau(1)}(I - P_{1, \dots, n})\| \leq \\ & \|P_{\tau(i)} \dots P_{\tau(\lfloor \frac{i}{n} \rfloor n + 1)}\| \left(\prod_{l=1}^{\lfloor \frac{i}{n} \rfloor} \|L_l\| \right) \|I - P_{1, \dots, n}\| \leq \\ & \beta^{n-1} (\beta^n)^{|B(\tau, \lfloor \frac{i}{n} \rfloor)|} q^{|A(\tau, \lfloor \frac{i}{n} \rfloor)|} \|I - P_{1, \dots, n}\|. \end{aligned}$$

As a result, by the definition of k_τ , for every $i \geq nk_\tau$ the following holds

$$\|P_{\tau(i)} \dots P_{\tau(1)}(I - P_{1, \dots, n})\| \leq \|I - P_{1, \dots, n}\| \beta^{n-1} (\beta^{n(1-\lambda)} q^\lambda)^{\lfloor \frac{i}{n} \rfloor}.$$

Recall that q was chosen to ensure that $\beta^{n(1-\lambda)} q^\lambda < 1$ and therefore the right-hand side of the above inequality tends to 0 (exponentially fast) as i tends to infinity. \square

REFERENCES

- [1] I. Amemiya and T. Andô. Convergence of random products of contractions in Hilbert space. *Acta Sci. Math. (Szeged)*, 26:239–244, 1965.
- [2] C. Badea, S. Grivaux, and V. Müller. The rate of convergence in the method of alternating projections. *Algebra i Analiz*, 23(3):1–30, 2011.
- [3] Catalin Badea and Yuri I. Lyubich. Geometric, spectral and asymptotic properties of averaged products of projections in Banach spaces. *Studia Math.*, 201(1):21–35, 2010.
- [4] J. Dye, M. A. Khamsi, and S. Reich. Random products of contractions in Banach spaces. *Trans. Amer. Math. Soc.*, 325(1):87–99, 1991.
- [5] John Dye. Convergence of random products of compact contractions in Hilbert space. *Integral Equations Operator Theory*, 12(1):12–22, 1989.
- [6] John M. Dye, Tadeusz Kuczumow, Pei-Kee Lin, and Simeon Reich. Convergence of unrestricted products of nonexpansive mappings in spaces with the Opial property. *Nonlinear Anal.*, 26(4):767–773, 1996.
- [7] John M. Dye and Simeon Reich. On the unrestricted iteration of projections in Hilbert space. *J. Math. Anal. Appl.*, 156(1):101–119, 1991.
- [8] John M. Dye and Simeon Reich. Unrestricted iterations of nonexpansive mappings in Banach spaces. *Nonlinear Anal.*, 19(10):983–992, 1992.
- [9] Israel Halperin. The product of projection operators. *Acta Sci. Math. (Szeged)*, 23:96–99, 1962.
- [10] Eva Kopecká and Vladimír Müller. A product of three projections. *Studia Math.*, 223(2):175–186, 2014.
- [11] Eva Kopecká and Adam Paszkiewicz. Strange product of projections. <http://arxiv.org/pdf/1508.05029v1.pdf>, 2015.
- [12] Michel L. Lapidus. Generalization of the Trotter-Lie formula. *Integral Equations Operator Theory*, 4(3):366–415, 1981.
- [13] Izhar Oppenheim. Vanishing of cohomology with coefficients in representations on Banach spaces of groups acting on buildings. <http://arxiv.org/abs/1512.08188>, 2015.
- [14] Izhar Oppenheim. Averaged projections, angles between groups and strengthening of property (T). *Mathematische Annalen*, To appear., 2016.
- [15] M. I. Ostrovskii. Topologies on the set of all subspaces of a Banach space and related questions of Banach space geometry. *Quaestiones Math.*, 17(3):259–319, 1994.
- [16] Adam Paszkiewicz. The amemiya-ando conjecture falls. <http://arxiv.org/pdf/1203.3354v1.pdf>, 2012.
- [17] Evgeniy Pustynnik, Simeon Reich, and Alexander J. Zaslavski. Convergence of non-cyclic infinite products of operators. *J. Math. Anal. Appl.*, 380(2):759–767, 2011.
- [18] Evgeniy Pustynnik, Simeon Reich, and Alexander J. Zaslavski. Convergence of non-periodic infinite products of orthogonal projections and nonexpansive operators in Hilbert space. *J. Approx. Theory*, 164(5):611–624, 2012.
- [19] Evgeniy Pustynnik, Simeon Reich, and Alexander J. Zaslavski. Inner inclination of subspaces and infinite products of orthogonal projections. *J. Nonlinear Convex Anal.*, 14(3):423–436, 2013.
- [20] Simeon Reich. A limit theorem for projections. *Linear and Multilinear Algebra*, 13(3):281–290, 1983.
- [21] Makoto Sakai. Strong convergence of infinite products of orthogonal projections in Hilbert space. *Appl. Anal.*, 59(1-4):109–120, 1995.

- [22] John von Neumann. On rings of operators. Reduction theory. *Ann. of Math. (2)*, 50:401–485, 1949.

DEPARTMENT OF MATHEMATICS, BEN-GURION UNIVERSITY OF THE NEGEV, BE'ER SHEVA 84105,
ISRAEL

E-mail address: `izharo@bgu.ac.il`