Excitation Entanglement Entropy in 2d Conformal Field Theories

M.M. Sheikh-Jabbari[‡], and H. Yavartanoo[†]

[‡] School of Physics, Institute for Research in Fundamental Sciences (IPM), P.O.Box 19395-5531, Tehran, Iran

[†] State Key Laboratory of Theoretical Physics, Institute of Theoretical Physics,

Chinese Academy of Sciences, Beijing 100190, China

(Ω Dated: March 7, 2022)

We analyze how excitations affect the entangl in a 2d conformal field theory (CFT) using the well as direct CFT computations. We introdu-difference between the entanglement entropy ge denoted by h. The excitation entanglement ent-quantity (independent of the cutoff), and hence excitation entanglement entropy is governed by a one point function of the energy momentum ten-analyze low and high temperature behavior of th $\Delta_h S$ grows as function of temperature. We p-entanglement entropy, that although $\Delta_h S$ can b positive. We also discuss the mutual and multip. Entanglement is one of the specific characteristics of quantum systems. From a theoretical perspective, given a system described by a normalized density matrix ρ (Tr $\rho = 1$), entanglement can be quantified in various dif-ferent measures built from ρ. Entanglement entropy (EE) is the simplest of such measures and is defined as the von Neumann entropy of the system $S_{EE} = -\text{Tr}\rho \ln \rho. \qquad (1)$ Entanglement entropy has emerged as a valuable tool for probing quantum systems in diverse situations, from con-densed matter physics [1–3] to quantum gravity [4–6]. Nonetheless, calculation of EE for a general quantum me-chanical system remains a daunting task. In (local) quantum field theory settings there is a one-to-one map between the Hilbert space of the theory and field configurations on a constant time slice. One may ask for entanglement between any two regions A and B on a We analyze how excitations affect the entanglement entropy for an arbitrary entangling interval in a 2d conformal field theory (CFT) using the holographic entanglement entropy techniques as well as direct CFT computations. We introduce the excitation entanglement entropy $\Delta_h S$, the difference between the entanglement entropy generic excitations and their arbitrary descendants denoted by h. The excitation entanglement entropy, unlike the entanglement entropy, is a finite quantity (independent of the cutoff), and hence a good physical observable. We show that the excitation entanglement entropy is governed by a second order differential equation sourced by the one point function of the energy momentum tensor computed in the excited background state. We analyze low and high temperature behavior of the excitation entanglement entropy and show that $\Delta_h S$ grows as function of temperature. We prove an "integrated positivity" for the excitation entanglement entropy, that although $\Delta_h S$ can be positive or negative, its average value is always positive. We also discuss the mutual and multipartite information with generic excitations.

$$S_{EE} = -\mathrm{Tr}\rho \ln\rho. \tag{1}$$

for entanglement between any two regions A and B on a constant time slice separated by a region ∂A . The entanglement would then depend on the shape of the regions A and B, as well as the field profile in these regions. The latter may be viewed as excitations on the vacuum field configuration. In principle one may try to compute (1)using standard path integral techniques, perturbatively in the coupling [2, 3]. These, except for free field theories with simple shapes for ∂A , are formidable computations.

However, in two dimensional conformal field theories (2d CFT's), where the entangling region A is a one dimensional interval, invoking the infinite dimensional conformal symmetry simplifies the calculations and EE can be computed analytically $S = \frac{c}{3} \ln \frac{L}{\epsilon}$, where L is the length of entangling region, ϵ is the cutoff regulating the ultraviolet divergences and c is the central charge [7, 8]. This is conveniently computed by employing the replica trick [2, 7], using

$$S_A = \lim_{n \to 1} \frac{1}{1-n} \ln \operatorname{Tr} \rho_A^n.$$
(2)

Computation of the entanglement entropy in 2d CFT's for a given entangling interval may be extended to cases where we have excitations in the interval. These excitations may be caused by presence of a primary operator or descendants of primaries and the field theory could be at zero or non-zero temperature or angular (chemical) potential [9–14]. Cases where we have time dependent excitations (e.g. quenches) has also been studied and time dependence and dynamics of EE has been analyzed [15– 19]. In this Letter we analyze further the computation of EE for generic excited states in 2d CFT's.

The AdS/CFT duality [20] (or more practical versions of it, the gauge/gravity correspondence) may be used as a tool to compute EE, especially for strongly coupled field theories. In general there is the proposal by Ryu-Takayanagi (RT) for computing the EE holographically using the AdS/CFT techniques [21, 22]. According to the RT proposal computation of EE is reduced to calculating the area of a surface which has the minimum area among surfaces attached to the boundary of the asymptotically AdS bulk at the boundary of entangling region ∂A . The (holographic) entanglement entropy is given in terms of the area of this minimal surface $m(\partial A)$:

$$S_{HEE} = \frac{Area(m(\partial A))}{4G_N},\tag{3}$$

where G_N is the Newton constant. The RT proposal which was initially made for static gravity backgrounds have been extended to stationary cases [23].

The RT proposal for computing HEE becomes particularly simple for AdS_3 geometry [21, 22]: the minimal surface, which is a one-dimensional curve in this case, becomes the geodesic connecting the two spacelike separated points at the AdS_3 boundary. The two points at the boundary are the endpoints of the entangling interval in the dual 2d

CFT. This geodesic length is logarithmically divergent, due to the near boundary behavior of the metric. One may then regulate this length by a cutoff scale ϵ . If the entangling interval at the boundary has length L, then

$$S_{HEE} = \frac{\text{Length}_{geodesic}}{4G_3} = \frac{\ell}{2G_3} \ln \frac{L}{\epsilon}, \qquad (4)$$

where ℓ is the AdS₃ radius and G_3 is the 3d Newton constant. If the presumed dual 2d CFT is at the Brown-Henneaux central charge c [24]

$$c = \frac{3\ell}{2G_3},\tag{5}$$

the above exactly matches the 2d CFT direct computations. The above analysis may be repeated for other known locally AdS_3 geometries [23], like BTZ black holes [25], successfully reproducing the entanglement entropy of a 2d CFT at non-zero temperature [7].

Here we analyze holographic computation of the EE for general asymptotic AdS_3 geometry, the Bañados geometries [26], associated with generic excited states in the dual 2d CFT. The pure AdS_3 Einstein gravity is described by

$$\mathcal{S} = \frac{1}{16\pi G_3} \int d^3x \sqrt{-g} (R + \frac{2}{\ell^2}) \implies R_{\mu\nu} + \frac{2}{\ell^2} g_{\mu\nu} = 0.$$
(6)

All solutions to (6) are locally AdS_3 geometries and are fully specified by the falloff (near boundary) behavior of the solutions and possibly other topological data [27]. The most general solution obeying Brown-Henneaux boundary conditions [24], are the Bañados geometries [26]

$$ds^{2} = \ell^{2} \frac{dz^{2}}{z^{2}} - \left(\frac{\ell^{2}}{z} dx^{+} - zL_{-} dx^{-}\right) \left(\frac{\ell^{2}}{z} dx^{-} - zL_{+} dx^{+}\right), \quad (7)$$

where $x^{\pm} \in [0, 2\pi]$ and $L_{\pm} = L_{\pm}(x^{\pm})$ are two smooth and periodic, but otherwise arbitrary, functions. In the coordinate system adopted here the causal boundary of these geometries is located at z = 0 and is (a part of) a cylinder $\mathbb{R} \times S^1$ spanned by x^{\pm} , where \mathbb{R} is the time like direction which is along $x^+ + x^-$ and S^1 is parametrized by $x^+ - x^-$. The above family contain the well-known BTZ black hole solutions [25] as a special class associated with constant and positive $L_{\pm} \equiv \mathbb{T}^2_{\pm}$. The inner and outer horizons radii ρ_{\pm} of the BTZ black hole are given by

$$\rho_{\pm} = \ell(\mathfrak{T}_{+} \pm \mathfrak{T}_{-}). \tag{8}$$

Bañados geometries and their geometric properties have been studied in some detail [28, 29], where it was shown that these geometries have generically two global (compact) U(1) Killing vectors. Moreover, it was shown in [30] (see also [29]) that there are two class of conserved charges associated with the above geometries: the charges associated with the two U(1)'s, the J_{\pm} charges, and the $L_n, \bar{L}_n, n \in \mathbb{Z}$ charge which form two copies of Virasoro algebra both at the Brown-Henneaux central charge (5). Importantly, J_{\pm} commute with L_n, \bar{L}_n [30]. Moreover, Bañados geometries are in one-to-one relation with representations of the left and right Virasoro groups, $Vir_+ \otimes$ Vir_{-} [29]. These representations are usually called Virasoro coadjoint orbits [31, 32]. The "thermodynamic" properties and quantities of the geometries (7) are shared among all geometries associated with the same orbit [29].

Appearance of Virasoro algebra suggests presence of a 2d CFT dual with central charge c (5) to the gravity theory (6). According to the standard AdS/CFT dictionary [20] the Bañados geometries are then dual to excited states in the presumed dual CFT and that the expectation value of the left and right components of the energy momentum tensor of the 2d CFT $T(x^+)$, $\bar{T}(x^-)$ are related to the functions L_{\pm} :

$$\langle T(x^+)\rangle = \frac{c}{6}L_+, \qquad \langle \bar{T}(x^-)\rangle = \frac{c}{6}L_-.$$
 (9)

The special case of $L_+ = L_- = 0$ corresponds to massless BTZ, and the vacuum state of the presumed dual 2d CFT on the plane (if we ignore the periodicity in x^{\pm}).

The primary operators which satisfy the unitarity bound, in the large c limit where we can trust classical gravity description, correspond to geometries with constant L_{\pm} with $L_{\pm} \geq -1/4$ and Bañados geometries in Virasoro coadjoint orbits with constant representative L_{\pm} correspond to generic descendants of these primaries in the presumed dual 2d CFT [29, 30]. Note that the relation between generic excitations and the geometry is not one-to-one. As (9) shows, all 2d CFT excitations with equal energy-momentum expectation values correspond to the same geometry. With the same token, any observable which is built from the geometry, like the HEE, would at most be able to encode information about the geometry, in the case of Bañados metrics; that is the information in functions L_{\pm} .

In this work we focus on generic excitations associated with constant representative orbits and study how the change in the excitations affects the entanglement entropy.

Holographic computation of the Entanglement Entropy. Consider a 2d CFT on a cylinder parametrized by x^{\pm} :

$$ds^{2} = -\ell^{2} dx^{+} dx^{-} = \ell^{2} (-dt^{2} + d\phi^{2}), \ \phi \in [0, 2\pi], \quad (10)$$

and an entangling interval whose endpoints are two spacelike separated points given by x_1^{\pm} and x_2^{\pm} . For holographic computation of the entanglement entropy we follow similar idea used in [15, 22, 23, 33]. To calculate the HEE associated with this interval for a generic excited state the expectation value of energy momentum tensor of which is given by (9), we note that all geometries of the form (7) are locally AdS₃. This means that there should be a coordinate transformation which locally brings the metric (7) to a similar metric with $L_{\pm} = 0$. One may readily check that such a coordinate transformation is of the form [34]

$$x^{+} \to y^{+} = \int \frac{dx^{+}}{\psi^{2}} - \frac{z^{2}\phi'}{\psi^{2}\phi(1 - z^{2}/z_{H}^{2})},$$

$$x^{-} \to y^{-} = \int \frac{dx^{-}}{\phi^{2}} - \frac{z^{2}\psi'}{\phi^{2}\psi(1 - z^{2}/z_{H}^{2})},$$

$$z \to w = \frac{z}{\psi\phi(1 - z^{2}/z_{H}^{2})},$$

(11)

where ψ, ϕ are solutions to the Hill's equation [28, 29]

$$\psi'' - L_+\psi = 0, \quad \phi'' - L_-\phi = 0.$$
 (12)

Each of the above two equations have two solutions, which may be denoted by ψ_1, ψ_2 and ϕ_1, ϕ_2 and one may choose the normalization

$$\psi'_1\psi_2 - \psi'_2\psi_1 = 1, \quad \phi'_1\phi_2 - \phi'_2\phi_1 = 1.$$
 (13)

In the above transformations (11) there are four choices for ψ 's and ϕ 's and four associated z_H^2 's:

$$z_{H}^{2} = \frac{\psi_{i}}{\psi_{i}'} \frac{\phi_{j}}{\phi_{j}'}, \quad i, j = 1, 2.$$
(14)

The z_H 's are radii of the Killing horizons of the Bañados geometry [29]. As expected, the coordinate transformation (11) is not globally defined, does not respect periodicity of x^{\pm} . Moreover, it is defined on one side of the inner and outer horizons.

To compute the HEE, we only need to know the length of geodesics anchored at the boundary at x_1^{\pm} , x_2^{\pm} , note that this length is invariant under local diffeomorphisms and that it only depends on the separation of the endpoints at the boundary (at z = 0). Therefore, for the HEE it suffices to know the near boundary behavior of (11). Such a coordinate transformation at z = 0 is like a conformal map at the boundary with [35]

$$x^+ \to H_+(x^+) = \frac{\psi_1}{\psi_2}, \quad x^- \to H_-(x^-) = \frac{\phi_1}{\phi_2}.$$
 (15)

Under this map

$$L_{\pm}(x^{\pm}) \to L_{\pm}^{H}(x^{\pm}) = H_{\pm}^{\prime 2}L_{\pm}(H_{\pm}) - \mathcal{S}[H_{\pm}; x^{\pm}],$$
 (16)

where $S[H_{\pm}; x^{\pm}]$ is the Schwarz derivative

$$\mathcal{S}[H;x] = \frac{H'''}{2H'} - \frac{3H''^2}{4H'^2},\tag{17}$$

and (10) is mapped to

$$ds^2 = -\ell^2 H'_+ H'_- \ dx^+ dx^-. \tag{18}$$

Here primes denote derivative w.r.t. the argument. One may then check that $S[H_{\pm}; x^{\pm}] = -L_{\pm}(x^{\pm})$ and hence (16) yields, $L^H_+(H_+(x^+)) = 0$, $L^H_-(H_-(x^-)) = 0$.

One can use (4) to readily compute the HEE for generic Bañados geometry

$$S_{HEE}(x_1, x_2) = \frac{c}{6} \ln \left(\frac{\mathcal{L}_+(x_1^+, x_2^+) \mathcal{L}_-(x_1^-, x_2^-)}{\epsilon^2} \right), \quad (19)$$

where

$$\mathcal{L}_{+}(x_{1}^{+}, x_{2}^{+}) = \psi_{1}(x_{1}^{+})\psi_{2}(x_{2}^{+}) - \psi_{2}(x_{1}^{+})\psi_{1}(x_{2}^{+}),$$

$$\mathcal{L}_{-}(x_{1}^{-}, x_{2}^{-}) = \phi_{1}(x_{1}^{-})\phi_{2}(x_{2}^{-}) - \phi_{2}(x_{1}^{-})\phi_{1}(x_{2}^{-}).$$
(20)

Recalling the fact that $H'_{+} = 1/\psi_2^2, H'_{-} = 1/\phi_2^2$, due to the conformal factor $H'_{+}H'_{-}$ in the metric (18) we have adjusted the cutoff ϵ with the mapping $x \to H(x)$. [36]. We also note that $S_{HEE}(x_1, x_2) = S_{HEE}(x_2, x_1)$ which readily follows from (20).

For the vacuum case $L_{\pm} = 0$, $\psi_1 = x^+, \psi_2 = 1$ (and similarly for ϕ 's) and therefore, $\mathcal{L} = x_1 - x_2$. As another example, let us consider constant positive $L_{\pm} = \mathcal{T}^2_{\pm}$ corresponding to BTZ black hole. For this case

$$\psi_{1} = \frac{1}{\sqrt{2\mathcal{T}_{+}}} e^{\mathcal{T}_{+}x^{+}}, \quad \psi_{2} = \frac{1}{\sqrt{2\mathcal{T}_{+}}} e^{-\mathcal{T}_{+}x^{+}},$$

$$\phi_{1} = \frac{1}{\sqrt{2\mathcal{T}_{-}}} e^{\mathcal{T}_{-}x^{-}}, \quad \phi_{2} = \frac{1}{\sqrt{2\mathcal{T}_{-}}} e^{-\mathcal{T}_{-}x^{-}}.$$
(21)

Then (19) simplifies to

$$S_{HEE} = \frac{c}{6} \ln \left(\frac{\sinh(\mathcal{T}_+ R_+)}{\mathcal{T}_+ \epsilon} \cdot \frac{\sinh(\mathcal{T}_- R_-)}{\mathcal{T}_- \epsilon} \right), \qquad (22)$$

where

$$R_{\pm} = \min(\Delta x^{\pm}, 2\pi - \Delta x^{\pm}), \quad \Delta x^{\pm} = |x_1^{\pm} - x_2^{\pm}|, \quad (23)$$

is the length of the entangling region (in units of AdS₃ radius ℓ). For an entangling region with the two endpoints of the same time coordinate $R_+ = R_- = R$, R being the length of entangling interval, this result of course matches that of [22, 23, 33] once we replace \mathcal{T}_{\pm} with the inverse temperature $\beta_{\pm} = \pi/\mathcal{T}_{\pm}$.

Excitation Entanglement Entropy. Descendants of any given primary operator are constructed from the primary by the action of a conformal transformations. In the terminology of Virasoro coadjoint orbits, i.e. primary operators correspond to constant representative orbits [29, 32]. At the level of the 3d dual geometries at least in the large c limit, as pointed out, primary operators in the 2d CFT correspond to geometries with constant L_{\pm} and the descendants are related through coordinate transformation which at the AdS₃ boundary are given as [29, 30]

$$x^{\pm} \to \tilde{x}^{\pm} = h^{\pm}(x^{\pm}), \ h_{\pm}(x^{\pm} + 2\pi) = h_{\pm}(x^{\pm}) + 2\pi, \ (24)$$

and $h'_{\pm} > 0$. Under this map L_{\pm} transform as (16) while $\psi_1, \psi_2 \to \tilde{\psi}_1, \tilde{\psi}_2$ where

$$\tilde{\psi}_1 = \frac{1}{\sqrt{h'_+}} \psi_1(h_+(x^+)), \ \tilde{\psi}_2 = \frac{1}{\sqrt{h'_+}} \psi_2(h_+(x^+)), \quad (25)$$

and similarly for ϕ 's. The descendants of primaries are hence specified by L_{\pm} functions of the form

$$L_{\pm} = h_{\pm}^{\prime 2} \mathfrak{T}_{\pm}^2 - \mathfrak{S}[h_{\pm}; x^{\pm}].$$
 (26)

One may now readily check how HEE (19) changes under conformal transformations (24) and hence obtain the HEE for the excitation generated by conformal transformation h,

$$S_{HEE}^{(h)}(x_1, x_2) = S_{HEE}(h(x_1), h(x_2)) - \Delta S_{EE}$$

$$\Delta S_{EE} = \frac{c}{12} \ln(h'_+(x_1^+)h'_+(x_2^+)h'_-(x_1^-)h'_-(x_2^-)).$$
(27)

We then define the excitation entanglement entropy

$$\Delta_h S(x_1, x_2) \equiv S_{HEE}^{(h)}(x_1, x_2) - S_{HEE}(x_1, x_2), \qquad (28)$$

i.e. $\Delta_h S$ computes the difference between the entanglement entropy of a (primary) excitation and its conformal descendants. For a 2d CFT, from (27)

$$\Delta_h S(x_1, x_2) = \frac{c}{6} \ln \left(\mathfrak{X}_+ \cdot \mathfrak{X}_- \right) \tag{29}$$

where

$$\mathfrak{X}_{\pm} = \frac{\mathcal{L}_{\pm}(h_{\pm}(1), h_{\pm}(2))}{\mathcal{L}_{\pm}(x_{1}^{\pm}, x_{2}^{\pm})\sqrt{h_{\pm}(1)h_{\pm}'(2)}},$$
(30)

where $X_{\pm}(\alpha) = X_{\pm}(x_{\alpha}^{\pm})$. The $\Delta_h S$ encodes both the information from the background (i.e. $L_{\pm}(x^{\pm})$) and the excitations h^{\pm} functions. We note that the cutoff dependence has been dropped from the excitation entanglement entropy (29). Moreover, the functions h^{\pm} and L_{\pm} can in principle be extracted if we know $\Delta_h S(x_1, x_2)$ for all x_1, x_2 .

As an example one may compute for the BTZ black hole and its conformal descendants. Using (20), (21) and (25), we obtain

$$\Delta_h S^{\text{BTZ}}(x_1, x_2) = \frac{c}{6} \ln \left(\mathcal{X}_+^{\text{BTZ}} \cdot \mathcal{X}_-^{\text{BTZ}} \right), \qquad (31)$$

with

Ì

$$\mathfrak{X}_{\pm}^{\text{BTZ}} = \frac{\sinh(\mathfrak{T}_{\pm}\sqrt{h'_{\pm}(1)h'_{\pm}(2)}R_{h_{\pm}})}{\sqrt{h'_{\pm}(1)h'_{\pm}(2)}\sinh(\mathfrak{T}_{\pm}R_{\pm})},\tag{32}$$

where R_{\pm} is given in (23) and

$$R_{h_{\pm}} = \frac{\min(\Delta h_{\pm}, 2\pi - \Delta h_{\pm})}{\sqrt{h'_{\pm}(1)h'_{\pm}(2)}}, \ \Delta h_{\pm} \equiv h_{\pm}(1) - h_{\pm}(2), \ (33)$$

where Δh_{\pm} shows the angular distance of the two endpoints of the entangling interval while $R_{h_{\pm}}$ is its physical length. (Note that after the map (24) the metric of the 2d boundary cylinder gets a conformal factor and hence the length of the space-like geodesic connecting them is not simply difference of the *h* functions at two end points.) For the special case of vacuum (massless BTZ) descendants with $T_{\pm} = 0$, the above reduces to

$$\chi_{\pm}^{\text{vac.desc.}} = \frac{R_{h\pm}}{R_{\pm}}.$$
(34)

This matches with the results of [15].

Comparison to 2d CFT results. The above holographic results may also be directly verified by 2d CFT computations. To this end, we use the standard analysis of [2, 7] and the replica trick. Although one may perform this analysis in a more general form, using a generic density matrix ρ with a primary, thermal or non-thermal excitation, for illustrative purposes and since it yields the same results, here we present the computation for a pure state excitation $\rho = |\Psi\rangle\langle\Psi|$ where Ψ denotes a generic of 2d CFT. We compute $\text{Tr}\rho^n$ for the interval A whose endpoints are denoted by coordinates z_1, z_2 (as usual we are considering the Euclidean 2d CFT on C-plane with complex coordinates z):

$$\operatorname{Tr} \rho_{A}^{n} = \frac{\langle \Psi | \sigma_{n}(z_{1}) \tilde{\sigma}_{n}(z_{2}) | \Psi \rangle_{\Sigma_{n}}}{(\langle \Psi | \Psi \rangle)^{n}}, \qquad (35)$$

where $\sigma, \tilde{\sigma}$ are the twist operators and Σ_n is the standard *n*-sheeted plane [7].

Any given state $|\Psi\rangle$ which is a part of a unitary representation (orbit) of the 2d conformal group and other states in the same family/orbit are related by a unitary transformation U generated by conformal transformations (descendant generating transformation):

$$|\Psi_{(h)}\rangle = U_h |\Psi\rangle \tag{36}$$

The conformal transformation U_h is associated with the conformal map (24) $z \to h(z), \bar{z} \to \bar{h}$. (Note that we are considering Euclidean 2d CFT on the plane and to compare it with our holographic gravity analysis we should replace z with x^+ , \bar{z} with x^- , h_+ with h(z) and h_- with \bar{h} .)

Then, as discussed in [15, 37],

$$\operatorname{Tr}(\rho_{(\mathrm{h}) \mathrm{A}}^{n}) = \frac{\langle \Psi | (U_{h}^{\dagger})^{n} \sigma_{n}(z_{1}) U_{h}^{n} (U_{h}^{\dagger})^{n} \tilde{\sigma}_{n}(z_{2}) U_{h}^{n} | \Psi \rangle_{\Sigma_{n}}}{(\langle \Psi | \Psi \rangle)^{n}}, \quad (37)$$

where

$$U_h^{\dagger}\sigma_n(z)U_h = h'(z)^{nh_{\sigma}}\bar{h}'(\bar{z})^{nh_{\sigma}}\sigma_n(h(z)), \qquad (38)$$

and $h_{\sigma} = \frac{c}{12}(n - \frac{1}{n})$. Following the lines of [15, 37] and recalling (2), above hence yields

$$S^{(h)}(z_1, z_2) = S(h(z_1), h(z_2)) - \frac{c}{6} \ln |h_1' h_2'|, \qquad (39)$$

where $h'_i = h'(z_i)$. We stress that (39) may be used to relate the entanglement entropy of any excitation (on the right-hand-side) and that of its conformal descendants generated by U_h (on the left-hand-side). (39) is of course the same result as (27) and hence leads to the same expression for the excitation entanglement entropy. For the particular case of vacuum excitations this result is precisely the well-known result of [8] (see also [15]) and is exactly what we have computed in (31) with (34).

Low and high temperature behavior. In real situations we usually do not have access to systems at zero temperature, while probing entanglement at low temperatures and studying low temperature behavior of the entanglement could be possible. Such an analysis for 2d CFT with a mass gap has been presented in [14]. As discussed in [29], the BTZ geometries associated with (21) and their conformal descendants have the same left and right temperatures \mathcal{T}_{\pm} . A system at temperature T is associated with $2/T = 1/\mathcal{T}_{+} + 1/\mathcal{T}_{-}$. Therefore, low temperature system corresponds to \mathcal{T}_{-} finite and $\mathcal{T}_{+}\Delta x \ll 1$, which is dual to near extremal BTZ black hole. The leading behavior of $\Delta_h S$ (32) is

$$\Delta_h S^{\text{low-temp.}} - \Delta_h S^{T=0} = \frac{c}{144} T^2 \big((\Delta h_+)^2 - (\Delta x_+)^2 \big), \quad (40)$$

where $\Delta_h S^{T=0} = \Delta_h S^{\text{BTZ}}$ at $\mathcal{T}_+ = 0$. As we see the deviation from the zero temperature is positive (recall that $h'_{\pm} > 0$) and starts from temperature-squared. That is, with the same excitation function h, increasing the temperature will increase the excitation entanglement entropy.

This is in contrast with the behavior of the thermal entropy of 2d CFT's which is linear in temperature. It is also different than the behavior in 2d CFT's with mass gap [14].

At high temperature one may again expand (32) for large T_{\pm} to obtain

$$\Delta_h S \sim \frac{c}{6} \left[\mathcal{T}_+ \left(\Delta h_+ - \Delta x_+ \right) + \mathcal{T}_- \left(\Delta h_- - \Delta x_- \right) \right].$$
(41)

As expected, the above is positive, that is the descendants (excited states) have bigger entanglement entropy than their primary operator excitations. Moreover, we have a linear temperature dependence. This linear behavior is expected, as at high temperature the entanglement entropy and the thermal entropy should show the same temperature dependence, e.g. see [23].

Thermal vs. excitation entanglement entropy. In our analysis both x^{\pm} and h_{\pm} are parametrizing coordinates on a circle, therefore, $\Delta x_{\pm}, \Delta h_{\pm} \in [0, 2\pi]$. As we have already used and discussed in the expression for entanglement entropy like (22) and (32) $R_{\pm}, R_{h_{\pm}}$ which are the physical length of the entangling interval should be used (see also [19]). This matches with the fact that with pure state excitations the entanglement entropy of a region Aand its complement are equal [38]. One can prove that in presence of thermal excitations, the difference between the entanglement entropy of the two "complementary" regions, the two entangling intervals which appear once we specify two points on the circle, is the thermal entropy of the corresponding black hole background/non-zero temperature 2d CFT system [23]. In our case one may see this in the $\Delta_h S$: Let us take $\Delta x_{\pm} = 2\pi$. Then we learn that $\Delta h_{\pm} = 2\pi$ (recall (24)). Therefore, for this case, recalling (29), $\Delta_h S = -\frac{c}{6} \ln(h'_+ h'_-)$ and is independent of temperature \mathcal{T}_{\pm} . (Note that the ratio of the physical length of the two entangling intervals associated with excitations his $h'_{+}h'_{-}$.) Recall that $\Delta_h S$ is the difference between the entanglement entropy of an excitation and the corresponding descendant and that, all of the these states are associated with the same temperature \mathcal{T}_{\pm} which is an orbit invariant quantity [29]. Note also that this $\Delta_h S$ can be positive or negative.

Small excitations. For small excitations around a thermal background with temperatures \mathcal{T}_{\pm} , i.e. for $h^{\pm}(x) = x^{\pm} + \epsilon_{\pm}(x)$, up to first order in ϵ we get:

$$\Delta_{\epsilon}S = \frac{c}{6} \left[\frac{\Im_{+}\Delta x^{+}}{\tanh(\Im_{+}\Delta x^{+})} \frac{\Delta \epsilon^{+}}{\Delta x^{+}} - \frac{1}{2} (\epsilon'_{+}(1) + \epsilon'_{+}(2)) + x^{-}\operatorname{sector} \right],$$

$$(42)$$

where $\epsilon(\alpha) = \epsilon(x_{\alpha})$ and $\Delta \epsilon^{+} = \epsilon_{+}(2) - \epsilon_{+}(1)$. For high temperature regime of $\Im \Delta x \gg 1$ this matches with (41) and for $\Im \Delta x \ll 1$ with the excitations above the vacuum (34),

$$\Delta_{\epsilon} S = \frac{c}{3} \operatorname{Re} \left(\frac{\epsilon(z_1) - \epsilon(z_2)}{z_1 - z_2} - \frac{1}{2} \epsilon'(z_1) - \frac{1}{2} \epsilon'(z_2) \right).$$

Note that for arbitrary $z_1, z_2, \Delta_{\epsilon}S$ is not necessarily positive definite, even when the expectation value of energy momentum tensors associated with the excitations $\delta_{\epsilon}T = -\frac{c}{12}\epsilon_{+}^{\prime\prime\prime}, \delta_{\epsilon}\bar{T} = -\frac{c}{12}\epsilon_{-}^{\prime\prime\prime}$ are positive.

Arbitrary excitation for small entangling interval. Another interesting limit to study is the arbitrary excitation for small entangling interval, i.e. when $x_1^{\pm} = x^{\pm}, x_2^{\pm} = x^{\pm} + \delta x^{\pm}$. For this case (29) yields

$$\delta \Delta_h S(x) = \frac{c}{36} \left(\Delta_{h_+} L_+ \cdot (\delta x^+)^2 + \Delta_{h_-} L_- \cdot (\delta x^-)^2 \right), \quad (43)$$

where $\Delta_h L$ is variation of L under conformal transformation,

$$\Delta_{h_{\pm}} L_{\pm} = (h_{\pm}^{\prime 2} - 1) \mathcal{T}_{\pm}^2 - \mathcal{S}[h_{\pm}; x^{\pm}].$$
(44)

Recalling (9), one may rewrite (43) as

$$\left(\partial_{x^{\pm}}^2 \Delta_h S(x,y)\right)_{y=x} = \frac{1}{3} \Delta_{h_{\pm}} \langle T_{\pm}(x) \rangle.$$
(45)

The above equation directly relates the variation in the entanglement entropy to the variation in the expectation value of the energy momentum tensor in the 2d CFT. Recalling the periodicity conditions on $\Delta_h S$ and h and

$$\Delta_h S(x,y) = \Delta_h S(y,x),$$

$$\Delta_h S(x,y)|_{x=y} = 0, \ \left(\partial_{x^{\pm}} \Delta_h S(x,y)\right)_{x=y} = 0,$$
(46)

(45) together with (9) and (44) completely determines $\Delta_h S(x, y)$. In fact, the solution is obtained to be precisely the one in (32). Note that while (45) is written only in terms of 2d CFT quantities, it has a natural interpretation in terms of first law of entanglement and how it is related to perturbation of Einstein equation in the bulk [10, 39, 40].

Integrated positivity. For generic h and L, $\Delta_h L$ and hence $\delta \Delta_h S(x)$, do not have a definite sign for generic hand for all values of x. Nonetheless, for the constant representative orbits one can prove that [31, 32]

$$\int_0^{2\pi} dx^{\pm} \Delta_{h_{\pm}} L_{\pm} \ge 0, \qquad (47)$$

and hence we have an *integrated positivity* for $\delta \Delta_h S(x)$, i.e. the average value $\partial_x^2 \Delta_h S(x, y)|_{y=x}$ is non-negative.

Similarly, for generic entangling interval Δx^{\pm} , as one can see from explicit expressions (32), $\Delta_h S(x, x + \Delta x)$ do not have a definite sign. Nonetheless one can prove integrated positivity relation for generic case given in (31) with (32):

$$\int_{0}^{2\pi} dx^{\pm} \Delta_h S(x, x + \Delta x) \ge 0.$$
(48)

We note that the above integral is basically giving the average of the excitation entanglement entropy which is potentially a good physical observables in backgrounds where translation symmetry is broken by the excitations.

Multi-partite information and for arbitrary excitations. Besides the entanglement entropy there are other information theoretic measures involving more than one entangling interval. The simplest such quantity defined between two entangling regions A, B (A, B may have overlapping regions or not) is the mutual information defined as [41]

$$I(A:B) = S_A + S_B - S_{A\cup B} \tag{49}$$

where S on the RHS is the entanglement entropy. In the regions A, B we may have various excitations, e.g. excitations by an operator or by its conformal descendants. Using (27), we find that

$$I(A:B)_{des}(x_1^{\pm}, x_2^{\pm}) = I(A:B)_{pri.}(h^{\pm}(x_1), h^{\pm}(x_2)).$$
 (50)

The above shows how the mutual information with excitations and their descendants are related to each other. Note that in the mutual information the ΔS_{EE} (27) drops out. Similar result and statement can also be made for multipartite information with a similar result as in (50).

Concluding remarks and discussion. We have studied, holographically and within 2d CFT, the entanglement entropy for a class of excitations and their descendants which can be represented through the expectation value of the energy momentum tensor (9), at zero or finite temperature. Although we mainly considered $L_{\pm} = \mathcal{T}_{\pm}^2 \geq 0$ cases, all of our results are also true for $-1/4 \leq L_{\pm} \leq 0$, by taking \mathcal{T} to be imaginary.

In particular, by introducing excitation entanglement entropy $\Delta_h S$ we quantified how the entanglement entropy changes as we turn on conformal excitations. $\Delta_h S$ is cutoff independent and finite, and can be a good physical measure for the system. Our results are compatible with and extends the results of [9].

As discussed $\Delta_h S$ encodes all the information about the excitation function h and the expectation value of the energy-momentum tensor of the excitation operator L_{\pm} . Therefore, one can in principle read off all the information about the geometry from $\Delta_h S$. To obtain our results we used the map (11). This coordinate transformation maps the outside horizon region of the Bañados geometries [29] to AdS₃ in Poincaré patch. So, we would expect to be able to reconstruct the outside horizon geometry using the $\Delta_h S$, in the same manner outlined in [5].

The excitation entanglement entropy $\Delta_h S$ is closely related to the geometry (and metric) of the kinematic space, the geometry of CFT intervals. In this regard (45) and (46) may be viewed as equations for metric of the kinematic space and the integrated positivity (48) as a statement about positive norms on the kimentic space. Such analyses has been initiated in [42]. See also [43].

Relative entropy [44] is another information theoretic measure which may be used to compare an excitation due to a primary operator and it descendants. It is interesting to compute and compare relative and our excitation entanglement entropy for the cases we analyzed here.

One may study how local operations, e.g. local projections measurements, affect the excitation entanglement entropy. This question can be posed and answered within the 2d CFT or the dual AdS_3 gravity setting. It would be interesting to extend the corresponding analysis [45] for We would like to especially thank Joan Simon for his role in the development of our analysis and results and for long discussions on various stages of this work. We would also like to that Pawel Caputa for collaboration in early stages of this work and Jan de Boer and Tadashi Takayanagi for comments on the draft. The work of M.M. Sh-J is supported in part by Allameh Tabatabaii Prize Grant of Boniad Melli Nokhbegan of Iran, the SarAmadan grant of Iranian vice presidency in science and technology and the ICTP network project NET-68 and the ICTP Simons fellowship. We would like to thank ICTP, Trieste for the workshop on aspects of 3d gravity and the hospitality in the course of this project.

Appendix: Proof of integrated positivity (48).

Denoting $I^{\pm}(\Delta x) = \int_{0}^{2\pi} dx^{\pm} \Delta_{h} S(x, x + \Delta x)$, we have already seen that for small intervals $\Delta x^{\pm} \ll 1$, (47) yields $I_{\pm} \geq 0$. To prove this statement for arbitrary intervals Δx^{\pm} , it is enough to show that $\frac{dI^{\pm}}{d\Delta x} \geq 0$. From (32) and (31) we get

$$\frac{dI^{\pm}}{d\Delta x} = \int_0^{2\pi} \left[\frac{\mathfrak{T}_{\pm} h'_{\pm} (x^{\pm} + \Delta x^{\pm})}{\tanh \mathfrak{T}_{\pm} \Delta h_{\pm}} - \frac{\mathfrak{T}_{\pm}}{\tanh \mathfrak{T}_{\pm} \Delta x^{\pm}} \right] dx^{\pm}.$$

To show that the right hand side of above equation is positive we show that

$$\int_0^{2\pi} \frac{\tanh \mathfrak{T}_{\pm} \Delta x^{\pm}}{\tanh \mathfrak{T}_{\pm} \Delta h_{\pm}} h'_{\pm} (x^{\pm} + \Delta x^{\pm}) \, dx^{\pm} \ge 2\pi.$$
 (51)

Noting that functions h'_{\pm} and Δh_{\pm} are positive everywhere and that $\tanh x$ is positive and increasing for x > 0, then

$$\int_{0}^{2\pi} \frac{\tanh \mathfrak{T}_{\pm} \Delta h_{\pm}}{\tanh \mathfrak{T}_{\pm} \Delta x^{\pm}} \leq \int_{0}^{2\pi} \frac{\Delta h_{\pm}}{\Delta x^{\pm}} = 2\pi.$$
(52)

To prove the above, we have used the fact that

$$\int_{0}^{2\pi} h'_{\pm} dx^{\pm} = \int_{0}^{2\pi} \frac{\Delta h_{\pm}}{\Delta x^{\pm}} = 2\pi,$$

and that for any two positive 2π -periodic functions f, g,

$$\int_{0}^{2\pi} \frac{f}{g} dx \ge 2\pi \frac{\int_{0}^{2\pi} f dx}{\int_{0}^{2\pi} g dx}.$$

- P. Calabrese and J. L. Cardy, Int. J. Quant. Inf. 4, 429 (2006) [quant-ph/0505193].
- [2] P. Calabrese and J. L. Cardy, J. Stat. Mech. 0406, P06002 (2004) [hep-th/0405152].
- [3] H. Casini and M. Huerta, J. Phys. A 42, 504007 (2009) [arXiv:0905.2562 [hep-th]].
- [4] L. Bombelli, R. K. Koul, J. Lee and R. D. Sorkin, Phys. Rev. D 34, 373 (1986).
 M. Srednicki, Phys. Rev. Lett. 71, 666 (1993) [hep-th/9303048].

- [5] M. Van Raamsdonk, Gen. Rel. Grav. 42, 2323 (2010) [Int. J. Mod. Phys. D 19, 2429 (2010)] [arXiv:1005.3035 [hep-th]].
- [6] S. N. Solodukhin, Living Rev. Rel. 14, 8 (2011) [arXiv:1104.3712 [hep-th]].
- [7] P. Calabrese and J. Cardy, J. Phys. A 42, 504005 (2009)
 [arXiv:0905.4013 [cond-mat.stat-mech]].
- [8] C. Holzhey, F. Larsen and F. Wilczek, Nucl. Phys. B 424, 443 (1994) [hep-th/9403108].
- [9] F. C. Alcaraz, M. I. Berganza and G. Sierra, Phys. Rev. Lett. 106, 201601 (2011) [arXiv:1101.2881 [cond-mat.statmech]].
- [10] J. Bhattacharya, M. Nozaki, T. Takayanagi and T. Ugajin, Phys. Rev. Lett. **110**, no. 9, 091602 (2013) [arXiv:1212.1164].
- [11] P. Caputa, M. Nozaki and T. Takayanagi, PTEP 2014, 093B06 (2014) [arXiv:1405.5946 [hep-th]].
- [12] W. Z. Guo and S. He, JHEP **1504**, 099 (2015) [arXiv:1501.00757 [hep-th]].
 B. Chen, W. Z. Guo, S. He and J. q. Wu, JHEP **1510**, 173 (2015) [arXiv:1507.01157 [hep-th]].
 P. Caputa and A. Veliz-Osorio, Phys. Rev. D **92**, no. 6, 065010 (2015) [arXiv:1507.00582 [hep-th]].
- [13] M. Nozaki, T. Numasawa and T. Takayanagi, Phys. Rev. Lett. **112**, 111602 (2014) [arXiv:1401.0539 [hep-th]].
- [14] J. Cardy and C. P. Herzog, Phys. Rev. Lett. 112, no. 17, 171603 (2014) [arXiv:1403.0578 [hep-th]].
- [15] M. M. Roberts, JHEP **1212** (2012) 027 [arXiv:1204.1982 [hep-th]].
- [16] G. Wong, I. Klich, L. A. Pando Zayas and D. Vaman, JHEP
 1312, 020 (2013) [arXiv:1305.3291 [hep-th]].
 S. R. Das, D. A. Galante and R. C. Myers, Phys. Rev. Lett.
 112, 171601 (2014) [arXiv:1401.0560 [hep-th]].
 P. Caputa, J. Simón, A. Štikonas, T. Takayanagi and
 K. Watanabe, JHEP 1508, 011 (2015) [arXiv:1503.08161 [hep-th]].
- [17] P. Caputa, J. Simón, A. Štikonas and T. Takayanagi, JHEP 1501, 102 (2015) [arXiv:1410.2287 [hep-th]].
- [18] M. Nozaki, T. Numasawa and T. Takayanagi, JHEP 1305, 080 (2013) [arXiv:1302.5703 [hep-th]].
- [19] C. T. Asplund, A. Bernamonti, F. Galli and T. Hartman, JHEP **1502**, 171 (2015) [arXiv:1410.1392 [hep-th]].
- [20] O. Aharony, S. S. Gubser, J. M. Maldacena, H. Ooguri and Y. Oz, Phys. Rept. **323** (2000) 183 [hep-th/9905111].
- [21] S. Ryu and T. Takayanagi, Phys. Rev. Lett. 96, 181602 (2006) [hep-th/0603001].
- [22] T. Nishioka, S. Ryu and T. Takayanagi, J. Phys. A 42, 504008 (2009) [arXiv:0905.0932 [hep-th]].
- [23] V. E. Hubeny, M. Rangamani and T. Takayanagi, JHEP 0707, 062 (2007) [arXiv:0705.0016 [hep-th]].
- [24] J. D. Brown and M. Henneaux, Commun. Math. Phys. 104, 207 (1986).
- [25] M. Banados, C. Teitelboim and J. Zanelli, Phys. Rev. Lett.
 69, 1849 (1992), [hep-th/9204099].
 M. Banados, M. Henneaux, C. Teitelboim and J. Zanelli, Phys. Rev. D 48, 1506 (1993) [gr-qc/9302012].
- [26] M. Bañados, "Three-Dimensional Quantum Geometry and Black Holes," hep-th/9901148.

- [27] For example see, J. D. Brown, "Lower Dimensional Gravity," Singapore, World Sscientific (1988).
 S. Carlip, "Quantum gravity in 2+1 dimensions," Cambridge, UK: Univ. Pr. (1998).
- [28] M. M. Sheikh-Jabbari and H. Yavartanoo, JHEP 1407, 104 (2014) [arXiv:1404.4472 [hep-th]];
- [29] M. M. Sheikh-Jabbari and H. Yavartanoo, arXiv:1603.05272 [hep-th].
- [30] G. Compère, P. J. Mao, A. Seraj and M. M. Sheikh-Jabbari, JHEP 1601, 080 (2016) [arXiv:1511.06079 [hep-th]].
- [31] E. Witten, Commun. Math. Phys. **114**, 1 (1988).
- [32] J. Balog, L. Feher and L. Palla, Int. J. Mod. Phys. A 13, 315 (1998) [hep-th/9703045].
- [33] V. E. Hubeny, H. Maxfield, M. Rangamani and E. Tonni, JHEP **1308**, 092 (2013) [arXiv:1306.4004 [hep-th]].
 H. Maxfield, JHEP **1504**, 031 (2015) [arXiv:1412.0687 [hep-th]].
- [34] The coordinate transformations (11) were also given in [15] directly in terms of L_{\pm} , and not in terms of ψ, ϕ functions.
- [35] There are in fact four choices for H_{\pm} , associated with the four possible choices in (11). These are explicitly $H_{+} = -\psi_2/\psi_1$ or $H_{+} = \psi_1/\psi_2$ and similarly for H_{-} . It is straightforward to see that our results and analysis for (excitation) entanglement entropy is independent of these choices.
- [36] Although we are using the mapping to AdS₃ in Poincare patch, one could have directly performed the HRT proposal [23] to obtain the same result as (19).
- [37] G. Mandal, R. Sinha and N. Sorokhaibam, JHEP 1501, 036 (2015), arXiv:1405.6695 [hep-th].
- [38] H. Araki and E. H. Lieb, Commun. Math. Phys. 18, 160 (1970).
- [39] N. Lashkari, M. B. McDermott and M. Van Raamsdonk, JHEP **1404**, 195 (2014) [arXiv:1308.3716 [hep-th]].
 T. Faulkner, M. Guica, T. Hartman, R. C. Myers and M. Van Raamsdonk, JHEP **1403** (2014) 051 [arXiv:1312.7856 [hep-th]].
- [40] J. Bhattacharya, V. E. Hubeny, M. Rangamani and T. Takayanagi, Phys. Rev. D 91, no. 10, 106009 (2015) [arXiv:1412.5472 [hep-th]].
- [41] M. Nielsen and I. Chuang, Quantum computation and quantum information, Cambridge University Press, 2000.
- [42] B. Czech, L. Lamprou, S. McCandlish and J. Sully, arXiv:1512.01548 [hep-th].
 B. Czech, L. Lamprou, S. McCandlish, B. Mosk and J. Sully, arXiv:1604.03110 [hep-th].
- [43] J. de Boer, M. P. Heller, R. C. Myers and Y. Neiman, Phys. Rev. Lett. **116**, no. 6, 061602 (2016) [arXiv:1509.00113 [hep-th]].

C. T. Asplund, N. Callebaut and C. Zukowski, arXiv:1604.02687 [hep-th].

M. J. S. Beach, J. Lee, C. Rabideau and M. Van Raamsdonk, arXiv:1604.05308 [hep-th].

- [44] D. D. Blanco, H. Casini, L. Y. Hung and R. C. Myers, JHEP **1308**, 060 (2013) [arXiv:1305.3182 [hep-th]].
 G. Sárosi and T. Ugajin, arXiv:1603.03057 [hep-th].
- [45] T. Numasawa, N. Shiba, T. Takayanagi and K. Watanabe, arXiv:1604.01772 [hep-th].