

A Linear-Time Algorithm for the Weighted Paired-Domination Problem on Block Graphs*

Ching-Chi Lin[†]

Cheng-Yu Hsieh[‡]

December 9, 2021

Abstract

In a graph $G = (V, E)$, a vertex subset $S \subseteq V(G)$ is said to be a dominating set of G if every vertex not in S is adjacent to a vertex in S . A dominating set S of G is called a paired-dominating set of G if the induced subgraph $G[S]$ contains a perfect matching. In this paper, we propose an $O(n + m)$ -time algorithm for the weighted paired-domination problem on block graphs using dynamic programming, which strengthens the results in [Theoret. Comput. Sci., 410(47–49):5063–5071, 2009] and [J. Comb. Optim., 19(4):457–470, 2010]. Moreover, the algorithm can be completed in $O(n)$ time if the block-cut-vertex structure of G is given.

Keywords: Weighted paired-domination problem, perfect matching, block graph, dynamic programming.

*This work is partially supported by the National Science Council under the Grants No. NSC-102-2221-E-019-038-, and NSC-103-2221-E-019-034-.

[†]Department of Computer Science and Engineering, National Taiwan Ocean University, Keelung 20224, Taiwan. Corresponding author. Email: lincc@mail.ntou.edu.tw

[‡]Department of Computer Science and Information Engineering, National Taiwan University, Taipei 10617, Taiwan. Email: r01922114@ntu.edu.tw

1 Introduction

In a graph $G = (V, E)$, a vertex subset $S \subseteq V(G)$ is said to be a *dominating set* of G if every vertex not in S is adjacent to a vertex in S . Let $G[S]$ denote the subgraph of G induced by a subset S of $V(G)$. A dominating set S of G is called a *paired-dominating set* if the induced subgraph $G[S]$ contains a perfect matching. The *paired-domination problem* involves finding a paired-dominating set S of G such that the cardinality of S is minimized. Suppose that, for each $v \in V(G)$, we have a weight $w(v)$ specifying the cost for adding v to S . The *weighted paired-domination problem* is to find a paired-dominating set S whose $w(S) = \sum_{v \in S} w(v)$ is minimized.

The domination problem has been extensively studied in the area of algorithmic graph theory for several decades; see [2, 7, 9–13] for books and survey papers. It has many applications in the real world such as location problems, communication networks, and kernels of games [10]. Depending on the requirements of different types of applications, there are several variants of the domination problem, such as the independent domination, connected domination, total domination, and perfect domination problems [2, 7, 13, 23]. These problems have been proved to be NP-complete and have polynomial-time algorithms on some special classes of graphs. In particular, Haynes and Slater [8] introduced the concept of paired-domination motivated by security concerns. In a museum protection program, beside the requirement that each region has a guard in it or is in the protection range of some guard, the guards must be able to back each other up.

The *paired-domination number* $\gamma_p(G)$ is the minimum cardinality of a paired-dominating set. In [8], Haynes and Slater showed that the problem of determining whether $\gamma_p(G) \leq c$ is NP-complete on general graphs and gave a lower bound of $n/\Delta(G)$ for $\gamma_p(G)$, where c is a positive even integer, n is the number of the vertices in G , and $\Delta(G)$ is the maximum degree of G . Recently, many studies have been made for this problem in proving NP-completeness, providing approximation algorithms, and finding polynomial-time algorithms on some special classes of graphs. Here, we only mention some related results. For more

detailed information regarding this problem, please refer to [15]. Chen *et al.* [5] demonstrated that the paired-domination problem is also NP-complete on bipartite graphs, chordal graphs, and split graphs. In [3], Chen *et al.* proposed an approximation algorithm with ratio $\ln(2\Delta(G)) + 1$ for general graphs and showed that the problem is APX-complete, i.e., has no PTAS. Panda and Pradhan [21] strengthened the results in [5] by showing that the problem is also NP-complete for perfect elimination bipartite graphs.

Meanwhile, polynomial-time algorithms have been studied intensively on some special classes of graphs such as tree graphs [22], weighted tree graphs [3], inflated tree graphs [16], convex bipartite graph [14, 20], permutation graphs [6, 17, 18], strongly chordal graphs [4], interval graphs [5] and circular-arc graphs [19]. Especially, Chen *et al.* [5] introduced an $O(m+n)$ -time algorithm for block graphs, a proper superfamily of tree graphs. In this paper, we propose an $O(n+m)$ -time algorithm for the weighted paired-domination problem on block graphs using dynamic programming, which strengthens the results in [3, 5]. Moreover, the algorithm can be completed in $O(n)$ time if the block-cut-vertex structure of G is given. Notice that the block-cut-vertex structure of a block graph G can be constructed in $O(n+m)$ time by the depth first search algorithm [1].

The remainder of this paper is organized as follows. In Section 2, given the block-cut-vertex structure of a block graph G , we employ dynamic programming to present an $O(n)$ -time algorithm for finding a minimum-weight paired-dominating set of G . In Section 3, the correctness proof and complexity analysis of the algorithm are provided. Section 4 contains some concluding remarks and future work.

2 The Proposed Algorithm for Block Graphs

In this section, given a weighted block graph G with the block-cut-vertex structure G^* of G , we propose an $O(n)$ -time algorithm that determines a minimum-weight paired-dominating set of G using dynamic programming. Since a graph G containing isolated vertices has no paired-dominating set, we suppose that G is a connected graph without isolated vertices in the rest of this paper. First, we introduce some preliminaries for block graphs.




Figure 1: (a) A block graph G . (b) The corresponding block-cut-vertex graph G^* for the block graph G in (a). In particular, Algorithm 1 considered the blocks of G in turn according to the ordering B_1, B_2, \dots, B_8 .

For any connected graph G , a vertex $x \in V(G)$ is called a *cut-vertex* of G , if $G - x$ contains more than one connected component. A *block* is a maximal connected subgraph without a cut-vertex. A graph G is called a *block graph*, if every block in G is a complete graph. Notice that block graphs are a proper superfamily of tree graphs and a proper subfamily of chordal graphs. Suppose G has blocks B_1, B_2, \dots, B_x and cut vertices c_1, c_2, \dots, c_y . We define the *block-cut-vertex graph* $G^* = (V, E)$ of G , where

$$V(G^*) = \{B_1, B_2, \dots, B_x, c_1, c_2, \dots, c_y\}; \text{ and}$$

$$E(G^*) = \{(B_i, c_j) \mid c_j \in B_i, 1 \leq i \leq x, 1 \leq j \leq y\}.$$

Consequently, the graph G^* is a tree and the leaves in G^* are precisely the blocks with exactly one cut-vertex in G . A block containing exactly one cut-vertex in G is called a *pendant block*. It should be noted that, by using the depth first search algorithm, one can recognize the block graphs and construct the block-cut-vertex graphs G^* , both in $O(n + m)$ time [1]. Figure 1 shows an illustrative example, in which Figure 1(b) depicts the corresponding block-cut-vertex graph G^* for the block graph G in Figure 1(a). Clearly, G has 8

blocks B_1, B_2, \dots, B_8 and 6 cut vertices c_1, c_2, \dots, c_6 . Moreover, the pendant blocks of G are B_1, B_2, B_4, B_5 , and B_7 .

2.1 The algorithm

In this subsection, given the block-cut-vertex structure of a weighted block graph G , we propose an $O(n)$ -time algorithm for finding a minimum-weight paired-dominating set of G . Before describing the approach in detail, four notations $D(H, u)$, $P(H, u)$, $P'(H, u)$, and $\bar{P}(H, u)$ are defined below, where H is a subgraph of G and $u \in V(H)$. The notations are introduced for the purpose of describing the recursive formulations used in developing dynamic programming algorithms.

$D(H, u)$: A minimum-weight dominating set of H containing u , and $H[D(H, u) - u]$ has a perfect matching.

$P(H, u)$: A minimum-weight paired-dominating set of H containing u .

$P'(H, u)$: A minimum-weight paired-dominating set of H not containing u .

$\bar{P}(H, u)$: A minimum-weight paired-dominating set of $H - u$, and u is not dominated by $\bar{P}(H, u)$.

Clearly, either $P(G, u)$ or $P'(G, u)$ is a minimum-weight paired-dominating set of G . For ease of subsequent discussion, $D(H, u)$, $P(H, u)$, $P'(H, u)$, and $\bar{P}(H, u)$ are called a κ_1 -paired-dominating set, κ_2 -paired-dominating set, κ_3 -paired-dominating set, and κ_4 -paired-dominating set of H with respect to u , respectively. Suppose that H is a weighted block graph and B is a block of H with $V(B) = \{u_1, u_2, \dots, u_k\}$. The following lemma shows an useful property which help us design efficient algorithms.

Lemma 1 *Suppose that H is a weighted block graph and B is a block of H with $V(B) = \{u_1, u_2, \dots, u_k\}$. If G_i is a maximal connected subgraph in $(H - B) \cup \{u_i\}$ for $1 \leq i \leq k$, then G_i and G_j have disjoint vertex sets for $i \neq j$.*




Figure 2: A weighted block graph $H = B \cup G_1 \cup G_2 \cup \dots \cup G_k$.

Proof. Suppose to the contrary that $V(G_i) \cap V(G_j) \neq \emptyset$. Then $G[B \cup G_i \cup G_j]$ is a connected subgraph of H without a cut-vertex, this contradicts our assumption that B is a maximal connected subgraph without a cut-vertex. \square

Refer to Figure 2 for an illustrative example. In order to obtain a minimum-weight paired-dominating set of G , we use dynamic programming to iteratively determine $D(H, u_1)$, $P(H, u_1)$, $P'(H, u_1)$, and $\bar{P}(H, u_1)$ in a bottom-up manner. One block is considered in each iteration of the loop. Suppose the dominating sets $D(G_i, u_i)$, $P(G_i, u_i)$, $P'(G_i, u_i)$, and $\bar{P}(G_i, u_i)$ have been determined in the previous iterations and are recorded in u_i for $1 \leq i \leq k$. We shall show that $D(H, u_1)$, $P(H, u_1)$, $P'(H, u_1)$, and $\bar{P}(H, u_1)$ can all be determined in $O(k)$ time in Section 3. With the aid of this result, we now propose the main algorithm of this paper. Notice that during the computation, the block-cut-vertex structure G^* of the block graph G can be exploited to get the relationship among blocks, which help us to apply dynamic programming.

The algorithm first sets the *current graph* $G' = G$ and the set of *processed blocks* $W = \emptyset$. Further, it initially assigns $D(G[\{v\}], v) = \{v\}$, $P(G[\{v\}], v) = \Delta$, $P'(G[\{v\}], v) = \Delta$ and $\bar{P}(G[\{v\}], v) = \emptyset$ to each vertex $v \in V(G)$. Specially, we use Δ to denote the empty set with a weight of infinity, i.e., $\Delta = \emptyset$ and $w(\Delta) = \infty$. The algorithm then iteratively processes blocks in the repeat loop. During each iteration of the loop, we remove a pendant block B in the current graph G' and determines the dominating sets $D(H, u)$, $P(H, u)$, $P'(H, u)$,

and $\bar{P}(H, u)$, where u is the cut vertex and H is the connected component containing u in $G[B \cup W]$. After the execution of the repeat loop, we have only one block left, i.e., the current graph G' is a block and G^* is a vertex. With the information determined in the repeat loop, we now can find the two paired-dominating sets $P(G, u)$ and $P'(G, u)$, where u is an arbitrary vertex in G' . Finally, the output S is selected from $P(G, u)$ and $P'(G, u)$ based on the weights of the sets. The steps of the algorithm are detailed below.

Algorithm 1 Finding a paired-dominating set on weighted block graphs

Input: A weighted block graph G with the block-cut-vertex structure G^* of G .

Output: A minimum-weight paired-dominating set S of G .

```

1: let  $G' \leftarrow G$  and  $W \leftarrow \emptyset$ ;
2: for each  $v \in V(G)$  do
3:   let  $D(G[\{v\}], v) \leftarrow \{v\}$  and  $P(G[\{v\}], v) \leftarrow \Delta$ ;
4:   let  $P'(G[\{v\}], v) \leftarrow \Delta$  and  $\bar{P}(G[\{v\}], v) \leftarrow \emptyset$ ;
5: end for
6: repeat
7:   arbitrarily choose a leaf  $v_B$  in  $G^*$ ;
8:   let  $B$  be the corresponding pendant block of  $v_B$  in  $G'$ ;
   suppose that  $V(B) = \{u_1, u_2, \dots, u_k\}$ , where  $u_1$  is the cut vertex and  $G_i$  is the connected component in  $G[W]$  such that  $V(B) \cap V(G_i) = u_i$  for  $1 \leq i \leq k$ ;
9:   let  $H \leftarrow B \cup G_1 \cup G_2 \cup \dots \cup G_k$ ;
10:  find  $D(H, u_1), P(H, u_1), P'(H, u_1), \bar{P}(H, u_1)$  by using the dominating sets  $D(G_i, u_i), P(G_i, u_i), P'(G_i, u_i)$ , and  $\bar{P}(G_i, u_i)$  determined in the previous iterations;
11:  record the results  $D(H, u_1), P(H, u_1), P'(H, u_1), \bar{P}(H, u_1)$  in vertex  $u_1$ ;
12:  let  $G' \leftarrow G' - \{u_2, \dots, u_k\}$  and  $W \leftarrow W \cup B$ ;
13:  suppose  $v_c$  is the neighbor of  $v_B$  in  $G^*$ ;
   let  $G^* \leftarrow G^* - \{v_B, v_c\}$  if  $v_c$  is a leaf in  $G^* - v_B$ , and let  $G^* \leftarrow G^* - v_B$  otherwise;
14: until  $G^*$  itself is a vertex
15: find  $P(G, u)$  and  $P'(G, u)$ , where  $u$  is an arbitrary vertex in  $G'$ ;
16: let  $S \leftarrow P(G, u)$  if  $w(P(G, u)) < w(P'(G, u))$ , and let  $S \leftarrow P'(G, u)$  otherwise;
17: return  $S$ .
```

For an illustrative example, consider the block graph G in Figure 1. In the beginning, the algorithm sets the default values to each vertex v in G . Then, by the rules of removing blocks and recording results, one block is removed from G' for each iteration of the repeat loop. The blocks in G are removed with respect to the ordering B_1, B_2, \dots, B_8 .




Figure 3: The intermediate execution steps of Algorithm 1. The blocks in G are removed with respect to the ordering B_1, B_2, \dots, B_8 .

Figure 3(a) depicts the case that block $B_1 = G[\{v_{10}, v_{14}\}]$ is selected in the first iteration. One can see that $H = G[\{v_{10}, v_{14}\}] \cup G[\{v_{10}\}] \cup G[\{v_{14}\}]$. Four dominating sets $D(H, v_{10}), P(H, v_{10}), P'(H, v_{10}), \bar{P}(H, v_{10})$ are recorded in vertex v_{10} by the rules of determining and recording results in Steps 10 and 11. Please refer to Figure 3(b) for the result of removing v_{14} from G' .

Similarly, Figure 3(c) depicts the case that block $B_3 = G[\{v_6, v_{10}, v_{11}\}]$ is selected in the third iteration with $H = G[\{v_6, v_{10}, v_{11}\}] \cup G[\{v_6\}] \cup G[\{v_{10}, v_{14}\}] \cup G[\{v_{11}, v_{15}\}]$. Again, by the rules of determining and recording results, vertices v_{10} and v_{11} were removed from G' and the corresponding results are recorded in vertex v_6 . Refer to Figure 3(d) for an illustrative example. After removing blocks B_1, B_2, \dots, B_7 , we have exactly one block B_8 left in G' , i.e., G^* now is a vertex, then the algorithm exits the repeat loop. In Step 15, the two dominating sets $P(G, v_1)$ and $P'(G, v_1)$ are determined by using a similar method of the arguments in Steps 10 and 11. Refer to Figure 3(e) for an illustrative example. Clearly, either $P(G, u)$ or $P'(G, u)$ is a minimum-weight paired-dominating set S of G depending on which has the smaller total weight.

In next section, given the dominating sets $D(G_i, u_i), P(G_i, u_i), P'(G_i, u_i)$, and $\bar{P}(G_i, u_i)$ for $1 \leq i \leq k$, four dynamic programming procedures are proposed in Subsections 3.1–3.4, which can determine $D(H, u_1), P(H, u_1), P'(H, u_1)$, and $\bar{P}(H, u_1)$ in $O(k)$ time, respectively. Clearly, the proposed procedures ensure the correctness of the algorithm. For the complexity analysis, suppose that G has blocks B_1, B_2, \dots, B_x . Since the dynamic programming procedures can be completed in $O(k)$ time, Steps 10 and 15 can be implemented in $O(|V(B_1)| + |V(B_2)| + \dots + |V(B_x)|)$ time. Recall that all the vertices in B_i are deleted from G' except the cut vertex in each iteration of the repeat loop. This implies that $|V(B_1)| + |V(B_2)| + \dots + |V(B_x)| = n + (x - 1)$.

Notice that by using the depth first search algorithm, one can computer a vertex ordering u_1, u_2, \dots, u_h of a h -vertices tree graph T in $O(h)$ time such that u_i is a leaf in $T[\{u_1, u_2, \dots, u_i\}]$ for $1 \leq i \leq h$. Therefore, with an $O(x)$ -time preprocessing of G^* , it takes $O(1)$ time to implement Step 7 for each iteration of the repeat loop, i.e., determining v_B .

Meanwhile, since G has at most $n - 1$ blocks, i.e., $x \leq n - 1$, the repeat loop and Step 15 can be done, both in $O(n)$ time. Further, the other steps can be completed in $O(n)$ time as well. Consequently, we obtain the main result of this paper.

Theorem 2 *Given a weighted block graph G with the block-cut-vertex structure G^* of G , a paired-dominating set of G can be determined by Algorithm 1 in $O(n)$ time.*

3 Finding $D(H, u_1)$, $P(H, u_1)$, $P'(H, u_1)$, and $\bar{P}(H, u_1)$

Suppose that H is a weighted block graph and B is a block of H with $V(B) = \{u_1, u_2, \dots, u_k\}$. For $1 \leq i \leq k$, let G_i be a maximal connected subgraph in $(H - B) \cup \{u_i\}$. By Lemma 1, we have $V(G_i) \cap V(G_j) = \emptyset$ for $i \neq j$, refer to Figure 2 for an illustrative example. In this section, given the dominating sets $D(G_i, u_i)$, $P(G_i, u_i)$, $P'(G_i, u_i)$, and $\bar{P}(G_i, u_i)$ for $1 \leq i \leq k$, we shall show that the four dominating sets $D(H, u_1)$, $P(H, u_1)$, $P'(H, u_1)$, and $\bar{P}(H, u_1)$ can be determined in $O(k)$ time.

First, some notations are introduced below, for the purpose of describing the procedures. For a set S of sets of vertices, $F(S)$ denotes the set with minimum weight in S . Let S_i^* be the set of vertices such that $S_i^* = F(\{D(G_i, u_i), P(G_i, u_i), P'(G_i, u_i), \bar{P}(G_i, u_i)\})$ for $2 \leq i \leq k$. We use α to denote the index in $\{2, 3, \dots, k\}$ such that $S_\alpha^* \neq D(G_\alpha, u_\alpha)$ and $w(D(G_\alpha, u_\alpha)) - w(S_\alpha^*)$ is minimized, and β to denote the index in $\{2, 3, \dots, k\}$ such that $S_\beta^* = D(G_\beta, u_\beta)$ and $w(F(\{P(G_\beta, u_\beta), P'(G_\beta, u_\beta), \bar{P}(G_\beta, u_\beta)\})) - w(S_\beta^*)$ is minimized. Further, let r denote the number of S_i^* such that $S_i^* = D(G_i, u_i)$, i.e., $r = |\{S_i^* \mid S_i^* = D(G_i, u_i) \text{ and } 2 \leq i \leq k\}|$.

3.1 Determination of $D(H, u_1)$

We first recall that $D(H, u_1)$ is a minimum-weight dominating set of H containing u_1 , and $H[D(H, u_1) - u_1]$ has a perfect matching. By the definition of $D(H, u_1)$, the only potential candidate for being a dominating set of G_1 is $D(G_1, u_1)$. Hence, in order to obtain $D(H, u_1)$, we first construct a dominating set $X = D(G_1, u_1) \cup S_2^* \cup S_3^* \cup \dots \cup S_k^*$. We will show that if r is even, then $S = X$ is a κ_1 -dominating set of H with respect

to u_1 . Otherwise, for the purpose of satisfying the requirement that $H[X - u_1]$ has a perfect matching with minimum cost, we can either replace S_α^* with $D(G_\alpha, u_\alpha)$, or replace S_β^* with $F(\{P(G_\beta, u_\beta), P'(G_\beta, u_\beta), \bar{P}(G_\beta, u_\beta)\})$. For the former case, a dominating set $X^+ = (X - S_\alpha^*) \cup D(G_\alpha, u_\alpha)$ is created. On the other hand, a dominating set $X^- = (X - S_\beta^*) \cup F(\{P(G_\beta, u_\beta), P'(G_\beta, u_\beta), \bar{P}(G_\beta, u_\beta)\})$ is built for the latter case. The output $S = F(\{X^+, X^-\})$ is selected from X^+ and X^- based on the weights of the sets. Similarly, we will show that S is a κ_1 -dominating set of H with respect to u_1 . The following is a formal description of the procedure.

Procedure 2 Finding $D(H, u_1)$

Input: A weighted block graph H and a block B of H with $V(B) = \{u_1, u_2, \dots, u_k\}$.

Dominating sets $D(G_i, u_i)$, $P(G_i, u_i)$, $P'(G_i, u_i)$, and $\bar{P}(G_i, u_i)$ for $1 \leq i \leq k$.

Output: A κ_1 -paired-dominating set S of H with respect to u_1 .

- 1: determine S_i^* for $2 \leq i \leq k$;
 - 2: determine α, β , and r ;
 - 3: let $X \leftarrow D(G_1, u_1) \cup S_2^* \cup S_3^* \cup \dots \cup S_k^*$;
 - 4: let $X^+ \leftarrow (X - S_\alpha^*) \cup D(G_\alpha, u_\alpha)$;
 - 5: let $X^- \leftarrow (X - S_\beta^*) \cup F(\{P(G_\beta, u_\beta), P'(G_\beta, u_\beta), \bar{P}(G_\beta, u_\beta)\})$;
 - 6: **if** r is even, **then** let $S \leftarrow X$; **otherwise**, let $S \leftarrow F(\{X^+, X^-\})$;
 - 7: **return** S .
-

Lemma 3 *Given the dominating sets $D(G_i, u_i)$, $P(G_i, u_i)$, $P'(G_i, u_i)$, and $\bar{P}(G_i, u_i)$ for $1 \leq i \leq k$, Procedure 2 outputs a κ_1 -paired-dominating set S of H with respect to u_1 in $O(k)$ time.*

Proof. We first introduce data structures which enable us to compute $D(H, u_1)$ in $O(k)$ time. For each element of array *select*, with $2 \leq i \leq k$, *select*[i] is used to denote the selection of vertices set for S_i^* , i.e., *select*[i] = 1 if $S_i^* = D(G_i, u_i)$, \dots , *select*[i] = 4 if $S_i^* = \bar{P}(G_i, u_i)$. Further, variable s is used to denote the selection of vertices set for S , i.e., $s = 1$ if $S = X$, $s = 2$ if $S = X^+$, and $s = 3$ if $S = X^-$. Meanwhile, variable w is used to denote the weight of S . With the aid of above data structures and variables α, β , and r , the procedure certainly

can be implemented in $O(k)$ time. To prove that S is a κ_1 -paired-dominating set of H with respect to u_1 , it suffices to show that the output S is a minimum-weight dominating set of H such that $u_1 \in S$ and $H[S - u_1]$ has a perfect matching. By the definition of $D(H, u_1)$, the only potential candidate for being a dominating set of G_1 is $D(G_1, u_1)$. Hence, we have $D(G_1, u_1) \subseteq S$. Since $u_1 \in D(G_1, u_1)$ and B is a clique, all the three sets X, X^+ and X^- are dominating sets of H . Therefore, it remains to show that the weight $w(S)$ of S is minimized subject to the condition that $H[S - u_1]$ contains a perfect matching.

Notice that, for $2 \leq i \leq k$, both $G_i[D(G_i, u_i) - u_i]$ and $G_i[P(G_i, u_i)]$ contain perfect matchings and $u_i \notin P'(G_i, u_i) \cup \bar{P}(G_i, u_i)$. Therefore, to satisfy the condition that $H[S - u_1]$ contains a perfect matching, the cardinality of $\{u_i \mid D(G_i, u_i) \in S \text{ and } 2 \leq i \leq k\}$ must be even. Hence, if r is even, then $H[X - u_1]$ contains a perfect matching and the weight $w(X)$ of X is minimized, as an immediate consequence of the selections of S_i^* . Next, suppose that r is odd. To minimize the cost, we can replace one S_i^* with $D(G_i, u_i)$ or replace one S_j^* with $P(G_j, u_j), P'(G_j, u_j)$, or $\bar{P}(G_j, u_j)$, where $S_i^* \neq D(G_i, u_i)$, $S_j^* = D(G_j, u_j)$, and $2 \leq i, j \leq k$. For the former case, a dominating set $X^+ = (X - S_\alpha^*) \cup D(G_\alpha, u_\alpha)$ is created. On the other hand, a dominating set $X^- = (X - S_\beta^*) \cup F(\{P(G_\beta, u_\beta), P'(G_\beta, u_\beta), \bar{P}(G_\beta, u_\beta)\})$ is built for the latter case. And, we select S from X^+ and X^- based on the weights of the sets. As a consequence of selections of S_α^* , S_β^* , and S_i^* for $2 \leq i \leq k$, one can verify that $S = F(\{X^+, X^-\})$ is a minimum-weight dominating set of H such that $H[S - u_1]$ contains a perfect matching. \square

3.2 Determination of $P(H, u_1)$

Notice that $P(H, u_1)$ is a minimum-weight dominating set of H containing u_1 and $H[P(H, u_1)]$ has a perfect matching. Therefore, either $D(G_1, u_1) \subseteq P(H, u_1)$ or $P(G_1, u_1) \subseteq P(H, u_1)$ is a dominating set of G_1 . In order to obtain $P(H, u_1)$, we construct the six dominating sets X, X^+, X^-, Y, Y^+ , and Y^- of H . The dominating sets X, X^+ and X^- are created for the situation when $D(H, u_1)$ is a dominating set of G_1 . Meanwhile, the dominating sets Y, Y^+ and Y^- are built for the situation when $P(G_1, u_1)$ is a dominating set

of G_1 , where $Y = P(G_1, u_1) \cup S_2^* \cup S_3^* \cup \dots \cup S_k^*$, $Y^+ = (Y - S_\alpha^*) \cup D(G_\alpha, u_\alpha)$, and $Y^- = (Y - S_\beta^*) \cup F(\{P(G_\beta, u_\beta), P'(G_\beta, u_\beta), \bar{P}(G_\beta, u_\beta)\})$.

If r is even, then all the induced subgraphs $H[X^+]$, $H[X^-]$ and $H[Y]$ contain perfect matchings. The output $S = F(\{X^+, X^-, Y\})$ is selected from X^+ , X^- and Y based on the weights of the sets. We will show that S is a κ_2 -dominating set of H with respect to u_1 . Similarly, if r is odd, then all the induced subgraphs $H[X]$, $H[Y^+]$ and $H[Y^-]$ contain perfect matchings. And, we will show that the output $S = F(\{X, Y^+, Y^-\})$ is a κ_2 -dominating set of H with respect to u_1 in this situation. The procedure is detailed below.

Procedure 3 Finding $P(H, u_1)$

Input: A weighted block graph H and a block B of H with $V(B) = \{u_1, u_2, \dots, u_k\}$.

Dominating sets $D(G_i, u_i)$, $P(G_i, u_i)$, $P'(G_i, u_i)$, and $\bar{P}(G_i, u_i)$ for $1 \leq i \leq k$.

Output: A κ_2 -paired-dominating set S of H with respect to u_1 .

- 1: determine S_i^* for $2 \leq i \leq k$;
 - 2: determine α, β , and r ;
 - 3: find the dominating sets X, X^+ and X^- as described in Procedure 2;
 - 4: let $Y \leftarrow P(G_1, u_1) \cup S_2^* \cup S_3^* \cup \dots \cup S_k^*$;
 - 5: let $Y^+ \leftarrow (Y - S_\alpha^*) \cup D(G_\alpha, u_\alpha)$;
 - 6: let $Y^- \leftarrow (Y - S_\beta^*) \cup F(\{P(G_\beta, u_\beta), P'(G_\beta, u_\beta), \bar{P}(G_\beta, u_\beta)\})$;
 - 7: **if** r is even, **then** let $S \leftarrow F(\{X^+, X^-, Y\})$; **otherwise**, let $S \leftarrow F(\{X, Y^+, Y^-\})$;
 - 8: **return** S .
-

Lemma 4 *Given the dominating sets $D(G_i, u_i)$, $P(G_i, u_i)$, $P'(G_i, u_i)$, and $\bar{P}(G_i, u_i)$ for $1 \leq i \leq k$, Procedure 3 outputs a κ_2 -paired-dominating set S of H with respect to u_1 in $O(k)$ time.*

Proof. By using a similar method of the arguments in Lemma 3, one can show that the procedure can be completed in $O(k)$ time. To prove the correctness of the procedure, it suffices to show that the output S is a minimum-weight dominating set of H such that $u_1 \in S$ and $H[S]$ contains a perfect matching. Further, since $v_1 \in D(G_1, u_1) \cap P(G_1, u_1)$ and B is a clique, all X, X^+, X^-, Y, Y^+ , and Y^- are dominating sets of H . Thus, it remains to

show that the weight $w(S)$ of S is minimized subject to the condition that $H[S]$ contains a perfect matching.

Notice that, for $2 \leq i \leq k$, both $G_i[D(G_i, u_i) - u_i]$ and $G_i[P(G_i, u_i)]$ contain perfect matchings and $u_i \notin P'(G_i, u_i) \cup \bar{P}(G_i, u_i)$. We first consider the situation when r is even. For the case when $D(G_1, u_1)$ is a dominating set of G_1 , in order to satisfy the condition that $H[X]$ contains a perfect matching with minimum cost, we can either replace S_α^* with $D(G_\alpha, u_\alpha)$ or replace S_β^* with $F(\{P(G_\beta, u_\beta), P'(G_\beta, u_\beta), \bar{P}(G_\beta, u_\beta)\})$. Thus, X^+ and X^- are the two potential candidates for S when $D(G_1, u_1) \subseteq P(H, u_1)$. For the case when $P(G_1, u_1)$ is a dominating set of G_1 , $H[Y]$ contains a perfect matching. We select $S = F(\{X^+, X^-, Y\})$ from X^+ , X^- and Y based on the weights of the sets. As a consequence of selections of S_α^* , S_β^* , and S_i^* for $2 \leq i \leq k$, one can verify that the output S is a minimum-weight dominating set of H such that $H[S]$ contains a perfect matching. Using a similar method of the above arguments, one can show that the correctness also holds for the situation when r is odd. \square

3.3 Determination of $P'(H, u_1)$

Recall that $P'(H, u_1)$ is a minimum-weight dominating set of H not containing u_1 and $H[P'(H, u_1)]$ has a perfect matching. Therefore, either $P'(G_1, u_1) \subseteq P'(H, u_1)$ or $\bar{P}(G_1, u_1) \subseteq P'(H, u_1)$ is a dominating set of G_1 . For ease of subsequent discussion, we consider the two cases $P'(G_1, u_1) \subseteq P'(H, u_1)$ and $\bar{P}(G_1, u_1) \subseteq P'(H, u_1)$, respectively, in the rest of this subsection. More concretely, a paired-dominating set Q_1 is created for the former situation. Meanwhile, a paired-dominating set Q_2 is built for the latter situation. Clearly, $P'(H, u_1)$ can be selected from Q_1 and Q_2 based on the weights of the sets.

3.3.1 Finding Q_1

Below we present an $O(k)$ -time procedure for finding Q_1 . The procedure solves the problem by considering eight cases C_1, C_2, \dots, C_8 depending on S_i^* and r . For $1 \leq i \leq 8$, the case $C_i = (c_1, c_2, c_3, c_4, c_5)$ is an ordered 5-tuple. For $1 \leq j \leq 5$, $c_j = 1$ if condition D_j holds, and $c_j = 0$ otherwise. Further, $c_j = *$ means “do not care”, i.e., condition D_j is not a factor

in this case. The five conditions D_1, D_2, \dots, D_5 are defined as follows:

$$D_1 : S_i^* = P(G_i, u_i) \text{ for some } 2 \leq i \leq k.$$

$$D_2 : r \text{ is odd.}$$

$$D_3 : r \text{ is equal to 1.}$$

$$D_4 : r \text{ is equal to 0.}$$

$$D_5 : S_i^* = \bar{P}(G_i, u_i) \text{ for some } 2 \leq i \leq k.$$

Then, we define the cases $C_1 = (1, 1, *, *, *)$, $C_2 = (1, 0, *, *, *)$, $C_3 = (0, 1, 1, *, 1)$, $C_4 = (0, 1, 1, *, 0)$, $C_5 = (0, 1, 0, *, *)$, $C_6 = (0, 0, *, 1, 1)$, $C_7 = (0, 0, *, 1, 0)$, and $C_8 = (0, 0, *, 0, *)$. For example, case C_1 represents the situation when there exists an index ℓ such that $S_\ell^* = P(G_\ell, u_\ell)$ with $2 \leq \ell \leq k$ and r is an odd number. Further, case C_7 represents the situation when there exists no index ℓ such that $S_\ell^* = P(G_\ell, u_\ell)$, or $S_\ell^* = \bar{P}(G_\ell, u_\ell)$ and $r = 0$, i.e., $S_i^* = P'(G_i, u_i)$ for $2 \leq i \leq k$. Moreover, one can verify that all the possible combinations of the five conditions have been considered.

Next, some notations and paired-dominating sets are introduced. Let α' be the index in $\{2, 3, \dots, k\} - \{\alpha\}$ such that $S_{\alpha'}^* \neq D(G_{\alpha'}, u_{\alpha'})$ and $w(D(G_{\alpha'}, u_{\alpha'})) - w(S_{\alpha'}^*)$ is minimized. Let γ be the index in $\{2, 3, \dots, k\}$ such that $S_\gamma^* \neq P(G_\gamma, u_\gamma)$ and $w(P(G_\gamma, u_\gamma)) - w(S_\gamma^*)$ is minimized. Let $I = \{i \mid S_i^* = \bar{P}(G_i, u_i) \text{ and } 2 \leq i \leq k\}$. We define the following paired-dominating sets of H , which are the potential candidates for Q_1 .

$$Z_1 = P'(G_1, u_1) \cup S_2^* \cup S_3^* \cup \dots \cup S_k^*.$$

$$Z_1^+ = (Z_1 - S_\alpha^*) \cup D(G_\alpha, u_\alpha).$$

$$Z_1^- = (Z_1 - S_\beta^*) \cup F(\{P(G_\beta, u_\beta), P'(G_\beta, u_\beta), \bar{P}(G_\beta, u_\beta)\}).$$

$$T_1 = (Z_1 - S_\gamma^*) \cup P(G_\gamma, u_\gamma).$$

$$T_2 = (Z_1 - S_\alpha^* - S_{\alpha'}^*) \cup D(G_\alpha, u_\alpha) \cup D(G_{\alpha'}, u_{\alpha'}).$$

$$T_3 = (Z_1 - \cup_{i \in I} S_i^*) \cup (\cup_{i \in I} P'(G_i, u_i)).$$

$$T_4 = (Z_1 - S_\beta^*) \cup P(G_\beta, u_\beta).$$

$$T_5 = (Z_1 - S_\beta^*) \cup F(\{P(G_\beta, u_\beta), P'(G_\beta, u_\beta)\}).$$

$$\begin{aligned}
T_6 &= (Z_1 - S_\gamma^* - S_\beta^*) \cup P(G_\gamma, u_\gamma) \cup F(\{P'(G_\beta, u_\beta), \bar{P}(G_\beta, u_\beta)\}). \\
T_7 &= (Z_1 - S_\gamma^* - S_\beta^*) \cup P(G_\gamma, u_\gamma) \cup \bar{P}(G_\beta, u_\beta). \\
T_8 &= (Z_1 - \cup_{i \in I} S_i^* - S_\beta^*) \cup (\cup_{i \in I} P'(G_i, u_i)) \cup P'(G_\beta, u_\beta).
\end{aligned}$$

As mentioned earlier, we solve the problem by considering the eight cases C_1, C_2, \dots, C_8 . The relations between the cases C_1, C_2, \dots, C_8 and the dominating sets $Z_1, Z_1^+, Z_1^-, T_1, \dots, T_8$ are detailed in Procedure 4. We will prove its correctness and analyze its running time in Lemma 5.

Procedure 4 Finding Q_1

Input: A weighted block graph H and a block B of H with $V(B) = \{u_1, u_2, \dots, u_k\}$.

Dominating sets $D(G_i, u_i)$, $P(G_i, u_i)$, $P'(G_i, u_i)$, and $\bar{P}(G_i, u_i)$ for $1 \leq i \leq k$.

Output: A minimum-weight dominating set S of H such that $u_1 \notin S$, $H[S]$ has a perfect matching, and $P'(G_1, u_1) \subseteq S$.

- 1: determine the paired-dominating sets $Z_1, Z_1^+, Z_1^-, T_1, \dots, T_8$;
 - 2: **if** C_1 or C_5 holds, **then** let $S \leftarrow F(\{Z_1^+, Z_1^-\})$;
 - 3: **if** C_2 or C_7 or C_8 holds, **then** let $S \leftarrow Z_1$;
 - 4: **if** C_3 holds, **then** let $S \leftarrow F(\{Z_1^+, T_4, T_6, T_8\})$;
 - 5: **if** C_4 holds, **then** let $S \leftarrow F(\{Z_1^+, T_5, T_7\})$;
 - 6: **if** C_6 holds, **then** let $S \leftarrow F(\{T_1, T_2, T_3\})$;
 - 7: **return** S .
-

Lemma 5 *Given the dominating sets $D(G_i, u_i)$, $P(G_i, u_i)$, $P'(G_i, u_i)$, and $\bar{P}(G_i, u_i)$ for $1 \leq i \leq k$, Procedure 4 outputs a minimum-weight dominating set S of H such that $u_1 \notin S$, $H[S]$ has a perfect matching, and $P'(G_1, u_1) \subseteq S$. Moreover, the procedure can be completed in $O(k)$ time.*

Proof. By using a similar method of the arguments in Lemma 3, one can show that all the paired-dominating sets $Z_1, Z_1^+, Z_1^-, T_1, \dots, T_8$ can be constructed in $O(k)$ time. Hence, the procedure certainly runs in $O(k)$ time. Further, one can verify that all the possible combinations of conditions D_1, D_2, \dots, D_5 have been considered in cases C_1, C_2, \dots, C_8 . Hence, to

prove the correctness of the procedure, it suffices to show that each step of the procedure is correct.

First, we consider cases C_1 and C_2 . In both of these cases, there exists an index ℓ such that $S_\ell^* = P(G_\ell, u_\ell)$. Therefore, Z_1, Z_1^+ , and Z_1^- are dominating sets of H . It follows that, if r is even, then Z_1 is a minimum-weight dominating set of H such that $u_1 \notin Z_1$ and $H[Z_1]$ has a perfect matching due to the selections of S_i^* for $2 \leq i \leq k$. So, we have $S = Z_1$ for case C_2 . On the other hand, if r is odd, then in order to satisfy the condition that $H[Q_1]$ contains a perfect matching with minimum cost, we can either replace S_α^* with $D(G_\alpha, u_\alpha)$, or replace S_β^* with $F(\{P(G_\beta, u_\beta), P'(G_\beta, u_\beta), \bar{P}(G_\beta, u_\beta)\})$. This implies that we have $S = F(\{Z_1^+, Z_1^-\})$ for case C_1 .

Next, we consider cases C_3, C_4 , and C_5 . Notice that in all three cases, there exists no index ℓ such that $S_\ell^* = P(G_\ell, u_\ell)$ and r is an odd number. Moreover, for any paired-dominating set Q_1 of H , we have either $S_i^* = P'(G_i, u_i)$ for $2 \leq i \leq k$ or $V(B) \cap V(Q_1) \neq \emptyset$, where $B = H[\{u_1, u_2, \dots, u_k\}]$. In case C_3 , a paired-dominating set T_8 is created for the former. Meanwhile, paired-dominating sets Z_1^+, T_4 , and T_6 are built for the latter. As a consequence of $r = 1$, in order to ensure $H[Q_1]$ contains a perfect matching when $V(B) \cap V(Q_1) \neq \emptyset$, we replace S_α^* with $D(G_\alpha, u_\alpha)$ in Z_1^+ , replace S_β^* with $P(G_\beta, u_\beta)$ in T_4 , and replace S_γ^* and S_β^* with $P(G_\gamma, u_\gamma)$ and $F(\{P'(G_\beta, u_\beta), \bar{P}(G_\beta, u_\beta)\})$ in T_6 , respectively. Under the premise of minimizing weight, one can verify that Z_1^+, T_4 , and T_6 are exactly the three potential candidates for Q_1 . In case C_4 , we have $S_\beta^* = D(G_\beta, u_\beta)$ and $S_i^* = P'(G_i, u_i)$ for $2 \leq i \leq k$ and $i \neq \beta$. Using a similar method of the above arguments, one can show that $S = F(\{Z_1^+, T_5, T_7\})$ is true for case C_4 . In case C_5 , we have $r \geq 3$. Therefore, for the same reasons as case C_1 , we have $S = F(\{Z_1^+, Z_1^-\})$ for case C_5 .

Finally, we consider cases C_6, C_7 , and C_8 . Notice that in all three cases, there exists no index ℓ such that $S_\ell^* = P(G_\ell, u_\ell)$ and r is an even number. In case C_6 , we have either $S_i^* = P'(G_i, u_i)$ or $S_i^* = \bar{P}(G_i, u_i)$, where $1 \leq i \leq k$. To ensure $H[Q_1]$ contains a perfect matching, we replace S_γ^* with $P(G_\gamma, u_\gamma)$ in T_1 , replace S_α^* and $S_{\alpha'}^*$ with $D(G_\alpha, u_\alpha)$ and $D(G_{\alpha'}, u_{\alpha'})$ in T_2 , and replace S_i^* with $P'(G_i, u_i)$ for all $i \in I$ in T_3 , respectively. Under the

premise of minimizing the weight $w(S)$, one can verify that T_1 , T_2 , and T_3 are exactly the three potential candidates for Q_1 . Notice that, in case C_7 , $S_i^* = P'(G_i, u_i)$ for $2 \leq i \leq k$. Further, $r \geq 2$ is an even number in case C_8 . Thus, in both of these cases, we have $S = Z_1$ for the same reasons as case C_2 . \square

3.3.2 Finding Q_2

In the following, we present a procedure to find the paired-dominating set Q_2 . Similar to Procedure 4, the procedure solves the problem by considering six cases $C_9, C_{10}, \dots, C_{14}$. For $9 \leq i \leq 14$, the case $C_i = (c_1, c_2, c_3, c_4)$ is an ordered 4-tuple. Further, the value of c_k has the same definition as before for $1 \leq k \leq 4$. Then, we define $C_9 = (1, 1, *, *)$, $C_{10} = (1, 0, *, *)$, $C_{11} = (0, 1, 1, *)$, $C_{12} = (0, 1, 0, *)$, $C_{13} = (0, 0, *, 1)$, and $C_{14} = (0, 0, *, 0)$. Again, one can verify that all the possible combinations of the four conditions have been considered in cases $C_9, C_{10}, \dots, C_{14}$. The paired-dominating sets $Z_2, Z_2^+, Z_2^-, T_9, \dots, T_{12}$ of H are defined below, which are the potential candidates for Q_2 .

$$\begin{aligned}
Z_2 &= \bar{P}(G_1, u_1) \cup S_2^* \cup S_3^* \cup \dots \cup S_k^*. \\
Z_2^+ &= (Z_1 - S_\alpha^*) \cup D(G_\alpha, u_\alpha). \\
Z_2^- &= (Z_1 - S_\beta^*) \cup F(\{P(G_\beta, u_\beta), P'(G_\beta, u_\beta), \bar{P}(G_\beta, u_\beta)\}). \\
T_9 &= (Z_2 - S_\gamma^*) \cup P(G_\gamma, u_\gamma). \\
T_{10} &= (Z_2 - S_\alpha^* - S_{\alpha'}^*) \cup D(G_\alpha, u_\alpha) \cup D(G_{\alpha'}, u_{\alpha'}). \\
T_{11} &= (Z_2 - S_\beta^*) \cup P(G_\beta, u_\beta). \\
T_{12} &= (Z_2 - S_\gamma^* - S_\beta^*) \cup P(G_\gamma, u_\gamma) \cup F(\{P'(G_\beta, u_\beta), \bar{P}(G_\beta, u_\beta)\}).
\end{aligned}$$

Moreover, the relations between the cases $C_9, C_{10}, \dots, C_{14}$ and the paired-dominating sets $Z_2, Z_2^+, Z_2^-, T_9, \dots, T_{12}$ are detailed in Procedure 5.

Lemma 6 *Given the dominating sets $D(G_i, u_i)$, $P(G_i, u_i)$, $P'(G_i, u_i)$, and $\bar{P}(G_i, u_i)$ for $1 \leq i \leq k$, Procedure 5 outputs a minimum-weight dominating set S of H such that $u_1 \notin S$, $H[S]$ has a perfect matching, and $\bar{P}(G_1, u_1) \subseteq S$. Moreover, the procedure can be completed in $O(k)$ time.*

Procedure 5 Finding Q_2

Input: A weighted block graph H and a block B of H with $V(B) = \{u_1, u_2, \dots, u_k\}$.

Dominating sets $D(G_i, u_i)$, $P(G_i, u_i)$, $P'(G_i, u_i)$, and $\bar{P}(G_i, u_i)$ for $1 \leq i \leq k$.

Output: A minimum-weight dominating set S of H such that $u_1 \notin S$, $H[S]$ has a perfect matching, and $\bar{P}(G_1, u_1) \subseteq S$.

- 1: determine the paired-dominating sets $Z_2, Z_2^+, Z_2^-, T_9, \dots, T_{12}$;
 - 2: **if** C_9 or C_{12} holds, **then** let $S \leftarrow F(\{Z_2^+, Z_2^-\})$;
 - 3: **if** C_{10} or C_{14} holds, **then** let $S \leftarrow Z_2$;
 - 4: **if** C_{11} holds, **then** let $S \leftarrow F(\{Z_2^+, T_{11}, T_{12}\})$;
 - 5: **if** C_{13} holds, **then** let $S \leftarrow F(\{T_9, T_{10}\})$;
 - 6: **return** S .
-

Proof. By using a similar method of the arguments in Lemma 3, one can show that each step of the procedure can be completed in $O(k)$ time. Therefore, the procedure runs in $O(k)$ time. Further, one can verify that all the possible combinations of conditions D_1, D_2, D_3 , and D_4 have been considered in cases $C_9, C_{10}, \dots, C_{14}$. Hence, to prove the correctness of the procedure, it suffices to show that each step of the procedure is correct.

First, we consider cases C_9 and C_{10} . In both of these cases, there exists an index ℓ such that $S_\ell^* = P(G_\ell, u_\ell)$. Therefore, for the same reasons as cases C_1 and C_2 in Procedure 4, we have $S = F(\{Z_2^+, Z_2^-\})$ for case C_9 and $S = Z_2$ for case C_{10} , respectively. Next, we consider cases C_{11} , and C_{12} . Notice that in both cases, there exists no index ℓ such that $S_\ell^* = P(G_\ell, u_\ell)$ and r is an odd number. Since we have $r = 1$ in case C_{11} , in order to satisfy the condition that $H[Q_2]$ contains a perfect matching with minimum cost, we replace S_α^* with $D(G_\alpha, u_\alpha)$ in Z_2^+ , replace S_β^* with $P(G_\beta, u_\beta)$ in T_{11} , and replace S_γ^* and S_β^* with $P(G_\gamma, u_\gamma)$ and $F(\{P'(G_\beta, u_\beta), \bar{P}(G_\beta, u_\beta)\})$ in T_{12} , respectively. Under the premise of minimizing weight, one can verify that Z_2^+, T_{11} , and T_{12} are exactly the three potential candidates for Q_2 . In case C_{12} , we have $r \geq 3$. Therefore, for the same reasons as case C_9 , we have $S = F(\{Z_2^+, Z_2^-\})$ for case C_{12} .

Finally, we consider cases C_{13} , and C_{14} . Notice that in both cases, there exists no index ℓ such that $S_\ell^* = P(G_\ell, u_\ell)$ and r is an even number. In case C_{13} , either $S_i^* = P'(G_i, u_i)$

or $S_i^* = \bar{P}(G_i, u_i)$ for $1 \leq i \leq k$. Therefore, to satisfy the condition that $H[Q_2]$ contains a perfect matching, we replace S_γ^* with $P(G_\gamma, u_\gamma)$ in T_9 , and replace S_α^* and $S_{\alpha'}^*$ with $D(G_\alpha, u_\alpha)$ and $D(G_{\alpha'}, u_{\alpha'})$ in T_{10} , respectively. Again, under the premise of minimizing the weight, one can verify that T_9 and T_{10} are exactly the two potential candidates for Q_2 . Notice that, in case C_{14} , $r \geq 2$ is an even number. Thus, we have $S = Z_2$ for the same reasons as case C_2 in Procedure 4. \square

Combining Lemmas 5 and 6, we obtain the following result.

Lemma 7 *Given the dominating sets $D(G_i, u_i)$, $P(G_i, u_i)$, $P'(G_i, u_i)$, and $\bar{P}(G_i, u_i)$ for $1 \leq i \leq k$, a κ_3 -paired-dominating set $P'(H, u_1)$ can be determined in $O(k)$ time.*

3.4 Determination of $\bar{P}(H, u_1)$

Remember that $\bar{P}(H, u_1)$ is a minimum-weight dominating set of $H - u_1$ and u_1 is not dominated by $\bar{P}(H, u_1)$. Hence, by the definition of $\bar{P}(H, u_1)$, the only composition is $\bar{P}(H, u_1) = \bar{P}(G_1, u_1) \cup P'(G_2, u_2) \cup \dots \cup P'(G_k, u_k)$. This implies that, given the dominating sets $D(G_i, u_i)$, $P(G_i, u_i)$, $P'(G_i, u_i)$, and $\bar{P}(G_i, u_i)$ for $1 \leq i \leq k$, a κ_4 -paired-dominating set $\bar{P}(H, u_1)$ can be determined in $O(k)$ time. Thus, we have the following result.

Lemma 8 *Given the dominating sets $D(G_i, u_i)$, $P(G_i, u_i)$, $P'(G_i, u_i)$, and $\bar{P}(G_i, u_i)$ for $1 \leq i \leq k$, a κ_4 -paired-dominating set $\bar{P}(H, u_1)$ can be determined in $O(k)$ time.*

4 Conclusion and Future Work

In this paper, we have presented an optimal algorithm for finding a paired-dominating set of a weighted block graph G . The algorithm uses dynamic programming to iteratively determine $D(H, u)$, $P(H, u)$, $P'(H, u)$, and $\bar{P}(H, u)$ in a bottom-up manner, where H is a subgraph of G and $u \in V(H)$ is a cut vertex of G . When the graph is given in an adjacency list representation, our algorithm runs in $O(n + m)$ time. Moreover, the algorithm can be completed in $O(n)$ time if the block-cut-vertex structure of G is given.

Below we present some open problems related to the paired-domination problem. It is known that distance-hereditary graphs is a proper superfamily of block graphs. Therefore, it is interesting to study the time complexity of paired-domination problem in distance-hereditary graphs. In [3], Chen *et al.* proposed an approximation algorithm with ratio $\ln(2\Delta(G)) + 1$ for general graphs and showed that the problem is APX-complete, i.e., has no PTAS. Thus, it would be useful if we could develop an approximation algorithm for general graphs with constant ratio. Meanwhile, it would be desirable to show that the problem remains NP-complete in planar graphs and design an approximation algorithm.

References

- [1] A. V. Aho, J. E. Hopcroft, and J. D. Ullman. *The design and analysis of computer algorithms*. Addison-Wesley Publishing Co., Reading, Mass.-London-Amsterdam, 1974.
- [2] G. J. Chang. Algorithmic aspects of domination in graphs. In *Handbook of Combinatorial Optimization*, pages 339–405. Springer-Verlag, New York, second edition, 2013.
- [3] L. Chen, C. Lu, and Z. Zeng. Hardness results and approximation algorithms for (weighted) paired-domination graphs. *Theoret. Comput. Sci.*, 410(47-49):5063–5071, 2009.
- [4] L. Chen, C. Lu, and Z. Zeng. A linear-time algorithm for paired-domination problem in strongly chordal graphs. *Inform. Process. Lett.*, 110(1):20–23, 2009.
- [5] L. Chen, C. Lu, and Z. Zeng. Labelling algorithms for paired-domination problems in block and interval graphs. *J. Comb. Optim.*, 19(4):457–470, 2010.
- [6] T. C. E. Cheng, L. Kang, and E. Shan. A polynomial-time algorithm for the paired-domination problem on permutation graphs. *Discrete Appl. Math.*, 157(2):262–271, 2009.
- [7] W. Goddard and M. A. Henning. Independent domination in graphs: a survey and recent results. *Discrete Math.*, 313(7):839–854, 2013.
- [8] T. Haynes and P. Slater. Paired-domination in graphs. *Networks*, 32:199–206, 1998.
- [9] T. W. Haynes, S. T. Hedetniemi, and P. J. Slater. *Domination in Graphs: Advanced Topics*. Marcel Dekker Inc., New York, 1998.
- [10] T. W. Haynes, S. T. Hedetniemi, and P. J. Slater. *Fundamentals of Domination in Graphs*. Marcel Dekker Inc., New York, 1998.
- [11] S. T. Hedetniemi and R. C. Laskar. Bibliography on domination in graphs and some basic definitions of domination parameters. *Discrete Math.*, 86(1-3):257–277, 1990.
- [12] S. T. Hedetniemi and R. C. Laskar, editors. *Topics on domination*. Annals of Discrete Mathematics. North-Holland Publishing Co., Amsterdam, 1991.

- [13] M. A. Henning. A survey of selected recent results on total domination in graphs. *Discrete Math.*, 309(1):32–63, 2009.
- [14] R.-W. Hung. Linear-time algorithm for the paired-domination problem in convex bipartite graphs. *Theory Comput. Syst.*, 50(4):721–738, 2012.
- [15] L. Kang. Variations of dominating set problem. In *Handbook of Combinatorial Optimization*, pages 3363–3394. Springer-Verlag, New York, second edition, 2013.
- [16] L. Kang, M. Y. Sohn, and T. C. E. Cheng. Paired-domination in inflated graphs. *Theor. Comput. Sci.*, 320(2-3):485–494, 2004.
- [17] E. Lappas, S. D. Nikolopoulos, and L. Palios. An $O(n)$ -time algorithm for the paired-domination problem on permutation graphs. In *Combinatorial algorithms*, volume 5874 of *Lecture Notes in Comput. Sci.*, pages 368–379. Springer, Berlin, 2009.
- [18] E. Lappas, S. D. Nikolopoulos, and L. Palios. An $O(n)$ -time algorithm for the paired domination problem on permutation graphs. *European J. Combin.*, 34(3):593–608, 2013.
- [19] C.-C. Lin and H.-L. Tu. A linear-time algorithm for paired-domination on circular-arc graphs. *Theoret. Comput. Sci.*, 591:99–105, 2015.
- [20] B. S. Panda and D. Pradhan. A linear time algorithm for computing a minimum paired-dominating set of a convex bipartite graph. *Discrete Appl. Math.*, 161(12):1776–1783, 2013.
- [21] B. S. Panda and D. Pradhan. Minimum paired-dominating set in chordal bipartite graphs and perfect elimination bipartite graphs. *J. Comb. Optim.*, 26(4):770–785, 2013.
- [22] H. Qiao, L. Kang, M. Cardei, and D.-Z. Du. Paired-domination of trees. *J. of Global Optimization*, 25(1):43–54, 2003.
- [23] C.-C. Yen and R. C. T. Lee. The weighted perfect domination problem and its variants. *Discrete Appl. Math.*, 66(2):147–160, 1996.