

On the Approximation of Laplacian Eigenvalues in Graph Disaggregation

Xiaozhe Hu^a, John C. Urschel^{b,c,d*}, Ludmil T. Zikatanov^{d,e}

^a*Department of Mathematics, Tufts University, Medford, MA, USA* ^b*Baltimore Ravens, NFL, Owings Mills, MD, USA*; ^c*Department of Mathematics, Massachusetts Institute of Technology, Cambridge, MA, USA*. ^d*Department of Mathematics, Penn State University, University Park, PA, USA*; ^e*Institute for Mathematics and Informatics, Bulgarian Academy of Sciences, Sofia, Bulgaria*

(Received 00 Month 20XX; final version received 00 Month 20XX)

Graph disaggregation is a technique used to address the high cost of computation for power law graphs on parallel processors. The few high-degree vertices are broken into multiple small-degree vertices, in order to allow for more efficient computation in parallel. In particular, we consider computations involving the graph Laplacian, which has significant applications, including diffusion mapping and graph partitioning, among others. We prove results regarding the spectral approximation of the Laplacian of the original graph by the Laplacian of the disaggregated graph. In addition, we construct an alternate disaggregation operator whose eigenvalues interlace those of the original Laplacian. Using this alternate operator, we construct a uniform preconditioner for the original graph Laplacian.

Keywords: Spectral Graph Theory; Graph Laplacian; Disaggregation; Spectral Approximation; Preconditioning

AMS Subject Classification: 05C85; 65F15; 65F08; 68R10

1. Introduction

A variety of real-world graphs, including web networks [1], social networks [14], and bioinformatics networks [9], exhibit a degree power law. Namely, the fraction of nodes of degree k , denoted by $P(k)$, follows a power distribution of the form $P(k) \sim k^{-\gamma}$, where γ is typically in the range $2 < \gamma < 3$. Networks of this variety are often referred to as scale-free networks. The pairing of a few high-degree vertices with many low-degree vertices on large scale-free networks makes computations such as Laplacian matrix-vector products and solving linear and eigenvalue equations challenging. The computation of the minimal nontrivial eigenpair can become prohibitively expensive. This eigenpair has many important applications, such as diffusion mapping and graph partitioning [2, 12, 15, 16].

Breaking the few high degree nodes into multiple smaller degree nodes is a way to address this issue, especially when large-scale parallel computers are available. This technique, called graph disaggregation, was introduced by Kuhleemann and

*Corresponding author. Email: urschel@mit.edu




Figure 1.1. Example of disagggregation: original graph (left); disagggregate using cycle (middle); disagggregate using clique (right).

Vassilevski [6, 10]. In this process, each of the high-degree vertices of the network is replaced by a graph, such as a cycle or a clique, where each incident edge of the original node now connects to a node of the cycle or clique (see Figure 1.1).

Independently, Lee, Peng, and Spielman investigated the concept of graph disagggregation, referred to as vertex splitting, in the setting of combinatorial spectral sparsifiers [11]. They proved results for graphs disagggregated from complete graphs and expanders, and used the Schur complement of the disagggregated Laplacian with respect to the disagggregated vertices to approximate the original Laplacian. The basic motivating assumption in such constructions is that the spectral structure of the graph Laplacian induced by the disagggregated graph approximates the spectral structure of the original graph well.

In [6, 10] Kuhleemann and Vassilevski took a numerical approach. We extend, expand upon, and prove precise and rigorous theoretical results regarding this technique. First, we look at the case of a single disagggregated vertex and establish bounds on the error in spectral approximation with respect to the Laplacians of the original and disagggregated graph, as well as results related to the Cheeger constant. We investigate a conjecture made in [10] and give strong theoretical evidence that it does not hold in general. Then, we treat the more general case of disagggregation of multiple vertices and prove analogous results. Finally, we construct an alternative disagggregation operator whose eigenvalues interlace with those of the original graph Laplacian, and, hence, provide excellent approximation to the spectrum of the latter. We then use this new disagggregation operator to construct a uniform preconditioner for the graph Laplacian of the original graph. We prove that the preconditioned graph Laplacian can be made arbitrarily close to the identity operator if we require that the weights of the internal disagggregated edges are sufficiently large.

2. Single Vertex Disagggregation

Consider a weighted, connected, undirected graph $G = (V, E, \omega)$, $|V| = n$. Let $e = (i, j)$ denote an edge that connects vertices i and j , and $\langle \cdot, \cdot \rangle$ and $\| \cdot \|$ denote the standard ℓ^2 -inner product and the corresponding induced norm. The associated weighted graph Laplacian $A \in \mathbb{R}^{n \times n}$ is given by

$$\langle A\mathbf{u}, \mathbf{v} \rangle = \sum_{e=(i,j) \in E} \omega_e(u_i - u_j)(v_i - v_j), \quad \omega_e = (-a_{ij}),$$

where we denote the (i, j) -th element of A by a_{ij} . Without loss of generality, let us disaggregate the last vertex \mathbf{v}_n of the graph \mathbf{G} . Then, the Laplacian can be written in the following block form

$$A = \begin{pmatrix} A_0 & -\mathbf{a}_n \\ -\mathbf{a}_n^T & a_{nn} \end{pmatrix},$$

where a_{nn} is the degree of \mathbf{v}_n . Here, we assume that the graph is simply connected and the associated Laplacian A has eigenvalues

$$0 = \lambda_1(A) < \lambda_2(A) \leq \cdots \leq \lambda_n(A)$$

and corresponding eigenvectors

$$\mathbf{1}_n = \boldsymbol{\varphi}^{(1)}(A), \boldsymbol{\varphi}^{(2)}(A), \dots, \boldsymbol{\varphi}^{(n)}(A), \quad \text{where } \mathbf{1}_n = \underbrace{(1, \dots, 1)}_n^T.$$

The eigenpair $(\lambda_2(A), \boldsymbol{\varphi}^{(2)}(A))$ has special significance, and therefore $\lambda_2(A)$ is referred to as the *algebraic connectivity*, denoted $a(\mathbf{G})$, and $\boldsymbol{\varphi}^{(2)}(A)$ is referred to as the *Fiedler vector*.

We can also write a given nontrivial eigenpair $(\lambda(A), \boldsymbol{\varphi}(A))$, $\lambda(A) \neq 0$, $\|\boldsymbol{\varphi}(A)\| = 1$, in block notation, namely

$$\boldsymbol{\varphi}(A) = \begin{pmatrix} \boldsymbol{\varphi}_0 \\ \varphi_n \end{pmatrix}.$$

We have the relations

$$\begin{aligned} \langle \boldsymbol{\varphi}_0, \mathbf{1}_{n_0} \rangle + \varphi_n &= 0, \\ A_0 \boldsymbol{\varphi}_0 - \varphi_n \mathbf{a}_n &= \lambda(A) \boldsymbol{\varphi}_0, \\ a_{nn} \varphi_n - \mathbf{a}_n^T \boldsymbol{\varphi}_0 &= \lambda(A) \varphi_n, \end{aligned}$$

where $n_0 = n - 1$. Suppose that the vertex \mathbf{v}_n is disaggregated into d vertices, with an unspecified connected structure between the disaggregated elements. We will denote this graph by \mathbf{G}_D . This induces a disaggregated graph Laplacian $A_D \in \mathbb{R}^{N \times N}$, $N = n_0 + d$, with eigenvalues $0 = \lambda_1(A_D) < \lambda_2(A_D) \leq \cdots \leq \lambda_N(A_D)$ and corresponding eigenvectors $\mathbf{1}_N = \boldsymbol{\varphi}^{(1)}(A_D), \boldsymbol{\varphi}^{(2)}(A_D), \dots, \boldsymbol{\varphi}^{(N)}(A_D)$. We can write A_D in block form

$$A_D = \begin{pmatrix} A_0 & -A_{0n} \\ -A_{0n}^T & A_n \end{pmatrix}.$$

We have the relations

$$\begin{aligned} a_{nn} &= \mathbf{a}_n^T \mathbf{1}_{n_0}, \\ A_{0n}^T \mathbf{1}_{n_0} &= A_n \mathbf{1}_d, \\ \mathbf{a}_n &= A_0 \mathbf{1}_{n_0} = A_{0n} \mathbf{1}_d. \end{aligned}$$

Let us introduce the prolongation operator $P : \mathbb{R}^n \rightarrow \mathbb{R}^N$,

$$P = \begin{pmatrix} I_{n_0 \times n_0} & 0 \\ 0 & \mathbf{1}_d \end{pmatrix}. \quad (2.1)$$

The following result is immediate.

LEMMA 1 *Let A and A_D be the graph Laplacian of the original graph G and the disaggregated and simply connected graph G_D , respectively. If P is defined as (2.1), then we have*

$$A = P^T A_D P.$$

We aim to show that the algebraic connectivity of A_D is bounded away from the algebraic connectivity of the original graph A . To do so, suppose we have an eigenpair (λ, φ) of the Laplacian of the original graph G . We prolongate the eigenvector φ to the disaggregated graph G_D and obtain an approximate eigenvector by the procedure

$$\tilde{\varphi} = P\varphi - s\mathbf{1}_N = \begin{pmatrix} \varphi_0 \\ \varphi_n \mathbf{1}_d \end{pmatrix} - s\mathbf{1}_N, \quad \text{where } s = \frac{d-1}{N} \varphi_n. \quad (2.2)$$

This gives $\langle \tilde{\varphi}, \mathbf{1}_N \rangle = 0$.

We consider $\tilde{\varphi}$ to be an approximation of φ on the non-trivial eigenspace of the disaggregated operator A_D . We have the following relation between the eigenvalue λ of A and the Rayleigh quotient of $\tilde{\varphi}$ with respect to A_D .

LEMMA 2 *Let (λ, φ) , $\|\varphi\| = 1$, be an eigenpair of the graph Laplacian A associated with a simply connected graph G , and $\tilde{\varphi}$ be defined by (2.2). We have*

$$\text{RQ}(\tilde{\varphi}) := \frac{\langle A_D \tilde{\varphi}, \tilde{\varphi} \rangle}{\langle \tilde{\varphi}, \tilde{\varphi} \rangle} = \frac{\lambda}{1 + \frac{(d-1)n}{N} \varphi_n^2}.$$

Proof. We have

$$\begin{aligned} \langle \tilde{\varphi}, \tilde{\varphi} \rangle &= \langle P\varphi - s\mathbf{1}_N, P\varphi - s\mathbf{1}_N \rangle = \langle P\varphi, P\varphi \rangle - 2s\langle P\varphi, \mathbf{1}_N \rangle + s^2\langle \mathbf{1}_N, \mathbf{1}_N \rangle \\ &= \langle \varphi_0, \varphi_0 \rangle + \varphi_n^2 \langle \mathbf{1}_d, \mathbf{1}_d \rangle - 2s(\langle \varphi_0, \mathbf{1}_{n_0} \rangle + \varphi_n \langle \mathbf{1}_d, \mathbf{1}_d \rangle) + s^2 N \\ &= \langle \varphi_0, \varphi_0 \rangle + \varphi_n^2 + (d-1)\varphi_n^2 - 2s(d-1)\varphi_n + s^2 N \\ &= 1 + \left[(d-1) - 2\frac{(d-1)^2}{N} + \frac{(d-1)^2}{N} \right] \varphi_n^2 \\ &= 1 + \frac{(d-1)n}{N} \varphi_n^2 \end{aligned}$$

and

$$\begin{aligned} \langle A_D \tilde{\varphi}, \tilde{\varphi} \rangle &= \langle A_D(P\varphi - s\mathbf{1}_N), P\varphi - s\mathbf{1}_N \rangle \\ &= \langle A_D P\varphi, P\varphi \rangle - 2s\langle A_D \mathbf{1}_N, P\varphi \rangle + s^2 \langle A_D \mathbf{1}_N, \mathbf{1}_N \rangle \\ &= \langle P^T A_D P\varphi, \varphi \rangle = \langle A\varphi, \varphi \rangle = \lambda. \end{aligned}$$

This completes the proof. \square

The following result quickly follows by applying Lemma 2 to the Fiedler vector.

THEOREM 1 *Let $\varphi = (\varphi_0, \varphi_n)^T$, $\|\varphi\| = 1$, be the Fiedler vector of the graph Laplacian A associated with a simply connected graph G . Let A_D be the graph Laplacian corresponding to the disaggregated and simply connected graph G_D resulting from disaggregating one vertex into $d > 1$ vertices. We have*

$$\frac{a(G)}{a(G_D)} \geq 1 + \frac{(d-1)n}{N} \varphi_n^2.$$

Proof. Noting that $\tilde{\varphi}$ is orthogonal to $\mathbf{1}_N$, we have

$$a(G_D) \leq \frac{\langle A_D \tilde{\varphi}, \tilde{\varphi} \rangle}{\langle \tilde{\varphi}, \tilde{\varphi} \rangle} = \frac{\lambda}{1 + \frac{(d-1)n}{N} \varphi_n^2} = \frac{a(G)}{1 + \frac{(d-1)n}{N} \varphi_n^2},$$

which completes the proof. \square

If the characteristic value of the disaggregated vertex is non-zero, then the algebraic connectivity of the disaggregated graph stays bounded away from that of the original graph, independent of the structure of A_n . Therefore, as the weight on the internal edges approaches infinity, the approximation stays bounded away.

In [10], the authors made the following conjecture.

CONJECTURE 1 *Under certain conditions the Laplacian eigenvalues of the graph Laplacian of the disaggregated graph approximate the eigenvalues of the graph Laplacian of the original graph, provided that the weight on the internal edges of the disaggregation is chosen to be large enough.*

Theorem 1 directly implies that Conjecture 1 is false when the characteristic value of the disaggregated vertex is non-zero, which, for a random power law graph, occurs with probability one.

We also have the following result, providing an estimate of how close the approximation $\tilde{\varphi}$ is to the invariant subspace with respect to A_D .

LEMMA 3 *Let A and A_D be the original graph and disaggregated graph, (λ, φ) be an eigenpair of the graph Laplacian A associated with a simply connected graph G , and $\tilde{\varphi}$ be defined by (2.2). We have*

$$\|A_D \tilde{\varphi} - \text{RQ}(\tilde{\varphi}) \tilde{\varphi}\| \leq \left(\|A_{0n}^T (\mathbf{1}_{n_0} - \varphi_0/\varphi_n)\| + \frac{\sqrt{dn(d+n)}}{N} \lambda + \frac{dn}{N} \lambda |\varphi_n| \right) |\varphi_n|.$$

Proof. We recall that

$$\|\tilde{\varphi}\| = \left(1 + \frac{(d-1)n}{N} \varphi_n^2 \right)^{1/2}$$

and

$$|\lambda - \text{RQ}(\tilde{\varphi})| = \frac{\frac{(d-1)n}{N}\varphi_n^2}{1 + \frac{(d-1)n}{N}\varphi_n^2}\lambda.$$

We also have

$$\begin{aligned} A_D \tilde{\varphi} &= A_D (P\varphi - s\mathbf{1}_N) = A_D P\varphi = \begin{pmatrix} A_0\varphi_0 - \varphi_n A_{0n}\mathbf{1}_d \\ \varphi_n A_n\mathbf{1}_d - A_{0n}^T\varphi_0 \end{pmatrix} = \begin{pmatrix} A_0\varphi_0 - \varphi_n \mathbf{a}_n \\ A_{0n}^T(\varphi_n \mathbf{1}_{n_0} - \varphi_0) \end{pmatrix} \\ &= \lambda P\varphi + \begin{pmatrix} \mathbf{0} \\ A_{0n}^T(\varphi_n \mathbf{1}_{n_0} - \varphi_0) - \lambda \varphi_n \mathbf{1}_d \end{pmatrix} = \lambda \tilde{\varphi} + s\lambda \mathbf{1}_N + \begin{pmatrix} \mathbf{0} \\ A_{0n}^T(\varphi_n \mathbf{1}_{n_0} - \varphi_0) - \lambda \varphi_n \mathbf{1}_d \end{pmatrix} \\ &= \lambda \tilde{\varphi} + \frac{\lambda \varphi_n}{N} \left[(d-1) \begin{pmatrix} \mathbf{1}_0 \\ \mathbf{0} \end{pmatrix} - n \begin{pmatrix} \mathbf{0} \\ \mathbf{1}_d \end{pmatrix} \right] + \begin{pmatrix} \mathbf{0} \\ A_{0n}^T(\varphi_n \mathbf{1}_{n_0} - \varphi_0) \end{pmatrix}, \end{aligned}$$

giving

$$\begin{aligned} \|A_D \tilde{\varphi} - \text{RQ}(\tilde{\varphi})\tilde{\varphi}\| &\leq \|A_D \tilde{\varphi} - \lambda \tilde{\varphi}\| + |\lambda - \text{RQ}(\tilde{\varphi})| \|\tilde{\varphi}\| \\ &\leq \|A_{0n}^T(\varphi_n \mathbf{1}_{n_0} - \varphi_0)\| + \frac{\sqrt{(d-1)^2 n_0 + dn^2}}{N} |\varphi_n| \lambda + \frac{\frac{(d-1)n}{N}\varphi_n^2}{\sqrt{1 + \frac{(d-1)n}{N}\varphi_n^2}} \varphi_n^2 \lambda \\ &\leq \left(\|A_{0n}^T(\mathbf{1}_{n_0} - \varphi_0/\varphi_n)\| + \frac{\sqrt{dn(d+n)}}{N} \lambda + \frac{dn}{N} \lambda |\varphi_n| \right) |\varphi_n|. \end{aligned}$$

□

In many applications, we are only concerned with minimal Laplacian eigenpairs. For minimal eigenvalues of scale-free graphs, we have $\lambda = O(1)$ and $\varphi_n = O(N^{-1/2})$. In this way, often the largest source of error comes from the term $\|A_{0n}^T(\varphi_n \mathbf{1}_{n_0} - \varphi_0)\|$. Heuristically, the error of this term is typically best controlled when d is relatively small and each new disaggregate is connected to roughly the same number of exterior vertices.

Next, we consider the eigenvalues of the normalized Laplacian. This gives us insight into how the Cheeger constant changes after disaggregating a vertex. Again, suppose we have an eigenpair (ν, ϕ) of the normalized graph Laplacian $D^{-1}A$, where D is the degree matrix of G , namely $D = \text{diag}(a_{11}, a_{22}, \dots, a_{nn})$. Again, we prolongate the eigenvector ϕ and obtain an approximate eigenvector of the disaggregated normalized graph Laplacian $D_D^{-1}A_D$, where D_D is the degree matrix of G_D , in a similar fashion

$$\tilde{\phi} = P\phi - s\mathbf{1}_N, \quad \text{where } s = \frac{\langle P\phi, \mathbf{1}_N \rangle_{D_D}}{\langle \mathbf{1}_N, \mathbf{1}_N \rangle_{D_D}}. \quad (2.3)$$

We may write D_D in the following way

$$\begin{aligned} D_D &= \text{diag}(a_{1,1}^D, \dots, a_{n_0,n_0}^D, a_{n,n}^D, a_{n+1,n+1}^D, \dots, a_{N,N}^D) \\ &= \text{diag}(a_{1,1}^D, \dots, a_{n_0,n_0}^D, \omega_n, \omega_{n+1}, \dots, \omega_N) + \text{diag}(0, \dots, 0, d_n^{ex}, d_{n+1}^{ex}, \dots, d_N^{ex}) \\ &=: D_D^1 + D_D^{ex}, \end{aligned}$$

where $\omega_n, \omega_{n+1}, \dots, \omega_N$ are the weights of the edges incident with vertex v_n on the original graph (note $\sum_{i=n}^N \omega_i = a_{n,n}$) and $d_i^{ex} = a_{i,i}^D - \omega_i$, $i = n, n+1, \dots, N$. We may also rewrite the shift s as

$$s = \frac{\langle P\phi, \mathbf{1}_N \rangle_{D_D}}{\langle \mathbf{1}_N, \mathbf{1}_N \rangle_{D_D}} = \frac{\sum_{i=n}^N d_i^{ex} \phi_n}{\sum_{i=1}^N a_{i,i}^D}.$$

Let $\omega_{\text{total}}(\mathbf{H})$ denote the total weights of a graph \mathbf{H} , and let \mathbf{G}_a be the disaggregated local subgraph. Similarly, we consider $\tilde{\phi}$ as an approximation of the eigenvectors of the disaggregated normalized graph Laplacian. We have the following lemma.

THEOREM 2 *Let (ν, ϕ) be an eigenpair of the normalized graph Laplacian associated with a simply connected graph \mathbf{G} and $\tilde{\phi}$ be defined by (2.3). We have*

$$\frac{\langle A_D \tilde{\phi}, \tilde{\phi} \rangle}{\langle D_D \tilde{\phi}, \tilde{\phi} \rangle} = \frac{\nu}{1 + \frac{2\omega_{\text{total}}(\mathbf{G}) \omega_{\text{total}}(\mathbf{G}_a)}{\omega_{\text{total}}(\mathbf{G}_D)} \phi_n^2}, \quad (2.4)$$

and

$$\nu_2^D = \alpha \nu_2, \quad \alpha = \left(1 + \frac{2\omega_{\text{total}}(\mathbf{G}) \omega_{\text{total}}(\mathbf{G}_a)}{\omega_{\text{total}}(\mathbf{G}_D)} \phi_n^2 \right)^{-1} \leq 1. \quad (2.5)$$

Proof. We have

$$\langle A_D \tilde{\phi}, \tilde{\phi} \rangle = \langle A_D(P\phi - s\mathbf{1}_N), P\phi - s\mathbf{1}_N \rangle = \langle A_D P\phi, \phi \rangle = \langle A\phi, \phi \rangle = \nu$$

and

$$\begin{aligned} \langle D_D \phi, \phi \rangle &= \langle D_D P\phi, P\phi \rangle - s \langle D_D \mathbf{1}_N, P\phi \rangle \\ &= \langle D_D^1 P\phi, P\phi \rangle + \langle D_D^{ex} P\phi, P\phi \rangle - s \langle D_D \mathbf{1}_N, P\phi \rangle \\ &= \langle D\phi, \phi \rangle + \sum_{i=n}^N d_i^{ex} \phi_n^2 - \frac{\left(\sum_{i=n}^N d_i^{ex} \phi_n \right)^2}{\sum_i a_{i,i}^D} \\ &= 1 + \frac{\left(\sum_i a_{i,i}^D - \sum_{i=n}^N d_i^{ex} \right) \sum_{i=n}^N d_i^{ex}}{\sum_i a_{i,i}^D} \phi_n^2 \\ &= 1 + \frac{\left(\sum_{i=1}^n a_{i,i} \right) \left(\sum_{i=n}^N d_i^{ex} \right)}{\sum_{i=1}^N a_{i,i}^D} \phi_n^2. \end{aligned}$$

Noting that $\sum_{i=1}^N a_{i,i} = 2\omega_{\text{total}}(\mathbf{G})$, $\sum_{i=1}^N a_{i,i}^D = 2\omega_{\text{total}}(\mathbf{G}_D)$, and $\sum_{i=n}^N d_i^{ex} = 2\omega_{\text{total}}(\mathbf{G}_a)$, we obtain (2.4). Moreover, (2.5) follows directly from (2.4). \square

The Cheeger constant of a weighted graph is defined as follows [4, 8]

$$h(\mathbf{G}) = \min_{\emptyset \neq U \subset V} \frac{|E(U, \bar{U})|}{\min(\text{vol}(U), \text{vol}(\bar{U}))},$$

where

$$\bar{U} = V \setminus U, \quad \text{vol}(U) = \sum_{i \in U} \delta_i, \quad \delta_i = \sum_{e=(i,j) \in E} \omega_e, \quad |E(U, \bar{U})| = \sum_{\substack{e=(i,j) \in E \\ i \in U, j \in \bar{U}}} \omega_e.$$

Due to the Cheeger inequality [4, 8], the Cheeger constant $h(\mathbf{G})$ and ν_2 are related as follows

$$1 - \sqrt{1 - h(\mathbf{G})^2} \leq \nu_2 \leq 2h(\mathbf{G}). \quad (2.6)$$

THEOREM 3 *For the Cheeger constant of the original graph \mathbf{G} and the disaggregated and simply connected graph \mathbf{G}_D , we have*

$$h(\mathbf{G}_D) \leq \sqrt{1 - (1 - 2\alpha h(\mathbf{G}))^2},$$

where α is defined by (2.5). If $h(\mathbf{G}) \geq \frac{4\alpha}{4\alpha^2+1}$, then $h(\mathbf{G}_D) \leq h(\mathbf{G})$.

Proof. Based on (2.6) and (2.5), we have

$$h(\mathbf{G}_D) \leq \sqrt{1 - (1 - \nu_2^D)^2} = \sqrt{1 - (1 - \alpha\nu_2)^2} \leq \sqrt{1 - (1 - 2\alpha h(\mathbf{G}))^2}.$$

Basic algebra shows that $h(\mathbf{G}_D) \leq h(\mathbf{G})$ if $h(\mathbf{G}) \geq \frac{4\alpha}{4\alpha^2+1}$. □

3. Graph Disaggregation

We now move on to the more general case of multiple disaggregated vertices. Without loss of generality, for a graph Laplacian $A \in \mathbb{R}^{n \times n}$, suppose we are disaggregating the first m vertices. This gives us the disaggregated Laplacian $A_D \in \mathbb{R}^{N \times N}$. Here, N is the number of vertices in the disaggregated graph, given by

$$N = n - m + \sum_{k=1}^m d_k = n - m + n_d, \quad n_d = \sum_{k=1}^m d_k.$$

Note that we have m groups of vertices associated with the disaggregation, which we can also number consecutively

$$\{1, \dots, N\} = \{\underbrace{1, \dots, d_1}_{d_1}, \dots, \underbrace{d_1 + 1, \dots, d_1 + d_2}_{d_2}, \dots, n_d + 1, \dots, N\}. \quad (3.1)$$

Similar to the case of a single disaggregated vertex, we can establish a relationship between A and A_D through a prolongation matrix $P: \mathbb{R}^n \rightarrow \mathbb{R}^N$, given by

$$P = \begin{pmatrix} P_m & 0 \\ 0 & I_{n_0 \times n_0} \end{pmatrix}, \quad \text{where } n_0 = (n - m). \quad (3.2)$$

Here, $P_m \in \mathbb{R}^{n_d \times m}$, and

$$P_m = \begin{pmatrix} \mathbf{1}_{d_1} & 0 & \dots & 0 \\ 0 & \mathbf{1}_{d_2} & \dots & 0 \\ \vdots & \vdots & \ddots & \vdots \\ 0 & 0 & \dots & \mathbf{1}_{d_m} \end{pmatrix}.$$

Note that $P_m^T P_m = \text{diag}(d_1, \dots, d_m)$. We have the following lemma, which can be easily verified by simple algebraic calculation.

LEMMA 4 *Let A and A_D be the graph Laplacian of the original graph G and the disaggregated and simply connected graph G_D , respectively. If P is defined as (3.2), then we have*

$$A = P^T A_D P. \quad (3.3)$$

If we look at the disaggregated graph Laplacian A_D directly, we can obtain a similar bound on the algebraic connectivity as shown in Theorem 1. Let (λ, φ) be an eigenpair of A . We can define an approximated eigenvector of A_D by prolongating φ as follows

$$\tilde{\varphi} = P\varphi - s\mathbf{1}_N, \quad \text{where } s = \frac{1}{N} \sum_{i=1}^m (d_i - 1)\varphi_i. \quad (3.4)$$

It is easy to check that $\langle \tilde{\varphi}, \mathbf{1}_N \rangle = 0$. Now we have the following lemma about the Rayleigh quotient of $\tilde{\varphi}$ with respect to A_D .

LEMMA 5 *Let (λ, φ) be an eigenpair of the graph Laplacian A associated with a simply connected graph G and $\tilde{\varphi}$ be defined by (3.4). We have*

$$\text{RQ}(\tilde{\varphi}) := \frac{\langle A_D \tilde{\varphi}, \tilde{\varphi} \rangle}{\langle \tilde{\varphi}, \tilde{\varphi} \rangle} \leq \frac{\lambda}{1 + \frac{1}{N} \sum_{i=1}^m (d_i - 1)(n + n_d - md_i)\varphi_i^2}.$$

Moreover, if for $d_i^{\max} = \max_i d_i$, we have $n + n_d - md_i^{\max} > 0$, then $\text{RQ}(\tilde{\varphi}) < \lambda$.

Proof. We note that

$$\begin{aligned} \langle A_D \tilde{\varphi}, \tilde{\varphi} \rangle &= \langle A_D(P\varphi - s\mathbf{1}_N), P\varphi - s\mathbf{1}_N \rangle = \langle A_D P\varphi, P\varphi \rangle - 2s\langle A_D \mathbf{1}_N, P\varphi \rangle + s^2\langle A_D \mathbf{1}_N, \mathbf{1}_N \rangle \\ &= \langle P^T A_D P\varphi, \varphi \rangle = \langle A\varphi, \varphi \rangle = \lambda. \end{aligned}$$

Denoting φ by $\varphi = (\varphi_1, \varphi_2, \dots, \varphi_m, \varphi_0)^T$, we have

$$\begin{aligned}
\langle \tilde{\varphi}, \tilde{\varphi} \rangle &= \langle P\varphi - s\mathbf{1}_N, P\varphi - s\mathbf{1}_N \rangle = \langle P\varphi, P\varphi \rangle - 2s\langle P\varphi, \mathbf{1}_N \rangle + s^2\langle \mathbf{1}_N, \mathbf{1}_N \rangle \\
&= \langle \varphi_0, \varphi_0 \rangle + \sum_{i=1}^m \varphi_i^2 \langle \mathbf{1}_{d_i}, \mathbf{1}_{d_i} \rangle - 2s \left(\langle \varphi_0, \mathbf{1}_{n_0} \rangle + \sum_{i=1}^m \varphi_i \langle \mathbf{1}_{d_i}, \mathbf{1}_{d_i} \rangle \right) + s^2 N \\
&= \langle \varphi_0, \varphi_0 \rangle + \sum_{i=1}^m \varphi_i^2 + \sum_{i=1}^m (d_i - 1) \varphi_i^2 - 2s \sum_{i=1}^m (d_i - 1) \varphi_i + s^2 N \\
&= 1 + \sum_{i=1}^m (d_i - 1) \varphi_i^2 - \frac{1}{N} \left(\sum_{i=1}^m (d_i - 1) \varphi_i \right)^2 \\
&\geq 1 + \sum_{i=1}^m (d_i - 1) \varphi_i^2 - \frac{m}{N} \sum_{i=1}^m (d_i - 1)^2 \varphi_i^2 \\
&= 1 + \sum_{i=1}^m (d_i - 1) \frac{N - m(d_i - 1)}{N} \varphi_i^2 \\
&= 1 + \frac{1}{N} \sum_{i=1}^m (d_i - 1)(n + n_d - md_i) \varphi_i^2.
\end{aligned}$$

This completes the proof. \square

From the Rayleigh quotient and applying the above lemma to the Fiedler vector, we have the following theorem concerning the algebraic connectivity.

THEOREM 4 *Let φ be the Fiedler vector of the graph Laplacian A associated with a simply connected graph G and A_D be the graph Laplacian corresponding to the disaggregated and simply connected graph G_D . Suppose we have disaggregated m vertices and each of those vertices are disaggregated into $d_i > 1$ vertices, $i = 1, 2, \dots, m$. We have*

$$\frac{a(G)}{a(G_D)} \geq 1 + \frac{1}{N} \sum_{i=1}^m (d_i - 1)(n + n_d - md_i) \varphi_i^2.$$

Proof. The proof is similar to the proof of Theorem 1 and uses Lemma 5. \square

It is possible to perform a more careful estimate, using the fact that disaggregating m vertices at once is equivalent to disaggregating m vertices one by one. Denote $n_0 = n$ and $n_i = n_{i-1} - 1 + d_i$, $i = 1, 2, \dots, m$. Note that $n_i = n - i + \sum_{k=1}^i d_k$, $i = 1, 2, \dots, m$ and $N = n_m$. Recursively applying Theorem 1, we have the following result.

THEOREM 5 *Let φ be the Fiedler vector of the graph Laplacian A associated with a simply connected graph G and A_D be the graph Laplacian corresponding to the disaggregated and simply connected graph G_D , respectively. Suppose we disaggregated m vertices and each of those are disaggregated into $d_i > 1$ vertices, $i = 1, 2, \dots, m$. We have*

$$\frac{a(G)}{a(G_D)} \geq \prod \left(1 + \frac{(d_i - 1)n_{i-1}}{n_i} \varphi_i^2 \right).$$

Proof. Let G_D^i be the resulting graph after disaggregating the i -th vertex in the graph G_D^{i-1} and note that $G_D = G_D^m$. We have

$$\frac{a(G)}{a(G_D)} = \frac{a(G)}{a(G_D^1)} \times \frac{a(G_D^1)}{a(G_D^2)} \times \cdots \times \frac{a(G_D^{m-1})}{a(G_D^m)}.$$

The result follows immediately from applying Theorem 1 on each pair of graphs G_D^i and G_D^{i+1} . \square

REMARK 1 *A direct consequence of Theorem 5 is $a(G_D) \leq a(G)$.*

Similarly, we also have the following result concerning the minimal eigenvalue of the normalized graph Laplacian after disaggregating several vertices. Denote $G_D^0 = G$, and denote the graph after disaggregating vertex i by G_D^i . Note that $G_D^m = G_D$. The local subgraph corresponding to disaggregating vertex i is denoted by G_a^i .

THEOREM 6 *Let ν_2 be the second smallest eigenvalue of the normalized graph Laplacian associated with a simply connected graph G and ν_2^D be the second smallest eigenvalue of the normalized graph Laplacian corresponding to the disaggregated and simply connected graph G_D . Suppose we disaggregated m vertices and each of them are disaggregated into $d_i > 1$ vertices, $i = 1, 2, \dots, m$. We have*

$$\frac{\nu_2}{\nu_2^D} \geq \prod_{i=1}^m \left(1 + \frac{2\omega_{total}(G_D^{i-1})\omega_{total}(G_a^i)}{\omega_{total}(G_D^i)} \phi_i^2 \right).$$

Consequently, we have $\nu_2^D = \alpha \nu_2$, where

$$\alpha := \left[\prod_{i=1}^m \left(1 + \frac{2\omega_{total}(G_D^{i-1})\omega_{total}(G_a^i)}{\omega_{total}(G_D^i)} \phi_i^2 \right) \right]^{-1} \leq 1. \quad (3.5)$$

Proof. The result follows by applying Theorem 2 recursively. \square

Based on the estimates on the eigenvalues of normalized graph Laplacian, we can estimate the Cheeger constants as follows.

THEOREM 7 *For the Cheeger constant of the original graph G and the disaggregated and simply connected graph G_D , we have*

$$h(G_D) \leq \sqrt{1 - (1 - 2\alpha h(G))^2},$$

where α is defined by (3.5). If $h(G) \geq \frac{4\alpha}{4\alpha^2+1}$, then $h(G_D) \leq h(G)$.

Proof. The proof is the same as the proof of Theorem 3. \square

4. Preconditioning Using Disaggregated Graph

We aim to show eigenvalue interlacing between A and a new operator, which is obtained by scaling A_D appropriately.

We can rescale P by introducing $\tilde{P} = D_s P$, where

$$D_s = \text{diag}(d_1^{-1/2} I_{d_1 \times d_1}, \dots, d_m^{-1/2} I_{d_m \times d_m}, I_{n_0 \times n_0}) \quad (4.1)$$

is a diagonal scaling matrix, giving us $\tilde{P}^T \tilde{P} = I$. Based on the scaled prolongation, we are able to show the eigenvalues of diagonal scaled matrix

$$\tilde{A}_D := D_s^{-1} A_D D_s^{-1} \quad (4.2)$$

interlaces with A . First, let us recall the interlacing theorem.

THEOREM 8 (Interlacing Theorem [5], Vol. 1, Chap. I) *Let $S \in \mathbb{R}^{N \times n}$ be such that $S^T S = I_{n \times n}$, $n < N$ and let $B \in \mathbb{R}^{N \times N}$ be symmetric, with eigenvalues $\lambda_1 \leq \lambda_2 \leq \dots \leq \lambda_N$. Define $A = S^T B S$ and let A have eigenvalues $\mu_1 \leq \mu_2 \leq \dots \leq \mu_m$. Then $\lambda_i \leq \mu_i \leq \lambda_{N-m+i}$.*

From here, we have the following.

THEOREM 9 *Let A have eigenvalues $\lambda_1(A) \leq \lambda_2(A) \leq \dots \leq \lambda_N(A)$ and $\tilde{A}_D = D_s^{-1} A_D D_s^{-1}$ have eigenvalues $\lambda_1(\tilde{A}_D) \leq \lambda_2(\tilde{A}_D) \leq \dots \leq \lambda_N(\tilde{A}_D)$. Then*

$$\lambda_i(\tilde{A}_D) \leq \lambda_i(A) \leq \lambda_{N-n+i}(\tilde{A}_D).$$

Proof. From the above Lemma, we have

$$A = P^T A_D P = P^T D_s D_s^{-1} A_D D_s^{-1} D_s P = \tilde{P}^T D_s^{-1} A_D D_s^{-1} \tilde{P} = \tilde{P}^T \tilde{A}_D \tilde{P}.$$

As $\tilde{P}^T \tilde{P} = I_{n \times n}$, by the Interlacing Theorem 8, the eigenvalues of A and \tilde{A}_D interlace. \square

We now discuss how to use the disaggregated graph G_D to solve the graph Laplacian on the original graph G . Here, we will use \tilde{A}_D as the auxiliary problem and design a preconditioner based on the Fictitious Space Lemma [13] and auxiliary space framework [18]. Because A and \tilde{A}_D are both symmetric positive semi-definite, we first state the refined version of the Fictitious Space Lemma proposed in [7].

THEOREM 10 (Theorem 6.3 and 6.4 in [7]) *Let \tilde{V} and V be two Hilbert spaces and $\Pi : \tilde{V} \mapsto V$ be a surjective map. Suppose that $\tilde{\mathcal{A}} : \tilde{V} \mapsto \tilde{V}'$ and $\mathcal{A} : V \mapsto V'$ are symmetric semi-definite operators. Moreover, suppose*

$$\Pi(N(\tilde{\mathcal{A}})) = N(\mathcal{A}), \quad (4.3)$$

$$\|\Pi \tilde{v}\|_A \leq c_1 \|\tilde{v}\|_{\tilde{\mathcal{A}}}, \quad \forall \tilde{v} \in \tilde{V}, \quad (4.4)$$

$$\text{for any } v \in V \text{ there exists } \tilde{v} \in \tilde{V} \text{ such that } \Pi \tilde{v} = v \text{ and } \|\tilde{v}\|_{\tilde{\mathcal{A}}} \leq c_0 \|v\|_A, \quad (4.5)$$

then for any symmetric positive definite operator $\tilde{\mathcal{B}} : \tilde{V}' \mapsto \tilde{V}$, we have that for $\mathcal{B} = \Pi \tilde{\mathcal{B}} \Pi^T$,

$$\kappa(\mathcal{B}\mathcal{A}) \leq \left(\frac{c_1}{c_0} \right)^2 \kappa(\tilde{\mathcal{B}}\tilde{\mathcal{A}}).$$

Applying the above theory to our disaggregation framework, we take $\mathcal{A} = A$, $\tilde{\mathcal{A}} = \tilde{A}_D$, and $\Pi = \tilde{P}^T$. Noting that the null space of \tilde{A}_D is spanned by $D_s \mathbf{1}_N$, we have

$$\tilde{P}^T D_s \mathbf{1}_N = P^T D_s^2 \mathbf{1}_N = \mathbf{1}_n$$

which verifies (4.3). Naturally, \tilde{P}^T is surjective. Using a preconditioner \tilde{B}_D of \tilde{A}_D , we can define a preconditioner

$$B = \tilde{P}^T \tilde{B}_D \tilde{P}$$

for A . We give the following results concerning the quality of the preconditioner B .

COROLLARY 1 *Let A be the graph Laplacian corresponding to the graph \mathbf{G} and A_D be the graph Laplacian corresponding to the disaggregated and simply connected graph \mathbf{G}_D . Let D_s be defined by (4.1) and \tilde{A}_D be defined by (4.2). If*

$$\|\tilde{P}^T \tilde{\mathbf{v}}\|_A \leq c_1 \|\tilde{\mathbf{v}}\|_{\tilde{A}_D}, \quad \forall \tilde{\mathbf{v}} \in \tilde{V} \quad (4.6)$$

and for any $\mathbf{v} \in V$, there exist a $\tilde{\mathbf{v}} \in \tilde{V}$ such that $\tilde{P}^T \tilde{\mathbf{v}} = \mathbf{v}$ and

$$\|\tilde{\mathbf{v}}\|_{\tilde{A}_D} \leq c_0 \|\mathbf{v}\|_A. \quad (4.7)$$

Then for the preconditioner $B = \tilde{P}^T \tilde{B}_D \tilde{P}$, we have

$$\kappa(BA) \leq \left(\frac{c_1}{c_0} \right)^2 \kappa(\tilde{B}_D \tilde{A}_D).$$

We need to verify that conditions (4.6) and (4.7) hold for $\tilde{P} = D_s P$. For condition (4.7), we choose $\tilde{\mathbf{v}} = \tilde{P}^T \mathbf{v}$ for any $\mathbf{v} \in V$, giving $\tilde{P}^T \tilde{\mathbf{v}} = \tilde{P}^T \tilde{P} \mathbf{v} = \mathbf{v}$ since $\tilde{P}^T \tilde{P} = I$. Note that

$$\|\tilde{\mathbf{v}}\|_{\tilde{A}_D}^2 = \langle \tilde{A}_D \tilde{P} \mathbf{v}, \tilde{P} \mathbf{v} \rangle = \|\mathbf{v}\|_A^2, \quad (4.8)$$

which implies condition (4.7) holds with $c_0 = 1$.

To show that condition (4.6) holds, we use the following result.

LEMMA 6 *Let $A \in \mathbb{R}^{n \times n}$ be a graph Laplacian corresponding to a connected graph with n vertices. For all $i \in \{1, \dots, n\}$ and $\mathbf{u} \in \mathbb{R}^n$ we have*

$$\frac{1}{n}(\mathbf{u}, \mathbf{1}_n) - (\mathbf{u}, \mathbf{e}_i) = \frac{1}{n}(A_i^{-1} \mathbf{1}_n, A\mathbf{u}),$$

where $A_i = (A + \mathbf{e}_i \mathbf{e}_i^T)$.

Proof. First, we note that for all $i \in \{1, \dots, n\}$ the matrices A_i are invertible, because they all are irreducibly diagonally dominant M -matrices. We refer to Varga [17] for this classical result.

Next, observe that $A_i \mathbf{1}_n = \mathbf{e}_i$, and hence, $A_i^{-1} \mathbf{e}_i = \mathbf{1}_n$. Therefore, we have that

$$\begin{aligned} (A_i^{-1} \mathbf{1}_n, A\mathbf{u}) &= (A_i^{-1} \mathbf{1}_n, (A_i - \mathbf{e}_i \mathbf{e}_i^T) \mathbf{u}) = (\mathbf{1}_n, \mathbf{u}) - (\mathbf{u}, \mathbf{e}_i)(A_i^{-1} \mathbf{1}_n, \mathbf{e}_i) \\ &= (\mathbf{1}_n, \mathbf{u}) - (\mathbf{u}, \mathbf{e}_i)(\mathbf{1}_n, A_i^{-1} \mathbf{e}_i) = (\mathbf{1}_n, \mathbf{u}) - (\mathbf{u}, \mathbf{e}_i)(\mathbf{1}_n, \mathbf{1}_n). \end{aligned}$$

As $(\mathbf{1}_n, \mathbf{1}_n) = n$, this completes the proof. \square

The result shown in Lemma 6 is also found in [3, Lemma 3.2], but is included for completeness.

We now apply Lemma 6 to each disaggregated local subgraph $G_a^k = (V_a^k, E_a^k, \omega_a^k)$, $k = 1, 2, \dots, m$, with $\mathbf{u} = \tilde{\mathbf{v}}_k$, the restriction of $\tilde{\mathbf{v}}$ on G_a^k . For $j \in V_a^k$, we have,

$$\tilde{v}_j = \frac{1}{d_k} \sum_{p \in V_a^k} \tilde{v}_p - \frac{1}{d_k} \langle L_{k,j}^{-1} \mathbf{1}_{d_k}, L_k \tilde{\mathbf{v}}_k \rangle, \quad (4.9)$$

where L_k is the unweighted graph Laplacian of the local graph G_a^k and $L_{k,j}$ is defined in accordance with Lemma 6: $L_{k,j} = L_k + \mathbf{e}_j^k (\mathbf{e}_j^k)^T$, for $j \in V_a^k$. Setting $W_k^j := \frac{1}{d_k^2} \|L_{k,j}^{-1} \mathbf{1}_{d_k}\|_{L_k}^2$, $j \in V_a^k$, and denoting

$$\begin{aligned} E_D^0 &:= \{e = (i, j) \in E_D, i, j \in V^0\}, \\ E_D^1 &:= \{e = (i, j) \in E_D, i \in V^0, j \in V_a^k, k = 1, 2, \dots, m\}, \\ E_D^2 &:= \{e = (i, j) \in E_D, i \in V_a^k, j \in V_a^\ell, k, \ell = 1, 2, \dots, m, k \neq \ell\}, \end{aligned}$$

we are ready to present the following lemma related to the condition (4.6).

LEMMA 7 *For each disaggregated local subgraph G_a^k , if, for an edge $e' = (p, q) \in E_a^k$, we assign a weight $\omega_{e'}$ such that*

$$\omega_{e'} \geq W_{e'} := (1 + \epsilon^{-1}) \left[\sum_{\substack{e=(i,j) \in E_D^1 \\ i \in V^0, j \in V_a^k}} \omega_e W_k^j + 2 \sum_{\ell=1}^m \left(\sum_{\substack{e=(i,j) \in E_D^2 \\ i \in V_a^k, j \in V_a^\ell}} \omega_e W_k^i + \sum_{\substack{e=(i,j) \in E_D^2 \\ i \in V_a^\ell, j \in V_a^k}} \omega_e W_k^j \right) \right], \quad (4.10)$$

then we have

$$\|\tilde{P}^T \tilde{\mathbf{v}}\|_A^2 \leq (1 + \epsilon) \|\tilde{\mathbf{v}}\|_{\tilde{A}_D}^2, \quad \forall \tilde{\mathbf{v}} \in \tilde{V}, \quad (4.11)$$

where $\epsilon > 0$.

Proof. We denote $\tilde{\mathbf{u}} = \tilde{P}^T \tilde{P} \tilde{\mathbf{v}}$, and we have,

$$\begin{aligned}
 \|\tilde{P}^T \tilde{\mathbf{v}}\|_A^2 &= \langle \tilde{A}_D \tilde{\mathbf{u}}, \tilde{\mathbf{u}} \rangle = \sum_{e=(i,j) \in E_D} \omega_e (\tilde{u}_i - \tilde{u}_j)^2 \\
 &= \sum_{e=(i,j) \in E_D^0} \omega_e (\tilde{u}_i - \tilde{u}_j)^2 + \sum_{k=1}^m \sum_{e=(i,j) \in E_a^k} \omega_e (\tilde{u}_i - \tilde{u}_j)^2 \\
 &\quad + \sum_{e=(i,j) \in E_D^1} \omega_e (\tilde{u}_i - \tilde{u}_j)^2 + \sum_{e=(i,j) \in E_D^2} \omega_e (\tilde{u}_i - \tilde{u}_j)^2 \\
 &=: I_0 + I_1 + I_2.
 \end{aligned}$$

Here, we have set $I_0 = \sum_{e=(i,j) \in E_D^0} \omega_e (\tilde{v}_i - \tilde{v}_j)^2$,

$$I_1 = \sum_{k=1}^m \sum_{\substack{e=(i,j) \in E_D^1 \\ i \in V^0, j \in V_a^k}} \omega_e \left(\tilde{v}_i - \frac{1}{d_k} \sum_{p \in V_a^k} \tilde{v}_p \right)^2,$$

and

$$I_2 = \sum_{k=1}^m \sum_{\ell=1}^m \sum_{\substack{e=(i,j) \in E_D^2 \\ i \in V_a^k, j \in V_a^\ell}} \omega_e \left(\frac{1}{d_k} \sum_{p \in V_a^k} \tilde{v}_p - \frac{1}{d_\ell} \sum_{q \in V_a^\ell} \tilde{v}_q \right)^2.$$

Next, we estimate I_1 and I_2 on the right-hand side. For $e = (i, j) \in E_D^1$, $i \in V^0$ and $j \in V_a^k$, using (4.9), we have

$$\begin{aligned}
 \left(\tilde{v}_i - \frac{1}{d_k} \sum_{p \in V_a^k} \tilde{v}_p \right)^2 &= \left(\tilde{v}_i - \tilde{v}_j - \frac{1}{d_k} \langle L_{k,j}^{-1} \mathbf{1}_{d_k}, L_k \tilde{\mathbf{v}}_k \rangle \right)^2 \\
 &\leq (1 + \epsilon) (\tilde{v}_i - \tilde{v}_j)^2 + (1 + \epsilon^{-1}) \frac{1}{d_k^2} \|L_{k,j}^{-1} \mathbf{1}_{d_k}\|_{L_k}^2 \|\tilde{\mathbf{v}}_k\|_{L_k}^2 \\
 &= (1 + \epsilon) (\tilde{v}_i - \tilde{v}_j)^2 + \sum_{e'=(p,q) \in E_a^k} \left[(1 + \epsilon^{-1}) W_k^j \right] (\tilde{v}_p - \tilde{v}_q)^2.
 \end{aligned}$$

Then

$$\begin{aligned}
 I_1 &\leq \sum_{k=1}^m \sum_{\substack{e=(i,j) \in E_D^1 \\ i \in V^0, j \in V_a^k}} \omega_e \left\{ (1+\epsilon) (\tilde{v}_i - \tilde{v}_j)^2 + \sum_{e'=(p,q) \in E_a^k} \left[(1+\epsilon^{-1}) W_k^j \right] (\tilde{v}_p - \tilde{v}_q)^2 \right\} \\
 &= (1+\epsilon) \sum_{k=1}^m \sum_{\substack{e=(i,j) \in E_D^1 \\ i \in V^0, j \in V_a^k}} \omega_e (\tilde{v}_i - \tilde{v}_j)^2 \\
 &\quad + \sum_{k=1}^m \sum_{e'=(p,q) \in E_a^k} \left[(1+\epsilon^{-1}) \sum_{\substack{e=(i,j) \in E_D^1 \\ i \in V^0, j \in V_a^k}} \omega_e W_k^j \right] (\tilde{v}_p - \tilde{v}_q)^2.
 \end{aligned}$$

Next, using (4.9), for $e = (i, j) \in E_D^2$, $i \in V_a^k$ and $j \in V_a^\ell$ we have

$$\begin{aligned}
 \left(\frac{1}{d_k} \sum_{p \in V_a^k} \tilde{v}_p - \frac{1}{d_\ell} \sum_{q \in V_a^\ell} \tilde{v}_q \right)^2 &= \left(\tilde{v}_i - \tilde{v}_j + \frac{1}{d_k} \langle L_{k,i}^{-1} \mathbf{1}_{d_k}, L_k \tilde{\mathbf{v}}_k \rangle - \frac{1}{d_\ell} \langle L_{\ell,j}^{-1} \mathbf{1}_{d_\ell}, L_\ell \tilde{\mathbf{v}}_\ell \rangle \right)^2 \\
 &\leq (1+\epsilon) (\tilde{v}_i - \tilde{v}_j)^2 + 2(1+\epsilon^{-1}) \frac{1}{d_k^2} \|L_{k,i}^{-1} \mathbf{1}_{d_k}\|_{L_k}^2 \|\tilde{\mathbf{v}}_k\|_{L_k}^2 \\
 &\quad + 2(1+\epsilon^{-1}) \frac{1}{d_\ell^2} \|L_{\ell,j}^{-1} \mathbf{1}_{d_\ell}\|_{L_\ell}^2 \|\tilde{\mathbf{v}}_\ell\|_{L_\ell}^2 \\
 &= (1+\epsilon) (\tilde{v}_i - \tilde{v}_j)^2 + \sum_{e'=(p,q) \in E_a^k} [2(1+\epsilon^{-1}) W_k^i] (\tilde{v}_p - \tilde{v}_q)^2 \\
 &\quad + \sum_{e'=(p,q) \in E_a^\ell} [2(1+\epsilon^{-1}) W_\ell^j] (\tilde{v}_p - \tilde{v}_q)^2.
 \end{aligned}$$

Then

$$\begin{aligned}
 I_2 &\leq \sum_{k=1}^m \sum_{\ell=1}^m \sum_{\substack{e=(i,j) \in E_D^2 \\ i \in V_a^k, j \in V_a^\ell}} \omega_e \left\{ (1+\epsilon) (\tilde{v}_i - \tilde{v}_j)^2 + \sum_{e'=(p,q) \in E_a^k} [2(1+\epsilon^{-1}) W_k^i] (\tilde{v}_p - \tilde{v}_q)^2 \right. \\
 &\quad \left. + \sum_{e'=(p,q) \in E_a^\ell} [2(1+\epsilon^{-1}) W_\ell^j] (\tilde{v}_p - \tilde{v}_q)^2 \right\}.
 \end{aligned}$$

Therefore, we have

$$\begin{aligned}
 I_2 \leq & (1 + \epsilon) \sum_{k=1}^m \sum_{\ell=1}^m \sum_{\substack{e=(i,j) \in E_D^2 \\ i \in V_a^k, j \in V_a^\ell}} \omega_e (\tilde{v}_i - \tilde{v}_j)^2 \\
 & + \sum_{k=1}^m \sum_{e'=(p,q) \in E_a^k} \left[\sum_{l=1}^m \sum_{\substack{e=(i,j) \in E_D^2 \\ i \in V_a^k, j \in V_a^\ell}} 2(1 + \epsilon^{-1}) \omega_e W_k^i (\tilde{v}_p - \tilde{v}_q)^2 \right. \\
 & \left. + \sum_{\ell=1}^m \sum_{e'=(p,q) \in E_a^\ell} \left[\sum_{k=1}^m \sum_{\substack{e=(i,j) \in E_D^2 \\ i \in V_a^k, j \in V_a^\ell}} 2(1 + \epsilon^{-1}) \omega_e W_\ell^j (\tilde{v}_p - \tilde{v}_q)^2 \right] \right]
 \end{aligned}$$

Hence,

$$\begin{aligned}
 I_2 \leq & (1 + \epsilon) \sum_{k=1}^m \sum_{\ell=1}^m \sum_{\substack{e=(i,j) \in E_D^2 \\ i \in V_a^k, j \in V_a^\ell}} \omega_e (\tilde{v}_i - \tilde{v}_j)^2 \\
 & + \sum_{k=1}^m \sum_{e'=(p,q) \in E_a^k} \left[2(1 + \epsilon^{-1}) \left(\sum_{\ell=1}^m \sum_{\substack{e=(i,j) \in E_D^2 \\ i \in V_a^k, j \in V_a^\ell}} \omega_e W_k^i + \sum_{\ell=1}^m \sum_{\substack{e=(i,j) \in E_D^2 \\ i \in V_a^\ell, j \in V_a^k}} \omega_e W_k^j \right) (\tilde{v}_p - \tilde{v}_q)^2 \right]
 \end{aligned}$$

Now, we use the definition of $W_{e'}$ (4.10) and the estimates on I_1 and I_2 to obtain that

$$\begin{aligned}
 \|\tilde{P}^T \tilde{\mathbf{v}}\|_A^2 \leq & \sum_{e \in E_D^0} \omega_e (\tilde{v}_i - \tilde{v}_j)^2 + (1 + \epsilon) \sum_{k=1}^m \sum_{\substack{e=(i,j) \in E_D^1 \\ i \in V^0, j \in V_a^k}} \omega_e (\tilde{v}_i - \tilde{v}_j)^2 \\
 & + (1 + \epsilon) \sum_{k=1}^m \sum_{\ell=1}^m \sum_{\substack{e=(i,j) \in E_D^2 \\ i \in V_a^k, j \in V_a^\ell}} \omega_e (\tilde{v}_i - \tilde{v}_j)^2 + \sum_{k=1}^m \sum_{e'=(p,q) \in E_a^k} W_{e'} (\tilde{v}_p - \tilde{v}_q)^2.
 \end{aligned}$$

Due to (4.10), we have that $\omega_{e'} \geq W_{e'}$ and (4.11) follows. This completes the proof. \square

Lemma 7 shows that the constant c_1 can be made arbitrarily close to 1 if the weights on the internal edges of the disaggregation are chosen to be large enough. As an immediate consequence, we have the following theorem for the preconditioner B .

THEOREM 11 *Under the assumptions of Corollary 1 and Lemma 7, for the preconditioner $B = \tilde{P}^T \tilde{B}_D \tilde{P}$, we have*

$$\kappa(BA) \leq (1 + \epsilon) \kappa(\tilde{B}_D \tilde{A}_D). \quad (4.12)$$

Proof. The relation (4.12) follows from Corollary 1 since $c_0 = 1$ in (4.8) and $c_1 = (1 + \epsilon)^{1/2}$ in Lemma 7. \square

Finally, since $\tilde{A}_D := D_s^{-1} A_D D_s^{-1}$, if we have a preconditioner B_D for A_D and define $\tilde{B}_D = D_s B_D D_s$, then it is easy to verify that $\kappa(\tilde{B}_D \tilde{A}_D) = \kappa(B_D A_D)$. We have the following theorem showing that the preconditioned operator BA has a condition number comparable to the condition number of $B_D A_D$.

THEOREM 12 *Under the assumptions of Corollary 1 and Lemma 7 and let $\tilde{B}_D = D_s B_D D_s$, for the preconditioner $B = \tilde{P}^T \tilde{B}_D \tilde{P}$, we have*

$$\kappa(BA) \leq (1 + \epsilon) \kappa(B_D A_D). \quad (4.13)$$

Proof. (4.13) follows from Theorem 11 and the fact that $\kappa(\tilde{B}_D \tilde{A}_D) = \kappa(B_D A_D)$. \square

Clearly, Theorems 4.12 and 4.13 imply that, when the weights on the internal edges of the disaggregation are chosen to be large enough, preconditioners for disaggregated graph provide effective preconditioners for the original graph, which indirectly supports the technique suggested in [10].

Acknowledgements

The authors would like to thank Louisa Thomas for improving the style of the presentation. The work of Ludmil Zikatanov was supported in part by the National Science Foundation under grants DMS-1418843 and DMS-1522615 and by the Department of Mathematics at Tufts University.

References

- [1] Albert-László Barabási and Réka Albert. Emergence of scaling in random networks. *Science*, 286(5439):509–512, 1999.
- [2] Stephen Barnard and Horst Simon. Fast multilevel implementation of recursive spectral bisection for partitioning unstructured problems. *Concurrency: Practice and Experience*, 6(2):101–117, 1994.
- [3] James Brannick, Yao Chen, Johannes Kraus, and Ludmil Zikatanov. Algebraic multilevel preconditioners for the graph Laplacian based on matching in graphs. *SIAM Journal on Numerical Analysis*, 51(3):1805–1827, 2013.
- [4] Fan Chung. *Spectral graph theory*, volume 92. American Mathematical Soc., 1997.
- [5] Richard Courant and David Hilbert. *Methoden der mathematischen Physik*. Berlin, 1924.
- [6] Pasqua D’Ambra and Panayot Vassilevski. Compatible matching adaptive AMG preconditioners for Laplacian matrices on general graphs. Tech. Rep. LLNL-TR-676601, Lawrence Livermore National Laboratory, 2015.
- [7] Blanca Ayuso de Dios, Franco Brezzi, L. Donatella Marini, Jinchao Xu, and Ludmil T. Zikatanov. A simple preconditioner for a discontinuous Galerkin method for the Stokes problem. *Journal of Scientific Computing*, 58(3):517–547, 2014.
- [8] Shmuel Friedland and Reinhard Nabben. On Cheeger-type inequalities for weighted graphs. *Journal of Graph Theory*, 41(1):1–17, 2002.

- [9] Yongmei Ji, Xing Xu, and Gary Stormo. A graph theoretical approach for predicting common RNA secondary structure motifs including pseudoknots in unaligned sequences. *Bioinformatics*, 20(10):1591–1602, 2004.
- [10] Verena Kuhlemann and Panayot Vassilevski. Improving the communication pattern in matrix-vector operations for large scale-free graphs by disaggregation. *SIAM Journal on Scientific Computing*, 35(5):S465–S486, 2013.
- [11] Yin Tat Lee, Richard Peng, and Daniel A Spielman. Sparsified cholesky solvers for sdd linear systems. *arXiv preprint arXiv:1506.08204*, 2015.
- [12] Boaz Nadler, Stephane Lafon, Ronald Coifman, and Ioannis Kevrekidis. Diffusion maps, spectral clustering and eigenfunctions of fokker-planck operators. In *in Advances in Neural Information Processing Systems 18*, pages 955–962. MIT Press, 2005.
- [13] Sergey Nepomnyaschikh. Decomposition and fictitious domains methods for elliptic boundary value problems. In *Fifth International Symposium on Domain Decomposition Methods for Partial Differential Equations (Norfolk, VA, 1991)*, pages 62–72. SIAM, Philadelphia, PA, 1992.
- [14] Mark E.J. Newman. Finding community structure in networks using the eigenvectors of matrices. *Phys. Rev. E (3)*, 74(3):036104, 19, 2006.
- [15] John C Urschel, Jinchao Xu, Xiaozhe Hu, and Ludmil Zikatanov. A cascadic multigrid algorithm for computing the fiedler vector of graph Laplacians. *Journal of Computational Mathematics*, 33(2), 2015.
- [16] John C Urschel and Ludmil Zikatanov. Spectral bisection of graphs and connectedness. *Linear Algebra and its Applications*, 449:1–16, 2014.
- [17] Richard S. Varga. *Matrix iterative analysis*, volume 27 of *Springer Series in Computational Mathematics*. Springer-Verlag, Berlin, expanded edition, 2000.
- [18] Jinchao Xu. The auxiliary space method and optimal multigrid preconditioning techniques for unstructured grids. *Computing*, 56(3):215–235, 1996.