

# ENUMERATIONS OF VERTICES AMONG ALL ROOTED ORDERED TREES WITH LEVELS AND DEGREES

SEN-PENG EU, SEUNGHYUN SEO, AND HEESUNG SHIN<sup>†</sup>

**ABSTRACT.** In this paper we enumerate and give bijections for the following four sets of vertices among rooted ordered trees of a fixed size: (i) first-children of degree  $k$  at level  $\ell$ , (ii) non-first-children of degree  $k$  at level  $\ell - 1$ , (iii) leaves having  $k - 1$  elder siblings at level  $\ell$ , and (iv) non-leaves of outdegree  $k$  at level  $\ell - 1$ . Our results unite and generalize several previous works in the literature.

## 1. INTRODUCTION

Let  $\mathcal{T}_n$  be the set of *rooted ordered trees* with  $n$  edges. It is well known that the cardinality of  $\mathcal{T}_n$  is the  $n$ -th *Catalan number*

$$\text{Cat}_n = \frac{1}{n+1} \binom{2n}{n}.$$

For example, there are 5 rooted ordered trees with 3 edges, see Figure 1. Clearly the number of vertices among trees in  $\mathcal{T}_n$  is

$$(n+1) \text{Cat}_n = \binom{2n}{n}.$$

Given a rooted ordered tree, a vertex  $v$  is a *child* of a vertex  $u$  and  $u$  is the *parent* of  $v$  if  $v$  is directly connected to  $u$  when moving away from the root. A vertex without children is called a *leaf*. Note that by definition the root is not a child. Vertices with the same parent are called *siblings*. Since siblings are linearly ordered, when drawing trees the siblings are put in the left-to-right order. Siblings to the left of  $v$  are called an *elder* siblings of  $v$ . The leftmost vertex among siblings is called the *first-child*. In Figure 1, there are 10 first-children as well as 10 leaves, which is precisely a half of all 20 vertices in trees in  $\mathcal{T}_3$ .

In 1999, Shapiro [Sha99] proved the following using generating functions.

**Theorem 1.** *For any positive integer  $n$ , the following four sets of vertices among trees in  $\mathcal{T}_n$  are equinumerous:*

- (i) *first-children,*
- (ii) *non-first-children,*
- (iii) *leaves, and*
- (iv) *non-leaves.*


Here, the cardinality of each set is

$$\frac{1}{2} \binom{2n}{n},$$

---

*Date:* November 11, 2021.

<sup>†</sup>Corresponding author.

FIGURE 1. The five trees in  $\mathcal{T}_3$ 

which is a half of the number of vertices among trees in  $\mathcal{T}_n$ .

Seo and Shin [SS02] gave an involution proving (iii) and (iv) are equinumerous. The equality with the other two sets is easily seen: (i) and (iv) are equinumerous since each non-leaf has its unique first-child and the union of (i) and (ii) are the same with (iii) and (iv).

**1.1. Degree and outdegree.** The *degree* of a vertex is the number of edges incident to it. As every edge has a natural outward direction away from the root, we can have the notion of the *outdegree* of a vertex  $v$ , which is the number of edges starting at  $v$  and pointing away from the root. Since each vertex has a degree and each non-leaf has a positive outdegree, Theorem 1 can be restated as follows.

**Theorem 1'** (Shapiro, 1999). *Let  $n \geq 1$ . Among trees in  $\mathcal{T}_n$ , the number of vertices of positive degree equals twice the number of vertices of positive outdegree.*

In 2004, Eu, Liu, and Yeh [ELY04] proved combinatorially the following by constructing a two-to-one correspondence which answered a problem posed by Deutsch and Shapiro [DS01, p. 259].

**Theorem 2** (Eu, Liu, and Yeh, 2004). *Let  $n \geq 1$ . Among trees in  $\mathcal{T}_n$ , the number of vertices of odd degree equals twice the number of vertices of odd positive outdegree.*

In light of these two results, it is natural to ask if more can be said. In Corollary 6 we will prove that for any positive integer  $k$  the number of vertices of degree  $k$  always equals *twice* the number of that of outdegree  $k$  among trees in  $\mathcal{T}_n$ .

**1.2. Even and odd level.** A vertex  $v$  in a rooted tree  $T$  is at *level*  $\ell$  if the distance (number of edges) from the root to  $v$  is  $\ell$ . By an involution, the following result was obtained by Chen, Li, and Shapiro [CLS07].

**Theorem 3** (Chen, Li, and Shapiro, 2007). *The number of vertices at odd levels equals the number of vertices at even levels among trees in  $\mathcal{T}_n$ .*

It is again natural to ask if more can be said. In Corollary 8 of this paper we will prove that for any positive integer  $k$  the number of vertices of degree  $k$  at odd levels always equals the number of vertices of degree  $k$  at even levels among trees in  $\mathcal{T}_n$ .

**1.3. Main result.** In this paper we generalize the above by regarding both degrees and levels simultaneously. We will give a combinatorial proof for the following main result.

**Theorem 4.** *Given  $n \geq 1$ , for any two positive integers  $k$  and  $\ell$ , there are one-to-one correspondences between the following four sets of vertices among trees in  $\mathcal{T}_n$ :*

- (i) *first-children of degree  $k$  at level  $\ell$ ,*
- (ii) *non-first-children of degree  $k$  at level  $(\ell - 1)$ ,*
- (iii) *leaves having exactly  $(k - 1)$  elder siblings at level  $\ell$ , and*
- (iv) *non-leaves of outdegree  $k$  at level  $(\ell - 1)$ .*

*Here, the cardinality of each set is*

$$\frac{k + 2\ell - 2}{2n - k} \binom{2n - k}{n + \ell - 1}. \quad (1)$$

The rest of the paper is organized as follows. In Section 2 we derive several corollaries from Theorem 4, including generalizations of Theorem 1 through 3. The proof of Theorem 4 is given in the next two sections: In Section 3 we construct bijections between these four sets while in Section 4 we show combinatorially that the cardinality is given by (1).

## 2. COROLLARIES

In 2012, Cheon and Shapiro [CS12, Example 2.2] gave a formula for the number of vertices of outdegree  $k$  by generating function arguments. Summing over  $\ell \geq 1$  in Theorem 4 yields the following, in which the fourth item recovers bijectively the result of Cheon and Shapiro.

**Corollary 5.** *Given  $n \geq 1$  and  $k \geq 1$ , there are one-to-one correspondences between the following four sets of vertices among trees in  $\mathcal{T}_n$ :*

- (i) *first-children of degree  $k$ ,*
- (ii) *non-first-children of degree  $k$ ,*
- (iii) *leaves having exactly  $(k - 1)$  elder siblings, and*
- (iv) *non-leaves of outdegree  $k$ .*

*Here, the cardinality of each set is*

$$\binom{2n - k - 1}{n - 1}.$$

*Proof.* The result follows by a telescoping summation on  $\ell \geq 1$  using the formula

$$\frac{k + 2\ell - 2}{2n - k} \binom{2n - k}{n + \ell - 1} = \binom{2n - k - 1}{n + \ell - 2} - \binom{2n - k - 1}{n + \ell - 1}.$$

□

The result mentioned in Subsection 1.1 can be obtained from (i), (ii), and (iv) of Corollary 5.

**Corollary 6.** *Given  $n \geq 1$  and  $k \geq 1$ , the number of vertices of degree  $k$  equals twice the number of vertices of outdegree  $k$  among trees in  $\mathcal{T}_n$ .*

Note that Corollary 6 is a refinement of Theorem 2. Summing over all  $k$  in Theorem 4 yields the following.

**Corollary 7.** *Given  $n \geq 1$  and  $\ell \geq 1$ , there are one-to-one correspondences between the following four sets of vertices among trees in  $\mathcal{T}_n$ :*

- (i) *first-children at level  $\ell$ ,*
- (ii) *non-first-children at level  $(\ell - 1)$ ,*
- (iii) *leaves at level  $\ell$ , and*
- (iv) *non-leaves at level  $(\ell - 1)$ .*

*Here, the cardinality of each set is*

$$\frac{\ell}{n} \binom{2n}{n+\ell}.$$

*Proof.* The result follows by a telescoping summation on  $k \geq 1$  using the following formula,

$$\frac{k+2\ell-2}{2n-k} \binom{2n-k}{n+\ell-1} = \frac{k+2\ell-1}{2n-k+1} \binom{2n-k+1}{n+\ell} - \frac{k+2\ell}{2n-k} \binom{2n-k}{n+\ell}.$$

□

The result mentioned in Subsection 1.2 can be obtained from (i) and (ii) in Theorem 4.

**Corollary 8.** *Given  $n \geq 1$ , the number of vertices of degree  $k$  at odd levels equals the number of vertices of degree  $k$  at even levels among trees in  $\mathcal{T}_n$ .*

### 3. PROOF OF THEOREM 4: BIJECTIONS



Denote  $\mathcal{TV}_n$  by the set of pairs  $(T, v)$  satisfying  $T \in \mathcal{T}_n$  and  $v \in V(T)$ . Given positive integers  $n, k$ , and  $\ell$ , we let

- (i)  $\mathcal{A}$  denote the set of  $(T, v) \in \mathcal{TV}_n$  such that  $v$  is a first-child of degree  $k$  at level  $\ell$  in  $T$ ,
- (ii)  $\mathcal{B}$  denote the set of  $(T, v) \in \mathcal{TV}_n$  such that  $v$  is a non-first-child of degree  $k$  at level  $(\ell - 1)$  in  $T$ ,
- (iii)  $\mathcal{C}$  denote the set of  $(T, v) \in \mathcal{TV}_n$  such that  $v$  is a leaf having exactly  $(k - 1)$  elder siblings at level  $\ell$  in  $T$ , and
- (iv)  $\mathcal{D}$  denote the set of  $(T, v) \in \mathcal{TV}_n$  such that  $v$  is a non-leaf of outdegree  $k$  at level  $(\ell - 1)$  in  $T$ .

**(i)  $\Leftrightarrow$  (iv).** A bijection from  $\mathcal{A}$  to  $\mathcal{D}$  is constructed as follows: given  $(T, v) \in \mathcal{A}$ , find the parent vertex  $u$  of  $v$ . As the Figure 2, consider a subtree  $D_v$  of  $v$  and another subtree  $D_u$  of  $u$  on the right of the edge  $(u, v)$ . By interchanging  $D_v$  and  $D_u$  we get the new tree  $T'$  with the vertex  $u$  such that

$$\text{outdeg}(T', u) = \deg(T, v) = k, \quad \text{lev}(T', u) = \text{lev}(T, v) - 1 = \ell - 1,$$

where  $\text{lev}(T, v)$  (resp.  $\deg(T, v)$ ,  $\text{outdeg}(T, v)$ ) means the level (resp. degree, outdegree) of a vertex  $v$  in a tree  $T$ . Thus,  $(T', u)$  belongs to  $\mathcal{D}$ . Since this interchanging action is reversible, it is a one-to-one correspondence.


 FIGURE 2. A bijection from  $\mathcal{A}$  to  $\mathcal{D}$ 

 FIGURE 3. A bijection from  $\mathcal{B}$  to  $\mathcal{D}$  if  $v$  is not the root of  $T$ 

**(ii)  $\Leftrightarrow$  (iv).** A bijection from  $\mathcal{B}$  to  $\mathcal{D}$  is constructed as follow: given  $(T, v) \in \mathcal{B}$ , we perform the following action:

(a) if  $v$  is the root of  $T$ , the pair  $(T, v)$  also belongs to  $\mathcal{D}$  due to

$$\text{outdeg}(T, v) = \deg(T, v) = k.$$

(b) if  $v$  is not the root of  $T$ , find the parent vertex  $u$  of  $v$ , the first-child  $w$  of  $u$  and the edge  $e = (u, w)$ . As the Figure 3, cut and paste the edge  $e$  and the subtree  $D_w$  consisting of all descendants of  $w$  from  $u$  to  $v$  such that the vertex  $w$  is the first-child of  $v$ . We obtain the tree  $T'$  and the vertex  $v$  in  $T'$  satisfies

$$\text{outdeg}(T', v) = \deg(T, v) = k, \quad \text{lev}(T', v) = \text{lev}(T, v) = \ell - 1.$$



Thus,  $(T', v)$  belongs to  $\mathcal{D}$ .

Since this action is reversible, it is a one-to-one correspondence.

**(iii)  $\Leftrightarrow$  (iv).** A bijection from  $\mathcal{C}$  to  $\mathcal{D}$  is constructed as follows: given  $(T, v) \in \mathcal{C}$ , find the parent vertex  $u$  of  $v$ . As the Figure 4, consider the subtree  $D_{uv}$  of  $u$  on the right of the edge  $(u, v)$ . By cutting and pasting the subtrees  $D_{uv}$  from  $u$  to  $v$  we get the new tree  $T'$  with the vertex  $u$  such that

$$\text{outdeg}(T', u) = \text{eld}(T, v) + 1 = k, \quad \text{lev}(T', u) = \text{lev}(T, v) - 1 = \ell - 1,$$

where  $\text{eld}(T, v)$  means the number of elder siblings of a vertex  $v$  in a tree  $T$ . Thus,  $(T', u)$  belongs to  $\mathcal{D}$ . Since this interchanging action is reversible, it is a one-to-one correspondence.

FIGURE 4. A bijection from  $\mathcal{C}$  to  $\mathcal{D}$ FIGURE 5. The bijection  $\varphi$ 

## 4. PROOF OF THEOREM 4: ENUMERATION

By putting  $\ell + 1$  in place of  $\ell$  in (iv) of Theorem 4, it suffices to show that for any two nonnegative integers  $k$  and  $\ell$ , the number of vertices of outdegree  $k$  at level  $\ell$  among trees in  $\mathcal{T}_n$  is

$$\frac{k + 2\ell}{2n - k} \binom{2n - k}{n + \ell}.$$

The following lemma gives a cumulative counting in  $k$  and  $\ell$ .

**Lemma 9** (Main lemma). *Given  $n \geq 1$ , for any two nonnegative integers  $k$  and  $\ell$ , the number of vertices of outdegree at least  $k$  and at level at least  $\ell$  among trees in  $\mathcal{T}_n$  is*

$$\binom{2n - k}{n + \ell}. \quad (2)$$

*Proof.* Let  $\mathcal{V}$  be the set of  $(T, v) \in \mathcal{TV}_n$  such that  $v$  is a vertex of outdegree at least  $k$  and at level at least  $\ell$  in  $T$ . Let  $\mathcal{L}$  be the set of *lattice paths* of length  $(2n - k)$  from  $(k, k)$  to  $(2n, -2\ell)$ , consisting of  $(n - k - \ell)$  up-steps along the vector  $(1, 1)$  and  $(n + \ell)$  down-steps along the vector  $(1, -1)$ . Since

$$\#\mathcal{L} = \binom{2n - k}{n - k - \ell, n + \ell} = \binom{2n - k}{n + \ell},$$

it is enough to construct a bijection  $\Phi$  between  $\mathcal{V}$  and  $\mathcal{L}$ .

Before constructing the bijection we first introduce two well-known bijections  $\varphi$  and  $\psi$  between rooted ordered trees and Dyck paths.

The bijection  $\varphi$  corresponds a tree to a Dyck path by recording the steps when the tree is traversed in preorder. Here we record an up-step when we go down an edge and a down-step when going up. An example of the bijection  $\varphi$  is shown in the Figure 5.

The bijection  $\psi$  corresponds a tree to a Dyck path by recording the steps when the tree is traversed in preorder. Here, whenever we meet a vertex of




 FIGURE 6. The bijection  $\psi$ 


FIGURE 7. Tree decomposition

outdegree  $k$ , except the last leaf, we record  $k$  up-steps and one down-step. An example of the bijection  $\psi$  is shown in the Figure 6.

Using two bijections  $\varphi$  and  $\psi$ , we will construct another bijection  $\Phi$  between  $\mathcal{V}$  to  $\mathcal{L}$ : given  $(T, v) \in \mathcal{V}$ , let  $k' (\geq k)$  be the number of children of  $v$  in  $T$  and let  $\ell' (\geq \ell)$  be the level of  $v$  in  $T$ . We decompose the tree  $T$  into  $\ell' + 2$  subtrees: the subtree  $D_v$  consisting of descendants of  $v$ ,  $\ell'$  subtrees  $R_1, R_2, \dots, R_{\ell'}$  on the right hand side of the path from  $v$  to the root  $r$  of  $T$ , and the remaining tree  $L$  as illustrated in the Figure 7. Clearly, the outdegree of the root of  $D_v$  is  $k'$ . In preorder, the vertex  $v$  is the last leaf at level  $\ell'$  in the tree  $L$ .

From  $(T, v)$ , we define a lattice path  $P$  of length  $(2n + \ell' + 1)$  from  $(0, 0)$  to  $(2n + \ell' + 1, -\ell' - 1)$  by



$$P = \psi(D_v) \searrow \varphi(R_1) \searrow \varphi(R_2) \searrow \cdots \searrow \varphi(R_{\ell'}) \searrow \varphi(L),$$

where  $\searrow$  means a down-step. By convention, if  $\ell' = 0$ . We set

$$P = \psi(D_v) \searrow \varphi(\emptyset).$$

Especially, if  $\ell' > 0$ , the lattice path  $P$  always starts with  $k'$  consecutive up-steps and ends with one up-step and  $\ell'$  consecutive down-steps as the Figure 8.

When we draw the line  $y = -\ell - 1$ , because the  $y$ -coordinate of the lowest point of  $P$  is  $-\ell' - 1$ , this line should meet the lattice path  $P$ . Denote  $p$  by the first meeting point of  $P$  and  $y = -\ell - 1$ . Replacing the portion of  $P$  from  $p$  to  $(2n + \ell' + 1, -\ell' - 1)$  by its reflection about  $y = -\ell - 1$ , we obtain the lattice path  $\tilde{P}$  from  $(0, 0)$  to  $(2n + \ell' + 1, -2\ell + \ell' - 1)$ , which also always starts with  $k'$  consecutive up-steps and ends with one down-step and

FIGURE 8. Outline of a lattice path  $P$  induced from tree decompositionFIGURE 9. Outline of a lattice path  $\tilde{P}$  and  $\hat{P}$  by reflection about  $y = -\ell - 1$ 

$\ell'$  consecutive up-steps as the Figure 9, even if  $\ell' = 0$ . Note that, if  $\ell' = 0$ ,  $p$  should be the last point  $(2n + 1, -1)$  of  $P$  and  $\tilde{P} = P$ .

By removing the first  $k$  steps and the last  $(\ell' + 1)$  steps from  $\tilde{P}$ , we obtain the lattice path  $\hat{P}$  from  $(k, k)$  to  $(2n, -2\ell)$ . Thus  $\hat{P}$  belongs to  $\mathcal{L}$  and define  $\Phi(T, v) = \hat{P}$ . Since the reflection and the removal are reversible,  $\Phi$  is a bijection.  $\square$

**Theorem 10.** *Given  $n \geq 1$ , for any two nonnegative integers  $k$  and  $\ell$ , the number of vertices of outdegree  $k$  at level  $\ell$  among trees in  $\mathcal{T}_n$  is*

$$\frac{k + 2\ell}{2n - k} \binom{2n - k}{n + \ell}. \quad (3)$$

*Proof.* Using a sieve method, we obtain (3) from (2) by

$$\binom{2n - k}{n + \ell} - \binom{2n - k - 1}{n + \ell} - \binom{2n - k}{n + \ell + 1} + \binom{2n - k - 1}{n + \ell + 1}.$$

$\square$

Thus, obtaining (1) from (3) by changing the index  $\ell$ , we may count the set  $\mathcal{D}$  as (iv) of Theorem 4.

#### ACKNOWLEDGEMENTS

For the first author, this research was partially supported by Ministry of Science of Technology, Taiwan ROC under grants 104-2115-M-003-014-MY3. For the second author, this study has been worked with the support



of a research grant of Kangwon National University in 2015. For the third author, this work was supported by Inha University Research Grant.

# REFERENCES

- [CLS07] William Y. C. Chen, Nelson Y. Li, and Louis W. Shapiro. The butterfly decomposition of plane trees. *Discrete Appl. Math.*, 155(17):2187–2201, 2007.
- [CS12] Gi-Sang Cheon and Louis Shapiro. The uplift principle for ordered trees. *Appl. Math. Lett.*, 25(6):1010–1015, 2012.
- [DS01] Emeric Deutsch and Louis Shapiro. A survey of the Fine numbers. *Discrete Math.*, 241(1-3):241–265, 2001. Selected papers in honor of Helge Tverberg.
- [ELY04] Sen-Peng Eu, Shu-Chung Liu, and Yeong-Nan Yeh. Odd or even on plane trees. *Discrete Math.*, 281(1-3):189–196, 2004.
- [Sha99] Lou Shapiro. The higher you go, the odder it gets. In *Proceedings of the Thirtieth Southeastern International Conference on Combinatorics, Graph Theory, and Computing (Boca Raton, FL, 1999)*, volume 138, pages 93–96, 1999.
- [SS02] Seunghyun Seo and Heesung Shin. Two involutions on vertices of ordered trees. In *14th International Conference on Formal Power Series and Algebraic Combinatorics (FPSAC 2002)*. 2002.

(Sen-Peng Eu) DEPARTMENT OF MATHEMATICS, NATIONAL TAIWAN NORMAL UNIVERSITY, TAIPEI 116, TAIWAN, ROC  
*E-mail address:* `speu@math.ntnu.edu.tw`

(Seunghyun Seo) DEPARTMENT OF MATHEMATICS EDUCATION, KANGWON NATIONAL UNIVERSITY, CHUNCHEON 24341, KOREA  
*E-mail address:* `shyunseo@kangwon.ac.kr`

(Heesung Shin) DEPARTMENT OF MATHEMATICS, INHA UNIVERSITY, INCHEON 22212, KOREA  
*E-mail address:* `shin@inha.ac.kr`