

THE COHOMOLOGY OF $(S(n, k))$ RELEVANT TO MORAVA STABILIZER ALGEBRA

LIMAN CHEN, XIANGJUN WANG AND XUEZHI ZHAO

ABSTRACT. In this paper we redefine a increasing filtration on the the Hopf algebra $S(n, k)$, From which we get a spectral sequence called May spectral sequence. As an application we computed $H^{*,*}S(n, n)$ at prime 2, $H^{*,*}S(3, 2)$ at prime 3 and $H^{*,*}S(4, 2)$ at prime $p \geq 5$

1. INTRODUCTION

In stable homotopy theory, the “chromatic” point of view plays an important role (*cf.* [3, 11, 14]). Fix a prime p . Let $E(n)_*$, $n \geq 0$ be the Johnson-Wilson homology theories and let L_n be localization functor with respect to $E(n)_*$. Then there are natural transformations $L_n X \rightarrow L_{n-1} X$, and the chromatic tower

$$\cdots \longrightarrow L_n X \longrightarrow L_{n-1} X \longrightarrow \cdots \longrightarrow L_2 X \longrightarrow L_1 X \longrightarrow L_0 X.$$

By the Hopkins-Ravenel chromatic convergence theorem, the homotopy inverse limit of this tower is the p -localization of X

$$X \longrightarrow \operatorname{Holim} L_n X.$$

Thus the homotopy groups $\pi_*(L_n X)$ is the part of homotopy groups $\pi_*(X)$ one could see from $E(n)_*$.

To determine the homotopy groups $\pi_*(L_n X)$, one has the Adams-Novikov spectral sequence based on the Brown-Peterson spectrum BP , whose E_2 -term is

$$E_2^{s,t} = Ext_{BP_* BP}^{s,t}(BP_*, BP_*(L_n X)).$$

(*cf.* [1, 10, 11, 14])

To determine the Adams-Novikov E_2 -term $Ext_{BP_* BP}^{s,t}(BP_*, BP_*(L_n X))$ one has the Bockstein spectral sequence. This is an argument based on the cohomology of the Morava stabilizer algebra $S(n)$ at each prime p (*cf.* [11, 16, 17, 19]). Here the Hopf algebra $S(n)$ is defined as

$$S(n) = Z/p \otimes_{K(n)_*} K(n)_* K(n) \otimes_{K(n)_*} Z/p,$$

where $K(n)_* = Z/p[v_n, v_n^{-1}]$,

$$K(n)_* K(n) = K(n)_* \otimes_{BP_*} BP_* BP \otimes_{BP_*} K(n)_* = K(n)_*[t_1, t_2, \dots]/(v_n t_s^{p^n} - v_n^{p^s} t_s),$$

$K(n)_*$ acts on Z/p by sending v_n to 1. Thus

$$S(n) = Z/p[t_1, t_2, \dots, t_s, \dots]/(t_s^{p^n} - t_s).$$

1991 *Mathematics Subject Classification.* Primary 55Q99; Secondary 55Q51.

Key words and phrases. Morava n -th K-theory, Morava stabilizer algebra, May spectral sequence.

The second author was supported by NSFC grant No. 11471167 and SRFPD No.20120031110025.

We write $S(n, k) = S(n)/(t_j : j < k) = Z/p[t_k, t_{k+1}, \dots, t_s, \dots]/(t_s^{p^n} - t_s)$. The Hopf algebra structure of $S(n)$ determines that of $S(n, k)$, while $S(n, 1) = S(n)$. Let $V(n-1)$ and $T(k-1)$ denote the Smith-Toda spectra and the Ravenel spectra respectively characterized by

$$\begin{aligned} BP_*V(n-1) &= BP_*/I_n = BP_*/(p, v_1, \dots, v_{n-1}) & \text{and} \\ BP_*T(k-1) &= BP_*[t_1, t_2, \dots, t_{k-1}]. \end{aligned}$$

If $L_nV(n-1) \wedge T(k-1)$ exist, (although $V(n-1)$ does not exist (*cf.* [6]), but $V(n-1) \wedge T(k-1)$ might exist), then by the change of rings theorem, the E_2 -term of the Adams-Novikov spectral sequence converging to $\pi_*(L_nV(n-1) \wedge T(k-1))$ is

$$\begin{aligned} &Ext_{BP_*BP}^{s,t}(BP_*, BP_*(L_nV(n-1) \wedge T(k-1))) \\ &\cong Ext_{S(n,k)}^{s,t}(Z/p, Z/p) \otimes K(n)_*[v_{n+1}, \dots, v_{n+k-1}]. \end{aligned}$$

In this paper, we will use $H^{s,*}S(n, k)$ to denote the Ext groups $Ext_{S(n,k)}^{s,*}(Z/p, Z/p)$. In [5, 13], Ravenel and Henn determined $H^{s,*}S(1)$, $H^{s,*}S(2)$ at all primes, and $H^{s,*}S(3)$ at the odd primes $p \geq 5$. $H^{s,*}S(n, k)$ is known from [11] for $k \geq n$ at odd primes and $k > n$ at the prime 2. In [15] Shimomura and Tokashiki computed $H^{s,*}S(n, n-1)$ at odd primes $p > 3$. In this paper we will be concentrated on the case $k \leq n$.

Consider the cohomology of the Hopf algebra $S(n, k)$ at all primes. In section 2 of this paper, we follow Ravenel's ideal (*cf.* [11] 3.2.5 Theorem), redefined the May filtration in $S(n, k)$ and its cobar complex $C^{s,*}S(n, k)$. This filtration induces a spectral sequence so called May spectral sequence $\{E_r^{s,*M}(n, k), d_r\}$ that converges to $H^{s,*}S(n, k)$. Then in section 3 we prove that the E_2 -term of the May spectral sequence is isomorphic to the cohomology of

$$\{E[h_{i,j}|k \leq i \leq s_0, j \in Z/n] \otimes P[b_{i,j}|k \leq i \leq s_0 - n, j \in Z/n], d_1\}$$

where $s_0 = \max\left\{\left\lceil \frac{2pn+p-2}{2(p-1)} \right\rceil, n+k-1\right\}$ and $\left\lceil \frac{2pn+p-2}{2(p-1)} \right\rceil$ is the integer part of $\frac{2pn+p-2}{2(p-1)}$. In particular, if

$$n+k-1 \geq \left\lceil \frac{2pn+p-2}{2(p-1)} \right\rceil,$$

the May's E_2 -term becomes the cohomology of

$$\{E[h_{i,j}|k \leq i \leq n+k-1, j \in Z/n], d_1\}.$$

The homological dimension of each element is given by

$$s(h_{i,j})=1, \quad s(b_{i,j})=2.$$

For the May differentials, one has $d_r : E_r^{s,t,M}S(n, k) \longrightarrow E_r^{s+1,t,M-r}S(n, k)$ and if $x \in E_r^{s,*}S(n, k)$ then

$$d_r(x \cdot y) = d_r(x) \cdot y + (-1)^s x \cdot d_r(y).$$

The first May differential d_1 is given by

$$d_1(h_{i,j}) = - \sum_{k \leq m \leq i-k} h_{m,j} h_{i-m,j+m}, \quad \text{and} \quad d_1(b_{i,j}) = 0.$$

We analyze the higher May differentials and give a collapse theorem in section 4. As an consequence we compute the cohomology of $S(n, n)$ at the prime 2, $S(3, 2)$ at the prime 3 and $S(4, 2)$ at the prime $p \geq 5$ in section 5.

2. THE MAY SPECTRAL SEQUENCE

Let p be a prime, $BP_* = Z_{(p)}[v_1, v_2, \dots]$ and $BP_*BP = BP_*[t_1, t_2, \dots]$. For the Hazewinkel's generators described inductively by $v_s = pm_s - \sum_{i=1}^{s-1} v_{s-i}^{p^i} m_i$ (cf. [4] and [10] 1.2), the diagonal map $\Delta : BP_*BP \rightarrow BP_*BP \otimes_{BP_*} BP_*BP$ is given by

$$\sum_{i+j=s} m_i(\Delta t_j) = \sum_{i+j+k=s} m_i t_j^{p^i} \otimes t_k^{p^{i+j}}.$$

One can easily prove that

$$\begin{aligned} \Delta(t_1) &= t_1 \otimes 1 + 1 \otimes t_1 & \text{and} \\ \Delta(t_2) &= \sum_{i+j=2} t_i \otimes t_j^{p^i} - v_1 b_{1,0}, \end{aligned}$$

where $p \cdot b_{1,0} = \Delta(t_1^p) - t_1^p \otimes 1 - 1 \otimes t_1^p$. Inductively define

$$p \cdot b_{s,k-1} = \Delta(t_s^{p^k}) - \sum_{i+j=s} t_i^{p^k} \otimes t_j^{p^{i+k}} + \sum_{0 < i < s} v_i^{p^k} b_{s-i,k+i-1},$$

one has

$$(2.1) \quad \Delta(t_{s+1}) = \sum_{i+j=s+1} t_i \otimes t_j^{p^i} - \sum_{0 < i < s+1} v_i b_{s+1-i,i-1}.$$

Thus for the n -th Morava K-theory $K(n) = Z/p[v_n, v_n^{-1}]$, the Hopf algebra

$$K(n)_* K(n) = K(n)_* \otimes_{BP_*} BP_*BP \otimes_{BP_*} K(n)_*$$

is isomorphic to

$$K(n)_* K(n) = Z/p[t_1, t_2, \dots, t_s, \dots] / (v_n t_s^{p^n} - v_n^{p^s} t_s).$$

And $S(n) = Z/p \otimes_{K(n)_*} K(n)_* K(n) \otimes_{K(n)_*} Z/p$ is isomorphic to

$$S(n) = Z/p[t_1, t_2, \dots, t_n, t_{n+1}, \dots] / (t_s^{p^n} - t_s).$$

The inner degree of t_s in $S(n)$ is

$$|t_s| \equiv 2(p-1)(1+p+\dots+p^{s-1}) \pmod{2(p-1)(1+p+\dots+p^{n-1})}$$

because v_n is sent to 1. The structure map $\Delta : S(n) \rightarrow S(n) \otimes S(n)$ acts on t_s as follows

$$\begin{aligned} \Delta(t_s) &= \sum_{0 \leq i \leq s} t_i \otimes t_{s-i}^{p^i} & \text{for } s \leq n \\ \Delta(t_s) &= \sum_{0 \leq i \leq s} t_i \otimes t_{s-i}^{p^i} - b_{s-n,n-1} & \text{for } s > n \end{aligned}$$

where $b_{i,j} = \sum_{0 < m < p} \binom{p}{m} / p \cdot t_i^{mp^j} \otimes t_i^{(p-m)p^j}$ at odd primes and $b_{i,j} = t_i^{2^j} \otimes t_i^{2^j}$ at the prime 2. For the integer $k \geq 1$, let $S(n, k) = S(n) / (t_s | s < k)$. We have

$$S(n, k) = Z/p[t_k, t_{k+1}, \dots, t_{n+k}, t_{n+k+1}, \dots] / (t_s^{p^n} - t_s),$$

the structure map $\Delta : S(n, k) \longrightarrow S(n, k) \otimes S(n, k)$ acts on t_s as

$$(2.2) \quad \begin{aligned} \Delta(t_s) &= 1 \otimes t_s + \sum_{k \leq i \leq s-k} t_i \otimes t_{s-i}^{p^i} + t_s \otimes 1 & \text{for } s \leq n+k-1, \\ \Delta(t_s) &= 1 \otimes t_s + \sum_{k \leq i \leq s-k} t_i \otimes t_{s-i}^{p^i} + t_s \otimes 1 - b_{s-n, n-1} & \text{for } s \geq n+k. \end{aligned}$$

In the resulting May spectral sequence, we want to have the 0-th May differential is

$$d_0(t_s^{p^j}) = 0$$

and the first May differential is given by

$$d_1(t_s) = t_k \otimes t_{s-k}^{p^k} + t_{k+1} \otimes t_{s-k-1}^{p^{k+1}} + \cdots + t_{s-k} \otimes t_k^{p^{s-k}} \quad \text{for } s \leq n+k-1,$$

and for $s \geq n+k$

$$d_1(t_s) = \begin{cases} t_k \otimes t_{s-k}^{p^k} + \cdots + t_{s-k} \otimes t_k^{p^{s-k}} & \text{if the May filtration } M(t_k \otimes t_{s-k}^{p^k}) > M(b_{s-n, n-1}), \\ -b_{s-n, n-1} & \text{if the May filtration } M(t_k \otimes t_{s-k}^{p^k}) \leq M(b_{s-n, n-1}). \end{cases}$$

So we define the May filtration on $S(n, k)$ as:

Definition 2.3 In the Hopf algebra $S(n, k)$, we define May filtration M as follows:

- (1) For $k \leq s \leq n+k-1$, set the May filtration of $t_s^{p^j}$ as $M(t_s^{p^j}) = 2s-1$.
- (2) For $n+k \leq s$, inductively set the May filtration of $t_s^{p^j}$ as

$$M(t_s^{p^j}) = \max\{2s-1, pM(t_{s-n}^{p^{j+n-1}}) + 1\}.$$

- (3) For the monomial t_s^j , express j by the p -adic number as $j = j_0 + j_1p + \cdots + j_mp^m$, where $0 \leq j_i < p$. Set the May filtration of t_s^j as

$$M(t_s^j) = \sum_{0 \leq i \leq m} j_i M(t_s^{p^i}).$$

- (4) And for $t_{s_1}^{j_1} \cdot t_{s_2}^{j_2} \cdots t_{s_m}^{j_m}$, where $s_i \neq s_j$ define its May filtration as

$$M(t_{s_1}^{j_1} \cdot t_{s_2}^{j_2} \cdots t_{s_m}^{j_m}) = \sum_{1 \leq i \leq m} M(t_{s_i}^{j_i}).$$

Lemma 2.4 Let $s_0 = \max \left\{ \left\lfloor \frac{2pn+p-2}{2(p-1)} \right\rfloor, n+k-1 \right\}$ where $\left\lfloor \frac{2pn+p-2}{2(p-1)} \right\rfloor$ is the integer part of $\frac{2pn+p-2}{2(p-1)}$. Then the May filtration of $t_s^{p^j}$ satisfies

- (1) $M(t_s^{p^j}) > M(t_{s-1}^{p^j}) + 1$ and
- (2) For $s \leq s_0$, the May filtration $M(t_s^{p^j}) = 2s-1$.
- (3) For $s > s_0$, $pM(t_{s-n}^{p^j}) + 1 \geq 2s-1$ and the May filtration $M(t_s^{p^j}) = pM(t_{s-n}^{p^j}) + 1$.

Proof. 1) If $s_0 = \max \left\{ \left\lfloor \frac{2pn+p-2}{2(p-1)} \right\rfloor, n+k-1 \right\} = n+k-1$. From its definition, we see that for $s \leq n+k-1 = s_0$, the May filtration of $t_s^{p^j}$ is $2s-1$ and $M(t_s^{p^j}) > M(t_{s-1}^{p^j}) + 1$. From $n+k-1 \geq \left\lfloor \frac{2pn+p-2}{2(p-1)} \right\rfloor$, one sees that

$$s_0 + 1 = n+k > \frac{2pn+p-2}{2(p-1)} \quad \text{and} \quad p(2k-1) + 1 > 2(n+k) - 1.$$

Thus from $M(t_k^{p^j}) = 2k - 1$, one knows that the May filtration of $t_{n+k}^{p^j}$ is $pM(t_k^{p^j}) + 1$ and

$$M(t_{n+k}^{p^j}) = p(2k - 1) + 1 > 2(n + k) - 1 = M(t_{n+k-1}^{p^j}) + 1.$$

Inductively suppose that $M(t_s^{p^j}) > M(t_{s-1}^{p^j}) + 1$ and for $s_0 < s \leq m$,

$$pM(t_{s-n}^{p^j}) + 1 > 2s - 1,$$

so the May filtration $M(t_s^{p^j}) = pM(t_{s-n}^{p^j}) + 1$. Then from $M(t_{m+1-n}^{p^j}) > M(t_{m-n}^{p^j}) + 1$ one get

$$\begin{aligned} pM(t_{m+1-n}^{p^j}) + 1 &> p(M(t_{m-n}^{p^j}) + 1) + 1 = pM(t_{m-n}^{p^j}) + p + 1 \\ &> 2m - 1 + p \geq 2(m + 1) - 1. \end{aligned}$$

The May filtration of $t_{m+1}^{p^j}$ is $pM(t_{m+1-n}^{p^j}) + 1$.

If $s_0 = \left\lfloor \frac{2pn + p - 2}{2(p-1)} \right\rfloor > n + k - 1$, then for $k \leq s \leq s_0$, $s \leq \frac{2pn + p - 2}{2(p-1)}$. This implies

$$p(2(s - n) - 1) + 1 \leq 2s - 1.$$

From $\frac{2pn + p - 2}{2(p-1)} \leq 2n$ we see that $s - n \leq n \leq n + k - 1$. Thus the May filtration

$M(t_{s-n}^{p^j}) = 2(s - n) - 1$ and $pM(t_{s-n}^{p^j}) + 1 < 2s - 1$. This implies that the May filtration of $t_s^{p^j}$ is $2s - 1$ and $M(t_s^{p^j}) > M(t_{s-1}^{p^j}) + 1$.

Notice that $s_0 + 1 = \left\lfloor \frac{2pn + p - 2}{2(p-1)} \right\rfloor + 1 > \frac{2pn + p - 2}{2(p-1)}$, this implies

$$p(2(s_0 + 1 - n) - 1) + 1 > 2(s_0 + 1) - 1.$$

The May filtration of $t_{s_0+1-n}^{p^j}$ is $2(s_0 + 1 - n) - 1$, so the May filtration

$$M(t_{s_0+1}^{p^j}) = pM(t_{s_0+1-n}^{p^j}) + 1.$$

Similarly, by induction we get the Lemma. \square

Example: The May filtration in $S(4, 2)$ is given by:

	t_2	t_3	t_4	t_5	t_6	t_7	t_8	t_9	t_{10}	\dots
$p = 2$	3	5	7	9	11	13	15	19	23	\dots
$p = 3$	3	5	7	9	11	16	22	28	34	\dots
$p \geq 5$	3	5	7	9	$3p + 1$	$5p + 1$	$7p + 1$	$9p + 1$	$p(3p + 1) + 1$	\dots

Let $F^{*,M}(n, k)$ be the sub-module of $S(n, k)$ generated by the elements with May filtration $\leq M$. Set $E^{*,M}(n, k) = F^{*,M}(n, k)/F^{*,M-1}(n, k)$. One can see from Lemma 2.4 that

$$(2.5) \quad E^{*,*}(n, k) \cong \bigotimes_{k \leq s} T[t_s^{p^j} | j \in Z/n]$$

is a bigraded Hopf algebra, where $T[\quad]$ denote the truncated polynomial algebra of height p on the indicated generators. The structure map

$$\Delta : E^{*,*}(n, k) \longrightarrow E^{*,*}(n, k) \otimes E^{*,*}(n, k)$$

acts the the generators $t_s^{p^j}$ as $\Delta(t_s^{p^j}) = 1 \otimes t_s^{p^j} + t_s^{p^j} \otimes 1$.

Let $C^{s,t}S(n, k) = \otimes^s \bar{S}(n, k)$ denote the cobar construction of $S(n, k)$ where $\bar{S}(n, k) = \text{Ker } \epsilon$ denote the augmentation ideal of $S(n, k)$. The differential

$$d : C^{s,t}S(n, k) \longrightarrow C^{s+1,t}S(n, k)$$

is given on the generators as

$$(2.6) \quad d(\alpha_1 \otimes \cdots \otimes \alpha_s) = \sum_{1 \leq i \leq s} (-1)^i \alpha_1 \otimes \cdots \otimes (\Delta(\alpha_i) - \alpha_i \otimes 1 - 1 \otimes \alpha_i) \otimes \cdots \otimes \alpha_s.$$

In general, the generator $\alpha_1 \otimes \alpha_2 \otimes \cdots \otimes \alpha_s$ of $C^{s,t}S(n, k)$ is denoted by $[\alpha_1 | \alpha_2 | \cdots | \alpha_s]$. For the generator $[\alpha_1 | \alpha_2 | \cdots | \alpha_s]$, define its May filtration as

$$M([\alpha_1 | \alpha_2 | \cdots | \alpha_s]) = M(\alpha_1) + M(\alpha_2) + \cdots + M(\alpha_s).$$

Let $FC^{*,*,M}S(n, k)$ denote the sub-complex of $C^{*,*}S(n, k)$ generated by the elements with May filtration $\leq M$. Then we get a short exact sequence

$$(2.7) \quad 0 \longrightarrow FC^{*,*,M-1}S(n, k) \longrightarrow FC^{*,*,M}S(n, k) \longrightarrow E_0^{*,*,M}S(n, k) \longrightarrow 0$$

of cochain complexes. The cochain complex

$$E_0^{*,*,M}S(n, k) = FC^{*,*,M}S(n, k) / FC^{*,*,M-1}S(n, k)$$

is isomorphic to the cobar complex of $E^{*,*}(n, k)$ given in (2.5). Let $E_1^{*,*,M}S(n, k)$ be the homology of $(E_0^{*,*,M}S(n, k), d_0)$. Then (2.7) gives rise to a spectral sequence (so called the May spectral sequence)

$$\{E_r^{s,t,M}(n, k), d_r\}$$

that converges to

$$H^{s,t}(C^{*,*}S(n, k), d) = Ext_{S(n, k)}^{s,t}(Z/p, Z/p).$$

Theorem 2.8 *For $k \leq n$ the Hopf algebra $S(n, k)$ can be given an increasing filtration as in Definition (2.3). The associated bigraded Hopf algebra $E^{*,*}(n, k)$ is primitively generated with the algebra structure of (2.5). In the associated spectral sequence, the E_1 -term $E_1^{s,t,M}S(n, k)$ is isomorphic to*

$$E[h_{i,j} | k \leq i, j \in Z/n] \otimes P[b_{i,j} | k \leq i, j \in Z/n].$$

The homological dimension of each element is given by $s(h_{i,j}) = 1$, $s(b_{i,j}) = 2$ and the degree is given by

$$\begin{aligned} h_{i,j} &\in E_1^{1,2(p^i-1)p^j,*}(n, k), \\ b_{i,j} &\in E_1^{2,2(p^i-1)p^{j+1},*}(n, k) \end{aligned}$$

here $h_{i,j}$ corresponds to $t_i^{p^j}$ and $b_{i,j}$ corresponds to $\sum \binom{p}{m} / pt_i^{mp^j} \otimes t_i^{(p-m)p^j}$. One has $d_r : E_r^{s,t,M}S(n, k) \longrightarrow E_r^{s+1,t,M-r}S(n, k)$ and if $x \in E_r^{s,,*}S(n, k)$ then*

$$d_r(x \cdot y) = d_r(x) \cdot y + (-1)^s x \cdot d_r(y).$$

In the E_1 -term of this spectral sequence, we have the following relations:

$$h_{i,j} \cdot h_{i_1,j_1} = -h_{i_1,j_1} h_{i,j} \quad h_{i,j} \cdot b_{i_1,j_1} = b_{i_1,j_1} \cdot h_{i,j} \quad b_{i,j} \cdot b_{i_1,j_1} = b_{i_1,j_1} \cdot b_{i,j}$$

Proof. It is a routine calculation in homology algebra that for the truncated polynomial algebra $\Gamma = T[x]$ with $|x| \equiv 0 \pmod{2}$ and x primitive,

$$Ext_{\Gamma}(Z/p, Z/p) = E[h] \otimes P[b]$$

where $h \in Ext^1$ is represented in the cobar complex by x and $b \in Ext^2$ is represented by $\sum \binom{p}{m}/p(x^m \otimes x^{(p-m)})$ ($b = h^2$ represented by $x \otimes x$ at the prime 2). Notice that the E_0 -term of the spectral sequence is isomorphic to the cobar complex of $E^{*,M}(n, k)$. Then from (2.5) we see that

$$H^{s,*,M}(E_0^{*,t,M}S(n, k), d_0) = Ext_{E^{*,*}(n, k)}^{s,t}(Z/p, Z/p) = \bigotimes_{k \leq s} Ext_{T[t_s^p]}^{*,*}(Z/p, Z/p)$$

Thus the May's E_1 -term

$$E_1^{s,t,M}S(n, k) = E[h_{i,j} | k \leq i, j \in Z/n] \otimes P[b_{i,j} | k \leq i, j \in Z/n].$$

Notice that $d_0(t_i^{p^j} \cdot t_{i_1}^{p^{j_1}}) = -t_i^{p^j} \otimes t_{i_1}^{p^{j_1}} - t_{i_1}^{p^{j_1}} \otimes t_i^{p^j}$, we get $h_{i,j}h_{i_1,j_1} = -h_{i_1,j_1}h_{i,j}$. In a similar way, one can proof that $h_{i,j} \cdot b_{i_1,j_1} = b_{i_1,j_1} \cdot h_{i,j}$ and $b_{i,j} \cdot b_{i_1,j_1} = b_{i_1,j_1} \cdot b_{i,j}$ (cf. [6] Lemma 3.4 and 3.8). \square

3. THE FIRST MAY DIFFERENTIALS

Now suppose $k \leq n$, then $s_0 \leq 2n$. From (2.2) and Lemma 2.4 one has

$$(3.1) \quad \begin{aligned} d_1(h_{i,j}) &= - \sum_{k \leq r \leq i-k} h_{r,j} h_{i-r,j+r} && \text{for } i \leq s_0 \\ d_1(h_{i,j}) &= b_{i-n,j+n-1} && \text{for } s_0 < i. \end{aligned}$$

Thus for $i > s_0 - n$, $b_{i,j}$ is the boundary of the first May differentials. Recall from [10] Theorem 4.3.22, in the cobar complex of BP/I_n one has

$$(3.2) \quad d(b_{i,j}) = \sum_{0 < r < i} \left(b_{r,j} \otimes t_{i-r}^{p^{r+j+1}} - t_r^{p^{j+1}} \otimes b_{i-r,r+j} \right)$$

Thus for $i \leq s_0 - n$, the first May differential $d_1(b_{i,j}) = 0$. This implies:

Theorem 3.3 *Let $k \leq n$ and s_0 be given in Lemma 2.4. The E_2 -term of the May spectral sequence is isomorphic to the cohomology of*

$$\tilde{E}_1^{*,*,*}S(n, k) = E[h_{i,j} | k \leq i \leq s_0, j \in Z/n] \otimes P[b_{i,j} | k \leq i \leq s_0 - n, j \in Z/n].$$

The first May differential are given by

$$\begin{aligned} d_1(h_{i,j}) &= - \sum_{k \leq r \leq i-k} h_{r,j} h_{i-r,r+j} && \text{for } i \leq s_0 \\ d_1(b_{i,j}) &= 0 && \text{for } k \leq i \leq s_0 - n \end{aligned}$$

At the prime $p = 2$, $s_0 = 2n$. The reduced May E_1 -term becomes

$$\tilde{E}_1^{*,*,*}S(n, k) = E[h_{i,j} | n < i \leq 2n, j \in Z/n] \otimes P[h_{i,j} | k \leq i \leq n, j \in Z/n]$$

and the first May differential of $h_{2n,j}$ is given by

$$d_1(h_{2n,j}) = - \sum_{k \leq i \leq 2n-k} h_{i,j} h_{2n-i,i+j} + h_{n,j+n-1}^2$$

Proof. We define a filtration in the May's E_1 -term

$$E_1^{*,*,*}S(n, k) = E[h_{i,j}|k \leq i, j \in Z/n] \otimes P[b_{i,j}|k \leq i, j \in Z/n]$$

as follows: for each $s \geq k$, define

$$F^s(n, k) = \begin{cases} E[h_{i,j}|k \leq i \leq s] & \text{for } k \leq s \leq n+k-1 \\ E[h_{i,j}|k \leq i \leq s] \otimes P[b_{i,j}|k \leq i \leq s-n] & \text{for } n+k-1 < s. \end{cases}$$

From (3.1), we see that for each $s \geq k$, $F^s(n, k)$ is a sub-complex of $E_1^{*,*,*}S(n, k)$ that satisfies

$$F^{s_0}(n, k) = \tilde{E}_1^{*,*,*}(n, k) = E[h_{i,j}|k \leq i \leq s_0, j \in Z/n] \otimes P[b_{i,j}|k \leq i \leq s_0-n, j \in Z/n]$$

and

$$F^k(n, k) \hookrightarrow F^{k+1}(n, k) \hookrightarrow \cdots \hookrightarrow F^s(n, k) \hookrightarrow F^{s+1}(n, k) \hookrightarrow \cdots \hookrightarrow E_1^{*,*,*}(n, k).$$

Indeed, for $s > n+k-1$,

$$F^s(n, k) = F^{s-1}(n, k) \bigotimes (E[h_{s,j}|j \in Z/n] \otimes P[b_{s-n,j}|j \in Z/n]).$$

For $s > s_0$ one has $d_1(h_{s,j}) = b_{s-n,j+n-1}$. Thus

$$E[h_{s,j}|j \in Z/n] \otimes P[b_{s-n,j}|j \in Z/n]$$

is a sub-complex of $F^s(n, k)$ whose cohomology is Z/p concentrated at dimensional 0. This implies

$$H^*F^{s_0}(n, k) \cong H^*F^{s_0+1}(n, k) \cong \cdots \cong H^*F^s(n, k) \cong \cdots \cong H^*E_1^{*,*,*}S(n, k)$$

At prime $p = 2$, $s_0 = \left\lfloor \frac{2 \times 2n}{2} \right\rfloor = 2n > n+k-1$. The first May differentials are deduced from (2.2). \square

As a corollary one can easily see that if $\frac{2pn+p-2}{2(p-1)} \leq n+k-1$, then the reduced May's E_1 -term becomes

$$\tilde{E}_1^{*,*,*}S(n, k) = E[h_{i,j}|k \leq i \leq n+k-1, j \in Z/n].$$

Theorem 3.4 *If $\frac{2pn+p-2}{2(p-1)} \leq n+k-1$, then the cohomology of $S(n, k)$ is of dimensional n^2 .*

4. THE HIGHER MAY DIFFERENTIALS IN THE MSS FOR $S(n, k)$

From (3.2) we see that the first non-trivial May differential of $b_{i,j}$ appears at

$$(4.1) \quad d_r(b_{i,j}) = \begin{cases} 0 & \text{if } i < 2k. \\ b_{i-k,j}h_{k,j+k+1} - h_{k,j+1}b_{i-k,j+k} & \text{if } i \geq 2k. \end{cases}$$

In [6] (2.10) and (2.11), a collapse theorem is given for the higher May differentials in the exterior part $E[h_{i,j}|i > 0, j \geq 0]$ of the MSS for the steenrod algebra A at odd primes. In this section, we will give a similar collapse theorem for the higher May differentials of $E_r^{*,*,*}S(n, k)$.

Let p be an odd prime. We define a Hopf algebra $T(n, k)$ as

$$(4.2) \quad T(n, k) = P[\xi_i|k \leq i \leq n+k-1].$$

The inner degree of ξ_i is defined to be $|\xi_i| = 2(p-1)(1+p+\dots+p^{i-1})$ and the structure map $\Delta : T(n, k) \longrightarrow T(n, k) \otimes T(n, k)$ acts on ξ_i by

$$\Delta(\xi_i) = \xi_i \otimes 1 + \sum_{k \leq r \leq i-k} \xi_r \otimes \xi_{i-r}^{p^r} + 1 \otimes \xi_i.$$

There is a Hopf algebra reduction homomorphism $\Phi : T(n, k) \rightarrow S(n, k)$ which send ξ_i to t_i . The image of Φ is $P[t_i | k \leq i \leq n+k-1] / (t_i^{p^n} - t_i)$ and $\text{Ker } \Phi$ is the idea generated by $(\xi_i^{p^n} - \xi_i)$. Further more the homomorphism Φ also induces homomorphism in cobar complexes and cohomologies

$$\Phi : \text{Ext}_{T(n, k)}^{s, *} (Z/p, Z/p) \longrightarrow \text{Ext}_{S(n, k)}^{s, *} (Z/p, Z/p)$$

Similar to that of definition 2.3, we set May filtration on $T(n, k)$ as

$$M(\xi_i^{p^j}) = 2i - 1$$

and let $F^{*, M}T(n, k)$ be the sub-module of $T(n, k)$ generated by the elements with May filtration $\leq M$. Then $E^{*, M}T(n, k) = F^{*, M}T(n, k) / F^{*, M-1}T(n, k)$ becomes a bigraded Hopf algebra with the structure of

$$E^{*, *}T(n, k) = \bigotimes T[\xi_i^{p^j} | k \leq i \leq n+k-1, j \geq 0]$$

and $\Delta(\xi_i^{p^j}) = \xi_i^{p^j} \otimes 1 + 1 \otimes \xi_i^{p^j}$.

Consider the cobar construction $C^{s, *}T(n, k)$ of $T(n, k)$. Similarly for the generator $[\beta_1 | \beta_2 | \dots | \beta_s]$ of $C^{s, *}T(n, k)$ define its May filtration as

$$M([\beta_1 | \beta_2 | \dots | \beta_s]) = M(\beta_1) + M(\beta_2) + \dots + M(\beta_s)$$

and let $FC^{*, *, M}T(n, k)$ denote the sun-complex generated by elements with May filtration $\leq M$. We get a spectral sequence $\{E_r^{s, *, M}T(n, k), d_r\}$ with E_0 -term

$$E_0^{*, *, M}T(n, k) = FC^{*, *, M}T(n, k) / FC^{*, *, M-1}T(n, k)$$

which is isomorphic to the cobar complex of $E^{*, *}T(n, k)$. The E_1 -term of this spectral sequence is isomorphic to

$$(4.3) \quad E_1^{*, *, *}T(n, k) = E[h'_{i,j} | k \leq i \leq n+k-1, j \geq 0] \otimes P[b'_{i,j} | k \leq i \leq n+k-1, j \geq 0].$$

Noticed that the reduction map $\Phi : T(n, k) \rightarrow S(n, k)$ is May filtration preserving, it induces a homomorphism of May spectral sequences

$$\Phi : E_r^{*, *, *}T(n, k) \longrightarrow E_r^{*, *, *}S(n, k).$$

Theorem 4.4 *The reduction map $\Phi : T(n, k) \longrightarrow S(n, k)$ induces a homomorphism between May spectral sequences $\Phi : E_1^{*, *, *}T(n, k) \longrightarrow E_1^{*, *, *}S(n, k)$ which sends $h'_{i,j}$ and $b'_{i,j}$ to $h_{i,j}$ and $b_{i,j}$ respectively. It sends infinite cocycles of $E_r^{*, *, *}T(n, k)$ to that of $E_r^{*, *, *}S(n, k)$.*

Similar to [6] (2.10) and (2.11) we give a collapse theorem in the MSS for $T(n, k)$. To the generators $h'_{i,j}, b'_{i,j} \in E_1^{*, *, *}T(n, k)$ define their index as

$$SI(h'_{i,j}) = SI(b'_{i,j}) = i.$$

Given a monomial $g = x_1 x_2 \dots x_m \in E_1^{*, *, *}T(n, k)$ where each x_i is of the generators $h'_{i,j}$ or $b'_{i,j}$, define its sum of index as

$$(4.5) \quad SI(g) = SI(x_1) + SI(x_2) + \dots + SI(x_m).$$

For example the sum of index of $h'_{4,0} h'_{3,0} b'_{2,1}$ is 9

We use $s(x)$ to denote the homological dimension of x . Noticed that the May filtration of $h'_{i,j}$, $b'_{i,j}$ satisfies

$$\begin{aligned} M(h'_{i,j}) &= 2i - 1 = 2SI(h'_{i,j}) - 1 = 2SI(h'_{i,j}) - s(h'_{i,j}) \\ M(b'_{i,j}) &= p(2i - 1) > 2SI(b'_{i,j}) - 2 = 2SI(b'_{i,j}) - s(b'_{i,j}) \end{aligned}$$

we see that for the monomial $g = x_1 x_2 \cdots x_m \in E_1^{s,*,*}T(n, k)$ of homological dimension s , its May filtration satisfies

$$\begin{aligned} (4.6) \quad M(g) &= M(x_1) + M(x_2) + \cdots + M(x_m) \\ &\geq 2SI(x_1) - s(x_1) + 2SI(x_2) - s(x_2) + \cdots + 2SI(x_m) - s(x_m) \\ &= 2SI(g) - s \end{aligned}$$

and the equality holds if and only if g is a monomial in $E[h'_{i,j} | k \leq i \leq n + k - 1, j \geq 0]$.

Given an integer $t = 2(p - 1)(c_0 + c_1 p + \cdots + c_m p^m)$ with $0 \leq c_i < p$, we define its sum of degree as

$$Sd(t) = c_0 + c_1 + \cdots + c_m$$

and for an element $g \in E_1^{s,*,*}T(n, k)$, express its inner degree $|g|$ as $|g| = 2(p - 1)(c_0 + c_1 p + \cdots + c_m p^m)$, where $0 \leq c_i < p$ and define its sum of degree to be

$$(4.7) \quad Sd(g) = Sd(|g|) = c_0 + c_1 + \cdots + c_m.$$

Then from

$$\begin{aligned} |h'_{i,j}| &= 2(p - 1)(p^j + p^{j+1} + \cdots + p^{i+j-1}) \\ |b'_{i,j}| &= 2(p - 1)(p^{j+1} + p^{j+2} + \cdots + p^{i+j}) \end{aligned}$$

we see that $SI(h'_{i,j}) = Sd(h'_{i,j})$, $SI(b'_{i,j}) = Sd(b'_{i,j})$. But for the reason of the p -adic numbers one has

$$(4.8) \quad SI(x_1 x_2 \cdots x_s) \geq Sd(x_1 x_2 \cdots x_m).$$

Theorem 4.9 *In the May spectral sequence for $T(n, k)$,*

(1) *If the inner degree $t = 2(p - 1)(c_0 + c_1 p + \cdots + c_m p^m)$ and the May filtration*

$$M < 2Sd(t) - s = 2(c_0 + c_1 + \cdots + c_m) - s,$$

then the May's E_1 -term $E_1^{s,t,M}T(n, k) = 0$.

(2) *If a cocycle $g \in E[h'_{i,j} | k \leq i \leq n + k - 1, j \geq 0]$ in the exterior part of May's E_1 -term satisfies $SI(g) = Sd(g)$, then it is an infinite cocycle in the MSS for $T(n, k)$ and $\Phi(g)$ is an infinite cocycle in the MSS for $S(n, k)$.*

Proof. (1) follows from (4.6) and (4.8).

Suppose $g \in E[h'_{i,j} | k \leq i \leq n + k - 1, j \geq 0]$ is a cocycle in the exterior part of May's E_1 -term $E_1^{s,t,M}T(n, k)$ that satisfies $SI(g) = Sd(g)$. Then its May filtration $M = 2SI(g) - s = 2Sd(t) - s$. Consider the higher May differentials

$$d_r : E_r^{s,t,M}T(n, k) \rightarrow E_r^{s+1,t,M-r}T(n, k),$$

we see that $M - r < 2Sd(t) - (s + 1)$ for $r > 1$. Thus the target $E_1^{s+1,t,M-r}T(n, k)$ and then $E_r^{s+1,t,M-r}T(n, k)$ is zero. \square

Example Let $p \geq 5$. The E_2 -term of the May spectral sequence for $H^{*,*}S(4, 2)$ is isomorphic to the homology of

$$E[h_{2,j}, h_{3,j}, h_{4,j}, h_{5,j} | j \in \mathbb{Z}/4]$$

with first May differentials

$$\begin{aligned} d_1(h_{2,j}) &= 0, & d_1(h_{3,j}) &= 0 \\ d_1(h_{4,j}) &= h_{2,j}h_{2,j+2}, & d_1(h_{5,j}) &= h_{2,j}h_{3,j+2} + h_{3,j}h_{2,j+3}. \end{aligned}$$

So $h_{5,0}h_{4,0}h_{3,0}h_{2,0}$ is a cohomology class in May's E_2 -term.

To prove that $h_{5,0}h_{4,0}h_{3,0}h_{2,0}$ is a infinite cocycle in the MSS $E_r^{4,*}S(4, 2)$, consider the MSS for $T(4, 2)$. $h_{5,0}h_{4,0}h_{3,0}h_{2,0}$ is a 4-dimensional cocycle in the exterior part of May's E_1 -term $E_1^{4,t,M}T(4, 2)$.

$$\deg(h'_{5,0}h'_{4,0}h'_{3,0}h'_{2,0}) = 2(p-1)(4+4p+3p^2+2p^3+p^4),$$

$$SI(h'_{5,0}h'_{4,0}h'_{3,0}h'_{2,0}) = 14 = Sd(h'_{5,0}h'_{4,0}h'_{3,0}h'_{2,0}).$$

Thus it is an infinite cocycle in the MSS for $T(4, 2)$ and $h_{5,0}h_{4,0}h_{3,0}h_{2,0} = \Phi(h'_{5,0}h'_{4,0}h'_{3,0}h'_{2,0})$ is an infinite cocycle in the MSS for $S(4, 2)$.

5. THE COHOMOLOGY OF $S(n, n)$ AT $p = 2$ AND OF $S(3, 2)$ AT $p = 3$

As an application of Theorem 3.3, we will compute $H^{*,*}S(n, n)$ at $p = 2$, $H^{*,*}S(3, 2)$ at prime $p = 3$ and $H^{*,*}S(4, 2)$ at prime $p \geq 5$ in this section.

5.1. The cohomology of $S(n, n)$ at prime two. Consider the cohomology of $S(n, n)$ at $p = 2$. The reduced Mays E_1 -term becomes

$$\tilde{E}_1^{*,*,*}S(n, n) = P[h_{n,j} | j \in \mathbb{Z}/n] \otimes E[h_{s,j} | n < s \leq 2n, j \in \mathbb{Z}/n]$$

(cf. Theorem 3.3). Noticed that the only non-trivial first May differential is

$$(5.1) \quad d_1(h_{2n,j}) = h_{n,j-1}^2 + h_{n,j}^2,$$

We see that the E_2 -term is the tensor product of $E[h_{s,j} | n < s < 2n]$ and the cohomology of

$$\{P[h_{n,j} | j \in \mathbb{Z}/n] \otimes E[h_{2n,j} | j \in \mathbb{Z}/n], d_1\}.$$

Lemma 5.2 *The May's E_2 -term $E_2^{*,*,*}S(n, n)$ at $p = 2$ is isomorphic to the tensor product of $E[h_{s,j} | n < s < 2n, j \in \mathbb{Z}/n]$ and $E[h_{n,j}, \rho_{2n} | j \in \mathbb{Z}/n] \otimes P[h_{n,n-1}]$, where $\rho_{2n} = \sum_{0 \leq j < n} h_{2n,j}^2$ and $h_{n,j}^2 = h_{n,n-1}^2$.*

Proof. We define $b_{n,j} = h_{n,j}^2 + h_{n,j+1}^2$ for $0 \leq j \leq n-2$ and define $b_{n,n-1} = h_{n,n-1}^2$. It is easy to see that $P[h_{n,j} | j \in \mathbb{Z}/n]$ could be divided as the tensor product of $P[b_{n,j} | 0 \leq j < n]$ and $E[h_{n,j} | j \in \mathbb{Z}/n]$ as $\mathbb{Z}/2$ -modules.

$$P[h_{n,j} | 0 \leq j \leq n-1] = P[b_{n,j} | 0 \leq j \leq n-1] \otimes E[h_{n,j} | 0 \leq j \leq n-1].$$

From (5.1) we see that

$$d_1(h_{2n,j}) = \begin{cases} b_{n,j-1} & \text{if } 1 \leq j < n \\ \sum_{0 \leq i < n-2} b_{n,i} & \text{if } j = n. \end{cases}$$

The cohomology of $\{P[h_{n,j} | j \in \mathbb{Z}/n] \otimes E[h_{2n,j} | j \in \mathbb{Z}/n], d_1\}$ is isomorphic to the tensor product of $E[h_{n,j} | j \in \mathbb{Z}/n]$ and the cohomology of

$$P[b_{n,j} | 0 \leq j < n] \otimes E[h_{2n,j} | j \in \mathbb{Z}/n].$$

The generator of $P[b_{n,j}|j \in Z/n]$ are denoted as

$$b_{n,i_1}^{s_1} b_{n,i_2}^{s_2} \cdots b_{n,i_m}^{s_m}$$

subject to $s_i > 0$, $0 \leq i_1 < i_2 < \cdots < i_m < n$ and the generators of $E[h_{2n,j}|j \in Z/n]$ are denoted as

$$h_{2n,j_k} \cdots h_{2n,j_2} h_{2n,j_1}$$

subject to $n \geq j_k > \cdots > j_2 > j_1 > 0$.

For the generators of $E[h_{2n,j}|0 < j \leq n] \otimes P[b_{n,j}|0 \leq j < n]$ described as above, one has

(1) For $j_1 > i_1 + 1$,

$$h_{2n,j_k} \cdots h_{2n,j_2} h_{2n,j_1} b_{n,i_1}^{s_1} b_{n,i_2}^{s_2} \cdots b_{n,i_m}^{s_m}$$

is the leading term of the first May differential

$$d_1(h_{2n,j_k} \cdots h_{2n,j_2} h_{2n,j_1} h_{2n,i_1-1} b_{n,i_1}^{s_1-1} b_{n,i_2}^{s_2} \cdots b_{n,i_m}^{s_m}).$$

While for $i_1 < n - 1$,

$$b_{n,i_1}^{s_1} b_{n,i_2}^{s_2} \cdots b_{n,i_m}^{s_m}$$

is the leading term of the first May differential

$$d_1(h_{2n,i_1+1} b_{n,i_1}^{s_1-1} b_{n,i_2}^{s_2} \cdots b_{n,i_m}^{s_m}).$$

(2) For $j_1 \leq i_1 + 1$ and $j_1 < n$, the leading term of the first May differential

$$d_1(h_{2n,j_k} \cdots h_{2n,j_2} h_{2n,j_1} b_{n,i_1}^{s_1} b_{n,i_2}^{s_2} \cdots b_{n,i_m}^{s_m})$$

is

$$h_{2n,j_k} \cdots h_{2n,j_2} h_{2n,j_1-1} b_{n,i_1}^{s_1} b_{n,i_2}^{s_2} \cdots b_{n,i_m}^{s_m}.$$

(3) For $j_1 = n = i_1 + 1$,

$$d_1(h_{2n,n} b_{n,n-1}^{s_1} \cdot x) = \sum_{i=0}^{n-2} b_{n,i} b_{n,n-1}^{s_1} \cdot x = d_1\left(\sum_{i=0}^{n-2} h_{2n,i+1} b_{n,n-1}^{s_1} \cdot x\right)$$

Thus the cohomology of $E[h_{2n,j}|j \in Z/n] \otimes P[b_{n,j}|j \in Z/n]$ is isomorphic to $E[\rho_{2n}|j \in Z/n] \otimes P[b_{n,n-1}]$, where $\rho_{2n} = \sum_{0 \leq i < n} h_{2n,i}$. The Lemma follows. \square

Theorem 5.3 *The May E_∞ -term $E_\infty^{*,*}S(n, n)$ is isomorphic to its E_2 -term. Thus the cohomology of $S(n, n)$ at prime 2 isomorphic to the tensor product of $E[h_{s,j}|n < s < 2n]$ and $E[h_{n,j}, \rho_{2n}|j \in Z/p] \otimes P[h_{n,0}]$*

Proof. It is easy to see from (2.2) that for $n \leq s < 2n$, $h_{s,j}$ is an infinite cocycle. From $d(t_{2n} + t_{2n}^2 + \cdots + t_{2n}^{2^{n-1}}) = 0$ we get the infinite cocycle ρ_{2n} . The Theorem follows. \square

5.2. The cohomology of $S(3, 2)$ at prime 3. Now consider the cohomology of $S(3, 2)$ at prime $p = 3$. From Lemma 2.4 we see that the $s_0 = 4$. Thus from Theorem 3.3 we see that the reduced May's E_1 -term is

$$\tilde{E}_1^{*,*}S(3, 2) = E[h_{2,j}, h_{3,j}, h_{4,j}|j \in Z/3]$$

and the first May differentials are given by

$$(5.4) \quad \begin{aligned} d_1(h_{2,j}) &= 0 & d_1(h_{3,j}) &= 0 & \text{and} \\ d_1(h_{4,j}) &= -h_{2,j}h_{2,j+2}. \end{aligned}$$

The May's E_2 -term is isomorphic to

$$E_2^{*,*,*}S(3, 2) = H^{*,*,*}(E[h_{2,j}, h_{4,j}|j \in Z/3], d_1) \otimes E[h_{3,j}|j \in Z/3]$$

Lemma 5.5 *The May E_2 -term for $H^*S(3, 2)$ at $p = 3$ has Poincare series $(x^6 + 3x^5 + 6x^4 + 9x^3 + 6x^2 + 3x + 1)(x + 1)^3$. It is the tensor product of $E[h_{3,j}|j \in Z/3]$ and the $Z/3$ module \mathcal{C} generated by the following element.*

The generators of \mathcal{C}							
Dimension	0	1	2	3	4	5	6
Generators	1	$h_{2,j}$	g_j k_j	l_j l'_j $k_j h_{2,j}$	$g_j g_{j+1}$ $k_j k_{j+1}$	$g_j l_{j+1}$	A

where $j \in Z/3$, $g_j = h_{4,j}h_{2,j}$, $k_j = h_{4,j}h_{2,j+2}$, $l_j = h_{4,j}h_{4,j+1}h_{2,j}$ and

$$l'_j = h_{4,j}h_{4,j+1}h_{2,j+1} + h_{4,j+1}h_{4,j+2}h_{2,j}$$

$$A = h_{4,0}h_{4,1}h_{4,2}h_{2,0}h_{2,1}h_{2,2} = -g_0g_1g_2.$$

Proof. From (5.4), it is easy to see that $d_1(h_{4,j}h_{2,j}) = 0$, $d_1(h_{4,j}h_{2,j+2}) = 0$ and from $d_1(h_{4,j+1}) = h_{2,j+1}h_{2,j+3} = h_{2,j+1}h_{2,j}$ we see that $d_1(h_{4,j}h_{4,j+1}h_{2,j}) = 0$. These gives the cohomology classes g_j , k_j and l_j . From

$$d_1(h_{4,j}h_{4,j+1}h_{2,j+1}) = -h_{2,j}h_{2,j+2}h_{4,j+1}h_{2,j+1} = -h_{4,j+1}h_{2,j}h_{2,j+1}h_{2,j+2}$$

$$d_1(h_{4,j+1}h_{4,j+2}h_{2,j}) = h_{4,j+1}h_{2,j+2}h_{2,j+4}h_{2,j} = h_{4,j+1}h_{2,j}h_{2,j+1}h_{2,j+2}$$

we get l'_j . A routine computation shows that $H^*(E[h_{2,j}, h_{4,j}|j \in Z/3]) = \mathcal{C}$. \square

Theorem 5.6 *The May E_2 -term for $H^{*,*}S(3, 2)$ at $p = 3$ is the E_∞ -term, thus $H^{*,*}S(3, 2)$ is the tensor product of $E[h_{3,j}|j \in Z/3]$ and \mathcal{C} .*

Proof. It is easy to see that $h_{2,j}$ and $h_{3,j}$ are infinite cycles. To prove that all the higher May differentials are trivial, consider the May filtration of each generator in the E_2 -term and the differentials

$$d_r : E_r^{s,t,M}S(3, 2) \longrightarrow E_r^{s+1,t,M-r}S(3, 2).$$

One has

$$\begin{aligned} g_j &\in E_2^{2,*,10}S(3, 2) & k_j &\in E_2^{2,*,10}S(3, 2) \\ l_j &\in E_2^{3,*,17}S(3, 2) & l'_j &\in E_2^{3,*,17}S(3, 2) \\ h_{2,j} &\in E_2^{1,*,3}S(3, 2) & h_{3,j} &\in E_2^{1,0,5}S(3, 2). \end{aligned}$$

The May filtration of g_j and k_j are 10. Beside, it is easy to check that each generator in the 3rd dimension $E_2^{3,*,*}S(3, 2)$ listed as below

$$l_j, \quad l'_j, \quad k_j h_{2,j}, \quad g_j h_{3,i}, \quad k_j h_{3,i}, \quad h_{3,i} h_{3,j} h_{2,k}, \quad h_{3,0} h_{3,1} h_{3,2}$$

has May filtration ≥ 10 . Thus g_j and k_j are infinite cycles. Similarly one can prove that l_j and l'_j are infinite cycles. This complete the proof. \square

5.3. The cohomology of $S(4, 2)$ at the primes $p > 3$. In this case, $s_0 = 5$ and the reduced May's E_1 -term is

$$\tilde{E}_1^{*,*,*} S(4, 2) = E[h_{2,j}, h_{3,j}, h_{4,j}, h_{5,j} | j \in Z/4].$$

To compute the E_2 -term, we set a filtration on the exterior algebra $E[h_{2,j}, h_{3,j}, h_{4,j}, h_{5,j} | j \in Z/4]$ as follows:

$$F^k = \bigoplus_{0 \leq r \leq k} Z/p\{h_{5,j_1} \cdots h_{5,j_r}\} \otimes E[h_{2,j}, h_{3,j}, h_{4,j} | j \in Z/4]$$

where $h_{5,j_1} \cdots h_{5,j_r}$'s are the generators of the r -dimensional module of the exterior algebra $E[h_{5,j} | j \in Z/4]$. This filtration gives rise to a spectral sequence with

$$E_0^k = F^k / F^{k-1} = Z/p\{h_{5,j_1} \cdots h_{5,j_k}\} \otimes E[h_{2,j}, h_{3,j}, h_{4,j} | j \in Z/4].$$

The E_1 -term of this spectral sequence is

$$E_1^k = Z/p\{h_{5,j_1} \cdots h_{5,j_r}\} \otimes H^* E[h_{2,j}, h_{3,j}, h_{4,j} | j \in Z/4],$$

and the differentials are given by

$$d_r : E_r^k \longrightarrow E_r^{k-r}.$$

By a routine computation, we get

Theorem 5.7 *The cohomology of $E[h_{2,j}, h_{3,j}, h_{4,j} | j \in Z/4]$ is the tensor product of $E[h_{3,j}, \rho_0, \rho_1]$ and \aleph , where*

$$\rho_0 = h_{4,0} + h_{4,2}, \quad \rho_1 = h_{4,1} + h_{4,3},$$

and \aleph is the direct sum of the modules generated by the following cohomology classes:

$$\begin{array}{llll} 1; & h_{2,j}; & e_j = h_{2,j}h_{2,j+1}, & g_j = h_{4,j}h_{2,j}; \\ h_{2,j}g_{j+1}, & h_{2,j}g_{j+2}, & h_{2,j}g_{j+3}; & \\ g_jg_{j+1}, & e_jg_{j+2}; & h_{2,j}g_{j+1}g_{j+2}; & e_0g_2g_3 \end{array}$$

with $j \in Z/4$. Beside, we also have the following relations:

$$h_{2,i}h_{2,i+2} = 0, \quad h_{2,i}g_{i+2} = h_{2,i+2}g_i, \quad h_{2,i}g_{i+2}g_{i+3} = h_{2,i+2}g_{i+3}g_i.$$

With the add of a personal computer, we compute that

Theorem 5.8 *The cohomology of the exterior algebra $E[h_{2,j}, h_{3,j}, h_{4,j}, h_{5,j}]$ has Poincaré series*

$$(1+t)^4(1+6t+18t^2+59t^3+92t^4+176t^5+161t^6+176t^7+92t^8+59t^9+18t^{10}+6t^{11}+t^{12}).$$

The ranks at each cohomological dimension are listed as

$$\begin{array}{cccccccccccc} 0, & 1, & 2, & 3, & 4, & 5, & 6, & 7, & 8, & 9, & \cdots & 16, \\ 1, & 10, & 48, & 171, & 461, & 976, & 1671, & 2303, & 2558, & 2303, & \cdots & 1 \end{array}$$

From the collapse Theorem 4.9, we claim that the MSS for the cohomology of $S(4, 2)$ collapse at E_2 -term.

REFERENCES

- [1] J. F. Adams, *Stable Homotopy and Generalised Homology*. University of Chicago Press, Chicago 1974.
- [2] H. Cartan and S. Eilenberg, *Homological Algebra*, Princeton University Press 1956.
- [3] P. Goerss, H. W. Henn, M. Mahowald and C. Rezk, A resolution of the $K(2)$ -local sphere at the prime 3, *Ann. Math.* **162** (2005) 777-822.
- [4] M. Hazewinkel, A universal formal group law and complex cobordism, *Bull. A. M. S.* **81** (1975), 930-933.
- [5] H. W. Henn, Centralization of abelian p -subgroups and mod- p cohomology of profinite groups, *Duck Math. J.* **91** (1998), 561-585.
- [6] S. Nave, Lee, The Smith-Toda complex $V((p+1)/2)$ does not exist, *Ann. Math.* **171** (2010) 491-509.
- [7] X. Liu and X. Wang, A four-filtered May spectral sequence and its applications, *Acta. Math. Sin. (English Ser.)* **24** (2008) 1507-1524.
- [8] J. P. May, The cohomology of restricted Lie algebras and of Hopf algebras. *J. Algebra* **3** (1966) 123-146.
- [9] J. P. May, The cohomology of restricted Lie algebras and of Hopf algebras; application to the Steenrod algebra. *Princeton Univ., 1964 Thesis*.
- [10] H. Miller, D. C. Ravenel and S. Wilson, Periodic phenomena in the Adams-Novikov spectral sequence *Ann. of Math.* **106** (1977), 469-516.
- [11] D. C. Ravenel, *Complex Cobordism and Stable Homotopy Groups of Spheres*. Academic Press, New York (1986).
- [12] D. C. Ravenel, *Nilpotence and Periodicity in Stable Homotopy Theory*, Ann. of Math. Studies **128**, Princeton Univ. Press, Princeton, NJ, 1992.
- [13] D. C. Ravenel, The cohomology of the Morava stabilizer algebras *Math. Z.* **152** (1977), 287-297.
- [14] D. C. Ravenel, Localization with respect to certain periodic homology theories. *Amer. J. Math.* **106** (1984) 351-414.
- [15] Shimomura, K. and Tokashiki, S., The cohomology of $S(n, n-1)$ relevant to the Morava Stabilizer algebra at odd prime, *Kochi Journal of Mathematics* **7** (2012), 109-118.
- [16] K. Shimomura and X. Wang, The homotopy groups $\pi_*(L_2S^0)$ at the prime 3, *Topology* **34** (2002), 1183-1198.
- [17] K. Shimomura and X. Wang, The Adams-Novikov E_2 -term for $\pi_*(L_2S^0)$ at the prime 2, *Math. Z.* **241** (2002) 271-311.
- [18] Tangora, M. C. On the cohomology of the Steenrod algebra, *Math. Z.* **116** (1970) 18-64.
- [19] X. Wang, $\pi_*(L_2T(1)/(v_1))$ and its applications in computing $\pi_*(L_2T(1))$ at the prime two, *Forum Math.* **19** (2007), 127-147.

SCHOOL OF MATHEMATICAL SCIENCE, NANKAI UNIVERSITY, TIANJIN 300071, P. R. CHINA
E-mail address: chenlimanstar1@163.com

SCHOOL OF MATHEMATICAL SCIENCE AND LPMC, NANKAI UNIVERSITY, TIANJIN 300071, P. R. CHINA
E-mail address: xjwang@nankai.edu.cn

DEPARTMENT OF MATHEMATICS & INSTITUTE OF MATHEMATICS AND INTERDISCIPLINARY SCIENCE, CAPITAL NORMAL UNIVERSITY, BEIJING 100048, P. R. CHINA
E-mail address: zhaoxve@mail.cnu.edu.cn