

Laurent phenomenon algebras and the discrete BKP equation

Naoto Okubo

Graduate School of Mathematical Sciences, the University of Tokyo,
3-8-1 Komaba, Tokyo 153-8914, Japan

Abstract

We construct the Laurent phenomenon algebras the cluster variables of which satisfy the discrete BKP equation and other difference equations obtained by its reduction. These Laurent phenomenon algebras are constructed from seeds with a generalization of mutation-period property. We show that a reduction of a seed corresponds to a reduction of a difference equation.

1 Introduction

In this article, we deal with the Laurent phenomenon algebras introduced by Lam and Pylyavskyy [9]. The Laurent phenomenon algebra is a generalization of the cluster algebra [6] with Laurent phenomenon of cluster variables. A Laurent phenomenon algebra is a commutative ring described by cluster variables. A generating set of the Laurent phenomenon algebra is defined by mutation, which is a transformation of a seed consisting of a set of cluster variables and exchange polynomials. The Laurent phenomenon is the property that all cluster variables obtained by mutations of an initial seed are Laurent polynomials in cluster variables of the initial seed. It is known that cluster variables of suitable cluster algebras can satisfy the Hirota-Miwa equation [3], the discrete KdV equation [1], the discrete Toda equation [2], Soms-4, and Somos-5 [4], when the initial seed includes suitable periodic quivers [7]. Difference equations obtained by a cluster algebra are restricted to the form:

$$x'_k x_k = \prod_{i \in I} x_i^{a_i} + \prod_{j \in J} x_j^{b_j}. \quad (1.1)$$

However, the discrete BKP equation

$$x_{n+1}^{m+1, l+1} x_n^{m, l} = x_n^{m+1, l+1} x_{n+1}^{m, l} + x_{n+1}^{m+1, l} x_n^{m, l+1} + x_{n+1}^{m, l+1} x_n^{m+1, l} \quad (1.2)$$

does not have the form given by (1.1). Cluster variables of a Laurent phenomenon algebra obtained from a mutation of an initial seed can satisfy the difference equations of more general form as

$$x'_k x_k = P \in \mathbb{Z}[x_i | i \in I]. \quad (1.3)$$

It is known that cluster variables of a Laurent phenomenon algebra can satisfy the Somos-6 [4] and some related difference equations, when the initial seed includes suitable exchange

polynomials [9, 10]. These seeds have the property called mutation-period. This property is an analogue of the mutation-periodic quiver of cluster algebras. Several results concerning period-1 seeds have been reported in [10]. In this paper, we construct the Laurent phenomenon algebras whose cluster variables satisfy the discrete BKP equation [3], the Somos-7, and several other difference equations. The seed which gives the discrete BKP equation has infinite rank. This seed has the property of a generalization of the mutation-period. The seeds which give the Somos-6, Somos-7, and related 2-dimensional difference equations are obtained from reductions of the seed that gives the discrete BKP equation. This reduction corresponds to a reduction from the discrete BKP equation to these difference equations.

2 Laurent phenomenon algebras

In this section, we briefly explain the notion of Laurent phenomenon algebra which we use in the following sections.

2.1 Definition of Laurent phenomenon algebra

Let $\mathbf{x} = (x_0, x_1, \dots, x_{N-1})$ be an N -tuple of variables. Let $\mathbf{F} = (F_0, F_1, \dots, F_{N-1})$ be an N -tuple of polynomials in $\mathbb{Z}[x_0, x_1, \dots, x_{N-1}]$. We assume that these polynomials satisfy the following conditions:

- (LP1) $F_i \in \mathbb{Z}[x_0, x_1, \dots, x_{N-1}]$ is irreducible and is not divisible by any x_j .
- (LP2) F_i does not depend on x_i .

Each x_i is called a cluster variable and each F_i is called an exchange polynomial. The pair $t = (\mathbf{x}, \mathbf{F})$ is called a seed. We will often express a seed as

$$t = \{(x_0, F_0), (x_1, F_1), \dots, (x_{N-1}, F_{N-1})\}. \quad (2.1)$$

The number of cluster variables N in a seed t is called the rank of the seed t or the rank of the Laurent phenomenon algebra. For Laurent polynomials $F, G \in \mathbb{Z}[x_0^\pm, x_1^\pm, \dots, x_{N-1}^\pm]$ and a cluster variable x_i , let $F|_{x_i \leftarrow G}$ be the polynomial in which we substituted x_i for G in F . A mutation is a particular transformation of seeds.

Definition 2.1 [9] *The mutation of the seed t at x_k is a transformation from t to t' ,*

$$\mu_k : t = \{(x_0, F_0), (x_1, F_1), \dots, (x_{N-1}, F_{N-1})\} \mapsto t' = \{(x'_0, F'_0), (x'_1, F'_1), \dots, (x'_{N-1}, F'_{N-1})\}, \quad (2.2)$$

that are defined via the following sequence of steps:

1. Let $\hat{F}_k \in \mathbb{Z}[x_0^\pm, x_1^\pm, \dots, x_{k-1}^\pm, x_{k+1}^\pm, \dots, x_{N-1}^\pm]$ be the unique Laurent polynomial satisfying following conditions:

- There exist $a_0, a_1, \dots, a_{N-1} \in \mathbb{Z}_{\leq 0}$ such that

$$\hat{F}_k = x_0^{a_0} x_1^{a_1} \dots x_{k-1}^{a_{k-1}} x_{k+1}^{a_{k+1}} \dots x_{N-1}^{a_{N-1}} F_k. \quad (2.3)$$

- For any $j \neq k$,

$$\hat{F}_k \Big|_{x_j \leftarrow F_j/x_j} \in \mathbb{Z} [x_0^\pm, x_1^\pm, \dots, x_{j-1}^\pm, x_j^\pm, x_{j+1}^\pm, \dots, x_{N-1}^\pm] \quad (2.4)$$

and is not divisible by F_j .

2. New cluster variable x'_i is defined as:

$$\begin{aligned} x'_i &= x_i \quad (i \neq k), \\ x'_k &= \hat{F}_k/x_k. \end{aligned} \quad (2.5)$$

3. If F_i does not depend on x_k , then we define $F'_i = F_i$. In the following steps, we assume that F_i depends on x_k .

4. We define the Laurent polynomial G_i by

$$G_i = F_i \Big|_{x_k \leftarrow \frac{\hat{F}_k|_{x_i \leftarrow 0}}{x'_k}}. \quad (2.6)$$

Note that we can show that if F_i depends on x_k , then $a_i = 0$. Therefore, $\hat{F}_k \Big|_{x_i \leftarrow 0}$ is well defined.

5. We define H_i to be the result of removing all common factors in

$$\mathbb{Z}[x_0, x_1, \dots, x_{k-1}, x_{k+1}, \dots, x_{i-1}, x_{i+1}, \dots, x_{N-1}] \text{ with } \hat{F}_k \Big|_{x_i \leftarrow 0} \text{ from } G_i.$$

6. Let M_i be the unique Laurent monomial of $x'_0, x'_1, \dots, x'_{N-1}$ satisfying following conditions:

- $M_i H_i \in \mathbb{Z}[x'_0, x'_1, \dots, x'_{N-1}]$.
- $M_i H_i$ is not divisible by any x'_j .
- $M_i H_i$ does not depend on x'_i .

New exchange polynomial F'_i is defined as $F'_i = M_i H_i$.

Proposition 2.2 [9] For any $k \in \{0, 1, \dots, N-1\}$ and any seed t , it holds that $\mu_k^2(t) = t$.

Proposition 2.3 [9] Suppose that we mutate at x_k . Then F'_i depends on x'_k if and only if F_i depends on x_k .

Definition 2.4 [9] Let us fix a seed $t = (\mathbf{x}, \mathbf{F})$. This seed is called an initial seed. Let $X(t)$ be the set of all the cluster variables obtained by iterative mutations to the initial seed t . Let $\mathcal{A}(t)$ be a Laurent phenomenon algebra defined as

$$\mathcal{A}(t) = \mathbb{Q}[x \in X(t)] \subset \mathbb{Q}(\mathbf{x}). \quad (2.7)$$

Theorem 2.5 [9] Let $t = (\mathbf{x}, \mathbf{F})$ be a initial seed. If $x \in X(t)$, then $x \in \mathbb{Z}[\mathbf{x}^\pm]$.

2.2 Period-1 seeds

For a polynomial

$$f(x_0, x_1, \dots, x_{N-1}) \in \mathbb{Z}[x_0, x_1, \dots, x_{N-1}] \quad (2.8)$$

and for $j \geq 1$, we define

$$u^j(f) = f(x_j, x_{j+1}, \dots, x_{j+N-1}) \in \mathbb{Z}[x_j, x_{j+1}, \dots, x_{j+N-1}]. \quad (2.9)$$

We also abbreviate $u := u^1$.

Definition 2.6 [10] *A seed*

$$t = \{(x_0, F_0), (x_1, F_1), \dots, (x_{N-1}, F_{N-1})\} \quad (2.10)$$

is called a period-1 seed if

$$\mu_0(t) = \{(x_N, u(F_{N-1})), (x_1, u(F_0)), (x_2, u(F_1)), \dots, (x_{N-1}, u(F_{N-2}))\} \quad (2.11)$$

holds, where $x_N = x'_0$ is the new cluster variable obtained by the mutation at x_0 .

We assume that

$$t_0 = \{(x_0, F_0), (x_1, F_1), \dots, (x_{N-1}, F_{N-1})\} \quad (2.12)$$

is a period-1 seed. We inductively define the seed t_n by $t_n = \mu_{n-1}(t_{n-1})$. The new cluster variable x'_n in $\mu_n(t_n)$ is expressed as x_{n+N} . Hereafter, we write this process as $x_n \rightarrow x_{n+N}$. We have

$$t_n = \{(x_n, u^n(F_0)), (x_{n+1}, u^n(F_1)), \dots, (x_{n+N-1}, u^n(F_{N-1}))\}. \quad (2.13)$$

If an exchange polynomial of a period-1 seed satisfies $\hat{F}_0 = F_0$, then the cluster variables satisfy $x_{n+N}x_n = u^n(F_0)$. In the following sections, all exchange polynomials satisfy $\hat{F}_i = F_i$. We put $F_0 = f(x_1, x_2, \dots, x_{N-1})$. All cluster variables satisfy the difference equation

$$x_{n+N}x_n = f(x_{n+1}, x_{n+2}, \dots, x_{n+N-1}). \quad (2.14)$$

Several results about a period-1 seed have already known.

- All the rank 2 or 3 period-1 seeds have been obtained [10].
- An example of a rank 6 period-1 seed has been obtained [9]. The cluster variables of this seed satisfy the difference equation

$$x_{n+6}x_n = x_{n+5}x_{n+1} + x_{n+4}x_{n+2} + x_{n+3}^2. \quad (2.15)$$

This difference equation is called Somos-6.

- Several rank N period-1 seeds have been obtained [10]. For example, the cluster variables of these seeds satisfy the following difference equations:

$$x_{n+N}x_n = \prod_{i=1}^{N-1} x_{n+i}^{a_i} + 1 \quad (a_i \in \mathbb{Z}_{>0}), \quad (2.16)$$

$$x_{n+N}x_n = x_{n+1}x_{n+N-1} + A \sum_{i=1}^{N-1} x_{n+i} + B \quad (A, B \in \mathbb{Z}). \quad (2.17)$$

3 Seed of the discrete BKP equation

3.1 Seed of the discrete BKP equation and its mutation

In this section, we consider the infinite rank seed. For $i \leq j$ ($i, j \in \mathbb{Z}$), we put $[i, j] = \{i, i+1, \dots, j\}$. For a polynomial $P \in \mathbb{Z}[x_n^{m,l} | n, m, l \in \mathbb{Z}]$, let $s_i^{j,k}(P)$ be the polynomial in which we substituted all variables $x_n^{m,l}$ for $x_{n+i}^{m+j, l+k}$ in P . We take

$$t_0 = \left\{ \left(x_{i-2j-k}^{j-k,k}, F_{i-2j-k}^{j-k,k} \right) \mid i \in [0, 5], j, k \in \mathbb{Z} \right\} \quad (3.1)$$

as an initial seed, where exchange polynomials are defined as

$$\begin{aligned} F_0^{0,0} &= x_0^{1,1} x_1^{0,0} + x_0^{1,0} x_1^{0,1} + x_0^{0,1} x_1^{1,0}, \\ F_1^{0,0} &= x_0^{1,0} x_1^{0,1} x_2^{0,0} + x_0^{0,1} x_1^{1,0} x_2^{0,0} + x_0^{0,0} x_1^{1,0} x_2^{0,1} + x_0^{0,0} x_1^{0,1} x_2^{1,0}, \\ F_2^{0,0} &= x_1^{1,0} x_2^{-1,0} x_3^{0,1} x_3^{0,0} + x_1^{0,1} x_2^{-1,0} x_3^{1,0} x_3^{0,0} \\ &\quad + x_1^{0,0} x_2^{-1,1} x_3^{1,0} x_3^{0,0} + x_1^{0,0} x_2^{-1,0} x_3^{0,1} x_3^{1,0} + x_1^{0,0} x_2^{0,1} x_3^{1,0} x_3^{-1,0}, \\ F_3^{0,0} &= x_4^{-1,0} x_3^{1,0} x_3^{0,-1} x_2^{0,0} + x_4^{0,-1} x_3^{1,0} x_3^{-1,0} x_2^{0,0} \\ &\quad + x_4^{0,0} x_3^{1,-1} x_3^{-1,0} x_2^{0,0} + x_4^{0,0} x_3^{1,0} x_3^{0,-1} x_2^{-1,0} + x_4^{0,0} x_3^{0,-1} x_3^{-1,0} x_2^{1,0}, \\ F_4^{0,0} &= x_5^{-1,0} x_4^{0,-1} x_3^{0,0} + x_5^{0,-1} x_4^{-1,0} x_3^{0,0} + x_5^{0,0} x_4^{-1,0} x_3^{0,-1} + x_5^{0,0} x_4^{0,-1} x_3^{-1,0}, \\ F_5^{0,0} &= x_5^{-1,-1} x_4^{0,0} + x_5^{-1,0} x_4^{0,-1} + x_5^{0,-1} x_4^{-1,0}, \end{aligned} \quad (3.2)$$

and

$$F_{i-2j-k}^{j-k,k} = s_{-2j-k}^{j-k,k} (F_i^{0,0}). \quad (3.3)$$

For example, $F_0^{1,1}$ is defined as

$$F_0^{1,1} = F_{5-2 \cdot 2-1}^{2-1,1} = s_{-5}^{1,1} (F_5^{0,0}) = x_0^{0,0} x_{-1}^{1,1} + x_0^{0,1} x_{-1}^{1,0} + x_0^{1,0} x_{-1}^{0,1}. \quad (3.4)$$

We define $\tilde{u} := s_1^{0,0}$ and the set of cluster variables X_i as

$$X_i := \left\{ x_{i-2j-k}^{j-k,k} \mid j, k \in \mathbb{Z} \right\}. \quad (3.5)$$

A mutation $m_i^{j,k}$ ($i, j, k \in \mathbb{Z}$) denotes the mutation at $x_{i-2j-k}^{j-k,k}$. We define the iteration of the mutations ν_i^n ($i \in \mathbb{Z}, n \in \mathbb{Z}_{\geq 0}$) by

$$\begin{aligned} \nu_i^n &= m_i^{n-1,-1} \circ \dots \circ m_i^{1,-n+1} \circ m_i^{0,-n} \circ m_i^{-1,-n+1} \circ \dots \circ m_i^{-n+1,-1} \circ m_i^{-n,0} \\ &\quad \circ m_i^{-n+1,1} \circ \dots \circ m_i^{-1,n-1} \circ m_i^{0,n} \circ m_i^{1,n-1} \circ \dots \circ m_i^{n-1,1} \circ m_i^{n,0} \quad (n \neq 0), \\ \nu_i^0 &= m_i^{0,0}. \end{aligned} \quad (3.6)$$

This ν_i^n is the iteration of the mutations at each and all $x_{i-2j-k}^{j-k,k} \in X_i$ ($j+k=n$) just once. We define the iteration of the mutations $\tilde{\mu}_i$ ($i \in \mathbb{Z}$) by

$$\tilde{\mu}_i = \dots \circ \nu_i^2 \circ \nu_i^1 \circ \nu_i^0. \quad (3.7)$$

This $\tilde{\mu}_i$ is the iteration of the mutations at each and all $x \in X_i$ just once. We also use the notation $x \rightarrow y$ that means a new cluster variable x' is denoted by y , i.e., $y = x'$.

Proposition 3.1 We define the seed t_1 by $t_1 = \tilde{\mu}_0(t_0)$. When we put $x_{-2j-k}^{j-k,k} \rightarrow x_{1-2j-k}^{1+j-k,1+k}$, it holds that

$$t_1 = \left\{ \left(x_{1+i-2j-k}^{j-k,k}, \tilde{u} \left(F_{i-2j-k}^{j-k,k} \right) \right) \middle| i \in [0, 5], j, k \in \mathbb{Z} \right\}. \quad (3.8)$$

The new cluster variables $x_{1-2j-k}^{1+j-k,1+k}$ satisfy

$$x_{1-2j-k}^{1+j-k,1+k} x_{-2j-k}^{j-k,k} = F_{-2j-k}^{j-k,k}. \quad (3.9)$$

Proof If $F_n^{m,l}$ does not depend on $x_{-2j-k}^{j-k,k}$, then it holds $(F_n^{m,l})' = F_n^{m,l}$, when we mutate at $x_{-2j-k}^{j-k,k} \in X_0$. The dependence on $x \in X_0$ of the exchange polynomials is as follows:

$$\begin{aligned} F_{-2j-k}^{j-k,k} & \text{ does not depend on } x \in X_0. \\ F_{1-2j-k}^{j-k,k} & \text{ depends on only } x_{-2j-k}^{j-k,k} \in X_0. \\ F_{2-2j-k}^{j-k,k} & \text{ depends on only } x_{-2(j-1)-k}^{(j-1)-k,k} \in X_0. \\ F_{3-2j-k}^{j-k,k} & \text{ depends on only } x_{-2(j-1)-(k-1)}^{(j-1)-(k-1),k-1}, x_{-2(j-1)-k}^{(j-1)-k,k} \in X_0. \\ F_{4-2j-k}^{j-k,k} & \text{ depends on only } x_{-2(j-1)-(k-1)}^{(j-1)-(k-1),k-1} \in X_0. \\ F_{5-2j-k}^{j-k,k} & \text{ depends on only } x_{-2(j-2)-(k-1)}^{(j-2)-(k-1),k-1} \in X_0. \end{aligned} \quad (3.10)$$

For each $i \in [0, 5]$, we consider the change of $(x_{i-2j-k}^{j-k,k}, F_{i-2j-k}^{j-k,k})$ by the mutations of initial seed t_0 at each and all $x \in X_0$ just once.

- The case of $i = 0$

By (3.10), the exchange polynomials $F_{-2j-k}^{j-k,k}$ do not change by the mutations at $x \in X_0$. For each seed

$$\begin{aligned} \tilde{t} &= \left\{ \left(\tilde{x}_{i-2j-k}^{j-k,k}, \tilde{F}_{i-2j-k}^{j-k,k} \right) \middle| i \in [0, 5], j, k \in \mathbb{Z} \right\} \\ &= \left\{ \left(\tilde{x}_{-2j-k}^{j-k,k}, F_{-2j-k}^{j-k,k} \right), \left(\tilde{x}_{i-2j-k}^{j-k,k}, \tilde{F}_{i-2j-k}^{j-k,k} \right) \middle| i \in [1, 5], j, k \in \mathbb{Z} \right\} \end{aligned} \quad (3.11)$$

obtained by the mutations at $x \in X_0$, if $F_{-2j-k}^{j-k,k} \Big|_{\tilde{x}_n^{m,l} \leftarrow \tilde{F}_n^{m,l}/x}$ is not divisible by $\tilde{F}_n^{m,l}$ for any $(n, m, l) \neq (-2j-k, j-k, k)$, then it holds that $\hat{F}_{-2j-k}^{j-k,k} = F_{-2j-k}^{j-k,k}$. In fact, all exchange polynomials which appear below satisfy $\hat{F}_{-2j-k}^{j-k,k} = F_{-2j-k}^{j-k,k}$. By the mutation at $x_{-2j-k}^{j-k,k}$, the change of the cluster variable $x_{-2j-k}^{j-k,k}$ is

$$x_{-2j-k}^{j-k,k} \xrightarrow{\mu_{-2j-k}^{j-k,k}} x_{1-2j-k}^{1+j-k,1+k} = \hat{F}_{-2j-k}^{j-k,k} / x_{-2j-k}^{j-k,k} = F_{-2j-k}^{j-k,k} / x_{-2j-k}^{j-k,k} \quad (3.12)$$

and other cluster variables $x_{-2j'-k'}^{j'-k',k'}$ ($(j', k') \neq (j, k)$) do not change, where $\mu_n^{m,l}$ ($n, m, l \in \mathbb{Z}$) is the mutation at $x_n^{m,l}$. Therefore, we have

$$\begin{aligned} \left(x_{-2j-k}^{j-k,k}, F_{-2j-k}^{j-k,k} \right) & \rightarrow \left(x_{1-2j-k}^{1+j-k,1+k}, F_{-2j-k}^{j-k,k} \right) \\ &= \left(x_{6-2(j+2)-(k+1)}^{(j+2)-(k+1),k+1}, \tilde{u} \left(F_{5-2(j+2)-(k+1)}^{(j+2)-(k+1),k+1} \right) \right) \end{aligned} \quad (3.13)$$

by the mutations of initial seed t_0 at each and all $x \in X_0$ just once.

- The case of $i = 1$

By (3.10), the exchange polynomials $F_{1-2j-k}^{j-k,k}$ do not change until we execute the mutation at $x_{-2j-k}^{j-k,k}$. Let λ_1 and λ_2 be the iteration of mutations before and after the mutation at $x_{-2j-k}^{j-k,k}$ respectively. We have

$$\left(F_{1-2j-k}^{j-k,k}, F_{-2j-k}^{j-k,k}\right) \xrightarrow{\lambda_1} \left(F_{1-2j-k}^{j-k,k}, F_{-2j-k}^{j-k,k}\right). \quad (3.14)$$

We mutate at $x_{-2j-k}^{j-k,k}$ and calculate $\left(F_{1-2j-k}^{j-k,k}\right)'$ from $F_{1-2j-k}^{j-k,k}$ and $F_{-2j-k}^{j-k,k}$. It holds that

$$\left(F_{1-2j-k}^{j-k,k}, F_{-2j-k}^{j-k,k}\right) \xrightarrow{\mu_{-2j-k}^{j-k,k}} \left(\tilde{u}\left(F_{-2j-k}^{j-k,k}\right), F_{-2j-k}^{j-k,k}\right). \quad (3.15)$$

$\tilde{u}\left(F_{-2j-k}^{j-k,k}\right)$ does not depend on any $x \in X_0$. We have

$$\tilde{u}\left(F_{-2j-k}^{j-k,k}\right) \xrightarrow{\lambda_2} \tilde{u}\left(F_{-2j-k}^{j-k,k}\right). \quad (3.16)$$

The cluster variable $x_{1-2j-k}^{j-k,k}$ does not change by the mutations at $x \in X_0$. Therefore, we have

$$\left(x_{1-2j-k}^{j-k,k}, F_{1-2j-k}^{j-k,k}\right) \rightarrow \left(x_{1-2j-k}^{j-k,k}, \tilde{u}\left(F_{-2j-k}^{j-k,k}\right)\right) \quad (3.17)$$

by mutations of the initial seed t_0 at each and all $x \in X_0$ just once.

- The cases of $i = 2, 4, 5$

The cases for $i = 2, 4, 5$ are the same as the case of $i = 1$. It holds that

$$\left(x_{i-2j-k}^{j-k,k}, F_{i-2j-k}^{j-k,k}\right) \rightarrow \left(x_{i-2j-k}^{j-k,k}, \tilde{u}\left(F_{-1+i-2j-k}^{j-k,k}\right)\right) \quad (i = 2, 4, 5) \quad (3.18)$$

by the mutations of initial seed t_0 at each and all $x \in X_0$ just once.

- The case of $i = 3$

By (3.10), the exchange polynomials $F_{3-2j-k}^{j-k,k}$ do not change until we execute the mutation at $x_{3-2j-k}^{j-k,k-1}$ or $x_{2-2j-k}^{-1+j-k,k}$. We consider two cases in which the order of the mutations at $x_{3-2j-k}^{j-k,k-1}$ and $x_{2-2j-k}^{-1+j-k,k}$ is different from each other.

- Suppose that we mutate at $x_{3-2j-k}^{j-k,k-1}$ before the mutation at $x_{2-2j-k}^{-1+j-k,k}$. Let λ_1, λ_2 , and λ_3 be the iteration of mutations before the mutation at $x_{3-2j-k}^{j-k,k-1}$, after the mutation at $x_{3-2j-k}^{j-k,k-1}$ and before the mutation at $x_{2-2j-k}^{-1+j-k,k}$, and after the mutation at $x_{2-2j-k}^{-1+j-k,k}$ respectively. We have

$$\left(F_{3-2j-k}^{j-k,k}, F_{3-2j-k}^{j-k,k-1}, F_{2-2j-k}^{-1+j-k,k}\right) \xrightarrow{\lambda_1} \left(F_{3-2j-k}^{j-k,k}, F_{3-2j-k}^{j-k,k-1}, F_{2-2j-k}^{-1+j-k,k}\right). \quad (3.19)$$

We mutate at $x_{3-2j-k}^{j-k,k-1}$ and calculate $\left(F_{3-2j-k}^{j-k,k}\right)'$ from $F_{3-2j-k}^{j-k,k}$ and $F_{3-2j-k}^{j-k,k-1}$. Then we have

$$\begin{aligned} & \left(F_{3-2j-k}^{j-k,k}, F_{3-2j-k}^{j-k,k-1}, F_{2-2j-k}^{-1+j-k,k}\right) \\ & \xrightarrow{\mu_{3-2j-k}^{j-k,k-1}} \left(s_{-2j-k}^{j-k,k}(P_1), F_{3-2j-k}^{j-k,k-1}, F_{2-2j-k}^{-1+j-k,k}\right), \end{aligned} \quad (3.20)$$

$$P_1 := x_2^{-1,0} x_3^{1,0} x_4^{0,0} + x_2^{1,0} x_3^{-1,0} x_4^{0,0} + x_2^{0,0} x_3^{-1,0} x_4^{1,0} + x_2^{0,0} x_3^{1,0} x_4^{-1,0}. \quad (3.21)$$

However, $s_{-2j-k}^{j-k,k}(P_1)$ depends on only $x_{2-2j-k}^{-1+j-k,k} \in X_0$. Hence, $s_{-2j-k}^{j-k,k}(P_1)$ does not change before the mutation at $x_{2-2j-k}^{-1+j-k,k}$. We have

$$\left(s_{-2j-k}^{j-k,k}(P_1), F_{2-2j-k}^{-1+j-k,k}\right) \xrightarrow{\lambda_2} \left(s_{-2j-k}^{j-k,k}(P_1), F_{2-2j-k}^{-1+j-k,k}\right). \quad (3.22)$$

Mutating at $x_{2-2j-k}^{-1+j-k,k}$ and calculating $\left(s_{-2j-k}^{j-k,k}(P_1)\right)'$ from $s_{-2j-k}^{j-k,k}(P_1)$ and $F_{2-2j-k}^{-1+j-k,k}$, we find

$$\left(s_{-2j-k}^{j-k,k}(P_1), F_{2-2j-k}^{-1+j-k,k}\right) \xrightarrow{\mu_{2-2j-k}^{-1+j-k,k}} \left(\tilde{u}\left(F_{2-2j-k}^{j-k,k}\right), F_{2-2j-k}^{-1+j-k,k}\right). \quad (3.23)$$

Since $\tilde{u}\left(F_{2-2j-k}^{j-k,k}\right)$ does not depend on any $x \in X_0$, we have

$$\tilde{u}\left(F_{2-2j-k}^{j-k,k}\right) \xrightarrow{\lambda_3} \tilde{u}\left(F_{2-2j-k}^{j-k,k}\right). \quad (3.24)$$

– Suppose that we mutate at $x_{2-2j-k}^{-1+j-k,k}$ before the mutation at $x_{3-2j-k}^{j-k,k-1}$. Let λ_1, λ_2 , and λ_3 be the iteration of mutations before the mutation at $x_{2-2j-k}^{-1+j-k,k}$, after the mutation at $x_{2-2j-k}^{-1+j-k,k}$ and before the mutation at $x_{3-2j-k}^{j-k,k-1}$, and after the mutation at $x_{3-2j-k}^{j-k,k-1}$ respectively. We have

$$\left(F_{3-2j-k}^{j-k,k}, F_{3-2j-k}^{j-k,k-1}, F_{2-2j-k}^{-1+j-k,k}\right) \xrightarrow{\lambda_1} \left(F_{3-2j-k}^{j-k,k}, F_{3-2j-k}^{j-k,k-1}, F_{2-2j-k}^{-1+j-k,k}\right). \quad (3.25)$$

By mutation at $x_{2-2j-k}^{-1+j-k,k}$ and calculation of $\left(F_{3-2j-k}^{j-k,k}\right)'$ from $F_{3-2j-k}^{j-k,k}$ and $F_{2-2j-k}^{-1+j-k,k}$, we find that

$$\begin{aligned} &\left(F_{3-2j-k}^{j-k,k}, F_{3-2j-k}^{j-k,k-1}, F_{2-2j-k}^{-1+j-k,k}\right) \\ &\xrightarrow{\mu_{2-2j-k}^{-1+j-k,k}} \left(s_{-2j-k}^{j-k,k}(P_2), F_{3-2j-k}^{j-k,k-1}, F_{2-2j-k}^{-1+j-k,k}\right), \end{aligned} \quad (3.26)$$

$$\begin{aligned} P_2 := & x_2^{1,0} x_3^{-1,0} x_3^{0,-1} x_4^{0,0} + x_2^{0,1} x_3^{-1,0} x_3^{0,-1} x_4^{1,0} x_4^{0,0} \\ & + x_2^{0,0} x_3^{-1,0} x_3^{0,1} x_4^{1,-1} x_4^{0,0} + x_2^{0,0} x_3^{-1,1} x_3^{0,-1} x_4^{1,0} x_4^{0,0} \\ & + x_2^{0,0} x_3^{-1,0} x_3^{0,1} x_4^{1,0} x_4^{0,-1} + x_2^{0,0} x_3^{0,1} x_3^{0,-1} x_4^{1,0} x_4^{-1,0}. \end{aligned} \quad (3.27)$$

Since $s_{-2j-k}^{j-k,k}(P_2)$ depends on only $x_{3-2j-k}^{j-k,k-1} \in X_0$, $s_{-2j-k}^{j-k,k}(P_2)$ does not change before the mutation at $x_{3-2j-k}^{j-k,k-1}$. We have

$$\left(s_{-2j-k}^{j-k,k}(P_2), F_{3-2j-k}^{j-k,k-1}\right) \xrightarrow{\lambda_2} \left(s_{-2j-k}^{j-k,k}(P_2), F_{3-2j-k}^{j-k,k-1}\right). \quad (3.28)$$

Mutation at $x_{3-2j-k}^{j-k,k-1}$ and calculation of $\left(s_{-2j-k}^{j-k,k}(P_2)\right)'$ from $s_{-2j-k}^{j-k,k}(P_2)$ and $F_{3-2j-k}^{j-k,k-1}$ give

$$\left(s_{-2j-k}^{j-k,k}(P_2), F_{3-2j-k}^{j-k,k-1}\right) \xrightarrow{\mu_{3-2j-k}^{j-k,k-1}} \left(\tilde{u}\left(F_{2-2j-k}^{j-k,k}\right), F_{3-2j-k}^{j-k,k-1}\right). \quad (3.29)$$

Since $\tilde{u} \left(F_{2-2j-k}^{j-k,k} \right)$ does not depend on any $x \in X_0$ we have

$$\tilde{u} \left(F_{2-2j-k}^{j-k,k} \right) \xrightarrow{\lambda_3} \tilde{u} \left(F_{2-2j-k}^{j-k,k} \right). \quad (3.30)$$

The cluster variable $x_{3-2j-k}^{j-k,k}$ does not change by the mutations at $x \in X_0$ and we have

$$\left(x_{3-2j-k}^{j-k,k}, F_{3-2j-k}^{j-k,k} \right) \rightarrow \left(x_{3-2j-k}^{j-k,k}, \tilde{u} \left(F_{2-2j-k}^{j-k,k} \right) \right) \quad (3.31)$$

by the mutations of initial seed t_0 at each and all $x \in X_0$ just once.

Therefore, it holds that

$$\begin{aligned} t_0 &= \left\{ \left(x_{-2j-k}^{j-k,k}, F_{-2j-k}^{j-k,k} \right), \left(x_{i-2j-k}^{j-k,k}, F_{i-2j-k}^{j-k,k} \right) \middle| i \in [1, 5], j, k \in \mathbb{Z} \right\} \\ \rightarrow t_1 &= \left\{ \left(x_{6-2(j+2)-(k+1)}^{(j+2)-(k+1),k+1}, \tilde{u} \left(F_{5-2(j+2)-(k+1)}^{(j+2)-(k+1),k+1} \right) \right), \right. \\ &\quad \left. \left(x_{i-2j-k}^{j-k,k}, \tilde{u} \left(F_{-1+i-2j-k}^{j-k,k} \right) \right) \middle| i \in [1, 5], j, k \in \mathbb{Z} \right\}, \end{aligned} \quad (3.32)$$

which does not depend on the order of mutations. ■

Note that the iteration of the mutations $\tilde{\mu}_0$ does not depend on the order of the mutations.

Theorem 3.2 We defined the seed t_n by $t_n = \tilde{\mu}_{n-1}(t_{n-1})$. When we put $x_{n-1-2j-k}^{j-k,k} \rightarrow x_{n-2j-k}^{1+j-k,1+k}$, it holds that

$$t_n = \left\{ \left(x_{n+i-2j-k}^{j-k,k}, \tilde{u}^n \left(F_{i-2j-k}^{j-k,k} \right) \right) \middle| i \in [0, 5], j, k \in \mathbb{Z} \right\}. \quad (3.33)$$

All cluster variables $x_n^{m,l}$ ($n, m, l \in \mathbb{Z}$) satisfy

$$x_{n+1}^{m+1,l+1} x_n^{m,l} = x_n^{m+1,l+1} x_{n+1}^{m,l} + x_{n+1}^{m+1,l} x_n^{m,l+1} + x_{n+1}^{m,l+1} x_n^{m+1,l}. \quad (3.34)$$

Proof Eqs. (3.33) and (3.34) are obtained by the shift of the suffixes in Eqs. (3.8) and (3.9). Therefore, proof of Theorem 3.2 is the same as that of Proposition 3.1. Equation (3.34) is obtained by the definition of the new cluster variable

$$x_{1+n-2j-k}^{1+j-k,1+k} = \tilde{u}^n \left(F_{-2j-k}^{j-k,k} \right) / x_{n-2j-k}^{j-k,k} \quad (3.35)$$

by the mutation of the seed t_n at $x_{n-2j-k}^{j-k,k}$. ■

Equation (3.34) is called the discrete BKP equation [3]. The discrete BKP equation is the relation among 8 point of Figure 1. By Proposition 3.1, it holds

$$\tilde{\mu}_0(t_0) = \left\{ \left(x_{1+i-2j-k}^{j-k,k}, \tilde{u} \left(F_{i-2j-k}^{j-k,k} \right) \right) \middle| i \in [0, 5], j, k \in \mathbb{Z} \right\}. \quad (3.36)$$

The initial seed t_0 satisfies the condition which is similar to the one in the Definition 2.6. Therefore, the initial seed t_0 is regarded as a generalization of period-1 seed. By Theorem 2.5, all $x_n^{m,l}$ are Laurent polynomials of the initial values in X_0 .




Figure 1: The discrete BKP equation

3.2 Another seed of the discrete BKP equation

Let us take

$$t_0 = \left\{ \left(x_{i-4j-2k}^{j,k}, F_{i-4j-2k}^{j,k} \right) \middle| i \in [0, 6], j, k \in \mathbb{Z} \right\} \quad (3.37)$$

as an initial seed, where exchange polynomials are defined as

$$\begin{aligned} F_0^{0,0} &= x_0^{1,1} x_1^{0,0} + x_0^{1,0} x_1^{0,1} + x_0^{0,1} x_1^{1,0}, \\ F_1^{0,0} &= x_0^{1,0} x_1^{0,1} x_2^{0,0} + x_0^{0,1} x_1^{1,0} x_2^{0,0} + x_0^{0,0} x_1^{1,0} x_2^{0,1} + x_0^{0,0} x_1^{0,1} x_2^{1,0}, \\ F_2^{0,0} &= x_1^{1,0} x_2^{0,1} x_3^{0,-1} x_3^{0,0} + x_1^{0,1} x_2^{0,-1} x_3^{1,0} x_3^{0,0} \\ &\quad + x_1^{0,0} x_2^{0,1} x_3^{1,-1} x_3^{0,0} + x_1^{0,0} x_2^{0,-1} x_3^{1,0} x_3^{0,1} + x_1^{0,0} x_2^{0,1} x_3^{1,0} x_3^{0,-1}, \\ F_3^{0,0} &= x_2^{0,-1} x_3^{0,1} x_4^{0,0} + x_2^{0,1} x_3^{0,-1} x_4^{0,0} + x_2^{0,0} x_3^{0,-1} x_4^{0,1} + x_2^{0,0} x_3^{0,1} x_4^{0,-1}, \\ F_4^{0,0} &= x_5^{-1,0} x_4^{0,-1} x_4^{0,1} x_3^{0,0} + x_5^{0,-1} x_4^{0,1} x_4^{-1,0} x_3^{0,0} \\ &\quad + x_5^{0,0} x_4^{0,-1} x_4^{-1,1} x_3^{0,0} + x_5^{0,0} x_4^{0,1} x_4^{-1,0} x_3^{0,-1} + x_5^{0,0} x_4^{0,-1} x_4^{-1,0} x_3^{0,1}, \\ F_5^{0,0} &= x_6^{-1,0} x_5^{0,-1} x_4^{0,0} + x_6^{0,-1} x_5^{-1,0} x_4^{0,0} + x_6^{0,0} x_5^{-1,0} x_4^{0,-1} + x_6^{0,0} x_5^{0,-1} x_4^{-1,0}, \\ F_6^{0,0} &= x_6^{-1,-1} x_5^{0,0} + x_6^{-1,0} x_5^{0,-1} + x_6^{0,-1} x_5^{-1,0}, \end{aligned} \quad (3.38)$$

and

$$F_{i-4j-2k}^{j,k} = s_{-4j-2k}^{j,k} (F_i^{0,0}). \quad (3.39)$$

Now we define $\tilde{u} = s_1^{0,0}$ and the set of cluster variables X_i as

$$X_i := \left\{ x_{i-4j-2k}^{j,k} \middle| j, k \in \mathbb{Z} \right\}. \quad (3.40)$$

Theorem 3.3 *We define the seed t_n as the seed obtained by the iteration of the mutations of the seed t_{n-1} at each and all $x \in X_{n-1}$ just once. When we put $x_{n-1-4j-2k}^{j,k} \rightarrow x_{n-4j-2k}^{1+j,1+k}$, we find*

$$t_n = \left\{ \left(x_{n+i-4j-2k}^{j,k}, \tilde{u}^n \left(F_{i-4j-2k}^{j,k} \right) \right) \middle| i \in [0, 6], j, k \in \mathbb{Z} \right\}, \quad (3.41)$$

which does not depend on the order of mutations. All cluster variables $x_n^{m,l}$ ($n, m, l \in \mathbb{Z}$) satisfy the discrete BKP equation (3.34).

Proof *Proof of Theorem 3.3 can be done in a similar manner to those of Proposition 3.1 and Theorem 3.2.*

4 Several difference equations associated with reductions of the seed

In this section, we show that several seeds which give the difference equations obtained by imposing constraints on the BKP equation and discuss the relation to the reductions for initial seeds and polynomials in Laurent phenomenon algebra.

4.1 2-dimensional difference equation (1)

We take

$$t_0 = \{ (x_{i-2j}^j, F_{i-2j}^j) \mid i \in [0, 5], j \in \mathbb{Z} \} \quad (4.1)$$

as an initial seed, where exchange polynomials are defined as

$$\begin{aligned} F_0^0 &= x_1^2 x_1^0 + x_0^1 x_2^1 + (x_1^1)^2, \\ F_1^0 &= x_0^1 x_2^1 x_2^0 + (x_1^1)^2 x_2^0 + x_0^0 x_1^1 x_3^1 + x_0^0 (x_2^1)^2, \\ F_2^0 &= x_1^1 x_2^{-1} x_3^1 x_3^0 + (x_2^1)^2 x_2^{-1} x_3^0 + x_1^0 (x_3^0)^2 x_2^1 + x_1^0 x_2^{-1} (x_3^1)^2 + x_1^0 x_3^1 x_2^1 x_3^{-1}, \\ F_3^0 &= x_4^{-1} x_3^1 x_2^{-1} x_2^0 + (x_3^{-1})^2 x_3^1 x_2^0 + x_4^0 (x_2^0)^2 x_3^{-1} + x_4^0 x_3^1 (x_2^{-1})^2 + x_4^0 x_2^{-1} x_3^{-1} x_2^1, \\ F_4^0 &= x_5^{-1} x_3^{-1} x_3^0 + (x_4^{-1})^2 x_3^0 + x_5^0 x_4^{-1} x_2^{-1} + x_5^0 (x_3^{-1})^2, \\ F_5^0 &= x_4^{-2} x_4^0 + x_5^{-1} x_3^{-1} + (x_4^{-1})^2, \end{aligned} \quad (4.2)$$

and

$$F_{i-2j}^j = s_{-2j}^j (F_i^0). \quad (4.3)$$

We define $\tilde{u} := s_1^0$ and the set of cluster variables X_i by

$$X_i := \{ x_{i-2j}^j \mid j \in \mathbb{Z} \}. \quad (4.4)$$

Proposition 4.1 *We define the seed t_n as the seed obtained by the iteration of the mutations of the seed t_{n-1} at each and all $x \in X_{n-1}$ just once. When we put $x_{n-1-2j}^j \rightarrow x_{n+1-2j}^{2+j}$, we have*

$$t_n = \{ (x_{n+i-2j}^j, \tilde{u}^n (F_{i-2j}^j)) \mid i \in [0, 5], j \in \mathbb{Z} \}, \quad (4.5)$$

which does not depend on the order of mutations. All cluster variables x_n^m ($n, m \in \mathbb{Z}$) satisfy

$$x_{n+2}^{m+2} x_n^m = x_{n+1}^{m+2} x_{n+1}^m + (x_{n+1}^{m+1})^2 + x_{n+2}^{m+1} x_n^{m+1}. \quad (4.6)$$

2-dimensional equation (4.6) is the relation among 7 points shown in Fig. 2. Note that 2-dimensional difference equation (4.6) is obtained from the discrete BKP equation (3.34) by imposing the reduction condition

$$x_n^{m,l+1} = x_{n+1}^{m+1,l}, \quad x_n^m := x_n^{m,0}. \quad (4.7)$$

The initial seed (4.1) and the exchange polynomials (4.2) are obtained from the initial seed (3.1) and the exchange polynomials (3.2) by imposing the same condition (4.7). Therefore, the reduction of the seed corresponds to the reduction of the difference equation. Proof of Proposition 4.1 is the same as those of Proposition 3.1 and Theorem 3.2.




Figure 2: 2-dimensional difference equation (1)

4.2 Somos-6

We impose the reduction condition

$$x_n^{m+1} = x_{n+2}^m, \quad x_n := x_n^0 \quad (4.8)$$

to the initial seed (4.1) and exchange polynomials (4.2). Then we obtain the seed and exchange polynomials

$$t_0 = \{(x_i, F_i) | i \in [0, 5]\}, \quad (4.9)$$

$$\begin{aligned} F_0 &= x_5 x_1 + x_2 x_4 + (x_3)^2, \\ F_1 &= (x_2)^2 x_4 + (x_3)^2 x_2 + x_0 x_3 x_5 + x_0 (x_4)^2, \\ F_2 &= (x_3)^2 x_0 x_5 + (x_4)^2 x_0 x_3 + x_1 (x_3)^2 x_4 + x_1 x_0 (x_5)^2 + (x_1)^2 x_5 x_4, \\ F_3 &= (x_2)^2 x_5 x_0 + (x_1)^2 x_5 x_2 + x_4 (x_2)^2 x_1 + x_4 x_5 (x_0)^2 + (x_4)^2 x_0 x_1, \\ F_4 &= (x_3)^2 x_1 + (x_2)^2 x_3 + x_5 x_2 x_0 + x_5 (x_1)^2, \\ F_5 &= x_0 x_4 + x_3 x_1 + (x_2)^2. \end{aligned} \quad (4.10)$$

This seed is a period-1 seed. This seed has already been obtained in [9].

Proposition 4.2 [9] *We define the seed t_n by $t_n = \mu_{n-1}(t_{n-1})$. When we put $x_{n-1} \rightarrow x_{n+5}$, it holds that*

$$t_n = \{(x_{n+i}, u^n(F_i)) | i \in [0, 5]\}. \quad (4.11)$$

All cluster variables x_n ($n \in \mathbb{Z}$) satisfy

$$x_{n+6} x_n = x_{n+5} x_{n+1} + (x_{n+3})^2 + x_{n+4} x_{n+2}. \quad (4.12)$$

Difference equation (4.12) is called the Somos-6 [4]. The Somos-6 (4.12) is obtained from 2-dimensional difference equation (4.6) by imposing the reduction condition (4.8).

4.3 2-dimensional difference equation (2)

We take

$$t_0 = \{(x_{i-4j}^j, F_{i-4j}^j) | i \in [0, 6], j \in \mathbb{Z}\} \quad (4.13)$$




Figure 3: 2-dimensional difference equation (2)

as an initial seed, where exchange polynomials are defined as

$$\begin{aligned}
F_0^0 &= x_2^1 x_1^0 + x_0^1 x_3^0 + x_2^0 x_1^1, \\
F_1^0 &= x_0^1 x_3^0 x_2^0 + (x_2^0)^2 x_1^1 + x_0^0 x_1^1 x_4^0 + x_0^0 x_3^0 x_2^1, \\
F_2^0 &= x_1^1 x_4^0 x_0^0 x_3^0 + (x_3^0)^2 x_0^0 x_2^1 + x_1^0 x_4^0 x_0^1 x_3^0 + x_1^0 x_0^0 x_2^1 x_5^0 + (x_1^0)^2 x_4^0 x_2^1, \\
F_3^0 &= x_0^0 x_5^0 x_4^0 + (x_4^0)^2 x_1^0 + x_2^0 x_1^0 x_6^0 + (x_2^0)^2 x_5^0, \\
F_4^0 &= x_5^1 x_2^0 x_6^0 x_3^0 + (x_3^0)^2 x_6^0 x_4^{-1} + x_5^0 x_2^0 x_6^{-1} x_3^0 + x_5^0 x_6^0 x_4^{-1} x_1^0 + (x_5^0)^2 x_2^0 x_4^{-1}, \\
F_5^0 &= x_6^{-1} x_3^0 x_4^0 + (x_4^0)^2 x_5^{-1} + x_6^0 x_5^{-1} x_2^0 + x_6^0 x_3^0 x_4^{-1}, \\
F_6^0 &= x_4^{-1} x_5^0 + x_6^{-1} x_3^0 + x_4^0 x_5^{-1},
\end{aligned} \tag{4.14}$$

and

$$F_{i-4j}^j = s_{-4j}^j (F_i^0). \tag{4.15}$$

We define $\tilde{u} = s_1^0$ and the set of cluster variables X_i as

$$X_i := \{x_{i-4j}^j \mid j \in \mathbb{Z}\}. \tag{4.16}$$

Proposition 4.3 *We define the seed t_n as the seed obtained by the iteration of the mutations of the seed t_{n-1} at each and all $x \in X_{n-1}$ just once. Putting $x_{n-1-4j}^j \rightarrow x_{n+2-4j}^{j+1}$, we find*

$$t_n = \{(x_{n+i-4j}^j, \tilde{u}^n (F_{i-4j}^j)) \mid i \in [0, 6], j \in \mathbb{Z}\}, \tag{4.17}$$

which does not depend on the order of mutations. All cluster variables x_n^m ($n, m \in \mathbb{Z}$) satisfy

$$x_{n+3}^{m+1} x_n^m = x_{n+2}^{m+1} x_{n+1}^m + x_{n+1}^{m+1} x_{n+2}^m + x_{n+3}^m x_n^{m+1}. \tag{4.18}$$

2-dimensional equation (4.18) is the relation among 8 points shown in Fig. 3.

Note that 2-dimensional difference equation (4.18) is obtained from the discrete BKP equation (3.34) by imposing the reduction condition

$$x_n^{m,l+1} = x_{n+2}^{m,l}, \quad x_n^m := x_n^{m,0}. \tag{4.19}$$

The initial seed (4.13) and the exchange polynomials (4.14) are obtained from the initial seed (3.37) and the exchange polynomials (3.38) by imposing the same condition (4.19).

4.4 Somos-7

We take

$$t_0 = \{(x_i, F_i) \mid i \in [0, 6]\} \tag{4.20}$$

as an initial seed, where exchange polynomials are defined as

$$\begin{aligned}
F_0 &= x_6x_1 + x_4x_3 + x_2x_5 \\
F_1 &= x_4x_3x_2 + (x_2)^2x_5 + x_0x_5x_4 + x_0x_3x_6 \\
F_2 &= x_5x_4x_0x_3 + (x_3)^2x_0x_6 + x_1(x_4)^2x_3 + x_1x_0x_6x_5 + (x_1)^2x_4x_6 \\
F_3 &= x_0x_5x_4 + (x_4)^2x_1 + x_2x_1x_6 + (x_2)^2x_5 \\
F_4 &= x_1x_2x_6x_3 + (x_3)^2x_6x_0 + x_5(x_2)^2x_3 + x_5x_6x_0x_1 + (x_5)^2x_2x_0 \\
F_5 &= x_2x_3x_4 + (x_4)^2x_1 + x_6x_1x_2 + x_6x_3x_0 \\
F_6 &= x_0x_5 + x_2x_3 + x_4x_1.
\end{aligned} \tag{4.21}$$

Proposition 4.4 *We define the seed t_n by $t_n = \mu_{n-1}(t_{n-1})$. We put $x_{n-1} \rightarrow x_{n+6}$ and we have*

$$t_n = \{(x_{n+i}, u^n(F_i)) \mid i \in [0, 6]\}. \tag{4.22}$$

All cluster variables x_n ($n \in \mathbb{Z}$) satisfy

$$x_{n+7}x_n = x_{n+6}x_{n+1} + x_{n+5}x_{n+2} + x_{n+3}x_{n+4}. \tag{4.23}$$

Difference equation (4.23) is called the Somos-7 [4]. Note that the Somos-7 (4.23) is obtained from the 2-dimensional difference equation (4.18) by imposing the reduction condition

$$x_n^{m+1} = x_{n+4}^m, \quad x_n := x_n^0. \tag{4.24}$$

The initial seed (4.20) and the exchange polynomials (4.21) are obtained from the initial seed (4.13) and the exchange polynomials (4.14) by imposing the same condition (4.24).

5 Conclusion

We have shown that cluster variables can satisfy the discrete BKP equation, the 2-dimensional difference equations of its reductions, and Somos-7, if we take appropriate initial seeds in Laurent phenomenon algebras. These initial seeds are obtained from reductions of the seed of the discrete BKP equation. It is known that cluster variables of suitable cluster algebras can satisfy the bilinear form of some q -discrete Painlevé equations [5], when the initial seed includes appropriate periodic quivers [8]. However, we have not obtained the q -discrete Painlevé equations of type $A_4^{(1)}$ and $E_n^{(1)}$ from cluster algebras. To clarify the relation between these equations and Laurent phenomenon algebras is one of the problems we wish to address in the future.

References

- [1] R. Hirota, “Nonlinear Partial Difference Equations. I. A Difference Analogue of the Korteweg-de Vries Equation”, Journal of the Physical Society of Japan, **43**, (1977), 1424-1433.
- [2] R. Hirota, “Discrete Analogue of a Generalized Toda Equation”, JPSJ, **50**, (1981), 3785-3791.

- [3] T. Miwa, “On Hirota’s difference equations”, Proc. Japan Acad. Ser. A Math. Sci. **58**, (1982), 912.
- [4] D. Gale, “Mathematical entertainments: the strange and surprising saga of the somos sequences”, Math. Intelligencer **13**, (1991), 4042.
- [5] H. Sakai, “Rational surfaces associated with affine root systems and geometry of the Painlevé equations”, Communications in Mathematical Physics, **220**, (2001), 165-229.
- [6] S. Fomin and A. Zelevinsky, “Cluster algebras IV: Coefficients”, Compositio Mathematica **143**, (2007), 112-164.
- [7] N. Okubo, “Discrete integrable systems and cluster algebras”, RIMS Kokyuroku Bessatsu, Research Institute for Mathematical Sciences, **B41**, (2013), 25-42.
- [8] N. Okubo, “Bilinear equations and q -discrete Painlevé equations satisfied by variables and coefficients in cluster algebras”, J. Phys. A: Math. Theor. **48**, (2015), 355201.
- [9] T. Lam, P. Pylyavskyy, “Laurent Phenomenon Algebras”, preprint, arXiv: 1206.2611.
- [10] J. Alman, C. Cuenca, J. Huang, “Laurent Phenomenon Sequences”, preprint, arXiv: 1309.0751.