

The variable metric forward-backward splitting algorithm under mild differentiability assumptions

Saverio Salzo

*LCSL, Istituto Italiano di Tecnologia and Massachusetts Institute of Technology
Via Morego 30, 16163 Genova, Italy
email: saverio.salzo@iit.it*

Abstract

We study the variable metric forward-backward splitting algorithm for convex minimization problems without the standard assumption of the Lipschitz continuity of the gradient. In this setting, we prove that, by requiring only mild assumptions on the smooth part of the objective function and using several types of line search procedures for determining either the gradient descent stepsizes, or the relaxation parameters, one still obtains weak convergence of the iterates and convergence in the objective function values. Moreover, the $o(1/k)$ convergence rate in the function values is obtained if slightly stronger differentiability assumptions are added. We also illustrate several applications including problems that involve Banach spaces and functions of divergence type.

Keywords. Convex optimization, forward-backward algorithm, variable metric, inexact line search methods, quasi-Fejér sequences, global convergence, convergence rates.

AMS Mathematics Subject Classification: 65K05, 90C25, 90C30

1 Introduction

The forward-backward splitting algorithm [20] is nowadays a well-established and widely used first order optimization method that is well suited for an objective function composed by a smooth convex function plus a (possibly nonsmooth) convex simple function. This algorithm has been studied in a number of works [2, 5, 13, 20, 22] which prove weak convergence of the iterates as well as $o(1/k)$ convergence rate in function values. The *variable metric* version of the forward-backward method aims at accelerating the convergence of the standard algorithm. It was first proposed in [15] and its global convergence property has been established in full generality in [19] where, under an appropriate monotonicity condition on the metrics, the authors prove weak convergence of the iterates. The same algorithm is also analyzed in [16, 24] for the minimization of *non convex* composite functions and global convergence is achieved by assuming the Kurdyka-Łojasiewicz property, together with compactness conditions, and suitably controlling the behavior of the variable metrics.

In this context, a fundamental and commonly adopted assumption is that the gradient of the smooth part is Lipschitz continuous on the entire space. However, there are a number of applications in which this condition is not satisfied: for instance, in inverse problems when the data fidelity term is based on Banach norms [13, 39] or Bregman distances (e.g., the Kullback-Leibler divergence, which is the appropriate choice when the data are corrupted by Poisson noise [11, 38]).

1.1 Objective and main contribution

In this paper we address the convergence analysis of the variable metric forward-backward splitting algorithm in infinite dimensional Hilbert spaces without the assumption of the Lipschitz continuity of the gradient and using different types of line search procedures. This study provides global convergence guarantees, both in terms of convergence of the iterates and rates of convergence in function values, and shows that the scope of applicability of the algorithm is significantly wider than that for which it was originally devised, up to cover problems involving Banach spaces and objective functions of divergence type. Our analysis is based on a general convergence principle for abstract variable metric descent algorithms which blends the concept of quasi-Fejer sequence with that of a sufficient decreasing condition. This principle simultaneously drives the convergence in the iterates and the convergence in the objective function values. Moreover, we provide a unifying view on several inexact line search procedures that have been proposed in literature in the context of projected/proximal gradient-type algorithms clarifying the relationships among them. We finally remark that, even under standard differentiability assumptions, we advance the related state of the art, since we provide rate of convergence in function values in infinite dimensional setting and we consider an alternative hypothesis on the metrics apart that of monotonicity made in [9, 19].

Our contribution is detailed below. We consider the problem

$$\underset{x \in \mathcal{H}}{\text{minimize}} \quad f(x) + g(x), \quad (\text{P})$$

where, \mathcal{H} is a real Hilbert space and

- H1** $f: \mathcal{H} \rightarrow]-\infty, +\infty]$ and $g: \mathcal{H} \rightarrow]-\infty, +\infty]$ are proper convex and lower semicontinuous functions with $\text{dom } g \subset \text{dom } f$;
- H2** f is Gâteaux differentiable on $\text{dom } g$ and ∇f is uniformly continuous on any weakly compact subset of $\text{dom } g$.

For that problem, we study the following algorithm [19]. Let $x_0 \in \text{dom } g$ and set

$$\begin{aligned} & \text{for } k = 0, 1, \dots \\ & \left[\begin{array}{l} \text{choose } \gamma_k \in \mathbb{R}_{++} \\ y_k = \text{prox}_{\gamma_k g}^k(x_k - \gamma_k \nabla^k f(x_k)) \\ \text{choose } \lambda_k \in]0, 1] \\ x_{k+1} = x_k + \lambda_k(y_k - x_k), \end{array} \right. \quad (\text{VM-FBS}) \end{aligned}$$

where ∇^k and prox^k denote the gradient operator and the proximity operator with respect to a given family of scalar products as specified by the following assumption.

- H3** $(\langle \cdot | \cdot \rangle_k)_{k \in \mathbb{N}}$ is a sequence of possibly varying scalar products (metrics) on \mathcal{H} , with induced norms $(\|\cdot\|_k)_{k \in \mathbb{N}}$ and associated positive operators $(W_k)_{k \in \mathbb{N}}$ (i.e., for every $k \in \mathbb{N}$, $W_k: \mathcal{H} \rightarrow \mathcal{H}$ is such that $\langle \cdot | \cdot \rangle_k = \langle \cdot | W_k \cdot \rangle$), and

$$\exists (\nu, \mu) \in \mathbb{R}^2, \quad 0 < \nu \leq \mu, \quad (\forall k \in \mathbb{N}) \quad \nu \|\cdot\|^2 \leq \|\cdot\|_k^2 \leq \mu \|\cdot\|^2. \quad (1.1)$$

Moreover, since we are dropping out the assumption of Lipschitz continuity of ∇f , we rely on inexact line search methods for determining the parameters γ_k and λ_k in (VM-FBS). We set, for every $k \in \mathbb{N}$ and every $x \in \text{dom } g$, $\gamma > 0$, and $\lambda \in]0, 1]$

$$J_k(x, \gamma, \lambda) = x + \lambda(\text{prox}_{\gamma g}^k(x - \gamma \nabla^k f(x)) - x),$$

so that $x_{k+1} = J_k(x_k, \gamma_k, \lambda_k)$. Then, the gradient descent *stepsizes* $(\gamma_k)_{k \in \mathbb{N}}$ and the *relaxation* parameters $(\lambda_k)_{k \in \mathbb{N}}$ are chosen according to one of the following rules:

LS1 $0 < \inf_{k \in \mathbb{N}} \lambda_k \leq \sup_{k \in \mathbb{N}} \lambda_k \leq 1$. Let $\delta, \theta \in]0, 1[$, $\bar{\gamma} > 0$, and, $\forall k \in \mathbb{N}$,

$$\gamma_k = \max \left\{ \gamma \in \mathbb{R}_{++} \mid (\exists i \in \mathbb{N})(\gamma = \bar{\gamma}\theta^i) \right. \\ \left. f(J_k(x_k, \gamma, \lambda_k)) - f(x_k) - \langle J_k(x_k, \gamma, \lambda_k) - x_k \mid \nabla f(x_k) \rangle \leq \frac{\delta}{\gamma \lambda_k} \|J_k(x_k, \gamma, \lambda_k) - x_k\|_k^2 \right\}.$$

LS2 $0 < \inf_{k \in \mathbb{N}} \gamma_k \leq \sup_{k \in \mathbb{N}} \gamma_k < +\infty$. Let $\delta, \theta \in]0, 1[$, $\bar{\lambda} \in]0, 1]$, and, $\forall k \in \mathbb{N}$,

$$\lambda_k = \max \left\{ \lambda \in \mathbb{R}_{++} \mid (\exists i \in \mathbb{N})(\lambda = \bar{\lambda}\theta^i) \right. \\ \left. f(J_k(x_k, \gamma_k, \lambda)) - f(x_k) - \langle J_k(x_k, \gamma_k, \lambda) - x_k \mid \nabla f(x_k) \rangle \leq \frac{\delta}{\gamma_k \lambda} \|J_k(x_k, \gamma_k, \lambda) - x_k\|_k^2 \right\}.$$

LS3 $0 < \inf_{k \in \mathbb{N}} \gamma_k \leq \sup_{k \in \mathbb{N}} \gamma_k < +\infty$. Let $\theta, \delta \in]0, 1[$, $\bar{\lambda} \in]0, 1]$, and, $\forall k \in \mathbb{N}$,

$$\lambda_k = \max \left\{ \lambda \in \mathbb{R}_{++} \mid (\exists i \in \mathbb{N})(\lambda = \bar{\lambda}\theta^i) \right. \\ \left. (f + g)(J_k(x_k, \gamma_k, \lambda)) - (f + g)(x_k) \leq (1 - \delta)\lambda(g(y_k) - g(x_k) + \langle y_k - x_k \mid \nabla f(x_k) \rangle) \right\}.$$

LS4 $0 < \inf_{k \in \mathbb{N}} \lambda_k \leq \sup_{k \in \mathbb{N}} \lambda_k \leq 1$. Let $\delta, \theta \in]0, 1[$, $\bar{\gamma} > 0$, and, $\forall k \in \mathbb{N}$,

$$\gamma_k = \max \left\{ \gamma \in \mathbb{R}_{++} \mid (\exists i \in \mathbb{N})(\gamma = \bar{\gamma}\theta^i) \right. \\ \left. \|\nabla^k f(J_k(x_k, \gamma, \lambda_k)) - \nabla^k f(x_k)\|_k \leq \frac{\delta}{\gamma \lambda_k} \|J_k(x_k, \gamma, \lambda_k) - x_k\|_k \right\}.$$

We remark that **LS1** and **LS4** search for an appropriate stepsize parameter γ_k before setting y_k , and choose a priori the relaxation parameters λ_k 's; while **LS2** and **LS3**, after computing y_k with an a priori choice of γ_k , search for a suitable relaxation parameter λ_k . Note that **LS1** and **LS2** ask for the descent lemma (Fact 2.6) for f to be locally satisfied, and **LS4** attempts to locally fulfill a Lipschitz condition for ∇f .

Our analysis guarantees that, under the mild hypotheses **H1–H3** and additional suitable assumptions on the metrics — either **H4** or **H5** in Section 3.3 — (which are in line with those of [9, 10, 19]), each of the line search rule above makes algorithm (VM-FBS) capable of generating a minimizing sequence for $f + g$ that also weakly converges to a solution of problem (P). More precisely, denoting by S_* the set of solutions of (P), we prove that

- if $S_* = \emptyset$, then $(f + g)(x_k) \rightarrow \inf_{\mathcal{H}}(f + g)$ and $\|x_k\| \rightarrow +\infty$.
- if $S_* \neq \emptyset$, then $(x_k)_{k \in \mathbb{N}}$ and $(y_k)_{k \in \mathbb{N}}$ weakly converge to the same point in S^* and $(f + g)(y_k) \rightarrow \inf_{\mathcal{H}}(f + g)$; if, in addition, ∇f is Lipschitz continuous on the weakly compact subsets of $\text{dom } g$, then $(\gamma_k)_{k \in \mathbb{N}}$ and $(\lambda_k)_{k \in \mathbb{N}}$ are bounded away from zero and $((f + g)(x_k) - \inf_{\mathcal{H}}(f + g)) = o(1/k)$.

As a consequence, the above conclusions are also valid when ∇f is L -Lipschitz continuous on $\text{dom } g$ and the sequences $(\gamma_k)_{k \in \mathbb{N}}$ and $(\lambda_k)_{k \in \mathbb{N}}$ are chosen a priori (without backtracking) provided that they are bounded away from zero and $\sup_{k \in \mathbb{N}} \gamma_k \lambda_k / \nu_k < 2/L$, ν_k being the minimum eigenvalue of the metric $\langle \cdot \mid \cdot \rangle_k$. This result is not covered by the convergence analysis in [19, 20, 21], since they are

based on the theory of fixed point algorithms for averaged nonexpansive operators and the Baillon-Haddad theorem [3], which requires ∇f to have full domain. Moreover, a coupling of the parameters γ_k , λ_k , and ν_k is disclosed which somehow complements the analogue result in [21].

Finally we show that the above results hold also for general placements of the domains of f and g (thus even if $\text{dom } g \not\subset \text{dom } f$) at the cost of requiring coercivity-type conditions and adding a further line search procedure that carry the iterates inside the domain of f before executing any of **LS1–LS4**. This further generalization allows to treat, e.g., linear inverse problems with Poisson noise, that requires smooth terms of divergence type.

1.2 Comparison with related works

In literature concerning the forward-backward algorithm, the problem of removing the requirement of the Lipschitz continuous gradient for the smooth part has been receiving growing attention during the last years. Currently there are two streams of research on this issue. The first one focuses on a restricted class of smooth functions that do not enjoy Lipschitz continuous gradient, but possess other special properties. The work [40] belongs to this kind of studies: for \mathcal{H} finite dimensional, it analyses the case that f is a smooth self-concordant function and addresses both global and local convergence. The second research line considers a wide class of smooth functions (e.g., continuously differentiable) and introduce line searches to determine the parameters of the algorithm. Our work is within this stream. In the following we discuss two aspects.

The forward-backward algorithm without the Lipschitz assumption

The literature on the forward-backward algorithm in the absence of the Lipschitz assumption is scarce. The pioneering work by Tseng and Yun [42] is the first that considers a variable metric forward-backward algorithm in finite dimensional spaces, where the smooth part is only continuously differentiable (possibly non-convex). They proposes a general Armijo-type line search rule and prove that cluster points of the generated sequence are stationary points. Special instances of this general line search are also employed in the recent works [6, 9] which advances the theory for the convex case by addressing global convergence of the iterates and rate of convergence in function values without the Lipschitz assumption. However, these studies are not completely satisfying since in [6] the proposed line searches are not quite suitable for the forward-backward algorithm (as we discuss below), the $o(1/k)$ convergence rate in function values is obtained only in finite dimension, and the differentiability assumptions are not completely relaxed, so that, e.g., functions of divergent type may remain out of scope; while in [9] the analysis is conducted in finite dimensional spaces and demands $\text{dom } g$ to be closed and still the Lipschitz assumption for the $O(1/k)$ rate of convergence in function values. On the other hand, the special case of the *gradient projection method* [27, 31, 29] have been studied for long time by requiring just the continuity of the gradient of the objective function and using different types of line searches for determining the step lengths [7, 8, 14, 25, 34]. In particular, for the convex and finite dimensional case, [29] proves convergence of the iterates using two types of Armijo line searches, while in the recent [10], both convergence of the iterates and a $O(1/k)$ convergence rate in function values are proved for the variable metric (scaled) version, assuming coercivity of the objective function and local Lipschitz continuity of the gradient.

Line search methods for proximal gradient-type methods

Line search methods for gradient-type algorithms date several decades ago [7, 14, 25, 26]. In the following we position **LS1–LS4** with respect to the more recent literature. We notice that, when g is

zero, **LS1**, **LS2**, and **LS3** reduce to the classical Armijo line search along the steepest descent direction [8, 12, 35]. Moreover, when g is the indicator of a closed convex set, **LS3** reduces to the Armijo line search along the feasible direction commonly used in gradient projection methods [8, 10, 29]. **LS1** (with $\delta = 1/2$ and $\lambda_k \equiv 1$) has been proposed for the first time in [5], where the authors provide the rate of convergence in function values of the forward-backward algorithm with no relaxation under the assumption that ∇f is everywhere defined and globally Lipschitz continuous. In that case the line search was introduced to cope with situations in which the Lipschitz constant of ∇f was unknown or expensive to compute. **LS3** (with $\gamma_k \equiv 1$) is a special instance of the general Armijo rule proposed by Tseng and Yun [42] — which we do not explicitly treat here, but is encompassed by our analysis (see Remark 3.14). **LS3** has also been employed in [30] within a proximal Newton-type method for convex minimization problems of type (P) in finite dimensional spaces and under Lipschitz assumption. We recall that (VM-FBS) can be seen as a proximal quasi-Newton method: indeed the y_k can be equivalently computed as

$$y_k = \operatorname{argmin}_{z \in \mathcal{H}} \langle z - x_k \mid \nabla f(x_k) \rangle + \frac{1}{2\gamma_k} \langle z - x_k \mid W_k(z - x_k) \rangle + g(z).$$

Unfortunately the proof of global convergence in [30, Theorem 3.1] is not correct. There, it is only proved the descent property and that $x_{k+1} - x_k \rightarrow 0$, and mistakenly inferred from this that $(x_k)_{k \in \mathbb{N}}$ is convergent. Finally, **LS4** was originally employed by Tseng in [41] for the more general problem of finding zeros of the sum of two maximal monotone operators. This line search rule has been also recently studied in [6] (with $\delta \leq 1/2$) in conjunction with the (stationary metric) forward-backward splitting algorithm for convex minimization problems without the assumption of Lipschitz continuous gradient. However, we stress that procedure **LS4**, at each iteration, calls for multiple evaluations of the gradient — a fact that may lead to significantly increase the computational cost of the algorithm — and, more importantly it may determine shorter stepsizes than those computed by the other line search rules (see Remarks 3.21 and 3.25). In this regard, we note that the procedure proposed in [41] is designed for general Lipschitz (or even continuous) monotone operators, not specifically for gradient operators. By contrast, **LS1–LS3** seem more appropriate to exploit the fact that we are dealing with gradient operators: we demonstrate indeed that they provide larger stepsizes that are consistent with those permitted under the standard Lipschitz assumptions (this issue parallels that between Lipschitz continuity and cocoercivity). In [6] a further line search is also analyzed which is in between **LS2** and **LS3** with $\delta = 1/2$, but again it leads to determine reduced step lengths (see Remark 3.14).

1.3 Outline of the paper

Section 2 contains notations and basic concepts and facts. In Section 3 we first give preliminary results concerning the differentiability assumptions and the well-posedness of the line searches **LS1–LS4**; then we present an abstract principle which is at the basis of the global convergence properties of variable metric descent algorithms; and finally we study the convergence of algorithm (VM-FBS) in conjunction with the proposed line search procedures. Section 4 shows that the convergence results can be extended to situations in which the domain of g is not contained in the domain of f , relaxing one requirement in **H1**. Finally, in Section 5 we present examples of problems of type (P), where the gradient of the smooth part is not Lipschitz continuous, that can be tackled by the proposed algorithm.

2 Basic definitions and facts

Throughout the paper the notation we employ is standard and as in [2]. We assume that \mathcal{H} is a real Hilbert space with scalar product $\langle \cdot | \cdot \rangle$ and norm $\|\cdot\|$. If $\langle \cdot | \cdot \rangle_1$ is another scalar product on \mathcal{H} , its associated positive operator is $V_1: \mathcal{H} \rightarrow \mathcal{H}$ such that, for every $(x, y) \in \mathcal{H}^2$, $\langle x | y \rangle_1 = \langle x | V_1 y \rangle$. We set $\mathbb{R}_+ = [0, +\infty[$ and $\mathbb{R}_{++} =]0, +\infty[$ and we denote by ℓ_+^1 the set of summable sequences in \mathbb{R}_+ . Moreover, for every $(x, y) \in \mathcal{H}^2$, we set $[x, y] = \{x + t(y - x) \mid t \in [0, 1]\}$. Let $h: \mathcal{H} \rightarrow]-\infty, +\infty]$ be a proper function. We set $\operatorname{argmin}_{\mathcal{H}} h = \{x \in \mathcal{H} \mid h(x) = \inf_{\mathcal{H}} h\}$, and when it is a singleton, its unique element, by an abuse of notation, is still denoted by $\operatorname{argmin}_{\mathcal{H}} h$.

We recall two fundamental facts about monotone sequences and Fejér sequences.

Fact 2.1 ([22, Lemma 3]). *Let $(\rho_k)_{k \in \mathbb{N}} \in \mathbb{R}_+^{\mathbb{N}}$ and $(\alpha_k)_{k \in \mathbb{N}} \in \mathbb{R}_{++}^{\mathbb{N}}$ be such that*

$$(\forall k \in \mathbb{N}) \quad \rho_{k+1} \leq \rho_k \quad \text{and} \quad \sum_{k \in \mathbb{N}} \alpha_k \rho_k < +\infty.$$

Then, for every $k \in \mathbb{N}$, $\rho_k \leq (\sum_{i=0}^k \alpha_i)^{-1} \sum_{k \in \mathbb{N}} \alpha_k \rho_k$ and $\rho_k = o(1 / \sum_{i=\lceil k/2 \rceil}^k \alpha_i)$. In particular, if $(\alpha_k)_{k \in \mathbb{N}} \notin \ell_+^1$, then $\rho_k \rightarrow 0$, and if $\inf_{k \in \mathbb{N}} \alpha_k > 0$, then $\rho_k = o(1/k)$.

Definition 2.2 ([18, Definition 3.1(ii)]). Let $(|\cdot|_k)_{k \in \mathbb{N}}$ be a sequence of Hilbert norms on \mathcal{H} such that, for some $\nu > 0$, $\nu \|\cdot\|^2 \leq |\cdot|_k^2$, for every $k \in \mathbb{N}$. Let $S \subset \mathcal{H}$ be a nonempty set. A sequence $(x_k)_{k \in \mathbb{N}}$ in \mathcal{H} is a *quasi-Fejér sequence with respect to S relative to $(|\cdot|_k)_{k \in \mathbb{N}}$* if there exist $(\varepsilon_k)_{k \in \mathbb{N}}$ and $(\eta_k)_{k \in \mathbb{N}}$ in ℓ_+^1 such that

$$(\forall x \in S)(\forall k \in \mathbb{N}) \quad |x_{k+1} - x|_{k+1}^2 \leq (1 + \eta_k) |x_k - x|_k^2 + \varepsilon_k. \quad (2.1)$$

Fact 2.3 ([18, Lemma 2.3, Proposition 3.2, and Theorem 3.3]). *Let $(|\cdot|_k)_{k \in \mathbb{N}}$ be a sequence of Hilbert norms on \mathcal{H} with associated positive operators $(V_k)_{k \in \mathbb{N}}$. Suppose that, for some $\nu > 0$, $\nu \|\cdot\|^2 \leq |\cdot|_k^2$ for every $k \in \mathbb{N}$. Consider the following statements.*

- (a) *There exists $(\eta_k)_{k \in \mathbb{N}} \in \ell_+^1$ such that, for every $k \in \mathbb{N}$, $|\cdot|_{k+1}^2 \leq (1 + \eta_k) |\cdot|_k^2$.*
- (b) *There exists a positive operator V such that, for every $x \in \mathcal{H}$, $V_k x \rightarrow Vx$.*

Then (a) \Rightarrow (b).¹ Moreover, let $S \subset \mathcal{H}$ be a nonempty set, and let $(x_k)_{k \in \mathbb{N}} \in \mathcal{H}^{\mathbb{N}}$ be a quasi Fejér sequence with respect to S relative to $(|\cdot|_k)_{k \in \mathbb{N}}$. Then, if (b) holds,

- (i) *$(x_k)_{k \in \mathbb{N}}$ is bounded and, for every $y \in S$, $(|x_k - y|_k)_{k \in \mathbb{N}}$ is convergent.*
- (ii) *$(x_k)_{k \in \mathbb{N}}$ is weakly convergent to a point of S if and only if every weak sequential cluster point of $(x_k)_{k \in \mathbb{N}}$ belongs to S .*

Definition 2.4. Let $h: \mathcal{H} \rightarrow]-\infty, +\infty]$ be proper and convex and let $x \in \operatorname{dom} h$. The function h is said to be *Gâteaux differentiable* at $x \in \operatorname{dom} h$ if there exists $u \in \mathcal{H}$, such that, for every $d \in \mathcal{H}$, $\lim_{t \rightarrow 0} (h(x + td) - h(x))/t = \langle d | u \rangle$ (see [2, p. 243]); in this case $\partial h(x) = \{u\}$ and the unique element of $\partial h(x)$ is denoted by $\nabla h(x)$. Moreover h is *Gâteaux differentiable on $A \subset \mathcal{H}$* if it is Gâteaux differentiable at every point of A .

Remark 2.5. If $h \in \Gamma_0(\mathcal{H})$ and it is Gâteaux differentiable at $x \in \operatorname{dom} h$, then $x \in \operatorname{int} \operatorname{dom} h$ and h is continuous on $\operatorname{int} \operatorname{dom} h \subset \operatorname{dom} \partial h$ [2].

¹ The condition $\sup_{k \in \mathbb{N}} \|V_k\| < +\infty$ required in [18] is not necessary, since it is a consequence of $|\cdot|_{k+1}^2 \leq (1 + \eta_k) |\cdot|_k^2$ and $\prod_{k=0}^{+\infty} (1 + \eta_k) < +\infty$.

Fact 2.6 (Descent Lemma). *Let $h: \mathcal{H} \rightarrow]-\infty, +\infty]$ be proper and convex. Suppose that h is Gâteaux differentiable with L -Lipschitz continuous gradient on the segment $[x, y] \subset \text{int dom } h$. Then, we have $h(y) - h(x) - \langle y - x | \nabla h(x) \rangle \leq (L/2) \|x - y\|^2$.*

Suppose that **H3** holds. Let $k \in \mathbb{N}$ and let W_k be the positive operator associated to $\langle \cdot | \cdot \rangle_k$. We set $\mu_k = \sup_{\|x\|=1} \langle x | W_k x \rangle = \|W_k\|$ and $\nu_k = \inf_{\|x\|=1} \langle x | W_k x \rangle = \|W_k^{-1}\|^{-1}$, being respectively the maximum and minimum eigenvalue of W_k . Then

$$\nu \|\cdot\|^2 \leq \nu_k \|\cdot\|^2 \leq \|\cdot\|_k^2 \leq \mu_k \|\cdot\|^2 \leq \mu \|\cdot\|^2. \quad (2.2)$$

Let $h \in \Gamma_0(\mathcal{H})$ and let $x \in \text{dom } h$. We denote by $\partial^k h$ the subdifferential of h at x in the metric $\langle \cdot | \cdot \rangle_k$ and we have, for every $u \in \mathcal{H}$,

$$u \in \partial^k h(x) \Leftrightarrow (\forall y \in \mathcal{H}) \ h(y) \geq h(x) + \langle y - x | W_k u \rangle \Leftrightarrow W_k u \in \partial h(x). \quad (2.3)$$

Moreover, if h is Gâteaux differentiable at x we denote by $\nabla^k h$ the gradient of h at x in the metric $\langle \cdot | \cdot \rangle_k$, and we have $\nabla h(x) = W_k \nabla^k h(x)$ and

$$(\forall d \in \mathcal{H}) \quad \langle d | \nabla h(x) \rangle = \lim_{t \rightarrow 0} \frac{h(x + td) - h(x)}{t} = \langle d | \nabla^k h(x) \rangle_k. \quad (2.4)$$

Fact 2.7. *Assume that **H3** holds. Let $k \in \mathbb{N}$ and let ν_k and μ_k be the minimum and maximum eigenvalue of W_k . Let $h \in \Gamma_0(\mathcal{H})$ and suppose that h is Gâteaux differentiable on a set $C \subset \mathcal{H}$. Let $x, y \in C$. Then*

$$\frac{1}{\sqrt{\mu_k}} \|\nabla h(x) - \nabla h(y)\| \leq \|\nabla^k h(x) - \nabla^k h(y)\|_k \leq \frac{1}{\sqrt{\nu_k}} \|\nabla h(x) - \nabla h(y)\|. \quad (2.5)$$

Let $h \in \Gamma_0(\mathcal{H})$. The *proximity operator* of h is the map $\text{prox}_h: \mathcal{H} \rightarrow \mathcal{H}$ such that, for every $x \in \mathcal{H}$, $\text{prox}_h(x) = \arg\min_{y \in \mathcal{H}} h(y) + (1/2) \|y - x\|^2$. Moreover

$$(\forall x \in \mathcal{H})(\forall z \in \mathcal{H}) \quad z = \text{prox}_h x \iff x - z \in \partial h(z). \quad (2.6)$$

The following result can be partially derived from the asymptotic behavior of the resolvent of maximal monotone operators [2, Theorem 23.47]. We also provide the bound (2.7), by slightly modifying the proof of [28, Proposition 4.1.5, Chap. XV].

Fact 2.8. *Let $h \in \Gamma_0(\mathcal{H})$ and let $\gamma > 0$. Then, for every $u \in \text{dom } h^*$, we have*

$$(\forall x \in \text{dom } h) \quad \|\text{prox}_{\gamma h}(x) - x\|^2 \leq 2\gamma \left(h(x) + h^*(u) - \langle x | u \rangle + \gamma \frac{\|u\|^2}{2} \right). \quad (2.7)$$

In particular, for every $x \in \text{dom } h$, $\text{prox}_{\gamma h}(x) \rightarrow x$ as $\gamma \rightarrow 0^+$.

Proof. Let $x \in \text{dom } h$ and set, for the sake of brevity, $p_\gamma = \text{prox}_{\gamma h}(x)$. It follows from (2.6) that $(x - p_\gamma)/\gamma \in \partial h(p_\gamma)$, hence

$$h(p_\gamma) + \frac{1}{\gamma} \|x - p_\gamma\|^2 = \left\langle x - p_\gamma, \frac{x - p_\gamma}{\gamma} \right\rangle + h(p_\gamma) \leq h(x). \quad (2.8)$$

Let $u \in \text{dom } h^*$. Then, since $\langle p_\gamma, u \rangle - h^*(u) \leq h(p_\gamma)$, we have

$$\begin{aligned} \frac{1}{2\gamma} \|p_\gamma - x + \gamma u\|^2 - \frac{\gamma}{2} \|u\|^2 + \langle x, u \rangle - h^*(u) + \frac{1}{2\gamma} \|p_\gamma - x\|^2 \\ = \langle p_\gamma, u \rangle - h^*(u) + \frac{1}{\gamma} \|p_\gamma - x\|^2 \leq h(x). \end{aligned}$$

Hence (2.7) follows. □

3 Convergence analysis

In this section we first discuss the hypotheses and the well-posedness of the procedures [LS1–LS4](#). Then, we give a general convergence principle for abstract variable metric descent algorithm (Theorem [3.10](#)). Finally, we study the role and relationships among the proposed line search rules (Proposition [3.13](#)) and prove the convergence properties of algorithm (VM-FBS) (Theorem [3.18](#)).

3.1 Preliminary results

We examine assumption **H2** and its consequences.

- Remark 3.1.** (i) If $\text{dom } g = \mathcal{H}$, **H2** is equivalent to requiring that f is Fréchet differentiable on \mathcal{H} and that ∇f is uniformly continuous on bounded sets (see Corollary [3.4\(ii\)](#)).
- (ii) **H2** is satisfied if ∇f is Lipschitz continuous on the weak compacts of $\text{dom } g$.
- (iii) By Remark [2.5](#), **H2** implies $\text{dom } g \subset \text{int dom } f$ and f is continuous on $\text{dom } g$.
- (iv) **H2** implies that the function $(\nabla f)|_{\text{dom } g}: \text{dom } g \rightarrow \mathcal{H}$ is continuous — in the relative topology of $\text{dom } g$ (see Corollary [3.4\(i\)](#) below).
- (v) Since continuity on compact sets yields uniform continuity (Heine-Cantor theorem), if \mathcal{H} is finite dimensional, hypothesis **H2** turns to require that f is Gâteaux differentiable on $\text{dom } g$ and that $(\nabla f)|_{\text{dom } g}: \text{dom } g \rightarrow \mathcal{H}$ is continuous (with respect to the relative topology of $\text{dom } g$).

The following lemmas are at the basis of our convergence analysis.

Lemma 3.2. *Let $(x_k)_{k \in \mathbb{N}}$ and $(\tilde{x}_k)_{k \in \mathbb{N}}$ be two sequences in \mathcal{H} , let $\bar{x} \in \mathcal{H}$ and suppose that $x_k \rightharpoonup \bar{x}$ and $\tilde{x}_k \rightharpoonup \bar{x}$. Then $\{\bar{x}\} \cup \bigcup_{k \in \mathbb{N}} [x_k, \tilde{x}_k]$ is weakly compact. In particular $\bigcup_{k \in \mathbb{N}} [x_k, \bar{x}]$ is weakly compact.*

Proof. Let us denote by \mathcal{H}_w the space \mathcal{H} endowed with the weak topology. We recall that \mathcal{H}_w is a locally convex space [\[37\]](#). Set $A = \{\bar{x}\} \cup \bigcup_{k \in \mathbb{N}} [x_k, \tilde{x}_k]$ and let $(U_i)_{i \in I}$ be an open covering of A in \mathcal{H}_w . Then there exists $i_* \in I$ such that $\bar{x} \in U_{i_*}$. Thus, since U_{i_*} is a weak neighborhood of \bar{x} , there exists a convex neighborhood V of the origin in \mathcal{H}_w such that $\bar{x} + V + V \subset U_{i_*}$. Since $\bar{x} + V$ is a weak neighborhood of \bar{x} , $x_k \rightharpoonup \bar{x}$ and $\tilde{x}_k - x_k \rightharpoonup 0$, there exists $\nu \in \mathbb{N}$ such that for every integer $k > \nu$, we have $x_k \in \bar{x} + V$ and $\tilde{x}_k - x_k \in V$; hence, for every $t \in [0, 1]$, $x_k + t(\tilde{x}_k - x_k) \in \bar{x} + V + V \subset U_{i_*}$. Moreover, for every integer $k \leq \nu$, since $[x_k, \tilde{x}_k]$ is weakly compact, there exists a finite $I_k \subset I$ such that $[x_k, \tilde{x}_k] \subset \bigcup_{i \in I_k} U_i$. Eventually, setting $\tilde{I} = \bigcup_{k=0}^{\nu} I_k$ (which is finite), we have $A \subset U_{i_*} \cup \bigcup_{i \in \tilde{I}} U_i$. The second part of the statement follows from the first part by just taking $\tilde{x}_k = \bar{x}$, for every $k \in \mathbb{N}$. \square

Lemma 3.3. *Let Ω be an open subset of \mathcal{H} and let $f: \Omega \rightarrow \mathbb{R}$. Suppose that f is Gâteaux differentiable on a convex set $C \subset \Omega$. Let $(x_k)_{k \in \mathbb{N}}$ and $(\tilde{x}_k)_{k \in \mathbb{N}}$ be sequences in C and let $\bar{x} \in C$ be such that $x_k \rightharpoonup \bar{x}$ and $\tilde{x}_k - x_k \rightarrow 0$. Then the following hold:*

- (i) *Suppose that ∇f is uniformly continuous on any weak compact of C . Then*
- (a) $\|\nabla f(\tilde{x}_k) - \nabla f(x_k)\| \rightarrow 0$;
 - (b) $(\forall \varepsilon > 0)(\exists \delta > 0)(\forall k \in \mathbb{N})$
 $\|\tilde{x}_k - x_k\| \leq \delta \Rightarrow |f(\tilde{x}_k) - f(x_k) - \langle \tilde{x}_k - x_k | \nabla f(x_k) \rangle| \leq \varepsilon \|\tilde{x}_k - x_k\|$.
- (ii) *Suppose that ∇f is Lipschitz continuous on any weakly compact subset of C . Then there exists $L > 0$ such that, for every $k \in \mathbb{N}$, $\|\nabla f(\tilde{x}_k) - \nabla f(x_k)\| \leq L \|\tilde{x}_k - x_k\|$ and $|f(\tilde{x}_k) - f(x_k) - \langle \tilde{x}_k - x_k | \nabla f(x_k) \rangle| \leq L \|\tilde{x}_k - x_k\|^2 / 2$.*

Proof. It follows from Lemma 3.2 that $A = \{\bar{x}\} \cup \bigcup_{k \in \mathbb{N}} [x_k, \tilde{x}_k]$ is weakly compact, and moreover $A \subset C$.

(i): That $\|\nabla f(\tilde{x}_k) - \nabla f(x_k)\| \rightarrow 0$ follows from the fact that ∇f is uniformly continuous on A , that, for every $k \in \mathbb{N}$, $\tilde{x}_k, x_k \in A$, and that $\tilde{x}_k - x_k \rightarrow 0$. Let $\varepsilon > 0$. Then, since ∇f is uniformly continuous on A , there exists $\delta > 0$ such that, for every $x, y \in A$, $\|x - y\| \leq \delta \Rightarrow \|\nabla f(y) - \nabla f(x)\| \leq \varepsilon$. Let $k \in \mathbb{N}$ be such that $\|\tilde{x}_k - x_k\| \leq \delta$. Then $[x_k, \tilde{x}_k] \subset A$ and the function $t \mapsto f(x_k + t(\tilde{x}_k - x_k))$ is differentiable with derivative $t \mapsto \langle \tilde{x}_k - x_k | \nabla f(x_k + t(\tilde{x}_k - x_k)) \rangle$, which is continuous (for ∇f is uniformly continuous on A). Therefore, we have

$$\begin{aligned} & |f(\tilde{x}_k) - f(x_k) - \langle \tilde{x}_k - x_k | \nabla f(x_k) \rangle| \\ &= \left| \int_0^1 \langle \tilde{x}_k - x_k | \nabla f(x_k + t(\tilde{x}_k - x_k)) - \nabla f(x_k) \rangle dt \right| \leq \varepsilon \|\tilde{x}_k - x_k\|. \end{aligned}$$

(ii): Since ∇f is Lipschitz continuous on A , there exists $L > 0$ such that, for every $x, y \in A$, $\|\nabla f(x) - \nabla f(y)\| \leq L\|x - y\|$. Let $k \in \mathbb{N}$. Then $\|\nabla f(\tilde{x}_k) - \nabla f(x_k)\| \leq L\|\tilde{x}_k - x_k\|$. Moreover, since $[x_k, \tilde{x}_k] \subset A$, arguing as before, we have

$$|f(\tilde{x}_k) - f(x_k) - \langle \tilde{x}_k - x_k | \nabla f(x_k) \rangle| \leq \int_0^1 Lt \|\tilde{x}_k - x_k\|^2 dt = \frac{L}{2} \|\tilde{x}_k - x_k\|^2.$$

□

Corollary 3.4. *Let Ω be a nonempty open subset of \mathcal{H} and let $f: \Omega \rightarrow \mathbb{R}$. Suppose that f is Gâteaux differentiable on a nonempty convex set $C \subset \Omega$ and that ∇f is uniformly continuous on any weakly compact subset of C . Then*

- (i) $(\nabla f)|_C: C \rightarrow \mathcal{H}$ is continuous (in the relative topology of C).
- (ii) for every $\bar{x} \in C$

$$\lim_{\substack{x \rightarrow \bar{x} \\ x \in C, x \neq \bar{x}}} \frac{|f(x) - f(\bar{x}) - \langle x - \bar{x} | \nabla f(\bar{x}) \rangle|}{\|x - \bar{x}\|} = 0.$$

Proof. (i): For every $\bar{x} \in C$ and every $(\tilde{x}_k)_{k \in \mathbb{N}}$ in C such that $\tilde{x}_k \rightarrow \bar{x}$, it follows from Lemma 3.3(i)(a) (with $x_k = \bar{x}$, for every $k \in \mathbb{N}$) that $\|\nabla f(\tilde{x}_k) - \nabla f(\bar{x})\| \rightarrow 0$.

(ii): Let $\bar{x} \in C$. Then for every $(\tilde{x}_k)_{k \in \mathbb{N}}$ in $C \setminus \{\bar{x}\}$ such that $\tilde{x}_k \rightarrow \bar{x}$, Lemma 3.3(i)(b) (with $x_k \equiv \bar{x}$) yields $|f(\tilde{x}_k) - f(\bar{x}) - \langle \tilde{x}_k - \bar{x} | \nabla f(\bar{x}) \rangle| / \|\tilde{x}_k - \bar{x}\| \rightarrow 0$. □

Lemma 3.5. *Assume that H1 holds and that f is Gâteaux differentiable on $\text{dom } g$. Let $x \in \text{dom } g$ and set, for every $\gamma \in \mathbb{R}_{++}$ and $\lambda \in]0, 1]$,*

$$J(x, \gamma, \lambda) = x + \lambda(\text{prox}_{\gamma g}(x - \gamma \nabla f(x)) - x). \quad (3.1)$$

Then the following hold.

- (i) Let $\lambda \in]0, 1]$. Then for every $(\gamma_1, \gamma_2) \in \mathbb{R}_{++}^2$,

$$\gamma_1 \leq \gamma_2 \Rightarrow \|J(x, \gamma_1, \lambda) - x\| \leq \|J(x, \gamma_2, \lambda) - x\| \leq \frac{\gamma_2}{\gamma_1} \|J(x, \gamma_1, \lambda) - x\|.$$

- (ii) Let $\gamma \in \mathbb{R}_{++}$. Then for every $(\lambda_1, \lambda_2) \in]0, 1]^2$,

$$\lambda_1 \leq \lambda_2 \Rightarrow \|J(x, \gamma, \lambda_1) - x\| \leq \|J(x, \gamma, \lambda_2) - x\| = \frac{\lambda_2}{\lambda_1} \|J(x, \gamma, \lambda_1) - x\|.$$

(iii) $\forall \lambda \in]0, 1] \lim_{\gamma \rightarrow 0^+} J(x, \gamma, \lambda) = x$ and $\forall \gamma \in \mathbb{R}_{++} \lim_{\lambda \rightarrow 0^+} J(x, \gamma, \lambda) = x$.

(iv) Let $u \in \text{dom } g^*$. Then for every $x \in \text{dom } g$ and every $\gamma > 0$,

$$\|J(x, \gamma, 1) - x\| \leq \gamma \|\nabla f(x)\| + \sqrt{2\gamma(g(x) + g^*(u) - \langle x | u \rangle + \gamma \|u\|^2 / 2)}. \quad (3.2)$$

Proof. (i): It is a consequence of the fact that $\gamma \mapsto \|J(x, \gamma, 1) - x\|$ is increasing and that $\gamma \mapsto \|J(x, \gamma, 1) - x\| / \gamma$ is decreasing: see [14, Lemma 2.2] for the projection case and [42, Lemma 3] and [6, Lemma 2.4] for the general case.

(ii): It is trivial.

(iii): We prove the first part. Let $\gamma \in \mathbb{R}_{++}$. Then, since $\text{prox}_{\gamma g}$ is non-expansive

$$\begin{aligned} \frac{\|J(x, \gamma, \lambda) - x\|}{\lambda} &\leq \|\text{prox}_{\gamma g}(x - \gamma \nabla f(x)) - \text{prox}_{\gamma g}(x)\| + \|\text{prox}_{\gamma g}(x) - x\| \\ &\leq \gamma \|\nabla f(x)\| + \|\text{prox}_{\gamma g}(x) - x\|. \end{aligned} \quad (3.3)$$

Therefore, we derive from Fact 2.8 that $\|J(x, \gamma, \lambda) - x\| \rightarrow 0$ as $\gamma \rightarrow 0^+$.

(iv): It follows from (3.3) and Fact 2.8. \square

Finally the following lemma addresses the well-posedness of the definitions of the various proposed line search procedures.

Lemma 3.6. Assume that **H1** holds and that f is Gâteaux differentiable on $\text{dom } g$. Let J as in (3.1) and let $x \in \text{dom } g$. Suppose that $x \notin \text{argmin}(f + g)$. Then

(i) If $(\nabla f)|_{\text{dom } g} : \text{dom } g \rightarrow \mathcal{H}$ is continuous at x , then

$$(\forall \lambda > 0) \lim_{\gamma \rightarrow 0^+} \frac{\gamma \|\nabla f(J(x, \gamma, \lambda)) - \nabla f(x)\|}{\|J(x, \gamma, \lambda) - x\|} = 0. \quad (3.4)$$

(ii) If ∇f is uniformly continuous on any weakly compact subsets of $\text{dom } g$, then

$$(\forall \lambda > 0) \lim_{\gamma \rightarrow 0^+} \frac{\gamma |f(J(x, \gamma, \lambda)) - f(x) - \langle J(x, \gamma, \lambda) - x | \nabla f(x) \rangle|}{\|J(x, \gamma, \lambda) - x\|^2} = 0 \quad (3.5)$$

and

$$(\forall \gamma > 0) \lim_{\lambda \rightarrow 0^+} \frac{\lambda |f(J(x, \gamma, \lambda)) - f(x) - \langle J(x, \gamma, \lambda) - x | \nabla f(x) \rangle|}{\|J(x, \gamma, \lambda) - x\|^2} = 0. \quad (3.6)$$

(iii) Let $\gamma > 0$ and $y \in \text{dom } g$. Suppose that $y - x$ is a descent direction for $f + g$ at x , that is $(f + g)'(x, y - x) < 0$. Then, for every $\delta \in]0, 1[$ there exists $\lambda_0 \in]0, 1]$ such that, for every $\lambda \in]0, \lambda_0]$,

$$(f + g)(x + \lambda(y - x)) - (f + g)(x) \leq (1 - \delta)\lambda(g(y) - g(x) + \langle y - x | \nabla f(x) \rangle).$$

Proof. (i)-(ii): We first note that, using (2.6), for every $\gamma > 0$, we have

$$x \in \text{argmin}(f + g) \iff -\nabla f(x) \in \partial g(x) \iff x = \text{prox}_{\gamma g}(x - \gamma \nabla f(x)).$$

Let $\lambda > 0$. Then, since $x \notin \text{argmin}(f + g)$, we have $\|J(x, \gamma, \lambda) - x\| \neq 0$, for every $\gamma > 0$. Next, since $\gamma \mapsto \|J(x, \gamma, \lambda) - x\| / \gamma$ is decreasing, $\lim_{\gamma \rightarrow 0^+} \|J(x, \gamma, \lambda) - x\| / \gamma = \sup_{\gamma > 0} \|J(x, \gamma, \lambda) - x\| / \gamma > 0$, and hence there exists $(\gamma_0, M) \in \mathbb{R}_{++}^2$ such that

$$\frac{\gamma}{\|J(x, \gamma, \lambda) - x\|} \leq M \quad \forall \gamma \in]0, \gamma_0]. \quad (3.7)$$

Moreover $\lim_{\gamma \rightarrow 0^+} J(x, \gamma, \lambda) = x$. Thus, if $(\nabla f)|_{\text{dom } g}$ is continuous at x (case (i)),

$$\lim_{\gamma \rightarrow 0^+} \|\nabla f(J(x, \gamma, \lambda)) - \nabla f(x)\| = 0, \quad (3.8)$$

otherwise, if ∇f is uniformly continuous on any compact subsets of $\text{dom } g$ (case (ii)), we derive from Corollary 3.4(ii) that

$$\lim_{\gamma \rightarrow 0^+} \frac{|f(J(x, \gamma, \lambda)) - f(x) - \langle J(x, \gamma, \lambda) - x | \nabla f(x) \rangle|}{\|J(x, \gamma, \lambda) - x\|} = 0. \quad (3.9)$$

Then, (3.4) follows from (3.8) and (3.7), whereas (3.5) follows from (3.9) and (3.7).

Since $\lim_{\lambda \rightarrow 0^+} J(x, \gamma, \lambda) = x$ and $(J(x, \gamma, \lambda) - x)/\lambda = \|\text{prox}_{\gamma g}(x - \gamma \nabla f(x)) - x\| \neq 0$, equation (3.6) follows from Corollary 3.4(ii), as done before.

(iii): We have

$$\frac{(f + g)(x + \lambda(y - x)) - (f + g)(x)}{\lambda} \rightarrow (f + g)'(x, y - x) \text{ as } \lambda \rightarrow 0 \quad (3.10)$$

and, since for every $z \in \text{dom } g$, $g'(x, z - x) \leq g(z) - g(x)$ [2, Proposition 17.2],

$$(f + g)'(x, y - x) = g'(x, y - x) + \langle y - x | \nabla f(x) \rangle \leq g(y) - g(x) + \langle y - x | \nabla f(x) \rangle. \quad (3.11)$$

Thus, if $-\infty < (f + g)'(x, y - x) < 0$, we have $(f + g)'(x, y - x) < (1 - \delta)(f + g)'(x, y - x)$ and hence, by (3.11),

$$(f + g)'(x, y - x) < (1 - \delta)(g(y) - g(x) + \langle y - x | \nabla f(x) \rangle). \quad (3.12)$$

Otherwise, if $(f + g)'(x, y - x) = -\infty$, clearly (3.12) still holds. Therefore, in any case, it follows from (3.10) and (3.12) that there exists $\lambda_0 > 0$ such that, $\forall \lambda \in]0, \lambda_0]$,

$$\frac{(f + g)(x + \lambda(y - x)) - (f + g)(x)}{\lambda} \leq (1 - \delta)(g(y) - g(x) + \langle y - x | \nabla f(x) \rangle).$$

□

Remark 3.7. (i) In view of Lemma 3.6(i)-(iii) (applied with respect to each metric $\langle \cdot | \cdot \rangle_k$) and (2.4), the line searches LS1, LS2, and LS4 are well-defined for every $\delta, \theta \in]0, 1[$ and every $\bar{\lambda} \in]0, 1]$ and $\bar{\gamma} > 0$; and, since $y_k - x_k$ is a descent direction for $f + g$, when $y_k \neq x_k$ (see the subsequent Lemma 3.12), LS3 is well-defined too.

(ii) The line search methods we presented have different computational costs. LS2 and LS3 are the cheapest one since they require just one evaluation of ∇f and prox_g ; LS1 requires multiple evaluation of the proximity operator of g , therefore it is feasible when computing prox_g is cheap. LS4 is the most costly since it demands also to compute ∇f multiple times.

3.2 An abstract convergence principle

We present an abstract convergence theorem underlying the different versions of the variable metric forward-backward splitting algorithm we will consider. It uses the property (b) below that blends the concept of quasi-Fejér sequence with that of a sufficient decreasing condition. This result has the same flavor of that given in [1, Section 2.3].

Proposition 3.8. Let $(|\cdot|_k)_{k \in \mathbb{N}}$ be a sequence of Hilbert norms on \mathcal{H} such that the sequence of the associated positive operators is strongly (that is pointwise) convergent in $(\mathcal{H}, \|\cdot\|)$. Suppose that there exists $\nu > 0$ such that, for every $k \in \mathbb{N}$, $\nu \|\cdot\|^2 \leq |\cdot|_k^2$. Let $h \in \Gamma_0(\mathcal{H})$ and let $(x_k)_{k \in \mathbb{N}}$ be a sequence in $\text{dom } h$. Set $S_* = \text{argmin}_{\mathcal{H}} h$ and $S = \{x \in \mathcal{H} \mid h(x) \leq \inf_{k \in \mathbb{N}} h(x_k)\}$. Consider the following properties

- (a) $(h(x_k))_{k \in \mathbb{N}}$ is decreasing;
- (b) There exist $(\alpha_k)_{k \in \mathbb{N}}$ in $\mathbb{R}_{++}^{\mathbb{N}}$, with $\sup_{k \in \mathbb{N}} \alpha_k < +\infty$, and $(\eta_k)_{k \in \mathbb{N}}$ and $(\varepsilon_k)_{k \in \mathbb{N}}$ in ℓ_+^1 , such that

$$(\forall x \in \text{dom } h)(\forall k \in \mathbb{N}) \quad |x_{k+1} - x|_{k+1}^2 \leq (1 + \eta_k) |x_k - x|_k^2 + 2\alpha_k (h(x) - h(x_{k+1})) + \varepsilon_k. \quad (3.13)$$

- (c) There exist $(y_k)_{k \in \mathbb{N}}$ and $(v_k)_{k \in \mathbb{N}}$ in $\mathcal{H}^{\mathbb{N}}$ such that, $\forall k \in \mathbb{N}$, $v_k \in \partial h(y_k)$ and for every weakly convergent subsequence $(x_{n_k})_{k \in \mathbb{N}}$ of $(x_k)_{k \in \mathbb{N}}$, $x_{n_k} - y_{n_k} \rightarrow 0$ and $v_{n_k} \rightarrow 0$.

Then the following hold.

1. Suppose that (a) is satisfied.
 - (i) If $(x_k)_{k \in \mathbb{N}}$ admits a bounded subsequence, then $\inf_{k \in \mathbb{N}} h(x_k) > -\infty$ and $S \neq \emptyset$.²
 - (ii) If $\inf_{k \in \mathbb{N}} h(x_k) = -\infty$, then $\|x_k\| \rightarrow +\infty$ and $h(x_k) \rightarrow \inf_{\mathcal{H}} h$.
2. Suppose that (a) and (b) are satisfied.
 - (iii) If $\inf_{k \in \mathbb{N}} h(x_k) > -\infty$, then $\sum_{k \in \mathbb{N}} \|x_{k+1} - x_k\|^2 < +\infty$.
 - (iv) If $S \neq \emptyset$, then $(x_k)_{k \in \mathbb{N}}$ is a quasi-Fejér sequence with respect to S relative to the sequence of norms $(|\cdot|_k)_{k \in \mathbb{N}}$.
 - (v) Suppose that $S_* \neq \emptyset$. If $\sum_{k \in \mathbb{N}} \alpha_k = +\infty$, then $h(x_k) \rightarrow \inf_{\mathcal{H}} h$. If $\inf_{k \in \mathbb{N}} \alpha_k > 0$, then $(h(x_k) - \inf_{\mathcal{H}} h) = o(1/k)$.
3. Suppose that (a), (b), and (c) are satisfied.
 - (vi) If $\bar{x} \in \mathcal{H}$ and $(x_{n_k})_{k \in \mathbb{N}}$ is a subsequence of $(x_k)_{k \in \mathbb{N}}$ such that $x_{n_k} \rightharpoonup \bar{x}$, then $\bar{x} \in S_*$ and $h(y_{n_k}) \rightarrow \inf_{\mathcal{H}} h$.
 - (vii) Suppose that $S_* \neq \emptyset$. Then $(x_k)_{k \in \mathbb{N}}$ and $(y_k)_{k \in \mathbb{N}}$ converge weakly to the same point in S_* and $h(y_k) \rightarrow \inf_{\mathcal{H}} h$.
 - (viii) Suppose that $S_* = \emptyset$. Then $\|x_k\| \rightarrow +\infty$ and $h(x_k) \rightarrow \inf_{\mathcal{H}} h$.

Proof. (i): Suppose that $(x_k)_{k \in \mathbb{N}}$ has a bounded subsequence. Then there exists a subsequence $(x_{n_k})_{k \in \mathbb{N}}$ of $(x_k)_{k \in \mathbb{N}}$ and $\bar{x} \in \mathcal{H}$ such that $x_{n_k} \rightharpoonup \bar{x}$. Since h is lower semicontinuous and $(h(x_k))_{k \in \mathbb{N}}$ is decreasing, $-\infty < h(\bar{x}) \leq \liminf_k h(x_k) = \inf_{k \in \mathbb{N}} h(x_k)$.

(ii): Since $\inf_{k \in \mathbb{N}} h(x_k) = -\infty$ and $(h(x_k))_{k \in \mathbb{N}}$ is decreasing, then $h(x_k) \rightarrow -\infty = \inf_{\mathcal{H}} h$. Moreover, it follows from (i) that $(x_k)_{k \in \mathbb{N}}$ does not have any bounded subsequence, hence $\liminf_k \|x_k\| \rightarrow +\infty$.

(iii): Taking $x = x_k$ in (3.13), we have

$$(\forall k \in \mathbb{N}) \quad \nu \|x_{k+1} - x_k\|^2 \leq |x_{k+1} - x_k|_{k+1}^2 \leq 2\alpha_k (h(x_k) - h(x_{k+1})) + \varepsilon_k. \quad (3.14)$$

Now note that, if $\inf_{k \in \mathbb{N}} h(x_k) > -\infty$, the sequence $(h(x_k) - h(x_{k+1}))_{k \in \mathbb{N}}$ is summable (and positive). Therefore, since $(\alpha_k)_{k \in \mathbb{N}}$ is bounded and $(\varepsilon_k)_{k \in \mathbb{N}}$ is summable, the right hand side of (3.14) is summable.

² $S_* \subset S$ and, since h is a proper function, $S \neq \emptyset \Rightarrow \inf_{k \in \mathbb{N}} h(x_k) > -\infty$.

(iv): Suppose that $S \neq \emptyset$ and let $x_* \in S$. Then it follows from (3.13) with $x = x_*$ that, for every $k \in \mathbb{N}$, $|x_{k+1} - x_*|_{k+1}^2 \leq (1 + \eta_k) |x_k - x_*|_k^2 + \varepsilon_k$ (see Definition 2.2).

(v): Let $x_* \in S_*$. It follows from (iv) that $(x_k)_{k \in \mathbb{N}}$ is a quasi-Fejer sequence with respect to S_* relative to $(|\cdot|_k)_{k \in \mathbb{N}}$, hence Fact 2.3(i) yields that $(|x_k - x_*|_k^2)_{k \in \mathbb{N}}$ is bounded. It follows from (3.13) with $x = x_*$ that, for every $k \in \mathbb{N}$

$$2\alpha_k(h(x_{k+1}) - h(x_*)) \leq |x_k - x_*|_k^2 - |x_{k+1} - x_*|_{k+1}^2 + \eta_k |x_k - x_*|_k^2 + \varepsilon_k. \quad (3.15)$$

Thus, $(h(x_{k+1}) - \inf_{\mathcal{H}} h)_{k \in \mathbb{N}}$ is a sequence in \mathbb{R}_+ which is decreasing and moreover, since $(|x_k - x_*|_k^2)_{k \in \mathbb{N}}$ is bounded, the right hand side of (3.15) is summable, and hence the sequence $(\alpha_k(h(x_{k+1}) - \inf_{\mathcal{H}} h))_{k \in \mathbb{N}}$ is summable. The statement follows from Fact 2.1.

(vi): Let $\bar{x} \in \mathcal{H}$ and let $(x_{n_k})_{k \in \mathbb{N}}$ be a subsequence of $(x_k)_{k \in \mathbb{N}}$ such that $x_{n_k} \rightharpoonup \bar{x}$. Since $y_{n_k} - x_{n_k} \rightarrow 0$, we have $y_{n_k} \rightarrow \bar{x}$. Then, since, for every $k \in \mathbb{N}$, $v_{n_k} \in \partial h(y_{n_k})$, $v_{n_k} \rightarrow 0$, and ∂h is demiclosed [2], then we have $0 \in \partial h(\bar{x})$, hence $\bar{x} \in S_*$. Moreover, $v_{n_k} \in \partial h(y_{n_k})$ yields,

$$(\forall k \in \mathbb{N}) \quad h(y_{n_k}) \leq h(\bar{x}) + \langle y_{n_k} - \bar{x} | v_{n_k} \rangle$$

and $\langle y_{n_k} - \bar{x} | v_{n_k} \rangle \rightarrow 0$, for $y_{n_k} \rightarrow \bar{x}$ and $v_{n_k} \rightarrow 0$. Thus, by the lower semicontinuity of h , $h(\bar{x}) \leq \liminf_k h(y_{n_k}) \leq \limsup_k h(y_{n_k}) \leq h(\bar{x})$, that is $h(y_{n_k}) \rightarrow h(\bar{x}) = \inf_{\mathcal{H}} h$.

(vii): Suppose $S_* \neq \emptyset$. Then, by (iv), $(x_k)_{k \in \mathbb{N}}$ is a quasi-Fejér sequence with respect to S_* relative to $(|\cdot|_k)_{k \in \mathbb{N}}$. Moreover, (vi) yields that every weak sequential cluster point of $(x_k)_{k \in \mathbb{N}}$ belongs to S_* . Thus, it follows from Fact 2.3(ii) that $(x_k)_{k \in \mathbb{N}}$ converges weakly to a point in S_* . Then, by applying (vi) and property (c) to the entire sequence $(x_k)_{k \in \mathbb{N}}$ we derive that $h(y_k) \rightarrow \inf_{\mathcal{H}} h$ and $y_k - x_k \rightarrow 0$.

(viii): It follows from (vi) that $(x_k)_{k \in \mathbb{N}}$ does not have any bounded subsequence. Therefore $\liminf_k \|x_k\| = +\infty$. If it was $\inf_k h(x_k) > \inf_{\mathcal{H}} h$, then the set S would be nonempty and (iv) would yield that $(x_k)_{k \in \mathbb{N}}$ is a quasi-Fejér sequence and hence bounded. Thus, necessarily $h(x_k) \rightarrow \inf_k h(x_k) = \inf_{\mathcal{H}} h$. \square

Remark 3.9. If in (3.13) we consider stationary metrics and replace $h(x_{k+1})$ by $h(x_k)$, we obtain the notion of *modified Fejér sequences* introduced in [32]. The authors show that that concept is useful to analyze the convergence in function values of splitting algorithms. However the convergence of the iterates is not studied.

Now we give the general theorem of convergence for variable metric algorithms.

Theorem 3.10. *Under the assumption of Proposition 3.8, suppose that (a) in Proposition 3.8 is satisfied and that, if $\inf_{k \in \mathbb{N}} h(x_k) > -\infty$, conditions (b) and (c) in Proposition 3.8 are satisfied for some $(\alpha_k)_{k \in \mathbb{N}}$ and $(y_k)_{k \in \mathbb{N}}$. Then the following hold.*

- (i) *If $\inf_{k \in \mathbb{N}} h(x_k) > -\infty$, then $\sum_{k \in \mathbb{N}} \|x_{k+1} - x_k\|^2 < +\infty$.*
- (ii) *Suppose that $S_* \neq \emptyset$. Then*
 - (a) *$(x_k)_{k \in \mathbb{N}}$ and $(y_k)_{k \in \mathbb{N}}$ weakly converge to the same point in S_* .*
 - (b) *$h(y_k) \rightarrow \inf_{\mathcal{H}} h$.*
 - (c) *If $\sum_{k \in \mathbb{N}} \alpha_k = +\infty$, then $h(x_k) \rightarrow \inf_{\mathcal{H}} h$.*
 - (d) *If $\inf_{k \in \mathbb{N}} \alpha_k > 0$, then $h(x_k) - \inf_{\mathcal{H}} h = o(1/k)$.*
- (iii) *If $S_* = \emptyset$, then $\|x_k\| \rightarrow +\infty$ and $h(x_k) \rightarrow \inf_{\mathcal{H}} h$.*

Proof. If $\inf_{k \in \mathbb{N}} h(x_k) = -\infty$, we are in the case $S_* = \emptyset$ and the statement follows from Proposition 3.8(ii). If $\inf_{k \in \mathbb{N}} h(x_k) > -\infty$, then conditions (a), (b) and (c) in Proposition 3.8 are satisfied and the conclusions follow from Proposition 3.8. \square

3.3 Convergence theorems

In this section we finally address the convergence of (VM-FBS) with line searches [LS1–LS4](#). In addition to **H1–H3**, we will also consider one of the following assumptions on the metrics.

H4 There exists $(\eta_k)_{k \in \mathbb{N}} \in \ell_+^1$ such that, for every $k \in \mathbb{N}$, $\|\cdot\|_{k+1}^2 \leq (1 + \eta_k) \|\cdot\|_k^2$.

H5 $\sum_{k \in \mathbb{N}} (\mu_k - \nu_k) < +\infty$, where, $\forall k \in \mathbb{N}$, ν_k and μ_k are respectively the minimum and maximum eigenvalue of the positive operator W_k associated to $\langle \cdot | \cdot \rangle_k$.

Remark 3.11.

- (i) **H4** can be equivalently written as $W_{k+1} \preceq (1 + \eta_k)W_k$ and it was considered in [36] for the proximal point algorithm and in [9, 19] for the forward-backward algorithm. In view of Fact 2.3, **H4** implies that W_k strongly converges to some positive operator.
- (ii) **H5** encompasses and generalizes the condition assumed in [10] for the scaled gradient projection method, where the W_k 's are indeed forced to converge to the identity operator at certain rate. By contrast, we stress that **H5** does not implies that the W_k 's strongly converge: just take $W_k = \mu_k \text{Id}$ with $(\mu_k)_{k \in \mathbb{N}}$ a non convergent bounded sequence in $[\varepsilon, +\infty[$ for some $\varepsilon > 0$. However, **H5** implies that, as $k \rightarrow +\infty$, W_k takes the form of a multiple of the identity operator, but the multiplicative constant may continue to vary with k .

The following result is fundamental and analyzes just one step of (VM-FBS). So, we can avoid to refer to the variable metric and state the result in the metric $\langle \cdot | \cdot \rangle$ of the space \mathcal{H} . Items (i)–(ii) below are standard and appear explicitly in [42, Lemma 1].

Lemma 3.12. *Assume **H1** and that f is Gâteaux differentiable on $\text{dom } g$. Let $k \in \mathbb{N}$ and let $\gamma_k \in \mathbb{R}_{++}$ and $\lambda_k \in]0, 1]$. Let $x_k \in \text{dom } g$ and set $y_k = \text{prox}_{\gamma_k g}(x_k - \gamma_k \nabla f(x_k))$ and $x_{k+1} = J(x_k, \gamma_k, \lambda_k) = x_k + \lambda_k(y_k - x_k)$. Then the following hold.*

- (i) $g(y_k) - g(x_k) + \langle y_k - x_k | \nabla f(x_k) \rangle \leq -\frac{\|y_k - x_k\|^2}{\gamma_k}$.
- (ii) $(f + g)'(x_k, y_k - x_k) \leq -\frac{\|y_k - x_k\|^2}{\gamma_k}$; in particular if $y_k \neq x_k$, then $y_k - x_k$ is a descent direction for $f + g$.
- (iii) $\forall x \in \text{dom } g$,

$$\begin{aligned} \|x_{k+1} - x\|^2 &\leq \|x_k - x\|^2 + 2\gamma_k \lambda_k ((f + g)(x) - (f + g)(x_k)) \\ &\quad - 2\gamma_k \lambda_k (g(y_k) - g(x_k) + \langle y_k - x_k | \nabla f(x_k) \rangle) - \|x_{k+1} - x_k\|^2. \end{aligned}$$

Proof. (i): By the definition of y_k and (2.6) we derive that $(x_k - y_k)/\gamma_k - \nabla f(x_k) \in \partial g(y_k)$, and hence that, for every $x \in \text{dom } g$,

$$\begin{aligned} g(x) &\geq g(y_k) + \left\langle x - x_k \left| \frac{x_k - y_k}{\gamma_k} - \nabla f(x_k) \right. \right\rangle + \left\langle x_k - y_k \left| \frac{x_k - y_k}{\gamma_k} - \nabla f(x_k) \right. \right\rangle \\ &= g(y_k) + \frac{1}{\gamma_k} \langle x - x_k | x_k - y_k \rangle - \langle x - y_k | \nabla f(x_k) \rangle + \frac{\|x_k - y_k\|^2}{\gamma_k}. \end{aligned} \tag{3.16}$$

Taking $x = x_k$ in the above inequality, (i) follows.

(ii): It follows from (i) and the fact that $g'(x_k, y_k - x_k) \leq g(y_k) - g(x_k)$.

(iii): Let $x \in \text{dom } g$. Since f is convex, $f(x) - f(x_k) \geq \langle x - x_k | \nabla f(x_k) \rangle$. Thus, it follows from (3.16) that

$$(f + g)(x) - (f + g)(x_k) \geq g(y_k) - g(x_k) + \langle y_k - x_k | \nabla f(x_k) \rangle + \frac{1}{\gamma_k} \langle x - x_k | x_k - y_k \rangle + \frac{\|y_k - x_k\|^2}{\gamma_k}.$$

Now, multiplying the above inequality by $2\gamma_k\lambda_k$ we obtain

$$2\gamma_k\lambda_k((f + g)(x) - (f + g)(x_k)) \geq 2\gamma_k\lambda_k(g(y_k) - g(x_k) + \langle y_k - x_k | \nabla f(x_k) \rangle) + 2\langle x - x_k | x_k - x_{k+1} \rangle + 2\frac{\|x_k - x_{k+1}\|^2}{\lambda_k}.$$

Finally, since $2\langle x - x_k | x_k - x_{k+1} \rangle = \|x_{k+1} - x\|^2 - \|x_k - x\|^2 - \|x_{k+1} - x_k\|^2$,

$$2\gamma_k\lambda_k((f + g)(x) - (f + g)(x_k)) \geq 2\gamma_k\lambda_k(g(y_k) - g(x_k) + \langle y_k - x_k | \nabla f(x_k) \rangle) + \|x_{k+1} - x\|^2 - \|x_k - x\|^2 - \|x_{k+1} - x_k\|^2 + 2\frac{\|x_{k+1} - x_k\|^2}{\lambda_k} \quad (3.17)$$

and hence, since $2/\lambda_k \geq 2$, the statement follows. \square

In view of Proposition 3.8(b) and Lemma 3.12(iii), it is clear that to obtain convergence of algorithm (VM-FBS), we need to ensure that the positive quantity

$$- (g(y_k) - g(x_k) + \langle y_k - x_k | \nabla f(x_k) \rangle) \quad (3.18)$$

is summable. Now we show how the various line search methods are related to each other and the role that they play in making (3.18) summable.

Proposition 3.13. *Under the same hypotheses of Lemma 3.12, let $\delta \in]0, 1[$ and consider the following statements*

- (a) $\|\nabla f(x_{k+1}) - \nabla f(x_k)\| \leq \frac{\delta}{\gamma_k\lambda_k} \|x_{k+1} - x_k\|.$
- (b) $f(x_{k+1}) - f(x_k) - \langle x_{k+1} - x_k | \nabla f(x_k) \rangle \leq \frac{\delta}{\gamma_k\lambda_k} \|x_{k+1} - x_k\|^2.$
- (c) $(f + g)(x_{k+1}) - (f + g)(x_k) \leq (1 - \delta)\lambda_k(g(y_k) - g(x_k) + \langle y_k - x_k | \nabla f(x_k) \rangle).$

Then the following hold.

- (i) (a) \Rightarrow (b) \Rightarrow (c).
- (ii) If (c) holds, then, for every $x \in \text{dom } g$,

$$\|x_{k+1} - x\|^2 \leq \|x_k - x\|^2 + 2\gamma_k\lambda_k((f + g)(x) - (f + g)(x_k)) + \frac{2\gamma_k}{1 - \delta}((f + g)(x_k) - (f + g)(x_{k+1})) - \|x_{k+1} - x_k\|^2. \quad (3.19)$$

- (iii) If (c) holds, then

$$(1 - \delta)\|x_{k+1} - x_k\|^2 \leq \gamma_k((f + g)(x_k) - (f + g)(x_{k+1}));$$

in particular $(f + g)(x_{k+1}) \leq (f + g)(x_k).$

Proof. (i): Suppose that (a) holds. Then, using the convexity of f , we derive that

$$\begin{aligned} f(x_k) &\geq f(x_{k+1}) + \langle x_k - x_{k+1} | \nabla f(x_{k+1}) - \nabla f(x_k) \rangle + \langle x_k - x_{k+1} | \nabla f(x_k) \rangle \\ &\geq f(x_{k+1}) - \frac{\delta}{\gamma_k \lambda_k} \|x_k - x_{k+1}\|^2 + \langle x_k - x_{k+1} | \nabla f(x_k) \rangle. \end{aligned}$$

hence (b) holds. Now, suppose that (b) holds. Then, since $g(x_{k+1}) - g(x_k) \leq \lambda_k(g(y_k) - g(x_k))$, we have

$$\begin{aligned} (f+g)(x_{k+1}) - (f+g)(x_k) &\leq g(x_{k+1}) - g(x_k) + \langle x_{k+1} - x_k | \nabla f(x_k) \rangle + \frac{\delta}{\gamma_k \lambda_k} \|x_{k+1} - x_k\|^2 \\ &\leq \lambda_k \left(g(y_k) - g(x_k) + \langle y_k - x_k | \nabla f(x_k) \rangle + \frac{\delta}{\gamma_k} \|y_k - x_k\|^2 \right) \quad (3.20) \\ &= (1 - \delta) \lambda_k \left(g(y_k) - g(x_k) + \langle y_k - x_k | \nabla f(x_k) \rangle \right) \\ &\quad + \delta \lambda_k \left(g(y_k) - g(x_k) + \langle y_k - x_k | \nabla f(x_k) \rangle + \frac{1}{\gamma_k} \|y_k - x_k\|^2 \right). \end{aligned}$$

Therefore, recalling Lemma 3.12(i), (c) follows.

(ii): It follows from (c) that

$$-\lambda_k \left(g(y_k) - g(x_k) + \langle y_k - x_k | \nabla f(x_k) \rangle \right) \leq \frac{(f+g)(x_k) - (f+g)(x_{k+1})}{1 - \delta}$$

and hence, multiplying by $2\gamma_k$, the statement follows from Lemma 3.12(iii).

(iii): It follows from (3.19), by taking $x = x_k$. \square

Remark 3.14. Another condition that is in between (b) and (c) of Proposition 3.13 is the following

$$(f+g)(x_{k+1}) - (f+g)(x_k) \leq \sigma \lambda_k \left(g(y_k) - g(x_k) + \langle y_k - x_k | \nabla f(x_k) \rangle + \frac{\beta}{\gamma_k} \|y_k - x_k\|^2 \right), \quad (3.21)$$

where $(\sigma, \beta) \in [0, 1]^2$, $0 < (1 - \beta)\sigma < 1$, and $1 - \delta = (1 - \beta)\sigma$. More precisely, since

$$\begin{aligned} g(y_k) - g(x_k) + \langle y_k - x_k | \nabla f(x_k) \rangle + \frac{1 - (1 - \beta)\sigma}{\gamma_k} \|y_k - x_k\|^2 \\ = \sigma \left(g(y_k) - g(x_k) + \langle y_k - x_k | \nabla f(x_k) \rangle + \frac{\beta}{\gamma_k} \|y_k - x_k\|^2 \right) \\ + (1 - \sigma) \left(g(y_k) - g(x_k) + \langle y_k - x_k | \nabla f(x_k) \rangle + \frac{1}{\gamma_k} \|y_k - x_k\|^2 \right) \\ \leq \sigma \left(g(y_k) - g(x_k) + \langle y_k - x_k | \nabla f(x_k) \rangle + \frac{\beta}{\gamma_k} \|y_k - x_k\|^2 \right), \end{aligned}$$

if (b) holds with $\delta = 1 - (1 - \beta)\sigma$, then, it follows from (3.20) that (3.21) holds. Moreover, it follows from (3.20) and Lemma 3.12(i) that (3.21) \Rightarrow (c) with $\delta = 1 - (1 - \beta)\sigma$. Note that (3.21) includes (c) by choosing $\beta = 0$ and $\sigma = \delta$. Condition (3.21) (with $0 < \sigma < 1$ and $0 \leq \beta < 1$) is at the basis of the Armijo line search proposed by Tseng and Yun in [42], which is also adopted in [9] with $\beta \leq 1/2$. Finally, the second line search method in [6] corresponds to (3.21) with $\sigma = 1$, $\beta = \delta = 1/2$, and $\lambda_k \equiv 1$, and hence it leads to values of λ_k smaller than necessary: by choosing δ close to 1, a larger step along $y_k - x_k$ is obtained. In view of Proposition 3.13, Proposition 3.17, Corollary 3.15 below, and the discussion above we can claim that our convergence results hold also for the Armijo-type rule considered in [9, 42].

The following result treats the four line search methods in a unifying manner.

Corollary 3.15. *Assume that **H1** and **H3** hold and that f is Gâteaux differentiable on $\text{dom } g$. Let $(x_k)_{k \in \mathbb{N}}$ and $(y_k)_{k \in \mathbb{N}}$ be defined according to (VM-FBS) for some sequences $(\gamma_k)_{k \in \mathbb{N}}$ and $(\lambda_k)_{k \in \mathbb{N}}$. Suppose that one of (a), (b), or (c) in Proposition 3.13 is satisfied in the metric $\langle \cdot | \cdot \rangle_k$, for every $k \in \mathbb{N}$. Then, the following hold.*

- (i) *The sequence $((f + g)(x_k))_{k \in \mathbb{N}}$ is decreasing.*
- (ii) *If $\sup_{k \in \mathbb{N}} \gamma_k < +\infty$ and $\inf_{k \in \mathbb{N}} (f + g)(x_k) > -\infty$, then*

$$\frac{1}{\sup_{k \in \mathbb{N}} \gamma_k} \sum_{k \in \mathbb{N}} \|x_{k+1} - x_k\|_k^2 \leq \sum_{k \in \mathbb{N}} \frac{\|x_{k+1} - x_k\|_k^2}{\gamma_k} < +\infty.$$

- (iii) *If **H4** holds, then, for every $x \in \text{dom } g$ and for every $k \in \mathbb{N}$,*

$$\begin{aligned} \|x_{k+1} - x\|_{k+1}^2 &\leq (1 + \eta_k) \|x_k - x\|_k^2 + 2\gamma_k \lambda_k (1 + \eta_k) ((f + g)(x) - (f + g)(x_k)) \\ &\quad + \frac{2\gamma_k (1 + \eta_k)}{1 - \delta} ((f + g)(x_k) - (f + g)(x_{k+1})). \end{aligned}$$

- (iv) *If **H5** holds, then, for every $x \in \text{dom } g$ and for every $k \in \mathbb{N}$,*

$$\begin{aligned} \|x_{k+1} - x\|^2 &\leq \left(1 + \frac{\mu_k - \nu_k}{\nu}\right) \|x_k - x\|^2 + 2\frac{\gamma_k \lambda_k}{\nu_k} ((f + g)(x) - (f + g)(x_k)) \\ &\quad + \frac{2\gamma_k}{\nu(1 - \delta)} ((f + g)(x_k) - (f + g)(x_{k+1})). \end{aligned}$$

Proof. (i): Invoking Proposition 3.13(iii) for each metric $\langle \cdot | \cdot \rangle_k$, we derive that, for every $k \in \mathbb{N}$, $(f + g)(x_{k+1}) \leq (f + g)(x_k)$.

(ii): Since $\bar{\gamma} := \sup_{k \in \mathbb{N}} \gamma_k < +\infty$, we derive from Proposition 3.13(iii), applied with respect to each metric $\langle \cdot | \cdot \rangle_k$, that, for every $k \in \mathbb{N}$

$$\begin{aligned} \frac{1 - \delta}{\bar{\gamma}} \sum_{i=0}^k \|x_{i+1} - x_i\|_i^2 &\leq (1 - \delta) \sum_{i=0}^k \frac{\|x_{i+1} - x_i\|_i^2}{\gamma_i} \leq \sum_{i=0}^k ((f + g)(x_i) - (f + g)(x_{i+1})) \\ &\leq (f + g)(x_0) - \inf_{k \in \mathbb{N}} (f + g)(x_k) < +\infty. \end{aligned}$$

(iii): Let $k \in \mathbb{N}$ and let $x \in \text{dom } g$. It is enough to note that, since $-(f + g)(x_k) \leq -(f + g)(x_{k+1})$, Proposition 3.13(ii), applied with respect to the metric $\langle \cdot | \cdot \rangle_k$, yields

$$\begin{aligned} \|x_{k+1} - x\|_k^2 &\leq \|x_k - x\|_k^2 + 2\gamma_k \lambda_k ((f + g)(x) - (f + g)(x_{k+1})) \\ &\quad + \frac{2\gamma_k}{1 - \delta} ((f + g)(x_k) - (f + g)(x_{k+1})). \end{aligned} \tag{3.22}$$

Hence, multiplying by $(1 + \eta_k)$ and taking into account that $\|\cdot\|_{k+1}^2 \leq (1 + \eta_k) \|\cdot\|_k^2$ the statement follows.

(iv): It follows from (3.22) and (2.2) that

$$\begin{aligned} \nu_k \|x_{k+1} - x\|^2 &\leq \mu_k \|x_k - x\|^2 + 2\gamma_k \lambda_k ((f + g)(x) - (f + g)(x_{k+1})) \\ &\quad + \frac{2\gamma_k}{1 - \delta} ((f + g)(x_k) - (f + g)(x_{k+1})). \end{aligned} \tag{3.23}$$

Hence, dividing (3.23) by ν_k , and noting that $(f + g)(x_k) - (f + g)(x_{k+1}) \geq 0$, that $\nu \leq \nu_k$, and $\mu_k/\nu_k = 1 + (\mu_k - \nu_k)/\nu_k \leq 1 + (\mu_k - \nu_k)/\nu$, the statement follows. \square

Remark 3.16. Under the hypotheses of Corollary 3.15, if $(\gamma_k)_{k \in \mathbb{N}}$ is bounded and $\inf_{k \in \mathbb{N}}(f+g)(x_k) > -\infty$, then the following hold:

- (i) If **H4** is satisfied, then setting $\bar{\gamma} = \sup_{k \in \mathbb{N}} \gamma_k$ and $\bar{\eta} = \sup_{k \in \mathbb{N}} \eta_k$, condition (b) in Proposition 3.8 is fulfilled with $h = f + g$, $(\alpha_k)_{k \in \mathbb{N}} = (\gamma_k \lambda_k (1 + \eta_k))_{k \in \mathbb{N}}$, $(|\cdot|_k)_{k \in \mathbb{N}} = (\|\cdot\|_k)_{k \in \mathbb{N}}$, and

$$(\varepsilon_k)_{k \in \mathbb{N}} = 2\bar{\gamma}(1 + \bar{\eta})/(1 - \delta)((f + g)(x_k) - (f + g)(x_{k+1}))_{k \in \mathbb{N}}.$$

- (ii) If **H5** is satisfied, then setting $\bar{\gamma} = \sup_{k \in \mathbb{N}} \gamma_k$, condition (b) in Proposition 3.8 is fulfilled with $h = f + g$, $(\alpha_k)_{k \in \mathbb{N}} = (\gamma_k \lambda_k / \nu_k)_{k \in \mathbb{N}}$, $(\eta_k)_{k \in \mathbb{N}} = \nu^{-1}(\mu_k - \nu_k)_{k \in \mathbb{N}}$, $(|\cdot|_k)_{k \in \mathbb{N}} \equiv \|\cdot\|$, and

$$(\varepsilon_k)_{k \in \mathbb{N}} = 2\bar{\gamma}/(\nu(1 - \delta))((f + g)(x_k) - (f + g)(x_{k+1}))_{k \in \mathbb{N}}.$$

To finish our convergence analysis it remains to verify (c) in Proposition 3.8.

Proposition 3.17. Assume that **H1–H3** hold and define $(x_k)_{k \in \mathbb{N}}$ and $(y_k)_{k \in \mathbb{N}}$ as in (VM-FBS) with some $(\gamma_k)_{k \in \mathbb{N}}$ and $(\lambda_k)_{k \in \mathbb{N}}$. Let $\delta > 0$ and consider the following properties:

$$(a') \quad (\exists \theta \in]0, 1[)(\exists \sigma \in \mathbb{R}_{++})(\forall k \in \mathbb{N}) \quad \gamma_k < \sigma \Rightarrow$$

$$\|\nabla^k f(J_k(x_k, \gamma_k/\theta, \lambda_k)) - \nabla^k f(x_k)\|_k > \frac{\delta\theta}{\gamma_k \lambda_k} \|J_k(x_k, \gamma_k/\theta, \lambda_k) - x_k\|_k.$$

$$(b') \quad (\exists \theta \in]0, 1[)(\exists \sigma \in \mathbb{R}_{++})(\forall k \in \mathbb{N}) \quad \gamma_k < \sigma \Rightarrow$$

$$f(J_k(x_k, \gamma_k/\theta, \lambda_k)) - f(x_k) - \langle J_k(x_k, \gamma_k/\theta, \lambda_k) - x_k \mid \nabla f(x_k) \rangle > \frac{\delta\theta}{\gamma_k \lambda_k} \|J_k(x_k, \gamma_k/\theta, \lambda_k) - x_k\|_k^2.$$

$$(b'') \quad (\exists \theta \in]0, 1[)(\exists \sigma \in]0, 1])(\forall k \in \mathbb{N}) \quad \lambda_k \leq \sigma\theta \Rightarrow$$

$$f(J_k(x_k, \gamma_k, \lambda_k/\theta)) - f(x_k) - \langle J_k(x_k, \gamma_k, \lambda_k/\theta) - x_k \mid \nabla f(x_k) \rangle > \frac{\delta\theta}{\gamma_k \lambda_k} \|J_k(x_k, \gamma_k, \lambda_k/\theta) - x_k\|_k^2.$$

$$(c') \quad (\exists \theta \in]0, 1[)(\exists \sigma \in]0, 1])(\forall k \in \mathbb{N}) \quad \gamma_k < \sigma \Rightarrow$$

$$\begin{aligned} & (f + g)(J_k(x_k, \gamma_k/\theta, \lambda_k)) - (f + g)(x_k) \\ & > (1 - \delta)\lambda_k(g(J_k(x_k, \gamma_k/\theta, 1)) - g(x_k) + \langle J_k(x_k, \gamma_k/\theta, 1) - x_k \mid \nabla f(x_k) \rangle). \end{aligned}$$

$$(c'') \quad (\exists \theta \in]0, 1[)(\exists \sigma \in]0, 1])(\forall k \in \mathbb{N}) \quad \lambda_k \leq \sigma\theta \Rightarrow$$

$$(f + g)(J_k(x_k, \gamma_k, \lambda_k/\theta)) - (f + g)(x_k) > (1 - \delta)(\lambda_k/\delta)(g(y_k) - g(x_k) + \langle y_k - x_k \mid \nabla f(x_k) \rangle).$$

Then the following hold.

- (i) (c') \Rightarrow (b') \Rightarrow (a') and (c'') \Rightarrow (b'').
(ii) Suppose that $((f + g)(x_k))_{k \in \mathbb{N}}$ is decreasing, $\|x_{k+1} - x_k\|_k \rightarrow 0$, $\sup_{k \in \mathbb{N}} \gamma_k < +\infty$, and that either $\inf_{k \in \mathbb{N}} \lambda_k > 0$ and (a') hold, or that $\inf_{k \in \mathbb{N}} \gamma_k > 0$ and (b'') hold. Then, there exists $(v_k)_{k \in \mathbb{N}} \in \mathcal{H}^{\mathbb{N}}$ such that, $\forall k \in \mathbb{N}$, $v_k \in \partial(f + g)(y_k)$ and for every weakly convergent subsequence $(x_{n_k})_{k \in \mathbb{N}}$ of $(x_k)_{k \in \mathbb{N}}$, $y_{n_k} - x_{n_k} \rightarrow 0$ and $v_{n_k} \rightarrow 0$.

Proof. (i): It follows from Proposition 3.13(i), applied for each metric $\langle \cdot | \cdot \rangle_k$, by using a simple contradiction argument.

(ii): Let, for every $k \in \mathbb{N}$, $y_k = \text{prox}_{\gamma_k g}^k(x_k - \gamma_k \nabla^k f(x_k))$. Then, for every $k \in \mathbb{N}$, $(x_k - y_k)/\gamma_k + \nabla^k f(y_k) - \nabla^k f(x_k) \in \partial^k(f + g)(y_k)$, hence, by (2.3),

$$(\forall k \in \mathbb{N}) \quad v_k := W_k \frac{x_k - y_k}{\gamma_k} + \nabla f(y_k) - \nabla f(x_k) \in \partial(f + g)(y_k). \quad (3.24)$$

Let $(x_{n_k})_{k \in \mathbb{N}}$ be a subsequence of $(x_k)_{k \in \mathbb{N}}$ such that $x_{n_k} \rightharpoonup \bar{x}$ for some $\bar{x} \in \mathcal{H}$. We note that, since $((f+g)(x_k))_{k \in \mathbb{N}}$ is decreasing $\lim(f+g)(x_{n_k}) = \inf_k(f+g)(x_{n_k}) < +\infty$, hence it follows from the lower semicontinuity of $f+g$ that $\bar{x} \in \text{dom}(f+g) \subset \text{dom } g$.

Suppose first that (a') is satisfied and that $\inf_{k \in \mathbb{N}} \lambda_k > 0$. Let, for every $k \in \mathbb{N}$, $\tilde{\gamma}_k = \gamma_k/\theta$ and $\tilde{x}_k = J_k(x_k, \tilde{\gamma}_k, \lambda_k)$. Then, for every $k \in \mathbb{N}$, by Lemma 3.5(i)

$$\|J_k(x_k, \tilde{\gamma}_k, \lambda_k) - x_k\|_k \leq \frac{\tilde{\gamma}_k}{\gamma_k} \|J_k(x_k, \gamma_k, \lambda_k) - x_k\|_k = \frac{1}{\theta} \|x_{k+1} - x_k\|_k \quad (3.25)$$

and $\|x_{k+1} - x_k\|_k = \|J_k(x_k, \gamma_k, \lambda_k) - x_k\|_k \leq \|J_k(x_k, \tilde{\gamma}_k, \lambda_k) - x_k\|_k$; hence, recalling also (2.2),

$$\sqrt{\nu} \frac{\|y_k - x_k\|}{\gamma_k} \leq \frac{\|y_k - x_k\|_k}{\gamma_k} = \frac{\|x_{k+1} - x_k\|_k}{\gamma_k \lambda_k} \leq \frac{\|\tilde{x}_k - x_k\|_k}{\gamma_k \lambda_k}. \quad (3.26)$$

Moreover, since (a') is satisfied, then

$$(\forall k \in \mathbb{N}) \max \left\{ \|\nabla^k f(\tilde{x}_k) - \nabla^k f(x_k)\|_k, \frac{\delta \theta}{\sigma} \frac{\|\tilde{x}_k - x_k\|_k}{\lambda_k} \right\} \geq \delta \theta \frac{\|\tilde{x}_k - x_k\|_k}{\lambda_k \gamma_k}. \quad (3.27)$$

Now, since $\|x_{k+1} - x_k\|_k \rightarrow 0$, we derive from (3.25) and (2.2), that $\tilde{x}_k - x_k \rightarrow 0$ and since $x_{n_k} \rightharpoonup \bar{x}$, it follows from Lemma 3.3(i) that $\|\nabla f(\tilde{x}_{n_k}) - \nabla f(x_{n_k})\| \rightarrow 0$, hence, by Fact 2.7 and the fact that $\nu \leq \nu_k$, we have

$$\|\nabla^k f(\tilde{x}_{n_k}) - \nabla^k f(x_{n_k})\|_k \rightarrow 0. \quad (3.28)$$

Moreover,

$$\frac{\|\tilde{x}_k - x_k\|_k}{\lambda_k} \leq \frac{\|\tilde{x}_k - x_k\|_k}{\inf_{k \in \mathbb{N}} \lambda_k} \rightarrow 0. \quad (3.29)$$

Thus, (3.28), (3.29), and (3.27) imply $\|\tilde{x}_{n_k} - x_{n_k}\|_k / (\gamma_{n_k} \lambda_{n_k}) \rightarrow 0$ and so, by (3.26),

$$\frac{\|y_{n_k} - x_{n_k}\|}{\sup_{k \in \mathbb{N}} \gamma_k} \leq \frac{\|y_{n_k} - x_{n_k}\|}{\gamma_{n_k}} \rightarrow 0. \quad (3.30)$$

Finally, since $y_{n_k} - x_{n_k} \rightarrow 0$ and $x_{n_k} \rightharpoonup \bar{x}$, by another application of Lemma 3.3(i), we have that $\|\nabla f(y_{n_k}) - \nabla f(x_{n_k})\| \rightarrow 0$. This together with (3.30), (3.24) and the fact that $\sup_{k \in \mathbb{N}} \|W_k\| \leq \mu$, gives $v_{n_k} \rightarrow 0$.

Now suppose that (b'') is satisfied and that $\inf_{k \in \mathbb{N}} \gamma_k > 0$. Set, for every $k \in \mathbb{N}$, $\tilde{\lambda}_k = \lambda_k/\theta$ and $\tilde{x}_k = J_k(x_k, \gamma_k, \tilde{\lambda}_k)$. Then, by Lemma 3.5(ii), for every $k \in \mathbb{N}$,

$$\|J_k(x_k, \gamma_k, \tilde{\lambda}_k) - x_k\|_k = \frac{\tilde{\lambda}_k}{\lambda_k} \|J_k(x_k, \gamma_k, \lambda_k) - x_k\|_k = \frac{1}{\theta} \|x_{k+1} - x_k\|_k \quad (3.31)$$

and $\|x_{k+1} - x_k\|_k = \|J_k(x_k, \gamma_k, \lambda_k) - x_k\|_k \leq \|J_k(x_k, \gamma_k, \tilde{\lambda}_k) - x_k\|_k$; hence

$$\sqrt{\nu} \frac{\|y_k - x_k\|}{\gamma_k} \leq \frac{\|y_k - x_k\|_k}{\gamma_k} = \frac{\|x_{k+1} - x_k\|_k}{\gamma_k \lambda_k} \leq \frac{\|\tilde{x}_k - x_k\|_k}{\gamma_k \lambda_k}. \quad (3.32)$$

Moreover, since (b'') is satisfied,

$$(\forall k \in \mathbb{N}) \max \left\{ \frac{|f(\tilde{x}_k) - f(x_k) - \langle \tilde{x}_k - x_k, \nabla f(x_k) \rangle_k|}{\|\tilde{x}_k - x_k\|_k}, \frac{\delta}{\sigma} \frac{\|\tilde{x}_k - x_k\|_k}{\gamma_k} \right\} > \delta \theta \frac{\|\tilde{x}_k - x_k\|_k}{\gamma_k \lambda_k}. \quad (3.33)$$

Now, since $\|x_{k+1} - x_k\|_k \rightarrow 0$, we derive from (3.31) and (2.2), that $\tilde{x}_k - x_k \rightarrow 0$ and, since $x_{n_k} \rightharpoonup \bar{x}$, it follows from Lemma 3.3(i) that

$$\frac{|f(\tilde{x}_{n_k}) - f(x_{n_k}) - \langle \tilde{x}_{n_k} - x_{n_k} | \nabla f(x_{n_k}) \rangle|}{\|\tilde{x}_{n_k} - x_{n_k}\|} \rightarrow 0.$$

Since $\langle \tilde{x}_{n_k} - x_{n_k} | \nabla f(x_{n_k}) \rangle = \langle \tilde{x}_{n_k} - x_{n_k} | \nabla^k f(x_{n_k}) \rangle_k$ and $\nu^{1/2} / \|\tilde{x}_{n_k} - x_{n_k}\|_k \leq 1 / \|\tilde{x}_{n_k} - x_{n_k}\|$, we have

$$\frac{|f(\tilde{x}_{n_k}) - f(x_{n_k}) - \langle \tilde{x}_{n_k} - x_{n_k} | \nabla^k f(x_{n_k}) \rangle_k|}{\|\tilde{x}_{n_k} - x_{n_k}\|_k} \rightarrow 0. \quad (3.34)$$

Moreover,

$$\frac{\|\tilde{x}_k - x_k\|_k}{\gamma_k} \leq \frac{\|\tilde{x}_k - x_k\|_k}{\inf_{k \in \mathbb{N}} \gamma_k} \rightarrow 0. \quad (3.35)$$

Now it follows from (3.33), (3.34), (3.35), and (3.32) that

$$\frac{\|y_{n_k} - x_{n_k}\|}{\sup_{k \in \mathbb{N}} \gamma_k} \leq \frac{\|y_{n_k} - x_{n_k}\|}{\gamma_{n_k}} \rightarrow 0 \quad (3.36)$$

and, since $x_{n_k} \rightharpoonup \bar{x}$, we derive again from Lemma 3.3(i) that

$$\|\nabla f(y_{n_k}) - \nabla f(x_{n_k})\| \rightarrow 0. \quad (3.37)$$

Therefore, since $\sup_{k \in \mathbb{N}} \|W_k\| \leq \mu$, (3.24), (3.36), and (3.37), yields $v_{n_k} \rightarrow 0$. \square

We finally present the main convergence theorem.

Theorem 3.18. *Assume that **H1**–**H3** hold and that either of the two **H4** or **H5** hold. Let $(x_k)_{k \in \mathbb{N}}$ and $(y_k)_{k \in \mathbb{N}}$ be generated by algorithm (VM-FBS) using one of the line search procedures **LS1**–**LS4** for determining the parameters $(\gamma_k)_{k \in \mathbb{N}}$ and $(\lambda_k)_{k \in \mathbb{N}}$. Set $S_* = \operatorname{argmin}_{\mathcal{H}}(f + g)$. Then the following hold.*

- (i) $((f + g)(x_k))_{k \in \mathbb{N}}$ is decreasing.
- (ii) If $\inf_{k \in \mathbb{N}}(f + g)(x_k) > -\infty$, then $\sum_{k \in \mathbb{N}} \|x_{k+1} - x_k\|^2 < +\infty$.
- (iii) Suppose that $S_* \neq \emptyset$. Then
 - (a) $(x_k)_{k \in \mathbb{N}}$ and $(y_k)_{k \in \mathbb{N}}$ weakly converge to the same point in S_* .
 - (b) $(f + g)(y_k) \rightarrow \inf_{\mathcal{H}}(f + g)$.
 - (c) If $\sum_{k \in \mathbb{N}} \gamma_k \lambda_k = +\infty$, then $(f + g)(x_k) \rightarrow \inf_{\mathcal{H}}(f + g)$.
 - (d) If $\inf_{k \in \mathbb{N}} \gamma_k \lambda_k > 0$, then $((f + g)(x_k) - \inf_{\mathcal{H}}(f + g)) = o(1/k)$.
- (iv) If $S_* = \emptyset$, then $\|x_k\| \rightarrow +\infty$ and $(f + g)(x_k) \rightarrow \inf_{\mathcal{H}}(f + g)$.

Proof. Let $h = f + g$. For any proposed line search method, one of the properties (a), (b), or (c) in Proposition 3.13 is satisfied for every $k \in \mathbb{N}$ with respect to $\langle \cdot | \cdot \rangle_k$, and the corresponding property (a'), (b')–(b''), or (c'') in Proposition 3.17 is satisfied too. Therefore, by Corollary 3.15(i), $((f + g)(x_k))_{k \in \mathbb{N}}$ is decreasing and (a) in Proposition 3.8 is fulfilled. Moreover, Remark 3.16 ensures that if $\inf_{k \in \mathbb{N}}(f + g)(x_k) > -\infty$, then (b) in Proposition 3.8 is fulfilled for $(\alpha_k)_{k \in \mathbb{N}} = (\beta_k \gamma_k \lambda_k)_{k \in \mathbb{N}}$ (with $(\beta_k)_{k \in \mathbb{N}} \in \mathbb{R}_{++}^{\mathbb{N}}$ such that $0 < \inf_{k \in \mathbb{N}} \beta_k \leq \sup_{k \in \mathbb{N}} \beta_k < +\infty$). Finally, Corollary 3.15(ii) and Proposition 3.17(ii) implies that if $\inf_{k \in \mathbb{N}}(f + g)(x_k) > -\infty$, then (c) in Proposition 3.8 is fulfilled too. Then the statements follow from Theorem 3.10. \square

Remark 3.19. In the proof of Theorem 3.18, we showed that when **H5** is in force, we apply Proposition 3.8 using the metric of the Hilbert space \mathcal{H} , instead of the variable metrics $(\langle \cdot | \cdot \rangle_k)_{k \in \mathbb{N}}$ of **H3** (see Remark 3.16(ii)). This is why if we assume **H5** we do not require the convergence of the W_k 's.

Corollary 3.20. Assume that **H1**–**H3** hold and that either **H4** or **H5** hold. Let $(x_k)_{k \in \mathbb{N}}$ be generated by algorithm (VM-FBS) using either **LS1** or **LS4** with $(\lambda_k)_{k \in \mathbb{N}} \equiv 1$ (no relaxation). Set $S_* = \operatorname{argmin}_{\mathcal{H}}(f + g)$. Then the following hold.

- (i) $((f + g)(x_k))_{k \in \mathbb{N}}$ is decreasing and $(f + g)(x_k) \rightarrow \inf_{\mathcal{H}}(f + g)$.
- (ii) If $\inf_{\mathcal{H}}(f + g) > -\infty$, then $\sum_{k \in \mathbb{N}} \|x_{k+1} - x_k\|^2 < +\infty$.
- (iii) Suppose that $S_* \neq \emptyset$. Then
 - (a) $(x_k)_{k \in \mathbb{N}}$ weakly converges to a point in S_* .
 - (b) If $\inf_{k \in \mathbb{N}} \gamma_k > 0$, then $((f + g)(x_k) - \inf_{\mathcal{H}}(f + g)) = o(1/k)$.
- (iv) If $S_* = \emptyset$, then $\|x_k\| \rightarrow +\infty$.

Remark 3.21. In [6, Theorem 4.2 and Theorem 4.3] the same line search **LS4** is studied for the stationary (metric) forward-backward algorithm. However, even in this setting, the corresponding results given in Theorem 3.18 are more general and stronger. More precisely in [6, Method 1]: a) no relaxation is allowed, that is $\lambda_k \equiv 1$; b) δ is required to be strictly less than $1/2$ (this halves the stepsizes compared with those of **LS4** — see also Remark 3.25(i)); c) ∇f is required to be uniformly continuous on any bounded subsets of $\operatorname{dom} g$ and to map bounded sets into bounded sets; d) the little- o rate of convergence is provided only for \mathcal{H} finite-dimensional.

Proposition 3.17 and Theorem 3.10 allow also to obtain new convergence results for the standard variable metric forward-backward algorithm (without backtracking) [19] by requiring the Lipschitz continuity of the gradient on the domain of g only.

Theorem 3.22. Assume that **H1** and **H3** hold and that either **H4** or **H5** hold. Suppose f is Gâteaux differentiable on $\operatorname{dom} g$ and ∇f is L -Lipschitz continuous on $\operatorname{dom} g$ for some $L \in \mathbb{R}_+$. Define $(x_k)_{k \in \mathbb{N}}$ and $(y_k)_{k \in \mathbb{N}}$ as in (VM-FBS) and suppose $(\lambda_k)_{k \in \mathbb{N}}$ and $(\gamma_k)_{k \in \mathbb{N}}$ are chosen a priori in a such way that $\inf_{k \in \mathbb{N}} \lambda_k > 0$, $\inf_{k \in \mathbb{N}} \gamma_k > 0$, and $\sup_{k \in \mathbb{N}} \gamma_k \lambda_k / \nu_k < 2/L$. Then the conclusions of Theorem 3.18 hold.

Proof. Assumption **H2** is fulfilled too. Set $h = f + g$ and $(\alpha_k)_{k \in \mathbb{N}} = (\beta_k \gamma_k \lambda_k)_{k \in \mathbb{N}}$, where β_k is equal to $(1 + \eta_k)$ if **H4** holds, and $1/\nu_k$ if **H5** holds. Since $\sup_{k \in \mathbb{N}} \gamma_k \lambda_k / \nu_k < 2/L$, there exists $\delta \in]0, 1[$ such that $\sup_{k \in \mathbb{N}} \gamma_k \lambda_k / \nu_k \leq 2\delta/L$. Hence, since ∇f is L -Lipschitz continuous on $\operatorname{dom} g$ and $\{x_k \mid k \in \mathbb{N}\} \subset \operatorname{dom} g$, Fact 2.6 ensures that

$$(\forall k \in \mathbb{N}) \quad f(x_{k+1}) - f(x_k) - \langle x_{k+1} - x_k \mid \nabla f(x_k) \rangle \leq \frac{L}{2} \|x_{k+1} - x_k\|^2.$$

Since $\langle x_{k+1} - x_k \mid \nabla f(x_k) \rangle = \langle x_{k+1} - x_k \mid \nabla^k f(x_k) \rangle_k$ and $L/(2\nu_k) \leq \delta/(\gamma_k \lambda_k)$, by (2.2), we have

$$(\forall k \in \mathbb{N}) \quad \frac{L}{2} \|x_{k+1} - x_k\|^2 \leq \frac{L}{2\nu_k} \|x_{k+1} - x_k\|_k^2 \leq \frac{\delta}{\gamma_k \lambda_k} \|x_{k+1} - x_k\|_k^2.$$

Thus, (b) in Proposition 3.13 is satisfied for every $k \in \mathbb{N}$ with respect to $\langle \cdot | \cdot \rangle_k$. Moreover, since $\inf_{k \in \mathbb{N}} \gamma_k > 0$, condition (b') in Proposition 3.17 is also trivially satisfied. Then, by Corollary 3.15, Remark 3.16, and Proposition 3.17, we have that (a) in Proposition 3.8 is fulfilled and that, if $\inf_{k \in \mathbb{N}} h(x_k) > -\infty$, conditions (b) and (c) in Proposition 3.8 are fulfilled too. Then the statements follow from Theorem 3.10. \square

Remark 3.23.

- (i) Theorem 3.22 provides a worst case rate of convergence which is new.
- (ii) Theorem 3.22 shows that the gradient descent stepsize parameter γ_k , the relaxation parameter λ_k , and the minimum eigenvalues of the metric $\langle \cdot | \cdot \rangle_k$ are linked together by the condition $\sup_{k \in \mathbb{N}} \gamma_k \lambda_k / \nu_k < 2/L$. Thus, if λ_k is reduced (or ν_k is increased), one is allowed to enlarge the stepsize γ_k , which may therefore exceed $2/L$. This result complements that in [21] (for stationary metrics), where the parameters γ_k 's and λ_k 's appear linked too, but in that case, it is the relaxation parameter that can go beyond the usual bound 1.
- (iii) In Theorem 3.22 ∇f is not required to have full domain. Thus, the above result is not covered by the convergence theory developed in [19, 20, 21], since there a full domain of the gradient is required by the application of the Baillon-Haddad theorem. This aspect has been also noted in [16].

In view of Theorem 3.18(iii)(d) it is important to know conditions that guarantees that $(\gamma_k \lambda_k)_{k \in \mathbb{N}}$ remains bounded away from zero, since in such case (VM-FBS) has $o(1/k)$ rate of convergence in function values. We now addresses this issue.

Proposition 3.24. *Assume that **H3** and either **H4** or **H5** hold. Let $f, g \in \Gamma_0(\mathcal{H})$ with $\text{dom } g \subset \text{dom } f$ and suppose that f is Gâteaux differentiable on $\text{dom } g$.*

- (i) *Suppose that ∇f is globally Lipschitz continuous on $\text{dom } g$ with constant L and that $(\gamma_k)_{k \in \mathbb{N}}$, and $(\lambda_k)_{k \in \mathbb{N}}$ are generated through algorithm (VM-FBS) using any of the line seaches **LS1–LS4**. Then*
 - *for **LS1**, we have $\inf_{k \in \mathbb{N}} \gamma_k \geq \min\{\bar{\gamma}, 2\delta\theta\nu/(L \sup_{k \in \mathbb{N}} \lambda_k)\}$;*
 - *for **LS2** or **LS3**, we have $\inf_{k \in \mathbb{N}} \lambda_k \geq \min\{\bar{\lambda}, 2\delta\theta\nu/(L \sup_{k \in \mathbb{N}} \gamma_k)\}$.*
 - *for **LS4**, we have $\inf_{k \in \mathbb{N}} \gamma_k \geq \min\{\bar{\gamma}, \delta\theta\nu/(L \sup_{k \in \mathbb{N}} \lambda_k)\}$.*
- (ii) *Suppose that $S_* = \text{argmin}_{\mathcal{H}}(f + g) \neq \emptyset$ and that ∇f is Lipschitz continuous on any weakly compact subset of $\text{dom } g$. Let $(\gamma_k)_{k \in \mathbb{N}}$, and $(\lambda_k)_{k \in \mathbb{N}}$ be defined according to algorithm (VM-FBS) using any of the line search procedures **LS1–LS4**. Then $\inf_{k \in \mathbb{N}} \gamma_k > 0$ and $\inf_{k \in \mathbb{N}} \lambda_k > 0$.*

Proof. We first remark that in both statements, **H1** and **H2** are fulfilled.

(i): First we consider the case that $(x_k)_{k \in \mathbb{N}}$, $(\gamma_k)_{k \in \mathbb{N}}$, and $(\lambda_k)_{k \in \mathbb{N}}$ are generated using either **LS1**, **LS2** or **LS3**. Let $k \in \mathbb{N}$. Since ∇f is L -Lipschitz continuous on $\text{dom } g$, we derive from the descent lemma (Fact 2.6) that for every $\gamma > 0$

$$f(J(x_k, \gamma, \lambda_k)) - f(x_k) - \langle J(x_k, \gamma, \lambda_k) - x_k | \nabla f(x_k) \rangle \leq \frac{L}{2} \|J_k(x_k, \gamma, \lambda_k) - x_k\|^2 \quad (3.38)$$

and for every $\lambda \in]0, 1]$,

$$f(J_k(x_k, \gamma_k, \lambda)) - f(x_k) - \langle J(x_k, \gamma_k, \lambda) - x_k | \nabla f(x_k) \rangle \leq \frac{L}{2} \|J_k(x_k, \gamma_k, \lambda) - x_k\|^2. \quad (3.39)$$

Moreover, again by (2.2),

$$(\forall (\gamma, \lambda) \in \mathbb{R}_{++} \times]0, 1]) \quad \frac{L}{2} \|J_k(x_k, \gamma, \lambda) - x_k\|^2 \leq \frac{L}{2\nu_k} \|J_k(x_k, \gamma, \lambda) - x_k\|_k^2. \quad (3.40)$$

Define

$$\bar{\gamma}_k = \max \{ \gamma \in \mathbb{R}_{++} \mid (\exists i \in \mathbb{N})(\gamma = \bar{\gamma}\theta^i) \ L \leq 2\nu_k\delta/(\gamma\lambda_k) \}, \quad (3.41)$$

$$\bar{\lambda}_k = \max \{ \lambda \in]0, 1] \mid (\exists i \in \mathbb{N})(\lambda = \bar{\lambda}\theta^i) \ L \leq 2\delta\nu_k/(\gamma_k\lambda) \}. \quad (3.42)$$

It follows from (3.38), (3.40), and the definition of γ_k in LS1 that $\gamma_k \geq \bar{\gamma}_k$. Moreover, by (3.41), we have that if $\bar{\gamma}_k < \bar{\gamma}$, then $L > 2\delta\theta\nu_k/(\bar{\gamma}_k\lambda_k)$, hence $\bar{\gamma}_k > 2\delta\theta\nu_k/(L\lambda_k)$. Therefore $\bar{\gamma}_k \geq \min\{\bar{\gamma}, 2\delta\theta\nu/(L\sup_{k \in \mathbb{N}} \lambda_k)\}$. Similarly, it follows from (3.39), (3.40), and the definition of λ_k in LS2 or in LS3, and Proposition 3.13(i) (invoked for the metric $\langle \cdot | \cdot \rangle_k$), that $\lambda_k \geq \bar{\lambda}_k$. Moreover, by (3.42), we have that if $\bar{\lambda}_k < \bar{\lambda}$, then $L > 2\delta\theta\nu_k/(\gamma_k\bar{\lambda}_k)$, hence $\bar{\lambda}_k > 2\delta\theta\nu_k/(L\gamma_k)$. Therefore, $\bar{\gamma}_k \geq \min\{\bar{\lambda}, 2\delta\theta\nu/(L\sup_{k \in \mathbb{N}} \gamma_k)\}$.

We now consider the case that $(x_k)_{k \in \mathbb{N}}$, $(\gamma_k)_{k \in \mathbb{N}}$, and $(\lambda_k)_{k \in \mathbb{N}}$ are defined according to LS4. Let $k \in \mathbb{N}$. Then, for every $\gamma > 0$, $\|\nabla f(J_k(x_k, \gamma, \lambda_k)) - \nabla f(x_k)\| \leq L \|J_k(x_k, \gamma, \lambda_k) - x_k\|$, and, by (2.2) and Fact 2.7,

$$\|\nabla^k f(J_k(x_k, \gamma)) - \nabla^k f(x_k)\|_k \leq \frac{L}{\nu_k} \|J_k(x_k, \gamma) - x_k\|_k. \quad (3.43)$$

Now, define

$$\bar{\gamma}_k = \max \{ \gamma \in \mathbb{R}_{++} \mid (\exists i \in \mathbb{N})(\gamma = \bar{\gamma}\theta^i) \ L \leq \nu_k\delta/(\gamma\lambda_k) \}. \quad (3.44)$$

It follows from (3.43) and the definition of γ_k in LS4 that $\gamma_k \geq \bar{\gamma}_k$. Moreover, by (3.44), if $\bar{\gamma}_k < \bar{\gamma}$, then $L > \delta\theta\nu_k/(\bar{\gamma}_k\lambda_k)$, hence $\bar{\gamma}_k > \delta\theta\nu_k/(L\lambda_k)$. Therefore $\bar{\gamma}_k \geq \min\{\bar{\gamma}, \delta\theta\nu/(L\sup_{k \in \mathbb{N}} \lambda_k)\}$.

(ii): Assume first that $(x_k)_{k \in \mathbb{N}}$, $(\gamma_k)_{k \in \mathbb{N}}$, and $(\lambda_k)_{k \in \mathbb{N}}$ are defined according to either LS1 or LS4. Since $S_* \neq \emptyset$, $\inf_{\mathcal{H}}(f+g) > -\infty$. Besides, we derive from Theorem 3.18(iii) that $x_k \rightharpoonup \bar{x} \in S^*$. Let, for every $k \in \mathbb{N}$, $\tilde{\gamma}_k = \gamma_k/\theta$ and $\tilde{x}_k = J(x_k, \tilde{\gamma}_k, \lambda_k)$. Then, by Lemma 3.5(i) (applied to each metric $\langle \cdot | \cdot \rangle_k$), for every $k \in \mathbb{N}$

$$\|J_k(x_k, \tilde{\gamma}_k, \lambda_k) - x_k\|_k \leq \frac{\tilde{\gamma}_k}{\gamma_k} \|J_k(x_k, \gamma_k, \lambda_k) - x_k\|_k = \frac{1}{\theta} \|x_{k+1} - x_k\|_k. \quad (3.45)$$

Moreover, according to the definition of γ_k in LS1 and LS4, we have respectively

$$(\forall k \in \mathbb{N}) \ \gamma_k < \bar{\gamma} \Rightarrow f(\tilde{x}_k) - f(x_k) - \langle \tilde{x}_k - x_k | \nabla^k f(x_k) \rangle_k > \delta \frac{\|\tilde{x}_k - x_k\|_k^2}{\tilde{\gamma}_k \lambda_k} \quad (3.46)$$

or

$$(\forall k \in \mathbb{N}) \ \gamma_k < \bar{\gamma} \Rightarrow \|\nabla^k f(\tilde{x}_k) - \nabla^k f(x_k)\|_k > \delta \frac{\|\tilde{x}_k - x_k\|_k}{\tilde{\gamma}_k \lambda_k}. \quad (3.47)$$

Now, by Theorem 3.18(ii) and (2.2), $\|x_{k+1} - x_k\|_k \rightarrow 0$, hence, by (3.45) and (2.2), $\tilde{x}_k - x_k \rightarrow 0$. Then, since $x_k \rightharpoonup \bar{x}$, Lemma 3.3(ii) yields that $\exists L > 0$ such that, $\forall k \in \mathbb{N}$, $\|\nabla f(\tilde{x}_k) - \nabla f(x_k)\| \leq L \|\tilde{x}_k - x_k\|$ and $f(\tilde{x}_k) - f(x_k) - \langle \tilde{x}_k - x_k | \nabla f(x_k) \rangle \leq L \|\tilde{x}_k - x_k\|^2/2$. The above inequalities, in view of (2.2), (2.4), and Fact 2.7, imply

$$(\forall k \in \mathbb{N}) \quad \|\nabla f(\tilde{x}_k) - \nabla f(x_k)\| \leq \frac{L}{\nu_k} \|\tilde{x}_k - x_k\| \quad (3.48)$$

$$(\forall k \in \mathbb{N}) \quad f(\tilde{x}_k) - f(x_k) - \langle \tilde{x}_k - x_k | \nabla^k f(x_k) \rangle_k \leq \frac{L}{2\nu_k} \|\tilde{x}_k - x_k\|_k^2. \quad (3.49)$$

Thus, (3.49) and (3.46) yield that, $\forall k \in \mathbb{N}$, $\gamma_k < \bar{\gamma} \Rightarrow L/(2\nu_k) \geq \delta/(\tilde{\gamma}_k\lambda_k) \Rightarrow \gamma_k = \tilde{\gamma}_k\theta \geq 2\delta\theta\nu_k/(L\lambda_k)$. Thus, $\forall k \in \mathbb{N}$, $\gamma_k \geq \min\{\bar{\gamma}, 2\delta\theta\nu/(L\sup_{k \in \mathbb{N}} \lambda_k)\} > 0$. Moreover, it follows from (3.48) and (3.47) that, $\forall k \in \mathbb{N}$, $\forall \gamma_k < \sigma \Rightarrow L/\nu_k \geq \delta/(\tilde{\gamma}_k\lambda_k) \Rightarrow \gamma_k = \tilde{\gamma}_k\theta\nu_k \geq \delta\theta/(L\lambda_k)$. Thus, $\forall k \in \mathbb{N}$, $\gamma_k \geq \min\{\bar{\gamma}, \delta\theta\nu/(L\sup_{k \in \mathbb{N}} \lambda_k)\} > 0$.

The case LS2 or LS3 is treated in the same way. \square

Remark 3.25.

- (i) The results given in Proposition 3.24, shows that when ∇f has some kind of Lipschitz continuity property, the stepsizes determined by LS4 may be half of those determined by the other line search methods. In particular, if ∇f is L -Lipschitz continuous on $\text{dom } g$ and $(\gamma_k)_{k \in \mathbb{N}}$ are defined according to LS4, we can make $\inf_{k \in \mathbb{N}} \gamma_k$ (by choosing $\lambda_k \equiv 1$, and δ and θ sufficiently close to 1) arbitrarily close to ν/L ; whereas using LS1, LS2, or LS3, $\inf_{k \in \mathbb{N}} \gamma_k$ can approach $2\nu/L$. This latter result is in line with the state of the art convergence theory on forward-backward splitting algorithm (with $\nu = 1$) [13, 20, 22]. This suggests that LS4 is not quite appropriate for (VM-FBS).
- (ii) Similar results recently appeared under stronger hypotheses and for more specific cases in [6] and [10]. In particular in [6, Proposition 4.4] (for stationary metrics) it is proved that if f is globally Lipschitz continuous and LS4 is used, $\inf_{k \in \mathbb{N}} \gamma_k$ can (only) reach $1/(2L)$. Moreover, it is also showed that $\inf_{k \in \mathbb{N}} \gamma_k > 0$, if \mathcal{H} is finite-dimensional and ∇f is locally Lipschitz continuous around any point of S_* . However, if \mathcal{H} is finite dimensional and ∇f is locally Lipschitz continuous around any point of $\text{dom } g$, then ∇f is Lipschitz continuous on any (weakly) compact subset of $\text{dom } g$ (recall also Remark 3.1(i)). Therefore Proposition 3.24(ii) encompasses [6, Proposition 4.4(ii)]. On the other hand in [10] a variable metric projected gradient method is studied for finite dimensional convex problems using the corresponding specialization of LS3. In Proposition 2.2 they prove that the λ_k 's are bounded away from zero provided that ∇f is locally Lipschitz continuous and additionally that f is coercive.

4 Dealing with general placements of the domains

This section provides a slight variation of algorithm (VM-FBS) that can handle more general configurations of the domains of f and g . This will be done at the expense of additional assumptions. We recall that in **H2** it is required that $\text{dom } g \subset \text{dom } f$; however that condition is not always appropriate when f is taken of divergence type [11, 38]. Here we replace assumptions **H1** and **H2** by the following **H1'**–**H3'**. For every $h: \mathcal{H} \rightarrow]-\infty, +\infty]$ and $\alpha \in \mathbb{R}$, we set $\{h \leq \alpha\} = \{x \in \mathcal{H} \mid h(x) \leq \alpha\}$. Moreover, we define the distance between subsets A and B of \mathcal{H} as $d(A, B) = \inf \{ \|x - y\| \mid x \in A \text{ and } y \in B \}$.

- H1'** $f: \mathcal{H} \rightarrow]-\infty, +\infty]$ and $g: \mathcal{H} \rightarrow]-\infty, +\infty]$ are proper convex and lower semicontinuous functions, bounded from below and such that $\text{dom } g \cap \text{int dom } f \neq \emptyset$;
- H2'** f is Gâteaux differentiable on $\text{dom } g \cap \text{int dom } f$, ∇f is uniformly continuous on any weakly compact subset of $\text{dom } g \cap \text{int dom } f$, and ∇f is bounded on any sublevel sets of $f + g$.
- H3'** for every $x \in \text{int dom } f \cap \text{dom } g$, $\{f + g \leq (f + g)(x)\} \subset \text{int dom } f \cap \text{dom } g$ and $d(\{f + g \leq (f + g)(x)\}, \mathcal{H} \setminus \text{int dom } f) > 0$.

The following result clarifies the role of the above hypotheses.

Proposition 4.1. *Suppose that **H1'**–**H3'** hold. Let $x_0 \in \text{int dom } f \cap \text{dom } g$ and $K_0 = \{f + g \leq (f + g)(x_0)\}$. Then there exist constants C_1 and C_2 in \mathbb{R}_+ , such that,*

$$(\forall x \in K_0)(\forall \gamma \in \mathbb{R}_{++}) \quad \|J(x, \gamma, 1) - x\| \leq C_1 \gamma + C_2 \sqrt{\gamma}. \quad (4.1)$$

Moreover, setting $\delta_0 = d(K_0, \mathcal{H} \setminus \text{int dom } f) > 0$, we have

$$(\forall x \in K_0)(\forall \gamma > 0) \quad J(x, \gamma, 1) \notin \text{dom } f \Rightarrow \gamma \geq c_0 := c(\delta_0) > 0,$$

where $c(\cdot)$ is the inverse of the strictly increasing function $\gamma \mapsto C_1 \gamma + C_2 \sqrt{\gamma}$.

Proof. We first prove that g is bounded on K_0 . Indeed, let $\vartheta \in \mathbb{R}$ be such $\vartheta \leq f$. Then, $\forall x \in K_0$, $\vartheta + g(x) \leq (f + g)(x) \leq (f + g)(x_0)$, hence $g(x) \leq (f + g)(x_0) - \vartheta$. Concerning the first part of the statement, we note that, since g is bounded from below, we have $g^*(0) < +\infty$, hence in (3.2) of Lemma 3.5(iv), we can take $u = 0$, obtaining $\|J(x, \gamma, 1) - x\| \leq \gamma \|\nabla f(x)\| + (2\gamma)^{1/2} \sqrt{g(x) + g^*(0)}$. Since, by **H2'**, ∇f is bounded on K_0 and g is bounded on K_0 too, (4.1) follows. Moreover, if $x \in K_0$ and $J(x, \gamma, 1) \notin \text{dom } f$, then $\delta_0 \leq \|J(x, \gamma, 1) - x\| \leq C_1\gamma + C_2\sqrt{\gamma}$ and hence $c(\delta_0) \leq \gamma$. \square

Proposition 4.2. *Let \mathcal{G} be a real Hilbert space, let $A: \mathcal{H} \rightarrow \mathcal{G}$ be a non-zero bounded linear operator, let $h \in \Gamma_0(\mathcal{G})$ and set $f = h \circ A$. Let $g \in \Gamma_0(\mathcal{H})$ and suppose that $\text{dom } g \cap \text{int dom } f \neq \emptyset$ and g is bounded from below. Then, the following hold.*

- (i) *f is bounded from above on the sublevel sets of $f + g$. Moreover, if f is bounded from below, then g is bounded from above on the sublevel sets of $f + g$.*
- (ii) *Suppose that $\text{dom } f \neq \mathcal{H}$, and that,*
 - (a) *for every $\alpha \in \mathbb{R}$, $\{h \leq \alpha\} \subset \text{int dom } h$ and $d(\{h \leq \alpha\}, \mathcal{G} \setminus \text{int dom } h) > 0$.**Then **H3'** holds.*
- (iii) *Suppose that \mathcal{G} is finite dimensional, h is coercive, and $\forall \alpha \in \mathbb{R}$, $\{h \leq \alpha\} \subset \text{int dom } h$. Then f is bounded from below and (a) is satisfied; hence **H1'** and **H3'** hold.*
- (iv) *In addition to the assumptions in (iii), suppose that $\text{dom } h = \text{int dom } h$ and that ∇h is continuously differentiable on $\text{dom } h$. Then **H2'** holds.*

Proof. (i): Let $\alpha \in \mathbb{R}$ and let $\beta \in \mathbb{R}$ be such that $\beta \leq g$. Then, for every $x \in \{f + g \leq \alpha\}$, $f(x) + \beta \leq (f + g)(x) \leq \alpha$, hence $f(x) \leq \alpha - \beta$. As regards the second part, let $\vartheta \in \mathbb{R}$ be such $\vartheta \leq f$. Then, for every $x \in \{f + g \leq \alpha\}$, $\vartheta + g(x) \leq (f + g)(x) \leq \alpha$, hence $g(x) \leq \alpha - \vartheta$.

(ii): Let $x \in \text{int dom } f \cap \text{dom } g$ and set $K = \{f + g \leq (f + g)(x)\}$ and $U = \text{int dom } h$. It follows from (i) that $\exists \alpha \in \mathbb{R}$ such that, $f \leq \alpha$ on K . Then $K \subset \{f \leq \alpha\} \subset A^{-1}(\{h \leq \alpha\}) \subset A^{-1}(U)$, which is open (and contained in $\text{dom } f$), hence $\{f \leq \alpha\} \subset A^{-1}(U) \subset \text{int dom } f$. Let $\delta = d(\{h \leq \alpha\}, \mathcal{G} \setminus U) > 0$. Now, let $x' \in \{f \leq \alpha\}$ and $x'' \in \mathcal{H} \setminus \text{int dom } f$. Then $Ax' \in \{h \leq \alpha\}$ and $Ax'' \notin U$, hence $\delta \leq \|Ax' - Ax''\| \leq \|A\| \|x' - x''\|$. Thus, $d(K, \mathcal{H} \setminus \text{int dom } f) \geq d(\{f \leq \alpha\}, \mathcal{H} \setminus \text{int dom } f) \geq \delta / \|A\| > 0$.

(iii): Since the range of A , $R(A)$, is closed in \mathcal{G} , $h + \iota_{R(A)}$ is lower semicontinuous and coercive, hence it has a minimizer, say $\bar{y} \in R(A)$. Then, taking $\bar{x} \in \mathcal{H}$ such that $A\bar{x} = \bar{y}$, we have, for every $x \in \mathcal{H}$, $h(A\bar{x}) \leq h(Ax) = f(x)$. Thus, f is bounded from below. Let $\alpha \in \mathbb{R}$. Since $\{h \leq \alpha\}$ is compact and $d(\cdot, \mathcal{G} \setminus \text{int dom } h)$ is continuous and strictly positive on $\{h \leq \alpha\}$, then $d(\{h \leq \alpha\}, \mathcal{G} \setminus \text{int dom } h) > 0$.

(iv): Clearly $\text{dom } f = A^{-1}(\text{dom } h) = \text{int dom } f$ and f is continuously differentiable on $\text{dom } f$ and, for every $x \in \text{dom } f$, $\nabla f(x) = A^* \nabla h(Ax)$. Let $\alpha \in \mathbb{R}$. It follows from (i) that there exists $\beta \in \mathbb{R}$ such that $f \leq \beta$ on $\{f + g \leq \alpha\}$. Then, for every $x \in \{f + g \leq \alpha\}$, $h(Ax) \leq \beta$, therefore $Ax \in \{h \leq \beta\}$. Since ∇h is continuous on the compact $\{h \leq \beta\}$, there exists $\eta \in \mathbb{R}_+$ such that $\|\nabla h\| \leq \eta$ on $\{h \leq \beta\}$. Thus $\|\nabla f(x)\| \leq \|A^*\| \|\nabla h(Ax)\| \leq \|A^*\| \eta$. This proves that ∇f is bounded on the sublevel sets of $f + g$. Let K be a weakly compact subset of $\text{int dom } f = A^{-1}(\text{int dom } h)$. Since A is weak-to-weak continuous, $A(K)$ is a (weakly) compact subset of $\text{int dom } h$. Thus, by the Heine-Cantor theorem, ∇h is uniformly continuous on $A(K)$. Hence $\nabla f = A^* \circ \nabla h \circ A|_{\text{dom } f}$ is uniformly continuous on K . \square

Example 4.3. Let $n \in \mathbb{N}$, $n \geq 1$. Let φ be a Legendre function on \mathbb{R}^n [4] which is twice continuously differentiable on $\text{int dom } \varphi$, and let

$$D: \mathbb{R}^n \times \mathbb{R}^n \rightarrow [0, +\infty]: (x, y) \mapsto \begin{cases} \varphi(x) - \varphi(y) - \langle \nabla \varphi(y) | x - y \rangle & \text{if } y \in \text{int dom } \varphi \\ +\infty & \text{otherwise} \end{cases} \quad (4.2)$$

be the associated Bregman distance. Suppose that, for every $x \in \text{int dom } \varphi$, $D(x, \cdot) \in \Gamma_0(\mathbb{R}^n)$ and $D(x, \cdot)$ is coercive — this case is studied in [4] (see in particular Lemma 2.6) and occurs, e.g., for the Kullback-Leibler divergence, where $\varphi(x) = \sum_{i=1}^n x_i \log x_i - x_i$. Let $b \in \text{int dom } \varphi$, let $A: \mathcal{H} \rightarrow \mathbb{R}^n$ be a bounded linear operator, and set $f = D(b, A \cdot)$. Thus, in virtue of Proposition 4.2(iii)-(iv), if $g \in \Gamma_0(\mathcal{H})$ is bounded from below and such that $A^{-1}(\text{int dom } \varphi) \cap \text{dom } g \neq \emptyset$, assumptions **H1'**–**H3'** are satisfied.

Under assumptions **H1'**–**H3'**, we can modify algorithm (VB-FBS), by adding a further line search for computing y_k . More precisely, for **LS2** and **LS3**, the sequence $(\gamma_k)_{k \in \mathbb{N}}$ cannot be chosen a priori anymore, but it has to be computed by the following procedure. Let $\bar{\gamma} > 0$, let $x_0 \in \text{int dom } f \cap \text{dom } g$, and set $K_0 = \{f + g \leq (f + g)(x_0)\}$. Then, for every $k \in \mathbb{N}$, assume that $x_k \in K_0$, and compute

$$\gamma_k = \max \left\{ \gamma \in \mathbb{R}_{++} \mid (\exists i \in \mathbb{N})(\gamma = \bar{\gamma} \theta^i) \quad J(x_k, \gamma, 1) \in \text{dom } f \right\}. \quad (4.3)$$

Note that, since $x_k \in \text{int dom } f$ and $J(x_k, \gamma, 1) \rightarrow x_k$ as $\gamma \rightarrow 0$, the procedure (4.3) is well-defined. Moreover, because of Proposition 4.1, if $\gamma_k \leq \bar{\gamma} \theta$, then $J(x_k, \gamma_k/\theta, 1) \notin \text{dom } f$ and hence $\gamma_k \geq c_0 \theta$. Therefore $\inf_k \gamma_k > 0$. Procedure (4.3) ensures that $y_k \in \text{dom } f \cap \text{dom } g$ and the subsequent line search makes sense. Moreover, Proposition 3.13 yields that $(f + g)(x_{k+1}) \leq (f + g)(x_k)$, hence $x_{k+1} \in K_0$ and the algorithm can continue. Concerning **LS1** and **LS4**, the γ_k computed by (4.3) is meant to replace $\bar{\gamma}$ in **LS1** and **LS4**, meaning that, for every $k \in \mathbb{N}$, they will do backtracking on γ starting from the output of (4.3). This will make sense of the subsequent procedures **LS1** and **LS4**. Again Proposition 3.13 proves that the next step is descendent and hence $x_{k+1} \in K_0$. Note that in **LS1**, if $(\lambda_k)_{k \in \mathbb{N}} \equiv 1$, then one can perform **LS1** only, since it will automatically search for a point in $\text{dom } f$.

5 Applications

In this section we illustrate several models that can be tackled by the proposed algorithm. In particular we show that its scope of applicability encompasses problems that involve Banach spaces or functions of divergence type. To that purpose we recall few facts.

Fact 5.1. *Let $A: \mathcal{H} \rightarrow \mathcal{B}$ be a bounded linear operator between a real Hilbert space and a real Banach space. Let $\varphi: \mathcal{B} \rightarrow \mathbb{R}$ be a differentiable function and suppose that its derivative φ' is α -Hölder continuous on bounded sets, for some $\alpha \in]0, 1]$, that is, for every bounded set $Y \subset \mathcal{B}$, there exists $C \in \mathbb{R}_+$ such that*

$$(\forall (y_1, y_2) \in Y^2) \quad \|\varphi'(y_1) - \varphi'(y_2)\|_{\mathcal{B}^*} \leq C \|y_1 - y_2\|_{\mathcal{B}}^\alpha.$$

Then, $f = \varphi \circ A: \mathcal{H} \rightarrow \mathbb{R}$ is differentiable and $\nabla f = A^ \circ \varphi' \circ A$, where $A^*: \mathcal{B}^* \rightarrow \mathcal{H}$ is the adjoint of A . Moreover, ∇f is α -Hölder continuous on the bounded sets of \mathcal{H} .*

The following result is in [39, Corollary 2.44 and Theorem 2.53(f)]

Fact 5.2. *Let \mathcal{B} be a uniformly smooth Banach space, let $p \in]1, +\infty[$ and set $\varphi = (1/p) \|\cdot\|_{\mathcal{B}}^p$. Then φ is differentiable and $\varphi' = J_{\mathcal{B}, p}$ is the p -duality mapping of \mathcal{B} , which is uniformly continuous on bounded sets. Moreover, if \mathcal{B} has modulus of smoothness of power type $q \in]1, 2]$, then, φ' is $(p-1)$ -Hölder continuous, if $p \leq q$, and $(q-1)$ -Hölder continuous on bounded sets, if $p > q$.*

In the following we give a prominent example in which the duality map of the involved Banach space is explicitly computable.

Remark 5.3. Let $(\Omega, \mathfrak{A}, \mu)$ be a σ -finite measure space and let $p \in]1, +\infty[$. Then $L^p(\Omega, \mu)$ is uniformly smooth with modulus of smoothness of power type $\min\{2, p\}$ [33]. Therefore, it follows from Fact 5.2 that the function $\varphi = (1/p) \|\cdot\|_p^p$ is differentiable and φ' is $(p-1)$ -Hölder continuous on $L^p(\Omega, \mu)$, if $p \leq 2$, and Lipschitz continuous on the bounded sets of $L^p(\Omega, \mu)$, if $p > 2$. Moreover, for every $x \in L^p(\Omega, \mu)$, $\varphi'(x) \in L^{p^*}(\Omega, \mu)$ and $\varphi'(x): \Omega \rightarrow \mathbb{R}: \omega \mapsto |x(\omega)|^{p-1} \text{sign}(x(\omega))$.

It follows from Fact 5.2 and Fact 5.1 that the following general optimization problem is of the form (P) and hypotheses **H1** and **H2** are satisfied.

Problem 5.4. Let $A: \mathcal{H} \rightarrow \mathcal{B}$ be a bounded linear operator between a real Hilbert space and a real uniformly smooth Banach space. Let $g \in \Gamma_0(\mathcal{H})$ and $b \in \mathcal{B}$. Then

$$\underset{x \in \mathcal{H}}{\text{minimize}} \quad \frac{1}{p} \|Ax - b\|_{\mathcal{B}}^p + g(x) \quad (p > 1).$$

Remark 5.5. In Problem 5.4, we have $\nabla((1/p) \|A \cdot - b\|_{\mathcal{B}}^p)(x) = A^* J_{\mathcal{B}, p}(Ax - b)$. In this case the gradient descent step in (VM-FBS) resembles the Landweber step in Banach spaces [13, 39].

Based on Remark 5.3, we give some significant instances of Problem 5.4.

Example 5.6. Let \mathcal{H} be a real Hilbert space and let $(e_k)_{k \in \mathbb{K}} \in \mathcal{H}^{\mathbb{K}}$ be an orthonormal basis of \mathcal{H} . Let $(\Omega, \mathfrak{A}, \mu)$ be a σ -finite measure space and let $p \in]1, +\infty[$. Let $A: \mathcal{H} \rightarrow L^p(\Omega, \mu)$ be a bounded linear operator, let $b \in L^p(\Omega, \mu)$, and let $(g_k)_{k \in \mathbb{N}}$ be a sequence of functions in $\Gamma_0(\mathbb{R})$ such that $g_k \geq g_k(0) = 0$, for every $k \in \mathbb{N}$. Then

$$\underset{x \in \mathcal{H}}{\text{minimize}} \quad \frac{1}{p} \|Ax - b\|_p^p + \sum_{k \in \mathbb{N}} g_k(\langle e_k | x \rangle). \quad (5.1)$$

Denoting by f and g respectively the first and second term in (5.1), we have $\nabla f(x) = A^*u$, where $u: \Omega \rightarrow \mathbb{R}: \omega \mapsto |(Ax)(\omega) - b(\omega)|^{p-1} \text{sign}((Ax)(\omega) - b(\omega))$, and the proximity operator can be computed component-wise [20], that is

$$\text{prox}_{\gamma g}(x) = (\text{prox}_{\gamma g_k}(\langle e_k | x \rangle))_{k \in \mathbb{N}}.$$

Moreover, it follows from Remark 5.3 that ∇f is $(p-1)$ -Hölder continuous if $p \leq 2$ and Lipschitz continuous on bounded sets if $p > 2$. Therefore, it follows from Theorem 3.18 and Proposition 3.24(ii) that, in this case, the sequence generated by (VM-FBS) is weakly convergent to a solution of (5.1) and converges in function values to the corresponding minimum. Moreover, if $p \geq 2$, the convergence in function values boasts a rate of $o(1/k)$. This example covers the class of problems approached by the iterative shrinkage/thresholding algorithm [23, 5], but here a more general discrepancy term is used. A special case of (5.1) is

$$\min_{x \in \ell^2(\mathbb{K})} \frac{1}{p} \|Ax - b\|_p^p + \|x\|_1, \quad (5.2)$$

where, \mathbb{K} is a countable set, $A: \ell^2(\mathbb{K}) \rightarrow \mathbb{R}^n$ is a bounded linear operator, and $b \in \mathbb{R}^n$. Note that here $\text{dom} \|\cdot\|_1 = \ell^1(\mathbb{K})$, which is not closed in $\ell^2(\mathbb{K})$. We highlight that problems of type (5.2) arise in function interpolation (from discrete data) and non parametric function estimation (support vector regression).

We end the section by showing a prototype of problems where $\text{dom } g \not\subset \text{dom } f$ and hypotheses **H1'**–**H3'** in Section 4 are met (recall Example 4.3).

Problem 5.7. Let \mathcal{H} be a real Hilbert space, let $n \in \mathbb{N}$ with $n \geq 1$, and let φ be a Legendre function on \mathbb{R}^n such that φ is twice continuously differentiable on $\text{int dom } \varphi$ and its associated Bregman distance D (see (4.2)) satisfies the condition: $\forall z \in \text{int dom } \varphi, D(z, \cdot) \in \Gamma_0(\mathbb{R}^n)$ and $D(z, \cdot)$ is coercive. Let $A: \mathcal{H} \rightarrow \mathbb{R}^n$ be a bounded linear operator, let $b \in \text{int dom } \varphi$, and let $g \in \Gamma_0(\mathcal{H})$ be such that $A^{-1}(\text{int dom } \varphi) \cap \text{dom } g \neq \emptyset$ and g is bounded from below. Then

$$\underset{x \in \mathcal{H}}{\text{minimize}} \quad D(b, Ax) + g(x).$$

Examples of Problem 5.7 are provided in the following.

Example 5.8. Let $n \in \mathbb{N}$ with $n \geq 1$ and let $D(z, y) = \sum_{i=1}^n z_i \log(z_i/y_i) + y_i - z_i$ be the Kullback-Leibler divergence. Then, let $A \in \mathbb{R}_+^{n \times n}$, $b \in \mathbb{R}_{++}^n$ and solve

$$\min_{x \in \mathbb{R}_+^n} D(b, Ax) + TV(x). \quad (5.3)$$

where TV is the (discrete) total variation. Note that in this case $g = \iota_{\mathbb{R}_+^n} + TV$. Moreover $\text{dom } g \not\subset \text{dom } D(b, \cdot) \circ A$ and $\nabla(D(b, \cdot) \circ A)$ is only locally Lipschitz continuous on its domain. Then, it follows from Theorem 3.18 and Proposition 3.24(ii) that (VM-FBS) (with the additional line search presented in Section 4) provides a sequence which converges to a solution of (5.3) and converges in functional values to the related minimum at rate $o(1/k)$. Problem (5.3) is of the type considered in [11, Section 4.2], but here the introduction of the background signal is avoided — provided that the sought signal x^* satisfies $Ax^* > 0$. Another instance of Problem 5.7 is

$$\min_{\substack{x \in \mathbb{R}_+^n \\ \beta \in \mathbb{R}_+}} D(b, Ax + \beta \mathbf{1}) + \|x\|_1,$$

where the signal and the background are sought and $\mathbf{1}$ is the vector of \mathbb{R}^n of all ones. Here again the domain of the map $(x, \beta) \mapsto D(b, Ax + \beta \mathbf{1})$ is not contained in $\mathbb{R}_+^n \times \mathbb{R}_+$.

References

- [1] H. Attouch, J. Bolte, and B.F. Svaiter, Convergence of descent methods for semi-algebraic and tame problems: proximal algorithms, forward-backward splitting, and regularized Gauss-Seidel method. *Math. Program.*, 137: 1-2, 91–129. 2013.
- [2] H. H. Bauschke and P. L. Combettes, *Convex Analysis and Monotone Operator Theory in Hilbert Spaces*. Springer, New York 2011.
- [3] H. H. Bauschke and P. L. Combettes, The Baillon-Haddad theorem revisited. *J. Convex Anal.*, 17: 3&4, 781–787. 2010.
- [4] H. H. Bauschke, P. L. Combettes and D. Noll, Joint minimization with alternating Bregman proximity operators. *Pac. J. Optim.*, 2, 401–424, 2006.
- [5] A. Beck and M. Teboulle, A fast iterative shrinkage-thresholding algorithm for linear inverse problems. *SIAM J. Imaging Sciences*, 2: 1, 183–202. 2009.
- [6] J. Y. Bello Cruz and T. T. A. Nghia, On the convergence of the proximal forward-backward splitting method with linesearch. arXiv:1501.02501v3, 2015.
- [7] D. P. Bertsekas, On the Goldstein-Levitin-Polyak gradient projection method. *IEEE Transactions on Automatic Control*, AC-21: 2, 174–184. 1976.

- [8] D. P. Bertsekas, *Nonlinear Programming*. Athena Scientific, Belmont 1995.
- [9] S. Bonettini, L. Loris, F. Porta, and M. Prato, Variable metric inexact line-search based methods for nonsmooth optimization, *SIAM J. Optim.*, 26: 2, 891–921. 2016.
- [10] S. Bonettini and M. Prato, New convergence results for the scaled gradient projection method, *Inverse Problems*, 31: 9. 2015.
- [11] S. Bonettini and V. Ruggiero, On the convergence of primal-dual hybrid gradient algorithms for total variation image restoration. *J. Math. Imaging Vis.*, 44: 3, 236–253. 2012.
- [12] R. Burachik, L. M. Graña Drummond, A. N. Iusem, and B. F. Svaiter, Full convergence of the steepest descent method with inexact line searches. *Optimization*, 32: 2, 137–146. 1995.
- [13] K. Bredies, A forward-backward splitting algorithm for the minimization of non-smooth convex functionals in Banach space. *Inverse Problems*, 25. 2009.
- [14] P. H. Calamai and J. J. Moré, Projected gradient methods for linearly constrained problems. *Math. Program.*, 39: 1, 93–116. 1987.
- [15] G. H-G. Chen and R. T. Rockafellar, Convergence rates in forward-backward splitting. *SIAM J. Optim.*, 7: 2, 421–444. 1997.
- [16] E. Chouzenoux, J.-C. Pesquet, and A. Repetti, Variable metric forward–backward algorithm for minimizing the sum of a differentiable function and a convex function. *J. Optim. Theory Appl.*, 162: 1, 107–132. 2014.
- [17] P. L. Combettes, Quasi-Fejérian analysis of some optimization algorithms. In *Inherently Parallel Algorithms in Feasibility and Optimization and Their Applications*, (D. Butnariu, Y. Censor, and S. Reich, Eds.), pp. 115–152. New York: Elsevier, 2001.
- [18] P. L. Combettes and B. C. Vũ, Variable metric quasi-Fejér monotonicity. *Nonlinear Analysis: Theory, Methods & Applications*, 78, 17–31. 2013.
- [19] P. L. Combettes and B. C. Vũ, Variable metric forward-backward splitting with applications to monotone inclusions in duality. *Optimization*, 63(9), 1289–1318. 2014.
- [20] P. L. Combettes and V. R. Wajs, Signal recovery by proximal forward-backward splitting. *Multiscale Modeling and Simulation*, 4:1168–1200, 2005.
- [21] P. L. Combettes and I. Yamada, Compositions and convex combinations of averaged nonexpansive operators. *J. Math. Anal. Appl.*, 425(1):55–70, 2015.
- [22] D. Davis and Y. Yin, Convergence rate analysis of several splitting schemes. arXiv:14064834v3, 2015.
- [23] I. Daubechies, M. Defrise, C. De Mol, An iterative thresholding algorithm for linear inverse problems with a sparsity constraint. *Comm. Pure Appl. Math.*, 57(11), 1413–1457, 2004.
- [24] P. Frankel, G. Garrigos, and J. Peypouquet, Splitting methods with variable metric for Kerdyka-Łojasiewicz functions and general convergence rates. *J. Optim. Theory Appl.*, 165, 874–900, 2015.
- [25] E. M. Gafni and D. P. Bertsekas, Convergence of a gradient projection method. *Technical report LIDS-P-1201, Laboratory for Information and Decision System*, 1982.
- [26] E. M. Gafni and D. P. Bertsekas, Two-metric projection methods for constrained optimization. *SIAM J. Control Optim.*, 22(6), 936–964, 1984.
- [27] A. A. Goldstein, Convex programming in Hilbert spaces. *Bull. Amer. Math. Soc.*, 70(5), 709–710, 1964.

- [28] J.-B. Hiriart-Urruty and C. Lemaréchal, *Convex Analysis and Minimization Algorithms II*. Springer, Berlin 1996.
- [29] A. N. Iusem, On the convergence properties of the projected gradient method for convex optimization. *Comput. Appl. Math.*, 22(1), 37–52, 2003.
- [30] J. D. Lee, Y. Sun, and M. A. Saunders, Proximal Newton-type methods for minimizing composite functions. *SIAM J. Optim.*, 24(3), 1420–1443, 2014.
- [31] E. S. Levitin and B. T. Polyak, Constrained minimization problems. *USSR Comput. Math. Math. Phys.*, 6, 1–50, 1966 (originally in *Zh. Vychisl. Mat. i Mat. Fiz.*, 6, 787–823, 1965).
- [32] J. Lin, L. Rosasco, S. Villa, D-X Zhou, Modified Fejér sequences and applications. arXiv:1510.04641, 2015.
- [33] L. Lindenstrauss and L. Tzafriri, *Classical Banach Spaces II*. Springer, Berlin 1979.
- [34] G. P. McCormick and R. A. Tapia, The gradient projection method under mild differentiability conditions. *SIAM J. Control.*, 10(1), 93–98, 1972.
- [35] J. Nocedal and S. J. Wright, *Numerical Optimization 2nd ed.* Springer, New York 2006.
- [36] L. A. Parente, P. A. Lotito, and M. V. Solodov, A Class of Inexact Variable Metric Proximal Point Algorithms. *SIAM J. Optim.*, 19(1), 240–260, 2008.
- [37] M. Reed and B. Simon, *Methods of Modern Mathematical Physics, Vol 1*. Academic Press, San Diego, 1980.
- [38] S. Salzo, S. Masecchia, A. Verri, and A. Barla, Alternating Proximal Regularized Dictionary Learning. *Neural Comput.*, 26:12, 2014.
- [39] T. Schuster, B. Kaltenbacher, B. Hofmann, and K. S. Kazimierski, *Regularization Methods in Banach Spaces*. De Gruyter, Berlin, 2012.
- [40] Q. Tran-Dinh, A. Kyrillidis, and V. Cevher, Composite self-concordant minimization. *J. Mach. Learn. Res.*, 16: 371–416, 2015.
- [41] P. Tseng, A modified forward-backward splitting method for maximal monotone mapping. *SIAM J. Control Optim.*, 38:431–446, 2000.
- [42] P. Tseng and S. Yun A coordinate gradient descent method for nonsmooth separable minimization. *Math. Program.*, 117:387–423, 2009.