

**$z$ -CLASSES IN FINITE GROUPS OF CONJUGATE TYPE  $(n, 1)$** 

SHIVAM ARORA AND KRISHNENDU GONGOPADHYAY

**ABSTRACT.** Two elements in a group  $G$  are said to  *$z$ -equivalent* or to be in the same  *$z$ -class* if their centralizers are conjugate in  $G$ . In [KKJ16], it was proved that a non-abelian  $p$ -group  $G$  can have at most  $\frac{p^k-1}{p-1} + 1$  number of  $z$ -classes, where  $|G/Z(G)| = p^k$ . In this note, we characterize the  $p$ -groups of conjugate type  $(n, 1)$  attaining this maximal number. As a corollary, we characterize  $p$ -groups having prime order commutator subgroup and maximal number of  $z$ -classes.

## 1. INTRODUCTION

Two elements in a group  $G$  are said to  *$z$ -equivalent* or to be in the same  *$z$ -class* if their centralizers are conjugate in  $G$ . The  $z$ -equivalence is an equivalence relation in  $G$  and give a partition of  $G$  into disjoint subsets. It is coarser than the conjugacy relation. An infinite group may have infinitely many conjugacy classes, but often the number of  $z$ -classes is finite and this finiteness of  $z$ -classes give a rough idea about dynamical types in a homogeneous space on which  $G$  acts, see [Kul07]. There have been some recent work where  $z$ -classes in various groups have been investigated, see [Gou06, Kul08, Sin08, GK09, Gon13, GK16]. We note here that a closely related problem is to compute the *genus number* that has been investigated in [Bos13]. Jadhav and Kitture has classified  $z$ -classes in  $p$ -groups of order up to  $p^5$ , [JK14].

In a recent work Jadhav, Kitture and Kulkarni [KKJ16] has investigated  $z$ -classes in finite  $p$ -groups. They proved that a non-abelian  $p$ -group has at least  $p + 2$   $z$ -classes and characterized the groups that attains this lower bound. They also proved that a non-abelian  $p$ -group can have at most  $\frac{p^k-1}{p-1} + 1$  number of  $z$ -classes, where  $|G/Z(G)| = p^k$ . They proved the following theorem that gives necessary conditions for a  $p$ -group  $G$  to attain this maximal number of  $z$ -classes.

**Theorem JKK.** [KKJ16] *Let  $G$  be a non-abelian group with  $|G/Z(G)| = p^k$ . If the number of  $z$ -classes in  $G$  is  $\frac{p^k-1}{p-1} + 1$  then either  $G/Z(G) \cong C_p \times C_p$  or the following holds:*

- (1)  $G$  has no abelian subgroup of index  $p$ .
- (2)  $G/Z(G)$  is an elementary abelian  $p$ -group.

---

2010 *Mathematics Subject Classification.* 20D15 (Primary); 20E45 (Secondary).

*Key words and phrases.* conjugacy classes of centralizers,  $z$ -classes,  $p$ -groups, extraspecial groups.

The authors further gave a counterexample to show that conditions (1) and (2) are not sufficient to attain this bound. It would be interesting to obtain sufficient conditions for a group  $G$  to attain the maximal number of  $z$ -classes and to characterize such groups.

In this note, we characterize groups of type  $(n, 1)$  attaining this bound. To state our result, we recall the following definition by Ito [Ito53].

**Definition.** [Ito53] *Let  $G$  be a finite group. Let  $n_1, n_2, \dots, n_r$ , where  $n_1 > n_2 > \dots > n_r = 1$ , be all the numbers which are the indices of the centralizers of elements of  $G$  in  $G$ . The vector  $(n_1, n_2, \dots, n_r)$  is called conjugate type vector of  $G$ . A group with the conjugate type vector  $(n_1, n_2, \dots, n_r)$  is called a group of type  $(n_1, n_2, \dots, n_r)$ .*

In [Ito53], Ito characterizes the groups of type  $(n, 1)$ . Any such group is nilpotent and  $n = p^a$  for some prime  $p$ . A group of type  $(p^a, 1)$  is the direct product of a  $p$ -group of the same type and of an abelian group. Thus, the study of groups of type  $(n, 1)$  can be reduced to that of  $p$ -groups of type  $(p^a, 1)$ . Ito showed that such a group  $G$  contains an abelian normal subgroup  $A$  such that  $G/A$  is of exponent  $p$ .

Here we prove the following.

**Theorem 1.1.** *Let  $G$  be a non-abelian group of type  $(n, 1)$  and  $[G : Z(G)] = p^k$ . Then  $G$  has  $\frac{p^k - 1}{p - 1} + 1$   $z$ -classes if and only if*

- (1)  $G/Z(G)$  is elementary abelian, and
- (2) for all  $x \in G \setminus Z(G)$ ,  $Z(C_G(x)) = \langle x, Z(G) \rangle$ .

As an application of the above result we have the following.

**Corollary 1.2.** *Let  $G$  be a non-abelian group with  $|G/Z(G)| = p^k$ , ( $k \geq 2$ ) and  $|G'| = p$ . Then the number of  $z$ -classes in  $G$  is  $\frac{p^k - 1}{p - 1} + 1$  if and only if  $G$  is isoclinic to an extraspecial  $p$ -group.*

We prove these results in Section 3.

We note that Corollary 1.2 may be compared with Theorem 5.3.1 of Kitture [Kit14, Section 5.3]. We thank Kitture for letting us know about his result and many comments. We are also thankful to Silvio Dolfi for useful comments and suggestions on a first draft of this note.

## 2. PRELIMINARIES

We recall here a few basic facts from Kulkarni [Kul07] that will be used in the proof.

Let  $G$  be a group. We denote  $C_G(x)$  to be the centralizer of an element  $x$  in  $G$ .

**Definition 2.1.** *Let  $G$  be a group. Two elements  $x, y \in G$  are said to be  $z$ -equivalent if their centralizers are conjugate, i.e. there exist  $g \in G$  such that,*

$$g^{-1}C_G(x)g = C_G(y).$$

Let  $G$  be a group acting on a set  $X$ . For  $x \in X$  let  $G(x)$  denote the orbit of  $x$  and  $G_x$  be the stabilizer subgroup of  $x$ :

$$G(x) = \{y \in X : \text{there exists } g \in G \text{ and } y = gx\}.$$

$$G_x = \{g \in G : gx = x\}.$$

We have a partition of  $X$  as:

$$X = \bigcup_{x \in X} G(x).$$

Let  $y \in G(x)$  be some element, then for some  $g \in G$  we have  $y = gx$  and  $G_y = gG(x)g^{-1}$ .

For subgroups  $A$  and  $B$  in  $G$  write  $A \sim_g B$  if they are conjugate. This gives us the following.

**Definition 2.2.** Let  $x, y \in X$  be any two element, they are said to be in same orbit class if  $G_x \sim_g G_y$ . Denote this equivalence relation as  $x \sim_0 y$  and denote  $R(x)$  as the equivalence class of  $x$ .

This gives another partition of  $X$ :

$$X = \bigcup_{G_x \sim_g G_y} R(x) = \bigcup_{G_x = G_y} R(x).$$

The second inequality is easy to note, since if  $x \sim_0 y$  then for some  $z \in X$ , we have  $C_G(x) = C_G(z)$ .

Let  $F_x$  denotes the points fixed by  $G_x$  i.e.

$$F_x = \{y \in X : G_y \supset G_x\}.$$

Let  $F'_x$  denotes the set  $F'_x = \{y \in X : G_y = G_x\}$ . Define  $W_x = N_x/G_x$ , where  $N_x$  is the normalizer of the  $G_x$  in  $G$ . Then we have the following theorem(refer [Kul07]).

**Theorem 2.3.** Let  $G$  act on a set  $X$ . Then the map  $\phi : G/G_x \times F'_x \rightarrow R(x)$ ,  $\phi(gG_x, y) = gy$  is well defined and induces a bijection,  $\bar{\phi} : (G/G_x \times F'_x)/W_x \rightarrow R(x)$ .

When we take our space  $X = G$ , the group itself, then  $R(x)$  in above notation becomes the  $z$ -classes and the above theorem gives us the following equation, which is used to calculate the size of a  $z$ -class: for any  $x \in G$  we have

$$|z - \text{class of } x| = [G : N_G(C_G(x))] \cdot |F'_x|,$$

where  $C_G(x)$  is the centralizer of  $x$  and  $F'_x = \{y \in G : C_G(y) = C_G(x)\}$ .

The following theorem states that that number of  $z$ -classes is invariant under the isoclinism. For basic notions on isoclinism, see [Hal40, KKJ16].

**Theorem 2.4.** [KKJ16] Let  $G_1$  and  $G_2$  be isoclinic groups, with an isoclinism  $(\phi, \psi)$ . Then the isoclinism  $(\phi, \psi)$  induces a bijection between the  $z$ -classes in  $G_1$  and  $G_2$ .

Finally we note the following theorem by Hall [Hal40] that is used in proof of Theorem 1.2.

**Theorem 2.5.** Every group is isoclinic to a group whose center is contained in the commutator subgroup.

## 3. PROOF OF THE MAIN RESULTS

It was proved in [KKJ16, Lemma 3.1] that if  $G$  is a finite group with  $|G/Z(G)| = p^k$ , then  $G$  is isoclinic to a finite  $p$ -group. Since number of  $z$ -classes is invariant under isoclinism, hence it is enough to prove our results assuming that  $G$  is a  $p$ -group.

**Lemma 3.1.** *Let  $G$  be a non-abelian  $p$ -group with  $[G : Z(G)] = p^k$ . If*

- (1)  $G/Z(G)$  is elementary abelian, and
- (2) for all  $z \in G \setminus Z(G)$ ,  $Z(C_G(x)) = \langle x, Z(G) \rangle$ ,

*then  $G$  has  $\frac{p^k-1}{p-1} + 1$   $z$ -classes.*

*Proof.* Let  $x \in G \setminus Z(G)$ . Since  $G/Z(G)$  is abelian, hence  $C_G(x) \trianglelefteq G$ , i.e.  $[G : N(C_G(x))] = 1$ . Consider

$$F'_x = \{y \in G : C_G(y) = C_G(x)\}.$$

By definition  $F'_x \subseteq Z(C_G(x)) \setminus Z(G) = \langle x, Z(G) \rangle \setminus Z(G)$ . Obviously  $C_G(x) = C_G(xt)$  for  $t \in Z(G)$ . Also since exponent of  $G/Z(G)$  is  $p$ , thus  $C_G(x) = C_G(x^i)$  for any  $i \in \{1, 2, \dots, p-1\}$ . Hence  $\langle x, Z(G) \rangle \setminus Z(G) \subseteq F'_x$ . Hence

$$F'_x = Z(C_G(x)) \setminus Z(G) = \langle x, Z(G) \rangle \setminus Z(G) = xZ(G) \cup \dots \cup x^{p-1}Z(G).$$

This implies,  $|F'_x| = (p-1)|Z(G)|$ . Therefore,

$$|z\text{-class of } x| = [G : N_G(C_G(x))] \cdot |F'_x| = 1 \cdot (p-1)|Z(G)|.$$

Thus  $G \setminus Z(G)$  contributes to a total  $\frac{p^k-1}{p-1}$  number of  $z$ -classes of  $G$ . Clearly,  $Z(G)$  is also a  $z$ -class. Hence,  $G$  has  $\frac{p^k-1}{p-1} + 1$   $z$ -classes.  $\square$

**Corollary 3.2.** *Let  $G$  be a non-abelian  $p$ -group with  $|G/Z(G)| = p^k$ . If*

- (1)  $G$  has no abelian subgroup of order exceeding  $p|Z(G)|$ .
- (2)  $G/Z(G)$  is an elementary abelian  $p$ -group.

*Then  $G$  has  $\frac{p^k-1}{p-1} + 1$  number of  $z$ -classes.*

*Proof.* We claim  $C_G(x) = \langle x, Z(G) \rangle$ , and the rest follow from Lemma 3.1.

Clearly  $\langle x, Z(G) \rangle \subseteq C_G(x)$ . Let us assume  $y \notin \langle x, Z(G) \rangle$  and  $y \in C_G(x)$ . But then  $\langle x, y, Z(G) \rangle$  is an abelian subgroup and  $|\langle x, y, Z(G) \rangle| > p|Z(G)|$ . Thus  $C_G(x) = \langle x, Z(G) \rangle$ .  $\square$

## 3.1. Proof of Theorem 1.1.

*Proof.* Let  $G$  be a non-abelian group of type  $(n, 1)$  and  $G$  has  $\frac{p^k-1}{p-1} + 1$   $z$ -classes, where  $[G : Z(G)] = p^k$ . (1) follows from the Theorem JKK. We prove (2).

Let  $x \in G \setminus Z(G)$ , then for any  $t \in Z(G)$ ,  $C_G(x) = C_G(xt)$ . Also for any  $m$  relatively prime to order of  $x$ , we have

$$C_G(x) = C_G(\langle x \rangle) = C_G(\langle x^m \rangle)$$

Thus  $z$ -class of  $x$  contains,

$$xZ(G) \cup x^2Z(G) \cup \dots \cup x^{p-1}Z(G),$$

and hence, for all  $x \in G \setminus Z(G)$ , the size of  $z$  class is at least  $(p-1)|Z(G)|$ . Since  $Z(G)$  is also a  $z$ -class, hence it follows from the hypothesis that  $G \setminus Z(G)$  constitute exactly  $\frac{p^k-1}{p-1}$   $z$ -classes. Now,  $|G \setminus Z(G)| = |G| - |Z(G)|$ . Hence, we must have for  $x \in G \setminus Z(G)$ ,

$$|z\text{-class of } x| = (p-1)|Z(G)|.$$

Now, it is clear that for  $x \in G \setminus Z(G)$ ,  $\langle x, Z(G) \rangle \subseteq Z(C_G(x))$ . Now let  $y \in Z(C_G(x))$  such that  $y \notin \langle x, Z(G) \rangle$ . Then a simple observation shows that  $C_G(y) \supseteq C_G(x)$ . But since  $G$  is of type  $(n, 1)$  we have  $|C_G(x)| = |C_G(y)|$ , thus we must have  $C_G(x) = C_G(y)$ . This implies  $y \in (z\text{-class of } x)$  and since  $y \notin \langle x, Z(G) \rangle$ , thus  $|z\text{-class of } x| > (p-1)|Z(G)|$ , which is a contradiction. Thus our assumption is incorrect and  $Z(C_G(x)) = \langle x, Z(G) \rangle$  for all  $x \in G \setminus Z(G)$ .

The converse part follows from Lemma 3.1. This completes the proof.  $\square$

### 3.2. Proof of Corollary 1.2.

*Proof.* : Without loss of generality, using Theorem 2.5, we assume up to isoclinism that,  $G$  is a group such that  $Z(G) \subseteq G'$ . Let  $G$  has  $\frac{p^k-1}{p-1} + 1$   $z$ -classes. Since  $|G'| = p$  and in a non-abelian  $p$ -group  $|Z(G)| > 1$ , it follows that  $G' = Z(G)$ ; hence,  $|G'| = |Z(G)| = p$ . Therefore  $G$  and  $G/Z(G)$  have the same exponent. Thus  $G$  is an extra-special  $p$ -group.

Conversely, let  $G$  be an extraspecial  $p$ -group. So,  $|Z(G)| = |G'| = p$ . For  $x \in G \setminus Z(G)$ , define the map:

$$\begin{aligned} \phi_x : G &\rightarrow G' \\ g &\mapsto [x, g] \end{aligned}$$

As  $G/Z(G)$  is elementary abelian,  $\phi_x$  is a homomorphism with  $\ker(\phi_x) = C_G(x)$  and since  $|G'| = p$ , the map is also surjective. Hence we get

$$G/C_G(x) \cong G'$$

Thus each  $C_G(x)$  is a maximal subgroup of index  $p$  in  $G$ . This shows that extraspecial  $p$ -groups are of type  $(n, 1)$ .

Now, we claim that  $Z(C_G(x)) = \langle x, Z(G) \rangle$ . This follows from the fact that  $G/Z(G)$  can be equipped with an alternating non-degenerate form induced by the commutator, see the proof of [Cra08, Theorem 3.14] or [Wil09, p. 83]. The claim follows from [Cra08, Lemma 3.11] noting that  $Z(C_G(x))/Z(G)$  is the radical of codimension one subspace  $C_G(x)/Z(G)$ .

The result now follows from Theorem 1.1.  $\square$

### REFERENCES

- [Bos13] Anirban Bose. On the genus number of algebraic groups. *J. Ramanujan Math. Soc.*, 28(4):443–482, 2013.
- [Cra08] David A. Craven. The theory of  $p$ -groups. <http://web.mat.bham.ac.uk/D.A.Craven/docs/lectures/pgroups.pdf>, 2008.
- [GK09] Krishnendu Gongopadhyay and Ravi S. Kulkarni.  $z$ -classes of isometries of the hyperbolic space. *Conform. Geom. Dyn.*, 13:91–109, 2009.

- [GK16] Krishnendu Gongopadhyay and Ravi S. Kulkarni. The  $z$ -classes of isometries. *Journal of Indian Mathematical Society*, 81(3-4), 2016.
- [Gon13] Krishnendu Gongopadhyay. The  $z$ -classes of quaternionic hyperbolic isometries. *J. Group Theory*, 16(6):941–964, 2013.
- [Gou06] Rony Gouraige. *Z-classes in central simple algebras*. ProQuest LLC, Ann Arbor, MI, 2006. Thesis (Ph.D.)—City University of New York.
- [Hal40] P. Hall. The classification of prime-power groups. *Journal für die reine und angewandte Mathematik*, 182:130–141, 1940.
- [Ito53] Noboru Ito. On finite groups with given conjugate types I. *Nagoya Math. J.*, 6:17–28, 1953.
- [JK14] Vikas S. Jadhav and Rahul D. Kitture.  $z$ -classes in  $p$ -groups less than equal to  $p^5$ . *Bull. Allahabad. Soc.*, 29:179–194, 2014.
- [Kit14] Rahul D. Kitture.  *$z$ -Classes in finite  $p$ -groups*. 2014. Thesis (Ph.D.)—University of Pune.
- [KKJ16] Ravindra Kulkarni, Rahul Dattatraya Kitture, and Vikas S. Jadhav.  $z$ -classes in groups. *Journal of Algebra and Its Applications*, 15(6):165013 (16 pages), 2016.
- [Kul07] Ravi S. Kulkarni. Dynamical types and conjugacy classes of centralizers in groups. *J. Ramanujan Math. Soc.*, 22(1):35–56, 2007.
- [Kul08] Ravi S. Kulkarni. Dynamics of linear and affine maps. *Asian J. Math.*, 12(3):321–343, 2008.
- [Sin08] Anupam Singh. Conjugacy classes of centralizers in  $G_2$ . *J. Ramanujan Math. Soc.*, 23(4):327–336, 2008.
- [Wil09] Robert A. Wilson. *The finite simple groups*, volume 251 of *Graduate Texts in Mathematics*. Springer-Verlag London, Ltd., London, 2009.

INDIAN INSTITUTE OF SCIENCE EDUCATION AND RESEARCH (IISER) MOHALI, KNOWLEDGE CITY, , SECTOR 81, S.A.S. NAGAR, PUNJAB 140306, INDIA  
*E-mail address:* shivam.iiserm@gmail.com

INDIAN INSTITUTE OF SCIENCE EDUCATION AND RESEARCH (IISER) MOHALI, KNOWLEDGE CITY, SECTOR 81, S.A.S. NAGAR, PUNJAB 140306, INDIA  
*E-mail address:* krishnendug@gmail.com, krishnendu@iisermohali.ac.in