

VALUED MODULES OVER SKEW POLYNOMIAL RINGS I

GÖNENÇ ONAY

ABSTRACT. We introduce a notion of valued module which is suitable to study valued fields of positive characteristic. Then we built-up a robust theory of henselianity in the language of valued modules and prove Ax-Kochen Ershov type results.

1. INTRODUCTION

A valued abelian group, is an abelian group M together with a linearly ordered set Δ and a function $v : M \rightarrow \Delta$, such that

- a. $v(x \pm y) \geq \min\{v(x), v(y)\}$
- b. $v(x) = \infty \Leftrightarrow x = 0$

where ∞ is the maximum of Δ .

A module such that the underlying abelian group is valued will be called a valued module. Depending on the context one wants to study, various compatibility assumptions relating the valuation v and the action of scalars are considered in the literature. The first article on the subject that the author is aware of is due to Fleischer ([5]); it extends the notion of Krull valuation to modules where $v(x.r) \geq v(x)$ for any scalar r and proves the equivalence of maximality and (pseudo-) convergence of pseudo-Cauchy series. In Rohwer's thesis ([14]) modules are also used to understand the model theory of valued fields of positive characteristic or more generally valued difference fields. In more recent articles of Point and Bélair [1] (resp. Maalouf [9]) valued modules (resp. vector spaces) are studied.

Similarly as in [14], we are interested in modules which come from the theory of valued fields in characteristic $p > 0$. The rings that we will consider are unitary and arise as K -subalgebras of the endomorphisms ring of a field K generated by a distinguished endomorphism φ over K .

Definition 1.1. *Let K be a field and φ a ring endomorphism of K . The ring $K[t; \varphi]$ is given as a set by formal sums $\sum_{i \in \mathbb{N}} t^i a_i$ where t is the indeterminate, $a_i \in K$ and $\{i \in \mathbb{N} \mid a_i \neq 0\}$ is finite. The addition is defined term by term and the multiplication obeys the (non-)commutation rule: $at = t\varphi(a)$. More precisely,*

- (1) $\sum_{i \in \mathbb{N}} t^i a_i + \sum_{i \in \mathbb{N}} t^i b_i = \sum_{i \in \mathbb{N}} t^i (a_i + b_i)$
- (2) $(\sum_{i \in \mathbb{N}} t^i a_i) (\sum_{i \in \mathbb{N}} t^i b_i) = \sum_{i \in \mathbb{N}} t^i c_i$ where $c_i = \sum_{k+l=i} \varphi^l(a_k) b_l$.

Most often for fixed K and φ we will refer to this ring $K[t; \varphi]$ as R . Any field K of characteristic of $p > 0$ has a natural module structure over its ring of additive polynomials: these are polynomials of the form $\sum_{i=0}^n a_i X^{p^i}$, where the ring operations are given by usual addition and (right-)composition. This ring is isomorphic to the ring $K[t; \text{Frob}]$ where Frob is the map $x \mapsto x^p$. It is widely believed that serious problems in the model theory of valued fields in positive characteristic arise from additive polynomials (see for example [18]). Studying and

The author would like to thank *Equipe de Logique Mathématiques de Paris* for supporting his stay during September 2013 where the current work is mainly completed. Françoise Delon read this paper several times, made lot of valuable comments, to whom the author is very much grateful. Many thanks go to Piotr Kowalski for his final remarks.

axiomatizing the underlying ultrametric module structure can be considered as an abstract analysis that isolates this phenomenon. Note also that any valued field together with an arbitrary automorphism inherits an analogue module structure.

After the famous Ax-Kochen and Ershov (A-K, E) theorem which provides a corrected form of a conjecture of Artin¹, there were similar theorems permitting to understand the first order theories of some other class of valued fields (see [17]) for a survey). In this paper, we establish (A-K, E) type results by proving a relative elimination of quantifiers for modules called *henselian divisible valued R -modules*. We first observe the theory of that divisible valued modules is not suitable to reach our aim (cf. counter example 3.26 and the remark which follows). Thus we add an extra assumption on the valuation, which comes from an interpretation of Hensel's lemma as a local inversion theorem. More precisely, in this context, there is an abstract analogue of the maximal ideal of a valuation ring; call this $M_{>\theta}$ and then our assumption asserts that the multiplication by some special elements of R (we will call them *separable polynomials*) induces a bijection $M_{>\theta} \rightarrow M_{>\theta}$.

Fix $R = K[K; \varphi]$ as above for given φ and K . In our model theoretical setting valued R -modules are considered in the two sorted first order language

$$L := \{0, +, -, (.r)_{r \in R}\} \cup \{<, \tau, \infty, (R_n)_{n \in \mathbb{N}}\} \cup \{v\}$$

where

- (1) the *module sort* language is $L_{Mod}(R) := \{0, +, -, \{.r\}_{r \in R}\}$ (which is the usual language of modules),
- (2) the *value set sort* language is $L_V := \{<, \tau, \infty, (R_n)_{n \in \mathbb{N}}\}$ with τ a unary function symbol and R_n a unary predicate symbol for each n ,
- (3) v is a unary function symbol to be interpreted as the valuation.

We describe now the content of our paper.

In section 2, we consider pure R -modules and describe the completions of the theory of divisible R -modules as follows:

Theorem 1.2. *Any complete theory of non zero divisible R -modules admits elimination of quantifiers and is obtained by specifying for each irreducible $q \in R$, the number of elements annihilated by q .*

In the Section 3, we introduce valued R -modules (M, Δ, v) which by definition satisfy the following compatibility properties:

- (1) $v(x.\lambda) = v(x)$ for all unit λ in R .
- (2) The map $x \rightarrow v(x.t)$ induces a strictly increasing function $\tau : \Delta \rightarrow \Delta$ and there exists at most one value $\theta \in \Delta$, such that, if $r \in R \setminus \{0\}$ and $x \in M$ with $v(x) \neq \theta$, then $v(x.r) = \tau^k(v(x))$ for some $k \in \mathbb{N}$ depending only on r and $v(x)$.

Note that any valued field (U, v) of characteristic $p > 0$ inherits naturally a valued $K[t; Frob]$ -module structure where $K \subseteq U$ is trivially valued by v and the function τ is $\gamma \mapsto p\gamma$.

We then investigate the L -theory of *henselian* and *henselian divisible R -modules*. We have the following main theorem which enables us to recover the theory of a divisible henselian valued R -module from the theory of its value set (in L_V) and from the theory of its torsion submodule (in $L_{Mod}(R)$) which, in some sense, plays the role of the *residue field*.

¹Artin conjectured that the field \mathbb{Q}_p is C_2 (i.e. every homogeneous polynomial of degree d with $> d^2$ variables has a non trivial zero) for any prime p . This is false by a result of Terjanian (see [16]).

Theorem 1.3 (A-K,E \equiv). *Let (F, v) and (G, w) be two non zero henselian divisible modules such that F_{tor} and G_{tor} are elementary equivalent as $L_{\text{Mod}}(R)$ -structures and $v(F)$ and $v(G)$ are elementary equivalent in the language L_V . Then (F, v) and (G, v) are elementary equivalent as L -structures.*

This article corresponds to Chapters Two and Three of author's thesis (cf. [11]).

2. DIVISIBLE R -MODULES

First, we quickly summarize some basic algebraic facts about the rings over which we will consider our modules. One can refer to [2] section 2.1 for more details.

Let K be a field and φ a *field endomorphism* of K . Set $R := K[t; \varphi]$, the ring of φ -twisted polynomials in the variable t (see Definition 1.1 in Introduction). We will mean by **polynomial** any element of this ring. Each $r \in R$ can be written uniquely as $\sum_{i \in \mathbb{N}} t^i a_i$, with $I := \{i \mid a_i \neq 0\}$ finite. The ring R is equipped with the degree function defined by $\deg(r) = \max I$ (with the convention that $\max \emptyset = -\infty$). We have $\deg(pq) = \deg(qp) = \deg(p) + \deg(q)$. Note that R is a domain, i.e. has no zero divisors. The set of units of R is K^\times and together with the function \deg , R is right euclidean (and hence right principal)²: for all $q, q' \in R$ with $q' \neq 0$ there exists $r, r' \in R$ such that $q = q'r' + r$ where $\deg(r) < \deg(q')$ and r is unique up to right multiplication by a unit. We say that q' divides q if q is a right multiple of q' , that is if $r = 0$ above. For any non zero a and b there exists a unique monic common divisor of maximal degree, denoted by $\gcd(a, b)$.

Furthermore, by the commutation rule $at = t\varphi(a)$, any non zero element q can be written as

$$(1) \quad q = t^n q_1 \dots q_s$$

where t and the q_i are irreducible. Note that R is in particular a right Ore ring, that is every two non zero elements a and b have a non zero common right multiple. Taking into account the degree function there is a common right multiple of least degree unique up to a unit: we denote it $\text{lcm}(a, b)$ as usual. As any Ore ring, R is embedded into a smallest (right-)division ring, called its skew field of fractions.

Notation 2.1. We will denote by D the skew field of fractions of R .

Any R -module that we will consider is a right R -module, and we will simply say R -module.

The following result is an easy observation as in the case of a torsion free divisible abelian groups:

Lemma 2.2. *Any torsion free divisible module over a Ore ring S has a unique vector space structure over the skew field of fraction of S . In particular, every divisible torsion free R -module is a D -vector space.*

Notation 2.3. Let M be an R -module. We denote by M_{tor} the set

$$\{x \in M \mid \exists r \in R \setminus \{0\} \text{ such that } x.r = 0\}.$$

Lemma 2.4. *If M is a divisible R -module then M_{tor} is a divisible submodule of M .*

Proof. Since R is a Ore ring, M_{tor} is a submodule (cf. [3] proposition 3.5). Moreover, if $0 \neq x \in M_{\text{tor}}$, with $q \in R \setminus \{0\}$ such that $x.q = 0$ then, for all $r \in R \setminus \{0\}$ and for all $y \in M$ such that $y.r = x$, we have $y.r.q = 0$. Therefore $y \in M_{\text{tor}}$, which shows that M_{tor} is divisible. \square

² Note that this ring is also left euclidean whenever φ is an automorphism.

Remark 2.5. Any torsion free non zero divisible R -module is infinite since it is a D -vector space.

Furthermore if M is a non zero divisible R -module then for every $r \neq 0$, $M.r = M \neq 0$. Hence $M_{\text{tor}}.r \neq 0$ when $r \neq 0$ and $M_{\text{tor}} \neq 0$. In particular, M_{tor} is infinite if it is non zero, since otherwise, the least common multiple of annihilators of non zero elements of M_{tor} would annihilate M_{tor} .

Recall that a module I over a ring S is said to be injective if for any given inclusion of S -modules $N \subseteq M$, and any homomorphism $f : N \rightarrow I$, f admits an extension to M .

In the following lemma we state that in the case of R -modules the notions of injectivity and divisibility coincide, and we also summarize well-known facts about injective modules that we will frequently use in the rest of this paper.

Lemma 2.6. Let S be any ring.

- (1) A divisible module over a right principal ring is injective, and an injective module over a domain is divisible. In particular for every R -module M , M is divisible if and only if it is injective.
- (2) Any injective S -module is a direct summand in every S -module which contains it.
- (3) Any S -module M has an extension $N \supseteq M$ maximal with the property

$$X \cap M \neq 0 \text{ for any non zero submodule } X \subseteq N.$$

N is called an injective hull of M and is unique up to an M -isomorphism of S -modules.

- (4) Let $N \supseteq M$ be R -modules. If N is an injective hull of M and $x \in N \setminus \{0\}$, then there exists $r \in R$ such that $x.r \in M \setminus \{0\}$.

Proof. We refer to [7] (p. 156-164). The last assertion is Exercise 3. page 164. \square

Definition 2.7. For R -modules $M \subseteq N$, where N is divisible, we call divisible closure of M any injective hull of M inside N .

Remark 2.8. (1) Let $M \subseteq E \subseteq N$ be R -modules, with E and N divisible. Then E is a divisible closure of M , if and only if for all $x \in E$ there exists a non zero $r \in R$ such that $x.r \in E$. Moreover note that E can be different from

$$\{x \in N \mid \exists r \in R, r \neq 0 \text{ \& } x.r \in M\}.$$

- (2) By lemma 2.6, divisible closures of an R -module are isomorphic as pure modules. When we will consider valued modules we will see that the various divisible closures are generally neither isomorphic nor even elementary equivalent as valued modules (see Proposition 3.26 and Corollary 3.27).

The following observation is essential:

Lemma 2.9. Let M be an R -module, $A \subseteq M$ a submodule of M and $x \in M \setminus A$ such that $x.r \in A$ for some $r \in R \setminus \{0\}$. Then there exists $q \in R$ of minimal degree such that $x.q \in A$. For such a q and for all r such that $x.r \in A$, q divides r .

Proof. Let $I := \{r \in R \mid x.r \in A\}$; I is a right ideal, therefore principal. Hence there is a generator of I , that we can take as q . \square

Definition 2.10. Any polynomial q as above is called a minimal polynomial of x over A . In particular, for all $x \in M_{\text{tor}} \setminus \{0\}$, there exists monic polynomials of minimal degree such that $x.q = 0$. It is called a minimal polynomial of x .

Remark 2.11. A minimal polynomial of x over A divides any minimal polynomial of x . If $x \in M \setminus A$ and $x.r \in A$ with r irreducible then r is a minimal polynomial of x over A . Minimal polynomials are unique up to a unit.

Definition 2.12. For any $r \in R$ and any R -module M , the annihilator set of r in M , is the set $\text{ann}_M(r) := \{x \in M \mid x.r = 0\}$ and its elements are called zeros or roots of r .

Notation 2.13. For an R -module M , let η_M be the cardinal-valued function defined for all $q \in R$, by $\eta_M(q) := |\text{ann}_M(q)|$.

Proposition 2.14. Let M and N be divisible R -modules satisfying $\eta_M \leq \eta_N$. Suppose $\eta_M(q)$ is finite for all non zero q and $f : A \rightarrow N$ is an R -module embedding where $A \subseteq M_{\text{tor}}$ is a submodule. Then f extends to an R -module embedding of M_{tor} into N_{tor} . In particular if $\text{ann}_M(q) = \text{ann}_N(q) < \infty$ for all non zero $q \in R$, then M_{tor} and N_{tor} are isomorphic.

Proof. Note that the range of f is in N_{tor} . Let $x \in A \setminus M_{\text{tor}}$ and q be a minimal polynomial of x over A . Write $q = q_1 \dots q_k$ where the q_i are irreducible. Since q is a minimal polynomial of x over A , $z := x.q_1 \dots q_{k-1} \notin A$. Set $r := q_k$. Since r is irreducible it is a minimal polynomial of z over A .

Claim : There exists $y \in N_{\text{tor}} \setminus f(A)$ such that $y.r = f(z.r)$.

Proof of the Claim. Set $b := f(z.r)$. Since N_{tor} is divisible it contains some b_1 such that $b_1.r = b$. Suppose $b_1 \in f(A)$. Then $a_1 := f^{-1}(b_1) \in A$. It follows that $(a_1 - z).r = 0$. Note that $(a_1 - z) \notin A$ and r has the same number of zeros in A and $f(A)$. Since $\eta_M(r)$ is finite and $\eta_M(r) \leq \eta_N(r)$ by hypothesis, there exists $b_0 \in N \setminus f(A)$ such that $b_0.r = 0$. Now $b_0 + b_1 \notin f(A)$ and $(b_0 + b_1).r = b.$

Take such an y , then since r is irreducible it is a minimal polynomial of y over A . Now it is easy to check that the map $A + z.R \rightarrow f(A) + y.R$ sending y to z is an isomorphism of R -modules.

Using Zorn Lemma we can extend f to M_{tor} . □

2.1. Complete theories of divisible R -modules. Let S be a ring. Let $L_{\text{Mod}}(S) := \{0, +, -\} \cup \{.s \mid s \in S\}$ be the language of (right-) S -modules. An equation is an $L_{\text{Mod}}(S)$ -formula $\gamma(x_1, \dots, x_m)$ of the form

$$\gamma(\bar{x}) : x_1.r_1 + \dots + x_m.r_m = 0.$$

A primitive positive (p.p.) formula $\phi(x_1, \dots, x_m)$ of the language of S -modules is a formula of the form

$$\exists \bar{y} \ (\gamma_1(\bar{x}, \bar{y}) \wedge \dots \wedge \gamma_n(\bar{x}, \bar{y}))$$

where the γ_i are equations. For instance, by choosing $\gamma_1(x, y) = x.1 + (-y).q = 0$ and $\gamma_2(x, y) = x.r + y.0 = 0$, the formula $\phi(x) : \exists y \ (y.q = x \wedge x.r = 0)$ is a p.p. formula. Note also that a p.p. formula can be written as

$$\exists y_1, \exists y_2, \dots, \exists y_k \ (y_1, \dots, y_k).A = (x_1, \dots, x_n).B$$

where A and B are matrices with coefficients in S . Such formulas define subgroups, called p.p. subgroups. It is a well known fact that modulo a complete theory of S -modules every formula is equivalent to a boolean combination of p.p. formulas. See [13] for more details. Moreover:

Proposition 2.15. The complete theory of a S -module $\langle M, +, -, 0, \{.s\}_{s \in S} \rangle$ is given by the p.p. indexes (these are cardinalities in $\mathbb{N} \cup \{\infty\}$ of relative quotients of p.p. subgroups of M).

In our case we have a simpler description of p.p. definable sets.

Proposition 2.16. *Modulo the theory of divisible R -modules, any p.p. formula is equivalent to a conjunction of atomic formulas. Consequently, a p.p. formula with only one free variable is equivalent to one of the form $x.q = 0$ for some $q \in R$.*

Proof. Let $\phi(x_1, \dots, x_n) := \exists y_1, \dots, y_k \ \overline{y}.A = \overline{x}.B$ be a p.p. formula. By Proposition 6.1 of [3] there exist invertible matrices P and Q , where P has coefficients in $\{0, 1\}$, such that $A' := PAQ$ is of the form $(A_1, 0)$, where A_1 is a $k \times l$ lower triangular matrix in which each coefficient on the diagonal is non zero³ and 0 is $k \times (m - l)$ zero matrix. Thus ϕ is equivalent to the formula:

$$\exists \overline{y} \ \overline{y}.PAQ = \overline{x}.BQ.$$

So, we can replace A by A' and B by BQ and assume that A is of the form $(A_1, 0)$, as described above. It follows that ϕ is of the form $\phi_1(\overline{x}) \wedge \phi_2(\overline{x})$, where ϕ_1 can be written as

$$\exists \overline{y} \ \overline{y}.A_1 = \left(\sum_{i=1}^n x_i.b_{i,1}, \dots, \sum_{i=1}^n x_i.b_{i,l} \right)$$

and ϕ_2 as

$$\left(\sum_{i=1}^n x_i.b_{i,l+1}, \dots, \sum_{i=1}^n x_i.b_{i,m} \right) = (0, \dots, 0).$$

Since the diagonal coefficients of A_1 are all non-zero, any divisible R -module satisfy $\forall \overline{x} \ \phi_1(\overline{x})$. Hence ϕ is equivalent to ϕ_2 . In particular, if x is a single variable, we have:

$$\phi(x) \longleftrightarrow \bigwedge_{j=1}^m x.b_{1,j} = 0 \longleftrightarrow x.\gcd(b_{1,1}, \dots, b_{1,m}) = 0.$$

□

Corollary 2.17. *Completions of the theory of divisible non zero R -modules admit elimination of quantifiers. In addition if M is a divisible R -module such that $M_{\text{tor}} \neq 0$ then M_{tor} is an elementary substructure of M .*

Proof. The first assertion follows directly from the above proposition and the fact that modulo a complete theory of R -modules, any formula is equivalent to a boolean combination of p.p. formulas.

For the second assertion, first observe that M_{tor} is a pure submodule of M since it is divisible (lemma 2.4). Thus, by [15], it is enough to show that M_{tor} and M are elementary equivalent. To obtain this we will show that for p.p. formulas $\phi(x)$ and $\psi(x)$ with $\phi(M) \subseteq \psi(M)$, we have $\psi(M)/\phi(M) = \psi(M_{\text{tor}})/\phi(M_{\text{tor}})$. By the above proposition $\phi(x)$ and $\psi(x)$ are respectively equivalent to $x.r = 0$ and $x.q = 0$ for some $r, q \in R$. If r and q are both non zeros, then there is nothing to show. If $r = 0$ then $\phi(M) = M = \psi(M)$ necessarily. Hence it is enough to consider the non trivial case where $r \neq 0$ & $q = 0$. But $M_{\text{tor}}/\phi(M_{\text{tor}})$ is infinite in this case. In fact, if not, then choosing a set of representatives x_1, \dots, x_n of M modulo $\phi(M)$ and non zero polynomials r_1, \dots, r_n such that $x_i.r_i = 0$, we have $M_{\text{tor}}.\text{lcm}(r, r_1, \dots, r_n) = 0$. This is impossible by Remark 2.5. □

To prove Theorem 1.2 we will study relative quotients of p.p. subgroups of a divisible R -module M . They are of the form $\text{ann}_M(r)/\text{ann}_M(s)$ by elimination of quantifiers.

Set $K_0 := \{x \in K \mid \varphi(x) = x\}$. Remark that every annihilator set $\text{ann}_M(r)$ is a K_0 -vector space. Until the end of this section, the symbol “ \simeq ” denotes isomorphism between K_0 -vector spaces. The following lemmas establish a generalization

³By definition a lower triangular matrix A (possibly not square) is such that the coefficient a_{ij} is 0 whenever $j > i$, and its diagonal is the sequence (a_{ii}) .

of lemmas 2.9 and 2.10 from [6] where it is assumed that φ is onto, which means R is both right and left euclidean.

Lemma 2.18. *Let M be a divisible R -module. For all $0 \neq q = q_1 \dots q_k \in R$,*

$$|ann_M(q)| = \prod_{i=1}^k |ann_M(q_i)|,$$

where $|ann_M(q_i)|$ is the cardinality of $ann_M(q_i)$.

Proof. If $q = sr$ then multiplication by s induces an one to one K_0 -linear transformation

$$ann_M(q)/ann_M(s) \longrightarrow ann_M(r)$$

which is in fact also onto by divisibility of M . The result clearly follows by induction on k . \square

Lemma 2.19. *Let T_0 be a complete theory of non zero divisible R -modules. For all $s, q \in R \setminus \{0\}$ such that $T_0 \models \forall x (x.s = 0 \rightarrow x.q = 0)$, there exists $r \in R$ such that*

$$(ann_N(q)/ann_N(s)) \simeq ann_N(r)$$

for all $N \models T_0$.

Proof. Let s and q be as in the statement of the lemma. If s divides q then the result follows by the proof of the above lemma. Also if $ann_N(s) = 0$ the assertion is trivial. Hence we assume that $ann_N(s) \neq 0$ for a model (hence for all models) N of T_0 . If s is irreducible, then s is (up to a unit) the minimal polynomial of a non zero root of q hence divides q . Now, consider the general case. We proceed by induction: Suppose that for all non zero polynomials h, g such that $ann_N(h) \supseteq ann_N(g) \neq 0$ and g can be written as a product of $n - 1$ irreducible polynomials, there exist h', g', r' such that $g' = h'r'$, and

$$ann_N(g)/ann_N(h) \simeq ann_N(g')/ann_N(h') \simeq ann_N(r')$$

for all $N \models T_0$. Consider the case $s = s_1 \dots s_n$ and set $s' = s_2 \dots s_n$. Since $ann_N(s) \neq 0$, s_1 or s' has a non-zero root in N .

Case 1 : $ann_N(s_1) \neq 0$. First, s_1 divides q since it is irreducible and hence is -up to a unit- the minimal polynomial of any non zero $x \in ann_N(s_1) \subseteq ann_N(q)$. Thus we have $q = s_1 r$ and, by the fact that N is divisible, $ann_N(s') \subseteq ann_N(r)$ for some $r \in R$. If $ann_N(s') = 0$ then we have: $ann_N(q)/ann_N(s) \simeq ann_N(q)/ann_N(s_1) \simeq ann_N(r)$. Otherwise, by induction hypothesis there are s'_t and r' such that s'_t divides r' and

$$ann_N(r')/ann_N(s'_t) \simeq ann_N(r)/ann_N(s').$$

Now the map “ $x \mapsto x.s_1 \bmod ann_N(s')$ ” establishes a morphism of K_0 -vector spaces from $ann_N(q)$ onto $ann_N(r)/ann_N(s')$ with kernel $ann_N(s)$.

Case 2 : $ann_N(s_1) = 0$. Then, $ann_N(s') \neq 0$,

$$ann_N(s).s_1 = ann_N(s') \subseteq ann_N(q).s_1$$

and the action of s_1 induces an isomorphism K_0 -vector spaces:

$$ann_N(q) \rightarrow ann_N(q).s_1 \quad .$$

Therefore

$$ann_N(q)/ann_N(s) \simeq ann_N(q).s_1/ann_N(s').$$

By lemma 2.16 the p.p. formula $\exists y \ y.s_1 = x \wedge y.q = 0$, defining $ann_N(q).s_1$, is equivalent to a formula of the form: $x.q_1 = 0$ for some $q_1 \in R$. By applying the induction hypothesis to (s', q_1) we get r as required.

It is clear from the proof that r does only depend on T_0 , s and q . \square

Proof of Theorem 1.2. The theorem follows from 2.15, 2.17 and 2.19. \square

We get by quantifier elimination the following consequence of Theorem 1.2.

Corollary 2.20. *Suppose $\eta_M = \eta_N$. Let f be a partial isomorphism between submodules $A \subseteq M$ and $B \subseteq N$ where N is $|M|^+$ -saturated. Then the restriction of f to $A \cap M_{\text{tor}}$ admits an extension to an embedding of M_{tor} in N_{tor} .*

In the rest of this section we will describe the *building blocks* of divisible R -modules and improve Theorem 1.2.

Notation 2.21. Let T_d denote the theory of divisible R -modules.

Definition 2.22. *A module M is said to be indecomposable if there are no submodules N_1, N_2 , both non zeros, such that $M = N_1 \oplus N_2$.*

Lemma 2.23. *Let N_1 and N_2 be indecomposable models of T_d and $q \in R$ an irreducible polynomial such that $\text{ann}_{N_i}(q)$ is non-trivial for $i = 1, 2$. Then N_1 and N_2 are isomorphic.*

Proof. Let $a \in \text{ann}_{N_1}(q)$ and $b \in \text{ann}_{N_2}(q)$. Then N_1 is a divisible closure of $a.R$ and N_2 is a divisible closure of $b.R$. So it suffices to show that $a.R$ is isomorphic to $b.R$. In fact, the application $a.r \mapsto b.r$ is an isomorphism of R -module since for all $r \in R \setminus \{0\}$, $a.r = 0$ if and only if q divides r , if and only if $b.r = 0$. \square

Lemma 2.24. *Let N be an indecomposable model of T_d , $r \in R$ a irreducible polynomial such that $\text{ann}_N(r) \neq 0$. Then, for all irreducible $q \in R$ we have: $\text{ann}_N(q) \neq 0$ if and only if, there exist $\lambda, \mu \in K \setminus \{0\}$ such that $q = \lambda r \mu$; in which case $\text{ann}_N(r) \simeq \text{ann}_N(q)$.*

Proof. Let $0 \neq a \in \text{ann}_N(r)$ and $0 \neq b \in \text{ann}_N(q)$. Then N is a divisible closure of both $a.R$ and $b.R$. Let r_0 be of minimal degree such that $0 \neq a.r_0 \in b.R$. So r_0 divides r . Since r is irreducible, $a.r_0 \neq 0$ implies $r_0 \in K$. Therefore $a \in b.R$, i.e. $a = bs$ for some $s \in R \setminus \{0\}$; take such an s of minimal degree. Then s divides q . Since q is irreducible and $bs \neq 0$, $s \in K$. Moreover, the fact that $b.sr = 0$ implies q divides sr . Since $s \in K$, sr is also irreducible. Therefore $\deg(q) = \deg(r)$ and $sr = q\nu$ with $s, \nu \in K$.

Conversely, suppose λ, μ are as in the statement. Then the map $x \mapsto x.\lambda$ establishes an isomorphism of K_0 -vector space between $\text{ann}_N(q)$ and $\text{ann}_N(r)$. \square

Definition 2.25. *Let $r, q \in R$ be irreducible polynomials. We say that r and q are K -conjugate if there exist $\lambda, \mu \in K \setminus \{0\}$ such that $q = \lambda r \mu$.*

Remark 2.26. *K -conjugation is an equivalence relation.*

Notation 2.27. For the rest of this section we fix a set \mathcal{P} of representatives of irreducible polynomials modulo K -conjugation. Note that the map by 2.18, the map η_M is entirely determined by its restriction on \mathcal{P} for any divisible R -module M .

Lemma 2.28. *Let $r \in \mathcal{P}$ and N be an indecomposable model of T_d containing a non zero root of r . Then for all $q, s \in R \setminus \{0\}$ such that $\text{ann}_N(q) \supseteq \text{ann}_N(s)$, if $|\text{ann}_N(q)/\text{ann}_N(s)|$ is finite, then it is equal to $|\text{ann}_N(r)|^k$, for some $0 \neq k \in \mathbb{N}$.*

Proof. By Lemma 2.19 there exists $r' \in R$ such that

$$\text{ann}_N(q)/\text{ann}_N(s) \simeq \text{ann}_N(r').$$

Now, $r' = r_0 \dots r_n$ with the r_i irreducible. We have on one hand

$$|\text{ann}_N(r')| = \prod_{i \leq n} |\text{ann}_N(r_i)|$$

by 2.18, and on the other hand, each $|\text{ann}_N(r_i)| \in \{|\text{ann}_N(r)|, 1\}$ by Lemma 2.24. \square

Proposition 2.29. *Any divisible R -module is the direct sum of indecomposable divisible submodules.*

Proof. Any injective module over a noetherian ring is the direct sum of indecomposable submodules (see [8] corollary 7.3) and every direct summand of a divisible R -module is divisible. \square

Notation 2.30. Let $M \models T_d$ and $q \in R$ be an irreducible polynomial. We denote by M_q the sum of all divisible indecomposable submodules of M_{tor} containing at least one non zero root of q .

Proposition 2.31. *One has $M_{\text{tor}} = \bigoplus_{q \in \mathcal{P}} M_q$.*

Proof. It is enough to see that if $N \subseteq M_{\text{tor}}$ is a divisible indecomposable submodule of M_{tor} then N is a divisible closure of some non zero element $a \in M$ annihilated by an irreducible polynomial q . First, since N is indecomposable, it is a divisible closure of any of its non zero elements. Then, if q_x is the minimal polynomial of a non zero $x \in N$, q_x can be written as $q_1 \dots q_k$, where the q_i are irreducible. Hence $a := x \cdot (q_1 \dots q_{k-1})$ is annihilated by q_k . \square

Hence we obtain the following improvement of theorem 1.2.

Corollary 2.32. *Completions of $T_d \cup \{\exists x \ x \neq 0\}$ are obtained by specifying for each $q \in \mathcal{P}$, $|\text{ann}(q)| \in \{|K_0|^n \mid n \in \mathbb{N}\} \cup \{\infty\}$, and they admit elimination of quantifiers.*

2.2. The case of $K[t; \text{Frob}]$ -modules. Let K be a field of characteristic $p > 0$ and Frob be the Frobenius endomorphism of K . We say that K is p -closed if every polynomial of the form $X^{p^n} + X^{p^{n-1}}a_{n-1} + \dots + X^p a_1 + X a_0 + b \in K[X]$ has a root in K . This amounts to say that K is divisible as a $K[t; \text{Frob}]$ -module. We have also the following result which was first shown by Whaples (cf. [19]) and then a more elementary proof was provided by Delon in her thesis (cf. [4]).

Proposition 2.33. *A field of characteristic $p > 0$ is p -closed if and only if it has no extension of degree divisible by p .*

Corollary 2.34. *If L is a field of characteristic $p > 0$, p -closed, then every algebraic extension of L is p -closed.*

Theorem 2.35. *Let $K \subseteq F$ be an extension of fields of characteristic $p > 0$. Then the torsion submodule F_{tor} of the $K[t; \text{Frob}]$ -module F is equal to the algebraic closure of K in F . In addition, two p -closed algebraic extensions F_1 and F_2 of K are elementary equivalent as $K[t; \text{Frob}]$ -modules if and only if F_1 and F_2 are isomorphic as $K[t; \text{Frob}]$ -modules.*

Proof. Any algebraic element over K is a zero of an additive polynomial by the fact that any polynomial divides an additive polynomial (see [12]). This shows the first assertion.

Since F_1 and F_2 are p -closed, by Corollary 2.17, their theories as $K[t; \text{Frob}]$ -modules are given by the theories of their torsion submodules. Since F_1 and F_2 are algebraic over K , $F_1 = F_{1, \text{tor}}$ and $F_2 = F_{2, \text{tor}}$. By Proposition 2.14, they are isomorphic. \square

Corollary 2.36. *Consider a field K of characteristic $p > 0$, p -closed and the following tower of extensions:*

$$K \subseteq L \subsetneq M \subseteq K^{\text{alg}}.$$

Then $L \neq M$ as $K[t; \text{Frob}]$ -modules.

3. VALUED MODULES

We are going to generalize the following situation:

Let (U, v) be a valued field of equal characteristic $p > 0$ with $K \subseteq U$ a subfield, on which the valuation v is trivial. We consider the valued group $(U, +, v)$ together with the right action of the ring $R = K[t; \text{Frob}]$ of additive polynomials over K . Consider $\tau : v(U) \rightarrow v(U)$, $\gamma \mapsto p\gamma$. The valued module structure of (U, v) is the two-sorted structure

$$\langle K, +, (\cdot)_{r \in R}, v(U), \leq, \tau, v, \infty \rangle$$

where $\langle U, +, (\cdot)_{r \in R} \rangle$ is the (right)- R -module structure on $(U, +)$ and $\langle v(U), \leq, \tau, \infty \rangle$ is the ordered value set structure of $v(U)$ equipped with τ .

Since v is trivial on K , for all $x \in U$ and $a \in U$, we have $v(x^{p^m}a) = v(x^{p^m}) = p^m v(x)$ and if $v(x) \neq 0$ then we have $v(x^{p^m}) \neq v(x^{p^n})$ whenever $n \neq m$. Hence by ultrametric inequality, for an additive polynomial $P(X) := \sum_{i \in I} X^{p^i} a_i$ ($a_i \neq 0$) over K , for all x with $v(x) \neq 0$ we have

$$v(P(x)) = \min_{i \in I} \{p^i v(x)\} = \begin{cases} \tau^{\min I}(v(x)) & \text{if } v(x) > 0 \\ \tau^{\max I}(v(x)) & \text{if } v(x) < 0 \end{cases}.$$

3.1. τ -chains. Let L_{V_0} be the language $\{\leq, \tau, \infty\}$.

Definition 3.1. A τ -chain is a L_{V_0} -structure Δ satisfying the following axioms:

1. Δ is linearly ordered by \leq with ∞ being its maximum,
2. τ is strictly increasing on Δ and $\tau(\infty) = \infty$,
3. $\forall \gamma \neq \infty \forall \delta \ (\tau(\gamma) \leq \gamma \wedge \delta < \gamma) \rightarrow \tau(\delta) < \delta$.

Example 3.2. For all integer $n \geq 1$ the structure $\langle \mathbb{Z} \cup \{\infty\}, \leq, x \mapsto nx \rangle$ satisfies the above axioms.

Remark that axiom 3 imply its dual:

$$(2) \quad \forall \gamma \forall \delta \neq \infty \ (\tau(\gamma) \geq \gamma \wedge \gamma < \delta) \rightarrow \tau(\delta) > \delta.$$

We denote:

- $\Delta_+ := \{\gamma \in \Delta \mid \tau(\gamma) > \gamma\} \cup \{\infty\}$
- $\Delta_- := \{\gamma \in \Delta \mid \tau(\gamma) < \gamma\}$

Thus, Δ_+ is a final segment and Δ_- is an initial segment of Δ . In addition, by Axiom 3 there exists at most one fixed point of τ other than ∞ : if it exists it is the unique point θ of Δ such that $\Delta_- < \theta < \Delta_+$. We define:

- the predicate $\delta < \theta$, saying $\delta \in \Delta_-$ and,
- the predicate $\delta > \theta$, saying $\delta \in \Delta_+$.

Similarly the expressions $\delta \leq \theta$ or $\delta \geq \theta$, does not mean that the cut (Δ_-, Δ_+) is realized in Δ .

3.2. (K -trivially) valued modules. Recall that a valued abelian group is a structure given by

- an abelian group M ,
- a linear order Δ with a maximum element ∞ ,
- a surjective map $v : M \rightarrow \Delta$ such that
 - for all $x \in M$, $v(x) = \infty$ if and only if $x = 0$
 - for all $x, y \in M$, $v(x \pm y) \geq \min\{v(x), v(y)\}$.

These axioms imply that for all $x, y \in M$, $v(x \pm y) = \min\{v(x), v(y)\}$, whenever $v(x) \neq v(y)$, and $v(x) = v(-x)$.

In a valued group (G, v) , the valuation v induces a topology, a basis of which is given by the *open balls*: these are subsets of G of the form $\{x \in G \mid v(x - a) > \gamma\}$, where $a \in G$ (center) and $\gamma \in v(G)$ (radius). We define *closed balls* as usual by changing $>$ to \geq .

Notation 3.3. In (G, v) , for all $\gamma \in v(G)$ we denote by $G_{\geq \gamma}$ (respectively $G_{> \gamma}$) the closed ball (respectively the open ball) centered at 0 with radius γ .

For the rest of this article we fix a field K and we let $R := K[t; \varphi]$ with $\varphi \in \text{End}(K, +, \times, 1, 0)$.

Definition 3.4. Let (M, v) be a valued abelian group with $v : M \rightarrow \Delta$. A K -trivially valued module structure on (M, v) is given by a right R -module structure on M such that

1. the function $x \mapsto x.t$ is injective,
2. Δ is a τ -chain,
3. $\forall x \in M, v(x.\lambda) = v(x)$ for all $\lambda \in K^\times$
4. $\forall x \in M, v(x.t) = \tau(v(x))$.

The name K -trivial comes from Axiom 3. In [11] we consider also different (“non trivial”) actions of K but in the present paper we will only deal with K -trivially valued modules. Therefore we permit our self to omit the expression K -trivial and say only **valued module** until the end of this paper.

Remark 3.5. When we deal with a valued module (M, v) , we write $M_{> \theta}$ or $M_{\geq \theta}$ independently of the existence of θ .

Remark 3.6. One has $v(x) > \theta$ if and only if $v(x.t) > v(x)$, and $v(x) < \theta$ if and only if $v(x.t) < v(x)$.

Definition 3.7. A polynomial $r \in R$ such that t does not divide r is called *separable*.

Remark 3.8. Any irreducible $r \neq t$ is separable. An R -module is *divisible* if and only if it is divisible by separable polynomials and by t , in which case we will say *t -divisible* for short.

Notation 3.9. Using Equation 1 we observe that any non zero polynomial $q \in R$ can be written in the form $q = t^n s$ where s separable. We set

$$\deg_{is}(q) := n.$$

Lemma 3.10. Let (M, v) be a valued module and $r = \sum_{i \in I} t^i a_i \in R \setminus \{0\}$. Then, for all $x \in M$, we have $v(x.r) \geq \min_{i \in I} \{v(x.t^i a_i)\} = \min_{i \in I} \{\tau^i(v(x))\}$. In addition for all $x \in M$,

1. if $v(x) > \theta$ then $v(x.r) = \tau^{\deg_{is}(r)}(v(x)) = \min_{i \in I} \{\tau^i(v(x))\}$,
2. if $v(x) < \theta$ then $v(x.r) = \tau^{\deg(r)}(v(x)) = \min_{i \in I} \{\tau^i(v(x))\}$,
3. if $v(x.r) > \min_{i \in I} \{\tau^i(v(x))\}$ and $x \neq 0$ then $v(x) = \theta$.

Proof. Let $x \in M \setminus \{0\}$. By the valued abelian group structure of (M, v) we have: $v(\sum x.t^i a_i) \geq v(x.t^i a_i)$ for all $i \in I$. Then we have $v(x.t^i a_i) = v(x.t^i) = \tau^i(v(x))$ by Axioms 3 and 4 of Definition 3.4.

1. If $v(x) > \theta$ then $v(x.t) > v(x)$. Now,

$$v(x.t^i a_i) = v(x.t^i) > v(x.t^j) = v(x.t^j a_j)$$

whenever $i > j$. Hence, $v(x.r) = v(\sum x.t^i a_i) = \min_i \{v(x.t^i)\} = \tau^k(v(x))$, where $k = \deg_{is}(r)$.

2. In this case we have $v(x.t) < v(x)$ and hence $v(x.t^i a_i) < v(x.t^j a_j)$ whenever $i > j$. Thus, $v(x.r) = v(\sum x.t^i a_i) = \min_i \{v(x.t^i)\} = \tau^n(v(x))$ where $n = \deg(r)$.
 3. By points 1. and 2. above, x can only have valuation θ . \square

Corollary 3.11. *The subsets $M_{>\theta}$ and $M_{\geq\theta}$ are R -submodules of M .*

Definition 3.12. *Let (G, v) be an abelian valued group. A subgroup of G is called convex if it is the inverse image under v of a non empty final segment of $v(G)$.*

Lemma 3.13. *A (closed or open) ball centered at 0 is a convex subgroup. If (M, v) is a valued module then $M_{\geq\theta}$ and $M_{>\theta}$ are convex subgroups of M .*

Proof. A ball centered at 0 with radius γ is the inverse image under v of the final segment $(\gamma, \infty]$ or $[\gamma, \infty]$. Hence $M_{\geq\theta}$ and $M_{>\theta}$ are convex subgroups if $\theta \in v(M)$. Otherwise, 3.5, $M_{\geq\theta} = M_{>\theta} = v^{-1}(\Delta_+)$. \square

Lemma 3.14. *If H is a convex subgroup of an abelian valued group (G, v) then the quotient G/H is valued by v_H , defined by:*

$$v_H(g + H) := \begin{cases} v(g) & \text{if } g \notin H \\ \infty & \text{otherwise} \end{cases}.$$

Proof. It is sufficient to verify that v_H is well defined. Let I_H be the final segment of $v(G)$ such that $H = v^{-1}(I_H)$. If $g \notin H$ and $g' \notin H$ are two representatives of the same class, then $v(g) \notin I_H$, $v(g') \notin I_H$ and $g - g' \in H$. Thus $v(g - g') \in I_H$ and $v(g - g') > v(g)$ necessarily. It follows that $v(g) = v(g')$. \square

Corollary 3.15. *Let (M, v) be a valued module. Then $M/M_{\geq\theta}$ is canonically equipped with a valued module structure.*

Definition 3.16. *A valued module (M, v) will be called regular if, for all $x \in M$ and non zero $r = \sum_{i=0}^n t^i a_i \in R$, we have*

$$v(x.r) = \min_i \{v(x.t^i)\} = \min_i \{\tau^i(v(x))\}.$$

If the above equality holds for a pair (x, r) we say that x is regular for r (otherwise irregular for r) and x is said to be regular if it is regular for all r (otherwise irregular).

Remark 3.17. *By 3.10 x is irregular if and only if $v(x) = \theta$ and $v(x.r) \in M_{>\theta}$ for some non zero r .*

Remark 3.18. *Regular modules are necessarily torsion free.*

The following lemma follows directly by Corollary 3.15 and by Lemma 3.10.

Lemma 3.19. *Let (M, v) be a valued module. The submodule $(M_{>\theta}, v)$ and the quotient $(M/M_{\geq\theta}, v_{M_{\geq\theta}})$ are regular valued modules.*

Definition 3.20. *We say that a sequence of submodules $(A_i)_{i \in I}$ of M is valuation independent if, for any sequence $(x_i)_{i \in I}$ with $x_i \in A_i$ and for all finite J with $J \subseteq I$, we have*

$$v(\sum_{i \in J} x_i) = \min\{v(x_i) \mid i \in J\}.$$

Remark 3.21. *If $(A_i)_{i \in I}$ are torsion free and valuation independent then any sequence $(x_i)_{i \in I}$ with $x_i \in A_i$, is R -linearly independent.*

Fact 3.22. *Let (M, v) be a valued module, $A \subseteq M$ be a regular submodule and B a submodule of M_{tor} . Then A and B are valuation independent.*

Proof. Let $x \in M_{\text{tor}}$, and $a \in A$. Without loss of generality we may assume both x and a are non zero. If $v(a - x) > v(x) = v(a)$ then $v(a) = \theta$ and since a is regular $v(a.r) = \theta$. But now, for some non zero r such that $x.r = 0$ we have $\theta = v(a.r) < v((a - x).r)$, a contradiction. \square

3.3. Henselian valued modules.

Definition 3.23. A valued module is said to be *henselian* if it satisfies the following axiom scheme:

$$H : \forall x \ v(x) > \theta \rightarrow \exists y \ x = y.r \wedge v(y) > \theta$$

for all separable $r \in R$.

We denote by T_h the theory of henselian valued modules.

Lemma 3.24. Let $(K \subseteq U, v)$ be an extension of valued fields of characteristic $p > 0$, where v is trivial on K . Then (U, v) canonically inherits a valued $K[t; \text{Frob}]$ -module structure. In addition, if v is henselian on the field U , then (U, v) is a henselian valued module. Moreover, if U is perfect, then the maximal ideal \mathcal{M}_U associated to v is a divisible $K[t; \text{Frob}]$ -module.

Proof. As usual we interpret $x.t$ as x^p , τ as the map $\tau : v(U) \rightarrow v(U)$, $\gamma \mapsto p\gamma$, and θ as $0 \in v(U)$ which makes (U, v) a valued module. Suppose now that (U, v) is henselian as a valued field. Let $q = t^n a_n + \dots + a_0$ be separable and $y \in U$ such that $v(y) > 0$. Let Q be the additive polynomial associated to q : i.e. $Q(X) = a_n X^{p^n} + \dots + a_0 X$. Since $a_n, \dots, a_0 \in K$, they have common valuation 0, and since q is separable, $a_0 \neq 0$. Then, by setting $F(X) := Q(X) - y$, we have $F'(0) = a_0$ and $F(0) = -y$. Since v is henselian, there exists $z \in U$ of valuation > 0 such that $F(z) = 0$. Thus $Q(z) = y$ and $v(z) > 0$. In other words, in the language of valued modules, we have $z.q = y$ and $v(z) > \theta$.

Finally, remark that $U_{>\theta} = \mathcal{M}_U$ and if U is perfect, then $U_{>\theta}$ is divisible by t . Since by the above paragraph it is also divisible by separable polynomials, it is divisible. \square

Lemma 3.25. Any torsion free henselian valued module is regular.

Proof. Suppose for a contradiction that we have a henselian torsion free module (M, v) , with an element $x \in M$ irregular. Then for some non zero $r \in R$, $v(x.r) > \theta$ and $v(x) = \theta$. Write $r = t^n s$ with s separable. By henselianity there exists $y \in M$, of valuation $> \theta$ such that $y.s = x.r$. Then $y - x.t^n$ is non zero but annihilated by s . Contradiction. \square

It is a trivial fact that algebraically closed valued fields are henselian. We could look to the notion of divisibility for valued modules as an analogue of the notion of algebraic closeness for valued fields. But there exist divisible valued modules which are not henselian as it is showed in the following proposition.

Proposition 3.26. There exist non henselian divisible valued modules.

Proof. Let U be an algebraically closed field of characteristic $p > 0$, v a non trivial valuation on U and K a trivially valued subfield. We consider U with its $K[t; \text{Frob}]$ valued module structure.

Let $y \in U$, of valuation > 0 . Then, by 3.10, y is not a torsion element. Since 1 is annihilated by $(t - 1)$, it is a torsion element (indeed $1.t = 1^p = 1$). Consider, the submodule $A := (1 + y).R$. It is torsion free. Set $x = (1 + y).(t - 1) \in A$. Since $1.(t - 1) = 0$, we have also $x = y.(t - 1)$. On the other hand, since U is algebraically closed, it is divisible as an R -module. Hence U contains a divisible closure of A . Let the module B be such a closure. By Lemma 2.6 (point 4) B is a torsion free

module. Thus $1 \notin B$, hence $y \notin B$: B can not be henselian since y is the unique element of U of valuation > 0 such that $y.(t-1) = x$. \square

Corollary 3.27. *Take x as above. Then $x.R$ has two divisible closures non elementary equivalent as valued modules.*

Proof. Since $v(x) > \theta$ and $U_{>\theta}$ is divisible, $x.R$ has a divisible closure inside $U_{>\theta}$, which is necessarily henselian. This can not be elementary equivalent to B , since B is not henselian. \square

3.4. Henselian divisible valued modules. The results of this subsection contain the essential information that will be used to establish an Ax-Kochen and Ershov principle.

Lemma 3.28. *If (M, v) is a henselian divisible valued module, $M_{\geq \theta}$ is divisible and $M_{>\theta}$ is divisible torsion free. Hence $M_{\geq \theta}$ is a direct summand in M , and $M_{>\theta}$ is a direct summand in $M_{\geq \theta}$.*

Proof. Let $r \in R \setminus \{0\}$, $x \in M_{\geq \theta}$ and $y \in M$ such that $y.r = x$. Then by 3.10 $y \in M_{\geq \theta}$. Thus $M_{\geq \theta}$ is a divisible submodule of M .

Now we show that $M_{>\theta}$ is divisible. Let x be of valuation $> \theta$ and $r \in R$, non zero. Write $r = t^n.r'$ with r' separable, then there exists, by axiom H , $y \in M_{>\theta}$ such that $x = y.r'$. By t -divisibility of M , we have $y = y'.t^n$ for some $y' \in M_{>\theta}$, hence $y'.r = x$. Moreover $M_{>\theta}$ is torsion free since it is a regular submodule. \square

Corollary 3.29. *Let (M, v) be a henselian divisible valued module. If $x \in M$ is irregular then there exists a unique couple $(x_{tor}, x_{>\theta})$, with $x_{tor} \in M_{tor}$ and $x_{>\theta} \in M_{>\theta}$ such that*

$$x = x_{tor} + x_{>\theta}.$$

As a consequence,

- (1) *for all $x \in M$ and $q = \sum_{i \in I} t^i a_i \in R \setminus \{0\}$, there exists regular $y \in M$ such that $x = y.q$ and hence $v(x) = \tau^k(v(y))$, where $k = \deg_{is}(q)$ if $v(x) > \theta$, $k = \deg(q)$ if $v(x) < \theta$, and $k = 0$ if $v(x) = \theta$; in any case*

$$v(x) = \min_{i \in I} \{v(y.t^i a_i)\} = \min_{i \in I} \{\tau^i(v(y))\},$$

- (2) *$0 \neq x \in M$ is regular for r (resp. regular) if and only if for all $a \in \text{ann}_M(r)$ (resp. for all $a \in M_{tor}$), $v(x-a) = \min\{v(x), \theta\}$.*

Proof. Take an irregular $x \in M$. Let $r \in R \setminus \{0\}$ be such that $x.r \in M_{>\theta}$ (given by 3.17). Since $M_{>\theta}$ is divisible, $x.r = y.r$ for some $y \in M_{>\theta}$ and y is necessarily regular. Since $M_{tor} \cap M_{>0} = 0$ the couple $(x-y, y)$ is unique as required. Then it follows from 3.10 that $v(y.r) = \tau^{\deg_{is}(r)}(v(y))$. The first consequence mentioned follows then by 3.10 and the second from the proof of the first one. \square

Theorem 3.30. *Let (M, v) be a henselian divisible valued module.*

1. *M_{tor} embeds in $M_{\geq \theta}/M_{>\theta}$ and for any such embedding the image of M_{tor} is a direct summand.*
2. *The R -modules $M_{\geq \theta}/(M_{tor} + M_{>\theta})$, $M_{>\theta}$ and $M/M_{\geq \theta}$ are torsion free and divisible.*
3. *M can be written as a direct sum:*

$$M_{tor} \oplus M_{\theta} \oplus M_{>\theta} \oplus M_{-},$$

where M_{θ} is isomorphic to $M_{\geq \theta}/(M_{tor} + M_{>\theta})$ and (M_{-}, v) is isomorphic as a valued module to $(M/M_{\geq \theta}, v_{M_{\geq \theta}})$. This decomposition is valuation independent and each member of this decomposition, except M_{tor} , is a regular D -vector space.

Proof. 1. Since all non-zero elements of M_{tor} are of valuation θ , the canonical surjection $(M_{\geq \theta} \rightarrow M_{\geq \theta}/M_{> \theta})$ induces an embedding of M_{tor} in $M_{\geq \theta}/M_{> \theta}$. Since M_{tor} is divisible its image in $M_{\geq \theta}/M_{> \theta}$ is a direct summand.

2. The fact that $M_{> \theta}$ is torsion free and divisible is given by 3.28. The divisibility of $M/M_{\geq \theta}$ is induced by the divisibility of $M_{\geq \theta}$, it is torsion free by 3.19 since it is a regular valued module. Also $M_{\geq \theta}/(M_{tor} + M_{> \theta})$ is divisible by divisibility of M . We will now show that $M_{\geq \theta}/(M_{tor} + M_{> \theta})$ is torsion free. Let $x \in M_{\geq \theta}$ such that $x.r \in M_{tor} + M_{> \theta}$ for some $r \neq 0$. Since M_{tor} is divisible, we have $x.r - z'.r = (x - z').r \in M_{> \theta}$ for some $z' \in M_{tor}$. If $x - z' \notin M_{> \theta}$ then $x - z'$ is irregular. In this case, by Corollary 3.29, $x - z' \in M_{tor} + M_{> \theta}$ and hence $x \in M_{tor} + M_{> \theta}$.

3. Since $M_{\geq \theta}$ and $M_{> \theta}$ are both divisible, $M \simeq M_{\geq \theta} \oplus M/M_{\geq \theta}$ and

$$M_{\geq \theta} \simeq M_{> \theta} \oplus M_{\geq \theta}/M_{> \theta}$$

and we get $M_{\geq \theta}/M_{> \theta} \simeq M_{tor} \oplus (M_{\geq \theta}/(M_{tor} + M_{> \theta}))$.

Take a direct summand M_- of $M_{\geq \theta}$ in M . If $x \in M_- \setminus \{0\}$ then $v(x) < \theta$ and by the definition of the quotient valuation $v_{M_{\geq \theta}}, (M_-, v)$ and $(M/M_{\geq \theta}, v_{M_{\geq \theta}})$ are isomorphic as valued modules.

It remains to show that this decomposition is valuation independent. For this it is enough to see that M_{tor} and M_{θ} are valuation independent. Suppose that there exist $x \in M_{tor}$, $y \in M_{\geq \theta} \setminus (M_{tor} + M_{> \theta})$ with $v(x - y) > \theta$. Then, for some $r \in R \setminus \{0\}$ annihilating x , we have $v((x - y).r) = v(y.r) > \theta$: this is impossible since $M_{\geq \theta}/(M_{tor} + M_{> \theta})$ is torsion free. \square

Corollary 3.31. *For all $i \in \{\theta, > \theta, -\}$, for all $x, y \in M_i \setminus \{0\}$ the following are equivalent:*

- $v(x - y) > v(x) = v(y)$,
- there is some $r \in R \setminus \{0\}$ such that $v(x.r - y.r) > v(x.r) = v(y.r)$,
- for all $r \in R \setminus \{0\}$ we have $v(x.r - y.r) > v(x.r) = v(y.r)$.

Proof. Follows by the fact that each M_i is regular. \square

Remark 3.32. *Let $(K \subseteq U, v)$ be an extension of fields of characteristic $p > 0$, where U is algebraically closed and $|K|^+$ -saturated and v is trivial on K . Then every non zero element of the set U_{θ} realizes the generic type of the valuation ring of U , i.e. the type $\{x \in \mathcal{O}_U \wedge x \notin B \mid B \subsetneq \mathcal{O}_U \text{ is } K\text{-definable}\}$. Note that here definable means definable in the language $\{0, 1, +, -, \times, \mathcal{O}\}$ of valued fields.*

3.5. Embedding theorems and A-K,E principles. Let (M, v) be a valued module. If A is a submodule of M , Corollary 3.27 shows that the various divisible closures of A in M , while being isomorphic as R -modules, may not be elementary equivalent as valued modules. This is the most important phenomenon to which we will pay attention.

Definition 3.33. *Let (M, v) be a divisible valued module and $A \subseteq M$ a submodule of M . Then we define*

$$\hat{A} := \{y \in M \mid y.r \in A \text{ for some } r \in R \setminus \{0\}\}.$$

Lemma 3.34. *The submodule \hat{A} is the unique divisible closure of $A + M_{tor}$ inside M . In particular, for all $r \in R \setminus \{0\}$ and all $x \in M$, if $x.r \in \hat{A}$ then $x \in \hat{A}$. Moreover, if (M, v) is henselian then \hat{A} is henselian.*

Proof. By construction, \hat{A} is a divisible submodule and all divisible submodules of M containing A and M_{tor} contains \hat{A} . This gives the uniqueness. Also by the definition of \hat{A} , if $x \in \hat{A}$ and $r \in R \setminus \{0\}$, \hat{A} contain all y such that $x = y.r$ since

\hat{A} contains M_{tor} . Now, if (M, v) is henselian and $x \in \hat{A}$ of valuation $> \theta$, then \hat{A} contains the unique element $y_{>\theta} \in M_{>\theta}$ such that $y_{>\theta}.r = x$. This shows that \hat{A} is henselian. \square

Remark 3.35. *The set $v(\hat{A})$ is an L_{V_0} -substructure of $v(M)$, it is the closure of $v(A)$ under the function τ^{-1} .*

Remark 3.36. *Consider a henselian divisible valued module (M, v) . Then,*

- (1) $M_{\geq \theta}/M_{>\theta} = 0$ or infinite,
- (2) if $\gamma \neq \theta$ and r is non zero with $\deg_{is}(r) = k$ and $\deg(r) = n$ then
 - (a) if $\gamma > \theta$ the multiplication by r

$$.r : M_{\geq \gamma} \rightarrow M_{\geq \tau^k(\gamma)}$$

is a bijection inducing a bijection

$$M_{\geq \gamma}/M_{>\gamma} \rightarrow M_{\geq \tau^k(\gamma)}/M_{>\tau^k(\gamma)},$$

- (b) if $\gamma < \theta$ the multiplication by r

$$.r : M_{\geq \gamma} \rightarrow M_{\geq \tau^n(\gamma)}$$

is a bijection inducing a bijection

$$M_{\geq \gamma}/M_{>\gamma} \rightarrow M_{\geq \tau^n(\gamma)}/M_{>\tau^n(\gamma)}.$$

Proof. If $\theta \in v(M)$ then M_θ or M_{tor} is non empty, divisible and embedded into $M_{\geq \theta}$. The second assertion follows from divisibility and regularity of any element of valuation $\neq \theta$. \square

– For each $n \in \mathbb{N}$, we add a unary predicate R_n into our language L_{V_0} and let L_V be this enrichment of L_{V_0} . We denote $L_0 := L_{Mod}(R) \cup L_{V_0} \cup \{v\}$ and $L := L_{Mod}(R) \cup L_V \cup \{v\}$. In every valued module (M, v) and for all $\gamma \in v(M) \setminus \{\infty\}$, $R_n(\gamma)$ will be interpreted by the equivalence

$$R_n(\gamma) \Leftrightarrow |M_{\geq \gamma}/M_{>\gamma}| \geq n.$$

Proposition 3.37. *Let (M, v) and (N, w) be divisible henselian valued modules such that $\eta_M = \eta_N$ and N is $|M|^+$ -saturated. Let $(A, \Delta_1) \subseteq (M, v(M))$ and $(B, \Delta_2) \subseteq (N, v(N))$ be L -substructures of M and N respectively, which are L -isomorphic via $\mathbf{f} = (f, f_v)$. Then there is an L -embedding $\hat{\mathbf{f}} = (\hat{f}, \hat{f}_v)$ of $(\hat{A}, v(\hat{A}) \cup \Delta_1)$ to $(N, v(N))$ extending \mathbf{f} , having range $(\hat{B}, v(\hat{B}) \cup \Delta_2)$.*

Proof. By Remark 3.35 and the fact that τ is strictly increasing, it is easy to see that f_v extends uniquely to an L_{V_0} -embedding

$$\hat{f}_v : v(\hat{A}) \cup \Delta_1 \rightarrow w(\hat{B}) \cup \Delta_2.$$

Now we will extend f to $\hat{f} : \hat{A} \rightarrow N$ such that (\hat{f}, \hat{f}_v) is an L -embedding. Take a decomposition

$$\hat{A} = M_{tor} \oplus \hat{A}_\theta \oplus \hat{A}_- \oplus \hat{A}_{>\theta}$$

as given by Theorem 3.30.

Let $I := \{\theta, >\theta, -\}$ and for $i \in I$ denote by f_i the restriction of f to $A \cap \hat{A}_i$. By the definition of \hat{A} and by the fact that \hat{A}_i is torsion free, if $x \in \hat{A}_i$ then there is $r \in R \setminus \{0\}$ such that $x.r \in A$ and $x.r \neq 0$ whenever $x \neq 0$. Therefore, each \hat{A}_i admits a D -vector space basis consisting of elements of A . Then, f_i extends uniquely to an isomorphism of D -vector spaces $\hat{f}_i : \hat{A}_i \rightarrow \hat{f}_i(\hat{A}_i) \subseteq N$. Now we let $C := \bigoplus_{i \in I} \hat{A}_i$, define $\hat{f}_C : C \rightarrow N$ as

$$x = x_\theta + x_- + x_{>\theta} \mapsto \hat{f}_\theta(x_\theta) + \hat{f}_-(x_-) + \hat{f}_{>\theta}(x_{>\theta})$$

and

$$A_t := \{x \in M_{tor} \mid \exists c \in C, x + c \in A\}.$$

Note that A_t is a submodule of M_{tor} . We will see that the restriction of f to $A \cap M_{tor}$ admits a unique extension h to A_t such that the map

$$x = x_{tor} + x_\theta + x_- + x_{>\theta} \mapsto h(x_{tor}) + \hat{f}_C(x - x_{tor})$$

defined on $A_t \oplus C$ is a monomorphism of R -modules extending f . Let $x \in A_t \setminus A$ and $c \in C$ be such that $x + c \in A$. We have to check that

$$x \mapsto h(x) := f(x + c) - \hat{f}_C(c)$$

is well defined. Let $d \in C$ be such that $x + d \in A$. Note that $d - c \in A$. Now we have

$$\begin{aligned} f(x + d) - \hat{f}_C(d) - (f(x + c) - \hat{f}_C(c)) \\ = f(d - c) - \hat{f}_C(d - c) = f(d - c) - f(d - c) = 0. \end{aligned}$$

By the definition of h , h is a morphism of R -modules. Moreover h is injective: If for some $c \in C$, $x + c \in A$ and $f(x + c) - \hat{f}_C(c) = 0$ then $\hat{f}_C(c) \in B$ hence $c \in A$. It follows that $x \in A$ and $f(x) = 0$ hence $x = 0$.

Set $\hat{B}_i := \hat{f}(\hat{A}_i)$. By construction the \hat{B}_i are valuation independent, in particular, their sum is direct. Define $E := \bigoplus_{i \in I} \hat{B}_i$. We will now check that the range of h is exactly

$$B_t := \{y \in N_{tor} \mid \exists z \in E, y + z \in B\}.$$

Let $y \in B_t$ and $z \in E$ be such that $y + z \in B$. Let r be the minimal polynomial of y . Then $z.r \in B$ since $y.r = 0$ and $(y + z).r \in B$. Let $c' \in C$ be the image of z under \hat{f}_C^{-1} and $x \in A$ be the image of $y + z$ under f^{-1} . Then $x_0 := x - c'$ is annihilated by r and $h(x_0) = y$.

Since $\eta_M = \eta_N$ and N is $|M|^+$ -saturated, by 2.20 we can extend h to an elementary $L_{Mod}(R)$ -embedding $M_{tor} \rightarrow N$, again denoted by h , having its range equal to N_{tor} . Hence we can define an $L_{Mod}(R)$ -embedding $\hat{A} \rightarrow \hat{B}$, denoted by \hat{f} , by setting

$$\hat{f}(a) := h(a_{tor}) + \hat{f}_C(c)$$

where $a = a_{tor} + a_\theta + a_- + a_{>\theta} \in \hat{A}$ with $c = a_\theta + a_- + a_{>\theta}$.

Note now that each \hat{B}_i is regular. In fact, if $\hat{f}_i(x) = y \in \hat{B}_i$ is irregular then, there exists a non zero $y_0 \in N_{tor}$ and $y_{>\theta} \in N_{>\theta}$ such that $y = y_0 + y_{>\theta}$. Let r be the minimal polynomial of y_0 , then $y.r = y_{>\theta}.r \in B$. Since $\hat{B}_{>\theta}$ is torsion free, $y_{>\theta} = y$. Contradiction.

Denote by $\hat{f}_{v,i}$ the restriction of \hat{f}_v on $v(\hat{A}_i)$. Since each \hat{B}_i is regular the $(\hat{f}_i, \hat{f}_{v,i})_i$ are L_0 -embeddings. Moreover since by Fact 3.22 the decomposition $N_{tor} \oplus E$ is valuation independent, we have $w(h(x) + \hat{f}_C(x)) = \min\{w(h(x)), w(\hat{f}_C(x))\}$ for all $x \in \hat{A}$. Hence (\hat{f}, \hat{f}_v) is an L_0 -embedding. Now it is in fact an L -embedding since \mathbf{f} is an L -embedding and, by Remark 3.35, $v(\hat{A})$ is the closure of $v(A)$ by τ^{-1} and by Remark 3.36 each quotient $\hat{A}_{\geq \gamma} / \hat{A}_{> \gamma}$ is entirely determined by a quotient of the form $A_{\geq \tau^k(\gamma)} / A_{> \tau^k(\gamma)}$. \square

Notation 3.38. For $a \in M$, where M is valued by v , and $\gamma \in v(M)$, we denote by $M_\gamma(a)$ the quotient of the closed ball $\{u \in M \mid v(u - a) \geq \gamma\}$ given by the equivalence relation

$$u \sim u' \Leftrightarrow v(u - u') > \gamma.$$

Note that

$$M_\gamma(a) = |M_{\geq \gamma} / M_{> \gamma}|.$$

Proposition 3.39. *Let M and N be as before, A be a submodule of M such that $v(A) = v(M)$ and $\mathbf{f} = (f, f_v) : A \rightarrow N$ an L -embedding of A into N . Suppose that N is $|M|^+$ -saturated. Then we can extend \mathbf{f} to an L -embedding of M into N .*

Proof. Note first that we can extend \mathbf{f} on \hat{A} by Proposition 3.37. So we assume $\hat{A} = A$ with image $B = \hat{B}$. Take $x \in M \setminus A$. It suffices to extend \mathbf{f} to $A \oplus x.R$. In that case, by the above proposition, we can extend \mathbf{f} to the divisible closure (unique by the fact that $A \supseteq M_{\text{tor}}$) of $A \oplus x.R$. Thus, by transfinite induction, \mathbf{f} extends to M .

Set

$$p(Y) := \{\gamma_a = w(Y - b) \mid \gamma_a = f_v(v(x - a)) \text{ and } b = f(a), a \in A\}.$$

We first show that if $y \in N$ realizes $p(Y)$, then we can extend \mathbf{f} to an L -embedding $: A \oplus x.R \rightarrow N$ sending x to y . Take $y \in N$ realizing $p(Y)$. Then for all $r \in R \setminus \{0\}$, $y.r \notin B$. We set $\tilde{f}(a + x.r) = f(a) + y.r$. Since $v(A) = v(M)$ it suffices to see that \tilde{f} is compatible with f_v , i.e. to check that one has $f_v(v(a - x.r)) = w(f(a) - y.r)$, for all $a \in A$ and $r \in R \setminus \{0\}$.

Claim: For $a \in A$ and $r \neq 0$, $x - a$ is regular for r if and only if $y - f(a)$ is regular for r .

Proof of the Claim. If $y - f(a)$ is irregular for some r , then for some $b \in \text{ann}_N(r) \subseteq B$,

$$w(y - f(a) - b) > \theta$$

by 3.10 (Consequence 2). Hence $v(x - a - f^{-1}(b)) > \theta$ by the choice of y . This means $x - a$ is irregular by 3.10. The inverse can be proven in the same way.

Let $a \in A$, $r \in R \setminus \{0\}$. Take a' such that $x.r - a = (x - a').r$ and $x - a'$ regular for r . Hence $f(a) = f(a').r$ and $f(a') - y$ is regular for r by the Claim above. Since f_v commutes with τ we have $f_v(v(a - x.r)) = w(f(a) - y.r)$ as required.†

It remains to prove that $p(Y)$ is realized in N . We will show that it is finitely consistent. Let $\alpha = \{a_1, \dots, a_m\} \subseteq A$. We will find $y \in N$ such that for all $1 \leq i \leq m$, $\gamma_i := f_v(v(x - a_i))$ is equal to $w(y - f(a_i))$. Let \sim be the equivalence relation on α defined by $a_i \sim a_j$ if and only if $\gamma_i = \gamma_j$. We will first observe that, without loss of generality, we can assume that there is only one class under this equivalence relation: In fact, it is enough to consider $\beta \subseteq \alpha$ be the equivalence class of $\gamma := \max\{\gamma_i\}_{i \in \{1, \dots, m\}}$. If $a \in \beta$ and $a' \notin \beta$ then

$$\begin{aligned} f_v(v(x - a')) &= f_v(v(x - a + a - a')) = f_v(v(a - a')) = w(f(a) - f(a')) \\ &= w(f(a) - y + y - f(a')) = w(f(a') - y). \end{aligned}$$

Set $\delta := f_v^{-1}(\gamma)$, with the assumption above, it is the common value of the $v(x - a)$ for $a \in \alpha$. Now, we can choose an element $y \in N$ such that for all $a \in \alpha$

$$w(y - f(a)) = \gamma$$

since $|M_\delta(a)| \leq |N_\gamma(f(a))|$, f_v preserves the predicate R_n and N is $|M|^+$ -saturated. \square

To prove Theorem 1.3 we recall the following general result.

Proposition 3.40. *Let \mathcal{L} be a language containing a constant symbol, \mathcal{T} an \mathcal{L} -theory and Θ a set of formulas of \mathcal{L} closed under boolean combinations and containing all quantifier free formulas. Suppose that for all $M, N \models \mathcal{T}$ where N is $|M|^+$ -saturated and for all substructure $A \subseteq M$, whenever $f : A \rightarrow N$ is an \mathcal{L} -embedding preserving formulas of Θ , there exists an \mathcal{L} -embedding $g : M \rightarrow N$ which extends f and preserves Θ . Then every formula of \mathcal{L} is equivalent to a formula of Θ .*

Proof. If Θ is the set of all quantifier-free formulas this is a well known fact (cf. [10] proposition 4.3.28). The general case reduces to this case after adding to the language the predicates P_ϕ for each $\phi \in \Theta$ and enriching the theory \mathcal{T} by the set of sentences $\{\forall x \ P_\phi(x) \leftrightarrow \phi(x) \mid \phi \in \Theta\}$. \square

Let $(F, v(F))$ be a non zero henselian divisible R -module. Let T_v be the complete theory of $v(F)$ in the language L_V and Tor_F be the $L_{\text{Mod}}(R)$ -theory of non zero divisible R -modules with the extra statements of the form $\eta_F(r) = n$ or sets of statements expressing $|\eta_F(r)| = \infty$ where

$$\eta_F(r) = |\{x \in F \mid x.r = 0\}|.$$

By 1.2 this is the complete theory of (the pure module) F and also of F_{tor} if it is non zero (by 2.4). Similarly to the case of valued fields, Tor_F can be seen as *the residual theory* of (F, v) since F_{tor} embeds in $F_{\geq \theta}/F_{> \theta}$.

We set $\mathbf{T} := T_h \cup T_v \cup \text{Tor}_F$ and Θ_v the set of L -formulas of the form:

$$\varphi(\bar{x}) \wedge Q\bar{y}_1 \ \psi(\bar{y}_1, \bar{y}_2, v(t_1(\bar{x})), \dots, v(t_k(\bar{x})),$$

where:

- T_h is the theory of henselian valued R -modules,
- φ is quantifier-free in $L_{\text{Mod}}(R)$, ψ is quantifier-free in L_V and Q is a sequence of quantifiers over \bar{y}_1 ,
- \bar{x} is a tuple of variables of the module sort, \bar{y}_1, \bar{y}_2 are tuples of variables of the value set sort, and the t_i are $L_{\text{Mod}}(R)$ -terms.

Theorem 1.3 follows now by the following theorem:

Theorem 3.41. *The theory \mathbf{T} is complete and eliminates quantifiers on the module sort: any formula of L is equivalent modulo \mathbf{T} to a formula of Θ_v .*

Proof. Take $(M, v), (N, w) \models \mathbf{T}$ such that N is $|M|^+$ -saturated. Let $\mathbf{f} = (f, f_v) : (A, \Delta) \rightarrow N$ be an L -embedding preserving formulas of Θ_v where (A, Δ) is a substructure of (M, v) . Note that the condition $\theta \in v(M)$ is described by the L_{V_0} -theory of $v(F)$. Furthermore, since f_v preserves in particular all L_V -formulas, f_v is partial elementary and extends to an elementary embedding of $v(M)$ into $v(N)$ by saturation hypothesis. This extension will be noted as f_v as well. Hence we may assume $\Delta = v(M)$ (but v is not necessarily surjective).

On the other hand by Proposition 3.37, there is an L -embedding $\hat{\mathbf{f}} = (\hat{f}, \hat{f}_v)$ extending \mathbf{f} to $(\hat{A}, v(M))$. Note that any element $\gamma \in v(\hat{A}) \setminus \{\theta\}$ is $v(A)$ -definable by a formula of the form $\tau^k(\gamma) = \gamma_a$ for some integer k and some $\gamma_a \in v(A)$. This implies that $\hat{\mathbf{f}}$ preserves the formulas from Θ_v . Thus from now on we can assume that $A = \hat{A}$.

– **Extending f to an (U, v) such that $v(U) = v(M)$.** Let $\gamma \in v(M) \setminus A$, $x \in M$ of valuation γ and $y \in N$ such that $w(y) = g_v(\gamma)$. Since $v(A)$ is closed by τ and τ^{-1} , we have, for all non zero $r \in R$, $v(x.r) \notin v(A)$. It follows that $v(a + x.r) = \min\{v(a), v(x.r)\}$ for all $r \in R$ and $a \in A$. Note that x is regular. In fact, either M is torsion-free and hence M is regular by Lemma 3.25, or, $M_{\text{tor}} \subseteq A$ and $\gamma \neq \theta$. Thus $v(x.r) = \tau^k(v(x))$ for some integer k depending only on γ . Taking into account that f_v is an elementary function, the map g defined by

$$g(a + x.r) := f(a) + y.r \quad (r \in R, a \in A)$$

yields that $\mathbf{g} = (g, f_v)$ is a partial L -isomorphism between $(A \oplus x.R, v)$ and $(B \oplus y.R, w)$ and \mathbf{g} preserves the set Θ_v by the above discussion. By Proposition 3.37, \mathbf{g} extends to $A + x.R$. Hence, by using the same argument, we can extend \mathbf{g} to a model U such that $v(U) = v(M)$.

Now, by Proposition 3.39 we can extend \mathbf{g} to M and hence we get quantifier elimination up to Θ_v by Theorem 3.40.

It follows that any L -sentence is equivalent modulo \mathbf{T} to an L_V -sentence. Since by the definition of \mathbf{T} any two models have L_V -elementary equivalent value sets, they are in fact L -elementary equivalent. This gives the completeness of \mathbf{T} as required. \square

Corollary 3.42. *Theorem 1.3 is a consequence of the completeness of \mathbf{T} . Moreover,*

- (A-K,E \preceq): *Let $(M \subseteq N, v)$ be an extension of non-zero henselian divisible valued modules such that the inclusion $(M_{\text{tor}} \subseteq N_{\text{tor}})$ is $L_{\text{Mod}}(R)$ -elementary and the inclusion $(v(M) \subseteq v(N))$ is L_V -elementary. Then the inclusion $(M \subseteq N, v)$ is L -elementary.*

REFERENCES

- [1] Luc Bélair and Françoise Point. Quantifier elimination in valued Ore modules. *The Journal of Symbolic Logic*, 75(03):1007–1034, 2010.
- [2] Paul Moritz Cohn. *Skew Fields*. Springer, 2003.
- [3] Pilar Dellunde, Françoise Delon, and Françoise Point. The theory of modules of separably closed fields 1. *The Journal of Symbolic Logic*, 67(03):997–1015, 2002.
- [4] Françoise Delon. *Quelques propriétés des corps valués en théorie des modèles*. Thèse d’Etat. Université Paris VII, 1982.
- [5] Isidore Fleischer. Maximality and ultracompleteness in normed modules. *Proceedings of the American Mathematical Society*, 9(1):151–157, 1958.
- [6] Ehud Hrushovski and Françoise Point. On von Neumann regular rings with an automorphism. *Journal of Algebra*, 315(1):76–120, 2007.
- [7] Nathan Jacobson. *Basic Algebra II: Second Edition*. Dover Books on Mathematics. Dover Publications, 2012.
- [8] Tsit-Yuen Lam. *Lectures on modules and rings*, volume 189. Springer Science & Business Media, 2012.
- [9] Fares Maalouf. Espaces vectoriels C-minimaux. *The Journal of Symbolic Logic*, 75(02):741–758, 2010.
- [10] David Marker. *Model Theory: An Introduction*. Springer Science & Business Media, 2002.
- [11] Gönenç Onay. *Modules valués: en vue d’applications à la théorie des corps valués de caractéristique positive*. PhD thesis, Paris 7, 2011.
- [12] Oystein Ore. On a special class of polynomials. *Transactions of the American Mathematical Society*, 35(3):559–584, 1933.
- [13] Mike Prest. Model theory and modules. *Handbook of algebra*, 3:227–253, 2003.
- [14] Thomas Rohwer. *Valued difference fields as modules over twisted polynomial rings*. PhD thesis, University of Illinois at Urbana-Champaign, 2003.
- [15] Gabriel Sabbagh. Sous-modules purs, existentiellement clos et élémentaires. *Comptes Rendus de l’Académie des Sciences de Paris*, 272(A):1289–1292, 1971.
- [16] Guy Terjanian. Un contre-exemple à une conjecture d’Artin. *Comptes Rendus de l’Académie des Sciences de Paris*, 262:612, 1966.
- [17] Lou van den Dries. Classical model theory of fields. In Deirdre Haskell, Anand Pillay, and Charles Steinhorn, editors, *Model Theory, Algebra, and Geometry*, volume 39, pages 37–52. MSRI Publications, 2000.
- [18] Franz viktor Kuhlmann. Additive polynomials and their role in the model theory of valued fields, model theory of valued fields. In *Logic in Tehran, Proceedings of the Workshop and Conference on Logic, Algebra, and Arithmetic, held October 18-22, 2003. Lecture Notes in Logic 26*, pages 160–203, 2006.
- [19] G. Whaples. Galois cohomology of additive polynomial and n -th power mappings of fields. *Duke Mathematical Journal*, 24(2):143–150, 1957.

MATHEMATICS DEPARTMENT, FACULTY OF SCIENCE AND LETTERS MIMAR SINAN FINE ARTS
UNIVERSITY BOMONTI CAMPUS 34380 ISTANBUL
E-mail address: gonenc.onay@msgsu.edu.tr