

THE FRÖLICHER-NIJENHUIS BRACKET AND THE GEOMETRY OF G_2 -AND $\text{Spin}(7)$ -MANIFOLDS

KOTARO KAWAI, HÔNG VÂN LÊ, AND LORENZ SCHWACHHÖFER

ABSTRACT. We extend the characterization of the integrability of an almost complex structure J on differentiable manifolds via the vanishing of the Frölicher-Nijenhuis bracket $[J, J]^{FN}$ to an analogous characterization of torsion-free G_2 -structures and torsion-free $\text{Spin}(7)$ -structures. We also explain the Fernández-Gray classification of G_2 -structures and the Fernández classification of $\text{Spin}(7)$ -structures in terms of the Frölicher-Nijenhuis bracket.

1. INTRODUCTION

A G_2 -structure on a 7-dimensional manifold M^7 is a 3-form $\varphi \in \Omega^3(M^7)$ which at each point $p \in M^7$ is contained in a certain open subset of $\Lambda^3 T_p^* M^7$; similarly, a $\text{Spin}(7)$ -structure on an 8-dimensional manifold M^8 is given by a 4-form $\Phi \in \Omega^4(M^8)$ which at each point is contained in a certain subset of $\Lambda^4 T_p^* M^8$. Such structures induce both an orientation and a Riemannian metric on the underlying manifold, denoted by g_φ and g_Φ , respectively, and Φ is self-dual w.r.t. this metric.

Manifolds with G_2 -structures have first been investigated by Fernández and Gray [FG1982], and $\text{Spin}(7)$ -structures by Fernández [Fernandez1986] who showed that the covariant derivatives $\nabla\varphi \in \Omega^1(M^7, \Lambda^3 T^* M^7)$ and $\nabla\Phi \in \Omega^1(M^8, \Lambda^4 T^* M^8)$, respectively, decompose into four irreducible components in case of G_2 -structures and into two irreducible components in the case of $\text{Spin}(7)$ -structures. Thus, the conditions of the vanishing of some of these components yield $2^4 = 16$ classes of G_2 -structure and $2^2 = 4$ classes of $\text{Spin}(7)$ -structures, respectively, and the underlying geometries were discussed in [FG1982] and [Fernandez1986]; see also Section 5 below.

A G_2 -structure ($\text{Spin}(7)$ -structure, respectively) is called *torsion-free*, if φ (Φ , respectively) is parallel. As it turns out, the parallelity of φ and Φ , respectively, is equivalent to φ and Φ being harmonic forms.

Alternatively, G_2 - and $\text{Spin}(7)$ -structures may be characterized via certain (2-fold or 3-fold) cross products on the tangent bundle. These are given as

Date: March 7, 2022.

2010 Mathematics Subject Classification. 53C25, 53C29.

Key words and phrases. Frölicher-Nijenhuis bracket, G_2 -manifold, $\text{Spin}(7)$ -manifold, Fernández-Gray's classification, Fernández's classification.

The first named author is supported Grant-in-Aid for JSPS fellows (26-7067), the second named author is partially supported by RVO: 67985840.

the sections

$$Cr_\varphi := \delta_{g_\varphi} \varphi \in \Omega^2(M^7, TM^7), \quad \chi_\varphi := -\delta_{g_\varphi} * \varphi \in \Omega^3(M^7, TM^7)$$

in case of G_2 -structures, and as

$$P_\Phi := -\delta_{g_\Phi} \Phi \in \Omega^3(M^8, TM^8),$$

where $\delta_g : \Omega^{k+1}(M) \rightarrow \Omega^k(M, TM)$ is the contraction of a differential form with the Riemannian metric g and have natural interpretations via octonian multiplication. The triple cross product $\chi \in \Omega^3(M^7, TM^7)$ on a manifold with a G_2 -structure has been introduced by Harvey-Lawson [HL1982] and was used in many papers on deformation of associative submanifolds, see e.g. [McLean1998], [Kawai2014], [LV2016]. The 3-fold cross product P on \mathbb{R}^8 has been first explicitly constructed by Brown and Gray [BG1967]. They also proved that (up to the G_2 -action) there are exactly two non-equivalent 3-fold cross products on $\mathbb{R}^8 = \mathbb{O}$. In [HL1982] Harvey and Lawson intensively used the 3-fold cross product on \mathbb{R}^8 which is related to the Cayley 4-form and hence is invariant under the action of $\text{Spin}(7)$. Fernandez showed the uniqueness of a $\text{Spin}(7)$ -invariant 4-form on \mathbb{R}^8 (up to a multiplicative constant) and used the associated 3-fold cross product to classify $\text{Spin}(7)$ -structures on 8-manifolds [Fernandez1986].

In this article, we view these cross products as elements of the *Frölicher-Nijenhuis Lie algebra* $\Omega^*(M, TM)$. Namely, it was shown by Frölicher-Nijenhuis in [FN1956] that $\Omega^*(M, TM)$ can be given the structure of a graded Lie algebra using the *Frölicher-Nijenhuis bracket* $[\cdot, \cdot]^{FN}$ in a natural way. Thus, given a manifold with a G_2 -structure (M^7, φ) , we may consider the Frölicher-Nijenhuis brackets

$$[Cr_\varphi, \chi_\varphi]^{FN} \in \Omega^5(M^7, TM^7), \quad [\chi_\varphi, \chi_\varphi]^{FN} \in \Omega^6(M^7, TM^7),$$

(observe that $[Cr, Cr]^{FN} = 0$ due to graded skew-symmetry), and analogously, for a manifold with a $\text{Spin}(7)$ -structure (M^8, Φ) we may consider

$$[P_\Phi, P_\Phi]^{FN} \in \Omega^6(M^8, TM^8).$$

These brackets may be regarded as a natural generalization of the Nijenhuis tensor of an almost complex structure J . Indeed, regarding such a structure as an element $J \in \Omega^1(M, TM)$, it turns out that $[J, J]^{FN} \in \Omega^2(M, TM)$ coincides – up to a constant multiple – with the Nijenhuis tensor of J , whence J is integrable if and only if $[J, J]^{FN} = 0$ [FN1956b].

Our main result is that the Frölicher-Nijenhuis bracket also characterizes the torsion-freeness of G_2 - and $\text{Spin}(7)$ -structures, respectively. Namely, we show the following.

Theorem 1.1. *Let (M^7, φ) be a manifold with a G_2 -structure and the associated Riemannian metric $g = g_\varphi$, and let ∇ be the Levi-Civita connection of g . Then for every $p \in M^7$ the following are equivalent.*

- (1) *The G_2 -structure is torsion-free at p , i.e., $(\nabla\varphi)_p = 0$.*
- (2) *$[Cr_\varphi, \chi_\varphi]_p^{FN} = 0 \in \Lambda^5 T_p^* M^7 \otimes T_p M^7$.*

$$(3) [\chi_\varphi, \chi_\varphi]_p^{FN} = 0 \in \Lambda^6 T_p^* M^7 \otimes T_p M^7.$$

In fact, we show in Theorem 3.7 that $(\nabla\varphi)_p$ is characterized by either $[\chi_\varphi, \chi_\varphi]_p^{FN}$, or by the projection of $[Cr_\varphi, \chi_\varphi]_p^{FN}$ onto a subspace isomorphic to $T_p M^7 \otimes T_p M^7$.

Theorem 1.2. *Let (M^8, Φ) be a manifold with a $\text{Spin}(7)$ -structure, and let ∇ be the Levi-Civita connection of the associated Riemannian metric $g = g_\Phi$. Then for every $p \in M^8$ the following are equivalent.*

- (1) *The $\text{Spin}(7)$ -structure is torsion-free at p , i.e., $(\nabla\Phi)_p = 0$.*
- (2) $[P_\Phi, P_\Phi]_p^{FN} = 0 \in \Lambda^6 T_p^* M^8 \otimes T_p M^8$.

Namely, we show in Theorem 4.6 that $(\nabla\Phi)_p$ is characterized by the projection of $[P_\Phi, P_\Phi]_p^{FN}$ onto a subspace isomorphic to $W_p^7 \otimes T_p M^8$ for some rank-7 bundle $\Lambda^6 T^* M^8 \supset W^7 \rightarrow M^8$.

These explicit descriptions also allow us to give a complete characterization of the 16 cases of G_2 -structures in terms of $[Cr_\varphi, \chi_\varphi]_p^{FN}$ and $[\chi_\varphi, \chi_\varphi]_p^{FN}$, and of the 4 classes of $\text{Spin}(7)$ -structures in terms of $[P_\Phi, P_\Phi]_p^{FN}$; cf. Section 5.

Our paper is organized as follows. In Section 2 we recall the Frölicher-Nijenhuis bracket on $\Omega^*(M, TM)$. Then we turn to the case of G_2 -structures in Section 3, characterizing the torsion endomorphism and showing the results that lead us to Theorem 1.1. In Section 4, we repeat this discussion for the case of $\text{Spin}(7)$ -structures which leads to Theorem 1.2. Finally, the characterization of the 16 classes of G_2 -structures and the 4 classes of $\text{Spin}(7)$ -structures in terms of the Frölicher-Nijenhuis bracket is given in Section 5. The appendix then contains the proofs of some identities on representations of G_2 and $\text{Spin}(7)$ which are used throughout the paper.

2. PRELIMINARIES

2.1. The Frölicher-Nijenhuis bracket. Let M be a manifold and $(\Omega^*(M), \wedge) = (\bigoplus_{k \geq 0} \Omega^k(M), \wedge)$ be the graded algebra of differential forms. We shall use superscripts to indicate the degree of a form, i.e., α^k denotes an element of $\Omega^k(M)$.

Evidently, contraction $\iota_X : \Omega^k(M) \rightarrow \Omega^{k-1}(M)$ with a vector field $X \in \mathfrak{X}(M)$ is a derivation of degree -1 . More generally, for $K \in \Omega^k(M, TM)$ we define $\iota_K \alpha^l$ as the *contraction of K with $\alpha^l \in \Omega^l(M)$* pointwise by

$$\iota_{\kappa^k \otimes X} \alpha^l := \kappa^k \wedge (\iota_X \alpha^l) \in \Omega^{k+l-1}(M),$$

where $\kappa^k \in \Omega^k(M)$ and $X \in \mathfrak{X}(M)$ is a vector field, and this is a derivation of $\Omega^*(M)$ of degree $l-1$. Thus, the *Nijenhuis-Lie derivative along $K \in \Omega^k(M, TM)$* defined as

$$(2.1) \quad \mathcal{L}_K(\alpha^l) := [\iota_K, d](\alpha^l) = \iota_K(d\alpha^l) + (-1)^k d(\iota_K \alpha^l) \in \Omega^{k+l}(M)$$

is a derivation of $\Omega^*(M)$ of degree k .

Observe that for $k = 0$ in which case $K \in \Omega^0(M, TM)$ is a vector field, both ι_K and \mathcal{L}_K coincide with the standard notion of contraction with and Lie derivative along a vector field.

In [FN1956] [FN1956b], it was shown that $\Omega^*(M, TM)$ can be given a unique Lie algebra structure, called the *Frölicher-Nijenhuis bracket* and denoted by $[\cdot, \cdot]^{FN}$, such that \mathcal{L} defines an action of $\Omega^*(M, TM)$ on $\Omega^*(M)$, that is,

$$(2.2) \quad \mathcal{L}_{[K_1, K_2]^{FN}} = [\mathcal{L}_{K_1}, \mathcal{L}_{K_2}] =: \mathcal{L}_{K_1} \circ \mathcal{L}_{K_2} - (-1)^{|K_1||K_2|} \mathcal{L}_{K_2} \circ \mathcal{L}_{K_1}.$$

It is given by the following formula for $\alpha^k \in \Omega^k(M)$, $\beta^l \in \Omega^l(M)$, $X_1, X_2 \in \mathfrak{X}(M)$ [KMS1993, Theorem 8.7 (6), p. 70]:

$$(2.3) \quad \begin{aligned} [\alpha^k \otimes X_1, \beta^l \otimes X_2]^{FN} &= \alpha^k \wedge \beta^l \otimes [X_1, X_2] \\ &+ \alpha^k \wedge \mathcal{L}_{X_1} \beta^l \otimes X_2 - \mathcal{L}_{X_2} \alpha^k \wedge \beta^l \otimes X_1 \\ &+ (-1)^k \left(d\alpha^k \wedge (\iota_{X_1} \beta^l) \otimes X_2 + (\iota_{X_2} \alpha^k) \wedge d\beta^l \otimes X_1 \right). \end{aligned}$$

In particular, for a vector field $X \in \mathfrak{X}(M)$ and $K \in \Omega^*(M, TM)$ we have [KMS1993, Theorem 8.16 (5), p. 75]

$$\mathcal{L}_X(K) = [X, K]^{FN},$$

that is, the Frölicher-Nijenhuis bracket with a vector field coincides with the Lie derivative of the tensor field $K \in \Omega^*(M, TM)$ which means that $\exp(tX) : \Omega^*(M, TM) \rightarrow \Omega^*(M, TM)$ is the action induced by (local) diffeomorphisms of M .

Example 2.1. Let $A \in \Omega^1(M, TM)$ be an endomorphism field on M . Then [KMS1993, Remark 8.17, p. 75]

$$[A, A]^{FN} = 2[A, A]_N,$$

where $[A, A]_N$ is the Nijenhuis tensor of A . W.r.t. a local frame (e_i) with dual frame (e^i) we can write $A = e^i \otimes Ae_i$, whence

$$(2.4) \quad \begin{aligned} \mathcal{L}_A \alpha^k &\stackrel{(2.1)}{=} e^i \wedge (\iota_{Ae_i} d\alpha^k) - d(e^i \wedge (\iota_{Ae_i} \alpha^k)) \\ &= A \cdot d\alpha^k - d(A \cdot \alpha^k), \end{aligned}$$

where we denote by \cdot the pointwise action of $A_p \in \text{End}(T_p M)$ on $\Lambda^k T_p^* M$. Observe that by (2.2) we have $\mathcal{L}_{[A, A]^{FN}} = 2(\mathcal{L}_A)^2$, so that the derivation $\mathcal{L}_A : \Omega^k(M) \rightarrow \Omega^{k+1}(M)$ is a differential iff $[A, A]_N = 0$.

For instance, if $A = Id$ then $I \cdot \alpha^k = e^i \wedge (\iota_{e_i} \alpha^k) = k\alpha^k$, so that

$$\mathcal{L}_I \alpha^k \stackrel{(2.4)}{=} I \cdot d\alpha^k - d(I \cdot \alpha^k) = (k+1)d\alpha^k - d(k\alpha^k) = d\alpha^k.$$

To see another example, let $A = J$ be an almost complex structure. Then $[J, J]^{FN} = 2[J, J]_N = 0$ iff J is integrable, and in this case one calculates from (2.4) that $\mathcal{L}_J = -d^c = i(\partial - \bar{\partial})$ is the negative of the complex differential, where $d = \partial + \bar{\partial}$ is the decomposition into the holomorphic and

anti-holomorphic part of d . In particular, $H_J^*(\Omega^*(M)) \cong H_{dR}^*(M)$ coincides with the deRham cohomology.

We end this section by providing a formula for the Frölicher-Nijenhuis bracket for those types of forms which we shall be concerned with. Recall from the introduction that on a Riemannian manifold (M, g) we define the map

$$(2.5) \quad \delta = \delta_g : \Lambda^k V^* \longrightarrow \Lambda^{k-1} V^* \otimes V, \quad \delta_g(\alpha^k) := (\iota_{e_i} \alpha^k) \otimes (e^i)^\#,$$

taking the sum over some basis (e_i) of $T_p M$ with dual basis (e^i) of $T_p^* M$. This implies that to each $\Psi \in \Omega^{k+1}(M)$ we may associate a section $\delta_g(\Psi) \in \Omega^k(M, TM)$.

Proposition 2.2. *Let (M, g) be an n -dimensional Riemannian manifold of dimension n and let $\Psi_l \in \Omega^{k_l+1}(M)$, $l = 1, 2$. Moreover, let*

$$K_l := \delta_g(\Psi_l) \in \Omega^{k_l}(M, TM)$$

with the map δ_g from (2.5).

Then the Frölicher-Nijenhuis bracket at $p \in M$ is given as

$$\begin{aligned} [K_1, K_2]_p^{FN} = & \left((\iota_{e_i} \Psi_1) \wedge (\iota_{e_j} \nabla_{e_i} \Psi_2) - (-1)^{k_1} (\iota_{e_j} \iota_{e_i} \Psi_1) \wedge e^k \wedge \iota_{e_i} \nabla_{e_k} \Psi_2 \right. \\ & \left. - (\iota_{e_j} \nabla_{e_i} \Psi_1) \wedge (\iota_{e_i} \Psi_2) - (-1)^{k_1} e^k \wedge \iota_{e_i} \nabla_{e_k} \Psi_1 \wedge (\iota_{e_j} \iota_{e_i} \Psi_2) \right) \otimes (e^j)^\#, \end{aligned}$$

where (e_i) is an arbitrary basis of $T_p M$ with dual basis (e^i) of $T_p^* M$. In particular, if $K_1 = K_2 =: K$ and $k_1 = k_2$ is odd, then

$$[K, K]_p^{FN} = 2 \left((\iota_{e_i} \Psi) \wedge (\iota_{e_j} \nabla_{e_i} \Psi) + (\iota_{e_j} \iota_{e_i} \Psi) \wedge e^k \wedge \iota_{e_i} \nabla_{e_k} \Psi \right) \otimes (e^j)^\#.$$

Remark 2.3. If $K_1 = K_2 = K$ and $k_1 = k_2$ is even, then $[K, K]^{FN} = 0$ due to the graded skew symmetry of the bracket. Furthermore, observe that $(e^j)^\# = e_j$ in case (e_i) is an orthonormal basis.

Proof. Evidently, if this formula holds for *some* basis (e_j) with dual basis (e^j) , then it holds for *any* basis. Therefore, it suffices to show the assertion for an orthonormal basis (e_j) in which case $(e^j)^\# = e_j$.

Choose geodesic normal coordinates (x^i) around $p \in M$ in such a way that $(\partial_i)_p := (\partial/\partial x^i)_p$ is an orthonormal basis of $T_p M$. The dual basis of ∂_i is dx^i , whence $(dx^i)^\# = g^{ij} \partial_j$. Thus,

$$K_l = (\iota_{\partial_i} \Psi_l) \otimes (dx^i)^\# = g^{ij} (\iota_{\partial_i} \Psi_l) \otimes \partial_j.$$

Thus, by (2.3)

$$\begin{aligned}
[K_1, K_2]^{FN} &= [g^{ij}(\iota_{\partial_i} \Psi_1) \otimes \partial_j, g^{rs}(\iota_{\partial_r} \Psi_2) \otimes \partial_s]^{FN} \\
&= (g^{ij}(\iota_{\partial_i} \Psi_1) \wedge \mathcal{L}_{\partial_j}(g^{rs}(\iota_{\partial_r} \Psi_2)) \otimes \partial_s \\
&\quad - \mathcal{L}_{\partial_s}(g^{ij}(\iota_{\partial_i} \Psi_1)) \wedge g^{rs}(\iota_{\partial_r} \Psi_2) \otimes \partial_j \\
&\quad + (-1)^{k_1} d(g^{ij}(\iota_{\partial_i} \Psi_1)) \wedge \iota_{\partial_j}(g^{rs}(\iota_{\partial_r} \Psi_2)) \otimes \partial_s \\
&\quad + (-1)^{k_1} (\iota_{\partial_s}(g^{ij}(\iota_{\partial_i} \Psi_1)) \wedge d(g^{rs}(\iota_{\partial_r} \Psi_2)) \otimes \partial_j).
\end{aligned}$$

Since at p , $g_{ij} = g^{ij} = \delta_{ij}$, $\partial_r g_{ij} = 0$, $\mathcal{L}_{\partial_j} \Psi = \nabla_{e_j} \Psi$, $\nabla_{\partial_i} \partial_j = 0$, and $\partial_j = (e^j)^\#$, the asserted formula follows. \square

3. CROSS PRODUCTS AND G_2 -STRUCTURES

3.1. G_2 -structures and associated cross products. In this section we collect some basic facts on G_2 -structures, see e.g. [Humphreys1978], [Bryant1987], [FG1982], [HL1982] for references.

Let M be an oriented 7-manifold. A G_2 -structure on M is a 3-form $\varphi \in \Omega^3(M)$ such that at each $p \in M$ there is a positively oriented basis (e_i) of $T_p M$ with dual basis (e^i) such that

$$(3.1) \quad \varphi_p = e^{123} + e^{145} + e^{167} + e^{246} - e^{257} - e^{347} - e^{356},$$

where $e^{i_1 \dots i_k}$ is short for $e^{i_1} \wedge \dots \wedge e^{i_k}$. We call such a basis a G_2 -frame. The stabilizer of φ_p is isomorphic to the exceptional group G_2 , and there is a unique G_2 -invariant Riemannian metric g_φ on M such that each G_2 -frame is orthonormal. In particular, the Hodge-dual of φ w.r.t. g_φ is given by

$$(3.2) \quad *_\varphi \varphi = e^{4567} + e^{2367} + e^{2345} + e^{1357} - e^{1346} - e^{1256} - e^{1247}.$$

The set of G_2 -frames yields a principal G_2 -bundle

$$G_2(M) = G_2(M, \varphi) \longrightarrow M,$$

whence for each G_2 -module V we denote by

$$(3.3) \quad V(M) := G_2(M) \times_{G_2} V \longrightarrow M$$

the associated vector bundle over M . For instance,

$$V_7(M) \cong TM \cong T^*M.$$

Definition 3.1. [HL1982] Let (M, φ) be an oriented manifold with a G_2 -structure φ and the induced Riemannian metric $g = g_\varphi$. Then the TM -valued forms $Cr_\varphi \in \Omega^2(M, TM)$ and $\chi_\varphi \in \Omega^3(M, TM)$ are defined by

$$Cr_\varphi := \delta_{g_\varphi}(\varphi) \quad \text{and} \quad \chi_\varphi := -\delta_g(*\varphi),$$

and are called the *2-fold and 3-fold cross product on M* , respectively. That is, for $x, y, z, w \in TM$ we have

$$g_\varphi(Cr_\varphi(x, y), z) = \varphi(x, y, z), \quad g_\varphi(\chi_\varphi(x, y, z), w) = *\varphi(x, y, z, w).$$

We shall usually suppress the indices φ for g , Cr and χ if it is clear from the context which G_2 -structure φ is used.

Remark 3.2. If we use the G_2 -structure to identify each $T_p M \cong Im\mathbb{O}$ with the imaginary octonions, then Cr and χ can be interpreted w.r.t. the octonian product $\cdot : \mathbb{O} \times \mathbb{O} \rightarrow \mathbb{O}$ as

$$Cr(x, y) := (x \cdot y)_{Im\mathbb{O}} \quad \text{and} \quad \chi(x, y, z) := ((x \cdot y) \cdot z - x \cdot (y \cdot z))_{Im\mathbb{O}}.$$

We summarize important known facts about the decomposition of tensor products of G_2 -modules into irreducible summands which are well known, see e.g. [Kar2005, Section 2]. We denote by V_k the k -dimensional irreducible G_2 -module if there is a unique such module. For instance, V_7 is the irreducible 7-dimensional G_2 -module from above, and $V_7^* \cong V_7$. For its exterior powers, we obtain the decompositions

$$(3.4) \quad \begin{aligned} \Lambda^0 V_7 &\cong \Lambda^7 V_7 \cong V_1, & \Lambda^2 V_7 &\cong \Lambda^5 V_7 \cong V_7 \oplus V_{14}, \\ \Lambda^1 V_7 &\cong \Lambda^6 V_7 \cong V_7, & \Lambda^3 V_7 &\cong \Lambda^4 V_7 \cong V_1 \oplus V_7 \oplus V_{27}, \end{aligned}$$

where $\Lambda^k V_7 \cong \Lambda^{7-k} V_7$ due to the Hodge isomorphism. We denote by $\Lambda_l^k V_7 \subset \Lambda^k V_7$ the subspace isomorphic to V_l in the above notation. Evidently, $\Lambda_1^3 V_7$ and $\Lambda_1^4 V_7$ are spanned by φ and $*\varphi$, respectively. For the decompositions of $\Lambda^2 V_7$ and $\Lambda^5 V_7$ the following descriptions are well known.

$$(3.5) \quad \begin{aligned} \Lambda_7^2 V_7 &= \{\iota_v \varphi \mid v \in V_7\}, \\ \Lambda_{14}^2 V_7 &= \{\alpha^2 \in \Lambda^2 V_7 \mid *\varphi \wedge \alpha^2 = 0\}, \\ \Lambda_1^3 V_7 &= \mathbb{R}\varphi, \\ \Lambda_7^3 V_7 &= \{\iota_v *\varphi \mid v \in V_7\}, \\ \Lambda_1^4 V_7 &= \mathbb{R}*\varphi, \\ \Lambda_7^4 V_7 &= \varphi \wedge V_7 = \{\varphi \wedge v \mid v \in V_7\}, \\ \Lambda_7^5 V_7 &= *\varphi \wedge V_7 = \{*\varphi \wedge v \mid v \in V_7\}, \\ \Lambda_{14}^5 V_7 &= \{\alpha^5 \in \Lambda^5 V_7 \mid \alpha^5 \wedge (\iota_v \varphi) = 0 \text{ for all } v \in V_7\}. \end{aligned}$$

We also point out that all representations of G_2 are of real type, meaning that for any real irreducible representation V of G_2 the complexified space $V^{\mathbb{C}} := V \otimes \mathbb{C}$ is (complex) irreducible; equivalently, a real irreducible representation of G_2 does not admit a G_2 -invariant complex structure.

These decompositions are used in the appendix to obtain many formulas which will be used in the sequel.

3.2. The torsion of manifolds with a G_2 -structure. Let (M, φ) be a manifold with a G_2 -structure with the corresponding Riemannian metric $g = g_\varphi$, and let ∇ be the Levi-Civita connection of g . In general, φ and $*\varphi$ will not be parallel w.r.t. ∇ , and the failure of their parallelity can be

described in the following way which is essentially a reformulation of the intrinsic torsion of a G_2 -structure discussed in [Bryant2005] and [Kar2005].

Proposition 3.3. *Let (M, φ) be a manifold with a G_2 -structure with associated Riemannian metric $g = g_\varphi$ and Levi-Civita connection ∇ . Then there is a section $T \in \Omega^1(M, TM) = \Gamma(\text{End}(TM))$ such that for all $v \in TM$ we have*

$$(3.6) \quad \nabla_v \varphi = \iota_{T(v)} * \varphi \quad \text{and} \quad \nabla_v * \varphi = -(T(v))^\flat \wedge \varphi.$$

Thus, the section $T \in \Omega^1(M, TM)$ measures how φ fails to be parallel, and this has been described in Fernández and Gray [FG1982] by slightly different means. In fact, it contains the same information as the intrinsic torsion of the G_2 -structure in the sense of [Bryant2005], whence we use the following terminology.

Definition 3.4. Let (M, φ) be a manifold with a G_2 -structure. The section $T \in \Omega^1(M, TM)$ for which (3.6) holds is called the *torsion endomorphism* of the G_2 -structure.

For an orthonormal frame (e_i) of $T_p M$ we define the coefficients of T by

$$(3.7) \quad t_{ij} := \langle T(e_i), e_j \rangle, \quad \text{so that} \quad T(e_i) = t_{ij} e_j.$$

Furthermore, we define the form

$$(3.8) \quad \tau := t_{ij} e^{ij} = \frac{1}{2}(t_{ij} - t_{ji})e^{ij} = e^i \wedge T(e_i)^\flat \in \Lambda^2 V_7^*.$$

For the exterior derivatives of φ and $*\varphi$, we have

$$(3.9) \quad d\varphi_p = T_p^\top(e_i)^\flat \wedge (\iota_{e_i} * \varphi_p) \quad \text{and} \quad d * \varphi_p = -\tau_p \wedge \varphi_p,$$

where we sum over an orthonormal basis (e_i) of $T_p M$ in the first equation and where T_p^\top denotes the transpose matrix of T_p . In particular, it is now a straightforward calculation to show that (M, φ) is torsion free at $p \in M$ (i.e., $T_p = 0$) iff $d\varphi_p = 0$ and $d * \varphi_p = 0$ (cf. [FG1982]).

3.3. The Frölicher-Nijenhuis brackets on a manifold with a G_2 -structure. In this section, we shall compute part of their Frölicher-Nijenhuis brackets of the sections $Cr = \delta_g \varphi \in \Omega^2(M, TM)$ and $\chi = -\delta_g * \varphi \in \Omega^3(M, TM)$. from Definition 3.1 on a manifold M with a G_2 -structure φ .

The Frölicher-Nijenhuis bracket $[Cr, Cr]^{FN}$ vanishes identically due to the graded skew-symmetry of the bracket. On the other hand, the Frölicher-Nijenhuis brackets $[Cr, \chi]^{FN}$ and $[\chi, \chi]^{FN}$ are elements of $\Omega^5(M, TM)$ and $\Omega^6(M, TM)$, respectively.

Due to the decomposition $\Lambda^5 V_7 = \Lambda_7^5 V_7 \oplus \Lambda_{14}^5 V_7$ as a G_2 -module, we may decompose

$$\Omega^5(M, TM) = \Gamma(M, \Lambda_7^5 T^* M \otimes TM) \oplus \Gamma(M, \Lambda_{14}^5 T^* M \otimes TM),$$

and we denote the projections onto the two summands by π_7 and π_{14} , respectively. We now wish to show Theorem 1.1 from the introduction. In order to work towards the proof, we first calculate $\pi_7([Cr, \chi]^{FN})$.

Proposition 3.5. *Let (M, g, φ) be a manifold with a G_2 -structure and let $T \in \Omega^1(M, TM)$ be its torsion endomorphism. Then for each $p \in M$,*

$$(3.10) \quad \pi_7([Cr, \chi]_p^{FN}) = 2 * \varphi \wedge \left((T_p^\top - 2T_p - \text{tr}(T_p)) e_i \right)^b \otimes e_i,$$

summing over an orthonormal basis (e_i) of $T_p M$, where T^\top denotes the transpose of T . In particular, $\pi_7([Cr, \chi]_p^{FN}) = 0$ if and only if $T_p = 0$ if and only if $[Cr, \chi]_p^{FN} = 0$.

Proof. We fix $p \in M$ and use normal coordinates around p . Then in order to calculate $[Cr, \chi]_p^{FN}$ we apply Proposition 2.2 to $\Psi_1 = \varphi$ and $\Psi_2 = *\varphi$ and obtain

$$-[Cr, \chi]_p^{FN} = [(\iota_{e_i} \varphi) \otimes e_i, (\iota_{e_j} *\varphi) \otimes e_j]_p^{FN} =: \beta_j \otimes e_j,$$

where

$$(3.11) \quad \begin{aligned} \beta_j &= (\iota_{e_k} \varphi) \wedge (\iota_{e_j} \nabla_{e_k} *\varphi) - (\iota_{e_j} \iota_{e_k} \varphi) \wedge e^l \wedge (\iota_{e_k} \nabla_{e_l} *\varphi) \\ &\quad - (\iota_{e_j} \nabla_{e_k} \varphi) \wedge (\iota_{e_k} *\varphi) - e^l \wedge (\iota_{e_k} \nabla_{e_l} \varphi) \wedge (\iota_{e_j} \iota_{e_k} *\varphi). \end{aligned}$$

Decomposing $\Lambda^5 T_p^* M$ according to (3.5), we write $\beta_j = *\varphi \wedge v_j^b + \beta_j^{14}$ with $v_j \in T_p M$ and $\beta_j^{14} \in \Lambda_{14}^5 T_p^* M$, so that $\pi_7([Cr, \chi]_p^{FN}) = *\varphi \wedge v_j^b \otimes e_j$. Let

$$(3.12) \quad b_{ij} = *((\iota_{e_i} \varphi) \wedge \beta_j).$$

Then as $(\iota_{e_i} \varphi) \wedge \beta_j^{14} = 0$ by (3.5) and $(\iota_{e_i} \varphi) \wedge *\varphi \wedge v_j^b = 3\langle e_i, v_j \rangle \text{vol}$ by (6.1), it follows that

$$(3.13) \quad \pi_7(-[Cr, \chi]_p^{FN}) = \frac{1}{3} b_{ij} (*\varphi \wedge e^i) \otimes e_j.$$

In order to determine the coefficients b_{ij} , we decompose β_j into the four summands from (3.11). Then from the first summand we get

$$(3.14) \quad \begin{aligned} (\iota_{e_i} \varphi) \wedge (\iota_{e_k} \varphi) \wedge (\iota_{e_j} \nabla_{e_k} *\varphi) &\stackrel{(3.6)}{=} -(\iota_{e_i} \varphi) \wedge (\iota_{e_k} \varphi) \wedge (\iota_{e_j} (T(e_k)^b \wedge \varphi)) \\ &= -(\iota_{e_i} \varphi) \wedge (\iota_{e_k} \varphi) \wedge (t_{kj} \varphi - T(e_k)^b \wedge (\iota_{e_j} \varphi)) \\ &\stackrel{(6.3), (6.4)}{=} -6t_{kj} \delta_{ik} \text{vol} + 2(\delta_{ik} t_{kj} + t_{ki} \delta_{kj} + t_{kk} \delta_{ij}) \text{vol} \\ &= 2(t_{ji} - 2t_{ij} + \text{tr}(T) \delta_{ij}) \text{vol}. \end{aligned}$$

From the second summand we obtain

$$\begin{aligned}
& -(\iota_{e_i}\varphi) \wedge (\iota_{e_j}\iota_{e_k}\varphi) \wedge e^l \wedge (\iota_{e_k}\nabla_{e_l} * \varphi) \\
& \stackrel{(3.6)}{=} (\iota_{e_i}\varphi) \wedge (\iota_{e_j}\iota_{e_k}\varphi) \wedge e^l \wedge (\iota_{e_k}(T(e_l)^\flat \wedge \varphi)) \\
& = (\iota_{e_i}\varphi) \wedge (\iota_{e_j}\iota_{e_k}\varphi) \wedge e^l \wedge (t_{lk}\varphi - T(e_l)^\flat \wedge (\iota_{e_k}\varphi)) \\
& \stackrel{(6.6)}{=} t_{lk} \left(2(\delta_{kl}\delta_{ji} - \delta_{jl}\delta_{ki})\text{vol} - 2e^{jkl} \wedge \varphi \right) \\
& \quad - (\iota_{e_i}\varphi) \wedge (\iota_{e_j}\iota_{e_k}\varphi) \wedge e^l \wedge T(e_l)^\flat \wedge (\iota_{e_k}\varphi) \\
& = 2(\text{tr}(T)\delta_{ij} - t_{ji})\text{vol} - 2e^{ij} \wedge \tau \wedge \varphi \\
& \quad - \frac{1}{2}(\iota_{e_i}\varphi) \wedge \iota_{e_j}((\iota_{e_k}\varphi) \wedge (\iota_{e_k}\varphi)) \wedge \tau \\
& \stackrel{(6.12)}{=} 2(\text{tr}(T)\delta_{ij} - t_{ji})\text{vol} - 2e^{ij} \wedge \tau \wedge \varphi \\
& \quad - 3(\iota_{e_i}\varphi) \wedge (\iota_{e_j} * \varphi) \wedge \tau \\
& \stackrel{(6.10)}{=} 2(\text{tr}(T)\delta_{ij} - t_{ji})\text{vol} - 2e^{ij} \wedge \tau \wedge \varphi \\
& \quad - 3(-2 * e^{ji} + e^{ji} \wedge \varphi) \wedge \tau \\
& = 2(\text{tr}(T)\delta_{ij} - t_{ji})\text{vol} + e^{ij} \wedge \tau \wedge \varphi \\
& \quad + 6t_{kl}(\delta_{jk}\delta_{il} - \delta_{jl}\delta_{ik})\text{vol} \\
(3.15) \quad & = 2(\text{tr}(T)\delta_{ij} + 2t_{ji} - 3t_{ij})\text{vol} + e^{ij} \wedge \tau \wedge \varphi.
\end{aligned}$$

From the third term in (3.11) we get

$$\begin{aligned}
& -(\iota_{e_i}\varphi) \wedge (\iota_{e_j}\nabla_{e_k}\varphi) \wedge (\iota_{e_k} * \varphi) \\
& \stackrel{(3.6)}{=} -(\iota_{e_i}\varphi) \wedge (\iota_{e_j}\iota_{T(e_k)} * \varphi) \wedge (\iota_{e_k} * \varphi) \\
& \stackrel{(6.7)}{=} -2(t_{kk}\delta_{ji} - \delta_{jk}t_{ki})\text{vol} - e^j \wedge T(e_k)^\flat \wedge e^{ki} \wedge \varphi \\
(3.16) \quad & = -2(\text{tr}(T)\delta_{ij} - t_{ji})\text{vol} - e^{ij} \wedge \tau \wedge \varphi.
\end{aligned}$$

Finally, from the last term in (3.11) we get

$$\begin{aligned}
& -(\iota_{e_i}\varphi) \wedge e^l \wedge (\iota_{e_k}\nabla_{e_l}\varphi) \wedge (\iota_{e_j}\iota_{e_k} * \varphi) \\
& \stackrel{(3.6)}{=} -(\iota_{e_i}\varphi) \wedge e^l \wedge (\iota_{e_k}\iota_{T(e_l)} * \varphi) \wedge (\iota_{e_j}\iota_{e_k} * \varphi) \\
& \stackrel{(6.13)}{=} 2(\iota_{e_i}\varphi) \wedge e^l \wedge (\iota_{T(e_l)}\varphi) \wedge (\iota_{e_j}\varphi) \\
& \stackrel{(6.4)}{=} 4(t_{li}\delta_{lj} + \delta_{li}t_{lj} + t_{ll}\delta_{ij})\text{vol} \\
(3.17) \quad & = 4(t_{ji} + t_{ij} + \text{tr}(T)\delta_{ij})\text{vol}.
\end{aligned}$$

Thus, adding (3.14) through (3.17), we get from (3.11) that

$$\begin{aligned}
b_{ij}\text{vol} &= (\iota_{e_i}\varphi) \wedge \beta_j = 2(t_{ji} - 2t_{ij} + \text{tr}(T)\delta_{ij})\text{vol} \\
&\quad + 2(\text{tr}(T)\delta_{ij} + 2t_{ji} - 3t_{ij})\text{vol} + e^{ij} \wedge \tau \wedge \varphi \\
&\quad - 2(\text{tr}(T)\delta_{ij} - t_{ji})\text{vol} - e^{ij} \wedge \tau \wedge \varphi \\
&\quad + 4(t_{ji} + t_{ij} + \text{tr}(T)\delta_{ij})\text{vol} \\
&= 6(2t_{ji} - t_{ij} + \text{tr}(T)\delta_{ij})\text{vol},
\end{aligned}$$

and hence, (3.13) implies (3.10).

Thus, $\pi_7([Cr, \chi]_p^{FN}) = 0$ iff $T_p^\top - 2T_p - \text{tr}(T_p)Id = 0$. Taking the trace, this implies that $\text{tr}(T_p) - 2\text{tr}(T_p) - 7\text{tr}(T_p) = 0$ and hence, $\text{tr}(T_p) = 0$, and $T_p^\top - 2T_p = 0$ evidently implies that $T_p = 0$. That is, $\pi_7([Cr, \chi]_p^{FN}) = 0$ iff $T_p = 0$, showing the last statement. \square

Next, let us consider the bracket $[\chi, \chi]^{FN}$.

Proposition 3.6. *Let (M, g, φ) be a manifold with a G_2 -structure and let $T \in \Omega^1(M, TM)$ be its torsion endomorphism with the associated form $\tau \in \Omega^2(M)$ from (3.8). Then for each $p \in M$,*

$$(3.18) \quad [\chi, \chi]_p^{FN} = -4 * (T_p + T_p^\top)(e_i) \otimes e_i + 6e^i \wedge \tau_p \wedge \varphi \otimes e_i,$$

summing over an orthonormal basis (e_i) of T_pM . In particular, $[\chi, \chi]_p^{FN} = 0$ if and only if $T_p = 0$.

Proof. According to Proposition 2.2 we have

$$[\chi, \chi]_p^{FN} = \gamma_j \otimes e_j,$$

where

$$(3.19) \quad \gamma_j = 2((\iota_{e_k} * \varphi) \wedge (\iota_{e_j} \nabla_{e_k} * \varphi) + (\iota_{e_j} \iota_{e_k} * \varphi) \wedge e^l \wedge (\iota_{e_k} \nabla_{e_l} * \varphi)).$$

Now let $c_{ij} := *(e^i \wedge \gamma_j) = \langle e_i, (*\gamma_j)^\flat \rangle$. Then

$$(3.20) \quad [\chi, \chi]_p^{FN} = c_{ij} * e^i \otimes e_j,$$

In order to evaluate the coefficients c_{ij} , we consider the two summands in (3.19) separately, and obtain from the first one

$$\begin{aligned}
e^i \wedge (\iota_{e_k} * \varphi) \wedge (\iota_{e_j} \nabla_{e_k} * \varphi) &\stackrel{(3.6)}{=} -e^i \wedge (\iota_{e_k} * \varphi) \wedge (\iota_{e_j} (T(e_k)^\flat \wedge \varphi)) \\
&= -t_{kj} e^i \wedge (\iota_{e_k} * \varphi) \wedge \varphi + e^i \wedge (\iota_{e_k} * \varphi) \wedge T(e_k)^\flat \wedge (\iota_{e_j} \varphi) \\
&\stackrel{(6.2), (6.8)}{=} -4t_{kj} \delta_{ik} \text{vol} + 2(\delta_{ik} t_{kj} - t_{kk} \delta_{ij}) \text{vol} + T(e_k)^\flat \wedge e^{ikj} \wedge \varphi \\
(3.21) \quad &= -2(t_{ij} + \text{tr}(T)\delta_{ij}) \text{vol} + e^{ij} \wedge \tau \wedge \varphi.
\end{aligned}$$

From the second summand in (3.19) we calculate

$$\begin{aligned}
& e^i \wedge (\iota_{e_j} \iota_{e_k} * \varphi) \wedge e^l \wedge (\iota_{e_k} \nabla_{e_l} * \varphi) \\
& \stackrel{(3.6)}{=} -e^{il} \wedge (\iota_{e_j} \iota_{e_k} * \varphi) \wedge (\iota_{e_k} (T(e_l)^b \wedge \varphi)) \\
& = -t_{lk} e^{il} \wedge (\iota_{e_j} \iota_{e_k} * \varphi) \wedge \varphi + e^i \wedge (\iota_{e_j} \iota_{e_k} * \varphi) \wedge \tau \wedge (\iota_{e_k} \varphi) \\
(6.9), (6.14) \quad & \stackrel{=}{=} -t_{lk} (2(\delta_{lj} \delta_{ik} - \delta_{ij} \delta_{lk}) \text{vol} - e^{iljk} \wedge \varphi) + 3e^{ij} \wedge \tau \wedge \varphi \\
(3.22) \quad & = -2(t_{ji} - \text{tr}(T)\delta_{ij}) \text{vol} + 2e^{ij} \wedge \tau \wedge \varphi.
\end{aligned}$$

Thus, adding (3.21) and (3.22), equation (3.19) yields

$$\begin{aligned}
c_{ij} \text{vol} = e^i \wedge \gamma_j &= 2 \left(-2(t_{ij} + \text{tr}(T)\delta_{ij}) \text{vol} + e^{ij} \wedge \tau \wedge \varphi \right. \\
& \quad \left. - 2(t_{ji} - \text{tr}(T)\delta_{ij}) \text{vol} + 2e^{ij} \wedge \tau \wedge \varphi \right) \\
&= 2(-2(t_{ij} + t_{ji}) + 3\langle *e^i, e^j \wedge \tau \wedge \varphi \rangle) \text{vol},
\end{aligned}$$

and from this and (3.20), the formula (3.18) follows.

In order to show the last statement, observe that $[\chi, \chi]_p^{FN} = 0$ iff $c_{ij} = 0$ for all i, j . Since then $c_{ij} + c_{ji} = -8(t_{ij} + t_{ji})$, it follows that $t_{ij} + t_{ji} = 0$ and hence $0 = c_{ij} = 6 * (e^{ij} \wedge \tau \wedge \varphi)$ for all i, j which implies that $\tau = t_{kl} e^{kl} = 0$ and hence, $t_{kl} = t_{lk}$. All of this together implies that $t_{ij} = 0$ for all i, j , and hence, $T_p = 0$ as asserted. \square

We are now ready to show the following result which immediately implies Theorem 1.1 from the introduction.

Theorem 3.7. *Let (M^7, φ) be a manifold with a G_2 -structure with associated metric $g = g_\varphi$, let ∇ be the Levi-Civita connection of g , and let $T \in \Omega^1(M^7, TM^7)$ be its torsion endomorphism defined in Definition 3.4. Then for every $p \in M^7$ the following are equivalent.*

- (1) $T_p = 0 \in T_p^* M^7 \otimes T_p M^7$.
- (2) The G_2 -structure is torsion-free at p , i.e., $(\nabla \varphi)_p = 0$.
- (3) $\pi_7([Cr, \chi]_p^{FN}) = 0 \in \Lambda_7^5 T_p^* M^7 \otimes T_p M^7$.
- (4) $[Cr, \chi]_p^{FN} = 0 \in \Lambda^5 T_p^* M^7 \otimes T_p M^7$.
- (5) $[\chi, \chi]_p^{FN} = 0 \in \Lambda^6 T_p^* M^7 \otimes T_p M^7$.

Proof. The equivalence of the first two statements is well known, see e.g. [FG1982]. Proposition 3.5 shows the equivalence of the first and the third, whereas Proposition 3.6 shows the equivalence of the first and the last statement. That $(\nabla \varphi)_p = 0$ implies $[Cr, \chi]_p^{FN} = 0$ is immediate from the formula of the bracket in Proposition 2.2, and obviously, $[Cr, \chi]_p^{FN} = 0$ implies $\pi_7([Cr, \chi]_p^{FN}) = 0$. \square

4. CROSS PRODUCTS AND Spin(7)-STRUCTURES

4.1. **Spin(7)-structures and associated cross products.** The exposition in this section mainly follows the references [Bryant1987], [Fernandez1986], [HL1982].

Let M be an oriented 8-manifold. A Spin(7)-*structure on M* is a 4-form $\Phi \in \Omega^4(M)$ such that at each $p \in M$ there is a positively oriented basis $(e_\mu)_{\mu=0}^7$ of T_pM with dual basis $(e^\mu)_{\mu=0}^7$ such that $\Phi_p \in \Lambda^4 T_pM$ is of the form

$$(4.1) \quad \Phi_p := e^{0123} + e^{0145} + e^{0167} + e^{0246} - e^{0257} - e^{0347} - e^{0356} \\ + e^{4567} + e^{2367} + e^{2345} + e^{1357} - e^{1346} - e^{1256} - e^{1247}.$$

Throughout this section, we shall use Greek indices μ, ν, \dots to run over $0, \dots, 7$, whereas Latin indices i, j, \dots range over $1, \dots, 7$.

A basis (e_μ) of T_pM whose dual basis (e^μ) satisfies (4.1) is called a Spin(7)-*frame*. Observe that if we define for a Spin(7)-frame (e_μ) the forms φ_p and $*_7\varphi_p$ on $V_p := \text{span}(e_i)_{i=1}^7 \subset T_pM$ as in (3.1) and (3.2), then

$$\Phi_p = e^0 \wedge \varphi_p + *_7\varphi_p.$$

The stabilizer of Φ_p is the group Spin(7) acting on T_pM via the spinor representation, and there is a unique Spin(7)-invariant Riemannian metric g_Φ on M such that each Spin(7)-frame is orthonormal. In particular, Φ is self-dual w.r.t. g_Φ . The set of all Spin(7)-frames forms a principal Spin(7)-bundle

$$\text{Spin}(7)_M = \text{Spin}(7)_{(M, \Phi)} \longrightarrow M,$$

and again, for each Spin(7)-module W we obtain the associated vector bundle

$$(4.2) \quad W(M) := \text{Spin}(7)_M \times_{\text{Spin}(7)} W \longrightarrow M.$$

For instance, if we denote the k -dimensional irreducible Spin(7)-module by W_k (in case the dimension uniquely specifies this module), then

$$W_8(M) \cong TM \cong T^*M.$$

It is well known that the action of Spin(7) on W_8 is transitive on the unit sphere $S^7 \subset W_8$, and the stabilizer of an element is isomorphic to $G_2 \subset \text{Spin}(7)$. In analogy of the products Cr and χ on manifolds with a G_2 -structure in Definition 3.1, we define on a Spin(7)-manifold M a triple product as follows.

Definition 4.1. Let (M, Φ) be manifold with a Spin(7)-structure, and let $g = g_\Phi$ be the induced Riemannian metric. Then the TM -valued form $P = P_\Phi \in \Omega^3(M, TM)$ is defined by

$$P_\Phi := -\delta_{g_\Phi}(\Phi),$$

and is called the *3-fold cross product on M* . That is, for $x, y, z, w \in TM$ we have

$$(4.3) \quad g(P(x, y, z), w) = \Phi(x, y, z, w).$$

We shall usually suppress the indices Φ for g and P if it is clear from the context which Spin(7)-structure Φ is used.

4.2. Spin(7)-representations. In this section, we shall discuss the decomposition of symmetric and anti-symmetric powers of W_8 as Spin(7)-modules. For its exterior powers, we obtain the decompositions

$$(4.4) \quad \begin{aligned} \Lambda^0 W_8 &\cong \Lambda^8 W_8 \cong W_1, & \Lambda^2 W_8 &\cong \Lambda^6 W_8 \cong W_7 \oplus W_{21}, \\ \Lambda^1 W_8 &\cong \Lambda^7 W_8 \cong W_8, & \Lambda^3 W_8 &\cong \Lambda^5 W_8 \cong W_8 \oplus W_{48}, \\ \Lambda^4 W_8 &\cong W_1 \oplus W_7 \oplus W_{27} \oplus W_{35} \end{aligned}$$

where $\Lambda^k W_8 \cong \Lambda^{8-k} W_8$ via the Hodge-*. Again, we denote by $\Lambda_l^k W_8 \subset \Lambda^k W_8$ the subspace isomorphic to W_l in the above notation.

Moreover, there are also irreducible decompositions of the symmetric powers of W_7 and W_8 as

$$(4.5) \quad \odot^2 W_7 \cong W_1 \oplus W_{27}, \quad \odot^2 W_8 \cong W_1 \oplus W_{35}$$

into the induced Spin(7)-invariant metric and the trace free symmetric tensors; see [Humphreys1978].

Lemma 4.2. *Let $e^0 \in W_8$ be a unit vector, let $V_7 := e_0^\perp$ on which Spin(7) acts as the double cover of $SO(7)$, so that $V_7 \cong W_7$ as a Spin(7)-module. Then the following maps are Spin(7)-equivariant embeddings.*

$$(4.6) \quad \begin{aligned} \lambda^2(v) &:= e^0 \wedge v^b + (i_v \varphi) \\ \lambda^k : W_7 &\longrightarrow \Lambda^k W_8, & \lambda^4(v) &:= e^0 \wedge (i_v * \varphi) - v^b \wedge \varphi \\ \lambda^6(v) &:= \Phi \wedge \lambda^2(v) = 3 * \lambda^2(v) \end{aligned}$$

Here $*$ and $*_7$ denote the Hodge-* in W_8 and V_7 , respectively.

Proof. The decompositions in (4.5) imply that there are Spin(7)-equivariant maps $\lambda^k : W_7 \rightarrow \Lambda^k W_8$, and these are unique up to rescaling.

The equivariance of λ^2 follows from [SW2017, p.68], and thus, $\Phi \wedge \lambda^2(v) \in \Lambda_7^6 W_8$, whence $\Phi \wedge \lambda^2(v) = 3 * \lambda^2(v)$ follows from [SW2017, Theorem 9.8]. This shows the statement on λ^6 .

By [SW2017, Theorem 9.8], $\Lambda_7^4 W_8$ is the infinitesimal orbit of Φ under the action of $\mathfrak{so}(W_8) \cong \Lambda^2 W_8$. That is,

$$\Lambda_7^4 W_8 = \{(u^b \wedge v^b) \cdot \Phi \mid u, v \in W_8\} = \{u^b \wedge (i_v \Phi) - v^b \wedge (i_u \Phi) \mid u, v \in W_8\}.$$

Setting $u := e_0$ and picking $v \in e_0^\perp \cong W_7$ for a Spin(7)-frame (e_μ) , it follows that the image of λ^4 equals $\Lambda_7^4 W_8$, and since λ^4 is evidently G_2 -equivariant, it must coincide with the Spin(7)-equivariant map $W_7 \rightarrow \Lambda_7^4 W_8$. \square

From this lemma, we obtain the following descriptions of the decompositions, which essentially recapitulates [SW2017, Theorem 9.8].

$$\begin{aligned}
\Lambda_7^k W_8 &= \{\lambda^k(v) \mid v \in V_7\} \quad \text{for } k = 2, 4, 6, \\
\Lambda_{21}^2 W_8 &= \{\alpha^2 \in \Lambda^2 W_8 \mid \alpha^2 \wedge \Phi \wedge \lambda^2(v) = 0 \text{ for all } v \in V_7\}, \\
\Lambda_{21}^6 W_8 &= \{\alpha^6 \in \Lambda^6 W_8 \mid \alpha^6 \wedge \lambda^2(v) = 0 \text{ for all } v \in V_7\}, \\
\Lambda_8^3 W_8 &= \{\iota_a \Phi \mid a \in W_8\}, \\
\Lambda_8^5 W_8 &= \{a^\flat \wedge \Phi \mid a \in W_8\}, \\
(4.7) \quad \Lambda_{48}^3 W_8 &= \{\alpha^3 \in \Lambda^3 W_8 \mid \Phi \wedge \alpha^3 = 0\}, \\
\Lambda_{48}^5 W_8 &= \{\alpha^5 \in \Lambda^3 W_8 \mid \Phi \wedge * \alpha^5 = 0\}, \\
\Lambda_1^4 W_8 &= \mathbb{R} \Phi, \\
\Lambda_{27}^4 W_8 &= \text{span}\{\lambda^2(v) \wedge \lambda^2(w) \mid v, w \in V_7, \langle v, w \rangle = 0\} \\
\Lambda_{35}^4 W_8 &= \{\alpha^4 \mid * \alpha^4 = -\alpha^4\}.
\end{aligned}$$

We also recall the decomposition of the tensor product

$$(4.8) \quad \text{Lin}(W_8, W_7) := W_8^* \otimes W_7 = W_8 \oplus W_{48}.$$

Here, the summand isomorphic to W_8 is given as

$$(4.9) \quad \{(\iota_a \lambda^2(e_i)) \otimes e_i \mid a \in W_8\},$$

where the sum is taken over an orthonormal basis (e_i) of $V_7 \cong W_7$. Finally, we define the $\text{Spin}(7)$ -invariant tensor $\sigma \in (W_8 \otimes W_7 \otimes W_8 \otimes W_7)^*$ by

$$(4.10) \quad \sigma(a, u, b, v) := \frac{1}{2} * (a^\flat \wedge b^\flat \wedge \lambda^4(u) \wedge \lambda^2(v)).$$

Contraction with the inner products on W_7 and W_8 induces a $\text{Spin}(7)$ -equivariant map

$$\begin{aligned}
(4.11) \quad \phi_\sigma &: \text{Lin}(W_8, W_7) \longrightarrow \text{Lin}(W_8, W_7) \\
\phi_\sigma(A)(a) &:= \sigma(a, A(e_\mu), e_\mu, e_i) e_i.
\end{aligned}$$

We calculate

$$\begin{aligned}
\lambda^4(u) \wedge \lambda^2(v) &= (e^0 \wedge (\iota_u *_{7} \varphi) - u^\flat \wedge \varphi) \wedge (e^0 \wedge v^\flat + \iota_v \varphi) \\
&= e^0 \wedge (\iota_u *_{7} \varphi) \wedge (\iota_v \varphi) + e^0 \wedge u^\flat \wedge v^\flat \wedge \varphi - u^\flat \wedge \varphi \wedge (\iota_v \varphi) \\
(4.12) \quad &\stackrel{(6.10), (6.11)}{=} 2u^\flat \wedge v^\flat \wedge (e^0 \wedge \varphi - *_{7} \varphi) - 2e^0 \wedge (*_{7}(u^\flat \wedge v^\flat)).
\end{aligned}$$

Lemma 4.3. *The map ϕ_σ has eigenvalues -1 and 6 with multiplicity 48 and 8 , respectively.*

Proof. Observe that the $\text{Spin}(7)$ -invariant inner products on W_7 and W_8 induce an inner product on $\text{Lin}(W_8, W_7) = W_8^* \otimes W_7$ for which $(e^\mu \otimes e_i)$ is

an orthonormal basis whenever (e_μ) is an orthonormal basis of W_8 so that $V_7 = e_0^\perp$ is spanned by (e_i) . This induced inner product satisfies

$$\langle \phi_\sigma(e^\mu \otimes e_i), e^\nu \otimes e_j \rangle_{Lin(W_8, W_7)} = \langle \phi_\sigma(e^\mu \otimes e_i)(e_\nu), e_j \rangle_{W_7} = \sigma(e_\nu, e_i, e_\mu, e_j),$$

and since $\sigma(e_\mu, e_i, e_\mu, e_j) = 0$, it follows that the matrix representation of ϕ_σ w.r.t. the basis $(e^\mu \otimes e_i)$ has 0's on the diagonal, whence $tr(\phi_\sigma) = 0$. Furthermore, ϕ_σ is self-adjoint since $\sigma(a, u, b, v) = \sigma(b, v, a, u)$ by (4.10) and (4.12), whence has real eigenvalues.

Decomposing $Lin(W_8, W_7) \stackrel{(4.8)}{\cong} W_8 \oplus W_{48}$, (4.9) implies that the elements in the summand congruent to W_8 are given by the maps

$$A_a : W_8 \longrightarrow W_7, \quad A_a(b) := \langle (\iota_a \lambda^2(e_i))^\#, b \rangle e_i$$

for a fixed $a \in W_8$. In order to calculate $\phi_\sigma(A_a)$, observe that $Spin(7)$ acts transitively on the unit sphere, whence we may assume w.l.o.g. that $a = e_0$, so that

$$A_{e_0}(b) = \langle e_i, b \rangle e_i = pr_{e_0^\perp}(b),$$

where $pr_{e_0^\perp} : W_8 \rightarrow e_0^\perp = V_7$ is the orthogonal projection. Thus,

$$\begin{aligned} \phi_\sigma(A_{e_0})(b) &= \sigma(b, A_{e_0}(e_\mu), e_\mu, e_i) e_i = \sigma(b, e_j, e_j, e_i) e_i \\ &\stackrel{(4.12)}{=} * \left(b^\flat \wedge e^j \wedge \left(e^j \wedge e^i \wedge (e^0 \wedge \varphi - *_7 \varphi) - e^0 \wedge (*_7 e^{ji}) \right) \right) e_i \\ &= * \left(e^0 \wedge b^\flat \wedge e^j \wedge (*_7 e^{ij}) \right) e_i \\ &= (1 - \delta_{ij}) * (e^0 \wedge b^\flat \wedge *_7 e^i) e_i \\ &= 6 \langle b, e_i \rangle e_i = 6A_{e_0}(b), \end{aligned}$$

so that $\phi_\sigma(A_a) = 6A_a$ for all A_a . By Schur's lemma and since ϕ_σ is self-adjoint, $\phi_\sigma|_{W_{48}} = cId_{W_{48}}$ for some $c \in \mathbb{R}$, whence

$$0 = tr(\phi_\sigma) = 6 \dim W_8 + c \dim W_{48},$$

and from this, $c = -1$ and the lemma follows. \square

For a manifold with a $Spin(7)$ -structure (M, Φ) and induced metric $g = g_\Phi$, the covariant derivative $g_\Phi \nabla_v \Phi$ w.r.t. the Levi-Civita connection is contained in the infinitesimal orbit of $\mathfrak{so}(T_p M, g_p)$ [Bryant1987] and hence in $\Lambda_7^4 T^* M$. That is, there is a section $T \in \Omega^1(M, W_7(M)) = \Gamma(Lin(TM, W_7(M)))$ such that

$$(4.13) \quad \nabla_v \Phi = \lambda^4(T(v)) = e^0 \wedge (\iota_{T(v)} * \varphi) - (T(v))^\flat \wedge \varphi$$

with the map $\lambda^4 : W_7 \rightarrow \Lambda_7^4 T_p M$ from (4.6). In analogy to Definition 3.4, we use the following terminology.

Definition 4.4. Let (M, Φ) be a manifold with a $Spin(7)$ -structure. The section $T \in \Omega^1(M, W_7(M))$ for which (4.13) holds is called the *torsion endomorphism* of the $Spin(7)$ -structure.

4.3. The Frölicher-Nijenhuis brackets on a manifold with a $\text{Spin}(7)$ -structure. Recall the section $P = -\delta_g \Phi \in \Omega^3(M, TM)$ on a manifold with a $\text{Spin}(7)$ -structure (M, Φ) from Definition 4.1. We wish to relate its Frölicher-Nijenhuis bracket to its torsion. In order to do this, recall that $[P, P]^{FN} \in \Omega^6(M, TM)$.

Due to the decomposition $\Lambda^6 W_8 = \Lambda_7^6 W_8 \oplus \Lambda_{21}^6 W_8$ as a G_2 -module, we may decompose

$$\Omega^6(M, TM) = \Gamma(M, \Lambda_7^6 T^* M \otimes TM) \oplus \Gamma(M, \Lambda_{21}^6 T^* M \otimes TM),$$

and we denote the projections onto the two summands by π_7 and π_{21} , respectively.

Proposition 4.5. *Let (M, Φ) be a manifold with a $\text{Spin}(7)$ -structure with the torsion endomorphism $T \in \Omega^1(M, W_7(M))$ from (4.13), and let $P = -\delta_g \Phi \in \Omega^3(M, TM)$ be as before. Then for $p \in M$,*

$$(4.14) \quad \pi_7([P, P]_p^{FN}) = -\frac{2}{3} \Phi \wedge \lambda^2 \left((4T_p + \phi_\sigma(T_p))(e_\mu) \right) \otimes e_\mu.$$

In particular, $\pi_7([P, P]_p^{FN}) = 0$ iff $T_p = 0$.

Proof. By Proposition 2.2, $[P, P]_p^{FN} = \gamma_\mu \otimes e_\mu$, where $\gamma_\mu \in \Lambda^6 T_p^* M$ is given by

$$(4.15) \quad \gamma_\mu = 2((\iota_{e_\nu} \Phi) \wedge \iota_{e_\mu} (\nabla_{e_\nu} \Phi) + (\iota_{e_\mu} \iota_{e_\nu} \Phi) \wedge e^\rho \wedge (\iota_{e_\nu} \nabla_{e_\rho} \Phi)).$$

If we decompose $\gamma_\mu = \Phi \wedge \lambda^2(v_\mu) + \gamma_\mu^{21}$ with $\gamma_\mu^{21} \in \Lambda_{21}^6 T_p^* M$, then for any $v \in V_7 = e_0^\perp$ we have $\gamma_\mu^{21} \wedge \lambda^2(v) = 0$ by (4.7) and hence,

$$\begin{aligned} \gamma_\mu \wedge \lambda^2(v) &= \Phi \wedge \lambda^2(v_\mu) \wedge \lambda^2(v) \\ &= (e^0 \wedge \varphi + * \gamma \varphi) \wedge (e^0 \wedge v_\mu^b + \iota_{v_\mu} \varphi) \wedge (e^0 \wedge v^b + \iota_v \varphi) \\ &= e^0 \wedge \varphi \wedge (\iota_{v_\mu} \varphi) \wedge (\iota_v \varphi) + * \gamma \varphi \wedge e^0 \wedge v_\mu^b \wedge (\iota_v \varphi) \\ &\quad + * \gamma \varphi \wedge (\iota_{v_\mu} \varphi) \wedge e^0 \wedge v^b \\ &\stackrel{(6.3), (6.1)}{=} 6 \langle v_\mu, v \rangle \text{vol} + 3 \langle v_\mu, v \rangle \text{vol} + 3 \langle v_\mu, v \rangle \text{vol} = 12 \langle v_\mu, v \rangle \text{vol}. \end{aligned}$$

Thus,

$$(4.16) \quad \pi_7([P, P]_p^{FN}) = \frac{1}{12} * (\gamma_\mu \wedge \lambda^2(e_i)) \Phi \wedge \lambda^2(e_i) \otimes e_\mu.$$

For arbitrary $v \in V_7 = e_0^\perp$ we compute

$$\begin{aligned}
\gamma_\mu \wedge \lambda^2(v) &= 2((\iota_{e_\nu} \Phi) \wedge \iota_{e_\mu} (\nabla_{e_\nu} \Phi) \wedge \lambda^2(v)) \\
&\quad + (\iota_{e_\mu} \iota_{e_\nu} \Phi) \wedge e^\rho \wedge (\iota_{e_\nu} \nabla_{e_\rho} \Phi) \wedge \lambda^2(v)) \\
&= 2((\iota_{e_\nu} \Phi) \wedge \iota_{e_\mu} \lambda^4(T_p(e_\nu)) \wedge \lambda^2(v)) \\
&\quad + (\iota_{e_\mu} \iota_{e_\nu} \Phi) \wedge e^\rho \wedge (\iota_{e_\nu} \lambda^4(T_p(e_\rho))) \wedge \lambda^2(v)) \\
&\stackrel{(6.17), (6.18)}{=} 2((-4\delta_{\mu\nu} \langle T(e_\nu), v \rangle \text{vol} + e^{\nu\mu} \wedge \lambda^4(T_p(e_\nu)) \wedge \lambda^2(v)) \\
&\quad + (-12\delta_{\mu\rho} \langle T_p(e_\rho), v \rangle \text{vol} + e^{\rho\mu} \wedge \lambda^4(T_p(e_\rho)) \wedge \lambda^2(v)) \\
&= 2(-16 \langle T_p(e_\mu), v \rangle \text{vol} - 2e^{\mu\nu} \wedge \lambda^4(T_p(e_\nu)) \wedge \lambda^2(v)) \\
&= 2\left(-16 \langle T_p(e_\mu), v \rangle - 4\sigma(e_\mu, T_p(e_\nu), e_\nu, v)\right) \text{vol} \\
&\stackrel{(4.11)}{=} -8 \langle (4T_p + \phi_\sigma(T_p))(e_\mu), v \rangle \text{vol},
\end{aligned}$$

and this together with (4.16) implies (4.14) and completes the proof. \square

With this, we are now ready to prove the following which immediately implies Theorem 1.2 from the introduction.

Theorem 4.6. *Let (M^8, Φ) be a manifold with a $\text{Spin}(7)$ -structure (M^8, Φ) , let ∇ be the Levi-Civita connection of $g = g_\Phi$, and let $T \in \Omega^1(M^8, W_7(M^8))$ be its torsion endomorphism defined in Definition 4.4. Then for every $p \in M^8$ the following are equivalent.*

- (1) $T_p = 0 \in T_p^* M^8 \otimes W^7(M)_p$.
- (2) The $\text{Spin}(7)$ -structure is torsion-free at p , i.e., $(\nabla \Phi)_p = 0$.
- (3) $\pi_7([P, P]_p^{FN}) = 0 \in \Lambda_7^6 T_p^* M^8 \otimes T_p M^8$.
- (4) $[P, P]_p^{FN} = 0 \in \Lambda^6 T_p^* M^8 \otimes T_p M^8$.

Proof. The equivalence of the first two statements was shown in [Fernandez1986]. Also, $T_p = 0$ implies $(\nabla \Phi)_p = 0$, whence by Proposition 2.2, $[P, P]_p^{FN} = 0$, and this trivially implies $\pi_7([P, P]_p^{FN}) = 0$.

By (4.14), $\pi_7([P, P]_p^{FN}) = 0$ iff $4T_p + \phi_\sigma(T_p) = 0$, and since ϕ_σ does not have -4 as an eigenvalue by Lemma 4.3, this implies that $T_p = 0$. \square

5. THE 16 CLASSES OF G_2 - AND 4 CLASSES OF $\text{Spin}(7)$ -STRUCTURES

In this section, we shall interpret the classification of G_2 -structures and of $\text{Spin}(7)$ -structures ([FG1982] and [Fernandez1986]) in terms of the Frölicher-Nijenhuis bracket.

For the G_2 -case, this classification is given by determining which components of the torsion endomorphism T vanish, where T is regarded as a section of the endomorphism bundle $\text{End}(TM^7)$ which is G_2 -equivariantly isomorphic to

$$(5.1) \quad V_7(M^7) \otimes V_7(M^7) \cong V_1(M^7) \oplus V_7(M^7) \oplus V_{14}(M^7) \oplus V_{27}(M^7).$$

Since this decomposition has 4 summands, the classification consists of $2^4 = 16$ cases.

Observe that both $\Lambda_7^5 T^* M^7 \otimes TM^7$ and $\Lambda^6 T^* M^7 \otimes TM^7$ are G_2 -equivariantly isomorphic to $V_7(M^7) \otimes V_7(M^7)$, where explicit isomorphisms are given by

$$\begin{aligned} K : \Lambda_7^5 T^* M^7 \otimes TM^7 &\ni (*\varphi \wedge v^b) \otimes w \longmapsto v^b \otimes w \in T^* M^7 \otimes TM^7 \\ L : \Lambda^6 T_p^* M^7 \otimes TM^7 &\ni (*v^b) \otimes w \longmapsto v^b \otimes w \in T^* M^7 \otimes TM^7. \end{aligned}$$

If (M^7, φ) is a manifold with a G_2 -structure and the cross products Cr and χ , then we define the sections

$$K_{\pi_7([Cr, \chi]^{FN})}, L_{[\chi, \chi]^{FN}} \in \Gamma(\text{End}(TM)).$$

Therefore, by Propositions 3.5, 3.6 there are G_2 -equivariant vector bundle isomorphisms

$$\tau_1, \tau_2 : \text{End}(TM^7) \longrightarrow \text{End}(TM^7)$$

such that for the torsion endomorphism $T \in \Gamma(\text{End}(TM))$ we have

$$(5.2) \quad \tau_1(T) = K_{\pi_7([Cr, \chi]^{FN})} \quad \text{and} \quad \tau_2(T) = L_{[\chi, \chi]^{FN}},$$

where by a slight abuse of notation we denote the map $\tau_i : \Gamma(\text{End}(TM)) \rightarrow \Gamma(\text{End}(TM))$ applying τ_i pointwise by the same symbol.

For an element $A = a_{ij} e^i \otimes e_j \in \text{End}(V_7)$ let us denote its skew-symmetrization by

$$\sigma_A := a_{ij} e^{ij} \in \Lambda^2 V_7^*.$$

With this notation, it follows from (3.10) and (3.18) that τ_1 and τ_2 take the form

$$\begin{aligned} \tau_1(T) &= 2(T - 2T^\top - \text{tr}(T)id), \\ \tau_2(T) &= -4(T + T^\top) + 6 * (e^i \wedge \sigma_T \wedge \varphi) \otimes e_i, \end{aligned}$$

summing over some basis (e_i) with dual basis (e^i) .

The G_2 -equivariance of τ_1 and τ_2 and (5.2) now implies that the $V_k(M)$ -component of T vanishes if and only if the $V_k(M)$ -component of $K_{\pi_7([Cr, \chi]^{FN})}$ vanishes if and only if the $V_k(M)$ -component of $L_{[\chi, \chi]^{FN}}$ vanishes. Since the cases in the Fernandez-Gray classification are determined by the vanishing of the components of T , we obtain the interpretation of these cases given in Table 1.

The interpretation of manifolds (M^8, Φ) with a Spin(7)-structure is analogous. Again, the torsion T and the projection $\pi_7([P, P]^{FN})$ are sections of the Spin(7)-equivariantly isomorphic bundles $T^* M^8 \otimes W_7(M^8)$ and $\Lambda_7^6 T^* M^8 \otimes TM^8$, respectively, with an explicit identification given by

$$H : \Lambda_7^6 T^* M^8 \otimes W_7(M^8) \ni \Phi \wedge (\lambda^2(a)) \otimes v \longmapsto v^b \otimes a \in T^* M^8 \otimes W_7(M^8),$$

TABLE 1.

Classes	Relation on $A \in \{T, K_{\pi_7([Cr, \chi]^{FN})}, L_{[\chi, \chi]^{FN}}\}$
$V_1(M) \oplus V_7(M) \oplus V_{14}(M) \oplus V_{27}(M)$	no relation
$V_7(M) \oplus V_{14}(M) \oplus V_{27}(M)$	$tr(A) = 0$
$V_1(M) \oplus V_{14}(M) \oplus V_{27}(M)$	$\sigma_A \in \Omega_{14}^2(M)$
$V_1(M) \oplus V_7(M) \oplus V_{27}(M)$	$\sigma_A \in \Omega_7^2(M)$
$V_1(M) \oplus V_7(M) \oplus V_{14}(M)$	$A + A^\top - \frac{2}{7}tr(A)id_{TM} = 0$
$V_{14}(M) \oplus V_{27}(M)$	$\sigma_A \in \Omega_{14}^2(M), \quad tr(A) = 0$
$V_7(M) \oplus V_{27}(M)$	$\sigma_A \in \Omega_7^2(M), \quad tr(A) = 0$
$V_7(M) \oplus V_{14}(M)$	$A + A^\top = 0$
$V_1(M) \oplus V_{27}(M)$	$A - A^\top = 0$
$V_1(M) \oplus V_{14}(M)$	$A + A^\top - \frac{2}{7}tr(A)id_{TM} = 0, \quad \sigma_A \in \Omega_{14}^2(M)$
$V_1(M) \oplus V_7(M)$	$A + A^\top - \frac{2}{7}tr(A)id_{TM} = 0, \quad \sigma_A \in \Omega_7^2(M)$
$V_{27}(M)$	$A - A^\top = 0, \quad tr(A) = 0$
$V_{14}(M)$	$A + A^\top = 0, \quad \sigma_A \in \Omega_{14}^2(M)$
$V_7(M)$	$A + A^\top = 0, \quad \sigma_A \in \Omega_7^2(M)$
$V_1(M)$	$A = \frac{1}{7}tr(A)id_{TM}$
$\{0\}$	$A = 0$

and if (M^8, Φ) is a manifold with a Spin(7)-structure and the 3-fold product P , then by (4.14)

$$H_{\pi_7([P, P]^{FN})} = \tau_3(T), \quad \text{where} \quad \tau_3(T) = -\frac{2}{3}(4T + \phi_\sigma(T)).$$

Here, by abuse of notation we regard ϕ_σ as the pointwise application of the map from (4.11) to sections of $W_7(M^8) \otimes T^*M^8 \cong Lin(W_8(M^8), W_7(M^8))$.

By (4.8), $W_7(M^8) \otimes T^*M^8$ can be decomposed as $W_8(M^8) \oplus W_{48}(M^8)$ whence by the Spin(7)-equivariance of τ_3 , the $W_k(M^8)$ -component of T vanishes if and only if the $W_k(M^8)$ -component of $H_{\pi_7([P, P]^{FN})}$ does. Since the classification of Fernández [Fernandez1986] into $2^2 = 4$ different cases is given by the vanishing of the components of the torsion T , it follows that these cases can be also interpreted by the vanishing of the components of $H_{\pi_7([P, P]^{FN})}$, which leads to the interpretation of the classes of Spin(7)-manifolds given in Table 2, where $pr_k : W_7(M^8) \otimes T^*M^8 \rightarrow W_k(M^8)$ is the canonical projection.

6. APPENDIX

In this appendix, we shall collect some of the formulas which we needed in the calculations in this paper. Most of them are known and can be found in a similar form e.g. in [SW2017, Lemma 4.37], but we shall collect them here for the reader's convenience.

TABLE 2.

Classes	Relation on $A \in \{T, H_{\pi_7}(\{P, P\}^{FN})\}$
$W_8 \oplus W_{48}$	no relation
W_{48}	$pr_8(A) = 0$
W_8	$pr_{48}(A) = 0$
$\{0\}$	$A = 0$

Lemma 6.1. *For all $u, v, w, r \in V_7$ and any orthonormal basis (e_i) of V_7 the following identities hold.*

$$(6.1) \quad u^\flat \wedge (\iota_v \varphi) \wedge * \varphi = 3 \langle u, v \rangle \text{vol}$$

$$(6.2) \quad u^\flat \wedge (\iota_w * \varphi) \wedge \varphi = 4 \langle u, w \rangle \text{vol}$$

$$(6.3) \quad (\iota_u \varphi) \wedge (\iota_v \varphi) \wedge \varphi = 6 \langle u, v \rangle \text{vol}$$

$$(6.4) \quad u^\flat \wedge (\iota_v \varphi) \wedge (\iota_w \varphi) \wedge (\iota_r \varphi) = 2 \left(\langle v, w \rangle \langle u, r \rangle \right. \\ \left. + \langle u, v \rangle \langle w, r \rangle + \langle u, w \rangle \langle v, r \rangle \right) \text{vol}$$

$$(6.5) \quad (\iota_u \iota_v * \varphi) \wedge w^\flat \wedge * \varphi = -2u^\flat \wedge v^\flat \wedge w^\flat \wedge * \varphi$$

$$(6.6) \quad (\iota_u \iota_v \varphi) \wedge w^\flat \wedge (\iota_r \varphi) \wedge \varphi = 2(\langle v, w \rangle \langle u, r \rangle - \langle u, w \rangle \langle v, r \rangle) \text{vol} \\ - 2u^\flat \wedge v^\flat \wedge w^\flat \wedge r^\flat \wedge \varphi$$

$$(6.7) \quad (\iota_u \iota_v * \varphi) \wedge (\iota_w * \varphi) \wedge (\iota_r \varphi) = 2(\langle v, w \rangle \langle u, r \rangle - \langle u, w \rangle \langle v, r \rangle) \text{vol} \\ + u^\flat \wedge v^\flat \wedge w^\flat \wedge r^\flat \wedge \varphi$$

$$(6.8) \quad u^\flat \wedge v^\flat \wedge (\iota_w * \varphi) \wedge (\iota_r \varphi) = 2(\langle v, w \rangle \langle u, r \rangle - \langle u, w \rangle \langle v, r \rangle) \text{vol} \\ + u^\flat \wedge v^\flat \wedge w^\flat \wedge r^\flat \wedge \varphi$$

$$(6.9) \quad u^\flat \wedge v^\flat \wedge (\iota_w \iota_r * \varphi) \wedge \varphi = 2(\langle v, w \rangle \langle u, r \rangle - \langle u, w \rangle \langle v, r \rangle) \text{vol} \\ - u^\flat \wedge v^\flat \wedge w^\flat \wedge r^\flat \wedge \varphi$$

$$(6.10) \quad (\iota_u * \varphi) \wedge (\iota_v \varphi) = -2 * (u^\flat \wedge v^\flat) + u^\flat \wedge v^\flat \wedge \varphi$$

$$(6.11) \quad \varphi \wedge (\iota_u \varphi) = 2u^\flat \wedge * \varphi$$

$$(6.12) \quad (\iota_{e_i} \varphi)^2 = 6 * \varphi$$

$$(6.13) \quad (\iota_u \iota_{e_i} * \varphi) \wedge (\iota_v \iota_{e_i} * \varphi) = 2(\iota_u \varphi) \wedge (\iota_v \varphi)$$

$$(6.14) \quad (\iota_u \iota_{e_i} * \varphi) \wedge (\iota_{e_i} \varphi) = 3u^\flat \wedge \varphi$$

Proof. For the proof of these identities, observe that the left hand side of each equation is a G_2 -invariant element of some tensor power of V_7 , and therefore it has to be a linear combination of the summands on the right hand side; the coefficients of this linear combination then can be determined by using the explicit formulas for φ and $*\varphi$ in (3.1) and (3.2).

To pick one explicit example which is not among the identities shown in [SW2017], observe that the left hand side of (6.4) is an element of $(V_7 \otimes \odot^3 V_7)^{G_2}$. Since $\odot^3 V_7 \cong V_7 \oplus V_{77}$, we have $\dim(V_7 \otimes \odot^3 V_7)^{G_2} = 1$, and there is one G_2 -invariant element of $V_7 \otimes \odot^3 V_7$ given by deriving the square of the scalar product which lies in $\odot^4 V_7$. Thus,

$$(6.15) \quad u^b \wedge (\iota_v \varphi) \wedge (\iota_w \varphi) \wedge (\iota_r \varphi) = c \left(\langle v, w \rangle \langle u, r \rangle + \langle u, v \rangle \langle w, r \rangle + \langle u, w \rangle \langle v, r \rangle \right) \text{vol.}$$

for some constant $c \in \mathbb{R}$. Now setting $u = v = w = r =: e_1$ and using (3.1) implies that $c = 2$, showing (6.4).

The remaining identities are shown in a similar fashion. \square

The following two decompositions of G_2 - and $\text{Spin}(7)$ -representations is also well known, cf. [SW2017, Theorem 8.5, 9.8], [Kar2005, (4.7), (4.8)].

Lemma 6.2. *Decompose $\Lambda^2 V_7^* = \Lambda_7^2 V_7^* \oplus \Lambda_{14}^2 V_7^*$ according to (3.5). Then*

$$\begin{aligned} \Lambda_7^2 V_7^* &= \{\alpha^2 \in \Lambda^2 V_7^* \mid *(\alpha^2 \wedge \varphi) = 2\alpha^2\}, \quad \text{and} \\ \Lambda_{14}^2 V_7^* &= \{\alpha^2 \in \Lambda^2 V_7^* \mid *(\alpha^2 \wedge \varphi) = -\alpha^2\}. \end{aligned}$$

In particular,

$$(6.16) \quad \begin{aligned} \Lambda_7^2 V_7^* &= \{\alpha^2 + *(\alpha^2 \wedge \varphi) \mid \alpha^2 \in \Lambda^2 V_7^*\}, \quad \text{and} \\ \Lambda_{14}^2 V_7^* &= \{2\alpha^2 - *(\alpha^2 \wedge \varphi) \mid \alpha^2 \in \Lambda^2 V_7^*\}. \end{aligned}$$

Lemma 6.3. *Decompose $\Lambda^2 W_8^* = \Lambda_7^2 W_8^* \oplus \Lambda_{21}^2 W_8^*$ according to (4.4). Then*

$$\begin{aligned} \Lambda_7^2 W_8^* &= \{\alpha^2 \in \Lambda^2 W_8^* \mid \Phi \wedge \alpha^2 = 3 * \alpha^2\}, \quad \text{and} \\ \Lambda_{21}^2 W_8^* &= \{\alpha^2 \in \Lambda^2 W_8^* \mid \Phi \wedge \alpha^2 = - * \alpha^2\}. \end{aligned}$$

We shall also need the following result.

Lemma 6.4. *For all $u, v \in W_7$ and $a, b \in W_8$ the following formulas hold:*

$$(6.17) \quad \begin{aligned} (\iota_a \Phi) \wedge (\iota_b \lambda^4(u)) \wedge \lambda^2(v) &= \\ -4 \langle a, b \rangle_{W_8} \langle u, v \rangle_{W_7} \text{vol} + a^b \wedge b^b \wedge \lambda^4(u) \wedge \lambda^2(v), \end{aligned}$$

$$(6.18) \quad \begin{aligned} a^b \wedge (\iota_b \iota_{e_\mu} \Phi) \wedge (\iota_{e_\mu} \lambda^4(u)) \wedge \lambda^2(v) &= \\ -12 \langle a, b \rangle_{W_8} \langle u, v \rangle_{W_7} \text{vol} + a^b \wedge b^b \wedge \lambda^4(u) \wedge \lambda^2(v) \end{aligned}$$

where in (6.18) the sum is taken over an orthonormal basis (e_μ) of W_8 .

Proof. By (4.4) and (4.5), the decomposition of $W_8 \otimes W_8$ and $W_7 \otimes W_7$ into $\text{Spin}(7)$ -irreducible summands yields

$$\begin{aligned} W_8 \otimes W_8 &= \odot^2 W_8 \oplus \Lambda^2 W_8 \cong (W_1 \oplus W_{35}) \oplus (W_7 \oplus W_{21}), \\ W_7 \otimes W_7 &= \odot^2 W_7 \oplus \Lambda^2 W_7 \cong (W_1 \oplus W_{27}) \oplus W_{21}, \end{aligned}$$

so that there are two inequivalent summands in common and hence, the space of $\text{Spin}(7)$ -invariant tensors in $W_8 \otimes W_8 \otimes W_7 \otimes W_7$ is 2-dimensional.

Since the left hand sides of (6.17) and (6.18) describe such tensors, it follows that there must be constants $c_1, \dots, c_4 \in \mathbb{R}$ such that

$$(6.19) \quad \begin{aligned} & (\iota_a \Phi) \wedge (\iota_b \lambda^4(u)) \wedge \lambda^2(v) = \\ & c_1 \langle a, b \rangle_{W_8} \langle u, v \rangle_{W_7} \text{vol} + c_2 a^b \wedge b^b \wedge \lambda^4(u) \wedge \lambda^2(v), \end{aligned}$$

$$(6.20) \quad \begin{aligned} & a^b \wedge (\iota_b \iota_{e_\mu} \Phi) \wedge (\iota_{e_\mu} \lambda^4(u)) \wedge \lambda^2(v) = \\ & c_3 \langle a, b \rangle_{W_8} \langle u, v \rangle_{W_7} \text{vol} + c_4 a^b \wedge b^b \wedge \lambda^4(u) \wedge \lambda^2(v). \end{aligned}$$

In order to determine these constants, we first let $a = b := e_0$, so that

$$(6.2) \quad \begin{aligned} (\iota_{e_0} \Phi) \wedge (\iota_{e_0} \lambda^4(u)) \wedge \lambda^2(v) &= \varphi \wedge (\iota_u * \varphi) \wedge (e^0 \wedge v^b + (\iota_v \varphi)) \\ &= -e^0 \wedge v^b \wedge (\iota_u * \varphi) \wedge \varphi \\ &\stackrel{(6.2)}{=} -4 \langle u, v \rangle \text{vol}, \end{aligned}$$

and from this, $c_1 = -4$ follows. Whence if we let $a := e_0$ and $b, u, v \in W_7 = e_0^\perp$ then

$$(6.5) \quad \begin{aligned} (\iota_{e_0} \Phi) \wedge (\iota_b \lambda^4(u)) \wedge \lambda^2(v) &\stackrel{(4.6)}{=} \varphi \wedge (-e^0 \wedge (\iota_b \iota_u * \varphi) - \iota_b (u^b \wedge \varphi)) \\ &\quad \wedge (e^0 \wedge v^b + (\iota_v \varphi)) \\ &= -\varphi \wedge e^0 \wedge (\iota_b \iota_u * \varphi) \wedge (\iota_v \varphi) \\ &\quad + \varphi \wedge u^b \wedge (\iota_b \varphi) \wedge e^0 \wedge v^b \\ &\stackrel{(6.11)}{=} 2e^0 \wedge (\iota_b \iota_u * \varphi) \wedge v^b \wedge * \varphi \\ &\quad + 2b^b \wedge * \varphi \wedge u^b \wedge e^0 \wedge v^b \\ &\stackrel{(6.5)}{=} -2e^0 \wedge b^b \wedge u^b \wedge v^b \wedge * \varphi \\ &\stackrel{(4.12)}{=} e^0 \wedge b^b \wedge \lambda^4(u) \wedge \lambda^2(v), \end{aligned}$$

so that $c_2 = 1$ follows. Now substituting $a = b := e_0$ and $u = v := e_1$ into (6.20) and using the index i to run from 1 to 7 yields

$$(6.3), (6.12) \quad \begin{aligned} e^0 \wedge (-\iota_{e_i} \varphi) \wedge (\iota_{e_i} (-e^1 \wedge \varphi)) \wedge (\iota_{e_1} \varphi) &= e^0 \wedge (\iota_{e_i} \varphi) \wedge (\delta_{1i} \varphi - e^1 \wedge (\iota_{e_i} \varphi)) \wedge (\iota_{e_1} \varphi) \\ &= e^0 \wedge (\iota_{e_1} \varphi) \wedge (\iota_{e_1} \varphi) \wedge \varphi \\ &\quad - e^0 \wedge e^1 \wedge (\iota_{e_i} \varphi) \wedge (\iota_{e_i} \varphi) \wedge \iota_{e_1} \varphi \\ &\stackrel{(6.1)}{=} 6 \text{vol} - 6e^0 \wedge e^1 \wedge \iota_{e_1} \varphi \wedge * \varphi \\ &\stackrel{(6.1)}{=} -12 \text{vol}, \end{aligned}$$

so that $c_3 = -12$ follows. Finally, for $a := e_0$, $u := e_1$ and $b, v \in W_7 = e_0^\perp$ (6.20) reads

$$\begin{aligned}
& e^0 \wedge (\iota_b \iota_{e_\mu} \Phi) \wedge (\iota_{e_\mu} \lambda^4(e_1)) \wedge \lambda^2(v) \\
&= e^0 \wedge (\iota_b \varphi) \wedge (\iota_{e_1} * \varphi) \wedge (\iota_v \varphi) \\
&\quad + e^0 \wedge (\iota_b \iota_{e_i} * \varphi) \wedge (-\iota_{e_i}(e^1 \wedge \varphi)) \wedge (\iota_v \varphi) \\
&\stackrel{(*)}{=} 0 + e^0 \wedge (\iota_b \iota_{e_i} * \varphi) \wedge (-\delta_{1i} \varphi + e^1 \wedge (\iota_{e_i} \varphi)) \wedge (\iota_v \varphi) \\
&= -e^0 \wedge (\iota_b \iota_{e_1} * \varphi) \wedge \varphi \wedge (\iota_v \varphi) \\
&\quad + e^{01} \wedge (\iota_b \iota_{e_i} * \varphi) \wedge (\iota_{e_i} \varphi) \wedge (\iota_v \varphi) \\
(6.11), (6.14) \quad &\stackrel{=}{=} -2e^0 \wedge (\iota_b \iota_{e_1} * \varphi) \wedge v^b \wedge * \varphi + 3e^{01} \wedge b^b \wedge \varphi \wedge (\iota_v \varphi) \\
(6.11), (6.5) \quad &\stackrel{=}{=} 4e^0 \wedge b^b \wedge e^1 \wedge v^b \wedge * \varphi + 6e^{01} \wedge b^b \wedge v^b \wedge * \varphi \\
&= -2e^0 \wedge b^b \wedge e^1 \wedge v^b \wedge * \varphi \\
(4.12) \quad &\stackrel{=}{=} e^0 \wedge b^b \wedge \lambda^4(e_1) \wedge \lambda^2(v),
\end{aligned}$$

so that $c_4 = 1$ follows. At $(*)$ we have used that the map

$$(u, v, w) \longmapsto * \left((\iota_u \varphi) \wedge (\iota_v * \varphi) \wedge (\iota_w \varphi) \right)$$

is a G_2 -invariant element of $W_7 \otimes \odot^2 W_7$, and since by (4.5) $\odot^2 W_7 \cong W_1 \oplus W_{27}$ has no irreducible component isomorphic to W_7 , there is no such element other than 0. \square

Acknowledgement. A part of this project has been discussed during HVL's visit to the Osaka City University in December 2015. She thanks Professor Ohnita for his invitation to Osaka and his hospitality. HVL and LS also thank the Max Planck Institute for Mathematics in the Sciences in Leipzig for its hospitality during extended visits. We also thank the referee for many helpful comments which enabled us to significantly improve the manuscript.

REFERENCES

- [BG1967] R. B. BROWN AND A. GRAY, Vector cross product, *Comment. Math. Helv.* 42(1967), 222-236.
- [Bryant1987] R. BRYANT, Metrics with exceptional holonomy, *Ann. of Math.* 126(1987), 525-576.
- [Bryant2005] R. BRYANT, Some remarks on G_2 -structures, *Proceedings of Gökova Geometry-Topology Conference 2005*, 75-109, *Gökova Geometry/Topology Conference (GGT)*, Gökova, 2006.
- [Fernandez1986] M. FERNANDEZ, A Classification of Riemannian Manifolds with Structure Group $\text{Spin}(7)$. *Annali di Mat. Pura Appl.* 143(1986), pp. 101-122.
- [FG1982] M. FERNÁNDEZ AND A. GRAY, Riemannian manifolds with structure group G_2 , *Annali di Mat. Pura Appl.* 32 (1982), 19-45.
- [FN1956] A. FRÖLICHER, A. NIJENHUIS, Theory of vector-valued differential forms. I. Derivations of the graded ring of differential forms., *Indag. Math.* 18 (1956), 338-359.

- [FN1956b] A. FRÖLICHER, A. NIJENHUIS, Some new cohomology invariants for complex manifolds. I, II, Indag. Math. 18 (1956), 540-552, 553-564.
- [HL1982] R. HARVEY AND H. B. LAWSON, Calibrated geometry, Acta Math. 148(1982), 47-157.
- [Humphreys1978] J. HUMPHREYS, Introduction to Lie algebras and representation theory, second printing, Graduate Texts in Mathematics, 9. Springer-Verlag, New York-Berlin, 1978.
- [Kar2005] S. KARIGIANNIS, Deformations of G_2 and Spin(7)-structures, Can. J. Math. 57(2005), 1012-1055.
- [Kawai2014] K. KAWAI, Deformations of homogeneous associative submanifolds in nearly parallel G_2 -manifolds, to appear in Asian J. Math., arXiv:1407.8046.
- [KMS1993] I. KOLAR, P. W. MICHOR AND J. SLOVAK, Natural operators in differential geometry, Springer 1993.
- [LV2016] H. V. LÊ AND J. VANŽURA, McLean's second variation formula revisited, J. Geom. Phys. 113(2017), 188-196, arXiv:1605.01267.
- [McLean1998] R. MCLEAN, Deformations of Calibrated Submanifolds, Comm. in Analysis and Geom. 6 (1998), 705-747.
- [SW2017] D. SALAMON, T. WALPUSKI, Notes on the octonions, arXiv: 1005.2820.

GRADUATE SCHOOL OF MATHEMATICAL SCIENCES, UNIVERSITY OF TOKYO, 3-8-1, KOMABA, MEGURO, TOKYO 153-8914, JAPAN

E-mail address: `kkawai@ms.u-tokyo.ac.jp`

CURRENT ADDRESS: GAKUSHUIN UNIVERSITY, 1-5-1, MEJIRO, TOSHIMA, TOKYO, 171-8588, JAPAN

E-mail address: `kkawai@math.gakushuin.ac.jp`

INSTITUTE OF MATHEMATICS CAS , ŽITNA 25, 11567 PRAHA 1, CZECH REPUBLIC

E-mail address: `hvle@math.cas.cz`

FAKULTÄT FÜR MATHEMATIK, TECHNISCHE UNIVERSITÄT DORTMUND, VOGELPOTH-SWEG 87, 44221 DORTMUND, GERMANY

E-mail address: `lschwach@math.tu-dortmund.de`