

Intelligent Initialization and Adaptive Thresholding for Iterative Matrix Completion; Some Statistical and Algorithmic Theory for *Adaptive-Impute*

Juhee Cho, Donggyu Kim, and Karl Rohe*

Department of Statistics, University of Wisconsin-Madison

September 15, 2018

Abstract

Over the past decade, various matrix completion algorithms have been developed. Thresholded singular value decomposition (SVD) is a popular technique in implementing many of them. A sizable number of studies have shown its theoretical and empirical excellence, but choosing the right threshold level still remains as a key empirical difficulty. This paper proposes a novel matrix completion algorithm which iterates thresholded SVD with theoretically-justified and data-dependent values of thresholding parameters. The estimate of the proposed algorithm enjoys the minimax error rate and shows outstanding empirical performances. The thresholding scheme that we use can be viewed as a solution to a non-convex optimization problem, understanding of whose theoretical convergence guarantee is known to be limited. We investigate this problem by introducing a simpler algorithm, generalized-*softImpute*, analyzing its convergence behavior, and connecting it to the proposed algorithm.

Keywords: *softImpute*, generalized-*softImpute*, non-convex optimization, thresholded singular value decomposition

*This research is supported by NSF grant DMS-1309998 and ARO grant W911NF-15-1-0423.

1 Introduction

Matrix completion appears in a variety of areas where it recovers a low-rank or approximately low-rank matrix from a small fraction of observed entries such as collaborative filtering (Rennie and Srebro (2005)), computer vision (Weinberger and Saul (2006)), positioning (Montanari and Oh (2010)), and recommender systems (Bennett and Lanning (2007)). Early work in this field was done by Achlioptas and McSherry (2001), Azar et al. (2001), Fazel (2002), Srebro et al. (2004), and Rennie and Srebro (2005). Later, Candès and Recht (2009) introduced the technique of matrix completion by minimizing the nuclear norm under convex constraints. This opened up a significant overlap with compressed sensing (Candès et al. (2006), Donoho (2006)) and led to accelerated research in matrix completion. They and others (Candès and Recht (2009), Candès and Tao (2010), Keshavan et al. (2010), Gross (2011), Recht (2011)) showed that the technique can exactly recover a low-rank matrix in the noiseless case. Many of the following works showed the approximate recovery of the low-rank matrix with the presence of noise (Candès and Plan (2010), Negahban and Wainwright (2011), Koltchinskii et al. (2011), Rohde and Tsybakov (2011)). Several other papers studied matrix completion in various settings (e.g. Davenport et al. (2014), Negahban and Wainwright (2012)) and proposed different estimation procedures of matrix completion (Srebro et al. (2004), Keshavan et al. (2009), Koltchinskii (2011), Cai and Zhou (2013), Chatterjee (2014)) than the ones by Candès and Recht (2009). In addition to the theoretical advances, a large number of algorithms have emerged (e.g. Rennie and Srebro (2005), Cai et al. (2010), Keshavan et al. (2009), Mazumder et al. (2010), Hastie et al. (2014)). An overview is well summarized in Mazumder et al. (2010) and Hastie et al. (2014).

Many of matrix completion algorithms employ thresholded singular value decomposition (SVD) which soft- or hard- thresholds the singular values. The statistical literature has responded by investigating its theoretical optimality and strong empirical performances. However, a key empirical difficulty of employing thresholded SVD for matrix completion is to find the right way and level of threshold. Depending on the choice of the thresholding scheme, the rank of the estimated low-rank matrix and predicted values for unobserved entries can widely change. Despite its importance, we lack understanding on how to choose

the threshold level and what bias or error we eliminate by thresholding.

We propose a novel iterative matrix completion algorithm, *Adaptive-Impute*, which recovers the underlying low-rank matrix from a few noisy entries via differentially and adaptively thresholded SVD. Specifically, the proposed *Adaptive-Impute* algorithm differentially thresholds the singular values and adaptively updates the threshold levels on every iteration. As was the case with adaptive Lasso (Zou (2006)) and adaptive thresholding for sparse covariance matrix estimation (Cai and Liu (2011)), the proposed thresholding scheme gives *Adaptive-Impute* stronger empirical performances than the thresholding scheme that uses a single thresholding parameter for all singular values throughout the iterations (e.g. *softImpute* (Mazumder et al. (2010))). Although *Adaptive-Impute* employs multiple thresholding parameters changing over iterations, we suggest specified values for the thresholding parameters that are theoretically-justified and data-dependent. Hence, *Adaptive-Impute* is free of the tuning problems associated with the choice of threshold levels. Its single tuning parameter is the rank of the resulting estimator. We suggest a way to choose the rank based on singular value gaps (for details, see Section 5.2). This novel threshold scheme of *Adaptive-Impute* makes its estimation via non-convex optimization, understanding of whose theoretical guarantees is known to be limited. However, to solve this problem and help understand the convergence behavior of *Adaptive-Impute*, we introduce a simpler algorithm than *Adaptive-Impute*, *generalized-softImpute*, and derive a sufficient condition under which it converges. Then, we prove that *Adaptive-Impute* behaves almost the same as *generalized-softImpute*. Numerical experiments and a real data analysis in Section 5 suggest superior performances of *Adaptive-Impute* over the existing *softImpute*-type algorithms.

The rest of this paper is organized as follows. Section 2 describes the model setup. Section 3 introduces the proposed algorithm *Adaptive-Impute*. Section 4 introduces a *generalized-softImpute*, a simpler algorithm than *Adaptive-Impute*. Section 5 presents numerical experiment results. Section 6 concludes the paper with discussion. All proofs are collected in Section 7.

2 The model setup

Suppose that we have an $n \times d$ matrix of rank r ,

$$M_0 = U\Lambda V^T, \quad (1)$$

where by SVD, $U = (U_1, \dots, U_r) \in \mathbb{R}^{n \times r}$, $V = (V_1, \dots, V_r) \in \mathbb{R}^{d \times r}$, $\Lambda = \text{diag}(\lambda_1, \dots, \lambda_r) \in \mathbb{R}^{r \times r}$, and $\lambda_1 \geq \dots \geq \lambda_r \geq 0$. The entries of M_0 are corrupted by noise $\epsilon \in \mathbb{R}^{n \times d}$ whose entries are i.i.d. sub-Gaussian random variables with mean zero and variance σ^2 . Hence, we can only observe $M_F = M_0 + \epsilon$. However, oftentimes in real world applications, not all entries of M_F are observable. So, define $y \in \mathbb{R}^{n \times d}$ such that $y_{ij} = 1$ if the (i, j) -th entry of M_F is observed and $y_{ij} = 0$ if it is not observed. The entries of y are assumed to be i.i.d. Bernoulli(p) and independent of the entries of ϵ . Then, the partially-observed noisy low-rank matrix $M \in \mathbb{R}^{n \times d}$ is written as

$$M_{ij} = y_{ij}M_{Fij} = \begin{cases} M_{0ij} + \epsilon_{ij} & \text{if observed } (y_{ij} = 1) \\ 0 & \text{otherwise } (y_{ij} = 0). \end{cases}$$

Throughout the paper, we assume that $r \ll d \leq n$ and the entries of M_0 are bounded by a positive constant L in absolute value. In this paper, we develop an iterative algorithm to recover M_0 from M and investigate its theoretical properties and empirical performances.

3 Adaptive-Impute algorithm

3.1 Initialization

We first introduce some notation. Let a set Ω contain indices of the observed entries, $y_{ij} = 1 \Leftrightarrow (i, j) \in \Omega$. Then, for any matrix $A \in \mathbb{R}^{n \times d}$, denote by $\mathcal{P}_\Omega(A)$ the projection of A onto Ω and by $\mathcal{P}_\Omega^\perp(A)$ the projection of A onto the complement of Ω ;

$$[\mathcal{P}_\Omega(A)]_{ij} = \begin{cases} A_{ij} & \text{if } (i, j) \in \Omega \\ 0 & \text{if } (i, j) \notin \Omega \end{cases} \quad \text{and} \quad [\mathcal{P}_\Omega^\perp(A)]_{ij} = \begin{cases} 0 & \text{if } (i, j) \in \Omega \\ A_{ij} & \text{if } (i, j) \notin \Omega. \end{cases}$$

That is, $\mathcal{P}_\Omega(A) + \mathcal{P}_\Omega^\perp(A) = A$. We let $\mathbf{u}_i(A)$ denote the i -th left singular vector of A , $\mathbf{v}_i(A)$ the i -th right singular vector of A , and $\lambda_i(A)$ the i -th singular value of A such that

$\lambda_1(A) \geq \dots \geq \lambda_d(A)$. The squared Frobenius norm is defined by $\|A\|_F^2 = \text{tr}(A^T A)$, the trace of $A^T A$, and the nuclear norm by $\|A\|_* = \sum_{i=1}^d \lambda_i(A)$, the sum of the singular values of A . For a symmetric matrix $A \in \mathbb{R}^{n \times n}$, $\text{diag}(A)$ represents a matrix with diagonal elements of A on the diagonal and zeros elsewhere.

Many of the iterative matrix completion algorithms (e.g. Cai et al. (2010), Mazumder et al. (2010), Keshavan et al. (2009), Chatterjee (2014)) in the current literature initialize with M , where the unobserved entries begin at zero. This initialization works well with algorithms that are based on convex optimization or that are robust to the initial. However, for algorithms that are based on non-convex optimization or that are sensitive to the initial, filling the unobserved entries with zeros may not be a good choice. Cho et al. (2016) proposed a one-step consistent estimator, \hat{M} , that attains the minimax error rate (Koltchinskii et al. (2011)), r/pd , and requires only two eigendecompositions. *Adaptive-Impute* employs the entries of this one-step consistent estimator instead of zeros as initial values of the unobserved entries. Algorithm 1 describes how to compute the initial \hat{M} of *Adaptive-Impute*. The following theorem shows that \hat{M} achieves the minimax error rate.

Algorithm 1 Initialization (Cho et al. (2016))

Require: M , y , and r

$$\hat{p} \leftarrow \frac{1}{nd} \sum_{i=1}^n \sum_{j=1}^d y_{ij}$$

$$\Sigma_{\hat{p}} \leftarrow M^T M - (1 - \hat{p}) \text{diag}(M^T M)$$

$$\Sigma_{t\hat{p}} \leftarrow M M^T - (1 - \hat{p}) \text{diag}(M M^T)$$

$$\hat{V}_i \leftarrow \mathbf{v}_i(\Sigma_{\hat{p}}), \quad \forall i \in \{1, \dots, r\}$$

$$\hat{U}_i \leftarrow \mathbf{u}_i(\Sigma_{t\hat{p}}), \quad \forall i \in \{1, \dots, r\}$$

$$\tilde{\alpha} \leftarrow \frac{1}{d-r} \sum_{i=r+1}^d \lambda_i(\Sigma_{\hat{p}})$$

$$\hat{\tau}_i \leftarrow \lambda_i(\Sigma_{\hat{p}}) - \frac{1}{\hat{p}} \sqrt{\lambda_i(\Sigma_{\hat{p}}) - \tilde{\alpha}}, \quad \forall i \in \{1, \dots, r\}$$

$$\hat{\lambda}_i \leftarrow \lambda_i(\Sigma_{\hat{p}}) - \hat{\tau}_i, \quad \forall i \in \{1, \dots, r\}$$

$$\hat{s} = (\hat{s}_1, \dots, \hat{s}_r) \leftarrow \arg \min_{s \in \{-1, 1\}^r} \left\| \mathcal{P}_\Omega \left(\sum_{i=1}^r s_i \hat{\lambda}_i \hat{U}_i \hat{V}_i^T - M \right) \right\|_F^2$$

$$\hat{M} \leftarrow \sum_{i=1}^r \hat{s}_i \hat{\lambda}_i \hat{U}_i \hat{V}_i^T$$

return \hat{M}

Assumption 1.

(1) $pd / \log n \rightarrow \infty$ and $n, d \rightarrow \infty$ with $d \leq n \leq e^{d^\beta}$, where $\beta < 1$ free of n , d , and p ;

(2) $\lambda_i = b_i \sqrt{nd}$ for all $i = 1, \dots, r$, where $\{b_i\}_{i=1, \dots, r}$ are positive bounded values;

(3) $b_i > b_{i+1}$ for all $i = 1, \dots, r$, where $b_{r+1} = 0$;

(4) $\lim_{n, d \rightarrow \infty} \mathbb{P} \left(\min_{s \in \{-1, 1\}^r} \|\mathcal{P}_\Omega(\sum_{i=1}^r s_i \hat{\lambda}_i \hat{U}_i \hat{V}_i^T - M)\|_F^2 < \|\mathcal{P}_\Omega(\sum_{i=1}^r s_{0i} \hat{\lambda}_i \hat{U}_i \hat{V}_i^T - M)\|_F^2 \right) = 0$,
 where $s = (s_1, \dots, s_r)$ and $s_{0i} = \text{sign}(\langle \hat{V}_i, V_i \rangle) \text{sign}(\langle \hat{U}_i, U_i \rangle)$ for $i = 1, \dots, r$.

Remark 1. Under the setting where the rank r is fixed as in this paper, Assumption 1(2) implies that the underlying low-rank matrix M_0 is dense. More specifically, note that the squared Frobenius norm indicates both the sum of all squared entries of a matrix and the sum of its singular values squared. Also, note that $\|M_0\|_F^2 = \sum_{i=1}^r \lambda_i^2(M_0) = cnd$ for some constant $c > 0$ by Assumption 1(2). Thus, the sum of all squared entries of M_0 has an order nd . This means that a non-vanishing proportion of entries of M_0 contains non-vanishing signals with dimensionality (see Fan et al. (2013)). For more discussion, see Remark 2 in Cho et al. (2016).

Remark 2. The singular vectors, $\{\hat{U}_i\}_{i=1}^r$ and $\{\hat{V}_i\}_{i=1}^r$, that compose \hat{M} are consistent estimators of U and V up to signs (for details, see Cho et al. (2016)). Hence, when combining them with $\{\hat{\lambda}_i\}_{i=1}^r$ to reconstruct \hat{M} , a sign problem happens. Assumption 1(4) assures that as n and d increase, the probability of choosing different signs than the true signs, $\{s_{0i}\}_{i=1}^r$, goes to zero. Given the asymptotic consistency of $\{\hat{U}_i\}_{i=1}^r$, $\{\hat{V}_i\}_{i=1}^r$, and $\{\hat{\lambda}_i\}_{i=1}^r$, this is not an unreasonable assumption to make.

Proposition 3.1. (Theorem 4.4 in Cho et al. (2016)) Under Assumption 1 and the model setup in Section 2, \hat{M} is a consistent estimator of M_0 . In particular,

$$\frac{1}{nd} \|\hat{M} - M_0\|_F^2 = o_p \left(\frac{h_n}{pd} \right),$$

where h_n diverges very slowly with the dimensionality, for example, $\log(\log d)$.

Remark 3. Since h_n in Proposition 3.1 can be any quantity that diverges slowly with the dimensionality, the convergence rate of \hat{M} can be thought of as $1/pd$. Under the setting where the rank of M_0 is fixed as in this paper, it is matched to the minimax error rate, r/pd , found in Koltchinskii et al. (2011).

Using \hat{M} to initialize *Adaptive-Impute* has two major advantages. First, since \hat{M} is already a consistent estimator of M_0 achieving the minimax error rate, it allows a series of the iterates of *Adaptive-Impute* coming after \hat{M} to be also consistent estimators of M_0 achieving the minimax error rate (see Theorem 3.1). Second, because *Adaptive-Impute* is based on a non-convex optimization problem (see Section 4), its convergence may depend on initial values. \hat{M} provides *Adaptive-Impute* a suitable initializer.

3.2 Adaptive thresholds

To motivate the novel thresholding scheme of *Adaptive-Impute*, we first consider the case where a fully-observed noisy low-rank matrix is available. Specifically, suppose that the probability of observing each entry, p , is 1 and thus $M_F = M_0 + \epsilon$ is observed. Under the model setup in Section 2 we can easily show that

$$\mathbb{E}(M_F^T M_F) = M_0^T M_0 + n\sigma^2 I_d \quad \text{and} \quad \mathbb{E}(M_F M_F^T) = M_0 M_0^T + d\sigma^2 I_n, \quad (2)$$

where I_d and I_n are identity matrices of size d and n , respectively. This shows that the eigenvectors of $\mathbb{E}(M_F^T M_F)$ and $\mathbb{E}(M_F M_F^T)$ are the same as the right and left singular vectors of M_0 . Also, the top r eigenvalues of $\mathbb{E}(M_F^T M_F)$ consist of the squared singular values of M_0 and a noise, $n\sigma^2$, the latter of which is the same as the average of the bottom $d - r$ eigenvalues of $\mathbb{E}(M_F^T M_F)$. In light of this, we want the estimator of M_0 based on M_F to keep the first r singular vectors of M_F as they are, but adjust the bias occurring in the singular values of M_F . Thus, the resulting estimator is

$$\hat{M}^F = \sum_{i=1}^r \sqrt{\lambda_i^2(M_F) - \alpha} \mathbf{u}_i(M_F) \mathbf{v}_i(M_F)^T, \quad \text{where } \alpha = \frac{1}{d-r} \sum_{i=r+1}^d \lambda_i^2(M_F). \quad (3)$$

A simple extension of Proposition 3.1 shows that \hat{M}^F achieves the best possible minimax error rate of convergence, $1/d$, since $p = 1$.

Now consider the cases where a partially-observed noisy low-rank matrix M is available. For each iteration $t \geq 1$, we fill out the unobserved entries of M with the corresponding entries of the previous iterate Z_t , treat the completed matrix $\widetilde{M}_t = \mathcal{P}_\Omega(M) + \mathcal{P}_\Omega^\perp(Z_t)$ as if it is a fully-observed matrix M_F , and find the next iterate Z_{t+1} in the same way that we

found \hat{M}^F from M_F in (3);

$$Z_{t+1} = \sum_{i=1}^r \sqrt{\lambda_i^2(\tilde{M}_t) - \tilde{\alpha}_t} \mathbf{u}_i(\tilde{M}_t) \mathbf{v}_i(\tilde{M}_t)^T, \quad \text{where } \tilde{\alpha}_t = \frac{1}{d-r} \sum_{i=r+1}^d \lambda_i^2(\tilde{M}_t). \quad (4)$$

Note that the difference in (4) from (3) is in the usage of \tilde{M}_t instead of M_F . Hence, the performance of *Adaptive-Impute* may depend on how close $\mathcal{P}_\Omega(Z_t)$ is to $\mathcal{P}_\Omega(M_0)$. Algorithm 2 summarizes these computing steps of *Adaptive-Impute* continued from Algorithm 1.

Algorithm 2 *Adaptive-Impute*

Require: M , y , r , and $\varepsilon > 0$

$Z_1 \leftarrow \hat{M}$ # from Algorithm 1

repeat for $t = 1, 2, \dots$

$\tilde{M}_t \leftarrow \mathcal{P}_\Omega(M) + \mathcal{P}_\Omega^\perp(Z_t)$

$V_i^{(t)} \leftarrow \mathbf{v}_i(\tilde{M}_t), \quad \forall i \in \{1, \dots, r\}$

$U_i^{(t)} \leftarrow \mathbf{u}_i(\tilde{M}_t), \quad \forall i \in \{1, \dots, r\}$

$\tilde{\alpha}_t \leftarrow \frac{1}{d-r} \sum_{i=r+1}^d \lambda_i^2(\tilde{M}_t)$

$\tau_{t,i} \leftarrow \lambda_i(\tilde{M}_t) - \sqrt{\lambda_i^2(\tilde{M}_t) - \tilde{\alpha}_t}, \quad \forall i \in \{1, \dots, r\}$ # Adaptive thresholds

$\lambda_i^{(t)} \leftarrow \lambda_i(\tilde{M}_t) - \tau_{t,i} \left(= \sqrt{\lambda_i^2(\tilde{M}_t) - \tilde{\alpha}_t} \right), \quad \forall i \in \{1, \dots, r\}$

$Z_{t+1} \leftarrow \sum_{i=1}^r \lambda_i^{(t)} U_i^{(t)} V_i^{(t)T}$

$t \leftarrow t + 1$

until $\|Z_{t+1} - Z_t\|_F^2 / \|Z_t\|_F^2 \leq \varepsilon$

return Z_{t+1}

The following theorem illustrates that the iterates of *Adaptive-Impute* retain the statistical performance of the initializer \hat{M} .

Assumption 2. For all $i = 1, \dots, r$, $\text{sign}(\langle \mathbf{u}_i(\tilde{M}_t), U_i \rangle) = \text{sign}(\langle \mathbf{v}_i(\tilde{M}_t), V_i \rangle)$.

Theorem 3.1. Under Assumptions 1-2 and the model setup in Section 2, we have for any fixed value of t ,

$$\frac{1}{nd} \|Z_t - M_0\|_F^2 = o_p\left(\frac{h_n}{pd}\right), \quad \text{as } n, d \rightarrow \infty \text{ with any } h_n \rightarrow \infty$$

where h_n diverges very slowly with the dimensionality, for example, $\log(\log d)$.

Remark 4. Similarly as in Remark 3, since h_n is a quantity diverging very slowly, the convergence rate of Z_t can be thought of as $1/pd$ which is matched to the minimax error rate, r/pd (Koltchinskii et al. (2011)).

3.3 Non-convexity of *Adaptive-Impute*

We can view *Adaptive-Impute* as an estimation method via non-convex optimization.

For $t \geq 1$, define

$$\tau_{t,i} = \begin{cases} \lambda_i(\widetilde{M}_t) - \sqrt{\lambda_i^2(\widetilde{M}_t) - \widetilde{\alpha}_t}, & i \leq r \\ \lambda_{r+1}(\widetilde{M}_t), & i > r \end{cases}, \quad (5)$$

where $\widetilde{\alpha}_t = \frac{1}{d-r} \sum_{i=r+1}^d \lambda_i^2(\widetilde{M}_t)$ and $\widetilde{M}_t = \mathcal{P}_\Omega(M) + \mathcal{P}_\Omega^\perp(Z_t)$. Then, in each iteration *Adaptive-Impute* provides a solution to the problem

$$\min_{Z \in \mathbb{R}^{n \times d}} \frac{1}{2nd} \|\widetilde{M}_t - Z\|_F^2 + \sum_{i=1}^d \frac{\tau_{t,i}}{\sqrt{nd}} \frac{\lambda_i(Z)}{\sqrt{nd}}. \quad (6)$$

Note that the threshold parameters, $\tau_{t,i}$, have dependence on both the i -th singular value and the t -th iteration. The following theorem provides an explicit solution to (6).

Theorem 3.2. *Let X be an $n \times d$ matrix and let $n \geq d$. The optimization problem*

$$\min_Z \frac{1}{2nd} \|X - Z\|_F^2 + \sum_{i=1}^d \frac{\tau_i}{\sqrt{nd}} \frac{\lambda_i(Z)}{\sqrt{nd}} \quad (7)$$

has a solution which is given by

$$\hat{Z} = \Phi(\Delta - \boldsymbol{\tau})_+ \Psi^T, \quad (8)$$

where $\Phi\Delta\Psi^T$ is the SVD of X , $\boldsymbol{\tau} = \text{diag}(\tau_1, \dots, \tau_d) \in \mathbb{R}^{d \times d}$, $(\Delta - \boldsymbol{\tau})_+ = \text{diag}((\lambda_1(X) - \tau_1)_+, \dots, (\lambda_d(X) - \tau_d)_+) \in \mathbb{R}^{d \times d}$, and $c_+ = \max(c, 0)$ for any $c \in \mathbb{R}$.

Remark 5. To see how Theorem 3.2 provides a solution to (6), let $X = \widetilde{M}_t$ and $\tau_i = \tau_{t,i}$ as specified in (5). Then, (6) and (7) become the same and \hat{Z} in (8) gives the explicit form of the $(t+1)$ -th iterate, Z_{t+1} , in Algorithm 2.

If all of the thresholding parameters in (6) are equal such that $\tau = \tau_{t,1} = \dots = \tau_{t,d}$ for all $1 \leq i \leq d$ and $t \geq 1$, the optimization problem (6) becomes equivalent to that of *softImpute* (Mazumder et al. (2010)) and Theorem 3.2 provides an iterative solution to it. While *softImpute* requires finding the right value of a thresholding parameter τ by using a cross validation (CV) technique which is time-consuming and often does not have a straightforward validation criteria, *Adaptive-Impute* suggests specific values of the thresholding levels as in (5). The novel thresholding scheme of *Adaptive-Impute* together with the rank constraint results in superior empirical performances over the existing *softImpute*-type algorithms (see Section 5).

The thresholding scheme of *Adaptive-Impute* can be viewed as a solution to a non-convex optimization problem since at every iteration it differentially and adaptively thresholds the singular values. As Hastie and others alluded to a similar issue for matrix completion methods via non-convex optimization in Hastie et al. (2014), it is hard to provide a direct convergence guarantee of *Adaptive-Impute*. So, in the following section we introduce a generalized-*softImpute* algorithm, simpler than *Adaptive-Impute* and yet still non-convex, and investigate its asymptotic convergence. It hints at the convergent behavior of *Adaptive-Impute* in the asymptotic sense.

4 Generalized *softImpute*

Generalized-*softImpute* is an algorithm which iteratively solves the problem,

$$\min_{Z \in \mathbb{R}^{n \times d}} Q_\tau(Z|Z_t^g) := \frac{1}{2nd} \|\mathcal{P}_\Omega(M) + \mathcal{P}_\Omega^\perp(Z_t^g) - Z\|_F^2 + \sum_{i=1}^d \frac{\tau_i}{\sqrt{nd}} \frac{\lambda_i(Z)}{\sqrt{nd}}, \quad (9)$$

to ultimately solve the optimization problem,

$$\min_{Z \in \mathbb{R}^{n \times d}} f_\tau(Z) := \frac{1}{2nd} \|\mathcal{P}_\Omega(M) - \mathcal{P}_\Omega(Z)\|_F^2 + \sum_{i=1}^d \frac{\tau_i}{\sqrt{nd}} \frac{\lambda_i(Z)}{\sqrt{nd}}. \quad (10)$$

Note that generalized-*softImpute* differentially penalizes the singular values, but the thresholding parameters do not change over iterations. The iterative solutions of generalized-*softImpute* are denoted by $Z_{t+1}^g := \arg \min_{Z \in \mathbb{R}^{n \times d}} Q_\tau(Z|Z_t^g)$ for $t \geq 1$ and Theorem 3.2 provides a closed form of Z_{t+1}^g . If $\tau_i = \tau$ for all $1 \leq i \leq d$, generalized-*softImpute* will

be equivalent to *softImpute* and both (9) and (10) become convex problems. However, by differentially penalizing the singular values, *generalized-softImpute* ends up solving a non-convex optimization problem. Theorem 4.1 below shows that despite the non-convexity of *generalized-softImpute*, the iterates of *generalized-softImpute*, $\{Z_t^g\}_{t \geq 1}$, converge to a solution of problem (10) under certain conditions.

Assumption 3. Let $\widetilde{M}_t^g = \mathcal{P}_\Omega(M) + \mathcal{P}_\Omega^\perp(Z_t^g)$ and $D_t^g := \widetilde{M}_t^g - Z_{t+1}^g$. Then,

$$\frac{1}{nd} \|D_t^g - D_{t+1}^g\|_F^2 + \frac{2}{nd} \langle D_t^g - D_{t+1}^g, Z_{t+1}^g - Z_{t+2}^g \rangle \geq 0 \quad \text{for all } t \geq 1.$$

Theorem 4.1. Let Z_∞ be a limit point of the sequence Z_t^g . Under Assumption 3, if the minimizer Z^s of (10) satisfies

$$Z^s \in \left\{ Z \in \mathbb{R}^{n \times d} : \sum_{i=1}^d \tau_i \lambda_i(Z) \geq \sum_{i=1}^d \tau_i \lambda_i(Z_\infty) + \langle (Z - Z_\infty), D_\infty \rangle \right\}, \quad (11)$$

we have $f_\tau(Z_\infty) = f_\tau(Z^s)$ and $\lim_{t \rightarrow \infty} f_\tau(Z_t^g) = f(Z^s)$.

Remark 6. If $\tau_i = \tau$ for all i as in case of *softImpute*, Assumption 3 and (11) are always satisfied because $\frac{1}{\tau} D_t^g$ belongs to the sub-gradient of $\|Z_{t+1}^g\|_*$.

Remark 7. If Z^s is unique, then *generalized-softImpute* finds the global minimum point of (10) by Theorem 4.1.

Generalized-softImpute resembles *Adaptive-Impute* in a sense that both of them employ different thresholding parameters on $\lambda_i(Z)$'s. However, *Adaptive-Impute* updates these tuning parameters every iteration while *generalized-softImpute* does not. The following lemmas show that despite this difference, the convergent behavior of *Adaptive-Impute* is asymptotically close to that of *generalized-softImpute*.

Lemma 4.1. Under Assumptions 1-2 and the model setup in Section 2, we have

$$\left| \frac{\tau_{t,i}}{\sqrt{nd}} - \frac{\tau_{t+1,i}}{\sqrt{nd}} \right| = o_p \left(\sqrt{\frac{h_n}{pd}} \right) \quad \text{for } i = 1, \dots, d,$$

where $\tau_{t,i}$ is defined in (5).

Lemma 4.2. Let $D_t := \widetilde{M}_t - Z_{t+1}$, where \widetilde{M}_t and Z_t are as defined in Algorithm 2. Then, under Assumptions 1-2 and the model setup in Section 2, we have

$$\frac{1}{nd} \|D_t - D_{t+1}\|_F^2 + \frac{2}{nd} \langle D_t - D_{t+1}, Z_{t+1} - Z_{t+2} \rangle + o_p\left(\frac{h_n}{pd}\right) \geq 0.$$

Lemma 4.1 shows that for large n and d , thresholding parameters of *Adaptive-Impute* are stable between iterations so that *Adaptive-Impute* behaves similarly to generalized-*softImpute*. Lemma 4.2 shows how Assumption 3 is adapted in *Adaptive-Impute*. It implies a possibility of *Adaptive-Impute* satisfying Assumption 3 asymptotically. Although this still does not provide a guarantee of convergence of *Adaptive-Impute*, numerical results below support this possibility.

5 Numerical results

In this section, we conducted simulations and a real-data analysis to compare *Adaptive-Impute* for estimating M_0 with the four different versions of *softImpute*:

1. *Adaptive-Impute*: the proposed algorithm, as summarized in Algorithm 2;
2. *softImpute*: the original *softImpute* algorithm (Mazumder et al. (2010));
3. *softImpute-Rank*: *softImpute* with rank restriction (Hastie et al. (2014));
4. *softImpute-ALS*: *Maximum-Margin Matrix Factorization* (Hastie et al. (2014));
5. *softImpute-ALS-Rank*: *rank-restricted Maximum-Margin Matrix Factorization* in Algorithm 3.1 (Hastie et al. (2014)).

SoftImpute algorithms were implemented with the R package, `softImpute` (Hastie and Mazumder (2015)). The R code for *Adaptive-Impute* is available at https://github.com/chojuhee/hello-world/blob/master/adaptiveImpute_Rfunction. In this R code, we made two adjustments from Algorithms 1 and 2 for technical reasons. First, in almost all real world applications that needed matrix completion, the entries of M_0 are bounded below and above by constants L_1 and L_2 such that

$$L_1 \leq M_{0ij} \leq L_2$$

and smaller or larger values than the constants do not make sense. So, after each iteration of *Adaptive-Impute*, $t \geq 1$, we replace the values of Z_t that are smaller than L_1 with L_1 and the values of Z_t that are greater than L_2 with L_2 . Second, the cardinality of the set, $\{-1, 1\}^r$, that we search over to find \hat{s} in Algorithm 1 increases exponentially. Hence, finding \hat{s} easily becomes a computational bottleneck of *Adaptive-Impute* or is even impossible for large r . We suggest two possible solutions to this problem. One solution is to find \hat{s} by computing $\hat{s}_i = \text{sign}(\langle \hat{V}_i, \mathbf{v}_i(M) \rangle) \text{sign}(\langle \hat{U}_i, \mathbf{u}_i(M) \rangle)$ for $i = 1, \dots, r$. Note that if we use V_i and U_i instead of $\mathbf{v}_i(M)$ and $\mathbf{u}_i(M)$, this gives us the true sign s_0 under Assumption 1. The other solution is to use a linear regression. Let a vector of the observed entries of M be the dependent variable and let a vector of the corresponding entries of $\hat{\lambda}_i \hat{U}_i \hat{V}_i^T$ be the i -th column of the design matrix for $i = 1, \dots, r$. Then, we set \hat{s} to be the coefficients of the regression line whose intercept is forced to be 0. The difference in the results of these two methods are negligible. In the following experiment, we only reported the results of the former solution for simplicity, while the R code provided in https://github.com/chojuhee/hello-world/blob/master/adaptiveImpute_Rfunction are written for both solutions.

5.1 Simulation study

To create $M_0 = AB^T \in \mathbb{R}^{n \times d}$, we sampled $A \in \mathbb{R}^{n \times r}$ and $B \in \mathbb{R}^{d \times r}$ to contain i.i.d. uniform $[-5, 5]$ random variables and a noise matrix $\epsilon \in \mathbb{R}^{n \times d}$ to contain i.i.d. $\mathcal{N}(0, \sigma^2)$. Then, each entry of $M_0 + \epsilon$ was observed independently with probability p . Across simulations, $n = 1700$, $d = 1000$, $r \in \{5, 10, 20, 50\}$, σ varies from 0.1 to 50, and p varies from 0.1 to 0.9. For each simulation setting, the data was sampled 100 times and the errors were averaged.

To evaluate performance of the algorithms, we measured three different types of errors; test, training, and total errors; the test error, $\text{Test}(\hat{M}) = \|\mathcal{P}_\Omega^\perp(\hat{M} - M_0)\|_F^2 / \|\mathcal{P}_\Omega^\perp(M_0)\|_F^2$, represents the distance between the estimate \hat{M} and the parameter M_0 measured on the unobserved entries, the training error, $\text{Training}(\hat{M}) = \|\mathcal{P}_\Omega(\hat{M} - M_0)\|_F^2 / \|\mathcal{P}_\Omega(M_0)\|_F^2$, the distance measured on the observed entries, and the total error, $\text{Total}(\hat{M}) = \|\hat{M} - M_0\|_F^2 / \|M_0\|_F^2$, the distance measured on all entries. For ease of comparison, Figure 1

and 3 plot the relative efficiencies with respect to *softImpute*-Rank. For example, the relative test efficiency of *Adaptive-Impute* with respect to *softImpute*-Rank is defined as $\text{Test}(\hat{M}_{rank})/\text{Test}(\hat{M}_{adapt})$, where \hat{M}_{adapt} is an estimate of *Adaptive-Impute* and \hat{M}_{rank} is an estimate of *softImpute-Rank*. The relative total and training efficiencies with respect to *softImpute*-Rank are defined similarly.

We used the best tuning parameter for the algorithms in comparison. Specifically, for algorithms with rank restriction (including *Adaptive-Impute*), we provided the true rank (i.e. 5, 10, 20, or 50). For *softImpute*-type algorithms, an oracle tuning parameter was chosen to minimize the total error.

Figure 1 shows the change of the relative efficiencies as the probability of observing each entry, p , increases with $\sigma = 1$. Three columns of plots in Figure 1 correspond to three different types of errors and four rows of plots to four different values of the rank. In all cases, *Adaptive-Impute* outperforms the competitors and works especially better when p is small. Among *softImpute*-type algorithms, the algorithms with rank constraint (i.e. *softImpute*-Rank and *softImpute*-ALS-Rank) perform better than the ones without (i.e. *softImpute* and *softImpute*-ALS). Figure 2 shows the change of the absolute errors that are used to compute relative efficiencies in Figure 1 as the probability of observing each entry, p , increases.

Figure 3 shows the change of the log relative efficiencies as the standard deviation (SD) of each entry of ϵ , σ , increases with $p = 0.1$. When the noise level is under 15, *Adaptive-Impute* outperforms the competitors, but when the noise level is over 15, *softImpute*-type algorithms start to outperform *Adaptive-Impute*. Hence, *softImpute*-type algorithms are more robust to large noises than *Adaptive-Impute*. It may be because when there exist large noises dominating the signals, the conditions for convergence presented in Section 4 are not satisfied. In real life applications, however, it is not common to observe such large noises that dominate the signals. Figure 4 shows the change of the absolute errors that are used to compute relative efficiencies in Figure 3.

Figure 5 shows convergence of the iterates of *Adaptive-Impute* to the underlying low-rank matrix over iterations; that is, the change of $\log \text{Total}(Z_t)$, $\text{Training}(Z_t)$, and $\text{Test}(Z_t)$ errors as t increases. Across all plots, $n = 1700$, $d = 1000$, $p = 0.1$, and the errors were




Figure 1: The relative efficiency plotted against the probability of observing each entry, p , when $\sigma = 1$. Training errors are measured over the observed entries, test errors over the unobserved entries, and total errors over all entries.




Figure 2: Change of the absolute errors when the probability of observing each entry, p increases and $\sigma = 1$.




Figure 3: The log relative efficiency plotted against the SD of each entry of ϵ when $p = 0.1$. Training errors are measured over the observed entries, test errors over the unobserved entries, and total errors over all entries.




Figure 4: Change of the absolute errors when the SD of each entry of ϵ increases and $p = 0.1$.

averaged over 100 replicates. In all cases, we observe that *Adaptive-Impute* converges well. Particularly, the smaller value of noise and/or rank is, the faster *Adaptive-Impute* converges.

5.2 A real data example

We applied *Adaptive-Impute* and the competing methods to a real data, MovieLens 100k (GroupLens (2015)). We used 5 training and 5 test data sets from 5-fold CV which are publicly available in GroupLens (2015). For the rank used in *Adaptive-Impute* and *soft-Impute*-type algorithms with rank constraint, we chose 3 based on a scree plot (Figure 6). Lemma 2 in Cho et al. (2016) provides justification of using the scree plot and the singular value gap to choose the rank. For the thresholding parameters for *softImpute*-type algorithms, we chose the optimal values which result in the smallest test errors. The test errors were measured by normalized mean absolute error (NMAE) (Herlocker et al. (2004)),

$$\frac{1}{(M_{max} - M_{min})|\Omega_{test}|} \sum_{(i,j) \in \Omega_{test}} |\hat{M}_{ij} - M_{ij}|,$$

where the set Ω_{test} contains indices of the entries in test data, $|\Omega_{test}|$ is the cardinality of Ω_{test} , $M_{max} = \max\{\{M_{i,j}\} \setminus 0\}$ is the largest entry of M , and $M_{min} = \min\{\{M_{i,j}\} \setminus 0\}$ is the smallest entry of M .

Figure 7 summarizes the resulting NMAEs. Five points in the x-axis correspond to the 5-fold CV test data, the y-axis represents the values of NMAE, and the five different lines on the plane correspond to the 5 different algorithms in comparison. We observe that *Adaptive-Impute* outperforms all of the other algorithms. Specifically, the test errors of *Adaptive-Impute* reduce those of *softImpute*-type algorithms by 6%-16%. Among *softImpute*-type algorithms, the ones with rank constraint (i.e. *softImpute*-Rank and *softImpute*-ALS-Rank) performs better than the ones without (i.e. *softImpute* and *softImpute*-ALS). This is the same result to the simulation results.




Figure 5: Convergence of the iterates of *Adaptive-Impute* to the underlying low-rank matrix. In all plots, $n = 1700$, $d = 1000$, $p = 0.1$, and all points were averaged over 100 replicates.




Figure 6: Log of the top 50 singular values of the MovieLens 100k data matrix (GroupLens (2015)).

6 Discussion

Choosing the right thresholding parameter for matrix completion algorithms using thresholded SVD often poses empirical challenges. This paper proposed a novel thresholded SVD algorithm for matrix completion, *Adaptive-Impute*, which employs a theoretically-justified and data-dependent set of thresholding parameters. We established its theoretical guarantees on statistical performance and showed its strong performances in both simulated and real data. It provides understanding on the effects of thresholding and the right threshold level. Yet, there is a newly open problem. Although we proposed a reasonable remedy in the paper, the choice of the rank of the underlying low-rank matrix is of another great practical interest. To estimate the rank and completely automate the entire procedure of *Adaptive-Impute* would be a potential direction for future research.

7 Proofs

Denote by C and C_1 generic constants whose values are free of n and p and may change from appearance to appearance. Also, denote by $\|v\|_2$ the ℓ_2 -norm for any vector $v \in \mathbb{R}^d$




Figure 7: The NMAEs of *Adaptive-Impute* and its competitors measured in 5-fold CV test data from MovieLens 100k (GroupLens (2015)).

and by $\|A\|_2$ the spectral norm, the largest singular value of A , for any matrix $A \in \mathbb{R}^{n \times d}$.

7.1 Proof of Theorem 3.1

Proof of Theorem 3.1. We have

$$\begin{aligned}
 \widetilde{M}_t &= \mathcal{P}_\Omega(M) + \mathcal{P}_\Omega^\perp(Z_t) \\
 &= y \cdot (M_0 + \epsilon) + (1_n 1_d^T - y) \cdot Z_t \\
 &= M_0 + y \cdot \epsilon + (1_n 1_d^T - y) \cdot \eta_t,
 \end{aligned}$$

where 1_n and 1_d are vectors of length n and d , respectively, filled with ones and $\eta_t = Z_t - M_0$, and, $A \cdot B = (A_{ij}B_{ij})_{1 \leq i \leq n, 1 \leq j \leq d}$ for any A and $B \in \mathbb{R}^{n \times d}$. Assume that

$$\frac{1}{\sqrt{nd}} \|\eta_t\|_F = o_p \left(\sqrt{\frac{h_n}{pd}} \right). \quad (12)$$

Then, simple algebraic manipulations show for large n

$$\begin{aligned} \frac{1}{\sqrt{nd}} \|\eta_{t+1}\|_F &= \frac{1}{\sqrt{nd}} \|Z_{t+1} - M_0\|_F \\ &= \frac{1}{\sqrt{nd}} \left\| \sum_{i=1}^r \sqrt{\lambda_i^2(\widetilde{M}_t) - \widetilde{\alpha}_t} \mathbf{u}_i(\widetilde{M}_t) \mathbf{v}_i(\widetilde{M}_t)^T - \sum_{i=1}^r \lambda_i U_i V_i^T \right\|_F \\ &\leq \frac{C}{\sqrt{nd}} \sum_{i=1}^r \left\{ \left| \sqrt{\lambda_i^2(\widetilde{M}_t) - \widetilde{\alpha}_t} - \lambda_i \right| \|U_i V_i^T\|_F \right. \\ &\quad \left. + \lambda_i \left\| \left(\mathbf{u}_i(\widetilde{M}_t) - U_i \mathcal{O}_i \right) V_i^T \right\|_F + \lambda_i \left\| U_i \left(\mathbf{v}_i(\widetilde{M}_t) - V_i \mathcal{Q}_i \right)^T \right\|_F \right\} \\ &\leq C \sum_{i=1}^r \left\{ \frac{1}{\sqrt{nd}} \left| \sqrt{\lambda_i^2(\widetilde{M}_t) - \widetilde{\alpha}_t} - \lambda_i \right| \right. \\ &\quad \left. + \frac{\lambda_i}{\sqrt{nd}} \left\| \mathbf{u}_i(\widetilde{M}_t) - U_i \mathcal{O}_i \right\|_F + \frac{\lambda_i}{\sqrt{nd}} \left\| \mathbf{v}_i(\widetilde{M}_t) - V_i \mathcal{Q}_i \right\|_F \right\}, \quad (13) \end{aligned}$$

where \mathcal{O}_i and \mathcal{Q}_i are in $\{-1, 1\}$ and minimize $\left\| \mathbf{u}_i(\widetilde{M}_t) - U_i \mathcal{O}_i \right\|_F$ and $\left\| \mathbf{v}_i(\widetilde{M}_t) - V_i \mathcal{Q}_i \right\|_F$, respectively.

To find the order of (13), first consider the term $\left\| \mathbf{v}_i(\widetilde{M}_t) - V_i \mathcal{Q}_i \right\|_F$. By Davis-Kahan Theorem (Theorem 3.1 in Li (1998b)) and Proposition 2.2 in Vu and Lei (2013),

$$\left\| \mathbf{v}_i(\widetilde{M}_t) - V_i \mathcal{Q}_i \right\|_F \leq \frac{\frac{1}{nd} \left\| \left(\widetilde{M}_t^T \widetilde{M}_t - [M_0^T M_0 + np\sigma^2 I] \right) V_i \right\|_F}{\left| \frac{1}{nd} \left(\lambda_i^2 + np\sigma^2 - \lambda_{i+1}^2(\widetilde{M}_t) \right) \right|}. \quad (14)$$

Consider the numerator of (14). We have

$$\begin{aligned} &\frac{1}{nd} \left\| \left(\widetilde{M}_t^T \widetilde{M}_t - [M_0^T M_0 + np\sigma^2 I] \right) V_i \right\|_F \\ &\leq \frac{1}{nd} \left\{ \left\| (y \cdot \epsilon)^T (y \cdot \epsilon) V_i - np\sigma^2 V_i \right\|_F + \left\| [(1_n 1_d^T - y) \cdot \eta_t]^T [(1_n 1_d^T - y) \cdot \eta_t] V_i \right\|_F \right. \\ &\quad + \left\| M_0^T (y \cdot \epsilon) V_i \right\|_F + \left\| (y \cdot \epsilon)^T M_0 V_i \right\|_F \\ &\quad + \left\| M_0^T [(1_n 1_d^T - y) \cdot \eta_t] V_i \right\|_F + \left\| [(1_n 1_d^T - y) \cdot \eta_t]^T M_0 V_i \right\|_F \\ &\quad \left. + \left\| [(1_n 1_d^T - y) \cdot \eta_t]^T (y \cdot \epsilon) V_i \right\|_F + \left\| (y \cdot \epsilon)^T [(1_n 1_d^T - y) \cdot \eta_t] V_i \right\|_F \right\} \end{aligned}$$

$$\begin{aligned}
&= \frac{1}{nd} \left\{ O_p \left(p\sqrt{nd} \right) + o_p \left(\frac{nh_n}{p} \right) + O_p \left(\sqrt{pdn^2} \right) + O_p \left(\sqrt{pd^2n} \right) \right. \\
&\quad \left. + o_p \left(\sqrt{\frac{h_n dn^2}{p}} \right) + o_p \left(\sqrt{\frac{h_n dn^2}{p}} \right) + o_p \left(\sqrt{h_n n^2} \right) + o_p \left(\sqrt{h_n dn^2} \right) \right\} \\
&= o_p \left(\sqrt{\frac{h_n}{pd}} \right), \tag{15}
\end{aligned}$$

where the first equality holds due to (1), Assumption 1(2), (12), and (16) and (17) below.

We have

$$\begin{aligned}
&\mathbb{E} \left\| (y \cdot \epsilon)^T (y \cdot \epsilon) V_i - np\sigma^2 V_i \right\|_F^2 \\
&= \mathbb{E} \left\{ \sum_{h=1}^d \left[\sum_{k=1}^n \sum_{j=1}^d (y_{kh} y_{kj} \epsilon_{kh} \epsilon_{kj} V_{ij} - p\sigma^2 V_{ih} \mathbb{1}_{(j=h)}) \right]^2 \right\} \\
&= \sum_{h=1}^d \sum_{k=1}^n \sum_{j=1}^d \mathbb{E} \left(y_{kh} y_{kj} \epsilon_{kh} \epsilon_{kj} V_{ij} - p\sigma^2 V_{ih} \mathbb{1}_{(j=h)} \right)^2 \\
&= \sum_{k=1}^n \sum_{j \neq h}^d V_{ij}^2 \mathbb{E} (y_{kh}^2 y_{kj}^2 \epsilon_{kh}^2 \epsilon_{kj}^2) + \sum_{k=1}^n \sum_{j=h}^d V_{ij}^2 \mathbb{E} (y_{kj}^2 \epsilon_{kj}^2 - p\sigma^2)^2 \\
&= O(p^2 nd), \tag{16}
\end{aligned}$$

where V_{ij} is the j -th element of V_i . Similarly, we have

$$\mathbb{E} \|(y \cdot \epsilon) V_i\|_F^2 = O(pn), \mathbb{E} \|U_i^T (y \cdot \epsilon)\|_F^2 = O(pd), \text{ and } \mathbb{E} \|y \cdot \epsilon\|_F^2 = O(pnd). \tag{17}$$

Consider the denominator of (14). By Weyl's theorem (Theorem 4.3 in Li (1998a)), we have

$$\begin{aligned}
&\max_{1 \leq i \leq d} \frac{1}{nd} |\lambda_i^2 + np\sigma^2 - \lambda_i^2(\widetilde{M}_t)| \\
&\leq \frac{1}{nd} \left\| \widetilde{M}_t^T \widetilde{M}_t - [M_0^T M_0 + np\sigma^2 I] \right\|_2 \\
&\leq \frac{1}{nd} \left\{ \left\| (y \cdot \epsilon)^T (y \cdot \epsilon) - np\sigma^2 I \right\|_2 + \left\| [(1_n 1_d^T - y) \cdot \eta_t]^T [(1_n 1_d^T - y) \cdot \eta_t] \right\|_2 \right. \\
&\quad \left. + 2 \|M_0^T (y \cdot \epsilon)\|_2 + 2 \|M_0^T [(1 - y) \cdot \eta_t]\|_2 + 2 \left\| (y \cdot \epsilon)^T [(1_n 1_d^T - y) \cdot \eta_t] \right\|_2 \right\} \\
&= \frac{1}{nd} \left\{ O_p \left(p\sqrt{nd^2} \right) + o_p \left(\frac{nh_n}{p} \right) + O_p \left(\sqrt{p dn^2} \right) + o_p \left(\sqrt{\frac{dn^2 h_n}{p}} \right) + o_p \left(\sqrt{dn^2 h_n} \right) \right\} \\
&= o_p(1), \tag{18}
\end{aligned}$$

where the last two lines holds similarly to (15).

Thus, by (18) and (15),

$$\|\mathbf{v}_i(\widetilde{M}_t) - V_i \mathcal{O}_i\|_F = o_p \left(\sqrt{\frac{h_n}{pd}} \right). \quad (19)$$

Secondly, similar to the proof of (19), we can show $\|\mathbf{u}_i(\widetilde{M}_t) - U_i \mathcal{O}_i\|_F = o_p \left(\sqrt{h_n/pd} \right)$.

Lastly, consider the term $\frac{1}{\sqrt{nd}} \left| \sqrt{\boldsymbol{\lambda}_i^2(\widetilde{M}_t) - \widetilde{\boldsymbol{\alpha}}_t} - \lambda_i \right|$. By Taylor's expansion, there is λ_*^2 between $\boldsymbol{\lambda}_i^2(\widetilde{M}_t) - \widetilde{\boldsymbol{\alpha}}_t$ and λ_i^2 such that

$$\begin{aligned} & \frac{1}{\sqrt{nd}} \left| \sqrt{\boldsymbol{\lambda}_i^2(\widetilde{M}_t) - \widetilde{\boldsymbol{\alpha}}_t} - \lambda_i \right| \\ &= \frac{1}{\sqrt{nd}} \left| \frac{1}{2\lambda_*} \left(\boldsymbol{\lambda}_i^2(\widetilde{M}_t) - \widetilde{\boldsymbol{\alpha}}_t - \lambda_i^2 \right) \right| \\ &\leq \frac{1}{2\lambda_* \sqrt{nd}} \left| \boldsymbol{\lambda}_i^2(\widetilde{M}_t) - (\lambda_i^2 + np\sigma^2) \right| + \frac{1}{2\lambda_* \sqrt{nd}} |\widetilde{\boldsymbol{\alpha}}_t - np\sigma^2|. \end{aligned} \quad (20)$$

We need to find the convergence rates of $\frac{1}{nd} \left| \boldsymbol{\lambda}_i^2(\widetilde{M}_t) - (\lambda_i^2 + np\sigma^2) \right|$ and $\frac{1}{nd} |\widetilde{\boldsymbol{\alpha}}_t - np\sigma^2|$. Let $V_c = (V_{r+1}, \dots, V_d) \in \mathbb{R}^{d \times (d-r)}$ be a matrix such that $V_c^T V_c = I_{d-r}$ and $V^T V_c = 0_{r \times (d-r)}$ and let $\widetilde{V}_t = (\mathbf{v}_1(\widetilde{M}_t), \dots, \mathbf{v}_d(\widetilde{M}_t)) \in \mathbb{R}^{d \times r}$ and $\widetilde{V}_{tc} = (\mathbf{v}_{r+1}(\widetilde{M}_t), \dots, \mathbf{v}_d(\widetilde{M}_t)) \in \mathbb{R}^{d \times (d-r)}$ so that $\widetilde{V}_t^T \widetilde{V}_{tc} = 0_{r \times (d-r)}$. Also, let $\mathcal{O} = \text{diag}(\mathcal{O}_1, \dots, \mathcal{O}_r)$ and $\mathcal{O}_c = \text{diag}(\mathcal{O}_{r+1}, \dots, \mathcal{O}_d)$, where

$$\mathcal{O}_i := \arg \min_{o \in \{-1, 1\}} \left\| V_i o - \mathbf{v}_i(\widetilde{M}_t) \right\|_2^2 \quad \text{for } i = 1, \dots, d.$$

Then, we have

$$\begin{aligned} & \frac{1}{nd} \left| \boldsymbol{\lambda}_i^2(\widetilde{M}_t) - (\lambda_i^2 + np\sigma^2) \right| \\ &= \frac{1}{nd} \left| \mathbf{v}_i(\widetilde{M}_t)^T \widetilde{M}_t^T \widetilde{M}_t \mathbf{v}_i(\widetilde{M}_t) - V_i^T (M_0^T M_0 + np\sigma^2 I) V_i \right| \\ &\leq \frac{1}{nd} \left| V_i^T \left[\widetilde{M}_t^T \widetilde{M}_t - (M_0^T M_0 + np\sigma^2 I) \right] V_i \right| + \frac{1}{nd} \left| \mathbf{v}_i(\widetilde{M}_t)^T \widetilde{M}_t^T \widetilde{M}_t \mathbf{v}_i(\widetilde{M}_t) - V_i^T \widetilde{M}_t^T \widetilde{M}_t V_i \right| \\ &\leq o_p \left(\sqrt{\frac{h_n}{pd}} \right) + \frac{1}{nd} \left| \mathbf{v}_i(\widetilde{M}_t)^T \widetilde{M}_t^T \widetilde{M}_t \mathbf{v}_i(\widetilde{M}_t) - V_i^T \widetilde{M}_t^T \widetilde{M}_t V_i \right| \\ &= o_p \left(\sqrt{\frac{h_n}{pd}} \right), \end{aligned} \quad (21)$$

where the second inequality can be derived by the similar way to the proof of (15), and the last equality is due to (22) below. Simple algebraic manipulations show

$$\frac{1}{nd} \left| \mathbf{v}_i(\widetilde{M}_t)^T \widetilde{M}_t^T \widetilde{M}_t \mathbf{v}_i(\widetilde{M}_t) - V_i^T \widetilde{M}_t^T \widetilde{M}_t V_i \right|$$

$$\begin{aligned}
&= \frac{1}{nd} \left| \left[V_i \mathcal{O}_i - \mathbf{v}_i(\widetilde{M}_t) \right]^T \widetilde{M}_t^T \widetilde{M}_t \left[V_i \mathcal{O}_i - \mathbf{v}_i(\widetilde{M}_t) \right] + 2\boldsymbol{\lambda}_i^2(\widetilde{M}_t) \left[V_i \mathcal{O}_i - \mathbf{v}_i(\widetilde{M}_t) \right]^T \mathbf{v}_i(\widetilde{M}_t) \right| \\
&\leq \frac{2\boldsymbol{\lambda}_1^2(\widetilde{M}_t)}{nd} \left\| V_i \mathcal{O}_i - \mathbf{v}_i(\widetilde{M}_t) \right\|_2^2 \\
&= o_p \left(\frac{h_n}{pd} \right), \tag{22}
\end{aligned}$$

where the last equality is due to (18) and (19). Also,

$$\begin{aligned}
&\frac{1}{nd} |\widetilde{\boldsymbol{\alpha}}_t - np\sigma^2| \\
&= \frac{1}{nd} \left| \frac{1}{d-r} \sum_{j=r+1}^d \mathbf{v}_j(\widetilde{M}_t)^T \widetilde{M}_t^T \widetilde{M}_t \mathbf{v}_j(\widetilde{M}_t) - np\sigma^2 \right| \\
&\leq \frac{1}{nd} \left| \frac{1}{d-r} \sum_{j=r+1}^d V_j^T \left[\widetilde{M}_t^T \widetilde{M}_t - (M_0^T M_0 + np\sigma^2 I) \right] V_j \right| \\
&\quad + \frac{1}{nd} \left| \frac{1}{d-r} \sum_{j=r+1}^d \left[\mathbf{v}_j(\widetilde{M}_t)^T \widetilde{M}_t^T \widetilde{M}_t \mathbf{v}_j(\widetilde{M}_t) - V_j^T \widetilde{M}_t^T \widetilde{M}_t V_j \right] \right| \\
&= o_p \left(\sqrt{\frac{h_n}{pd}} \right) + \frac{1}{nd} \left| \frac{1}{d-r} \sum_{j=r+1}^d \left[\mathbf{v}_j(\widetilde{M}_t)^T \widetilde{M}_t^T \widetilde{M}_t \mathbf{v}_j(\widetilde{M}_t) - V_j^T \widetilde{M}_t^T \widetilde{M}_t V_j \right] \right| \\
&= o_p \left(\sqrt{\frac{h_n}{pd}} \right), \tag{23}
\end{aligned}$$

where the second equality can be derived by the similar way to the proof of (15), and the last equality is due to (24) below. Similar to the proof of (22), we have

$$\begin{aligned}
&\frac{1}{nd(d-r)} \sum_{j=r+1}^d \left| \mathbf{v}_j(\widetilde{M}_t)^T \widetilde{M}_t^T \widetilde{M}_t \mathbf{v}_j(\widetilde{M}_t) - V_j^T \widetilde{M}_t^T \widetilde{M}_t V_j \right| \\
&\leq \frac{1}{nd(d-r)} \sum_{j=r+1}^d 2\boldsymbol{\lambda}_1^2(\widetilde{M}_t) \left\| V_j \mathcal{O}_j - \mathbf{v}_j(\widetilde{M}_t) \right\|_2^2 \\
&\leq \frac{2\boldsymbol{\lambda}_1^2(\widetilde{M}_t)}{nd(d-r)} \left\| V_c \mathcal{O}_c - \widetilde{V}_{tc} \right\|_F^2 \\
&\leq \frac{4\boldsymbol{\lambda}_1^2(\widetilde{M}_t)}{nd(d-r)} \left\| V_c V_c^T - \widetilde{V}_{tc} \widetilde{V}_{tc}^T \right\|_F^2 \\
&= \frac{4\boldsymbol{\lambda}_1^2(\widetilde{M}_t)}{nd(d-r)} \left\| V V^T - \widetilde{V}_t \widetilde{V}_t^T \right\|_F^2 \\
&\leq \frac{4\boldsymbol{\lambda}_1^2(\widetilde{M}_t)}{nd(d-r)} \left\| V \mathcal{O} - \widetilde{V}_t \right\|_F^2 \\
&= \frac{4\boldsymbol{\lambda}_1^2(\widetilde{M}_t)}{nd(d-r)} \sum_{i=1}^r \left\| V_i \mathcal{O}_i - \mathbf{v}_i(\widetilde{M}_t) \right\|_2^2
\end{aligned}$$

$$= o_p \left(\frac{h_n}{pd^2} \right), \quad (24)$$

where the fourth and sixth lines are due to Proposition 2.2 in Vu and Lei (2013), and the last line holds from (19).

The three results above (20), (21), and (23) give $\frac{1}{\sqrt{nd}} \left| \sqrt{\boldsymbol{\lambda}_i^2(\widetilde{M}_t) - \widetilde{\boldsymbol{\alpha}}_t - \lambda_i} \right| = o_p \left(\sqrt{\frac{h_n}{pd}} \right)$.

Therefore, combining the results above, we have that $\frac{1}{\sqrt{nd}} \|\eta_{t+1}\|_F$ in (13) is $o_p \left(\sqrt{\frac{h_n}{pd}} \right)$. Since $\frac{1}{\sqrt{nd}} \|\eta_1\|_F = o_p \left(\sqrt{\frac{h_n}{pd}} \right)$ by Proposition 3.1, we have $\frac{1}{\sqrt{nd}} \|\eta_t\|_F = o_p \left(\sqrt{\frac{h_n}{pd}} \right)$ for any fixed t by mathematical induction. \square

7.2 Proof of Theorem 3.2

Proof of Theorem 3.2. We have

$$\begin{aligned} & \min_Z \frac{1}{2nd} \|X - Z\|_F^2 + \sum_{i=1}^d \frac{\tau_i}{\sqrt{nd}} \frac{\boldsymbol{\lambda}_i(Z)}{\sqrt{nd}} \\ &= \min_Z \frac{1}{2nd} \left\{ \|X\|_F^2 - 2 \sum_{i=1}^d \tilde{\lambda}_i \cdot \tilde{u}_i^T X \tilde{v}_i + \sum_{i=1}^d \tilde{\lambda}_i^2 \right\} + \frac{1}{nd} \sum_{i=1}^d \tau_i \tilde{\lambda}_i, \end{aligned} \quad (25)$$

where $\tilde{\lambda}_i = \boldsymbol{\lambda}_i(Z)$, $\tilde{u}_i = \mathbf{u}_i(Z)$, and $\tilde{v}_i = \mathbf{v}_i(Z)$. Minimizing (25) is equivalent to minimizing

$$-2 \sum_{i=1}^d \tilde{\lambda}_i \cdot \tilde{u}_i^T X \tilde{v}_i + \sum_{i=1}^d \tilde{\lambda}_i^2 + \sum_{i=1}^d 2\tau_i \tilde{\lambda}_i,$$

with respect to $\tilde{\lambda}_i$, \tilde{u}_i , and \tilde{v}_i , $i = 1, \dots, d$, under the conditions that $(\tilde{u}_1, \dots, \tilde{u}_d)^T (\tilde{u}_1, \dots, \tilde{u}_d) = I_d$, $(\tilde{v}_1, \dots, \tilde{v}_d)^T (\tilde{v}_1, \dots, \tilde{v}_d) = I_d$, and $\tilde{\lambda}_1 \geq \tilde{\lambda}_2 \geq \dots \geq \tilde{\lambda}_d \geq 0$. Thus, we have

$$\begin{aligned} & \min_{\tilde{\lambda}_i \geq 0, \tilde{u}_i, \tilde{v}_i, i=1, \dots, d} -2 \sum_{i=1}^d \tilde{\lambda}_i \cdot \tilde{u}_i^T X \tilde{v}_i + \sum_{i=1}^d \tilde{\lambda}_i^2 + \sum_{i=1}^d 2\tau_i \tilde{\lambda}_i \\ &= \min_{\tilde{\lambda}_i \geq 0, i=1, \dots, d} -2 \sum_{i=1}^d \tilde{\lambda}_i \cdot \boldsymbol{\lambda}_i(X) + \sum_{i=1}^d \tilde{\lambda}_i^2 + \sum_{i=1}^d 2\tau_i \tilde{\lambda}_i \\ &= \min_{\tilde{\lambda}_i \geq 0, i=1, \dots, d} \sum_{i=1}^d \left\{ \tilde{\lambda}_i^2 - 2\tilde{\lambda}_i [\boldsymbol{\lambda}_i(X) - \tau_i] \right\}, \end{aligned} \quad (26)$$

where the first equality is due to the facts that $\tilde{\lambda}_1 \geq \dots \geq \tilde{\lambda}_d \geq 0$, and for every i , the problem

$$\max_{\|u_i\|_2^2 \leq 1, \|v_i\|_2^2 \leq 1} u_i^T X v_i \quad \text{such that} \quad u_i \perp \{\tilde{u}_1^*, \dots, \tilde{u}_{i-1}^*\}, v_i \perp \{\tilde{v}_1^*, \dots, \tilde{v}_{i-1}^*\}$$

is solved by $\tilde{u}_i^*, \tilde{v}_i^*$, the left and right singular vectors of X corresponding to the i -th largest singular value of X . Note that $\tilde{u}_i = \tilde{u}_i^*$. Since (26) is a quadratic function of $\tilde{\lambda}_i$, the solution to the problem (26) is then $\tilde{\lambda}_i = (\boldsymbol{\lambda}_i(X) - \tau_i)_+$. \square

7.3 Proof of Theorem 4.1

To ease the notation, we drop the superscript ‘g’ in Z_t^g, \widetilde{M}_t^g , and D_t^g in this section.

Lemma 7.1. *Let $Z_{t+1} := \arg \min_{Z \in \mathbb{R}^{n \times d}} Q_\tau(Z|Z_t)$ in (9). Then, under Assumption 3, we have*

$$\|Z_{t+1} - Z_t\|_F^2 \rightarrow 0 \quad \text{as } t \rightarrow \infty.$$

Proof of Lemma 7.1. By the construction of D_t ,

$$(\widetilde{M}_{t-1} - \widetilde{M}_t) - (Z_t - Z_{t+1}) - (D_{t-1} - D_t) = 0.$$

Thus, we have

$$\langle \widetilde{M}_{t-1} - \widetilde{M}_t, Z_t - Z_{t+1} \rangle - \langle Z_t - Z_{t+1}, Z_t - Z_{t+1} \rangle - \langle D_{t-1} - D_t, Z_t - Z_{t+1} \rangle = 0 \quad (27)$$

and

$$\langle \widetilde{M}_{t-1} - \widetilde{M}_t, \widetilde{M}_{t-1} - \widetilde{M}_t \rangle - \langle Z_t - Z_{t+1}, \widetilde{M}_{t-1} - \widetilde{M}_t \rangle - \langle D_{t-1} - D_t, \widetilde{M}_{t-1} - \widetilde{M}_t \rangle = 0. \quad (28)$$

Add (28) and (27), and

$$\begin{aligned} 0 &= \|\widetilde{M}_{t-1} - \widetilde{M}_t\|_F^2 - \|Z_t - Z_{t+1}\|_F^2 - \langle D_{t-1} - D_t, Z_t + \widetilde{M}_{t-1} - (Z_{t+1} + \widetilde{M}_t) \rangle \\ &= \|\widetilde{M}_{t-1} - \widetilde{M}_t\|_F^2 - \|Z_t - Z_{t+1}\|_F^2 - \|D_{t-1} - D_t\|_F^2 - 2\langle D_{t-1} - D_t, Z_t - Z_{t+1} \rangle. \end{aligned} \quad (29)$$

Under Assumption 3, (29) gives

$$\|Z_t - Z_{t+1}\|_F^2 \leq \left\| \widetilde{M}_{t-1} - \widetilde{M}_t \right\|_F^2,$$

and thus

$$\begin{aligned} \|Z_{t+1} - Z_t\|_F^2 &\leq \left\| \widetilde{M}_{t-1} - \widetilde{M}_t \right\|_F^2 \\ &\leq \left\| \mathcal{P}_\Omega^\perp(Z_{t-1} - Z_t) \right\|_F^2 \\ &\leq \|Z_t - Z_{t-1}\|_F^2 \end{aligned} \quad (30)$$

for all $t \geq 1$. This proves that the sequence $\{\|Z_{t+1} - Z_t\|_F^2\}$ converges (since it is decreasing and bounded below).

The convergence of $\{\|Z_{t+1} - Z_t\|_F^2\}$ gives

$$\|Z_{t+1} - Z_t\|_F^2 - \|Z_t - Z_{t-1}\|_F^2 \rightarrow 0 \text{ as } t \rightarrow \infty.$$

Then, by (30),

$$\begin{aligned} 0 &\geq \|\mathcal{P}_\Omega^\perp(Z_t - Z_{t-1})\|_F^2 - \|Z_t - Z_{t-1}\|_F^2 \\ &\geq \|Z_{t+1} - Z_t\|_F^2 - \|Z_t - Z_{t-1}\|_F^2 \\ &\rightarrow 0 \text{ as } t \rightarrow \infty, \end{aligned}$$

which implies

$$\|\mathcal{P}_\Omega^\perp(Z_t - Z_{t-1})\|_F^2 - \|Z_t - Z_{t-1}\|_F^2 \rightarrow 0 \Rightarrow \|\mathcal{P}_\Omega(Z_t - Z_{t-1})\|_F^2 \rightarrow 0. \quad (31)$$

Furthermore, similarly to the proof of Lemma 2 in Mazumder et al. (2010), we can show

$$f_\tau(Z_t) \geq Q_\tau(Z_{t+1}|Z_t) \geq Q_\tau(Z_{t+1}|Z_{t+1}) = f_\tau(Z_{t+1}) \geq 0 \quad (32)$$

for every fixed $\tau_1, \dots, \tau_d > 0$ and $t \geq 1$. Thus, we have

$$Q_\tau(Z_{t+1}|Z_t) - Q_\tau(Z_{t+1}|Z_{t+1}) \rightarrow 0 \text{ as } t \rightarrow \infty,$$

which implies

$$\|\mathcal{P}_\Omega^\perp(Z_t - Z_{t+1})\|_F^2 \rightarrow 0 \text{ as } t \rightarrow \infty.$$

The above along with (31) gives

$$\|Z_{t+1} - Z_t\|_F^2 \rightarrow 0 \text{ as } t \rightarrow \infty.$$

□

Proof of Theorem 4.1. By the construction of D_t , we have

$$0 = \left(\widetilde{M}_t - Z_{t+1}\right) - D_t \text{ for all } t \geq 1.$$

Since Z_∞ is a limit point of the sequence Z_t , there exists a subsequence $\{n_t\} \subset \{1, 2, \dots\}$ such that $Z_{n_t} \rightarrow Z_\infty$ as $t \rightarrow \infty$. By Lemma 7.1, this subsequence Z_{n_t} satisfies

$$Z_{n_t} - Z_{n_t+1} \rightarrow 0$$

which implies

$$\mathcal{P}_\Omega^\perp(Z_{n_t}) - Z_{n_{t+1}} \rightarrow \mathcal{P}_\Omega^\perp(Z_\infty) - Z_\infty = -\mathcal{P}_\Omega(Z_\infty).$$

Hence,

$$D_{n_t} = (\mathcal{P}_\Omega(M) + \mathcal{P}_\Omega^\perp(Z_{n_t})) - Z_{n_{t+1}} \rightarrow \mathcal{P}_\Omega(M) - \mathcal{P}_\Omega(Z_\infty) = D_\infty. \quad (33)$$

Due to (11) and (33), we have

$$\begin{aligned} f_\tau(Z^s) &\geq f_\tau(Z_\infty) - \frac{1}{nd} \langle Z^s - Z_\infty, \mathcal{P}_\Omega(M) - \mathcal{P}_\Omega(Z_\infty) - D_\infty \rangle \\ &= f_\tau(Z_\infty). \end{aligned}$$

Since $f_\tau(Z^s) \leq f_\tau(Z_\infty)$ by definition of Z^s , we have $f_\tau(Z^s) = f_\tau(Z_\infty)$. Lastly, by (32), we have $\lim_{t \rightarrow \infty} f_\tau(Z_t) = f(Z^s)$. \square

7.4 Proofs of Lemmas 4.1-4.2

Proof of Lemma 4.1. For $i = 1, \dots, r$, we have

$$\begin{aligned} &\left| \frac{\tau_{t,i}}{\sqrt{nd}} - \frac{\tau_{t+1,i}}{\sqrt{nd}} \right| \\ &= \frac{1}{\sqrt{nd}} \left| \lambda_i(\widetilde{M}_t) - \sqrt{\lambda_i^2(\widetilde{M}_t) - \widetilde{\alpha}_t} - \lambda_i(\widetilde{M}_{t+1}) + \sqrt{\lambda_i^2(\widetilde{M}_{t+1}) - \widetilde{\alpha}_{t+1}} \right| \\ &\leq \frac{1}{\sqrt{nd}} \left| \lambda_i(\widetilde{M}_t) - \left(\sqrt{\lambda_i^2 - np\sigma^2} \right) \right| + \frac{1}{\sqrt{nd}} \left| \sqrt{\lambda_i^2(\widetilde{M}_t) - \widetilde{\alpha}_t} - \lambda_i^2 \right| \\ &\quad + \frac{1}{\sqrt{nd}} \left| \lambda_i(\widetilde{M}_{t+1}) - \left(\sqrt{\lambda_i^2 - np\sigma^2} \right) \right| + \frac{1}{\sqrt{nd}} \left| \sqrt{\lambda_i^2(\widetilde{M}_{t+1}) - \widetilde{\alpha}_{t+1}} - \lambda_i^2 \right| \\ &= (I) + (II) + (III) + (IV). \end{aligned}$$

Then, by (21) and (23), we have

$$\begin{aligned} (I) &= \frac{1}{\sqrt{nd}} \left| \lambda_i(\widetilde{M}_t) - \left(\sqrt{\lambda_i^2 - np\sigma^2} \right) \right| \\ &= \frac{1}{2\lambda_*\sqrt{nd}} \left| \lambda_i^2(\widetilde{M}_t) - (\lambda_i^2 - np\sigma^2) \right| \\ &\leq \frac{1}{2\lambda_*\sqrt{nd}} \left| \lambda_i^2(\widetilde{M}_t) - \widetilde{\alpha}_t - \lambda_i^2 \right| + \frac{1}{2\lambda_*\sqrt{nd}} \left| \widetilde{\alpha}_t - np\sigma^2 \right| \\ &= o_p \left(\sqrt{\frac{h_n}{pd}} \right), \end{aligned}$$

where the second equality holds for some λ_* between $\lambda_i(\widetilde{M}_t)$ and $\sqrt{\lambda_i^2 - np\sigma^2}$ by Taylor's expansion. We can similarly show that $(III) = o_p \left(\sqrt{h_n/pd} \right)$. Both of (II) and (IV) are also $o_p \left(\sqrt{h_n/pd} \right)$ by (20) and (21). \square

Proof of Lemma 4.2. From Theorem 3.1 and the construction of D_t in Assumption 3, we have

$$\begin{aligned}
& \left| \frac{1}{nd} \langle D_t - D_{t+1}, Z_{t+1} - Z_{t+2} \rangle \right| \\
& \leq \frac{1}{nd} \|D_t - D_{t+1}\|_F \|Z_{t+1} - Z_{t+2}\|_F \\
& \leq \frac{1}{nd} \left\| \widetilde{M}_t - Z_{t+1} - (\widetilde{M}_{t+1} - Z_{t+2}) \right\|_F \|Z_{t+1} - Z_{t+2}\|_F \\
& \leq \frac{1}{nd} \left\{ \left\| \widetilde{M}_t - \widetilde{M}_{t+1} \right\|_F + \|Z_{t+1} - Z_{t+2}\|_F \right\} \|Z_{t+1} - Z_{t+2}\|_F \\
& = \frac{1}{nd} \left\{ \left\| \mathcal{P}_\Omega^\perp(Z_t - Z_{t+1}) \right\|_F + \|Z_{t+1} - Z_{t+2}\|_F \right\} \|Z_{t+1} - Z_{t+2}\|_F \\
& \leq \frac{1}{nd} \left\{ \|Z_t - Z_{t+1}\|_F + \|Z_{t+1} - Z_{t+2}\|_F \right\} \|Z_{t+1} - Z_{t+2}\|_F \\
& \leq \frac{1}{nd} \left\{ \|Z_t - M_0\|_F + 2\|Z_{t+1} - M_0\|_F + \|Z_{t+2} - M_0\|_F \right\} \\
& \quad \times \left\{ \|Z_{t+1} - M_0\|_F + \|Z_{t+2} - M_0\|_F \right\} \\
& = o_p\left(\frac{h_n}{pd}\right).
\end{aligned}$$

□

References

- Achlioptas, D. and F. McSherry (2001). Fast computation of low rank matrix approximations. In *Proceedings of the thirty-third annual ACM symposium on Theory of computing*, pp. 611–618. ACM.
- Azar, Y., A. Fiat, A. Karlin, F. McSherry, and J. Saia (2001). Spectral analysis of data. In *Proceedings of the thirty-third annual ACM symposium on Theory of computing*, pp. 619–626. ACM.
- Bennett, J. and S. Lanning (2007). The netflix prize. In *Proceedings of KDD cup and workshop*, Volume 2007, pp. 35.
- Cai, J.-F., E. J. Candès, and Z. Shen (2010). A singular value thresholding algorithm for matrix completion. *SIAM Journal on Optimization* 20(4), 1956–1982.
- Cai, T. and W. Liu (2011). Adaptive thresholding for sparse covariance matrix estimation. *Journal of the American Statistical Association* 106(494), 672–684.

- Cai, T. T. and W.-X. Zhou (2013). Matrix completion via max-norm constrained optimization. *arXiv preprint arXiv:1303.0341*.
- Candès, E. J. and Y. Plan (2010). Matrix completion with noise. *Proceedings of the IEEE* 98(6), 925–936.
- Candès, E. J. and B. Recht (2009). Exact matrix completion via convex optimization. *Foundations of Computational mathematics* 9(6), 717–772.
- Candès, E. J., J. Romberg, and T. Tao (2006). Robust uncertainty principles: Exact signal reconstruction from highly incomplete frequency information. *Information Theory, IEEE Transactions on* 52(2), 489–509.
- Candès, E. J. and T. Tao (2010). The power of convex relaxation: Near-optimal matrix completion. *Information Theory, IEEE Transactions on* 56(5), 2053–2080.
- Chatterjee, S. (2014). Matrix estimation by universal singular value thresholding. *The Annals of Statistics* 43(1), 177–214.
- Cho, J., D. Kim, and K. Rohe (2016). Asymptotic theory for estimating the singular vectors and values of a partially-observed low rank matrix with noise.
- Davenport, M. A., Y. Plan, E. van den Berg, and M. Wootters (2014). 1-bit matrix completion. *Information and Inference* 3(3), 189–223.
- Donoho, D. L. (2006). Compressed sensing. *Information Theory, IEEE Transactions on* 52(4), 1289–1306.
- Fan, J., Y. Liao, and M. Mincheva (2013). Large covariance estimation by thresholding principal orthogonal complements. *Journal of the Royal Statistical Society: Series B (Statistical Methodology)* 75(4), 603–680.
- Fazel, M. (2002). *Matrix rank minimization with applications*. Ph. D. thesis, PhD thesis, Stanford University.
- Gross, D. (2011). Recovering low-rank matrices from few coefficients in any basis. *Information Theory, IEEE Transactions on* 57(3), 1548–1566.

- GroupLens (2015). Movielens100k @MISC. <http://grouplens.org/datasets/movielens/>.
- Hastie, T. and R. Mazumder (2015). softimpute @MISC. <https://cran.r-project.org/web/packages/softImpute/index.html>.
- Hastie, T., R. Mazumder, J. Lee, and R. Zadeh (2014). Matrix completion and low-rank svd via fast alternating least squares. *arXiv preprint arXiv:1410.2596*.
- Herlocker, J. L., J. A. Konstan, L. G. Terveen, and J. T. Riedl (2004). Evaluating collaborative filtering recommender systems. *ACM Transactions on Information Systems (TOIS)* 22(1), 5–53.
- Keshavan, R., A. Montanari, and S. Oh (2009). Matrix completion from noisy entries. In *Advances in Neural Information Processing Systems*, pp. 952–960.
- Keshavan, R. H., A. Montanari, and S. Oh (2010). Matrix completion from a few entries. *Information Theory, IEEE Transactions on* 56(6), 2980–2998.
- Koltchinskii, V. (2011). Von neumann entropy penalization and low-rank matrix estimation. *The Annals of Statistics* 39(6), 2936–2973.
- Koltchinskii, V., K. Lounici, and A. B. Tsybakov (2011). Nuclear-norm penalization and optimal rates for noisy low-rank matrix completion. *The Annals of Statistics* 39(5), 2302–2329.
- Li, R.-C. (1998a). Relative perturbation theory: I. eigenvalue and singular value variations. *SIAM Journal on Matrix Analysis and Applications* 19(4), 956–982.
- Li, R.-C. (1998b). Relative perturbation theory: II. eigenspace and singular subspace variations. *SIAM Journal on Matrix Analysis and Applications* 20(2), 471–492.
- Mazumder, R., T. Hastie, and R. Tibshirani (2010). Spectral regularization algorithms for learning large incomplete matrices. *The Journal of Machine Learning Research* 11, 2287–2322.

- Montanari, A. and S. Oh (2010). On positioning via distributed matrix completion. In *Sensor Array and Multichannel Signal Processing Workshop (SAM), 2010 IEEE*, pp. 197–200. IEEE.
- Negahban, S. and M. J. Wainwright (2011). Estimation of (near) low-rank matrices with noise and high-dimensional scaling. *The Annals of Statistics* 39(2), 1069–1097.
- Negahban, S. and M. J. Wainwright (2012). Restricted strong convexity and weighted matrix completion: Optimal bounds with noise. *The Journal of Machine Learning Research* 13(1), 1665–1697.
- Recht, B. (2011). A simpler approach to matrix completion. *The Journal of Machine Learning Research* 12, 3413–3430.
- Rennie, J. D. and N. Srebro (2005). Fast maximum margin matrix factorization for collaborative prediction. In *Proceedings of the 22nd international conference on Machine learning*, pp. 713–719. ACM.
- Rohde, A. and A. B. Tsybakov (2011). Estimation of high-dimensional low-rank matrices. *The Annals of Statistics* 39(2), 887–930.
- Srebro, N., J. Rennie, and T. S. Jaakkola (2004). Maximum-margin matrix factorization. In *Advances in neural information processing systems*, pp. 1329–1336.
- Vu, V. Q. and J. Lei (2013). Minimax sparse principal subspace estimation in high dimensions. *The Annals of Statistics* 41(6), 2905–2947.
- Weinberger, K. Q. and L. K. Saul (2006). Unsupervised learning of image manifolds by semidefinite programming. *International Journal of Computer Vision* 70(1), 77–90.
- Zou, H. (2006). The adaptive lasso and its oracle properties. *Journal of the American statistical association* 101(476), 1418–1429.