

Remarks on GJMS operator of order six

Xuezhang Chen^{*} and Fei Hou[†]

^{*} [†]Department of Mathematics & IMS, Nanjing University, Nanjing 210093, P. R. China

Abstract

We study analysis aspects of the sixth order GJMS operator P_g^6 . Under conformal normal coordinates around a point, the expansions of Green's function of P_g^6 with pole at this point are presented. As a starting point of the study of P_g^6 , we manage to give some existence results of prescribed Q -curvature problem on Einstein manifolds. One among them is that for $n \geq 10$, let (M^n, g) be a closed Einstein manifold of positive scalar curvature and f a smooth positive function in M . If the Weyl tensor is nonzero at a maximum point of f and f satisfies a vanishing order condition at this maximum point, then there exists a conformal metric $\tilde{g}$ of g such that its Q -curvature $Q_{\tilde{g}}^6$ equals f .

2010 MSC: Primary 53A30, 58J05, 53C21; Secondary 35J35, 35J08, 35B50.

Keywords: Sixth order GJMS operator, Prescribed Q -curvature problem, Green's function, Mountain pass critical points.

1 Introduction

Recently, some remarkable developments have been achieved in the existence theory of positive constant Q -curvature problem associated to Paneitz-Branson operator. One key ingredient in such works is that a strong maximum principle for the fourth order Paneitz-Branson operator is discovered under a hypothesis on the positivity of some conformal invariants or Q -curvature of the background metric. The readers are referred to [8, 9, 10, 13] and the references therein. This naturally stimulates us to study GJMS operator of order six and its associated Q -curvature problem, the analogue to the Yamabe problem and Q -curvature problem for Paneitz-Branson operator. Except for the aforementioned cases, due to the lack of a maximum principle for higher order elliptic equations in general, the existence theory of such problems needs to be developed. Until an analogue of Aubin's result in [2] for the Yamabe problem is verified in Proposition 3.1 below, by adapting some ideas for Paneitz-Branson operator from [4, 3] we establish some existence results of prescribed Q -curvature problem on Einstein manifolds, in which case the sixth order GJMS operator is of constant coefficients.

The conformally covariant GJMS operators with principle part $(-\Delta_g)^k$, $k \in \mathbb{N}$ are discovered by Graham-Jenne-Mason-Sparling [6]. In particular, the GJMS operator of order six and the associated Q -curvature are given as follows (cf. [11, 17]): on manifolds (M^n, g) of dimension $n \geq 3$ and $n \neq 4$, denote by $\sigma_k(A_g)$ the k -th elementary symmetric function of Schouten

^{*}X. Chen: xuezhangchen@nju.edu.cn; [†]F. Hou: houfeimath@gmail.com.

tensor $A_{ij} = \frac{1}{n-2}(R_{ij} - \frac{R_g}{2(n-1)}g_{ij})$. Denote by

$$C_{ijk} = \nabla_k A_{ij} - \nabla_j A_{ik}, \quad B_{ij} = \Delta_g A_{ij} - \nabla^k \nabla_j A_{ik} - A^{kl} W_{kijl} = \nabla^k C_{ijk} - A^{kl} W_{kijl}$$

the Cotton tensor and Bach tensor, respectively. Let

$$\begin{aligned} T_2 &= (n-2)\sigma_1(A_g)g - 8A_g = -\frac{8}{n-2}\text{Ric}_g + \frac{n^2 - 4n + 12}{2(n-1)(n-2)}R_g g; \\ T_4 &= -\frac{3n^2 - 12n - 4}{4}\sigma_1(A_g)^2 g + 4(n-4)|A|_g^2 g + 8(n-2)\sigma_1(A_g)A_g \\ &\quad + (n-6)\Delta_g \sigma_1(A_g)g - 48A_g^2 - \frac{16}{n-4}B_g; \\ v_6 &= -\frac{1}{8}\sigma_3(A_g) - \frac{1}{24(n-4)}\langle B, A \rangle_g. \end{aligned}$$

Then, the Q -curvature Q_g^6 is defined by

$$\begin{aligned} Q_g^6 &= -3!2^6 v_6 - \frac{n+2}{2}\Delta_g(\sigma_1(A_g)^2) + 4\Delta_g|A|_g^2 - 8\delta(A_g d\sigma_1(A_g)) + \Delta_g^2 \sigma_1(A_g) \\ &\quad - \frac{n-6}{2}\sigma_1(A_g)\Delta_g \sigma_1(A_g) - 4(n-6)\sigma_1(A_g)|A|_g^2 + \frac{(n-6)(n+6)}{4}\sigma_1(A_g)^3 \end{aligned} \quad (1.1)$$

and the GJMS operator of sixth order P_g^6 is given by¹

$$-P_g^6 = \Delta_g^3 + \Delta_g \delta T_2 d + \delta T_2 d \Delta_g + \frac{n-2}{2}\Delta_g(\sigma_1(A_g)\Delta_g) + \delta T_4 d - \frac{n-6}{2}Q_g^6, \quad (1.2)$$

where $-\delta d = \Delta_g$. The operator P_g^6 is conformally covariant in the sense that if $\tilde{g} = u^{\frac{4}{n-6}}g$, $0 < u \in C^\infty(M)$ with $n \geq 3$ and $n \neq 4, 6$,

$$u^{\frac{n+6}{n-6}}P_{\tilde{g}}^6\varphi = P_g^6(u\varphi) \quad (1.3)$$

and in dimension 6,

$$P_{e^{2u}g}^6\varphi = e^{-6u}P_g^6\varphi$$

for all $\varphi \in C^\infty(M)$. When (M, g) is Einstein, P_g^6 is of constant coefficients, explicitly,

$$\begin{aligned} Q_g^6 &= \frac{n^4 - 20n^2 + 64}{32n^2(n-1)^3}R_g^3, \\ -P_g^6 &= \Delta_g^3 + \frac{-3n^2 + 6n + 32}{4n(n-1)}R_g\Delta_g^2 + \frac{3n^4 - 12n^3 - 52n^2 + 128n + 192}{16n^2(n-1)^2}R_g^2\Delta_g - \frac{n-6}{2}Q_g^6. \end{aligned}$$

Obviously, when $n \geq 7$, Q_g^6 is a positive constant whenever the scalar curvature R_g is positive. Through a direct computation, the GJMS operator P_g^6 has the following factorization:

$$P_g^6 = \left(-\Delta_g + \frac{(n-6)(n+4)}{4n(n-1)}R_g\right) \left(-\Delta_g + \frac{(n-4)(n+2)}{4n(n-1)}R_g\right) \left(-\Delta_g + \frac{n-2}{4(n-1)}R_g\right). \quad (1.4)$$

¹The definition of P_g^6 differs from the formula (10.15) in [11] by a minus sign.

In general, as shown in [5] and [7], on Einstein manifolds the GJMS operator of order $2k$ for all positive integers k satisfies the above property as

$$P_g^{2k} = \prod_{i=1}^k \left(-\Delta_g + \frac{R_g}{4n(n-1)}(n+2i-2)(n-2i) \right).$$

In particular, choose $M^n = S^n$, $g = g_{S^n}$, then

$$\begin{aligned} Q_{S^n}^6 &= \frac{n(n^4-20n^2+64)}{32}, \\ P_{S^n}^6 &= -\Delta_{S^n}^3 - \frac{-3n^2+6n+32}{4}\Delta_{S^n}^2 - \frac{3n^4-12n^3-52n^2+128n+192}{16}\Delta_{S^n} + \frac{n-6}{2}Q_{S^n}^6 \\ &= \left(-\Delta_{S^n} + \frac{(n-6)(n+4)}{4} \right) \left(-\Delta_{S^n} + \frac{(n-4)(n+2)}{4} \right) \left(-\Delta_{S^n} + \frac{n(n-2)}{4} \right). \end{aligned}$$

From now on, we set $P_g = P_g^6$ and $Q_g = Q_g^6$ in the whole paper unless stated otherwise. Then, for any $\varphi \in H^3(M, g)$, we get

$$\begin{aligned} &\int_M \varphi P_g \varphi d\mu_g \\ &= \int_M \left(|\nabla \Delta \varphi|_g^2 - 2T_2(\nabla \Delta \varphi, \nabla \varphi) - \frac{n-2}{2}\sigma_1(A)(\Delta_g \varphi)^2 - T_4(\nabla \varphi, \nabla \varphi) + \frac{n-6}{2}Q_g \varphi^2 \right) d\mu_g. \end{aligned}$$

As a starting point of the study on the sixth order GJMS operator, we obtain some existence results of conformal metrics with positive Q -curvature candidates on closed Einstein manifolds under some additional natural assumptions.

Theorem 1.1. *Suppose (M^n, g) is a closed Einstein manifold of dimension $n \geq 10$ and has positive scalar curvature. Let f be a smooth positive function in M . Assume the Weyl tensor W_g is nonzero at a maximum point p of f and f satisfies the vanishing order condition at p :*

$$\begin{cases} \Delta_g f(p) = 0, & \text{if } n = 10, \\ \nabla^k f(p) = 0, k = 2, 3, 4, & \text{if } n \geq 11. \end{cases} \quad (1.5)$$

Then there exists a smooth solution to the Q -curvature equation

$$P_g u = f u^{\frac{n+6}{n-6}}, \quad u > 0 \text{ in } M.$$

We remark that the condition (1.5) imposed on the Q -curvature candidates f is conformally invariant. The condition that (M, g) is Einstein is only used to seek a positive solution. Theorem 1.1 is a special case of a generalized Theorem 3.1.

This paper is organized as follows. In section 2, the expansions of Green's function for P_g when $n \geq 7$ are presented under conformal normal coordinates around a point. The technique used here is basically inspired by Lee-Parker [12], see also [10]. The complicate computations of the term $P_g(r^{6-n})$ are left to Appendix A, where r is the geodesic distant from this point. In section 3, we prove an analogue (cf. Proposition 3.1) of Aubin's result for any closed manifold of dimension $n \geq 10$, which is not locally conformal flat. Based on this result, using the Mountain Pass Lemma we state in Theorem 3.1 some results of prescribed Q -curvature problem associated to the sixth order GJMS operator on Einstein manifolds. Then our main Theorem 1.1 directly follows from Theorem 3.1.

Acknowledgments: The first author is supported through NSFC (No.11201223), A Foundation for the Author of National Excellent Doctoral Dissertation of China (No.201417) and Program for New Century Excellent Talents in University (NCET-13-0271). We thank the referee for careful reading and for helpful advice which improves the exposition.

2 Expansion of Green's function of P_g

Based on the survey paper by Lee-Parker [12] on the Yamabe problem, the method of deriving expansions of Green's function of P_g is more or less standard except for careful computations on some lower order terms involved in P_g . One may also refer to [10] for the Paneitz-Branson operator case. Green's functions of conformally covariant operators play an important role in the solvability of the constant curvature problems, for instance, the Yamabe problem (cf. [12] etc.) and the constant Q -curvature problem for Paneitz-Branson operator (cf.[3, 4, 8, 10], etc.). Especially, F. Hang and P. Yang [10] set up a dual variational method of the minimization for the Paneitz-Branson functional to seek a positive maximizer of the dual functional, such a scheme heavily relies on the positivity and expansion of its Green's function. We expect that the expansion of Green's function for P_g^6 will be useful to some possible future applications.

Throughout the whole paper, we use the following notation: $2^\sharp = \frac{2n}{n-6}$, $\omega_n = \text{vol}(S^n, g_{S^n})$ and when $n > 6$, $c_n = \frac{1}{8(n-2)(n-4)(n-6)\omega_{n-1}}$. For $m \in \mathbb{Z}_+$, let

$$\mathcal{P}_m := \{\text{homogeneous polynomials in } \mathbb{R}^n \text{ of degree } m\}$$

and

$$\mathcal{H}_m := \{\text{harmonic polynomials in } \mathbb{R}^n \text{ of degree } m\}.$$

Moreover, $\mathcal{P}_m$ has the following decomposition (cf. [16], P. 68-70)

$$\mathcal{P}_m = \bigoplus_{k=0}^{\lfloor \frac{m}{2} \rfloor} (r^{2k} \mathcal{H}_{m-2k}).$$

Proposition 2.1. *Assume $n > 6$ and $\ker P_g = 0$. Let $G_p(x)$ be the Green's function of sixth order GJMS operator at the pole $p \in M^n$ with the property that $P_g G_p = c_n \delta_p$ in the sense of distribution. Then, under the conformal normal coordinates around p with conformal metric g , $G_p(x)$ has the following expansions:*

(a) *If n is odd, then*

$$G_p(x) = r^{6-n} \left(1 + \sum_{k=1}^n \psi_k \right) + A + O(r),$$

where A is a constant and $\psi_k \in \mathcal{P}_k$.

(b) *If n is even, then*

$$G_p(x) = r^{6-n} \left(1 + \sum_{k=1}^n \psi_k \right) + r^{6-n} \left(\sum_{k=n-4}^n \varphi_k \right) \log r + r^{6-n} \left(\sum_{k=n-4}^n \varphi'_k \right) \log^2 r$$

$$+r^{6-n}\left(\sum_{k=n-2}^n \varphi_k''\right) \log^3 r + \varphi_n''' \log^4 r + A + O(r),$$

where A is a constant and $\psi_k, \varphi_k, \varphi_k', \varphi_k'', \varphi_k''' \in \mathcal{P}_k$.

Moreover, we may restate some of the above results in another way.

(c) If $n = 7, 8, 9$ or M is conformally flat near p , then

$$G_p(x) = c_n r^{6-n} + A + O(r),$$

where A is a constant.

(d) If $n = 10$, then

$$G_p(x) = c_n r^{-4} + \frac{1}{17280} |W(p)|^2 \log r + O(1).$$

(e) If $n \geq 11$, then

$$G_p(x) = c_n r^{6-n} + \psi_4 r^{6-n} + O(r^{11-n}),$$

where $\psi_4 \in \mathcal{P}_4$ and

$$\begin{aligned} \psi_4(x) = & \frac{1}{135(n-2)} \left[\sum_{k,l} (W_{iklj}(p) x^i x^j)^2 - \frac{r^2}{n+4} \sum_{k,l,s} ((W_{ikls}(p) + W_{ilks}(p)) x^i)^2 \right. \\ & \left. + \frac{3}{2(n+4)(n+2)} |W(p)|^2 r^4 \right] \\ & + \frac{3n-20}{270(n+4)(n-4)(n-8)} r^2 \left[\sum_{k,l,s} ((W_{ikls}(p) + W_{ilks}(p)) x^i)^2 - \frac{3}{n} |W(p)|^2 r^2 \right] \\ & - \frac{5n^2 - 66n + 224}{120(n-8)(n-4)} r^2 \left[\sigma_1(A)_{,ij}(p) x^i x^j + \frac{|W(p)|^2}{12n(n-1)} r^2 \right] \\ & \left. + \frac{3n^4 - 16n^3 - 164n^2 + 400n + 2432}{576(n+4)(n+2)n(n-1)} |W(p)|^2 r^4 \right\}. \end{aligned}$$

Before starting to derive the expansion of Green's function of P_g , we first need to introduce some notation. For $\alpha \in \mathbb{R}$, set

$$\begin{aligned} A_\alpha &= r^2 \Delta_0 + 2\alpha r \partial_r + \alpha(\alpha + n - 2), \\ A_{\alpha,g} &= r^2 \Delta_g + 2\alpha r \partial_r + \alpha(\alpha + n - 2), \end{aligned}$$

where Δ_0 denotes the Euclidean Laplacian, and

$$B_\alpha = \frac{\partial}{\partial \alpha} A_\alpha = 2r \partial_r + 2\alpha + n - 2.$$

For $k \in \mathbb{Z}_+$, a straightforward computation yields (also see [10, Lemma 2.4])

$$A_\alpha(\varphi \log^k r) = A_\alpha \varphi \log^k r + k B_\alpha \varphi \log^{k-1} r + k(k-1) \varphi \log^{k-2} r.$$

From this, for $\alpha, \beta, \gamma \in \mathbb{R}$ we get

$$\begin{aligned}
& A_\gamma A_\beta A_\alpha (\varphi \log^k r) \\
&= A_\gamma A_\beta A_\alpha \varphi \log^k r \\
&+ k(B_\gamma A_\beta A_\alpha + A_\gamma B_\beta A_\alpha + A_\gamma A_\beta B_\alpha) \varphi \log^{k-1} r \\
&+ k(k-1)(A_\beta A_\alpha + B_\gamma B_\beta A_\alpha + B_\gamma A_\beta B_\alpha + A_\gamma B_\beta B_\alpha + A_\gamma A_\alpha + A_\gamma A_\beta) \varphi \log^{k-2} r \\
&+ k(k-1)(k-2)(B_\beta A_\alpha + A_\beta B_\alpha + B_\gamma A_\alpha + B_\gamma B_\beta B_\alpha + B_\gamma A_\beta + A_\gamma B_\alpha + A_\gamma B_\beta) \varphi \log^{k-3} r \\
&+ k(k-1)(k-2)(k-3)(A_\alpha + A_\beta + A_\gamma + B_\gamma B_\beta + B_\gamma B_\alpha + B_\beta B_\alpha) \varphi \log^{k-4} r \\
&+ k(k-1)(k-2)(k-3)(k-4)(B_\alpha + B_\beta + B_\gamma) \varphi \log^{k-5} r \\
&+ k(k-1)(k-2)(k-3)(k-4)(k-5) \varphi \log^{k-6} r.
\end{aligned} \tag{2.1}$$

A direct computation yields

$$\begin{aligned}
\Delta_0(r^\alpha \varphi) &= r^{\alpha-2} A_\alpha \varphi, \\
\Delta_0^2(r^\alpha \varphi) &= \Delta_0(r^{\alpha-2} A_\alpha \varphi) = r^{\alpha-4} A_{\alpha-2} A_\alpha \varphi, \\
\Delta_0^3(r^\alpha \varphi) &= r^{\alpha-6} A_{\alpha-4} A_{\alpha-2} A_\alpha \varphi.
\end{aligned}$$

In particular,

$$\Delta_0^3(r^{6-n} \varphi) = r^{-n} A_{2-n} A_{4-n} A_{6-n} \varphi.$$

Define

$$M_g := \Delta_g \delta T_2 d + \delta T_2 d \Delta_g + \frac{n-2}{2} \Delta_g (\sigma_1(A) \Delta_g) + \delta T_4 d,$$

then rewrite (1.2) as $-P_g = (\Delta_g)^3 + M_g - \frac{n-6}{2} Q_g$. Notice that

$$\begin{aligned}
A_{\alpha,g} &= A_\alpha + r^2 (\Delta_g - \Delta_0) = A_\alpha + r^2 \partial_i ((g^{ij} - \delta^{ij}) \partial_j), \\
-P_g(r^\alpha \varphi) &= r^{\alpha-6} (A_{\alpha-4} A_{\alpha-2} A_\alpha \varphi + K_\alpha \varphi),
\end{aligned}$$

where

$$\begin{aligned}
K_\alpha \varphi &= r^2 (\Delta_g - \Delta_0) A_{\alpha-2} A_\alpha \varphi + A_{\alpha-4} (r^2 (\Delta_g - \Delta_0)) A_\alpha \varphi + A_{\alpha-4} A_{\alpha-2} (r^2 (\Delta_g - \Delta_0)) \varphi \\
&+ r^{6-\alpha} M_g(r^\alpha \varphi) - \frac{n-6}{2} r^6 Q_g \varphi.
\end{aligned} \tag{2.2}$$

We first state the expression of $P_g(r^{6-n})$ and leave complicate computations to Appendix A.

Lemma 2.1. *Under conformal normal coordinates around p with metric g , there holds*

$$\begin{aligned}
& -P_g(r^{6-n}) \\
&= -c_n \delta_p + (n-6) r^{-n} \left\{ \frac{64(n-4)}{9} \left[\sum_{k,l} (W_{iklj}(p) x^i x^j)^2 - \frac{r^2}{n+4} \sum_{k,l,s} ((W_{ikls}(p) + W_{ilks}(p)) x^i)^2 \right. \right. \\
&\quad \left. \left. + \frac{3}{2(n+4)(n+2)} |W(p)|^2 r^4 \right] + \frac{16(3n-20)}{9(n+4)} r^2 \left[\sum_{k,l,s} ((W_{ikls}(p) + W_{ilks}(p)) x^i)^2 - \frac{3}{n} |W(p)|^2 r^2 \right] \right\}
\end{aligned}$$

$$\begin{aligned}
& -4(5n^2 - 66n + 224)r^2 \left[\sigma_1(A)_{,ij}(p)x^i x^j + \frac{|W(p)|^2}{12n(n-1)}r^2 \right] \\
& + \frac{3n^4 - 16n^3 - 164n^2 + 400n + 2432}{3(n+4)(n+2)n(n-1)} |W(p)|^2 r^4 \Big\} + O(r^{5-n}),
\end{aligned}$$

where W_{ijkl} is the Weyl tensor of metric g and each term in square brackets on the right hand side of the identity is harmonic polynomial.

Consequently, we rewrite the above equation in Lemma 2.1 as

$$P_g(r^{6-n}) = c_n \delta_p + r^{-n} f$$

with $f = O(r^4)$.

Observe that for $i = 0, 1, \dots, [\frac{m}{2}]$,

$$A_\alpha|_{r^{2i}\mathcal{H}_{m-2i}} = (\alpha + 2i)(2m - 2i + \alpha + n - 2)$$

and

$$B_\alpha|_{r^{2i}\mathcal{H}_{m-2i}} = 2m + 2\alpha + n - 2,$$

then it yields

$$\begin{aligned}
& A_{2-n}A_{4-n}A_{6-n}|_{r^{2i}\mathcal{H}_{m-2i}} \\
& = (6 - n + 2i)(4 - n + 2i)(2 - n + 2i)(2m + 4 - 2i)(2m + 2 - 2i)(2m - 2i). \quad (2.3)
\end{aligned}$$

We start to find a formal asymptotic solution like $G_p(x) = r^{6-n}(1 + \sum_{k=1}^n \psi_k) + \varphi$ with $\psi_k \in \mathcal{P}_k$. If we can find $\bar{\psi} = \sum_{k=1}^n \psi_k$ such that

$$A_{2-n}A_{4-n}A_{6-n}\bar{\psi} + K_{6-n}\bar{\psi} + f = O(r^{n+1}), \quad (2.4)$$

the regularity theory for elliptic equations gives that there exists a solution $\varphi \in C_{\text{loc}}^{6,\alpha}$ for any $0 < \alpha < 1$ to

$$P_g(\varphi) = -r^{-n}(A_{2-n}A_{4-n}A_{6-n}\bar{\psi} + K_{6-n}\bar{\psi} + f) \in C_{\text{loc}}^\alpha.$$

Thus it only remains to seek $\bar{\psi}$ satisfying (2.4) via induction. For any nonnegative integer k , it is not hard to see from the definition (2.2) for K_{6-n} that $K_{6-n}\varphi \in \mathcal{P}_{k+2}$ when $\varphi \in \mathcal{P}_k$. We first set $\psi_1 = \psi_2 = \psi_3 = 0$ by (2.4) and define

$$f_3 = f = O(r^4).$$

Case 1. n is odd.

If we have found $\psi_1, \dots, \psi_k$ for $3 \leq k \leq n-1$ with $\psi_k \in \mathcal{P}_k$ and

$$f_k = A_{2-n}A_{4-n}A_{6-n} \left(\sum_{i=1}^k \psi_i \right) + K_{6-n} \left(\sum_{i=1}^k \psi_i \right) + f := b_{k+1} + O(r^{k+2}),$$

then it follows from (2.3) that $A_{2-n}A_{4-n}A_{6-n}$ is invertible on $\mathcal{P}_{k+1}$ for $0 \leq k \leq n-1$. Thus there exists a unique $\psi_{k+1} \in \mathcal{P}_{k+1}$ such that

$$A_{2-n}A_{4-n}A_{6-n}\psi_{k+1} + b_{k+1} = 0.$$

This implies that

$$\begin{aligned} f_{k+1} &= A_{2-n}A_{4-n}A_{6-n} \left(\sum_{i=1}^{k+1} \psi_i \right) + K_{6-n} \left(\sum_{i=1}^{k+1} \psi_i \right) + f \\ &= f_k + A_{2-n}A_{4-n}A_{6-n}\psi_{k+1} + K_{6-n}\psi_{k+1} \\ &= O(r^{k+2}). \end{aligned}$$

This finishes the induction and assertion (a) follows.

Case 2. n is even and not less than 10.

Since $A_{2-n}A_{4-n}A_{6-n}$ is invertible on $\mathcal{P}_k$ for $0 \leq k \leq n-7$, by the same induction in Case 1, we may find $\psi_1, \dots, \psi_{n-7}$ such that

$$\begin{aligned} f_{n-7} &= A_{2-n}A_{4-n}A_{6-n} \left(\sum_{k=1}^{n-7} \psi_k \right) + K_{6-n} \left(\sum_{k=1}^{n-7} \psi_k \right) + f = O(r^{n-6}) \\ &:= b_{n-6} + O(r^{n-5}). \end{aligned}$$

Let $\psi_{n-6}^{(0)} = \alpha_{n-6}^{(0)}(x) + \beta_{n-6}^{(0)}(x) \log r$, where $\alpha_{n-6}^{(0)}(x) \in \mathcal{P}_{n-6} \setminus r^{n-6}\mathcal{H}_0$ and $\beta_{n-6}^{(0)}(x) \in r^{n-6}\mathcal{H}_0$, then it yields from (2.1) that

$$\begin{aligned} &A_{2-n}A_{4-n}A_{6-n}\psi_{n-6}^{(0)} \\ &= A_{2-n}A_{4-n}A_{6-n}\alpha_{n-6}^{(0)} + (B_{2-n}A_{4-n}A_{6-n} + A_{2-n}B_{4-n}A_{6-n} + A_{2-n}A_{4-n}B_{6-n})\beta_{n-6}^{(0)}. \end{aligned}$$

Notice that for $0 \leq i \leq \frac{n-8}{2}$, there hold

$$A_{2-n}A_{4-n}A_{6-n}|_{r^{2i}\mathcal{H}_{m-2i}} \neq 0$$

by (2.3) and

$$(B_{2-n}A_{4-n}A_{6-n} + A_{2-n}B_{4-n}A_{6-n} + A_{2-n}A_{4-n}B_{6-n})|_{r^{n-6}\mathcal{H}_0} = 8(n-2)(n-4)(n-6) \neq 0.$$

Hence there exists a unique $\psi_{n-6}^{(0)} \in \mathcal{P}_{n-6} + \mathcal{P}_{n-6} \log r$ to

$$A_{2-n}A_{4-n}A_{6-n}\psi_{n-6}^{(0)} + b_{n-6} = 0.$$

This indicates that

$$\begin{aligned} f_{n-6} &= f_{n-7} + A_{2-n}A_{4-n}A_{6-n}\psi_{n-6}^{(0)} + K_{6-n}\psi_{n-6}^{(0)} \\ &= O(r^{n-5}) + \left(K_{6-n}\beta_{n-6}^{(0)} \right) \log r \end{aligned}$$

$$:= b_{n-5} + O(r^{n-4}) \log r + O(r^{n-4}).$$

Let $\psi_{n-5}^{(0)} = \alpha_{n-5}^{(0)} + \beta_{n-5}^{(0)} \log r$, where $\alpha_{n-5}^{(0)} \in \mathcal{P}_{n-5} \setminus r^{n-6}\mathcal{H}_1$ and $\beta_{n-5}^{(0)} \in r^{n-6}\mathcal{H}_1$, then it yields

$$\begin{aligned} & A_{2-n}A_{4-n}A_{6-n}\psi_{n-5}^{(0)} \\ &= A_{2-n}A_{4-n}A_{6-n}\alpha_{n-5}^{(0)} + (B_{2-n}A_{4-n}A_{6-n} + A_{2-n}B_{4-n}A_{6-n} + A_{2-n}A_{4-n}B_{6-n})\beta_{n-5}^{(0)}. \end{aligned}$$

By similar arguments, there exists a unique $\psi_{n-5}^{(0)} \in \mathcal{P}_{n-5} + r^{n-6}\mathcal{H}_1 \log r$ such that

$$A_{2-n}A_{4-n}A_{6-n}\psi_{n-5}^{(0)} + b_{n-5} = 0.$$

This implies that

$$\begin{aligned} f_{n-5} &= f_{n-6} + A_{2-n}A_{4-n}A_{6-n}\psi_{n-5}^{(0)} + K_{6-n}\psi_{n-5}^{(0)} \\ &= O(r^{n-4}) \log r + O(r^{n-4}) := b_{n-4}^{(1)} \log r + O(r^{n-4}) + O(r^{n-3}) \log r. \end{aligned}$$

Choose $\psi_{n-4}^{(1)} = \alpha_{n-4}^{(1)} \log r + \beta_{n-4}^{(1)} \log^2 r \in \mathcal{P}_{n-4} \log r + (r^{n-6}\mathcal{H}_2 + r^{n-4}\mathcal{H}_0) \log^2 r$, then (2.1) gives

$$\begin{aligned} & A_{2-n}A_{4-n}A_{6-n}\psi_{n-4}^{(1)} \\ &= \left[A_{2-n}A_{4-n}A_{6-n}\alpha_{n-4}^{(1)} + 2(B_{6-n}A_{4-n}A_{2-n} + A_{6-n}B_{4-n}A_{2-n} + A_{6-n}A_{4-n}B_{2-n})\beta_{n-4}^{(1)} \right] \log r \\ & \quad + A_{2-n}A_{4-n}A_{6-n}\beta_{n-4}^{(1)} \log^2 r + O(r^{n-4}). \end{aligned}$$

Since

$$\begin{aligned} (B_{6-n}A_{4-n}A_{2-n} + A_{6-n}B_{4-n}A_{2-n} + A_{6-n}A_{4-n}B_{2-n})|_{r^{n-6}\mathcal{H}_2} &= 8(n+2)n(n-2) \neq 0; \\ (B_{6-n}A_{4-n}A_{2-n} + A_{6-n}B_{4-n}A_{2-n} + A_{6-n}A_{4-n}B_{2-n})|_{r^{n-4}\mathcal{H}_0} &= -4n(n-2)(n-4) \neq 0 \end{aligned}$$

and $A_{2-n}A_{4-n}A_{6-n}|_{r^{2i}\mathcal{H}_{n-4-2i}} \neq 0$ for $0 \leq i \leq \frac{n-8}{2}$, then there exists a unique $\psi_{n-4}^{(1)}$ such that

$$A_{2-n}A_{4-n}A_{6-n}\alpha_{n-4}^{(1)} + 2(B_{6-n}A_{4-n}A_{2-n} + A_{6-n}B_{4-n}A_{2-n} + A_{6-n}A_{4-n}B_{2-n})\beta_{n-4}^{(1)} + b_{n-4}^{(1)} = 0$$

and

$$\begin{aligned} f_{n-4}^{(1)} &= f_{n-5} + A_{2-n}A_{4-n}A_{6-n}\psi_{n-4}^{(1)} + K_{6-n}\psi_{n-4}^{(1)} \\ &= O(r^{n-4}) + O(r^{n-3}) \log r + O(r^{n-2}) \log^2 r \\ &:= b_{n-4}^{(0)} + O(r^{n-3}) \log r + O(r^{n-3}) + O(r^{n-2}) \log^2 r. \end{aligned}$$

Choose $\psi_{n-4}^{(0)} \in \mathcal{P}_{n-4} + (r^{n-6}\mathcal{H}_2 + r^{n-4}\mathcal{H}_0) \log r$ to remove the term $b_{n-4}^{(0)}$ and set

$$\begin{aligned} f_{n-4}^{(0)} &= f_{n-4}^{(1)} + A_{2-n}A_{4-n}A_{6-n}\psi_{n-4}^{(0)} + K_{6-n}\psi_{n-4}^{(0)} \\ &= O(r^{n-3}) \log r + O(r^{n-3}) + O(r^{n-2}) \log^2 r. \end{aligned}$$

By similar arguments and (2.1), we get

$$\begin{aligned}
\psi_{n-3}^{(1)} &\in \mathcal{P}_{n-3} \log r + (r^{n-6} \mathcal{H}_3 + r^{n-4} \mathcal{H}_1) \log^2 r; \\
\psi_{n-3}^{(0)} &\in \mathcal{P}_{n-3} + (r^{n-6} \mathcal{H}_3 + r^{n-4} \mathcal{H}_1) \log r; \\
\psi_{n-2}^{(i)} &\in \mathcal{P}_{n-2} \log^i r + (r^{n-6} \mathcal{H}_4 + r^{n-4} \mathcal{H}_2 + r^{n-2} \mathcal{H}_0) \log^{i+1} r, \quad \text{for } i = 0, 1, 2; \\
\psi_{n-1}^{(i)} &\in \mathcal{P}_{n-1} \log^i r + (r^{n-6} \mathcal{H}_5 + r^{n-4} \mathcal{H}_3 + r^{n-2} \mathcal{H}_1) \log^{i+1} r, \quad \text{for } i = 0, 1, 2; \\
\psi_n^{(i)} &\in \mathcal{P}_n \log^i r + (r^{n-6} \mathcal{H}_6 + r^{n-4} \mathcal{H}_4 + r^{n-2} \mathcal{H}_2) \log^{i+1} r, \quad \text{for } i = 0, 1, 2, 3.
\end{aligned}$$

Now we set

$$\psi_{n-6} = \psi_{n-6}^{(0)}, \psi_{n-5} = \psi_{n-5}^{(0)}, \psi_{n-4} = \psi_{n-4}^{(0)} + \psi_{n-4}^{(1)}, \psi_{n-3} = \psi_{n-3}^{(0)} + \psi_{n-3}^{(1)}$$

and

$$\psi_{n-2} = \sum_{i=0}^2 \psi_{n-2}^{(i)}, \quad \psi_{n-1} = \sum_{i=0}^2 \psi_{n-1}^{(i)}, \quad \psi_n = \sum_{i=0}^3 \psi_n^{(i)}.$$

Eventually, we obtain

$$\begin{aligned}
f_n &= A_{2-n} A_{4-n} A_{6-n} \left(\sum_{k=1}^n \psi_k \right) + K_{6-n} \left(\sum_{k=1}^n \psi_k \right) + f \\
&= O(r^{n+1}) (\log^3 r + \log^2 r + \log r + 1) + O(r^{n+2}) \log^4 r.
\end{aligned}$$

Hence, $r^{-n} f_n \in C^\alpha$ for any $0 < \alpha < 1$. This finishes the induction and we obtain assertion (b) as desired.

Case 3. $n = 8$.

Notice that

$$P_g(G_p(x) - c_n r^{-2}) = O(r^{-4}) \in L^p,$$

for some $\frac{8}{5} < p < 2$, then it follows from regularity theory of elliptic equations that $G_p(x) - c_n r^{-2} \in C_{\text{loc}}^{6-\frac{8}{p}}$. From this, we have

$$G_p(x) = c_n r^{-2} + A + O(r).$$

Case 4. M is locally conformal flat.

One may choose g flat near p and $P_g = -\Delta_0^3$. Hence, one has $P_g(G(x) - c_n r^{6-n}) = 0$ and then $G_p(x) - c_n r^{6-n}$ is smooth near p .

Therefore, the assertion (c) follows from cases 1,3,4. In some special cases, the leading term ψ_4 can be computed with the help of Lemma 2.1. The proof of Proposition 2.1 is complete.

3 $n \geq 10$ and not locally conformally flat

Similar to the Yamabe constant, for $n \geq 3$ and $n \neq 4, 6$, we define

$$Y_6^+(M, g) = \inf_{0 < u \in H^3(M, g)} \frac{\int_M u P_g u d\mu_g}{\left(\int_M u^{\frac{2n}{n-6}} d\mu_g \right)^{\frac{n-6}{n}}}.$$

It follows from (1.3) that $Y_6^+(M, g)$ is a conformal invariant. However, due to the lack of a maximum principle for higher order elliptic equations in general, we first study another conformally invariant quantity

$$Y_6(M, g) = \inf_{u \in H^3(M, g) \setminus \{0\}} \frac{\int_M u P_g u d\mu_g}{\left(\int_M |u|^{\frac{2n}{n-6}} d\mu_g \right)^{\frac{n-6}{n}}}.$$

In particular, we have $Y_6(S^n) = Y_6^+(S^n) = \frac{n-6}{2} Q_{S^n} \omega_n^{\frac{6}{n}}$. For $w \in C_c^\infty(\mathbb{R}^n)$, let

$$\|w\|_{\mathcal{D}^{3,2}} := \sum_{|\beta|=3} \|D^\beta w\|_{L^2(\mathbb{R}^n)} \approx \|\nabla \Delta w\|_{L^2(\mathbb{R}^n)}$$

and let $\mathcal{D}^{3,2}(\mathbb{R}^n)$ denote the completion of $C_c^\infty(\mathbb{R}^n)$ under this norm. The equivalence of the above last two norms can be easily deduced by the formula (3.4) below. We first recall an optimal Euclidean Sobolev inequality (cf. [15, p.154-165], [14]).

Lemma 3.1. *For $n \geq 7$, the following sharp Sobolev embedding inequality holds*

$$Y_6(S^n) \left(\int_{\mathbb{R}^n} |w|^{\frac{2n}{n-6}} dy \right)^{\frac{n-6}{n}} \leq \int_{\mathbb{R}^n} |\nabla \Delta w|^2 dy \text{ for all } w \in \mathcal{D}^{3,2}(\mathbb{R}^n).$$

The equality holds if and only if $w(y) = \left(\frac{2}{1+|y|^2} \right)^{\frac{n-6}{2}}$ up to any nonzero constant multiple, as well as all translations and dilations.

Proposition 3.1. *On a closed Riemannian manifold (M^n, g) of dimension $n \geq 10$, if there exists $p \in M^n$ such that the Weyl tensor $W_g(p) \neq 0$, then $Y_6(M^n) < Y_6(S^n)$.*

Proof. Recall the definition of P_g :

$$-P_g = \Delta_g^3 + \Delta_g \delta T_2 d + \delta T_2 d \Delta_g + \frac{n-2}{2} \Delta_g (\sigma_1(A) \Delta_g) + \delta T_4 d - \frac{n-6}{2} Q_g,$$

then for all $\varphi \in H^3(M, g)$,

$$\begin{aligned} \int_M \varphi P_g \varphi d\mu_g &= \int_M |\nabla \Delta \varphi|_g^2 d\mu_g - 2 \int_M T_2(\nabla \varphi, \nabla \Delta \varphi) d\mu_g - \frac{n-2}{2} \int_M \sigma_1(A) (\Delta \varphi)^2 d\mu_g \\ &\quad - \int_M T_4(\nabla \varphi, \nabla \varphi) d\mu_g + \frac{n-6}{2} \int_M Q_g \varphi^2 d\mu_g. \end{aligned}$$

Fix $\rho > 0$ small and choose test functions

$$\varphi(x) = \eta_\rho(x) u_\epsilon(x), \quad u_\epsilon(x) = \left(\frac{2\epsilon}{\epsilon^2 + |x|^2} \right)^{\frac{n-6}{2}}, \quad \epsilon > 0,$$

where $r = |x| = d_g(x, p)$ and

$$\eta_\rho \in C_c^\infty, \quad 0 \leq \eta_\rho \leq 1, \quad \eta_\rho \equiv 1 \text{ in } B_\rho \text{ and } \eta_\rho \equiv 0 \text{ in } B_{2\rho}^c.$$

It is known from Lee-Park [12] that up to a conformal factor, under conformal normal coordinates around p with metric g , for all $N \geq 5$ there hold

$$\sigma_1(A_g)(p) = 0, \quad \sigma_1(A_g)_{,i}(p) = 0, \quad \Delta_g \sigma_1(A_g)(p) = -\frac{|W(p)|_g^2}{12(n-1)}$$

and

$$\sqrt{\det g} = 1 + O(r^N).$$

Our purpose is to estimate

$$\int_M \varphi P_g \varphi d\mu_g \quad \text{and} \quad \int_M \varphi^{\frac{2n}{n-6}} d\mu_g.$$

A direct computation shows

$$u'_\epsilon = -(n-6)u_\epsilon \frac{r}{\epsilon^2 + r^2}, \quad u''_\epsilon = -(n-6)u_\epsilon \frac{\epsilon^2 - (n-5)r^2}{(\epsilon^2 + r^2)^2}$$

and

$$\begin{aligned} \Delta_0 u_\epsilon &= -(n-6) \frac{u_\epsilon}{(\epsilon^2 + r^2)^2} (n\epsilon^2 + 4r^2), \\ (\Delta_0 u_\epsilon)' &= (n-6)(n-4) \frac{u_\epsilon r}{(\epsilon^2 + r^2)^3} [(n+2)\epsilon^2 + 4r^2]. \end{aligned}$$

We start with $\int_M |\nabla \Delta \varphi|_g^2 d\mu_g$ and divide its integral into two parts $\int_M = \int_{B_\rho} + \int_{M \setminus \overline{B_\rho}}$. Compute with

$$\begin{aligned} \int_{B_\rho} |\nabla \Delta \varphi|_g^2 d\mu_g &= \int_{B_\rho} g^{ij} (\Delta \varphi)_{,i} (\Delta \varphi)_{,j} d\mu_g \\ &= \int_{B_\rho} (\delta^{ij} + O(r^2)) (\Delta_0 \varphi + O(r^{N-1}) \varphi')_{,i} (\Delta_0 \varphi + O(r^{N-1}) \varphi')_{,j} (1 + O(r^N)) dx \\ &= \int_{B_\rho} |(\nabla \Delta)_0 \varphi|^2 dx + \int_{B_\rho} (\Delta_0 \varphi)' (O(r^{N-2}) \varphi' + O(r^{N-1}) \varphi'') dx \end{aligned}$$

and

$$\begin{aligned} \int_{\mathbb{R}^n \setminus \overline{B_\rho}} |(\nabla \Delta)_0 \varphi|^2 dx &= (n-6)^2 (n-4)^2 \int_{\mathbb{R}^n \setminus \overline{B_\rho}} \frac{u_\epsilon^2 r^2}{(\epsilon^2 + r^2)^6} [(n+2)\epsilon^2 + 4r^2]^2 dx \\ &\leq C \int_{\rho/\epsilon}^\infty \sigma^{5-n} d\sigma = O(\epsilon^{n-6}). \end{aligned}$$

Similarly, we estimate

$$\int_{M \setminus \overline{B_\rho}} |\nabla \Delta \varphi|_g^2 d\mu_g = O(\epsilon^{n-6}).$$

Thus, we obtain

$$\int_M |\nabla \Delta \varphi|_g^2 d\mu_g = \int_{\mathbb{R}^n} |\nabla \Delta_0 u_\epsilon|^2 dx + O(\epsilon^{n-6}).$$

Secondly, we compute

$$\begin{aligned} & \int_{B_\rho} \sigma_1(A) (\Delta \varphi)^2 d\mu_g \\ &= \int_{B_\rho} \left(\frac{1}{2} \sigma_1(A)_{,ij}(p) x^i x^j + O(r^3) \right) (\Delta_0 \varphi + O(r^{N-1}) \varphi')^2 (1 + O(r^N)) dx \\ &= \int_{B_\rho} \frac{1}{2n} \Delta \sigma_1(A)(p) |x|^2 (\Delta_0 \varphi)^2 dx + \int_{B_\rho} O(r^3) \frac{u_\epsilon^2}{(\epsilon^2 + r^2)^4} (n\epsilon^2 + 4r^2)^2 dx \\ &= -\frac{(n-6)^2 |W(p)|^2}{24n(n-1)} \omega_{n-1} \int_0^\rho \frac{(n\epsilon^2 + 4r^2)^2}{(\epsilon^2 + r^2)^4} u_\epsilon^2 r^{n+1} dr + \int_{B_\rho} \frac{O(r^3) u_\epsilon^2}{(\epsilon^2 + r^2)^2} dx \end{aligned}$$

and for some large enough N

$$\begin{aligned} \int_{B_{2\rho} \setminus \overline{B_\rho}} \sigma_1(A) (\Delta \varphi)^2 d\mu_g &\leq C \int_{B_{2\rho} \setminus \overline{B_\rho}} |\Delta_0 \varphi + O(r^{N-1}) \varphi'|^2 (1 + O(r^N)) dx \\ &\leq C \int_{B_{2\rho} \setminus \overline{B_\rho}} \left[(\Delta_0 \varphi)^2 + O(r^{2(N-1)}) |\varphi'|^2 \right] dx \\ &\leq C \int_{B_{2\rho} \setminus \overline{B_\rho}} \left(u_\epsilon \Delta_0 \eta_\rho + 2 \nabla u_\epsilon \cdot \nabla \eta_\rho + \eta_\rho \Delta_0 u_\epsilon \right)^2 dx + O(\epsilon^{n-6}) \\ &\leq C \int_\rho^{2\rho} \frac{(n\epsilon^2 + 4r^2)^2}{(\epsilon^2 + r^2)^4} u_\epsilon^2 r^{n-1} dr + O(\epsilon^{n-6}) \\ &\stackrel{\sigma=\frac{r}{\epsilon}}{\leq} C \epsilon^2 \int_{\rho/\epsilon}^{2\rho/\epsilon} \frac{(n + 4\sigma^2)^2 \sigma^{n-1}}{(1 + \sigma^2)^{n-2}} d\sigma + O(\epsilon^{n-6}) \\ &\leq C \epsilon^2 \left(\frac{\rho}{\epsilon} \right)^{8-n} + O(\epsilon^{n-6}) = O(\epsilon^{n-6}). \end{aligned}$$

Observe that

$$\int_{B_\rho} \frac{r^3 u_\epsilon^2}{(\epsilon^2 + r^2)^2} dx = \begin{cases} O(\epsilon^4) & \text{if } n = 10, \\ O(\epsilon^5 |\log \epsilon|) & \text{if } n = 11, \\ O(\epsilon^5) & \text{if } n \geq 12. \end{cases} \quad (3.1)$$

Hence,

$$\begin{aligned} & -\frac{n-2}{2} \int_M \sigma_1(A) (\Delta \varphi)^2 d\mu_g \\ &= \frac{(n-6)^2 (n-2) |W(p)|^2}{48n(n-1)} \omega_{n-1} \int_0^\rho \frac{(n\epsilon^2 + 4r^2)^2}{(\epsilon^2 + r^2)^4} u_\epsilon^2 r^{n+1} dr + \begin{cases} O(\epsilon^4) & \text{if } n = 10, \\ O(\epsilon^5 |\log \epsilon|) & \text{if } n = 11, \\ O(\epsilon^5) & \text{if } n \geq 12. \end{cases} \end{aligned}$$

Thirdly, we compute $\int_M T_2(\nabla \varphi, \nabla \Delta \varphi) d\mu_g$.

$$\int_{B_\rho} T_2(\nabla \varphi, \nabla \Delta \varphi) d\mu_g = \int_{B_\rho} \left[(n-2) \sigma_1(A) \langle \nabla \varphi, \nabla \Delta \varphi \rangle - 8 A_{ij} \varphi_{,i} (\Delta \varphi)_{,j} \right] d\mu_g.$$

Observe that $u_{\epsilon,i} = \frac{x^i}{r} u'_\epsilon$ and $(\Delta_0 u_\epsilon)_{,i} = \frac{x^i}{r} (\Delta_0 u_\epsilon)'$. Then we get

$$\begin{aligned}
& (n-2) \int_{B_\rho} \sigma_1(A) \langle \nabla \varphi, \nabla \Delta \varphi \rangle d\mu_g \\
&= (n-2) \int_{B_\rho} \left(\frac{1}{2} \sigma_1(A)_{,ij}(p) x^i x^j + O(r^3) \right) g^{kl} \varphi_{,k} (\Delta \varphi)_{,l} d\mu_g \\
&= (n-2) \int_{B_\rho} \left(\frac{1}{2} \sigma_1(A)_{,ij}(p) x^i x^j + O(r^3) \right) (\delta^{kl} + O(r^2)) \varphi_{,k} (\Delta_0 \varphi + O(r^{N-1}) \varphi')_{,l} d\mu_g \\
&= \frac{n-2}{2} \int_{B_\rho} \frac{1}{n} \Delta \sigma_1(A)(p) |x|^2 \varphi_{,i} (\Delta_0 \varphi)_{,i} dx + \int_{B_\rho} O(r^3) |\varphi'| |(\Delta_0 \varphi)'| dx \\
&= -\frac{(n-2)|W(p)|^2}{24n(n-1)} \int_{B_\rho} \left\{ -(n-6)^2(n-4) \frac{u_\epsilon^2 r^4}{(\epsilon^2 + r^2)^4} [(n+2)\epsilon^2 + 4r^2] \right\} dx \\
&\quad + \int_{B_\rho} \frac{O(r^3) u_\epsilon^2}{(\epsilon^2 + r^2)^2} dx \\
&= \frac{(n-2)(n-4)(n-6)^2}{24n(n-1)} |W(p)|^2 \int_{B_\rho} \frac{r^4}{(\epsilon^2 + r^2)^4} u_\epsilon^2 [(n+2)\epsilon^2 + 4r^2] dx \\
&\quad + \int_{B_\rho} \frac{O(r^3) u_\epsilon^2}{(\epsilon^2 + r^2)^2} dx,
\end{aligned}$$

and

$$\begin{aligned}
& -8 \int_{B_\rho} A_{ij} \varphi_{,i} (\Delta \varphi)_{,j} d\mu_g \\
&= -8 \int_{B_\rho} \left(A_{ij,k}(p) x^k + \frac{1}{2} A_{ij,kl}(p) x^k x^l + O(r^3) \right) \varphi_{,i} (\Delta_0 \varphi + O(r^{N-1}) \varphi')_{,j} d\mu_g \\
&= -4 \int_{B_\rho} A_{ij,kl}(p) x^k x^l x^i x^j \left[-(n-4)(n-6)^2 \frac{u_\epsilon^2}{(\epsilon^2 + r^2)^4} [(n+2)\epsilon^2 + 4r^2] \right] dx \\
&\quad + \int_{B_\rho} O(r^3) |\varphi'| |(\Delta_0 \varphi)'| dx \\
&= 4(n-4)(n-6)^2 \int_{B_\rho} \left[-\frac{2}{9} \frac{1}{n-2} \sum_{k,l} (W_{iklj}(p) x^i x^j)^2 - \frac{\sigma_1(A)_{,ij}(p) x^i x^j r^2}{n-2} \right] \\
&\quad \times \frac{u_\epsilon^2}{(\epsilon^2 + r^2)^4} [(n+2)\epsilon^2 + 4r^2] dx + \int_{B_\rho} \frac{O(r^3) u_\epsilon^2}{(\epsilon^2 + r^2)^2} dx \\
&= -\frac{8(n-4)(n-6)^2}{9(n-2)} \int_{B_\rho} \left[\sum_{k,l} (W_{iklj}(p) W_{sklt}(p) x^i x^j x^s x^t) \frac{u_\epsilon^2}{(\epsilon^2 + r^2)^4} [(n+2)\epsilon^2 + 4r^2] \right] dx \\
&\quad - \frac{4(n-4)(n-6)^2}{n(n-2)} \int_{B_\rho} \frac{\Delta \sigma_1(A)(p) r^4}{(\epsilon^2 + r^2)^4} u_\epsilon^2 [(n+2)\epsilon^2 + 4r^2] dx + \int_{B_\rho} \frac{O(r^3) u_\epsilon^2}{(\epsilon^2 + r^2)^2} dx \\
&= -\frac{(n-4)(n-6)^2}{(n-1)n(n+2)} \omega_{n-1} |W(p)|^2 \int_0^\rho \frac{r^{n+3} u_\epsilon^2}{(\epsilon^2 + r^2)^4} [(n+2)\epsilon^2 + 4r^2] dr \\
&\quad + \int_{B_\rho} \frac{O(r^3) u_\epsilon^2}{(\epsilon^2 + r^2)^2} dx,
\end{aligned}$$

where the last identity follows from

$$\begin{aligned}
& \sum_{k,l} W_{iklj}(p) W_{sklt}(p) \int_{B_\rho} x^i x^j x^s x^t \frac{u_\epsilon^2}{(\epsilon^2 + r^2)^4} [(n+2)\epsilon^2 + 4r^2] dx \\
&= \sum_{k,l} W_{iklj}(p) W_{sklt}(p) \int_{\mathbb{S}^{n-1}} \xi^i \xi^j \xi^s \xi^t d\mu_{\mathbb{S}^{n-1}} \int_0^\rho r^{n+3} \frac{u_\epsilon^2}{(\epsilon^2 + r^2)^4} [(n+2)\epsilon^2 + 4r^2] dr \\
&= \frac{\omega_{n-1}}{n(n+2)} \sum_{k,l} W_{iklj}(p) W_{sklt}(p) [\delta_{ij}\delta_{st} + \delta_{is}\delta_{jt} + \delta_{it}\delta_{js}] \int_0^\rho \frac{r^{n+3} u_\epsilon^2}{(\epsilon^2 + r^2)^4} [(n+2)\epsilon^2 + 4r^2] dr \\
&= \frac{\omega_{n-1}}{n(n+2)} \left[|W(p)|^2 + W_{iklj}(p) W_{jkli}(p) \right] \int_0^\rho r^{n+3} \frac{u_\epsilon^2}{(\epsilon^2 + r^2)^4} [(n+2)\epsilon^2 + 4r^2] dr \\
&= \frac{3}{2} \frac{\omega_{n-1}}{n(n+2)} |W(p)|^2 \int_0^\rho r^{n+3} \frac{u_\epsilon^2}{(\epsilon^2 + r^2)^4} [(n+2)\epsilon^2 + 4r^2] dr.
\end{aligned}$$

Then we have

$$\begin{aligned}
& -2 \int_{B_\rho} T_2(\nabla\varphi, \nabla\Delta\varphi) d\mu_g \\
&= -\frac{(n^2 - 28)(n-4)(n-6)^2}{12n(n-1)(n+2)} |W(p)|^2 \omega_{n-1} \int_0^\rho r^{n+3} \frac{u_\epsilon^2}{(\epsilon^2 + r^2)^4} [(n+2)\epsilon^2 + 4r^2] dr \\
& \quad + \int_{B_\rho} \frac{O(r^3)u_\epsilon^2}{(\epsilon^2 + r^2)^2} dx.
\end{aligned}$$

By a similar argument, one has

$$\begin{aligned}
\left| \int_{B_{2\rho} \setminus \overline{B_\rho}} T_2(\nabla\varphi, \nabla\Delta\varphi) \right| &\leq C \int_{B_{2\rho} \setminus \overline{B_\rho}} |\nabla\varphi| |\nabla\Delta\varphi| d\mu_g \\
&\leq C \int_{B_{2\rho} \setminus \overline{B_\rho}} |u'_\epsilon| |(\Delta u_\epsilon)'| dx + O(\epsilon^{n-6}) = O(\epsilon^{n-6}).
\end{aligned}$$

Fourthly, we compute $\int_M T_4(\nabla\varphi, \nabla\varphi) d\mu_g$.

$$\begin{aligned}
& (n-6) \int_{B_\rho} \Delta\sigma_1(A) |\nabla\varphi|_g^2 d\mu_g \\
&= (n-6) \int_{B_\rho} (\Delta\sigma_1(A)(p) + O(r)) (|\varphi'|^2 + O(r^2)|\varphi|^2) dx \\
&= -(n-6)^3 \frac{|W(p)|^2}{12(n-1)} \int_{B_\rho} \frac{u_\epsilon^2 r^2}{(\epsilon^2 + r^2)^2} dx + \int_{B_\rho} \frac{O(r^3)u_\epsilon^2}{(\epsilon^2 + r^2)^2} dx.
\end{aligned}$$

Using (A.5), we get

$$\begin{aligned}
& -\frac{16}{n-4} \int_{B_\rho} B_{ij\varphi, i\varphi, j} d\mu_g = -\frac{16}{n-4} \int_{B_\rho} (n-6)^2 u_\epsilon^2 \frac{B_{ij} x^i x^j}{(\epsilon^2 + r^2)^2} dx \\
&= -\frac{16(n-6)^2}{n-4} \int_{B_\rho} \left[-\frac{2}{9} \frac{1}{n-2} \sum_{k,l,s} [(W_{ikls}(p) + W_{ilks}(p)) x^i]^2 \right.
\end{aligned}$$

$$\begin{aligned}
& + \frac{1}{12(n-2)(n-1)} |W(p)|^2 r^2 - \frac{7n-8}{n-2} \sigma_1(A)_{,ij}(p) x^i x^j + O(r^3) \Big] \frac{u_\epsilon^2}{(\epsilon^2 + r^2)^2} dx \\
& = - \frac{16(n-6)^2}{n-4} \left[-\frac{2}{3n(n-2)} + \frac{1}{12(n-2)(n-1)} + \frac{7n-8}{12(n-2)(n-1)n} \right] |W(p)|^2 \int_{B_\rho} \frac{r^2 u_\epsilon^2}{(\epsilon^2 + r^2)^2} dx \\
& \quad + \int_{B_\rho} \frac{O(r^3) u_\epsilon^2}{(\epsilon^2 + r^2)^2} dx \\
& = \int_{B_\rho} \frac{O(r^3) u_\epsilon^2}{(\epsilon^2 + r^2)^2} dx,
\end{aligned}$$

where the second identity follows from

$$\sum_{i,k,l,s} (W_{ikls}(p) + W_{ilks}(p))^2 = 2|W(p)|^2 + 2 \sum_{i,k,l,s} W_{ikls}(p) W_{ilks}(p) = 3|W(p)|^2,$$

in view of

$$\begin{aligned}
0 & = W_{ikls}(W_{ilks} + W_{iksl} + W_{islk}) \\
& = W_{ikls}W_{ilks} + W_{ikls}W_{iksl} + W_{ikls}W_{islk} \\
& = 2W_{ikls}W_{ilks} - |W|^2 \quad \text{at } p.
\end{aligned}$$

Also we have

$$\int_{B_{2\rho} \setminus \overline{B_\rho}} T_4(\nabla \varphi, \nabla \varphi) d\mu_g \leq C \int_{B_{2\rho} \setminus \overline{B_\rho}} |\nabla \varphi|_g^2 d\mu_g = O(\epsilon^{n-6}).$$

Hence, collecting the above terms together with (3.1), we obtain

$$\begin{aligned}
& - \int_M T_4(\nabla \varphi, \nabla \varphi) d\mu_g \\
& = - (n-6) \int_{B_\rho} \Delta \sigma_1(A) |\nabla \varphi|_g^2 d\mu_g + \frac{16}{n-4} \int_{B_\rho} B_{ij} \varphi_{,i} \varphi_{,j} d\mu_g + O(\epsilon^{n-6}) \\
& = (n-6)^3 \frac{|W(p)|^2}{12(n-1)} \int_{B_\rho} \frac{u_\epsilon^2 r^2}{(\epsilon^2 + r^2)^2} dx + \begin{cases} O(\epsilon^4) & \text{if } n = 10, \\ O(\epsilon^5 |\log \epsilon|) & \text{if } n = 11, \\ O(\epsilon^5) & \text{if } n \geq 12. \end{cases}
\end{aligned}$$

Finally, we compute $\frac{n-6}{2} \int_M Q_g \varphi^2 d\mu_g$. By the definition (1.1) of Q_g , integration by parts gives

$$\begin{aligned}
& \frac{n-6}{2} \int_M Q_g \varphi^2 d\mu_g \\
& = \frac{n-6}{2} \int_M \Delta^2 \sigma_1(A) \varphi^2 d\mu_g + \int_{B_\rho} \frac{O(r^3) u_\epsilon^2}{(\epsilon^2 + r^2)^2} dx + O(\epsilon^{n-6}) \\
& = \frac{n-6}{2} \int_M \Delta \sigma_1(A) \Delta \varphi^2 d\mu_g + \int_{B_\rho} \frac{O(r^3) u_\epsilon^2}{(\epsilon^2 + r^2)^2} dx + O(\epsilon^{n-6}) \\
& = - \frac{(n-6)^2 |W(p)|^2}{12(n-1)} \omega_{n-1} \int_0^\rho \frac{u_\epsilon^2 r^{n-1}}{(\epsilon^2 + r^2)^2} [(n-10)r^2 - n\epsilon^2] dr + \begin{cases} O(\epsilon^4) & \text{if } n = 10, \\ O(\epsilon^5 |\log \epsilon|) & \text{if } n = 11, \\ O(\epsilon^5) & \text{if } n \geq 12, \end{cases}
\end{aligned}$$

by (3.1), where the last identity follows from

$$\begin{aligned}
& \frac{n-6}{2} \int_{B_\rho} \Delta \sigma_1(A) \Delta \varphi^2 d\mu_g \\
&= \frac{n-6}{2} \int_{B_\rho} (\Delta \sigma_1(A)(p) + O(r)) (\Delta_0 \varphi^2 + O(r^{N-1})(\varphi^2)') dx \\
&= \frac{n-6}{2} \Delta \sigma_1(A)(p) \int_{B_\rho} 2(\varphi \Delta_0 \varphi + |\nabla \varphi|_0^2) dx + \int_{B_\rho} \frac{O(r) u_\epsilon^2}{\epsilon^2 + r^2} dx \\
&= -\frac{(n-6)^2 |W(p)|^2}{12(n-1)} \omega_{n-1} \int_0^\rho \frac{u_\epsilon^2 r^{n-1}}{(\epsilon^2 + r^2)^2} [(n-10)r^2 - n\epsilon^2] dr + \int_{B_\rho} \frac{O(r) u_\epsilon^2}{\epsilon^2 + r^2} dx
\end{aligned}$$

and the first identity follows from

$$\left| \int_{B_{2\rho} \setminus \overline{B_\rho}} Q_g \varphi^2 d\mu_g \right| \leq C \int_{B_{2\rho} \setminus \overline{B_\rho}} u_\epsilon^2 dx = O(\epsilon^{n-6}).$$

Therefore collecting all the above terms together, we obtain

$$\int_M \varphi P_g \varphi d\mu_g = \int_{\mathbb{R}^n} |\nabla \Delta_0 u_\epsilon|^2 dx + A_{n,\rho,\epsilon} |W(p)|^2 \omega_{n-1} + O(\epsilon^{\min\{n-6,5\}}),$$

where $A_{n,\rho,\epsilon}$ is a constant given by

$$\begin{aligned}
& (n-6)^2 \left\{ \frac{n-2}{48n(n-1)} \int_0^\rho \frac{(n\epsilon^2 + 4r^2)^2}{(\epsilon^2 + r^2)^4} u_\epsilon^2 r^{n+1} dr + \frac{n-6}{12(n-1)} \int_0^\rho \frac{u_\epsilon^2 r^{n+1}}{(\epsilon^2 + r^2)^2} dr \right. \\
& - \frac{1}{12(n-1)} \int_0^\rho \frac{u_\epsilon^2 r^{n-1}}{(\epsilon^2 + r^2)^2} [(n-10)r^2 - n\epsilon^2] dr \\
& \left. - \frac{(n^2 - 28)(n-4)}{12n(n-1)(n+2)} \int_0^\rho r^{n+3} \frac{u_\epsilon^2}{(\epsilon^2 + r^2)^4} [(n+2)\epsilon^2 + 4r^2] dr \right\} \\
&= 2^{n-6} \frac{(n-6)^2}{12(n-1)} \epsilon^4 \left\{ \frac{n-2}{4n} \int_0^{\rho/\epsilon} \frac{(n+4\sigma^2)^2}{(1+\sigma^2)^4} (1+\sigma^2)^{-(n-6)} \sigma^{n+1} d\sigma \right. \\
& + (n-6) \int_0^{\rho/\epsilon} \frac{1}{(1+\sigma^2)^2} (1+\sigma^2)^{-(n-6)} \sigma^{n+1} d\sigma \\
& - \int_0^{\rho/\epsilon} \frac{1}{(1+\sigma^2)^2} (1+\sigma^2)^{-(n-6)} \sigma^{n-1} [(n-10)\sigma^2 - n] d\sigma \\
& \left. - \frac{(n^2 - 28)(n-4)}{n(n+2)} \int_0^{\rho/\epsilon} \frac{\sigma^{n+3}}{(1+\sigma^2)^4} (1+\sigma^2)^{-(n-6)} [(n+2) + 4\sigma^2] d\sigma \right\},
\end{aligned}$$

where $r = \epsilon\sigma$. When $n = 10$, we claim that the leading term of the constant in the brace on the right hand side of the above identity:

$$\begin{aligned}
& \frac{1}{5} \int_0^{\rho/\epsilon} \frac{(4\sigma^2 + 10)^2}{(1+\sigma^2)^4} (1+\sigma^2)^{-4} \sigma^{11} d\sigma + \int_0^{\rho/\epsilon} \frac{1}{(1+\sigma^2)^2} (1+\sigma^2)^{-4} (4\sigma^2 + 10) \sigma^9 d\sigma \\
& - \frac{18}{5} \int_0^{\rho/\epsilon} \frac{1}{(1+\sigma^2)^4} (1+\sigma^2)^{-4} (4\sigma^2 + 12) \sigma^{13} d\sigma
\end{aligned}$$

is a negative constant multiple of $|\log \epsilon|$. To see this, it is obviously true for the third term and the first two terms equal

$$\begin{aligned} & \frac{1}{5} \int_0^{\rho/\epsilon} \left\{ \sigma^2[(4\sigma^2 + 10)^2 - 18\sigma^2(4\sigma^2 + 12)] + 5(4\sigma^2 + 10)(1 + \sigma^2)^2 \right\} (1 + \sigma^2)^{-8} \sigma^9 d\sigma \\ &= \frac{1}{5} \int_0^{\rho/\epsilon} (-36\sigma^6 - 46\sigma^4 + 220\sigma^2 + 50)(1 + \sigma^2)^{-8} \sigma^9 d\sigma, \end{aligned}$$

whose leading term is also a negative constant multiple of $|\log \epsilon|$. For $n \geq 11$, let $t = \sigma^2$, the limit of the coefficient of $|W(p)|^2 \omega_{n-1}$ as $\epsilon \rightarrow 0$ is

$$\begin{aligned} & 2^{n-7} \frac{(n-6)^2}{12(n-1)} \epsilon^4 \left\{ \frac{n-2}{4n} \int_0^\infty \frac{(n+4t)^2}{(1+t)^{n-2}} t^{\frac{n}{2}} dt \right. \\ & + (n-6) \int_0^\infty \frac{1}{(1+t)^{n-4}} t^{\frac{n}{2}} dt - \int_0^\infty \frac{(n-10)t - n}{(1+t)^{n-4}} t^{\frac{n}{2}-1} dt \\ & \left. - \frac{(n^2-28)(n-4)}{n(n+2)} \int_0^\infty \frac{(n+2)+4t}{(1+t)^{n-2}} t^{\frac{n}{2}+1} dt \right\}. \end{aligned}$$

With the help of Beta function: $\int_0^\infty \frac{x^{\alpha-1}}{(1+x)^{\alpha+\beta}} dx = B(\alpha, \beta) = \frac{\Gamma(\alpha)\Gamma(\beta)}{\Gamma(\alpha+\beta)}$ for $\text{Re}(\alpha) > 0, \text{Re}(\beta) > 0$, we have

$$\begin{aligned} & \frac{n-2}{4n} \int_0^\infty \frac{(n+4t)^2}{(1+t)^{n-2}} t^{\frac{n}{2}} dt \\ &= \frac{n-2}{4n} \int_0^\infty \frac{(n-4)^2 + 8(n-4)(1+t) + 16(1+t)^2}{(1+t)^{n-2}} t^{\frac{n}{2}} dt \\ &= \frac{n-2}{4n} \left[(n-4)^2 B\left(\frac{n}{2}+1, \frac{n}{2}-3\right) + 8(n-4) B\left(\frac{n}{2}+1, \frac{n}{2}-4\right) + 16 B\left(\frac{n}{2}+1, \frac{n}{2}-5\right) \right], \\ & (n-6) \int_0^\infty \frac{1}{(1+t)^{n-4}} t^{\frac{n}{2}} dt = (n-6) B\left(\frac{n}{2}+1, \frac{n}{2}-5\right), \\ & - \int_0^\infty \frac{(n-10)t - n}{(1+t)^{n-4}} t^{\frac{n}{2}-1} dt = -(n-10) B\left(\frac{n}{2}+1, \frac{n}{2}-5\right) + n B\left(\frac{n}{2}, \frac{n}{2}-4\right), \\ & - \frac{(n^2-28)(n-4)}{n(n+2)} \int_0^\infty \frac{(n+2)+4t}{(1+t)^{n-2}} t^{\frac{n}{2}+1} dt \\ &= - \frac{(n^2-28)(n-4)}{n(n+2)} \int_0^\infty \frac{4(1+t)^2 + (n-6)(1+t) - (n-2)}{(1+t)^{n-2}} t^{\frac{n}{2}} dt \\ &= - \frac{4(n^2-28)(n-4)}{n(n+2)} B\left(\frac{n}{2}+1, \frac{n}{2}-5\right) - \frac{(n^2-28)(n-4)(n-6)}{n(n+2)} B\left(\frac{n}{2}+1, \frac{n}{2}-4\right) \\ & + \frac{(n^2-28)(n-4)(n-2)}{n(n+2)} B\left(\frac{n}{2}+1, \frac{n}{2}-3\right). \end{aligned}$$

Hence, the above limit of the coefficient of $|W(p)|^2 \omega_{n-1}$ is rewritten as

$$2^{n-7} \frac{(n-6)^2}{12(n-1)} \epsilon^4 \left\{ n B\left(\frac{n}{2}, \frac{n}{2}-4\right) \right.$$

$$\begin{aligned}
& + B\left(\frac{n}{2} + 1, \frac{n}{2} - 3\right) \left[\frac{n-2}{4n} (n-4)^2 + \frac{(n^2-28)(n-4)(n-2)}{n(n+2)} \right] \\
& + B\left(\frac{n}{2} + 1, \frac{n}{2} - 4\right) \left[\frac{2(n-2)(n-4)}{n} - \frac{(n^2-28)(n-4)(n-6)}{n(n+2)} \right] \\
& + B\left(\frac{n}{2} + 1, \frac{n}{2} - 5\right) \left[\frac{4(n-2)}{n} - n + 10 + n - 6 - \frac{4(n^2-28)(n-4)}{n(n+2)} \right] \Big\} \\
& = 2^{n-7} \frac{(n-6)^2}{12(n-1)} B\left(\frac{n}{2} + 1, \frac{n}{2} - 5\right) \epsilon^4 \Big\{ (n-10) \\
& + \frac{(n-2)(\frac{n}{2}-4)(\frac{n}{2}-5)}{4n(n+2)(n-3)} (5n^2 - 2n - 120) + \frac{\frac{n}{2}-5}{n(n+2)} (-n^3 + 8n^2 + 28n - 176) \\
& + \frac{4}{n(n+2)} (-n^3 + 6n^2 + 30n - 116) \Big\}, \tag{3.2}
\end{aligned}$$

where we have used some elementary identities

$$B\left(\frac{n}{2} + 1, \frac{n}{2} - 3\right) = \frac{\Gamma(\frac{n}{2} + 1)\Gamma(\frac{n}{2} - 3)}{\Gamma(n-2)} = \frac{(\frac{n}{2}-4)(\frac{n}{2}-5)}{(n-3)(n-4)} B\left(\frac{n}{2} + 1, \frac{n}{2} - 5\right),$$

$$B\left(\frac{n}{2} + 1, \frac{n}{2} - 4\right) = \frac{\frac{n}{2}-5}{n-4} B\left(\frac{n}{2} + 1, \frac{n}{2} - 5\right),$$

$$B\left(\frac{n}{2}, \frac{n}{2} - 4\right) = \frac{\Gamma(\frac{n}{2})\Gamma(\frac{n}{2} - 4)}{\Gamma(n-4)} = \frac{n-10}{n} B\left(\frac{n}{2} + 1, \frac{n}{2} - 5\right).$$

The constant in the last brace of (3.2) when $n \geq 11$ is

$$\begin{aligned}
& n - 10 + \frac{1}{16n(n+2)(n-3)} \Big\{ (n-2)(n-8)(n-10)(5n^2 - 2n - 120) \\
& + 8(n-3)[(n-10)(-n^3 + 8n^2 + 28n - 176) + 8(-n^3 + 6n^2 + 30n - 116)] \Big\} \\
& = n - 10 + \frac{1}{16n(n+2)(n-3)} \left[-3n^5 + 2n^4 + 228n^3 - 264n^2 - 1760n - 768 \right] \\
& = \frac{-3n^5 + 18n^4 + 52n^3 - 200n^2 - 800n - 768}{16n(n+2)(n-3)} < 0.
\end{aligned}$$

On the other hand, we have

$$\int_M \varphi^{\frac{2n}{n-6}} d\mu_g = \int_{B_\rho} u_\epsilon^{\frac{2n}{n-6}} d\mu_g + \int_{B_{2\rho} \setminus \overline{B}_\rho} \varphi^{\frac{2n}{n-6}} d\mu_g = \int_{\mathbb{R}^n} u_\epsilon^{\frac{2n}{n-6}} dx + O(\epsilon^n).$$

Therefore, putting these facts together, we conclude by Lemma 3.1 that

$$\begin{aligned}
& \frac{\int_M \varphi P_g \varphi d\mu_g}{\left(\int_M \varphi^{\frac{2n}{n-6}} d\mu_g \right)^{\frac{n-6}{n}}} \\
& = Y_6(S^n) + A_{n,\rho,\epsilon} |W(p)|^2 \omega_{n-1} + \begin{cases} O(\epsilon^4) & \text{if } n = 10, \\ O(\epsilon^5 |\log \epsilon|) & \text{if } n = 11, \\ O(\epsilon^5) & \text{if } n \geq 12, \end{cases}
\end{aligned}$$

$$= \begin{cases} Y_6(S^n) - C_n |W(p)|^2 \epsilon^4 |\log \epsilon| + O(\epsilon^4) & \text{if } n = 10, \\ Y_6(S^n) - C_n |W(p)|^2 \epsilon^4 + o(\epsilon^4) & \text{if } n \geq 11, \end{cases}$$

for some positive constant $C_n > 0$. Consequently, choosing ϵ sufficiently small, we obtain $Y_6(M^n) < Y_6(S^n)$. This finishes the proof. $\square$

Given a smooth positive function f in M^n , we define a “free” energy functional by

$$E_f[u] = \frac{1}{2} \int_M u P_g u d\mu_g - \frac{1}{2^\sharp} \int_M f |u|^{2^\sharp} d\mu_g.$$

Let $u_{,i}$ or $\nabla_i u$ denote the covariant derivatives of u with respect to metric g and R_{ijk}^l be Riemannian curvature tensor of metric g . Notice that

$$\nabla_j \nabla_i \nabla^i u = \nabla_i \nabla_j \nabla^i u + R_{ij}^k \nabla_k u = \nabla_i \nabla^i \nabla_j u - R_j^k \nabla_k u,$$

there holds

$$\int_M |\nabla \Delta u|_g^2 d\mu_g = \int_M |\Delta \nabla_j u - R_j^k \nabla_k u|_g^2 d\mu_g. \quad (3.3)$$

Under g -normal coordinates around a point, one gets

$$\begin{aligned} & \frac{1}{2} \Delta_g |\nabla^2 u|_g^2 \\ &= |\nabla^3 u|_g^2 + \langle \nabla \Delta \nabla_i u, \nabla \nabla^i u \rangle_g + u_{,ij} (R_{ijk}^l u_{,lk} + R_j^l u_{,il} + R_{ijk,k}^l u_{,l} + R_{ijk}^l u_{,lk}). \end{aligned}$$

Integrating the above identity over M to show

$$\int_M |\Delta \nabla u|_g^2 d\mu_g = \int_M |\nabla^3 u|_g^2 d\mu_g + \int_M O(|\text{Rm}| |\nabla^2 u|_g + |\nabla \text{Rm}| |\nabla u|_g) |\nabla^2 u|_g d\mu_g. \quad (3.4)$$

From (3.3) and (3.4), it yields that the following two norms are equivalent:

$$\begin{aligned} \|u\|_{H^3} &:= \left(\int_M (|\nabla \Delta u|_g^2 d\mu_g + |\nabla^2 u|_g^2 + |\nabla u|_g^2 + u^2) d\mu_g \right)^{\frac{1}{2}} \\ &\approx \left(\int_M (|\nabla^3 u|_g^2 d\mu_g + |\nabla^2 u|_g^2 + |\nabla u|_g^2 + u^2) d\mu_g \right)^{\frac{1}{2}}, \quad u \in H^3(M, g). \end{aligned}$$

Let $\|\cdot\|_p$ denote the norm of $L^p(M, g)$ for $1 \leq p \leq \infty$.

A sequence $\{u_k\}$ in $H^3(M, g)$ is called a Palais-Smale (P-S) $_\beta$ sequence for E_f if $E_f[u_k] \rightarrow \beta \in \mathbb{R}$ and $DE_f[u_k] \rightarrow 0$ as $k \rightarrow \infty$. The energy E_f satisfies (P-S) $_\beta$ condition if any Palais-Smale sequence of E_f has a strongly convergent subsequence. We call that P_g is coercive if there exists a constant $\mu(g) > 0$ such that

$$\int_M \psi P_g \psi d\mu_g \geq \mu(g) \int_M \psi^2 d\mu_g, \quad \text{for all } \psi \in H^3(M, g).$$

Remark 3.1. If (M, g) is Einstein and of positive constant scalar curvature, from the factorization (1.4) of P_g , the coercivity of P_g is automatically satisfied.

As an application, we adapt some arguments in Esposito-Robert [4] to show some existence results of prescribed Q -curvature equation, whose solution may change signs due to the lack of maximum principles (in general).

Theorem 3.1. *Let (M^n, g) be a smooth closed manifold of dimension $n \geq 10$ and f be a smooth positive function in M^n . Suppose the Weyl tensor W_g is nonzero at a maximum point of f and f satisfies the vanishing order condition (1.5) at this maximum point. Assume P_g is coercive, then there exists a nontrivial $C^{6,\mu}$ ($0 < \mu < 1$) solution to*

$$P_g u = f|u|^{2^\sharp-2}u \text{ in } M. \quad (3.5)$$

In addition, assume (M, g) is Einstein and of positive scalar curvature, then there exists a smooth solution to the Q -curvature equation

$$P_g u = f u^{\frac{n+6}{n-6}}, u > 0 \text{ in } M. \quad (3.6)$$

Proof. By the assumptions, there exists $p \in M$ such that $f(p) = \max_{x \in M^n} f(x)$, $W_g(p) \neq 0$ and the vanishing order condition (1.5) of f is true at p . Let

$$\gamma_\epsilon(t) = t \frac{\varphi}{\|f^{\frac{1}{2^\sharp}} \varphi\|_{2^\sharp}},$$

where $\varphi = \eta_\rho u_\epsilon$ is the test function chosen in Proposition 3.1. By choosing t_0 large enough, we get $E[\gamma_\epsilon(t_0)] < 0$. Let

$$\Gamma = \left\{ \gamma(t) \in C([0, t_0], H^3(M, g)); \gamma(0) = 0, \gamma(t_0) = t_0 \frac{\varphi}{\|f^{\frac{1}{2^\sharp}} \varphi\|_{2^\sharp}} \right\}.$$

From the coercivity of P_g and Sobolev embedding theorem, we have

$$E_f\left[\frac{\varphi}{\|f^{\frac{1}{2^\sharp}} \varphi\|_{2^\sharp}}\right] = \frac{1}{2} \frac{\int_M \varphi P_g \varphi d\mu_g}{\|f^{\frac{1}{2^\sharp}} \varphi\|_{2^\sharp}^2} - \frac{1}{2^\sharp} \geq \frac{1}{2}C - \frac{1}{2^\sharp}.$$

It only suffices to estimate the term:

$$\begin{aligned} \int_M f \varphi^{\frac{2n}{n-6}} d\mu_g &= \int_{B_\rho} \left[f(p) + \sum_{k=2}^4 \frac{1}{k!} \partial_{i_1 \dots i_k} f(p) x^{i_1} \dots x^{i_k} + O(|x|^5) \right] u_\epsilon^{2^\sharp} dx + O(\epsilon^n) \\ &= f(p) \int_{\mathbb{R}^n} u_\epsilon^{\frac{2n}{n-6}} dx + \begin{cases} O(\epsilon^4) & \text{if } n = 10, \\ o(\epsilon^4) & \text{if } n \geq 11, \end{cases} \end{aligned}$$

where the second equality follows from the vanishing order condition (1.5) of f at p . From this and some existing estimates in the proof of Proposition 3.1, we conclude that there exist some sufficiently small $\epsilon > 0$ and a constant $C'_n > 0$ such that

$$\sup_{t \geq 0} E_f[\gamma_\epsilon(t)] = E_f[\gamma_\epsilon(t^*)] = \frac{3}{n} \left(\frac{\int_M \varphi P_g \varphi d\mu_g}{\|f^{\frac{1}{2^\sharp}} \varphi\|_{2^\sharp}^2} \right)^{\frac{2^\sharp}{2^\sharp-2}}$$

$$\leq \begin{cases} \frac{3}{n}(\max_M f)^{\frac{6-n}{6}} Y_6(S^n)^{\frac{n}{6}} - C'_n |W(p)|^2 \epsilon^4 |\log \epsilon| + O(\epsilon^4) & \text{if } n = 10, \\ \frac{3}{n}(\max_M f)^{\frac{6-n}{6}} Y_6(S^n)^{\frac{n}{6}} - C'_n |W(p)|^2 \epsilon^4 + o(\epsilon^4) & \text{if } n \geq 11, \end{cases}$$

where $t^* = \left(\frac{\int_M \varphi P_g \varphi d\mu_g}{\|f^{\frac{1}{2^\sharp}} \varphi\|_{2^\sharp}^2} \right)^{\frac{1}{2^\sharp-2}}$. Then it follows from Mountain Pass Lemma (cf. [1] or [4, Proposition 1]) that

$$\beta = \inf_{\gamma \in \Gamma} \sup_{0 \leq t \leq t_0} E_f[\gamma(t)] \leq \sup_{t \geq 0} E_f[\gamma_\epsilon(t)] < \frac{3}{n} Y_6(S^n)^{\frac{n}{6}} (\max_M f)^{\frac{6-n}{6}}$$

is a critical value of E_f and there exists a (P-S) $_\beta$ sequence $\{u_k\}$ of E_f in $H^3(M, g)$.

Next we claim that E_f satisfies (P-S) $_\beta$ condition. For the above (P-S) $_\beta$ sequence $\{u_k\}$ satisfying $E_f[u_k] \rightarrow \beta$ and $DE_f[u_k] \rightarrow 0$ as $k \rightarrow \infty$, there holds

$$2\beta + o(\|u_k\|_{H^3}) = 2E_f[u_k] - \langle DE_f[u_k], u_k \rangle = \frac{6}{n} \int_M f |u_k|^{2^\sharp} d\mu_g.$$

Together with the coercivity of P_g , one has

$$\mu(g) \|u_k\|_{H^3} \leq 2E_f[u_k] + \frac{2}{2^\sharp} \int_M f |u_k|^{2^\sharp} d\mu_g \leq C + o(\|u_k\|_{H^3}).$$

From this, we get $\{u_k\}$ is bounded in $H^3(M, g)$. Then up to a subsequence, as $k \rightarrow \infty$, $u_k \rightharpoonup u$ in $H^3(M, g)$ and $u_k \rightarrow u$ in $L^p(M, g)$ for $1 \leq p < 2^\sharp$. It is easy to verify that u is a weak solution to (3.5), that is, for all $\psi \in H^3(M, g)$,

$$\int_M \psi P_g u d\mu_g = \int_M f |u|^{2^\sharp-2} u \psi d\mu_g.$$

Choosing $\psi = u$, one has

$$\int_M u P_g u d\mu_g = \int_M f |u|^{2^\sharp} d\mu_g,$$

whence

$$E_f[u] = \frac{3}{n} \int_M f |u|^{2^\sharp} d\mu_g \geq 0.$$

Applying Brezis-Lieb lemma to

$$\begin{aligned} \int_M |\nabla \Delta u_k|_g^2 d\mu_g &= \int_M |\nabla \Delta u|_g^2 d\mu_g + \int_M |\nabla \Delta (u - u_k)|_g^2 d\mu_g + o(1), \\ \int_M f |u_k|^{2^\sharp} d\mu_g &= \int_M f |u|^{2^\sharp} d\mu_g + \int_M f |u - u_k|^{2^\sharp} d\mu_g + o(1), \end{aligned}$$

we have

$$\begin{aligned} E_f[u_k] - E_f[u] &= \frac{1}{2} \int_M |\nabla \Delta (u - u_k)|_g^2 d\mu_g - \frac{1}{2^\sharp} \int_M f |u - u_k|^{2^\sharp} d\mu_g + o(1) \\ &= E_f[u - u_k] + o(1). \end{aligned}$$

Since $DE_f[u_k] \rightarrow 0$ in $(H^3(M, g))'$, we have

$$\begin{aligned} o(1) &= \langle u_k - u, DE_f[u_k] \rangle = \langle u_k - u, DE_f[u_k] - DE_f[u] \rangle \\ &= \int_M |\nabla \Delta(u - u_k)|_g^2 d\mu_g - \int_M f|u - u_k|^{2^\sharp} d\mu_g + o(1). \end{aligned}$$

Thus, we obtain

$$\frac{3}{n} \int_M |\nabla \Delta(u - u_k)|_g^2 d\mu_g + o(1) = E_f[u_k - u] = E_f[u_k] - E_f[u] + o(1) \leq E_f[u_k] + o(1) \rightarrow \beta,$$

as $k \rightarrow \infty$, which yields

$$\int_M |\nabla \Delta(u - u_k)|_g^2 d\mu_g \leq \frac{n}{3} \beta + o(1). \quad (3.7)$$

Mimicking a cut-and-paste argument as in [3], we obtain that given $\epsilon > 0$, there exists a constant $B_\epsilon > 0$ such that

$$\left(\int_M |\psi|^{2^\sharp} d\mu_g \right)^{\frac{2}{2^\sharp}} \leq (1 + \epsilon) Y_6(S^n)^{-1} \int_M (|\nabla \Delta \psi|_g^2 + |\nabla^2 \psi|_g^2 + |\nabla \psi|_g^2) d\mu_g + B_\epsilon \int_M \psi^2 d\mu_g,$$

for all $\psi \in H^3(M, g)$. Choosing $\psi = u_k - u$ and k sufficiently large to get

$$\left(\int_M |u - u_k|^{2^\sharp} d\mu_g \right)^{\frac{2}{2^\sharp}} \leq (1 + \epsilon) Y_6(S^n)^{-1} \int_M |\nabla \Delta(u - u_k)|_g^2 d\mu_g + o(1).$$

Hence we have

$$\begin{aligned} o(1) &= \int_M |\nabla \Delta(u - u_k)|_g^2 d\mu_g - \int_M f|u - u_k|^{2^\sharp} d\mu_g \\ &\geq \int_M |\nabla \Delta(u - u_k)|_g^2 d\mu_g \left[1 - (\max_M f)(1 + \epsilon)^{\frac{2^\sharp}{2}} Y_6(S^n)^{-\frac{2^\sharp}{2}} \left(\int_M |\nabla \Delta(u - u_k)|_g^2 d\mu_g \right)^{\frac{6}{n-6}} \right]. \end{aligned}$$

From (3.7) and $\beta < \frac{3}{n} Y_6(S^n)^{\frac{n}{6}} (\max_M f)^{\frac{6-n}{6}}$, choosing ϵ sufficiently small, we get

$$o(1) \geq C \int_M |\nabla \Delta(u - u_k)|_g^2 d\mu_g.$$

Combining the above inequality and the coercivity of P_g to show that $u_k \rightarrow u$ in $H^3(M, g)$. Using the regularity result in Lemma 3.2 below, we know that $u \in C^{6,\mu}(M)$ for any $0 < \mu < 1$.

In addition, assume (M, g) is Einstein and has positive constant scalar curvature, we define the modified energy in $H^3(M, g)$ by

$$E_f^+[u] = \frac{1}{2} \int_M u P_g u d\mu_g - \frac{1}{2^\sharp} \int_M f u_+^{2^\sharp} d\mu_g,$$

where $u_+ = \max\{u, 0\}$. Using the above similar arguments associated with Mountain Pass Lemma and mimicking what we did in Lemma 3.2 below for E_f^+ , we get that there exists a nontrivial C^6 -solution u to

$$P_g u = f u_+^{\frac{n+6}{n-6}} \text{ in } M. \quad (3.8)$$

Since P_g is coercive by Remark 3.1, testing equation (3.8) with $u_- = \min\{u, 0\}$ we conclude that $u \geq 0$ in M . Together with R_g being a positive constant and the factorization (1.4) of GJMS operator:

$$\left(-\Delta_g + \frac{(n-6)(n+4)}{4n(n-1)}R_g\right)\left(-\Delta_g + \frac{(n-4)(n+2)}{4n(n-1)}R_g\right)\left(-\Delta_g + \frac{n-2}{4(n-1)}R_g\right)u \geq 0$$

and $u \not\equiv 0$ in M , we employ the maximum principle twice and strong maximum principle once for elliptic equations of second order to show that u is a positive solution to the equation (3.6). From this and Schauder estimates for elliptic equations, we conclude that $u \in C^\infty(M)$. This completes the proof. $\square$

We are now concerned with the regularity of mountain pass critical points for E .

Lemma 3.2. *Let (M, g) be a smooth closed Riemannian manifold of dimension $n \geq 7$. Assume $u \in H^3(M, g)$ is a weak solution of equation (3.5). Then $u \in C^{6, \mu}(M)$ for any $0 < \mu < 1$.*

Proof. Rewrite $P_g = (-\Delta_g)^3 - M_g + \frac{n-6}{2}Q_g$ by (1.2). Let $u \in H^3(M, g)$ be a weak solution of equation (3.5) and rewrite this equation as

$$\begin{aligned} (-\Delta_g + 1)^3 u &= M_g u + 3\Delta_g^2 u - 3\Delta_g u + (1 - \frac{n-6}{2}Q_g)u + f|u|^{2^\sharp-2}u \\ &:= b + f|u|^{2^\sharp-2}u, \end{aligned} \quad (3.9)$$

where $b \in H^{-1}(M, g)$. By the Sobolev embedding theorem we have $u \in L^{2^\sharp}(M, g)$ and $|u|^{2^\sharp-2} \in L^{\frac{n}{6}}(M, g)$. Given $\epsilon > 0$, there exist a $K_\epsilon > 0$ and a decomposition of $f|u|^{2^\sharp-2} = h_\epsilon + \eta_\epsilon$ with $\|h_\epsilon\|_{\frac{n}{6}} \leq \epsilon$, $\|\eta_\epsilon\|_\infty \leq K_\epsilon$. Inspired by the arguments in [4, Proposition 3], for $s > 1$ we define an operator

$$H_\epsilon : v \in L^s(M, g) \rightarrow (-\Delta_g + 1)^{-3}(h_\epsilon v) \in L^s(M, g).$$

Indeed, from Sobolev embedding theorem, the standard $W^{2,p}$ -regularity theory of the elliptic operator $-\Delta_g + 1$ and Hölder's inequality, we have

$$\begin{aligned} \|H_\epsilon v\|_s &\leq C \|(-\Delta_g + 1)^{-3}(h_\epsilon v)\|_{W^{6, \frac{ns}{n+6s}}} \leq C \|h_\epsilon v\|_{\frac{ns}{n+6s}} \\ &\leq C \|h_\epsilon\|_{\frac{n}{6}} \|v\|_s \leq C\epsilon \|v\|_s, \end{aligned}$$

where the constant C is independent of u . Choose $\epsilon > 0$ small enough, then the norm of H_ϵ on space $L^s(M, g)$ satisfies

$$\|H_\epsilon\|_{L^s \rightarrow L^s} \leq C\epsilon \leq \frac{1}{2}.$$

With the help of the operator H_ϵ , we rewrite equation (3.9) as

$$(Id - H_\epsilon)u = (-\Delta_g + 1)^{-3}(b + \eta_\epsilon u),$$

then it is easy to show $Id - H_\epsilon : L^s \rightarrow L^s$ is bounded and invertible. We intend to prove $u \in H^6(M, g)$. To see this, notice that $(-\Delta_g + 1)^{-3}(b + \eta_\epsilon u) \in H^5(M, g)$ since $b + \eta_\epsilon u \in H^{-1}(M, g)$. In the following, we first show $u \in H^4(M, g)$. Applying the Sobolev embedding

theorem and the L^s boundedness of the operator $(Id - H_\varepsilon)^{-1}$ to show that if $n \leq 10$, $u \in L^p(M, g)$ for all $p > 1$, and if $n > 10$, $u \in L^{\frac{2n}{n-10}}(M, g)$. In the latter case we have $|u|^{2^\sharp-2}u \in L^{\frac{2n(n-6)}{(n+6)(n-10)}}(M, g)$. From equation (3.9), we get

$$(-\Delta_g + 1)^2 u = (-\Delta_g + 1)^{-1} b + (-\Delta_g + 1)^{-1} (f|u|^{2^\sharp-2}u).$$

From $(-\Delta_g + 1)^{-1}(|u|^{2^\sharp-2}u) \in W^{2, \frac{2n(n-6)}{(n+6)(n-10)}}(M, g) \hookrightarrow L^2(M, g)$ and $(-\Delta_g + 1)^{-1}b \in L^2(M, g)$, we have $u \in H^4(M, g)$ in both cases. Repeat the above step with $u \in H^4(M, g)$ and $b \in L^2(M, g)$ in this situation. Notice that $(-\Delta_g + 1)^{-3}(b + \eta_\epsilon u) \in H^6(M, g)$, similar arguments in the above step show that if $n \leq 12$, $u \in L^p(M, g)$ for all $p > 1$ and if $n > 12$, $u \in L^{\frac{2n}{n-12}}(M, g)$. In the latter case, we get $|u|^{2^\sharp-2}u \in L^2(M, g)$ due to $\frac{2n(n-6)}{(n+6)(n-12)} > 2$. Hence we obtain $u \in H^6(M, g)$.

Finally we start with the classical bootstrap. We now construct a non-decreasing sequence $s_k \in \mathbb{R} \cup \{+\infty\}$ such that $u \in W^{6, s_k}(M, g)$ for all $k \in \mathbb{N}$. Set $s_0 = 2$, and find $k \geq 0$ such that $u \in W^{6, s_k}(M, g)$. Next we will define s_{k+1} by induction. The Sobolev embedding theorem yields

$$b \in L^{\frac{ns_k}{n-2s_k}}(M, g),$$

with the convention that $\frac{ns_k}{n-2s_k} = +\infty$ if $s_k \geq \frac{n}{2}$, and

$$|u|^{2^\sharp-2}u \in L^{\frac{ns_k(n-6)}{(n-6s_k)(n+6)}}(M, g),$$

with the convention that $\frac{ns_k}{n-6s_k} = +\infty$ if $s_k \geq \frac{n}{6}$. In view of equation (3.9), we have $u \in W^{6, s_{k+1}}(M, g)$ with $s_{k+1} = \min\{\frac{ns_k}{n-2s_k}, \frac{ns_k(n-6)}{(n-6s_k)(n+6)}\}$. If $s_k \in \mathbb{R}$ for all $k \in \mathbb{N}$, it must hold that $s_k \rightarrow +\infty$. Then we have $u \in W^{6, p}(M, g)$ for all $1 \leq p < +\infty$. If $s_k = +\infty$ for all $k \geq k_0 + 1$, then $s_{k_0} \geq \frac{n}{6}$, whence $b \in L^{\frac{n}{4}}(M, g)$ and $|u|^{2^\sharp-2}u \in L^q(M, g)$ for all $1 \leq q < +\infty$. The equation (3.9) leads to $u \in W^{6, \frac{n}{4}}(M)$. Repeating the argument twice, we obtain $u \in W^{6, p}(M, g)$ for all $1 \leq p < +\infty$. From this and the Sobolev embedding theorem, we have $u \in C^{5, \nu}(M)$ for all $0 < \nu < 1$. By the regularity theory for the classical solution of the elliptic operator $-\Delta_g + 1$, we get $u \in C^{6, \mu}(M)$ for some $0 < \mu < 1$. This completes the proof. $\square$

A Appendix: Proof of Lemma 2.1

As in Proposition 3.1, one may employ all computations under conformal normal coordinates of metric g around a point in M . From Lee-Park [12] that up to a conformal factor, under g -conformal normal coordinates around this point, for all $N \geq 5$ there hold

$$\sigma_1(A_g) = 0, \quad \sigma_1(A_g)_{,i} = 0, \quad \Delta_g \sigma_1(A_g) = -\frac{|W|_g^2}{12(n-1)}$$

at this point and

$$\sqrt{\det g} = 1 + O(r^N) \text{ near this point.}$$

To simplify the notation, we will omit the subscript g . Notice that

$$-P_g(r^{6-n}) = \left[\Delta^3 + \Delta \delta T_2 d + \delta T_2 d \Delta + \frac{n-2}{2} \Delta(\sigma_1(A) \Delta) + \delta T_4 d - \frac{n-6}{2} Q_g \right] (r^{6-n})$$

$$:= \sum_{k=1}^6 I_k.$$

Next, we begin to estimate all terms I_1 - I_6 .

For I_1 , let $u = u(r)$ be a radial function, we have

$$\begin{aligned} \Delta u(r) &= \Delta_0 u(r) + O(r^{N-1})u'; \\ \Delta^2 u(r) &= \Delta_0(\Delta_0 u(r) + O(r^{N-1})u') + O(r^{N-1})(\Delta_0 u(r) + O(r^{N-1})u')' \\ &= \Delta_0^2 u(r) + O(r^{N-1})u''' + O(r^{N-2})u'' + O(r^{N-3})u'; \\ \Delta^3 u(r) &= \Delta_0^3 u(r) + O(r^{N-1})u^{(5)} + O(r^{N-2})u^{(4)} + O(r^{N-3})u''' \\ &\quad + O(r^{N-4})u'' + O(r^{N-5})u'. \end{aligned}$$

Hence we obtain

$$I_1 = \Delta^3(r^{6-n}) = -c_n \delta_p + O(r^{N-n}).$$

To estimate I_2 . Notice that

$$\begin{aligned} I_2 &= \Delta \delta T_2 d(r^{6-n}) \\ &= -\Delta[(T_2)_{ij}(r^{6-n})_{,j}],_i \\ &= -\Delta[(T_2)_{ij,i}(r^{6-n})_{,j} + (T_2)_{ij}(r^{6-n})_{,ji}]. \end{aligned}$$

Using

$$\begin{aligned} (r^{6-n})_{,j} &= (6-n)r^{4-n}x^j, \\ (r^{6-n})_{,ji} &= (4-n)(6-n)r^{2-n}x^i x^j + (6-n)r^{4-n}\delta_{ij} + O(r^{6-n}), \end{aligned} \tag{A.1}$$

one has

$$\begin{aligned} (T_2)_{ij,i}(r^{6-n})_{,j} &= (n-10)\sigma_1(A)_{,j}(6-n)r^{4-n}x^j \\ &= (n-10)(6-n)\sigma_1(A)_{,j}x^j r^{4-n} \end{aligned}$$

and

$$\begin{aligned} &(T_2)_{ij}(r^{6-n})_{,ji} \\ &= [(n-2)\sigma_1(A)g_{ij} - 8A_{ij}](6-n)[(4-n)r^{2-n}x^i x^j + r^{4-n}\delta_{ij} + O(r^{6-n})] \\ &= (6-n)[4(n-4)\sigma_1(A)r^{4-n} - 8(4-n)A_{ij}x^i x^j r^{2-n}] + O(r^{7-n}). \end{aligned}$$

Hence, we obtain

$$\begin{aligned} I_2 &= - (6-n)\Delta[(n-10)\sigma_1(A)_{,j}x^j r^{4-n} + 4(n-4)\sigma_1(A)r^{4-n} - 8(4-n)A_{ij}x^i x^j r^{2-n}] \\ &= (n-6)\{(n-10)[4(4-n)\sigma_1(A)_{,j}x^j r^{2-n} + 2(4-n)\sigma_1(A)_{,jk}x^j x^k r^{2-n} \\ &\quad + \sigma_1(A)_{,jkk}x^j r^{2-n} + 2\Delta\sigma_1(A)r^{4-n}] + O(r^{5-n}) \end{aligned}$$

$$\begin{aligned}
& + 4(n-4)[\Delta\sigma_1(A)r^{4-n} + 2(4-n)\sigma_1(A)_{,k}x^k r^{2-n} + 2(4-n)\sigma_1(A)r^{2-n}] \\
& + 8(n-4)[4(2-n)A_{ij}x^i x^j r^{-n} + \Delta A_{ij}x^i x^j r^{2-n} + 4\sigma_1(A)_{,i}x^i r^{2-n} + 2\sigma_1(A)r^{2-n}]\} \\
& = (n-6)\left\{ -4(n-4)(3n-26)\sigma_1(A)_{,j}x^j r^{2-n} + 6(n-6)\Delta\sigma_1(A)r^{4-n} \right. \\
& \quad - 2(n-10)(n-4)\sigma_1(A)_{,jk}x^j x^k r^{2-n} + (n-10)\sigma_1(A)_{,jkk}x^j r^{4-n} + O(r^{5-n}) \\
& \quad \left. - 8(n-6)(n-4)\sigma_1(A)r^{2-n} - 32(n-4)(n-2)A_{ij}x^i x^j r^{-n} + 8(n-4)\Delta A_{ij}x^i x^j r^{2-n} \right\} \\
& = (n-6)\left\{ -4(n-4)(3n-26)\sigma_1(A)_{,ij}(p)x^i x^j r^{2-n} - \frac{n-6}{2(n-1)}|W(p)|^2 r^{4-n} \right. \\
& \quad - 2(n-10)(n-4)\sigma_1(A)_{,ij}(p)x^i x^j r^{2-n} - 4(n-6)(n-4)\sigma_1(A)_{,ij}(p)x^i x^j r^{2-n} \\
& \quad \left. - 16(n-4)(n-2)A_{ij,kl}(p)x^i x^j x^k x^l r^{-n} + 8(n-4)\Delta A_{ij}x^i x^j r^{2-n} \right\} + O(r^{5-n}) \\
& = (n-6)\left\{ -2(n-4)(9n-74)\sigma_1(A)_{,ij}(p)x^i x^j r^{2-n} - \frac{n-6}{2(n-1)}|W(p)|^2 r^{4-n} \right. \\
& \quad \left. - 16(n-4)(n-2)A_{ij,kl}(p)x^i x^j x^k x^l r^{-n} + 8(n-4)\Delta A_{ij}x^i x^j r^{2-n} \right\} + O(r^{5-n}).
\end{aligned}$$

To estimate

$$\begin{aligned}
I_3 & = \delta T_2 d\Delta(r^{6-n}) \\
& = -[(T_2)_{ij}(\Delta r^{6-n})_{,j}],_i \\
& = -(T_2)_{ij,i}(\Delta r^{6-n})_{,j} - (T_2)_{ij}(\Delta r^{6-n})_{,ji}.
\end{aligned}$$

Recall that $T_2 = (n-2)\sigma_1(A)g - 8A$, then

$$(T_2)_{ij,i} = (n-10)\sigma_1(A)_{,j}.$$

Observe that

$$\begin{aligned}
\Delta r^{6-n} & = 4(6-n)r^{4-n} + O(r^{N+4-n}), \\
(\Delta r^{6-n})_{,j} & = 4(6-n)(4-n)x^j r^{2-n} + O(r^{N+3-n}),
\end{aligned}$$

and

$$(\Delta r^{6-n})_{,ji} = 4(6-n)(4-n)[(2-n)x^i x^j r^{-n} + r^{2-n}\delta_{ij}] + O(r^{4-n}),$$

then it yields

$$\begin{aligned}
& (T_2)_{ij}(\Delta r^{6-n})_{,ji} \\
& = 4(n-6)(n-4)[(n-2)\sigma_1(A)g_{ij} - 8A_{ij}][(2-n)x^i x^j r^{-n} + r^{2-n}\delta_{ij} + O(r^{4-n})] \\
& = 4(n-6)(n-4)[-(n-2)^2\sigma_1(A)r^{2-n} + n(n-2)\sigma_1(A)r^{2-n} \\
& \quad + 8(n-2)r^{-n}A_{ij}x^i x^j - 8\sigma_1(A)r^{2-n}] + O(r^{5-n}) \\
& = 4(n-6)(n-4)\left[2(n-6)\sigma_1(A)r^{2-n} + 8(n-2)r^{-n}A_{ij}x^i x^j\right] + O(r^{5-n}) \\
& = 4(n-6)(n-4)[(n-6)\sigma_1(A)_{,ij}(p)x^i x^j r^{2-n} + 4(n-2)r^{-n}(A_{ij,kl}(p)x^i x^j x^k x^l)] + O(r^{5-n}).
\end{aligned}$$

Hence, we obtain

$$\begin{aligned}
I_3 &= -4(n-6)(n-4) \left[(n-6)\sigma_1(A)_{,ij}(p)x^i x^j r^{2-n} + 4(n-2)r^{-n}(A_{ij,kl}(p)x^i x^j x^k x^l) \right] \\
&\quad - 4(n-6)(n-4)(n-10)r^{2-n}\sigma_1(A)_{,i}x^i + O(r^{5-n}) \\
&= -8(n-8)(n-6)(n-4)\sigma_1(A)_{,ij}(p)x^i x^j r^{2-n} \\
&\quad - 16(n-6)(n-4)(n-2)r^{-n}(A_{ij,kl}(p)x^i x^j x^k x^l) + O(r^{5-n}).
\end{aligned}$$

We now compute

$$\begin{aligned}
I_4 &= \frac{n-2}{2} \Delta(\sigma_1(A) \Delta(r^{6-n})) \\
&= 2(n-2)(6-n) \Delta(\sigma_1(A) r^{4-n}) + O(r^{N+4-n}) \\
&= 2(n-2)(6-n) r^{2-n} [\Delta \sigma_1(A) r^2 + 2(4-n)\sigma_1(A)_{,i}x^i + 2(4-n)\sigma_1(A)] + O(r^{N+2-n}) \\
&= 2(n-2)(n-6) r^{2-n} \left[\frac{1}{12(n-1)} |W(p)|^2 r^2 + 3(n-4)\sigma_1(A)_{,ij}(p)x^i x^j \right] + O(r^{5-n}).
\end{aligned}$$

For I_5 , from (A.1) we have

$$\begin{aligned}
I_5 &= \delta T_4 d(r^{6-n}) \\
&= -((T_4)_{ij} r^{6-n})_{,j}, i \\
&= -(T_4)_{ij,i} (r^{6-n})_{,j} - (T_4)_{ij} (r^{6-n})_{,ji} \\
&= (n-6) [r^{4-n} (T_4)_{ij,i} x^j - (n-4) r^{2-n} (T_4)_{ij} x^i x^j + r^{4-n} \text{tr}(T_4)] \\
&:= (n-6) [I_1^{(5)} + I_2^{(5)} + I_3^{(5)}].
\end{aligned}$$

Also from [12], there holds

$$\text{Sym} \left(R_{kl,ij} + \frac{2}{9} R_{nkml} R_{nijm} \right) (p) = 0 \quad \text{and} \quad R_{ij}(p) = 0,$$

then

$$R_{kl,ij}(p) x^i x^j x^k x^l = -\frac{2}{9} W_{nkml}(p) W_{nijm}(p) x^i x^j x^k x^l.$$

Thus we have

$$A_{kl,ij}(p) x^i x^j x^k x^l = -\frac{2}{9} \frac{1}{n-2} \sum_{k,l} (W_{iklj}(p) x^i x^j)^2 - \frac{\sigma_1(A)_{,ij}(p) x^i x^j r^2}{n-2}. \quad (\text{A.2})$$

To estimate $I_3^{(5)}$. From the definition of T_4 , one gets

$$\begin{aligned}
\text{tr}(T_4) &= -\frac{3n^3-12n^2-36n+64}{4} \sigma_1(A)^2 + 4(n^2-4n-12)|A|^2 + n(n-6)\Delta \sigma_1(A) \\
&= -\frac{n(n-6)}{12(n-1)} |W(p)|^2 + O(r).
\end{aligned}$$

Thus one obtains

$$I_3^{(5)} = -\frac{n(n-6)}{12(n-1)} |W(p)|^2 r^{4-n} + O(r^{5-n}).$$

For the term $I_1^{(5)}$, it is easy to see

$$I_1^{(5)} = r^{4-n}(T_4)_{ij,i}x^j = O(r^{5-n}).$$

It remains to estimate the term $I_2^{(5)}$, one has

$$(T_4)_{ij}x^ix^j = (n-6)\Delta\sigma_1(A)r^2 - \frac{16}{n-4}B_{ij}x^ix^j + O(r^4). \quad (\text{A.3})$$

Notice that

$$\begin{aligned} B_{ij}x^ix^j &= [C_{ijk,k} - A_{kl}W_{kijl}]x^ix^j = [(A_{ij,k} - A_{ik,j})_{,k} - A_{kl}W_{kijl}]x^ix^j \\ &= [\Delta A_{ij} - A_{ik,jk} + O(r)]x^ix^j \end{aligned}$$

and

$$\begin{aligned} \Delta(A_{ij}x^ix^j) &= (A_{ij,k}x^ix^j + A_{ij}(x^i\delta_{jk} + x^j\delta_{ik}))_{,k} \\ &= (\Delta A_{ij})x^ix^j + 2A_{ij,k}(x^i\delta_{jk} + x^j\delta_{ik}) + 2\sigma_1(A) + O(r^3) \\ &= (\Delta A_{ij})x^ix^j + 4\sigma_1(A)_{,i}x^i + 2\sigma_1(A) + O(r^3). \end{aligned}$$

By (A.2), one gets

$$\begin{aligned} (\Delta A_{ij})x^ix^j &= \Delta(A_{ij}x^ix^j) - 4\sigma_1(A)_{,i}x^i - 2\sigma_1(A) + O(r^3) \\ &= \Delta\left[\frac{1}{2}A_{ij,kl}(p)x^ix^jx^kx^l + O(r^5)\right] - 4[\sigma_1(A)_{,ij}(p)x^ix^j + O(r^3)] \\ &\quad - \sigma_1(A)_{,ij}(p)x^ix^j + O(r^3) \\ &= \Delta\left[-\frac{1}{9}\frac{1}{n-2}\sum_{k,l}(W_{iklj}(p)x^ix^j)^2 - \frac{\sigma_1(A)_{,ij}(p)x^ix^jr^2}{2(n-2)}\right] \\ &\quad - 5\sigma_1(A)_{,ij}(p)x^ix^j + O(r^3) \\ &= -\frac{2}{9}\frac{1}{n-2}\sum_{k,l,s}[(W_{ikls}(p) + W_{ilks}(p))x^i]^2 + \frac{1}{12(n-2)(n-1)}|W(p)|^2r^2 \\ &\quad - 6\frac{n-1}{n-2}\sigma_1(A)_{,ij}(p)x^ix^j + O(r^3), \end{aligned} \quad (\text{A.4})$$

where the last identity follows from the following two estimates:

$$\begin{aligned} \Delta(\sigma_1(A)_{,ij}(p)x^ix^jr^2) &= \Delta(\sigma_1(A)_{,ij}(p)x^ix^j)r^2 + 2\nabla_s(\sigma_1(A)_{,ij}(p)x^ix^j)\nabla_sr^2 \\ &\quad + (\sigma_1(A)_{,ij}(p)x^ix^j)\Delta r^2 \\ &= 2\Delta\sigma_1(A)(p)r^2 + 8\sigma_1(A)_{,ij}(p)x^ix^j + 2n\sigma_1(A)_{,ij}(p)x^ix^j + O(r^3) \\ &= -\frac{1}{6(n-1)}|W(p)|^2r^2 + 2(n+4)\sigma_1(A)_{,ij}(p)x^ix^j + O(r^3) \end{aligned}$$

and

$$\Delta\sum_{k,l}(W_{iklj}(p)x^ix^j)^2 = 2\sum_{k,l,s}[W_{iklj}(p)(x^i\delta_{js} + x^j\delta_{is})]^2 = 2\sum_{k,l,s}[(W_{ikls}(p) + W_{ilks}(p))x^i]^2,$$

which follows from

$$\Delta \left[\sum_{k,l} (W_{iklj}(p) x^i x^j)^2 \right] = 2 \sum_{k,l} [(W_{iklj}(p) x^i x^j) \Delta(W_{sklt}(p) x^s x^t) + |\nabla(W_{iklj}(p) x^i x^j)|^2]$$

and $\Delta(W_{sklt}(p) x^s x^t) = (W_{sklt}(p)(x^s \delta_{it} + x^t \delta_{is}))_{,i} = 2W_{sklt}(p) \delta_{st} = 0$. Using $A_{ik,jk} = A_{ik,kj} + R_{ijk}^l A_{lk} + R_{kjk}^l A_{il} = \sigma_1(A)_{,ij} + R_{lik} A_{lk} + R_{lj} A_{il}$, one has

$$\begin{aligned} & A_{ik,jk} x^i x^j \\ &= \sigma_1(A)_{,ij} x^i x^j + R_{lik} A_{lk} x^i x^j + R_{lj} A_{il} x^i x^j \\ &= (\sigma_1(A)_{,ij}(p) + O(r)) x^i x^j + (W_{lik}(p) + O(r))(A_{lk,m}(p) x^m + O(r^2)) x^i x^j + O(r^4) \\ &= \sigma_1(A)_{,ij}(p) x^i x^j + O(r^3). \end{aligned}$$

Thus, one obtains

$$\begin{aligned} B_{ij} x^i x^j &= -\frac{2}{9} \frac{1}{n-2} \sum_{k,l,s} [(W_{ikls}(p) + W_{ilks}(p)) x^i]^2 + \frac{1}{12(n-2)(n-1)} |W(p)|^2 r^2 \\ &\quad - \frac{7n-8}{n-2} \sigma_1(A)_{,ij}(p) x^i x^j + O(r^3). \end{aligned} \tag{A.5}$$

Inserting the above equations into (A.3), one gets

$$\begin{aligned} & (T_4)_{ij} x^i x^j \\ &= -\frac{n-6}{12(n-1)} |W(p)|^2 r^2 + \frac{32}{9(n-4)(n-2)} \sum_{k,l,s} [(W_{ikls}(p) + W_{ilks}(p)) x^i]^2 \\ &\quad - \frac{4}{3(n-4)(n-2)(n-1)} |W(p)|^2 r^2 + \frac{16(7n-8)}{(n-4)(n-2)} \sigma_1(A)_{,ij}(p) x^i x^j + O(r^3), \end{aligned}$$

whence

$$\begin{aligned} I_2^{(5)} &= r^{2-n} \left[\frac{(n-6)(n-4)}{12(n-1)} |W(p)|^2 r^2 - \frac{32}{9(n-2)} \sum_{k,l,s} ((W_{ikls}(p) + W_{ilks}(p)) x^i)^2 \right. \\ &\quad \left. + \frac{4}{3(n-2)(n-1)} |W(p)|^2 r^2 - \frac{16(7n-8)}{n-2} \sigma_1(A)_{,ij}(p) x^i x^j \right] + O(r^{5-n}). \end{aligned}$$

Combining all the terms together, one has

$$\begin{aligned} I_5 &= \left[-\frac{n^2-8n+8}{3(n-1)(n-2)} |W(p)|^2 r^{4-n} - \frac{32}{9(n-2)} \sum_{k,l,s} ((W_{ikls}(p) + W_{ilks}(p)) x^i)^2 r^{2-n} \right. \\ &\quad \left. - \frac{16(7n-8)}{n-2} \sigma_1(A)_{,ij}(p) x^i x^j r^{2-n} \right] (n-6) + O(r^{5-n}). \end{aligned}$$

Finally, from the definition of Q_g in (1.1), it is not hard to show that

$$I_6 = -\frac{n-6}{2} Q_g r^{6-n} = O(r^{6-n}).$$

Therefore, collecting all the terms I_1 - I_6 together with (A.2) and (A.4), we conclude that

$$\begin{aligned}
& -P_g(r^{6-n}) \\
& = -c_n \delta_p + (n-6) \left[-\frac{16}{9} \sum_{k,l,s} ((W_{ikls}(p) + W_{ilks}(p))x^i)^2 r^{2-n} - \frac{2(n-8)}{3(n-1)} |W(p)|^2 r^{4-n} \right. \\
& \quad \left. + \frac{64(n-4)}{9} \sum_{k,l} (W_{iklj}(p) x^i x^j)^2 r^{-n} - 4(5n^2 - 66n + 224) \sigma_1(A)_{,ij}(p) x^i x^j r^{2-n} \right] + O(r^{5-n}) \\
& = -c_n \delta_p + (n-6) r^{-n} \left\{ \frac{64(n-4)}{9} \left[\sum_{k,l} (W_{iklj}(p) x^i x^j)^2 - \frac{r^2}{n+4} \sum_{k,l,s} ((W_{ikls}(p) + W_{ilks}(p))x^i)^2 \right. \right. \\
& \quad \left. \left. + \frac{3}{2(n+4)(n+2)} |W(p)|^2 r^4 \right] + \frac{16(3n-20)}{9(n+4)} r^2 \left[\sum_{k,l,s} ((W_{ikls}(p) + W_{ilks}(p))x^i)^2 - \frac{3}{n} |W(p)|^2 r^2 \right] \right. \\
& \quad \left. - 4(5n^2 - 66n + 224) r^2 \left[\sigma_1(A)_{,ij}(p) x^i x^j + \frac{|W(p)|^2}{12n(n-1)} r^2 \right] \right. \\
& \quad \left. + \frac{3n^4 - 16n^3 - 164n^2 + 400n + 2432}{3(n+4)(n+2)n(n-1)} |W(p)|^2 r^4 \right\} + O(r^{5-n}),
\end{aligned}$$

where each term in square brackets on the right hand side of the last identity is harmonic polynomial. This finishes the proof of Lemma 2.1.

References

- [1] A. Ambrosetti and P. Rabinowitz, *Dual variational methods in critical point theory and applications*, J. Functional Analysis 14 (1973), 349-381.
- [2] T. Aubin, *Équations différentielles non linéaires et problème de Yamabe concernant la courbure scalaire*, J. Math. Pures Appl. (9) 55 (1976), no. 3, 269-296.
- [3] Z. Djadli, E. Hebey and M. Ledoux, *Paneitz-type operators and applications*, Duke Math. J. 104 (2000), no. 1, 129-169.
- [4] P. Esposito and F. Robert, *Mountain pass critical points for Paneitz-Branson operators*, Calc. Var. Partial Differential Equations 15 (2002), no. 4, 493-517.
- [5] C. Fefferman and Graham, *The Ambient metric*, Annals of Mathematics Studies, 178, Princeton University Press, Princeton, NJ, 2012. x+113 pp.
- [6] C. Graham, R. Jenne, L. Mason and G. Sparling, *Conformally invariant powers of the Laplacian. I. Existence*, J. London Math. Soc. (2) 46 (1992), no. 3, 557-565.
- [7] A. Gover, *Laplacian operators and Q-curvature on conformally Einstein manifolds*, Math. Ann. 336 (2006), no. 2, 311-334.
- [8] M. Gursky, F. Hang and Y. Lin, *Riemannian manifolds with positive Yamabe invariant and Paneitz operator*, preprint (2015), arXiv:1502.01050.

- [9] M. Gursky and A. Malchiodi, *A strong maximum principle for the Paneitz operator and a non-local flow for the Q -curvature*, to appear in JEMS, arXiv:1401.3216.
- [10] F. Hang and P. Yang, *Q curvature on a class of manifolds with dimension at least five*, preprint (2014), arXiv:1411.3926.
- [11] A. Juhl, *Explicit formulas for GJMS-operators and Q -curvatures*, Geom. Funct. Anal. 23 (2013), no. 4, 1278-1370.
- [12] J. Lee and T. Parker, *The Yamabe problem*, Bull. Amer. Math. Soc. (N.S.) 17 (1987), no. 1, 37-91.
- [13] Y. Li and J. Xiong, *Compactness of conformal metrics with constant Q -curvature. I*, preprint (2015), arXiv:1506.00739.
- [14] E. Lieb, *Sharp constants in the Hardy-Littlewood-Sobolev and related inequalities*, Ann. of Math. (2) 118 (1983), 349-374.
- [15] P. L. Lions, *The concentration-compactness principle in the calculus of variations: The limit case. Part I*, Rev. Mat. Iberoamericana 1 (1985), 145-201.
- [16] E. Stein, *Singular integrals and differentiability properties of functions*, Princeton University Press, Princeton, New Jersey, 1970.
- [17] V. Wünsch, *On conformally invariant differential operators*, Math. Nachr. 129 (1986), 269-281.