

Mean-correction and Higher Order Moments for a Stochastic Volatility Model with Correlated Errors

Sujay Mukhoti and Pritam Ranjan

Operations Management and Quantitative Techniques,
Indian Institute of Management Indore, M.P., India, 453556
(sujaym@iimidr.ac.in, pritamr@iimidr.ac.in)

Abstract

In an efficient stock market, the log-returns and their time-dependent variances are often jointly modelled by stochastic volatility models (SVMs). Many SVMs assume that errors in log-return and latent volatility process are uncorrelated, which is unrealistic. It turns out that if a non-zero correlation is included in the SVM (e.g., Shephard (2005)), then the expected log-return at time t conditional on the past returns is non-zero, which is not a desirable feature of an efficient stock market. In this paper, we propose a mean-correction for such an SVM for discrete-time returns with non-zero correlation. We also find closed form analytical expressions for higher moments of log-return and its lead-lag correlations with the volatility process. We compare the performance of the proposed and classical SVMs on S&P 500 index returns obtained from NYSE.

KEY WORDS: Leverage effect, Martingale difference, Skewness, Volatility asymmetry.

1 Introduction

Over the last few decades different aspects of stock price movements in discrete time have been the focus of numerous research avenues. Suppose P_t denotes the price of a stock at time t , then the continuously compounded return or log-return (here onwards referred to as return) of the stock is defined as $r_t = \log(P_t/P_{t-1})$. A stock market is said to be efficient if the price of a stock contains every available information about it. In such a market the risk involved in investing on a stock is measured by the standard deviation of r_t , often termed as the *volatility* of the stock in finance literature. It has been noted that volatility varies over time (Engle 1982). Stochastic Volatility Models (SVMs) is a popular class of models for describing the time-varying volatility of stock returns (Shephard 2005).

Although there are a plethora of SVMs for describing the stock returns, one of the simplest yet most popular discrete-time SVM is given by Taylor (1982), where the return process r_t is a non-linear product of two independent stochastic processes, viz. an i.i.d. error process ϵ_t , and a latent volatility process h_t , which is further modelled as an $AR(1)$. That is,

$$\begin{aligned} r_t &= \exp\left\{\frac{h_t}{2}\right\} \epsilon_t \\ h_t &= \alpha + \phi(h_{t-1} - \alpha) + \sigma\eta_t, \quad \forall t = 1, 2, \dots, \end{aligned} \tag{1}$$

where $\alpha = E(h_t)$ is the long-range volatility, ϕ is the stationarity parameter, σ measures the variability of the volatility process h_t , and ϵ_t and η_t are uncorrelated i.i.d. $N(0, 1)$ errors. Hereafter this model will be referred as SVM_0 .

As in (1), many of the new generation SVMs which are being used in the finance literature assume that ϵ_t and η_t are independent $N(0, 1)$ errors. In reality, however, ϵ_t and η_t are often correlated (Harvey & Siddique 1999). Though discrete-time SVMs with non-zero $corr(\epsilon_t, \eta_t)$ have been developed earlier and are being used, they assume that h_{t+1} (instead of h_t as in (1)) depends on η_t via $AR(1)$ (see *e.g.* Meyer & Yu (2000) and Berg, Meyer & Yu (2004)). In this paper, we focus on the SVM presented in (1) with correlated errors (denoted as SVM_ρ). That is, the additional assumption in (1) is $corr(\epsilon_t, \eta_t) = \rho$.

It turns out that introducing a non-zero correlation between η_t and ϵ_t in (1) has an adverse effect on the admissibility of the SVM from an efficient market's viewpoint. In particular, the conditional expectation of r_t given the past data, $E[r_t | \mathcal{F}_{t-1}]$, is not zero, where \mathcal{F}_{t-1} is the space (σ -field) generated with r_1, \dots, r_{t-1} . This zero conditional expectation of the return is a necessary requirement for an *efficient market hypothesis* (EMH) (see Yu (2005) for a review).

In this paper, we propose a mean-correction for SVM_ρ - model (1) with correlated errors, such that $E[r_t | \mathcal{F}_{t-1}]$ becomes zero and the corrected SVM would satisfy EMH. The proposed mean-corrected model is denoted by $SVM_{\rho\mu}$. Further, Black (1976) mentioned that, usually, the amount of increment in volatility due to price fall is larger than the magnitude of reduction in the volatility due to price increase. In turn, this indicates the volatility of positive returns, $var(r_t | r_t > 0)$, is less than the volatility of the negative returns, $var(r_t | r_t < 0)$ resulting in skewness in return distribution. Moreover, the kurtosis quantifies the proportion of extreme values, that occur during crashes, explained by the model. We find the closed form expressions for the higher-order moments and the lead-lag correlation of the underlying

return process. These descriptive statistics indicate the influence of past/future volatility on today's return.

The remainder of the article is organized as follows. Section 2 presents the main results: $SVM_{\rho\mu}$ - the mean-corrected SVM with non-zero correlation that satisfies EMH, and the closed form analytical expressions for the higher order moments and lead-lag correlation for the proposed model. For the returns of S&P 500 NSYE, Section 3 presents a comparison between the standard zero correlation model (1) and the ones with non-zero correlation. Finally Section 4 outlines the concluding remarks and a few possible future directions.

2 Main Results

For this section, we assume that the error terms ϵ_t and η_t in (1) have not only a constant correlation ρ and i.i.d. $N(0, 1)$ marginals, but they also follow a bivariate normal distribution. The proposed mean-corrected model ($SVM_{\rho\mu}$) contains an additional term μ , i.e.,

$$\begin{aligned} r_t &= \mu + \exp\left\{\frac{h_t}{2}\right\} \epsilon_t \\ h_t &= \alpha + \phi(h_{t-1} - \alpha) + \sigma\eta_t, \quad \forall t = 1, 2, \dots, T. \end{aligned} \quad (2)$$

Theorem 1 establishes the value of μ for which the proposed mean-corrected model (2) gives zero conditional expectation $E[r_t | \mathcal{F}_{t-1}]$ and hence satisfy EMH. Later in this section, we derive closed form expressions for the higher-order moments, i.e., variance, skewness, and kurtosis of r_t , and lead-lag correlations between r_t and $h_{t\pm k}$.

Theorem 1 *For $SVM_{\rho\mu}$ in (2) with $|\phi| \leq 1, \sigma > 0$ and $-\infty < \alpha < \infty$, if (ϵ_t, η_t) follows a standard bivariate normal distribution with correlation ρ , the mean term*

$$\mu = -\frac{\rho\sigma}{2} \exp\left\{\frac{\alpha}{2} + \frac{\sigma^2}{8(1-\phi^2)}\right\} \quad (3)$$

gives $E[r_t | \mathcal{F}_{t-1}] = 0$ and vice-versa.

Proof The conditional expected return $E[r_t | \mathcal{F}_{t-1}] = 0$ gives

$$\begin{aligned} -\mu &= E\left[\exp\left\{\frac{h_t}{2}\right\} \epsilon_t\right] = E\left[\exp\left\{\frac{\alpha + \phi(h_{t-1} - \alpha) + \sigma\eta_t}{2}\right\} \epsilon_t\right] \\ &= \exp\left\{\frac{\alpha}{2}\right\} \times E\left[\exp\left\{\frac{\phi\sigma}{2} \sum_{j=1}^{\infty} \phi^{j-1} \eta_{t-j}\right\}\right] \times E\left[\exp\left\{\frac{\sigma\eta_t}{2}\right\} \epsilon_t\right]. \end{aligned} \quad (4)$$

Since (ϵ_t, η_t) follows a standard bivariate normal with correlation ρ , the condition distribution of $\epsilon_t \mid \eta_t$ is given by $N(\rho\eta_t, 1 - \rho^2)$. This conditional normal distribution and the moment generating function (mgf) of a normal distribution simplifies the third term in (4) as

$$E \left[\exp \left\{ \frac{\sigma\eta_t}{2} \right\} \epsilon_t \right] = E_{\eta_t} \left[\exp \left\{ \frac{\sigma\eta_t}{2} \right\} \rho\eta_t \right] = \frac{\rho\sigma}{2} \exp \left\{ \frac{\sigma^2}{8} \right\}, \quad (5)$$

and the second term to

$$\prod_{j=1}^{\infty} E \left[\exp \left\{ \frac{\sigma\phi^j}{2} \eta_{t-j} \right\} \right] = \exp \left\{ \frac{\sigma^2}{8} \sum_{j=1}^{\infty} \phi^{2j} \right\} = \exp \left\{ \frac{\sigma^2\phi^2}{8(1-\phi^2)} \right\}. \quad (6)$$

Hence the final expression for μ follows from (4)-(6). \square

Yu (2005) tried to compute $E[r_t \mid \mathcal{F}_{t-1}]$, but the final expression appears to be incorrect. Note that the proposed mean-correction (in Theorem 2.1) makes the model (2) usable in the stock market, as it now satisfies EMH (in particular, $E[r_t \mid \mathcal{F}_{t-1}] = 0$). Further, the proof of the above theorem prohibits the usage of heavy-tail distributions (like t distribution) as the volatility error distribution (Wang, Chan & Choy 2011) as its moment generating function would not exist resulting in in-existence of expected returns. In Section 3, we discuss the usage of this model for the index returns of S&P500 index of New York Stock Exchange (NYSE) observed during 1st April, 2002 - 30th March, 2006.

2.1 Higher-order moments

For additional key features on the distribution of returns, we estimate higher order moments, in particular, variance, skewness and kurtosis conditional on \mathcal{F}_{t-1} .

Theorem 2 *For SVM $_{\rho\mu}$ in (2), if Theorem 2.1 holds, then the variance of returns conditional on \mathcal{F}_{t-1} is given by*

$$V(r_t \mid \mathcal{F}_{t-1}) = \exp \left\{ \alpha + \frac{\sigma^2}{2(1-\phi^2)} \right\} \left(1 + \rho^2\sigma^2 - \frac{\rho^2\sigma^2}{4} \exp \left\{ -\frac{\sigma^2}{4(1-\phi^2)} \right\} \right). \quad (7)$$

Proof Following the definition of variance,

$$\begin{aligned}
V(r_t | \mathcal{F}_{t-1}) &= E[r_t^2 | \mathcal{F}_{t-1}] - \mu^2 \\
&= E[\exp\{h_t\}\epsilon_t^2] - \mu^2 \\
&= \exp\{\alpha\} \times E\left[\exp\left\{\sigma \sum_{j=1}^{\infty} \phi^j \eta_{t-j}\right\}\right] \times E[\exp\{\sigma\eta_t\}\epsilon_t^2] - \mu^2 \\
&= \exp\left\{\alpha + \frac{\sigma^2\phi^2}{2(1-\phi^2)}\right\} (1 + \rho^2\sigma^2) \exp\left\{\frac{\sigma^2}{2}\right\} - \mu^2 \quad (\text{as in (4)-(6)}).
\end{aligned}$$

The final result follows by substituting the value of μ from Theorem 2.1. \square

The expressions of the conditional mean and variance are the most crucial components in finding the skewness and kurtosis statistics. For $SVM_{\rho\mu}$ in (2), under the same conditions as in Theorem 2.2, the skewness conditional on \mathcal{F}_{t-1} is measured by $\mu_3/(Var(r_t | \mathcal{F}_{t-1}))^{3/2}$, where

$$\begin{aligned}
\mu_3 &= \frac{3\rho\sigma}{2} \exp\left\{\frac{3\alpha}{2} + \frac{9\sigma^2}{8(1-\phi^2)}\right\} \left[3 + \frac{9\sigma^2\rho^2}{4} + \frac{\rho^2\sigma^2}{6} \exp\left\{-\frac{3\sigma^2}{4(1-\phi^2)}\right\}\right. \\
&\quad \left. - (1 + \rho^2\sigma^2) \exp\left\{-\frac{\sigma^2}{2(1-\phi^2)}\right\}\right]. \tag{8}
\end{aligned}$$

The proof of (8) starts with $\mu_3 = E[r_t^3 | \mathcal{F}_{t-1}]$, and proceeds in the exact same manner as in Theorems 2.1 and 2.2. Similarly the closed form expression of kurtosis can also be found as $\mu_4/(Var(r_t | \mathcal{F}_{t-1}))^2$, where

$$\begin{aligned}
\mu_4 &= \exp\left\{2\alpha + \frac{2\sigma^2}{(1-\phi^2)}\right\} \times \left[\frac{3}{2}\rho^2\sigma^2(1 + \sigma^2\rho^2) \exp\left\{\frac{-5\sigma^2}{4(1-\phi^2)}\right\}\right. \\
&\quad \left.+ (3 + 24\rho^2\sigma^2 + 16\rho^4\sigma^4) - \frac{3}{16}\rho^4\sigma^4 \exp\left\{\frac{-3}{2} \frac{\sigma^2}{(1-\phi^2)}\right\}\right. \\
&\quad \left.- 9\rho^2\sigma^2 \left(1 + \frac{3}{4}\rho^2\sigma^2\right) \exp\left\{\frac{-3\sigma^2}{4(1-\phi^2)}\right\}\right]. \tag{9}
\end{aligned}$$

As expected all four descriptive statistics found here depends heavily on $corr(\epsilon_t, \eta_t) = \rho$. On a closer inspection of these statistics, we see that $\rho = 0$ (i.e., the classical SVM by (Taylor 1982)) gives $\mu = 0, \mu_3 = 0$,

$$Var(r_t | \mathcal{F}_{t-1}) = \exp\left\{\alpha + \frac{\sigma^2}{2(1-\phi^2)}\right\} \quad \text{and} \quad \mu_4 = 3 \exp\left\{2\alpha + \frac{2\sigma^2}{(1-\phi^2)}\right\}.$$

Note that the simplified expressions found here are consistent with the ones reported by Ghysels, Harvey & Renault (1996), and hence the proposed model $SVM_{\rho\mu}$ (2) is a general-

ization of the classical model SVM_0 in (1). Next we investigate the conditional (on \mathcal{F}_{t-1}) dependence between the current returns and past, current and future volatility.

2.2 Lead-lag correlations

In this section, we wish to estimate three quantities: (1) dependence between the current returns and current volatility, $corr(r_t, h_t | \mathcal{F}_{t-1})$, (2) the potential influence of current returns on future volatility, $corr(r_t, h_{t+k} | \mathcal{F}_{t-1})$, and (c) the influence of past volatility on current returns, $corr(r_t, h_{t-k} | \mathcal{F}_{t-1})$. Though empirical estimation of such quantities is not uncommon, e.g., in Bollerslev, Litvinova & Tauchen (2006), our aim is to find closed analytical expression for these descriptive measures under $SVM_{\rho\mu}$ specification.

Since $var(r_t | \mathcal{F}_{t-1})$ is given by (7) and $var(h_t | \mathcal{F}_{t-1}) = \sigma^2 / (1 - \phi^2)$, we only need to find the expressions for the conditional covariances. First, we recall that under the proposed model, the conditional means are $E(r_t | \mathcal{F}_{t-1}) = 0$ and $E(h_t | \mathcal{F}_{t-1}) = \alpha$. Now, if we assume that $corr(r_t, h_t | \mathcal{F}_{t-1}) = \sigma_{rh}$, then

$$cov(r_t, h_{t+1}) = E[r_t(h_t - \alpha)] = E[r_t(\phi(h_t - \alpha) + \sigma\eta_{t+1})] = \phi E[r_t(h_t - \alpha)] = \phi\sigma_{rh},$$

which further implies that $cov(r_t, h_{t+k}) = \phi^k \sigma_{rh}$ for $k \geq 1$. By applying the key mathematical techniques (i.e., properties of expectation, normal mgf and the expansion of $h_t = \alpha + \sigma \sum_{j=1}^{\infty} \eta_{t-j} \phi^j$) used in proving results of Section 2.1, one can easily show that

$$\begin{aligned} \sigma_{rh} = cov(r_t, h_t | \mathcal{F}_{t-1}) &= \rho\sigma \exp \left\{ \frac{\alpha}{2} + \frac{\sigma^2}{8(1-\phi^2)} \right\} \times \left\{ 1 + \frac{\sigma^2}{4(1-\phi^2)} \right\}, \\ cov(r_t, h_{t-k} | \mathcal{F}_{t-1}) &= \sigma_{rh} \cdot \phi^k \cdot \left[\frac{\sigma^2}{4(1-\phi^2)} \right] / \left[1 + \frac{\sigma^2}{4(1-\phi^2)} \right]. \end{aligned}$$

Clearly, both the lead ($cov(r_t, h_{t+k})$) and lag ($cov(r_t, h_{t-k})$) covariances are smaller than the contemporaneous covariance $cov(r_t, h_{t-k} | \mathcal{F}_{t-1})$. The contemporaneous correlation can be interpreted as *feedback effect* of volatility change on future returns, whereas the impact of return change on future volatility is termed as *leverage effect*. Bekaert & Wu (2000) found that volatility feedback effect is stronger than leverage effect. The closed form expressions we have derived above provide a theoretical proof of the mentioned findings under $SVM_{\rho\mu}$ specification. Moreover, Bollerslev et al. (2006) have empirically observed that the lag-correlation with lag h is smaller than lead correlation with lead h which we have established theoretically. Further note that all these covariances and hence correlations vanish if $\rho =$

$\text{corr}(\epsilon_t, \eta_t) = 0$. Next, we compare the goodness of fit of the three stochastic volatility models, SVM_0 (classical - with zero correlation), SVM_ρ (with correlation ρ) and $SVM_{\rho\mu}$ (mean-corrected with correlation ρ), for a real data on returns.

3 Example: S&P 500 NYSE

In this paper, we compare the performance of the three models ($SVM_0, SVM_\rho, SVM_{\rho\mu}$) on the index returns of Standard and Poor 500 index (S&P500) obtained from New York Stock Exchange during April 01, 2002 – March 30, 2006. We selected this period to avoid extreme behaviour during “2000 – 2002 dot-com bubble” and “2008 Lehman Brothers’ crash”. Figure 1 displays the time-plot of the returns of 1008 trading days (less than the total number of calendar days).




Figure 1: Time plot of S&P500 returns during April 01, 2002 – March 30, 2006

From Figure 1 one can infer that the volatility is relatively high during September 2003 and June 2004, whereas during October 2004 to April 2005, the volatility is relatively lower than usual. A few descriptive statistics of the observed returns are as follows:

We follow Meyer & Yu (2000), and use the same Markov Chain Monte Carlo (MCMC) algorithm implemented in Just Another Gibbs Sampler (JAGS) for fitting the classical model

$$\begin{aligned} \text{mean} &= 0.0014, & \text{variance} &= 0.0005, \\ \text{skewness} &= 0.0329, & \text{kurtosis} &= 10.9813. \end{aligned}$$

SVM_0 . For fitting the other two models, SVM_ρ and $SVM_{\rho\mu}$, we slightly modify the JAGS code to include the $\text{corr}(\epsilon_t, \eta_t) = \rho$ and μ (derived in Theorem 2.1). For implementing SVM_0 in JAGS, the hierarchical model structure is characterized by

$$r_t \mid (h_t, h_{t-1}, \dots, h_1, h_0; \alpha, \phi, \sigma) \sim N(0, \exp\{h_t\}),$$

$$\text{and } h_t \mid (h_{t-1}, \dots, h_1, h_0; \alpha, \phi, \sigma) \sim N(\alpha + \phi(h_{t-1} - \alpha), \sigma^2).$$

For SVM_ρ , the mean and variance of the conditional distribution of r_t changes to

$$r_t \mid (h_t, \dots, h_0; \alpha, \phi, \sigma) \sim N\left(\frac{\rho e^{h_t/2}}{\sigma}(h_t - \alpha - \phi(h_{t-1} - \alpha)), e^{h_t}(1 - \rho^2)\right),$$

and the conditional distribution of h_t remains the same. Similarly, the implementation of the mean-corrected model $SVM_{\rho\mu}$ is characterized by updating the mean and variance of the conditional distribution of r_t to

$$r_t \mid (h_t, \dots, h_0; \alpha, \phi, \sigma) \sim N\left(\mu + \frac{\rho e^{h_t/2}}{\sigma}(h_t - \alpha - \phi(h_{t-1} - \alpha)), e^{h_t}(1 - \rho^2)\right).$$

The parameters of interest are $(\alpha, \phi, \rho, \sigma) = \Theta$ (say). We use the same prior (including the hyperparameters) for α, ϕ and σ as in Meyer & Yu (2000), and a non-informative $Unif(-1, 1)$ prior for the correlation parameter. The posterior of Θ and $\mathcal{H} = \{h_t, h_{t-1}, \dots\}$ given the data $\{r_t, r_{t-1}, \dots\}$ is obtained via JAGS. We set the total length of chains to be 180,000, out of which 30,000 was the burn-in, and from the remaining 150,000 posterior realizations (with the thinning of every 50th realization) were used (i.e., 3000 realizations in total) to obtain the plug-in estimates of the parameters. The thinning process facilitates a safeguard against the chain dependency in the sampling process. Figure 2 shows the density plots of the posterior distribution of Θ for the three models, SVM_0 , SVM_ρ and $SVM_{\rho\mu}$. We have not included the traceplots, as all parameters converge nicely and the plots do not reveal anything extra. The plug-in estimates of the parameters are obtained via posterior mean and variance (summarized in Table 1).

Table 1 shows that the posterior estimates of the parameters in SVM_0 , SVM_ρ and $SVM_{\rho\mu}$ are similar. Further, the near-unity estimate of ϕ indicates presence of strong volatility clustering. The estimate of the correlation parameter ρ is small yet positive, which is

Table 1: Plug-in estimators of $\Theta = (\alpha, \phi, \sigma, \rho)$ for the three models. The numbers in parentheses show the standard deviation of the posterior realizations.

Parameter	SVM_0	SVM_ρ	$SVM_{\rho\mu}$
α	-7.88 (0.1837)	-7.87 (0.2077)	-7.88 (0.192)
ϕ	0.96 (0.016)	0.97 (0.014)	0.96 (0.014)
σ	0.2 (0.04)	0.177 (0.034)	0.18 (0.038)
ρ		0.1185 (0.1362)	0.105 (0.1278)

similar to the findings of French, Schwert & Stambaugh (1987) and Campbell & Hentschel (1992). This may be taken as an indication of no significant effect of current return on future volatility.

Figure 2 shows that the posterior distributions of the parameters for SVM_0 , SVM_ρ and $SVM_{\rho\mu}$ are different in their kurtosis. A general pattern that can be noticed is that posterior distributions of the parameters under SVM_0 are more leptokurtic compared to their counterparts under other two models except for α . Importantly, too strong volatility clustering is more probable under SVM_ρ and $SVM_{\rho\mu}$ compared to SVM_0 . In case of variance of volatility, posterior distribution under SVM_0 indicates higher values compared to the other two models. Comparing the posterior distributions of ρ under SVM_ρ and $SVM_{\rho\mu}$, the former shows higher probability of being positive valued relative to the latter.

As per Figure 2(d), ρ is very small (close to zero), and thus, it is expected that the proposed model would not provide significant additional strength in modelling the returns data.

We now compare the three models using the descriptive measures (mean, variance, skewness and kurtosis), three lead-lag correlations, mean deviance over the posterior distribution, and the mean square prediction error (MSPE): $\sum_{t=1}^T \hat{r}_t^2/T$. The deviance function, suggested by (Dempster 1974), is

$$D(\Theta) = -2 \log f(r | \Theta, \mathcal{H}) + 2 \log g(r),$$




Figure 2: Posterior distribution of Θ for the three models. The black solid curves represent SVM_0 , blue dashed curves are from SVM_ρ model, and the red dotted curves are obtained from the proposed model $SVM_{\rho\mu}$.

where $f(r | \Theta, \mathcal{H})$ is the likelihood for a given realization of Θ and \mathcal{H} , and $g(r)$ is the normalizing constant. Table 2 presents the plug-in values of these “goodness of fit” measures for the three models.

Since $\rho \approx 0.1$ (very small), the estimated mean is also small $\mu = -4.05 \cdot 10^{-6}$. Thus

Table 2: Goodness of fit measures for the true data and the three models.

GOF measure	True data	SVM_0	SVM_ρ	$SVM_{\rho\mu}$
Mean	0.0014	0	–	-4.05×10^{-6}
Variance	0.0005	0.0005	0.0005	0.0005
Skewness	0.0329	0	0.0856	0.0769
Kurtosis	10.981	5.196	5.105	5.076
$corr(r_t, h_t)$			0.0305	0.0276
$corr(r_t, h_{t-10})$				0.0053
Deviance		-5019	-5033	-5043
MSPE ($\times 10^{-7}$)		0.178	12.94	9.298

all three models would behave very similarly (which is reflected in the estimated moments under the three models). Surprisingly plug-in estimates of kurtosis obtained from all three models under-estimates the kurtosis measured from the data. Deviance values indicate that $SVM_{\rho\mu}$ provides a slightly better fit compared to the other two models. On the other hand, MSPE values indicate that the basic SVM provides better prediction among the three models. Though the numerical results presented through the S& P 500 NYSE example do not provide sufficient evidence for $SVM_{\rho\mu}$ giving additional information than SVM_0 , it certainly is the generalization of SVM_0 and an example with large $\rho = corr(\epsilon_t, \eta_t)$ might have given more convincing evidence.

4 Concluding Remarks

In this paper, we have proposed a mean-correction for the SVM with correlation between ϵ_t and η_t . This mean-correction step enables the conditional expected return to be zero, which is a necessary condition for a good SVM (i.e., a model that adhere to the EMH). We have also found the closed form analytical expression for the higher moments of returns and lead-lag correlation between the return and volatility.

From S&P500 example, we see that most of the empirical observations on statistical properties of returns are reflected through all the three models. However, $SVM_{\rho\mu}$ gives a slightly better fit to the data (in terms of average deviance) compared to the classical model

SVM_0 as well as SVM_ρ . A close look at this research endeavour generates several interesting and challenging research problems.

First, the estimated error correlation ρ turns out to be positive despite the fact that return and its volatility move in opposite directions (Nelson 1991). Glosten, Jagannathan & Runkle (1993) attributed this discrepancy due to mis-specification in the underlying SVM, which is caused by not accounting for the size discrepancy in volatility change due to up or down movement of price. The authors have shown that if the size discrepancy is accounted for then ρ becomes negative. This result demonstrates that ρ alone can not explain the asymmetric response of return to its volatility sufficiently. As we have pointed out in the introduction that this size discrepancy can be interpreted as different conditional variances (or volatility) for positive and negative returns, which leads to skewed return distribution instead of a Gaussian one, a new model can be developed by extending $SVM_{\rho\mu}$ in the line of Abanto-Valle, Bandyopadhyay, Lachos & Enriquez (2010).

Second, the observed kurtosis from the data is not completely explained by the model based estimates of kurtosis. Indeed, the significant difference between empirical kurtosis and the model based estimates again suggests non-normality of the return error distribution. The problem can be tackled in two ways- (1) introducing jumps in returns or (2) allowing the return error to be heavy-tailed (*e.g.* Student's t). Notice, adding a jump to the return only explains transient changes (as seen on 8th & 9th August, 2002) and does not cause the return distribution to change permanently whereas jump in both return and volatility explains persistent effects of extreme values (*e.g.* September, 2003 – June, 2004). $SVM_{\rho\mu}$ can further be generalized by including jumps in return and volatility (Eraker, Johanners & Polson 2003) following the 1st line of argument and using skew Student's- t distributions following the 2nd line of argument (Abanto-Valle, Lachos & Dey 2015).

Although continuous time stochastic volatility has been studied extensively in the literature, the comparatively new discrete-time SVM brings out new interesting features such as leverage effect and feedback effect which occurs due to lagged reaction between return and its volatility. In this paper we have established that the empirically observed pattern of leverage effect and lagged correlations (Bollerslev et al. 2006) are explained by $SVM_{\rho\mu}$. In particular, we have shown that the correlation between current return and future volatility is maximum in magnitude at lead 0 (or contemporaneously) and the future leverage effects disappear exponentially with the lead time. Indeed, strong volatility clustering effect in-

icates more persistent leverage effect. It may also be noted that the existing practice of assuming $h_{t+1} = \alpha + \phi(h_t - \alpha) + \sigma\eta_t$ (instead of h_t) and $\text{corr}(\epsilon_t, \eta_t) = \rho$ for a correct SVM specification would not support the empirical observation on contemporaneous correlation.

Mean correction to the contemporaneously correlated SVM has a very important application. Zero conditional expected returns is also referred to as the martingale difference property, which is a necessary and sufficient condition for *no arbitrage* - which further leads to the existence of option pricing kernel (Back (1991), Shephard & Andersen (2009)). That is, we believe that the proposed mean-correction strategy can also be used in option pricing.

References

- Abanto-Valle, C. A., Bandyopadhyay, D., Lachos, V. H., & Enriquez, I. (2010), “Robust Bayesian analysis of heavy-tailed stochastic volatility models using scale mixtures of normal distributions,” *Computational Statistics and Data Analysis*, 54(12), 2883–2898.
- Abanto-Valle, C., Lachos, V., & Dey, D. K. (2015), “Bayesian estimation of a skew-student-t stochastic volatility model,” *Methodology and Computing in Applied Probability*, 17(3), 721–738.
- Back, K. (1991), “Asset pricing for general processes,” *Journal of Mathematical Economics*, 20(4), 371–395.
- Bekaert, G., & Wu, G. (2000), “Asymmetric volatility and risk in equity markets,” *The review of financial studies*, 13, 1–42.
- Berg, A., Meyer, R., & Yu, J. (2004), “Deviance Information Criterion for Comparing Stochastic Volatility Models,” *Journal of Business and Economic Statistics*, 22(1), 107–120.
- Black, F. (1976), “The pricing of commodity contracts,” *Journal of Financial Economics*, 3, 167–179.
- Bollerslev, T., Litvinova, J., & Tauchen, G. (2006), “Leverage and feedback effects in high frequency data,” *Journal of Financial Econometrics*, 4, 353–384.

- Campbell, J. Y., & Hentschel, L. (1992), “No news is good news: An asymmetric model of changing volatility in stock returns,” *Journal of Financial Economics*, 31(3), 281 – 318.
- Dempster, A. P. (1974), “The direct use of likelihood for significance testing,” *Proceedings of conference on foundational questions in statistical inference*, University of Aarhus, 335–352.
- Engle, R. F. (1982), “Autoregressive Conditional Heteroscedasticity with Estimates of the Variance of United Kingdom Inflation,” *Econometrica*, 50(4), 987–1007.
- Eraker, B., Johanners, M., & Polson, N. G. (2003), “The impact of jumps in volatility and returns,” *Journal of Finance*, 53(3), 1269–1300.
- French, R. K., Schwert, G. W., & Stambaugh, F. R. (1987), “Expected Stock Returns and Volatility,” *Journal of Financial Economics*, 19, 3–29.
- Ghysels, E., Harvey, A. C., & Renault, E. (1996), “5 Stochastic volatility,” in *Statistical Methods in Finance*, Vol. 14 of *Handbook of Statistics*, New York: Elsevier, pp. 119 – 191.
- Glosten, L. R., Jagannathan, R., & Runkle, D. E. (1993), “On the Relation between the Expected Value and the Volatility of the Nominal Excess Return on Stocks,” *The Journal of Finance*, 48(5), 1779–1801.
- Harvey, C., & Siddique, A. (1999), “Autoregressive conditional skewness,” *Journal of Finance and Quantitative Analysis*, 34, 465–487.
- Meyer, R., & Yu, J. (2000), “BUGS for a Bayesian analysis of stochastic volatility models,” *Econometrics Journal*, 3(2), 198–215.
- Nelson, B. D. (1991), “Conditional heteroskedasticity in asset pricing: a new approach,” *Econometrica*, 59, 347–370.
- Shephard, N., & Andersen, G. (2009), “Stochastic Volatility: Origins and Overview,” in *Handbook of Financial Time Series*, eds. T. Mikosch, J.-P. Krei, R. A. Davis, & T. G. Andersen, Berlin Heidelberg: Springer, pp. 237–254.

- Shephard, N., ed (2005), *Stochastic Volatility: Selected Readings*, Oxford: Oxford University Press.
- Taylor, S. J. (1982), “Financial returns modeled by the product of two stochastic processes a study of daily sugar prices 1961-79,” in *Time Series Analysis: Theory and Practice 1*, ed. O. D. Anderson, Amsterdam: North-Holland, pp. 203–226.
- Wang, J. J. J., Chan, J. S. K., & Choy, S. T. B. (2011), “Stochastic volatility models with leverage and heavy-tailed distributions: A Bayesian approach using scale mixtures,” *Computational Statistics and Data Analysis*, 55(1), 852–862.
- Yu, J. (2005), “On leverage in a stochastic volatility model,” *Journal of Econometrics*, 127(2), 165–178. cited By (since 1996) 37.