

# NOTE ON RESTRICTION MAPS OF CHOW RINGS TO WEYL GROUP INVARIANTS

NOBUAKI YAGITA

ABSTRACT. Let  $G$  be an algebraic group over  $\mathbb{C}$  corresponding a compact simply connected Lie group. When  $H^*(G)$  has  $p$ -torsion, we see  $\rho_{CH}^* : CH^*(BG) \rightarrow CH^*(BT)^{W_G(T)}$  is always not surjective. We also study the algebraic cobordism version  $\rho_\Omega^*$ . In particular when  $G = Spin(7)$  and  $p = 2$ , we see each Griffiths element in  $CH^*(BG)$  is detected by an element in  $\Omega^*(BT)$ ,

## 1. INTRODUCTION

Let  $p$  be a prime number. Let  $G$  be a compact Lie group and  $T$  the maximal torus. Let us write  $H^*(-) = H^*(-; \mathbb{Z}_{(p)})$ , and  $BG, BT$  classifying spaces of  $G, T$ . Let  $W = W_G(T) = N_G(T)/T$  be the Weyl group and  $Tor \subset H^*(BG)$  be the ideal generated by torsion elements. Then we have the restriction map

$$\rho_H^* : H^*(BG) \rightarrow H^*(BG)/Tor \subset H^*(BT)^W$$

by using the Becker-Gottlieb transfer.

It is well known that when  $H^*(G)$  is  $p$ -torsion free (hence  $H^*(BG)$  is  $p$ -torsion free), then  $\rho_H^*$  is surjective. However when  $H^*(G)$  has  $p$ -torsion, there are cases that  $\rho_H^*$  are not surjective, which are founded by Feshbach [Fe].

Let us write by  $G_{\mathbb{C}}, T_{\mathbb{C}}$  the reductive group over  $\mathbb{C}$  and its maximal torus corresponding the Lie group  $G, T$ . Let us write simply  $CH^*(BG) = CH^*(BG_{\mathbb{C}})_{(p)}$ ,  $CH^*(BT) = CH^*(BT_{\mathbb{C}})_{(p)}$  the Chow rings of  $BG_{\mathbb{C}}$  and  $BT_{\mathbb{C}}$  localized at  $p$ . We consider the Chow ring version

$$\rho_{CH}^* : CH^*(BG) \rightarrow CH^*(BG)/Tor \subset CH^*(BT)^W \cong H^*(BT)^W.$$

Our first observation is

**Theorem 1.1.** *Let  $G$  be simply connected. If  $H^*(G)$  has  $p$ -torsion, then the map  $\rho_{CH}^*$  is always not surjective.*

---

2010 *Mathematics Subject Classification.* 55N20, 55R12, 55R40.

*Key words and phrases.* Chow ring, algebraic cobordism,  $BSpin(n)$ .

In the proof, we use an element  $x \in H^4(BG)$  with  $\rho_H(x) \notin \text{Im}(\rho_{CH}^*)$ . Hence  $x \notin \text{Im}(cl)$  for the cycle map  $cl : CH^*(BG) \rightarrow H^*(BG)$ . The corresponding element  $1 \otimes x \in CH^*(B\mathbb{G}_m \times BG)$  is the element founded as a counterexample for the integral Hodge and hence the integral Tate conjecture in [Pi-Ya].

Next, we consider elements in *Tor*. To study torsion elements, we consider the following restriction map

$$\text{res}_H : H^*(BG) \rightarrow \Pi_{A:\text{abelian} \subset G} H^*(BA)^{W_G(A)}.$$

There are cases such that  $\text{res}_H$  is not injective, while for many cases  $\text{res}_H$  are injective. We consider the Chow ring version ([To1,2]) of the above restriction map

$$\text{res}_{CH} : CH^*(BG) \rightarrow \Pi_{A:ab.} CH^*(BA)^{W_G(A)} \subset \Pi_{A:ab.} H^*(BA)^{W_G(A)}.$$

In general  $\text{res}_{CH}$  has non zero kernel. In particular, elements in  $\text{Ker}(cl)$  (i.e. Griffiths elements) for the cycle map  $cl : CH^*(BG) \rightarrow H^*(BG)$  are always in  $\text{Ker}(\text{res}_{CH})$ . Namely Griffiths elements are not detected by  $\text{res}_{CH}$ .

On the other hand, if the Totaro conjecture

$$CH^*(BG) \cong BP^*(BG) \otimes_{BP^*} \mathbb{Z}_{(p)}$$

(for the Brown-Peterson cohomology  $BP^*(-)$ ) is correct, then of course all elements in  $CH^*(BG)$  are detected by elements in  $BP^*(BG)$ . We show that there is a case that Griffiths elements are detected by  $\rho_\Omega^*$  the restriction for algebraic cobordism theory  $\Omega^*(-)$ .

Let  $\Omega^*(X) = MGL^{2*,*}(X) \otimes_{MU_{(p)}^*} BP^*$  be the  $BP$ -version of the algebraic cobordism defined by Voevodsky, Levine-Morel ([Vo1], [Le-Mo1,2]) such that  $CH^*(X) \cong \Omega^*(X) \otimes_{BP^*} \mathbb{Z}_{(p)}$ . In particular, we consider the case  $G = Spin(7)$  and  $p = 2$ . We note that there are (non zero) Griffiths elements in  $CH^*(BG)$ .

**Theorem 1.2.** *Let  $G = Spin(7)$  and  $p = 2$ . Then each Griffiths element (in  $CH^*(BG)$ ) is detected by an element in  $\Omega^*(BT)^W \cong BP^*(BT)^W$ .*

In §2 we study the map  $\rho_H^*$  for the ordinary cohomology theory, and recall Feshbach's result. In §3, we study the Chow ring version and show Theorem 1.1. In §4, we study the case  $G = Spin(n)$ . In §5, we study the  $BP^*$ -version and the algebraic cobordism version for the restriction  $\rho^*$ . In §6, we write down the case  $G = Spin(7)$  quite explicitly, and show Theorem 1.2. In the last section, we note some partial results for the exceptional group  $G = F_4$  and  $p = 3$ .

The author thanks Kirill Zainoulline to start considering this problem, and Masaki Kameko who gives many comments and suggestions, and lets the author know works by Benson-Wood and Feshbach.

## 2. COHOMOLOGY THEORY AND FESHBACH THEOREM

Let  $p$  be a prime number. Let  $G$  be a compact Lie group and  $T$  the maximal Torus. Then we have the restriction map

$$\rho_H^* : H^*(BG) \rightarrow H^*(BT)^W$$

where  $H^*(-) = H^*(-; \mathbb{Z}_{(p)})$ ,  $BG, BT$  are classifying spaces and  $W = W_G(T) = N_G(T)/T$  is the Weyl group.

It is well known that when  $H^*(G)$  is  $p$ -torsion free, then  $\rho_H^*$  is surjective (and hence is isomorphic). However when  $H^*(G)$  has  $p$ -torsion, there are cases that  $\rho_H^*$  are not surjective by Feshbach.

For a connected compact Lie group  $G$ , we have the Becker-Gottlieb transfer  $\tau : H^*(BT) \rightarrow H^*(BG)$  such that  $\tau\rho_H^* = \chi(G/T)$  for the Euler number  $\chi(-)$ , and  $\rho_H^*\tau(x) = \chi(G/T)x$  for  $x \in H^*(BT)^W$ . Let  $\chi(G/T) = N$  and  $Tor$  be the ideal of  $H^*(BG)$  generated by torsion elements. Then we have the injections

$$N \cdot H^*(BT)^W \subset H^*(BG)/Tor \subset H^*(BT)^W.$$

Feshbach found good criterion to see  $\rho_H^*$  is surjective.

**Theorem 2.1.** (Feshbach [Fe]) *The restriction  $\rho_H^*$  is surjective if and only if  $(H^*(BG)/Tor) \otimes \mathbb{Z}/p$  has not nonzero nilpotent elements.*

*Proof.* (Feshbach) First note that  $H^*(BT) \cong \mathbb{Z}_{(p)}[t_1, \dots, t_\ell]$  for  $|t_i| = 2$ . Hence if  $x^m = px'$  in  $H^*(BT)$ , then  $x = px''$  for  $x'' \in H^*(BT)$ . Moreover if  $x \in H^*(BT)^W$ , then so is  $x'$  since  $H^*(BT)$  is  $p$ -torsion free. Thus we see  $H^*(BT)^W \otimes \mathbb{Z}/p$  has no non zero nilpotent elements.

Assume that  $\rho_H^*$  is not surjective, and  $x \in H^*(BT)^W$  but  $x \notin Im(\rho_H^*)$ . Let  $s \geq 1$  be the smallest number such that  $p^s x = \rho_H^*(y)$  for some  $y \in H^*(BG)$ . Hence  $y \neq 0 \mod(p)$ . Then

$$\rho_H^*(y^N) = (p^s x)^N = p^{sN} x^N \in pN \cdot H^*(BT)^W \subset pIm(\rho_H^*).$$

This means  $y$  is a nonzero nilpotent element in  $(H^*(BG)/Tor) \otimes \mathbb{Z}/p$ .  $\square$

Using this theorem, Feshbach [Fe] showed  $\rho_H^*$  is surjective for  $G = G_2, Spin(n)$  for  $n \leq 10$ , and is not surjective for  $Spin(11), Spin(12)$ . Wood [Wo] showed that  $Spin(13)$  is not surjective but  $Spin(n)$  for  $14 \leq n \leq 18$  are surjective. Benson and Wood solved this problem completely, namely  $\rho^*$  is not surjective if and only if  $n \geq 11$  and  $n = 3, 4, 5 \mod(8)$ .

For odd prime, we consider  $\text{mod}(p)$  version

$$\rho_{H/p} : H^*(BG; \mathbb{Z}/p) \rightarrow H^*(BT; \mathbb{Z}/p)^W \cong (H^*(BT)/p)^W.$$

It is known that  $\rho_{H/p}^*$  are surjective when  $G = F_4$  for  $p = 3$  by Toda [Tod] using a completely different arguments. Also different arguments (but without computations of  $H^*(BT)^W$  for concrete cases), Kameko and Mimura [Ka-Mi] prove that  $\rho_{H/p}^*$  are surjective when  $G = E_6, E_7$  for  $p = 3$  and  $G = E_8$  for  $p = 5$ . (The only remain case is  $G = E_8$ ,  $p = 3$  for odd primes.)

Kameko-Mimura get more strong result. Recall the Milnor  $Q_i$  operation

$$Q_i : H^*(X; \mathbb{Z}/p) \rightarrow H^{*+2p^i-1}(X; \mathbb{Z}/p)$$

defined by  $Q_0 = \beta$  and  $Q_{i+1} = [P^{p^i}Q_i, Q_iP^{p^i}]$  for the Bockstein  $\beta$  and the reduced powers  $P^j$ .

**Theorem 2.2.** (*Kameko-Mimura [Ka-Mi]*) *Let  $G = F_4, E_6, E_7$  for  $p = 3$  or  $E_8$  for  $p = 5$ . Let us write a generator by  $x_4$  in  $H^4(BG) \cong \mathbb{Z}_{(p)}$ . Then we have*

$$H^*(BT; \mathbb{Z}/p)^W \cong H^{\text{even}}(BG; \mathbb{Z}/p)/(Q_1Q_2x_4).$$

**Corollary 2.3.** *For  $(G, p)$  in the above theorem,  $\rho_H^*$  is surjective.*

Since its  $Q_0$ -image is zero, we can identify the element  $Q_1Q_2(x_4)$  is in  $H^*(BG)$  and  $p$ -torsion. The above corollary is immediate from the following lemma.

**Lemma 2.4.** *If the composition*

$$(H^*(BG)/\text{Tor}) \otimes \mathbb{Z}/p \rightarrow H^*(BT)^W/p \rightarrow H^*(BT; \mathbb{Z}/p)^W$$

*is injective, then  $\rho_H^*$  is surjective.*

*Proof.* Let  $\rho_H^*$  be not surjective and  $y \in H^*(BT)^W$  with  $y \notin \text{Im}(\rho_H^*)$ . Then  $p^s y = \rho_H^*(x)$  for  $s \geq 1$  and an additive generator  $x \in H^*(BG)/\text{Tor}$ . Of course  $x = 0 \in (H^*(BT)/p)^W$ .  $\square$

### 3. CHOW RINGS

Let us write by  $G_{\mathbb{C}}, T_{\mathbb{C}}$  the reductive group over  $\mathbb{C}$  and its maximal torus corresponding the Lie group  $G$ . Let  $CH^*(BG) = CH^*(BG_{\mathbb{C}})_{(p)}$  be the Chow ring of  $BG_{\mathbb{C}}$  localized at  $p$ .

The arguments of Feshbach also work for Chow rings since the Becker-Gottlieb transfer is constructed by Totaro [To2].

**Lemma 3.1.** *The restriction map  $\rho_{CH}^*$  of Chow rings is surjective if and only if  $(CH^*(BG)/T) \otimes \mathbb{Z}/p$  has not nonzero nilpotent elements.*

However if  $H^*(G)$  has  $p$ -torsion and  $G$  is simply connected, then  $(CH^*(BG)/Tor) \otimes \mathbb{Z}/p$  always has non zero nilpotent elements. In fact,  $c_2 = px_4 \in CH^4(BG)$  in the proof of Theorem 3.3 below, is nilpotent in  $(CH^*(BG)/(Tor)) \otimes \mathbb{Z}/p$ . However from the proof of the above lemma, we note

**Corollary 3.2.** *If  $x \in CH^*(BT)^W$  but  $x \notin Im(\rho_{CH}^*)$ , then there is  $y \in CH^*(BG)$  such that  $\rho_{CH}^*(y) = p^s x$  for some  $s \geq 1$  and  $y$  is non zero nilpotent element in  $(CH^*(BG)/(Tor)) \otimes \mathbb{Z}/p$ .*

Voevodsky [Vo1,2] defined the Milnor operation  $Q_i$  on the mod  $p$  motivic cohomology (over a perfect field  $k$  of any  $ch(k)$ )

$$Q_i : H^{*,*'}(X; \mathbb{Z}/p) \rightarrow H^{*+2(p^n-1), *'+p^n-1}(X; \mathbb{Z}/p)$$

which is compatible with the usual topological  $Q_i$  by the realization map  $t_{\mathbb{C}} : H^{*,*'}(X; \mathbb{Z}/p) \rightarrow H^*(X(\mathbb{C}); \mathbb{Z}/p)$  when  $ch(k) = 0$ . In particular, note for smooth  $X$ ,

$$Q_i|CH^*(X)/p = Q_i|H^{2*,*}(X; \mathbb{Z}/p) = 0.$$

(See §2 in [Pi-Ya] for details.) We will prove the following theorem without using Feshbach theorem (Lemma 3.1).

**Theorem 3.3.** *Let  $G$  be simply connected and  $H^*(G)$  has  $p$ -torsion. Then the restriction map*

$$\rho_{CH}^* : CH^2(BG) \rightarrow CH^2(BT)^W$$

*is not surjective.*

*Proof.* (See §2, 3 in [Pi-Ya].) At first, we note that  $H^*(BT)^W \cong CH^*(BT)^W$  since  $H^*(BT) \cong CH^*(BT)$ . If  $H^*(G)$  has  $p$ -torsion, then  $G$  has a subgroup isomorphic to  $G_2$  (resp.  $F_4$ ,  $E_8$ ) for  $p = 2$  (resp.  $p = 3, 5$ ). (For details, see [Ya2] or §3 in [Pi-Ya].) We prove the theorem for  $p = 2$  but the other cases are proved similarly.

It is known that the inclusion  $G_2 \subset G$  induces a surjection  $H^4(BG) \rightarrow H^4(BG_2) \cong \mathbb{Z}_{(2)}$  and let us write by  $x_4$  its generator. Then it is also known  $Q_1 x_4 \neq 0$  in  $H^*(BG_2; \mathbb{Z}/2)$  where  $Q_1$  is the Milnor operation. Therefore  $x_4 \in H^4(BG_2)$  is not in the image of the cycle map

$$cl : CH^2(BG_2) \rightarrow H^4(BG_2).$$

On the other hand, the element  $2x_4$  is in  $Im(cl)$  because it is represented by the second Chern class  $c_2$ . Since  $\rho^* \otimes \mathbb{Q}$  is an isomorphism,  $\rho_H^*(x_4) \neq 0$ . But  $\rho_H^*(x_4)$  is not in the image  $\rho_{CH}^*$ .  $\square$

**Remark.** The condition of simply connected is necessary. By Vistoli ([Vi], [Ka-Mi]), it is known that  $\rho_{CH}^*$  is surjective for  $G = PGL(p)$ .

**Remark.** The above theorem is also proved by seeing that  $x_4$  is not generated by Chern classes, since  $CH^2(X)$  is always generated by Chern classes [To2].

Recall that for a smooth projective complex variety  $X$ , the integral Hodge conjecture is that the cycle map

$$cl_{/Tor} : CH^*(X) \rightarrow H^{2*}(X)/Tor \cap H^{*,*}(X)$$

is surjective where  $H^{*,*}(X) \subset H^{2*}(X; \mathbb{C})$  is the submodule generated by  $(*, *)$ -forms. Since  $px_4 = c_2$  in the proof of the above theorem and  $c_2 \in H^{*,*}(X)$ , we see  $x_4 \in H^{*,*}(X)$ .

We know [To1], [Pi-Ya] that  $B\mathbb{G}_m \times BG$  can be approximated by smooth projective varieties. Hence counterexamples for the integral Hodge conjecture with  $X = B\mathbb{G}_m \times BG$  give the examples such that  $\rho_{CH}^*$  is not surjective.

**Lemma 3.4.** *Let  $1 \otimes y \notin Im(cl_{/Tor}) \subset H^*(B\mathbb{G}_m \times BG)$  be a counterexample of the integral Hodge conjecture. Then it gives an example such that  $\rho_{CH}^*$  is not surjective, namely,  $\rho_H^*(y) \notin Im(\rho_{CH}^*)$ .*

*Proof.* First note that  $\rho_{H/Tor}^* : H^*(BG)/Tor \rightarrow H^*(BT)^W$  is injective. Since  $CH^*(BT)^W \cong H^*(BT)^W$ , we note  $\rho_{CH}^* = \rho_{H/Tor}^* cl_{/Tor}$ . Therefore  $y \notin Im(cl_{/Tor})$  implies that  $\rho_H^*(y) \notin Im(\rho_{H/Tor}^* cl_{/Tor}) = Im(\rho_{CH}^*)$ .  $\square$

For each prime  $p$ , there are counterexamples  $X = B\mathbb{G}_m \times BG$  for the integral Hodge conjecture, while they are not simply connected. Indeed, Kameko, Antineau and Tripaphy ([Ka1,2], [An], [Tr]) show this for  $G = (SL_p \times SL_p)/\mathbb{Z}/p$  and  $SU(p^2)/\mathbb{Z}/p$ . Hence they give the examples such that  $\rho_{CH}^*$  are not surjective for non simply connected and all  $p$  cases. They proved these facts by using Chern classes.

We also note its converse. Recall [Pi-Ya] that the integral Tate conjecture over a finite field  $k$  is the  $ch(k) > 0$  version of the integral Hodge conjecture.

**Lemma 3.5.** *Let  $x \in H^*(BT)^W$  such that  $x \notin \rho_{CH}^*$  but  $x = \rho_H^*(y)$ . Moreover let  $p^s y$  be represented by Chern class for  $s \geq 1$ . Then  $1 \otimes y \in H^*(B\mathbb{G}_m \times BG)$  gives a counterexample of the integral Hodge conjecture. It also gives a counterexample of the integral Tate conjecture for a finite field  $k$  of  $ch(k) \neq p$*

*Proof.* Since  $p^s y$  is represented by a Chern class, we see  $p^s y \in Im(cl)$ . Hence it is contained in  $H^{*,*}(B\mathbb{G}_m \times BG)$ . Hence so is  $y$ . Since  $x \notin \rho_{CH}^*$ , we see  $y \notin cl_{/Tor}$ . For arguments for the integral Tate conjecture see [Pi-Ya].  $\square$

4.  $Spin(n)$  FOR  $p = 2$ 

In this section, we study Chow rings for the cases  $G = Spin(n)$ ,  $p = 2$ . Recall that the  $mod(2)$  cohomology is given by Quillen [Qu1]

$$H^*(BSpin(n); \mathbb{Z}/2) \cong \mathbb{Z}/2[w_2, \dots, w_n]/J \otimes \mathbb{Z}/2[e]$$

where  $e = w_{2h}(\Delta)$  and  $J = (w_2, Q_0 w_2, \dots, Q_{h-2} w_2)$ . Here  $w_i$  is the Stiefel-Whitney class for the natural covering  $Spin(n) \rightarrow SO(n)$ . The number  $2^h$  is the Radon-Hurewitz number, dimension of the spin representation  $\Delta$  (which is the representation  $\Delta|_C \neq 0$  for the center  $C \cong \mathbb{Z}/2 \subset Spin(n)$ ). The element  $e$  is the Stiefel-Whitney class  $w_{2h}$  of the spin representation  $\Delta$ .

Hereafter this section we always assume  $G = Spin(n)$  and  $p = 2$ .

By Kono [Ko], it is known that  $H^*(BG; \mathbb{Z})$  has no higher 2-torsion, that is

$$H(H^*(BG; \mathbb{Z}/2); Q_0) \cong (H^*(BG)/Tor) \otimes \mathbb{Z}/2$$

where  $H(A; Q_0)$  is the homology of  $A$  with the differential  $d = Q_0$ .

For ease of arguments, let  $n$  be odd i.e.,  $n = 2k + 1$ . Let  $T'$  be a maximal Torus of  $SO(n)$  and  $W' = W_{SO(n)}(T')$  its Weyl group. Then  $W' \cong S_k^\pm$  is generated by permutations and change of signs so that  $|S_k^\pm| = 2^k k!$ . Hence we have

$$H^*(BT')^{W'} \cong \mathbb{Z}_{(2)}[p_1, \dots, p_k] \subset H^*(BT') \cong \mathbb{Z}_{(2)}[t_1, \dots, t_k], \quad |t_i| = 2$$

where the Pontriyagin class  $p_i$  is defined by  $\Pi_i(1 + t_i^2) = \sum_i p_i$ .

For the projection  $\pi : Spin(n) \rightarrow SO(n)$ , the maximal torus of  $Spin(n)$  is given  $\pi^{-1}(T')$  and  $W = W_{Spin(n)}(T) \cong W'$ . To seek the invariant  $H^*(BT)^W$  is not so easy, since the action  $W \cong S_k^\pm$  is not given by permutations and change of signs. Benson and Wood decided the  $H^*(BT')^{W'}$ -algebra structure of  $H^*(BT)^W$  (Theorem 7.1 in [Be-Wo]) and proved

**Theorem 4.1.** (*Benson-Wood*) *Let  $G = Spin(n)$  and  $p = 2$ . Then  $\rho_H^*$  is not surjective if and only if  $n \geq 11$  and  $n = 3, 4, 5 \pmod{8}$  (i.e., the quaternion case).*

Hereafter to study the Chow ring version, we assume  $Spin(n)$  is in real case [Qu1], that is  $n = 8\ell - 1, 8\ell, 8\ell + 1$  (hence  $\rho^*$  is surjective and  $h = 4\ell - 1, 4\ell - 1, 4\ell$  respectively).

In this case, it is known [Qu1] that each maximal elementary abelian 2-group  $A$  has  $rank_2 A = h + 1$  and

$$e|A = \Pi_{x \in H^1(B\bar{A}; \mathbb{Z}/2)}(z + x)$$

where we identify  $A \cong C \oplus \bar{A}$  and  $H^1(B\bar{A}; \mathbb{Z}/2) \cong \mathbb{Z}/2\{x_1, \dots, x_h\}$  is the  $\mathbb{Z}/2$  vector space generated by  $x_1, \dots, x_h$ , and

$$H^*(BC; \mathbb{Z}/2) \cong \mathbb{Z}/2[z], \quad H^*(B\bar{A}; \mathbb{Z}/2) \cong \mathbb{Z}/2[x_1, \dots, x_h].$$

The Dickson algebra is written as a polynomial algebra

$$\mathbb{Z}/2[x_1, \dots, x_h]^{GL_h(\mathbb{Z}/2)} \cong \mathbb{Z}/2[d_0, \dots, d_{h-1}].$$

where  $d_i$  is defined as

$$e|A = z^{2^h} + d_{h-1}z^{2^{h-1}} + \dots + d_0z.$$

Hence we can [Qu1] also identify  $d_i = w_{2^h-2^i}(\Delta) \in H^*(BSpin(n); \mathbb{Z}/2)$ .

**Lemma 4.2.** (*Lemma 2.1 in [Sc-Ya]*) *Milnor operations act on  $d_i$  by*

$$\begin{aligned} Q_{h-1}d_i &= d_0d_i, & Q_{j-1}d_j &= d_0, \text{ for } 1 \leq j, \\ Q_id_j &= 0 \text{ for } i < n-1 \text{ and } i \neq j-1. \end{aligned}$$

**Lemma 4.3.** (*Corollary 2.1 in [Sc-Ya]*) *We have*

$$Q_{h-1}e = d_0e \quad \text{and} \quad Q_ke = 0 \text{ for } 0 \leq k \leq h-2.$$

**Theorem 4.4.** *Let  $T \subset G = Spin(n)$  for  $n = 8\ell, 8\ell \pm 1$ . There is an  $e' \in CH^*(BT)^W$  such that  $e' \notin Im(\rho_{CH}^*)$  and  $\rho_H^*(e) = e' \bmod(2)$ .*

*Proof.* First note that  $e|C = z^{2^h}$ , which is not in the  $Q_0$ -image, and hence  $e$  itself is not. From the preceding Lemma 4.2 we see  $Q_0e = 0$ . By Kono's result, we see

$$0 \neq e \in H(H^*(BG; \mathbb{Z}/2); Q_0) \cong (H^*(BG)/Tor) \otimes \mathbb{Z}/2.$$

Take  $e'' \in H^*(BG)/Tor$  with that  $e'' = e \bmod(2)$ . Then

$$e' = \rho_H^*(e'') \neq 0 \text{ in } H^*(BG)/Tor \subset H^*(BT)^W.$$

From the preceding Lemma 4.2,  $Q_{h-1}(e) \neq 0$ . hence we see  $e' \notin \rho_{CH}^*$  by the existence of  $Q_i$  in the motivic cohomology by Voevodsky.  $\square$

Let  $\Delta_{\mathbb{C}}$  be the complex representation induced from the real representation  $\Delta$ . Then we see (see Theorem 4.2 in [Sc-Ya])

$$c_{2^{h-1}}(\Delta_{\mathbb{C}})|C = 2w_{2^h}|C = 2z^{2^h}.$$

Of course this element  $c_{2^{h-1}}(\Delta_{\mathbb{C}})$  is in the Chow ring  $CH^*(BG)$ . hence we see that we can take  $2e' \in Im(\rho_{CH}^*)$ .

From the result by Benson-Wood, we know  $\rho_H^*$  is surjective in this (real) case. Hence from Lemma 3.5 (or  $Q_{h-1}(e) \neq 0$ ), we have

**Corollary 4.5.** *Let  $X = B\mathbb{G}_m \times BSpin(n)$  with  $n = 8\ell, 8\ell \pm 1$ . The element  $1 \otimes e \in H^{2^h}(X) \cap H^{2^{h-1}, 2^{h-1}}(X)$  gives a counterexample for the integral Hodge and the integral Tate conjectures, namely  $1 \otimes e \notin Im(cl_{H/Tor})$ .*

## 5. COBORDISM

Let  $BP^*(X)$  be the Brown-Peterson cohomology theory with the coefficients ring  $BP^* = \mathbb{Z}_{(p)}[v_1, v_2, \dots]$  of degree  $|v_i| = -2(p^i - 1)$ . Let  $\Omega^*(X) = MGL^{2*,*}(X) \otimes_{MU^*} BP^*$  be the  $BP^*$ -version of the algebraic cobordism ([Vo1], [Le-Mo1,2]) such that  $\Omega^*(X) \otimes_{BP^*} \mathbb{Z}_{(p)} \cong CH^*(X)$ .

We consider the cobordism version of the map  $\rho_H^*$

$$\rho_\Omega^* : \Omega^*(BG) \rightarrow \Omega^*(BT)^W \cong BP^*(BT)^W.$$

Although  $\mathbb{A}^1$ -homotopy category has the Becker-Gottlieb transfer (this fact is announced in [Ca-Jo]), we see

$$\tau \cdot \rho_\Omega^* = \chi(G/T) \mod(v_1, v_2, \dots)$$

which is not  $\chi(G/T)$  in general. So we can not have the  $\Omega^*$ -version of Feshbach's theorem.

We are interesting in an element  $x \in \Omega^*(BT)^W$  which is in  $Im(\rho_\Omega^*)$ . Of course, it is torsion free in  $\Omega^*(X)$ , but there is a case such that

$$0 \neq x \in CH^*(BG)/p \cong \Omega^*(BG) \otimes_{BP^*} \mathbb{Z}/p$$

and  $x$  is  $p$ -torsion in  $CH^*(BG)$ .

**Lemma 5.1.** *Let  $f \in H^*(BT)^W$ ,  $f \neq 0 \mod(p)$ , and identify  $f \in gr\Omega^*(BT) \cong \Omega^* \otimes H^*(BT)$ . Let  $f \notin Im(\rho_\Omega^*)$  but  $v_m f \in Im(\rho_\Omega^*)$ . Then  $v_j f \in Im(\rho_\Omega^*)$  for all  $0 \leq j \leq m$ . Namely, there is  $c_j \in \Omega^*(BG)$  such that  $\rho_\Omega^*(c_j) = v_j f$ ,*

$$c_j \neq 0 \in \Omega^*(BG) \otimes_{BP^*} \mathbb{Z}/p \cong CH^*(BG)/p,$$

moreover  $pc_j = 0$  in  $CH^*(BG)$  for  $j > 0$ .

*Proof.* We consider the Landweber-Novikov cohomology operation  $r_a$  in  $gr\Omega^*(BT) \cong \Omega^* \otimes H^*(BT)$ . By Cartan formula,

$$r_a(v_m f) = \sum_{a=a'+a''} r_{a'}(v_m) r_{a''}(f).$$

Here  $r_{a''}(f) = 0$  for  $|a''| > 0$  in  $gr\Omega^*(BT) \cong \Omega^* \otimes H^*(BT)$ . We have operations  $r_{\beta_j}(v_m) = v_j$  for  $j \leq m$ , and we have the first statement.

From the assumption,  $f$  itself is not in the cycle map  $\rho_{\Omega^*}$ . Hence  $v_j f$  is a  $BP^*$ -module generator in  $\Omega^*(BT)^W \cap Im(\Omega^*(BG))$ . Hence it is also non zero in  $CH^*(BG)/p$ . Since  $pv_j f = v_j pf \in v_j Im(\Omega^*(BG))$ , we have  $pc_j = 0 \in CH^*(BG)$ .  $\square$

We consider the Atiyah-Hirzebruch spectral sequence (AHss)

$$E_2^{*,*'} \cong H^*(X; BP^{*'}) \implies BP^*(X)$$

It is known that

$$(*) \quad d_{2p^i-1}(x) = v_i \otimes Q_i(x) \mod(p, v_1, \dots, v_{i-1}).$$

In general, there are many other types of non zero differential. However we consider cases that differentials are only of this form.

**Lemma 5.2.** *Let  $X = BSpin(n)$  and  $n = 8\ell, 8\ell \pm 1$ . In AHss for  $BP^*(X)$ , assume all non zero differentials are of form  $(*)$ . Then  $pe, v_1e, \dots, v_{h-2}e$  are all permanent cycles.*

*Proof.* We use Lemma 4.2 in the preceding section. First recall  $Q_i(d_0) = 0$ ,  $Q_i(e) = 0$  for  $i < h-1$ . Therefore  $d_0e$  exists in  $E_{2h-1}$ .

Since  $Q_{j-1}d_j = d_n$  and  $Q_k(d_j) = 0$  for  $k < j-1$ , the differential in AHss is

$$d_{2j-1}(d_je) = v_{j-1} \otimes Q_{j-1}(d_je) = v_{j-1}d_0e.$$

Hence we have  $(2, v_1, v_2, \dots, v_{h-2})(d_0e) = 0$  in  $E_{2h-1}^{*,*}$ .

Now we study the differential

$$d_{2h-1}(e) = v_{h-1}Q_{h-1}(e) = v_{h-1}d_0e.$$

Note that  $e$  is  $BP^*$ -free in  $E_{2h-1}^{*,*}$ , since  $e|C = z^{2h}$  and  $e \notin \text{Im}(Q_i)$ . Hence we have

$$\text{Ker}(d_{2h-1}) \cap BP^*\{e\} \cong \text{Ideal}(2, v_1, \dots, v_{h-2})\{e\}.$$

(In this paper,  $R\{a, b, \dots\}$  means the  $R$ -free module generated by  $a, b, \dots$ ) By the assumption  $(*)$  for differentials,  $pe, v_1e, \dots, v_{h-2}e$  are all permanent cycles.  $\square$

For  $7 \leq n \leq 9$ , AHss converging  $BP^*(BSpin(n))$  is computed in [Ko-Ya], ([Sc-Ya] also), and it is known that  $(*)$  is satisfied.

**Corollary 5.3.** *For  $n = 7, 8$  (resp.  $n = 9$ ), the elements  $pe, v_1e$  (resp.  $pe, v_1e, v_2e$ ) are in  $\text{Im}(\rho_{BP}^*) \subset BP^*(BT)^W$  (but  $e$  itself is not).*

Let  $K(n)^*(X)$  be the Morava  $K$ -theory with the coefficients ring  $K(n)^* \cong \mathbb{Z}/p[v_n, v_n^{-1}]$ , and  $AK(n)^*(X) = AK(n)^{2*,*}(X)$  its algebraic version. Here we consider an assumption such that

$$(**) \quad AK(n)^*(BG) \rightarrow K(n)^*(BG) \quad \text{is surjective.}$$

It is known by Merkurjev (see [To1] for details) that  $AK^*(BG) \cong K^*(BG)$  for the algebraic  $K$ -theory  $AK^*(X)$  and the complex  $K$ -theory  $K^*(X)$ , which induces  $AK(1)^*(BG) \cong K(1)^*(BG)$ . Hence  $(**)$  is correct when  $n = 1$  for all  $G$ .

**Lemma 5.4.** *Let  $X = BSpin(n)$ ,  $n = 8\ell, 8\ell \pm 1$  and suppose  $(*), (**)$ . Moreover let  $h \geq 3$ . Then  $v_{h-2}e \in \text{Im}(\rho_\Omega^*)$ , and hence there is  $c_i \in CH^*(X)$  for  $0 \leq i \leq h-2$  in Lemma 5.1.*

*Proof.* First note  $0 \neq v_{h-2}e \in K(h-2)^*(X)$  (hence so is  $e$ ). On the other hand

$$AK(h-2)^*(X) \cong K(h-2)^* \otimes CH^*(X)/I$$

for some ideal  $I$  of  $CH^*(X)$ . Therefore there is an element  $c \in CH^*(X)$  which corresponds  $v_{h-2}^s e$  that is  $cl_\Omega(c) = v_{h-2}^s e$  for  $cl_\Omega : \Omega^*(X) \rightarrow BP^*(X)$ . Since  $e \notin Im(cl_\Omega)$ , we see  $s$  must be positive. The possibility of

$$|v_{h-2}^s e| = -2(2^{h-2} - 1)s + 2^h > 0$$

is  $s = 1$  or  $s = 2$ . Note  $|v_{h-2}^2 e| = 4$ . However it is known by Totaro (Theorem 15.1 in [To2]),

$$cl : CH^2(X) \rightarrow H^4(X) \text{ is injective.}$$

Hence  $s = 1$  and  $cl_\Omega(c) = v_{h-1}e$ .  $\square$

**Corollary 5.5.** *For  $X = BSpin(7)$ , there is an element  $c \in CH^3(X)$  such that  $c \neq 0 \in CH^*(X)/2$ ,  $cl(c) = 0$  and  $\rho_\Omega^*(c) \neq 0 \in \Omega^*(BT)^W$ .*

## 6. $Spin(7)$ FOR $p = 2$

Let  $G$  be a compact Lie group. Consider the restriction map

$$res_{H/p} : H^*(BG; \mathbb{Z}/p) \rightarrow Lim_{V:el.ab} H^*(BV; \mathbb{Z}/p)^{W_G(A)}$$

where  $W_G(A) = N_G(A)/C_G(A)$  and  $V$  ranges in the conjugacy classes of maximal elementary abelian  $p$ -groups. Quillen [Qu2] showed this  $res_{H/p}$  is an  $F$ -isomorphism (i.e. its kernel and cokernel are generated by nilpotent elements). We consider its integral version

$$res_H : H^*(BG) \rightarrow \Pi_{A:ab} H^*(BA)^{W_G(A)},$$

where  $A$  ranges in the conjugacy classes of maximal abelian subgroups of  $G$ .

Hereafter this section, we assume  $G = Spin(7)$  and  $p = 2$  and hence  $h = 3$ . The number of conjugacy classes of the maximal abelian subgroups of  $G$  is two, one is the torus  $T$  and the other is  $A' \cong (\mathbb{Z}/2)^4$  which is not contained in  $T$ . The Weyl group is  $W_G(A') \cong \langle U, GL_3(\mathbb{Z}/2) \rangle \subset GL_4(\mathbb{Z}/2)$  where  $U$  is the maximal unipotent group in  $GL_4(\mathbb{Z}/2)$ . It is well known

$$\begin{aligned} H^*(BG; \mathbb{Z}/2) &\cong H^*(BA'; \mathbb{Z}/2)^{W_G(A')} \\ &\cong \mathbb{Z}/2[w_4, w_6, w_7, w_8], \quad Q_0 w_6 = w_7 \end{aligned}$$

where  $w_i$  for  $i \leq 7$  (resp.  $i = 8$ ) are the Stiefel-Whitney class for the representation induced from  $Spin(7) \rightarrow SO(7)$  (resp. the spin representation  $\Delta$  and hence  $w_8(\Delta) = e$ ).

Since  $H^*(BG)$  has just 2-torsion by Kono, and hence the restriction map  $res_H$  injects  $Tor \subset H^*(BG)$  into  $H^*(BA'; \mathbb{Z}/2)^{W_G(A')}$ .

Next we consider the integral case. Also note  $H^*(BG)$  has just 2-torsion and

$$(H^*(BG)/Tor) \otimes \mathbb{Z}/2 \cong H(H^*(BG; \mathbb{Z}/2); Q_0).$$

Since  $Q_0 w_i = 0$  for  $i \neq 6$  and  $Q_0 w_6 = w_7$ , we have

$$H(H^*(BG; \mathbb{Z}/2); Q_0) \cong \mathbb{Z}/2[w_4, c_6, w_8] \quad c_6 = w_6^2.$$

Of course the right hand side ring has no nonzero nilpotent elements. Hence we see that  $\rho_H^*$  is surjective and

$$H^*(BT)^W \otimes \mathbb{Z}/2 \cong \mathbb{Z}/2[w_4, c_6, w_8].$$

The integral cohomogy is written as

$$H^*(BG) \cong \mathbb{Z}_{(2)}[w_4, c_6, w_8] \otimes (\mathbb{Z}_{(2)}\{1\} \oplus \mathbb{Z}/2[w_7]\{w_7\}).$$

In particular, we note  $res_H$  is injective.

Next we consider the Atiyah-Hirzebruch spectral sequence

$$E_2^{*,*'} \cong H^*(BG) \otimes BP^* \implies BP^*(BG).$$

Its differentials have forms of  $(*)$  in §5. Using  $Q_1(w_4) = x_7$ ,  $Q_1(e_7) = c_7$ ,  $Q_2(w_8) = w_7 w_8$  and  $Q_3(w_7 w_8) = c_7 c_8$ , we can compute the spectral sequence (while it is some what complicated)

$$\begin{aligned} gr BP^*(BG) &\cong BP^*[c_4, c_6, c_8]\{1, 2w_4, 2w_8, 2w_4 w_8, v_1 w_8\} \\ &\quad \oplus BP^*/(2, v_1, v_2)[c_4, c_6, c_7, c_8]\{c_7\}/(v_3 c_7 c_8). \end{aligned}$$

Hence  $BP^*(BG) \otimes_{BP^*} \mathbb{Z}_{(2)}$  is isomorphic to

$$\begin{aligned} &\mathbb{Z}_{(2)}^*[c_4, c_6, c_8]\{1, 2w_4, 2w_8, 2w_4 w_8, v_1 w_8\}/(2v_1 w_8) \\ &\quad \oplus \mathbb{Z}/2[c_4, c_6, c_7, c_8]\{c_7\}. \end{aligned}$$

On the other hand, the Chow ring of  $BG$  is given by Guillot [Gu, Ya]

$$CH^*(BG) \cong BP^*(BG) \otimes_{BP^*} \mathbb{Z}_{(2)}$$

$$\cong \mathbb{Z}_{(2)}[c_4, c_6, c_8] \otimes (\mathbb{Z}_{(2)}\{1, c'_2, c'_4 \cdot c'_6\} \oplus \mathbb{Z}/2\{\xi_3\} \oplus \mathbb{Z}/2[c_7]\{c_7\})$$

where  $cl(c_i) = w_i^2$ ,  $cl(c'_2) = 2w_4$ ,  $cl(c'_4) = 2w_8$ ,  $cl(c'_6) = 2w_4 w_8$ , and  $cl(\xi_3) = 0$ ,  $|\xi_3| = 6$ . Note  $cl_\Omega(\xi_3) = v_1 w_8$  in  $BP^*(BT)^W$ , and  $\xi_3 = c$  in Corollary 5.5. Hence we have

$$\begin{aligned} CH^*(BG)/Tor &\cong \mathbb{Z}_{(2)}[c_4, c_6, c_8]\{1, c'_2, c'_4 \cdot c'_6\} \\ &\subset \mathbb{Z}_{(2)}[w_4, c_6, w_8] \cong CH^*(BT)^W. \end{aligned}$$

In fact the nilpotent ideal in  $(CH^*(BG)/(Tor)) \otimes \mathbb{Z}/2$  is generated by  $c'_2, c'_4, c'_6$ .

Next we consider the Chow rings version for the restriction map

$$res_{CH} : CH^*(BG) \rightarrow \Pi_{A:ab}.CH^*(BA)^{W_G(A)}.$$

Recall  $CH^*(BA') \cong \mathbb{Z}_{(2)}[y_1, \dots, y_4]$  with  $cl(y_i) = x_i^2$ . Hence we have

$$(CH^*(BA')/2)^{W_G(A')} \cong \mathbb{Z}/2[c_4, c_6, c_7, c_8].$$

Since  $Tor$  is just 2-torsion, we have

**Lemma 6.1.** *For the torsion ideal  $Tor$  in  $CH^*(BG)$*

$$res_{CH}(Tor) \cong \mathbb{Z}/2[c_4, c_6, c_8, c_7]\{c_7\} \subset CH^*(BA').$$

Thus we see that  $Ker(res_{CH}) \cong \mathbb{Z}/2[c_4, c_6, c_8]\{\xi_3\}$ , which is the ideal of Griffiths elements. We write down the above results.

**Theorem 6.2.** *Let  $(G, p) = (Spin(7), 2)$ . Let  $Gri f$  be the ideal generated by Griffiths elements and  $D = \mathbb{Z}_{(2)}[c_4, c_6, c_8]$ . Then we have*

$$CH^*(BG)/Tor \cong D\{1, 2w_4, 2w_8, 2w_4w_8\}$$

$$\subset D\{1, w_4, w_8, w_4w_8\} \cong CH^*(BT)^W, \text{ with } w_i^2 = c_i,$$

$$Tor/Gri f \cong D/2[c_7]\{c_7\}, \quad Gri f \cong D/2\{\xi_3\}.$$

There is only one nonzero element  $\xi_3$  in  $CH^3(BG)/2$ , and this  $\xi_3 = c$  in Corollary 5.5 in the preceding section. Thus we see Theorem 1.2 in the introduction.

**Corollary 6.3.** *Take an element  $\xi \in \Omega^*(BG)$  such that  $\xi = \xi_3$  in  $\Omega^*(BG) \otimes_{BP^*} \mathbb{Z}_{(2)} \cong CH^*(BG)$ . We also identify  $c_i \in \Omega^*(BG)$ . Then*

$$\mathbb{Z}/2[c_4, c_6, c_8]\{\xi\} \subset \Omega^*(BT)^W/2.$$

**Theorem 6.4.** *Let  $G = Spin(7)$  and  $p = 2$ . We consider a (non natural) map*

$$CH^*(BG) \subset \Omega^*(BG) \rightarrow \Omega^*(BT)$$

(here the first inclusion is just one section of the projection  $\Omega^*(BG) \rightarrow CH^*(BG)$ ). Then the following restriction map is injective

$$res : CH^*(BG) \subset CH^*(BA') \times \Omega^*(BT).$$

**Corollary 6.5.** *Let  $J \subset BP^*$  is the ideal  $J = (2^2, 2v_1, v_1^2, v_2, \dots)$  so that  $BP^*/J \cong \mathbb{Z}/4\{1\} \oplus \mathbb{Z}/2\{v_1\}$ . Let  $D = \mathbb{Z}_{(2)}[c_4, c_6, c_8]$ . Then*

$$\Omega^*(BG)/J \cong (\Omega^*/J \otimes D\{1, c'_2, c'_4, c'_6, \xi_3\}/(2\xi_3 = v_1c'_4)) \oplus D/2[c_7]\{c_7\}.$$

7. THE EXCEPTIONAL GROUP  $F_4$ ,  $p = 3$ .

In this section, we assume  $(G, p) = (F_4, 3)$ . (However similar arguments also work for  $(G, p) = (E_6, 3), (E_7, 3)$  and  $(E_8, 5)$ , [Ka-Ya].) Toda computed the  $\text{mod}(3)$  cohomology of  $BF_4$ . (For details see [Tod].)

$$H^*(BG; \mathbb{Z}/3) \cong C \otimes D, \quad \text{where}$$

$$C = F\{1, x_{20}, x_{20}^2\} \oplus \mathbb{Z}/3[x_{26}] \otimes \Lambda(x_9) \otimes \{1, x_{20}, x_{21}, x_{26}\}$$

$$D = \mathbb{Z}_{(3)}[x_{36}, x_{48}], \quad F = \mathbb{Z}_{(3)}[x_4, x_8].$$

Using that  $H^*(BG)$  has no higher 3-torsion and  $Q_0x_8 = x_9$ ,  $Q_0x_{20} = x_{21}$ ,  $Q_0x_{25} = x_{26}$ , we can compute

$$H^*(BG) \cong D \otimes C' \quad \text{where}$$

$$C'/\text{Tor} \cong \mathbb{Z}_{(3)}\{x_4\} \oplus E, \quad \text{where } E = F\{ab|a, b \in \{x_4, x_8, x_{20}\}\}$$

$$C' \supset \text{Tor} \cong \mathbb{Z}/3[x_{26}]\{x_{26}, x_{21}, x_9, x_9x_{21}\}.$$

Note  $x_{26} = Q_2Q_1(x_4)$  in Theorem 2.2 and

$$H^*(BT; \mathbb{Z}/p)^W \cong H^{\text{even}}(BG; \mathbb{Z}/3)/(Q_2Q_1x_4) \cong D \otimes F\{1, x_{20}, x_{20}^2\}.$$

Hence we see

$$(H^*(BG)/\text{Tor}) \otimes \mathbb{Z}/3 \cong D/3 \otimes (\mathbb{Z}/3\{1, x_4\} \oplus E) \subset D/3 \otimes F\{1, x_{20}, x_{20}^2\}.$$

Thus from Lemma 2.3, we see  $\rho_H^*$  is surjective and

$$H^*(BT)^W \cong H^*(BG)/\text{Tor} \cong D \otimes (\mathbb{Z}_{(3)}\{1, x_4\} \oplus E).$$

Next we consider the Atiyah-Hirzebruch spectral sequence [Ko-Ya]

$$E_2^{*,*'} \cong H^*(BG) \otimes BP^* \implies BP^*(BG).$$

Its differentials have forms of  $(*)$  in §5. Using  $Q_1(x_4) = x_9$ ,  $Q_1(x_{20}) = x_{25}$ ,  $Q_1(x_{21}) = x_{26}$  and  $Q_2x_9 = x_{26}$  we can compute

$$\text{gr}BP^*(BG) \cong D \otimes (BP^* \otimes (\mathbb{Z}_{(3)}\{1, 3x_4\} \oplus E) \oplus BP^*/(3, v_1, v_2)[x_{26}]\{x_{26}\}).$$

Hence we have

$$BP^*(BG) \otimes_{BP^*} \mathbb{Z}_{(3)} \cong D \otimes (\mathbb{Z}_{(3)}\{1, 3x_4\} \oplus E \oplus \mathbb{Z}/3[x_{26}]\{x_{26}\}).$$

**Proposition 7.1.** *Let  $(G, p) = (F_4, 3)$  and  $\text{Tor} \supset \text{Grif}$  be the ideal generated by Griffiths elements. Then we have*

$$CH^*(BG)/\text{Tor} \subset D \otimes (\mathbb{Z}_{(3)}\{1, 3x_4\} \oplus E) \subset H^*(BG)/\text{Tor},$$

$$\text{Tor}/\text{Grif} \cong D \otimes \mathbb{Z}/3[x_{26}]\{x_{26}\}.$$

If Totaro's conjecture is correct, then  $\text{Grif} = \{0\}$  and the first inclusion is an isomorphism. From [Ya1], it is known that if  $x_8^2 \in \text{Im}(cl)$  for the cycle map  $cl$ , then we can show that  $cl$  itself is surjective. However it seems still unknown whether  $x_8^2 \in \text{Im}(cl)$  or not.

**Corollary 7.2.** *Let  $(G, p) = (F_4, 3)$ . If  $(**)$  in §5 is correct for some  $n \geq 2$ , then the cycle map  $CH^*(BG) \rightarrow BP^*(BG) \otimes_{BP^*} \mathbb{Z}_{(3)}$  is surjective and*

$$CH^*(BG)/Tor \cong D \otimes (\mathbb{Z}_{(3)}\{1, 3x_4\} \oplus E).$$

*Proof.* The corollary follows from  $|v_n x_8^2| = 16 - 2(3^n - 1) \leq 0$ .  $\square$

## REFERENCES

- [An] B. Antieau. On the integral Tate conjecture for finite fields and representation theory. *arXiv:1504.04879v2 [math.AG]* May (2015).
- [Be-Wo] D. Benson and J. Wood. Integral invariants and cohomology of  $BSpin(n)$ . *Topology*. **34** (1994), 13-28.
- [Ca-Jo] G. Carlsson and R. Joshua. Motivic Spanier-Whitehead duality and motivic Becker-Becker-Gottlieb transfer. *www.researchgate.net/* (2011).
- [Fe] M. Feshbach. The image of  $H^*(BG; \mathbb{Z})$  in  $H^*(BT; \mathbb{Z})$  for a compact Lie group with maximal torus  $T$ . *Topology*. **20** (1985), 93-95.
- [Gu] P. Guillot. The Chow rings of  $G_2$  and  $Spin(7)$ . *J. reine angew. Math.* **604** (2007), 137-158.
- [Ka1] M. Kameko. On the integral Tate conjecture over finite fields. *Math. Proc. Cambridge Philos. Soc.* **158** (2015), 531-546.
- [Ka2] M. Kameko. Representation theory and the cycle map of a classifying space. *arXiv:1601.06199v1 [math.AG]* Jan. (2016).
- [Ka-Mi] M. Kameko and M. Mimura, Weyl group invariants. *arXiv: 1202.6459v1 [math.AT]*, (2012).
- [Ka-Ya] M. Kameko and N. Yagita. The Brown-Peterson cohomology of the classifying spaces of the projective unitary group  $PU(p)$  and exceptional Lie groups. *Trans. Amer. Math. Soc.* **360** (2008), 2265-2284.
- [Ko] A. Kono, On the integral cohomology of  $BSpin(n)$ . *J. Math. Kyoto Univ.* **26** (1986), 333-337.
- [Ko-Ya] A. Kono and N. Yagita, Brown-Peterson and ordinary cohomology theories of classifying spaces for compact Lie groups. *Trans. Amer. Math. Soc.* **339** (1993), 781-798.
- [Le-Mo1] M. Levine and F. Morel, Cobordisme algébrique I, *C. R. Acad. Sci. Paris* **332** (2001), 723-728.
- [Le-Mo2] M. Levine and F. Morel, Cobordisme algébrique II, *C. R. Acad. Sci. Paris* **332** (2001), 815-820.
- [Pi-Ya] A. Pirutka and N. Yagita. Note on the counterexamples for the integral Tate conjecture over finite fields. *Document Math. Extra Vol. A. Merkurjev and A. Suslin's sixtieth birthday.* (2015), 501-511.
- [Qu1] D. Quillen, The mod 2 cohomology rings of extra-special 2-groups and the spinor groups, *Math. Ann.* **194** (1971), 197-212.
- [Qu2] D. Quillen, A cohomological criterion for  $p$ -nilpotence. *J. of Pure and Applied Algebra*. **1** (1971), 361-372.
- [Sc-Ya] B. Schuster and N. Yagita, Transfers of Chern classes in BP-cohomology and Chow rings, *Trans. Amer. Math. Soc.* **353** (2001), 1039-1054.
- [Tod] H. Toda, Cohomology mod(3) of the classifying space  $BF_4$  of the exceptional group  $F_4$ . *J. Math. Kyoto Univ.* **13** (1973) 97-115..

- [To1] B. Totaro, The Chow ring of classifying spaces, *Proc. of Symposia in Pure Math. "Algebraic K-theory" (1997:University of Washington, Seattle)* **67** (1999), 248-281.
- [To2] B. Totaro, Group cohomology and algebraic cycles, *Cambridge tracts in Math. (Cambridge Univ. Press)* **204** (2014).
- [Tr] A. Tripathy. Further counterexamples to the integral Hodge conjecture. *arXiv:1601.061/0v1 [math.AG]* Jan. (2016).
- [Vi] A. Vistoli. On the cohomolgy and the Chow ring of the classifying space  $PGL_p$ . *J. reine angew. Math.* **610** (2007), 181-227.
- [Vo1] V. Voevodsky, The Milnor conjecture, *www.math.uiuc.edu/K-theory/0170* (1996).
- [Vo2] V. Voevodsky, Motivic cohomology with  $\mathbb{Z}/2$  coefficient, *Publ. Math. IHES* **98** (2003), 59-104.
- [Wo] J. Wood, Nilpotent elements in the Bockstein spectral sequence for  $BSpin(n)$ . *Contemporary Math.* **146** (1993), 453-469.
- [Ya1] N. Yagita, The image of the cycle map of classifying space of the exceptional Lie group  $F_4$ . *J. Math. Kyoto Univ.* **44** (2004), 181-191.
- [Ya2] N. Yagita, Coniveau filtration of cohomology of groups. *Proc. London Math. Soc.* **101** (2010), 179-206.

FACULTY OF EDUCATION, IBARAKI UNIVERSITY, MITO, IBARAKI, JAPAN  
*E-mail address:* nobuaki.yagita.math@vc.ibaraki.ac.jp,