

ON SPLITS OF COMPUTABLY ENUMERABLE SETS

PETER A. CHOLAK

ABSTRACT. Our focus will be on the computably enumerable (c.e.) sets and trivial, non-trivial, Friedberg, and non-Friedberg splits of the c.e. sets. Every non-computable set has a non-trivial Friedberg split. Moreover, this theorem is uniform. V. Yu. Shavrukov recently answered the question which c.e. sets have a non-trivial non-Friedberg splitting and we provide a different proof of his result. We end by showing there is no uniform splitting of all c.e. sets such that all non-computable sets are non-trivially split and, in addition, all sets with a non-trivial non-Friedberg split are split accordingly.

1. TRIVIAL SPLITS

Given a c.e. set A , a *split* of A is a pair of c.e. sets A_0, A_1 such that $A_0 \sqcup A_1 = A$, \sqcup is disjoint union. If one of A_0 or A_1 is computable the splitting is *trivial*. If A_0 is computable then $A = A_0 \sqcup (\overline{A_0} \cap A)$.

It is straightforward to see that any splitting of a computable set is trivial. Given a c.e. set A , letting $A_0 = \emptyset$ and $A_1 = A$, provides a trivial splitting of A . We would like to avoid splits where one of the sets is finite. It is known that every infinite c.e. set A has an infinite computable subset R . This provides a trivial splitting of A , $A = R \sqcup (A \cap \overline{R})$, into two infinite c.e. sets assuming A is not computable.

Given this, Myhill asked

Question 1.1 (Myhill [9]). *Does every non-computable c.e. set have a non-trivial splitting?*

Myhill's Question was answered positively by Friedberg [5].

2000 *Mathematics Subject Classification*. Primary 03D25.

Draft as of February 2, 2022. We want to thank V. Yu. Shavrukov for allowing us to include his result, Theorem 3.8. Without it, this paper would look very different. This research was started while Cholak participated in the Buenos Aires Semester in Computability, Complexity and Randomness, 2013. Thanks to Rachel Epstein, Greg Igusa, Nathan Pierson, Mike Stob, and the referees for comments and suggestions. My interest in Friedberg splits was sparked in 1989 by Rod Downey. I cannot forgive him.

2. FRIEDBERG SPLITS

Most of this section is known but we wanted to provide an explicit proof of Corollary 2.7. This corollary will be useful later. One of the focuses of this paper is splitting procedures that always produce a non-trivial split when possible.

At this point we will fix the standard uniform enumeration $W_{e,s}$ of all c.e. sets with the convention that at stage s , there is at most one pair e, x where x enters W_e at stage s . Some details on how we can effectively achieve this enumeration can be found in Soare [12, Exercise I.3.11].

Every c.e. set has an index according to this fixed enumeration. For the sets that we construct we have to appeal to Kleene's Recursion Theorem to find this index. Moreover, by the standard trick of slowing down or pausing our construction, we can assume the enumerations of our fixed point W_e and our constructed set A are the same. Our construction, at times, will construct sets other than A . While we will focus on the constructed sets, the actual outcome of our constructions will be a uniform enumeration of all constructed sets. We will be using Kleene's Recursion Theorem with parameters to get a function from each constructed set to an index with the same enumeration for that set in the above enumeration.

By the Padding Lemma, we know that each c.e. set A has infinitely many indices. By Rice's Theorem, we know that for a given c.e. set A the set of indices (in this fixed enumeration) for A is not computable.

Also at this point we will fix the convention that A, B, W, X and Y always refer to c.e. sets with some fixed index in our given enumeration. Now we need the following.

Definition 2.1. A split $A_0 \sqcup A_1 = A$ is a *Friedberg split* of A iff, for all W , if $W - A$ is not c.e. then both $W - A_0$ and $W - A_1$ are not c.e. sets.

Lemma 2.2. *If A is not computable and $A_0 \sqcup A_1$ is a Friedberg split then the split is not trivial.*

Proof. $\mathbb{N} - A$ is not c.e. so $\mathbb{N} - A_0$ and $\mathbb{N} - A_1$ are not c.e. and hence A_0 and A_1 are not computable. \square

Definition 2.3. For $A = W_e$ and $B = W_i$,

$$A \setminus B = \{x \mid \exists s [x \in (W_{e,s} - W_{i,s})]\}$$

and $A \searrow B = A \setminus B \cap B$. (This is with respect to our given enumeration and hence this definition depends on our chosen enumeration.)

By the above definition, $A \setminus B$ is a c.e. set. $A \setminus B$ is the set of balls that enter A before they enter B . If $x \in A \setminus B$ then x may or may not enter B and if x does enter B , it only does so after x enters A (in terms of our enumeration). Since the intersection of two c.e. sets is c.e., $A \searrow B$ is a c.e. set. The c.e. set $A \searrow B$ is the c.e. set of balls that first enter A and then enter B (under the above enumeration). So $A \setminus B$ reads “ A before B ” and $A \searrow B$ reads “ A before B and then B ”.

Note that for all W , $W \setminus A = (W - A) \sqcup (W \searrow A)$. Since $W \setminus A$ is a c.e. set, if $W - A$ is not a c.e. set then $W \searrow A$ must be infinite. (This happens for all enumerations not just our given enumeration.)

Lemma 2.4 (Friedberg). *Assume $A = A_0 \sqcup A_1$, and, for all e , if $W_e \searrow A$ is infinite then both $W_e \searrow A_0$ and $W_e \searrow A_1$ are infinite. Then $A_0 \sqcup A_1$ is a Friedberg split of A .*

Proof. Assume that $W - A$ is not a c.e. set but $X = W - A_0$ is a c.e. set. $X - A = (W - A_0) - A = W - A$ is not a c.e. set. So $X \searrow A$ is infinite which implies that $X \searrow A_0$ is infinite but $X \searrow A_0 = (W - A_0) \searrow A_0 = \emptyset$. Contradiction. \square

Friedberg invented the priority method to split every c.e. set into two disjoint c.e. sets while meeting the hypothesis of the above lemma.

Theorem 2.5 (Friedberg). *Every non-computable set A has a Friedberg split.*

Proof. When a ball x enters A at stage s we add it to one of A_0 or A_1 but which one x enters is determined by priority. Our requirements are:

$\mathcal{P}_{e,i,k}$: if $W_e \searrow A$ is infinite then $|W_e \searrow A_i| \geq k$.

We say x meets $\mathcal{P}_{e,i,k}$ at stage s if $|W_e \searrow A_i| = k - 1$ by stage $s - 1$ and if we add x to A_i at stage s then $|W_e \searrow A_i| = k$ at stage s . Find the smallest $\langle e, i, k \rangle$ that x can meet and add x to A_i at stage s . If no such triple can be found, add x to A_0 at stage s . It is not hard to show that all the $\mathcal{P}_{e,i,k}$ are met. \square

Observe that the procedure in Theorem 2.5 is uniform. Given this we made the following definition and corollary.

Definition 2.6. A computable function h is a *splitting procedure* iff, for all e , if $h(e) = \langle e_0, e_1 \rangle$ then $W_{e_0} \sqcup W_{e_1}$ is a split of A and if W_e is not computable then this split is not trivial. If h is a splitting procedure, we say that $h(e)$ gives a split of W_e or splits W_e .

Corollary 2.7 (of Friedberg's Proof). *There is a splitting procedure h such that if W_e is not computable then $h(e)$ gives a Friedberg split of W_e .*

3. NON-TRIVIAL NON-FRIEDBERG SPLITS

The above section brings us to the following question:

Question 3.1. *When does a c.e. set have a non-trivial non-Friedberg split?*

This question was first asked, in a different form, as Question 1.4 in Cholak [1]. In [1], it was asked if there is a definable collection of c.e. sets such that for each set A in this collection the Friedberg splits of A are a proper subclass of the non-trivial splits of A . This question later appeared, in yet a different form, as Question 4.6 in the first unpublished version of Cholak, Gerdes, and Lange [3]. There it was suggested to compare the class of all c.e. sets all of whose non-trivial splits are Friedberg with the \mathcal{D} -maximal sets (defined below). As we will see in Theorem 3.8 every form of this question was answered by Shavrukov [11]. Shavrukov showed that a c.e. set A has a non-trivial non-Friedberg split iff A is not \mathcal{D} -maximal.

3.1. There are c.e. sets with non-trivial non-Friedberg Splits.

Let R be an infinite, coinfinite, computable set. There is a non-computable c.e. subset of R , call this set K_R . There is a non-computable c.e. subset of \overline{R} , call this set $K_{\overline{R}}$. Let $A = K_R \sqcup K_{\overline{R}}$. Then $K_R \sqcup K_{\overline{R}}$ is a non-trivial split of A . $R - A = R - K_R$ is not c.e. but $R - K_{\overline{R}} = R$ is a c.e. set. So this split is not Friedberg. Please note that the set A and its non-trivial non-Friedberg split are built simultaneously.

See Theorem 3.8 for more examples of sets with non-trivial non-Friedberg Splits. There are published examples of sets with non-trivial non-Friedberg splits. In Section 3.2 of Cholak, Gerdes, and Lange [3], a number of such sets are constructed. But, like in the construction in the above paragraph and Theorem 3.8, for the examples in [3] the set A and its non-trivial non-Friedberg split are built simultaneously.

3.2. There are c.e. sets without non-trivial non-Friedberg Splits.

For this we need the following definitions:

Definition 3.2. (1) $\mathcal{D}(A) = \{B \mid B - A \text{ is a c.e. set}\}$.
 (2) W is *complemented* modulo $\mathcal{D}(A)$ iff there is a c.e. Y such that $W \cup Y \cup A = \mathbb{N}$ and $(W \cap Y) - A$ is a c.e. set.

- (3) A is \mathcal{D} -hhsimple iff, for every c.e. W , W is complemented modulo $\mathcal{D}(A)$.
- (4) A c.e. set W is 0 modulo $\mathcal{D}(A)$ iff $W \in \mathcal{D}(A)$.
- (5) A c.e. set W is 1 modulo $\mathcal{D}(A)$ iff there is a Y such that $Y \cap A = \emptyset$ and $W \cup Y \cup A = \mathbb{N}$.
- (6) A non-computable set A is \mathcal{D} -maximal iff for every W , W is complemented modulo $\mathcal{D}(A)$ and either 0 or 1 modulo $\mathcal{D}(A)$.

Assume W is 0 modulo $\mathcal{D}(A)$. WLOG we can assume $W \cap A = \emptyset$. Then $W \cup \mathbb{N} \cup A = \mathbb{N}$ and $\mathbb{N} \cap W = W$ is disjoint from A . So W is complemented modulo $\mathcal{D}(A)$. If $W - A$ is not c.e. then W is not 0 modulo $\mathcal{D}(A)$. A c.e. set W is 0 modulo $\mathcal{D}(A)$ iff $W - A$ is a c.e. set. The set W is 1 modulo $\mathcal{D}(A)$ as witnessed by Y iff W is complemented by Y modulo $\mathcal{D}(A)$ and Y is 0 modulo $\mathcal{D}(A)$. We will not go through the details but the property of a set A being \mathcal{D} -maximal is definable in the c.e. sets, \mathcal{E} .

Lemma 3.3 (Cholak, Downey, Herrmann). *All non-trivial splits of a \mathcal{D} -maximal set A are Friedberg.*

Proof. Let $A_0 \sqcup A_1 = A$ be a non-trivial split of A . Assume that $W - A$ is not a c.e. set. So $W \cup A$ is 1 modulo $\mathcal{D}(A)$. Then, for some Y , $W \cup A \cup Y = \mathbb{N}$ and $Y \cap A = \emptyset$. If $W - A_0$ is c.e. then $A_0 \sqcup ((W - A_0) \cup A_1 \cup Y) = \mathbb{N}$ and hence A_0 is computable. Contradiction. \square

This result and the above proof explicitly appears in an earlier unpublished version of Cholak, Gerdes, and Lange [3] but not in the published version. It was first implicitly mentioned in Cholak, Downey, and Herrmann [2]. It follows a similar result about maximal sets in Downey and Stob [4].

3.3. The Herrmann and Kummer Splitting Theorem. Shortly we will need the following theorem.¹

Theorem 3.4 (Herrmann and Kummer Splitting Theorem). *Let A and B be c.e. sets such that $A \subseteq B$ and B is non-complemented modulo $\mathcal{D}(A)$. Then there are B_0 and B_1 such that B_i is non-complemented modulo $\mathcal{D}(A)$ and $B_0 \sqcup B_1 = B$.*

¹The Herrmann and Kummer Splitting Theorem appears, in a very different form, in Herrmann and Kummer [7]. This theorem appears in the only if direction of the proof of Theorem 2.4 of Herrmann and Kummer [7] starting on page 63 from the first full paragraph on that page. It is interesting enough to be isolated in its own right as a theorem.

Proof. As balls x enter B they will be enumerated into either B_0 or B_1 . So $B = B_0 \sqcup B_1$. Let Y_e, Z_j be two listings of all c.e. sets. We need to meet the requirements:

$\mathcal{R}_{e,j,i}$: either $B_i \cup A \cup Y_e \neq \mathbb{N}$ or $(B_i \cap Y_e) - A \neq Z_j$.

If we fail to meet this requirement then Y_e and Z_j witness that B_i is complemented modulo $\mathcal{D}(A)$.

We need a *disagreement* function. Let $l(e, j, i, s)$ be the least $x \leq s$ such that either $x \notin B_{i,s} \cup A_s \cup Y_{e,s}$, or $x \in ((B_{i,s} \cap Y_{e,s}) - A_s)$ iff $x \notin Z_{j,s}$. If x does not exist, let $l(e, j, i, s) = s$. The $\lim_s l(e, j, i, s)$ exists iff we will have met $\mathcal{R}_{e,j,i}$.

We will use l to define a *restraint* function, $r(e, j, i, -1) = \langle e, j, i \rangle$ and $r(e, j, i, s)$ is the max of $r(e, j, i, s-1)$ and $l(e, j, i, s)$. Again, the $\lim_s r(e, j, i, s)$ exists iff we will have met $\mathcal{R}_{e,j,i}$. Moreover $r(e, j, i, s)$ is a non-decreasing function in s .

When a ball x enters B at stage s find the least $\langle e, j, i \rangle$ such that $x \leq r(e, j, i, s)$ and add x to B_i .

Let $\langle e, j, i \rangle$ be the least triple such that $\lim_s r(e, j, i, s)$ does not exist. Let x be such that for all $\langle e', j', i' \rangle < \langle e, j, i \rangle$, $\lim_s l(e', j', i', s) < x$. Assume $i = 0$. Then B_1 is computable (for all $y > x$, after $r(e, j, 0, s) > y$, y cannot enter B_1), Y_e and Z_j witness that B_0 is complemented modulo $\mathcal{D}(A)$. Now $Y = Y_e \cap \overline{B_1}$ and Z_j witness that B is complemented modulo $\mathcal{D}(A)$. Contradiction. Similarly if $i = 1$. \square

This construction is uniform. Given an index for B we can uniformly get a split of B via the above theorem. Assume B is 0 modulo $\mathcal{D}(A)$ witnessed by the c.e. set $Z = B - A$. Then \mathbb{N} and Z witness that B and any splits of B are complemented modulo $\mathcal{D}(A)$. Let e' and j' be the least such that $Y_{e'} = \mathbb{N}$ and $Z_{j'} = Z$, and $l(e', j', i, s) = s$ (this last item just takes playing a little with the enumeration of these sets). For some $e \leq e', j \leq j'$ and i , $\lim_s r(e, j, i, s)$ does not exist and the argument above shows that the split is trivial. So if $B \subseteq A$ this split will be trivial. So this theorem does not give rise to a splitting procedure.

If B is not complemented modulo $\mathcal{D}(A)$ then it is open if the above split (as given above) is always Friedberg. We conjecture yes with the following evidence: We can combine the requirements \mathcal{P} from the proof of Theorem 2.5 with the one here to force the split to be a Friedberg split.

We also want to point out that the Herrmann and Kummer Splitting Theorem is very similar to the Owings Splitting Theorem. B is *non complemented modulo* A iff $B - A$ is not co-c.e. iff $\overline{B} \cup A$ is not c.e. The

following theorem is an easy corollary of the Owings Splitting Theorem, [10]. Also see Soare [12, X.2.5].

Theorem 3.5 (Owings). *Let A and B be c.e. sets such that $A \subseteq B$ and B is non-complemented modulo A . Then there are B_0 and B_1 such that B_i is non-complemented modulo A and $B_0 \sqcup B_1 = B$.*

We are not going to provide a proof. The standard proof is Soare [12, X.2.5]. What is not clear is whether this standard proof always provides a Friedberg split and, if $B \subseteq A$, whether the resulting split is non-trivial. We can arrange the enumeration (let $W_0 = \mathbb{N}$) such that if $B \subseteq A$ then the resulting split is non-trivial. But it is open what occurs when we use the standard enumeration. So it is unknown if the Owings Splitting Theorem gives a splitting procedure.

The Owings and the Herrmann and Kummer Splitting theorems are like Friedberg's in that all three are uniform, but unlike Friedberg's in that they do not necessarily provide non-trivial splits when possible. Herrmann and Kummer Splitting Theorem does not give rise to a splitting procedure. It is open if the Owings Splitting Theorem gives rise to a splitting procedure. Friedberg Splitting Theorem does give rise to a splitting procedure.

There is one more (little) known splitting theorem, Hammond [6], which extends all three of the splitting theorems above discussed in this subsection. Let \mathcal{E} be the collection of c.e. sets with inclusion, intersection, union, \emptyset and \mathbb{N} ; this is called the lattice of c.e. sets. An *ideal* of \mathcal{E} is a collection of sets \mathcal{I} such that $\emptyset \in \mathcal{I}$ and \mathcal{I} is closed under subset and inclusion. An ideal \mathcal{I} is Σ_3^0 if the relation $W_e \in \mathcal{I}$ is Σ_3^0 . \mathcal{F} , collection of all finite sets, is an Σ_3^0 ideal. For any A , so are $\mathcal{S}(A) = \{B \mid B \subseteq A\}$ and $\mathcal{D}(A)$. W is *complemented modulo \mathcal{I}* iff there is a Y such that $W \cup Y = \mathbb{N}$ and $W \cap Y$ is in \mathcal{I} . For any A , the Friedberg, Owings, and Herrmann and Kummer Splitting Theorems, respectively, imply any B which is non-complemented modulo \mathcal{F} , $\mathcal{S}(A)$, or $\mathcal{D}(A)$ can be split into B_0 and B_1 such that each B_i is non-complemented modulo \mathcal{F} , $\mathcal{S}(A)$, or $\mathcal{D}(A)$.

Theorem 3.6 (Hammond [6]). *Let \mathcal{I} be any Σ_3^0 ideal. If B is non-complemented modulo \mathcal{I} then B can be split into B_0 and B_1 such that each B_i is non-complemented modulo \mathcal{I} .*

We will not include a proof here. Unlike the other three splitting theorems discussed here the proof is not finite injury. It is uniform in \mathcal{I} . Since \mathcal{I} can equal $\mathcal{D}(A)$, it does not always give rise to a splitting procedure. What happens when \mathcal{I} is $\mathcal{S}(A)$ is open.

3.4. Shavrukov's Result. First we need to use the Herrmann and Kummer Splitting Theorem for the following corollary. The proof is not uniform.

Corollary 3.7. *For all non-computable non- \mathcal{D} -maximal A , there are disjoint X_0 and X_1 such that $X_i - A$ is not c.e. and $A \subseteq X_0 \sqcup X_1$.*

Proof. When A is not \mathcal{D} -hhsimple there is a c.e. X such that $A \subseteq X$ and X is not complemented modulo $\mathcal{D}(A)$. Apply the above Herrmann and Kummer Splitting Theorem to get $X_0 \sqcup X_1 = X$ where the X_i s are also not complemented modulo $\mathcal{D}(A)$. If $X_i - A$ is c.e. then X_i is 0 and hence complemented modulo $\mathcal{D}(A)$. Therefore $X_i - A$ is not a c.e. set.

Otherwise A is \mathcal{D} -hhsimple but not \mathcal{D} -maximal. So there must be a c.e. superset W of A which is not 0 or 1. So $W - A$ is not a c.e. set. There is a Y such that $W \cup Y = \mathbb{N}$, $(W \cap Y) - A$ is c.e. but $Y - A$ is not a c.e. set.

Let $X_0 = W \setminus Y$ and $X_1 = Y \setminus W$. Now $W = X_0 \cup (W \cap Y)$. So $W - A = (X_0 - A) \cup ((W \cap Y) - A)$. The set $(W \cap Y) - A$ is known to be c.e., so if $X_0 - A$ is c.e. then so is $W - A$. Therefore $X_0 - A$ is not a c.e. set. $Y = X_1 \cup (W \cap Y)$. So $Y - A = (X_1 - A) \cup ((W \cap Y) - A)$. $(W \cap Y) - A$ is known to be c.e., so if $X_1 - A$ is c.e. then so is $Y - A$. Therefore $X_1 - A$ is not a c.e. set. \square

Theorem 3.8 (Shavrukov). *All c.e. non-computable non- \mathcal{D} -maximal sets A have non-trivial non-Friedberg splits.*

Proof. By the above corollary, there are disjoint X_0 and X_1 such that $X_i - A$ is not c.e. and $A \subseteq X_0 \sqcup X_1$. If $X_i \cap A$ were computable then $X_i - A = X_i \cap \overline{(X_i \cap A)}$ is c.e. Therefore $X_0 \cap A, X_1 \cap A$ is a non-trivial split of A . $X_0 - A$ is not c.e. but $X_0 - (X_1 \cap A) = X_0$ is a c.e. set. Hence $X_0 \cap A, X_1 \cap A$ is a non-trivial non-Friedberg split. \square

Corollary 3.9 (Shavrukov). *All of A 's non-trivial splits are Friedberg iff A is \mathcal{D} -maximal.*

Again we want to thank V. Yu. Shavrukov for allowing us to include his results. The proof we presented here is very different than Shavrukov's, see [11]. Shavrukov's proof used the fact that every \mathcal{D} -hhsimple is not a diagonal. For the definition of a diagonal set see Kummer [8] and Herrmann and Kummer [7].

4. UNIFORM NON-TRIVIAL NON-FRIEDBERG SPLITS

The question we will answer in this section follows:

Question 4.1. *Is there a splitting procedure h such that all non- \mathcal{D} -maximal sets W_e are split by $h(e)$ into a non-trivial non-Friedberg split?*

The answer is no by the following theorem:

Theorem 4.2. *For every total computable h there is an e such that W_e is not computable and $h(e) = \langle e_0, e_1 \rangle$ then either*

- (1) W_{e_0}, W_{e_1} is not a split of W_e ,
- (2) $W_{e_0} \sqcup W_{e_1}$ is a trivial split of W_e , or
- (3) $W_{e_0} \sqcup W_{e_1}$ is a Friedberg split of W_e and W_e is not \mathcal{D} -maximal.

Moreover given an index for h we can effectively find e .

Hence if h is a splitting procedure then Case (3) applies. Actually, Case (3) applies infinitely often.

Corollary 4.3. *Let h be a splitting procedure. Then there is an infinite set J of indices that, for all $e \in J$, W_e has a non-Friedberg split but the split given by $h(e)$ is a Friedberg split.*

Proof of the Corollary. Let $h_0 = h$ and apply Theorem 4.2 to get e_0 . Only Case (3) can apply. So W_{e_0} has a non-trivial non-Friedberg split but $h(e_0)$ gives a Friedberg split. Inductively, assume for all $j \leq i$, that h_j and distinct e_j exist and that Case (3) applies to W_{e_j} . Let $W_{a_i} \sqcup W_{b_i}$ be a non-trivial non-Friedberg split of W_{e_i} . Let $h_{i+1}(e_i) = \langle a_i, b_i \rangle$ and if $e \neq e_i$ let $h_{i+1}(e) = h_i(e)$. Apply Theorem 4.2 to h_{i+1} to effectively get an e_{i+1} . Case (3) applies to e_{i+1} and $e_{i+1} \neq e_j$, for all $j \leq i$. Let J be the infinite set $\{e_i | i \in \omega\}$. \square

We can create a splitting procedure that is correct on infinite many indices of a non- \mathcal{D} -maximal set. Take $A_0 \sqcup A_1 = A = W_a \sqcup W_b = W_c$ to be a non-trivial non-Friedberg splitting of A . Using the padding lemma, let I be an infinite computable set of indices for A . Define $h(e)$ to be $\langle a, b \rangle$ if $e \in I$ and $h_F(e)$ otherwise, where h_F is from Corollary 2.7. By Rice's Theorem, I is not all indices for A . But the following is open.

Question 4.4. *Is there a splitting procedure h and a c.e. set A with a non-trivial non-Friedberg split such that, if $W_e = A$ then $h(e)$ gives a non-trivial non-Friedberg split of $W_e = A$?*

5. PROOF OF THEOREM 4.2

The goal of the rest of the paper is to provide a proof of the above Theorem 4.2. Assume that we are given h and we will construct A . Via the Recursion Theorem we can assume that $W_e = A$. Also assume that $h(e) = \langle e_0, e_1 \rangle$.

For our proof we will work using an oracle for certain Π_2^0 questions. Certainly $\mathbf{0}''$ works but is overkill. The index set of all infinite c.e. sets works nicely. We will use a tree argument to provide answers to our Π_2^0 questions. The tree will also provide a framework for our construction.

We will build A in pieces. First we will construct a Δ_3^0 list of pairwise disjoint computable sets R such that every c.e. set or its complement will be in the union of finitely many of these computable sets and the union of all them is \mathbb{N} . Inside each of these computable sets we will build a piece of A . The default is that A will be maximal inside each R but finite or cofinite inside R are also possible. The construction will ensure that the union of these pieces is a c.e. set A . If A is maximal in only finitely many of these computable sets then A will turn out to be \mathcal{D} -maximal.

We will *try* to construct infinite, coinfinite, computable sets R_i such that, for all j , either

$$(5.0.1) \quad W_j \subseteq^* \bigcup_{i \leq j} R_i \cup A,$$

or

$$(5.0.2) \quad W_j \cup \bigcup_{i \leq j} R_i \cup A =^* \mathbb{N}.$$

(We will remind the reader that $X =^* Y$ iff $(X - Y) \sqcup (Y - X)$ is finite.) Since these sets are meant to be computable we also have to build \overline{R}_i while we are building R_i . Assume that we have built the sets R_i up to j . The balls in $\bigcap_{i < j} \overline{R}_i$ have not yet been added to R_j or A . So our construction will ensure $(\bigcap_{i < j} \overline{R}_i) \setminus A$ is infinite. To build R_j ask if

$$(5.0.3) \quad P_j = (W_j \cap \bigcap_{i < j} \overline{R}_i) \setminus A$$

is infinite. This is a Π_2^0 question. If P_j is infinite, we will build \overline{R}_j as a subset of W_j , so that Equation 5.0.2 is satisfied. When we add balls from the set $\bigcap_{i < j} \overline{R}_i$ to R_j , we will make sure that there is at least one ball in $W_j \cap \bigcap_{i < j} \overline{R}_i$ currently uncommitted. We will add that ball to \overline{R}_j and the rest of the balls under consideration to R_j . We will do this infinitely often. In this case, we satisfy Equation 5.0.2. If P_j is finite, then, since $(\bigcap_{i < j} \overline{R}_i) \setminus A$ is infinite, we just build R_j and \overline{R}_j to be infinite within $\bigcap_{i < j} \overline{R}_i$ and Equation 5.0.1 is satisfied.

Now inside each R_i we will build A to be finite, cofinite, or maximal depending on various outcomes. The default will be for A to be maximal in R_i . To do this we use the construction presented in Soare [12, X.3.3] as a guide to work inside R_i . We will go over the details later. Since maximal sets are not computable, A will not be computable. Assume that A is maximal inside R_i and R_l , where $l \neq i$, then, since $A \cap R_l$ is a non-computable subset of $A \cap \overline{R_i}$, $A = (A \cap R_i) \sqcup (A \cap \overline{R_i})$ is a non-trivial non-Friedberg split of A . The details follow the construction in Subsection 3.1. Now by Theorem 3.8, A is not \mathcal{D} -maximal. If $W_{e_0} \sqcup W_{e_1}$ is not a split of A then we are done. So we may as well assume that $W_{e_0} \sqcup W_{e_1}$ is a split of A .

We will now consider how this split behaves inside each R_i . Since A is maximal inside R_i there are two choices either the split is trivial or Friedberg. We are going to ask an infinite series of questions designed to tell if the split inside R_i is trivial. The questions are is “ $W_k \sqcup (W_{e_0} \cap R_i) = R_i$ ” and is “ $W_k \sqcup (W_{e_1} \cap R_i) = R_i$ ”, for all k . Again these questions are Π_2^0 . A positive answer will tell us the split is trivial inside R_i and which set $W_{e_0} \cap R_i$ or $W_{e_1} \cap R_i$ is computable.

Assume that we get a positive answer and the information that the set $W_{e_0} \cap R_i$ is computable. In this case we will take the following action: Dump almost all of R_l , for $l < i$, into A and, for $l > i$, stop adding balls from R_l into A . In fact, stop building R_l . In this case, A is computable outside R_i and hence W_{e_0} must also be computable. So $W_{e_0} \sqcup W_{e_1}$ is a trivial split of A . We act similarly if $W_{e_1} \cap R_i$ is computable.

If none of the answers to these questions for each R_i is positive then $W_{e_0} \sqcup W_{e_1}$ is a non-trivial split of A . We know inside each R_i the split is Friedberg. We must show that globally the split is Friedberg. Let's consider W_j . If Equation 5.0.1 holds, then $W_j - A \subseteq^* \bigsqcup_{i \leq j} R_i$. So, if $W_j - A$ is not a c.e. set neither are $W_j - W_{e_0}$ and $W_j - W_{e_1}$. So assume Equation 5.0.2 holds, $W_j - A$ is not a c.e. set, but $W_j - W_{e_0}$ is a c.e. set. For any $n > j$, $(W_j - A) \cap R_n$ cannot be a c.e. set. But $(W_j - W_{e_0}) \cap R_n$ is a c.e. set. This contradicts that our split is Friedberg inside R_n . A similar argument works if Equation 5.0.2 holds, $W_j - A$ is not a c.e. set, but $W_j - W_{e_1}$ is a c.e. set. Our split is a Friedberg split.

With one positive answer, we must take action to ensure that our given split is trivial. One positive answer is a Σ_3^0 event. If all questions have negative answers then we have a Π_3^0 event and, in this case, our split is a Friedberg split.

5.1. Coding our Π_2^0 Questions via a Tree. We will work with the tree, $2^{<\omega}$. We consider the tree to grow downward. At the empty node,

λ , we will construct A and $\overline{R}_\lambda = \tilde{R}_\lambda = \mathbb{N}$. At nodes α of length $i^2 > 0$ we will construct R_α and \tilde{R}_α ($\overline{R}_\alpha = \sqcup_{\beta \subset \alpha} R_\beta \sqcup \tilde{R}_\alpha$.) We will call such nodes R -nodes. The idea is that if f is the true path, $|\alpha| = i^2$, and $\alpha \prec f$, then $R_\alpha = R_i$ and $\sqcup_{\beta \subset \alpha} R_\beta \sqcup \tilde{R}_\alpha = {}^*\mathbb{N}$. (We will start indexing the R_i at 1.)

Since we need to ask questions about the potential R_i 's we need the indices for the R_α 's. So the real outcome of our construction is a pair of functions g and \tilde{g} such that $W_{g(\lambda)} = A$, $W_{g(\alpha)} = R_\alpha$, and $W_{\tilde{g}(\alpha)} = \tilde{R}_\alpha$, for all α . Via the Recursion Theorem, we can assume we know g and \tilde{g} prior to the construction. We will use this knowledge to code our questions into the tree.

Let $|\gamma| = j^2 - 1$. Let $\delta \subset \gamma$ such that $|\delta| = (j-1)^2$. At γ we will code the question “Is $(W_j \cap \tilde{R}_\delta) \setminus A$ infinite?”. *Strictly* between two R -nodes of length j^2 and $(j+1)^2$ there are $((j+1)^2 - 1) - j^2 = 2j$ nodes. If $|\gamma| = j^2 + 2k - 2$, $1 \leq k \leq j$, $\beta \preceq \gamma$, and $|\beta| = k^2$, then at γ code the question “Does $W_j \sqcup (W_{e_0} \cap R_\beta) = R_\beta$?”. If $|\gamma| = j^2 + 2k - 1$, $1 \leq k \leq j$, $\beta \preceq \gamma$, and $|\beta| = k^2$, then at γ code the question “Does $W_j \sqcup (W_{e_1} \cap R_\beta) = R_\beta$?”. (The only difference in these two sentences is the length of γ differences by 1 and the second uses W_{e_1} rather than W_{e_0} .)

Via the use of the Recursion Theorem, as we discussed two paragraphs above, these are uniformly Π_2^0 questions. There is a uniform reduction from these questions to the index set of infinite c.e. sets or INF. So uniformly, for all γ , we can associate a c.e. *chip* set C_γ such that C_γ is infinite iff the question coded at γ has a positive answer.

Earlier we have called some nodes R -nodes. These were the nodes whose length is a perfect square. Other than the empty node, we will call the remaining nodes A -nodes; they provide answers to questions coded at α 's predecessor, $\alpha^- = \gamma$. We call an A -node α *positive* iff $\alpha \hat{=} 1 = \gamma$. Otherwise an A -node is negative.

We will inductively define the *true path*, f . λ is on f . Assume that $\alpha \preceq f$. If α is a positive A -node then $f = \alpha$. Otherwise, $\alpha \hat{=} 1 \prec f$ iff C_α is infinite and $\alpha \hat{=} 0 \prec f$ iff C_α is finite. Since nodes of length 0 and 1 are not A -nodes, there is always an R -node on the true path. Either all the A -nodes on f are negative or f is finite and ends with a positive A -node.

A key to the construction is the approximation to the true path at stage s , f_s . Define $f_0 = \lambda$, the empty node. Assume that $\alpha \subseteq f_{s+1}$ and $|\alpha| < s^2$. If α is a positive A -node, let $f_{s+1} = \alpha$. Assume that α is not a positive A -node. Let t be the greatest stage less than $s+1$ such

that $\alpha \subseteq f_t$. If no such stage exists, let $t = 0$. If $C_{\alpha,t} \neq C_{\alpha,s+1}$ then let $\alpha \hat{1} \subseteq f_{s+1}$. Otherwise, $\alpha \hat{0} \subseteq f_{s+1}$.

Since nodes of length s^2 are R -nodes, for $s > 0$, f_s always ends in an R -node or a positive A -node. We say $\alpha <_L \beta$ (or α is to the left of β) iff $\alpha \subsetneq \beta$ or there is a γ such that $\gamma \hat{1} \subseteq \alpha$ and $\gamma \hat{0} \subseteq \beta$. By induction on l , we can show that $\liminf_s f_s \restriction l = f \restriction l$ (the \liminf is measured w.r.t. $<_L$). So, $\liminf_s f_s = f$. If $f_s <_L \alpha$ then there is always a least (in terms of length) R -node or positive A -node, β , such that $\beta \subseteq f_s$ and $\beta <_L \alpha$.

5.2. Action on the Tree. We will use the tree and f_s to construct A , R_α , and \tilde{R}_α , for all α . We think of this construction as a pinball machine. Integers or balls enter at top node, λ , and move downwards and leftwards. The position of a ball, x , at the end of stage s is given by the function $\alpha(x, s)$. The movement on the tree is done such that the $\lim_s \alpha(x, s)$ exists. Let $\alpha(x) = \lim_s \alpha(x, s)$. Initially, $\alpha(x, s)$ is not defined (so x is not on the machine) and, unless explicitly changed, $\alpha(x, s)$ remains the same from stage to stage. For the balls on the machine, at every stage s , $\alpha(x, s)$ is always an R -node or a positive A -node, and $|\alpha(x, s)| \leq x^2$. (The bound x^2 was chosen since balls can only rest at R -nodes or positive A nodes and the length of R -nodes are perfect squares.) If a ball x enters A at stage s , x is removed from the tree at stage s and $\alpha(x, s)$ is undefined again.

Entering the machine and leftward movement is determined by f_{s+1} . Downward movement will be discussed later. Let β be the R -node of length 1 such that $\beta \subseteq f_{s+1}$. Let $\alpha(s, s+1) = \beta$. So all the balls on the machine at stage s are less than s . Assume that $\alpha(x, s) = \alpha$ and $f_{s+1} <_L \alpha$. Then there is always a least (in terms of length) R -node or positive A -node, β , such that $\beta \subseteq f_{s+1}$, $\beta \not\subseteq \alpha$, and $\beta <_L \alpha$. Let $\alpha(x, s+1) = \beta$. Since $|\alpha| \leq x^2 + 1$, the same is true for β . A ball x can only move leftward finitely many times. Since $\liminf_s f_s = f$, $\alpha(x) <_L f$ or $\alpha(x) \subseteq f$.

Assume that α is an R -node. So the length of α is j^2 for some j . Either $\alpha = \alpha^- \hat{1}$ or $\alpha = \alpha^- \hat{0}$. At $\gamma = \alpha^-$ we asked the question “Is $(W_j \cap \tilde{R}_\delta) \setminus A$ infinite?”, where δ is the greatest proper R -subnode of γ . If α ends with a 1, then α believes this set is infinite. If α ends with a 0 then α believes this set is finite. If α ends with a 1 let $P_\alpha = (W_j \cap \tilde{R}_\delta) \setminus A$. Otherwise, let $P_\alpha = \tilde{R}_\delta \setminus A$. We also defined P_α for positive A -nodes to be $P_\alpha = \tilde{R}_\delta \setminus A$, where $\delta \subset \alpha$ is the greatest R -node contained in α . α wants all balls in P_α to go through α . Moreover the construction of A inside R_α requires that α see fresh balls in P_α . So these α are allowed to pull balls in P_α .

We will now work on the remaining movement, pulling, on our pinball machine. An R -node or positive A -node α is allowed to pull balls from subnodes of α or nodes to the right of α . Pulling will be downward or leftward movement. The only downward movement allowed is done via pulling. When α can pull balls is controlled by f_s . When $\alpha \subseteq f_s$, α puts a request coded by s on a list denoted by \mathcal{P}_α at stage s . α can only pull balls when there is an unfulfilled request on the list. If α takes action (as described below) at stage s then the least request on \mathcal{P}_α has been fulfilled. If $f_s <_L \alpha$ then all the current requests at stage s on \mathcal{P}_α are declared fulfilled.

Let α be an R -node or A -node of length l and assume that there is an unfulfilled request on \mathcal{P}_α at stage s . Assume that there are two different balls, x_0 and x_1 , such that $x_i > l$, $x_i \in P_{\alpha,s}$, and either $\alpha <_L \alpha(x_i, s)$ (x_i is to the right of α) or $\alpha(x_i, s) \subset \alpha$ (x_i is above α). For leftmost α and the least such pair, at stage $s + 1$, take the following action: Let $\alpha(x_i, s + 1) = \alpha$ and, if α is a R -node, then put x_0 into $R_{\alpha,s+1}$ and put x_1 into $\tilde{R}_{\alpha,s+1}$. For all balls y , such that $|\alpha|^2 < y < \max x_i$, $y \in \tilde{R}_{\delta,s}$ (using the above notation for δ), and either $\alpha <_L \alpha(x_i, s)$ or $\alpha(x_i, s) \subset \alpha$, let $\alpha(y, s + 1) = \alpha$ and, if α is R -node, then add y to $R_{\alpha,s+1}$. This request is declared fulfilled.

There is just a little more to the construction of R_α . In the next section we will discuss the construction of A inside $R_\alpha \setminus A$. Recall earlier that we said that if a ball enters A it is removed from the machine. That means that none of the above balls added to R_α and \tilde{R}_α are in A . To make sure that R_α is computable when $\alpha \subset f$ we must be sure that almost all balls from \tilde{R}_δ enter R_α or \tilde{R}_α . Because of the construction to the right of the true path, infinitely many balls in \tilde{R}_δ might enter A before they enter R_α or \tilde{R}_α . The balls we are talking about are in the c.e. set $(\tilde{R}_\delta \searrow A) \setminus (R_\alpha \sqcup \tilde{R}_\alpha)$. The above action cannot add these balls to R_α or \tilde{R}_α . So we will simply add these balls to R_α . So the above set is equal to $A \searrow R_\alpha$.

Let's see inductively that for $\alpha \subset f$ and α is an R -node, that $R_\alpha \setminus A$ is infinite, $\tilde{R}_\alpha \setminus A$ is infinite, $\bigsqcup_{\beta \subseteq \alpha} R_\beta \sqcup \tilde{R}_\alpha =^* \mathbb{N}$, and $A \searrow R_\alpha$ is computable. Let δ be the greatest proper R -subnode of α . If no such node exists let $\delta = \lambda$. So by our inductive hypothesis $\tilde{R}_\delta \setminus A$ is infinite. Moreover, by the movement on the tree, only finite many of these balls are ever to the left of α . Ignore those balls. Since α is on the true path, infinitely many requests are placed on \mathcal{P}_α and only finitely many of them are fulfilled because $f_s <_L \alpha$. We claim all of the remaining requests are fulfilled. If not then all but finitely balls of P_α can be pulled by α and α will eventually pull two balls fulfilling the desired

request. So the action discussed two paragraphs above occurs infinitely often. We have ensured that $R_\alpha \setminus A$ and $\tilde{R}_\alpha \setminus A$ are infinite. The sets R_β , for $\beta \subseteq \alpha$, and \tilde{R}_α are all pairwise disjoint. By the action in the above paragraph the union of all these sets is almost everything. Since the disjoint union of these sets is almost everything, we also have that $A \searrow R_\alpha$ is computable. Moreover if α ends with a 1 then $\tilde{R}_\alpha \subseteq W_j$, where $|\alpha| = j^2$, and hence $W_j \cup A \cup \bigcup_{\beta \subseteq \alpha} R_\beta = {}^*\mathbb{N}$ and Equation 5.0.2 holds. If α ends with a 0 then $W_j \subseteq {}^*A \cup \bigcup_{\beta \subseteq \alpha} R_\beta$ and Equation 5.0.1 holds.

Assume f is finite. So $\alpha = f$ is a positive A node. Let γ be the greatest R -subnode of α . Let Z be the set of x such that there is a stage s where $\alpha(x, s) = \alpha$. Z is a c.e. set. Because $\alpha = f$ for almost all balls x in Z , $\alpha(x) = \alpha$. Almost all of the balls in Z never enter A . Z is the end of the line. Recall that $P_\alpha = \tilde{R}_\delta \setminus A$. By the pulling action almost all of the balls in P_α will enter Z . By the above paragraph, $\bigcup_{\beta \subseteq \delta} R_\beta \sqcup \tilde{R}_\delta = {}^*\mathbb{N}$. So Z and $A \searrow \tilde{R}_\delta$ are computable sets.

5.3. The construction of A . We will build A to be maximal inside $R_\alpha \setminus A$. Since $\alpha \subset f$, $R_\alpha \setminus A$ is an infinite computable set. Let $R = R_\alpha \setminus A$. We build $A \cap R$ stagewise based on the construction of a maximal set from Soare [12, Theorem X.3.3].

The main requirement is to ensure that, for all e ,

$$\mathcal{M}_e : \quad W_e \cap R \subseteq {}^*A \cap R \text{ or } (W_e \cap R) \cup (A \cap R) = R.$$

$\sigma(e, x, s) = \{i : i \leq e \wedge x \in W_{i,s}\}$ is the e -state of x at stage s . We will have a series of markers Γ_n^α with $a_n^{\alpha,s}$ denoting the position of Γ_n^α at stage s and such that $\overline{A}_s \cap R = \{a_0^{\alpha,s} < a_1^{\alpha,s} \dots\}$. Each marker Γ_e wants to move to maximize the e -state of $a_e^\alpha = \lim a_e^{\alpha,s}$.

Initially, we let $A_0 \cap R = \emptyset$ and define the $a_n^{\alpha,0}$ accordingly. At stage $s+1$, if there is a least e such that for some least i , $e < i < s$ and $\sigma(e, a_i^{\alpha,s}, s) > \sigma(e, a_e^{\alpha,s}, s)$, then we dump $a_e^{\alpha,s}, a_{e+1}^{\alpha,s}, \dots, a_{i-1}^{\alpha,s}$ into A at stage $s+1$. So $a_e^{\alpha,s+1} = a_i^{\alpha,s}$. Let's call this dumping the *original dumping*. If e does not exist do nothing.

Certain positive A -nodes γ below α can also dump balls from $R = R_\alpha \setminus A$ into A . Let γ be a positive A -node such that $\alpha \subset \gamma$ and at γ^- is coded the question “Is $W_j \sqcup (W_{e_0} \cap R_\beta) = R_\beta$ infinite?” or “Is $W_j \sqcup (W_{e_1} \cap R_\beta) = R_\beta$ infinite?”, for some j and some $\beta \neq \alpha$. γ believes that our split is trivial inside some R_β and wants to dump almost all of R_α into A . Let $t_{\gamma,s}$ be the maximum of $|\gamma|$ and the greatest stage t such that $t \leq s$ and $f_t <_L \gamma$. Assume $\gamma \subset f_{s+1}$, dump $a_s^{\alpha,t_{\gamma,s}}$ into A at stage $s+1$ (if the above movement of balls at stage $s+1$ has already

forced $a_s^{\alpha, t_{\gamma, s}} \neq a_{s+1}^{\alpha, t_{\gamma, s}}$ that is enough). Let's call this dumping, *extra dumping*.

The positive A -nodes to the left or to the right of the true path only dump $a_s^{\alpha, e}$ finitely often. The ones to the left of the true path are only on f_s finitely often and hence only dump finitely many balls from $R_\alpha \setminus A$ into A . If $f <_L \gamma$ then $\lim_s t_{\gamma, s}$ goes to infinity and γ can only dump each $a_s^{\alpha, e}$ into A finitely often.

Assume $\gamma = f_s$ is a positive A -node and $\alpha \subset \gamma$ and at γ^- is coded the question “Is $W_j \sqcup (W_{e_0} \cap R_\beta) = R_\beta$ infinite?” or “Is $W_j \sqcup (W_{e_1} \cap R_\beta) = R_\beta$ infinite?”, for some j and some $\beta \neq \alpha$. Then $\lim_s t_{\gamma, s}$ exists and almost all balls in R_α are dumped into A , i.e. $(R_\alpha \setminus A) \subseteq^* A$. For the rest of this section we will assume the above assumption is false.

So the extra dumping at most dumps each $a_s^{\alpha, e}$ into A finitely often. Assume that $a_s^{\alpha, e}$ will not be dumped after stage s via our extra dumping. Since the original dumping only dumps to increase the e -state and there are 2^e many e -states, the original dumping only dumps $a_s^{\alpha, e}$ finitely often. Hence $\lim_s a_s^{\alpha, e}$ exists and equals $a^{\alpha, e}$.

Now we are in a position to show that the requirements \mathcal{M}_e are met. Assume that \mathcal{M}_i holds for $i < e$ and there is an $(e - 1)$ -state τ such that almost all of $R - A$ are in state τ . Assume all balls greater than k in $R - A$ are in state τ . Let

$$M = \{x : \exists s, n [\sigma(e - 1, x, s) = \tau \wedge n \geq k \wedge x = a_s^{\alpha, n}]\}.$$

So $R - A \subseteq^* M$. Assume $(M \cap W_e) \setminus A$ is finite. Then $W_e \cap R \subseteq^* A \cap R$ and almost all balls in $R - A$ are in e -state τ . Now assume $(M \cap W_e) \setminus A$ is infinite. Let $n \geq k$ and $\sigma(e, a_s^{\alpha, n}, s) = \tau$. Since eventually there will be an m and stage t where $\sigma(e, a_t^{\alpha, m}, t) = \tau \cup \{e\}$, $a^{\alpha, n} \neq a_s^{\alpha, n}$. So $R - A \subseteq^* W_e$. So, A is maximal inside R_α .

5.4. Putting it all together. Recall that if $\alpha = f$ is a positive A -node then, for some j , some i , and some $\beta \subset f$, W_j witnesses that $W_{e_i} \cap A$ is a computable subset of R_β . In this case, a Σ_3^0 event occurs. By the work in the above paragraph, we know that $A \cap R_\beta$ is maximal in R_β and hence $A \cap R_\beta$ is not computable. So A is not computable. So if $W_{e_0} \sqcup W_{e_1}$ is not a split of A we are done. Assume otherwise. By our assumption, inside R_β , $W_{e_0} \sqcup W_{e_1}$ is the trivial split. Let δ be the greatest R -subnode of α . By work in the last paragraph of Section 5.2, there is a set Z , such that $\bigsqcup_{\gamma \subseteq \delta} R_\gamma \sqcup (A \setminus \bar{R}_\delta) \sqcup Z =^* \mathbb{N}$ and $Z \setminus A = \emptyset$. Now, by the above section, for all γ , such that $\gamma \subset \alpha$ and $\gamma \neq \beta$, $A \cap R_\gamma =^* R_\gamma$. Therefore outside of R_β , A is computable; i.e. $A \cap \bar{R}_\beta$ is computable. Any split of a computable set is trivial.

Therefore, $W_{e_0} \sqcup W_{e_1}$ is a trivial split of A . So, by Theorem 4.2, A is \mathcal{D} -maximal.

For the remaining part of this paper, assume that f is an infinite path through $2^{<\omega}$. So a Π_3^0 event occurs. In this case, by the above section, for all $\alpha \subset f$, where α is an R -node, $A \cap R_\alpha$ is not computable. So A is not computable. If $W_{e_0} \sqcup W_{e_1}$ is not a split of A we are done. Assume otherwise. Let α be any R -node where $\alpha \subset f$. Let $A = (A \cap R_\alpha) \sqcup (A \cap \overline{R_\alpha})$. There is an R -node $\beta \neq \alpha$ on the true path. $A \cap R_\beta$ is also not computable. Hence, $(A \cap \overline{R_\alpha})$ is not computable and $A = (A \cap R_\alpha) \sqcup (A \cap \overline{R_\alpha})$ is a non-trivial split of A . The split is not Friedberg, since $R_\alpha - A$ is not a c.e. set but $R_\alpha - (A \cap \overline{R_\alpha}) = R_\alpha$ is computable. Therefore, by Theorem 4.2, A is not \mathcal{D} -maximal.

It just remains to show that $W_{e_0} \sqcup W_{e_1}$ is a Friedberg split of A . We know that for all $\gamma \subset f$, $W_{e_0} \sqcup W_{e_1}$ is a Friedberg split of A inside R_γ . Since splits of maximal sets are either trivial or Friedberg, otherwise the above Σ_3^0 event occurs. We must show that globally the split is Friedberg. Let's consider W_j . Let $\alpha \subset f$ such that $|\alpha| = j^2$. By the work in the second to last paragraph of 5.2, either $W_j \subseteq^* A \cup \bigsqcup_{\beta \subseteq \alpha} R_\beta$ or $W_j \cup A \cup \bigsqcup_{\beta \subseteq \alpha} R_\beta =^* \mathbb{N}$. In the first case, if $W_j - A$ is not a c.e. set neither are $W_j - W_{e_0}$ and $W_j - W_{e_1}$. Assume that $W_j \cup A \cup \bigsqcup_{\beta \subseteq \alpha} R_\beta =^* \mathbb{N}$. Furthermore, assume $W_j - A$ is not a c.e. set, but $W_j - W_{e_0}$ is a c.e. set. For any γ , where $\alpha \subset \gamma \subset f$, $(W_j - A) \cap R_\gamma$ cannot be a c.e. set since this set contains $R_\gamma - A$ and A is maximal inside R_γ . But, since $W_j - W_{e_0}$ is a c.e. set, $(W_j - W_{e_0}) \cap R_\gamma$ is a c.e. set. This contradicts the fact that our split is Friedberg inside R_γ . A similar argument works if $W_j - A$ is not a c.e. set, but $W_j - W_{e_1}$ is a c.e. set. So our split is a Friedberg split. \square

REFERENCES

- [1] Peter Cholak. Some recent research directions in the computably enumerable sets. To appear in the book “The Incomputable” in the Springer/CiE book series, 2013. 3
- [2] Peter Cholak, Rod Downey, and Eberhard Herrmann. Some orbits for \mathcal{E} . *Ann. Pure Appl. Logic*, 107(1-3):193–226, 2001. ISSN 0168-0072. doi: 10.1016/S0168-0072(00)00060-9. URL [http://dx.doi.org/10.1016/S0168-0072\(00\)00060-9](http://dx.doi.org/10.1016/S0168-0072(00)00060-9). 3.2
- [3] Peter A. Cholak, Peter Gerdes, and Karen Lange. \mathcal{D} -maximal sets. *J. Symb. Log.*, 80(4):1182–1210, 2015. ISSN 0022-4812. URL [arXiv:1401.1266v1](https://arxiv.org/abs/1401.1266v1). The Url is to the first unpublished version of this paper. The first and last citation of this paper are actually to this version. 3, 3.1, 3.2

- [4] R. G. Downey and M. Stob. Automorphisms of the lattice of recursively enumerable sets: Orbits. *Adv. in Math.*, 92:237–265, 1992. [3.2](#)
- [5] Richard M. Friedberg. Three theorems on recursive enumeration. I. Decomposition. II. Maximal set. III. Enumeration without duplication. *J. Symb. Logic*, 23:309–316, 1958. ISSN 0022-4812. [1](#)
- [6] Todd Hammond. Friedberg splittings in Σ_3^0 quotient lattices of \mathcal{E} . *J. Symbolic Logic*, 64(4):1403–1406, 1999. ISSN 0022-4812. doi: 10.2307/2586786. URL <http://dx.doi.org/10.2307/2586786>. [3.3](#), [3.6](#)
- [7] Eberhard Herrmann and Martin Kummer. Diagonals and \mathcal{D} -maximal sets. *J. Symbolic Logic*, 59(1):60–72, 1994. [1](#), [3.4](#)
- [8] Martin Kummer. Diagonals and semihyperhypersimple sets. *J. Symbolic Logic*, 56(3):1068–1074, 1991. [3.4](#)
- [9] John Myhill. Problems. *J. Symb. Logic*, 21:215, 1956. This question was the eighth problem appearing in this section. [1.1](#)
- [10] James C. Owings. Recursion, metarecursion, and inclusion. *J. Symbolic Logic*, 32:173–178, 1967. [3.3](#)
- [11] V. Yu. Shavrukov. Friedberg splittings and \mathcal{D} -maximal sets. A letter from V. Yu. Shavrukov to Peter Cholak and James Schmerl, May 16 2015. [3](#), [3.4](#)
- [12] Robert I. Soare. *Recursively Enumerable Sets and Degrees*. Perspectives in Mathematical Logic, Omega Series. Springer–Verlag, Heidelberg, 1987. [2](#), [3.3](#), [3.3](#), [5](#), [5.3](#)

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF NOTRE DAME, NOTRE DAME, IN 46556-5683

E-mail address: Peter.Cholak.1@nd.edu

URL: <http://www.nd.edu/~cholak>