

# A dynamical presentation of the better than nice metric on the disc

Daniela Emmanuele and Marcos Salvai\*

## Abstract

This is a postprint of our paper *Force free Moebius motions of the circle* (J. Geom. Symmetry Phys. 27 (2012) 59–65), which we hadn't uploaded to arXiv previously. We would like to draw attention to the relationship with the article *A geometry where everything is better than nice*, by Larry Bates and Peter Gibson (to appear in Proc. Amer. Math. Soc.).

In our note we treat thoroughly a simple particular case of two previous, more substantial articles. We describe the force free Moebius motions of the circle, that is, the geodesics of the Lie group  $\mathcal{M} \simeq \mathrm{PSL}_2(\mathbb{R}) \simeq \mathrm{O}_o(1, 2)$  of Moebius transformations of the circle, equipped with the Riemannian metric given by the kinetic energy induced by the action. It turns up that  $\mathcal{M}$  decomposes as a Riemannian product  $S^1 \times \Delta$ , where  $\Delta$  is the unit disc endowed with a certain metric, which we now recognize as being essentially the one that is better than nice. Concerning geodesics, we only give the differential equation for their trajectories (using Clairaut's Theorem); we took pleasure in learning from the article by Bates and Gibson that they are actually hypocycloids.

MSC 2010: 70 K 25, 22 E 70, 53 C 22.

Key words: Force free motion, Moebius transformation, Riemannian metric, geodesic, Clairaut

## 1 Introduction

In the spirit of the classical description of the force free motions of a rigid body in Euclidean space using an invariant metric on  $\mathrm{SO}(3)$  [1, Appendix 2], the second author defined in [5] an appropriate metric on the Lorentz group  $\mathrm{SO}_o(n+1, 1)$  to study force free conformal motions of the sphere  $\mathbb{S}^n$ , obtaining a few explicit ones (only through the identity and those which can be described using the Lie structure

---

\*Partially supported by Conicet, Foncyt, Secyt Univ. Nac. Córdoba

of the configuration space). In this note, in the particular case  $n = 1$ , that is, Moebius motions of the circle, we obtain all force free motions.

This is an example of a situation in which using concepts of Physics one can state and solve a problem in Differential Geometry.

Notice that the canonical action of  $\mathrm{PSL}(2, \mathbb{R})$  on  $\mathbb{R}P^1 \cong \mathbb{S}^1$  is equivalent to the action of the group of Moebius transformations on the circle. Then, the results presented here, up to a double covering, also extend the case  $n = 1$  of [4], where force free projective motions of the sphere  $\mathbb{S}^n$  were studied.

This note, as well as [5, 4], is weakly related with mass transportation [7]. In our situation, the set of admitted mass distributions is finite dimensional, and also the allowed transport maps are very particular.

We would like to mention that the closure of  $\mathcal{M}$  in the space of all measurable functions of the circle into itself provides a compactification of  $\mathcal{M}$  as an open dense subset of the three-sphere, with a dynamical meaning related to generalized flows (see [6]).

## 1.1 Moebius motions of the circle

Let  $\mathbb{S}^1$  be the unit circle centered at zero in  $\mathbb{C}$  with the usual metric and let  $\mathcal{M}$  be the Lie group of Moebius transformations of the circle, that is, the group of Moebius transformations of the extended plane preserving the circle. It consists of maps of the form  $cT_\alpha$ , where  $c \in \mathbb{S}^1$  and

$$T_\alpha(z) = \frac{z + \alpha}{1 + \bar{\alpha}z} \quad (1)$$

for  $\alpha \in \mathbb{C}$ ,  $|\alpha| < 1$  and all  $z \in \mathbb{S}^1$ . Although we are interested in the action of  $\mathcal{M}$  on the circle, we recall that if the unit disc  $\Delta = \{z \in \mathbb{C}; |z| < 1\}$  carries the canonical Poincaré metric of constant negative curvature  $-1$  and  $\alpha \neq 0$ , then  $T_\alpha$  is the transvection translating the geodesic with end points  $\pm\alpha/|\alpha|$ , sending 0 to  $\alpha$ .

A *Moebius motion* of the circle is by definition a smooth curve in  $\mathcal{M}$ , thought of as a curve of diffeomorphisms of the circle. (Throughout the paper, smooth means of class  $C^\infty$ .)

In the next two subsections we recall, specialized for the circle, some definitions and statements given in [5] for conformal motions on the  $n$ -dimensional sphere.

## 1.2 The energy of Moebius motions of the circle

Suppose that the circle has initially a homogeneous distribution of mass of constant density 1 and that the particles are allowed to move only in such a way that two configurations differ in an element of  $\mathcal{M}$ . The configuration space may be naturally identified with  $\mathcal{M}$ .

Let  $\gamma : [t_0, t_1] \rightarrow \mathcal{M}$  be a Moebius motion of  $\mathbb{S}^1$ . The total kinetic energy  $E_\gamma(t)$  of the motion  $\gamma$  at the instant  $t$  is given by

$$E_\gamma(t) = \frac{1}{2} \int_{\mathbb{S}^1} |v_t(q)|^2 \rho_t(q) dm(q), \quad (2)$$

where integration is taken with respect to the canonical volume form of  $\mathbb{S}^1$  and, if  $q = \gamma(t)(p)$  for  $p \in \mathbb{S}^1$ , then

$$v_t(q) = \left. \frac{d}{ds} \right|_t \gamma(s)(p) \in T_q \mathbb{S}^1, \quad \rho_t(q) = 1 / \det(d\gamma(t)_p)$$

are the velocity of the particle  $q$  and the density at  $q$  at the instant  $t$ , respectively. Applying to (2) the formula for change of variables, one obtains

$$E_\gamma(t) = \frac{1}{2} \int_{\mathbb{S}^1} \left| \left. \frac{d}{ds} \right|_t \gamma(s)(p) \right|^2 dm(p). \quad (3)$$

The kinetic energy of  $\gamma$  is defined by

$$E(\gamma) = \int_{t_0}^{t_1} E_\gamma(t) dt.$$

The following definition is based on the principle of least action.

**Definition 1** *A smooth curve  $\gamma$  in  $\mathcal{M}$ , thought of as a Moebius motion of  $\mathbb{S}^1$ , is said to be force free if it is a critical point of the kinetic energy functional, that is,*

$$\left. \frac{d}{ds} \right|_0 E(\gamma_s) = 0$$

for any proper smooth variation  $\gamma_s$  of  $\gamma$  (here  $\gamma_s(t) = \Gamma(s, t)$ , where  $\Gamma : (-\varepsilon, \varepsilon) \times [t_0, t_1] \rightarrow \mathcal{M}$  is a smooth map, with  $\varepsilon > 0$ ,  $\Gamma(0, t) = \gamma(t)$  and  $\Gamma(s, t_i) = \gamma(t_i)$  for all  $s \in (-\varepsilon, \varepsilon)$ ,  $i = 0, 1$ ).

### 1.3 A Riemannian metric on the configuration space

Given  $g \in \mathcal{M}$  and  $X \in T_g \mathcal{M}$ , let us define the map  $\tilde{X} : \mathbb{S}^1 \rightarrow T\mathbb{S}^1$  by

$$\tilde{X}(q) = \left. \frac{d}{dt} \right|_0 \gamma(t)(q) \in T_{g(q)} \mathbb{S}^1, \quad (4)$$

where  $\gamma$  is any smooth curve in  $\mathcal{M}$  with  $\gamma(0) = g$  and  $\dot{\gamma}(0) = X$ . The map  $\tilde{X}$  is well-defined and smooth and it is a vector field on  $\mathbb{S}^1$  if and only if  $X \in T_e \mathcal{M}$ . Moreover,

$$X \mapsto \|X\|^2 = \frac{1}{2\pi} \int_{\mathbb{S}^1} \left| \tilde{X}(q) \right|^2 dm(q) \quad (5)$$

is a quadratic form on  $T_g \mathcal{M}$  and gives a Riemannian metric on  $\mathcal{M}$ .

**Remark 2** a) *The fundamental property of the metric (5) on  $\mathcal{M}$  is that a curve  $\gamma$  in  $\mathcal{M}$  is a geodesic if and only if (thought of as a Moebius motion) it is force free, since by (5) and (3),  $E_\gamma(t) = \pi \|\dot{\gamma}(t)\|^2$ .*

b) *The metric on  $\mathcal{M}$  is neither left nor right invariant, since we saw in [5] that it is not even complete.*

## 2 Force free Moebius motions of the circle

The next theorem describes completely the geometry of  $\mathcal{M}$  endowed with the metric (5) given by the kinetic energy. Recall from (1) that  $T_\alpha$  denotes the transvection associated with  $\alpha$  and that  $\Delta$  is the unit disc centered at zero in  $\mathbb{C}$ .

**Theorem 3** *Let  $ds^2$  be the metric on the disc  $\Delta$  given in polar coordinates  $(r, \theta)$  by*

$$ds^2 = \frac{2(dr^2 + r^2 d\theta^2)}{1 - r^2} \quad (6)$$

*and consider on  $\mathbb{S}^1 \times \Delta$  the Riemannian product metric, where  $\mathbb{S}^1$  has length  $2\pi$ . Then the map*

$$F : \mathbb{S}^1 \times \Delta \rightarrow \mathcal{M}, \quad F(u, \alpha) = uT_\alpha$$

*is an isometry.*

**Remark 4** a) *Note that the metric (6) on  $\Delta$  is not the canonical metric of constant negative curvature on  $\Delta$ . Indeed, the curvature function can be easily computed to be  $K(r, \theta) = -1/(1 - r^2)$ , in particular, it tends to  $-\infty$  as  $r \rightarrow 1^-$ . Also, the metric on  $\Delta$  is not complete, since the inextendible ray  $(0, 1) \ni r \mapsto T_r$  has length  $\pi/\sqrt{2}$ , since  $\left\|\frac{\partial}{\partial r}\right\|^2 = \frac{2}{1-r^2}$ .*

b) *In the higher dimensional situation [5] it is proven that the group  $\text{SO}(n)$  (with the metric induced from the one given by the kinetic energy) is totally geodesic in the group of directly conformal transformations of  $\mathbb{S}^n$ , but the author did not know whether this subgroup is a Riemannian factor, as it turned to be for  $n = 1$ . In the projective case [4],  $\text{SO}(n)$  is not even totally geodesic.*

**Proof of Theorem 3.** Let  $\mathbb{S}^1 \subset \mathcal{M}$  be the subgroup of isometries of the circle. The torus  $\mathbb{S}^1 \times \mathbb{S}^1$  acts on  $\mathcal{M}$  on the left by  $(u, v) \cdot g = ug\bar{v}$ , where  $(ug\bar{v})(z) = ug(z\bar{v})$  for any  $z \in \mathbb{S}^1$ . We know from the higher dimensional cases in [5] that this action is by isometries of  $\mathcal{M}$ , provided that this group is endowed with the metric (5).

We fix  $0 < r < 1$ . By the torus symmetry just described, it suffices to verify that  $dF_{(1,r)} : T_{(1,r)}(\mathbb{S}^1 \times \Delta) \rightarrow T_{F(1,r)}\mathcal{M}$  is a linear isometry. We put coordinates  $t \mapsto e^{it}$  on  $\mathbb{S}^1$  and  $(\rho, \theta) \mapsto \rho e^{i\theta}$  on  $\Delta$ . We denote  $\partial_x = \frac{d}{dx}$ . Let  $X, Y, Z$  be the images under

$dF_{(1,r)}$  of  $\partial_t, \partial_\rho, \partial_\theta$ , respectively. It suffices to show that  $\{X, Y, Z\}$  is an orthogonal basis of  $T_{F(1,r)}\mathcal{M}$  with

$$\|X\|^2 = 1, \quad \|Y\|^2 = \frac{2}{1-r^2}, \quad \|Z\|^2 = \frac{2r^2}{1-r^2}.$$

First, we compute  $\tilde{X}, \tilde{Y}$  and  $\tilde{Z}$  by their definition (4). In each case, we take the curve  $\gamma$  as the image under  $F$  of the coordinate curves in  $\mathbb{S}^1 \times \Delta$  through the point  $(1, r)$ . We have

$$\begin{aligned} \tilde{X}(z) &= \left. \frac{d}{dt} \right|_0 F(e^{it}, r)(z) = \left. \frac{d}{dt} \right|_0 e^{it} T_r(z) = i e^{it} T_r(z) \Big|_{t=0} = iT_r(z) = i \frac{z+r}{1+rz} \\ \tilde{Y}(z) &= \left. \frac{d}{d\rho} \right|_r F(1, \rho)(z) = \left. \frac{d}{d\rho} \right|_r T_\rho(z) = \left. \frac{d}{d\rho} \right|_r \frac{z+\rho}{1+\rho z} = \frac{1-z^2}{(1+rz)^2} \\ \tilde{Z}(z) &= \left. \frac{d}{d\theta} \right|_0 F(1, r e^{i\theta})(z) = \left. \frac{d}{d\theta} \right|_0 T_{r e^{i\theta}}(z) = \left. \frac{d}{d\theta} \right|_0 \frac{z+r e^{i\theta}}{1+r e^{-i\theta} z} = \frac{ri(1+2rz+z^2)}{(1+rz)^2}. \end{aligned}$$

Next we compute

$$2\pi \|X\|^2 = \int_{\mathbb{S}^1} |\tilde{X}(z)|^2 dm(z) = \int_{\mathbb{S}^1} |iT_r(z)|^2 dm(z) = \int_{\mathbb{S}^1} 1 dm(z) = 2\pi.$$

We have also

$$2\pi \|Y\|^2 = \int_{\mathbb{S}^1} |\tilde{Y}(z)|^2 dm(z) = \int_{\mathbb{S}^1} \left| \frac{1-z^2}{(1+rz)^2} \right|^2 dm(z).$$

Setting  $z = e^{is}$ , we have

$$2\pi \|Y\|^2 = \int_0^{2\pi} \frac{1}{ie^{is}} \left| \frac{1-e^{is^2}}{(1+re^{is})^2} \right|^2 ie^{is} ds = \int_{\mathbb{S}^1} \frac{1}{iz} \left| \frac{1-z^2}{(1+rz)^2} \right|^2 dz.$$

Now, the integrand is a complex analytic function inside the circle (observe that  $\bar{z} = 1/z$  for  $|z| = 1$ ), except for a simple pole at  $z = 0$  and a pole of order two at  $z = -r$ , with residues  $\frac{i}{r^2}$  and  $\frac{i(r^2+1)}{-r^2(1-r^2)}$ , respectively. One obtains that  $\|Y\|^2 = 2/(1-r^2)$ . In the same way one gets  $\|Z\|^2 = 2r^2/(1-r^2)$ .

We claim that the vectors  $X, Y, Z$  are pairwise orthogonal. Let  $h(U, V) = U\bar{V}$  denote the Hermitian inner product on  $\mathbb{C}$ . We compute

$$\int_{\mathbb{S}^1} h(\tilde{X}(z), \tilde{Y}(z)) dm(z) = \int_{\mathbb{S}^1} f(z) dz,$$

where  $f(z) = \frac{z^2-1}{z(1+rz)(z+r)}$  is a complex analytic function inside the circle, except for simple poles at  $z = 0$  and  $z = -r$ , with residues  $1/r$  and  $-1/r$ , respectively. Then,

$$\langle X, Y \rangle = \Re \int_{\mathbb{S}^1} h\left(\tilde{X}(z), \tilde{Y}(z)\right) dm(z) = 0.$$

Analogously, we find that  $\langle Y, Z \rangle = \langle X, Z \rangle = 0$ . □

**Corollary 5** *The force free Moebius motions of the circle, or equivalently, the geodesics of  $\mathcal{M}$ , are, via  $F$ , of the form  $\gamma = (\gamma_1, \gamma_2)$ , where  $\gamma_1$  parametrizes the circle with constant speed and  $\gamma_2$  is a geodesic in the disc  $\Delta$  whose trajectory coincides with the images of either  $c_1(\rho) = (\rho, \theta_0)$  or  $c_2(\theta) = (\rho(\theta), \theta)$ , where  $\rho$  satisfies the differential equation*

$$(\rho')^2 = \frac{\mu + \rho^2}{(1 - \rho^2)\rho^2} \tag{7}$$

for some constant  $\mu > -1$ .

Proof. Clearly, a geodesic of a Riemannian product projects to a geodesic in each factor. Besides, as the coefficients of the first fundamental form of  $\Delta$  depend only on  $\rho$ , the corresponding metric is Clairaut. Then, the trajectories of the geodesics of  $\Delta$  are, in polar form,

$$c_1(\rho) = (\rho, \theta_0) \quad \text{or} \quad c_2(\theta) = (\rho(\theta), \theta)$$

for some constant  $\theta_0$ , where  $\rho(\theta)$  satisfies Clairaut's differential equation, for some  $\lambda$ :

$$\lambda E^2(\rho) = E(\rho) + (\rho')^2 G(\rho).$$

Since in our case  $E(\rho) = \left\| \frac{\partial}{\partial r} \right\|^2 = \frac{2}{1-\rho^2}$  and  $G(\rho) = \left\| \frac{\partial}{\partial \theta} \right\|^2 = \frac{2\rho^2}{1-\rho^2}$ , the differential equation is equivalent to (7) for some constant  $\mu > -1$ . □

## References

- [1] Arnold V., *Mathematical methods of classical mechanics*, Graduate Texts in Math. 60, Springer, Berlin 1989.
- [2] Bates L. and Gibson P., *A geometry where everything is better than nice*, to appear in Proc. Amer. Math. Soc.
- [3] Emmanuele D. and Salvai M., *Force free Moebius motions of the circle*, J. Geom. Symmetry Phys. **27** (2012) 59–65

- [4] Lazarte M., Salvai M. and Will A., *Force free projective motions of the sphere*, J. Geom. Phys. **57** (2007) 2431–2436.
- [5] Salvai M., *Force free conformal motions of the sphere*, Diff. Geom. Appl. **16** (2002) 285–292.
- [6] Salvai, M., *A dynamical approach to compactify the three-dimensional Lorentz group*, J. Lie Theory **15** (2005) 335–339.
- [7] Villani, C., *Topics in optimal transportation*, Graduate Studies in Mathematics 58, American Mathematical Society (2003).

Daniela Emmanuele  
Departamento de Matemática, Escuela de Ciencias Exactas y Naturales,  
Facultad de Ciencias Exactas, Ingeniería y Agrimensura,  
Universidad Nacional de Rosario, Argentina  
emman@fceia.unr.edu.ar

Marcos Salvai  
CIEM - FaMAF,  
Conicet - Universidad Nacional de Córdoba, Argentina  
salvai@famaf.unc.edu.ar