# GEOMETRIC CONFIGURATION OF RIEMANNIAN SUBMANIFOLDS OF ARBITRARY CODIMENSION

## MOHAMED ABDELMALEK, MOHAMMED BENALILI, AND KAMIL NIEDZIAŁOMSKI

ABSTRACT. In this paper we study a geometric configuration of submanifolds of arbitrary codimension in an ambient Riemannian space. We obtain relations between the geometry of a q-codimension submanifold  $M^n$  along its boundary and the geometry of the boundary  $\Sigma^{n-1}$  of  $M^n$  as an hypersurface of a q-codimensional submanifold  $P^n$  in an ambient space  $\overline{M}^{n+q}$ . As a consequence of these geometric ralations we get that the ellipticity of the generalized Newton transformations implies the transversality of  $M^n$  and  $P^n$  in  $P^n$  is totally geodesic in  $\overline{M}^{n+q}$ .

#### 1. Introduction

Let  $\overline{M}^{n+q}$  be n+q-dimensional connected and orientable Riemannian manifold with metric  $\langle .,. \rangle$  and Levi-Civita connection  $\overline{\nabla}$ . Denote by  $P^n$  an oriented connected n- submanifold of  $\overline{M}^{n+q}$  and consider  $\Sigma^{n-1}$  an n-1-compact hypersurface of  $P^n$ . If  $\Psi: M^n \to \overline{M}^{n+q}$  is an oriented connected and compact submanifold of  $\overline{M}^{n+q}$  with boundary  $\partial M$ .  $M^n$  will be said submanifold of  $\overline{M}^{n+q}$  with boundary  $\Sigma^{n-1}$  if the restriction of  $\Psi$  to  $\partial M$  is a diffeomorphism onto  $\Sigma^{n-1}$ . A natural question would be: How can one describe the geometry of  $M^n$  along its boundary  $\partial M$  with respect to the geometry of the inclusion  $P^n \subset \overline{M}^{n+q}$ ? A partial answer to this question is given by the following formula, obtained in this paper, which holds along the boundary  $\partial M$ : for any multi-index  $u=(u_1,...,u_q)$  with length  $|u| \leq n-1$ ,

$$(1.1) \qquad \langle T_u \nu, \nu \rangle = \frac{1}{n-1-|u|} \sum_{l \le u} \binom{n-1-|l|}{|u|-l} \rho^l \mu^{u-l} \sigma_{|l|}(A_{\Sigma}).$$

where  $(T_u)_u$  stands for the family of the generalized Newton transformations introduced in [4] associated to the matrix  $A=(A_1,...,A_q);\ (A_\alpha)_{\alpha\in\{1,...,q\}}$  is a system of matrices of the shape operators corresponding to a normal basis to the manifold  $M^n$  and  $\tilde{\sigma}_u=\tilde{\sigma}_u(A_1|_{\Sigma},...,A_q|_{\Sigma})$  are the coefficients of the

<sup>2000</sup> Mathematics Subject Classification. 53A10, 53C42; Secondary 53A10.

Key words and phrases. Geometric Configuration, Generalized Newton Transformations, Ellipticity, Transversality.

Newton polynomial  $P_{\widetilde{A}}: \mathbb{R}^q \longrightarrow \mathbb{R}$  defined by

$$P_{\widetilde{A}}(t) = \sum_{|u| \le n-1} \widetilde{\sigma}_u t^u$$

where  $\widetilde{A} = (A_1|_{\Sigma},...,A_q|_{\Sigma})$ ,  $A_{\alpha}|_{\Sigma}$  is the restriction of  $A_{\alpha}$  to  $\Sigma^{n-1}$  and  $\sigma_r(A_{\Sigma})$  is the symmetric function coefficient of  $t^r$  in the characteristic polynomial of the matrix  $A_{\Sigma}$ .

In [2] Alías and Malacarne considered the above geometric configuration to the case of hypersurfaces where  $\Sigma_{n-1}$  is an (n-1)-dimensional compact submanifold contained in an hyperplane  $\Pi$  of  $\mathbb{R}^{n+1}$  and  $M^n$  stands for a smooth compact, connected and oriented manifold with boundary  $\partial M^n$ . Moreover  $M^n$  is an hypersurface of  $\mathbb{R}^{n+1}$  with boundary  $\Sigma_{n-1}$  in the sense that there exists  $\psi: M^n \to R^{n+1}$  an oriented hypersurface immersed in  $\mathbb{R}^{n+1}$  such that the restriction of  $\psi$  to the boundary  $\partial M^n$  is a diffeomorphism onto  $\Sigma_{n-1}$ . They showed that along the boundary  $\partial M^n$ , for every  $1 \le r \le n-1$ :

$$\langle T_r \nu, \nu \rangle = (-1)^r s_r \langle a, \nu \rangle^r$$

where  $\nu$  stands for the outward pointing unit conormal vector field along  $\partial M^n$  while  $T_r$  denotes the classical Newton transformation,  $a \in R^{n+1}$  such that  $\Pi = a^{\perp}$  and  $s_r$  is the r-th symmetric function of the principal curvatures of  $\Sigma_{n-1}$  with respect to  $\nu$ . In [3] Alías, de Lira and Malacarne studied the question in the context of an (n+1)-dimensional connected oriented ambient Riemannian manifold  $\overline{M}^{n+1}$ , they established that along the boundary  $\partial M^n$ , for every  $1 \leq r \leq n-1$ :

(1.3) 
$$\langle T_r \nu, \nu \rangle = (-1)^r s_r \langle \xi, \nu \rangle^r$$

where  $T_r$ ,  $\nu$ ,  $s_r$ , are as in relation /(1.2) where  $P^n \subset \overline{M}^{n+1}$  is an embedded totally geodesic submanifold instead of the hyperplane  $\Pi$  and  $\xi$  is a unitary normal vector field to  $P^n$ . Relation (1.3) shows that the ellipticity of the Newton transformation  $T_r$ , for some  $1 \leq r \leq n-1$  on  $M^n$ , implies the transversality of the hypersurfaces  $M^n$  and  $P^n$  along their boundary. Formula (1.3) was also obtained, in [1], by the two first authors in context of pseudo-Riemannian spaces. Moreover we deduce from relation (1.1) that the ellipticity of the generalized Newton transformation  $T_u$  implies the transversality of the q-codimension submanifolds  $M^n$  and  $P^n$  in case where  $P^n$  is totally geodesic submanifold of  $\overline{M}^{n+q}$ .

#### 2. Preliminaries

In this section, we will recall some properties of the generalized Newton transformations and we will show how our method works.

2.1. Generalized Newton Transformations. Let E be an n-dimensional real vector space and End(E) be the vector space of endomorphisms of E. Denote by  $\mathbb{N}$  the set of nonnegative integers and let  $\mathbb{N}^q$  be the one of multi- index  $u = (u_1, ..., u_q)$  with  $u_i \in \mathbb{N}$ . The length |u| of u is given by

 $|u| = u_1 + ... + u_q$ .  $End^q(E)$  stands for the vector space  $End(E) \times ... \times End(E)$  q-times. For  $A = (A_1, ..., A_q) \in End^q(E)$ ,  $t = (t_1, ..., t_q) \in R^q$  and  $u \in \mathbb{N}^q$ , we set

$$tA = t_1 A_1 + \dots + t_q A_q$$
  
$$t^u = t_1^{u_1} \dots t_q^{u_q}.$$

For  $\alpha \in \{1, ..., q\}$  we define ( see [4] ) the musical functions  $\alpha_{\flat} : \mathbb{N}^q \longrightarrow \mathbb{N}^q$  and  $\alpha^{\sharp} : \mathbb{N}^q \longrightarrow \mathbb{N}^q$  by

$$\alpha_{\flat}(i_1, \dots, i_q) = (i_1, \dots, i_{\alpha-1}, i_{\alpha} - 1, i_{\alpha+1}, \dots, i_q)$$

and

$$\alpha^{\sharp}(i_1, \dots, i_q) = (i_1, \dots, i_{\alpha-1}, i_{\alpha} + 1, i_{\alpha+1}, \dots, i_q)$$

It is clear that  $\alpha_b$  is the inverse map of  $\alpha^{\sharp}$ .

The generalized Newton transformation (GNT in brief) is a system of endomorphisms  $T_u = T_u(A)$ ,  $u \in \mathbb{N}^q$ , that satisfies the following recursive relations

$$T_0 = I$$
 where  $0 = (0, ..., 0),$  
$$T_u = \sigma_u I - \sum_{\alpha} A_{\alpha} T_{\alpha_{\flat}(u)}$$
 where  $|u| > 1$  
$$= \sigma_u I - \sum_{\alpha} T_{\alpha_{\flat}(u)} A_{\alpha}$$

where  $\sigma_u$  are the coefficients of the Newton polynomial  $P_A : \mathbb{R}^q \longrightarrow \mathbb{R}$  of A, given by

$$P_A(t) = \det(I + tA) = \sum_{|u| \le n} \sigma_u t^u$$

 $\sigma_u = \sigma_u(A_1, ..., A_q)$  depends only on  $A = (A_1, ..., A_q)$  and I is the identity map on E.

- 2.2. **The method.** We will describe how our method works.
- 2.2.1. Hypersurfaces' case. Let  $M^n$  be a n-submanifold of codimension one in  $\overline{M}^{n+1}$  of boundary  $\partial M$ . Assume the boundary  $\Sigma^{n-1} = \partial M$  is a codimension one in  $P^n \subset \overline{M}^{n+1}$ . Then we have the inclusions

$$\Sigma^{n-1}\subset M^n\subset \overline{M}^{n+1},\, \Sigma^{n-1}\subset P^n\subset \overline{M}^{n+1}.$$

Denote the corresponding shape operators, respectively, by

$$A_{\Sigma}$$
,  $A_{P}$ ,  $A_{\Sigma,P}$ ,  $A$ .

In our consideration we will need only  $A_{\Sigma}$ ,  $A_{P}$ , A. More precisely we will use

$$A_{\Sigma}$$
,  $A_P|_{\Sigma}$ ,  $A$ .

First two are represented by square matrices of dimension n-1 whereas the last one by a square matrix of dimension n. The intrinsic geometry of  $\Sigma^{n-1}$ 

in  $M^n$  is coded in the pair  $(A_{\Sigma}, A_P|_{\Sigma})$  and the geometry of  $M^n \subset \overline{M}^{n+1}$  is given by A. Therefore we will use the following Newton Transformation and the generalized Newton Transformations

$$T_{(k,l)} = T_{(k,l)}(A_{\Sigma}, A_P|_{\Sigma})$$
 and  $T_r = T_r(A)$ 

and corresponding symmetric functions

$$\sigma_{(k,l)} = \sigma_{(k,l)}(A_{\Sigma}, A_P|_{\Sigma})$$
 and  $\sigma_r = \sigma_r(A)$ .

The goal is to show that

(2.1) 
$$\langle T_r \nu, \nu \rangle = \sum_{k+l=r} \sigma_{(k,l)}$$

where  $\nu$  is the unit normal vector to  $\Sigma^{n-1}$  in  $M^n$ .

The only geometric considerations involved are the ones which lead to the formulas

$$\langle N, \nu \rangle = \langle \xi, N \rangle, \ \langle \eta, N \rangle = - \langle \xi, \nu \rangle$$

and

$$\langle Ae_i, e_j \rangle = -\langle A_{\Sigma}e_i, e_j \rangle \langle \xi, \nu \rangle + \langle A_Pe_i, e_j \rangle \langle \xi, N \rangle$$

where N is unit normal vector with the respect to inclusion  $M^n \subset \overline{M}^{n+1}$ ,  $\xi$  unit normal vector with respect to  $P^n \subset \overline{M}^{n+1}$  and  $\eta$  is the unit normal vector of  $\Sigma^{n-1} \subset P^n$  and  $(e_1, ..., e_{n-1})$  is a local orthonormal basis of  $T\Sigma^{n-1}$ , we may assume that this basis consists of eigenvectors of  $A_{\Sigma}$  i.e.  $A_{\Sigma}e_i = \tau_i e_i$ . In other words

$$A|_{\Sigma} = -\langle \xi, \nu \rangle A_{\Sigma} + \langle \xi, N \rangle A_{P}|_{\Sigma}.$$

Assuming P is totally umbilical in  $\overline{M}^{n+1}$ , we have  $A_P|_{\Sigma} = \lambda I_{T\Sigma^{n-1}}$ . Hence

(2.2) 
$$A|_{\Sigma} = -\langle \xi, \nu \rangle A_{\Sigma} + \lambda \langle \xi, N \rangle I_{T\Sigma^{n-1}}.$$

Denote by  $\widetilde{A}$  the matrix of A with the respect of the basis  $(e_1, ..., e_{n-1}, \nu)$  and by A the matrix of  $A|_{\Sigma}$ . Then

$$\widetilde{A} = \begin{pmatrix} A & B \\ B^{\mathsf{T}} & c \end{pmatrix}$$
, where  $B = \begin{pmatrix} \langle A\nu, e_1 \rangle \\ \vdots \\ \langle A\nu, e_{n-1} \rangle \end{pmatrix}$  and  $c = \langle A\nu, \nu \rangle$ .

Let us compare symmetric functions of  $\widehat{A}$  with symmetric functions of A. We have

$$P_{\widetilde{A}}(t) = \det \begin{pmatrix} I_{n-1} + tA & tB \\ tB^{\mathsf{T}} & 1 + tc \end{pmatrix} = (1 + tc - t^2B^{\mathsf{T}}(I_{n-1} + tA)^{-1}B)\det(I_{n-1} + tA)$$
$$= f(t)P_A(t),$$

where  $f(t) = 1 + tc - t^2 B^{\intercal} (I_{n-1} + tA)^{-1} B$  . Recall that

$$P_{\widetilde{A}}(t) = \sum_{j=0}^{n} (-1)^{j} \sigma_{j}(\widetilde{A}) t^{j}.$$

Hence

$$(-1)^{r} r! \sigma_{r}(\widetilde{A}) = \frac{d^{r}}{dt^{r}} P_{\widetilde{A}}(0) = \sum_{j=0}^{r} {r \choose j} \frac{d^{r-j}}{dt^{r-j}} P_{A}^{(r-j)}(0) f^{(j)}(0)$$
$$= \sum_{j=0}^{r} {r \choose j} (-1)^{r-j} (r-j)! \sigma_{r-j}(A) f^{(j)}(0).$$

It is not hard to see that

$$f(0) = 1, f'(0) = -c$$
 and  $f^{(j)}(0) = -j!B^{\mathsf{T}}A^{j-2}B$  for  $j \ge 2$ .

Therefore

(2.3) 
$$\sigma_r(\widetilde{A}) = \sigma_r(A) + c\sigma_{r-1}(A) - \sum_{j=2}^r (-1)^j (B^{\mathsf{T}} A^{j-2} B) \sigma_{r-j}(A).$$

Let us now move to symmetric functions of two matrices. Notice first that

$$\sigma_r(\widetilde{A} + \lambda I_n) = \sum_{j=0}^r \begin{pmatrix} n-j \\ r-j \end{pmatrix} \lambda^{r-j} \sigma_j(\widetilde{A}).$$

Indeed,  $P_{\widetilde{A}+\lambda I_n}(t) = (1+t(a_1+\lambda)...(1+t(a_n+\lambda))$  if  $a_1,...,a_n$  are the eigenvalues of  $\widetilde{A}$ . Notice moreover that

$$P_{\widetilde{A},\lambda I_n}(t) = \det\left(I_n + t\widetilde{A} + s\lambda I_n\right)$$
$$= (1 + ta_1 + s\lambda) \dots (1 + ta_n + s\lambda).$$

Thus

(2.4) 
$$\sigma_{(k,l)}(\widetilde{A},\lambda I_n) = \begin{pmatrix} n-k \\ l \end{pmatrix} \lambda^l \sigma_k(\widetilde{A}).$$

Hence

(2.5) 
$$\sigma_r(\widetilde{A} + \lambda I_n) = \sum_{j=0}^r \sigma_{(j,r-j)}(\widetilde{A}, \lambda I_n)$$

We need to show, by (2.5) and (2.1), that

(2.6) 
$$\langle T_r \nu, \nu \rangle = \widetilde{\sigma}_r (A) .$$

By the recurrence formula for  $T_r$  we have

$$\langle T_r \nu, \nu \rangle = \sigma_r \left( \widetilde{A} \right) - \left\langle T_{r-1} \nu, \widetilde{A} \nu \right\rangle$$
$$= \sigma_r \left( \widetilde{A} \right) - c \left\langle T_{r-1} \nu, \nu \right\rangle - \sum_{i=1}^{n-1} \left\langle T_{r-1} \nu, e_i \right\rangle \left\langle \widetilde{A} \nu, e_i \right\rangle.$$

We will show that

(2.7) 
$$\sum_{i=1}^{n-1} \langle T_k \nu, e_i \rangle \left\langle \widetilde{A} \nu, e_i \right\rangle = -\sum_{j=2}^k (-1)^j \left( B^{\mathsf{T}} A^{j-2} B \right) \sigma_{k-j}(A).$$

Indeed, assuming  $(e_i)$  is a basis consisting of eigenvectors with eigenvalues  $(\tau_i)_i$  by induction we have

$$\begin{split} \sum_{i=1}^{n-1} \left\langle T_k \nu, e_i \right\rangle \left\langle \widetilde{A} \nu, e_i \right\rangle &= -b_i^2 \sum_{j=2}^k \left( -j \right) \sigma_{k-j} \left( A \right) \tau_i^{j-2} \\ &= \sum_{j=2}^k (-1)^j \left( b_i \tau_i^{j-2} b_i \right) \sigma_{k-j} \left( A \right) \end{split}$$

where  $B^{\mathsf{T}} = (b_i)_i$ , which proves (2.7). Thus by induction and (2.3), we get

$$\langle T_r \nu, \nu \rangle = \sigma_r(\widetilde{A}) - c\sigma_{r-1}(A) + \sum_{j=2}^r (-1)^j \left( B^{\mathsf{T}} A^{j-2} B \right) \sigma_{r-j}(A) = \widetilde{\sigma}_r(A).$$

2.2.2. General case. We generalize considerations to any codimension q. We generalize equations (2.1) and (2.6). Let  $P^n$  and  $M^n$  be of codimension q in  $\overline{M}^{n+q}$ . We assume that  $P^n$  is totally umbilical in  $\overline{M}^{n+1}$ . As before  $\Sigma^{n-1} \subset P^n$  is a boundary of  $M^n$ . Then we have the following shape operators

$$A_{\Sigma}, A_{P}^{\xi_{1}}, ..., A_{P}^{\xi_{p}}, A^{N_{1}}, ..., A^{N_{q}},$$

corresponding to inclusions  $\Sigma^{n-1} \subset P^n$ ,  $P^n \subset \overline{M}^{n+q}$ , and  $M^n \subset \overline{M}^{n+q}$ , where  $(\xi_1, ..., \xi_q)$  are orthogonal to  $P^n$  and  $(N_1, ..., N_q)$  are orthogonal to  $M^n$ . Let

$$T_u = T_u (A^{N_1}, ..., A^{N_q}), \quad T_v = T_v (A_{\Sigma}, A^{\xi_1}|_{\Sigma}, ..., A^{\xi_q}|_{\Sigma})$$

and

$$\widetilde{T_u} = \widetilde{T}_u \left( A^{N_1}|_{\Sigma}, ..., A^{N_q}|_{\Sigma} \right)$$

where u is of length q and v of length q+1. We hope that the following relations hold

$$\langle T_u \nu; \nu \rangle = \widetilde{\sigma}_u$$

and

$$\widetilde{\sigma}_u = \sum_{|v|=|u|, \ v \text{ not increasing}} c_v \sigma_v$$

where  $c_v$  is a coefficient independent of  $(A^{N_1}|_{\Sigma},...,A^{N_q}|_{\Sigma})$ . Both of them imply

$$\langle T_u \nu; \nu \rangle = \sum_{|v|=|u|, \ v \text{ not increasing}} c_v \sigma_v.$$

The first equality is a generalization of (2.6), whereas the second one of (2.5). Moreover, the first one is purely algebraic and the second one uses correspondence between  $A_{\Sigma}$ ,  $A^{\xi_1}|_{\Sigma}$ ,...,  $A^{\xi_q}|_{\Sigma}$  and  $A^{N_1}|_{\Sigma}$ ,...,  $A^{N_q}|_{\Sigma}$  analogous to the relation (2.2) in the codimension one case.

Let us be more precise. First notice that

$$\overline{\nabla}_{e_i} e_j = \sum_{k=1}^{n-1} \langle \overline{\nabla}_{e_i} e_j, e_k \rangle e_k + \langle \overline{\nabla}_{e_i} e_j, \nu \rangle \nu + \sum_{\alpha=1}^q \langle \overline{\nabla}_{e_i} e_j, N_\alpha \rangle N_\alpha$$

$$= \sum_{k=1}^{n-1} \langle \overline{\nabla}_{e_i} e_j, e_k \rangle e_k + \langle \overline{\nabla}_{e_i} e_j, \nu \rangle \nu + \sum_{\alpha=1}^q \langle A^{N_\alpha} e_i, e_j \rangle N_\alpha$$

and

$$\overline{\nabla}_{e_i} e_j = \sum_{k=1}^{n-1} \langle \overline{\nabla}_{e_i} e_j, e_k \rangle e_k + \langle \overline{\nabla}_{e_i} e_j, \eta \rangle \eta + \sum_{\alpha=1}^q \langle \overline{\nabla}_{e_i} e_j, \xi_\alpha \rangle \xi_\alpha$$

$$= \sum_{k=1}^{n-1} \langle \overline{\nabla}_{e_i} e_j, e_k \rangle e_k + \langle A_{\Sigma} e_i, e_j \rangle \eta + \sum_{\alpha=1}^q \langle A^{\xi_\alpha} e_i, e_j \rangle \xi_\alpha$$

Thus

$$\langle \overline{\nabla}_{e_i} e_j, \nu \rangle \nu + \sum_{\alpha=1}^q \langle A^{N_\alpha} e_i, e_j \rangle N_\alpha = \langle A_\Sigma e_i, e_j \rangle \eta + \sum_{\alpha=1}^q \langle A^{\xi_\alpha} e_i, e_j \rangle \xi_\alpha$$

Hence

$$\begin{split} \left\langle A^{N_{\alpha}}e_{i},e_{j}\right\rangle &=\left\langle \eta,N_{\alpha}\right\rangle \left\langle A_{\Sigma}(e_{i}),e_{j}\right\rangle +\sum_{\beta=1}^{q}\left\langle \xi_{\beta},N_{\alpha}\right\rangle \left\langle \left(A^{\xi_{\beta}}\right)e_{i},e_{j}\right\rangle \\ &=\left\langle \eta,N_{\alpha}\right\rangle \left\langle A_{\Sigma}(e_{i}),e_{j}\right\rangle +\sum_{\beta=1}^{q}\left\langle \xi_{\beta},N_{\alpha}\right\rangle \left\langle A^{\xi_{\beta}}e_{i},e_{j}\right\rangle \end{split}$$

Assuming that  $P^n$  is totally umbilical, i.e.  $A^{\xi_{\alpha}} = \lambda_{\alpha} I_n$ , where  $I_n$  denotes the identity map of the tangent space  $T_p P^n$ , we get

$$A^{N_{\alpha}}|_{\Sigma} = \langle \eta, N_{\alpha} \rangle A_{\Sigma} + \sum_{\beta=1}^{q} \langle \xi_{\beta}, N_{\alpha} \rangle \lambda_{\beta} I_{n-1}.$$

Notice that it can be written in the form

(2.9) 
$$A^{N_{\alpha}}|_{\Sigma} = \langle \eta, N_{\alpha} \rangle A_{\Sigma} + \langle V, N_{\alpha} \rangle I_{n-1}, \text{ where } V = \sum_{\beta=1}^{q} \lambda_{\beta} \xi_{\beta}.$$

Moreover one can show that

$$\langle \eta, V \rangle = \det \left( \langle \xi_{\alpha}, N_{\beta} \rangle \right)_{\alpha, \beta} \text{ and } \langle \eta, N_{\alpha} \rangle = -\det C_{\alpha}$$

where  $C_{\alpha}$  is a matrix obtained from  $C=(\langle \xi_{\alpha}, N_{\beta} \rangle)_{\alpha,\beta}$  by replacing the  $\alpha$ -th column by  $\langle \xi_{\alpha}, \nu \rangle$ .

Hence we have

$$\widetilde{T}_{u} = \widetilde{T}_{u} \left( \rho_{1} A_{\Sigma} + \mu_{1} I_{n-1}, ..., \rho_{q} A_{\Sigma} + \mu_{q} I_{n-1} \right), \text{ where } \quad \rho_{\alpha} = \left\langle \eta, N_{\alpha} \right\rangle, \, \mu_{\alpha} = \left\langle V, N_{\alpha} \right\rangle.$$

Now if  $\{e_1, \ldots, e_{n-1}\}$  is a basis of eigenvectors of the shape operator  $A_{\Sigma}$  then, for every  $i \in \{1, \ldots, n-1\}$ ,

$$A_{\Sigma}(e_i) = \tau_i e_i.$$

where  $\tau_i$  are the corresponding eigenvalues.

By relation (2.9) we get

$$\langle A_{\alpha}(e_i), e_j \rangle = (\langle \eta, N_{\alpha} \rangle \tau_i + \langle V, N_{\alpha} \rangle) \, \delta_i^j$$
$$= \gamma_{i,\alpha} \delta_i^j,$$

where

$$\gamma_{i,\alpha} = \langle \eta, N_{\alpha} \rangle \, \tau_i + \langle V, N_{\alpha} \rangle \, .$$

Thus the matrix associated to the shape operator  $A_{\alpha}$  with respect to the basis  $\{e_1, \dots, e_{n-1}, \nu\}$  is given by

$$A_{\alpha} = \begin{pmatrix} \gamma_{1,\alpha} & 0 & \dots & 0 & \langle A_{\alpha}\nu, e_{1} \rangle \\ 0 & \gamma_{2,\alpha} & \dots & 0 & \vdots \\ \vdots & \vdots & \ddots & \vdots & \vdots \\ 0 & \dots & \ddots & \gamma_{n-1,\alpha} & \langle A_{\alpha}\nu, e_{n-1} \rangle \\ \langle A_{\alpha}\nu, e_{1} \rangle & \dots & \dots & \langle A_{\alpha}\nu, e_{n-1} \rangle & \langle A_{\alpha}\nu, \nu \rangle \end{pmatrix}.$$

#### 3. Algebraic Formulas

Let us prove the algebraic relation (2.8), to do so, we put

$$\rho_{\alpha} = \langle \eta, N_{\alpha} \rangle, \quad \mu_{\alpha} = \langle V, N_{\alpha} \rangle.$$

Moreover, we will write  $v \leq u$  for multi-indices  $v, u \in \mathbb{N}^q$  if the difference  $u - v \in \mathbb{N}^q$ . For a multi-index  $u = (u_1, ..., u_q) \in \mathbb{N}^q$  of length |u| = n, we set

$$\binom{n}{u} = \frac{n!}{u!} = \frac{n!}{u_1! \dots u_q!}.$$

**Proposition 1.** Let  $\tilde{A} = (A_1|_{\Sigma}, ..., A_q|_{\Sigma}) = (\rho_1 A_{\Sigma} + \mu_1 I, ..., \rho_q A_{\Sigma} + \mu_q I)$ . Put  $\tilde{\sigma}_u = \sigma_u(\tilde{A})$ . Then for every multi-index  $u \in \mathbb{N}^q$  we have

$$(3.1) \qquad \tilde{\sigma}_{u} = \frac{1}{(n-1-|u|)!} \sum_{l \leq u} \left( \begin{array}{c} |l| \\ l \end{array} \right) \rho^{l} \mu^{u-l} \left( \begin{array}{c} n-1-|l| \\ u-l \end{array} \right) \sigma_{|l|} \left( A_{\Sigma} \right).$$

*Proof.* First, by applying the following relation (see [6])

$$\sigma_u(aA_1,\ldots,A_q)=a^{u_1}\sigma_u(A_1,\ldots,A_q)$$

we obtain

(3.2)  $\tilde{\sigma}_u(\rho_1 A_{\Sigma} + \mu_1 I, \dots, \rho_q A_{\Sigma} + \mu_q I) = \rho^u \hat{\sigma}_u(A_{\Sigma} + \theta_1 I, \dots, A_{\Sigma} + \theta_q I)$  where

$$\theta_{\alpha} = \frac{\mu_{\alpha}}{\rho_{\alpha}}$$
 and  $\rho^{u} = \rho_{1}^{u_{1}}...\rho_{q}^{u_{q}}$ .

Consider the GNT  $T_u = T_u(\hat{A})$ , with

$$\hat{A} = (A_{\Sigma} + \theta_1 I, \dots, A_{\Sigma} + \theta_q I).$$

The characteristic polynomial associate to  $\hat{A}$  is then

$$P_{\hat{A}}(t) = \sum_{|u| \le n-1} \hat{\sigma}_u t^u$$

Moreover, we have

(3.3) 
$$P_{\hat{A}}(t) = \det \left( I + \sum_{\alpha=1}^{q} t_{\alpha} \left( A_{\Sigma} + \theta_{\alpha} \right) I \right)$$

(3.4) 
$$= \prod_{j=1}^{n-1} (1 + \tau_j (t_1 + ... + t_q) + t_1 \theta_1 + ... + t_q \theta_q).$$

By expanding the linear factorization of a monic polynomial we obtain

$$P_{\hat{A}}(t) = \sum_{j=0}^{n-1} (1 + t_1 \theta_1 + \dots + t_q \theta_q)^{n-1-j} (t_1 + \dots + t_q)^j \sigma_j (A_{\Sigma})$$

and by the multinomial theorem we get

$$P_{\hat{A}}(t) = \sum_{j=0}^{n-1} \sum_{k=0}^{n-1-j} \sum_{|v|=k} \sum_{|l|=j} \frac{1}{(n-1-j-k)!} \binom{j}{l} \binom{n-1-j}{v} \sigma_j(A_{\Sigma}) \theta^v t^{v+l}.$$

where  $l=(l_1,...,l_q),\ v=(v_1,...,v_q),\ \theta=(\theta_1,...,\theta_q)$  and  $t=(t_1,...,t_q).$  Let u=v+l so j=|u|-|v| and

$$P_{\hat{A}}(t) = \sum_{|u| \le n-1} \sum_{l \le u} \frac{1}{(n-1-|u|)!} \binom{|l|}{l} \binom{n-1-|l|}{u-l} \theta^{u-l} \sigma_{|l|} (A_{\Sigma}) t^{u}.$$

Thus the coefficient before  $t^u$  is given by

$$\hat{\sigma}_u(A_{\Sigma}+ heta_1I,\ldots,A_{\Sigma}+ heta_qI)=rac{1}{(n-1-|u|)!}\sum_{l\leq u}\left(egin{array}{c}|l|\line l\end{array}
ight) heta^{u-l}\left(egin{array}{c}n-1-|l|\u-l\end{array}
ight)\sigma_{|l|}\left(A_{\Sigma}
ight).$$

Consequently

$$\tilde{\sigma}_u(\rho_1 A_{\Sigma} + \mu_1 I, \dots, \rho_q A_{\Sigma} + \mu_q I) =$$

$$(3.5) \qquad \frac{1}{(n-1-|u|)!} \sum_{l \leq u} \begin{pmatrix} |l| \\ l \end{pmatrix} \rho^{l} \mu^{u-l} \begin{pmatrix} n-1-|l| \\ u-l \end{pmatrix} \sigma_{|l|} \left(A_{\Sigma}\right)$$

On the other hand

Denote by  $\bar{A} = (A_{\Sigma}, \mu_1 I, \dots, \mu_q I)$  and  $\bar{\sigma}_{(k,v)} = \sigma_{(k,v)}(\bar{A})$ . We have

$$\sigma_{(k,v)}(\bar{A}) = \mu^v \sigma_{(k,v)}(A_{\Sigma}, I, \dots, I).$$

Now, we use the following formula (see [6])

$$\sigma(v,k)(B_1,\ldots,B_q,I) = \binom{n-1-|v|}{k}\sigma_v(B_1,\ldots,B_q)$$

consecutively q-times and we get

$$\sigma_{(j,v)}(A_{\Sigma}, I, \dots, I) = \binom{n-1-|v|+v_1}{v_1} \dots \binom{n-1-|v|+|v|}{v_q} \sigma_j(A_{\Sigma})$$
$$= \frac{1}{(n-1-|v|-j)!} \binom{n-1-j}{v} \sigma_j(A_{\Sigma}).$$

Taking j = |u| - |v|, we obtain

$$= \frac{1}{(n-1-|u|)!} \binom{n-1-|u|+|v|}{v} \sigma_{|u|-|v|}(A_{\Sigma})$$

and if we let l = u - v, we have

(3.6) 
$$\sigma_{(|l|,u-l)}(A_{\Sigma},I,\ldots,I) = \frac{1}{(n-1-|u|)!} \binom{n-1-|l|}{u-l} \sigma_{|l|}(A_{\Sigma})$$

So by relations (3.2) and (3.6), we get

(3.7) 
$$\tilde{\sigma}_{u} = \frac{1}{(n-1-|u|)!} \sum_{l \leq u} \begin{pmatrix} |l| \\ l \end{pmatrix} \rho^{l} \mu^{u-l} \begin{pmatrix} n-1-|l| \\ u-l \end{pmatrix} \sigma_{|l|} (A_{\Sigma})$$

Notice, that in the proof of the above Proposition, we assumed  $\rho_{\alpha} \neq 0$  for all  $\alpha$ , this allowed us to defined the constants  $\theta_{\alpha}$ . The assumption  $\rho_{\alpha} \neq 0$  is not necessary. Since if there exists  $\alpha \in \{1,..,q\}$  such that  $\rho_{\alpha} = 0$ . then (see [6])

$$\begin{split} \tilde{\sigma}_{u} &= \tilde{\sigma}_{u}(\rho_{1}A_{\Sigma} + \mu_{1}I, \dots, \rho_{q}A_{\Sigma} + \mu_{q}I) \\ &= \tilde{\sigma}_{u}(\rho_{1}A_{\Sigma} + \mu_{1}I, \dots, \rho_{i-1}A_{\Sigma} + \mu_{i-i}I, \mu_{i}I, \rho_{i+1}A_{\Sigma} + \mu_{i+1}I, \dots, \rho_{q}A_{\Sigma} + \mu_{q}I) \\ &= \mu_{i}^{u_{i}} \tilde{\sigma}_{\tilde{u}}(\rho_{1}A_{\Sigma} + \mu_{1}I, \dots, \rho_{i-1}A_{\Sigma} + \mu_{i-i}I, \rho_{i+1}A_{\Sigma} + \mu_{i+1}I, \dots, \rho_{q}A_{\Sigma} + \mu_{q}I), \end{split}$$

where

$$\tilde{u} = (u_1, \dots, u_{i-1}, u_{i+1}, \dots, u_q)$$

and we may apply the above Proposition to  $\tilde{\sigma}_{\tilde{u}}$ .

**Proposition 2.** Let  $\tilde{A} = (A_1|_{\Sigma}, ..., A_q|_{\Sigma})$ , where  $A_{\alpha}|_{\Sigma} = (\rho_{\alpha}A_{\Sigma} + \mu_{\alpha}I, and \overline{A} = (A_{\Sigma}, \mu_1 I, ..., \mu_q I)$ . For any multi-index  $u \in \mathbb{N}^q$ , put  $\tilde{\sigma}_u = \sigma_u(\tilde{A})$  and  $\overline{\sigma}_u = \sigma_u(\overline{A})$ . Then we have

(3.8) 
$$\tilde{\sigma}_u = \sum_{l < u} \begin{pmatrix} |l| \\ l \end{pmatrix} \rho^l \overline{\sigma}_{(|l|, u - l)}.$$

*Proof.* First notice that

$$\overline{\sigma}_{(j,v)} = \sigma_{(j,v)}(\overline{A}) = \mu^v \sigma_{(j,v)}(A_{\Sigma}, I, \dots, I).$$

Using the following formula (see [6])

$$\sigma(v,j)(B_1,\ldots,B_q,I) = \binom{n-1-|v|}{j}\sigma_v(B_1,\ldots,B_q)$$

consecutively q -times, we obtain

$$\sigma_{(j,v)}(A_{\Sigma}, I, \dots, I) = \binom{n-1-|v|+v_1}{v_1} \dots \binom{n-1-|v|+|v|}{v_q} \sigma_j(A_{\Sigma})$$
$$= \frac{1}{(n-1-|v|-j)!} \binom{n-1-j}{v} \sigma_j(A_{\Sigma})$$

or by putting l = u - v

$$\sigma_{(|l|,u-l)} = \frac{1}{(n-1-|u|)!} \begin{pmatrix} n-1-|l| \\ u-l \end{pmatrix} \sigma_{|l|}(A_{\Sigma})$$

so

$$\overline{\sigma}_{(|l|,u-l)} = \frac{\mu^{u-l}}{(n-1-|u|)!} \left( \begin{array}{c} n-1-|l| \\ u-l \end{array} \right) \sigma_{|l|}(A_{\Sigma})$$

Thus by (3.1) we get (3.8).

Let us now state the relation between symmetric functions corresponding to the families  $(A_{\alpha})$  and  $(A_{\alpha}|\Sigma)$ . Notice that there are of different sizes.

**Proposition 3.** Let  $A = (A_1, ..., A_q)$  and  $\tilde{A} = (A_1|_{\Sigma}, ..., A_q|_{\Sigma})$ . Put  $\sigma_u = \sigma_u(A)$  and  $\tilde{\sigma}_u = \sigma_u(\tilde{A})$ . Then

$$\sigma_{u} = \tilde{\sigma}_{u} + \sum_{\alpha} C_{\alpha} \tilde{\sigma}_{\alpha_{\flat}(v)}$$

$$+ \sum_{\alpha^{\sharp} \beta^{\sharp}(0) \leq w \leq u} \sum_{\alpha, \beta} (-1)^{|w| - |v| + 1} \binom{|u| - |w|}{u - w} B_{\alpha}^{\top} \tilde{A}^{u - w} B_{\beta} \tilde{\sigma}_{\alpha_{\flat} \beta_{\flat}(w)}$$

where  $B_{\alpha}^{\top} = (\langle A_{\alpha}v, e_1 \rangle, ..., \langle A_{\alpha}v, e_{n-1} \rangle)$  and  $C_{\alpha} = \langle A_{\alpha}v, v \rangle$ .

*Proof.* First calculate the characteristic polynomial  $P_A(t)$  of  $A = (A_1, ..., A_q)$ . By definition, we have

$$P_A(t) = \det(I + \sum_{\alpha} t_{\alpha} A_{\alpha})$$

$$= \det\left(I_{n-1} + \sum_{\alpha} t_{\alpha} A_{\alpha}|_{\Sigma}\right) \det\left(C - B^{\top} \left(I_{n-1} + \sum_{\alpha} t_{\alpha} A_{\alpha}|_{\Sigma}\right)^{-1} B\right)$$

where  $B^{\top}$  and C are respectively given by

$$B^{\top} = \left(\sum_{\alpha} t_{\alpha} \langle A_{\alpha} v, e_{1} \rangle, ..., \sum_{\alpha} t_{\alpha} \langle A_{\alpha} v, e_{n-1} \rangle\right) = \sum_{\alpha} t_{\alpha} B_{\alpha}^{\top}$$

and

$$C = 1 + \sum_{\alpha} t_{\alpha} \langle A_{\alpha} v, v \rangle.$$

To simplify the expressions, we put

$$M = \left(I_{n-1} + \sum_{\alpha} t_{\alpha} A_{\alpha}|_{\Sigma}\right).$$

and

$$f(t) = \det(C - B^{\top} M^{-1} B).$$

Moreover it is clear that

$$B^{\top} M^{-1} B = \sum_{\alpha,\beta} t_{\alpha} t_{\beta} B_{\alpha}^{\top} M^{-1} B_{\beta} ,$$

therefore we get

$$f(t) = \left(1 + \sum_{\alpha} t_{\alpha} C_{\alpha} - \sum_{\alpha, \beta} t_{\alpha} t_{\beta} B_{\alpha}^{\top} M^{-1} B_{\beta}\right).$$

As it is easily seen that

$$P_A(t) = f(t)P_{\tilde{A}}(t).$$

where  $P_A(t)$  and  $P_{\tilde{A}}(t)$  are defined by

$$P_A(t) = \sum_{|u| \le n} \sigma_u t^u$$

and

$$P_{\tilde{A}}(t) = \sum_{|u| \le n-1} \tilde{\sigma}_u t^u.$$

Thus we get

$$\frac{\partial^u}{\partial t^u} P_A(t)\mid_{t=0} = u! \sigma_u, \quad \frac{\partial^u}{\partial t^u} P_{\tilde{A}}(t)\mid_{t=0} = u! \tilde{\sigma}_u.$$

Similarly we obtain

$$\frac{\partial^u}{\partial t^u} \left( P_{\tilde{A}}(t).f(t) \right) \mid_{t=0} = \sum_{v \leq u} \binom{u}{v} \left( \frac{\partial^v}{\partial t^v} P_{\tilde{A}}(t).\frac{\partial^{u-v}}{\partial t^{u-v}} f(t) \right) \mid_{t=0},$$

where

$$\binom{u}{v} = \frac{u!}{v!} = \frac{u_1! \dots u_1!}{v_1! \dots v_q!}.$$

Now we will compute  $\frac{\partial^{u-v}}{\partial t^{u-v}} f(t)$ . For v=u, it is checked

$$\frac{\partial^{u-v}}{\partial t^{u-v}} f(t) \mid_{t=0} = f(0, ..., 0) = 1.$$

If  $v = \alpha_{\flat}(u)$ , we have

$$\frac{\partial^{u-v}}{\partial t^{u-v}}f(t)\mid_{t=0}=\frac{\partial}{\partial t_{\alpha}}f(t)\mid_{t=0}=C_{\alpha}$$

and for any  $v \leq \alpha_{\flat}\beta_{\flat}(u)$ , we obtain

$$\frac{\partial^{u-v}}{\partial t^{u-v}} f(t) \mid_{t=0} = \frac{\partial^{u-v}}{\partial t^{u-v}} \left( 1 + \sum_{\alpha} t_{\alpha} C_{\alpha} - \sum_{\alpha,\beta} t_{\alpha} t_{\beta} B_{\alpha}^{\top} \left( I_{n-1} + \sum_{\alpha} t_{\alpha} A_{\alpha} \mid_{\Sigma} \right)^{-1} B_{\beta} \right) \mid_{t=0}.$$

Taking  $|t| < \varepsilon$  for some small enough  $\varepsilon$  we get

$$\left\| \sum_{\alpha} t_{\alpha} A_{\alpha} |_{\Sigma} \right\| \le 1.$$

Thus

$$\left(I_{n-1} + \sum_{\alpha} t_{\alpha} A_{\alpha}|_{\Sigma}\right)^{-1} = \sum_{k=0}^{\infty} (-1)^{k} \left(\sum_{\alpha} t_{\alpha} A_{\alpha}|_{\Sigma}\right)^{k}$$

Since the matrices  $A_{\alpha}|_{\Sigma}$  are diagonal, then they commute. Therefore, we may write

$$\left(\sum_{\alpha} t_{\alpha} A_{\alpha}|_{\Sigma}\right)^{k} = \sum_{w_{1} + \dots + w_{n} = k} {k \choose w} t^{w} \tilde{A}^{w}.$$

Hence

$$\left(I_{n-1} + \sum_{\alpha} t_{\alpha} A_{\alpha}|_{\Sigma}\right)^{-1} = \sum_{k=0}^{\infty} (-1)^k \left(\sum_{w_1 + \dots + w_q = k} {k \choose w} t^w \tilde{A}^w\right)$$

Which gives

$$\frac{\partial^{u-v}}{\partial t^{u-v}} f(t) \mid_{t=0} = \frac{\partial^{u-v}}{\partial t^{u-v}} \left( -\sum_{\alpha,\beta} t_{\alpha} t_{\beta} B_{\alpha}^{\top} \sum_{k=0}^{\infty} (-1)^{k} \left( \sum_{w_{1}+\ldots+w_{q}=\mid w\mid=k} \binom{\mid w\mid}{w} t^{w} \tilde{A}^{w} \right) B_{\beta} \right) \mid_{t=0}$$

$$= \frac{\partial^{u-v}}{\partial t^{u-v}} \left( -\sum_{\alpha,\beta} B_{\alpha}^{\top} \sum_{k=0}^{\infty} (-1)^{k} \left( \sum_{w_{1}+\ldots+w_{q}=\mid w\mid=k} \binom{\mid w\mid}{w} t^{w+\alpha^{\sharp}\beta^{\sharp}(0)} \tilde{A}^{w} \right) B_{\beta} \right) \mid_{t=0}$$

$$= -\sum_{\alpha,\beta} \sum_{v \leq \alpha_{\flat}\beta_{\flat}(u)} B_{\alpha}^{\top} (-1)^{\mid u\mid-\mid v\mid-2} (u-v)! \binom{\mid u\mid-\mid v\mid-2}{u-\alpha^{\sharp}\beta^{\sharp}(v)} \tilde{A}^{u-\alpha^{\sharp}\beta^{\sharp}(v)} B_{\beta}$$

$$= \sum_{\alpha,\beta} \sum_{v \leq \alpha_{\flat}\beta_{\flat}(u)} B_{\alpha}^{\top} (-1)^{\mid u\mid-\mid v\mid-1} (u-v)! \binom{\mid u\mid-\mid v\mid-2}{u-\alpha^{\sharp}\beta^{\sharp}(v)} \tilde{A}^{u-\alpha^{\sharp}\beta^{\sharp}(v)} B_{\beta} .$$

Finally we obtain

$$u!\sigma_{u} = u!\tilde{\sigma}_{u} + \sum_{\alpha} (\alpha_{\flat}(u))! \binom{u}{\alpha_{\flat}(u)} C_{\alpha}\tilde{\sigma}_{\alpha_{\flat}(u)}$$
$$+ \sum_{\alpha} \sum_{\beta} \binom{u}{v} (-1)^{|u|-|v|-1} (u-v)! v! \binom{|u|-|v|-2}{u-\alpha_{\flat}\beta_{\flat}(v)} B_{\alpha}\tilde{\sigma}_{v}$$

or equivalently,

$$\sigma_{u} = \tilde{\sigma}_{u} + \sum_{\alpha} C_{\alpha} \tilde{\sigma}_{\alpha_{\flat}(u)} + \sum_{\alpha,\beta} \sum_{0 \leq v \leq \alpha_{\flat},\beta_{\flat}(u)} (-1)^{|u|-|v|-1} \binom{|u|-|v|-2}{u-\alpha^{\sharp}\beta^{\sharp}(v)} B_{\alpha}^{\top} \tilde{A}^{u-\alpha^{\sharp}\beta^{\sharp}(v)} B_{\beta} \tilde{\sigma}_{v}.$$

Hence

$$\sigma_{u} = \tilde{\sigma}_{u} + \sum_{\alpha} C_{\alpha} \tilde{\sigma}_{\alpha_{\flat}(u)} + \sum_{\alpha^{\sharp} \beta^{\sharp}(0) \leq w \leq u} \sum_{\alpha,\beta} (-1)^{|w| - |v| + 1} \binom{|u| - |w|}{u - w} B_{\alpha} \tilde{A}^{u - w} B_{\beta} \tilde{\sigma}_{\alpha_{\flat} \beta_{\flat}(w)}.$$

#### 4. Generalized Newton transformation on the boundary

We use the same notations as in previous sections. In this section we give the expression of the GNT  $T_u = T_u(\tilde{A})$ , where  $\tilde{A} = (A_1|\Sigma, \dots, A_q|\Sigma)$ , on the boundary  $\Sigma^{n-1}$  of  $M^n$ . Recall that

$$A_{\alpha}|_{\Sigma} = \rho_{\alpha}A_{\Sigma} + \mu_{\alpha}I,$$

where

$$\rho_{\alpha} = \langle \eta, N_{\alpha} \rangle, \quad \mu_{\alpha} = \langle V, N_{\alpha} \rangle, \quad V = \sum_{\alpha=1}^{q} \lambda_{\alpha} \xi_{\alpha}.$$

**Proposition 4.** Let  $\overline{M}^{n+q}$  be an (n+q)-Riemannian manifold and  $P^n \subset \overline{M}^{n+q}$  an oriented totally umbilical n-submanifold of  $\overline{M}^{n+q}$ . Denote by  $\Sigma^{n-1} \subset P^n$  an (n-1)-compact hypersurface of  $P^n$ . Let  $\Psi: M^n \longrightarrow \overline{M}^{n+q}$  be an oriented connected and compact submanifold of  $\overline{M}^{n+q}$  with boundary  $\Sigma^{n-1} = \Psi(\partial M)$ . Then along the boundary  $\partial M$ , we have

(4.1) 
$$\langle T_u \nu, \nu \rangle = \tilde{\sigma}_u(A_1|_{\Sigma}, ..., A_q|_{\Sigma}).$$

*Proof.* We make a recursive proof. Assume that (4.1) holds for any multiindex v < u. We have by the recurrence definition of  $T_u$ 

$$\begin{split} \langle T_{u}\nu,\nu\rangle &= \sigma_{u} \left\langle \nu,\nu\right\rangle - \sum_{\alpha} \left\langle A_{\alpha}T_{\alpha_{\flat}(u)}\nu,\nu\right\rangle \\ &= \sigma_{u} - \sum_{\alpha} \left\langle T_{\alpha_{\flat}(u)}\nu,A_{\alpha}\nu\right\rangle \\ &= \sigma_{u} - \sum_{\alpha} \left\langle T_{\alpha_{\flat}(u)}\nu,\nu\right\rangle \left\langle A_{\alpha}\nu,\nu\right\rangle - \sum_{\alpha,i} \left\langle T_{\alpha_{\flat}(u)}\nu,e_{i}\right\rangle \left\langle A_{\alpha}e_{i},\nu\right\rangle. \end{split}$$

Put

$$C_{\alpha} = \langle A_{\alpha} \nu, \nu \rangle , b_{i,\alpha} = \langle A_{\alpha} e_i, \nu \rangle$$

then

$$\langle T_u \nu, \nu \rangle = \sigma_u - \sum_{\alpha} C_{\alpha} \tilde{\sigma}_{\alpha_{\flat}(u)} - \sum_{\alpha, i} b_{i,\alpha} \langle T_{\alpha_{\flat}(u)} \nu, e_i \rangle.$$

Let us compute  $\langle T_u \nu, e_i \rangle$  for any multi-index  $u \in \mathbb{N}^q$ . Notice that

$$A_{\alpha}e_i = \rho_{\alpha}A_{\Sigma}e_i + \mu_{\alpha}e_i + b_{i,\alpha}\nu$$

assuming that  $\{e_1, ..., e_{n-1}\}$  is an orthonormal basis of  $T_p\Sigma^{n-1}$  consisting of eigenvectors of  $A_{\Sigma}$  (with eigenvalues  $\tau_i$ ). Thus

$$A_{\alpha}e_i = \gamma_{i,\alpha}e_i + b_{i,\alpha}\nu$$

where

$$\gamma_{i,\alpha} = \rho_{\alpha} \tau_i + \mu_{\alpha}$$
.

We will show inductively that

$$(4.2) \qquad \langle T_u \nu, e_i \rangle = \sum_{\alpha} \sum_{\alpha^{\sharp}(0) \le w \le u} (-1)^{|u| - |w| + 1} \binom{|u| - |w|}{u - w} b_{i,\alpha} \gamma_i^{u - w} \tilde{\sigma}_{\alpha_{\flat}(w)}$$

where

$$\gamma_i = (\gamma_{i,1}, ..., \gamma_{i,q})$$

Indeed, for  $u = \beta^{\sharp}(0)$  we have,

$$\begin{split} \langle T_{u}\nu, e_{i} \rangle &= \sigma_{\beta^{\sharp}(0)} \langle \nu, e_{i} \rangle - \sum_{\alpha} \left\langle A_{\alpha} T_{\alpha_{\flat}(\beta^{\sharp}(0))} \nu, \nu \right\rangle \\ &= -\sum_{\alpha} \left\langle A_{\alpha} \nu, \nu \right\rangle \\ &= -b_{i,\beta} \\ &= \sum_{\alpha} \sum_{\alpha^{\sharp}(0) \leq w \leq \beta^{\sharp}(0)} (-1)^{\left|\beta^{\sharp}(0)\right| - |w| + 1} \binom{\left|\beta^{\sharp}(0)\right| - |w|}{\beta^{\sharp}(0) - w} b_{i,\alpha} \gamma_{i}^{\beta^{\sharp}(0) - w} \tilde{\sigma}_{\alpha_{\flat}(w)}, \end{split}$$

since the sum reduces to one element for  $\alpha = \beta$ .

Assume that (4.2) holds for all multi index v < u. Then again by the recursive definition of  $T_u$  we get

$$\begin{split} \langle T_{u}\nu,e_{i}\rangle &= \sigma_{u} \left\langle \nu,e_{i}\right\rangle - \sum_{\alpha} \left\langle A_{\alpha}T_{\alpha_{\flat}(u)}\nu,e_{i}\right\rangle \\ &= -\sum_{\alpha} \left\langle T_{\alpha_{\flat}(u)}\nu,A_{\alpha}e_{i}\right\rangle \\ &= -\sum_{\alpha} \left\langle T_{\alpha_{\flat}(u)}\nu,\nu\right\rangle \left\langle A_{\alpha}e_{i},\nu\right\rangle - \sum_{\alpha,j} \left\langle T_{\alpha_{\flat}(u)}\nu,e_{i}\right\rangle \left\langle A_{\alpha}e_{i},e_{j}\right\rangle \\ &= -\sum_{\alpha} \left\langle T_{\alpha_{\flat}(u)}\nu,e_{i}\right\rangle \gamma_{i,\alpha} - \sum_{\alpha} b_{i,\alpha}\tilde{\sigma}_{\alpha_{\flat}(u)} \\ &= -\sum_{\alpha} \gamma_{i,\alpha} \left(\sum_{\beta} \sum_{\beta^{\sharp}(0) \leq w \leq \alpha_{\flat}(u)} (-1)^{|u|-|w|} \binom{|u|-|w|-1}{\alpha_{\flat}(u)-w} b_{i,\beta} \gamma_{i}^{\alpha_{\flat}(u)-w} \tilde{\sigma}_{\beta_{\flat}(w)} \right) \\ &- \sum_{\alpha} b_{i,\alpha}\tilde{\sigma}_{\alpha_{\flat}(u)}. \end{split}$$

Clearly

$$\gamma_{i,\alpha}.\gamma_i^{\alpha_{\flat}(u)-w} = \gamma_i^{u-w}$$

Notice that taking w = u we get the last sum. Thus we obtain

$$\langle T_u \nu, e_i \rangle = \sum_{\alpha, \beta} \left( \sum_{\beta^{\sharp}(0) \le w \le u} (-1)^{|u| - |w| + 1} \binom{|u| - |w| - 1}{\alpha_{\flat}(u) - w} b_{i,\beta} \gamma_i^{u - w} \tilde{\sigma}_{\beta_{\flat}(w)} \right).$$

Since

$$\sum_{\alpha} \binom{|u| - |w| - 1}{\alpha_{\flat}(u) - w} = \binom{|u| - |w|}{u - w}$$

we deduce

$$\langle T_u \nu, e_i \rangle = \sum_{\beta} \left( \sum_{\beta \sharp (0) \le w \le u} (-1)^{|u| - |w| + 1} \binom{|u| - |w|}{u - w} b_{i,\beta} \gamma_i^{u - w} \tilde{\sigma}_{\beta_{\flat}(w)} \right)$$

which end the proof of (4.2).

Now, we may prove (4.1). First, we have

$$\sum_{\alpha,i} \left\langle T_{\alpha_{\flat}(u)} \nu, e_i \right\rangle b_{i,\alpha} = \sum_{\alpha,\beta,i} \left( \sum_{\beta^{\sharp}(0) \leq w \leq \alpha_{\flat}(u)} (-1)^{|u|-|w|} \binom{|u|-|w|-1}{\alpha_{\flat}(u)-w} b_{i,\beta} \gamma_i^{\alpha_{\flat}(u)-w} \tilde{\sigma}_{\beta_{\flat}(w)} \right)$$

Replacing w by  $\alpha_{\flat}\beta_{\flat}(w)$ , we obtain

$$\sum_{\alpha,i} \left\langle T_{\alpha_{\flat}(u)} \nu, e_i \right\rangle b_{i,\alpha} = \sum_{\alpha,\beta,i} \left( \sum_{\alpha^{\sharp}\beta^{\sharp}(0) \leq w \leq u} (-1)^{|u|-|w|+1} \binom{|u|-|w|}{u-w} b_{i,\alpha} \gamma_i^{u-w} b_{i,\beta} \tilde{\sigma}_{\alpha_{\flat}\beta_{\flat}(w)} \right)$$

Noticing that

$$\sum_{i} b_{i,\alpha} \gamma_i^{u-w} b_{i,\beta} = B_{\alpha}^{\top} \tilde{A}^{u-w} B_{\beta}$$

we infer

$$\sum_{\alpha,i} \left\langle T_{\alpha_{\flat}(v)} \nu, e_i \right\rangle b_{i,\alpha} = \sum_{\alpha,\beta} \sum_{\alpha^{\sharp} \beta^{\sharp}(0) \leq w \leq u} (-1)^{|u|-|w|+1} \binom{|u|-|w|}{u-w} B_{\alpha}^{\top} \tilde{A}^{u-w} B_{\beta}.\tilde{\sigma}_{\alpha_{\flat}\beta_{\flat}(w)}$$

Summing up all the above considerations we get

$$\begin{split} \left\langle T_{u}\nu,\nu\right\rangle &=\sigma_{u}-\sum_{\alpha}\left\langle T_{\alpha_{\flat}(u)}\nu,\nu\right\rangle \left\langle A_{\alpha}\nu,\nu\right\rangle -\sum_{\alpha,i}\left\langle T_{\alpha_{\flat}(u)}\nu,e_{i}\right\rangle \left\langle A_{\alpha}e_{i},\nu\right\rangle \\ &=\sigma_{u}-\sum_{\alpha}C_{\alpha}\tilde{\sigma}_{\alpha_{\flat}(u)}-\sum_{\alpha,\beta}\sum_{\alpha^{\sharp}\beta^{\sharp}(0)\leq w\leq u}(-1)^{|u|-|w|+1}\binom{|u|-|w|}{u-w}B_{\alpha}^{\intercal}\tilde{A}^{u-w}B_{\beta}.\tilde{\sigma}_{\alpha_{\flat}\beta_{\flat}(w)} \end{split}$$

or, equivalently,

$$\sigma_{u} = \tilde{\sigma}_{u} + \sum_{\alpha} C_{\alpha} \tilde{\sigma}_{\alpha_{\flat}(u)} + \sum_{\alpha,\beta} \sum_{\alpha \notin \beta \notin \{0\} \le w \le u} (-1)^{|u| - |w| + 1} \binom{|u| - |w|}{u - w} B_{\alpha}^{\top} \tilde{A}^{u - w} B_{\beta} . \tilde{\sigma}_{\alpha_{\flat}\beta_{\flat}(w)}$$

Applying Proposition (3), we obtain

$$\langle T_u \nu, \nu \rangle = \tilde{\sigma}_u.$$

By Proposition (1) we obtain the following expression of  $\langle T_u \nu, \nu \rangle$  in terms of symmetric functions of the shape operator  $A_{\Sigma}$ .

**Corollary 1.** With the conditions of Proposition (4), we have

(4.3) 
$$\langle T_u \nu, \nu \rangle = \frac{1}{n-1-|u|} \sum_{l < u} \binom{n-1-|l|}{|u|-l} \rho^l \mu^{u-l} \sigma_{|l|}(A_{\Sigma}).$$

The relation (4.3) will be more simple if we suppose that the embedding  $P^n \subset \overline{M}^{n+q}$  is totally geodesic.

**Corollary 2.** With the conditions of Proposition (4) and assuming that  $P^n \subset \overline{M}^{n+q}$  is totally geosedic, then for every multi-index u with length  $|u| \leq n-1$ , we have

$$\langle T_u \nu, \nu \rangle = \rho^u \sigma_{|u|} (A_{\Sigma})$$

*Proof.* It suffices to use (4.3) with  $\mu_{\alpha} = 0$ .

#### 5. Transversality of Submanifolds

The formula for the generalized Newton transformation implies the relation between transversality of  $M^n$  and  $P^n$  and ellipticity of  $T_u$  provided that  $P^n$  is totally geodesic in  $\overline{M}^{n+1}$ . This generalizes the result in ([2]) to any arbitrary codimension.

**Theorem 1.** With the conditions in Corollary (2) the submanifolds  $M^n$  and  $P^n$  are transversal along  $\partial M$  provided that for some multi-index u of length  $1 \leq |u| \leq n-1$ , the generalized Newton transformation  $T_u$  is positive definite on  $M^n$ .

*Proof.* Saying that  $M^n$  and  $P^n$  are not transversal means that there exist  $p \in \partial M$  such that for every  $\alpha \in \{1, ..., q\}$  we have

$$\rho_u = \langle \eta, N_\alpha \rangle = 0 \text{ at } p.$$

Therefore, if we suppose that for all  $p \in \overline{M}^{n+q}$ ,  $T_u$  is positive definite, then by Corollary 7,  $\rho^u(p) \neq 0$ . Thus

$$\langle \eta, N_{\alpha} \rangle \neq 0$$
,

hence  $M^n$  and  $P^n$  are transversal.

### REFERENCES

- [1] M. Abdelmalek, M. Benallili, Transversality versus ellipticity on pseudo-Riemannian manifolds, International Journal of Geometric Methods in Modern Physics, Volume 12, Issue, 7, (2015), 12pp..
- [2] L. J. Alias, S. de Lira, J. M. Malacarne, Constant higher-order mean curvature hypersurfaces in Riemannian spaces, J Inst. of Math. Jussieu, 5(4), 527-562 (2006).
- [3] L. J. Alias, A.G. Colares, Uniqueness of spacelike hypersurfaces with constant higher order mean curvature in generalized Robertson-Walker spacetimes, Math. Proc. Camb. Phil. Soc. 143, 703-729 (2007).

#### 18MOHAMED ABDELMALEK, MOHAMMED BENALILI, AND KAMIL NIEDZIAŁOMSKI

- [4] K. Andrzejewski, W. Kozlowski, K. Niedzialomski, Generalized Newton transformation and its applications to extrinsic geometry, arXiv:1211.4754.
- [5] K. Andrzejewski, P. Walczak, The Newton transformations and new integral formulae for foliated manifolds, Ann. Glob. Anal.Geom, Vol. 37,103-111 (2010).
- [6] L. Cao, H. Li, r-minimal submanifolds in space forms, Ann Glob Anal Geom. Vol. 32, 311-341 (2007)..
- [7] V. Rovenski, P. G. Walczak, Integral formulae on foliated symmetric spaces, Math. Ann. Vol. 352, (2012) 223-237.

Current address: Laboratoire Systèmes Dynamiques et Applications, Université Aboubakr Belkaid, Tlemcen and Preparatory, School of Economy; Business and Management Sciences,, Tlemcen Algeria.

E-mail address: abdelmalekmhd@yahoo.fr

Current address: Laboratoire Syetèmes Dynamiques et Applications, Dept. Mathematics, Faculty of Sciences, University Abou-bakr Belkaïd Tlemcen, Algeria.

 $E ext{-}mail\ address: m\_benalili@mail.univ-tlemcen.dz}$ 

Current address: Department of Mathematics and Computer Science, University of Lodz, Banacha 22, 90-238 Lódź, Poland.

E-mail address: kamiln@math.uni.lodz.pl