

CHARACTERIZATION OF THE RIESZ EXPONENTIAL FAMILY ON HOMOGENEOUS CONES

HIDEYUKI ISHI AND BARTOSZ KOŁODZIEJEK

ABSTRACT. In the paper we present a characterization theorem of the Riesz measure and a Wishart exponential family on homogeneous cones through the invariance property of a natural exponential family under the action of the triangular group.

1. INTRODUCTION

Following Casalis [3], we consider natural exponential family (NEF), which is invariant under a subgroup G of the general linear group of a finite dimensional linear space. NEF is considered on arbitrary finite dimensional linear space \mathbb{E} . Under a weak assumption on G (see Theorem 2.1), the generating measure of such family is of the form

$$e^{-\langle \theta_0, x \rangle} \mu_0(dx)$$

for some θ_0 in the dual space \mathbb{E}^* , where μ_0 is a G -invariant measure.

We apply Theorem 2.1 to the problem of a characterization of the Riesz measure on a homogeneous cone through the invariance property of NEF under the action of the triangular group and we solve this problem in Theorem 3.5. Since NEF generated by the Riesz measure consists of Wishart distributions, Theorem 3.5 provides also a characterization of a Wishart distribution on homogeneous cones [1, 11, 27]. We would like to mention that this problem was announced by Letac in [25, Section 4], who pointed out that the natural framework for such characterizations is indeed homogeneous cone.

There are essentially two types of characterizations of NEFs. The first one is connected with the central object of NEF, that is, the variance function. The aim is then to describe a generating measure of a NEF with given variance function. Much has been done in this direction, but still much more is unknown. A generating measure is known for $\mathbb{E} = \mathbb{R}$, when the variance function is a quadratic polynomial [29], cubic polynomial [15, 28] and of some more sophisticated forms, like $P\Delta + Q\sqrt{\Delta}$, where P , Δ and Q are quadratic polynomials [26]. For $\mathbb{E} = \mathbb{R}^d$ we have to emphasize [4] with its deep connection between homogeneous quadratic variance functions and Euclidean Jordan algebras, simple quadratic [5] and simple cubic variance functions [16], just to name a few.

The second type of characterizations of NEFs is through the invariance properties under some group action. We quote some results on invariant NEFs under a given subgroup G of the general affine group: one parameter group [7], group of rotations [33], Moebius group [23], the identity component $\text{Aut}_0(\text{Sym}_+(N, \mathbb{R}))$ of the linear automorphism group of the cone $\text{Sym}_+(N, \mathbb{R}) \subset \text{Sym}(N, \mathbb{R})$ [25], triangular group of a simple Euclidean Jordan algebra [13] and its modification [14]. Here $\text{Sym}(N, \mathbb{R})$ stands for the symmetric $N \times N$ matrices with real entries and $\text{Sym}_+(N, \mathbb{R})$ is the cone of positive definite real $N \times N$ matrices. In the last three quoted papers the characterizations were carried out by showing that the variance function (which uniquely determines NEF) coincides with the variance function of some Riesz measure (or its image by the involution $x \mapsto -x$) on $\text{Sym}_+(N, \mathbb{R})$ ([25]) and symmetric cones ([13, 14]).

2010 *Mathematics Subject Classification.* Primary 60B11; secondary 62E10.

Key words and phrases. natural exponential families, Riesz measure, Wishart distribution, general affine group.

In the present paper we solve the problem raised by Letac [25] on homogeneous cones. We use a matrix realization of homogeneous cones, which proves here very useful and is more accessible to the reader who is not familiar with homogeneous cones, clans and T -algebras. We would like to emphasize that homogeneous cones are of great importance in statistics despite their theoretical character. Particularly, homogeneous cones appear very naturally in statistics in the context of graphical Gaussian models [27]. The aforementioned matrix realization of homogeneous cones allows to describe many of the colored graphical Gaussian models [17]. The formula for variance function of $F(\mathcal{R}_s)$ on homogeneous cones is the topic of our joint paper with Piotr Graczyk [12]. At last we would like to point out that this work is closely related to [30, 31], where NEFs and information theory are treated on homogeneous Hessian structures.

The paper is organized as follows. The concept of natural exponential families is introduced in the next section. In Section 3.2 we give an application of Theorem 2.1 to a characterization of the Riesz measure on homogeneous cone (Theorem 3.5). A crucial role in the proof of characterization is played by a matrix realization of any homogeneous cone ([19]), which is explained in Section 3.1. Section 3 ends with some comments.

2. NATURAL EXPONENTIAL FAMILIES

In this section we will introduce all the necessary facts on NEFs that will be needed later on. It is however worth mentioning that the standard reference book on exponential families is [2].

Let \mathbb{E} be a finite dimensional real linear space and \mathbb{E}^* its dual space. The coupling of $\theta \in E^*$ and $x \in E$ is denoted by $\langle \theta, x \rangle$. Let μ be a positive Radon measure on \mathbb{E} . We define its Laplace transform $L_\mu: \mathbb{E}^* \rightarrow (0, \infty]$ by

$$L_\mu(\theta) = \int_{\mathbb{E}} e^{\langle \theta, x \rangle} \mu(dx).$$

By $\Theta(\mu)$ we will denote the interior of the set $\{\theta \in \mathbb{E}^*: L_\mu(\theta) < \infty\}$. Hölder's inequality implies that the set $\Theta(\mu)$ is convex and the cumulant function

$$k_\mu(\theta) = \log L_\mu(\theta)$$

is convex on $\Theta(\mu)$ and it is strictly convex if and only if μ is not concentrated on some affine hyperplane of \mathbb{E} . Let $\mathcal{M}(\mathbb{E})$ be the set of positive Radon measures on \mathbb{E} such that $\Theta(\mu)$ is not empty and μ is not concentrated on some affine hyperplane of \mathbb{E} .

For $\mu \in \mathcal{M}(\mathbb{E})$ we define the *natural exponential family (NEF) generated by μ* (denoted by $F(\mu)$) as the set of probability measures of the form

$$P(\theta, \mu)(dx) = e^{\langle \theta, x \rangle - k_\mu(\theta)} \mu(dx), \quad \theta \in \Theta(\mu).$$

Let us note that $F(\mu) = F(\mu')$ if and only if $\mu'(dx) = e^{\langle a, x \rangle + b} \mu(dx)$ for some $a \in \mathbb{E}^*$ and $b \in \mathbb{R}$.

We will now describe the action of elements from the general linear group $GL(\mathbb{E})$ on a NEF. The identity element of $GL(\mathbb{E})$ will be denoted by Id . Let $F = F(\mu)$ be a NEF on \mathbb{E} . Let $g_*\mu$ denote the image measure of μ by g and let $g.F(\mu)$ stand for the family of image measures $g_*P(\theta, \mu)$ ($\theta \in \Theta(\mu)$). Then, for any $g \in GL(\mathbb{E})$, we have $g.F(\mu) = F(g_*\mu)$.

We say that a measure μ_0 is *invariant under a subgroup G* of $GL(\mathbb{E})$ if for all $g \in G$ there exists a constant $c_g > 0$ for which $\mu_0(gA) = c_g \mu_0(A)$ for any measurable set $A \subset \mathbb{E}$. This condition is equivalent to

$$(2.1) \quad L_{\mu_0}(g^*\theta) = c_g^{-1} L_{\mu_0}(\theta), \quad \theta \in \Theta(\mu_0).$$

Note that the correspondence $G \ni g \mapsto c_g \in \mathbb{R}$ is a character of the group G .

Let $\mu \in \mathcal{M}(\mathbb{E})$. Observe that the condition $g.F(\mu) = F(\mu)$ implies that for any $g \in G$ there exists $a(g) \in \Theta(\mu) \subset \mathbb{E}^*$ and $b(g) \in \mathbb{R}$ such that

$$(2.2) \quad g_*\mu(dx) = e^{\langle a(g), x \rangle + b(g)} \mu(dx).$$

Then, Casalis [3] showed that (see [3, Theorem 2.2]) functions a and b satisfy the following system of equations: for any $(g, g') \in G^2$

$$\begin{aligned} a(gg') &= (g^*)^{-1}a(g') + a(g), \\ b(gg') &= b(g) + b(g'). \end{aligned}$$

Let us assume that G contains $c\text{Id}$ for some $c \neq 1$. Then, we obtain for any $g \in G$,

$$\begin{aligned} a(c\text{Id } g) &= \frac{1}{c}a(g) + a(c\text{Id}), \\ a(g \ c\text{Id}) &= (g^*)^{-1}a(c\text{Id}) + a(g). \end{aligned}$$

Equating the right hand sides of above formulas, we arrive at

$$a(g) = \theta_0 - (g^*)^{-1}\theta_0, \quad g \in G,$$

with $\theta_0 = \frac{c}{1-c}a(c\text{Id})$. Define $\mu_0(dx) = e^{\langle \theta_0, x \rangle} \mu(dx)$. Then, (2.2) implies that μ_0 is G -invariant. Thus, we obtain the following

Theorem 2.1 *Let G be a subgroup of $GL(\mathbb{E})$ and let $F = F(\mu)$ be a G -invariant NEF on \mathbb{E} , that is, for any $g \in G$ one has $g.F = F$. If G contains $c\text{Id}$ for some $c \neq 1$, then there exist $\theta_0 \in \mathbb{E}^*$ and a G -invariant measure μ_0 such that*

$$\mu(dx) = e^{-\langle \theta_0, x \rangle} \mu_0(dx).$$

In such case we have also $F = F(\mu_0)$.

3. CHARACTERIZATION OF THE RIESZ MEASURE ON HOMOGENEOUS CONES

3.1. Matrix realization of homogeneous cones. Let V be a real linear space and Ω a regular open convex set in V containing no line. The cone Ω is said to be *homogeneous* if the linear automorphism group $G(\Omega) = \{g \in GL(V) : g\Omega = \Omega\}$ acts transitively on Ω , that is, for any x and y in Ω there exists $g \in G(\Omega)$ such that $y = gx$.

We will now give a very useful representation of homogeneous cones following [21, Section 3]. For a symmetric matrix $x \in \text{Sym}(N, \mathbb{R})$, we denote by \mathfrak{x} the lower triangular matrix of size N defined by

$$(\mathfrak{x})_{ij} = \begin{cases} x_{ij}, & \text{if } i > j, \\ x_{ii}/2, & \text{if } i = j, \\ 0, & \text{if } i < j. \end{cases}$$

Then we have $x = \mathfrak{x} + \hat{x}$, where $\hat{x} = \mathfrak{x}^\top$ is the transpose of \mathfrak{x} . For $x, y \in \text{Sym}(N, \mathbb{R})$, we define

$$x \triangle y := \mathfrak{x}y + y\hat{x} \in \text{Sym}(N, \mathbb{R}).$$

Then, $(\text{Sym}(N, \mathbb{R}), \triangle)$ forms a non-associative algebra with unit element I_N . Let \mathcal{Z} be a subalgebra of $(\text{Sym}(N, \mathbb{R}), \triangle)$ and $H_{\mathcal{Z}}$ be the set of lower triangular matrices from elements of \mathcal{Z} with positive diagonal entries, that is, $H_{\mathcal{Z}} := \{\mathfrak{x} : \mathfrak{x} \in \mathcal{Z} \text{ and } x_{ii} > 0\}$.

Define $\Omega_{\mathcal{Z}} = \{x \in \mathcal{Z} : x \text{ is positive definite}\}$ and consider for any $T \in H_{\mathcal{Z}}$ the linear operators $\rho(T) : \mathcal{Z} \rightarrow \mathcal{Z}, x \mapsto \rho(T)x = TxT^\top$. It can be shown that $\rho(H_{\mathcal{Z}})$ acts on $\Omega_{\mathcal{Z}}$ transitively, which means that $\Omega_{\mathcal{Z}}$ is a homogeneous cone ([22, Theorem 3]). Furthermore, we have the following

Theorem 3.1 ([20]) *For a homogeneous cone $\Omega \subset V$, there exists a subalgebra $\mathcal{Z} \subset \text{Sym}(N, \mathbb{R})$ and a linear isomorphism $\phi: V \rightarrow \mathcal{Z}$ such that $\phi(\Omega) = \Omega_{\mathcal{Z}}$.*

We say that $\mathcal{Z} \subset \text{Sym}(N, \mathbb{R})$ admits a normal block decomposition if there exists a partition $N = n_1 + \dots + n_r$ and subspaces $\mathcal{V}_{lk} \subset \text{Mat}(n_l, n_k, \mathbb{R})$, $1 \leq k < l \leq r$, such that \mathcal{Z} is the set of symmetric matrices of the form

$$\begin{pmatrix} X_{11} & X_{21}^\top & \cdots & X_{r1}^\top \\ X_{21} & X_{22} & & X_{r2}^\top \\ \vdots & & \ddots & \\ X_{r1} & X_{r2} & & X_{rr} \end{pmatrix} \quad \left(\begin{array}{l} X_{ll} = x_{ll} I_{n_l}, \ x_{ll} \in \mathbb{R}, \ 1 \leq l \leq r \\ X_{lk} \in \mathcal{V}_{lk}, \ 1 \leq k < l \leq r \end{array} \right).$$

We will write $\mathcal{Z}_{\mathcal{V}}$ for this space.

Theorem 3.2 (i) ([22, Theorem 2]) *Let \mathcal{Z} be a subalgebra of $(\text{Sym}(N, \mathbb{R}), \Delta)$ with $I_N \in \mathcal{Z}$. Then there exists a permutation matrix w , such that $w\mathcal{Z}w^\top$ admits a normal block decomposition.*
(ii) ([22, Proposition 2]) *$\mathcal{Z}_{\mathcal{V}}$ is a subalgebra of $(\text{Sym}(N, \mathbb{R}), \Delta)$ if and only if the subspaces $\{\mathcal{V}_{lk}\}_{1 \leq k < l \leq r}$ satisfy the following conditions:*
(V1) $A \in \mathcal{V}_{lk}, B \in \mathcal{V}_{ki} \implies AB \in \mathcal{V}_{li}$ for any $1 \leq i < k < l \leq r$,
(V2) $A \in \mathcal{V}_{li}, B \in \mathcal{V}_{ki} \implies AB^\top \in \mathcal{V}_{lk}$ for any $1 \leq i < k < l \leq r$,
(V3) $A \in \mathcal{V}_{lk} \implies AA^\top \in \mathbb{R}I_{n_l}$ for any $1 \leq k < l \leq r$.

If $\mathcal{Z} = \mathcal{Z}_{\mathcal{V}}$ we shall write $\Omega_{\mathcal{V}}, H_{\mathcal{V}}$ for $\Omega_{\mathcal{Z}}$ and $H_{\mathcal{Z}}$, respectively.

Condition (V3) allows us to define an inner product on \mathcal{V}_{lk} , $1 \leq k < l \leq r$, by

$$AA^\top = (A|A)I_{n_l}, \quad A \in \mathcal{V}_{lk}.$$

We then define the *standard inner product* on $\mathcal{Z}_{\mathcal{V}}$ by

$$\langle x, y \rangle := \sum_{k=1}^r x_{kk} y_{kk} + 2 \sum_{1 \leq k < l \leq r} (X_{lk}|Y_{lk}), \quad x, y \in \mathcal{Z}_{\mathcal{V}}.$$

We identify $\mathcal{Z}_{\mathcal{V}}^*$ with $\mathcal{Z}_{\mathcal{V}}$ using the standard inner product. Note that $\langle \cdot, \cdot \rangle$ coincides with the trace inner product only if $n_1 = \dots = n_r = 1$.

Define a one-dimensional representation of $H_{\mathcal{V}}$ by

$$\chi_{\underline{s}}(T) := \prod_{k=1}^r t_{kk}^{2s_k},$$

where $\underline{s} = (s_1, \dots, s_r) \in \mathbb{C}^r$. We have (see e.g. [8, Lemma 2.4])

Lemma 3.1 *Let χ be a one-dimensional representation of $H_{\mathcal{V}}$. Then, there exists $\underline{s} \in \mathbb{C}^r$ such that*

$$\chi = \chi_{\underline{s}}.$$

This fact will be important later on.

For any open convex cone Ω we define the *dual cone* of Ω by

$$\Omega^* = \{\xi \in V^*: \langle \xi, x \rangle > 0 \ \forall x \in \bar{\Omega} \setminus \{0\}\},$$

where V^* is the dual space of V . If Ω is homogeneous, then so is Ω^* . Let $\Omega_{\mathcal{V}}^*$ denote the dual cone of $\Omega_{\mathcal{V}}$.

For $T \in H_{\mathcal{V}}$, we denote by $\rho^*(T)$ the adjoint operator of $\rho(T) \in GL(\mathcal{Z}_{\mathcal{V}})$ defined in such a way that $\langle \rho^*(T)\xi, x \rangle = \langle \xi, \rho(T)x \rangle$ for $x, \xi \in \mathcal{Z}_{\mathcal{V}}$. Then we see from [32, Chapter 1, Proposition 9] that for any $\xi \in \Omega_{\mathcal{V}}^*$, there exists a unique $T \in H_{\mathcal{V}}$ such that $\xi = \rho^*(T)I_N$.

Definition 3.1 Let $\Delta_{\underline{s}}^*: \Omega_{\mathcal{V}}^* \rightarrow \mathbb{C}$ be the function given by

$$\Delta_{\underline{s}}^*(\xi) = \Delta_{\underline{s}}^*(\rho^*(T)I_N) := \chi_{\underline{s}^*}(T),$$

where $\underline{s}^* = (s_r, \dots, s_1) \in \mathbb{C}^n$.

Example 3.1 Let

$$\mathcal{Z}_{\mathcal{V}} := \left\{ \begin{pmatrix} x_{11} & 0 & x_{31} \\ 0 & x_{22} & x_{32} \\ x_{31} & x_{32} & x_{33} \end{pmatrix} : x_{11}, x_{22}, x_{33}, x_{31}, x_{32} \in \mathbb{R} \right\}.$$

Conditions (V1)–(V3) are satisfied and we have $n_1 = n_2 = n_3 = 1$, $N = r = 3$. Then,

$$\Omega_{\mathcal{V}} = \mathcal{Z}_{\mathcal{V}} \cap \text{Sym}_+(3, \mathbb{R}) = \{x \in \mathcal{Z}_{\mathcal{V}} : x_{11} > 0, x_{22} > 0, \det x > 0 \in \mathbb{R}\}$$

and its dual cone is given by

$$\Omega_{\mathcal{V}}^* = \{\xi \in \mathcal{Z}_{\mathcal{V}} : \xi_{33} > 0, \xi_{11}\xi_{33} > \xi_{31}^2, \xi_{22}\xi_{33} > \xi_{32}^2\}.$$

The cone $\Omega_{\mathcal{V}}^*$ is called the Vinberg cone, while $\Omega_{\mathcal{V}}$ is called the dual Vinberg cone. The cones $\Omega_{\mathcal{V}}^*$ and $\Omega_{\mathcal{V}}$ are the lowest dimensional non-symmetric homogeneous cones. Moreover, for $\xi \in \Omega_{\mathcal{V}}^*$, we have

$$\Delta_{\underline{s}}^*(\xi) = \left(\frac{\xi_{11}\xi_{33} - \xi_{31}^2}{\xi_{33}} \right)^{s_3} \left(\frac{\xi_{22}\xi_{33} - \xi_{32}^2}{\xi_{33}} \right)^{s_2} \xi_{33}^{s_1}.$$

We see that for any $S \in H_{\mathcal{V}}$ and $\xi = \rho^*(T)I_N \in \Omega_{\mathcal{V}}^*$ we have

$$(3.1) \quad \Delta_{\underline{s}}^*(\rho^*(S)\xi) = \chi_{\underline{s}^*}(TS) = \chi_{\underline{s}^*}(T)\chi_{\underline{s}^*}(S) = \Delta_{\underline{s}}^*(\xi)\Delta_{\underline{s}}^*(\rho^*(S)I_N)$$

and, since any character of $H_{\mathcal{V}}$ is of the form $\chi_{\underline{s}^*}$, property (3.1) characterizes Δ^* (see [24]). Function Δ^* is sometimes termed a generalized power function. Its importance is emphasized by the following result [10, 18].

Theorem 3.3 There exists a positive measure $\mathcal{R}_{\underline{s}}$ on $\mathcal{Z}_{\mathcal{V}}$ with the Laplace transform $L_{\mathcal{R}_{\underline{s}}}(-\theta) = \Delta_{-\underline{s}^*}^*(\theta)$ for $\theta \in \Omega_{\mathcal{V}}^*$ if and only if $\underline{s} \in \Xi := \bigsqcup_{\underline{\varepsilon} \in \{0,1\}^r} \Xi(\underline{\varepsilon})$, where

$$\Xi(\underline{\varepsilon}) := \left\{ \underline{s} \in \mathbb{R}^r; \begin{array}{l} s_k > p_k(\underline{\varepsilon})/2 \text{ if } \varepsilon_k = 1 \\ s_k = p_k(\underline{\varepsilon})/2 \text{ if } \varepsilon_k = 0 \end{array} \right\}$$

and $p_k(\underline{\varepsilon}) = \sum_{i < k} \varepsilon_i \dim \mathcal{V}_{ki}$.

The measure $\mathcal{R}_{\underline{s}}$ with $\underline{s} \in \Xi$ has the support in $\overline{\Omega_{\mathcal{V}}}$ and is called the *Riesz measure*, while the set Ξ is called the *Gindikin-Wallach set*.

In order to define NEF generated by $\mathcal{R}_{\underline{s}}$ we have to know if $\mathcal{R}_{\underline{s}} \in \mathcal{M}(\mathcal{Z}_{\mathcal{V}})$, that is, if $\mathcal{R}_{\underline{s}}$ is not concentrated on some affine hyperplane of $\mathcal{Z}_{\mathcal{V}}$. The following result is a generalization of [13, Theorem 3.1]. Let us define for $\underline{\varepsilon} \in \{-1, 0, 1\}^r$,

$$E_{\underline{\varepsilon}} := \begin{pmatrix} \varepsilon_1 I_{n_1} & & \\ & \ddots & \\ & & \varepsilon_r I_{n_r} \end{pmatrix} \in \mathcal{Z}_{\mathcal{V}}$$

Theorem 3.4 The support of $\mathcal{R}_{\underline{s}}$ is not concentrated on any affine hyperplane in $\mathcal{Z}_{\mathcal{V}}$ if and only if $s_k > 0$ for all $k = 1, \dots, r$.

Proof. We write $\mathcal{O}_{\underline{\varepsilon}}$ for the $\rho(H_V)$ -orbit in \mathcal{Z}_V through $E_{\underline{\varepsilon}}$. Note that $\mathcal{O}_{(1,\dots,1)} = \Omega_V$. It is shown in [18, Theorem 6.2] that, if $\underline{s} \in \Xi(\underline{\varepsilon})$, then $\mathcal{R}_{\underline{s}}$ is a positive measure on $\mathcal{O}_{\underline{\varepsilon}}$, so that the support of $\mathcal{R}_{\underline{s}}$ coincides with the closure $\overline{\mathcal{O}_{\underline{\varepsilon}}}$ of $\mathcal{O}_{\underline{\varepsilon}}$. In particular, if for any $k = 1, \dots, r$,

$$(3.2) \quad s_k > p_k(1, \dots, 1)/2 = \frac{1}{2} \sum_{i < k} \dim \mathcal{V}_{ki},$$

then $\underline{s} \in \Xi(1, \dots, 1)$ and $\mathcal{R}_{\underline{s}}$ is a regular measure on the cone $\Omega_V = \rho(H_V)I_N$.

Now we show the ‘if’ part of the statement. Assume that $\underline{s} \in \Xi \cap \mathbb{R}_{>0}^r$. In view of (3.2), we see that there exists a positive integer m such that $m\underline{s} \in \Xi(1, \dots, 1)$. Then $\mathcal{R}_{m\underline{s}}$ is a regular measure, while $\mathcal{R}_{m\underline{s}}$ equals the convolution measure $\mathcal{R}_{\underline{s}} * \mathcal{R}_{\underline{s}} * \dots * \mathcal{R}_{\underline{s}}$ (m times). It follows that the support of $\mathcal{R}_{\underline{s}}$ is not concentrated on any affine hyperplane in \mathcal{Z}_V .

Next we show the ‘only if’ part. It suffices to show that, if $\underline{s} \in \Xi(\underline{\varepsilon})$ with $s_k = 0$ for some k , then $\text{supp } \mathcal{R}_{\underline{s}} = \overline{\mathcal{O}_{\underline{\varepsilon}}}$ is contained in the subspace $(\mathbb{R}E_k)^\perp := \{x \in \mathcal{Z}_V : x_{kk} = 0\}$ of \mathcal{Z}_V . Recalling the definition of $\Xi(\underline{\varepsilon})$, we see that

$$\varepsilon_k = 0 \quad \text{and} \quad p_k(\underline{\varepsilon}) = \sum_{i < k} \varepsilon_i \dim \mathcal{V}_{ki} = 0,$$

and the latter equality implies

$$\mathcal{V}_{ki} = \{0\} \text{ if } \varepsilon_i = 1.$$

Therefore, for any $x = \rho(T)E_{\underline{\varepsilon}} = TE_{\underline{\varepsilon}}T^\top \in \mathcal{O}_{\underline{\varepsilon}}$ with $T \in H_V$, we have

$$x_{kk} = \varepsilon_k(t_{kk})^2 + \sum_{i < k} \varepsilon_i \|T_{ki}\|^2 = 0,$$

which means that $\mathcal{O}_{\underline{\varepsilon}} \subset (\mathbb{R}E_k)^\perp$. Hence $\text{supp } \mathcal{R}_{\underline{s}} \subset (\mathbb{R}E_k)^\perp$ and the proof is completed. \square

For $\underline{\varepsilon} \in \{-1, 1\}^r$, consider $\mathcal{O}_{\underline{\varepsilon}}^* := \rho^*(H_V)E_{\underline{\varepsilon}}$. The set

$$(3.3) \quad \bigsqcup_{\underline{\varepsilon} \in \{-1, 1\}^r} \mathcal{O}_{\underline{\varepsilon}}^*$$

is dense in \mathcal{Z}_V and $\mathcal{O}_{\underline{\varepsilon}}^*$ are the only open orbits of $\rho^*(H_V)$ (see [9, p. 77]).

3.2. Characterization of the Riesz measure on homogeneous cone. In the following section we will give an application of Theorem 2.1 to a characterization of the Riesz measure on homogeneous cone. We generalize the results of [13], where the characterization of Riesz measure through invariance property of NEF on simple Euclidean algebras was considered.

We say that the subalgebra \mathcal{Z} of $(\text{Sym}(N, \mathbb{R}), \triangle)$ is *irreducible* if \mathcal{Z} is not equal to a direct sum of two non-trivial ideals.

Theorem 3.5 *Let $\mathbb{E} = \mathcal{Z}_V$ be an irreducible subalgebra of $(\text{Sym}(N, \mathbb{R}), \triangle)$ that admits a normal block decomposition and let $\mu \in \mathcal{M}(\mathbb{E})$. Assume that $F(\mu)$ is a NEF invariant by $G = \rho(H_V)$. Then there exist $\theta_0 \in \mathcal{Z}_V$, $a_0 \in \mathbb{R}$ and $\underline{s} \in \Xi \cap \mathbb{R}_{>0}^r$ such that*

$$\mu(dx) = e^{a_0 - \langle \theta_0, x \rangle} \mathcal{R}_{\underline{s}}(dx)$$

or

$$\mu(dx) = e^{a_0 - \langle \theta_0, x \rangle} \mathcal{R}_{\underline{s}}(-dx)$$

Thanks to Theorem 3.1, Theorem 3.5 provides a characterization of Riesz measures on homogeneous cones (note that the support of characterized measure is a homogeneous cone).

Proof. We have $\mathbb{R}_{>0}I_N \subset H_V$ and so $c\text{Id} \in \rho(H_V)$ for all $c > 0$. Theorem 2.1 implies that there exists θ_0 such that

$$\mu(dx) = e^{-\langle \theta_0, x \rangle} \mu_0(dx),$$

where (by (2.1))

$$(3.4) \quad L_{\mu_0}(\rho^*(T)\theta) = \chi(T)L_{\mu_0}(\theta), \quad (\theta, T) \in \Theta(\mu_0) \times H_V$$

with a certain character χ of H_V . We will determine L_{μ_0} and $\Theta(\mu_0)$. The proof is split into four steps:

- (i) There exists $\underline{\varepsilon} \in \{-1, 1\}^r$ such that $E_{\underline{\varepsilon}} \in \Theta(\mu_0)$. Moreover, if $E_{\underline{\varepsilon}}, E_{\underline{\varepsilon}'} \in \Theta(\mu_0)$, then $\underline{\varepsilon} = \underline{\varepsilon}'$.
- (ii) $\Theta(\mu_0) = \mathcal{O}_{\underline{\varepsilon}}^*$ for some $\underline{\varepsilon} \in \{-1, 1\}^r$.
- (iii) If $\Theta(\mu_0) = \mathcal{O}_{\underline{\varepsilon}}^*$ then $\underline{\varepsilon} = (-1, \dots, -1)$ or $\underline{\varepsilon} = (1, \dots, 1)$.
- (iv) μ is of the postulated form.

First step. By definition, the set $\Theta(\mu_0)$ is open and non-empty. Since $\rho^*(T)\Theta(\mu_0) = \Theta(\mu_0)$ for any $T \in H_V$ and the set (3.3) is dense in \mathcal{Z}_V , we have $E_{\underline{\varepsilon}} \in \mathcal{O}_{\underline{\varepsilon}}^* \subset \Theta(\mu_0)$ for some $\underline{\varepsilon} \in \{-1, 1\}^r$.

Assume now that $E_{\underline{\varepsilon}}, E_{\underline{\varepsilon}'} \in \Theta(\mu_0)$ and $\underline{\varepsilon} \neq \underline{\varepsilon}'$. Since $\Theta(\mu_0)$ is convex, we know that $\sigma := \frac{1}{2}(E_{\underline{\varepsilon}} + E_{\underline{\varepsilon}'}) \in \Theta(\mu_0)$. Define $I_0 := \{i \in \{1, \dots, r\} : \frac{\varepsilon_i + \varepsilon'_i}{2} = 0\}$. The set I_0 is not empty. Define

$$H_0 := \{T \in H_V : t_{ii} = 1 \text{ for any } i \notin I_0\}$$

If $T \in H_0$ is diagonal, then

$$\rho^*(T)\sigma = \sigma.$$

On the other hand, by (3.4) we obtain

$$L_{\mu_0}(\rho^*(T)\sigma) = \chi(T)L_{\mu_0}(\sigma),$$

which implies that $\chi(T) = 1$ for any $T \in H_0$ (note that $\chi(T)$ depends on T only through diagonal elements; see Lemma 3.1). Further, this implies that for any diagonal $T = \text{diag}(t_{11}, \dots, t_{rr}) \in H_0$,

$$L_{\mu_0}(\rho^*(T)E_{\underline{\varepsilon}}) = L_{\mu_0}(E_{\underline{\varepsilon}}),$$

where $\rho^*(T)E_{\underline{\varepsilon}} = \text{diag}(t_{11}\varepsilon_1, \dots, t_{rr}\varepsilon_r)$. This contradicts the assumption that μ_0 is not supported on any affine hyperplane of \mathcal{Z}_V . Indeed, function $(0, \infty) \ni t_{ii} \mapsto \int \exp(t_{ii}\varepsilon_i x_{ii}) \mu_0(dx)$ is constant if and only if the support of μ_0 is contained in the subspace of \mathcal{Z}_V whose (i, i) -components are zero for any $i \in I_0$.

Second step. Since $\Theta(\mu_0)$ contains only one open orbit, we have

$$\text{int}\{\Theta(\mu_0) \setminus \mathcal{O}_{\underline{\varepsilon}}^*\} = \emptyset.$$

This implies that

$$\Theta(\mu_0) \setminus \overline{\mathcal{O}_{\underline{\varepsilon}}^*} = \emptyset,$$

thus

$$\mathcal{O}_{\underline{\varepsilon}}^* \subset \Theta(\mu_0) \subset \overline{\mathcal{O}_{\underline{\varepsilon}}^*}$$

which proves the claim, since $\Theta(\mu_0)$ is open.

Third step. We will show that $\mathcal{O}_{\underline{\varepsilon}}^*$ is not convex unless all ε_k , $1 \leq k \leq r$, are simultaneously 1 or -1. Define

$$I^+(\underline{\varepsilon}) := \{i \in \{1, \dots, r\} : \varepsilon_i = 1\} \text{ and } I^-(\underline{\varepsilon}) := \{1, \dots, r\} \setminus I^+(\underline{\varepsilon})$$

Suppose that $I^+(\underline{\varepsilon})$ and $I^-(\underline{\varepsilon})$ are non-empty. Then there exists $k \in I^+(\underline{\varepsilon})$ and $l \in I^-(\underline{\varepsilon})$ such that the space \mathcal{V}_{lk} is not $\{0\}$ (without loss of generality we may assume that $l > k$). If not, \mathcal{Z}_V has the block form

$$w \left(\begin{array}{c|c} \mathcal{Z}_+ & 0 \\ \hline 0 & \mathcal{Z}_- \end{array} \right) w^\top,$$

for some permutation matrix w . This contradicts the assumption that \mathcal{Z}_V is irreducible.

Thus there exists $\mathcal{V}_{lk} \neq \{0\}$. We have $\varepsilon_k = 1$ and $\varepsilon_l = -1$. For $v \in \mathcal{V}_{lk}$ let $T(v)$ be the element of $H_{\mathcal{V}}$ such that $T_{lk}(v) = v$, $t_{ii}(v) = 1$ for $i = 1, \dots, r$ and $T_{ji}(v) = 0$ for all $(i, j) \neq (k, l)$, $1 \leq i < j \leq r$. Take any $v \in \mathcal{V}_{lk}$ with $(v|v) = 2$. Then

$$(3.5) \quad \frac{1}{2} (\rho^*(T(v))E_{\underline{\varepsilon}} + \rho^*(T(-v))E_{\underline{\varepsilon}}) = E_{\underline{\varepsilon}'},$$

where $\varepsilon'_i = \varepsilon_i$ for $i \neq k$ and $\varepsilon'_k = -1$. We will use the definition of $\rho^*(T)$, but there is other natural approach; see Remark 3.1 after the proof. For any $x \in \mathcal{Z}_{\mathcal{V}}$ consider the matrix

$$x_v := \frac{1}{2} (\rho(T(v))x + \rho(T(-v))x) = \frac{1}{2} (T(v)xT(v)^{\top} + T(-v)xT(-v)^{\top}).$$

It may be verified by direct calculation that matrices x and x_v differ only on their (l, l) -components, which in the latter case equals

$$(vv^{\top}x_{kk} + x_{ll})I_{n_l} = (2x_{kk} + x_{ll})I_{n_l}.$$

Thus,

$$\begin{aligned} \left\langle \frac{1}{2} (\rho^*(T(v))E_{\underline{\varepsilon}} + \rho^*(T(-v))E_{\underline{\varepsilon}}), x \right\rangle &= \left\langle E_{\underline{\varepsilon}'}, \frac{1}{2} (\rho(T(v))x + \rho(T(-v))x) \right\rangle \\ &= \sum_{i \notin \{k, l\}} x_{ii}\varepsilon_i + x_{kk} + (2x_{kk} + x_{ll})(-1) = \langle E_{\underline{\varepsilon}'}, x \rangle. \end{aligned}$$

But $\Theta(\mu_0)$ is convex, thus (3.5) implies that $E_{\underline{\varepsilon}'} \in \Theta(\mu_0)$. This contradicts the point (i). This means that $I^+(\underline{\varepsilon}) = \{1, \dots, r\}$ or $I^-(\underline{\varepsilon}) = \{1, \dots, r\}$ and so $-I_N$ or I_N belongs to $\Theta(\mu_0)$. Finally, by (ii) and the fact that $\rho^*(H_{\mathcal{V}})I_N = \Omega_{\mathcal{V}}^*$,

$$\Theta(\mu_0) = -\Omega_{\mathcal{V}}^* \quad \text{or} \quad \Theta(\mu_0) = \Omega_{\mathcal{V}}^*.$$

Fourth step. Let us first consider the case $\Theta(\mu_0) = -\Omega_{\mathcal{V}}^*$. Putting $\theta = \rho^*(S)(-I_N)$ for $S \in H_{\mathcal{V}}$, we obtain

$$L_{\mu_0}(-\rho^*(ST)I_N) = \chi(T)L_{\mu_0}(-\rho^*(S)I_N), \quad (S, T) \in H_{\mathcal{V}}^2,$$

which implies that for $a_0 = \log L_{\mu_0}(-I_N)$,

$$L_{\mu_0}(-\rho^*(T)I_N) = e^{a_0}\chi(T)$$

and $\chi(T)$ is a one-dimensional representation of $H_{\mathcal{V}}$. Thus, by Lemma 3.1, there exists $\underline{s} \in \Xi$ such that $\chi = \chi_{-\underline{s}}$ and then, by the definition of $\Delta_{\underline{s}}^*$,

$$L_{\mu_0}(-\theta) = e^{a_0}\Delta_{-\underline{s}}^*(\theta), \quad \theta \in \Omega_{\mathcal{V}}^* = -\Theta(\mu_0).$$

By Theorems 3.3 and 3.4 we see that $\underline{s} \in \Xi \cap \mathbb{R}_{>0}^r$ and that $\mu_0(dx) = e^{a_0}\mathcal{R}_{\underline{s}}(dx)$ and $\Theta(\mu_0) = -\Omega_{\mathcal{V}}^*$.

In the second case, when $\Theta(\mu_0) = \Omega_{\mathcal{V}}^*$, one shows similarly that $\mu_0(dx) = e^{a_0}\mathcal{R}_{\underline{s}}(-dx)$. \square

Remark 3.1 To show (3.5) we could switch to a matrix realization of the dual cone $\Omega_{\mathcal{V}}^*$, where (under suitable linear isomorphism) $\rho^*(T)$ is just a multiplication by some upper triangular matrix on the left and its transpose on the right.

3.3. Comments.

- (1) In [14], the authors considered characterization of the Riesz measure $\mathcal{R}_{\underline{s}}$ through the invariance property of NEF on a simple Euclidean algebra \mathbb{E} by some subgroup G of $GL(\mathbb{E})$. This subgroup was carefully chosen in order to ensure that some components of vector $\underline{s} \in \Xi$ are equal. Taking ordinary triangular group $\rho(H_{\mathcal{V}})$ imposes no additional conditions on these components. On the other hand if one considers the invariance property of NEF by the connected component containing identity of $G(\Omega)$ (this is in principle what Letac did on symmetric matrices in [25], but it is true on homogeneous cones also; see Comment (4) below), then all s_i have to be equal,

- $s_i = p$ for $1 \leq i \leq r$. Then $\Delta_{\underline{s}^*}^* = \det^p$ and p belongs to the set Λ called the Jorgensen set (see [6]).
- (2) Elements of $F(\mathcal{R}_{\underline{s}})$ are actually the Wishart distributions on homogeneous cones introduced in [1] (the subcase of $s_i = p$ for $1 \leq i \leq r$) and in [11].
 - (3) It should be stressed that our approach is very different from [13, 25], where the characterization of NEF was proved by showing that the variance function of $\rho(H_V)$ -invariant NEF coincides with the one of $F(\mathcal{R}_{\underline{s}})$ for some $\underline{s} \in \Xi$. In the present paper we do not even need to know what is the variance function of the Riesz measure on homogeneous cones. We perceive our approach as less technical and more natural.
 - (4) We can rephrase our result in terms of theory of homogeneous cones as follows. Let $\Omega \subset \mathbb{E}$ be an irreducible homogeneous cone, and $G \subset GL(\mathbb{E})$ a linear algebraic group acting on Ω transitively. Then we conclude that any G -invariant NEF is generated by a Riesz distributions on Ω or $-\Omega$ because G contains a triangular subgroup isomorphic to the group $\rho(H_V)$ discussed in the present paper (cf. [32, Chapter 1, Section 9]).

Acknowledgements. We would like to thank the anonymous referee for helpful comments. B. Kołodziejek was partially supported by NCN Grant No. 2012/05/B/ST1/00554. H. Ishi was partially supported by JSPS KAKENHI Grant Number 16K05174 and JST PRESTO.

REFERENCES

- [1] S. A. Andersson and G. G. Wojnar. Wishart distributions on homogeneous cones. *J. Theoret. Probab.*, 17(4):781–818, 2004.
- [2] O. Barndorff-Nielsen. *Information and exponential families in statistical theory*. Wiley Series in Probability and Statistics. John Wiley & Sons, Ltd., Chichester, 2014.
- [3] M. Casalis. Familles exponentielles naturelles sur \mathbb{R}^d invariantes par un groupe. *Int. Stat. Rev.*, 59(2):241–262, 1991.
- [4] M. Casalis. Les familles exponentielles à variance quadratique homogène sont des lois de Wishart sur un cône symétrique. *C. R. Acad. Sci. Paris Sér. I Math.*, 312(7):537–540, 1991.
- [5] M. Casalis. The $2d + 4$ simple quadratic natural exponential families on \mathbf{R}^d . *Ann. Statist.*, 24(4):1828–1854, 1996.
- [6] M. Casalis and G. Letac. Characterization of the Jørgensen set in generalized linear models. *Test*, 3(1):145–162, 1994.
- [7] P. S. Eriksen. $(k, 1)$ exponential transformation models. *Scand. J. Statist.*, 11(3):129–145, 1984.
- [8] G. Garrigós. Generalized Hardy spaces on tube domains over cones. *Colloq. Math.*, 90(2):213–251, 2001.
- [9] S. G. Gindikin. Analysis in homogeneous domains. *Russian Math. Surveys*, 19:1–89, 1964.
- [10] S. G. Gindikin. Invariant generalized functions in homogeneous domains. *Funkcional. Anal. i Prilozhen.*, 9(1):56–58, 1975.
- [11] P. Graczyk and H. Ishi. Riesz measures and Wishart laws associated to quadratic maps. *J. Math. Soc. Japan*, 66(1):317–348, 2014.
- [12] P. Graczyk, H. Ishi and B. Kołodziejek. Wishart laws and variance function on homogeneous cones. to appear in *Probab. Math. Statist.*, pages 1–25, 2019.
- [13] A. Hassairi and S. Lajmi. Riesz exponential families on symmetric cones. *J. Theoret. Probab.*, 14(4):927–948, 2001.
- [14] A. Hassairi and S. Lajmi. Classification of Riesz exponential families on a symmetric cone by invariance properties. *J. Theoret. Probab.*, 17(3):521–539, 2004.
- [15] A. Hassairi and M. Zarai. Characterization of the cubic exponential families by orthogonality of polynomials. *Ann. Probab.*, 32(3B):2463–2476, 2004.

- [16] A. Hassairi and M. Zarai. Characterization of the simple cubic multivariate exponential families. *J. Funct. Anal.*, 235(1):69–89, 2006.
- [17] S. Højsgaard and S. L. Lauritzen. Graphical Gaussian models with edge and vertex symmetries. *J. R. Stat. Soc. Ser. B Stat. Methodol.*, 70(5):1005–1027, 2008.
- [18] H. Ishi. Positive Riesz distributions on homogeneous cones. *J. Math. Soc. Japan*, 52(1):161–186, 2000.
- [19] H. Ishi. On symplectic representations of normal j -algebras and their application to Xu’s realizations of Siegel domains. *Differential Geom. Appl.*, 24(6):588–612, 2006.
- [20] H. Ishi. Representation of clans and homogeneous cones. *Vestn. Tambov. Univ.*, 16:1669–1675, 2011.
- [21] H. Ishi. Homogeneous cones and their applications to statistics. In *Modern methods of multivariate statistics*, volume 82, pages 135–154. Hermann, 2014.
- [22] H. Ishi. Matrix realization of a homogeneous cone. In *Geometric Science of Information*, volume 9389 of *Lecture Notes in Computer Science*, pages 248–256. Springer International Publishing, 2015.
- [23] J. L. Jensen. On the hyperboloid distribution. *Scand. J. Statist.*, 8(4):193–206, 1981.
- [24] B. Kołodziejek. Multiplicative Cauchy functional equation on symmetric cones. *Aequationes Math.*, 89(4):1075–1094, 2015.
- [25] G. Letac. A characterization of the Wishart exponential families by an invariance property. *J. Theoret. Probab.*, 2(1):71–86, 1989.
- [26] G. Letac. *Lectures on natural exponential families and their variance functions*, volume 50 of *Monografías de Matemática [Mathematical Monographs]*. Instituto de Matemática Pura e Aplicada (IMPA), Rio de Janeiro, 1992.
- [27] G. Letac and H. Massam. Wishart distributions for decomposable graphs. *Ann. Statist.*, 35(3):1278–1323, 2007.
- [28] G. Letac and M. Mora. Natural real exponential families with cubic variance functions. *Ann. Statist.*, 18(1):1–37, 1990.
- [29] C. N. Morris. Natural exponential families with quadratic variance functions. *Ann. Statist.*, 10(1):65–80, 1982.
- [30] H. Shima. Homogeneous Hessian manifolds. *Ann. Inst. Fourier (Grenoble)*, 30(3):91–128, 1980.
- [31] H. Shima. *The geometry of Hessian structures*. World Scientific Publishing Co. Pte. Ltd., Hackensack, NJ, 2007.
- [32] È. B. Vinberg. The theory of homogeneous convex cones. *Tr. Mosk. Mat. Obs.*, 12:303–358, 1963.
- [33] G. S. Watson. *Statistics on spheres*. University of Arkansas Lecture Notes in the Mathematical Sciences, 6. John Wiley & Sons, Inc., New York, 1983. A Wiley-Interscience Publication.

NAGOYA UNIVERSITY, GRADUATE SCHOOL OF MATHEMATICS, FURO-CHO, NAGOYA 464-8602, JAPAN /JST PRESTO, 4-1-8 HONCHO, KAWAGUCHI 332-0012, JAPAN.

E-mail address: `hideyuki@math.nahoya-u.ac.jp`

WARSAW UNIVERSITY OF TECHNOLOGY, FACULTY OF MATHEMATICS AND INFORMATION SCIENCE, KOSZYKOWA 75, 00-662 WARSAW, POLAND

E-mail address: `b.kolodziejek@mini.pw.edu.pl`