

# MODULAR CURVES OF PRIME-POWER LEVEL WITH INFINITELY MANY RATIONAL POINTS

ANDREW V. SUTHERLAND AND DAVID ZYWINA

ABSTRACT. For each open subgroup  $G$  of  $\mathrm{GL}_2(\hat{\mathbb{Z}})$  containing  $-I$  with full determinant, let  $X_G/\mathbb{Q}$  denote the modular curve that loosely parametrizes elliptic curves whose Galois representation, which arises from the Galois action on its torsion points, has image contained in  $G$ . Up to conjugacy, we determine a complete list of the 248 such groups  $G$  of prime power level for which  $X_G(\mathbb{Q})$  is infinite. For each  $G$ , we also construct explicit maps from each  $X_G$  to the  $j$ -line. This list consists of 220 modular curves of genus 0 and 28 modular curves of genus 1. For each prime  $\ell$ , these results provide an explicit classification of the possible images of  $\ell$ -adic Galois representations arising from elliptic curves over  $\mathbb{Q}$  that is complete except for a finite set of exceptional  $j$ -invariants.

## 1. INTRODUCTION

Let  $E$  be an elliptic curve defined over  $\mathbb{Q}$  and denote its  $j$ -invariant by  $j_E$ . For each positive integer  $N$ , let  $E[N]$  denote the  $N$ -torsion subgroup of  $E(\overline{\mathbb{Q}})$ , where  $\overline{\mathbb{Q}}$  is a fixed algebraic closure of  $\mathbb{Q}$ . The natural action of the absolute Galois group  $\mathrm{Gal}_{\mathbb{Q}} := \mathrm{Gal}(\overline{\mathbb{Q}}/\mathbb{Q})$  on  $E[N] \simeq (\mathbb{Z}/N\mathbb{Z})^2$  induces a Galois representation

$$\rho_{E,N}: \mathrm{Gal}_{\mathbb{Q}} \rightarrow \mathrm{GL}_2(\mathbb{Z}/N\mathbb{Z}).$$

After choosing compatible bases for the torsion subgroups  $E[N]$ , these representations determine a Galois representation

$$\rho_E: \mathrm{Gal}_{\mathbb{Q}} \rightarrow \mathrm{GL}_2(\hat{\mathbb{Z}})$$

whose composition with the projection  $\mathrm{GL}_2(\hat{\mathbb{Z}}) \rightarrow \mathrm{GL}_2(\mathbb{Z}/N\mathbb{Z})$  given by reduction modulo  $N$  is equal to  $\rho_{E,N}$ , for each  $N$ . The images of  $\rho_{E,N}$  and  $\rho_E$  are uniquely determined up to conjugacy in  $\mathrm{GL}_2(\mathbb{Z}/N\mathbb{Z})$  and  $\mathrm{GL}_2(\hat{\mathbb{Z}})$ , respectively. If  $E$  does not have complex multiplication (CM), then  $\rho_E(\mathrm{Gal}_{\mathbb{Q}})$  is an open subgroup of  $\mathrm{GL}_2(\hat{\mathbb{Z}})$ , by Serre's open image theorem [Ser72], hence of finite index in  $\mathrm{GL}_2(\hat{\mathbb{Z}})$ .

Let  $G$  be an open subgroup of  $\mathrm{GL}_2(\hat{\mathbb{Z}})$  that satisfies  $\det(G) = \hat{\mathbb{Z}}^\times$  and  $-I \in G$ . Let  $N$  be the least positive integer such that  $G$  is the inverse image of its image under the reduction map  $\mathrm{GL}_2(\hat{\mathbb{Z}}) \rightarrow \mathrm{GL}_2(\mathbb{Z}/N\mathbb{Z})$ ; we call  $N$  the *level* of  $G$ .

Associated to  $G$  is a modular curve  $X_G/\mathbb{Q}$ ; one can define  $X_G$  as the generic fiber of the smooth proper  $\mathbb{Z}[1/N]$ -scheme that is the coarse moduli space for the algebraic stack  $\mathcal{M}_{\overline{G}}[1/N]$  in the sense of [DR73, §IV], where  $\overline{G}$  denotes the image of  $G$  under reduction modulo  $N$ . See §2 for some background on  $X_G$  and an alternate description; in particular, it is a smooth projective geometrically integral curve defined over  $\mathbb{Q}$ .

When  $G = \mathrm{GL}_2(\hat{\mathbb{Z}})$ , the modular curve  $X_G$  is the  $j$ -line  $\mathbb{P}_{\mathbb{Q}}^1 = \mathbb{A}_{\mathbb{Q}}^1 \cup \{\infty\}$ . If  $G$  and  $G'$  are open subgroups of  $\mathrm{GL}_2(\hat{\mathbb{Z}})$  with  $\det(G) = \det(G') = \hat{\mathbb{Z}}^\times$  and  $-I \in G, G'$  such that  $G \subseteq G'$ , then there is a natural morphism  $X_G \rightarrow X_{G'}$  of degree  $[G' : G]$ . In particular, with  $G' = \mathrm{GL}_2(\hat{\mathbb{Z}})$ , we have a

---

2000 *Mathematics Subject Classification*. Primary 14G35; Secondary 11G05, 11F80.

The first author was supported by NSF grants DMS-1115455 and DMS-1522526.

morphism

$$\pi_G: X_G \rightarrow \mathbb{P}_{\mathbb{Q}}^1 = \mathbb{A}_{\mathbb{Q}}^1 \cup \{\infty\}$$

of degree  $[\mathrm{GL}_2(\hat{\mathbb{Z}}) : G]$  from  $X_G$  to the  $j$ -line.

The key property for our applications is that for an elliptic curve  $E/\mathbb{Q}$  with  $j_E \notin \{0, 1728\}$ , the group  $\rho_E(\mathrm{Gal}_{\mathbb{Q}})$  is conjugate in  $\mathrm{GL}_2(\hat{\mathbb{Z}})$  to a subgroup of  $G$  if and only if  $j_E$  is an element of  $\pi_G(X_G(\mathbb{Q}))$ ; see Proposition 2.7. This property requires  $-I \in G$ , since there is always an elliptic curve  $E$  with any given rational  $j$ -invariant such that  $-I \in \rho_E(\mathrm{Gal}_{\mathbb{Q}})$ ; it also requires  $\det(G) = \hat{\mathbb{Z}}^{\times}$ , since  $\det(\rho_E(\mathrm{Gal}_{\mathbb{Q}})) = \hat{\mathbb{Z}}^{\times}$ , and that  $G$  contain an element corresponding to complex conjugation.

We are interested in those groups  $G$  for which  $X_G$  has infinitely many rational points; equivalently, for which there are infinitely many elliptic curves  $E/\mathbb{Q}$ , with distinct  $j$ -invariants, such that  $\rho_E(\mathrm{Gal}_{\mathbb{Q}})$  is conjugate to a subgroup of  $G$ . We need only consider modular curves  $X_G$  of genus 0 or 1 since otherwise  $X_G(\mathbb{Q})$  is finite by Faltings' Theorem [Fal83].

In this article, we give an explicit description of all such subgroups  $G \subseteq \mathrm{GL}_2(\hat{\mathbb{Z}})$  for which the modular curve  $X_G$  has infinitely many rational points in the special case where the level  $N$  of  $G$  is a *prime power*; we also give an explicit model for  $X_G$  and the morphism  $\pi_G$ . We need only describe the groups  $G$  up to conjugacy in  $\mathrm{GL}_2(\hat{\mathbb{Z}})$ . For notational simplicity, we define the genus of  $G$  to be the genus of the corresponding curve  $X_G$ .

**Theorem 1.1.** *Up to conjugacy, there are 248 open subgroups  $G$  of  $\mathrm{GL}_2(\hat{\mathbb{Z}})$  of prime power level satisfying  $-I \in G$  and  $\det(G) = \hat{\mathbb{Z}}^{\times}$  for which  $X_G$  has infinitely many rational points. Of these 248 groups, there are 220 of genus 0 and 28 of genus 1.*

The 220 subgroups of genus 0 in Theorem 1.1 are given in Tables 1, 2 and 3 in Appendix A. For such a group  $G$  of genus 0, we also describe the morphism  $\pi_G$ . More precisely, we give a rational function  $J(t) \in \mathbb{Q}(t)$  such that the function field of  $X_G$  is of the form  $\mathbb{Q}(t)$  and the morphism from  $X_G$  to the  $j$ -line is given by the equation  $j = J(t)$ . In particular, if  $E/\mathbb{Q}$  is an elliptic curve with  $j_E \notin \{0, 1728\}$ , then  $\rho_E(\mathrm{Gal}_{\mathbb{Q}})$  is conjugate to a subgroup of  $G$  if and only if  $j_E = J(t_0)$  for some  $t_0 \in \mathbb{Q} \cup \{\infty\}$ .

The 28 subgroups of genus 1 in Theorem 1.1 are listed in Table 4 of Appendix A; their levels are all powers of 2 except for a group of level 11 whose image in  $\mathrm{GL}_2(\mathbb{Z}/11\mathbb{Z})$  is the normalizer of a non-split Cartan subgroup. For such a group  $G$  of genus 1, we give a Weierstrass model for  $X_G$  and the morphism  $\pi_G$  to the  $j$ -line.

**Example 1.2.** Up to conjugacy, there is a unique subgroup  $G \subseteq \mathrm{GL}_2(\hat{\mathbb{Z}})$  of genus 0 and level 27 given by Theorem 1.1. It has label 27A<sup>0</sup>-27a in our classification, and we may choose it so that the image of  $G$  in  $\mathrm{GL}_2(\mathbb{Z}/27\mathbb{Z})$  is generated by the matrices  $\begin{pmatrix} 1 & 1 \\ 0 & 1 \end{pmatrix}$ ,  $\begin{pmatrix} 2 & 1 \\ 9 & 5 \end{pmatrix}$  and  $\begin{pmatrix} 1 & 2 \\ 3 & 2 \end{pmatrix}$ . Using Table 1, associated to  $G$  is the rational function

$$J(t) = F_3(F_2(F_1(t))) = \frac{(t^3 + 3)^3(t^9 + 9t^6 + 27t^3 + 3)^3}{t^3(t^6 + 9t^3 + 27)},$$

where  $F_1(t) = t^3$ ,  $F_2(t) = t(t^2 + 9t + 27)$  and  $F_3(t) = (t+3)^3(t+27)/t$ . That  $J(t)$  is the composition of three rational functions reflects the fact that the morphism  $\pi_G$  factors as  $X_G \rightarrow X_{G'} \rightarrow X_{G''} \rightarrow \mathbb{P}_{\mathbb{Q}}^1$  for some groups  $G \subsetneq G' \subsetneq G'' \subsetneq \mathrm{GL}_2(\hat{\mathbb{Z}})$ . The groups  $G'$  and  $G''$  have labels 9B<sup>0</sup>-9a and 3B<sup>0</sup>-3a, respectively, and can also be found in Table 1.

*Remark 1.3.* In contrast to the case of prime power level, in general there are infinitely many open subgroups  $G$  of  $\mathrm{GL}_2(\hat{\mathbb{Z}})$  satisfying  $-I \in G$  and  $\det(G) = \hat{\mathbb{Z}}^{\times}$  for which the modular curve  $X_G$  has infinitely many rational points. Let us explicitly construct just one of several infinite families of such groups  $G$ .

Let  $D$  be the discriminant of a quadratic number field and let  $\chi_D: \hat{\mathbb{Z}}^\times \rightarrow \{\pm 1\}$  be the continuous quadratic character arising from the corresponding Dirichlet character. Let  $\varepsilon: \mathrm{GL}_2(\hat{\mathbb{Z}}) \rightarrow \{\pm 1\}$  be the character obtained by composing the reduction map  $\mathrm{GL}_2(\hat{\mathbb{Z}}) \rightarrow \mathrm{GL}_2(\mathbb{Z}/2\mathbb{Z})$  with the unique non-trivial homomorphism  $\mathrm{GL}_2(\mathbb{Z}/2\mathbb{Z}) \rightarrow \{\pm 1\}$ . Define the group

$$G_D := \{A \in \mathrm{GL}_2(\hat{\mathbb{Z}}) : \varepsilon(A) = \chi_D(\det(A))\};$$

it is an open subgroup of  $\mathrm{GL}_2(\hat{\mathbb{Z}})$  of index 2 containing  $-I$  with  $\det(G_D) = \hat{\mathbb{Z}}^\times$  whose level is  $|D|$  or  $2|D|$ , depending on whether  $D \equiv 0 \pmod{4}$  or  $D \equiv 1 \pmod{4}$ . For  $D \neq D'$ , the groups  $G_D$  and  $G_{D'}$  are not conjugate in  $\mathrm{GL}_2(\hat{\mathbb{Z}})$ .

The modular curve  $X_{G_D}$  has genus 0 and a rational point (it has a unique, hence rational, cusp); the function field of  $X_{G_D}$  is of the form  $\mathbb{Q}(t)$  with the map to the  $j$ -line given by  $J(t) = Dt^2 + 1728$ . Each  $X_{G_D}$  is a  $\mathbb{Q}(\sqrt{D})$ -twist of the modular curve  $X_G$  corresponding to the unique index 2 subgroup  $G \subseteq \mathrm{GL}_2(\hat{\mathbb{Z}})$  whose reduction has index 2 in  $\mathrm{GL}_2(\mathbb{Z}/2\mathbb{Z})$ ; it has label  $2A^0-2a$  in our classification and can be found in Table 3, along with its map to the  $j$ -line, which is  $J(t) = t^2 + 1728$ .

In general, if  $\Gamma \subseteq \mathrm{SL}_2(\mathbb{Z})$  is a fixed congruence subgroup of level  $N$  and index  $m$  containing  $-I$ , there will be infinitely many non-conjugate open subgroups  $G \subseteq \mathrm{GL}_2(\hat{\mathbb{Z}})$  of index  $M$  containing  $-I$  with  $\det(G) = \hat{\mathbb{Z}}^\times$  whose reductions modulo  $N$  coincide with that of  $\Gamma$  upon intersection with  $\mathrm{SL}_2(\mathbb{Z}/N\mathbb{Z})$ . The levels  $M$  of these groups  $G$  may be arbitrarily large multiples of  $N$  (and divisible by arbitrarily large primes). The corresponding modular curves  $X_G/\mathbb{Q}$  are non-isomorphic, but for each  $X_G$  there is a cyclotomic field  $\mathbb{Q}(\zeta_M)$  over which  $X_G$  becomes isomorphic to the modular curve  $X_\Gamma/\mathbb{Q}(\zeta_N)$  (the quotient of the extended upper half plane by the action of  $\Gamma$ ) after base change; as in our example, the  $X_G$  form an infinite family of twists.

1.1.  **$\ell$ -adic representations.** Fix a prime  $\ell$ . Define the set

$$\mathcal{J}_\ell := \bigcup_G (\pi_G(X_G(\mathbb{Q})) \cap \mathbb{Q})$$

of rational numbers, where  $G$  varies over the open subgroups of  $\mathrm{GL}_2(\hat{\mathbb{Z}})$  whose level is a power of  $\ell$  and satisfies  $-I \in G$  and  $\det(G) = \hat{\mathbb{Z}}^\times$ , and for which  $X_G(\mathbb{Q})$  is finite. Note that the set  $\mathcal{J}_\ell$  contains the 13  $j$ -invariants of CM elliptic curves over  $\mathbb{Q}$ : for  $n \geq 1$  each CM  $j$ -invariant corresponds to points on at least one of the modular curves  $X_S^+(\ell^n)$ ,  $X_{\mathrm{ns}}^+(\ell^n)$ ,  $X_0(\ell^n)$ , and for sufficiently large  $n$  these curves have genus at least 2, hence finitely many rational points (by Faltings' Theorem).

For an elliptic curve  $E/\mathbb{Q}$ , let

$$\rho_{E,\ell^\infty}: \mathrm{Gal}_{\mathbb{Q}} \rightarrow \mathrm{GL}_2(\mathbb{Z}_\ell)$$

be the representation describing the Galois action on the  $\ell$ -power torsion points; it is the composition of  $\rho_E$  with the natural projection  $\mathrm{GL}_2(\hat{\mathbb{Z}}) \rightarrow \mathrm{GL}_2(\mathbb{Z}_\ell)$ . After excluding a finite number of  $j$ -invariants, we will describe the possible images of the  $\ell$ -adic representation arising from elliptic curves over  $\mathbb{Q}$ . Denote by  $\pm\rho_{E,\ell^\infty}(\mathrm{Gal}_{\mathbb{Q}})$  the group generated by  $-I$  and  $\rho_{E,\ell^\infty}(\mathrm{Gal}_{\mathbb{Q}})$ .

The following theorem describes the possibilities for  $\pm\rho_{E,\ell^\infty}(\mathrm{Gal}_{\mathbb{Q}})$ , up to conjugacy, when  $j_E$  is not in the (finite!) set  $\mathcal{J}_\ell$ .

**Theorem 1.4.**

- (i) *The set  $\mathcal{J}_\ell$  is finite.*
- (ii) *If  $E/\mathbb{Q}$  is an elliptic curve with  $j_E \notin \mathcal{J}_\ell$ , then  $\pm\rho_{E,\ell^\infty}(\mathrm{Gal}_{\mathbb{Q}})$  is conjugate in  $\mathrm{GL}_2(\mathbb{Z}_\ell)$  to the  $\ell$ -adic projection of a unique group  $G$  from Theorem 1.1 with  $\ell$ -power level. Moreover,  $G$  does not have genus 1, level 16, and index 24 in  $\mathrm{GL}_2(\hat{\mathbb{Z}})$ .*
- (iii) *Let  $G$  be a group from Theorem 1.1 with  $\ell$ -power level that does not have genus 1, level 16, and index 24 in  $\mathrm{GL}_2(\hat{\mathbb{Z}})$ . Then there are infinitely many elliptic curves  $E/\mathbb{Q}$ , with distinct  $j$ -invariants, such that  $\pm\rho_{E,\ell^\infty}(\mathrm{Gal}_{\mathbb{Q}})$  is conjugate in  $\mathrm{GL}_2(\mathbb{Z}_\ell)$  to the  $\ell$ -adic projection of  $G$ .*

*Remark 1.5.*

- (i) Serre has asked whether  $\rho_{E,\ell}$  is surjective for all non-CM elliptic curves  $E/\mathbb{Q}$  and all primes  $\ell > 37$ , cf. [Ser81, p. 399]. For  $\ell > 37$ , this would imply that the set  $\mathcal{J}_\ell$  consists of only the 13  $j$ -invariants of CM elliptic curves over  $\mathbb{Q}$ .
- (ii) It would be nice to explicitly know the finite sets  $\mathcal{J}_\ell$ ; the proof that  $\mathcal{J}_\ell$  is finite relies on [Zyw15b], which is ineffective since it applies Faltings' Theorem several times.

Theorem 1.4 describes the subgroups of  $\mathrm{GL}_2(\mathbb{Z}_\ell)$ , up to conjugacy, that occur as  $\pm\rho_{E,\ell^\infty}(\mathrm{Gal}_{\mathbb{Q}})$  for infinitely many elliptic curves  $E/\mathbb{Q}$  with distinct  $j$ -invariants.

Theorem 1.4 also allows us to determine the subgroups of  $\mathrm{GL}_2(\mathbb{Z}_\ell)$ , up to conjugacy, that occur as  $\rho_{E,\ell^\infty}(\mathrm{Gal}_{\mathbb{Q}})$  for infinitely many elliptic curves  $E/\mathbb{Q}$  with distinct  $j$ -invariants. They are precisely the subgroups  $H$  of the  $\ell$ -adic projection  $G$  of a group from Theorem 1.4 with  $\ell$ -power level such that  $\pm H = G$ . Indeed if  $G = \pm\rho_{E,\ell^\infty}(\mathrm{Gal}_{\mathbb{Q}})$ , then for any such  $H$  there is a quadratic twist of  $E$  such that  $H$  is conjugate to  $\rho_{E',\ell^\infty}(\mathrm{Gal}_{\mathbb{Q}})$ , cf. [Zyw15a, §5.1] and [Sut16, §5.6]; when  $H$  is properly contained in  $G$  this quadratic twist is unique up to isomorphism and can be explicitly determined.

**Corollary 1.6.** *For  $\ell = 2, 3, 5, 7, 11, 13$  there are respectively 1201, 47, 23, 15, 2, 11 subgroups  $H$  of  $\mathrm{GL}_2(\mathbb{Z}_\ell)$  that arise as  $\rho_{E,\ell^\infty}(\mathrm{Gal}_{\mathbb{Q}})$  for infinitely many elliptic curves  $E/\mathbb{Q}$  with distinct  $j$ -invariants; for  $\ell > 13$  the only such subgroup is  $H = \mathrm{GL}_2(\mathbb{Z}_\ell)$ .*

A list of the groups  $H$  appearing in Corollary 1.6 can be found in electronic form at [SZ16].

**1.2. Overview.** We now give a brief overview of the contents of this paper. As already noted, the groups  $G$  from Theorem 1.1, along with the corresponding modular curves  $X_G$  and morphisms  $\pi_G$ , can be found in Appendix A

In §2, we review the background material we need concerning the modular curves  $X_G$ . If  $G$  has level  $N$ , then we can identify the function field of  $X_G$  with a subfield of the field  $\mathcal{F}_N$  of modular functions on  $\Gamma(N)$  whose Fourier coefficients lie in the cyclotomic field  $\mathbb{Q}(\zeta_N)$ . As a working definition of  $X_G$ , we define it in terms of its function field.

In §3, we determine up to conjugacy the open subgroups  $G$  of  $\mathrm{GL}_2(\hat{\mathbb{Z}})$  with genus at most 1 that satisfy  $\det(G) = \hat{\mathbb{Z}}^\times$ ,  $-I \in G$  and contain an element that “looks like complex conjugation”; this last condition is necessary, since otherwise  $X_G(\mathbb{R})$ , and therefore  $X_G(\mathbb{Q})$ , is empty. We are left with 220 groups of genus 0 and 250 groups of genus 1 that include all the groups that appear in Theorem 1.1. These computations make use of the tables of Cummins and Pauli of congruence subgroups of low genus [CP03].

Let  $\Gamma$  be a congruence subgroup of  $\mathrm{SL}_2(\mathbb{Z})$  and let  $X_\Gamma$  be the smooth compact Riemann surface obtained by taking the quotient of the complex upper-half plane by  $\Gamma$  and adjoining cusps. Assume further that  $X_\Gamma$  has genus 0. In §4, we describe how to explicitly construct a *hauptmodul* for  $\Gamma$ ; it is a meromorphic function  $h$  on  $X_\Gamma$  that has a unique pole at the cusp at  $\infty$ . We describe  $h$  in terms of *Siegel functions*; its Fourier coefficients are computable and lie in the field  $\mathbb{Q}(\zeta_N) \subseteq \mathbb{C}$ .

In §5, we prove the part of Theorem 1.1 concerning genus 0 groups. Let  $G$  be one of the genus 0 groups from §3 and let  $J(t) \in \mathbb{Q}(t)$  be the corresponding rational function from Appendix A. We need to verify that the function field  $\mathbb{Q}(X_G)$  of  $X_G$  is of the form  $\mathbb{Q}(f)$ , for some modular function  $f$  for which  $J(f)$  coincides with the modular  $j$ -function. Using our work in §4, we can construct an explicit modular function  $h$  such that  $\mathbb{Q}(\zeta_N)(X_G) = \mathbb{Q}(\zeta_N)(h)$ , along with a rational function  $J'(t) \in \mathbb{Q}(\zeta_N)(t)$  such that  $J'(h) = j$ . The function  $f$  must satisfy  $f = \psi(h)$  for some  $\psi(t) \in \mathbb{Q}(\zeta_N)(t)$  of degree 1, and therefore  $J'(h) = j = J(f) = J(\psi(h))$ ; this in turn implies that  $J'(t) = J(\psi(t))$ . We then directly test all the modular functions  $f := \psi(h)$ , where  $\psi(t) \in \mathbb{Q}(\zeta_N)(t)$  is one of the finitely many degree 1 rational functions that satisfy  $J'(t) = J(\psi(t))$ .

In §6, we prove the part of Theorem 1.1 concerning genus 1 groups. Let  $G$  be one of the genus 1 groups from §3. One can show that  $X_G$  has good reduction at all primes  $p \nmid N$  and its modular

interpretation gives a way to compute  $\#X_G(\mathbb{F}_p)$  directly from the group  $G$ , without requiring an explicit model. By computing  $\#X_G(\mathbb{F}_p)$  for enough primes  $p \nmid N$ , one can determine the Jacobian  $J_G$  of  $X_G$  up to isogeny. This allows us to compute the rank of  $J_G(\mathbb{Q})$  which is an isogeny invariant of  $J_G$ . We need only consider groups for which  $J_G(\mathbb{Q})$  has positive rank since otherwise  $X_G(\mathbb{Q})$  is finite; this leaves the 28 genus 1 groups in Theorem 1.1. These 28 groups  $G$  of genus 1 and a description of their morphisms  $\pi_G$  already appear in the literature; our contribution lies in proving that there are no others.

In §7, we complete the proof of Theorem 1.4, and in §8 we explain how we found the rational functions  $J(t) \in \mathbb{Q}(t)$  whose verification is described in §5.

Appendix A lists the 248 groups  $G$  that appear in Theorem 1.1, along with explicit maps from  $X_G$  to the  $j$ -line; for the 220 groups of genus 0 these are rational functions  $J(t)$ , and for the 28 groups of genus 1 these are morphisms  $J(x, y)$  from an explicit Weierstrass model for  $X_G$  as an elliptic curve of positive rank. One can use these maps to explicitly construct infinite families of elliptic curves  $E/\mathbb{Q}$  with distinct  $j$ -invariants whose  $\ell$ -adic Galois images match the groups  $G$  listed in Theorem 1.4 and the groups  $H$  listed in Corollary 1.6 by choosing appropriate quadratic twists.

**1.3. Related results.** Contemporaneous with our work, Rouse and Zureick-Brown independently computed explicit models for all modular curves  $X_G/\mathbb{Q}$  of 2-power level that have a non-cuspidal rational point, including all those for which  $X_G(\mathbb{Q})$  is infinite [RZB15]. The  $X_G$  of 2-power level in our list agree with theirs, although we generally obtain different (but isomorphic) models (N.B. our groups are transposed relative to theirs; in our choice of the isomorphism  $\text{Aut}(E[N]) \simeq \text{GL}_2(\mathbb{Z}/N\mathbb{Z})$  we view matrices in  $\text{GL}_2(\mathbb{Z}/N\mathbb{Z})$  as acting on the left, rather than the right).

**Acknowledgements.** We thank Jeremy Rouse and David Zureick-Brown for the feedback on an early draft of this article, and the referees for their careful review and helpful comments.

**Notation and Terminology.** For each integer  $n \geq 1$ , we denote by  $\zeta_n$  the  $n$ th root of unity  $e^{2\pi i/n}$  in  $\mathbb{C}$ , and let  $K_n := \mathbb{Q}(\zeta_n)$  denote the corresponding cyclotomic field. For any non-constant function  $f \in K(t)$ , where  $K$  is a field, the *degree* of  $f$  is its degree as a morphism  $\mathbb{P}_K^1 \rightarrow \mathbb{P}_K^1$ .

For any ring  $R$ , we denote by  $M_2(R)$  the ring of  $2 \times 2$  matrices with coefficients in  $R$ . We denote by  $\hat{\mathbb{Z}}$  the profinite completion of  $\mathbb{Z}$ , and view the profinite group

$$\text{GL}_2(\hat{\mathbb{Z}}) \simeq \varprojlim_N \text{GL}_2(\mathbb{Z}/N\mathbb{Z}) \simeq \prod_{\ell} \text{GL}_2(\mathbb{Z}_{\ell})$$

as a topological group in the profinite topology. If  $G$  is an open subgroup of  $\text{GL}_2(\hat{\mathbb{Z}})$ , we define its *level* to be the least positive integer  $N$  for which  $G$  is the inverse image of a subgroup of  $\text{GL}_2(\mathbb{Z}/N\mathbb{Z})$  under the natural projection  $\text{GL}_2(\hat{\mathbb{Z}}) \rightarrow \text{GL}_2(\mathbb{Z}/N\mathbb{Z})$ . If  $G$  is a subgroup of  $\text{GL}_2(\mathbb{Z}/N\mathbb{Z})$ , its level is defined to be the level of its inverse image in  $\text{GL}_2(\hat{\mathbb{Z}})$ , which is necessarily a divisor of  $N$ . For convenience we may identify the level  $N$  subgroups of  $\text{GL}_2(\mathbb{Z}/N\mathbb{Z})$  with their inverse images in  $\text{GL}_2(\hat{\mathbb{Z}})$ , and conversely. By the *genus* of an open subgroup  $G$  of  $\text{GL}_2(\hat{\mathbb{Z}})$  satisfying  $-I \in G$  and  $\det(G) = \hat{\mathbb{Z}}^{\times}$ , we mean the genus of the modular curve  $X_G$  defined in §2.

For sets  $S$  and  $T$  we use  $S - T$  to denote the set of elements that lie in  $S$  but not  $T$ .

## 2. MODULAR FUNCTIONS AND MODULAR CURVES

In this section, we summarize the background we need concerning modular curves.

**2.1. Congruence subgroups.** Fix a congruence subgroup  $\Gamma$  of  $\mathrm{SL}_2(\mathbb{Z})$ , i.e., a subgroup of  $\mathrm{SL}_2(\mathbb{Z})$  containing

$$\Gamma(N) := \{A \in \mathrm{SL}_2(\mathbb{Z}) : A \equiv I \pmod{N}\}$$

for some integer  $N \geq 1$ . The smallest such  $N$  is the *level* of  $\Gamma$ .

The group  $\Gamma$  acts on the complex upper half plane  $\mathbb{H}$  by linear fractional transformations, and the quotient  $Y_\Gamma = \Gamma \backslash \mathbb{H}$  is a smooth Riemann surface. By adding *cusps*, we can extend  $Y_\Gamma$  to a smooth compact Riemann surface  $X_\Gamma$ . We denote by  $X(N)$  the Riemann surface  $X_{\Gamma(N)}$ . The *genus* of  $\Gamma$  is the genus of the Riemann surface  $X_\Gamma$ .

**2.2. Cusps.** Define the extended upper half plane by  $\mathbb{H}^* := \mathbb{H} \cup \mathbb{P}^1(\mathbb{Q}) = \mathbb{H} \cup \mathbb{Q} \cup \{\infty\}$ . The action of  $\Gamma$  extends to  $\mathbb{H}^*$  and we can identify the quotient  $\Gamma \backslash \mathbb{H}^*$  with  $X_\Gamma$ . In particular, the cusps correspond to the  $\Gamma$ -orbits of  $\mathbb{Q} \cup \{\infty\}$ .

**Lemma 2.1.** *Let  $a/b$  and  $\alpha/\beta$  be elements of  $\mathbb{Q} \cup \{\infty\}$  with  $\gcd(a, b) = \gcd(\alpha, \beta) = 1$  (where we take  $\infty = \pm 1/0$ ). Then  $\Gamma \cdot a/b = \Gamma \cdot \alpha/\beta$  if and only if  $\gamma \begin{pmatrix} a \\ b \end{pmatrix} \equiv \pm \begin{pmatrix} \alpha \\ \beta \end{pmatrix} \pmod{N}$  for some  $\gamma \in \Gamma$ .*

*Proof.* For the case  $\Gamma = \Gamma(N)$ , see [Shi94, Lemma 1.42]. The general case follows easily.  $\square$

Let  $\pm\Gamma$  be the congruence subgroup generated by  $-I$  and  $\Gamma$ . From Lemma 2.1, we find that the cusps of  $X_\Gamma$  correspond with the orbits of  $\pm\Gamma$  on the set of  $\begin{pmatrix} a \\ b \end{pmatrix} \in (\mathbb{Z}/N\mathbb{Z})^2$  of order  $N$ . Using this, it is straightforward to find representatives of the cusps of  $X_\Gamma$ .

**2.3. Modular functions.** A *modular function* for  $\Gamma$  is a meromorphic function of  $X_\Gamma$ ; they correspond to meromorphic functions  $f$  of  $\mathbb{H}$  that satisfy  $f(\gamma\tau) = f(\tau)$  for all  $\gamma \in \Gamma$  and are meromorphic at the cusps. The function field  $\mathbb{C}(X_\Gamma)$  of  $X_\Gamma$  consists of the meromorphic functions of  $X_\Gamma$ .

Let  $\tau$  be a variable of the upper half plane. Let  $w$  be the width of the cusp at  $\infty$ , i.e., the smallest positive integer for which  $\begin{pmatrix} 1 & w \\ 0 & 1 \end{pmatrix}$  is an element of  $\Gamma$ ; it is a divisor of  $N$ . For any rational number  $m$ , define  $q^m := e^{2\pi im\tau}$ . Then any modular function  $f$  for  $\Gamma$  has a unique  $q$ -expansion

$$f(\tau) = \sum_{n \in \mathbb{Z}} c_n q^{n/w},$$

where the  $c_n$  are complex numbers that are 0 for all but finitely many  $n < 0$ . We will often refer to the  $c_n$  as the *coefficients* of  $f$ .

**2.4. Field of modular functions.** Fix a positive integer  $N$ . Denote by  $\mathcal{F}_N$  the field of meromorphic functions of the Riemann surface  $X(N)$  whose  $q$ -expansions have coefficients in  $K_N := \mathbb{Q}(\zeta_N)$ . For example,  $\mathcal{F}_1 = \mathbb{Q}(j)$ , where  $j$  is the modular  $j$ -invariant.

For  $f \in \mathcal{F}_N$  and  $\gamma \in \mathrm{SL}_2(\mathbb{Z})$ , let  $f|_\gamma \in \mathcal{F}_N$  denote the modular function satisfying  $f|_\gamma(\tau) = f(\gamma\tau)$ .

For each  $d \in (\mathbb{Z}/N\mathbb{Z})^\times$ , let  $\sigma_d$  be the automorphism of  $K_N$  satisfying  $\sigma_d(\zeta_N) = \zeta_N^d$ . We extend  $\sigma_d$  to an automorphism of the field  $\mathcal{F}_N$  by defining

$$\sigma_d(f) := \sum_n \sigma_d(c_n) q^{n/N},$$

where  $f$  has expansion  $\sum_n c_n q^{n/N}$ . We now recall some facts about the extension  $\mathcal{F}_N$  of  $\mathcal{F}_1 = \mathbb{Q}(j)$ .

**Proposition 2.2.** *The extension  $\mathcal{F}_N$  of  $\mathbb{Q}(j)$  is Galois. There is a unique isomorphism*

$$\theta_N : \mathrm{GL}_2(\mathbb{Z}/N\mathbb{Z}) / \{\pm I\} \xrightarrow{\sim} \mathrm{Gal}(\mathcal{F}_N / \mathbb{Q}(j))$$

such that the following hold for all  $f \in \mathcal{F}_N$ :

- (a) For  $g \in \mathrm{SL}_2(\mathbb{Z}/N\mathbb{Z})$ , we have  $\theta_N(g)f = f|_{\gamma^t}$ , where  $\gamma$  is any matrix in  $\mathrm{SL}_2(\mathbb{Z})$  that is congruent to  $g$  modulo  $N$  and  $\gamma^t$  is the transpose of  $\gamma$ .
- (b) For  $g = \begin{pmatrix} 1 & 0 \\ 0 & d \end{pmatrix} \in \mathrm{GL}_2(\mathbb{Z}/N\mathbb{Z})$ , we have  $\theta_N(g)f = \sigma_d(f)$ .

Moreover, the algebraic closure of  $\mathbb{Q}$  in  $\mathcal{F}_N$  is  $\mathbb{Q}(\zeta_N)$ ; it corresponds to the subgroup  $\mathrm{SL}_2(\mathbb{Z}/N\mathbb{Z}) / \{\pm I\}$ .

*Proof.* This is well known; see [KL81, Ch. 2 §2] for a summary (where the action given is a right action obtained as above but without the transpose in (a)).  $\square$

Throughout the rest of the paper, we let  $\mathrm{GL}_2(\mathbb{Z}/N\mathbb{Z})$  act on  $\mathcal{F}_N$  via the homomorphism  $\theta_N$  of Proposition 2.2. We set  $g_*(f) := \theta_N(g)(f)$  for  $g \in \mathrm{GL}_2(\mathbb{Z}/N\mathbb{Z})$  and  $f \in \mathcal{F}_N$ .

*Remark 2.3.* There are other natural actions of  $\mathrm{GL}_2(\mathbb{Z}/N\mathbb{Z})$  on  $\mathcal{F}_N$ ; for example, one could replace  $\gamma^t$  in condition (a) by  $\gamma^{-1}$  or just act on the right. Our choice is motivated by Proposition 2.6 below.

**2.5. Modular curves.** Let  $G$  be a subgroup of  $\mathrm{GL}_2(\mathbb{Z}/N\mathbb{Z})$  satisfying  $-I \in G$  and  $\det(G) = (\mathbb{Z}/N\mathbb{Z})^\times$ . Let  $\mathcal{F}_N^G$  be the subfield of  $\mathcal{F}_N$  fixed by the action of  $G$  from Proposition 2.2. Proposition 2.2 and the assumption  $\det(G) = (\mathbb{Z}/N\mathbb{Z})^\times$  imply that  $\mathbb{Q}$  is algebraically closed in  $\mathcal{F}_N^G$ .

The *modular curve*  $X_G$  associated with  $G$  is the smooth projective curve with function field  $\mathcal{F}_N^G$ . The curve  $X_G$  is defined over  $\mathbb{Q}$  and is geometrically irreducible. The inclusion of fields  $\mathcal{F}_N^G \supseteq \mathcal{F}_1 = \mathbb{Q}(j)$  gives rise to a non-constant morphism

$$\pi_G: X_G \rightarrow \mathrm{Spec} \mathbb{Q}[j] \cup \{\infty\} = \mathbb{P}_{\mathbb{Q}}^1$$

of degree  $[\mathrm{GL}_2(\mathbb{Z}/N\mathbb{Z}) : G]$ . Moreover, given another group  $G \subseteq G' \subseteq \mathrm{GL}_2(\mathbb{Z}/N\mathbb{Z})$ , the inclusion  $\mathcal{F}_N^{G'} \subseteq \mathcal{F}_N^G$  induces a nonconstant morphism  $X_G \rightarrow X_{G'}$  of degree  $[G' : G]$ . Composing  $X_G \rightarrow X_{G'}$  with  $\pi_{G'}$  gives the morphism  $\pi_G$ .

Let  $\Gamma$  be the congruence subgroup consisting of  $\gamma \in \mathrm{SL}_2(\mathbb{Z})$  for which  $\gamma^t$  modulo  $N$  lies in  $G \cap \mathrm{SL}_2(\mathbb{Z}/N\mathbb{Z})$ . The level of  $\Gamma$  divides, but need not equal,  $N$ .

**Lemma 2.4.**

- (i) *The field  $K_N(X_G)$ , i.e., the function field of the base extension of  $X_G$  to  $K_N$ , is the field consisting of  $f \in \mathcal{F}_N$  satisfying  $f|_\gamma = f$  for all  $\gamma \in \Gamma$ .*
- (ii) *The genus of the modular curve  $X_G$  is equal to the genus of  $\Gamma$ .*

*Proof.* Proposition 2.2 implies that  $K_N$  is algebraically closed in  $\mathcal{F}_N$  and that we have an isomorphism  $\mathrm{Gal}(\mathcal{F}_N/K_N(j)) \xrightarrow{\sim} \mathrm{SL}_2(\mathbb{Z}/N\mathbb{Z})/\{\pm I\}$ . Thus  $K_N(X_G)$  is the subfield of  $\mathcal{F}_N$  fixed by  $G \cap \mathrm{SL}_2(\mathbb{Z}/N\mathbb{Z})$ . Part (i) is now clear.

Since  $K_N$  is algebraically closed in  $\mathcal{F}_N$  and  $\mathbb{Q}$  is algebraically closed in  $\mathbb{Q}(X_G)$ , we have

$$[\mathbb{C} \cdot K_N(X_G) : \mathbb{C}(j)] = [K_N(X_G) : K_N(j)] = [\mathbb{Q}(X_G) : \mathbb{Q}(j)] = [\mathrm{GL}_2(\mathbb{Z}/N\mathbb{Z}) : G].$$

Since  $\det(G) = (\mathbb{Z}/N\mathbb{Z})^\times$ , we deduce that  $[\mathbb{C} \cdot K_N(X_G) : \mathbb{C}(j)] = [\mathrm{SL}_2(\mathbb{Z}) : \Gamma]$ .

Clearly each  $f \in K_N(X_G)$  is a modular function for  $\Gamma$ , thus  $\mathbb{C} \cdot K_N(X_G) \subseteq \mathbb{C}(X_\Gamma)$ . We in fact have  $\mathbb{C} \cdot K_N(X_G) = \mathbb{C}(X_\Gamma)$ , since  $[\mathbb{C} \cdot K_N(X_G) : \mathbb{C}(j)] = [\mathrm{SL}_2(\mathbb{Z}) : \Gamma] = [\mathbb{C}(X_\Gamma) : \mathbb{C}(j)]$ . The curve  $X_G$  has the same genus as the Riemann surface  $X_\Gamma$  because  $\mathbb{C}(X_G) = \mathbb{C}(X_\Gamma)$ .  $\square$

*Remark 2.5.* Another natural congruence subgroup to study is the congruence subgroup  $\Gamma'$  consisting of  $\gamma \in \mathrm{SL}_2(\mathbb{Z})$  such that  $\gamma$  modulo  $N$  lies in  $G \cap \mathrm{SL}_2(\mathbb{Z}/N\mathbb{Z})$ , which we use later in the paper. Observe that the congruence subgroups  $\Gamma$  and  $\Gamma'$  are conjugate in  $\mathrm{SL}_2(\mathbb{Z})$ ; indeed, we have  $B^{-1}\gamma B = (\gamma^t)^{-1}$  for all  $\gamma \in \mathrm{SL}_2(\mathbb{Z})$ , where  $B := \begin{pmatrix} 0 & 1 \\ -1 & 0 \end{pmatrix}$ . Thus  $\Gamma$  and  $\Gamma'$  have the same genus.

The following proposition is crucial to our application.

**Proposition 2.6.** *Let  $E$  be an elliptic curve defined over  $\mathbb{Q}$  with  $j_E \notin \{0, 1728\}$ . Then  $\rho_{E,N}(\mathrm{Gal}_{\mathbb{Q}})$  is conjugate in  $\mathrm{GL}_2(\mathbb{Z}/N\mathbb{Z})$  to a subgroup of  $G$  if and only if  $j_E$  belongs to  $\pi_G(X_G(\mathbb{Q}))$ .*

*Proof.* See §3 of [Zyw15a] for a proof.  $\square$

**2.6. Modular curves and open subgroups.** Fix an open subgroup  $G$  of  $\mathrm{GL}_2(\hat{\mathbb{Z}})$  that satisfies  $-I \in G$  and  $\det(G) = \hat{\mathbb{Z}}^\times$ . Let  $N \geq 1$  be an integer that is divisible by the level of  $G$ . Define the modular curve

$$X_G := X_{\bar{G}},$$

where  $\bar{G} \subseteq \mathrm{GL}_2(\mathbb{Z}/N\mathbb{Z})$  is the image of  $G$  modulo  $N$ . Observe that the modular curve  $X_G$  and its function field do not depend on the initial choice of  $N$ .

Every (open) subgroup  $G'$  of  $\mathrm{GL}_2(\hat{\mathbb{Z}})$  that contains  $G$  satisfies  $-I \in G'$  and  $\det(G') = \hat{\mathbb{Z}}^\times$ , and we have a morphism  $X_G \rightarrow X_{G'}$ . With  $G' = \mathrm{GL}_2(\hat{\mathbb{Z}})$ , we obtain a morphism  $\pi_G: X_G \rightarrow X_{G'} = \mathbb{P}_{\mathbb{Q}}^1$  to the  $j$ -line that agrees with  $\pi_{\bar{G}}$ . The following is equivalent to Proposition 2.6.

**Proposition 2.7.** *Let  $E$  be an elliptic curve defined over  $\mathbb{Q}$  with  $j_E \notin \{0, 1728\}$ . Then  $\rho_E(\mathrm{Gal}_{\mathbb{Q}})$  is conjugate in  $\mathrm{GL}_2(\hat{\mathbb{Z}})$  to a subgroup of  $G$  if and only if  $j_E$  belongs to  $\pi_G(X_G(\mathbb{Q}))$ .  $\square$*

**2.7. Complex conjugation.** Fix a subgroup  $G$  of  $\mathrm{GL}_2(\mathbb{Z}/N\mathbb{Z})$  satisfying  $-I \in G$  and  $\det(G) = (\mathbb{Z}/N\mathbb{Z})^\times$ . For our curve  $X_G$  to have rational points, we need  $G$  to contain an element that “looks like” complex conjugation.

**Lemma 2.8.** *For any elliptic curve  $E/\mathbb{Q}$  and integer  $N > 1$ , the group  $\rho_{E,N}(\mathrm{Gal}_{\mathbb{Q}})$  contains an element that is conjugate in  $\mathrm{GL}_2(\mathbb{Z}/N\mathbb{Z})$  to  $\begin{pmatrix} 1 & 0 \\ 0 & -1 \end{pmatrix}$  or  $\begin{pmatrix} 1 & 1 \\ 0 & -1 \end{pmatrix}$ .*

*Proof.* This follows from Proposition 3.5 of [Zyw15b] (and its proof for the cases  $j_E \in \{0, 1728\}$ ).  $\square$

Note that  $\begin{pmatrix} 1 & 0 \\ 0 & -1 \end{pmatrix}$  and  $\begin{pmatrix} 1 & 1 \\ 0 & -1 \end{pmatrix}$  are conjugate to each other in  $\mathrm{GL}_2(\mathbb{Z}/N\mathbb{Z})$  if  $N$  is odd. If  $G$  does not contain an element that is conjugate in  $\mathrm{GL}_2(\mathbb{Z}/N\mathbb{Z})$  to  $\begin{pmatrix} 1 & 0 \\ 0 & -1 \end{pmatrix}$  or  $\begin{pmatrix} 1 & 1 \\ 0 & -1 \end{pmatrix}$ , then  $X_G(\mathbb{Q})$  must be empty since  $X_G(\mathbb{R})$  is finite (by [Zyw15b, Proposition 3.5]), hence empty, since  $X_G$  is non-singular.

### 3. GROUP THEORETIC COMPUTATIONS

We define an *admissible group* to be an open subgroup  $G$  of  $\mathrm{GL}_2(\hat{\mathbb{Z}})$  for which the following conditions hold:

- $G$  has prime power level,
- $-I \in G$  and  $\det(G) = \hat{\mathbb{Z}}^\times$ ,
- $G$  contains an element that is conjugate in  $\mathrm{GL}_2(\hat{\mathbb{Z}})$  to  $\begin{pmatrix} 1 & 0 \\ 0 & -1 \end{pmatrix}$  or  $\begin{pmatrix} 1 & 1 \\ 0 & -1 \end{pmatrix}$ .

The condition  $\det(G) = \hat{\mathbb{Z}}^\times$  is needed for Proposition 2.7 since we have  $\det(\rho_E(\mathrm{Gal}_{\mathbb{Q}})) = \hat{\mathbb{Z}}^\times$ . If we were interested in elliptic curve defined over other number fields, then we could loosen this restriction which could increase the base field of the modular curve  $X_G$ .

The condition  $-I \in G$  is also needed in Proposition 2.7. For an elliptic curve  $E/\mathbb{Q}$ , there is a quadratic twist  $E'/\mathbb{Q}$ , which automatically has the same  $j$ -invariant as  $E$ , such that  $-I \in \rho_{E'}(\mathrm{Gal}_{\mathbb{Q}})$ .

The last condition on  $G$  is necessary in order for  $X_G(\mathbb{Q})$  to be non-empty, as explained in §2.7.

**Proposition 3.1.** *Let  $G$  be an admissible group of genus 0. The set  $X_G(\mathbb{Q})$  is infinite.*

*Proof.* We have  $X_G(\mathbb{R}) \neq \emptyset$  by Proposition 3.5 of [Zyw15b]. For primes  $p$  not dividing its prime power level the modular curve  $X_G$  has good reduction at  $p$  and  $X_G(\mathbb{Q}_p) \neq \emptyset$ , since the reduction of  $X_G$  to  $\mathbb{F}_p$  necessarily has rational points that can be lifted to  $\mathbb{Q}_p$  via Hensel’s lemma. Thus  $X_G$  has rational points locally at all but at most one place of  $\mathbb{Q}$ . The product formula for Hilbert symbols and the Hasse-Minkowski Theorem then imply that  $X_G$  has a rational point and is thus isomorphic to  $\mathbb{P}^1$  and has infinitely many rational points.  $\square$

*Remark 3.2.* As shown by Proposition 3.1, our three criteria for admissibility rule out genus 0 curves with no rational points. There are ten groups  $G$  of 2-power level that satisfy our first two criteria but not the third; these give rise to the ten pointless conics  $X_G$  found in [RZB15]. There are three such groups of 3-power level, three of 5-power level, and none of higher prime-power level.

Fix an integer  $g \geq 0$ . In this section, we explain how to enumerate all admissible subgroups  $G$  of  $\mathrm{GL}_2(\hat{\mathbb{Z}})$ , up to conjugacy, that have genus at most  $g$ . We shall apply these methods with  $g = 1$  to verify Theorem 3.3 below, and to find explicit representatives of these conjugacy classes of groups; Magma [BCP97] scripts that perform this enumeration can be found at [SZ16].

**Theorem 3.3.**

- (i) *Up to conjugacy in  $\mathrm{GL}_2(\hat{\mathbb{Z}})$ , there are 220 admissible subgroups of genus 0.*
- (ii) *Up to conjugacy in  $\mathrm{GL}_2(\hat{\mathbb{Z}})$ , there are 250 admissible subgroups of genus 1.*

*Remark 3.4.* The 220 admissible subgroups  $G$  of genus 0, up to conjugacy, are precisely those given in Tables 1–3. More precisely, for each entry of the table, we have an integer  $N$  and a set of generators that generates the image in  $\mathrm{GL}_2(\mathbb{Z}/N\mathbb{Z})$  of an admissible group of level  $N$  and genus 0.

*Remark 3.5.* The 28 admissible subgroups  $G$  of genus 1 that have infinitely many rational points, up to conjugacy, are precisely those given in Table 4, of which 27 have level 16 and 1 has level 11. The levels arising among the remaining 222 are 7, 8, 9, 11, 16, 17, 19, 27, 32, and 49.

For a fixed admissible group  $G$  of level  $N$ , let  $\Gamma$  be the congruence subgroup of  $\mathrm{SL}_2(\mathbb{Z})$  consisting of matrices whose image modulo  $N$  lies in the image of  $G$  modulo  $N$ ; the level of  $\Gamma$  necessarily divides  $N$ , and  $\Gamma$  contains  $-I$ . By Lemma 2.4(ii) and Remark 2.5, the modular curve  $X_G$  has the same genus as  $\Gamma$ .

The basic idea of our computation is to reverse the process above; we start with a congruence subgroup  $\Gamma$  of genus at most  $g$  and prime power level, and then enumerate the possible groups  $G$  that could produce  $\Gamma$ .

Let  $S_g$  be the set of congruence subgroups of  $\mathrm{SL}_2(\mathbb{Z})$  of prime power level that contain  $-I$  and have genus at most  $g$ . We know that the set  $S_g$  is finite from a theorem of Dennin [Den74]. When  $g \leq 24$ , and in particular, for  $g = 1$ , we can explicitly determine the elements of  $S_g$  from the tables of Cummins and Pauli [CP03], which are available online at [CP] (their methods can also be extended to larger  $g$ ).

Let  $L_g$  be the set of primes that divide the level of some congruence subgroup  $\Gamma \in S_g$ . The set  $L_g$  is finite, since  $S_g$  is finite, and we have  $L_1 = \{2, 3, 5, 7, 11, 13, 17, 19\}$ . If  $G$  is an admissible group of genus at most  $g$ , then its level must be a power of a prime  $\ell \in L_g$ . For the rest of the section, we fix a prime  $\ell \in L_g$ . Since  $L_g$  is finite, it suffices to explain how to compute the admissible groups  $G$  with genus at most  $g$  whose level is a power of  $\ell$ , and we need only consider levels strictly greater than 1 since  $\mathrm{GL}_2(\hat{\mathbb{Z}})$  is the only admissible group of level 1.

Fix a prime power  $N := \ell^n > 1$ , and consider any congruence subgroup  $\Gamma \in S_g$  whose level divides  $N$ . By enumerating subgroups of  $\mathrm{GL}_2(\mathbb{Z}/N\mathbb{Z})$  one can explicitly determine those subgroups  $G_N$  that satisfy the following conditions:

- (1)  $G_N$  has level  $N$ ,
- (2)  $G_N \cap \mathrm{SL}_2(\mathbb{Z}/N\mathbb{Z})$  is equal to the image of  $\Gamma$  modulo  $N$ ,
- (3)  $\det(G_N) = (\mathbb{Z}/N\mathbb{Z})^\times$ ,
- (4)  $G_N$  contains an element that is conjugate in  $\mathrm{GL}_2(\mathbb{Z}/N\mathbb{Z})$  to  $\begin{pmatrix} 1 & 0 \\ 0 & -1 \end{pmatrix}$  or  $\begin{pmatrix} 1 & 1 \\ 0 & -1 \end{pmatrix}$ .

Let  $H$  be the image of  $\Gamma$  in  $\mathrm{SL}_2(\mathbb{Z}/N\mathbb{Z})$ . The group  $H = G_N \cap \mathrm{SL}_2(\mathbb{Z}/N\mathbb{Z})$  is normal in  $G_N$  and hence  $G_N$  is a subgroup of the normalizer  $K$  of  $H$  in  $\mathrm{GL}_2(\mathbb{Z}/N\mathbb{Z})$ . So rather than searching for  $G_N$  in  $K$ , we can work in the quotient  $K/H$  where the image of  $G_N$  is an abelian group isomorphic to  $(\mathbb{Z}/N\mathbb{Z})^\times$ . Using Magma, we can efficiently enumerate all abelian subgroups  $A$  of  $K/H$  of order  $\#(\mathbb{Z}/N\mathbb{Z})^\times$ . For each such subgroup  $A$  we then test whether its inverse image  $G_N$  in  $K$  satisfies conditions (1)–(4) above.

Let  $G$  be the subgroup of  $\mathrm{GL}_2(\hat{\mathbb{Z}})$  consisting of those matrices whose image modulo  $N$  lies in a fixed group  $G_N$  satisfying the conditions (1)–(4). The group  $G$  is admissible of level  $N$  and has

genus at most  $g$ . Moreover, it is clear that every admissible group of level  $N$  and genus at most  $g$  arises in this manner.

Fix an integer  $e \geq 1$ . By applying the above method with  $1 \leq n \leq e$ , we obtain all admissible groups  $G$  of genus at most  $g$  and level dividing  $\ell^e$ . Our algorithm proceeds by applying this procedure to increasing values of  $e$ . In order for it to terminate we need to know that there are only finitely many admissible groups  $G$  of  $\ell$ -power level and genus at most  $g$ , and we need an explicit way to determine when we have reached an  $e$  that is large enough to guarantee that we have found them all. Proposition 3.6 below addresses both issues.

**Proposition 3.6.**

- (i) *There are only finitely many admissible groups  $G$  with genus at most  $g$  whose level is a power of  $\ell$ .*
- (ii) *Take any integer  $n \geq 2$  with  $n \neq 2$  if  $\ell = 2$ . Define  $N := \ell^n$ . Suppose that there is no subgroup  $G_N$  of  $\mathrm{GL}_2(\mathbb{Z}/N\mathbb{Z})$  that satisfies conditions (1)–(4) for some  $\Gamma \in S_g$  with level dividing  $N$ . Then any admissible group  $G$  of genus at most  $g$  with level a power of  $\ell$  has level at most  $N$ .*

The remainder of this section is devoted to proving Proposition 3.6. We will need the following basic lemma.

**Lemma 3.7.** *Let  $\ell$  be a prime and let  $G$  be an open subgroup of  $\mathrm{GL}_2(\mathbb{Z}_\ell)$ . For each integer  $m \geq 1$ , let  $i_m$  be the index of the image of  $G$  in  $\mathrm{GL}_2(\mathbb{Z}/\ell^m\mathbb{Z})$ . If  $i_{n+1} = i_n$  for an integer  $n \geq 1$ , with  $n \neq 1$  if  $\ell = 2$ , then  $[\mathrm{GL}_2(\mathbb{Z}_\ell) : G] = i_n$ .*

*Proof.* Since  $G$  is an open subgroup, it suffices to prove  $i_{m+1} = i_m$  for all  $m \geq n$ ; we proceed by induction on  $m$ . The base case is given, so we assume  $i_{m+1} = i_m$  for some  $m \geq n$ ; we need to show that  $i_{m+2} = i_{m+1}$ . Let  $G_m$  denote the image of  $G$  in  $\mathrm{GL}_2(\mathbb{Z}/\ell^m\mathbb{Z})$ . Reduction modulo  $\ell^m$  gives exact sequences related by inclusions

$$\begin{array}{ccccccc} 1 & \longrightarrow & K_{m+1} & \longrightarrow & \mathrm{GL}_2(\mathbb{Z}/\ell^{m+1}\mathbb{Z}) & \longrightarrow & \mathrm{GL}_2(\mathbb{Z}/\ell^m\mathbb{Z}) & \longrightarrow & 1 \\ & & \uparrow & & \uparrow & & \uparrow & & \\ 1 & \longrightarrow & H_{m+1} & \longrightarrow & G_{m+1} & \longrightarrow & G_m & \longrightarrow & 1. \end{array}$$

The inductive hypothesis  $i_{m+1} = i_m$  implies that the kernels  $H_{m+1}$  and  $K_{m+1}$  coincide; in particular,  $H_{m+1}$  is as large as possible (i.e., it has order  $\ell^4$ ). It thus suffices to show that the kernel  $H_{m+2}$  of the reduction map from  $G_{m+2}$  to  $G_{m+1}$  also has order  $\ell^4$ . We have  $|H_{m+2}| \leq \ell^4$ , so it suffices to give an injective map  $H_{m+1} \rightarrow H_{m+2}$ .

Let  $M$  be an element of  $G$  whose image in  $G_{m+1}$  lies in  $H_{m+1}$ ; then  $M = I + \ell^m A$  for some  $A \in \mathrm{M}_2(\mathbb{Z}_\ell)$ . Since  $m \geq 1$ , with  $m \geq 2$  if  $\ell = 2$ , we have

$$(1 + \ell^m A)^\ell = 1 + \binom{\ell}{1} \ell^m A + \binom{\ell}{2} \ell^{2m} A^2 + \dots \equiv 1 + \ell^{m+1} A \pmod{\ell^{m+2}}.$$

The  $\ell$ -power map thus induces an injection  $H_{m+1} \rightarrow H_{m+2}$ . □

*Remark 3.8.* Lemma 3.7 holds more generally. One can replace  $\mathrm{GL}_2(\mathbb{Z}_\ell)$  with the unit group of any (unital associative)  $\mathbb{Z}_\ell$ -algebra  $\mathcal{A}$  that is torsion-free and finitely-generated as a  $\mathbb{Z}_\ell$ -module (in the lemma,  $\mathcal{A} = \mathrm{M}_2(\mathbb{Z}_\ell)$ ); the proof is exactly the same.

*Proof of Proposition 3.6(i).* Let  $\mathcal{G}$  be the set of admissible groups of genus at most  $g$  whose level is a power of  $\ell$ . Note that if  $G'$  is a subgroup of  $\mathrm{GL}_2(\hat{\mathbb{Z}})$  containing some  $G \in \mathcal{G}$ , then  $G' \in \mathcal{G}$ . We wish to show that  $\mathcal{G}$  is finite.

We claim that any admissible group  $G$  has only finitely many maximal subgroups that are also admissible and whose level is a power of  $\ell$ . It suffices to show that an open subgroup  $H$  of  $\mathrm{GL}_2(\mathbb{Z}_\ell)$

has only finitely many open maximal subgroups. Let  $\Phi(H)$  be the Frattini subgroup of  $H$ ; it is the intersection of the maximal closed proper subgroups of  $H$ . By the proposition in §10.5 of [Ser97],  $\Phi(H)$  is an open subgroup of  $H$ . This proves the claim.

Now suppose that  $\mathcal{G}$  is infinite. The claim implies that  $\mathcal{G}$  contains an infinite descending chain  $G_1 \supsetneq G_2 \supsetneq G_3 \supsetneq \cdots$  (let  $G_1 = \mathrm{GL}_2(\hat{\mathbb{Z}}) \in \mathcal{G}$ , let  $G_2 \in \mathcal{G}$  be one of the finitely many maximal subgroups of  $G_1$  in  $\mathcal{G}$  that has infinitely many subgroups in  $\mathcal{G}$ , and continue in this fashion). For each  $i \geq 1$ , let  $\Gamma_i$  be the congruence subgroup associated to  $G_i$  (i.e.,  $\Gamma_i$  consists of the matrices in  $\mathrm{SL}_2(\mathbb{Z})$  whose image modulo  $N$  lies in the image modulo  $N$  of  $G_i$ , where  $N$  is the level of  $G_i$ ); then  $\Gamma_i \in S_g$ . Since  $[\mathrm{GL}_2(\hat{\mathbb{Z}}) : G_i] = [\mathrm{SL}_2(\mathbb{Z}) : \Gamma_i]$ , we have inclusions  $\Gamma_1 \supsetneq \Gamma_2 \supsetneq \Gamma_3 \supsetneq \cdots$ . This contradicts the finiteness of  $S_g$  and the proposition follows.  $\square$

*Proof of Proposition 3.6(ii).* Fix an integer  $n \geq 1$  as in the statement of part (ii). Suppose there is an integer  $m > n$  such that there is an admissible group  $G$  of level  $\ell^m$  and genus at most  $g$ .

With  $N := \ell^n$ , let  $G_N$  be the image of  $G$  in  $\mathrm{GL}_2(\mathbb{Z}/N\mathbb{Z})$ . The curve  $X_{G_N}$  has genus at most  $g$  since it is dominated by  $X_G$ . Therefore, conditions (2), (3) and (4) hold for some  $\Gamma \in S_g$  with level dividing  $N$ . Our assumption on  $n$  implies that the level of  $G_N$  is a proper divisor of  $N$ . This implies that the index  $i_n := [\mathrm{GL}_2(\mathbb{Z}/N\mathbb{Z}) : G_N]$  agrees with  $i_{n-1} := [\mathrm{GL}_2(\mathbb{Z}/\ell^{n-1}\mathbb{Z}) : G_{\ell^{n-1}}]$ , where  $G_{\ell^{n-1}}$  is the image of  $G$  in  $\mathrm{GL}_2(\mathbb{Z}/\ell^{n-1}\mathbb{Z})$ . Since  $i_n = i_{n-1}$ , Lemma 3.7 implies that  $[\mathrm{GL}_2(\mathbb{Z}) : G] = i_{n-1}$ . However, this means that  $G$  has level dividing  $\ell^{n-1}$  which is impossible since, by assumption,  $G$  has level  $\ell^m > \ell^{n-1}$ . Therefore, no such admissible group  $G$  exists.  $\square$

#### 4. CONSTRUCTION OF HAUPTMODULS

Fix a congruence subgroup  $\Gamma$  of genus 0 and level  $N$ . The function field of  $X_\Gamma$  is then of the form  $\mathbb{C}(h)$ , where the function  $h: X_\Gamma \rightarrow \mathbb{C} \cup \{\infty\}$  gives an isomorphism between  $X_\Gamma$  and the Riemann sphere; in particular,  $h$  has a unique (simple) pole.

We may choose  $h$  so that its unique pole is at the cusp  $\infty$ ; we will call such an  $h$  a *hauptmodul* of  $\Gamma$ . Every hauptmodul of  $\Gamma$  is then of the form  $ah + b$  for some complex numbers  $a \neq 0$  and  $b$ . For example, the familiar modular  $j$ -invariant

$$j(\tau) = q^{-1} + 744 + 196884q + 21493760q^2 + 864299970q^3 + \cdots$$

is a hauptmodul for  $\mathrm{SL}_2(\mathbb{Z})$ . If  $h$  is a hauptmodul for  $\Gamma$ , then we have an inclusion of function fields  $\mathbb{C}(j) \subseteq \mathbb{C}(h)$  and hence  $J(h) = j$  for a unique rational function  $J(h) \in \mathbb{C}(t)$ .

The main task of §4 is to describe how to find an *explicit* hauptmodul  $h$  of  $\Gamma$  in terms of Siegel functions when  $N$  is a prime power. Our  $h$  will have coefficients in  $K_N$ . In §4.4, we explain how to compute the rational function  $J(t)$  corresponding to  $h$ .

**4.1. Siegel functions.** Take any pair  $a = (a_1, a_2) \in \mathbb{Q}^2 - \mathbb{Z}^2$ . We define the *Siegel function*  $g_a(\tau)$  to be the holomorphic function  $\mathbb{H} \rightarrow \mathbb{C}^\times$  defined by the series

$$-q^{\frac{1}{2}B_2(a_1)} \cdot e(a_2(a_1 - 1)/2) \cdot (1 - e(a_2)q^{a_1}) \prod_{n=1}^{\infty} (1 - e(a_2)q^{n+a_1})(1 - e(-a_2)q^{n-a_1}),$$

where  $e(z) = e^{2\pi iz}$  and  $B_2(x) = x^2 - x + 1/6$ .

Recall that the *Dedekind eta function* is the holomorphic function on  $\mathbb{H}$  given by

$$\eta(\tau) := q^{1/24} \prod_{n=1}^{\infty} (1 - q^n).$$

For each  $\gamma = \begin{pmatrix} a & b \\ c & d \end{pmatrix} \in \mathrm{SL}_2(\mathbb{Z})$ , there is a unique 12-th root of unity  $\varepsilon(\gamma) \in \mathbb{C}^\times$  such that

$$(4.1) \quad \eta(\gamma\tau)^2 = \varepsilon(\gamma)(c\tau + d)\eta(\tau)^2.$$

We can characterize the map  $\varepsilon: \mathrm{SL}_2(\mathbb{Z}) \rightarrow \mathbb{C}^\times$  by the property that it is a homomorphism satisfying  $\varepsilon\left(\begin{pmatrix} 1 & 1 \\ 0 & 1 \end{pmatrix}\right) = \zeta_{12}$  and  $\varepsilon\left(\begin{pmatrix} 0 & 1 \\ -1 & 0 \end{pmatrix}\right) = \zeta_4$ , cf. [KL81, Ch. 3 §5]. Moreover, the kernel of  $\varepsilon$  is a congruence subgroup of level 12 and agrees with the commutator subgroup of  $\mathrm{SL}_2(\mathbb{Z})$ .

The following lemma gives several key properties of Siegel functions.

**Lemma 4.1.** *For any  $\gamma \in \mathrm{SL}_2(\mathbb{Z})$ ,  $a \in \mathbb{Q}^2 - \mathbb{Z}^2$ , and  $b \in \mathbb{Z}^2$ , the following hold:*

- (i)  $g_{-a} = -g_a$ ,
- (ii)  $g_{a+b} = (-1)^{b_1+b_2+b_1b_2} \cdot e((b_2a_1 - b_1a_2)/2) \cdot g_a$ ,
- (iii)  $g_a|_\gamma = \varepsilon(\gamma) \cdot g_{a\gamma}$ , where we view  $a$  as a row vector.

*Proof.* In [KL81, Ch. 2 §1], we see that  $g_a(\tau) = \mathfrak{k}_a(\tau)\eta(\tau)^2$ , where  $\mathfrak{k}_a(\tau)$  is a Klein form (with  $W = W_\tau$  in the notation of loc. cit.). Part (ii) follows directly from property K2 of §1 of [KL81, Ch. 2].

Take any  $\gamma \in \mathrm{SL}_2(\mathbb{Z})$  and let  $(c, d)$  be the last row of  $\gamma$ . From properties K0 and K1 of §1 of [KL81, Ch. 2], we find that

$$(4.2) \quad \mathfrak{k}_a(\gamma\tau) = (c\tau + d)^{-1} \mathfrak{k}_{a\gamma}(\tau)$$

From (4.1) and (4.2), we deduce that  $g_a(\gamma\tau) = \varepsilon(\gamma) \cdot g_{a\gamma}(\tau)$  which proves part (iii). Finally, part (i) follows from part (iii) with  $\gamma = -I$ , since  $\varepsilon(-I) = -1$ .  $\square$

For an integer  $N > 1$ , let  $\mathcal{A}_N$  be the set of pairs  $(a_1, a_2) \in N^{-1}\mathbb{Z}^2 - \mathbb{Z}^2$  that satisfy one of the following conditions:

- $0 < a_1 < 1/2$  and  $0 \leq a_2 < 1$ ,
- $a_1 = 0$  and  $0 < a_2 \leq 1/2$ ,
- $a_1 = 1/2$  and  $0 \leq a_2 \leq 1/2$ .

The set  $\mathcal{A}_N$  is chosen so that every non-zero coset of  $(N^{-1}\mathbb{Z}^2)/\mathbb{Z}^2$  is represented by an element of the form  $a$  or  $-a$  for a unique  $a \in \mathcal{A}_N$ . So for any  $a \in N^{-1}\mathbb{Z}^2 - \mathbb{Z}^2$ , we can use parts (i) and (ii) of Lemma 4.1 to show that

$$g_a = \epsilon \cdot \zeta \cdot g_{a'}$$

for an explicit sign  $\epsilon \in \{\pm 1\}$ ,  $N$ -th root of unity  $\zeta$ , and pair  $a' \in \mathcal{A}_N$ .

**4.2. Siegel orbits.** Now fix a congruence subgroup  $\Gamma$  of level  $N > 1$ . For each  $a \in \mathcal{A}_N$  and  $\gamma \in \mathrm{SL}_2(\mathbb{Z})$ , let  $a * \gamma$  be the unique element of  $\mathcal{A}_N$  such that  $a * \gamma$  or  $-a * \gamma$  lies in the coset  $a\gamma + \mathbb{Z}^2$ . The map

$$\mathcal{A}_N \times \mathrm{SL}_2(\mathbb{Z}) \rightarrow \mathcal{A}_N, \quad (a, \gamma) \mapsto a * \gamma$$

then gives a right action of  $\mathrm{SL}_2(\mathbb{Z})$  on  $\mathcal{A}_N$ . In particular, this gives a right action of  $\Gamma$  on  $\mathcal{A}_N$ .

Fix a  $\Gamma$ -orbit  $\mathcal{O}$  of  $\mathcal{A}_N$  and define

$$g_{\mathcal{O}} := \prod_{a \in \mathcal{O}} g_a;$$

it is a holomorphic function  $\mathbb{H} \rightarrow \mathbb{C}^\times$ .

**Lemma 4.2.** *The function  $g_{\mathcal{O}}^{12N}$  is a modular function for  $\Gamma$ . Every pole and zero of  $g_{\mathcal{O}}^{12N}$  on  $X_\Gamma$  is a cusp.*

*Proof.* Take any  $\gamma \in \Gamma$  and  $a \in \mathcal{A}_N$ . By Lemma 4.1(iii), we have  $g_a^{12N}|_\gamma = g_{a\gamma}^{12N}$ . We have  $a\gamma = \epsilon \cdot (a * \gamma + b)$  for some  $\epsilon \in \{\pm 1\}$  and  $b \in \mathbb{Z}^2$ . By parts (i) and (ii) of Lemma 4.1, we find that  $g_a^{12N}|_\gamma = g_{a\gamma}^{12N}$  is equal to  $g_{a * \gamma}^{12N}$ . Therefore,

$$g_{\mathcal{O}}^{12N}|_\gamma = \prod_{a \in \mathcal{O}} g_a^{12N}|_\gamma = \prod_{a \in \mathcal{O}} g_{a * \gamma}^{12N} = g_{\mathcal{O}}^{12N},$$

where the last equality uses the fact that the map  $\mathcal{O} \rightarrow \mathcal{O}$ ,  $a \mapsto a * \gamma$  is a bijection (since  $\mathcal{O}$  is a  $\Gamma$ -orbit). The remaining statement about the poles and zeros of  $g_{\mathcal{O}}^{12N}$  follows immediately since each  $g_a$  is holomorphic and non-zero on  $\mathbb{H}$ .  $\square$

Let  $P_1, \dots, P_r$  be the cusps of  $X_\Gamma$ . Choose a representative  $s_j \in \mathbb{Q} \cup \{\infty\}$  of each cusp  $P_j$  and a matrix  $A_j \in \mathrm{SL}_2(\mathbb{Z})$  satisfying  $A_j \cdot \infty = s_j$ . Let  $w_j$  be the *width* of the cusp  $P_j$ ; it is the smallest positive integer  $b$  such that  $A_j \begin{pmatrix} 1 & b \\ 0 & 1 \end{pmatrix} A_j^{-1}$  is an element of  $\Gamma$ .

For a non-zero meromorphic function  $f$  of  $\mathbb{H}$  given by a  $q$ -expansion, we define  $\mathrm{ord}_q(f)$  to be the smallest rational number  $m$  such that there is a non-zero term of the form  $q^m$  in the expansion of  $f$ . For each cusp  $P_j$ , define the map

$$v_{P_j}: \mathbb{C}(X_\Gamma)^\times \rightarrow \mathbb{Z}, \quad f \mapsto w_j \cdot \mathrm{ord}_q(f|_{A_j});$$

it is a surjective homomorphism and agrees with the valuation giving the order of vanishing of a function at  $P_j$ . We extend  $\mathrm{ord}_q$  and  $v_{P_j}$  by setting  $\mathrm{ord}_q(0) = +\infty$  and  $v_{P_j}(0) = +\infty$ .

We now give a computable expression for the divisor of  $g_{\mathcal{O}}^{12N}$  on  $X_\Gamma$ .

**Lemma 4.3.** *With notation as above, we have*

$$\mathrm{div}(g_{\mathcal{O}}^{12N}) = \sum_{j=1}^r \left( 6Nw_j \sum_{a \in \mathcal{O}} B_2(\langle (aA_j)_1 \rangle) \right) \cdot P_j,$$

where  $B_2(x) = x^2 - x + 1/6$ ,  $(aA_j)_1$  is the first coordinate of the row vector  $aA_j$ , and  $\langle x \rangle$  denotes the positive fractional part of the real number  $x$ , chosen so  $0 \leq \langle x \rangle < 1$  and  $x - \langle x \rangle \in \mathbb{Z}$ .

*Proof.* For any  $a \in (N^{-1}\mathbb{Z}^2) - \mathbb{Z}^2$ , we have  $\mathrm{ord}_q(g_a) = \frac{1}{2} \cdot B_2(\langle a_1 \rangle)$ , cf. [KL81, p. 31]. We have

$$v_{P_j}(g_{\mathcal{O}}^{12N}) = \sum_{a \in \mathcal{O}} v_{P_j}(g_a^{12N}) = \sum_{a \in \mathcal{O}} w_j \mathrm{ord}_q(g_a^{12N}|_{A_j}) = \sum_{a \in \mathcal{O}} w_j \mathrm{ord}_q(g_{aA_j}^{12N}),$$

where the last equality uses Lemma 4.1(iii). Therefore,

$$v_{P_j}(g_{\mathcal{O}}^{12N}) = \sum_{a \in \mathcal{O}} 12Nw_j \mathrm{ord}_q(g_{aA_j}) = 6Nw_j \sum_{a \in \mathcal{O}} B_2(\langle (aA_j)_1 \rangle).$$

Since the poles and zeros of  $g_{\mathcal{O}}^{12N}$  are all cusps, we have  $\mathrm{div}(g_{\mathcal{O}}^{12N}) = \sum_{i=1}^r v_{P_i}(g_{\mathcal{O}}^{12N}) \cdot P_i$ , and the lemma follows immediately.  $\square$

**4.3. Constructing hauptmoduls of prime power level.** Fix a congruence subgroup  $\Gamma$  of  $\mathrm{SL}_2(\mathbb{Z})$  of *prime power* level  $N > 1$  that has genus 0. Let  $P_1, \dots, P_r$  be the cusps of  $\Gamma$ ; we choose our cusps so that  $P_1$  is the cusp at  $\infty$ .

In this section, we explain how to construct an explicit hauptmodul of  $\Gamma$  whose  $q$ -expansion has coefficients in  $K_N$ . Moreover, our hauptmodul will be of the form

$$(4.3) \quad \sum_{i=1}^M \zeta_{2N^2}^{e_i} \prod_{a \in \mathcal{A}_N} g_a^{m_{a,i}}$$

with integers  $m_{a,i}$  and  $e_i$ .

**4.3.1. Case 1: multiple cusps.** First assume that  $\Gamma$  has at least two cusps. We will use the following lemma to construct a hauptmodul for certain genus 0 congruence subgroups.

Let  $\mathcal{O}_1, \dots, \mathcal{O}_n$  be the distinct  $\Gamma$ -orbits of  $\mathcal{A}_N$ . For each  $\mathcal{O}_i$ , define the divisor  $D_i := \mathrm{div}(g_{\mathcal{O}_i}^{12N})$  on  $X_\Gamma$ . By Lemma 4.3, the divisors  $D_1, \dots, D_n$  are supported on  $\{P_1, \dots, P_r\}$  and are straightforward to compute.

**Lemma 4.4.** *Suppose there is an  $n$ -tuple  $m \in \mathbb{Z}^n$  such that*

$$\sum_{i=1}^n m_i D_i = -12N \cdot P_1 + 12N \cdot P_2$$

Let  $0 \leq e < 2N^2$  be the integer satisfying  $e \equiv \sum_{i=1}^n m_i \sum_{a \in \mathcal{O}_i} Na_2(N - Na_1) \pmod{2N^2}$ . Then

$$h := \zeta_{2N^2}^e \prod_{i=1}^n g_{\mathcal{O}_i}^{m_i}$$

is a hauptmodul for  $\Gamma$  whose  $q$ -expansion has coefficients in  $K_N$ . On  $X_\Gamma$ , we have  $\operatorname{div}(h) = -P_1 + P_2$ .

*Proof.* Since  $X_\Gamma$  has genus 0, there is a meromorphic function  $f$  on  $X_\Gamma$  with  $\operatorname{div}(f) = -P_1 + P_2$ . Lemma 4.2 implies that  $f^{12N}/h^{12N}$  defines a function on  $X_\Gamma$ ; it has divisor

$$12N \operatorname{div}(f) - \sum_{i=1}^n m_i \operatorname{div}(g_{\mathcal{O}_i}^{12N}) = 12N(-P_1 + P_2) - \sum_{i=1}^n m_i D_i = 0,$$

where the last equality uses our assumption on  $m$ . Therefore,  $f^{12N}/h^{12N}$  is constant. Since  $f$  and  $h$  are meromorphic functions on the upper half-plane, we deduce that  $f/h$  is a (non-zero) constant. In particular,  $h$  is modular for  $\Gamma$  and  $\operatorname{div}(h) = -P_1 + P_2$ . The function  $h$  on  $X_\Gamma$  is a hauptmodul for  $\Gamma$  since its only pole is the simple pole at  $P_1$ , i.e., the cusp at  $\infty$ .

It remains to show that the coefficients of  $h$  lie in  $K_N$ . Take any  $a \in \mathcal{A}_N$ . From the series defining  $g_a$ , we find that it is equal to the root of unity  $e(a_2(a_1 - 1)/2) = \zeta_{2N^2}^{Na_2(Na_1 - N)}$  times a Laurent series in  $q^{1/(6N^2)}$  with coefficients in  $K_N$ . Set  $e' := \sum_{i=1}^n m_i \sum_{a \in \mathcal{O}_i} Na_2(Na_1 - N)$ . The coefficients of  $\zeta_{2N^2}^{-e'} \prod_{i=1}^n g_{\mathcal{O}_i}^{m_i}$  thus all lie in  $K_N$ . The lemma follows since  $e \equiv -e' \pmod{2N^2}$ .  $\square$

Using the Cummins-Pauli classification of genus 0 congruence subgroups [CP03], we have explicitly verified that the  $n$ -tuple  $m$  from Lemma 4.4 always exists. Using Lemma 4.3, the existence of  $m$  comes down to finding integral solutions to  $r$  linear equations with integer coefficients in  $n$  variables. Using Lemma 4.4, we can thus find an explicit hauptmodul for  $\Gamma$  of the form (4.3) with  $M = 1$  (we have  $m_{a,i} = m_i$  if  $a \in \mathcal{O}_i$ ).

*Remark 4.5.* One can also abstractly prove the existence of the  $n$ -tuple  $m$ . If  $N$  is an odd prime power, then any modular function of level  $N$  whose zeros and poles are all cusps can be expressed as a constant times a product of Siegel functions  $g_a$  with  $a \in N^{-1}\mathbb{Z}^2 - \mathbb{Z}$ ; cf. Theorem 1.1(i) of Chapter 5 of [KL81].

If  $N \geq 4$  is a power of 2, this can also be deduced from Theorem 1.1(i) of Chapter 5 of [KL81]. (One needs to be a little careful here since  $g_a$  has a different definition in [KL81, Ch. 4 §1] when  $2a \in \mathbb{Z}$ . For the alternate  $g_a$  from loc. cit. with  $2a \in \mathbb{Z}$ , one can express them as a constant times a product of Siegel functions  $g_{a'}$  with  $a' \in \mathcal{A}_4 \subseteq \mathcal{A}_N$ .)

The case  $N = 2$  can be handled directly. For example, one can show that

$$g_{(1/2,0)}^8 \cdot g_{(1/2,1/2)}^4 \quad \text{and} \quad g_{(1/2,0)}^{12} \cdot g_{(1/2,1/2)}^{12}$$

are hauptmoduls for  $\Gamma(2)$  and  $\Gamma_0(2)$ , respectively (note that  $\Gamma_{\text{ns}}(2)$  has a single cusp and does not fall into this case, it falls into case 2 below).

**4.3.2. Case 2: a single cusp and  $N \neq 11$ .** Now assume that  $X_\Gamma$  has a single cusp and that  $N \neq 11$ . There are no non-constant modular functions for  $\Gamma$  whose zeros and poles are only at the cusps of  $X_\Gamma$ . In particular, a hauptmodul of  $\Gamma$  will never be equal to a product of Siegel functions.

Using the Cummins-Pauli classification, we find that there is a congruence subgroup  $\Gamma'$  that is a proper normal subgroup of  $\Gamma$ , also of level  $N$  and containing  $-I$ , such that  $X_{\Gamma'}$  has genus 0 and has exactly  $[\Gamma : \Gamma']$  cusps. (This is where we use  $N \neq 11$ .)

Since  $X_{\Gamma'}$  has multiple cusps, we know from §4.3.1 how to construct a hauptmodul  $h'$  of  $\Gamma'$  with coefficients in  $K_N$  that is of the form (4.3). Using that  $\Gamma'$  is normal in  $\Gamma$ , we find that  $h'|_\gamma$  is

modular for  $\Gamma'$  for all  $\gamma \in \Gamma$  and the function depends only on the coset  $\Gamma' \cdot \gamma$ . Define

$$h := \sum_{\gamma \in \Gamma' \backslash \Gamma} h'|_{\gamma};$$

it is a modular function for  $\Gamma$ . Since  $X_{\Gamma}$  has only one cusp and  $X_{\Gamma'}$  has  $[\Gamma : \Gamma']$  cusps, we deduce that the modular functions  $\{h'|_{\gamma}\}_{\gamma \in \Gamma' \backslash \Gamma}$  on  $X_{\Gamma'}$  each have their unique (simple) pole at different cusps. This implies that  $h$  has a simple pole at the unique cusp of  $X_{\Gamma}$  and is holomorphic elsewhere. Therefore,  $h$  is a hauptmodul for  $\Gamma$ .

Since  $h'$  is modular for  $\Gamma(N)$  and has coefficients in  $K_N$ , so does  $h'|_{\gamma}$  for all  $\gamma \in \text{SL}_2(\mathbb{Z})$ ; see Proposition 2.2. Therefore, the coefficients of  $h$  lie in  $K_N$ .

Finally, it remains to show that  $h$  is of the form (4.3). It suffices to show that  $h'|_{\gamma}$  is of the form (4.3) for a fixed  $\gamma \in \Gamma$ . We know that  $h'$  is equal to some product  $\zeta_{2N^2}^e \prod_{a \in \mathcal{A}_N} g_a^{m_a}$ , so  $h|_{\gamma} = \varepsilon(\gamma)^b \zeta_{2N^2}^e \prod_{a \in \mathcal{A}_N} g_{a\gamma}^{m_a}$  with  $b := \sum_{a \in \mathcal{A}_N} m_a$  by Lemma 4.1(iii). Recall that for each  $a \in \mathcal{A}_N$ , there is a unique  $a * \gamma \in \mathcal{A}_N$  such that  $a\gamma$  lies in the same coset of  $(N^{-1}\mathbb{Z}^2)/\mathbb{Z}^2$  as  $a * \gamma$  or  $-a * \gamma$ . From Lemma 4.1(i) and (ii), the functions  $g_{a\gamma}^{m_a}$  and  $g_{a*\gamma}^{m_a}$  agree up to a multiplication by some computable root of unity  $-\zeta_N^e$ . Therefore,  $h|_{\gamma}$  is equal to  $\varepsilon(\gamma)^b$  times a function of the form (4.3) with  $M = 1$ .

It remains only to show that  $\varepsilon(\gamma)^b$  is a power of a  $2N^2$ -th root of unity. In [KL81, Ch. 3 §5], Kubert and Lang give a necessary and sufficient condition for the product  $\prod_{a \in \mathcal{A}_N} g_a^{m_a}$  to be modular for  $\Gamma(N)$ ; these conditions hold since  $h'$  is modular for  $\Gamma' \supseteq \Gamma(N)$ . If  $N$  is a power of a prime  $\ell \geq 5$ , then Theorem 5.2 of Chapter 3 of [KL81] implies that  $b \equiv 0 \pmod{12}$  and hence  $\varepsilon(\gamma)^b = 1$ . If  $N$  is a power of 3, then Theorem 5.3 of Chapter 3 of [KL81] implies that  $b \equiv 0 \pmod{4}$  and hence  $\varepsilon(\gamma)^b$  is a power of  $\zeta_3$ . If  $N$  is a power of 2, then Theorem 5.3 of Chapter 3 of [KL81] implies that  $b \equiv 0 \pmod{3}$  and hence  $\varepsilon(\gamma)^b$  is a power of  $\zeta_4$ . Therefore,  $\varepsilon(\gamma)^b$  is indeed a power of a  $2N^2$ -th root of unity.

4.3.3. *Case 3:  $N = 11$ .* The remaining case is when  $X_{\Gamma}$  has a single cusp and  $N = 11$ . We include this case only for completeness; we will not need it for our application.

Define the function

$$f(\tau) := \prod_{(a_1, a_2) \in B} g_{(a_1/11, a_2/11)}(\tau),$$

where

$$B := \left\{ \begin{array}{l} (0, 1), (0, 2), (0, 3), (1, 0), (1, 2), (1, 5), (1, 7), (2, 1), (2, 2), (2, 4), (2, 5), (2, 6), \\ (2, 7), (2, 8), (2, 9), (2, 10), (3, 0), (3, 2), (3, 4), (3, 5), (3, 6), (3, 8), (3, 10), (4, 0), \\ (4, 1), (4, 2), (4, 4), (4, 5), (4, 6), (5, 1), (5, 4), (5, 5), (5, 6), (5, 7), (5, 8), (5, 9) \end{array} \right\}.$$

One can verify that

$$\sum_{(a_1, a_2) \in B} a_1^2 \equiv \sum_{(a_1, a_2) \in B} a_2^2 \equiv \sum_{(a_1, a_2) \in B} a_1 a_2 \equiv 0 \pmod{11}$$

and that  $|B| = 36 \equiv 0 \pmod{12}$ . Theorem 5.2 of [KL81, Ch. 3 §5] implies that  $f$  is a modular function for  $\Gamma(11)$ . Using  $\sum_{(a_1, a_2) \in B} a_2/11 \cdot (a_1/11 - 1)/2 = -60/11$  and the  $q$ -expansion of Siegel functions from §4.1, we find that all the coefficients of  $f$  lie in  $K_{11}$ . Therefore,  $f \in \mathcal{F}_{11}$ .

Using that  $\Gamma(11)$  is normal in  $\Gamma$ , we find that  $f|_{\gamma}$  is modular for  $\Gamma(11)$  for all  $\gamma \in \Gamma$  and the function depends only on the coset  $\Gamma(11) \cdot \gamma$ . Define

$$h := \sum_{\gamma \in \Gamma(11) \backslash \Gamma} f|_{\gamma};$$

it is a modular function for  $\Gamma$ . That  $h$  is of the form (4.3) follows as in the previous case.

We claim that  $h$  is a hauptmodul for  $\Gamma$ . From our description of  $h$  in terms of Siegel functions, we find that  $h$  has no poles except possibly at the unique cusp (at  $\infty$ ). From Cummins and Pauli [CP03], there is a unique genus 0 congruence subgroup of  $\mathrm{SL}_2(\mathbb{Z})$  of level 11 up to conjugacy in  $\mathrm{GL}_2(\mathbb{Z})$  (the one labeled 11A<sup>0</sup>). We have computed all the possible  $\Gamma$  and shown that  $h$  has a simple pole at  $\infty$ , and is therefore a hauptmodul.

*Remark 4.6.* The set  $B$  comes from §5.3 of [CLY04]. In [CLY04], methods are given to compute hauptmoduls for genus 0 congruence subgroups (unfortunately, their hauptmodul tables are no longer available). They use “generalized Dedekind eta functions” which are essentially Siegel functions.

**4.4. Rational function  $J(t)$ .** For a hauptmodul  $h$  of  $\Gamma$ , there is a unique function  $J(t) \in \mathbb{C}(t)$  such that  $J(h) = j$ ; it has degree  $d := [\mathrm{SL}_2(\mathbb{Z}) : \pm\Gamma]$ .

Let us briefly explain how to compute  $J(t)$  assuming that one can compute sufficiently many terms of the expansion of  $f$ . Let  $K \subseteq \mathbb{C}$  be a field containing all the coefficients of  $h$ . Consider the equation

$$(4.4) \quad (a_d h^d + \dots + a_1 h + a_0) - j \cdot (b_d h^d + \dots + b_1 h + b_0) = 0$$

with unknowns  $a_i, b_i \in K$ , where  $d := [\mathrm{SL}_2(\mathbb{Z}) : \pm\Gamma]$ . Computing the  $q$ -expansion coefficients of the left-hand side of (4.4) yields a system of homogeneous linear equations in the unknowns  $a_i$  and  $b_i$ . The existence and uniqueness of  $J$  ensure that the solutions  $(a_1, \dots, a_d, b_1, \dots, b_d) \in K^{2d}$  form a one-dimensional subspace. By computing sufficiently many coefficients of (4.4) one can find a non-zero solution  $(a_1, \dots, a_d, b_1, \dots, b_d) \in K^{2d}$ , unique up to scaling by  $K^\times$ , and

$$J(t) = (a_d t^d + \dots + a_1 t + a_0) / (b_d t^d + \dots + b_1 t + b_0) \in K(t)$$

is then the unique rational function for which  $J(h) = j$ . Note that if the hauptmodul  $h$  is constructed as in the previous section then we will have  $J(t) \in K_N(t)$ , where  $N$  is the level of  $\Gamma$ .

## 5. MODULAR CURVES OF GENUS 0

Fix the following:

- An integer  $N > 1$  that is a prime power.
- A subgroup  $G$  of  $\mathrm{GL}_2(\mathbb{Z}/N\mathbb{Z})$  satisfying  $-I \in G$  and  $\det(G) = (\mathbb{Z}/N\mathbb{Z})^\times$ .
- A rational function  $J(t) \in \mathbb{Q}(t)$ .

In this section, we explain how to determine if the function field of  $X_G$  is of the form  $\mathbb{Q}(f)$  for some modular function  $f \in \mathcal{F}_N$  satisfying  $J(f) = j$ . We will use this to verify the entries of Tables 1–3.

If such an  $f$  exists, then  $X_G$  is isomorphic to  $\mathbb{P}_{\mathbb{Q}}^1$  and the morphism  $\pi_G: X_G \rightarrow \mathbb{P}_{\mathbb{Q}}^1$  is given by the relation  $j = J(f)$  in their function fields. We may assume the necessary condition that  $[\mathrm{GL}_2(\mathbb{Z}/N\mathbb{Z}) : G] = \deg \pi_G$  agrees with the degree of  $J(t)$ .

*Remark 5.1.* In §8 we explain how the  $J(t)$  listed in Tables 1–3, were actually found, which involves the use of a Monte Carlo algorithm and assumes the Generalized Riemann Hypothesis (GRH). The purpose of this section is to explain how we can unconditionally verify a given  $J(t)$ , regardless of how it was found.

**5.1. Construction of possible  $f$ .** Let  $\Gamma$  be the congruence subgroup consisting of  $\gamma \in \mathrm{SL}_2(\mathbb{Z})$  for which  $\gamma^t$  modulo  $N$  lies in  $G$  (equivalently, in  $G \cap \mathrm{SL}_2(\mathbb{Z}/N\mathbb{Z})$ ). By Lemma 2.4(ii), we may assume that  $\Gamma$  has genus 0 since otherwise  $X_G$  has positive genus and its function field cannot be of the form  $\mathbb{Q}(f)$ .

The group  $\Gamma$  acts on the right on the field  $\mathcal{F}_N$ ; let  $\mathcal{F}_N^\Gamma$  be subfield fixed by this action. By Lemma 2.4(i), we have  $K_N(X_G) = \mathcal{F}_N^\Gamma$ .

In §4.3, we described how to compute an explicit hauptmodul  $h$  for  $\Gamma$  such that coefficients of its  $q$ -expansion all lie in  $K_{N'} \subseteq K_N$ , where the level  $N'$  of  $\Gamma$  divides  $N$ . Therefore, we have

$$K_N(X_G) = \mathcal{F}_N^\Gamma = K_N(h).$$

Moreover, we can express  $h$  in terms of Siegel functions and hence we can compute as many of its coefficients as we desire. In §4.4, we described how to compute the unique rational function  $J'(t) \in K_N(t)$  for which  $j = J'(h)$ . The degree of  $J'(t)$  agrees with  $[\mathrm{SL}_2(\mathbb{Z}) : \Gamma] = [\mathrm{GL}_2(\mathbb{Z}/N\mathbb{Z}) : G]$ , thus  $J(t)$  and  $J'(t)$  have the same degree.

*Remark 5.2.* The rational function  $J'(t)$  gives a map to the  $j$ -line from  $X_\Gamma$ , which is defined over  $K_N = \mathbb{Q}(\zeta_N)$ , while the rational function  $J(t)$  gives a map to the  $j$ -line from  $X_G$ , which is defined over  $\mathbb{Q}$ . We use  $J'(t)$  in our procedure to verify  $J(t)$ , but note that  $J'(t)$  does not determine  $J(t)$ ; in general there will be multiple non-conjugate subgroups  $G$  corresponding to  $\Gamma$  and a different rational function  $J(t)$  for each of the corresponding  $X_G$  (in total we have 220 modular curves  $X_G$  of genus 0 corresponding to 73 modular curves  $X_\Gamma$ ).

**Lemma 5.3.** *The modular functions  $f \in K_N(X_G)$  that satisfy  $K_N(X_G) = K_N(f)$  and  $J(f) = j$  are precisely those of the form  $\psi(h)$ , where  $\psi(t) \in K_N(t)$  is a degree 1 function satisfying  $J'(t) = J(\psi(t))$ .*

*Proof.* First take any  $\psi(t) \in K_N(t)$  of degree 1 satisfying  $J'(t) = J(\psi(t))$ . Define  $f := \psi(h)$ . We have  $K_N(f) = K_N(h) = K_N(X_G)$ , since  $\psi$  has an inverse, and  $J(f) = J(\psi(h)) = J'(h) = j$ .

Now suppose that  $K_N(X_G) = K_N(f)$  for some  $f \in K_N(X_G)$  satisfying  $J(f) = j$ . Since  $K_N(f) = K_N(X_G) = K_N(h)$ , we have  $f = \psi(h)$  for a unique  $\psi(t) \in K_N(t)$  of degree 1. We then have  $j = J(f) = J(\psi(h))$  and therefore  $J'(t) = J(\psi(t))$ , since  $J'(t)$  is the unique element of  $K_N(t)$  that satisfies  $J'(h) = j$ .  $\square$

**5.2. Finding possible  $f$ .** Define  $\Psi$  to be the set of  $\psi(t) \in K_N(t)$  of degree 1 for which  $J'(t) = J(\psi(t))$ ; these  $\psi$  arise in Lemma 5.3. We now explain how to compute  $\Psi$ .

Choose three distinct elements  $\beta_1, \beta_2, \beta_3 \in K_N \cup \{\infty\}$ . For  $1 \leq i \leq 3$ , define the set

$$R_i := \{\alpha \in K_N \cup \{\infty\} : J'(\beta_i) = J(\alpha) \text{ and } \mathrm{ord}_{\beta_i}(J') = \mathrm{ord}_\alpha(J)\},$$

where  $\mathrm{ord}_{\beta_i}(J')$  is the order of vanishing of  $J'(t)$  at  $t = \beta_i$ . Let  $R$  be the set of triples  $\alpha = (\alpha_1, \alpha_2, \alpha_3) \in R_1 \times R_2 \times R_3$  such that  $\alpha_1, \alpha_2$  and  $\alpha_3$  are distinct. Let  $\psi_\alpha \in K_N(t)$  be the *unique* rational function of degree 1 such that  $\psi_\alpha(\beta_i) = \alpha_i$  for all  $1 \leq i \leq 3$ .

Take any  $\psi \in \Psi$ . We have  $J'(\beta_i) = J(\psi(\beta_i))$  and  $\mathrm{ord}_{\beta_i}(J') = \mathrm{ord}_{\psi(\beta_i)}(J)$  for each  $1 \leq i \leq 3$ . Therefore,  $\psi(\beta_i) \in R_i$  for each  $1 \leq i \leq 3$  and hence  $\psi = \psi_\alpha$  for some  $\alpha \in R$ . So we have

$$\Psi = \{\psi_\alpha : \alpha \in R, J'(t) = J(\psi(t))\}.$$

Since  $R$  is finite, this gives us a way to compute the (finite) set  $\Psi$ .

By Lemma 5.3, the set

$$\{\psi(h) : \psi \in \Psi\}$$

is the set of modular functions  $f \in K_N(X_G)$  that satisfy  $K_N(X_G) = K_N(f)$  and  $J(f) = j$ .

**5.3. Checking each  $f$ .** Let  $f$  be one of the finite number of functions that satisfy  $K_N(X_G) = K_N(f)$  and  $J(f) = j$ . We just saw how to compute all such  $f$ ; they are of the form  $\psi(h)$  for a degree 1 function  $\psi(t) \in K_N(t)$  and a modular function  $h$  satisfying  $K_N(X_G) = K_N(h)$  that is expressed in terms of Siegel functions. Recall from §2.4 that each  $A \in \mathrm{GL}_2(\mathbb{Z}/N\mathbb{Z})$  acts on  $\mathcal{F}_N$  via the isomorphism  $\theta_N : \mathrm{GL}_2(\mathbb{Z}/N\mathbb{Z})/\{\pm I\} \xrightarrow{\sim} \mathrm{Gal}(\mathcal{F}_N/\mathbb{Q}(j))$  of Proposition 2.2, and for  $f \in \mathcal{F}_N$  we use  $A_*(f) := \theta_N(A)(f)$  to denote this action.

**Lemma 5.4.**

- (i) We have  $\mathbb{Q}(X_G) = \mathbb{Q}(f)$  if and only if  $f \in \mathbb{Q}(X_G)$ .
- (ii) For a matrix  $A \in G$ , we have  $A_*(f) = f$  if and only if  $\text{ord}_q(A_*(f) - f) > 2w/N'$ , where  $w$  is the width of the cusp  $\infty$  of  $X_\Gamma$  and  $N'$  is the level of  $\Gamma$ .

*Proof.* We first prove part (i); only one implication needs proof. Suppose that  $f \in \mathbb{Q}(X_G)$ . Then  $\mathbb{Q}(f) \subseteq \mathbb{Q}(X_G)$  and it suffices to show that these two fields have the same degree over  $\mathbb{Q}(j)$ . This is true since we have been assuming that  $\deg \pi_G$  is equal to the degree of  $J(t)$ .

We now prove part (ii); only one implication needs proof. Suppose that  $\text{ord}_q(A_*(f) - f) > 2w/N'$ . As meromorphic functions on  $X_\Gamma$ ,  $f$  and  $A_*(f)$  have a unique (simple) pole since  $h$  has this property and  $\psi$  has degree 1. Therefore, the function  $A_*(f) - f$  on  $X_\Gamma$  is zero or has at most two poles (and hence at most two zeros). Our assumption  $\text{ord}_q(A_*(f) - f) > 2w/N'$  implies that  $A_*(f) - f$  has a zero of order 3 at the cusp  $\infty$  and thus  $A_*(f) - f = 0$ .  $\square$

By Lemma 5.4(i), we have  $\mathbb{Q}(X_G) = \mathbb{Q}(f)$  if and only if  $A_*(f) = f$  for all  $A \in G$  in a set of generators of  $G$ ; it suffices to consider  $A \in G$  for which  $\det(A)$  generate  $(\mathbb{Z}/N\mathbb{Z})^\times$  since  $h$  and hence  $f$  is fixed by  $G \cap \text{SL}_2(\mathbb{Z}/N\mathbb{Z})$ . It remains to describe how to determine whether  $A_*(f)$  is equal to  $f$ . By Lemma 5.4(ii), it suffices to compute enough terms of the  $q$ -expansion of  $A_*(f) - f$  to determine whether  $\text{ord}_q(A_*(f) - f) > 2w/N'$  holds.

Finally, let us briefly explain how to compute terms in the  $q$ -expansion of  $A_*(f) - f$ . Let  $d$  be an odd integer congruent to  $\det(A)$  modulo  $N$ . Choose a matrix  $\gamma \in \text{SL}_2(\mathbb{Z})$  so that  $A^t \equiv \begin{pmatrix} 1 & 0 \\ 0 & d \end{pmatrix} \gamma \pmod{N}$ . We thus have

$$(5.1) \quad A_*(f) - f = \sigma_d(f)|_\gamma - f = \sigma_d(\psi)(\sigma_d(h)|_\gamma) - \psi(h),$$

where  $\sigma_d(\psi)$  is the rational function with  $\sigma_d$  applied to the coefficients of its numerator and denominator. Our hauptmodul  $h$  is of the form  $\sum_{i=1}^M \zeta_{2N^2}^{e_i} \prod_{a \in \mathcal{A}_N} g_a^{m_{a,i}}$  for certain integers  $e_i$  and  $m_{a,i}$ , so

$$\sigma_d(h)|_\gamma = \sum_{i=1}^M \zeta_{2N^2}^{e_i d} \prod_{a \in \mathcal{A}_N} (\sigma_d(g_a)|_\gamma)^{m_{a,i}}.$$

From the series expansion of  $g_a$ , one easily checks that  $\sigma_d(g_{(a_1, a_2)}) = g_{(a_1, da_2)}$ . From Lemma 4.1(iii), we have  $\sigma_d(g_a)|_\gamma = \varepsilon(\gamma)g_{(a_1, da_2)\gamma}$  and hence

$$\sigma_d(h)|_\gamma = \sum_{i=1}^M \zeta_{2N^2}^{e_i d} \cdot \prod_{a \in \mathcal{A}_N} \varepsilon(\gamma)^{m_{a,i}} \cdot \prod_{a \in \mathcal{A}_N} g_{(a_1, da_2)\gamma}^{m_{a,i}}.$$

Thus by computing enough terms in the  $q$ -expansion of the functions  $\{g_a\}_{a \in \mathcal{A}_N}$ , we are able to compute the  $q$ -expansion of  $h$  and  $\sigma_d(h)|_\gamma$  to as many terms as we desire. This allows us to compute terms in the  $q$ -expansion of  $A_*(f) - f$  via (5.1).

*Remark 5.5.* Suppose that  $X_\Gamma$  has at least 3 cusps. We then have  $A_*(f) = f$  if and only if  $A_*(f)$  and  $f$  take the same the value at any three of the cusps (as in the proof of Lemma 5.4 this implies that  $A_*(f) - f$  has at least three zeros and hence is the zero function). In the case of at least 3 cusps, our hauptmodul  $h$  was given as a constant times a product of Siegel functions; so its value at the cusp  $\infty$  is determined by the first term of the  $q$ -expansion of  $h$ . The value at any other cusp  $c$  can be determined by the first term of the  $q$ -expansion of  $h|_\gamma$  with  $\gamma \in \text{SL}_2(\mathbb{Z})$  satisfying  $\gamma\infty = c$ . This approach is quicker since fewer terms of the  $q$ -expansions are required.

**5.4. Verifying the entries of our tables.** We now explain how to verify the validity of our genus 0 tables. `Magma` scripts that perform these verifications can be found at [SZ16].

Each row of Tables 1–3 gives a set of generators of a subgroup  $G$  of  $\mathrm{GL}_2(\mathbb{Z}/N\mathbb{Z})$  that satisfies  $-I \in G$  and  $\det(G) = (\mathbb{Z}/N\mathbb{Z})^\times$  for a prime power  $N$ . We may assume that  $N > 1$ . By composing rational maps, we obtain a corresponding rational function  $J(t) \in \mathbb{Q}(t)$ .

Using the earlier parts of §5, we can construct a modular function  $f \in \mathcal{F}_N$  such that  $\mathbb{Q}(X_G) = \mathbb{Q}(f)$  and  $J(f) = j$ . So  $X_G$  is isomorphic to  $\mathbb{P}_\mathbb{Q}^1$  and the morphism  $\pi_G: X_G \rightarrow \mathbb{P}_\mathbb{Q}^1$  is given by the relation  $j = J(f)$  in their function fields. (We also note that there is no harm in replacing  $G$  by a conjugate group; this is useful because one can reuse the hauptmodul computations for different groups in the tables).

Fix a group  $G \subseteq \mathrm{GL}_2(\mathbb{Z}/N\mathbb{Z})$  as above, and a modular function  $f \in \mathcal{F}_N$  satisfying  $\mathbb{Q}(X_G) = \mathbb{Q}(f)$  and  $J(f) = j$ .

Now fix another group  $G' \subseteq \mathrm{GL}_2(\mathbb{Z}/N'\mathbb{Z})$  from our table so that  $N$  divides  $N'$  and the image of  $G'$  in  $\mathrm{GL}_2(\mathbb{Z}/N\mathbb{Z})$  is conjugate to a subgroup of  $G$ . In the above computations, we have constructed a modular function  $f'$  satisfying  $\mathbb{Q}(X_{G'}) = \mathbb{Q}(f')$  and  $J'(f') = j$  for a rational function  $J'(t) \in \mathbb{Q}(t)$  also arising from the tables.

Take any subgroup  $\tilde{G} \subseteq \mathrm{GL}_2(\mathbb{Z}/N\mathbb{Z})$  conjugate to  $G'$  whose image modulo  $N$  lies in  $G$ . Choose any  $A \in \mathrm{GL}_2(\mathbb{Z}/N'\mathbb{Z})$  for which  $\tilde{G} := AG'A^{-1}$  and define  $\tilde{f} := A_*(f')$ . We have an inclusion of fields

$$\mathbb{Q}(\tilde{f}) = \mathbb{Q}(X_{\tilde{G}}) \supseteq \mathbb{Q}(X_G) = \mathbb{Q}(f).$$

The extension  $\mathbb{Q}(\tilde{f})/\mathbb{Q}(f)$  has degree  $i := [\mathrm{GL}_2(\mathbb{Z}/N'\mathbb{Z}):G']/[\mathrm{GL}_2(\mathbb{Z}/N\mathbb{Z}):G]$ . Therefore,  $\varphi(\tilde{f}) = f$  for a unique  $\varphi(t) \in \mathbb{Q}(t)$  of degree  $i$ . We can compute  $\varphi(t)$  using the method from §4.4; the coefficients of  $f$  and  $\tilde{f}$  can be computed as in §5.3.

The rational function  $\varphi$  is not unique, it depends on the choices of  $\tilde{G}$ ,  $f$ ,  $f'$ , and  $A$ . However, any other rational function occurring would be of the form  $\psi'(\varphi(\psi(t)))$ , where  $\psi, \psi' \in \mathbb{Q}(t)$  are degree 1 functions satisfying  $J(\psi(t)) = J(t)$  and  $J'(\psi'(t)) = J'(t)$ . Note that all the possible  $\psi$  and  $\psi'$  can be computed as in §5.2 (with  $J = J'$ ). We have checked that the rational function relating  $G$  and  $G'$  in our tables, when given, is indeed of the form  $\psi'(\varphi(\psi(t)))$ .

## 6. MODULAR CURVES OF GENUS 1

We now consider the open subgroups  $G$  of  $\mathrm{GL}_2(\hat{\mathbb{Z}})$  with genus 1 and prime power level  $N = \ell^e$  that satisfy  $-I \in G$  and  $\det(G) = \hat{\mathbb{Z}}^\times$ . We are interested in describing those  $G$  for which  $X_G(\mathbb{Q})$  is infinite. There is no harm in replacing  $G$  by a conjugate. So by Theorem 3.3(ii), there are 250 cases that need to be checked.

Let  $J_G$  be the Jacobian of the curve  $X_G$ . Using the methods of [Zyw15b], we can compute the rank of  $J_G(\mathbb{Q})$ . From [DR73, §IV], we find that the curve  $X_G$  has good reduction at all primes  $p \nmid N = \ell^e$ . Therefore,  $J_G$  is an elliptic curve defined over  $\mathbb{Q}$  whose conductor is a power of  $\ell$ . The primes  $\ell$  that arise are small enough to ensure that  $J_G$  is isomorphic to one of the elliptic curves in Cremona's tables [Cre14]; this gives a finite number of candidates for  $J_G$  up to isogeny.

For each prime  $p \nmid 6\ell$ , we can compute  $\#J_G(\mathbb{F}_p) = \#X_G(\mathbb{F}_p)$  from the modular interpretation of  $X_G$ , cf. [Zyw15b, §3.6] for details. In particular, we can compute  $\#J_G(\mathbb{F}_p)$  directly from the group  $G$  without computing a model for  $X_G$  (or its reduction modulo  $p$ ). By computing several values of  $\#J_G(\mathbb{F}_p)$  with  $p \neq \ell$ , we can quickly distinguish the isogeny class of  $J_G$  among the finite set of candidates. We then compute the rank of  $J_G(\mathbb{Q})$ , which we note is an isogeny invariant.

Running this procedure on each of the 250 genus 1 groups  $G$  given by Theorem 3.3, we find that  $J_G(\mathbb{Q})$  has rank 0 for 222 groups and  $J_G(\mathbb{Q})$  has positive rank for 28 groups; a `Magma` script that

performs this computation can be found in [SZ16]. We need only consider the 28 groups  $G$  for which  $J_G(\mathbb{Q})$  has positive rank, since  $X_G(\mathbb{Q})$  is finite if  $J_G(\mathbb{Q})$  has rank 0.

Now let  $G$  be one of the 28 groups for which  $J_G(\mathbb{Q})$  has positive rank; they are precisely the 28 genus 1 groups in Theorem 1.1 and can be found in Table 4 in Appendix A. For each of these groups  $G$ , if  $X_G(\mathbb{Q})$  is nonempty then it must be infinite, since the Abel-Jacobi map then gives a bijection from  $X_G(\mathbb{Q})$  to  $J_G(\mathbb{Q})$ . We initially verified that  $X_G(\mathbb{Q})$  is nonempty by finding an elliptic curve  $E/\mathbb{Q}$  with  $\rho_E(\text{Gal}_{\mathbb{Q}}) \subseteq G$  using an extension of the algorithm in [Sut16].

For each of these 28 groups  $G$ , a model for  $X_G$  and the morphism  $\pi_G$  can already be found in the literature (and are equivalent to the ones we give in Appendix A). For the 27 groups  $G$  of level 16 these curves and morphisms were constructed by Rouse and Zureick-Brown in [RZB15]; the models and morphisms we give in Table 4 for these groups are slightly different (we constructed them by taking fiber products of our genus 0 curves), but we have verified that they are isomorphic (note that their groups are transposed relative to ours). The remaining group  $G$  has level 11 and its image in  $\text{GL}_2(\mathbb{Z}/11\mathbb{Z})$  is the normalizer of a non-split Cartan subgroup. An explicit model for  $X_G = X_{\text{ns}}^+(11)$  and the morphism to the  $j$ -line can be found in [Hal98]; these are reproduced in Appendix A.

## 7. PROOF OF THEOREM 1.4

If  $\ell \leq 13$ , then the set  $\mathcal{J}_\ell$  is finite by [Zyw15b, Proposition 4.8]. If  $\ell > 13$ , this follows from [Zyw15b, Proposition 4.9]; note that  $\rho_{E, \ell^\infty}$  is surjective if and only if  $\rho_{E, \ell}$  is surjective, since  $\ell \geq 5$ , by [Ser89, IV Lemma 3]. This proves (i).

For a group  $G$  from Theorem 1.1, define the set

$$\mathcal{S}_G := \bigcup_{G'} \pi_{G', G}(X_{G'}(\mathbb{Q})),$$

where  $G'$  varies over the proper subgroups of  $G$  that are conjugate to one of groups in Theorem 1.1 of  $\ell$ -power level and  $\pi_{G', G}: X_{G'} \rightarrow X_G$  is the natural morphism induced by the inclusion  $G' \subseteq G$ . Note that this is a finite union.

Suppose first that  $G$  has genus 0. Then  $X_G \simeq \mathbb{P}_{\mathbb{Q}}^1$  and  $\mathcal{S}_G$  is a *thin* subset of  $X_G(\mathbb{Q})$ , in the language of [Ser97, §9]. The field  $\mathbb{Q}$  is Hilbertian, and in particular  $\mathbb{P}_1(\mathbb{Q}) \simeq X_G(\mathbb{Q})$  is not thin; this implies that the complement  $X_G(\mathbb{Q}) - \mathcal{S}_G$  cannot be thin and must be infinite.

Suppose that  $G$  has genus 1. If  $G$  does not have level 16 and index 24, then there are no proper subgroups  $G'$  of  $G$  that are conjugate to a group from Theorem 1.1, and therefore  $\mathcal{S}_G$  is empty and  $X_G(\mathbb{Q}) - \mathcal{S}_G$  is infinite.

Now suppose that  $G$  has genus 1, level 16, and index 24. There are 7 such  $G$ , labeled

$$16C^1-16c, 16C^1-16d, 16B^1-16a, 16B^1-16c, 16D^1-16d, 8D^1-16b, 8D^1-16c$$

and explicitly described in Table 4 of Appendix A. Each of these  $G$  contains either 2 or 4 index 2 subgroups  $G'$  that are conjugate to one of the groups in Theorem 1.1. In every case we have  $\mathcal{S}_G = X_G(\mathbb{Q})$ , so that  $X_G(\mathbb{Q}) - \mathcal{S}_G$  is empty; see Example 6.11 and Remark 6.3 in [RZB15].

Let  $E/\mathbb{Q}$  be an elliptic curve with  $j_E \notin \mathcal{J}_\ell$ . The group  $\pm\rho_{E, \ell^\infty}(\text{Gal}_{\mathbb{Q}})$  is conjugate in  $\text{GL}_2(\mathbb{Z}_\ell)$  to the  $\ell$ -adic projection of a unique group  $G$  from Theorem 1.1 with  $\ell$ -power level. Using Proposition 2.6, we can also characterize  $G$  as the unique group from Theorem 1.1 with  $\ell$ -power level such that  $j_E \in \pi_G(X_G(\mathbb{Q}) - \mathcal{S}_G)$ . Parts (ii) and (iii) follow by noting that  $\pi_G(X_G(\mathbb{Q}) - \mathcal{S}_G)$  is empty when  $G$  has genus 1, level 16, and index 24, and it is infinite otherwise.

## 8. HOW THE $J(t)$ WERE FOUND

Let  $G$  be one of the genus 0 subgroups of  $\mathrm{GL}_2(\hat{\mathbb{Z}})$  from Theorem 1.1; they are listed in Tables 1–3 in Appendix A and were determined using the algorithm described in §3. For each  $G$ , we also have a rational function  $J(t) \in \mathbb{Q}(t)$  such that the function field of  $X_G$  is of the form  $\mathbb{Q}(f)$  and  $j = J(f)$ , where  $j$  is the modular  $j$ -invariant; the verification of this property is described in §5.

In this section, we explain how we found  $J(t)$ ; note that the method we used to verify the correctness of  $J(t)$  does not depend on how it was found! None of our theorems depend on the techniques described in this section. All that matters is that they eventually produced functions  $J(t)$  whose correctness we could verify using the procedure described in §5.4.

We used an extension of the algorithm in [Sut16] to search for elliptic curves  $E/\mathbb{Q}$  for which  $\rho_E(\mathrm{Gal}_{\mathbb{Q}})$  is conjugate to a subgroup of  $G$ . This was initially done by simply checking elliptic curves in Cremona’s tables [Cre14] and the LMFDB [LMF15] (but see Remark 8.1 below). After enough searching, we find elliptic curves  $E_1, E_2, E_3$  defined over  $\mathbb{Q}$  with distinct  $j$ -invariants  $j_1, j_2, j_3$  for which we believe that  $\rho_{E_i}(\mathrm{Gal}_{\mathbb{Q}})$  is conjugate in  $\mathrm{GL}_2(\hat{\mathbb{Z}})$  to a subgroup of  $G$ ; in particular, we expect that  $j_1, j_2, j_3 \in \pi_G(X_G(\mathbb{Q}))$ . We ran the Monte Carlo algorithm in [Sut16] using parameters that ensure the error probability is less than  $2^{-100}$ , under the GRH.

Now suppose that  $j_1, j_2, j_3$  are indeed elements of  $\pi_G(X_G(\mathbb{Q}))$ . The curve  $X_G$  has genus 0 and rational points, so it is isomorphic to  $\mathbb{P}_{\mathbb{Q}}^1$ . We can choose an isomorphism  $X_G \simeq \mathbb{P}_{\mathbb{Q}}^1$  such that there are points  $P_1, P_2, P_3 \in X_G(\mathbb{Q})$  satisfying  $\pi_G(P_i) = j_i$  which map to  $0, 1, \infty$ , respectively. There is thus a rational function  $J(t) \in \mathbb{Q}(t)$  such that  $J(0) = j_1, J(1) = j_2, J(\infty) = j_3$  and such that  $\mathbb{Q}(X_G) = \mathbb{Q}(f)$  for a modular function  $f$  satisfying  $J(f) = j$ ; the function  $f$  is obtained by composing our isomorphism  $\mathbb{P}_{\mathbb{Q}}^1 \simeq X_G$  with  $\pi_G$ .

We can now find all such potential  $J$ . As explained in §5, we can construct a modular function  $h \in \mathcal{F}_N$  and a rational function  $J'(t) \in K_N(t)$  such that  $K_N(X_G) = K_N(h)$  and  $j = J'(h)$ , where  $N$  is the level of  $G$ . We thus have

$$J(t) = J'(\psi(t))$$

for some degree 1 function  $\psi(t) \in K_N(t)$  satisfying  $\psi(0) \in R_1, \psi(1) \in R_2$  and  $\psi(\infty) \in R_3$ , where  $R_i := \{\alpha \in K_N \cup \{\infty\} : J'(\alpha) = j_i\}$ . Since the sets  $R_i$  are finite and disjoint, there are only finitely many  $\psi(t) \in \mathbb{Q}(t)$  of degree 1 satisfying  $\psi(0) \in R_1, \psi(1) \in R_2, \psi(\infty) \in R_3$ . For each such  $\psi(t)$ , we check whether  $J'(\psi(t))$  lies in  $\mathbb{Q}(t)$ .

Consider any  $\psi$  as above for which  $J'(\psi(t)) \in \mathbb{Q}(t)$ . Set  $J(t) := J'(\psi(t))$  and  $f := \psi^{-1}(h) \in K_N(X_G)$ . We have  $J(f) = J'(h) = j$ . The field  $\mathbb{Q}(f)$  is thus the function field of a modular curve  $X_{G'}$ , where  $G'$  is an open subgroup of  $\mathrm{GL}_2(\hat{\mathbb{Z}})$  of level  $N$  satisfying  $\det(G') = \hat{\mathbb{Z}}^\times$  and  $-I \in G'$ ; it consists of matrices whose reduction modulo  $N$  fix  $f$ . We can then check whether  $G$  is equal to  $G'$ . Since  $[\mathrm{GL}_2(\hat{\mathbb{Z}}) : G] = \deg \pi_G = \deg J = [\mathrm{GL}_2(\hat{\mathbb{Z}}) : G']$ , it suffices to determine whether  $G$  is a subgroup of  $G'$ ; equivalently, whether  $G$  fixes  $f$ . A method for determining whether  $f$  is fixed by  $G$  is described in §5.3.

We will eventually find a  $\psi$  for which we have  $G = G'$  (provided that our initial  $j$ -invariants  $j_i$  are valid). This then proves that  $\mathbb{Q}(X_G) = \mathbb{Q}(f)$  for some  $f$  satisfying  $J(f) = j$ , where  $J(t) := J'(\psi(t)) \in \mathbb{Q}(t)$ .

Note this rational function  $J(t)$  is not unique since  $J(\varphi(t))$  would also work for any  $\varphi(t) \in \mathbb{Q}(t)$  of degree 1. Using similar reasoning, it is easy to determine if two  $J_1, J_2 \in \mathbb{Q}(t)$  satisfy  $J_2(t) = J_2(\varphi(t))$  for some degree 1 function  $\varphi \in \mathbb{Q}(t)$ . We have chosen our rational functions so that they are relatively compact when written down.

*Remark 8.1.* Having run this procedure to obtain functions  $J(t)$  for each of the groups  $G$  where we were able to find suitable  $E_1, E_2, E_3$  in Cremona’s tables, we then address the remaining groups  $G$  by picking a group  $G'$  that contains a subgroup conjugate to  $G$  for which we already know a function

$J'(t) \in \mathbb{Q}(t)$ ; such a  $G'$  existed for every  $G$  not addressed in our initial search of Cremona's tables. Using the function  $J'(t)$  we can quickly obtain a large list of elliptic curves  $E$  for which  $\rho_E(\text{Gal}_{\mathbb{Q}})$  is a subgroup of  $G'$ . By running the algorithm in [Sut16] on several thousand (or even millions) of these curves we are eventually able to find  $E_1, E_2, E_3$  with distinct  $j$ -invariants for which it is highly probable that  $\rho_{E_i}(\text{Gal}_{\mathbb{Q}})$  is actually conjugate to a subgroup of the smaller group  $G$  contained in  $G'$ . We then proceed as above to compute the function  $J(t)$  for  $G$ .

## APPENDIX A. TABLES

This section includes tables that list data for all the groups  $G$ , up to conjugacy in  $\text{GL}_2(\hat{\mathbb{Z}})$ , from Theorem 1.1. The genus 0 groups are given in Tables 1, 2 and 3. The genus 1 groups are given in Table 4.

We now describe how to read Tables 1–4. Each row corresponds to a unique group  $G$  from Theorem 1.1 up to conjugacy; it is given a unique label of the form  $MZ^g-Nz$ , where  $M, N$  and  $g$  are integers and  $Z$  and  $z$  are letters. The integers  $N$  and  $g$  are the level and genus of  $G$ , respectively. Let  $\Gamma$  be the congruence subgroup consisting of matrices in  $\text{SL}_2(\mathbb{Z})$  whose image modulo  $N$  lies in the image of  $G$  modulo  $N$ . The integer  $g$  is also the genus of the Riemann surface obtained by taking the quotient of the complex upper-half plane by the action of  $\Gamma$ . The integer  $M$  is the level of  $\Gamma$  and  $Z$  is an uppercase letter that distinguishes  $\Gamma$  up to conjugacy in  $\text{GL}_2(\mathbb{Z})$ ; the prefix  $MZ^g$  for  $\Gamma$  matches the label used by Cummins and Pauli [CP03]. The letter  $z$  is chosen so that the label  $MZ^g-Nz$  distinguishes  $G$  up to conjugacy in  $\text{GL}_2(\hat{\mathbb{Z}})$ . In some of the tables, we also number the rows.

For each row of these tables, there is a positive integer  $N$  and a set of matrices  $S$  in  $\text{GL}_2(\mathbb{Z}/N\mathbb{Z})$ ; the corresponding open subgroup  $G$  of  $\text{GL}_2(\hat{\mathbb{Z}})$  consists of the matrices  $A \in \text{GL}_2(\hat{\mathbb{Z}})$  whose image in  $\text{GL}_2(\mathbb{Z}/N\mathbb{Z})$  lies in the subgroup generated by  $S$ . The integer  $i$  in each row is the index  $[\text{GL}_2(\hat{\mathbb{Z}}) : G]$ .

For each row of a table, we also have a list of minimal supergroups up to conjugacy (given by labels or numberings of other rows); more precisely, the groups  $G'$  satisfying  $G \subsetneq G' \subseteq \text{GL}_2(\hat{\mathbb{Z}})$ , up to conjugacy in  $\text{GL}_2(\hat{\mathbb{Z}})$ , for which there are no subgroups strictly between  $G$  and  $G'$ .

If  $G$  has genus 0, then  $X_G$  is isomorphic to  $\mathbb{P}_{\mathbb{Q}}^1$ ; its function field is of the form  $\mathbb{Q}(t)$ . If  $G$  has genus 1, then the curve  $X_G$  is isomorphic to the elliptic curve in Table 5 given by the ‘‘curve’’ column of Table 4; its function field is of the form  $\mathbb{Q}(x, y)$ , where  $x$  and  $y$  are the given Weierstrass coordinates.

Now for simplicity, suppose that  $G$  has genus 0. The function  $F$  listed in the ‘‘map’’ column describes the morphism  $X_G \rightarrow X_{G'}$  for one of the corresponding minimal supergroups  $G'$  of  $G$  (or the first minimal supergroup listed if only one map is given). More precisely, after appropriately conjugating  $G'$  satisfying  $G \subsetneq G'$ , the morphism  $X_G \rightarrow X_{G'}$  is given by the rational function  $F(t) \in \mathbb{Q}(t)$ , i.e., the curves  $X_G$  and  $X_{G'}$  have function fields  $\mathbb{Q}(t)$  and  $\mathbb{Q}(u)$ , respectively, and are related by the equation  $u = F(t)$ . By composing these rational maps down to  $X_{\text{GL}_2(\hat{\mathbb{Z}})} = \mathbb{P}_{\mathbb{Q}}^1$  (i.e., to the group labeled 1A<sup>0</sup>-1a), we obtain from  $G$  a rational function

$$J(t) \in \mathbb{Q}(t);$$

in §5, we showed that  $J(t)$  describes the morphism  $\pi_G$  from  $X_G$  to the  $j$ -line. If  $J'(t) \in \mathbb{Q}(t)$  is the rational function arising from  $G'$  in the same manner, then we will have  $J(t) = J'(F(t))$ . The function  $J(t)$  is independent of any choice of supergroups.

Similar remarks and conventions hold when  $G$  has genus 1.

#	label	$i$	$N$	generators	map	sup
0	1A <sup>0</sup> -1a	1	1			
1	3A <sup>0</sup> -3a	3	3	$\begin{pmatrix} 0 & 1 \\ 2 & 0 \end{pmatrix}, \begin{pmatrix} 1 & 1 \\ 1 & 2 \end{pmatrix}, \begin{pmatrix} 1 & 0 \\ 0 & 2 \end{pmatrix}$	$t^3$	0
2	3B <sup>0</sup> -3a	4	3	$\begin{pmatrix} 0 & 1 \\ 2 & 1 \end{pmatrix}, \begin{pmatrix} 1 & 2 \\ 0 & 2 \end{pmatrix}$	$(t+3)^3(t+27)/t$	0
3	3C <sup>0</sup> -3a	6	3	$\begin{pmatrix} 0 & 1 \\ 2 & 0 \end{pmatrix}, \begin{pmatrix} 1 & 0 \\ 0 & 2 \end{pmatrix}$	$(t-9)(t+3)/t$	1
4	3D <sup>0</sup> -3a	12	3	$\begin{pmatrix} 2 & 0 \\ 0 & 2 \end{pmatrix}, \begin{pmatrix} 1 & 0 \\ 0 & 2 \end{pmatrix}$	$729/(t^3-27)$	2
					$-27(t-3)/(t^2+3t+9)$	3
5	9A <sup>0</sup> -9a	9	9	$\begin{pmatrix} 0 & 2 \\ 4 & 0 \end{pmatrix}, \begin{pmatrix} 1 & 1 \\ 4 & 5 \end{pmatrix}, \begin{pmatrix} 1 & 0 \\ 0 & 2 \end{pmatrix}$	$t^3+9t-6$	1
6	9B <sup>0</sup> -9a	12	9	$\begin{pmatrix} 1 & 1 \\ 0 & 1 \end{pmatrix}, \begin{pmatrix} 2 & 0 \\ 0 & 5 \end{pmatrix}, \begin{pmatrix} 1 & 0 \\ 0 & 2 \end{pmatrix}$	$t(t^2+9t+27)$	2
7	9C <sup>0</sup> -9a	12	9	$\begin{pmatrix} 2 & 0 \\ 0 & 5 \end{pmatrix}, \begin{pmatrix} 4 & 2 \\ 3 & 4 \end{pmatrix}, \begin{pmatrix} 1 & 0 \\ 0 & 2 \end{pmatrix}$	$t^3$	2
8	9D <sup>0</sup> -9a	18	9	$\begin{pmatrix} 2 & 0 \\ 0 & 5 \end{pmatrix}, \begin{pmatrix} 1 & 3 \\ 3 & 1 \end{pmatrix}, \begin{pmatrix} 0 & 2 \\ 4 & 0 \end{pmatrix}, \begin{pmatrix} 1 & 0 \\ 0 & 2 \end{pmatrix}$	$-27/t^3$	3
					$(t^2-3)/t$	5
9	9E <sup>0</sup> -9a	18	9	$\begin{pmatrix} 1 & 3 \\ 0 & 1 \end{pmatrix}, \begin{pmatrix} 2 & 1 \\ 1 & 1 \end{pmatrix}, \begin{pmatrix} 4 & 2 \\ 0 & 5 \end{pmatrix}$	$-9(t^3+3t^2-9t-3)/(8t^3)$	3
10	9F <sup>0</sup> -9a	27	9	$\begin{pmatrix} 0 & 2 \\ 4 & 1 \end{pmatrix}, \begin{pmatrix} 4 & 3 \\ 5 & 4 \end{pmatrix}, \begin{pmatrix} 4 & 5 \\ 0 & 5 \end{pmatrix}$	$\frac{3^7(t^2-1)^3(t^6+3t^5+6t^4+t^3-3t^2+12t+16)^3(2t^3+3t^2-3t-5)}{(t^3-3t-1)^9}$	0
11	9G <sup>0</sup> -9a	27	9	$\begin{pmatrix} 0 & 4 \\ 2 & 3 \end{pmatrix}, \begin{pmatrix} 5 & 1 \\ 1 & 4 \end{pmatrix}, \begin{pmatrix} 5 & 3 \\ 0 & 4 \end{pmatrix}$	$\frac{(t^3-9t-12)(9-3t^3)(5t^3+18t^2+18t+3)}{(t^3+3t^2-3)^3}$	1
12	9H <sup>0</sup> -9a	36	9	$\begin{pmatrix} 1 & 3 \\ 0 & 1 \end{pmatrix}, \begin{pmatrix} 5 & 0 \\ 3 & 2 \end{pmatrix}, \begin{pmatrix} 1 & 0 \\ 2 & 2 \end{pmatrix}$	$3(t^3+9)/t^3$	4
					$3t/(2t^2-3t+6)$	9
13	9H <sup>0</sup> -9b	36	9	$\begin{pmatrix} 1 & 3 \\ 0 & 1 \end{pmatrix}, \begin{pmatrix} 5 & 0 \\ 3 & 2 \end{pmatrix}, \begin{pmatrix} 2 & 1 \\ 0 & 1 \end{pmatrix}$	$3(t^3+9t^2-9t-9)/(t^3-9t^2-9t+9)$	4
14	9H <sup>0</sup> -9c	36	9	$\begin{pmatrix} 1 & 3 \\ 0 & 1 \end{pmatrix}, \begin{pmatrix} 5 & 0 \\ 3 & 2 \end{pmatrix}, \begin{pmatrix} 4 & 2 \\ 0 & 5 \end{pmatrix}$	$-6(t^3-9t)/(t^3+9t^2-9t-9)$	4
					$-(t^2+3)/(t^2+8t+3)$	9
15	9I <sup>0</sup> -9a	36	9	$\begin{pmatrix} 2 & 1 \\ 0 & 5 \end{pmatrix}, \begin{pmatrix} 1 & 2 \\ 3 & 2 \end{pmatrix}$	$-6(t^3-9t)/(t^3-3t^2-9t+3)$	6
16	9I <sup>0</sup> -9b	36	9	$\begin{pmatrix} 2 & 1 \\ 0 & 5 \end{pmatrix}, \begin{pmatrix} 4 & 0 \\ 3 & 5 \end{pmatrix}$	$-3(t^3+9t^2-9t-9)/(t^3+3t^2-9t-3)$	6
17	9I <sup>0</sup> -9c	36	9	$\begin{pmatrix} 2 & 2 \\ 0 & 5 \end{pmatrix}, \begin{pmatrix} 2 & 2 \\ 3 & 1 \end{pmatrix}$	$(t^3-6t^2+3t+1)/(t^2-t)$	6
18	9J <sup>0</sup> -9a	36	9	$\begin{pmatrix} 1 & 3 \\ 0 & 1 \end{pmatrix}, \begin{pmatrix} 2 & 2 \\ 3 & 8 \end{pmatrix}, \begin{pmatrix} 1 & 2 \\ 0 & 2 \end{pmatrix}$	$(t^3-3t+1)/(t^2-t)$	7
19	9J <sup>0</sup> -9b	36	9	$\begin{pmatrix} 1 & 3 \\ 0 & 1 \end{pmatrix}, \begin{pmatrix} 2 & 2 \\ 3 & 8 \end{pmatrix}, \begin{pmatrix} 2 & 1 \\ 0 & 1 \end{pmatrix}$	$-18(t^2-1)/(t^3-3t^2-9t+3)$	7
20	9J <sup>0</sup> -9c	36	9	$\begin{pmatrix} 1 & 3 \\ 0 & 1 \end{pmatrix}, \begin{pmatrix} 5 & 2 \\ 3 & 5 \end{pmatrix}, \begin{pmatrix} 4 & 0 \\ 0 & 5 \end{pmatrix}$	$3(t^3+3t^2-9t-3)/(t^3-3t^2-9t+3)$	7
21	27A <sup>0</sup> -27a	36	27	$\begin{pmatrix} 1 & 1 \\ 0 & 1 \end{pmatrix}, \begin{pmatrix} 2 & 1 \\ 9 & 5 \end{pmatrix}, \begin{pmatrix} 1 & 2 \\ 3 & 2 \end{pmatrix}$	$t^3$	6

TABLE 1. Genus zero groups of 3-power level

label	$i$	$N$	generators	map	supergroup
1A <sup>0</sup> -1a	1	1			
5A <sup>0</sup> -5a	5	5	$(\begin{smallmatrix} 2 & 1 \\ 0 & 3 \end{smallmatrix}), (\begin{smallmatrix} 1 & 2 \\ 2 & 0 \end{smallmatrix}), (\begin{smallmatrix} 1 & 1 \\ 0 & 2 \end{smallmatrix})$	$t^3(t^2 + 5t + 40)$	1A <sup>0</sup> -1a
5B <sup>0</sup> -5a	6	5	$(\begin{smallmatrix} 2 & 0 \\ 0 & 3 \end{smallmatrix}), (\begin{smallmatrix} 1 & 0 \\ 1 & 1 \end{smallmatrix}), (\begin{smallmatrix} 1 & 0 \\ 0 & 2 \end{smallmatrix})$	$(t^2 + 10t + 5)^3 / t$	1A <sup>0</sup> -1a
5C <sup>0</sup> -5a	10	5	$(\begin{smallmatrix} 3 & 1 \\ 0 & 2 \end{smallmatrix}), (\begin{smallmatrix} 1 & 2 \\ 2 & 0 \end{smallmatrix}), (\begin{smallmatrix} 2 & 2 \\ 2 & 1 \end{smallmatrix})$	$8000t^3(t+1)(t^2-5t+10)^3 / (t^2-5)^5$	1A <sup>0</sup> -1a
5D <sup>0</sup> -5a	12	5	$(\begin{smallmatrix} 4 & 0 \\ 1 & 4 \end{smallmatrix}), (\begin{smallmatrix} 1 & 0 \\ 0 & 2 \end{smallmatrix})$	$125t / (t^2 - 11t - 1)$	5B <sup>0</sup> -5a
5D <sup>0</sup> -5b	12	5	$(\begin{smallmatrix} 4 & 0 \\ 1 & 4 \end{smallmatrix}), (\begin{smallmatrix} 2 & 0 \\ 0 & 1 \end{smallmatrix})$	$(t^2 - 11t - 1) / t$	5B <sup>0</sup> -5a
5E <sup>0</sup> -5a	15	5	$(\begin{smallmatrix} 2 & 1 \\ 0 & 3 \end{smallmatrix}), (\begin{smallmatrix} 2 & 0 \\ 2 & 3 \end{smallmatrix}), (\begin{smallmatrix} 1 & 0 \\ 2 & 2 \end{smallmatrix})$	$(t+5)(t^2-5) / (t^2+5t+5)$	5A <sup>0</sup> -5a
5G <sup>0</sup> -5a	30	5	$(\begin{smallmatrix} 3 & 1 \\ 0 & 2 \end{smallmatrix}), (\begin{smallmatrix} 2 & 1 \\ 0 & 1 \end{smallmatrix})$	$125 / (t(t^4 + 5t^3 + 15t^2 + 25t + 25))$ $(t^2 + 5) / t$	5E <sup>0</sup> -5a 5B <sup>0</sup> -5a
5G <sup>0</sup> -5b	30	5	$(\begin{smallmatrix} 3 & 1 \\ 0 & 2 \end{smallmatrix}), (\begin{smallmatrix} 2 & 1 \\ 3 & 3 \end{smallmatrix})$	$-t(t^2 + 5t + 10) / (t^3 + 5t^2 + 10t + 10)$ $-5(t^2 + 4t + 5) / (t^2 + 5t + 5)$	5C <sup>0</sup> -5a 5E <sup>0</sup> -5a
5H <sup>0</sup> -5a	60	5	$(\begin{smallmatrix} 4 & 0 \\ 0 & 4 \end{smallmatrix}), (\begin{smallmatrix} 2 & 0 \\ 0 & 1 \end{smallmatrix})$	$-1 / t^5$ $\frac{-(t^4 - 2t^3 + 4t^2 - 3t + 1)}{t(t^4 + 3t^3 + 4t^2 + 2t + 1)}$ $5t / (t^2 - t - 1)$	5D <sup>0</sup> -5a 5D <sup>0</sup> -5b 5G <sup>0</sup> -5a
25A <sup>0</sup> -25a	30	25	$(\begin{smallmatrix} 2 & 2 \\ 0 & 13 \end{smallmatrix}), (\begin{smallmatrix} 4 & 1 \\ 3 & 1 \end{smallmatrix}), (\begin{smallmatrix} 2 & 3 \\ 0 & 6 \end{smallmatrix})$	$(t-1)(t^4 + t^3 + 6t^2 + 6t + 11)$	5B <sup>0</sup> -5a
25B <sup>0</sup> -25a	60	25	$(\begin{smallmatrix} 9 & 10 \\ 0 & 14 \end{smallmatrix}), (\begin{smallmatrix} 0 & 7 \\ 7 & 2 \end{smallmatrix}), (\begin{smallmatrix} 2 & 8 \\ 0 & 1 \end{smallmatrix})$	$-t^5$	5D <sup>0</sup> -5b
25B <sup>0</sup> -25b	60	25	$(\begin{smallmatrix} 9 & 10 \\ 0 & 14 \end{smallmatrix}), (\begin{smallmatrix} 0 & 7 \\ 7 & 2 \end{smallmatrix}), (\begin{smallmatrix} 4 & 1 \\ 0 & 7 \end{smallmatrix})$	$(1 - t^2) / t$ $\frac{-(t^4 - 2t^3 + 4t^2 - 3t + 1)}{t(t^4 + 3t^3 + 4t^2 + 2t + 1)}$ $(t^2 + 4t - 1) / (t^2 - t - 1)$	25A <sup>0</sup> -25a 5D <sup>0</sup> -5a 25A <sup>0</sup> -25a
7B <sup>0</sup> -7a	8	7	$(\begin{smallmatrix} 2 & 0 \\ 0 & 4 \end{smallmatrix}), (\begin{smallmatrix} 3 & 0 \\ 1 & 5 \end{smallmatrix}), (\begin{smallmatrix} 1 & 0 \\ 0 & 3 \end{smallmatrix})$	$(t^2 + 5t + 1)^3(t^2 + 13t + 49) / t$	1A <sup>0</sup> -1a
7D <sup>0</sup> -7a	21	7	$(\begin{smallmatrix} 2 & 3 \\ 0 & 3 \end{smallmatrix}), (\begin{smallmatrix} 2 & 4 \\ 4 & 5 \end{smallmatrix}), (\begin{smallmatrix} 3 & 1 \\ 0 & 4 \end{smallmatrix})$	$\frac{(2t-1)^3(t^2-t+2)^3(t^2+5t+4)^3(5t^2+2t-4)^3}{(t^3+2t^2-t-1)^7}$	1A <sup>0</sup> -1a
7E <sup>0</sup> -7a	24	7	$(\begin{smallmatrix} 6 & 0 \\ 1 & 6 \end{smallmatrix}), (\begin{smallmatrix} 1 & 0 \\ 0 & 3 \end{smallmatrix})$	$49(t^2 - t) / (t^3 - 8t^2 + 5t + 1)$	7B <sup>0</sup> -7a
7E <sup>0</sup> -7b	24	7	$(\begin{smallmatrix} 6 & 0 \\ 1 & 6 \end{smallmatrix}), (\begin{smallmatrix} 3 & 0 \\ 0 & 1 \end{smallmatrix})$	$(t^3 - 8t^2 + 5t + 1) / (t^2 - t)$	7B <sup>0</sup> -7a
7E <sup>0</sup> -7c	24	7	$(\begin{smallmatrix} 6 & 0 \\ 1 & 6 \end{smallmatrix}), (\begin{smallmatrix} 3 & 0 \\ 0 & 4 \end{smallmatrix})$	$-7(t^3 - 2t^2 - t + 1) / (t^3 - t^2 - 2t + 1)$	7B <sup>0</sup> -7a
7F <sup>0</sup> -7a	28	7	$(\begin{smallmatrix} 3 & 1 \\ 4 & 4 \end{smallmatrix}), (\begin{smallmatrix} 4 & 4 \\ 1 & 3 \end{smallmatrix}), (\begin{smallmatrix} 3 & 4 \\ 0 & 4 \end{smallmatrix})$	$\frac{t(t+1)^3(t^2-5t+1)^3(t^2-5t+8)^3(t^4-5t^3+8t^2-7t+7)^3}{(t^3-4t^2+3t+1)^7}$	1A <sup>0</sup> -1a
13A <sup>0</sup> -13a	14	13	$(\begin{smallmatrix} 2 & 0 \\ 0 & 7 \end{smallmatrix}), (\begin{smallmatrix} 1 & 0 \\ 1 & 1 \end{smallmatrix}), (\begin{smallmatrix} 1 & 0 \\ 0 & 2 \end{smallmatrix})$	$(t^2 + 5t + 13)(t^4 + 7t^3 + 20t^2 + 19t + 1)^3 / t$	1A <sup>0</sup> -1a
13B <sup>0</sup> -13a	28	13	$(\begin{smallmatrix} 3 & 0 \\ 0 & 9 \end{smallmatrix}), (\begin{smallmatrix} 4 & 0 \\ 1 & 10 \end{smallmatrix}), (\begin{smallmatrix} 1 & 0 \\ 0 & 2 \end{smallmatrix})$	$13t / (t^2 - 3t - 1)$	13A <sup>0</sup> -13a
13B <sup>0</sup> -13b	28	13	$(\begin{smallmatrix} 3 & 0 \\ 0 & 9 \end{smallmatrix}), (\begin{smallmatrix} 4 & 0 \\ 1 & 10 \end{smallmatrix}), (\begin{smallmatrix} 2 & 0 \\ 0 & 1 \end{smallmatrix})$	$(t^2 - 3t - 1) / t$	13A <sup>0</sup> -13a
13C <sup>0</sup> -13a	42	13	$(\begin{smallmatrix} 5 & 0 \\ 0 & 8 \end{smallmatrix}), (\begin{smallmatrix} 1 & 0 \\ 1 & 1 \end{smallmatrix}), (\begin{smallmatrix} 1 & 0 \\ 0 & 2 \end{smallmatrix})$	$13(t^2 - t) / (t^3 - 4t^2 + t + 1)$	13A <sup>0</sup> -13a
13C <sup>0</sup> -13b	42	13	$(\begin{smallmatrix} 5 & 0 \\ 0 & 8 \end{smallmatrix}), (\begin{smallmatrix} 1 & 0 \\ 1 & 1 \end{smallmatrix}), (\begin{smallmatrix} 2 & 0 \\ 0 & 1 \end{smallmatrix})$	$(t^3 - 4t^2 + t + 1) / (t^2 - t)$	13A <sup>0</sup> -13a
13C <sup>0</sup> -13c	42	13	$(\begin{smallmatrix} 5 & 0 \\ 0 & 8 \end{smallmatrix}), (\begin{smallmatrix} 1 & 0 \\ 1 & 1 \end{smallmatrix}), (\begin{smallmatrix} 2 & 0 \\ 0 & 3 \end{smallmatrix})$	$-(5t^3 - 7t^2 - 8t + 5) / (t^3 - 4t^2 + t + 1)$	13A <sup>0</sup> -13a

TABLE 2. Genus zero groups of  $\ell$ -power level for primes  $\ell > 3$











label	Weierstrass model
<b>121b1</b>	$y^2 + y = x^3 - x^2 - 7x + 10$
<b>128a1</b>	$y^2 = x^3 + x^2 + x + 1$
<b>128a2</b>	$y^2 = x^3 + x^2 - 9x + 7$
<b>256a1</b>	$y^2 = x^3 + x^2 - 3x + 1$
<b>256a2</b>	$y^2 = x^3 + x^2 - 13x - 21$
<b>256b1</b>	$y^2 = x^3 - 2x$
<b>256b2</b>	$y^2 = x^3 + 8x$

TABLE 5. Some elliptic curves

## REFERENCES

- [BCP97] Wieb Bosma, John Cannon, and Catherine Playoust, *The Magma algebra system. I. The user language*, J. Symbolic Comput. **24** (1997), no. 3-4, 235–265. Computational algebra and number theory (London, 1993). [↑3](#)
- [CLY04] Kok Seng Chua, Mong Lung Lang, and Yifan Yang, *On Rademacher’s conjecture: congruence subgroups of genus zero of the modular group*, J. Algebra **277** (2004), no. 1, 408–428. [↑4.6](#)
- [Cre14] John E. Cremona, *Elliptic curve data*, 2014. <http://homepages.warwick.ac.uk/staff/J.E.Cremona/ftp/data/>. [↑6, 8](#)
- [CP03] C. J. Cummins and S. Pauli, *Congruence subgroups of  $\mathrm{PSL}(2, \mathbb{Z})$  of genus less than or equal to 24*, Experiment. Math. **12** (2003), no. 2, 243–255. MR2016709 (2004i:11037) [↑1.2, 3, 4.3.1, 4.3.3, A](#)
- [CP] ———, *Database of congruence subgroups of  $\mathrm{PSL}(2, \mathbb{Z})$  of genus less than or equal to 24*. available at <http://www.uncg.edu/mat/faculty/pauli/congruence/>. [↑3](#)
- [DR73] P. Deligne and M. Rapoport, *Les schémas de modules de courbes elliptiques*, Modular functions of one variable, II (Proc. Internat. Summer School, Univ. Antwerp, Antwerp, 1972), 1973, pp. 143–316. Lecture Notes in Math., Vol. 349. MR0337993 (49 #2762) [↑1, 6](#)
- [Den74] Jr. Dennis Joseph B., *The genus of subfields of  $K(p^n)$* , Illinois J. Math. **18** (1974), 246–264. MR0342466 (49 #7212) [↑3](#)
- [Fal83] G. Faltings, *Endlichkeitssätze für abelsche Varietäten über Zahlkörpern*, Invent. Math. **73** (1983), no. 3, 349–366, DOI 10.1007/BF01388432 (German). MR718935 (85g:11026a) [↑1](#)
- [Hal98] Emmanuel Halberstadt, *Sur la courbe modulaire  $X_{nd\acute{e}p}(11)$* , Experiment. Math. **7** (1998), no. 2, 163–174 (French, with English and French summaries). MR1677158 (99m:11062) [↑6](#)
- [KL81] Daniel S. Kubert and Serge Lang, *Modular units*, Grundlehren der Mathematischen Wissenschaften [Fundamental Principles of Mathematical Science], vol. 244, Springer-Verlag, New York-Berlin, 1981. MR648603 (84h:12009) [↑2.4, 4.1, 4.1, 4.2, 4.5, 4.3.2, 4.3.3](#)
- [LMF15] LMFDB Collaboration, *The L-functions and Modular Forms Database*, 2015. <http://www.lmfdb.org>. [↑8](#)
- [RZB15] Jeremy Rouse and David Zureick-Brown, *Elliptic curves over  $\mathbb{Q}$  and 2-adic images of Galois*, Research in Number Theory **1** (2015). [↑1.3, 3.2, 6, 7](#)
- [Ser72] Jean-Pierre Serre, *Propriétés galoisiennes des points d’ordre fini des courbes elliptiques*, Invent. Math. **15** (1972), no. 4, 259–331. MR0387283 (52 #8126) [↑1](#)
- [Ser81] ———, *Quelques applications du théorème de densité de Chebotarev*, Inst. Hautes Études Sci. Publ. Math. **54** (1981), 323–401. MR644559 (83k:12011) [↑1](#)
- [Ser89] ———, *Abelian  $l$ -adic representations and elliptic curves*, Second Ed., Advanced Book Classics, Addison-Wesley Publishing Company Advanced Book Program, Redwood City, CA, 1989. With the collaboration of Willem Kuyk and John Labute. MR1043865 (91b:11071) [↑7](#)
- [Ser97] ———, *Lectures on the Mordell-Weil theorem*, Third Ed., Aspects of Mathematics, Friedr. Vieweg & Sohn, Braunschweig, 1997. Translated from the French and edited by Martin Brown from notes by Michel Waldschmidt, With a foreword by Brown and Serre. MR1757192 (2000m:11049) [↑3, 7](#)
- [Shi94] Goro Shimura, *Introduction to the arithmetic theory of automorphic functions*, Publications of the Mathematical Society of Japan, vol. 11, Princeton University Press, Princeton, NJ, 1994. Reprint of the 1971 original; Kanô Memorial Lectures, 1. MR1291394 (95e:11048) [↑2.2](#)
- [Sut16] Andrew V. Sutherland, *Computing images of Galois representations attached to elliptic curves*, Forum of Mathematics, Sigma **4** (2016), 79 pages. [↑1.1, 6, 8, 8.1](#)

- [SZ16] Andrew V. Sutherland and David Zywina, *Magma scripts associated to “Modular curves of prime-power level with infinitely many rational points”*, 2016. available at <http://math.mit.edu/~drew/SZ16>. ↑1.1, 3, 5.4, 6
- [Zyw15a] David Zywina, *On the possible images of the mod  $\ell$  representations associated to elliptic curves over  $\mathbb{Q}$* , 2015. arXiv:1508.07660v1 [math.NT]. ↑1.1, 2.5
- [Zyw15b] ———, *Possible indices for the Galois image of elliptic curves over  $\mathbb{Q}$* , 2015. arXiv:1508.07663v1 [math.NT]. ↑ii, 2.7, 3, 6, 7

DEPARTMENT OF MATHEMATICS, MASSACHUSETTS INSTITUTE OF TECHNOLOGY, CAMBRIDGE, MA 02139, USA

*Email address:* [drew@math.mit.edu](mailto:drew@math.mit.edu)

*URL:* <http://math.mit.edu/~drew>

DEPARTMENT OF MATHEMATICS, CORNELL UNIVERSITY, ITHACA, NY 14853, USA

*Email address:* [zywina@math.cornell.edu](mailto:zywina@math.cornell.edu)

*URL:* <http://www.math.cornell.edu/~zywina>