

There is no direct generalization of positive partial transpose criterion to the three-by-three case

Łukasz Skowronek^{1, a)}

Brygadziów 23A/3, 40-807 Katowice, Poland^{b)}

(Dated: 28 January 2022)

We show that there cannot exist a straightforward generalization of the famous positive partial transpose criterion to three-by-three systems. We call straightforward generalizations that use a finite set of positive maps and arbitrary local rotations of the tested two-partite state. In particular, we show that a family of extreme positive maps discussed in a paper by Ha and Kye¹, cannot be replaced by a finite set of witnesses in the task of entanglement detection in three-by-three systems. In a more mathematically elegant parlance, our result says that the convex cone of positive maps of the set of three-dimensional matrices into itself is not finitely generated as a mapping cone

I. INTRODUCTION

In multi-partite quantum systems, separable states are a generalization of pure product states to density matrices. They are defined as convex combinations of one-dimensional projections onto product states,

$$\rho = \sum_{i=1}^k \alpha_i |\phi_i^1\rangle \langle \phi_i^1| \otimes \dots \otimes |\phi_i^n\rangle \langle \phi_i^n|, \quad (1)$$

where $\sum_{i=1}^k \alpha_i = 1$ and $\forall_i \alpha_i > 0$.

Separable states exhibit fully classical correlations between the individual subsystems. Only states that are not separable may therefore be useful for tasks like quantum cryptography^{2,3}. Such non-separable states are called *entangled*. In the two-partite case, all entangled states have been shown to be capable of violating a Bell type inequality in a setting where separable states are useless⁴⁻⁶. On the other hand, correlations obtained for an entangled state may exhibit a hidden variable description⁷. Unlike for pure product states, it turns out to be very difficult to test whether a given density matrix is separable or entangled. A notable exception to this rule are two-partite density matrices for 2×2 and 2×3 systems, which turn out to be separable if and only if

$$(\text{id} \otimes \text{t}) \rho \geq 0, \quad (2)$$

where id and t stand for the identity and the transpose maps, respectively, and ρ denotes the density matrix in question. This is the famous positive partial transpose (PPT) criterion by Peres and Horodecki^{8,9}, which builds on the decomposability of positive maps of 2×2 matrices into 2×2 or 3×3 matrices^{10,11}.

It is well known that the PPT test is not a sufficient separability criterion for 3×3 systems. In the 3×3

case, there exist entangled states with positive partial transpose¹². They are examples of so-called *bound entangled states*.

In the present paper, we shall prove that no generalization of (2) of the form

$$(\text{id} \otimes \Phi_i) (A \otimes B) \rho (A \otimes B)^* \geq 0 \quad \forall_{i=1,2,\dots,m; A, B \in M_3} \quad (3)$$

can be a sufficient criterion for separability in 3×3 systems. Here, $\{\Phi_i\}_{i=1}^k$ stands for a finite set of positive maps and A, B denote arbitrary 3×3 matrices with complex entries. We denote the set of all such matrices by M_3 . We immediately see that the similarity transform by A is superfluous in (3) and we can therefore replace (3) with

$$(\text{id} \otimes \Phi_i) (\mathbb{I} \otimes B) \rho (\mathbb{I} \otimes B)^* \geq 0 \quad \forall_{i=1,2,\dots,m; B \in M_3} \quad (4)$$

We claim that no such criterion can be sufficient for separability in 3×3 systems. This result has been obtained by the author in spring 2011 and presented at a seminar at ICFO on 2011/05/09. Due to difficult personal conditions and taking up a job outside academia, the result has not been available in a polished written up form until now.

II. PROOF

Let us assume that Φ is an element of the set $\mathcal{M}_{3,3}$ of linear maps of the set of three-dimensional matrices M_3 into itself. It will be convenient for us to use the Jamiołkowski-Choi transform^{13,14} of Φ ,

$$C_\Phi = \sum_{i,j=1,2,3} e_{ij} \otimes \Phi(e_{ij}), \quad (5)$$

where e_{ij} are elements of a canonical basis of M_3 . The object C_Φ is called the *Choi matrix* of Φ after¹⁴, and the mapping $J : \mathcal{M}_{3,3} \ni \Phi \mapsto C_\Phi \in M_9$ is an isomorphism between the space of maps of the set of three-dimensional matrices into itself and the set of nine-dimensional matrices. This isomorphism is named the *Jamiołkowski isomorphism* after¹³. Interestingly, if we define a natural

^{a)}Electronic mail: lukasz.m.skowronek@gmail.com

^{b)}Former affiliation: Instytut Fizyki im. Smoluchowskiego, Uniwersytet Jagielloński, Reymonta 4, 30-059 Kraków, Poland

inner product in $\mathcal{M}_{3,3}$,

$$\langle \Phi, \Psi \rangle'' := \sum_{i,j=1,2,3} \text{Tr} (\Phi(e_{ij}) \Psi(e_{ij})^*), \quad (6)$$

J also turns out to be an isometry,

$$\langle \Phi, \Psi \rangle'' = \text{Tr} (C_\Phi C_\Psi^*) =: \langle C_\Phi, C_\Psi \rangle', \quad (7)$$

where the conjugation is taken with the assumption of orthonormality of the canonical basis of \mathbb{C}^3 underlying e_{ij} . The inner product $\langle \cdot, \cdot \rangle'$ is simply a Hilbert-Schmidt product in M_9 . Similarly, $\langle \cdot, \cdot \rangle''$ is a Hilbert-Schmidt product in $\mathcal{M}_{3,3}$.

We shall need the following properties of Jamiołkowski isomorphism and the Hilbert-Schmidt product in $\mathcal{M}_{3,3}$ (cf. e.g.¹⁵).

Proposition II.1. *Let $A = \sum_{i,j=1,2,3} A_{ij} e_{ij}$ be an element of M_3 and let $\text{Ad}_A : M_3 \ni X \mapsto AXA^* \in M_3$ be the conjugation map by A . Then we have*

$$J(\text{Ad}_A) = C_{\text{Ad}_A} = |\alpha\rangle \langle \alpha|, \quad (8)$$

where $\alpha := \sum_{i,j} A_{ij} e_i \otimes e_j$ for $\{e_i\}_{i=1}^3$ a canonical basis of \mathbb{C}^3 corresponding to the canonical basis $\{e_{ij}\}_{i,j=1,2,3}$ of M_3 .

In other words, arbitrary conjugation maps in M_3 map to arbitrary non-normalized one-dimensional projections in \mathbb{C}^3 .

Proposition II.2. *For all $\Phi, \Sigma, \Psi \in \mathcal{M}_{3,3}$, we have*

$$\langle \Phi \circ \Sigma, \Psi \rangle'' = \langle \Sigma, \Phi^* \circ \Psi \rangle'' \quad (9)$$

Proposition II.3. *For all $\Theta, \Phi, \Sigma, \Psi \in \mathcal{M}_{3,3}$, we have*

$$\langle \Theta \circ \Phi \circ \Sigma, \Psi \rangle'' = \langle \Phi, \Theta^* \circ \Psi \circ \Sigma^* \rangle'' \quad (10)$$

We also have the trivial

Proposition II.4. *For arbitrary $\Phi, \Psi \in \mathcal{M}_{3,3}$*

$$C_{\Phi \circ \Psi} = (\mathbb{I} \otimes \Phi) C_\Psi \quad (11)$$

and the slightly more involved

Proposition II.5. *For arbitrary $A, B \in M_3$ and $\Phi \in \mathcal{M}_{3,3}$, we have*

$$C_{\text{Ad}_A \circ \Phi \circ \text{Ad}_B} = \text{Ad}_{B^t \otimes A} C_\Phi \quad (12)$$

which we prove in Appendix A.

Let us note that the convex cone $\mathcal{P}_{3,3}$ of positive (i.e. positivity preserving) maps from M_3 into M_3 is a subset of the set $\mathcal{H}_{3,3}$ of Hermiticity preserving maps (cf. e.g.¹⁶). It is easy to see that $\mathcal{H}_{3,3}$ is a linear space over \mathbb{R} and $\langle \cdot, \cdot \rangle''$ induces a symmetric inner product in $\mathcal{H}_{3,3}$. Thus for every convex cone $\mathcal{K} \subset \mathcal{H}_{3,3}$, we can define the *dual cone* of \mathcal{K} ,

$$\mathcal{K}^\circ := \{ \Psi \in \mathcal{H}_{3,3} \mid \langle \Psi, \Phi \rangle'' \geq 0 \forall \Phi \in \mathcal{K} \} \quad (13)$$

Among the convex cones contained in $\mathcal{H}_{3,3}$, and more generally in $\mathcal{H}_{m,n}$ with m, n arbitrary, there is a distinguished class of so-called *mapping cones*, introduced by Størmer¹⁷.

Definition II.6. *Let $\mathcal{K} \subset \mathcal{P}_{3,3}$ be a cone. We call \mathcal{K} a mapping cone if \mathcal{K} is closed, not consisting only of the zero map, and*

$$\text{Ad}_A \circ \Phi \circ \text{Ad}_B \in \mathcal{K} \forall \Phi \in \mathcal{K} \forall A, B \in M_3 \quad (14)$$

In other words, \mathcal{K} is closed with respect to left and right multiplications by arbitrary conjugation maps. Note that convexity is not explicitly assumed in Definition II.6, however we shall always assume \mathcal{K} to be convex in what follows. The papers^{15,18–23} show several interesting properties of mapping cones in the finite-dimensional case, specifically concerning their duals and their relation to operator algebras. The cone of positive maps $\mathcal{P}_{m,n}$ is obviously the biggest mapping cone in $\mathcal{H}_{m,n}$, whereas its dual $\mathcal{SP}_{m,n}$, called the set of *superpositive maps*²⁴, is in a sense the smallest one, because it is contained in all convex mapping cones. As long as convex cones are concerned, which is the case in this paper and in¹⁵, we can rightly call $\mathcal{SP}_{m,n}$ the smallest mapping cone. It turns out that (cf. e.g.²¹) $\mathcal{SP}_{m,n}$ can be characterized as the set of convex combinations of the form

$$\sum_{i=1}^n \text{Ad}_{A_i}, \text{rk}(A_i) = 1 \quad (15)$$

where $\text{rk}(A)$ denotes the rank of a matrix A . From Proposition II.1, we easily see that, up to scaling, superpositive maps are in a one-to-one correspondence with separable states via Jamiołkowski isomorphism. It is also not difficult to deduce (for more details cf.^{13,21}) that the set of all positive maps Φ corresponds via Jamiołkowski isomorphism to the set of *block positive* Choi matrices, i.e. Choi matrices C_Φ such that

$$\langle x \otimes y \mid C_\Phi \mid x \otimes y \rangle > 0 \forall x, y \in \mathbb{C}^3. \quad (16)$$

Let us denote the set of separable states in 3×3 systems by $\text{Sep}_{3 \times 3}$. All the above facts give us enough information to prove the following equivalence.

Proposition II.7. *The nonexistence of a separability criterion of the form*

$$\rho \in \text{Sep}_{3 \times 3} \Leftrightarrow \forall_{i=1,2,\dots,k} \forall_{B \in M_3} (\mathbb{I} \otimes \Phi_i) \text{Ad}_{\mathbb{I} \otimes B} \rho \geq 0 \quad (17)$$

is equivalent to the fact that there does not exist a finite set of positive maps $\{\Xi_i\}_{i=1}^k \in \mathcal{P}_{3,3}$ such that

$$\mathcal{P}_{3,3} = \overline{\text{convhull}(\{\text{Ad}_X \circ \Xi_i \circ \text{Ad}_Y \mid i; X, Y \in M_3\})}, \quad (18)$$

where convhull denotes the convex hull of a set and the overline refers to topological closure.

Proof. Let us denote by Ψ the inverse Jamiołkowski transform of a density matrix $\rho \in M_9$. By Propositions II.1–II.5 and the fact that Jamiołkowski isomorphism is

an isometry, we have

$$\begin{aligned}
(\mathbb{I} \otimes \Phi_i) \text{Ad}_{\mathbb{I} \otimes B} \rho &\geq 0 \Leftrightarrow C_{\Phi_i \circ \text{Ad}_B \circ \Psi} \geq 0 \Leftrightarrow \\
&\Leftrightarrow \langle |\alpha\rangle \langle \alpha|, C_{\Phi_i \circ \text{Ad}_B \circ \Psi} \rangle' \geq 0 \forall \alpha \in \mathbb{C}^9 \Leftrightarrow \\
&\Leftrightarrow \langle C_{\text{Ad}_A}, C_{\Phi_i \circ \text{Ad}_B \circ \Psi} \rangle' \geq 0 \forall A \in M_3 \Leftrightarrow \\
&\Leftrightarrow \langle \text{Ad}_A, \Phi_i \circ \text{Ad}_B \circ \Psi \rangle'' \geq 0 \forall A \in M_3 \Leftrightarrow \\
&\Leftrightarrow \langle \text{Ad}_{B^*} \circ \Phi_i^* \circ \text{Ad}_A, \Psi \rangle'' \geq 0 \forall A \in M_3 \quad (19)
\end{aligned}$$

Let us assume that (17) is a sufficient criterion for separability of a state ρ . If we additionally assume that (18) does not hold for $\Xi_i = \Phi_i^*$, $i = 1, 2, \dots, k$, then by the identity (cf.²⁵ and a comment in¹⁵)

$$\mathcal{K}^{\circ\circ} = \mathcal{K} \quad (20)$$

applied to the closure of the convex hull of maps of the form $\text{Ad}_{B^*} \circ \Phi_i^* \circ \text{Ad}_A$, $A, B \in M_3$, we see that there must exist an element Ψ of $\mathcal{P}_{3,3} \setminus \mathcal{SP}_{3,3}$ such that

$$\langle \text{Ad}_{B^*} \circ \Phi_i^* \circ \text{Ad}_A, \Psi \rangle'' \geq 0 \forall i, A, B \in M_3 \quad (21)$$

We already know by (19) that (21) is equivalent to

$$(\mathbb{I} \otimes \Phi_i) \text{Ad}_{\mathbb{I} \otimes B} C_\Psi \geq 0 \forall i, B \in M_3 \quad (22)$$

this implies, up to scaling, $C_\Psi \in \text{Sep}_{3,3}$, which is a contradiction since $J^{-1}(\text{Sep}_{3,3}) \subset \mathcal{SP}_{3,3}$, where the last inclusion turns into an equality if we allow arbitrary scaling of the separable density matrices by non-negative factors. We are led to a conclusion that (17) cannot be a sufficient criterion for separability. If it is, then (18) must hold. Conversely, if the cone of positive maps can be written in the form (18), then by the property (20) applied to $\mathcal{K} = \mathcal{P}_{3,3}$ ($\mathcal{K}^\circ = \mathcal{SP}_{3,3}$), a sufficient criterion for a map Ψ to be superpositive, or in other words, for the Choi matrix C_Ψ to be a non-negative multiple of a separable density matrix, is (21) with Ξ_i substituted for Φ_i^* , X substituted for B^* and Y substituted for B . That is further equivalent to Equation (22) with Ξ_i^* in place of Φ_i and X^* substituted for B . Up to relabeling, this is the same as (17), which finishes the proof. \square

The main result of this paper is therefore equivalent to the fact that does not exist a finite set of generators $\{\Xi_i\}_{i=1}^k$ of $\mathcal{P}_{3,3}$ considered as a mapping cone. Let us assume for a moment that such set of generators exists. For $C \in M_9$, let us also denote by $\langle \cdot \otimes y | C | \cdot \otimes y \rangle$ and $\langle x \otimes \cdot | C | x \otimes \cdot \rangle$ matrices in M_3 with matrix elements $\sum_{k,l=1}^3 \bar{y}^k C_{ik,jl} y^l$ and $\sum_{k,l=1}^3 \bar{x}^k C_{ki,lj} x^l$, respectively. We have the following.

Proposition II.8. *Let $\{\Xi_i\}_{i=1}^k$ be a set of positive maps in $\mathcal{P}_{3,3}$ such that equality (18) holds. Then we can assume that:*

1. All Ξ_i 's are extreme.
2. $\Xi_i = \text{Ad}_X \circ \Xi_j \circ \text{Ad}_Y$ does not hold for any $i, j \in \{1, 2, \dots, k\}$, $X, Y \in M_3$ unless $i = j$.
3. For any Φ positive, extreme and such that the matrices $\langle \cdot \otimes y | C_\Phi | \cdot \otimes y \rangle$ and $\langle x \otimes \cdot | C_\Phi | x \otimes \cdot \rangle$ are of rank at least 2 for all $x, y \in \mathbb{C}^3$, $\Phi = \text{Ad}_X \circ \Xi_i \circ \text{Ad}_Y$ for some $i \in \{1, 2, \dots, k\}$ and $X, Y \in M_3$.
4. If Φ positive and extreme is not of the form $\Phi = \text{Ad}_Z$ or $\Phi = \text{Ad}_Z \circ \mathbf{t}$ for some $Z \in M_3$, i.e. Φ is not decomposable, the matrices X and Y in $\Phi = \text{Ad}_X \circ \Xi_i \circ \text{Ad}_Y$ must be invertible.
5. For any Φ positive, extreme and such that the matrices $\langle \cdot \otimes y | C_\Phi | \cdot \otimes y \rangle$ and $\langle x \otimes \cdot | C_\Phi | x \otimes \cdot \rangle$ are of rank at least 2 for all $x, y \in \mathbb{C}^3$, there is exactly one i such that $\Phi = \text{Ad}_X \circ \Xi_i \circ \text{Ad}_Y$ for some $X, Y \in M_3$.

Proof. Points 1) and 2) are obvious (in particular, point 2) simply states the fact that we can keep removing some Ξ_i 's until none of them are redundant as generators of the mapping cone (18)). Point 3) needs a longer proof that we provide in Appendix B. Point 4) is a consequence of the fact that positive maps in 2×2 and 2×3 systems are *decomposable*, i.e. they can be written in the form $\sum_i \text{Ad}_{X_i} + \sum_j \text{Ad}_{X'_j} \circ \mathbf{t}$ for some $X_i, X'_j \in M_3$ ^{10,11}. Non-invertible A or B would bring us back to this basic case, as it is easily seen from Proposition II.5. Point 5) follows because from point 3) we already know that there must exist $X, Y \in M_3$ and $i \in \{1, 2, \dots, k\}$ such that equality $\Phi = \text{Ad}_X \circ \Xi_i \circ \text{Ad}_Y$ holds. Moreover, by point 4) the matrices X and Y must be invertible. If i were not unique, we would have $\Phi = \text{Ad}_Z \circ \Xi_j \circ \text{Ad}_T = \text{Ad}_X \circ \Xi_i \circ \text{Ad}_Y$ for some $j \neq i$ and $Z, T \in M_3$ invertible. This would imply $\Xi_j = \text{Ad}_{Z^{-1}X} \circ \Xi_i \circ \text{Ad}_{YT^{-1}}$, contradicting point 2). \square

It is easy to see from the above proposition that the existence of an infinite family $\{\Phi_\alpha\}_{\alpha \in \mathcal{A}} \subset \mathcal{P}_{3,3}$ of positive, indecomposable, extreme maps such that $\text{rk}(\langle x \otimes \cdot | C_{\Phi_\alpha} | x \otimes \cdot \rangle) \geq 2$ and $\text{rk}(\langle \cdot \otimes y | C_{\Phi_\alpha} | \cdot \otimes y \rangle) \geq 2$ for all $x, y \in \mathbb{C}^3$ and

$$\neg(\exists_{\alpha_1 \neq \alpha_2} \exists_{X, Y \in M_3} \Phi_{\alpha_1} = \text{Ad}_X \circ \Phi_{\alpha_2} \circ \text{Ad}_Y) \quad (23)$$

would preclude the existence of a finite set of mapping cone generators of $\mathcal{P}_{3,3}$. It turns out that the family of extreme, indecomposable positive maps

$$\Phi_t : M_3 \ni [x_{ij}] \mapsto \begin{bmatrix} a_t x_{11} + b_t x_{22} + c_t x_{33} & -x_{12} & -x_{13} \\ -x_{21} & c_t x_{11} + a_t x_{22} + b_t x_{33} & -x_{23} \\ -x_{31} & -x_{32} & b_t x_{11} + c_t x_{22} + a_t x_{33} \end{bmatrix} \in M_3 \quad (24)$$

with $a_t = (1-t)^2/(1-t+t^2)$, $b_t = t^2/(1-t+t^2)$, $c_t = 1/(1-t+t^2)$, $t \in [0, 1)$, introduced by Ha and Kye¹, has the required properties. To show this, let us first prove the following facts about of Φ_t

Proposition II.9. *Let Φ_t denote positive maps of the form (24). Let $x = [x^1, x^2, x^3]$ and $y = [y^1, y^2, y^3]$ denote normalized vectors in \mathbb{C}^3 . Then*

1. $\det(\langle \cdot \otimes y | C_{\Phi_t} | \cdot \otimes y \rangle)$ only depends on $|y^1|$, $|y^2|$ and $|y^3|$
2. $\langle \cdot \otimes y | C_{\Phi_t} | \cdot \otimes y \rangle$ is singular if and only if

$$|y^1| = |y^2| = |y^3| = \frac{1}{\sqrt{3}} \quad (25)$$

or

$$|y^1| = 0, |y^2| = \sqrt{\frac{t}{1+t}}, |y^3| = \sqrt{\frac{1}{1+t}} \quad (26)$$

or

$$|y^1| = \sqrt{\frac{1}{1+t}}, |y^2| = 0, |y^3| = \sqrt{\frac{t}{1+t}} \quad (27)$$

or

$$|y^1| = \sqrt{\frac{t}{1+t}}, |y^2| = \sqrt{\frac{1}{1+t}}, |y^3| = 0 \quad (28)$$

3. $\langle \cdot \otimes y | C_{\Phi_t} | \cdot \otimes y \rangle$ is never of rank 1 or 0

4. For every y that satisfies one of the condition sets (25)-(28), there is, up to multiplication by a constant factor, exactly one x such that

$$\langle x \otimes y | C_{\Phi_t} | x \otimes y \rangle = 0 \quad (29)$$

holds

5. For y_1 and y_2 , $y_1 \neq y_2$, substituted in place of y in (29), the corresponding vectors x are never multiples of each other

6. $\langle x \otimes \cdot | C_{\Phi_t} | x \otimes \cdot \rangle$ is never of rank 1 or 0

Proof. The Choi matrix of Φ_t equals

$$\begin{bmatrix} a_t & 0 & 0 & 0 & -1 & 0 & 0 & 0 & -1 \\ 0 & c_t & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & b_t & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & b_t & 0 & 0 & 0 & 0 & 0 \\ -1 & 0 & 0 & 0 & a_t & 0 & 0 & 0 & -1 \\ 0 & 0 & 0 & 0 & 0 & c_t & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & c_t & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & b_t & 0 \\ -1 & 0 & 0 & 0 & -1 & 0 & 0 & 0 & a_t \end{bmatrix} \quad (30)$$

By an elementary calculation, one can show that

$$\langle \cdot \otimes y | C_{\Phi_t} | \cdot \otimes y \rangle = \begin{bmatrix} a_t |y^1|^2 + b_t |y^2|^2 + c_t |y^3|^2 & -\bar{y}^1 y^2 & -\bar{y}^1 y^3 \\ -y^1 \bar{y}^2 & c_t |y^1|^2 + a_t |y^2|^2 + b_t |y^3|^2 & -\bar{y}^2 y^3 \\ -y^1 \bar{y}^3 & -y^2 \bar{y}^3 & b_t |y^1|^2 + c_t |y^2|^2 + a_t |y^3|^2 \end{bmatrix} \quad (31)$$

It is now almost obvious that the determinant of the matrix in (31) is dependent only on the moduli of y^1 , y^2 and y^3 . Concretely, we have

$$\begin{aligned} F_t(y) &:= \det(\langle \cdot \otimes y | C_{\Phi_t} | \cdot \otimes y \rangle) = \\ &= a_t b_t c_t (|y^1|^6 + |y^2|^6 + |y^3|^6) + \\ &+ (a_t b_t^2 + b_t c_t^2 + c_t a_t^2 - c_t) (|y^2|^2 |y^3|^4 + |y^3|^2 |y^1|^4 + \\ &+ |y^1|^2 |y^2|^4) + (a_t c_t^2 + b_t a_t^2 + c_t b_t^2 - b_t) \cdot \\ &\cdot (|y^1|^2 |y^3|^4 + |y^2|^2 |y^1|^4 + |y^3|^2 |y^2|^4) + \\ &+ (a_t^3 + b_t^3 + c_t^3 + 3a_t b_t c_t - 2) |y^1|^2 |y^2|^2 |y^3|^2 \end{aligned} \quad (32)$$

where we introduced the notation $F_t(y)$ for later convenience. From formula (32), we see that point 1) of the proposition holds. If we note that $a_t + b_t + c_t = 2 \forall t \in [0, 1)$

and that the matrix

$$\begin{bmatrix} 2 & -1 & -1 \\ -1 & 2 & -1 \\ -1 & -1 & 2 \end{bmatrix} \quad (33)$$

is singular, we immediately see that $y^1 = y^2 = y^3 = 1/\sqrt{3}$ solves $F_t(y) = 0$. Because $F_t(y)$ only depends on the moduli of the coordinates of y , we see that all vectors y that fulfill equation (25) correspond to singular $\langle \cdot \otimes y | C_{\Phi_t} | \cdot \otimes y \rangle$. Indeed, for such vectors the matrix $\langle \cdot \otimes y | C_{\Phi_t} | \cdot \otimes y \rangle$ is a multiple of

$$\begin{bmatrix} 2 & -e^{i(\phi_2 - \phi_1)} & -e^{i(\phi_3 - \phi_1)} \\ -e^{i(\phi_1 - \phi_2)} & 2 & -e^{i(\phi_3 - \phi_2)} \\ -e^{i(\phi_1 - \phi_3)} & -e^{i(\phi_2 - \phi_3)} & 2 \end{bmatrix}, \quad (34)$$

where $y^1 = |y^1| e^{i\phi_1}$, $y^2 = |y^2| e^{i\phi_2}$ and $y^3 = |y^3| e^{i\phi_3}$. This matrix is singular.

We now show that there are no other y with non-vanishing y^1 , y^2 and y^3 such that $F_t(y) = 0$. Because

C_{Φ_t} is block positive (cf. eq. (16)), points at which $F_t(y)$ vanishes must be minima of F_t , considered as a function of the coordinates of y . More conveniently, if we define $l_1 := |y^1|^2$, $l_2 := |y^2|^2$ and $l_3 := |y^3|^2$ and write $F_t(y)$ as a function $F_t(l_1, l_2, l_3)$ of l_1 , l_2 and l_3 , singular C_{Φ_t} will correspond to minima of $F_t(l_1, l_2, l_3)$, as long as we are only interested in y 's with all coordinates non-vanishing (i.e. $l_i > 0 \forall i=1,2,3$). The partial derivatives of $F_t(l_1, l_2, l_3)$ with respect to l_1 , l_2 and l_3 equal

$$3A_t l_1^2 + B_t(2l_1 l_3 + l_2^2) + C_t(2l_1 l_2 + l_3^2) + D_t l_2 l_3 \quad (35)$$

$$3A_t l_2^2 + B_t(2l_2 l_1 + l_3^2) + C_t(2l_2 l_3 + l_1^2) + D_t l_3 l_1 \quad (36)$$

$$3A_t l_3^2 + B_t(2l_3 l_2 + l_1^2) + C_t(2l_3 l_1 + l_2^2) + D_t l_1 l_2 \quad (37)$$

respectively, where we have introduced the notation

$$A_t := a_t b_t c_t \quad (38)$$

$$B_t := a_t b_t^2 + b_t c_t^2 + c_t a_t^2 - c_t \quad (39)$$

$$C_t := a_t c_t^2 + b_t a_t^2 + c_t b_t^2 - b_t \quad (40)$$

$$D_t := a_t^3 + b_t^3 + c_t^3 + 3a_t b_t c_t - 3a_t - 2 \quad (41)$$

By equating the partial derivatives (35)-(37) to zero and summing up the resulting equations, we get

$$(3A_t + B_t + C_t)(l_1^2 + l_2^2 + l_3^2) + (2B_t + 2C_t + D_t)(l_1 l_2 + l_2 l_3 + l_3 l_1) = 0 \quad (42)$$

which must be fulfilled by squared moduli of coordinates of all vectors y such that $\det(\langle \cdot \otimes y | C_{\Phi_t} | \cdot \otimes y \rangle) = 0$. Because we already know that $l_1 = l_2 = l_3$ results in a singular $\langle \cdot \otimes y | C_{\Phi_t} | \cdot \otimes y \rangle$, we must have

$$3A_t + B_t + C_t = -2B_t - 2C_t - D_t \quad (43)$$

If we can prove that the above expressions are non-vanishing for all t , equation (42) will imply

$$l_1^2 + l_2^2 + l_3^2 = l_1 l_2 + l_2 l_3 + l_3 l_1 \quad (44)$$

which by Schwartz inequality can only hold if $l_1 = l_2 = l_3$. The expression $3A_t + B_t + C_t$ indeed does not vanish for any $t \in [0, 1)$. Elementary algebra shows that

$$\begin{aligned} 3A_t + B_t + C_t &= (a_t + b_t + c_t)(a_t b_t + b_t c_t + c_t a_t) + \\ &\quad - c_t - b_t = 2(a_t b_t + b_t c_t + c_t a_t) - c_t - b_t = \\ &= \frac{(1-t)^3}{(1-t+t^2)^2} \end{aligned} \quad (45)$$

which is positive for all $t \in [0, 1)$. In this way we have proved that the only choice of $l_1 \neq 0$, $l_2 \neq 0$, $l_3 \neq 0$ such that $F_t(l_1, l_2, l_3) = 0$ is $l_1 = l_2 = l_3$. This corresponds to normalized vectors y that fulfill equations (25). To prove point 2) of the proposition, let us now consider the case $l_1 l_2 l_3 = 0$. We shall focus on the case $l_1 = 0$, $l_2 = 1$, where we have given up the normalization assumption for

y for a while. For other combinations of vanishing/non-vanishing l_i , arguments follow analogously. In the case of our interest, the matrix $\langle \cdot \otimes y | C_{\Phi_t} | \cdot \otimes y \rangle$ equals

$$\begin{bmatrix} b_t + c_t l_3 & 0 & 0 \\ 0 & a_t + b_t l_3 & -e^{i(\phi_3 - \phi_2)} \sqrt{l_3} \\ 0 & -e^{i(\phi_2 - \phi_3)} \sqrt{l_3} & c_t + a_t l_3 \end{bmatrix}, \quad (46)$$

where ϕ_2 and ϕ_3 are defined as in equation (34). Because the above matrix is a direct sum of a 1×1 and 2×2 matrix, and the individual coefficients are of degree at most 1 in l_3 , it is very easy to find the values of l_3 corresponding to singular $\langle \cdot \otimes y | C_{\Phi_t} | \cdot \otimes y \rangle$. It turns out that the only solution when $t \in (0, 1)$ is

$$l_3 = \frac{1}{\sqrt{t}} \quad (47)$$

This corresponds to the following family of normalized vectors y ,

$$[y^1, y^2, y^3] = \left[0, \sqrt{\frac{t}{1+t}} e^{i\phi_2}, \sqrt{\frac{1}{1+t}} e^{i\phi_3} \right] \quad (48)$$

in accordance with formula (26). With little additional effort, one can show that formula (48) extends to the case of $t = 0$. By a permutation symmetry of C_{Φ_t} ,

$$(C_{\Phi_t})_{ij,kl} = (C_{\Phi_t})_{\gamma(i)\gamma(j),\gamma(k)\gamma(l)} \quad (49)$$

with γ equal to the three-cycle $1 \rightarrow 2 \rightarrow 3 \rightarrow 1$, we see that indeed formulas (27) and (28) describe the only remaining choices of the moduli of the coordinates of y that result in singular $\langle \cdot \otimes y | C_{\Phi_t} | \cdot \otimes y \rangle$. In other words, point 2) of the proposition holds. To prove point 3), one only needs to show that the matrices $\langle \cdot \otimes y | C_{\Phi_t} | \cdot \otimes y \rangle$ are of rank 2 for the three choices (25)-(28) of moduli of the coordinates of y referred to in point 2). We leave this as an elementary algebra exercise for the reader. Points 4), 5) and 6) follow because for every y such that $\det(\langle \cdot \otimes y | C_{\Phi_t} | \cdot \otimes y \rangle) = 0$, there exists, up to scaling, exactly one x such that $\langle x \otimes y | C_{\Phi_t} | x \otimes y \rangle = 0$ (cf. point 3)). Moreover, these x 's never repeat for different y 's. Using block positivity of C_{Φ_t} , we get

$$\forall x \in \mathbb{C}^3! \exists y \in \mathbb{C}^3 \langle x \otimes y | C_{\Phi_t} | x \otimes y \rangle = 0 \quad (50)$$

which immediately proves that $\text{rk}(\langle x \otimes \cdot | C_{\Phi_t} | x \otimes \cdot \rangle) \geq 2$ as well as the assertions of points 4) and 3). To see that the x 's really never repeat, one can simply determine the form of the vectors in the kernel of $\langle \cdot \otimes y | C_{\Phi_t} | \cdot \otimes y \rangle$ and see that they really never coincide for different y . For y that fulfills relationships (25), i.e.

$$y = [e^{i\phi_1}, e^{i\phi_2}, e^{i\phi_3}] / \sqrt{3} \quad (51)$$

the matrix $\langle \cdot \otimes y | C_{\Phi_t} | \cdot \otimes y \rangle$ takes the form (34). Elementary algebra shows that the kernel of this matrix consists of vectors proportional to

$$x = [e^{-i\phi_1}, e^{-i\phi_2}, e^{-i\phi_3}] / \sqrt{3} \quad (52)$$

Similarly, for y that fulfills the relationships (26), i.e.

$$y = \left[0, e^{i\phi_2} \sqrt{\frac{t}{1+t}}, e^{i\phi_3} \sqrt{\frac{1}{1+t}} \right] \quad (53)$$

the matrix $\langle \cdot \otimes y | C_{\Phi_t} | \cdot \otimes y \rangle$ takes the form (46). Again, elementary algebra shows that the kernel of this matrix consists of vectors proportional to

$$x = \left[0, e^{-i\phi_2} \sqrt{\frac{1}{1+t}}, e^{-i\phi_3} \sqrt{\frac{t}{1+t}} \right] \quad (54)$$

By the permutation symmetry (49), we also have

$$x = \left[e^{-i\phi_1} \sqrt{\frac{t}{1+t}}, 0, e^{-i\phi_3} \sqrt{\frac{1}{1+t}} \right] \quad (55)$$

corresponding to

$$y = \left[e^{i\phi_1} \sqrt{\frac{1}{1+t}}, 0, e^{i\phi_3} \sqrt{\frac{t}{1+t}} \right] \quad (56)$$

and

$$x = \left[e^{-i\phi_1} \sqrt{\frac{1}{1+t}}, e^{-i\phi_2} \sqrt{\frac{t}{1+t}}, 0 \right] \quad (57)$$

corresponding to

$$y = \left[e^{i\phi_1} \sqrt{\frac{t}{1+t}}, e^{i\phi_2} \sqrt{\frac{1}{1+t}}, 0 \right] \quad (58)$$

Note that the above x 's and y 's agree with formulas (8)-(11) in¹. It is now clear that the x 's given by formulas (52), (54), (55) and (57) can never be proportional to each other, as long as the corresponding y 's differ. As explained above, this implies points 4), 5) and 6) of the proposition. \square

Due to the above proposition and Proposition II.8, the nonexistence of matrices $R, S \in M_3$ such that

$$C_{\Phi_{t_1}} = \text{Ad}_{R \otimes S} C_{\Phi_{t_2}} \quad (59)$$

for arbitrary $t_1 \neq t_2$, $t_1, t_2 \in [0, 1)$ would imply the main result of this paper, cf. Proposition II.5 and the discussion above equation (23). In the following we show that indeed equality (59) cannot hold for any $R, S \in M_3$ and $t_1 \neq t_2$.

Proposition II.10. *The Choi matrices of the Ha-Kye maps¹ defined by formula (24) with $a_t = (1-t)^2/(1-t+t^2)$, $b_t = t^2/(1-t+t^2)$, $c_t = 1/(1-t+t^2)$, $t \in [0, 1)$ are not locally equivalent for different t , i.e. equality (59) does not hold for any $R, S \in M_3$ if $t_1 \neq t_2$, $t_1, t_2 \in [0, 1)$.*

Proof. By nondecomposability of C_{Φ_t} for all $t \in [0, 1)$, we get that the hypothetical matrices R, S in (59) would have to be invertible, cf. Proposition II.8. If the equality really held, the following equivalence would also need to be true

$$F_{t_1}(y) = 0 \Leftrightarrow F_{t_2}(S^*y) = 0 \quad (60)$$

cf. the definition of F_t , eq. (32). Let us consider $y = [e^{i\phi_1}, e^{i\phi_2}, e^{i\phi_3}] / \sqrt{3}$ for arbitrary ϕ_1, ϕ_2 and ϕ_3 , which is an admissible choice of y according to the proof of Proposition II.9. We now show that none of the moduli of the coordinates of S^*y can vanish for any choice of the phases ϕ_j , $j = 1, 2, 3$. Indeed, according to the proof of Proposition II.9, at most two of them could possibly vanish for some choice of ϕ_j 's to ensure $\det(\langle \cdot \otimes S^*y | C_{\Phi_{t_2}} | \cdot \otimes S^*y \rangle) = 0$. Let the non-vanishing coordinate be the third one and assume that the first one vanishes. Since S^* is non-singular, by changing one of the ϕ_j 's by a sufficiently small δ , we can make the first coordinate non-negative without changing the third coordinate significantly. Hence, the ratio of the modulus of the first coordinate to the modulus of the second coordinate would have to become different from zero. Thus, by continuity of the moduli of the coordinates of S^*y as functions of ϕ_j 's, the ratio of the first coordinate to the second coordinate would need to take an infinite number of values as a function of the δ . However, this is not possible if $\det(\langle \cdot \otimes S^*y | C_{\Phi_{t_2}} | \cdot \otimes S^*y \rangle)$ is assumed to vanish for all ϕ_j 's, since by the proof of Proposition II.9, only three values of the ratio are allowed, namely 0, $\sqrt{t_2}$ and 1, cf. equations (52)-(57). In this way we have shown that the moduli of the coordinates of S^*y for y of the form $[e^{i\phi_1}, e^{i\phi_2}, e^{i\phi_3}] / \sqrt{3}$ must all be non-vanishing, no matter what the choice of the ϕ_j 's is. Let us note that these coordinates equal

$$\frac{1}{\sqrt{3}} \sum_{l=1}^3 \overline{S_{lk}} e^{i\phi_l} \quad (61)$$

for $k = 1, 2, 3$. The numbers S_{lk} are the matrix coefficients of S . According to an argument presented in the proof of Proposition II.9, cf. equations (52)-(57), the only possibility for $\langle \cdot \otimes S^*y | C_{\Phi_{t_2}} | \cdot \otimes S^*y \rangle$ to be singular for all choices of the ϕ_j 's is when

$$\left| \sum_{l=1}^3 \overline{S_{l1}} e^{i\phi_l} \right| = \left| \sum_{l=1}^3 \overline{S_{l2}} e^{i\phi_l} \right| = \left| \sum_{l=1}^3 \overline{S_{l3}} e^{i\phi_l} \right| \quad (62)$$

for all choices of $\{\phi_j\}_{j=1,2,3}$. With little additional effort, one can show that the above equalities can only hold if the matrix S^* has exactly one nonzero entry in each row or that some of its rows must be proportional to each other (cf. Appendix C). Because in our setting S^* is necessarily non-singular, it must therefore be a permutation matrix multiplied by a non-singular diagonal matrix. We now show that the permutation part is necessarily a three-cycle or the identity permutation. To exclude

transpositions as the permutation part, it is enough to observe that the specific vectors y of the form (54), (55) and (57) with $t = t_1$ cannot all transform to some of their legitimate counterparts for $t = t_2$, i.e. some vectors of the form (54), (55) and (57), or possibly (52), with $t = t_2$. We provide a detailed argument for the transposition $2 \leftrightarrow 3$ and skip the other ones because they follow by trivial analogy. Let us therefore assume that

$$S^* = \begin{bmatrix} \zeta_1 & 0 & 0 \\ 0 & \zeta_2 & 0 \\ 0 & 0 & \zeta_3 \end{bmatrix} \begin{bmatrix} 1 & 0 & 0 \\ 0 & 0 & 1 \\ 0 & 1 & 0 \end{bmatrix} \quad (63)$$

where $\zeta_1, \zeta_2, \zeta_3 \in \mathbb{Z}$, $\zeta_1 \zeta_2 \zeta_3 \neq 0$. For y of the form (54) with $t = t_1$, we have

$$S^*y = \left[0, \zeta_2 e^{i\phi_2} \sqrt{\frac{1}{1+t_1}}, \zeta_3 e^{i\phi_3} \sqrt{\frac{t_1}{1+t_1}} \right] \quad (64)$$

According to Proposition II.9, the only possibility for $\langle \cdot \otimes S^*y | C_{\Phi_2} | \cdot \otimes S^*y \rangle$ to be singular is when

$$\zeta_2 e^{i\phi_2} \sqrt{\frac{1}{1+t_1}} = \alpha e^{i\phi'_2} \sqrt{\frac{t_2}{1+t_2}} \quad (65)$$

$$\zeta_3 e^{i\phi_3} \sqrt{\frac{t_1}{1+t_1}} = \alpha e^{i\phi'_3} \sqrt{\frac{1}{1+t_2}} \quad (66)$$

for some phases ϕ'_2, ϕ'_3 and a scaling factor $\alpha \in \mathbb{R} \setminus \{0\}$. In the above formula, we used the fact that at least one of the parameters t_1, t_2 is nonzero. Similarly, by considering the transforms S^*y for y of the form (55) and (57), we arrive at the following equations for t_1, t_2 and the ζ_i 's

$$\zeta_1 e^{i\phi_1} \sqrt{\frac{1}{1+t_1}} = \beta e^{i\phi'_1} \sqrt{\frac{t_2}{1+t_2}} \quad (67)$$

$$\zeta_2 e^{i\phi_2} \sqrt{\frac{t_1}{1+t_1}} = \beta e^{i\phi'_2} \sqrt{\frac{1}{1+t_2}} \quad (68)$$

and

$$\zeta_1 e^{i\phi_1} \sqrt{\frac{t_1}{1+t_1}} = \gamma e^{i\phi'_1} \sqrt{\frac{1}{1+t_2}} \quad (69)$$

$$\zeta_3 e^{i\phi_2} \sqrt{\frac{1}{1+t_1}} = \gamma e^{i\phi'_3} \sqrt{\frac{t_2}{1+t_2}} \quad (70)$$

where $\phi'_1, \phi'_2, \phi'_3$ and $\beta, \gamma \in \mathbb{R} \setminus \{0\}$ - arbitrary scaling factors.

By calculating the moduli of (66) and (70), we get the following conditions on $t_1 \in (0, 1)$, $t_2 \in (0, 1)$ and $\zeta_3 \neq 0$

$$|\zeta_3| \sqrt{\frac{1}{1+t_1}} = |\gamma| \sqrt{\frac{t_2}{1+t_2}} \quad (71)$$

$$|\zeta_3| \sqrt{\frac{t_1}{1+t_1}} = |\alpha| \sqrt{\frac{1}{1+t_2}} \quad (72)$$

Clearly, the above formulas can only hold if

$$t_1 t_2 = \left| \frac{\alpha}{\gamma} \right|^2 \quad (73)$$

In a similar way, one can obtain

$$t_1 t_2 = \left| \frac{\beta}{\alpha} \right|^2 \quad (74)$$

and

$$t_1 t_2 = \left| \frac{\gamma}{\beta} \right|^2 \quad (75)$$

The above formulas imply

$$t_1 t_2 = 1 \quad (76)$$

which is not true for any $t_1, t_2 \in (0, 1)$, $t_1 \neq t_2$. Thus matrices S^* of the form (63) are not good candidates for the transform S^* . By trivial analogy, the same conclusion holds when the particular transposition matrix in (63) is replaced with another one. Thus we have proved that the only possible form of the matrix S^* is

$$S^* = \begin{bmatrix} \zeta_1 & 0 & 0 \\ 0 & \zeta_2 & 0 \\ 0 & 0 & \zeta_3 \end{bmatrix} \begin{bmatrix} 0 & 0 & 1 \\ 1 & 0 & 0 \\ 0 & 1 & 0 \end{bmatrix} \quad (77)$$

or

$$S^* = \begin{bmatrix} \zeta_1 & 0 & 0 \\ 0 & \zeta_2 & 0 \\ 0 & 0 & \zeta_3 \end{bmatrix} \begin{bmatrix} 0 & 1 & 0 \\ 0 & 0 & 1 \\ 1 & 0 & 0 \end{bmatrix} \quad (78)$$

or

$$S^* = \begin{bmatrix} \zeta_1 & 0 & 0 \\ 0 & \zeta_2 & 0 \\ 0 & 0 & \zeta_3 \end{bmatrix} \quad (79)$$

It turns out that for matrices S^* of the form (77), (78) and (79), the equality (59) can only hold if $t_1 = t_2$. In the following, we prove this for S^* of the form (77). For the cases of (78) and (79), the reasoning follows analogously and we leave the details to the interested reader. Once we have convinced ourselves that matrices of the form (77), (78) and (79) are of no use in transforming different members of the family $\{\Phi_t\}_{t \in [0,1]}$ between each other, we will have completed the proof of the proposition.

Let us assume that S^* is of the form (77). We again consider the transforms S^*y for the characteristic vectors y of the form (54), (55) and (57) with $t = t_1$. Just as before, these must be again of the form (54), (55), (57), or possibly (52), for $t = t_2$. If y is of the form (54), the transform S^*y equals

$$S^*y = \left[\zeta_1 e^{i\phi_3} \sqrt{\frac{1}{1+t_1}}, 0, \zeta_3 e^{i\phi_2} \sqrt{\frac{t_1}{1+t_1}} \right] \quad (80)$$

Under the assumption of $t_1 \neq 0$ or $t_2 \neq 0$, which is true when $t_1 \neq t_2$, the requirement that $\det(\langle \cdot \otimes S^*y | C_{\Phi_2} | \cdot \otimes S^*y \rangle)$ vanishes can only be met if the S^*y given by (80) coincides with

$$\alpha \left[e^{i\phi'_1} \sqrt{\frac{1}{1+t_2}}, 0, e^{i\phi'_3} \sqrt{\frac{t_2}{1+t_2}} \right] \quad (81)$$

for some phases ϕ'_1 and ϕ'_3 and a scaling factor $\alpha \in \mathbb{R} \setminus \{0\}$. This clearly implies

$$\zeta_1 e^{i\phi_3} \sqrt{\frac{1}{1+t_1}} = \alpha e^{i\phi'_1} \sqrt{\frac{1}{1+t_2}} \quad (82)$$

$$\zeta_3 e^{i\phi_2} \sqrt{\frac{t_1}{1+t_1}} = \alpha e^{i\phi'_3} \sqrt{\frac{t_2}{1+t_2}} \quad (83)$$

Similarly, by considering the transforms S^*y for vectors y of the form (55) and (57), we arrive at the conditions

$$\zeta_1 e^{i\phi_3} \sqrt{\frac{t_1}{1+t_1}} = \beta e^{i\phi''_1} \sqrt{\frac{t_2}{1+t_2}} \quad (84)$$

$$\zeta_2 e^{i\phi_1} \sqrt{\frac{1}{1+t_1}} = \beta e^{i\phi''_2} \sqrt{\frac{1}{1+t_2}} \quad (85)$$

and

$$\zeta_2 e^{i\phi_1} \sqrt{\frac{t_1}{1+t_1}} = \gamma e^{i\phi'''_2} \sqrt{\frac{t_2}{1+t_2}} \quad (86)$$

$$\zeta_3 e^{i\phi_2} \sqrt{\frac{1}{1+t_1}} = \gamma e^{i\phi'''_3} \sqrt{\frac{1}{1+t_2}} \quad (87)$$

where $\phi''_1, \phi''_2, \phi'''_2$ and ϕ'''_3 denote some arbitrary phases and $\beta, \gamma \in \mathbb{R} \setminus \{0\}$ are arbitrary nonzero scaling factors. By considering the moduli of (82) and (83), we easily obtain

$$|\zeta_1|^2 t_2 = |\zeta_3|^2 t_1 \quad (88)$$

In a similar way, from equations (84)-(87) we can get

$$|\zeta_2|^2 t_2 = |\zeta_1|^2 t_1 \quad (89)$$

$$|\zeta_3|^2 t_2 = |\zeta_2|^2 t_1 \quad (90)$$

We can now multiply equalities (88)-(90) together and get

$$|\zeta_1|^2 |\zeta_2|^2 |\zeta_3|^2 t_2^3 = |\zeta_1|^2 |\zeta_2|^2 |\zeta_3|^2 t_1^3 \Rightarrow t_2 = t_1 \quad (91)$$

This clearly contradicts $t_1 \neq t_2$ and proves the main assertion of the proposition. \square

As we already explained above, the above proposition implies the main result of this paper. We briefly remind the reader why this is the case. By Proposition II.9, the family $\{\Phi_t\}_{t \in [0,1]}$ of extreme indecomposable maps of the form (24), as considered by Ha and Kye¹, fulfills the assumptions of point 5) of Proposition II.8. Therefore, if equality (18) holds for some positive maps $\{\Xi_i\}_{i=1}^k \subset \mathcal{P}_{3,3}$, there must exist at least two values $t_1, t_2 \in [0,1]$, $t_1 \neq t_2$ such that

$$\Phi_{t_1} = \text{Ad}_A \circ \Phi_{t_2} \circ \text{Ad}_B \quad (92)$$

for some non-singular $A, B \in M_3$ (actually, there must exist an infinite number of such pairs). By Proposition II.5, we know that (92) is equivalent to (59) with $R = B^t$, $S = A$. However, we know from Proposition II.10 that (59) does not hold for any $t_1 \neq t_2$, $R, S \in M_3$. Thus equality (18) cannot be true. By Proposition II.7,

this is equivalent to the non-existence of a necessary and sufficient separability criterion of the form (17). In this way we have arrived at the following conclusion

Theorem II.11. *There does not exist a separability criterion of the form*

$$\rho \in \text{Sep}_{3 \times 3} \Leftrightarrow \forall_{i=1,2,\dots,k} \forall_{B \in M_3} (\mathbb{I} \otimes \Phi_i) \text{Ad}_{\mathbb{I} \otimes B} \rho \geq 0 \quad (93)$$

for a finite set of positive maps $\{\Phi_i\}_{i=1,2,\dots,k}$.

For a more mathematically oriented audience, the same can be expressed, by Proposition II.7, as

Theorem II.12. *The cone $\mathcal{P}_{3,3}$ of positive maps from the set M_3 of three-dimensional matrices into itself is not finitely generated as a mapping cone. In other words, there does not exist a finite set of positive maps $\{\Xi_i\}_{i=1,2,\dots,k}$ such that*

$$\mathcal{P}_{3,3} = \overline{\text{convhull}(\{\text{Ad}_X \circ \Xi_i \circ \text{Ad}_Y | i; X, Y \in M_3\})}, \quad (94)$$

holds.

Some other noteworthy properties of mapping cones can be found in^{15,18–23}.

III. CONCLUSION

It is a generally accepted opinion that the detection of all density matrix entanglement in 3×3 and more complex systems is a difficult task. In this paper we have shown that for such systems there is no hope to have a sufficient and necessary criterion for separability of a state, similar in structure to the famous positive partial transpose for 2×2 and 2×3 systems^{8,9}. This is even true despite the fact that we allow arbitrary local transforms in formula (17). The argument we presented also suggests that in order to have a strong entanglement detection procedure, one should at least consider a criterion of the form (17) with $\{\Xi_i\}_{i=1}^k$ substituted with the whole Ha-Kye family of positive maps $\{\Phi_t\}_{t \in [0,1]}$ ¹.

From a more mathematics centered point of view, our result shows that the cone $\mathcal{P}_{3,3}$ of positive maps from the set of three-dimensional matrices into itself, is not finitely generated as a mapping cone. This is in contrast with $\mathcal{P}_{3,2}$, $\mathcal{P}_{2,3}$ and $\mathcal{P}_{2,2}$, where all positive maps turn out to be decomposable^{10,11}.

Remark III.1. *It is natural to expect that the cones $\mathcal{P}_{m,n}$ of positive maps with $m \geq 3$ and $n \geq 3$ are also not finitely generated as mapping cones, however the author has not yet found a proof of this fact. If it holds, we immediately obtain a generalization of the main result of this paper to $m \times n$ systems where $m, n \geq 3$.*

Remark III.2. *In the light of the proof presented above, it is natural to ask whether the whole family of Ha-Kye maps can provide us with a necessary and sufficient separability criterion of the form (17).*

ACKNOWLEDGEMENT

The result presented in this paper is a byproduct of work done together with Erling Størmer. The author is heartily indebted to Karol Życzkowski, whose emails finally convinced him write this paper up, even after a few years spent outside academia. Valuable comments by Martyna Rymer are gratefully acknowledged. This work has been financially supported by the NCN Grant no. 2015/18/A/ST2/00274.

Appendix A: Choi matrix change under multiplication by conjugation maps

We would like to show that

$$C_{\text{Ad}_A \circ \Phi \circ \text{Ad}_B} = \text{Ad}_{B^t \otimes A} C_\Phi \quad (\text{A1})$$

for arbitrary $A, B \in M_3$ and a positive map $\mathcal{P}_{3,3}$. As expected, this result holds for general $\mathcal{P}_{m,n}$ as well.

From the definition (5) of the Choi matrix, we immediately see that

$$C_{\text{Ad}_A \circ \Phi \circ \text{Ad}_B} = \text{Ad}_{\mathbb{I} \otimes A} C_{\Phi \circ \text{Ad}_B} \quad (\text{A2})$$

Thus, we only need to prove that

$$C_{\Phi \circ \text{Ad}_B} = \text{Ad}_{B^t \otimes \mathbb{I}} C_\Phi \quad (\text{A3})$$

We have

$$C_{\Phi \circ \text{Ad}_B} = (\mathbb{I} \otimes \Phi) \sum_{i,j} \sum_{k,l} e_{ij} \otimes B_{ki} \overline{B_{lj}} e_{kl}, \quad (\text{A4})$$

where B_{ij} are the matrix coefficients of B . We further have

$$\sum_{i,j} \sum_{k,l} e_{ij} \otimes B_{ki} \overline{B_{lj}} e_{kl} = \sum_{i,j} \sum_{k,l} B_{ki} \overline{B_{lj}} e_{ij} \otimes e_{kl}, \quad (\text{A5})$$

which is the same as

$$\text{Ad}_{B^t \otimes \mathbb{I}} \sum_{k,l} e_{kl} \otimes e_{kl} \quad (\text{A6})$$

Thus we can write

$$C_{\Phi \circ \text{Ad}_B} = (\mathbb{I} \otimes \Phi) \text{Ad}_{B^t \otimes \mathbb{I}} \sum_{k,l} e_{kl} \otimes e_{kl} \quad (\text{A7})$$

From this, we easily obtain

$$C_{\Phi \circ \text{Ad}_B} = \text{Ad}_{B^t \otimes \mathbb{I}} \sum_{k,l} e_{kl} \otimes \Phi(e_{kl}) = \text{Ad}_{B^t \otimes \mathbb{I}} C_\Phi \quad (\text{A8})$$

Appendix B: Extreme positive maps in a finitely generated mapping cone

In this appendix we prove that extreme positive maps $\Phi \in \mathcal{P}_{3,3}$ in a finitely generated closed mapping cone of

the form (18) must either be of the form $\text{Ad}_X \circ \Xi_i \circ \text{Ad}_Y$ for some $i \in \{1, 2, \dots, k\}$ and $X, Y \in M_3$ or there must exist $x \in \mathbb{C}^3$ or $y \in \mathbb{C}^3$ such that

$$\text{rk}(\langle x \otimes \cdot | C_\Phi | x \otimes \cdot \rangle) < 2 \quad (\text{B1})$$

or

$$\text{rk}(\langle \cdot \otimes y | C_\Phi | \cdot \otimes y \rangle) < 2 \quad (\text{B2})$$

Point 3) of Proposition II.8 trivially follows from this result.

Let us assume that Φ is an element of the closed convex cone (18). Equivalently, there exist sequences $\{i_j^l\}_{l=1}^{+\infty} \subset \mathbb{N}$, $\{X_j^l\}_{l=1}^{+\infty} \subset M_3$ and $\{Y_j^l\}_{l=1}^{+\infty} \subset M_3$, for $j = 1, 2, \dots, 82$ such that

$$\sum_{j=1}^{82} \text{Ad}_{X_j^l} \circ \Xi_{i_j^l} \circ \text{Ad}_{Y_j^l} \xrightarrow{l \rightarrow +\infty} \Phi \quad (\text{B3})$$

The possibility to set an upper border of 82 on summation by j follows from Carathéodory's theorem²⁵.

Note that every non-zero positive map Φ satisfies $\text{Tr}(C_\Phi) > 0$ and hence can be normalized to have trace of its Choi matrix equal to one. Moreover, positive maps Φ such that C_Φ have unit trace, form a compact subset of the subspace of all maps Ψ in $\mathcal{H}_{m,n}$ with the trace of C_Ψ equal to one. In that subspace, equation (B3) induces

$$\sum_{j=1}^{82} \lambda_j^l \frac{\text{Ad}_{X_j^l} \circ \Xi_{i_j^l} \circ \text{Ad}_{Y_j^l}}{\tau_j^l} \xrightarrow{l \rightarrow +\infty} \frac{\Phi}{\text{Tr}(C_\Phi)}, \quad (\text{B4})$$

where $\tau_j^l := \text{Tr}\left(C_{\text{Ad}_{X_j^l} \circ \Xi_{i_j^l} \circ \text{Ad}_{Y_j^l}}\right)$, $\lambda_j^l \geq 0$, the sum $\sum_{j=1}^{82} \lambda_j^l$ tends to 1 for $l \rightarrow +\infty$, and we are free to assume that $\tau_j^l > 0$ for all j and l . The last assumption is permissible because by setting X_j^l and Y_j^l equal to \mathbb{I} if initially $\text{Tr}\left(C_{\text{Ad}_{X_j^l} \circ \Xi_{i_j^l} \circ \text{Ad}_{Y_j^l}}\right) = 0$ and bringing the respective λ_j^l to zero, we can make all the τ_j^l 's positive, at the same time keeping the relationship (B4) a consequence of (B3). Moreover, we can assume that the largest singular values of X_j^l 's and Y_j^l 's are equal to one because it is always possible to rescale the X_j^l 's and Y_j^l 's and their singular values with no influence on $(\text{Ad}_{X_j^l} \circ \Xi_{i_j^l} \circ \text{Ad}_{Y_j^l}) / \tau_i^l$.

In summary, equation (B3) implies (B4) with λ_j^l 's and $(\text{Ad}_{X_j^l} \circ \Xi_{i_j^l} \circ \text{Ad}_{Y_j^l}) / \tau_i^l$'s contained in compact subsets of \mathbb{R} and $\mathcal{P}_{3,3}$. Therefore, there must exist limits

$$\frac{\text{Ad}_{X_j^{l_w}} \circ \Xi_{i_j^{l_w}} \circ \text{Ad}_{Y_j^{l_w}}}{\tau_i^{l_w}} \xrightarrow{w \rightarrow +\infty} \Sigma_j \in \mathcal{P}_{3,3} \quad (\text{B5})$$

$$\lambda_j^{l_w} \xrightarrow{w \rightarrow +\infty} \lambda_j \geq 0 \quad (\text{B6})$$

for certain subsequence $\{l_w\}_{w=1}^{+\infty} \subset \mathbb{N}$. Hence,

$$\frac{\Phi}{\text{Tr}(C_\Phi)} = \sum_{j=1}^{82} \lambda_j \Sigma_j \quad (\text{B7})$$

Obviously, for Φ extreme, we can assume that the above sum only contains a single nonzero term Σ , where

$$\Sigma = \lim_{w \rightarrow +\infty} \frac{\text{Ad}_{X^w} \circ \Xi_i \circ \text{Ad}_{Y^w}}{\text{Tr}(C_{\text{Ad}_{X^w} \circ \Xi_i \circ \text{Ad}_{Y^w}})} \quad (\text{B8})$$

for appropriately chosen X^w and Y^w with largest singular values equal to one. By an appropriate choice of a subsequence, we can also assure that i_w is a constant $i \in \{1, 2, \dots, k\}$. In the following, we shall simply assume that $i_w = i$ in equation (B8).

We now prove that the limit on the right-hand side of equation (B8) must either be of the form $\text{Ad}_X \circ \Xi_i \circ \text{Ad}_Y$ for some $X, Y \in M_3$ or equation (B1) or (B2) with Σ substituted for Φ must hold for some x or y in \mathbb{C}^3 .

Because X^w and Y^w can be assumed to belong to a compact set of matrices with largest singular value equal to one, there must exist a subsequence $\{w_t\}_{t=1}^{+\infty}$ and matrices X, Y with largest singular values equal to one, such that

$$X^{w_t} \xrightarrow{t \rightarrow +\infty} X \quad (\text{B9})$$

$$Y^{w_t} \xrightarrow{t \rightarrow +\infty} Y \quad (\text{B10})$$

Without loss of generality, we may assume that X^w and Y^w themselves converge to X and Y , respectively.

If $\lim_{w \rightarrow +\infty} \text{Tr}(C_{\text{Ad}_{X^w} \circ \Xi_i \circ \text{Ad}_{Y^w}}) = \tau > 0$, our main assertion is obviously true, because we obtain

$$\Sigma = \text{Ad}_{X/\sqrt[4]{\tau}} \circ \Xi_i \circ \text{Ad}_{Y/\sqrt[4]{\tau}} \quad (\text{B11})$$

and

$$\Phi = \text{Ad}_{\sqrt[4]{\frac{\sigma}{\tau}} X} \circ \Xi_i \circ \text{Ad}_{\sqrt[4]{\frac{\sigma}{\tau}} Y}, \quad (\text{B12})$$

where $\sigma := \text{Tr}(C_\Phi) > 0$. Therefore, we only need to consider the case of $\text{Tr}(C_{\text{Ad}_{X^w} \circ \Xi_i \circ \text{Ad}_{Y^w}})$ that tends to zero as $w \rightarrow +\infty$.

By Proposition II.5

$$C_{\text{Ad}_{X^w} \circ \Xi_i \circ \text{Ad}_{Y^w}} = \text{Ad}_{(Y^w)^t \otimes X^w} C_{\Xi_i} \quad (\text{B13})$$

Denote the second and the third singular value of $(Y^w)^t$ by σ_2^w, σ_3^w . Similarly, let the second and the third singular value of X^w be δ_2^w and δ_3^w . For convenience, we also define $\sigma_1^w = \delta_1^w := 1$. Let

$$(Y^w)^t = \sum_{i=1}^3 \sigma_i^w |u_i^w\rangle \langle v_i^w| \quad (\text{B14})$$

$$X^w = \sum_{i=1}^3 \delta_i^w |p_i^w\rangle \langle q_i^w| \quad (\text{B15})$$

be decompositions following directly from singular value decompositions of $(Y^w)^t$ and X^w . In particular, $\{u_i^w\}_{i=1}^3, \{v_i^w\}_{i=1}^3, \{p_i^w\}_{i=1}^3, \{q_i^w\}_{i=1}^3$ in equations (B14) and (B15) are orthonormal bases of \mathbb{C}^3 with elements corresponding to rows or columns of unitary matrices present in the respective singular value decompositions. We then have

$$\begin{aligned} C_{\text{Ad}_{X^w} \circ \Xi_i \circ \text{Ad}_{Y^w}} &= \\ &= \sum_{r,s,t,u=1}^3 \sigma_r^w \sigma_t^w \delta_s^w \delta_u^w \langle v_r^w \otimes q_s^w | C_{\Xi_i} | v_t^w \otimes q_u^w \rangle \cdot \\ &\quad \cdot |u_r^w \otimes p_s^w\rangle \langle u_t^w \otimes p_u^w| \quad (\text{B16}) \end{aligned}$$

With no loss of generality, we may assume that all the sequences $\{\sigma_i^w\}_{w=1}^{+\infty}, \{\delta_i^w\}_{w=1}^{+\infty}, \{u_i^w\}_{w=1}^{+\infty}, \{v_i^w\}_{w=1}^{+\infty}, \{p_i^w\}_{w=1}^{+\infty}, \{q_i^w\}_{w=1}^{+\infty}$ are convergent. This is so because there always exists their common convergent subsequence, which we can keep and remove the other terms. It is then easy to see that in an appropriate orthonormal basis, the (r, s, t, u) -th matrix element of $C_{\text{Ad}_{X^w} \circ \Xi_i \circ \text{Ad}_{Y^w}} / \text{Tr}(C_{\text{Ad}_{X^w} \circ \Xi_i \circ \text{Ad}_{Y^w}})$ equals

$$\lim_{w \rightarrow +\infty} \frac{\sigma_r^w \sigma_t^w \delta_s^w \delta_u^w \langle v_r^w \otimes q_s^w | C_{\Xi_i} | v_t^w \otimes q_u^w \rangle}{\sum_{h,j=1}^3 (\sigma_h^w)^2 (\delta_j^w)^2 \langle v_h^w \otimes q_j^w | C_{\Xi_i} | v_h^w \otimes q_j^w \rangle} \quad (\text{B17})$$

Note also that we are free to assume the existence of a subsequence $\{w_n\}_{n=1}^{+\infty}$ such that $\sigma_r^{w_n}, \sigma_t^{w_n}, \delta_s^{w_n}, \delta_u^{w_n} > 0$. Otherwise, we could choose one of the singular values to be equal to 0 all the time on an appropriate subsubsequence and the existence of x or y such that equations (B1) or (B2) hold would immediately follow (actually, C_Φ would turn out to be supported on a 2×3 or smaller subspace). As a consequence of that, with no loss of generality, we may assume $\sigma_r^w, \sigma_t^w, \delta_s^w, \delta_u^w > 0$.

The fact that the denominator of (B17) vanishes in the limit $w \rightarrow +\infty$, together with $\sigma_1 = \delta_1 = 1$ implies that $\langle v_1 \otimes q_1 | C_{\Xi_i} | v_1 \otimes q_1 \rangle = 0$. Let us now focus on the terms with σ_2^w , although the same argument goes for any other of the singular values $\sigma_2^w, \sigma_3^w, \delta_2^w$ and δ_3^w . If one of the corresponding matrix elements of Ξ_i , $\langle v_2 \otimes q_1 | C_{\Xi_i} | v_2 \otimes q_1 \rangle$, equals 0, the following lemma shows that the main assertion of this appendix is true because (B2) holds for some y .

Lemma B.1. *Let $\langle v_1 \otimes q_j | C_{\Xi_i} | v_1 \otimes q_j \rangle = 0$ and $\langle v_2 \otimes q_j | C_{\Xi_i} | v_2 \otimes q_j \rangle = 0$ for some $j \in \{1, 2, 3\}$. Then there exists $y \in \mathbb{C}^3$ such that*

$$\text{rk}(\langle \cdot \otimes y | C_\Phi | \cdot \otimes y \rangle) = \text{rk}(\langle \cdot \otimes y | C_\Sigma | \cdot \otimes y \rangle) < 2 \quad (\text{B18})$$

Proof. Let us decompose the vector q_j in the basis $\{q_u^w\}_{u=1}^3$,

$$q_j = \sum_{u=1}^3 \lambda_{ju}^w q_u^w \quad (\text{B19})$$

Now define

$$y^w := \frac{\sum_{u=1}^3 \frac{\lambda_{ju}^w}{\delta_u^w} p_u^w}{\sqrt{\sum_{u=1}^3 \left| \frac{\lambda_{ju}^w}{\delta_u^w} \right|^2}} \quad (\text{B20})$$

From equation (B16), we easily see that

$$\text{rk} \left(\langle \cdot \otimes y^w | \frac{C_{\text{Ad}_{X^w} \circ \Xi_i \circ \text{Ad}_{Y^w}}}{\text{Tr}(C_{\text{Ad}_{X^w} \circ \Xi_i \circ \text{Ad}_{Y^w}})} | \cdot \otimes y^w \rangle \right) < 2, \quad (\text{B21})$$

where we have used the fact that for the vanishing of $\langle x_1 \otimes y^w | X | x_1 \otimes y^w \rangle$ and $\langle x_2 \otimes y^w | X | x_2 \otimes y^w \rangle$ for a block positive matrix X implies $\text{rk}(\langle \cdot \otimes y^w | X | \cdot \otimes y^w \rangle) < 2$, as long as $x_1 \neq x_2$.

Since y^w 's belong to a compact subset of \mathbb{C}^3 consisting of normalized vectors, there is a subsequence $\{w_n\}_{n=1}^{+\infty}$ and a $y \in \mathbb{C}^3$, $\|y\| = 1$ such that $y^{w_n} \rightarrow y$ as $n \rightarrow +\infty$. Just as we did previously, with no loss of generality we may assume $y^w \rightarrow y$ for $w \rightarrow +\infty$. We thus have

$$\lim_{w \rightarrow +\infty} \frac{C_{\text{Ad}_{X^w} \circ \Xi_i \circ \text{Ad}_{Y^w}}}{\text{Tr}(C_{\text{Ad}_{X^w} \circ \Xi_i \circ \text{Ad}_{Y^w}})} = C_\Sigma \quad (\text{B22})$$

$$\lim_{w \rightarrow +\infty} y^w = y \quad (\text{B23})$$

Inequality (B18) now trivially follows from (B21), (B22), (B23) and the continuity of submatrix determinants. \square

As a consequence of the above lemma, the only case where we have not yet proved our assertion is that of $\langle v_2 \otimes q_1 | C_{\Xi_i} | v_2 \otimes q_1 \rangle > 0$. Because of the arbitrariness of the choice of i and the tensor product side in the above lemma, with no loss of generality we may assume that σ_2^w tends to zero in a *slowest* way among σ_2^w , σ_3^w , δ_2^w , δ_3^w (otherwise, simply begin the argument with a different singular value). By this, we mean the existence of a subsequence $\{w_n\}_{n=1}^{+\infty}$ and numbers $\alpha, \beta, \gamma > 0$ such that

$$\sigma_3^{w_n} < \alpha \sigma_2^{w_n} \quad (\text{B24})$$

$$\delta_2^{w_n} < \beta \sigma_2^{w_n} \quad (\text{B25})$$

$$\delta_3^{w_n} < \gamma \sigma_2^{w_n} \quad (\text{B26})$$

Just like we did earlier a few times, we may neglect terms not in w_n and assume that inequalities (B24)-(B26) hold with w substituted for w_n .

Because the denominator of (B17) contains the term $(\sigma_2^w)^2 \langle v_2 \otimes q_1 | C_{\Xi_i} | v_2 \otimes q_1 \rangle$, all the terms in the numerator that vanish faster than $(\sigma_2^w)^2$ result in zero terms in C_Σ and therefore also in C_Φ . This applies to all terms in (B17) with $s = u = 2$ and $r \neq 3$ or $t \neq 3$, which clearly implies

$$\text{rk}(\langle u_2 \otimes \cdot | C_\Phi | u_2 \otimes \cdot \rangle) = \text{rk}(\langle u_2 \otimes \cdot | C_\Sigma | u_2 \otimes \cdot \rangle) < 2, \quad (\text{B27})$$

Hence, the main assertion of this appendix also holds in the case of positive $\langle v_2 \otimes q_1 | C_{\Xi_i} | v_2 \otimes q_1 \rangle > 0$.

Note that despite the fact that our proof was carried out for Φ in $\mathcal{P}_{3,3}$, it is easily generalized to positive maps in $\mathcal{P}_{m,n}$ for m and n arbitrary, however the inequalities

(B1) and (B2) need to be replaced by the condition that the rank of $\langle x \otimes \cdot | C_\Phi | x \otimes \cdot \rangle$ or $\langle \cdot \otimes y | C_\Phi | \cdot \otimes y \rangle$ must be smaller than the maximum possible value minus one.

Appendix C: A class of moduli preserving linear transformations

In this appendix, we prove that $n \times n$ matrices X such that

$$\left| \sum_{l=1}^n X_{kl} e^{i\phi_l} \right| = \left| \sum_{t=1}^n X_{mt} e^{i\phi_t} \right| \quad \forall_{\{\phi_l\}_{l=1,2,\dots,n}} \quad \forall_{k,m \in \{1,2,\dots,n\}} \quad (\text{C1})$$

are necessarily singular or they only have a single nonzero entry in each row. To show this, it is enough to prove the following lemma.

Lemma C.1. *Let y and z be vectors in \mathbb{C}^n such that*

$$\left| \sum_{l=1}^n y^l e^{i\phi_l} \right| = \left| \sum_{t=1}^n z^t e^{i\phi_t} \right| \quad \forall_{\{\phi_l\}_{l=1,2,\dots,n}} \quad (\text{C2})$$

where y^l and z^t are the coefficients of the vectors y and z , respectively. Then y and z must either be proportional to each other or each of them must have exactly one nonzero coefficient.

Proof. We prove the lemma by induction on n . For $n = 1$, its assertion is obviously true. For $n > 1$, let us write the moduli in (C2) as

$$\left| y^1 e^{i\phi_1} + \sum_{l=2}^n y^l e^{i\phi_l} \right| = \left| z^1 e^{i\phi_1} + \sum_{t=2}^n z^t e^{i\phi_t} \right| \quad (\text{C3})$$

A quick thought reveals that there are only four cases where (C3) can hold for all possible choices of the ϕ_l 's. These are:

1.

$$y^1 = 0 \quad (\text{C4})$$

$$z^1 = 0 \quad (\text{C5})$$

$$\forall_{\{\phi_l\}_{l=2}^n} \left| \sum_{l=2}^n y^l e^{i\phi_l} \right| = \left| \sum_{l=2}^n z^l e^{i\phi_l} \right| \quad (\text{C6})$$

2.

$$y^1 = 0 \quad (\text{C7})$$

$$z^1 \neq 0 \quad (\text{C8})$$

$$\exists_{\alpha > 0} \forall_{\{\phi_l\}_{l=2}^n} \left| \sum_{l=2}^n y^l e^{i\phi_l} \right| = \alpha \quad (\text{C9})$$

$$\forall_{\{\phi_l\}_{l=2}^n} \left| \sum_{t=2}^n z^t e^{i\phi_t} \right| = 0 \quad (\text{C10})$$

3.

$$y^1 \neq 0 \quad (\text{C11})$$

$$z^1 = 0 \quad (\text{C12})$$

$$\exists_{\beta > 0} \forall_{\{\phi_t\}_{t=2}^n} \left| \sum_{t=2}^n z^t e^{i\phi_t} \right| = \beta \quad (\text{C13})$$

$$\forall_{\{\phi_l\}_{l=2}^n} \left| \sum_{l=2}^n y^l e^{i\phi_l} \right| = 0 \quad (\text{C14})$$

4.

$$y^1 \neq 0 \quad (\text{C15})$$

$$z^1 \neq 0 \quad (\text{C16})$$

$$\forall_{\{\phi_l\}_{l=2}^n} \frac{\sum_{l=2}^n y^l e^{i\phi_l}}{y^1} = \frac{\sum_{t=2}^n z^t e^{i\phi_t}}{z^1} \quad (\text{C17})$$

In case 1), the assertion of the lemma automatically holds by induction. In case 2), equation (C9) can only hold if exactly one of the numbers y^2, y^3, \dots, y^n is nonzero. Moreover, equation (C10) can only hold if all the numbers z^2, z^3, \dots, z^n are equal to zero. Altogether, y and z have exactly one nonzero coordinate, which fulfills the assertion of the lemma. Case 3) is symmetric to case 2). In case 4), we must have

$$y^l = \frac{y^1}{z^1} z^l \quad (\text{C18})$$

for $l = 2, 3, \dots, n$. However, equation (C18) obviously also holds when $l = 1$. In other words, the vectors y and z turn out to be proportional to each other, just as in the assertion of the lemma. \square

From the above lemma, the main conclusion of the appendix easily follows if we take as y and z arbitrary two rows of the matrix X . In case they turn out to be proportional to each other, the matrix turns out to be singular. Otherwise, there is exactly one nonzero entry in each of these rows. This argument can be repeated for all row pairs in X to yield the main result of this appendix.

¹K C Ha and S H Kye. Entanglement witnesses arising from exposed positive linear maps. *Open Syst. Inf. Dyn.*, 18:323–337, 2011.

²C H Bennett and G Brassard. Quantum cryptography: Public key distribution and coin tossing. In *Proceedings of the IEEE International Conference on Computers, Systems and Signal Processing*, pages 175–179, New York, 1984.

³N Gisin, G Ribordy, W Tittel, and H Zbinden. Quantum cryptography. *Rev. Mod. Phys.*, 74:145–195, 2002.

⁴L Masanes. All bipartite entangled states are useful for information processing. *Phys. Rev. Lett.*, 96:150501, 2006.

⁵L Masanes. Useful entanglement can be extracted from all non-separable states. *J. Math. Phys.*, 49:022102, 2008.

⁶L Masanes and Doherty A C Liang, Y C. All bipartite entangled states display some hidden nonlocality. *Phys. Rev. Lett.*, 100:090403, 2008.

⁷R Werner. Quantum states with Einstein-Podolsky-Rosen correlations admitting a hidden-variable model. *Phys. Rev. A*, 40:4277–4281, 1989.

⁸A Peres. Separability criterion for density matrices. *Phys. Rev. Lett.*, 77:1413–1415, 1996.

⁹M Horodecki, P Horodecki, and R Horodecki. Separability of mixed states: necessary and sufficient conditions. *Phys. Lett. A*, 223:1–8, 1996.

¹⁰E Størmer. Positive linear maps of operator algebras. *Acta. Math.*, 110:223–278, 1963.

¹¹S L Woronowicz. Positive maps of low dimensional matrix algebras. *Rep. Math. Phys.*, 10:165–183, 1976.

¹²P Horodecki. Separability criterion and inseparable mixed states with positive partial transposition. *Phys. Lett. A*, 232:333–339, 1997.

¹³A Jamiolkowski. Linear transformations which preserve trace and positive semidefiniteness of operators. *Rep. Math. Phys.*, 3:275–278, 1972.

¹⁴M D Choi. Completely positive maps on complex matrices. *Lin. Alg. Appl.*, 10:285–290, 1975.

¹⁵L Skowronek. Cones with a mapping cone symmetry in the finite-dimensional case. *Lin. Alg. Appl.*, 435:361–370, 2011.

¹⁶J De Pillis. Linear transformations which preserve hermitian and positive semidefinite operators. *Pacific J. Math.*, 23:129–137, 1967.

¹⁷E Størmer. Extension of positive maps into $B(\mathcal{H})$. *J. Func. Anal.*, 66:235–254, 1986.

¹⁸E Størmer. Separable states and positive maps. *J. Func. Anal.*, 254:2303–2312, 2008.

¹⁹E Størmer. Duality of cones of positive maps. *Münster J. Math.*, 2:299–310, 2009.

²⁰E Størmer. Separable states and positive maps II. *Math. Scand.*, 105:188–198, 2009.

²¹L Skowronek, E Størmer, and K Życzkowski. Cones of positive maps and their duality relations. *J. Math. Phys.*, 50:062106, 2009.

²²N Johnston and E Størmer. Mapping cones are operator systems. *Bull. Lond. Math. Soc.*, 44:738–748, 2012.

²³N Johnston, L Skowronek, and E Størmer. Generation of mapping cones from small sets. *Lin. Alg. Appl.*, 438:3062–3075, 2013.

²⁴T Ando. Cones and norms in the tensor product of matrix spaces. *Lin. Alg. Appl.*, 379:3–41, 2004.

²⁵R Rockafellar. *Convex analysis*. Princeton University Press, 1997.