

DEFORMATION THEORY OF SCALAR-FLAT KÄHLER ALE SURFACES

JIYUAN HAN AND JEFF A. VIACLOVSKY

ABSTRACT. We prove a Kuranishi-type theorem for deformations of complex structures on ALE Kähler surfaces. This is used to prove that for any scalar-flat Kähler ALE surface, all small deformations of complex structure also admit scalar-flat Kähler ALE metrics. A local moduli space of scalar-flat Kähler ALE metrics is then constructed, which is shown to be universal up to small diffeomorphisms (that is, diffeomorphisms which are close to the identity in a suitable sense). A formula for the dimension of the local moduli space is proved in the case of a scalar-flat Kähler ALE surface which deforms to a minimal resolution of \mathbb{C}^2/Γ , where Γ is a finite subgroup of $U(2)$ without complex reflections.

1. INTRODUCTION

This article is concerned with the following class of metrics:

Definition 1.1. *An ALE Kähler surface (X, g, J) is a Kähler manifold of complex dimension 2 with the following property. There exists a compact subset $K \subset X$ and a diffeomorphism $\psi : X \setminus K \rightarrow (\mathbb{R}^4 \setminus \overline{B})/\Gamma$, such that for each multi-index \mathcal{I} of order $|\mathcal{I}|$*

$$(1.1) \quad \partial^{\mathcal{I}}(\psi_*(g) - g_{Euc}) = O(r^{-\mu-|\mathcal{I}|}),$$

as $r \rightarrow \infty$, where Γ is a finite subgroup of $U(2)$ containing no complex reflections, B denotes a ball centered at the origin, and g_{Euc} denotes the Euclidean metric. The real number μ is called the order of g .

Remark 1.2. In this paper, henceforth Γ will always be a finite subgroup of $U(2)$ containing no complex reflections.

We are interested in the class of scalar-flat Kähler ALE metrics. These are interesting since they are *extremal* in the sense of Calabi [Cal85], and they arise as “bubbles” in gluing constructions for extremal Kähler metrics [ALM15, ALM16, AP06, APS11, BR15, RS05, RS09]. In the case of scalar-flat Kähler ALE metrics, it is known that there exists an ALE coordinate system for which the order of such a metric is at least 2 [LM08].

We note that for an ALE Kähler metric of order μ , there exist ALE coordinates for which

$$(1.2) \quad \partial^{\mathcal{I}}(J - J_{Euc}) = O(r^{-\mu-|\mathcal{I}|}),$$

for any multi-index \mathcal{I} as $r \rightarrow \infty$, where J_{Euc} is the standard complex structure on Euclidean space [HL16].

There are many known examples of scalar-flat Kähler ALE metrics. In the case that $\Gamma \subset SU(2)$, Kronheimer constructed and classified the hyperkähler ALE metrics [Kro89a, Kro89b]. Calderbank-Singer constructed a family of scalar-flat Kähler ALE metrics on the

The authors were partially supported by NSF Grant DMS-1405725. This paper was completed while the authors were in residence at Mathematical Sciences Research Institute in Berkeley, California. The authors would like to thank MSRI for their support, and for providing such an excellent working environment.

minimal resolution of any cyclic quotient singularity [CS04]. For the remaining subgroups of $U(2)$, the existence of scalar-flat Kähler metrics on the minimal resolution of \mathbb{C}^2/Γ , was shown by Lock-Viaclovsky [LV14].

The question we address in this paper is whether the scalar-flat Kähler property is preserved under small deformations of complex structure. In the cases where $\Gamma \subset SU(2)$, the hyperkähler quotient construction produces hyperkähler metrics for the minimal resolution complex structure as well as for all small deformations of the minimal resolution complex structure. In the case of the LeBrun negative mass metrics, using arguments from twistor theory, Honda has shown that all small deformations of the complex structure on $\mathcal{O}(-n)$ also admit scalar-flat Kähler metrics [Hon13, Hon14]. For the case of general Γ , in [LV14], employing Honda's result, it was shown that *some* of small deformations of complex structure admit scalar-flat Kähler ALE metrics.

Our main result in this paper shows that for *any* scalar-flat Kähler ALE surface, *all* small deformations of complex structure admit scalar-flat Kähler ALE metrics. Our proof is analytic in nature, and in particular, gives a new analytic proof of Honda's result on $\mathcal{O}(-n)$ mentioned above.

To state our result precisely, we next recall some basic facts regarding deformations of complex structures. For a complex manifold (X, J) , let $\Lambda^{p,q}$ denote the bundle of (p, q) -forms, and let Θ denote the holomorphic tangent bundle. The deformation complex corresponds to a real complex as shown in the commutative diagram

$$(1.3) \quad \begin{array}{ccccc} \Gamma(\Theta) & \xrightarrow{\bar{\partial}} & \Gamma(\Lambda^{0,1} \otimes \Theta) & \xrightarrow{\bar{\partial}} & \Gamma(\Lambda^{0,2} \otimes \Theta) \\ \downarrow Re & & \downarrow Re & & \downarrow Re \\ \Gamma(TX) & \xrightarrow{Z \mapsto -\frac{1}{2}J \circ \mathcal{L}_Z J} & \Gamma(End_a(TX)) & \xrightarrow{I \mapsto \frac{1}{4}J \circ N'_J(I)} & \Gamma(\{\Lambda^{0,2} \otimes \Theta \oplus \Lambda^{2,0} \otimes \bar{\Theta}\}_{\mathbb{R}}), \end{array}$$

where $\mathcal{L}_Z J$ is the Lie derivative of J ,

$$(1.4) \quad End_a(TX) = \{I \in End(TX) : IJ = -JI\},$$

and N'_J is the linearization of Nijenhuis tensor

$$(1.5) \quad N(X, Y) = 2\{[JX, JY] - [X, Y] - J[X, JY] - J[JX, Y]\}$$

at J . Each isomorphism Re is simply taking the real part of a section. If g is a Hermitian metric compatible with J , then let \square denote the $\bar{\partial}$ -Laplacian

$$(1.6) \quad \square \equiv \bar{\partial}^* \bar{\partial} + \bar{\partial} \bar{\partial}^*,$$

where $\bar{\partial}^*$ denotes the formal L^2 -adjoint. Each complex bundle in the diagram (1.3) admits a \square -Laplacian, and these correspond to real Laplacians on each real bundle in (1.3). We will use the same \square -notation for these real Laplacians.

To state our most general result, we need the following definition. This is necessary because there is a gauge freedom of Euclidean motions in the definition of ALE coordinates.

Definition 1.3. Let (X, g, J) be a Kähler ALE surface. For any bundle E in the diagram (1.3), and $\tau \in \mathbb{R}$, define

$$(1.7) \quad \mathcal{H}_\tau(X, E) = \{\theta \in \Gamma(X, E) : \square\theta = 0, \theta = O(r^\tau) \text{ as } r \rightarrow \infty\}.$$

Define

$$(1.8) \quad \mathbb{W} = \{Z \in \mathcal{H}_1(X, TX) \mid \mathfrak{L}_Z g = O(r^{-1}), \mathfrak{L}_Z J = O(r^{-3}), \text{ as } r \rightarrow \infty\}.$$

Finally, define the real subspace

$$(1.9) \quad \mathcal{H}_{ess}(X, End_a(TX)) \subset \mathcal{H}_{-3}(X, End_a(X))$$

(*ess* is short for essential) to be the L^2 -orthogonal complement in $\mathcal{H}_{-3}(X, End_a(X))$ of the subspace

$$(1.10) \quad \mathbb{V} = \{\theta \in \mathcal{H}_{-3}(X, End_a(TX)) \mid \theta = J \circ \mathfrak{L}_Z J, Z \in \mathbb{W}\}.$$

Our main result is the following.

Theorem 1.4. *Let (X, g, J) be a scalar-flat Kähler ALE surface. Let $-2 < \delta < -1$, $0 < \alpha < 1$, and k an integer with $k \geq 4$ be fixed constants. Let $B_{\epsilon_1}^1$ denote an ϵ_1 -ball in $\mathcal{H}_{ess}(X, End_a(TX))$, $B_{\epsilon_2}^2$ denote an ϵ_2 -ball in $\mathcal{H}_{-3}(X, \Lambda^{1,1})$ (both using the L^2 -norm). Then there exists $\epsilon_1 > 0$ and $\epsilon_2 > 0$ and a family \mathfrak{F} of scalar-flat Kähler metrics near g , parametrized by $B_{\epsilon_1}^1 \times B_{\epsilon_2}^2$, that is, there is a differentiable mapping*

$$(1.11) \quad F : B_{\epsilon_1}^1 \times B_{\epsilon_2}^2 \rightarrow Met(X),$$

into the space of smooth Riemannian metrics on X , with $\mathfrak{F} = F(B_{\epsilon_1}^1 \times B_{\epsilon_2}^2)$ satisfying the following “versal” property: there exists a constant $\epsilon_3 > 0$ such that for any scalar-flat Kähler metric $\tilde{g} \in B_{\epsilon_3}(g)$, there exists a diffeomorphism $\Phi : X \rightarrow X$, $\Phi \in C_{loc}^{k+1, \alpha}$, such that $\Phi^* \tilde{g} \in \mathfrak{F}$, where

$$(1.12) \quad B_{\epsilon_3}(g) = \{g' \in C_{loc}^{k, \alpha}(S^2(T^*X)) \mid \|g - g'\|_{C_{\delta}^{k, \alpha}(S^2(T^*X))} < \epsilon_3\}.$$

Remark 1.5. The norm in (1.12) is a certain weighted Hölder norm, see Section 2 for the precise definition. For a more precise description of the diffeomorphism Φ , see Theorem 9.2 below.

In order to state our next result, we must recall some facts about vector fields and diffeomorphisms. If (X, g) is an ALE metric, and Y is a vector field on X , the Riemannian exponential mapping $\exp_p : T_p X \rightarrow X$ induces a mapping

$$(1.13) \quad \Phi_Y : X \rightarrow X$$

by

$$(1.14) \quad \Phi_Y(p) = \exp_p(Y).$$

If $Y \in C_s^{k, \alpha}(TX)$ has sufficiently small norm, ($s < 0$ and k will be determined in specific cases) then Φ_Y is a diffeomorphism. We will use the correspondence $Y \mapsto \Phi_Y$ to parametrize a neighborhood of the identity, analogous to [Biq06].

Definition 1.6. We say that $\Phi : X \rightarrow X$ is a *small diffeomorphism* if Φ is of the form $\Phi = \Phi_Y$ for some vector field Y satisfying

$$(1.15) \quad \|Y\|_{C_{\delta+1}^{k+1, \alpha}} < \epsilon_4$$

for some $\epsilon_4 > 0$ sufficiently small which depends on ϵ_3 .

The family \mathfrak{F} is not necessarily “universal”, because some elements in \mathfrak{F} might be isometric. However, the following theorem shows that after taking a quotient by an action of the holomorphic isometries of the central fiber (X, g, J) , the family \mathfrak{F} is in fact universal (up to small diffeomorphisms).

Theorem 1.7. *Let (X, g, J) be as in Theorem 1.4, and let \mathfrak{G} denote the group of holomorphic isometries of (X, g, J) . Then there is an action of \mathfrak{G} on \mathfrak{F} with the following properties.*

- *Two metrics in \mathfrak{F} are isometric if they are in the same orbit of \mathfrak{G} .*
- *If two metrics in \mathfrak{F} are isometric by a small diffeomorphism then they must be the same.*

Since each orbit represents a unique isometry class of metric (up to small diffeomorphism), we will refer to the quotient $\mathfrak{M} = \mathfrak{F}/\mathfrak{G}$ as the “local moduli space of scalar-flat Kähler ALE metrics near g .” The local moduli space \mathfrak{M} is not a manifold in general, but its dimension is in fact well-defined, and we define

$$(1.16) \quad m = \dim(\mathfrak{M}).$$

Remark 1.8. We should point out that our local moduli space of metrics contains small rescalings, i.e, $g \mapsto \frac{1}{c}g(c \cdot, c \cdot)$ for c close to 1. If one considers scaled metrics as equivalent (which we do not), then the dimension would decrease by 1.

1.1. Deformations of the minimal resolution. As mentioned above, there are families of examples of scalar-flat Kähler ALE metrics on minimal resolutions of isolated quotient singularities. For convenience, we next recall the definition of a minimal resolution.

Definition 1.9. Let $\Gamma \subset \mathrm{U}(2)$ be as above. Then, a smooth complex surface X is called a *minimal resolution* of \mathbb{C}^2/Γ if there is a mapping $\pi : X \rightarrow \mathbb{C}^2/\Gamma$ such that the restriction $\pi : X \setminus \pi^{-1}(0) \rightarrow \mathbb{C}^2/\Gamma \setminus \{0\}$ is a biholomorphism, and $\pi^{-1}(0)$ is a divisor in X containing no -1 curves. The divisor $\pi^{-1}(0)$ is called the *exceptional divisor*.

In the cyclic case, the exceptional divisor is a string of rational curves with normal crossing singularities. In the case that Γ is non-cyclic, it was shown by Brieskorn, see [Bri68], that the exceptional divisor is a tree of rational curves with normal crossing singularities. There are three Hirzebruch-Jung strings attached to a single curve, called the *central rational curve*. The self-intersection number of this curve will be denoted $-b_\Gamma$, and the total number of rational curves will be denoted by k_Γ .

A specialized version of Theorem 1.4 is the following.

Theorem 1.10. *Let (X, g, J) be any scalar-flat Kähler ALE metric on the minimal resolution of \mathbb{C}^2/Γ , where $\Gamma \subset \mathrm{U}(2)$ is as above. Define*

$$(1.17) \quad j_\Gamma = 2 \sum_{i=1}^{k_\Gamma} (e_i - 1),$$

where $-e_i$ is the self-intersection number of the i th rational curve, and k_Γ is the number of rational curves in the exceptional divisor, and let

$$(1.18) \quad d_\Gamma = j_\Gamma + k_\Gamma.$$

Then there is a family, \mathfrak{F} , parametrized by a ball in \mathbb{R}^{d_Γ} , of scalar-flat Kähler metrics near g with the “versal” property stated in Theorem 1.4.

We let m_Γ denote the dimension of the moduli space near the minimal resolution of \mathbb{C}^2/Γ , where the dimension is defined in (1.16). Note $m_\Gamma < d_\Gamma$ due to the action of the automorphism group, and m_Γ can be explicitly computed for all groups Γ , but those computations will not be done here.

Remark 1.11. The dimension of the moduli space of hyperkähler metrics is known to be $3k - 3$ in the A_k, D_k and E_k cases for $k \geq 2$, and equal to 1 in the A_1 case [Kro86]. Our method of parametrizing by complex structures and Kähler classes overcounts in this case (i.e., F is not injective), since a hyperkähler metric is Kähler with respect to a 2-sphere of complex structures.

Our final result applies to a generic deformation of the minimal resolution.

Theorem 1.12. *Let (X, g, J) be any scalar-flat Kähler ALE surface which deforms to the minimal resolution of \mathbb{C}^2/Γ through a path (X, g_t, J_t) ($0 \leq t \leq 1$), where $g_1 = g$, g_0 is the minimal resolution, and $\|g_t - g_s\|_{C_\delta^{k,\alpha}(g_0)} \leq C \cdot |s - t|$ with $C > 0$ a uniform constant for any $0 \leq s, t \leq 1$, $k \geq 4$, $-2 < \delta < -1$. If $\mathfrak{G}(g) = \{e\}$ then the local moduli space \mathfrak{F} is smooth near g and is a manifold of dimension $m = m_\Gamma$.*

Remark 1.13. It was recently shown that Kähler ALE surface with group $\Gamma \subset \mathrm{U}(2)$ is birational to a deformation of \mathbb{C}^2/Γ [HRŠ16]. There are several possible components of the deformation of such a cone, here we consider surfaces in the “Artin component” of deformations of \mathbb{C}^2/Γ . We note however that there are some known examples of scalar-flat Kähler metrics on non-Artin components, which are free quotients of hyperkähler metrics of A_k -type, see for example [Šuv12].

We end the introduction with an outline of the paper. In Section 2, we begin with the definitions of the weighted Hölder spaces which will be used throughout the paper. Then, in Section 3 we give some analysis of the complex analytic compactifications, due to Hein-LeBrun-Maskit [HL16, LM08], of Kähler ALE spaces. In Section 4, we study the deformation of complex structures using an adaptation of Kuranishi’s theory to ALE spaces. The main point is that since the manifold is non-compact, the sheaf cohomology group $H^1(X, \Theta)$ should be replaced by an appropriate space of decaying harmonic forms. In Section 5, several key results about gauging and diffeomorphisms are proved, which are used to prove “versality” of the family constructed in Section 4. In Section 6, a refined gauging procedure is carried out, to construct the Kuranishi family of “essential” deformations. In Section 7, we generalize Kodaira-Spencer’s stability theorem for Kähler structures to the ALE setting, using some arguments of Biquard-Rollin [BR15]. In Section 8, we adapt the LeBrun-Singer-Simanca theory of deformations of extremal Kähler metrics to the ALE setting [LS93, LS94]. In Section 9, we prove the versal property of the family \mathfrak{F} , using a local slicing theorem, and prove Theorem 1.7. In Section 10, we restrict attention to the minimal resolution, and prove Theorems 1.10 and 1.12.

1.2. Acknowledgements. The authors would like to thank Olivier Biquard, Ronan Conlon, Akira Fujiki, Ryushi Goto, Hans-Joachim Hein, Nobuhiro Honda, and Claude LeBrun, for numerous helpful discussions on deformations of complex structures. Olivier Biquard, Joel Fine, and Jason Lotay provided crucial assistance with the slicing arguments. We would also like to thank Michael Lock for valuable discussions on properties of subgroups of $\mathrm{U}(2)$.

Finally, the authors would like to thank to the anonymous referee for detailed helpful remarks, and for suggesting numerous improvements to the paper.

2. NOTATION

We begin with the definition of weighted Hölder spaces, which will be used throughout the paper.

Definition 2.1. Let E be a tensor bundle on X , with Hermitian metric $\|\cdot\|_h$. Let φ be a smooth section of E . We fix a point $p_0 \in X$, and define $r(p)$ to be the distance between p_0 and p . Then define

$$(2.1) \quad \|\varphi\|_{C_\delta^0} := \sup_{p \in X} \left\{ \|\varphi(p)\|_h \cdot (1 + r(p))^{-\delta} \right\}$$

$$(2.2) \quad \|\varphi\|_{C_\delta^k} := \sum_{|\mathcal{I}| \leq k} \sup_{p \in X} \left\{ \|\nabla^{\mathcal{I}} \varphi(p)\|_h \cdot (1 + r(p))^{-\delta + |\mathcal{I}|} \right\},$$

where $\mathcal{I} = (i_1, \dots, i_n)$, $|\mathcal{I}| = \sum_{j=1}^n i_j$. Next, define

$$(2.3) \quad [\varphi]_{C_{\delta-\alpha}^\alpha} := \sup_{0 < d(x,y) < \rho_{inj}} \left\{ \min\{r(x), r(y)\}^{-\delta+\alpha} \frac{\|\varphi(x) - \varphi(y)\|_h}{d(x,y)^\alpha} \right\},$$

where $0 < \alpha < 1$, ρ_{inj} is the injectivity radius, and $d(x, y)$ is the distance between x and y . The meaning of the tensor norm is to use parallel transport along the unique minimal geodesic from y to x , and then take the norm of the difference at x . The weighted Hölder norm is defined by

$$(2.4) \quad \|\varphi\|_{C_\delta^{k,\alpha}} := \|\varphi\|_{C_\delta^k} + \sum_{|\mathcal{I}|=k} [\nabla^{\mathcal{I}} \varphi]_{C_{\delta-k-\alpha}^\alpha},$$

and the space $C_\delta^{k,\alpha}(X, E)$ is the closure of $\{\varphi \in C^\infty(X, E) : \|\varphi\|_{C_\delta^{k,\alpha}} < \infty\}$.

Remark 2.2. The dual space of Hölder space is not a Hölder space, but for the purpose of computing the dimension of cokernel of a Fredholm operator of order o ,

$$(2.5) \quad H : C_\delta^{k,\alpha} \rightarrow C_{\delta-o}^{k-o,\alpha},$$

we consider the adjoint operator as mapping between Hölder spaces

$$(2.6) \quad H^* : C_{-4-\delta+o}^{k,\alpha} \rightarrow C_{-4-\delta}^{k-o,\alpha},$$

since the adjoint weight to weight δ is $-4 - \delta$, and using elliptic regularity.

3. ALE KÄHLER SURFACES

In this section, we will prove several results about ALE Kähler surfaces which will be needed later. We note that the results in this section do not use the scalar-flat assumption.

In [LM08, HL16], LeBrun-Maskit and Hein-LeBrun analyzed the asymptotic behavior of the metric and complex structure near infinity of ALE Kähler surfaces. The next proposition gives a summary of their results. (See also [Li14] for other related results on complex analytic compactifications).

Proposition 3.1 (Hein-LeBrun-Maskit). *Let X_∞ be an end of an ALE Kähler surface (X, g, J) with metric g asymptotic to Euclidean metric at rate*

$$(3.1) \quad |\nabla_{Euc}^{\mathcal{I}}(g_{j,k} - \delta_{j,k})| = O(\rho^{-|\mathcal{I}|-1-\epsilon})$$

for $|\mathcal{I}| = 0, 1$ if $\epsilon > 1/2$ or for $|\mathcal{I}| = 0, \dots, 4$ if $\epsilon \in (0, 1/2]$; here ∇_{Euc} denotes the Euclidean derivative. Then there is a surface S containing an embedded holomorphic curve $C \simeq \mathbb{CP}^1$ with self-intersection 1, such that the universal cover \tilde{X}_∞ of X_∞ is biholomorphic to $S \setminus C$.

Let (X, g, J) be a Kähler ALE surface, then X has one end and can be analytically compactified to a smooth compact surface \hat{X} by adding a tree of rational curves E_∞ . The surface \hat{X} is a rational surface or a ruled surface. Also, X can be compactified to an orbifold surface \hat{X}_{orb} by adding a single rational curve at ∞ .

Furthermore, if the order of g satisfies $1 < \mu < 3$, then there exists an ALE coordinate, under which $|J - J_{Euc}| = O(r^{-3})$, $|g - g_{Euc}| = O(r^{-\mu})$.

Remark 3.2. The orbifold compactification \hat{X}_{orb} has cyclic singularities on the rational curve at ∞ . The smooth compactification \hat{X} is obtained from \hat{X}_{orb} by resolving these singularities.

Consider the space of holomorphic vector fields on X with at most growth of order τ

$$(3.2) \quad \mathcal{Z}_\tau(X, \Theta) = \{Z \in \Gamma(X, \Theta) \mid \bar{\partial}Z = 0, Z = O(r^\tau) \text{ as } r \rightarrow \infty\}.$$

The first vanishing result we need is the following.

Proposition 3.3. *Let (X, g, J) be an ALE Kähler surface. If $\tau < 0$ then*

$$(3.3) \quad \dim \mathcal{Z}_\tau(X, \Theta) = 0.$$

Proof. Let $X_\infty = X \setminus \overline{B(R)}$. Let R be sufficiently large, then by Proposition 3.1, the universal cover \tilde{X}_∞ can be analytically compactified by adding a rational curve C at infinity, where C has self-intersection +1. By [HL16], there exists a smooth map $[\xi_1, \xi_2, f]$ that maps $S = \tilde{X}_\infty \cup C$ to \mathbb{CP}^2 , which maps C holomorphically to $\{f = 0\}$ as a curve $\mathbb{CP}^1 \subset \mathbb{CP}^2$, where

$$(3.4) \quad \xi_j = \{\text{holomorphic part}\} + O(|f|^3),$$

as $f \rightarrow 0$. Let

$$(3.5) \quad z_j = x_j + \sqrt{-1}y_j = \frac{\xi_j}{f}, (j = 1, 2)$$

where $\{x_1, y_1, x_2, y_2\}$ gives an ALE coordinate on \tilde{X}_∞ , and $|J - J_{Euc}| = O(|z|^{-3})$ as $z \rightarrow \infty$. Let

$$(3.6) \quad (v, w) = \left(\frac{1}{z_1}, \frac{z_2}{z_1} \right)$$

where $\{v = 0\}$ represents the complement of one point in C .

First, let σ be a decaying holomorphic vector field on X , which can be lifted to $\tilde{\sigma}$ on \tilde{X}_∞ . Note that $\tilde{\sigma}$ can be extended to \mathbb{C}^2 smoothly by using a cut-off function. Since $\bar{\partial} - \bar{\partial}_{Euc} = O(|z|^{-3})$ and $\bar{\partial}\tilde{\sigma} = 0$ on \tilde{X}_∞ ,

$$(3.7) \quad \bar{\partial}_{Euc}\tilde{\sigma} = \sum_{i,j} h_{i,j} d\bar{z}_i \otimes \frac{\partial}{\partial z_j} = O(|z|^{-3}|\sigma|),$$

as $z \rightarrow \infty$. By using the $\bar{\partial}_{Euc}$ -Poincaré lemma, there exist

$$(3.8) \quad p_j = O(|z|^{-2}\sigma) \in C^\infty(\mathbb{C}^2), \quad \bar{\partial}_{Euc} p_j = \sum_i h_{i,j} d\bar{z}_i.$$

The formula for p_j can be written out explicitly as follows. Let

$$(3.9) \quad \begin{aligned} q_j &= \frac{1}{2\pi\sqrt{-1}} \int_{\mathbb{C}} h_{2,j}(z_1, \zeta_2) \frac{d\zeta_2 \wedge d\bar{\zeta}_2}{\zeta_2 - z_2} \\ q'_j &= \frac{1}{2\pi\sqrt{-1}} \int_{\mathbb{C}} (h_{1,j} - \bar{\partial}_1 q_j)(\zeta_1, z_2) \frac{d\zeta_1 \wedge d\bar{\zeta}_1}{\zeta_1 - z_1}, \end{aligned}$$

where the integral formula is valid since $h_{i,j} = o(|z|^{-3})$. Then $p_j = q_j + q'_j$. Then we have $\tilde{\sigma} - \sum_j p_j \frac{\partial}{\partial z_j}$ is a decaying $\bar{\partial}_{Euc}$ -holomorphic vector field on \mathbb{C}^2 , which must vanish identically. The growth rate of σ can be dropped by 2 iteratively by this argument, so we can assume that σ decays at any rate in the coordinates $\{z_1, z_2\}$.

On \tilde{X}_∞ , there exists a holomorphic $(2, 0)$ -form $\Omega = dz_1 \wedge dz_2 + O(|z|^{-2+\epsilon})$ as $|z| \rightarrow \infty$. Let $\tilde{\sigma}^* \in \Gamma(\tilde{X}_\infty, \Omega^1)$ denote the contraction of $\tilde{\sigma}$ with Ω . The section $\tilde{\sigma}^*$ is also holomorphic and we have $\tilde{\sigma}^* = O(|z|^{-10})$ as $z \rightarrow \infty$ since we can assume $\tilde{\sigma}$ decays to any order. The coordinate change gives: $dz_1 = \frac{1}{v^2} dv$, $dz_2 = \frac{1}{v} dw - \frac{w}{v^2} dv$, $|z|^2 = \frac{1}{|v|^2}(1 + |w|^2)$. Then $\tilde{\sigma}^* = O(|v|^8)$ near C and can be extended to C with $\tilde{\sigma}^* = 0$ on C . Consider the exact sequence

$$(3.10) \quad 0 \rightarrow N_C^* \rightarrow \Omega^1|_C \rightarrow \Omega^1(C) \rightarrow 0,$$

where $\Omega^1|_C$ is the restriction of the bundle Ω^1 on C , $\Omega^1(C)$ is the cotangent bundle of C and N_C is the normal bundle of C in S . Then we have $\Omega^1|_C = \mathcal{O}(-2) \oplus \mathcal{O}(-1)$. This implies $\tilde{\sigma}^*$ vanishes on any rational curve with self-intersection $+1$ on S . Since C has self-intersection $+1$, $H^1(C, N_C^* \otimes \Theta(C)) = 0$, so C is rigid in S . Then there is an open neighborhood $U \subset S$ of C , such that $\tilde{\sigma}^*$ vanishes identically over U . Then σ vanishes on an open subset of X , which implies that σ vanishes identically on X . \square

Next, we consider harmonic $(0, 2)$ -forms with values in the holomorphic tangent bundle.

Proposition 3.4. *Let X be a Kähler ALE surface. If $\delta < 0$, then $\mathcal{H}_\delta(X, \Lambda^{0,2} \otimes \Theta) = 0$.*

Proof. The proposition follows an argument in [LM08, Theorem 4.2], with minor modifications. For completeness, we give a proof here following their idea. Let $\sigma \in \mathcal{H}_\delta(X, \Lambda^{0,2} \otimes \Theta)$. Recall that the conjugated Hodge star operator $\bar{*}$ maps σ to a \square -harmonic form $\sigma^* \in \mathcal{H}_\delta(X, \Lambda^{2,0} \otimes \Omega^1)$. Let $\tilde{\sigma}^*$ be its lifting on \tilde{X}_∞ . Following the same notion as in the proof of Proposition 3.3, we have

$$(3.11) \quad \tilde{\sigma}^* = \sum_j a_j dz_1 \wedge dz_2 \otimes dz_j + O(|z|^{-3}|\tilde{\sigma}^*|),$$

where a_j are $\bar{\partial}_{Euc}$ -holomorphic functions and $a_j = O(|z|^\delta)$ as $z \rightarrow \infty$. As shown in the proof of Proposition 3.3, $a_j = 0$, so $\tilde{\sigma}^* = O(|z|^{\delta-3})$ as $z \rightarrow \infty$. Since the growth rate of $\tilde{\sigma}^*$ can be dropped by 3 iteratively, $\tilde{\sigma}^*$ can decay to any order under the coordinate of $\{z_1, z_2\}$, then $\tilde{\sigma}^*$ can be extended to C with $\tilde{\sigma}^* = 0$ on C . Since

$$(3.12) \quad \Omega^{2,0} \otimes \Omega^1|_C = (\Omega^1(C) \oplus N_C^*) \otimes (\Omega^{2,0}(C) \otimes N_C^*),$$

we have

$$(3.13) \quad \Omega^{2,0} \otimes \Omega^1|_C = (\mathcal{O}(-2) \oplus \mathcal{O}(-1)) \otimes (\mathcal{O}(-2) \otimes \mathcal{O}(-1)) = \mathcal{O}(-5) \oplus \mathcal{O}(-4).$$

This implies that $\tilde{\sigma}^*$ vanishes on any rational curve with self-intersection +1 on \tilde{X}_∞ . By the same argument as in Proposition 3.3, there is an open neighborhood $U \subset S$ of C , such that $\tilde{\sigma}^*$ vanishes identically over U . Then σ vanishes on an open subset of X , which implies that σ vanishes identically on X . \square

The next proposition shows that $\mathcal{H}_\delta(X, \Lambda^{0,1} \otimes \Theta)$ automatically has a faster decaying rate.

Proposition 3.5. *Let X be an ALE Kähler surface, and \hat{X} be its analytic compactification. Then for any $-3 < \tau < 0$ we have*

$$(3.14) \quad \mathcal{H}_\tau(X, \Lambda^{0,1} \otimes \Theta) = \mathcal{H}_{-3}(X, \Lambda^{0,1} \otimes \Theta),$$

and if Γ is not cyclic or $\Gamma = \mathbb{Z}/2\mathbb{Z}$, then

$$(3.15) \quad \mathcal{H}_\tau(X, \Lambda^{0,1} \otimes \Theta) = \mathcal{H}_{-4}(X, \Lambda^{0,1} \otimes \Theta).$$

Proof. First, we will show that any decaying harmonic element $\phi \in H^0(X, \Lambda^{0,1} \otimes \Theta)$ has a decay rate of at least $O(r^{-3})$ at infinity. Since X is Kähler, by [Mor07, Part 5], the operator \square admits an expansion near infinity of the form

$$(3.16) \quad \square = \frac{1}{2} \nabla^* \nabla + \mathfrak{R}$$

on $\Lambda^{0,1} \otimes \Theta$, where ∇ is the covariant derivative on $\Lambda^{0,1} \otimes \Theta$, and \mathfrak{R} is an operator given by curvature forms acting on the same bundle. The leading term of $\nabla^* \nabla$ is the Euclidean Laplacian Δ_{Euc} , so we have

$$(3.17) \quad \square = \frac{-1}{2} \Delta_{Euc} + Q,$$

where $Q = A(\nabla\phi) + Ric(\phi)$ denotes higher order terms. The element ϕ admits an expansion of the form

$$(3.18) \quad \phi = f + O(r^{-3+\epsilon}),$$

as $r \rightarrow \infty$, where f is of the form

$$(3.19) \quad f = \sum_{i,j} \frac{f_{i,j}}{r^2} d\bar{z}_i \otimes \frac{\partial}{\partial z_j},$$

and $f_{i,j}$ are constants. Since ϕ is \square -harmonic, both $\bar{\partial}f = 0$ and $\bar{\partial}^*f = 0$. It is easy to see that this implies that each $f_{i,j} = 0$. Consequently, $\phi = O(r^{-3+\epsilon})$ as $r \rightarrow \infty$. Since ϕ admits an expansion with harmonic leading term, we must have $\phi = O(r^{-3})$.

Furthermore, if Γ is not cyclic or $\Gamma = \mathbb{Z}/2\mathbb{Z}$, then Γ contains the element -1 . The leading term of ϕ is of the form

$$(3.20) \quad f = \sum_{k,l} \frac{A_{kl}}{r^4} d\bar{z}_k \otimes \frac{\partial}{\partial z_l},$$

where A_{kl} is a linear combination of $z_1, \bar{z}_1, z_2, \bar{z}_2$. Since any such nonzero f is not invariant under the action of -1 , we must have $f = 0$, and then $\phi = O(r^{-4})$. This finishes the proof of (3.14) and (3.15). \square

The following Proposition will be used in Section 7.

Proposition 3.6. *Let (X, g, J) be an ALE Kähler surface. Then*

$$(3.21) \quad b_1(X) = 0,$$

where $b_1(X)$ denotes the first Betti number of X . Furthermore, for $\tau < 0$,

$$(3.22) \quad \dim(\mathcal{H}_\tau(X, \Lambda^{0,2})) = \dim(\mathcal{H}_\tau(X, \Lambda^{2,0})) = 0.$$

Proof. Let U be a tubular neighborhood of E_∞ , $\hat{X} = X \cup E_\infty$ be the analytic compactification of X . By Proposition 3.1, \hat{X} is either a rational surface or a ruled surface, then $b^1(\hat{X}) = 0$. Since E_∞ is a tree of rational curves, $b^1(U) = 0$ and $b^1(X \cap U) = 0$. Then by the Mayer-Vietoris theorem, $b^1(X) = 0$.

Since $\dim(\mathcal{H}_\tau(X, \Lambda^{0,2})) = \dim(\mathcal{H}_\tau(X, \Lambda^{2,0}))$, we just need to show the latter one equals to zero. The proof is similar to the proof of Proposition 3.4, so we will skip some details. For any $\sigma \in \mathcal{H}_\tau(X, \Lambda^{2,0})$, let $\tilde{\sigma}$ be its lifting on \tilde{X}_∞ . Then

$$(3.23) \quad \tilde{\sigma} = a \cdot dz_1 \wedge dz_2 + O(|z|^{-3}\tilde{\sigma})$$

where $a = O(|z|^\tau)$ as $r \rightarrow \infty$ is a $\bar{\partial}_{Euc}$ -holomorphic function. Then $a = 0$, and $\tilde{\sigma} = O(|z|^{-3+\tau})$, which implies that $\tilde{\sigma}$ can decay to any order in the ALE coordinates $\{z_1, z_2\}$. Consequently, $\tilde{\sigma}$ can be extended to C . Furthermore,

$$(3.24) \quad \Omega^{2,0}|_C = \Omega^{2,0}(C) \otimes N_C^* = \mathcal{O}(-3).$$

This implies that $\tilde{\sigma}$ vanishes on an open neighborhood of C , so σ vanishes identically on X . \square

4. SMALL DEFORMATIONS OF COMPLEX STRUCTURE

First we need a fixed point theorem for operators on Banach spaces (see, for example [BR15]).

Lemma 4.1. *Let $F : \mathcal{B}_1 \rightarrow \mathcal{B}_2$ be a bounded differentiable operator between Banach spaces. In a small neighborhood of $0 \in \mathcal{B}_1$, $F(x) = F(0) + F'(0)x + Q(x)$, where*

$$(4.1) \quad \|Q(x) - Q(x')\|_{\mathcal{B}_2} \leq C_0 \cdot (\|x\|_{\mathcal{B}_1} + \|x'\|_{\mathcal{B}_1}) \cdot \|x - x'\|_{\mathcal{B}_1}.$$

Assume that $\|F(0)\|_{\mathcal{B}_2} \ll 1$. Then

- (i) *If $F'(0)$ is an isomorphism with right inverse G bounded by C_1 , then there exists a ball $B(0, s) \subset \mathcal{B}_1$ such that there exists a unique $x \in B(0, s)$ with $F(x) = 0$.*
- (ii) *If $F'(0)$ is Fredholm and surjective, with right inverse bounded by C_1 , then there is an $s > 0$, so that $F^{-1}(0) \cap B(0, s)$ is isomorphic to $U \subset \ker(F'(0))$, where U is a small neighborhood of the origin.*

The following lemma gives the properties of the linearized operator we will need in order to invoke Lemma 4.1.

Lemma 4.2. *Let (X, g, J) be a Kähler ALE surface. The linear operator*

$$(4.2) \quad P : C_{\delta-1}^{k,\alpha}(X, \Lambda^{0,1} \otimes \Theta) \xrightarrow{(\bar{\partial}^*, \bar{\partial})} C_{\delta-2}^{k-1,\alpha}(X, \Theta) \oplus C_{\delta-2}^{k-1,\alpha}(X, \Lambda^{0,2} \otimes \Theta)$$

is surjective and Fredholm, for some $k \geq 3$ and $\delta \in (-2, -1)$.

Proof. It is not hard to see that P is an elliptic operator, and that the indicial roots of P are integral. Consequently, by standard weighted space theory, P is Fredholm since δ is non-integral [LM85].

The cokernel is given by

$$(4.3) \quad \ker(P^*) = \{(\sigma_1, \sigma_2) \in C_{-2-\delta}^{k-1,\alpha}(\Theta) \oplus C_{-2-\delta}^{k-1,\alpha}(\Lambda^{0,2} \otimes \Theta) : \bar{\partial}^* \sigma_2 = \bar{\partial} \sigma_1\}.$$

Let $(\sigma_1, \sigma_2) \in \ker(P^*)$, then

$$(4.4) \quad \bar{\partial}^* \sigma_1 = 0, \quad \bar{\partial}^* \bar{\partial} \sigma_1 = \bar{\partial}^* \bar{\partial} \sigma_2 = 0,$$

which implies that σ_1 is \square -harmonic. From Proposition 3.3, $\sigma_1 = 0$. Then

$$(4.5) \quad \bar{\partial} \sigma_2 = 0, \quad \bar{\partial}^* \sigma_2 = 0,$$

and the vanishing of σ_2 follows from Proposition 3.4. Since $\ker(P^*) = 0$, and P is Fredholm, P is surjective. \square

For a fixed complex structure J , there is a nonlinear correspondence between sufficiently small sections of $\Gamma(X, \text{End}_a(TX))$ and almost complex structures near J given by

$$(4.6) \quad \begin{aligned} E_J : \Gamma(X, \text{End}_a(TX)) &\rightarrow \mathcal{A} \\ E_J(I) &= \left(J + \frac{1}{2}I\right) J \left(J + \frac{1}{2}I\right)^{-1}, \end{aligned}$$

where \mathcal{A} is the space of almost complex structures on X . If $\phi \in \Gamma(\Lambda^{0,1} \otimes \Theta)$ has sufficiently small norm, then the corresponding almost complex structure will be denoted by $J(\phi) = E_J(\text{Re}(\phi))$. Note that there is an expansion:

$$(4.7) \quad J(\phi) = J + \text{Re}(\phi) + Q$$

where $|Q| = O(|\phi|^2)$, as $|\phi| \rightarrow 0$.

Next we prove a weighted version of Kuranishi's theorem.

Theorem 4.3. *Let (X, J_0, g_0) be a Kähler ALE surface, and let B_{ϵ_1} be an ϵ_1 -ball around 0 in $\mathcal{H}_{-3}(X, \Lambda^{0,1} \otimes \Theta)$. Then for $\epsilon_1 > 0$ sufficiently small, there is a differentiable family of complex structures J_t for $t \in B_{\epsilon_1}$ such that*

$$(4.8) \quad J_t = J(\phi(t)), \quad \phi(t) \in C_{\delta-1}^{k,\alpha}(X, \Lambda^{0,1} \otimes \Theta), \quad \delta \in (-2, -1), \quad k \geq 3,$$

where $\phi(t)$ satisfies $\phi(t) = t + \phi(t)^\perp$, with $\phi(t)^\perp$ is L^2 -orthogonal to $\mathcal{H}_{-3}(X, \Lambda^{0,1} \otimes \Theta)$, and $\bar{\partial}^(\phi(t)) = 0$. Furthermore, there exists an $\epsilon' > 0$, such that if $J = J_0(\phi)$ is any integrable complex structure satisfying*

$$(4.9) \quad \|\phi\|_{C_\delta^{k,\alpha}} < \epsilon', \quad \text{and} \quad \bar{\partial}^*(\phi) = 0,$$

then ϕ is in the family J_t . Finally, for any $t \in B_{\epsilon_1}$, there exists a constant C such that $\|\phi(t)\|_{C_{-3}^{k,\alpha}} \leq C \cdot \epsilon_1$.

Proof. Define the operator

$$(4.10) \quad \begin{aligned} F : C_{\delta-1}^{k,\alpha}(X, \Lambda^{0,1} \otimes \Theta) &\rightarrow C_{\delta-2}^{k-1,\alpha}(X, \Theta) \oplus C_{\delta-2}^{k-1,\alpha}(X, \Lambda^{0,2} \otimes \Theta) \\ \phi &\mapsto (\bar{\partial}^* \phi, \bar{\partial} \phi + [\phi, \phi]), \end{aligned}$$

where $[\phi, \phi]$ is a globally defined operator, which can be expressed locally as

$$(4.11) \quad [\phi, \phi] = \frac{1}{2} \left[\sum_{i,j} \phi_{i,j} d\bar{z}_i \otimes \frac{\partial}{\partial z_j}, \sum_{k,l} \phi_{k,l} d\bar{z}_k \otimes \frac{\partial}{\partial z_l} \right]$$

$$(4.12) \quad = \sum_{i,j,k,l} \phi_{i,j} \frac{\partial \phi_{k,l}}{\partial z_j} d\bar{z}_i \wedge d\bar{z}_k \otimes \frac{\partial}{\partial z_l}.$$

Formally, $[\phi, \phi]$ can be written as $\phi * \nabla \phi$, where $*$ means a linear combination of quadratic terms, each involving some contraction of ϕ with $\nabla \phi$.

Clearly, F is a bounded differentiable mapping, and $F'(0) = (\bar{\partial}^*, \bar{\partial})$. Obviously, in a small neighborhood of 0 in $C_{\delta-1}^{k,\alpha}(X, \Lambda^{0,1} \otimes \Theta)$, F admits an expansion $F = F(0) + F'(0) + Q$. We next prove the estimate (4.1). Let r denote the radius. For any two elements $\phi, \phi' \in C_{\delta-1}^{k,\alpha}(\Lambda^{0,1} \otimes \Theta)$, we have

$$(4.13) \quad \begin{aligned} r^{2-\delta} |[\phi, \phi] - [\phi', \phi']| &= r^{2-\delta} |[\phi * \nabla \phi] - [\phi' * \nabla \phi']| \\ &= r^{2-\delta} |\phi * (\nabla \phi - \nabla \phi') - (\phi - \phi') * \nabla \phi'| \\ &\leq Cr^{\delta-1} \{ r^{1-\delta} |\phi| r^{2-\delta} |\nabla \phi - \nabla \phi'| + r^{1-\delta} |\phi - \phi'| r^{2-\delta} |\nabla \phi'| \} \\ &\leq C(r^{1-\delta} |\phi|)(r^{2-\delta} |\nabla(\phi - \phi')|) + (r^{2-\delta} |\nabla \phi'|)(r^{1-\delta} |\phi - \phi'|). \end{aligned}$$

Next, let $x \neq y \in X$, with $r(x) < r(y)$. Similarly, we have

$$(4.14) \quad \begin{aligned} &r(x)^{2-\delta} \frac{|([\phi, \phi] - [\phi', \phi'])(x) - ([\phi, \phi] - [\phi', \phi'])(y)|}{d(x, y)^\alpha} \\ &\leq r(x)^{\delta-1} \left\{ r(x)^{1-\delta} \frac{|\phi(x) - \phi(y)|}{d(x, y)^\alpha} |\nabla \phi(x) - \nabla \phi'(x)| r(x)^{2-\delta} \right\} + \\ &r(y)^{\delta-1} \left\{ r(x)^{2-\delta} \frac{|(\nabla \phi(x) - \nabla \phi'(x)) - (\nabla \phi(y) - \nabla \phi'(y))|}{d(x, y)^\alpha} |\phi(y)| r(y)^{1-\delta} \right\} + \\ &r(x)^{\delta-1} \left\{ r(x)^{1-\delta} \frac{|(\phi(x) - \phi'(x)) - (\phi(y) - \phi'(y))|}{d(x, y)^\alpha} |\nabla \phi'(x)| r(x)^{2-\delta} \right\} + \\ &r(y)^{\delta-1} \left\{ r(x)^{2-\delta} \frac{|\nabla \phi'(x) - \nabla \phi'(y)|}{d(x, y)^\alpha} |\phi(y) - \phi'(y)| r(y)^{1-\delta} \right\}. \end{aligned}$$

This shows that

$$(4.15) \quad \|[\phi, \phi] - [\phi', \phi']\|_{C_{\delta-2}^{0,\alpha}} \leq C_0 (\|\phi\|_{C_{\delta-1}^{1,\alpha}} + \|\phi'\|_{C_{\delta-1}^{1,\alpha}}) \|\phi - \phi'\|_{C_{\delta-1}^{1,\alpha}}.$$

The higher derivative terms can be handled similarly, to prove

$$(4.16) \quad \|[\phi, \phi] - [\phi', \phi']\|_{C_{\delta-2}^{j-1,\alpha}} \leq C_j (\|\phi\|_{C_{\delta-1}^{j,\alpha}} + \|\phi'\|_{C_{\delta-1}^{j,\alpha}}) \|\phi - \phi'\|_{C_{\delta-1}^{j,\alpha}}, \quad 1 \leq j \leq k$$

which shows that Q satisfies (4.1).

By Lemma 4.2, we know $P = F'(0)$ is Fredholm. We can choose a right inverse operator G such that the image of G is L^2 -orthogonal to $\mathcal{H}_{-3}(X, \Lambda^{0,1} \otimes \Theta)$. By Lemma 4.1 there

exists $s = \epsilon_1 > 0$, such that $F^{-1}(0)$ is locally isomorphic to an ϵ_1 -neighborhood of 0 in $\mathcal{H}_{\delta-1}(X, \Lambda^{0,1} \otimes \Theta)$, with the property that for each $t \in B_{\epsilon_1}$, there is a $\phi(t) = t + \phi(t)^\perp \in F^{-1}(0)$, where $\phi(t)^\perp$ is L^2 -orthogonal to $\mathcal{H}_{-3}(X, \Lambda^{0,1} \otimes \Theta)$, and $J_0(\phi(t))$ is the corresponding complex structure. It is a straightforward consequence of the implicit function theorem that the mapping $\psi : t \mapsto \phi(t)$ is differentiable. This finishes the proof of the existence of the family of complex structures.

By the fixed point Lemma 4.1, near 0, the zero set of $(\bar{\partial}^*, \bar{\partial} + [\cdot, \cdot])$ is locally bijective to the kernel of $(\bar{\partial}^*, \bar{\partial})$, which is $\mathcal{H}_{-3}(X, \Lambda^{0,1} \otimes \Theta)$. Thus there exists an $\epsilon'' > 0$ such that for any $\|\phi\|_{C_{\delta-1}^{k,\alpha}} < \epsilon''$, $\bar{\partial}^* \phi = 0$, $\bar{\partial} \phi + [\phi, \phi] = 0$, then $J = J_0(\phi)$ is in the family we just constructed.

Next we will show that $\phi(t) \in C_{-3}^{k,\alpha}(X, \Lambda^{0,1} \otimes \Theta)$ for $t \in B_{\epsilon_1}$. In fact, we will show that for any $\phi \in C_{\delta}^{k,\alpha}(X, \Lambda^{0,1} \otimes \Theta)$ that satisfies the system

$$(4.17) \quad \bar{\partial} \phi = -[\phi, \phi], \quad \bar{\partial}^* \phi = 0,$$

then $\phi \in C_{-3}^{k,\alpha}(X, \Lambda^{0,1} \otimes \Theta)$. To see this, (4.17) implies that

$$(4.18) \quad \|\square \phi\|_{C_{2\delta-2}^{k-2,\alpha}} = \|-\bar{\partial}^*[\phi, \phi]\|_{C_{2\delta-2}^{k-2,\alpha}} \leq C \cdot \|\phi\|_{C_{\delta}^{k,\alpha}}^2$$

Then outside of a compact subset,

$$(4.19) \quad \frac{1}{2} \Delta_{Euc} \phi = -\square \phi + \left(\frac{1}{2} \Delta_{Euc} + \square\right) \phi = O(r^{2\delta-2}),$$

which implies that

$$(4.20) \quad \phi = \sum_{i,j} \frac{a_{i,j}}{r^2} d\bar{z}_i \otimes \frac{\partial}{\partial z_j} + O(r^{-3+\epsilon}),$$

where $a_{i,j}$ are constants and $\frac{a_{i,j}}{r^2}$ is Δ_{Euc} -harmonic. By the same argument as in the proof of (3.14), we have $a_{i,j} = 0$. Then a similar argument shows that $\phi \in C_{-3}^{k,\alpha}(X, \Lambda^{0,1} \otimes \Theta)$.

Next we will show that, if $\phi \in C_{\delta}^{k,\alpha}(X, \Lambda^{0,1} \otimes \Theta)$, $\|\phi\|_{C_{\delta}^{k,\alpha}} < \epsilon'$, and satisfies the system (4.17), then $\|\phi\|_{C_{\delta-1}^{k,\alpha}} \leq C \cdot \epsilon'$ for some constant $C > 0$. By the argument above, $\phi \in C_{-3}^{k,\alpha}(X, \Lambda^{0,1} \otimes \Theta)$, and $\phi = \phi^h + \phi^\perp$, where $\phi^h \in \mathcal{H}_{-3}(X, \Lambda^{0,1} \otimes \Theta)$, and ϕ^h is L^2 -orthogonal to ϕ^\perp . By an argument similar to Corollary 1.16 in [Bar86], $\|\phi^\perp\|_{C_{\delta-1}^{k,\alpha}} \leq C \cdot \|\square \phi\|_{C_{\delta-3}^{k-2,\alpha}}$. Since ϕ^h is in a finite dimensional space, combined with the estimate (4.18), we can make ϵ' small enough such that $\|\phi\|_{C_{\delta-1}^{k,\alpha}} < \epsilon''$, and then ϕ is in the family J_t as shown above. The estimate of $\|\phi\|_{C_{-3}^{k,\alpha}}$ follows from finite-dimensionality of the kernel. \square

5. VERSALITY OF KURANISHI FAMILY

The next result shows that any nearby complex structure can be brought into the family J_t by a suitable diffeomorphism. Of course, the implicit function theorem requires a mapping between Banach spaces to be differentiable. The following lemma, inspired by [Biq06], will be used below to this end.

Lemma 5.1. *Let (X, g_0, J_0) be a Kähler ALE surface with $g_0, J_0 \in C^\infty$. Then for $\delta' < -1$, for g_0, J_0 with ALE asymptotic rate $O(r^{\delta'})$, there exists an $\epsilon_1 > 0$, such that for an ϵ_1 -ball*

$B_{\epsilon_1} \subset C_{\delta'+1}^{k+1,\alpha}(TX)$, the maps

$$(5.1) \quad B_{\epsilon_1} \rightarrow C_{\delta'}^{k,\alpha}(S^2(T^*X)), \quad Y \mapsto \Phi_{Y*}g_0 - g_0$$

$$(5.2) \quad B_{\epsilon_1} \rightarrow C_{\delta'}^{k,\alpha}(End(TX)), \quad Y \mapsto \Phi_{Y*}J_0 - J_0$$

are smooth.

Proof. Let $U \subset X$ be a normal coordinate chart with respect to g_0 , and let U', U'' be smaller charts, such that the ϵ_1 -neighborhood of U' is in U , and the ϵ_1 -neighborhood of U'' is in U' . The tangent bundle TU is isomorphic to $U \times \mathbb{R}^4$, with coordinate functions $(x, v) = (x_1, \dots, x_4, v_1, \dots, v_4)$. On U' , we have the Riemannian exponential map:

$$(5.3) \quad \begin{aligned} Exp : U' \times B_{\epsilon_1} \subset TU' &\rightarrow U \\ (x, v) &\rightarrow \gamma(x, v) \end{aligned}$$

where $\gamma(x, v)$ is a geodesic with initial position and derivative (x, v) . The j^{th} -coordinate function of Exp ($1 \leq j \leq 4$) is a smooth function over (x, v) , and has an expansion

$$(5.4) \quad Exp_j(x, v) = x_j + \sum_{k=1}^4 f_k(x)v_k + Q(x, v)$$

where $|Q(x, v)| \leq C(U, g_0) \cdot |v|^2$. Let $\xi \in C^{k+1,\alpha}(TU)$, and $\|\xi\|_{C^{k+1,\alpha}} < \epsilon_1$. The geodesic flow Φ_ξ is defined as

$$(5.5) \quad \begin{aligned} \Phi_\xi : U' &\rightarrow U \\ x &\rightarrow Exp(x, \xi(x)) \end{aligned}$$

Then $\Phi_\xi(x) = Exp(x, \xi(x))$ is $C^{k+1,\alpha}$ over x , since Exp is smooth and $\xi(x)$ is $C^{k+1,\alpha}$. Let ξ' be another vector field on U with $\|\xi'\|_{C^{k+1,\alpha}} < \epsilon_1$. Then by the expansion above, for each $1 \leq j \leq 4$, we have:

$$(5.6) \quad \|(\Phi_\xi)_j - (\Phi_{\xi'})_j - \sum_{k=1}^4 f_k \cdot (\xi_k - \xi'_k)\|_{C^{k+1,\alpha}(U'')} \leq C \cdot \|\xi - \xi'\|_{C^{k+1,\alpha}(U)}^2$$

This implies that each coordinate function of Φ is Fréchet differentiable as a map from $B_{\epsilon_1} \subset C^{k+1,\alpha}(TU)$ to $C^{k+1,\alpha}(U'')$.

Since $\Phi_{\xi_*}g_0$ is defined as $(\Phi_\xi^{-1})^*g_0$, without the loss of generality, we will do the analysis on $\Phi_\xi^*g_0 - g_0$. On the chart U'' , we have $\Phi_\xi^*g_0(x)(\partial_i, \partial_j) = g_0(\Phi_\xi(x))(\Phi_{\xi_*}\partial_i, \Phi_{\xi_*}\partial_j)$. Since g_0 is smooth over x , and $\Phi_{\xi_*}\partial_j = \partial_j\Phi_\xi$ is $C^{k,\alpha}$ over x , by using the argument above, we can see that $\Phi^*g_0 - g_0 : B_{\epsilon_1} \subset C^{k+1,\alpha}(TU) \rightarrow C^{k,\alpha}(S^2(T^*U''))$ is Fréchet differentiable.

For some large $R > 0$, denote $A_{R,2R}$ as an annulus in the ALE space X . By covering $A_{R,2R}$ with finite many normal coordinate charts, using the argument above for each chart, we have the map $\Phi^*g_0 : B_{\epsilon_1} \subset C^{k+1,\alpha}(TA_{R,2R}) \rightarrow C^{k,\alpha}(S^2(T^*A_{R-\epsilon_1,2R-2\epsilon_1}))$ is Fréchet differentiable. By a standard dilation argument, which dilates $A_{R,2R}$ to $A_{2^k R, 2^{k+1} R}$, we have $\Phi^*g_0 - g_0 : B_{\epsilon_1} \subset C_{\delta'+1}^{k+1,\alpha}(TA_{2^k R, 2^{k+1} R}) \rightarrow C_{\delta'}^{k,\alpha}(S^2(T^*A_{2^k(R-\epsilon_1), 2^{k+1}(R-\epsilon_1)}))$ is Fréchet differentiable. The constant $C(U, g_0)$ in each chart U can be chosen uniformly bounded on X . This implies that $\Phi^*g_0 - g_0 : B_{\epsilon_1} \subset C_{\delta'+1}^{k+1,\alpha}(TX) \rightarrow C_{\delta'}^{k,\alpha}(S^2(T^*X))$ is Fréchet differentiable. Furthermore, it

follows that for $Y, Y' \in B_{\epsilon_1}$, there exists a constant C ,

$$(5.7) \quad \|\Phi_Y^* g_0 - \Phi_{Y'}^* g_0 - \mathfrak{L}_{Y-Y'} g_0\|_{C_{\delta'}^{k,\alpha}} \leq C \cdot \|Y - Y'\|_{C_{\delta'+1}^{k+1,\alpha}} \cdot (\|Y\|_{C_{\delta'+1}^{k+1,\alpha}} + \|Y'\|_{C_{\delta'+1}^{k+1,\alpha}}).$$

By using higher order expansions similar to (5.4), we can furthermore show that $\Phi^* g_0 - g_0$ is indeed a smooth map. The proof for J_0 can proceed in a similar fashion. Specifically we have

$$(5.8) \quad \|\Phi_Y^* J_0 - \Phi_{Y'}^* J_0 - \mathfrak{L}_{Y-Y'} J_0\|_{C_{\delta'}^{k,\alpha}} \leq C \cdot \|Y - Y'\|_{C_{\delta'+1}^{k+1,\alpha}} \cdot (\|Y\|_{C_{\delta'+1}^{k+1,\alpha}} + \|Y'\|_{C_{\delta'+1}^{k+1,\alpha}}).$$

□

Let $\nabla^* : \Gamma(\text{End}_a(TX)) \rightarrow \Gamma(TX)$ be the adjoint operator of $\frac{-1}{2}J \circ \mathfrak{L}_* J$, which is defined as:

$$(5.9) \quad (\nabla^* A)^k = - \sum_{l,j} g^{lj} \nabla_l A_j^k.$$

Note that under the identification of $\text{End}_a(TX)$ with $\text{Re}(\Lambda^{0,1} \otimes \Theta)$ and TX with $\text{Re}(\Theta)$, this corresponds to the operator $\bar{\partial}^*$. In the following proofs, for the convenience of the notation, we will denote each element in $\Gamma(\text{End}_a(TX))$ as $\text{Re}(\phi)$, where $\phi \in \Gamma(\Lambda^{0,1} \otimes \Theta)$.

Lemma 5.2. *Let (X, J_0, g_0) be a Kähler ALE surface with $J_0, g_0 \in C^\infty$. Let $\delta' < -1, k \geq 3, 0 < \alpha < 1$. Then the following map \mathfrak{P} is smooth in an open neighborhood of $(0, 0, 0)$*

$$(5.10) \quad \mathfrak{P} : C_{\delta'}^{k,\alpha}(S^2(TX)) \times C_{\delta'}^{k,\alpha}(\text{End}(TX)) \times C_{\delta'}^{k,\alpha}(\text{End}_a(TX)) \rightarrow C_{\delta'-1}^{k-1,\alpha}(TX)$$

$$(h, w, \text{Re}(\phi)) \mapsto \nabla_{g_0+h}^*(E_{J_0+w}^{-1}(J_0(\phi)))$$

Proof. The divergence operator has the expansion formula

$$(5.11) \quad \nabla_{g_0+h} = \nabla_{g_0} + (g_0 + h)^{-1} * \nabla_{g_0} h,$$

[GV16, Formula 3.39], where $*$ means a linear combination of tensor contractions, and $(g_0 + h)^{-1}$ is analytic for smooth h . By the expansions (4.7),

$$(5.12) \quad E_{J_0+w}^{-1}(J_0(\phi)) = -w + \text{Re}(\phi) + Q'$$

where Q' represents the higher order terms, which is analytic for small w . Note that $\mathfrak{P}(0, 0, 0) = 0$. For $(h, w, \text{Re}(\phi)), (h', w', \text{Re}(\phi'))$ of sufficiently small norm, by the expansion above, there exists a constant C so that

$$(5.13) \quad \|\mathfrak{P}(h, w, \text{Re}(\phi)) - \mathfrak{P}(h', w', \text{Re}(\phi')) + \nabla_{g_0}^*(w - w') - \nabla_{g_0}^* \text{Re}(\phi - \phi')\|_{C_{\delta'-1}^{k-1,\alpha}} \leq$$

$$C \cdot (\|h\|_{C_{\delta'}^{k,\alpha}} + \|h'\|_{C_{\delta'}^{k,\alpha}} + \|w\|_{C_{\delta'}^{k,\alpha}} + \|w'\|_{C_{\delta'}^{k,\alpha}} + \|\text{Re}(\phi)\|_{C_{\delta'}^{k,\alpha}}^2 + \|\text{Re}(\phi')\|_{C_{\delta'}^{k,\alpha}}^2)$$

$$\cdot (\|h - h'\|_{C_{\delta'}^{k,\alpha}} + \|w - w'\|_{C_{\delta'}^{k,\alpha}} + \|\text{Re}(\phi - \phi')\|_{C_{\delta'}^{k,\alpha}}^2)$$

This implies \mathfrak{P} is differentiable, and satisfies the condition of Lemma 4.1. By higher order expansions of \mathfrak{P} over $(h, w, \text{Re}(\phi))$, we can furthermore show that \mathfrak{P} is smooth. □

Lemma 5.3. *Let (X, J_0, g_0) be a Kähler ALE surface with $J_0, g_0 \in C^\infty$. There exists an $\epsilon'_1 > 0$ such that for any complex structure $\|J_1 - J_0\|_{C_\delta^{k,\alpha}} < \epsilon'_1$, where $k \geq 3, \alpha \in (0, 1), \delta \in (-2, -1)$, there exists a unique diffeomorphism Φ , of the form Φ_Y for $Y \in C_{\delta+1}^{k+1,\alpha}(TX)$ such that $\Phi^*(J_1)$ is in the family J_t from Theorem 4.3.*

Proof. Let $J_1 = J_0(\phi)$, where $\|\phi\|_{C_\delta^{k,\alpha}} < \epsilon'_1$ for some small ϵ'_1 to be determined later, and $\bar{\partial}\phi + [\phi, \phi] = 0$. Define $\phi_Y \in \Gamma(\Lambda^{0,1} \otimes \Theta)$ by $\Phi_Y^* J_0(\phi) = E_{J_0}(Re(\phi_Y))$. Define the operator \mathfrak{N} by choosing the first parameter of \mathfrak{P} to be $(\Phi_Y)_* g_0 - g_0$, choosing the second parameter to be $(\Phi_Y)_* J_0 - J_0$, and letting $\delta' = \delta$ as the following

$$(5.14) \quad \begin{aligned} C_{\delta+1}^{k+1,\alpha}(TX) \times C_\delta^{k,\alpha}(X, End_a(TX)) &\rightarrow C_{\delta-1}^{k-1,\alpha}(TX) \\ (Y, Re(\phi)) &\mapsto \nabla_{\Phi_Y^* g_0}^*(E_{\Phi_Y^* J_0}^{-1}(J_0(\phi))), \end{aligned}$$

where Φ_Y is defined in (1.14), and $\nabla_{\Phi_Y^* g_0}^*$ is the divergence operator associated to metric $\Phi_Y^* g_0$. Note that

$$(5.15) \quad \nabla_{\Phi_Y^* g_0}^*(E_{\Phi_Y^* J_0}^{-1}(J_0(\phi))) = 0 \Leftrightarrow \nabla_{g_0}^* E_{J_0}^{-1}(\Phi_Y^*(J_0(\phi))) = 0 \Leftrightarrow \nabla_{g_0}^* Re(\phi_Y) = 0,$$

so a zero of \mathcal{N} satisfies the desired gauge condition. (The latter map has a regularity issue which is why we consider the former, see [Biq06].) Since \mathfrak{P} is smooth by Lemma 5.2, and since the mappings $Y \mapsto (\Phi_Y)_* g_0 - g_0$, and $Y \mapsto (\Phi_Y)_* J_0 - J_0$ are smooth by Lemma 5.1, the composition \mathfrak{N} is smooth in an open neighborhood of $(0, 0)$.

Consequently, at $(0, 0)$, the linearization of \mathfrak{N} restricted to the tangent space of the first variable is

$$(5.16) \quad \begin{aligned} D\mathfrak{N}_0 : C_{\delta+1}^{k+1,\alpha}(TX) &\rightarrow C_{\delta-1}^{k-1,\alpha}(TX) \\ Y &\mapsto \nabla_{g_0}^* \circ \frac{-1}{2} J \circ \mathfrak{L}_Y J_0 = \square_{g_0} Y, \end{aligned}$$

where the adjoint operator of $D\mathfrak{N}_0$ is:

$$(5.17) \quad \begin{aligned} (D\mathfrak{N}_0)^* : C_{-3-\delta}^{k+1,\alpha}(TX) &\rightarrow C_{-5-\delta}^{k-1,\alpha}(TX) \\ \eta &\mapsto \square_{g_0} \eta, \end{aligned}$$

The kernel of $(D\mathfrak{N}_0)^*$ consists of those $\eta \in C_{-3-\delta}^{k+1,\alpha}(TX)$ such that $\square_{g_0} \eta = 0$, which implies that η is the real part of a holomorphic vector field. From Proposition 3.3, $\eta = 0$. Since the cokernel of $D\mathfrak{N}_0$ is trivial, $D\mathfrak{N}_0$ is surjective. Note also that the kernel of $D\mathfrak{N}_0$ is trivial.

Of course, in a small neighborhood of $(0, 0)$ in $C_{\delta+1}^{k+1,\alpha}(TX) \times C_\delta^{k,\alpha}(X, End_a(TX))$, $D\mathfrak{N}$ is surjective. Since \mathfrak{N} is smooth, condition (4.1) is satisfied, so by Lemma 4.1, there exists a unique diffeomorphism Φ_Y such that $Re(\phi_Y)$ is divergence-free. Since ϕ_Y satisfies (4.9), by Theorem 4.3, $\Phi_Y^*(J_1)$ is in the family J_t . \square

6. ESSENTIAL DEFORMATIONS OF COMPLEX STRUCTURE

The arguments in this section are inspired by [CT94, Theorem 3.1], with some appropriate modifications for the smoothness arguments. Recall that the space \mathbb{W} is defined as

$$(6.1) \quad \mathbb{W} = \{Z \in \mathcal{H}_1(X, TX) \mid \mathfrak{L}_Z g_0 = O(r^{-1}), \mathfrak{L}_Z J_0 = O(r^{-3}), \text{ as } r \rightarrow \infty\}.$$

Clearly, \mathbb{W} is finite-dimensional. For $Z \in \mathbb{W}$, we define $\Upsilon_Z : X \rightarrow X$ to be time-one flow for the one-parameter group of diffeomorphisms generated by Z . Note that since \mathbb{W} is finite-dimensional the use of the one-parameter group here will not have the loss-of-regularity problem as in infinite dimensions. Furthermore, the time-one flow for vector fields in \mathbb{W} has the following nice property.

Lemma 6.1. *Fix an ALE coordinate under which $|J_0 - J_{Euc}| = O(r^{-3})$. For $Z \in \mathbb{W}$ sufficiently small, the time-one flow for the one-parameter group of diffeomorphisms generated by Z exists. Furthermore, there exist a matrix $A \in \mathrm{U}(2)$ and vector $\hat{e}_0 \in \mathbb{C}^2$, such that in ALE coordinates of order μ with $-2 < -\mu < -1$,*

$$(6.2) \quad d(\Upsilon_Z(x), Ax + \hat{e}_0) = O(r^{-\mu}),$$

as $r \rightarrow \infty$, for any $\epsilon > 0$.

Proof. First, note that $Z \in \mathbb{W}$ admits an expansion

$$(6.3) \quad Z = Z_1 + Z_0 + Z_\mu,$$

for any $\epsilon > 0$, where Z_1 is in the Lie algebra of $\mathrm{U}(2)$, Z_0 is vector field with constant coefficients, and $Z_\mu = O(r^{-\mu})$ as $r \rightarrow \infty$. To see this, by standard harmonic expansion for kernel elements of \square_{g_0} , we have that any $Z \in \mathbb{W}$ satisfies

$$(6.4) \quad Z = Z_1 + Z_0 + O(r^{-\mu}),$$

where Z_1 is any vector field which is homogeneous of degree one, and Z_0 is a vector field with constant coefficients. The conditions that $\mathfrak{L}_Z g_0 = O(r^{-1})$ and $\mathfrak{L}_Z J_0 = O(r^{-3})$ as $r \rightarrow \infty$ imply that $\mathfrak{L}_{Z_1} g_{Euc} = 0$ and $\mathfrak{L}_{Z_1} J_{Euc} = 0$, that is, Z_1 is in the Lie algebra of $\mathrm{U}(2)$. Next,

$$(6.5) \quad \mathfrak{L}_{Z_1+Z_0} J_0 = \mathfrak{L}_{Z_1+Z_0} (J_0 - J_{Euc}) = \mathfrak{L}_{Z_1+Z_0} (O(r^{-3})),$$

which implies that

$$(6.6) \quad \mathfrak{L}_{Z_0+Z_1} J_0 = O(r^{-3}),$$

as $r \rightarrow \infty$. Consequently,

$$(6.7) \quad \square_0(Z - Z_1 - Z_0) = O(r^{-4}).$$

Since there is no harmonic term of degree -1 , the claimed expansion (6.3) follows.

Clearly for Z sufficiently small in norm, the one-parameter group of diffeomorphisms which is defined by

$$(6.8) \quad x'(t) = Z(x(t)), \quad x(0) = x,$$

exists by the standard short-time existence theorem for ODEs. By the expansion (6.3), $x(t)$ will be given by a family of rotations in $\mathrm{U}(2)$, and translations plus decaying terms. \square

Remark 6.2. Lemma 6.1 has assumed an ALE coordinate with inverse cubic complex structure asymptotic rate. In the following, note that the statements Lemma 6.3 and Theorem 6.4 are independent of the ALE coordinate in the sense that, if they are true under the ALE coordinate (x^1, x^2, x^3, x^4) , then they are also true under the ALE coordinate (x'^1, x'^2, x'^3, x'^4) , as long as the two coordinates are comparable, i.e. $\frac{1}{C}|x| < |x'| < C|x|$ for some constant $C > 0$. Thus, in their proofs, we are free to choose an ALE coordinate such that $|J_0 - J_{Euc}| = O(r^{-3})$ by Proposition 3.1.

The next lemma states some useful properties of \mathbb{W} , which will be needed in the proof of the subsequent gauging theorem. For the simplicity of the statement of the theorem, define the norm $\|Z\|_{k+1, \alpha}$ for $Z \in \mathbb{W}$ as $\|Z\|_{k+1, \alpha} = \|Z\|_{C_1^{k+1, \alpha}}$ if $\lim_{r \rightarrow \infty} |r^{-1} \cdot Z| \neq 0$, $\|Z\|_{k+1, \alpha} = \|Z\|_{C^{k+1, \alpha}}$ if $\lim_{r \rightarrow \infty} |r^{-1} \cdot Z| = 0$. Note that if $Z \in \mathbb{W}$ is decaying, then, $Z \equiv 0$.

Lemma 6.3. *Let (X, J_0, g_0) be a Kähler ALE surface, the space \mathbb{W} satisfies the following properties:*

- *There exists a small neighborhood $U \subset \mathbb{W}$ of 0, and an $\epsilon' > 0$ such that if g_1 is any metric satisfying $\|g_1 - g_0\|_{C_\delta^{k,\alpha}} < \epsilon'$, then there exists a constant C , such that for any $Z \in U$,*

$$(6.9) \quad \|\Upsilon_Z^* g_1 - g_0\|_{C_\delta^{k,\alpha}} \leq C(\|Z\|_{k+1,\alpha} + \epsilon').$$

- *Given $\epsilon > 0$ sufficiently small, for any complex structure $J_1 = J_0(\phi)$ such that $\|\phi\|_{C_{-3+\epsilon}^{k,\alpha}} < \epsilon'$ in the family constructed in Theorem 4.3 for some sufficiently small ϵ' , there exists a small neighborhood $U \subset \mathbb{W}$ of 0, and a constant C , such that for any $Z, Z' \in U$,*

$$(6.10) \quad \|\Upsilon_Z^* J_1 - \Upsilon_{Z'}^* J_1 - \mathfrak{L}_{Z-Z'} J_1\|_{C_{-3+\epsilon}^{k,\alpha}} \leq C \cdot \|Z - Z'\|_{k+1,\alpha} \cdot (\|Z\|_{k+1,\alpha} + \|Z'\|_{k+1,\alpha}).$$

Furthermore, the following map

$$(6.11) \quad \begin{aligned} \Upsilon^* : \mathcal{H}_{-3}(X, \text{End}_a(TX)) \times \mathbb{W} &\rightarrow C_{-3+\epsilon}^{k,\alpha}(X, \text{End}_a(TX)) \\ (\text{Re}(t), Z) &\mapsto E_{J_0}^{-1}(\Upsilon_Z^* J_0(\phi(t))) \equiv \phi_Z(t) \end{aligned}$$

is smooth in an open neighborhood of $(0, 0)$.

Proof. Fix an ALE coordinate under which $|J_0 - J_{\text{Euc}}| = O(r^{-3})$. Under such a coordinate, we have the decomposition of Z as stated in Lemma 6.1. For simplicity, we assume $Z_1 \neq 0$ in the following, and $\|Z\|_{k+1,\alpha} = \|Z\|_{C_1^{k+1,\alpha}}$. (The proof for the case $Z_1 = 0$ is similar.)

First, we address the estimate (6.9). Let $\Upsilon_{A,\hat{\epsilon}_0}$ denote the mapping $x \mapsto Ax + \hat{\epsilon}_0$. Using Lemma 6.1,

$$\begin{aligned} |\Upsilon_Z^* g_1 - g_0| &= |\Upsilon_Z^* g_1 - g_1 + g_1 - g_0| \leq |\Upsilon_Z^* g_1 - g_1| + |g_1 - g_0| \\ &\leq |(\Upsilon_{A,\hat{\epsilon}_0} + O(r^{-2+\epsilon}))^*(\delta_{ij} + O(r^\delta)) - (\delta_{ij} + O(r^\delta))| + \epsilon' r^\delta \\ &\leq C\|Z\|_{C_1^{k+1,\alpha}} r^\delta + \epsilon' r^\delta + o(r^\delta) \\ &\leq C r^\delta (\|Z\|_{C_1^{k+1,\alpha}} + \epsilon'). \end{aligned}$$

as $r \rightarrow \infty$. Since Υ_Z is the time one flow of $Z \in \mathbb{W}$, and \mathbb{W} is finite-dimensional, a similar estimate holds on any compact subset of X , so the C_δ^0 part of the norm is bounded by the right hand side of (6.9). Higher regularity estimates are similar, and are omitted.

Next we will discuss (6.10). As in the first part, since \mathbb{W} is finite-dimensional, we only need to make estimates outside of a compact set $\overline{B_R(p_0)}$, where a global coordinate exists. Let $\gamma : [0, 1] \rightarrow X$ be the path $\gamma(t) = \Upsilon_{tZ}(x)$. First, we estimate

$$(6.12) \quad \begin{aligned} r(x)^{3-\epsilon} |\Upsilon_Z^* J_0 - J_0 - \mathfrak{L}_Z J_0|(x) &\leq C r^{3-\epsilon} \int_0^1 |\Upsilon_{tZ}^*(\mathfrak{L}_Z J_0)(\gamma(0)) - \mathfrak{L}_Z J_0(\gamma(0))| dt \\ &\leq C r^{3-\epsilon} \int_0^1 \int_0^t |\mathfrak{L}_Z(\mathfrak{L}_Z J_0)(\gamma(s))| ds dt. \end{aligned}$$

By the expansion (6.3), given $c > 0$, the estimates

$$(6.13) \quad (1 - c)r(\gamma(s)) \leq r(x) = r(\gamma(0)) \leq (1 + c)r(\gamma(s))$$

are satisfied for all Z sufficiently small in norm. Note that since Z_1 is a vector field corresponding to a rotation in $\mathfrak{u}(2)$, $|Z_1| \geq C \cdot r$ for some constant C . Also, since \mathbb{W} is finite dimensional, $|\nabla^Z Z| \leq C(|\mathcal{I}|) \cdot r^{-|\mathcal{I}|} \cdot (|Z| + 1)$ for some constant $C(|\mathcal{I}|) > 0$. Then

$$(6.14) \quad |\mathfrak{L}_Z(\mathfrak{L}_Z J_0)|(\gamma(s)) \leq Cr(\gamma(s))^{-5}(|Z(\gamma(s))| + 1)^2,$$

so we have

$$(6.15) \quad \begin{aligned} r(x)^{3-\epsilon} |\Upsilon_Z^* J_0 - J_0 - \mathfrak{L}_Z J_0|(x) &\leq C \int_0^1 \int_0^t r(\gamma(s))^{3-\epsilon} r(\gamma(s))^{-5} (|Z| + 1)^2(\gamma(s)) ds dt \\ &\leq C \int_0^1 \int_0^t r(\gamma(s))^{-2-\epsilon} (|Z| + 1)^2(\gamma(s)) ds dt \leq C \|Z\|_{C_1^{k+1,\alpha}}^2. \end{aligned}$$

Next, we estimate

$$(6.16) \quad \begin{aligned} r(x)^{3-\epsilon} |\Upsilon_Z^* \phi - \phi - \mathfrak{L}_Z \phi|(x) &\leq Cr^{3-\epsilon} \int_0^1 |\Upsilon_{tZ}^*(\mathfrak{L}_Z \phi)(\gamma(0)) - \mathfrak{L}_Z \phi(\gamma(0))| dt \\ &\leq Cr^{3-\epsilon} \int_0^1 \int_0^t |\mathfrak{L}_Z(\mathfrak{L}_Z \phi)(\gamma(s))| ds dt. \end{aligned}$$

Since $|\mathfrak{L}_Z(\mathfrak{L}_Z \phi)|(\gamma(s)) \leq Cr(\gamma(s))^{-5+\epsilon} (|Z| + 1)^2(\gamma(s))$, similarly, we have the estimate

$$(6.17) \quad \begin{aligned} r(x)^{3-\epsilon} |\Upsilon_Z^* \phi - \phi - \mathfrak{L}_Z \phi|(x) &\leq C \int_0^1 \int_0^t r(\gamma(s))^{3-\epsilon} (r(\gamma(s))^{-5+\epsilon} (|Z| + 1)^2(\gamma(s))) ds dt \\ &\leq C \int_0^1 \int_0^t r(\gamma(s))^{-2} (|Z| + 1)^2(\gamma(s)) ds dt \leq C \|Z\|_{C_1^{k+1,\alpha}}^2. \end{aligned}$$

By a similar calculation, for the higher order term Q in (4.7) we also have

$$(6.18) \quad r(x)^{3-\epsilon} |\Upsilon_Z^* Q - Q - \mathfrak{L}_Z Q|(x) \leq C \|Z\|_{C_1^{k+1,\alpha}}^2$$

It follows that

$$(6.19) \quad \begin{aligned} r(x)^{3-\epsilon} |\Upsilon_Z^* J_1 - J_1 - \mathfrak{L}_Z J_1|(x) &\leq r(x)^{3-\epsilon} \{ |\Upsilon_Z^* J_0 - J_0 - \mathfrak{L}_Z J_0|(x) + \\ &\quad |\Upsilon_Z^* \phi - \phi - \mathfrak{L}_Z \phi|(x) \} + |\Upsilon_Z^* Q - Q - \mathfrak{L}_Z Q|(x) \\ &\leq C \|Z\|_{C_1^{k+1,\alpha}}^2, \end{aligned}$$

where C is uniform for all $Z \in \mathbb{W}$ of sufficiently small norm. And similarly, for any Z, Z' in a small neighborhood of 0,

$$(6.20) \quad \Upsilon_Z^* J_1 - \Upsilon_{Z'}^* J_1 = \Upsilon_{Z'}^* ((\Upsilon_{Z'}^*)^{-1} \circ \Upsilon_Z^* J_1 - J_1)$$

$$(6.21) \quad = \Upsilon_{Z'}^* ((Id - \mathfrak{L}_{Z'} + R') \circ (Id + \mathfrak{L}_Z + R) J_1 - J_1)$$

where R, R' represents the corresponding higher order terms, and has estimate as in (6.19). Then we have

$$(6.22) \quad r^{3-\epsilon} |\Upsilon_Z^* J_1 - \Upsilon_{Z'}^* J_1 - \mathfrak{L}_{Z-Z'} J_1| \leq C \cdot \|Z - Z'\|_{C_1^{k+1,\alpha}} \cdot (\|Z\|_{C_1^{k+1,\alpha}} + \|Z'\|_{C_1^{k+1,\alpha}})$$

Higher order derivative estimates are similar, and are omitted, and (6.10) follows. Then by (5.12) and (6.10) we can see that Υ^* is differentiable, and satisfies the condition of

Lemma 4.1. A similar estimate of higher order expansions of Υ^* shows that it is a smooth map near $(0, 0)$. \square

Recall that

$$(6.23) \quad \mathbb{V} = \{\theta \in \mathcal{H}_{-3}(X, \text{End}_a(TX)) \mid \theta = \mathfrak{L}_Z J, Z \in \mathbb{W}\}.$$

The deformations of complex structure arising from elements of \mathbb{V} become trivial if one takes into account changes of coordinates at infinity, as the following theorem shows.

Theorem 6.4. *Let (X, J_0, g_0) be a Kähler ALE surface with $J_0 \in C^\infty$ and $g_0 \in C^\infty$. There exists an $\epsilon'_1 > 0$ such that for any complex structure $\|J_1 - J_0\|_{C_\delta^{k,\alpha}} < \epsilon'_1$, where $k \geq 3, \alpha \in (0, 1), \delta \in (-2, -1)$, there exists a diffeomorphism Φ , of the form $\Phi_{Y_1} \circ \Upsilon_Z \circ \Phi_{Y_2}$ for $Y_1, Y_2 \in C_{\delta+1}^{k+1,\alpha}(TX)$ and $Z \in \mathbb{W}$, such that $\nabla_{g_0}^*(E_{J_0}^{-1}(\Phi^*(J_1))) = 0$, and such that $E_{J_0}^{-1}(\Phi^*(J_1))$ is L^2 -orthogonal to \mathbb{V} . Furthermore, there exists a constant C so that*

$$(6.24) \quad \|E_{J_0}^{-1}(\Phi^* J_1)\|_{C_{-3}^{k,\alpha}} < C\epsilon'_1.$$

Proof. Let $J_1 = J_0(\phi)$, where $\phi \in C_\delta^{k,\alpha}(X, \Lambda^{0,1} \otimes \Theta)$, and $\bar{\partial}\phi + [\phi, \phi] = 0$. By Lemma 5.3 and Theorem 4.3, without loss of generality, we may assume that $\phi = \phi(t) \in C_{-3}^{k,\alpha}(X, \Lambda^{0,1} \otimes \Theta)$, where $t \in B_{\epsilon_1} \subset \mathcal{H}_{-3}(X, \Lambda^{0,1} \otimes \Theta)$. In the following of the proof, we will fixed the ALE coordinate as in Lemma 6.1, and again for simplicity, we will assume that $Z_1 \neq 0$.

Define the operator \mathfrak{N} as

$$(6.25) \quad \begin{aligned} & C_{-2+\epsilon}^{k+1,\alpha}(TX) \times \mathcal{H}_{-3}(X, \text{End}_a(TX)) \times \mathbb{W} \rightarrow C_{-4+\epsilon}^{k-1,\alpha}(TX) \times \mathbb{R}^m \\ & (Y, \text{Re}(t), Z) \mapsto \\ & \left\{ \nabla_{\Phi_{Y^*g_0}}^* E_{\Phi_{Y^*J_0}}^{-1}(\Upsilon_Z^* J_0(\phi(t))), \int \left\langle E_{\Phi_{Y^*J_0}}^{-1}(\Upsilon_Z^* J_0(\phi(t))), \Phi_{Y^*} v_1 \right\rangle_{\Phi_{Y^*g_0}} dV_{\Phi_{Y^*g_0}}, \right. \\ & \quad \left. \dots, \int \left\langle E_{\Phi_{Y^*J_0}}^{-1}(\Upsilon_Z^* J_0(\phi(t))), \Phi_{Y^*} v_m \right\rangle_{\Phi_{Y^*g_0}} dV_{\Phi_{Y^*g_0}} \right\}, \end{aligned}$$

where $0 < \epsilon \ll 1$, $\nabla_{\Phi_{Y^*g_0}}^*$ is the divergence operator associated to the metric $\Phi_{Y^*g_0}$, $m = \dim(\mathbb{V})$, $\{v_1, \dots, v_m\}$ is an orthonormal basis of \mathbb{V} . Note that the method of (5.15) is used here again to make the operator smooth. Also note that the map

$$(6.26) \quad (Y, \text{Re}(t), Z) \mapsto \nabla_{\Phi_{Y^*g_0}}^* E_{\Phi_{Y^*J_0}}^{-1}(\Upsilon_Z^* J_0(\phi(t)))$$

is defined by choosing the first parameter of \mathfrak{P} to be $(\Phi_Y)_*g_0 - g_0$, choosing the second parameter to be $(\Phi_Y)_*J_0 - J_0$, choosing the third parameter to be $\phi_Z(t)$ (defined in (6.11)), and letting $\delta' = -3 + \epsilon$. Since \mathfrak{P} is smooth by Lemma 5.2, and since the mappings $Y \mapsto (\Phi_Y)_*g_0 - g_0$, and $Y \mapsto (\Phi_Y)_*J_0 - J_0$ are smooth by Lemma 5.1, and since Υ^* is smooth by Lemma 6.3, we have that (6.26) is a smooth mapping. By applying the same argument to the inner products in \mathfrak{N} , we can show that \mathfrak{N} is smooth in an open neighborhood of $(0, 0, 0)$.

At $(0, 0, 0)$, the linearization of \mathfrak{N} restricted to the tangent spaces of the first and third variables is:

$$(6.27) \quad \begin{aligned} D\mathfrak{N}_0 : C_{-2+\epsilon}^{k+1,\alpha}(TX) \times \mathbb{W} &\rightarrow C_{-4+\epsilon}^{k-1,\alpha}(TX) \times \mathbb{R}^m \\ (Y, Z) &\mapsto \left\{ \nabla_{g_0}^* \mathfrak{L}_Y J_0 = \square_{g_0} Y, \int \left\langle \frac{-1}{2} J_0 \mathfrak{L}_Z J_0 - \frac{1}{2} J_0 \mathfrak{L}_Y J_0, v_1 \right\rangle_{g_0} dV_{g_0}, \right. \\ &\quad \left. \dots, \int \left\langle \frac{-1}{2} J_0 \mathfrak{L}_Z J_0 - \frac{1}{2} J_0 \mathfrak{L}_Y J_0, v_m \right\rangle_{g_0} dV_{g_0} \right\}. \end{aligned}$$

Restricting the domain and image, consider $D\mathfrak{N}_0$ mapping from $C_{-2+\epsilon}^{k+1,\alpha}(TX)$ to $C_{-4+\epsilon}^{k-1,\alpha}(TX)$. The adjoint of this restricted mapping, which we again denote by $D\mathfrak{N}_0$, is:

$$(6.28) \quad \begin{aligned} (D\mathfrak{N}_0)^* : C_{-2-\epsilon}^{k+1,\alpha}(TX) &\rightarrow C_{-2-\epsilon}^{k-1,\alpha}(TX) \\ \eta &\mapsto (\square_{g_0} \eta) \end{aligned}$$

The kernel of $(D\mathfrak{N}_0)^*$ consists of those $\eta \in C_{-2-\epsilon}^{k+1,\alpha}(TX)$ such that $(\square_{g_0} \eta) = 0$, which implies that η is the real part of a holomorphic vector field. From Proposition 3.3, $\eta = 0$. Since the cokernel of $D\mathfrak{N}_0$ is trivial, $D\mathfrak{N}_0$ is surjective on the restricted domain and range.

It follows that the full mapping $D\mathfrak{N}_0$ is surjective, since \mathbb{V} is generated by $\frac{-1}{2} J_0 \mathfrak{L}_Z J_0$ for $Z \in \mathbb{W}$. Clearly then, in a small neighborhood of $(0, 0, 0)$, $D\mathfrak{N}$ is surjective. Since the mapping \mathfrak{N} is smooth, Condition (4.1) is satisfied, so the existence of diffeomorphism Φ then follows from Lemma 4.1. Finally, the estimate (6.24) follows from estimates above. \square

7. STABILITY OF KÄHLER METRICS

In this section, we want to generalize Kodaira-Spencer's stability theory of Kähler metrics to ALE surfaces to the family of complex structures found above [KS60]. Specifically, we want to find a smooth family of Kähler forms on (X, J_t) for t small, where J_t for $t \in B_{\epsilon_1} \subset \mathcal{H}_{ess}(X, \Lambda^{0,1} \otimes \Theta)$ is the family of complex structures constructed in Section 4. In a slight abuse of notation, the notation X_t will stand for (X, J_t) . All the norms below in this section are based on the central metric g_0 .

Lemma 7.1. *The integer-valued function $\dim(\mathcal{H}_\tau(X_t, \Lambda^{p,q}))$ is upper semicontinuous at $t = 0$, for any $\tau \in (-2, 0)$.*

Proof. The Hodge Laplacian $\Delta_d = dd^* + d^*d$ is a Fredholm operator on $C_\tau^{k,\alpha}$ ($-2 < \tau < 0, k \geq 2$). The following is proved in [Bar86]: If P is a pseudo-differential operator such that $\|P - \frac{1}{2}\Delta_d\|_{op} < \epsilon$ for some ϵ sufficiently small, then $\ker(P) \leq \ker(\Delta_d)$, where $\|\cdot\|_{op}$ is the norm of operators from $C_\tau^{k,\alpha}$ to $C_{\tau-2}^{k-2,\alpha}$, which is defined as:

$$(7.1) \quad \|P\|_{op} = \sup\{\|P(u)\|_{C_{\tau-2}^{k-2,\alpha}} : \|u\|_{C_\tau^{k,\alpha}} = 1\}.$$

We may apply this to the operator $P = \square_t = \bar{\partial}_t^* \bar{\partial}_t + \bar{\partial}_t \bar{\partial}_t^*$, since we have $\|\square_t - \frac{1}{2}\Delta_d\|_{op} < \epsilon$ when $t \in \Delta_\gamma^{d_1}$ is small enough (when $t = 0$, $\square_0 = \frac{1}{2}\Delta_d$ by the Kähler identities). Then

$$(7.2) \quad \dim(\mathcal{H}_\tau(X_t, \Lambda^{p,q})) \leq \dim(\mathcal{H}_\tau(X_0, \Lambda^{p,q})).$$

\square

Let e_t^τ denote the dimension of the space of \square_t -harmonic forms with decay rate of τ . Note that by a similar argument as in Proposition 3.5, any such harmonic form must decay like $O(r^{-3})$ as $r \rightarrow \infty$. So by [Joy00, Sections 8.4 and 8.9], and Proposition 3.6, we have

$$(7.3) \quad b_0^{2,0} = b_0^{0,2} = b_0^{1,0} = b_0^{0,1} = 0.$$

From semi-continuity, it follows that

$$(7.4) \quad e_t^{2,0} = e_t^{0,2} = e_t^{1,0} = e_t^{0,1} = 0.$$

The proof of the following theorem is inspired by [BR15].

Theorem 7.2. *By choosing ϵ_1 small enough, for the family of deformation (X, J_t) , $t \in B_{\epsilon_1}$, there exists a smooth family of Kähler forms ω_t on X_t , and*

$$(7.5) \quad \|\omega_t - \omega_0\|_{C_\delta^{k,\alpha}} \leq C \cdot \|t\|_{C_{-3}^{k,\alpha}} \leq C \cdot \epsilon_1$$

for some constant C , and some $k \geq 3, \delta \in (-2, -1)$.

Furthermore, letting \mathfrak{G} denote the group of holomorphic isometries of (X, g_0, J_0) , the mapping

$$(7.6) \quad \begin{aligned} \mathfrak{N} : B_{\epsilon_1} &\rightarrow \Lambda^2(T^*X) \\ t &\mapsto \omega_t \end{aligned}$$

can be chosen equivariantly with respect to the action of \mathfrak{G} . That is, $\mathfrak{N}(\iota^*t) = \iota^*\mathfrak{N}(t)$, for all $\iota \in \mathfrak{G}$.

Proof. Let $g_t = \frac{1}{2}(g + g \circ J_t)$ be the Hermitian metric on X_t , where g is the Riemannian metric of the central fiber and J_t is the complex structure of X_t . Let $\square_t = \bar{\partial}_t^* \bar{\partial}_t + \bar{\partial}_t \bar{\partial}_t^*$ be the $\bar{\partial}_t$ -Laplacian defined with respect to g_t . Let $\omega'_t = \frac{\sqrt{-1}}{2}g_t(J_t, \cdot)$ be the corresponding $(1, 1)$ -form. We want to perturb ω'_t to acquire a Kähler form ω_t . We have $\omega'_t = \omega_t^h + \phi_t$, where ω_t^h is the \square_t -harmonic part of ω'_t , which is asymptotic to $dz_t \wedge d\bar{z}_t$ at the rate of $-\delta$, with $\|\phi_t\|_{C_\delta^{k,\alpha}} \leq C \|\square_t \omega'_t\|_{C_{\delta-2}^{k-2,\alpha}}$. We next show that this claimed decomposition is valid. Consider the following Fredholm operator

$$(7.7) \quad \square_t : C_\delta^{k,\alpha}(X_t, \Lambda^{1,1}) \rightarrow C_{\delta-2}^{k-2,\alpha}(X_t, \Lambda^{1,1}).$$

The kernel $\ker \square_t^* \subset C_{-2-\delta}^{k,\alpha}(X_t, \Lambda^{1,1})$ is of finite dimension. For any $\sigma \in \ker \square_t^*$, σ admits an expansion

$$(7.8) \quad \sigma = \sum_{k=1}^2 \frac{a_k}{r^2} dz_t \wedge d\bar{z}_t + O(r^{-3})$$

as $r \rightarrow \infty$, where a_k are constants. Since

$$(7.9) \quad \begin{aligned} 0 &= \int_X \langle \sigma, \square_t \sigma \rangle_{g_t} dV_{g_t} = \lim_{R \rightarrow \infty} \left\{ \int_{B(R)} (\langle \bar{\partial} \sigma, \bar{\partial} \sigma \rangle_{g_t} + \langle \bar{\partial}^* \sigma, \bar{\partial}^* \sigma \rangle_{g_t}) dV_{g_t} \right. \\ &\quad \left. + \int_{S(R)} (\langle \sigma, \bar{\partial} r \lrcorner \bar{\partial} \sigma \rangle_{g_t} + \langle \bar{\partial}^* \sigma, \bar{\partial} r \lrcorner \sigma \rangle_{g_t}) dA_{g_t} \right\} \\ &= \int_X (\langle \bar{\partial} \sigma, \bar{\partial} \sigma \rangle_{g_t} + \langle \bar{\partial}^* \sigma, \bar{\partial}^* \sigma \rangle_{g_t}) dV_{g_t}. \end{aligned}$$

This shows that $\bar{\partial}\sigma = 0$, $\bar{\partial}^*\sigma = 0$. Since also $\bar{\partial}_t - \bar{\partial}_{Euc} = O(r^{-3})$, we have $\bar{\partial}_{Euc}\sigma = O(r^{-5})$. Then $a_k = 0$ and $\sigma = O(r^{-3})$. This implies that $\square_t\omega'_t \in (\ker \square_t^*)^\perp \cap C_{\delta-2}^{k-2,\alpha}(X_t, \Lambda^{1,1})$, which is the L^2 -complement of $\ker \square_t^*$ in $C_{\delta-2}^{k-2,\alpha}(X_t, \Lambda^{1,1})$. Then there exists a $\phi_t \in C_\delta^{k,\alpha}(X_t, \Lambda^{1,1})$ such that $\square_t\phi_t = \square_t\omega'_t$ and $\|\phi_t\|_{C_\delta^{k,\alpha}} \leq C\|\square_t\omega'_t\|_{C_{\delta-2}^{k-2,\alpha}}$. Then we can let $\omega_t^h = \omega'_t - \phi_t$ which is \square_t -harmonic.

Proposition 7.3. *If $\square_t\omega_t^h = 0$, then $\bar{\partial}_t\omega_t^h = 0$.*

Proof. $\bar{\partial}_t\square_t\omega_t^h = \square_t\bar{\partial}_t\omega_t^h = 0$, then $\bar{\partial}_t\omega_t^h$ is \square_t -harmonic, with decay rate of $-(\delta-1)$. However, $e_t^{2,1} = e_t^{0,1} = 0$ by the upper semi-continuity and since the conjugate Hodge-star operator maps harmonic forms to harmonic forms. Then $\bar{\partial}_t\omega_t^h = 0$. \square

We want to find ω_t such that $d\omega_t = 0$; for this we need a lemma.

Lemma 7.4. *We have $\partial_t\omega_t^h = \bar{\partial}_ta_t$, where $a_t \in C_\delta^{k,\alpha}(X_t, \Lambda^{2,0})$, $k \geq 3$. Furthermore,*

$$(7.10) \quad \|a_t\|_{C_\delta^{k,\alpha}} \leq C\|\partial_t\omega_t^h\|_{C_{\delta-1}^{k-1,\alpha}} \leq C'\|t\|_{C_\delta^{k,\alpha}}$$

Proof. First note that

$$(7.11) \quad \bar{\partial}_t\partial_t\omega_t^h = -\partial_t\bar{\partial}_t\omega_t^h = 0.$$

Then

$$(7.12) \quad \partial_t\omega_t^h \in C_{\delta-1}^{k-1,\alpha}(X_t, \Lambda^{2,1}) = \mathcal{H}_{\delta-1}(X_t, \Lambda^{2,1}) \oplus \square_t C_{\delta+1}^{k+1,\alpha}(X_t, \Lambda^{2,1}).$$

(This decomposition follows since \square_t is Fredholm on $C_{\delta+1}^{k,\alpha}(X_t, \Lambda^{2,1})$.) Then

$$(7.13) \quad \partial_t\omega_t^h = h_t + (\bar{\partial}_t\bar{\partial}_t^* + \bar{\partial}_t^*\bar{\partial}_t)f_t$$

with $f_t \in C_{\delta+1}^{k+1,\alpha}(X_t, \Lambda^{2,1})$, where h_t is the harmonic part. Since $\bar{\partial}_t\partial_t\omega_t^h = 0$, it follows that $\square_t\bar{\partial}_t f_t = 0$. From injectivity of \square_t on $C_\delta^{k,\alpha}$, we have $\bar{\partial}_t f_t = 0$. Then

$$(7.14) \quad \partial_t\omega_t^h = h_t + \bar{\partial}_t\bar{\partial}_t^* f_t = h_t + \bar{\partial}_ta_t,$$

where $a_t = \bar{\partial}_t^* f_t \in C_\delta^{k,\alpha}(X_t, \Lambda^{2,0})$. Since $e_t^{2,1} = 0$, then $h_t = 0$, and $\partial_t\omega_t^h = \bar{\partial}_ta_t$. Since $\dim(\mathcal{H}_{\delta+1}(X_t, \Lambda^{2,1})) = 0$,

$$(7.15) \quad \square_t : C_{\delta+1}^{k+1,\alpha}(X_t, \Lambda^{2,1}) \rightarrow C_{\delta-1}^{k-1,\alpha}(X_t, \Lambda^{2,1})$$

is an isomorphism. Then $\|f_t\|_{C_{\delta+1}^{k+1,\alpha}} \leq C\|\partial_t\omega_t^h\|_{C_{\delta-1}^{k-1,\alpha}}$ implies that

$$(7.16) \quad \|a\|_{C_\delta^{k,\alpha}} \leq C\|\partial_t\omega_t^h\|_{C_{\delta-1}^{k-1,\alpha}} \leq C'\|t\|_{C_\delta^{k,\alpha}},$$

\square

Now we have $a_t \in C_\delta^{k,\alpha}(X_t, \Lambda^{2,0})$ and $\bar{\partial}_t\bar{a}_t = 0$. Let $\delta + 2$ not be a indicial root of \square_t , then

$$(7.17) \quad \square_t : C_{\delta+2}^{k+2,\alpha}(X_t, \Lambda^{0,2}) \rightarrow C_\delta^{k,\alpha}(X_t, \Lambda^{0,2})$$

is Fredholm. Since $\dim(\mathcal{H}_{-4-\delta}(X_t, \Lambda^{0,2})) = 0$, it follows that \square_t is surjective. We can therefore choose $f_t \in C_{\delta+2}^{k+2,\alpha}(X_t, \Lambda^{0,2})$ such that $\bar{a}_t = \square_t f_t = \bar{\partial}_t\bar{\partial}_t^* f_t$ with

$$(7.18) \quad \|f_t\|_{C_{\delta+2}^{k+2,\alpha}} \leq C\|\bar{a}_t\|_{C_\delta^{k,\alpha}} \leq C'\|t\|_{C_\delta^{k,\alpha}}.$$

Let $\bar{\beta}_t = \bar{\partial}_t^* f_t$, and let $\omega_t = \omega_t^h + \bar{\partial}_t\beta_t$. Then $d\omega_t = \partial_t\omega_t^h - \bar{\partial}_ta_t = 0$. Choose ϵ_1 to be small enough, then ω_t is a closed positive (1,1)-form. \square

8. DEFORMATIONS OF SCALAR-FLAT KÄHLER METRICS

Notice that the previous sections did not use the scalar-flat assumption; this section is where we start using it.

8.1. The linearized operator. In previous sections, we have shown that there exists a family of decaying deformation of complex structure with dimension equal to $\dim(\mathcal{H}_{ess}(X, \Lambda^{0,1} \otimes \Theta))$, and there exists a smooth family of Kähler forms along the deformation. In this section, we will show that near an initial scalar-flat Kähler ALE metric ω_0 , there is a smooth family of scalar-flat Kähler ALE metrics under these deformations. The arguments in this section are inspired by [LS93, LS94].

Denote $S(\omega_0 + \sqrt{-1}\partial\bar{\partial}f)$ as the scalar curvature of X with metric $\omega_0 + \sqrt{-1}\partial\bar{\partial}f$. We now consider S as mapping between weighted Hölder spaces, with some $0 < \epsilon \ll 1$

$$(8.1) \quad \begin{aligned} S : C_\epsilon^{k,\alpha}(X) &\rightarrow C_{\epsilon-4}^{k-4,\alpha}(X) \\ f &\mapsto S(\omega_0 + \sqrt{-1}\partial\bar{\partial}f). \end{aligned}$$

Its linearization and adjoint are maps between:

$$(8.2) \quad L : C_\epsilon^{k,\alpha}(X) \rightarrow C_{\epsilon-4}^{k-4,\alpha}(X)$$

$$(8.3) \quad L^* : C_{-\epsilon}^{k,\alpha}(X) \rightarrow C_{-4-\epsilon}^{k-4,\alpha}(X).$$

By direct calculation,

$$(8.4) \quad L(f) := D S(\omega_0 + \sqrt{-1}\partial\bar{\partial}f) = -(\Delta^2 f + R_{i,\bar{j}} \nabla^i \nabla^{\bar{j}} f).$$

Define $\bar{\partial}^\# f = g_0^{i,\bar{j}} \bar{\partial}_{\bar{j}} f$. Also by direct calculation,

$$(8.5) \quad (\bar{\partial}\bar{\partial}^\#)^*(\bar{\partial}\bar{\partial}^\#)f = \Delta^2 f + R_{i,\bar{j}} \nabla^i \nabla^{\bar{j}} f + \nabla_k S \nabla^k f.$$

Clearly then, when ω_0 is scalar-flat, $L(f) = -(\bar{\partial}\bar{\partial}^\#)^*(\bar{\partial}\bar{\partial}^\#)(f)$, see [LS94].

The following lemma shows that the linearized mapping is surjective. This can also be interpreted as the nondegeneracy of the Futaki invariant, see [Fut88] and [LS94, Section 3.2].

Lemma 8.1. *For $0 < \epsilon \ll 1$, $k \geq 4$, L is surjective.*

Proof. Obviously, L is an elliptic operator, and the indicial roots of L are the integers. Consequently, by standard weighted space theory, L is Fredholm since δ is non-integral [LM85].

Let $f \in \ker(L^*)$, then $f = O(r^{-\epsilon})$ and $(\bar{\partial}\bar{\partial}^\#)^*(\bar{\partial}\bar{\partial}^\#)f = 0$. Since

$$(8.6) \quad f(\bar{\partial}\bar{\partial}^\#)^*(\bar{\partial}\bar{\partial}^\#)f = o(r^{-4}),$$

integrating by parts, It follows that $\bar{\partial}\bar{\partial}^\# f = 0$, so $\bar{\partial}^\# f$ is a holomorphic vector field on X which has a decay rate of $-(\epsilon - 1)$, and Proposition 3.3 then implies that $\bar{\partial}^\# f = 0$. Since f is a real-valued function, f must be constant; but since $f = O(r^{-\epsilon})$, we have $f \equiv 0$. We have shown that $\ker(L^*) = \{0\}$, so L is surjective. \square

Proposition 8.2. *For $0 < \epsilon \ll 1$ and $k \geq 4$, $\ker(L)$ is spanned by the constants.*

Proof. The operator L admits an expansion near infinity of the form $L = \Delta_{Euc}^2 + Q$, where Q represents lower order terms. Let

$$(8.7) \quad N(L, \tau) = \text{Index}(L : C_\tau^{k,\alpha}(X) \rightarrow C_{\tau-4}^{k-4,\alpha}(X)).$$

Then, by the relative index theorem of Lockhart-McOwen,

$$(8.8) \quad N(L, \epsilon) - N(L, -\epsilon) = N(\Delta_{Euc}^2, \epsilon) - N(\Delta_{Euc}^2, -\epsilon) = 2.$$

see [LM85]. The above argument shows that $\text{Ker}(L) \cap C_{-\epsilon}^{k,\alpha}(X) = \{0\}$, which implies that

$$(8.9) \quad \ker(L) \cap C_{\epsilon}^{k,\alpha}(X) = \{\text{constants}\}.$$

□

8.2. Construction of \mathfrak{F} . Now we start the construction of the class \mathfrak{F} . First, we show a weighted version $\partial\bar{\partial}$ -lemma on ALE surfaces. This is similar to [Joy00, Theorem 8.4.4], but with slightly different assumptions.

Lemma 8.3. *Consider the Kähler ALE surface (X, J_0, g_0) . Let $\zeta \in C_{\delta}^{k,\alpha}(X, \Lambda^{1,1})$ be a real $(1, 1)$ form, for some $k \geq 2$ and $\delta \in (-2, -1)$, satisfying $d\zeta = 0$. Then there exists $\theta \in C_{\delta+2}^{k+2,\alpha}(X)$ such that $\zeta = \zeta_h + \sqrt{-1}\partial\bar{\partial}\theta$, and $\|\theta\|_{C_{\delta+2}^{k+2,\alpha}} \leq C \cdot \|\zeta - \zeta_h\|_{C_{\delta}^{k,\alpha}}$, where $\zeta_h \in \mathcal{H}_{\delta}(X, \Lambda^{1,1})$.*

Proof. Since ζ is closed, can be decomposed as $\zeta = \zeta_h + \partial\eta + \bar{\partial}\xi$, where $\zeta_h \in \mathcal{H}_{\delta}(X, \Lambda^{1,1})$, $\eta \in C_{\delta+1}^{k+1,\alpha}(X, \Lambda^{0,1})$, and $\xi \in C_{\delta+1}^{k+1,\alpha}(X, \Lambda^{1,0})$. Since ζ is a real form, we can assume that $\eta = \bar{\xi}$. Consider the operator \square and its Fredholm adjoint \square^* :

$$(8.10) \quad \begin{aligned} \square &: C_{\delta+3}^{k+3,\alpha}(X, \Lambda^{0,1}) \rightarrow C_{\delta+1}^{k+1,\alpha}(X, \Lambda^{0,1}) \\ \square^* &: C_{-\delta-3}^{k+3,\alpha}(X, \Lambda^{0,1}) \rightarrow C_{-\delta-5}^{k+1,\alpha}(X, \Lambda^{0,1}) \end{aligned}$$

This implies that \square has finite-dimensional cokernel. Then we have

$$(8.11) \quad \eta = \eta_h + \square\gamma = \eta_h + \bar{\partial}\bar{\partial}^*\gamma + \bar{\partial}^*\bar{\partial}\gamma,$$

where $\gamma \in C_{\delta+3}^{k+3,\alpha}(X, \Lambda^{0,1})$ and η_h is the \square -harmonic part. Without loss of generality, assume $\eta_h = 0$. Since g_0 is Kähler, $\partial\bar{\partial}^*(\bar{\partial}\gamma) = -\bar{\partial}\bar{\partial}^*(\bar{\partial}\gamma) = 0$, so we can assume that $\bar{\partial}\gamma = 0$. Then $\zeta = \zeta_h + \sqrt{-1}\partial\bar{\partial}\theta'$, where $\theta' = 2\text{Im}(\bar{\partial}^*\gamma) \in C_{\delta+2}^{k+2,\alpha}(X)$.

Now consider the Fredholm operator

$$(8.12) \quad F = \sqrt{-1}\partial\bar{\partial} : C_{\delta+2}^{k+2,\alpha}(X) \rightarrow C_{\delta}^{k,\alpha}(X, \Lambda^{1,1})$$

Let $(\ker F^*)^{\perp}$ be the L^2 -complement of $\ker F^*$ in $C_{\delta}^{k,\alpha}(X, \Lambda^{1,1})$. The argument above shows that $\ker F^* = \ker \square^*$. Then $\zeta - \zeta_h \in (\ker F^*)^{\perp}$. By the same argument as in the proof of Theorem 7.2, we can show that there exists a $\theta \in C_{\delta+2}^{k+2,\alpha}(X)$, such that $\sqrt{-1}\partial\bar{\partial}\theta = \zeta - \zeta_h$, and $\|\theta\|_{C_{\delta+2}^{k+2,\alpha}} \leq C\|\zeta - \zeta_h\|_{C_{\delta}^{k,\alpha}}$ for some constant C . □

Theorem 8.4. *Let (X, J_0, g_0) be a scalar-flat Kähler ALE metric on a surface X . Then there is family \mathfrak{F} of scalar-flat Kähler metrics near g_0 , parametrized by B , that is, there is a differentiable mapping*

$$(8.13) \quad F : B_{\epsilon_1}^1 \times B_{\epsilon_2}^2 \rightarrow \mathcal{M}(X),$$

*into the space of smooth Riemannian metrics $\mathcal{M}(X)$ with $\mathfrak{F} = F(B_{\epsilon_1}^1 \times B_{\epsilon_2}^2)$. Furthermore, letting \mathfrak{G} denote the group of holomorphic isometries of (X, J_0, g_0) , the mapping F is equivariant with respect to the action of \mathfrak{G} . That is, for $\iota \in \mathfrak{G}$, and $(t, \rho) \in B_{\epsilon_1}^1 \times B_{\epsilon_2}^2$, we have $F(\iota^*t, \iota^*\rho) = \iota^*F(t, \rho)$.*

Proof. Recall that

$$(8.14) \quad B_{\epsilon_1}^1 = B_{\epsilon_1}(0) \subset \mathcal{H}_{ess}(X_0, \text{End}_a(TX_0)); \quad B_{\epsilon_2}^2 = B_{\epsilon_2}(0) \subset \mathcal{H}_\delta(X_0, \Lambda^{1,1}) \simeq H^{1,1}(X_0).$$

Let t denote the parameter of $B_{\epsilon_1}^1$. By Section 7, $\dim(\mathcal{H}_\delta(X_t, \Lambda^{p,q}))$ (using the metric determined by ω_t) is upper semicontinuous, and

$$(8.15) \quad \begin{aligned} \dim(\mathcal{H}_\delta(X_0, \Lambda^{1,1})) &= \dim(H^{1,1}(X_0)) \\ \dim(\mathcal{H}_\delta(X_0, \Lambda^{2,0})) &= \dim(\mathcal{H}_\delta(X_0, \Lambda^{0,2})) = 0. \end{aligned}$$

So for ϵ_1 sufficiently small,

$$(8.16) \quad \dim(\mathcal{H}_\delta(X_t, \Lambda^{2,0})) = \dim(\mathcal{H}_\delta(X_t, \Lambda^{0,2})) = 0,$$

and also

$$(8.17) \quad \begin{aligned} \dim(\mathcal{H}_\delta(X_t, \Lambda^{2,0})) &\geq \dim(H^{2,0}(X_t)) \\ \dim(\mathcal{H}_\delta(X_t, \Lambda^{0,2})) &\geq \dim(H^{0,2}(X_t)). \end{aligned}$$

This implies that $\dim(H^2(X_t)) = \dim(H^{1,1}(X_t))$ which is a topological invariant. Then

$$(8.18) \quad \begin{aligned} \dim(H^2(X_t)) &\leq \dim(\mathcal{H}_\delta(X_t, \Lambda^{1,1})) \leq \dim(\mathcal{H}_\delta(X_0, \Lambda^{1,1})) = \\ &= \dim(H^2(X_0)) = \dim(H^2(X_t)). \end{aligned}$$

This implies that $\dim(\mathcal{H}_\delta(X_t, \Lambda^{1,1})) = \dim(H^2(X_t))$ is constant, so choose a smooth family of isomorphisms ψ_t which map $\mathcal{H}_\delta(X_0, \Lambda^{1,1})$ to $\mathcal{H}_\delta(X_t, \Lambda^{1,1})$ for t sufficiently small. Note that \mathfrak{G} acts on $\mathcal{H}_\delta(X_0, \Lambda^{1,1})$, and from Theorem 7.2, \mathfrak{G} also acts on $\mathcal{H}_\delta(X_t, \Lambda^{1,1})$. Clearly we can choose ψ_t to be equivariant with respect to these actions.

It has been shown in Section 7 that there is a smooth family of Kähler forms ω_t . Let $\rho \in B_{\epsilon_2}^2$, define

$$(8.19) \quad \omega(t, \rho) = \omega_t + \psi_t(\rho)$$

Now consider the mapping

$$(8.20) \quad \mathcal{S} : B_{\epsilon_1}^1 \times B_{\epsilon_2}^2 \times C_\epsilon^{k,\alpha}(X) \rightarrow C_{\epsilon-4}^{k-4,\alpha}(X),$$

defined by

$$(8.21) \quad \mathcal{S} : (t, \rho, f) \mapsto S(\omega(t, \rho) + \sqrt{-1}\partial_t\bar{\partial}_t f).$$

We endow the domain with the product norm, where $B_{\epsilon_1}^1$ and $B_{\epsilon_2}^2$ have the L^2 -norm. By direct calculation, the linearization of \mathcal{S} at 0 is:

$$(8.22) \quad D\mathcal{S} = (*, \quad -Ric_h, \quad L)$$

where Ric_h is the harmonic part of Ricci form. Since L is surjective as shown in Lemma 8.3, $D\mathcal{S}$ is surjective by the lemma above. Next, we recall the expansion of the curvature tensor

$$(8.23) \quad Rm(g_0 + h) = Rm(g_0) + (g_0 + h)^{-1} * \nabla^2 h + (g_0 + h)^{-2} * \nabla h * \nabla h,$$

where $*$ denotes a various tensor contractions, and h is a symmetric tensor such that $g_0 + h$ is a Riemannian metric [GV16, Formula 3.40]. In our case, h can be written as

$$(8.24) \quad h(\cdot, \cdot) = (\omega(t, \rho) + \sqrt{-1}\partial_t\bar{\partial}_t f)(-J_t, \cdot) - \omega(0, 0)(-J_0, \cdot).$$

Next, by (7.5), we have the estimates

$$\begin{aligned}
 (8.25) \quad |h| &\leq C(|t| + |\rho| + |\nabla^2 f|) \leq C(r^\delta \|t\|_{C_\delta^{k,\alpha}} + r^\delta \|\rho\|_{C_\delta^{k,\alpha}} + r^{-2+\epsilon} \|f\|_{C_\epsilon^{k,\alpha}}) \\
 |\nabla h| &\leq C(|\nabla t| + |\nabla \rho| + |\nabla^3 f|) \leq C(r^{-1+\delta} \|t\|_{C_\delta^{k,\alpha}} + r^{-1+\delta} \|\rho\|_{C_\delta^{k,\alpha}} + r^{-3+\epsilon} \|f\|_{C_\epsilon^{k,\alpha}}) \\
 |\nabla^2 h| &\leq C(|\nabla^2 t| + |\nabla^2 \rho| + |\nabla^4 f|) \leq C(r^{-2+\delta} \|t\|_{C_\delta^{k,\alpha}} + r^{-2+\delta} \|\rho\|_{C_\delta^{k,\alpha}} + r^{-4+\epsilon} \|f\|_{C_\epsilon^{k,\alpha}}).
 \end{aligned}$$

Then by (8.23), we have

$$\begin{aligned}
 (8.26) \quad S(g_0 + h) - DS \cdot h &= C_0(g_0, h) * Rm(g_0) * h * h \\
 &\quad + C_1(g_0, h) * h * \nabla^2 h + C_2(g_0, h) * \nabla h * \nabla h,
 \end{aligned}$$

where $C_0(g_0, h)$, $C_1(g_0, h)$, and $C_2(g_0, h)$ are bounded functions, when h is sufficiently small. Without loss of generality, we can assume that $-2 + \epsilon < \delta$. Then

$$\begin{aligned}
 (8.27) \quad |S(g_0 + h) - DS \cdot h| &\leq C\{r^{-2}|h|^2 + |h| \cdot |\nabla^2 h| + |\nabla h| \cdot |\nabla h|\} \\
 &\leq C \cdot r^{-2+2\delta} \cdot (\|t\|_{C_\delta^{k,\alpha}} + \|\rho\|_{C_\delta^{k,\alpha}} + \|f\|_{C_\epsilon^{k,\alpha}})^2
 \end{aligned}$$

which implies that

$$(8.28) \quad \|S(g_0 + h) - DS \cdot h\|_{C_{-4+\epsilon}^{k,\alpha}} \leq C(\|t\|_{C_\delta^{k,\alpha}} + \|\rho\|_{C_\delta^{k,\alpha}} + \|f\|_{C_\epsilon^{k,\alpha}})^2.$$

Furthermore,

$$\begin{aligned}
 (8.29) \quad \|S(g_0 + h) - S(g_0 + h') - DS \cdot (h - h')\|_{C_{-4+\epsilon}^{k,\alpha}} &\leq \\
 &C \cdot (\|t - t'\|_{C_\delta^{k,\alpha}} + \|\rho - \rho'\|_{C_\delta^{k,\alpha}} + \|f - f'\|_{C_\epsilon^{k,\alpha}}) \\
 &\quad \cdot (\|t\|_{C_\delta^{k,\alpha}} + \|\rho\|_{C_\delta^{k,\alpha}} + \|f\|_{C_\epsilon^{k,\alpha}} + \|t'\|_{C_\delta^{k,\alpha}} + \|\rho'\|_{C_\delta^{k,\alpha}} + \|f'\|_{C_\epsilon^{k,\alpha}}).
 \end{aligned}$$

Since $B_{\epsilon_1}^1$ and $B_{\epsilon_2}^2$ are finite-dimensional, we can replace the corresponding norms by the L^2 -norm. This shows that \mathcal{S} satisfies the condition (4.1). Then by Lemma 4.1, the zero set of \mathcal{S} is in one-to-one correspondence with the kernel of $\mathcal{S}'(0)$. Since any solution of the nonlinear equation can be written as a kernel element plus a unique element in the image of a bounded right inverse for the linearized operator, the zero set of \mathcal{S} can also be written as a graph over $B_{\epsilon_1}^1 \times B_{\epsilon_2}^2$. That is, for any $(t, \rho) \in B_{\epsilon_1}^1 \times B_{\epsilon_2}^2$, there exists a unique $f(t, \rho)$, up to constants, of

$$(8.30) \quad \mathcal{S}(t, \rho, f(t, \rho)) = 0,$$

so a family of scalar-flat Kähler metrics over $B_{\epsilon_1}^1 \times B_{\epsilon_2}^2$ can be constructed as

$$(8.31) \quad F : (t, \rho) \mapsto g(t, \rho) = \left(\omega(t, \rho) + \sqrt{-1} \partial_t \bar{\partial}_t f(t, \rho) \right) (-J_t \cdot, \cdot),$$

where $g(0, 0) = g_0$. It is a straightforward consequence of the implicit function theorem that the mapping F is differentiable. Then the image of F gives us the family of scalar-flat Kähler metrics \mathfrak{F} , and the construction is clearly equivariant with respect to the action of \mathfrak{G} .

To finish the proof, we show that metrics in \mathfrak{F} are smooth. By the result of [TV05a], any Kähler constant scalar curvature metric satisfies an equation of the form

$$(8.32) \quad \Delta Ric = Rm * Ric,$$

where the right hand side denotes quadratic curvature contractions involving the full curvature tensor Rm and the Ricci tensor. By a Moser iteration method and regularity bootstrap argument in harmonic coordinates, it follows that g is smooth, see [TV05b, Theorem 6.4]. \square

9. VERSALITY AND UNIQUENESS OF THE MODULI SPACE

In this section, all norms are defined based on the initial Kähler metric g_0 . We let (g_1, J_1) be any scalar-flat Kähler metric satisfying $\|g_1 - g_0\|_{C_\delta^{k,\alpha}} < \epsilon_3$, (for some $k \geq 4$, $\delta \in (-2, -1)$, and ϵ_3 will be determined later). We begin with the following lemma.

Lemma 9.1. *For ϵ_3 sufficiently small, if $\|g_1 - g_0\|_{C_\delta^{k,\alpha}} < \epsilon_3$, then there exists a diffeomorphism $\Phi_0 : X \rightarrow X$ and constants C_1, C_2 depending upon g_0, J_0 so that*

$$(9.1) \quad \|\Phi_0^* g_1 - g_0\|_{C_\delta^{k,\alpha}} < C_1 \epsilon_3$$

$$(9.2) \quad \|\Phi_0^* J_1 - J_0\|_{C_\delta^{k,\alpha}} < C_2 \epsilon_3.$$

Proof. Let ∇_0 denote the covariant derivative of g_0 , and Γ_0, Γ_1 denote the Christoffel symbols of g_0, g_1 , respectively. Then $\|\Gamma_1 - \Gamma_0\|_{C_\delta^{k-1,\alpha}} < C\epsilon_3$, so in particular $|\Gamma_1 - \Gamma_0| < \frac{C\epsilon_3}{(1+r)^{-\delta+1}}$ as $r \rightarrow \infty$.

Since (g_i, J_i) is Kähler, $\nabla_i J_i = 0$ for $i = 0, 1$. We now estimate $|J_1 - J_0|(p)$ along any geodesic ray γ starting at p_0 . From [HL16, Lemma 1.1], both J_1 and J_0 have a finite limit along γ as $t \rightarrow \infty$, and furthermore

$$(9.3) \quad J_i = (J_i)_0 + O(r^{-2}),$$

where $(J_i)_0$ is a constant complex structure on \mathbb{R}^4 for $i = 0, 1$. We are assuming that $(J_0)_0 = J_{Euc}$. Clearly, there exists a diffeomorphism $\Phi_0 : X \rightarrow X$ so that

$$(9.4) \quad \Phi_0^* J_1 = J_{Euc} + O(r^{-2}),$$

as $r \rightarrow \infty$ such that (9.1) is satisfied. This can be done, for example, by connecting a rotation defined on the complement of a large ball $X \setminus B(p_0, 2R)$ to the identity transformation on $B(p_0, R)$ by smooth path of rotations of each sphere on the annulus $A(R, 2R)$. We then have

$$(9.5) \quad \begin{aligned} |\Phi_0^* J_1 - J_0|(p) &= |\Phi_0^* J_1 - J_0|(p) - \lim_{t \rightarrow \infty} |\Phi_0^* J_1 - J_0|(\gamma(t)) \\ &\leq \int_{s=r}^{\infty} |\nabla_0(\Phi_0^* J_1 - J_0)(\gamma(s))| ds \\ &\leq \int_{s=r}^{\infty} |\nabla_0 \Phi_0^* J_1(\gamma(s))| ds \\ &< \int_{s=r}^{\infty} \frac{C \cdot \epsilon_3}{(1+s)^{-\delta+1}} ds \leq \frac{C' \cdot \epsilon_3}{(1+r)^{-\delta}}. \end{aligned}$$

Since this estimate is true along any ray, it follows that $\|\Phi_0^* J_1 - J_0\|_{C_\delta^0} < C'\epsilon_3$. We can estimate the higher regularities in the same way, and (9.2) follows. \square

We next prove the ‘‘versality’’ of our family \mathfrak{F} . That is, for g_1 as above, there exists a diffeomorphism Φ such that $\Phi^* g_1$ is in the class \mathfrak{F} .

Theorem 9.2. *Let $-2 < \delta < -1$ be fixed. There exists an $\epsilon_3 > 0$ such that for any scalar-flat Kähler metric (g_1, J_1) satisfying $\|g_1 - g_0\|_{C_\delta^{k,\alpha}} \leq \epsilon_3$, there exists a diffeomorphism $\Phi : X \rightarrow X$ of the form $\Phi_0 \circ \Phi_{Y_1} \circ \Upsilon_Z \circ \Phi_{Y_2}$ where Φ_0 is as in Lemma 9.1, $Y_1, Y_2 \in C_{\delta+1}^{k+1,\alpha}(TX)$, and $Z \in \mathbb{W}$, such that $\Phi^*g_1 \in \mathfrak{F}$. Furthermore, there exists a constant C so that*

$$(9.6) \quad \|\Phi^*g_1 - g_0\|_{C_\delta^{k,\alpha}} \leq C\epsilon_3.$$

Proof. Let Φ_0 denote the diffeomorphism from Lemma 9.1. Then $\Phi_0^*J_1$ satisfies the assumptions of Theorem 6.4, so there exists a diffeomorphism $\tilde{\Phi}$ satisfying the properties stated in that theorem. In particular, $\tilde{\Phi}$ is of the form $\tilde{\Phi} = \Phi_{Y_1} \circ \Upsilon_Z \circ \Phi_{Y_2}$, where $Y_1, Y_2 \in C_{\delta+1}^{k+1,\alpha}(TX)$, and $Z \in \mathbb{W}$. The estimate (9.6) for $\Phi_0 \circ \tilde{\Phi}$ is then proved as follows. For Φ_{Y_1} , we estimate

$$(9.7) \quad \|\Phi_{Y_1}^* \Phi_0^*g_1 - g_0\|_{C_\delta^{k,\alpha}} \leq \|\Phi_{Y_1}^* \Phi_0^*g_1 - \Phi_0^*g_1\|_{C_\delta^{k,\alpha}} + \|\Phi_0^*g_1 - g_0\|_{C_\delta^{k,\alpha}} \leq C\epsilon_3.$$

This estimate holds since $\Phi_0^*g_1$ is also a scalar-flat Kähler metric which is smooth by the last observation in the proof of Theorem 8.4, so when ϵ_3 is sufficiently small, Lemma 5.1 holds for g_1 . The same argument applies to Φ_{Y_2} . The estimate (9.6) then follows from (9.7), (6.9) and Lemma 9.1.

Let ω_1 denote the Kähler form of Φ^*g_1 . It was shown in Section 8 that $\dim(H^{1,1}(X_t))$ is locally constant, which implies $H^{1,1}(X_0) \simeq H^{1,1}(X_t)$ for any $t \in B^1$. Then there exists a $\rho \in \mathcal{H}_\delta(X_0, \Lambda^{1,1})$ such that $[\omega(\phi_{ess}, \rho)] = [\omega_1] \in H^{1,1}(X_t)$, where $\omega(\phi_{ess}, \rho)$ is defined in (8.19). Let $g' = F(\phi_{ess}, \rho)$, and let ω' denote the corresponding Kähler form. Then by Lemma 8.3, $\omega' - \omega_1 = \sqrt{-1}\bar{\partial}_t \partial_t^* f$ for some potential function $f \in C_{\delta+2}^{k+2,\alpha}(X)$, and $\|f\|_{C_{\delta+2}^{k+2,\alpha}} \leq C\|\omega' - \omega_1\|_{C_\delta^{k,\alpha}}$.

As shown in the proof of Theorem 8.4, in a neighborhood of g_0 , a scalar-flat Kähler metric is uniquely determined by the Kähler class and complex structure. This implies that $g' = g_1$. \square

This completes the proof of Theorem 1.4. Next, we will complete the proof of Theorem 1.7. From Theorem 8.4, the mapping F is \mathfrak{G} -equivariant. That is, for $\iota \in \mathfrak{G}$, and $(t, \rho) \in B_{e_1}^1 \times B_{e_2}^2$, we have $F(\iota^*t, \iota^*\rho) = \iota^*F(t, \rho)$. This clearly implies that two elements in the same orbit of \mathfrak{G} are isometric, which proves the first statement in Theorem 1.7.

For the second statement in Theorem 1.7, we need the following lemma, which says that in the non-hyperkähler case, the dimension of \mathfrak{F} is the same as the dimension of the parameter space.

Lemma 9.3. *Let (X, g, J) be as above. If g is not hyperkähler then F is injective, so $\dim(\mathfrak{F}) = d = \dim(B_{e_1}^1 \times B_{e_2}^2)$.*

Proof. If $F(\phi_1, \rho_1) = F(\phi_2, \rho_2) = \tilde{g}$, then the metric \tilde{g} is Kähler with respect to the two complex structures J_1 and J_2 , corresponding to ϕ_1 and ϕ_2 , respectively.

We have $J_i \in \text{End}(TX)$ ($i = 1, 2$) such that $J_i^2 = -1$. Locally, J_i can be considered as an purely imaginary Hamiltonian number. Define

$$(9.8) \quad J_3 = \frac{J_1 J_2 - J_2 J_1}{\|J_1 J_2 - J_2 J_1\|},$$

where for $p \in X$,

$$(9.9) \quad \|J_i(p)\| = \sup_{0 \neq v \in T_p X} \left\{ \frac{\|J_i(v)\|_h}{\|v\|_h} \right\}.$$

Then $J_3 \in \text{End}(TX)$. We claim that J_3 is well-defined, $J_3^2 = -1$, J_1, J_2, J_3 are linearly independent. Clearly,

$$(9.10) \quad (J_1 J_2 - J_2 J_1)(J_1 J_2 - J_2 J_1) = -2 + (J_1 J_2 J_1 J_2 - J_2 J_1 J_2 J_1).$$

Since $J_1 J_2 J_1 J_2 - J_2 J_1 J_2 J_1$ is real and has norm which is < 2 , we have that

$$(9.11) \quad (J_1 J_2 - J_2 J_1)(J_1 J_2 - J_2 J_1) \neq 0,$$

so J_3 is well-defined, and $J_3 \cdot J_3 = -1$. If we write J_1, J_2 locally as Hamiltonian numbers $a_1 I + b_1 J + c_1 K$, $a_2 I + b_2 J + c_2 K$ (where $a_i, b_i, c_i \in \mathbb{R}$, $I^2 = J^2 = K^2 = -1$), then

$$(9.12) \quad J_1 J_2 - J_2 J_1 = (b_0 c_1 - b_1 c_0)I + (c_0 a_1 - c_1 a_0)J + (a_0 c_1 - a_1 c_0)K,$$

which is linearly independent with J_1 and J_2 . Then J_1, J_2, J_3 are linearly independent. As a result, $\{J_1, J_2, J_3\}$ gives a hyperkähler structure. Then we have proved that, if $\Gamma \not\subset \text{SU}(2)$, then $J_1 = J_2$, then by the proof of Theorem 8.4, F is injective. \square

The next proposition immediately implies the second part of Theorem 1.7.

Proposition 9.4. *For any two elements $g_1, g_2 \in \mathfrak{F}$ associated to complex structures J_1, J_2 such that $E_{J_0}^{-1}(J_1), E_{J_0}^{-1}(J_2)$ are divergence-free, if there exists a small diffeomorphism Φ_Y which is induced by the exponential map of a vector field $Y \in C_{\delta+1}^{k+1, \alpha}(TX)$, such that $g_1 = \Phi_Y^* g_2$, then g_1, g_2 are the same.*

Proof. First, assume X is non-hyperkähler. Since $\Phi_Y^* g_2 = g_1$, by Lemma 9.3, $J_1 = \Phi_Y^* J_2$. However by Lemma 5.3, there is a unique small diffeomorphism that gauges $E_{J_0}^{-1}(J_2)$ to be divergence-free. Then $\Phi_Y = \text{Id}$, $g_1 = g_2$.

When X is hyperkähler, then by rotating the hyperkähler sphere, there exists a J'_1 such that J'_1 is compatible with g_1 and $(X, \Phi_Y^* J_2)$ is biholomorphic to (X, J'_1) . Then by the same argument above, $g_1 = g_2$. \square

Since \mathfrak{F} is of finite dimension, and \mathfrak{G} is a compact group action on \mathfrak{F} , the dimension m of $\mathfrak{M} = \mathfrak{F}/\mathfrak{G}$ is well-defined. In the non-hyperkähler case, by Lemma 9.3,

$$(9.13) \quad m = d - (\text{the dimension of a maximal orbit of } \mathfrak{G}).$$

(For the hyperkähler case, recall Remark 1.11.)

10. DEFORMATIONS OF THE MINIMAL RESOLUTION

In this section, we prove Theorems 1.10 and 1.12.

10.1. **Harmonic representation of $H^1(X, \Theta)$.** Let X denote the minimal resolution of \mathbb{C}^2/Γ , where Γ is a finite subgroup of $U(2)$ without complex reflections. In the following, we want to construct a weighted version of Hodge theory, that links the sheaf cohomology with the decaying harmonic forms.

The divisor $E = \cup_i E_i$ is a union of irreducible components which are rational curves, with only normal crossing singularities. Let $Der_E(X)$ denote the sheaf dual to logarithmic 1-forms along E (see [Kaw78]). We note that $Der_E(X)$ is a locally free sheaf of rank 2, see [Wah75]. Away from E , this is clear. If $p \in E_i$, we can choose a holomorphic coordinate chart $\{z_1, z_2\}$ such that near p , $E_i = \{z_1 = 0\}$. Then local sections of $Der_E(X)$ are generated by $\{z_1 \frac{\partial}{\partial z_1}, \frac{\partial}{\partial z_2}\}$.

The short exact sequence

$$(10.1) \quad 0 \rightarrow Der_E(X) \rightarrow \Theta_X \rightarrow \mathcal{O}_E(E) \rightarrow 0,$$

induces an exact sequence of cohomologies

$$(10.2) \quad 0 \rightarrow H^1(X, Der_E(X)) \rightarrow H^1(X, \Theta) \rightarrow H^1(E, \mathcal{O}_E(E)) \rightarrow H^2(X, Der_E(X)).$$

Since E is composed of rational curves whose self-intersection numbers are negative, we have $H^0(E, \mathcal{O}_E(E)) = 0$.

Proposition 10.1. *We have the vanishing result: $H^1(X, Der_E(X)) = 0$.*

Proof. This is a result which follows from work of [BKR88, Bri68, Lau73, Wah75]. \square

By Siu's vanishing theorem ([Siu69]), since X is a non-compact σ -compact complex manifold, for any coherent analytic sheaf \mathcal{F} on X , the top degree sheaf cohomology $H^2(X, \mathcal{F})$ is trivial. Consequently, $H^2(X, Der_E(X)) = 0$, which gives us an isomorphism $H^1(\Theta_X) = H^1(\mathcal{O}_E(E))$. Let $-e_j$ be the self-intersection number of each rational curve $E_j \subset E$, and let k_Γ be the number of rational curves (which is equal to b_2). Then

$$(10.3) \quad \dim(H^1(X, \Theta)) = \sum_{j=1}^{k_\Gamma} (e_j - 1).$$

Theorem 10.2. *Let (X, g, J) denote the minimal resolution of \mathbb{C}^2/Γ with any ALE Kähler metric g of order $\tau > 1$. Then*

$$(10.4) \quad H^1(X, \Theta) \cong \mathcal{H}_{-3}(X, \Lambda^{0,1} \otimes \Theta) \cong \mathcal{H}_{ess}(X, \Lambda^{0,1} \otimes \Theta)$$

Proof. First, we consider the case on $\mathbb{C}^2 \setminus \{0\}$, and we compute $H^1(\mathbb{C}^2 \setminus \{0\}, \Theta)$. The domain $\mathbb{C}^2 \setminus \{0\}$ can be covered by two charts: $U_1 = \{z_1 \neq 0\}$ and $U_2 = \{z_2 \neq 0\}$. Note that U_1, U_2 are each isomorphic to $\mathbb{C} \times \mathbb{C}^*$, and $U_1 \cap U_2$ is isomorphic to $\mathbb{C}^* \times \mathbb{C}^*$. Then $\theta_1 \in \Gamma(U_1, \Theta)$ can be expanded into a Laurent series

$$(10.5) \quad \theta_1 = \sum_{i \in \mathbb{Z}_{\geq 0}, j \in \mathbb{Z}} a_{i,j}^1 z_1^i z_2^j \frac{\partial}{\partial z_1} + b_{i,j}^1 z_1^i z_2^j \frac{\partial}{\partial z_2},$$

$\theta_2 \in \Gamma(U_2, \Theta)$ can be expanded into

$$(10.6) \quad \theta_2 = \sum_{i \in \mathbb{Z}, j \in \mathbb{Z}_{\geq 0}} a_{i,j}^2 z_1^i z_2^j \frac{\partial}{\partial z_1} + b_{i,j}^2 z_1^i z_2^j \frac{\partial}{\partial z_2},$$

$\theta_{1,2} \in \Gamma(U_1 \cap U_2, \Theta)$ can be expanded into

$$(10.7) \quad \theta_{1,2} = \sum_{i \in \mathbb{Z}, j \in \mathbb{Z}} a_{i,j}^{1,2} z_1^i z_2^j \frac{\partial}{\partial z_1} + b_{i,j}^{1,2} z_1^i z_2^j \frac{\partial}{\partial z_2}.$$

Then any $\theta_{1,2} \in \Gamma(U_1 \cap U_2, \Theta)$ which cannot be represented as $\theta_1 - \theta_2$ must be of the form

$$(10.8) \quad \theta_{1,2} = \sum_{i \in \mathbb{Z}_-, j \in \mathbb{Z}_-} a_{i,j}^{1,2} z_1^i z_2^j \frac{\partial}{\partial z_1} + b_{i,j}^{1,2} z_1^i z_2^j \frac{\partial}{\partial z_2}.$$

By Dolbeault's lemma, we have $H^1(X, \Theta) \simeq H^0(X, \Lambda^{0,1} \otimes \Theta)$ ([GH94]). Using this, we transform the Čech cohomology element (10.8) to an element of Dolbeault cohomology $\phi \in \Gamma(\mathbb{C}^2 \setminus \{0\}, \Lambda^{0,1} \otimes \Theta)$. Let ρ_1, ρ_2 be a partition of unity, where ρ_1 is supported in U_1 and $\rho_1 = 1$ for $|z_1| > 1$; ρ_2 is supported in U_2 and $\rho_2 = 1$ for $|z_2| > 1$; $\rho_1 + \rho_2 = 1$. Then define $\phi \in \Gamma(\mathbb{C}^2 \setminus \{0\}, \Lambda^{0,1} \otimes \Theta)$ as

$$(10.9) \quad \phi = \begin{cases} \bar{\partial}(\rho_2 \cdot \theta_{1,2}) & \text{on } U_1 \\ -\bar{\partial}(\rho_1 \cdot \theta_{1,2}) & \text{on } U_2 \end{cases}.$$

The form ϕ is well-defined since $\bar{\partial}(\rho_2 \cdot \theta_{1,2}) + \bar{\partial}(\rho_1 \cdot \theta_{1,2}) = \bar{\partial}(\theta_{1,2}) = 0$ on $U_1 \cap U_2$. Furthermore, ϕ is decaying at infinity, since the degrees appearing in (10.8) are all negative.

Now for any closed form $\phi \in H^0(X, \Lambda^{0,1} \otimes \Theta)$, since $X - E$ is biholomorphic to $\mathbb{C}^2 \setminus \{0\}/\Gamma$, ϕ corresponds with a Γ -equivariant form $\phi' \in H^0(\mathbb{C}^2 \setminus \{0\}, \Lambda^{0,1} \otimes \Theta)$. Then $\phi' = \bar{\partial}\tau + \psi$, with ψ a closed form and decaying at infinity by the argument above. Since Γ is a finite group, by averaging τ by the group action, we can assume that τ is Γ -invariant. Let ρ be a cutoff function on X which equals to 0 in $B(R)$ and 1 outside of $B(2R)$. Then $\phi - \bar{\partial}(\rho \cdot \tau)$ is a closed decaying form, which is in the same class as ϕ .

This shows that the natural mapping

$$(10.10) \quad \mathcal{H}_{-3}(X, \Lambda^{0,1} \otimes \Theta) \rightarrow H^1(X, \Theta)$$

is surjective. We next show that this mapping is injective.

Let $\phi = \bar{\partial}\eta \in \mathcal{H}_{-3}(X, \Lambda^{0,1} \otimes \Theta)$, where $\eta \in \Gamma(X, \Theta)$. Then $[\phi] = 0 \in H^1(X, \Theta)$. We want to prove $\phi = 0$ by showing that $\phi = \bar{\partial}\xi$ for some $\xi = O(r^{-2})$. Let χ be a smooth cutoff function defined on X with compact support that contains E . Then $\phi = \bar{\partial}(\chi\eta) + \bar{\partial}((1-\chi)\eta)$. Since π is biholomorphic on $X \setminus E$, $(1-\chi)\eta$ can be pushed down by π and extended to ζ on \mathbb{C}^2 such that $\zeta = 0$ in a neighborhood of the origin. Also $\bar{\partial}\zeta = O(r^{-3})$ on \mathbb{C}^2 by the decaying rate of ϕ . Exactly as in (3.9) above, by the Poincaré lemma, there exists a $\sigma \in C_{-2}^\infty(\mathbb{C}^2, \Theta)$ such that $\bar{\partial}\sigma = \bar{\partial}\zeta$. Average σ by the group action of Γ such that σ is Γ -invariant, then $\sigma(0) = 0$. Note that σ is holomorphic in a neighborhood of 0. By standard argument, we can lift up σ to $\tilde{\sigma} \in \Gamma(X, \Theta)$, where $\tilde{\sigma} = O(|z|^{-2})$. Then we have $\phi = \bar{\partial}(\chi\eta + \tilde{\sigma})$. Since $\chi\eta + \tilde{\sigma} = O(r^{-2})$, $\phi = 0$.

Finally, we are going to prove that

$$(10.11) \quad \mathcal{H}_{-3}(X, \Lambda^{0,1} \otimes \Theta) \simeq \mathcal{H}_{ess}(X, \Lambda^{0,1} \otimes \Theta),$$

It is clear that $\mathcal{H}_{ess}(X, \Lambda^{0,1} \otimes \Theta) \subset \mathcal{H}_{-3}(X, \Lambda^{0,1} \otimes \Theta)$. To show the isomorphism, we need to show that $\mathbb{V} = \{0\}$. Let $Y \in \mathbb{W}$, by definition, $Y \in \mathcal{H}_1(X, \Theta)$. Then Y has the following

expansion as $r \rightarrow \infty$:

$$(10.12) \quad Y = \sum_{i,j} a_{i,j} z_i \frac{\partial}{\partial z_j} + \sum_k b_k \frac{\partial}{\partial z_k} + O(r^{-3+\epsilon})$$

where $a_{i,j}, b_k$ are constants. The decaying rate of $O(r^{-3+\epsilon})$ comes from the fact that \square has no indicial roots between -2 and 0 , and there is no $\bar{\partial}$ -closed kernel element corresponding to the root of -2 .

If Γ is nontrivial, then $b_k = 0$. Denote

$$(10.13) \quad \pi : X \mapsto \mathbb{C}^2/\Gamma$$

as the minimal resolution of \mathbb{C}^2/Γ . Since $Z = \sum_{i,j} a_{i,j} z_i \frac{\partial}{\partial z_j}$ is a holomorphic vector field on $\mathbb{C}^2/\Gamma \setminus \{0\}$ and $\lim_{r \rightarrow 0} Z = 0$, by standard argument, Z can be lifted up to a holomorphic vector field \tilde{Z} on X . Then $Y - \tilde{Z} \in \mathcal{H}_{-3}(X, \Theta)$. By using integration by parts, $Y - \tilde{Z}$ is holomorphic, then Y is holomorphic, $\bar{\partial}Y = 0$, and $V = 0$. Then $\mathcal{H}_{-3}(X, \Lambda^{0,1} \otimes \Theta) \simeq \mathcal{H}_{ess}(X, \Lambda^{0,1} \otimes \Theta)$.

Finally, if Γ is trivial, then X is biholomorphic to \mathbb{C}^2 , and the coordinate vector fields extend to global holomorphic vector fields, so these do not give any non-trivial elements of \mathbb{V} . \square

10.2. Proof of Theorem 1.12.

Proof of Theorem 1.12. In the following, X_t will stand for the triple (X, g_t, J_t) . Since the index of a strongly continuous family of Fredholm operators is constant, the index $P(t)$ of the operator P_t defined in Lemma 4.2 is locally constant along a path of ALE Kähler metrics. Consequently, $P(t)$ is constant along the path X_t . By the same argument as in Lemma 4.2, the cokernel of P_t is trivial for all $t \in [0, 1]$. Therefore P_t are all surjective Fredholm operators, with $\dim(\ker P_t) = \dim(\ker P_0) = \dim(\ker P_1)$. Therefore if there exists a path that connects X_1 with a minimal resolution X_0 , then $\dim(\mathcal{H}_{-3}(X_1, \Lambda^{0,1} \otimes \Theta)) = \dim(\mathcal{H}_{-3}(X_0, \Lambda^{0,1} \otimes \Theta))$. Since $\mathfrak{G}(X_1) = \{e\}$,

$$(10.14) \quad \dim_{\mathbb{R}}(\mathcal{H}_{ess}(X_1, \Lambda^{0,1} \otimes \Theta)) = \dim_{\mathbb{R}}(\mathcal{H}_{-3}(X_0, \Lambda^{0,1} \otimes \Theta)) - \dim(\mathfrak{G}(X_0)) = j_{\Gamma}.$$

In the non-hyperkähler case, Lemma 9.3 implies that

$$(10.15) \quad m(X_1) = j_{\Gamma} + b_2(X) = m_{\Gamma},$$

and Theorem 1.7 implies that the local moduli space of scalar-flat Kähler ALE metrics near g_1 is a smooth manifold of dimension m_{Γ} . As remarked above, the hyperkähler moduli space is constructed globally by Kronheimer, and $m_{\Gamma} = 3k - 3$. \square

REFERENCES

- [ALM15] Claudio Arezzo, Riccardo Lena, and Lorenzo Mazzieri, *On the Kummer construction for KCSC metrics*, arXiv.org:1507.05105, 2015.
- [ALM16] ———, *On the resolution of extremal and constant scalar curvature Kähler orbifolds*, Int. Math. Res. Not. IMRN (2016), no. 21, 6415–6452.
- [AP06] Claudio Arezzo and Frank Pacard, *Blowing up and desingularizing constant scalar curvature Kähler manifolds*, Acta Math. **196** (2006), no. 2, 179–228.
- [APS11] Claudio Arezzo, Frank Pacard, and Michael Singer, *Extremal metrics on blowups*, Duke Math. J. **157** (2011), no. 1, 1–51.

- [Bar86] Robert Bartnik, *The mass of an asymptotically flat manifold*, Comm. Pure Appl. Math. **39** (1986), no. 5, 661–693.
- [Biq06] Olivier Biquard, *Asymptotically symmetric Einstein metrics*, SMF/AMS Texts and Monographs, vol. 13, American Mathematical Society, Providence, RI; Société Mathématique de France, Paris, 2006, Translated from the 2000 French original by Stephen S. Wilson.
- [BKR88] Kurt Behnke, Constantin Kahn, and Oswald Riemenschneider, *Infinitesimal deformations of quotient surface singularities*, Singularities (Warsaw, 1985), Banach Center Publ., vol. 20, PWN, Warsaw, 1988, pp. 31–66.
- [BR15] Olivier Biquard and Yann Rollin, *Smoothing singular extremal Kähler surfaces and minimal Lagrangians*, Advances in Math. **285** (2015), 980–1024.
- [Bri68] Egbert Brieskorn, *Rationale Singularitäten komplexer Flächen*, Invent. Math. **4** (1967/1968), 336–358.
- [Cal85] Eugenio Calabi, *Extremal Kähler metrics. II*, Differential geometry and complex analysis, Springer, Berlin, 1985, pp. 95–114.
- [CS04] David M. J. Calderbank and Michael A. Singer, *Einstein metrics and complex singularities*, Invent. Math. **156** (2004), no. 2, 405–443.
- [CT94] Jeff Cheeger and Gang Tian, *On the cone structure at infinity of Ricci flat manifolds with Euclidean volume growth and quadratic curvature decay*, Invent. Math. **118** (1994), no. 3, 493–571.
- [Fut88] Akito Futaki, *Kähler-Einstein metrics and integral invariants*, Lecture Notes in Mathematics, vol. 1314, Springer-Verlag, Berlin, 1988.
- [GH94] Phillip Griffiths and Joseph Harris, *Principles of algebraic geometry*, Wiley Classics Library, John Wiley & Sons, Inc., New York, 1994, Reprint of the 1978 original.
- [GV16] Matthew J. Gursky and Jeff A. Viaclovsky, *Critical metrics on connected sums of Einstein four-manifolds*, Adv. Math. **292** (2016), 210–315. MR 3464023
- [HL16] Hans-Joachim Hein and Claude LeBrun, *Mass in Kähler geometry*, Comm. Math. Phys. **347** (2016), no. 1, 183–221.
- [Hon13] Nobuhiro Honda, *Deformation of LeBrun’s ALE metrics with negative mass*, Comm. Math. Phys. **322** (2013), no. 1, 127–148.
- [Hon14] ———, *Scalar flat Kähler metrics on affine bundles over $\mathbb{C}P^1$* , SIGMA Symmetry Integrability Geom. Methods Appl. **10** (2014), Paper 046, 25.
- [HRŞ16] Hans-Joachim Hein, Rares Rasdeaconu, and Ioana Şuvaina, *On the classification of ALE Kähler manifolds*, arXiv:1610.05239, 2016.
- [Joy00] Dominic D. Joyce, *Compact manifolds with special holonomy*, Oxford Mathematical Monographs, Oxford University Press, Oxford, 2000.
- [Kaw78] Yujiro Kawamata, *On deformations of compactifiable complex manifolds*, Math. Ann. **235** (1978), no. 3, 247–265.
- [Kro86] Peter Benedict Kronheimer, *Instantons gravitationnels et singularités de Klein*, C. R. Acad. Sci. Paris Sér. I Math. **303** (1986), no. 2, 53–55.
- [Kro89a] P. B. Kronheimer, *The construction of ALE spaces as hyper-Kähler quotients*, J. Differential Geom. **29** (1989), no. 3, 665–683.
- [Kro89b] ———, *A Torelli-type theorem for gravitational instantons*, J. Differential Geom. **29** (1989), no. 3, 685–697.
- [KS60] K. Kodaira and D. C. Spencer, *On deformations of complex analytic structures. III. Stability theorems for complex structures*, Ann. of Math. (2) **71** (1960), 43–76.
- [Lau73] Henry B. Laufer, *Taut two-dimensional singularities*, Math. Ann. **205** (1973), 131–164.
- [Li14] Chi Li, *On sharp rates and analytic compactifications of asymptotically conical Kähler metrics*, arXiv.org:1405.2433, 2014.
- [LM85] Robert B. Lockhart and Robert C. McOwen, *Elliptic differential operators on noncompact manifolds*, Ann. Scuola Norm. Sup. Pisa Cl. Sci. (4) **12** (1985), no. 3, 409–447.
- [LM08] Claude LeBrun and Bernard Maskit, *On optimal 4-dimensional metrics*, J. Geom. Anal. **18** (2008), no. 2, 537–564.

- [LS93] Claude LeBrun and Michael Singer, *Existence and deformation theory for scalar-flat Kähler metrics on compact complex surfaces*, Invent. Math. **112** (1993), no. 2, 273–313.
- [LS94] C. LeBrun and S. R. Simanca, *Extremal Kähler metrics and complex deformation theory*, Geom. Funct. Anal. **4** (1994), no. 3, 298–336.
- [LV14] Michael T. Lock and Jeff A. Viaclovsky, *A smörgåsbord of scalar-flat Kähler ALE surfaces*, arXiv.org:1410.6461 (to appear in Crelle’s Journal), 2014.
- [Mor07] Andrei Moroianu, *Lectures on Kähler geometry*, London Mathematical Society Student Texts, vol. 69, Cambridge University Press, Cambridge, 2007.
- [RS05] Yann Rollin and Michael Singer, *Non-minimal scalar-flat Kähler surfaces and parabolic stability*, Invent. Math. **162** (2005), no. 2, 235–270.
- [RS09] ———, *Constant scalar curvature Kähler surfaces and parabolic polystability*, J. Geom. Anal. **19** (2009), no. 1, 107–136.
- [Siu69] Yum-tong Siu, *Analytic sheaf cohomology groups of dimension n of n -dimensional noncompact complex manifolds*, Pacific J. Math. **28** (1969), 407–411.
- [Şuv12] Ioana Şuvaina, *ALE Ricci-flat Kähler metrics and deformations of quotient surface singularities*, Ann. Global Anal. Geom. **41** (2012), no. 1, 109–123.
- [TV05a] Gang Tian and Jeff Viaclovsky, *Bach-flat asymptotically locally Euclidean metrics*, Invent. Math. **160** (2005), no. 2, 357–415.
- [TV05b] ———, *Moduli spaces of critical Riemannian metrics in dimension four*, Adv. Math. **196** (2005), no. 2, 346–372.
- [Wah75] Jonathan M. Wahl, *Vanishing theorems for resolutions of surface singularities*, Invent. Math. **31** (1975), no. 1, 17–41.

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF WISCONSIN, MADISON, WI, 53706
E-mail address: jiyuan@math.wisc.edu

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF CALIFORNIA, IRVINE, CA, 92697
E-mail address: jviaclov@uci.edu