

CONVERGENCE PROOF FOR THE MULTIGRID METHOD OF THE NONLOCAL MODEL *

MINGHUA CHEN[†] AND WEIHUA DENG[‡]

Abstract. Recently, nonlocal models attract the wide interests of scientist. They mainly come from two applied scientific fields: peridynamics and anomalous diffusion. Even though the matrices of the algebraic equation corresponding the nonlocal models are usually Toeplitz (denote a_0 as the principal diagonal element, a_1 as the trailing diagonal element, etc). There are still some differences for the models in these two fields. For the model of anomalous diffusion, a_0/a_1 is uniformly bounded; most of the time, a_0/a_1 of the model for peridynamics is unbounded as the stepsize h tends to zero. Based on the uniform boundedness of a_0/a_1 , the convergence of the two-grid method is well established [Chan, Chang, and Sun, *SIAM J. Sci. Comput.*, 19 (1998), pp. 516–529; Pang and Sun, *J. Comput. Phys.*, 231 (2012), pp. 693–703; Chen, Wang, Cheng, and Deng, *BIT*, 54 (2014), pp. 623–647]. This paper provides the detailed proof of the convergence of the two-grid method for the nonlocal model of peridynamics. Some special cases of the full multigrid and the V-cycle multigrid are also discussed. The numerical experiments are performed to verify the convergence.

Key words. multigrid method, nonlocal model, Toeplitz matrices

AMS subject classifications. 65M55

1. Introduction. Ranging from characterizing peridynamics [25] to anomalous diffusion [20], the nonlocal models have been builded in more and more scientific fields. The used nonlocal operators include nonlocal diffusion operators [15], fractional Laplacian operators [19], Riesz fractional derivative [9, 34], and the Riesz tempered fractional derivative [10, 11]. In the field of anomalous diffusion, the nonlocal operators are derived in both the time and space directions. For the peridynamics, the nonlocal operators are just applied in the space direction. Mathematically, the nonlocal operators corresponding to these two applied fields have close connections, being examined in [14, 15]. The nonlocal operator mentioned in this paper has a finite range of nonlocal interactions measured by a horizon parameter δ [16, 25]. When $\delta \rightarrow 0$, the nonlocal effect diminishes and the local or classical partial differential equation (PDE) models are recovered, if the latter are well-defined. For $\delta > 0$, compared with classical PDE models, the complexities are introduced by the nonlocal interactions and the matrices of the resulting discrete systems are no longer sparse. In a series of recent studies [28, 29], the robust discretizations of the nonlocal models have been well developed. Based on the fast Toeplitz solver, the direct solution method for the resulting algebraic equation is discussed in [30]. In this work, we focus on the efficiently iterative solvers, especially, providing the strict convergence proof for the algorithm.

As is well know that the structure and conditioning of the resulting coefficient matrix of the numerical scheme play a key role for the effectiveness of the linear solver. For the nonlocal models, the associated stiffness matrices tend to be dense, and its condition number depends on both the nonlocal interaction kernel and horizon parameter δ . When δ is fixed, the condition number is bounded even the stepsize h tends

*This work was supported by NSFC 11271173.

[†]Corresponding author. School of Mathematics and Statistics, Gansu Key Laboratory of Applied Mathematics and Complex Systems, Lanzhou University, Lanzhou 730000, P.R. China (Email: chenmh@lzu.edu.cn).

[‡] School of Mathematics and Statistics, Gansu Key Laboratory of Applied Mathematics and Complex Systems, Lanzhou University, Lanzhou 730000, P.R. China (Email: dengwh@lzu.edu.cn).

to zero [1, 35]. However, if δ depends on h , the condition number will tend to infinity as h becomes smaller and smaller. So, it is interesting/necessary to understand the performance of the linear solver for the different types of the horizon δ . In particular, finding the effective linear solver should be paid much more attention for the case that δ depends on h . This work focuses on the multigrid method (MGM) with uniform convergence rate for various types of the horizon δ . MGM has often been shown to be the most efficient iterative method for numerically solving the PDEs [4, 18]. For the uniform convergence of the V-cycle MGM, one can refer to [8, 31, 32, 33] for the second order elliptic operator and [13] for the block tridiagonal matrix. For the multilevel matrix algebras like circulant, tau, Hartely, the V-cycle convergence is theoretically obtained by using some special interpolation operators [2, 3]. For works on the so-called full MGM, i.e., recursive application of the two-grid method (TGM) procedure, see, e.g., [6, 17, 24].

For nonlocal models, peridynamics and anomalous diffusion are two of the most successful applied fields. The common feature of the stiffness matrix of the resulting algebraic equation from the models is to have Toeplitz structure. For the Toeplitz matrix, we denote a_0 as the principal diagonal element, a_1 as the trailing diagonal element, etc. For the stiffness matrix of nonlocal model describing anomalous diffusion, a_0/a_1 is bounded; using this attribute, the uniform convergence of the TGM is theoretically obtained [6, 12, 21]. However, most of the time, a_0/a_1 is unbounded for the stiffness matrix of the peridynamic model. So, some new ideas must be introduced for proving the uniform convergence of the TGM for the nonlocal peridynamic model. This paper provides the detailed proof of the uniform convergence of TGM with unbounded a_0/a_1 . Furthermore, the special cases of the full MGM and V-cycle MGM are also discussed. The performed numerical experiments show the effectiveness of the MGM.

The outline of this paper is as follows. In the next section, we discuss the recently introduced finite difference discretizations of the nonlocal operator. The MGM algorithms are introduced in Section 3. In Section 4, we study the uniform convergence estimates of the TGM for the nonlocal model. Convergence of the full MGM and V-cycle MGM in a special case is analyzed in Section 5. To show the effectiveness of the presented schemes, results of numerical experiments are reported in Section 6. Finally, we conclude the paper with some remarks.

2. Preliminaries: numerical scheme and multigrid method. Before delivering the detailed convergence proof of the TGM, in this section, we review and discuss the numerical discretization and multigrid method for the nonlocal model (2.3).

2.1. The nonlocal operator and discretization scheme. In this subsection, we introduce the discretization of the nonlocal operator proposed in [28] and make some discussions on the generating of the matrix elements and treating of the non-homogeneous boundaries. Let Ω be a finite bar in \mathbb{R} . Without loss of generality, we take $\Omega = (0, b)$, $b > 0$. For $u = u(x) : \Omega \rightarrow \mathbb{R}$, the nonlocal operator \mathcal{L}_δ is defined by [28],

$$(2.1) \quad \mathcal{L}_\delta u(x) = \int_{B_\delta(x)} (u(y) - u(x)) \gamma_\delta(|x - y|) dy \quad \forall x \in \Omega$$

with $B_\delta(x) = \{y \in \mathbb{R} : |y - x| < \delta\}$ denoting a neighborhood centered at x of radius δ , which is the horizon parameter and represents the size of nonlocality; the symmetric nonlocal kernel $\gamma_\delta(|x - y|) = 0$ if $y \notin B_\delta(x)$.

The operator \mathcal{L} is used in both the time-dependent nonlocal volume-constrained diffusion problem [15]

$$(2.2) \quad \begin{cases} u_t - \mathcal{L}_\delta u = f_\delta & \text{on } \Omega, t > 0, \\ u(x, 0) = u_0 & \text{on } \Omega \cup \Omega_{\mathcal{I}}, \\ u = g & \text{on } \Omega_{\mathcal{I}}, t > 0, \end{cases}$$

for the function $u = u(x, t)$ and its steady-state counterpart

$$(2.3) \quad \begin{cases} -\mathcal{L}_\delta u = f_\delta & \text{on } \Omega, \\ u = g & \text{on } \Omega_{\mathcal{I}}, \end{cases}$$

where $u = g$ denotes a volumetric constraint imposed on a volume $\Omega_{\mathcal{I}}$ that has a nonzero volume and is made to be disjoint from Ω . For 1D case, we use $\Omega_{\mathcal{I}} = (-\delta, 0) \cup (b, b + \delta)$.

Let γ_δ be nonnegative and radial, i.e., $\gamma_\delta = \gamma_\delta(|y - x|) \geq 0$. As in [28], we can rewrite (2.1) as

$$(2.4) \quad \mathcal{L}_\delta u(x) = \int_0^\delta (u(x+s) - 2u(x) + u(x-s))\gamma_\delta(s)ds,$$

which makes the nonlocal operator as a *continuum difference operator*, or rather an average of finite difference operators over a continuum scale $(0, \delta)$ [28]. Assuming that $u(x)$ is regular enough, from (2.4) there exists

$$\mathcal{L}_\delta u(x) = Cu''(x) + \mathcal{O}\left(\int_0^\delta s^4 \gamma_\delta(s)ds\right),$$

where, C , is assumed to be positive and independent of δ , i.e.,

$$0 < C = \int_0^\delta s^2 \gamma_\delta(s)ds < \infty.$$

Now, we introduce and discuss the discretization scheme of (2.3). Denote the ratio of the horizon δ and the mesh size h as

$$(2.5) \quad R = \frac{\delta}{h} > 0 \quad \text{and} \quad r = \lfloor R \rfloor,$$

which plays an important role in nonlocal diffusion models. Here $\lfloor R \rfloor$ denotes the greatest integer that is less than or equal to R . And we will use $\lceil R \rceil$ to denote the least integer that is greater than or equal to R .

Let $\Omega = (0, b)$ with $\delta < b$ and the mesh points $x_i = ih$, $h = b/(N+1)$, $i \in \Omega_N = \{-r, \dots, 0, 1, \dots, N+1, \dots, N+1+r\}$, where r is defined by (2.5); and u_i as the numerical approximation of $u(x_i)$ and $f_{\delta,i} = f_\delta(x_i)$. Denote $I_p = ((p-1)h, ph)$ for $1 \leq p \leq r$, and $I_{r+1} = (rh, Rh) = (rh, \delta)$, and the piecewise linear basis function is

$$\phi_p(x) = \begin{cases} \frac{x - x_{p-1}}{h} & x \in [x_{p-1}, x_p], \\ \frac{x_{p+1} - x}{h} & x \in [x_p, x_{p+1}] \quad \text{for } i \in \Omega_N, \\ 0 & \text{otherwise.} \end{cases}$$

Eq. (2.1) can be rewritten as [28]

$$(2.6) \quad \mathcal{L}_\delta u(x) = \sum_{p=0}^{r+1} \int_0^\delta \frac{u(x+s) - 2u(x) + u(x-s)}{s} \phi_p(s) s \gamma_\delta(s) ds,$$

and an asymptotically compatible discretization of the nonlocal operator \mathcal{L}_δ has the following form

$$(2.7) \quad \begin{aligned} \mathcal{L}_\delta^h u_i &= \sum_{p=1}^r \frac{u_{i-p} - 2u_i + u_{i+p}}{ph} \int_{I_p \cup I_{p+1}} \phi_p(s) s \gamma_\delta(s) ds \\ &+ \frac{u_{i-r-1} - 2u_i + u_{i+r+1}}{(r+1)h} \int_{I_{r+1}} \phi_{r+1}(s) s \gamma_\delta(s) ds. \end{aligned}$$

Note that the above integral over I_{r+1} automatically vanishes when $r = R$.

The discretization of (2.3) then has the following form

$$(2.8) \quad \begin{aligned} -\mathcal{L}_\delta^h u_i &= f_{\delta,i}, \quad i \in \{1, 2, \dots, N\}, \\ u_i &= g_i, \quad i \in \{-r, \dots, 0\} \cup \{N+1, \dots, N+r+1\} \end{aligned}$$

with the following sketch that characterizes different variables:

$$\begin{array}{cccccc} \underbrace{[x_{-r} \cdots x_{-1}, x_0]}_{\text{boundary points}} & \underbrace{[x_1, x_2 \cdots x_r]}_{\text{interface points}} & \underbrace{[x_{r+1} \cdots x_{N-r}]}_{\text{internal points}} & \underbrace{[x_{N-r+1} \cdots x_N]}_{\text{interface points}} & \underbrace{[x_{N+1} \cdots x_{N+r+1}]}_{\text{boundary points}} \\ \underbrace{[g_{-r} \cdots g_{-1}, g_0]}_{\text{boundary values}} & \underbrace{[u_1, u_2 \cdots u_r]}_{\text{interface values}} & \underbrace{[u_{r+1} \cdots u_{N-r}]}_{\text{internal values}} & \underbrace{[u_{N-r+1} \cdots u_N]}_{\text{interface values}} & \underbrace{[g_{N+1} \cdots g_{N+r+1}]}_{\text{boundary values}} \end{array}$$

For notational convenience, we let

$$(2.9) \quad \begin{aligned} U_\delta^h &= [u_1, u_2, \dots, u_N]^T, \quad F_\delta^h = [f_{\delta,1}, f_{\delta,2}, \dots, f_{\delta,N}]^T; \\ F_{\mathcal{V},\delta}^h &= [f_{\mathcal{V},1}, f_{\mathcal{V},2}, \dots, f_{\mathcal{V},r+1}, 0, \dots, 0, f_{\mathcal{V},N-r}, \dots, f_{\mathcal{V},N}]^T. \end{aligned}$$

Thus, the finite difference scheme (2.8) can be recast as

$$(2.10) \quad A_\delta^h U_\delta^h = F_\delta^h + F_{\mathcal{V},\delta}^h,$$

where the stiffness matrix $A_\delta^h = \{a_{i,j}\}_{i,j=1}^N$ has a banded structure given by

$$(2.11) \quad a_{i_1, j_1} = a_{i_2, j_2} \quad \text{for } |i_1 - j_1| = |i_2 - j_2| \leq r+1, \quad \text{and } a_{i,j} = 0 \quad \text{otherwise.}$$

We denote $a_k = a_{i,j}$ with $k = |i-j|$. The auxiliary vector $F_{\mathcal{V},\delta}^h$ can be determined by the following matrix form

$$(2.12) \quad \begin{bmatrix} f_{\mathcal{V},1} \\ f_{\mathcal{V},2} \\ \vdots \\ f_{\mathcal{V},r} \\ f_{\mathcal{V},r+1} \end{bmatrix} = \begin{bmatrix} g_0 & g_{-1} & \cdots & g_{1-r} & g_{-r} \\ 0 & g_0 & \cdots & \cdots & g_{1-r} \\ \cdots & \cdots & \cdots & \cdots & \cdots \\ 0 & \cdots & \cdots & g_0 & g_{-1} \\ 0 & 0 & \cdots & 0 & g_0 \end{bmatrix} \begin{bmatrix} a_1 \\ a_2 \\ \vdots \\ a_r \\ a_{r+1} \end{bmatrix},$$

and

$$(2.13) \quad \begin{bmatrix} f_{\mathcal{V},N} \\ f_{\mathcal{V},N-1} \\ \vdots \\ f_{\mathcal{V},N-r+1} \\ f_{\mathcal{V},N-r} \end{bmatrix} = \begin{bmatrix} g_{N+1} & g_{N+2} & \ddots & g_{N+r} & g_{N+r+1} \\ 0 & g_{N+1} & \ddots & \ddots & g_{N+r} \\ \ddots & \ddots & \ddots & \ddots & \ddots \\ 0 & \ddots & \ddots & g_{N+1} & g_{N+2} \\ 0 & 0 & \ddots & 0 & g_{N+1} \end{bmatrix} \begin{bmatrix} a_1 \\ a_2 \\ \vdots \\ a_r \\ a_{r+1} \end{bmatrix}.$$

In the following, we focus on the special case where the kernel $\gamma_\delta(s)$ is taken to be a constant, i.e., $\gamma_\delta(s) = 3\delta^{-3}$ [28]. More general kernel types [1, 15, 28] can be similarly studied. The entries of the stiffness matrix A_δ^h can be explicitly documented by

Case 1: $R \leq 1$.

$$(2.14) \quad a_{|i-j|} = a_{i,j} = \begin{cases} \frac{2}{h^2}, & j = i, \\ -\frac{1}{h^2}, & |j - i| = 1, \\ 0, & \text{otherwise.} \end{cases}$$

Case 2: $R > 1$. Let $p = |j - i| \geq 1$. Then

$$(2.15) \quad a_{|i-j|} = a_{i,j} = \begin{cases} -2 \sum_{p=1}^{r+1} a_{i,p}, & j = i, \\ -\frac{3}{h^2 R^3}, & p = 1 : r - 1, \\ -\frac{3r - 1 + (R - r)(r^2 + rR - 2R^2 + 3r + 3R)}{2h^2 R^3 r}, & p = r, \\ -\frac{(R - r)(2R^2 - rR - r^2)}{2h^2 R^3 (r + 1)}, & p = r + 1, \\ 0, & \text{otherwise.} \end{cases}$$

If we insert $R \leq 1$ into (2.15), which reduces to (2.14).

2.2. Multigrid method. Let the finest mesh points $x_i = ih$, $h = b/(N + 1)$. Define the multiple level of grids [13, 33]

$$(2.16) \quad \mathcal{M}_m = \left\{ x_i^m = \frac{i}{2^m} b, i = 1 : N_m \right\} \quad \text{with } N_m = 2^m - 1, m = 1 : J,$$

where \mathcal{M}_m represents not only the grid with grid spacing $h_m = 2^{(J-m)}h$, but also the space of vectors defined on that grid. The classical restriction operator I_m^{m-1} and prolongation operator I_{m-1}^m are, respectively, defined by

$$(2.17) \quad \nu^{m-1} = I_m^{m-1} \nu^m \quad \text{with } \nu_i^{m-1} = \frac{1}{4} (\nu_{2i-1}^m + 2\nu_{2i}^m + \nu_{2i+1}^m), \quad i = 1 : N_{m-1};$$

and

$$(2.18) \quad \nu^m = I_{m-1}^m \nu^{m-1} \quad \text{with} \quad I_{m-1}^m = 2(I_m^{m-1})^T.$$

We use the coarse grid operators defined by the Galerkin approach [23, p. 455]

$$(2.19) \quad A_{m-1} = I_{m-1}^{m-1} A_m I_{m-1}^m, \quad m = 1 : J;$$

and for all intermediate $(m, m-1)$ coarse grids we apply the correction operators [22, p. 87]

$$(2.20) \quad T^m = I_m - I_{m-1}^m A_{m-1}^{-1} I_m^{m-1} A_m = I_m - I_{m-1}^m P_{m-1}$$

with

$$P_{m-1} = A_{m-1}^{-1} I_m^{m-1} A_m.$$

We choose the damped Jacobi iteration matrix by [5, p. 9]

$$(2.21) \quad K_m = I - S_m A_m \quad \text{with} \quad S_m := S_{m,\omega} = \omega D_m^{-1}$$

with a weighting factor $\omega \in (0, 1/3]$, and D_m is the diagonal of A_m .

A multigrid process can be regarded as defining a sequence of operators $B_m : \mathcal{M}_m \mapsto \mathcal{M}_m$ which is an approximate inverses of A_m in the sense that $\|I - B_m A_m\|$ is bounded away from one. We list the following V-cycle multigrid algorithm [33]: Algorithm 1. If $m = 2$, the resulting Algorithm 1 is TGM [13, 33].

Algorithm 1 V-cycle Multigrid Algorithm: Define $B_1 = A_1^{-1}$. Assume that $B_{m-1} : \mathcal{M}_{m-1} \mapsto \mathcal{M}_{m-1}$ is defined. We shall now define $B_m : \mathcal{M}_m \mapsto \mathcal{M}_m$ as an approximate iterative solver for the equation associated with $A_m \nu^m = f_m$.

- 1: Pre-smooth: Let $S_{m,\omega}$ be defined by (2.21) and $\nu_0^m = 0, l = 1 : m_1$

$$\nu_l^m = \nu_{l-1}^m + S_{m,\omega_{pre}}(f_m - A_m \nu_{l-1}^m).$$

- 2: Coarse grid correction: $e^{m-1} \in \mathcal{M}_{m-1}$ is the approximate solution of the residual equation $A_{m-1} e = I_m^{m-1}(f_m - A_m \nu_{m_1}^m)$ by the iterator B_{m-1} :

$$e^{m-1} = B_{m-1} I_m^{m-1}(f_m - A_m \nu_{m_1}^m).$$

- 3: Post-smooth: $\nu_{m_1+1}^m = \nu_{m_1}^m + I_{m-1}^m e^{m-1}$ and $l = m_1 + 2 : m_1 + m_2$

$$\nu_l^m = \nu_{l-1}^m + S_{m,\omega_{post}}(f_m - A_m \nu_{l-1}^m).$$

- 4: Define $B_m f_m = \nu_{m_1+m_2}^m$.
-

3. Convergence of TGM for nonlocal model. Now, we start to prove the convergence of the TGM for nonlocal model. First, we give some Lemmas that will be used.

LEMMA 3.1. [7, p. 5] Given $n \times n$ symmetric matrices P and Q and let P' be a principal submatrix of P of order $n - 1$. Then, for $m = 1, 2, \dots, n$,

$$(3.1) \quad \lambda_k(P) + \lambda_1(Q - P) \leq \lambda_k(Q) \leq \lambda_k(P) + \lambda_n(Q - P),$$

$$(3.2) \quad \lambda_1(P) \leq \lambda_1(P') \leq \lambda_2(P) \leq \lambda_2(P') \leq \dots \leq \lambda_{n-1}(P') \leq \lambda_n(P),$$

$$(3.3) \quad \lambda_{\min}(P) = \lambda_1(P) = \min_{x \neq 0} \frac{x^T P x}{x^T x}, \quad \lambda_{\max}(P) = \lambda_n(P) = \max_{x \neq 0} \frac{x^T P x}{x^T x}.$$

DEFINITION 3.2. [7, p. 13] Let $n \times n$ Toeplitz matrix T_n be of the following form:

$$T_n = \begin{bmatrix} t_0 & t_{-1} & \cdots & t_{2-n} & t_{1-n} \\ t_1 & t_0 & t_{-1} & \cdots & t_{2-n} \\ \vdots & t_1 & t_0 & \ddots & \vdots \\ t_{n-2} & \cdots & \ddots & \ddots & t_{-1} \\ t_{n-1} & t_{n-2} & \cdots & t_1 & t_0 \end{bmatrix};$$

i.e., $t_{i,j} = t_{i-j}$ and T_n is constant along its diagonals. Assume that the diagonals $\{t_k\}_{k=-n+1}^{n-1}$ are the Fourier coefficients of a function f , i.e.,

$$t_k = \frac{1}{2\pi} \int_{-\pi}^{\pi} f(x) e^{-ikx} dx.$$

Then the function f is called the generating function of T_n .

LEMMA 3.3. [7, p. 13-15] (Grenander-Szegö theorem) Let T_n be given by above matrix with a generating function f , where f is a 2π -periodic continuous real-valued functions defined on $[-\pi, \pi]$. Let $\lambda_{\min}(T_n)$ and $\lambda_{\max}(T_n)$ denote the smallest and largest eigenvalues of T_n , respectively. Then we have

$$f_{\min} \leq \lambda_{\min}(T_n) \leq \lambda_{\max}(T_n) \leq f_{\max},$$

where f_{\min} and f_{\max} is the minimum and maximum values of $f(x)$, respectively. Moreover, if $f_{\min} < f_{\max}$, then all eigenvalues of T_n satisfy

$$f_{\min} < \lambda(T_n) < f_{\max},$$

for all $n > 0$. In particular, if $f_{\min} > 0$, then T_n is positive definite.

LEMMA 3.4. Let the discrete Laplacian-like operators $\{L_j\}_{j=1}^{N-1}$ be defined by

$$(3.4) \quad L_j = \begin{bmatrix} 2 & \overbrace{\cdots}^{j-1 \text{ zeros}} & -1 & & \\ \cdots & \ddots & \ddots & \ddots & \\ -1 & \ddots & \ddots & \ddots & -1 \\ & \ddots & \ddots & \ddots & \\ & & -1 & \ddots & 2 \end{bmatrix}_{N \times N} \quad \text{with} \quad 1 \leq j \leq N - 1.$$

Then, the smallest eigenvalues of L_j satisfy

$$\lambda_1(L_j) \geq 4 \sin^2 \left(\frac{\pi}{2(\lceil N/j \rceil + 1)} \right), \quad j = 1, 2, \dots, N - 1.$$

Moreover, if N/j is an integer

$$\lambda_k(L_j) = 4 \sin^2 \left(\frac{k\pi}{2(N/j+1)} \right), \quad k = 1, 2, \dots, N/j.$$

Proof. Let $\nu_1^{k,1} = [\nu_1^k, \nu_2^k, \dots, \nu_N^k]^T$ be the associated eigenvector with the tridiagonal matrix L_1 . It is well known that its eigenvalues are given by [26, p. 702]

$$\lambda_{k,1} = 4 \sin^2 \left(\frac{k\pi}{2(N+1)} \right), \quad k = 1, 2, \dots, N.$$

Define

$$\nu_i^{k,j} = [\underbrace{0 \cdots 0}_{i-1}, \nu_1^k, \underbrace{0 \cdots 0}_{j-i}, \underbrace{0 \cdots 0}_{i-1}, \nu_2^k, \underbrace{0 \cdots 0}_{j-i}, \dots, \underbrace{0 \cdots 0}_{i-1}, \nu_N^k, \underbrace{0 \cdots 0}_{j-i}]^T, \quad i = 1, 2, \dots, j.$$

Then, for the matrix L_j with dimension $(jN) \times (jN)$, we have

$$\{L_j\}_{(jN) \times (jN)} \nu_i^{k,j} = \lambda_{k,1} \nu_i^{k,j} = 4 \sin^2 \left(\frac{k\pi}{2(N+1)} \right) \nu_i^{k,j}, \quad i = 1, 2, \dots, j,$$

leading to all eigenvalues with multiplicity j and eigenvectors of $\{L_j\}_{(jN) \times (jN)}$. A dimension rescaling then shows that

$$(3.5) \quad \lambda_k(\{L_j\}_{N \times N}) := \lambda_{k,j} = 4 \sin^2 \left(\frac{k\pi}{2(N/j+1)} \right), \quad k = 1, 2, \dots, N/j$$

if N/j is an integer.

If N/j is not an integer, we extend N to \tilde{N} such that \tilde{N}/j is an integer, i.e.,

$$\tilde{N}/j := \lceil N/j \rceil = \frac{N + j - \text{mod}(N, j)}{j},$$

where $\text{mod}(N, j)$ means the remainder of division of N by j .

From (3.5) and (3.2), we obtain

$$\lambda_1(\{L_j\}_{\tilde{N} \times \tilde{N}}) = 4 \sin^2 \left(\frac{\pi}{2(\tilde{N}/j+1)} \right) = 4 \sin^2 \left(\frac{\pi}{2(\lceil N/j \rceil + 1)} \right),$$

and

$$\lambda_1(\{L_j\}_{N \times N}) \geq \lambda_1(\{L_j\}_{\tilde{N} \times \tilde{N}}) = 4 \sin^2 \left(\frac{\pi}{2(\lceil N/j \rceil + 1)} \right).$$

The proof is completed. \square

LEMMA 3.5. *Let the matrix A_δ^h be defined by (2.11) and (2.15) on a finite bar $\Omega = (0, b)$, $b > 0$. Let $\delta = ch^\beta$, $\beta \geq 0$, $h \rightarrow 0$ and $c > 0$. Then*

$$\lambda_{\min}(A_\delta^h) \geq \frac{1}{27b^2}.$$

Proof. According the definition of L_j given in Lemma 3.4, we can recast (2.11) with its elements defined by (2.15) as

$$(3.6) \quad A_\delta^h = -a_1 L_1 - a_2 L_2 \cdots - a_{r+1} L_{r+1} = - \sum_{j=1}^{r+1} a_j L_j,$$

where $\{a_j\}$ are entries on different off-diagonals. We should check the following two cases: $r \leq 1$ and $r \geq 2$.

Case 1: $r \leq 1$. From (2.14) and (3.6), we obtain

$$A_\delta^h = -a_1 L_1,$$

which means that

$$\lambda_{\min}(A_\delta^h) = -a_1 \lambda_{\min}(L_1) = \frac{1}{h^2} \cdot 4 \sin^2 \left(\frac{\pi}{2(N+1)} \right) = \frac{1}{h^2} \cdot 4 \sin^2 \left(\frac{\pi h}{2b} \right) = \frac{\pi^2}{b^2} + \mathcal{O}(h^2).$$

Case 2: $r \geq 2$. Using

$$\frac{2}{\pi} x \leq \sin(x) \leq x, \quad x \in \left[0, \frac{\pi}{2} \right],$$

we obtain

$$\begin{aligned} \lambda_{\min}(A_\delta^h) &\geq - \sum_{j=1}^{r+1} a_j \lambda_{\min}(L_j) \geq \frac{1}{h^2} \frac{3}{R^3} \sum_{j=1}^{r-1} 4 \sin^2 \left(\frac{\pi}{2(\lceil N/j \rceil + 1)} \right) \\ &\geq \frac{1}{h^2} \frac{3}{R^3} \sum_{j=1}^{r-1} 4 \sin^2 \left(\frac{\pi}{2(N/j + 2)} \right) \geq \frac{1}{h^2} \frac{3}{R^3} \sum_{j=1}^{r-1} 4 \sin^2 \left(\frac{j\pi}{6N} \right) \\ &\geq \frac{1}{h^2} \frac{3}{R^3} \sum_{j=1}^{r-1} 4 \sin^2 \left(\frac{j\pi h}{6b} \right) \geq \frac{4}{3b^2 R^3} \sum_{j=1}^{r-1} j^2 \\ &\geq \frac{4}{3b^2 (r+1)^3} \cdot \frac{1}{6} \cdot \frac{r+1}{3} \cdot \frac{r+1}{2} \cdot \frac{2(r+1)}{2} = \frac{1}{27b^2}. \end{aligned}$$

The proof is completed. \square

REMARK 3.1. From (3.6), we know that the discrete nonlocal operator A_δ^h can be viewed as the superposition of discrete Laplacian-like operators $\{L_j\}_{j=1}^{r+1}$; and it reduces to the classical discrete Laplacian operator when $r = 0$ or 1.

LEMMA 3.6. Let the matrix A_δ^h be defined by (2.11) and (2.15) on a finite bar $\Omega = (0, b)$, $b > 0$. Let $\delta = ch^\beta$, $\beta \geq 0$, $h \rightarrow 0$ and $c > 0$. Then, there exists the bound of the condition number

$$\text{cond}(A_\delta^h) = \frac{\lambda_{\max}(A_\delta^h)}{\lambda_{\min}(A_\delta^h)} \leq c_* \min\{\delta^{-2}, h^{-2}\},$$

where c_* is a positive constant.

Proof. Case 1: $\beta > 1$. From (2.5), there exists

$$R = \frac{\delta}{h} = ch^{\beta-1} \leq c_0 \quad \text{with } c_0 \text{ a constant.}$$

If $R \leq 1$, from (2.14) and (3.6), we obtain

$$A_\delta^h = -a_1 L_1,$$

which means that

$$\lambda_{\max}(A_\delta^h) = -a_1 \lambda_{\max}(L_1) = \frac{1}{h^2} \cdot 4 \sin^2 \left(\frac{N\pi}{2(N+1)} \right) \leq \frac{4}{h^2}.$$

If $1 < R \leq c_0$, according to (3.6), (3.1), (3.4) and (2.15), there exists

$$\lambda_{\max}(A_\delta^h) \leq -\sum_{j=1}^{r+1} a_j \lambda_{\max}(L_j) \leq \frac{1}{h^2} \frac{3}{R^3} \sum_{j=1}^{r+1} 4 \leq \frac{12(R+1)}{R^3} \frac{1}{h^2} \leq \frac{12(c_0+1)}{h^2}.$$

Case 2: $0 \leq \beta \leq 1$. Using (2.15), we obtain

$$(3.7) \quad a_0 = -2 \sum_{m=1}^{r+1} a_{i,m} \leq 2 \sum_{m=1}^{r+1} \frac{3}{h^2 R^3} \leq \frac{12}{\delta^2}.$$

From the Gerschgorin theorem [27, p. 133], the eigenvalues λ of the matrix A_δ^h satisfy

$$\lambda_{\max}(A_\delta^h) \leq 2a_0 \leq \frac{24}{\delta^2}.$$

From Lemma 3.5, and the discussions of Case 1 and Case 2, the desired result is obtained. \square

Since the matrix A_δ^h is symmetric positive definite, we can define the following inner products

$$(u, v)_D = (Du, v), \quad (u, v)_A = (Au, v), \quad (u, v)_{AD^{-1}A} = (Au, Av)_{D^{-1}},$$

where for convenience we have dropped the explicit notational dependence on h and δ so that $A := A_J = A_\delta^h$ and D is its diagonal. Here (\cdot, \cdot) is the usual Euclidean inner product.

LEMMA 3.7. [22, p. 84] *Let A_J be a symmetric positive definite. If $\eta \leq \omega(2 - \omega\eta_0)$ with $\eta_0 \geq \lambda_{\max}(D_J^{-1}A_J)$, then the damped Jacobi iteration with relaxation parameter $0 < \omega < 2/\eta_0$ satisfies*

$$(3.8) \quad \|K_J \nu^J\|_{A_J}^2 \leq \|\nu^J\|_{A_J}^2 - \eta \|A_J \nu^J\|_{D_J^{-1}}^2 \quad \forall \nu^J \in \mathcal{M}_J.$$

LEMMA 3.8. [22, p. 89] *Let A_J be a symmetric positive definite matrix and K_J satisfies (3.8) and*

$$(3.9) \quad \min_{\nu^{J-1} \in \mathcal{M}_{J-1}} \|\nu^J - I_{J-1}^J \nu^{J-1}\|_{D_J}^2 \leq \kappa \|\nu^J\|_{A_J}^2 \quad \forall \nu^J \in \mathcal{M}_J$$

with $\kappa > 0$ independent of ν^J . Then, $\kappa \geq \eta > 0$ and the convergence factor of TGM satisfies

$$\|K_J T^J\|_{A_J} \leq \sqrt{1 - \eta/\kappa} \quad \forall \nu^J \in \mathcal{M}_J.$$

We present the convergence results of TGM in Theorem 3.9 and Theorem 3.11. The first part of the proof of Theorem 3.9 follows the traditional idea [6, 12, 21], but just the convergence result for $\beta \geq 1$ is obtained; we use a different technique to prove the case $\beta = 0$. After using the new idea, the convergence results for $\beta \geq 0$ are got, being proposed in Theorem 3.11 .

THEOREM 3.9. *Let $A_J = A_\delta^h$ be defined by (2.10) and (2.15) on a finite bar $\Omega \in (0, b)$, where $\delta = ch^\beta$, $\beta \geq 0$, $h \rightarrow 0$ and $c > 0$. Then K_J satisfies (3.8) and the convergence factor of the TGM satisfies*

$$\|K_J T^J\|_{A_J} \leq \begin{cases} \sqrt{1 - \eta/c_0} < 1 & \text{with } \beta \geq 1, c_0 = \max(1, 2c), \\ \sqrt{1 - \eta c^2/(648b^2)} < 1 & \text{with } \beta = 0, \end{cases}$$

where $\eta \leq 2\omega(1 - \omega)$ with $0 < \omega < 1$.

Proof. Since $\lambda_{\max}(D_J^{-1} A_J) \leq \eta_0$ with $\eta_0 = 2$, it leads to $0 < \omega < 2/\eta_0 = 1$. From Lemma 3.7, we conclude that K_J satisfies (3.8) with $\eta \leq 2\omega(1 - \omega)$. Let

$$\nu^J = (\nu_1, \nu_2, \dots, \nu_N)^T \in \mathcal{M}_J, \quad \nu^{J-1} = (\nu_2, \nu_4, \dots, \nu_{N-1})^T \in \mathcal{M}_{J-1},$$

and $\nu_0 = \nu_{N+1} = 0$ with $N = 2^J - 1$ in (2.16). From [6, 12, 21] and (3.6), we have

$$(3.10) \quad \|\nu^J - I_{J-1}^J \nu^{J-1}\|_{D_J}^2 \leq a_0 \sum_{i=1}^N (\nu_i^2 - \nu_i \nu_{i+1}),$$

$$(3.11) \quad \sum_{i=1}^N \nu_i^2 \geq \sum_{i=1}^N |\nu_i \nu_{i+1}|,$$

and

$$(3.12) \quad \|\nu^J\|_{A_J}^2 = (\nu^J, A_J \nu^J) \geq (\nu^J, -a_1 L_1 \nu^J) = -2a_1 \sum_{i=1}^N (\nu_i^2 - \nu_i \nu_{i+1}),$$

where a_0 and a_1 are given in (2.15). According to (3.10) and (3.12), there exists

$$(3.13) \quad \|\nu^J - I_{J-1}^J \nu^{J-1}\|_{D_J}^2 \leq \frac{a_0}{-2a_1} \|\nu^J\|_{A_J}^2.$$

Next we prove that (3.9) holds.

Case 1: $\beta \geq 1$. Using (2.6), there exists

$$R = \frac{\delta}{h} = ch^{\beta-1} \leq c \text{ as } h \leq 1.$$

When $R \leq 1$, from (2.14), it leads to $\kappa = \frac{a_0}{-2a_1} = 1$; then from Lemma 3.8,

$$(3.14) \quad \|K_J T^J\|_{A_J} \leq \sqrt{1 - \eta}.$$

When $1 < R \leq c$, using (2.15) and (3.7), we have

$$(3.15) \quad \begin{aligned} a_1 &= -\frac{1}{h^2} \cdot \frac{3}{R^3} = -\frac{3}{\delta^3} h = -\frac{3}{c^3} h^{1-3\beta}, \\ a_0 &= -2 \sum_{m=1}^{r+1} a_{i,m} \leq \frac{12}{\delta^2} = \frac{12}{c^2} h^{-2\beta}. \end{aligned}$$

It leads to

$$(3.16) \quad \kappa = \frac{a_0}{-2a_1} \leq 2ch^{\beta-1},$$

i.e., $\kappa \leq 2c$ as $h \leq 1$. Thus we obtain

$$(3.17) \quad \|K_J T^J\|_{A_J} \leq \sqrt{1 - \eta/(2c)} < 1.$$

Combining (3.14) and (3.17), it yields

$$\|K_J T^J\|_{A_J} \leq \sqrt{1 - \eta/c_0} \quad \text{with } \beta \geq 1, c_0 = \max(1, 2c).$$

Case 2: $\beta = 0$, i.e., $\delta = c$. Since $\kappa \rightarrow \infty$ as $\beta = 0$ in the estimate (3.16), we need to look for an estimate of the other form. From (3.15), we obtain $a_0 \leq \frac{12}{c^2}$. Using (3.11), it yields

$$(3.18) \quad \|\nu^J\|^2 = \sum_{i=1}^N \nu_i^2 = \frac{1}{2} \sum_{i=1}^N \nu_i^2 + \frac{1}{2} \sum_{i=1}^N \nu_i^2 \geq \frac{1}{2} \sum_{i=1}^N (\nu_i^2 - \nu_i \nu_{i+1}).$$

From Lemma 3.5, we have

$$(3.19) \quad \|\nu^J\|_{A_J}^2 = (A_J \nu^J, \nu^J) \geq \lambda_{\min}(A_J) \|\nu^J\|^2 \geq \frac{1}{27b^2} \|\nu^J\|^2.$$

According to (3.10), (3.18), (3.19) and (3.15), we get

$$\|\nu^J - I_{J-1}^J \nu^{J-1}\|_{D_J}^2 \leq a_0 \sum_{i=1}^N (\nu_i^2 - \nu_i \nu_{i+1}) \leq 2a_0 \|\nu^J\|^2 \leq 54b^2 a_0 \|\nu^J\|_{A_J}^2 \leq \kappa \|\nu^J\|_{A_J}^2$$

with $\kappa = 54b^2 a_0 \leq \frac{648b^2}{c^2}$. Hence

$$\|K_J T^J\|_{A_J} \leq \sqrt{1 - \eta c^2 / (648b^2)} \quad \text{with } \beta = 0.$$

The proof is completed. \square

In the works [6, 12, 21], the convergence factor of the two-grid method is uniformly bounded below one independent of h by estimating $\kappa = \frac{a_0}{2|a_1|} < \infty$, $a_1 \neq 0$. Since $\kappa = \frac{a_0}{2|a_1|} \rightarrow \infty$ as $\beta \in [0, 1)$ in the estimate (3.16), next we need to use a different idea to prove the case: $\beta \geq 0$.

LEMMA 3.10. Let $A = \sum_{j=1}^n L_j$ and $B = nL_1$ with $n \geq 1$, where L_j are defined by

(3.4). Then $2A - B$ is a positive definite matrix.

Proof. The generating functions of A and B are

$$f_A(x) = 2n - 2 \sum_{k=1}^n \cos(kx) \quad \text{and} \quad f_B(x) = 2n(1 - \cos x),$$

respectively. Since $f_A(x)$ and $f_B(x)$ are the even function and 2π -periodic continuous real-valued functions defined on $[-\pi, \pi]$, we just need to consider on $[0, \pi]$. Moreover

$$(3.20) \quad 2f_A(x) - f_B(x) = 4ng(x)$$

with

$$(3.21) \quad g(x) = \cos^2 \frac{x}{2} - \frac{1}{n} \sum_{k=1}^n \cos(kx), \quad x \in [0, \pi].$$

Next we prove $g(x) \geq 0$. If $x = 0$, it yields $g(x) = 0$. Denote

$$(3.22) \quad \varphi_n(x) := \frac{1}{n} \sum_{k=1}^n \cos(kx) = \frac{\sin \frac{2n+1}{2}x - \sin \frac{x}{2}}{2n \sin \frac{x}{2}}, \quad x \in (0, \pi].$$

Case 1: $0 < \frac{2n+1}{2}x \leq \pi$. We can rewrite (3.22) as

$$\varphi_n(x) = \frac{x}{2 \sin \frac{x}{2}} \phi(y) \quad \text{with} \quad \phi(y) = \frac{\sin y - \sin \frac{x}{2}}{y - \frac{x}{2}}, \quad y = \frac{2n+1}{2}x, \quad x \in \left(\frac{x}{2}, \pi\right].$$

It is easy to prove that $\phi(y)$ decreases with respect to y , which implies

$$\varphi_n(x) \leq \varphi_{n-1}(x) \leq \cdots \leq \varphi_1(x) = \frac{\sin \frac{3}{2}x - \sin \frac{x}{2}}{2 \sin \frac{x}{2}} = \cos x < \cos^2 \frac{x}{2},$$

i.e., $g(x) > 0$.

Case 2: $\pi \leq \frac{2n+1}{2}x \leq 2\pi + \frac{x}{2}$. Since $\varphi_n(x) \leq 0 < \cos^2 \frac{x}{2}$, it yields $g(x) > 0$.

Case 3: $\frac{2n+1}{2}x \geq 2\pi + \frac{x}{2}$. Using (3.22), there exists

$$\varphi_n(x) \leq \frac{1 - \sin \frac{x}{2}}{2n \sin \frac{x}{2}} = \frac{\cos^2 \frac{x}{2}}{2n \sin \frac{x}{2} (1 + \sin \frac{x}{2})} < \cos^2 \frac{x}{2},$$

since

$$2n \sin \frac{x}{2} \left(1 + \sin \frac{x}{2}\right) \geq \frac{4\pi}{x} \sin \frac{x}{2} \geq \frac{4\pi}{x} \cdot \frac{2}{\pi} \cdot \frac{x}{2} = 4.$$

According to the above equations and Lemma 3.3, the desired result is obtained. \square

THEOREM 3.11. *Let $A_J = A_\delta^h$ be defined by (2.10) and (2.15) on a finite bar $\Omega \in (0, b)$, where $\delta = ch^\beta$, $\beta \geq 0$, $h \rightarrow 0$ and $c > 0$. Then K_J satisfies (3.8) and the convergence factor of the TGM satisfies*

$$\|K_J T^J\|_{A_J} \leq \sqrt{1 - \eta/6} < 1, \quad \beta \geq 0,$$

where $\eta \leq 2\omega(1 - \omega)$ with $0 < \omega < 1$.

Proof. By the proof of Theorem 3.9, we know that K_J satisfies (3.8) and

$$(3.23) \quad \|\nu^J - I_{J-1}^J \nu^{J-1}\|^2 \leq \sum_{i=1}^N (\nu_i^2 - \nu_i \nu_{i+1}) = \frac{1}{2} (L_1 \nu^J, \nu^J).$$

Case 1: $r \leq 1$. From (2.14) and (3.6), we obtain

$$A_J = A_\delta^h = -a_1 L_1,$$

which means that

$$(3.24) \quad \|\nu^J\|_{A_J}^2 = (A_J \nu^J, \nu^J) = (-a_1 L_1 \nu^J, \nu^J).$$

According to (3.23) and (3.24), there exists

$$\|\nu^J - I_{J-1}^J \nu^{J-1}\|_{D_J}^2 \leq \frac{a_0}{2} (L_1 \nu^J, \nu^J) = \frac{a_0}{-2a_1} \|\nu^J\|_{A_J}^2 = \|\nu^J\|_{A_J}^2.$$

Thus from Lemma 3.8, we obtain

$$\|K_J T^J\|_{A_J} \leq \sqrt{1-\eta} \quad \text{with } r \leq 1.$$

Case 2: $r \geq 2$. According to (3.6), (2.15), Lemma 3.10 and (3.23), we obtain

$$\begin{aligned} \|\nu^J\|_{A_J}^2 &= (A_J \nu^J, \nu^J) \geq -a_1 \left(\sum_{j=1}^{r-1} L_j \nu^J, \nu^J \right) \geq -\frac{a_1(r-1)}{2} (L_1 \nu^J, \nu^J) \\ &\geq -\frac{a_1(r+1)}{6} (L_1 \nu^J, \nu^J) \geq \frac{a_0}{12} (L_1 \nu^J, \nu^J) \geq \frac{a_0}{6} \|\nu^J - I_{J-1}^J \nu^{J-1}\|_{D_J}^2 \\ &= \frac{1}{6} \|\nu^J - I_{J-1}^J \nu^{J-1}\|_{D_J}^2. \end{aligned}$$

Thus from Lemma 3.8, we have

$$\|K_J T^J\|_{A_J} \leq \sqrt{1-\eta/6} \quad \text{with } r \geq 2.$$

The proof is completed. \square

4. Convergence of the full MGM and V-cycle MGM with $\delta = ch$. We extend the convergence results of TGM given in the above section to the full MGM and V-cycle MGM in case that $\delta = ch$ with c being an appropriate natural number. First, we introduce some lemmas. We will use the notion of M-matrix, which is a positive definite matrix with positive entries on the diagonal and nonpositive off-diagonal entries. And another notion is called weakly diagonal dominant [5, p. 3], if the diagonal element of a matrix is at least as large as the sum of the off-diagonal elements in the same row or column.

LEMMA 4.1 ([13]). *Let $A^{(1)} = \{a_{i,j}^{(1)}\}_{i,j=1}^\infty$ with $a_{i,j}^{(1)} = a_{|i-j|}^{(1)}$ be a symmetric Toeplitz matrix and $A^{(k)} = L_h^H A^{(k-1)} L_h^H$ with $L_h^H = 4I_k^{k-1}$ and $L_h^H = (L_h^H)^T$. Then $A^{(k)}$ can be computed by (4.1). Here*

$$(4.1) \quad \begin{aligned} a_0^{(k)} &= (4C_k + 2^{k-1})a_0^{(1)} + \sum_{m=1}^{2 \cdot 2^{k-1} - 1} {}_0C_m^k a_m^{(1)}; \\ a_1^{(k)} &= C_k a_0^{(1)} + \sum_{m=1}^{3 \cdot 2^{k-1} - 1} {}_1C_m^k a_m^{(1)}; \\ a_j^{(k)} &= \sum_{m=(j-2)2^{k-1}}^{(j+2)2^{k-1}-1} {}_jC_m^k a_m^{(1)} \quad \forall j \geq 2 \quad \forall k \geq 2 \end{aligned}$$

with $C_k = 2^{k-2} \cdot \frac{2^{2k-2}-1}{3}$. And

$${}_0C_m^k = \begin{cases} 8C_k - (m^2 - 1)(2^k - m) & \text{for } m = 1 : 2^{k-1}; \\ \frac{1}{3}(2^k - m - 1)(2^k - m)(2^k - m + 1) & \text{for } m = 2^{k-1} : 2 \cdot 2^{k-1} - 1; \end{cases}$$

$${}_1C_m^k =$$

$$\begin{cases} 2C_k + m^2 \cdot 2^{k-1} - \frac{2}{3}(m-1)m(m+1) & \text{for } m = 1 : 2^{k-1}; \\ 2C_k + (2^k - m)^2 \cdot 2^{k-1} - \frac{2}{3}(2^k - m - 1)(2^k - m)(2^k - m + 1) \\ - \frac{1}{6}(m - 2^{k-1} - 1)(m - 2^{k-1})(m - 2^{k-1} + 1) & \text{for } m = 2^{k-1} : 2 \cdot 2^{k-1}; \\ \frac{1}{6}(3 \cdot 2^{k-1} - m - 1)(3 \cdot 2^{k-1} - m)(3 \cdot 2^{k-1} - m + 1) & \text{for } m = 2 \cdot 2^{k-1} : 3 \cdot 2^{k-1} - 1; \end{cases}$$

and for $j \geq 2$,

$${}_jC_m^k = \begin{cases} \varphi_1 & \text{for } m = (j-2)2^{k-1} : (j-1)2^{k-1}; \\ \varphi_2 & \text{for } m = (j-1)2^{k-1} : j2^{k-1}; \\ \varphi_3 & \text{for } m = j2^{k-1} : (j+1)2^{k-1}; \\ \varphi_4 & \text{for } m = (j+1)2^{k-1} : (j+2)2^{k-1} - 1, \end{cases}$$

where

$$\varphi_1 = \frac{1}{6}(m - (j-2)2^{k-1} - 1)(m - (j-2)2^{k-1})(m - (j-2)2^{k-1} + 1);$$

$$\begin{aligned} \varphi_2 &= 2C_k + (m - (j-1)2^{k-1})^2 \cdot 2^{k-1} \\ &\quad - \frac{1}{6}(j2^{k-1} - m - 1)(j2^{k-1} - m)(j2^{k-1} - m + 1) \\ &\quad - \frac{2}{3}(m - (j-1)2^{k-1} - 1)(m - (j-1)2^{k-1})(m - (j-1)2^{k-1} + 1); \end{aligned}$$

$$\begin{aligned} \varphi_3 &= 2C_k + ((j+1)2^{k-1} - m)^2 \cdot 2^{k-1} \\ &\quad - \frac{1}{6}(m - j2^{k-1} - 1)(m - j2^{k-1})(m - j2^{k-1} + 1) \\ &\quad - \frac{2}{3}((j+1)2^{k-1} - m - 1)((j+1)2^{k-1} - m)((j+1)2^{k-1} - m + 1); \end{aligned}$$

$$\varphi_4 = \frac{1}{6}((j+2)2^{k-1} - m - 1)((j+2)2^{k-1} - m)((j+2)2^{k-1} - m + 1).$$

LEMMA 4.2. Let $A^{(1)} = \{a_{i,j}^{(1)}\}_{i,j=1}^\infty$ with $a_{i,j}^{(1)} = a_{|i-j|}^{(1)}$ be a weakly diagonally dominant symmetric Toeplitz M -matrix and $D_{(k)}$ be the diagonal of the matrix $A^{(k)}$, where $A^{(k)} = I_k^{k-1} A^{(k-1)} I_{k-1}^k$. Then

$$1 \leq \lambda_{\max} \left(D_{(k)}^{-1} A^{(k)} \right) < 3.$$

In particular,

$$1 \leq \lambda_{\max} \left(D_{(k)}^{-1} A^{(k)} \right) \leq 2 \quad \text{if } a_1^{(k)} \leq 0.$$

Proof. We take $A^{(k)} = L_h^H A^{(k-1)} L_H^h$ with $L_h^H = 4I_k^{k-1}$ and $L_H^h = (L_h^H)^T$, and denote

$$A^{(k)} = \{a_{i,j}^{(k)}\}_{i,j=1}^\infty \quad \text{with} \quad a_{i,j}^{(k)} = a_{|i-j|}^{(k)}, \quad \forall k \geq 1.$$

By mathematical induction, we prove the estimates

$$(4.2) \quad 2 \sum_{j=1}^n |a_j^{(s)}| \leq a_0^{(s)} \quad \text{if} \quad a_1^{(s)} \leq 0, \quad s \geq 2, n \geq 1;$$

and

$$(4.3) \quad 2 \sum_{j=1}^n |a_j^{(s)}| \leq \frac{2C_s + 6C_s + 2^{s-1}}{4C_s + 2^{s-1}} a_0^{(s)} < 2a_0^{(s)} \quad \text{if} \quad a_1^{(s)} \geq 0.$$

For $s = 2$ and $a_1^{(2)} \leq 0$. From (4.1) and (2.13) of [13], we get $a_j^{(2)} \leq 0$, $j \geq 2$ and

$$\begin{aligned} 2 \sum_{j=1}^n |a_j^{(2)}| &= -2 \sum_{j=1}^n a_j^{(2)} \\ &= -2a_0^{(1)} + 8 \sum_{j=1}^{2n-1} |2a_j^{(1)}| + 8a_1^{(1)} + 2a_2^{(1)} + 2a_{2n}^{(1)} + 8a_{2n+1}^{(1)} + 14a_{2n+2}^{(1)} \\ &\leq 6a_0^{(1)} + 8a_1^{(1)} + 2a_2^{(1)} = a_0^{(2)}, \end{aligned}$$

where we use the property of $A^{(1)}$, being a weakly diagonally dominant M-matrix.

For $s = 2$ and $a_1^{(2)} \geq 0$. According to (4.1) and (2.13) of [13], there exists $a_j^{(2)} \leq 0$, $j \geq 2$ and

$$\begin{aligned} 2 \sum_{j=1}^n |a_j^{(2)}| &= 2a_1^{(2)} - 2 \sum_{j=2}^n a_j^{(2)} \\ &= 2a_0^{(1)} + 8 \sum_{j=1}^{2n-1} |2a_j^{(1)}| + 24a_1^{(1)} + 26a_2^{(1)} + 16a_3^{(1)} + 4a_4^{(1)} \\ &\quad + 2a_{2n}^{(1)} + 8a_{2n+1}^{(1)} + 14a_{2n+2}^{(1)} \\ &\leq \frac{10}{6} (6a_0^{(1)} + 8a_1^{(1)} + 2a_2^{(1)}) = \frac{5}{3} a_0^{(2)}. \end{aligned}$$

Then (4.2) and (4.3) hold for $s = 2$. Suppose that (4.2) and (4.3) hold for $s = 2, 3, \dots, k-1$. Next we prove (4.3) holds for $s = k$.

Taking $a_1^{(k)} \geq 0$ and using (4.1) and $a_j^{(k)} \leq 0$, $j \geq 2$, we have

$$\begin{aligned}
2 \sum_{j=1}^n |a_j^{(k)}| &= 2a_1^{(k)} - 2 \sum_{j=2}^n a_j^{(k)} \\
&= 2C_k a_0^{(1)} + 2 \sum_{m=1}^{2^{k-1}-1} ({}_1C_m^k - {}_2C_m^k) a_m^{(1)} + 2 \sum_{m=2^{k-1}}^{2 \cdot 2^{k-1}-1} ({}_1C_m^k - {}_2C_m^k - {}_3C_m^k) a_m^{(1)} \\
&\quad + 2 \sum_{m=2 \cdot 2^{k-1}}^{3 \cdot 2^{k-1}-1} ({}_1C_m^k - {}_2C_m^k - {}_3C_m^k - {}_4C_m^k) a_m^{(1)} \\
&\quad - 2 \sum_{j=2}^{n-3} \sum_{m=(j+1)2^{k-1}}^{(j+2)2^{k-1}-1} ({}_jC_m^k + {}_{j+1}C_m^k + {}_{j+2}C_m^k + {}_{j+3}C_m^k) a_m^{(1)} \\
&\quad - 2 \sum_{m=(n-1)2^{k-1}}^{n2^{k-1}-1} ({}_{n-2}C_m^k + {}_{n-1}C_m^k + {}_nC_m^k) a_m^{(1)} \\
&\quad - 2 \sum_{m=n2^{k-1}}^{(n+1)2^{k-1}-1} ({}_{n-1}C_m^k + {}_nC_m^k) a_m^{(1)} - 2 \sum_{m=(n+1)2^{k-1}}^{(n+2)2^{k-1}-1} {}_nC_m^k a_m^{(1)} \\
&\leq \frac{2C_k + 6C_k + 2^{k-1}}{4C_k + 2^{k-1}} a_0^{(k)} < 2a_0^{(k)},
\end{aligned}$$

where we use ${}_jC_m^k + {}_{j+1}C_m^k + {}_{j+2}C_m^k + {}_{j+3}C_m^k = 6C_k + 2^{k-1}$ and ${}_1C_m^k - {}_2C_m^k + 6C_k + 2^{k-1} \geq {}_0C_m^k$, $m = 1 : 2^{k-1}$ and ${}_1C_m^k - {}_2C_m^k - {}_3C_m^k + 6C_k + 2^{k-1} \geq {}_0C_m^k$, $m = 2^{k-1} : 2 \cdot 2^{k-1}$ and the property of $A^{(1)}$. Similarly, we can prove (4.2). Then we obtain

$$1 \leq \lambda_{\max} \left(D_{(k)}^{-1} A^{(k)} \right) < 3, \quad a_1^{(k)} \geq 0 \quad \text{and} \quad 1 \leq \lambda_{\max} \left(D_{(k)}^{-1} A^{(k)} \right) \leq 2, \quad a_1^{(k)} \leq 0.$$

The proof is completed. \square

4.1. The operation count and storage requirement. We now discuss the computation count and the required storage for the MGM of nonlocal problem (2.10).

From (2.11), we know that the matrix $A_h = A_\delta^h$ is a symmetric banded Toeplitz matrix with bandwidth $r + 1$ (obviously less than N). Then, we only need to store the first column of A_h , which have N parameters, instead of the full matrix A_h with N^2 entries. From Lemma 4.2, we know that $\{A_k\}$ is the symmetric Toeplitz matrix with the grid sizes $\{2^{J-k}h\}_{k=1}^{J-1}$, i.e., \mathcal{M}_k requires $2^{J-k}N$ storage. Adding these terms together, we

$$\text{Storage} = \mathcal{O}(N) \cdot \left(1 + \frac{1}{2} + \frac{1}{2^2} + \dots + \frac{1}{2^{J-1}} \right) = \mathcal{O}(N).$$

As for operation counts, the matrix-vector product associated with the matrix A_h is a discrete convolution. While the cost of a direct product is $\mathcal{O}(rN)$, the cost of using the FFT would lead to $\mathcal{O}(N \log(N))$ [7]. Moreover, from (4.1), we know that the bandwidth of $\{A_k\}$ is not bigger than the bandwidth of A_h . Hence, with the change of r , we may adopt different strategies. Thus, the total per V-cycle MGM operation count is

$$\mathcal{O}(\min\{rN, N \log N\}) \cdot \left(1 + \frac{1}{2} + \dots + \frac{1}{2^{J-1}} \right) = \begin{cases} \mathcal{O}(N), & r \text{ is bounded,} \\ \mathcal{O}(N \log(N)), & \text{in the worst case.} \end{cases}$$

4.2. Convergence of the full MGM with $\delta = \mathbf{ch}$. We further consider the convergence of the full MGM (recursive application of the TGM procedure). More precisely we show that the constants η in Lemma 3.7 and κ in Lemma 3.8 do not depend on the levels; this level independence is crucial for the convergence theory of the full MGM [6, 24]. In the following, we consider the simple algebraic systems, but these algebraic arguments are mostly motivated from analytic considerations.

LEMMA 4.3. *Let $A^{(1)} = c_1 L_1 + c_2 L_2 + c_3 L_3$, $c_i > 0$, $i = 1, 2, 3$ and $A^{(k)} = I_k^{k-1} A^{(k-1)} I_{k-1}^k$. Then*

$$\|K_k T^k\|_{A_k} \leq \sqrt{1 - \eta/\kappa} < 1 \quad \forall 1 \leq k \leq J,$$

where $A_k = A^{(J-k+1)}$, $\kappa = \max\left\{1 + \frac{c_2 + c_3}{c_1}, 3\right\}$, and $\eta \leq 2\omega(1 - \omega)$ with $0 < \omega < 1$.

Proof. Let $A^{(k)} = L_h^H A^{(k-1)} L_h^h$ with $L_h^H = 4I_k^{k-1}$ and $L_h^h = (L_h^H)^T$, and $A^{(k)} = \{a_{i,j}^{(k)}\}_{i,j=1}^\infty$ with $a_{i,j}^{(k)} = a_{|i-j|}^{(k)}$, where

$$(4.4) \quad \begin{aligned} A^{(1)} = & c_1 \cdot \text{diag}(-1, 2, -1) + c_2 \cdot \text{diag}(-1, 0, 2, 0, -1) \\ & + c_3 \cdot \text{diag}(-1, 0, 0, 2, 0, 0, -1). \end{aligned}$$

According to Lemma 4.1 and above equations, we obtain

$$(4.5) \quad \begin{aligned} A^{(k)} = & c_1 \cdot \text{diag}\left(-\frac{d_1^{(k)}}{2}, d_1^{(k)}, -\frac{d_1^{(k)}}{2}\right) \\ & + c_2 \cdot \text{diag}\left(-1, -\frac{d_2^{(k)} - 2}{2}, d_2^{(k)}, -\frac{d_2^{(k)} - 2}{2}, -1\right) \\ & + c_3 \cdot \text{diag}\left(-4, -\frac{d_3^{(k)} - 8}{2}, d_3^{(k)}, -\frac{d_3^{(k)} - 8}{2}, -4\right) \\ = & \sigma_1^{(k)} L_1 + \sigma_2^{(k)} L_2, \end{aligned}$$

where

$$\sigma_1^{(k)} = c_1 \frac{d_1^{(k)}}{2} + c_2 \frac{d_2^{(k)} - 2}{2} + c_3 \frac{d_3^{(k)} - 8}{2}, \quad \sigma_2^{(k)} = c_2 + 4c_3, \quad 2 \leq k \leq J$$

with $d_1^{(k)} = 2^k$, $d_2^{(k)} = 2^{k+2} - 6$ and $d_3^{(k)} = 9 \cdot 2^k - 24$.

Using (3.16) and (4.4), there exists

$$\begin{aligned} \kappa &= \frac{a_0^{(k)}}{-2a_1^{(k)}} = \frac{2(\sigma_1^{(k)} + \sigma_2^{(k)})}{2\sigma_1^{(k)}} = 1 + \frac{c_2 + 4c_3}{\sigma_1^{(k)}} \\ &\leq 1 + \frac{c_2 + 4c_3}{\sigma_1^{(2)}} = 1 + \frac{c_2 + 4c_3}{2c_1 + 4c_2 + 2c_3} < 3, \quad 2 \leq k \leq J; \end{aligned}$$

and

$$\kappa = \frac{a_0^{(1)}}{-2a_1^{(1)}} = \frac{2(c_1 + c_2 + c_3)}{2c_1} = 1 + \frac{c_2 + c_3}{c_1}, \quad k = 1.$$

Combining the proof of Theorem 3.9, the desired results are obtained. \square

THEOREM 4.4. *Let $A_J = A_\delta^h$ be defined by (2.10) and (2.15) on a finite bar $\Omega \in (0, b)$, where $\delta = Rh$, $R = 3$, $h \rightarrow 0$. Then*

$$\|K_k T^k\|_{A_k} \leq \sqrt{1 - \eta/6} < 1, \quad 1 \leq k \leq J,$$

where $\eta \leq 2\omega(1 - \omega)$ with $0 < \omega < 1$.

Proof. According to Lemma 4.3 and Theorem 3.11, there exists

$$\|K_k T^k\|_{A_k} \leq \sqrt{1 - \eta/3} < 1, \quad 1 \leq k \leq J - 1,$$

and

$$\|K_J T^J\|_{A_J} \leq \sqrt{1 - \eta/6} < 1, \quad k = J.$$

The proof is completed. \square

4.3. Convergence of the V-cycle MGM with $\delta = \mathbf{h}$. In the special case, the convergence of the V-cycle MGM can also be simply obtained. Firstly, we have the following lemma.

LEMMA 4.5 ([13]). *Let the symmetric positive definite matrix A_k satisfy*

$$(4.6) \quad \frac{\omega}{\lambda_{\max}(A_k)} (\nu^k, \nu^k) \leq (S_k \nu^k, \nu^k) \leq (A_k^{-1} \nu^k, \nu^k) \quad \forall \nu^k \in \mathcal{B}_k,$$

and

$$(4.7) \quad \min_{\nu^{k-1} \in \mathcal{B}_{k-1}} \|\nu^k - I_{k-1}^k \nu^{k-1}\|_{A_k}^2 \leq m_0 \|A_k \nu^k\|_{D_k^{-1}}^2 \quad \forall \nu^k \in \mathcal{B}_k$$

with $m_0 > 0$ independent of ν^k . Then

$$\|I - B_k A_k\|_{A_k} \leq \frac{m_0}{2l\omega + m_0} < 1 \quad \text{with } 1 \leq k \leq K,$$

where the operator B_k is defined by the V-cycle method in Algorithm 1 and l is the number of smoothing steps.

We know that A_δ^h reduces to the second order elliptic operator when $R \leq 1$ in (2.10). From [13, 31, 32, 33], it is easy to check that (4.6) and (4.7) hold with $m_0 = 1$. Then we have the following results.

THEOREM 4.6. *Let $A_J = A_\delta^h$ be defined by (2.10) and (2.15) on a finite bar $\Omega \in (0, b)$, where $\delta = Rh$, $R \leq 1$, $h \rightarrow 0$. Then*

$$\|I - B_k A_k\|_{A_k} \leq \frac{1}{2l\omega + 1} < 1 \quad \text{with } 1 \leq k \leq J, \quad \omega \in (0, 1/2],$$

where the operator B_k is defined by the V-cycle method in Multigrid Algorithm 1 and l is the number of smoothing steps.

REMARK 4.1. *Using Lemma 4.2, we know that (4.6) holds for the general nonlocal models or fractional models [12, 21] with $\omega \in (0, 1/3]$, but it is not easy to check the condition (4.7).*

5. Numerical Results. We employ the V-cycle MGM described in Algorithm 1 to solve the steady-state nonlocal problem (2.3). The stopping criterion is taken as $\frac{\|r^{(i)}\|}{\|r^{(0)}\|} < 10^{-8}$, where $r^{(i)}$ is the residual vector after i iterations; and the number of iterations $(m_1, m_2) = (1, 2)$ and $(\omega_{pre}, \omega_{post}) = (1, 1/3)$. In all tables, N denotes the number of spatial grid points; and the numerical errors are measured by the l_∞ (maximum) norm, ‘Rate’ denotes the convergence orders. ‘CPU’ denotes the total CPU time in seconds (s) for solving the resulting discretized systems; and ‘Iter’ denotes the average number of iterations required to solve a general linear system $A_h u_h = f_h$ at each time level.

All numerical experiments are programmed in Matlab, and the computations are carried out on a laptop with the configuration: Inter(R) Core (tm) i3 CPU 2.27 GHZ and 2 GB RAM and a Windows 7 operating system.

EXAMPLE 5.1. *Consider the steady-state nonlocal problem*

$$-\mathcal{L}_\delta u(x) = -12x^2 + 12bx - 2b^2 - \frac{6}{5}\delta^2$$

with a finite domain $0 < x < b$, $b = 4$. The exact solution of the equation is $u(x) = x^2(b-x)^2$, and the boundary conditions $u = g$ on $\Omega_{\mathcal{I}}$.

Table 5.1: Using **Galerkin approach** $A_{k-1} = I_k^{k-1} A_k I_k^k$ computed by (3.6) to solve the resulting systems (2.10) with $h = 4/N$.

N	$\delta = 1$	Rate	Iter	CPU	$\delta = \sqrt{h}$	Rate	Iter	CPU
2^{10}	4.0638e-05		13	0.043 s	3.1010e-05		21	0.067 s
2^{11}	1.0169e-05	2.00	13	0.067 s	7.7246e-06	2.01	21	0.103 s
2^{12}	2.5461e-06	2.00	12	0.101 s	1.9262e-06	2.00	21	0.175 s
2^{13}	6.3918e-07	2.00	12	0.203 s	4.8200e-07	2.00	21	0.308 s
N	$\delta = 5h$	Rate	Iter	CPU	$\delta = h$	Rate	Iter	CPU
2^{10}	3.0396e-05		22	0.069 s	2.4416e-05		18	0.058 s
2^{11}	7.5840e-06	2.00	23	0.112 s	6.1057e-06	2.00	18	0.091 s
2^{12}	1.8943e-06	2.00	23	0.196 s	1.5310e-06	2.00	18	0.146 s
2^{13}	4.7244e-07	2.00	23	0.384 s	3.8268e-07	2.00	18	0.261 s

Table 5.2: Using **doubling the mesh size** $A_{k-1} = A_\delta^{2^{k-k+1}h}$ to solve the resulting systems (2.10) with $h = 4/N$.

N	$\delta = 1$	Rate	Iter	CPU	$\delta = \sqrt{h}$	Rate	Iter	CPU
2^{10}	4.0589e-05		42	0.133 s	3.0999e-05		56	0.175 s
2^{11}	1.0132e-05	2.00	40	0.195 s	7.7139e-06	2.01	54	0.259 s
2^{12}	2.5229e-06	2.01	39	0.312 s	1.9184e-06	2.01	54	0.430 s
2^{13}	6.2373e-07	2.02	38	0.542 s	4.7520e-07	2.01	53	0.757 s
N	$\delta = 5h$	Rate	Iter	CPU	$\delta = h$	Rate	Iter	CPU
2^{10}	3.0393e-05		54	0.169 s	2.4385e-05		47	0.160 s
2^{11}	7.5819e-06	2.00	54	0.261 s	6.0749e-06	2.01	47	0.242 s
2^{12}	1.8917e-06	2.00	53	0.422 s	1.4982e-06	2.02	47	0.372 s
2^{13}	4.7021e-07	2.01	52	0.752 s	3.7118e-07	2.01	47	0.665 s

We use two coarsening strategies: *Galerkin approach* and *doubling the mesh size*, respectively, to solve the resulting system (2.10). Tables 5.1 and 5.2 show that these two methods have almost the same error values and the numerically confirm that the numerical scheme has second-order accuracy and the computation cost is almost $\mathcal{O}(N \log N)$ operations.

6. Conclusions. There are already some theoretical convergence results for using the multigrid method to solve the PDEs, the algebraic system of which has the Toeplitz structure. We notice that the proofs are mainly based on the boundedness of a_0/a_1 , where a_0 and a_1 are, respectively, the principal diagonal element and the trailing diagonal element of the Toeplitz matrix. However, in the nonlocal system, most of the time the boundedness of a_0/a_1 does not hold again. In this work, we rewrite the corresponding symmetric Toeplitz matrix as a sum of a series of Laplacian-like matrices. Then based on the analysis of the Laplacian-like matrix, we present the strict proof of the uniform convergence of the TGM. And the convergence results of the full MGM and V-cycle MGM in a special case are also derived. For the framework of the uniform convergence of the V-cycle MGM, the condition (4.6) has been confirmed to hold for the class of weakly diagonally dominant symmetric Toeplitz M-matrices, in the future we will try to find the way to verify the condition (4.7).

Acknowledgments. The first author wishes to thank Qiang Du for his valuable comments while working in Columbia university.

REFERENCES

- [1] B. AKSOYLU AND Z. UNLU, *Conditioning analysis of nonlocal integral operators in fractional sobolev spaces*, SIAM J. Numer. Anal., **52** (2014), pp. 653–677.
- [2] A. ARICÒ AND M. DONATELLI, *A V-cycle multigrid for multilevel matrix algebras: proof of optimality*, Numer. Math., **105** (2007), pp. 511–547.
- [3] A. ARICÒ, M. DONATELLI, AND S. SERRA-CAPIZZANO, *V-cycle optimal convergence for certain (multilevel) structured linear systems*, SIAM J. Matrix Anal. Appl., **26** (2004), pp. 186–214.
- [4] J. H. BRAMBLE AND J. E. PASCIAK, *New convergence estimates for multigrid algorithms*, Math. Comp., **49** (1987), pp. 311–329.
- [5] W. L. BRIGGS, V. E. HENSON, AND S. F. MCCORMICK, *A Multigrid Tutorial*, SIAM, 2000.
- [6] R. H. CHAN, Q. S. CHANG, AND H. W. SUN, *Multigrid method for ill-conditioned symmetric toeplitz systems*, SIAM J. Sci. Comput., **19** (1998), pp. 516–529.
- [7] R. H. CHAN AND X. Q. JIN, *An Introduction to Iterative Toeplitz Solvers*, SIAM, 2007.
- [8] Q. S. CHANG AND R. Q. JIA, *A refined convergence analysis of multigrid algorithms for elliptic equations*, Appl. Anal., **13** (2015), pp. 255–290.
- [9] M. H. CHEN AND W. H. DENG, *Fourth order accurate scheme for the space fractional diffusion equations*, SIAM J. Numer. Anal., **52** (2014), pp. 1418–1438.
- [10] M. H. CHEN AND W. H. DENG, *Discretized fractional substantial calculus*, ESAIM: Math. Mod. Numer. Anal., **49** (2015), pp. 373–394.
- [11] M. H. CHEN AND W. H. DENG, *High order algorithms for the fractional substantial diffusion equation with truncated Levy flights*, SIAM J. Sci. Comput., **37** (2015), pp. A890–A917.
- [12] M. H. CHEN, Y. T. WANG, X. CHENG, AND W. H. DENG, *Second-order LOD multigrid method for multidimensional Riesz fractional diffusion equation*, BIT, **54** (2014), pp. 623–647.
- [13] M. H. CHEN AND W. H. DENG, *Multigrid method for symmetric Toeplitz block tridiagonal matrix: Convergence analysis & application*, arXiv:1602.08226.
- [14] O. DEFTERLI, M. D’ELIA, Q. DU, M. GUNZBURGER, R. LEHOUCQ, AND M. MEERSCHAERT, *Fractional diffusion on bounded domains*, FCAA, **18** (2015), pp. 342–360.
- [15] Q. DU, M. GUNZBURGER, R. LEHOUCQ, AND K. ZHOU, *Analysis and approximation of nonlocal diffusion problems with volume constraints*, SIAM Rev., **56** (2012), pp. 676–696.
- [16] Q. DU, M. GUNZBURGER, R. LEHOUCQ, AND K. ZHOU, *A nonlocal vector calculus, nonlocal volume-constrained problems, and nonlocal balance laws*, Math. Model. Methods Appl. Sci., **23** (2013), pp. 493–540.

- [17] G. FIORENTINO AND S. SERRA, *Multigrid methods for symmetric positive definite block Toeplitz matrices with nonnegative generating functions*, SIAM J. Sci. Comput., **17** (1996), pp. 1068–1081.
- [18] W. HACKBUSCH, *Multigrid Methods and Applications*, Springer-Verlag, Berlin, 1985.
- [19] Y. H. HUANG AND A. OBERMAN, *Numerical methods for the fractional Laplacian: a finite difference-quadrature approach*, SIAM J. Numer. Anal., **52** (2014), pp. 3056–3084.
- [20] R. METZLER AND J. KLAFTER, *The random walks guide to anomalous diffusion: A fractional dynamics approach*, Phys. Rep., **339** (2000), pp. 1–77.
- [21] H. PANG AND H. SUN, *Multigrid method for fractional diffusion equations*, J. Comput. Phys., **231** (2012), pp. 693–703.
- [22] J. RUGE AND K. STÜBEN, *Algebraic multigrid*, in *Multigrid Methods*, Ed: S. McCormick, pp. 73–130, SIAM, 1987.
- [23] Y. SAAD, *Iterative Methods for Sparse Linear Systems*, SIAM, 2003.
- [24] S. SERRA-CAPIZZANO, *Convergence analysis of two-grid methods for elliptic Toeplitz and PDEs matrix-sequences*, Numer. Math., **92** (2002), pp. 433–465.
- [25] S. A. SILLING, *Reformulation of elasticity theory for discontinuities and long-range forces*, J. Mech. Phys. Solids, **48** (2000), pp. 175–209.
- [26] J. STOER AND R. BULIRSCH, *Introduction to Numerical Analysis*, Springer, 2002.
- [27] J. W. THOMAS, *Numerical Partial Differential Equations: Finite Difference Methods*, Springer, 1995.
- [28] X. C. TIAN AND Q. DU, *Analysis and comparison of different approximations to nonlocal diffusion and linear peridynamic equations*, SIAM J. Numer. Anal., **51** (2013), pp. 3458–3482.
- [29] X. C. TIAN AND Q. DU, *Asymptotically compatible schemes and applications to robust discretization of nonlocal models*, SIAM J. Numerical Analysis, **52** (2014), pp. 1641–1665.
- [30] H. WANG AND H. TIAN, *A fast Galerkin method with efficient matrix assembly and storage for a peridynamic model*, J. Comput. Phys., **231** (2012), pp. 7730–7738.
- [31] J. XU, *Iterative methods by space decomposition and subspace correction*, SIAM Review, **34** (1992), pp. 581–613.
- [32] J. XU AND L. ZIKATANOV, *The method of alternating projections and the method of subspace corrections in Hilbert space*, J. Am. Math. Soc., **15** (2002), pp. 573–597.
- [33] J. XU, *An introduction to multilevel methods*, in: M. Ainsworth, J. Levesley, W. A. Light and M. Marletta (Eds.), *Wavelets, Multilevel Methods and Elliptic PDEs*, Leicester, 1996, Oxford University Press, New York, (1997), pp. 213–302.
- [34] Q. YANG, I. TURNER, F. LIU, AND M. ILIĆ, *Novel numerical methods for solving the time-space fractional diffusion equation in two dimensions*, SIAM J. Sci. Comput., **33** (2011), pp. 1159–1180.
- [35] K. ZHOU AND Q. DU, *Mathematical and numerical analysis of linear peridynamic models with nonlocal boundary conditions*, SIAM J. Numer. Anal., **48** (2010), pp. 1759–1780.