

One class of conservative difference schemes for solving molecular dynamics equations of motion

E. G. Nikonov

Laboratory of Information Technologies, Joint Institute for Nuclear Research, Dubna, Russia

Simulation of many-particle system evolution by molecular dynamics takes to decrease integration step to provide numerical scheme stability on the sufficiently large time interval. It leads to a significant increase of the volume of calculations. An approach for constructing symmetric symplectic numerical schemes with given approximation accuracy in relation to integration step, for solving molecular dynamics Hamiltonian equations, is proposed in this paper. Numerical experiments show that obtained under this approach symmetric symplectic third order scheme is more stable for integration step, time-reversible and conserves Hamiltonian of the system with more accuracy at a large integration interval than second order velocity Verlet numerical schemes.

Key words: Hamiltonian systems of equations, symplectic difference schemes, generating functions, molecular dynamics

PACS: 02.60.Cb 02.60.Jh 02.70.Bf 02.70.Ns

I. INTRODUCTION

Numerical schemes, which is using for solving systems of equations of many-particle dynamics, can have restrictions on a step and an interval of integration because if they increase, the numerical schemes become unstable and do not conserve integrals of motion. As a result, when we simulate many-particle system behaviour on the sufficiently large time interval we should decrease an integration step, which leads to considerable increasing of computation quantity.

II. HAMILTONIAN MOLECULAR DYNAMICS

A motion of N -particle system in a field with potential $V(\mathbf{q})$ can be described by the system of Hamiltonian equations [1]

$$\dot{\mathbf{p}} = -\frac{\partial H(\mathbf{p}, \mathbf{q})}{\partial \mathbf{q}}, \quad \dot{\mathbf{q}} = \frac{\partial H(\mathbf{p}, \mathbf{q})}{\partial \mathbf{p}}, \quad (1)$$

with initial conditions $\mathbf{p}(0) = \mathbf{p}^0$, $\mathbf{q}(0) = \mathbf{q}^0$.

Here $\mathbf{q} = (q_1, \dots, q_d)^T$ - particle coordinates, $\mathbf{p} = (p_1, \dots, p_d)^T$ - particle momentums, $d = 3N$ - a dimension of coordinate space and

$$H(\mathbf{p}, \mathbf{q}) = \frac{1}{2} \mathbf{p}^T M^{-1}(\mathbf{q}) \mathbf{p} + V(\mathbf{q})$$

is a separable Hamiltonian of the system with symmetric and positive definite mass matrix $M(\mathbf{q})$ and a field potential $V(\mathbf{q})$.

III. DESIGN OF CONSERVATIVE DIFFERENCE SCHEMES

An approach to design of conservative difference scheme for solving Hamiltonian equations (1) is based

on following stages. First is a choice of appropriate type of generating function S , which fixes a definite family of symplectic difference scheme [2]. Symplectic difference scheme corresponds to the canonical transformation of canonical variables [3]. Solution of Hamiltonian system (1) for a definite time moment can be represented as canonical transformation and that is why conserves the value of Hamiltonian [4]. Second is using of forward and backward Taylor expansion of coordinate and momentum on a time step for getting of a corresponding generating function and remaining values of coordinate and momentum. At last, obtained at the previous stage explicit and implicit schemes are used for constructing symmetric symplectic scheme for solving Hamiltonian equations (1).

$$S = \frac{h}{2} \left(\frac{\mathbf{q}^{k+1} - \mathbf{q}^k}{h} \right)^2 - \frac{h}{2} [V(\mathbf{q}^k) + V(\mathbf{q}^{k+1})] - \frac{h}{12} [\nabla V(\mathbf{q}^{k+1}) - \nabla V(\mathbf{q}^k)] \cdot (\mathbf{q}^{k+1} - \mathbf{q}^k),$$

$$\mathbf{p}^k = \frac{\mathbf{q}^{k+1} - \mathbf{q}^k}{h} - \frac{h}{12} [5\nabla V(\mathbf{q}^k) + \nabla V(\mathbf{q}^{k+1})] - \frac{h}{12} \mathcal{H}(V)(\mathbf{q}^k) \cdot (\mathbf{q}^{k+1} - \mathbf{q}^k),$$

$$\mathbf{p}^{k+1} = \frac{\mathbf{q}^{k+1} - \mathbf{q}^k}{h} + \frac{h}{12} [\nabla V(\mathbf{q}^k) + 5\nabla V(\mathbf{q}^{k+1})] -$$

$$-\frac{h}{12} \mathcal{H}(V)(\mathbf{q}^{k+1}) \cdot (\mathbf{q}^{k+1} - \mathbf{q}^k),$$

where $\mathcal{H}(V)(q)$ is Hessian matrix for $V(q)$.

IV. NUMERICAL EXPERIMENTS

Obtained by described above approach difference scheme of third order [5] was tested by solving the system of equations (1) with Hamiltonian

$$H(\mathbf{p}, \mathbf{q}) = \frac{1}{2}(p_1^2 + p_2^2) - \frac{1}{\sqrt{q_1^2 + q_2^2}},$$

so called Kepler two body problem, Results of numerical calculations are compared for the symmetric symplectic scheme of third order and well-known second order velocity Verlet scheme with a time step $h = 0.2$ and a time interval $[0, T]$, $T = 5000$. Approximate solution by Verlet scheme with a time step $h=0.02$ is used as an exact solution [5].




FIG. 1: Phase trajectories.




FIG. 2: Conservation of Hamiltonian.

V. RESULTS

A new approach for constructing symmetric symplectic numerical schemes for solving Hamiltonian systems of equations is proposed. The numerical schemes constructed by this approach produce more stable and accurate solution of Hamiltonian system (1) and better conserve the energy of a system on the large interval of numerical integration for a relatively large integration step in comparison with the Verlet method, which is usually using for solving equations of motion in molecular dynamics.

Acknowledgments. E.G.N. acknowledge Dr. B. Batgerel for computations and numerical results. The work was supported in part by the Russian Foundation for Basic Research, Grant No. 15-01-06055.

- [1] E. Hairer, C. Lubich, G. Wanner, Geometric Numerical Integration. Second edition, Springer, 2006..
 [2] F. Kang et al, J. Comp. Math., **11**, No.1, 71 (1989).
 [3] F.R. Gantmakher, Lectons on analytical mechanics,

- PhysMathLit,Moscow, 2001.
 [4] V.I. Arnold, Mathematical methods on Classical mechanics, Springer, New York, 1978.