

ON L^p -THEORY FOR PARABOLIC AND ELLIPTIC INTEGRO-DIFFERENTIAL EQUATIONS WITH SCALABLE OPERATORS IN THE WHOLE SPACE

R. MIKULEVIČIUS AND C. PHONSOM

ABSTRACT. Elliptic and parabolic integro-differential model problems are considered in the whole space. By verifying Hörmander condition, the existence and uniqueness is proved in L_p -spaces of functions whose regularity is defined by a scalable, possibly nonsymmetric, Levy measure. Some rough probability density function estimates of the associated Levy process are used as well.

CONTENTS

1. Introduction	2
2. Notation and Main Results	3
2.1. Example	6
3. Auxiliary results	8
3.1. Some L_p estimates	8
3.2. Density estimates	11
3.3. Estimates of ψ^π	18
4. Proof of the main results	20
4.1. Existence and uniqueness for smooth input functions	21
4.2. L_2 -estimates	23
4.3. Continuity of L^π and proof of Theorem 1	26
4.4. Proof of Theorem 2	30
5. Appendix	35
5.1. Vitali Lemma, maximal functions	36
5.2. Calderon-Zygmund decomposition	39
5.3. L_p -estimates	42
References	43

Date: April 26, 2016.

1991 Mathematics Subject Classification. 45K05, 60J75, 35B65.

Key words and phrases. non-local parabolic and elliptic integro-differential equations, Lévy processes.

1. INTRODUCTION

Let $\sigma \in (0, 2)$ and \mathfrak{A}^σ be the class of all nonnegative measures π on $\mathbf{R}_0^d = \mathbf{R}^d \setminus \{0\}$ such that $\int |y|^2 \wedge 1 d\pi < \infty$ and

$$\sigma = \inf \left\{ \alpha < 2 : \int_{|y| \leq 1} |y|^\alpha d\pi < \infty \right\}.$$

In addition we assume that for $\pi \in \mathfrak{A}^\sigma$,

$$\begin{aligned} \int_{|y| > 1} |y| d\pi &< \infty \text{ if } \sigma \in (1, 2), \\ \int_{R < |y| \leq R'} y d\pi &= 0 \text{ if } \sigma = 1 \text{ for all } 0 < R < R' < \infty. \end{aligned}$$

In this paper we consider the parabolic Cauchy problem with $\lambda \geq 0$

$$\begin{aligned} (1.1) \quad \partial_t u(t, x) &= Lu(t, x) - \lambda u(t, x) + f(t, x) \text{ in } E = [0, T] \times \mathbf{R}^d, \\ u(0, x) &= 0, \end{aligned}$$

and the elliptic problem with $\lambda > 0$,

$$(1.2) \quad \lambda u(x) - Lu(x) = g(x), x \in \mathbf{R}^d,$$

with $\lambda \geq 0$ and $\lambda > 0$ and integrodifferential operator

$$L\varphi(x) = L^\pi \varphi(x) = \int [\varphi(x+y) - \varphi(x) - \chi_\sigma(y) \nabla \varphi(x) y] \pi(dy), \varphi \in C_0^\infty(\mathbf{R}^d),$$

where $\pi \in \mathfrak{A}^\sigma$, $\chi_\sigma(y) = 0$ if $\sigma \in [0, 1)$, $\chi_\sigma(y) = 1_{\{|y| \leq 1\}}(y)$ if $\sigma = 1$ and $\chi_\sigma(y) = 1$ if $\sigma \in (1, 2)$. The symbol of L is

$$\psi(\xi) = \psi^\pi(\xi) = \int \left[e^{i\xi \cdot y} - 1 - i\chi_\sigma(y) \xi \cdot y \right] \pi(dy), \xi \in \mathbf{R}^d.$$

Note that $\pi(dy) = dy/|y|^{d+\sigma} \in \mathfrak{A}^\sigma$ and, in this case, $L = L^\pi = c(\sigma, d)(-\Delta)^{\sigma/2}$, where $(-\Delta)^{\sigma/2}$ is a fractional Laplacian. Let $\pi_0 \in \mathfrak{A}^\sigma$ and

$$(1.3) \quad c_1 |\psi^{\pi_0}(\xi)| \leq |\psi^\pi(\xi)| \leq c_2 |\psi^{\pi_0}(\xi)|, \xi \in \mathbf{R}^d,$$

for some $0 < c_1 \leq c_2$. Given $\pi_0 \in \mathfrak{A}^\sigma, p \in [1, \infty)$, we denote $H_p^{\pi_0}(\mathbf{R}^d)$ (resp. $\mathcal{H}_p^{\pi_0}(E)$) the closure in $L_p(\mathbf{R}^d)$ (resp. $L_p(E)$) of $C_0^\infty(\mathbf{R}^d)$ (resp. $C_0^\infty(E)$) with respect to the norm

$$|f|_{\pi_0, p} = |f|_{L_p(\mathbf{R}^d)} + |L^{\pi_0} f|_{L_p(\mathbf{R}^d)}, \text{ resp. } |g|_{\pi_0, p} = |g|_{L_p(E)} + |L^{\pi_0} g|_{L_p(E)}.$$

If $f_n \in C_0^\infty(\mathbf{R}^d)$, $f_n \rightarrow f$ and $L^\pi f_n \rightarrow g$ in $L_p(\mathbf{R}^d)$ we denote $g = L^\pi f$.

In this note, under certain "scalability" assumptions (see Assumption **D**(κ, l) below), we prove the existence and uniqueness of (1.1) and (1.2) in $H_p^{\pi_0}(\mathbf{R}^d)$ (resp. $\mathcal{H}_p^{\pi_0}(E)$). Moreover the following estimates hold:

$$(1.4) \quad |u|_{\mathcal{H}_p^{\pi_0}} \leq C |f|_{L_p(E)}, |u|_{H_p^{\pi_0}} \leq C |f|_{L_p(\mathbf{R}^d)}.$$

The symbol $\psi^\pi(\xi)$ is not smooth in ξ and the standard Fourier multiplier results do not apply in this case. In order to prove (1.4), we associate to L^π

a family of balls and verify Hörmander condition (see Theorem 5 and (5.5) below) for it, and apply Calderon-Zygmund theorem. As an example, we consider $\pi \in \mathfrak{A}^\sigma$ defined in radial and angular coordinates $r = |y|$, $w = y/r$, as

$$(1.5) \quad \pi(\Gamma) = \int_0^\infty \int_{|w|=1} \chi_\Gamma(rw) a(r, w) j(r) r^{d-1} S(dw) dr, \Gamma \in \mathcal{B}(\mathbf{R}_0^d),$$

where $S(dw)$ is a finite measure on the unit sphere on \mathbf{R}^d . In [9], the parabolic equation (1.1) was considered, with π in the form (1.5) with $a = 1$, $j(r) = r^{-d-\sigma}$, and such that

$$\begin{aligned} & \int_0^\infty \int_{|w|=1} \chi_\Gamma(rw) r^{-1-\sigma} \rho_0(w) S(dw) dr \\ & \leq \pi(\Gamma) = \int_0^\infty \int_{|w|=1} \chi_\Gamma(rw) r^{-1-\sigma} a(r, w) S(dw) dr \\ & \leq \int_0^\infty \int_{|w|=1} \chi_\Gamma(rw) r^{-1-\sigma} S(dw) dr, \Gamma \in \mathcal{B}(\mathbf{R}_0^d), \end{aligned}$$

and (1.3) holds with $\psi^{\pi_0}(\xi) = |\xi|^\sigma$, $\xi \in \mathbf{R}^d$. In this case, $\mathcal{H}_p^{\pi_0}(E) = H_p^\sigma(E)$ is the fractional Sobolev space. The solution estimate (1.4) for (1.1) was derived in [9], using L^∞ - BMO type estimate. In [3], the elliptic problem (1.2) was studied for π in the form (1.5) with $S(dw) = dw$ being a Lebesgue measure on the unit sphere in \mathbf{R}^d , with $0 < c_1 \leq a \leq c_2$, and a set of technical assumptions on $j(r)$. The inequality (1.4) for (1.2) was obtained using sharp function estimate based on the solution Hölder norm estimate (following the idea in [1], where (1.2) was considered in $H_p^\sigma(\mathbf{R}^d)$ with π as in (1.5) with $j(r) = r^{-d-\sigma}$ and $0 < c_1 \leq a \leq c_2$).

The note is organized as follows. In Section 2, the main theorem is stated, and an example of the form (1.5) considered. In Section 3, the essential technical results are presented. The main theorem is proved in Section 4.

2. NOTATION AND MAIN RESULTS

Denote $E = [0, T] \times \mathbf{R}^d$, $\mathbf{N} = \{0, 1, 2, \dots\}$, $\mathbf{R}_0^d = \mathbf{R}^d \setminus \{0\}$. If $x, y \in \mathbf{R}^d$, we write

$$x \cdot y = \sum_{i=1}^d x_i y_i, |x| = (x \cdot x)^{1/2}.$$

For a function $u(t, x)$ on E , we denote its partial derivatives by $\partial_t u(t, x) = \partial u / \partial t$, $\partial_i u = \partial u / \partial x_i$, and $D^\gamma u = \partial^{|\gamma|} u / \partial x_1^{\gamma_1} \dots \partial x_d^{\gamma_d}$, where multiindex $\gamma = (\gamma_1, \dots, \gamma_d) \in \mathbf{N}^d$, $\nabla u = (\partial_1 u, \dots, \partial_d u)$ denotes the gradient of u with respect to x . For $k \in \mathbf{N}$, we denote $D^k u = (\partial^\gamma u)_{|\gamma|=k}$.

Let $L_p(T) = L_p(E)$ is the space of p -integrable functions with norm, $p \geq 1$,

$$|f|_{L_p(T)} = \left(\int_0^T \int |f(t, x)|^p dx dt \right)^{1/p}, |f|_{L_\infty(T)} = \operatorname{ess\,sup}_{(t, x) \in E} |f(t, x)|.$$

Similar space of functions on \mathbf{R}^d is denoted $L_p(\mathbf{R}^d)$.

Let $\mathcal{S}(\mathbf{R}^d)$ be the Schwartz space of smooth real valued rapidly decreasing functions. For $s \in \mathbf{N}$, we define the Sobolev space $H_p^s(\mathbf{R}^d)$ (resp. $H_p^s(E)$) as closure of $C_0^\infty(\mathbf{R}^d)$ (resp. $C_0^\infty(E)$) with respect to the norm

$$|f|_{n,p} = \sum_{|\beta| \leq n} \left| D^\beta f \right|_{L_p(\mathbf{R}^d)}, \text{ resp. } |g|_{n,p} = \sum_{|\beta| \leq n} \left| D^\beta f \right|_{L_p(E)}.$$

For $\sigma \in (0, 2)$ and $v \in C_0^\infty(\mathbf{R}^d)$, we define the fractional Laplacian

$$\partial^\sigma v(x) = \int \nabla_y^\sigma v(x) \frac{dy}{|y|^{d+\sigma}}, x \in \mathbf{R}^d,$$

where

$$\nabla_y^\sigma v(x) = v(x+y) - v(x) - (\nabla v(x), y) \chi_\sigma(y)$$

with $\chi_\sigma(y) = 1_{\{|y| \leq 1\}} 1_{\sigma=1} + 1_{\{\sigma \in (1,2)\}}$ is the integrand in the definition of L^π .

Given $\pi_0 \in \mathfrak{A}^\sigma, p \in [1, \infty)$, we denote $H_p^{\pi_0}(\mathbf{R}^d)$ (resp. $\mathcal{H}_p^{\pi_0}(E)$) the closure in $L_p(\mathbf{R}^d)$ (resp. $L_p(E)$) of $C_0^\infty(\mathbf{R}^d)$ (resp. $C_0^\infty(E)$) with respect to the norm

$$|f|_{\pi_0,p} = |f|_{L_p(\mathbf{R}^d)} + |L^{\pi_0} f|_{L_p(\mathbf{R}^d)}, \text{ resp. } |g|_{\pi_0,p} = |g|_{L_p(E)} + |L^{\pi_0} g|_{L_p(E)}.$$

If $f_n \in C_0^\infty(\mathbf{R}^d)$, $f_n \rightarrow f$ and $L^\pi f_n \rightarrow g$ in $L_p(\mathbf{R}^d)$ we denote $g = L^\pi f$. Notice that $f_n \rightarrow 0$, $L^\pi f_n \rightarrow h$ in $L_p(\mathbf{R}^d)$ implies that $h = 0$. Indeed,

$$\int \varphi L^\pi f_n = \int f_n L^{\pi^*} \varphi \rightarrow 0, \varphi \in C_0^\infty(\mathbf{R}^d),$$

where $\pi^*(\Gamma) = \pi(-\Gamma)$, $\Gamma \in \mathcal{B}(\mathbf{R}_0^d)$, i.e. $\pi^* \in \mathfrak{A}^\sigma$ as well. Note that if $\pi \in \mathfrak{A}^\sigma$, then for any $f \in C_0^\infty(\mathbf{R}^d)$,

$$|L^\pi f|_{L_p(\mathbf{R}^d)} \leq \left| \int_{|y| \leq 1} \dots \right|_{L_p(\mathbf{R}^d)} + \left| \int_{|y| > 1} \dots \right|_{L_p(\mathbf{R}^d)} \leq C |f|_{2,p},$$

that is $H_p^2(\mathbf{R}^d) \subseteq H_p^\pi(\mathbf{R}^d)$ and the embedding is continuous. The same holds for $H_p^2(E) \subseteq H_p^\pi(E)$.

We denote $\mathfrak{A} = \cup_{\sigma \in (0,2)} \mathfrak{A}^\sigma$.

We denote Fourier transform and its inverse

$$\begin{aligned} \mathcal{F}v(\xi) &= \hat{v}(\xi) = \int v(x) e^{-i2\pi x \cdot \xi} dx, \xi \in \mathbf{R}^d, \\ \mathcal{F}^{-1}v(x) &= \int v(\xi) e^{i2\pi x \cdot \xi} d\xi, x \in \mathbf{R}^d, v \in \mathcal{S}(\mathbf{R}^d). \end{aligned}$$

We denote $C_b^\infty(E)$ the space of bounded infinitely differentiable in x functions whose derivatives are bounded.

$C = C(\cdot, \dots, \cdot)$ denotes constants depending only on quantities appearing in parentheses. In a given context the same letter is (generally) used to denote different constants depending on the same set of arguments.

We also introduce an auxiliary Levy measure μ_0 on \mathbf{R}_0^d such that the following assumption holds.

Assumption \mathbf{A}_0 . Let $\mu^0 \in \mathfrak{A}, \chi_{\{|y| \leq 1\}} \mu^0(dy) = \mu^0(dy)$, and

$$\begin{aligned} \int |y| \mu^0(dy) + \int |\xi|^2 [1 + \zeta(\xi)]^{d+3} \exp\{-\phi_0(\xi)\} d\xi &\leq N_0 \text{ if } \sigma \in (0, 1), \\ \int |y|^2 \mu^0(dy) + \int |\xi|^4 [1 + \zeta(\xi)]^{d+3} \exp\{-\phi_0(\xi)\} d\xi &\leq N_0 \text{ if } \sigma \in [1, 2), \end{aligned}$$

where

$$\begin{aligned} \phi_0(\xi) &= \int_{|y| \leq 1} [1 - \cos(2\pi\xi \cdot y)] \mu^0(dy), \\ \zeta(\xi) &= \int_{|y| \leq 1} \chi_\sigma(y) |y| [(|\xi| |y|) \wedge 1] \mu^0(dy), \xi \in \mathbf{R}^d. \end{aligned}$$

In addition, we assume that for any $\xi \in S_{d-1} = \{\xi \in \mathbf{R}^d : |\xi| = 1\}$,

$$\int_{|y| \leq 1} |\xi \cdot y|^2 \mu^0(dy) \geq c_1 > 0.$$

For $\pi \in \mathfrak{A} = \cup_{\sigma \in (0, 2)} \mathfrak{A}^\sigma$ and $R > 0$, we denote

$$\pi_R(\Gamma) = \int \chi_\Gamma(y/R) \pi(dy), \Gamma \in \mathcal{B}(\mathbf{R}_0^d).$$

Definition 1. We say that a continuous function $\kappa : (0, \infty) \rightarrow (0, \infty)$ is a scaling function if $\lim_{R \rightarrow 0} \kappa(R) = 0, \lim_{R \rightarrow \infty} \kappa(R) = \infty$ and there is a nondecreasing continuous function $l(\varepsilon), \varepsilon > 0$, such that $\lim_{\varepsilon \rightarrow 0} l(\varepsilon) = 0$ and

$$\kappa(\varepsilon r) \leq l(\varepsilon) \kappa(r), r > 0, \varepsilon > 0.$$

We call $l(\varepsilon), \varepsilon > 0$, a scaling factor of κ .

For a scaling function κ with a scaling factor l and $\pi \in \mathfrak{A}^\sigma$ we introduce the following

Assumption $\mathbf{D}(\kappa, l)$. (i) For every $R > 0$,

$$\tilde{\pi}_R(dy) = \kappa(R) \pi_R(dy) \geq 1_{\{|y| \leq 1\}} \mu^0(dy),$$

with $\mu^0 = \mu^{0;\pi}$ satisfying Assumption \mathbf{A}_0 . If $\sigma = 1$ we, in addition assume that $\int_{R < |y| \leq R'} y \mu^0(dy) = 0$ for any $0 < R < R' \leq 1$. Here $\tilde{\pi}_R(dy) = \kappa(R) \pi_R(dy)$.

(ii) There exist α_1 and α_2 and a constant $N > 0$ such that

$$\int_{|z| \leq 1} |z|^{\alpha_1} \tilde{\pi}_R(dz) + \int_{|z| > 1} |z|^{\alpha_2} \tilde{\pi}_R(dz) \leq N \quad \forall R > 0,$$

where $\alpha_1, \alpha_2 \in (0, 1]$ if $\sigma \in (0, 1)$; $\alpha_1, \alpha_2 \in (1, 2]$ if $\sigma \in (1, 2)$; $\alpha_1 \in (1, 2]$ and $\alpha_2 \in [0, 1]$ for $\sigma = 1$.

(iii) Let $\gamma(t) = \inf(s > 0 : l(s) > t)$, $t > 0$. With $I_1 = \{t > 0 : \gamma(t) \leq 1\}$, $I_2 = \{t > 0 : \gamma(t) > 1\}$ we have

$$\int_{I_1} [t\gamma(t)^{-\alpha_1} + 1_{\sigma \in (1,2)} \gamma(t)^{-1}] dt \leq N_1 < \infty,$$

and

$$\int_{I_2} [\gamma(t)^{-(1+\alpha_2)} + \gamma(t)^{-2\alpha_2}] dt \leq N_1 < \infty.$$

The main result of this paper for (1.2) is

Theorem 1. Let $p > 1, \pi_0, \pi \in \mathfrak{A}^\sigma, \lambda > 0$. Assume there is a scaling function κ such that $\mathbf{D}(\kappa, l)$ hold for both, π and π_0 .

Then for each $f \in L_p(\mathbf{R}^d)$ there is a unique $u \in H_p^{\pi_0}(\mathbf{R}^d)$ solving (1.1). Moreover, there is $C = C(d, p, \kappa, l, N_0, N, N_1, c_1)$ such that

$$\begin{aligned} |L^{\pi_0} u|_{L_p(\mathbf{R}^d)} &\leq C |f|_{L_p(\mathbf{R}^d)}, \\ |u|_{L_p(\mathbf{R}^d)} &\leq \frac{1}{\lambda} |f|_{L_p(\mathbf{R}^d)}. \end{aligned}$$

The main result for (1.1) is

Theorem 2. Let $p \in (1, \infty), \pi_0, \pi \in \mathfrak{A}^\sigma$. Assume there is a scaling function κ such that $\mathbf{D}(\kappa, l)$ hold for both, π and π_0 .

Then for each $f \in L_p(E)$ there is a unique $u \in \mathcal{H}_p^{\pi_0}(E)$ solving (1.1). Moreover, there is $C = C(d, p, \kappa, l, N_0, N, N_1, c_1)$ such that

$$\begin{aligned} |L^{\pi_0} u|_{L_p(E)} &\leq C |f|_{L_p(E)}, \\ |u|_{L_p(E)} &\leq \left(\frac{1}{\lambda} \wedge T \right) |f|_{L_p(E)}. \end{aligned}$$

Remark 1. Assumption $\mathbf{D}(\kappa, l)$ holds for both, π, π_0 , means that κ, l , and the parameters $\alpha_1, \alpha_2, N, N_1, N_0, c_1$ are the same.

2.1. Example. Let $\mu(dt)$ be a measure on $(0, \infty)$ such that $\int_0^\infty (1 \wedge t) \mu(dt) < \infty$, and let

$$\phi(r) = \int_0^\infty (1 - e^{-rt}) \mu(dt), r \geq 0,$$

be Bernstein function (see [4], [3]). Let

$$j(r) = \int_0^\infty (4\pi t)^{-\frac{d}{2}} \exp\left(-\frac{r^2}{4t}\right) \mu(dt), r > 0.$$

We consider $\pi \in \mathfrak{A} = \cup_{\sigma \in (0,2)} \mathfrak{A}^\sigma$ defined in radial and angular coordinates $r = |y|, w = y/r$, as

$$(2.1) \quad \pi(\Gamma) = \int_0^\infty \int_{|w|=1} \chi_\Gamma(rw) a(r, w) j(r) r^{d-1} S(dw) dr, \Gamma \in \mathcal{B}(\mathbf{R}_0^d),$$

where $S(dw)$ is a finite measure on the unite sphere on \mathbf{R}^d . If $S(dw) = dw$ is the Lebesgue measure on the unit sphere, then

$$\pi(\Gamma) = \pi^{J,a}(\Gamma) = \int_{\mathbf{R}^d} \chi_{\Gamma}(y) a(|y|, y/|y|) J(y) dy, \Gamma \in \mathcal{B}(\mathbf{R}_0^d),$$

where $J(y) = j(|y|)$, $y \in \mathbf{R}^d$. Let $\pi_0 = \pi^{J,1}$, i.e.,

$$(2.2) \quad \pi_0(\Gamma) = \int_{\mathbf{R}^d} \chi_{\Gamma}(y) J(y) dy, \Gamma \in \mathcal{B}(\mathbf{R}_0^d).$$

We assume

H. (i) There is $N > 0$ so that

$$N^{-1} \phi(r^{-2}) r^{-d} \leq j(r) \leq N \phi(r^{-2}) r^{-d}, r > 0.$$

(ii) There are $0 < \delta_1 \leq \delta_2 < 1$ and $N > 0$ so that for $0 < r \leq R$

$$N^{-1} \left(\frac{R}{r} \right)^{\delta_1} \leq \frac{\phi(R)}{\phi(r)} \leq N \left(\frac{R}{r} \right)^{\delta_2}.$$

G. There is $\rho_0(w)$, $|w| = 1$, such that $\rho_0(w) \leq a(r, w) \leq 1$, $r > 0$, $|w| = 1$, and for every $|\xi| = 1$,

$$\int_{|w|=1} |\xi \cdot w|^2 \rho_0(w) S(dw) \neq 0.$$

For example, in [4] and [3] among others the following specific Bernstein functions satisfying **H** are listed:

- (0) $\phi(r) = \sum_{i=1}^n r^{\alpha_i}$, $\alpha_i \in (0, 1)$, $i = 1, \dots, n$;
- (1) $\phi(r) = (r + r^{\alpha})^{\beta}$, $\alpha, \beta \in (0, 1)$;
- (2) $\phi(r) = r^{\alpha} (\ln(1 + r))^{\beta}$, $\alpha \in (0, 1)$, $\beta \in (0, 1 - \alpha)$;
- (3) $\phi(r) = [\ln(\cosh \sqrt{r})]^{\alpha}$, $\alpha \in (0, 1)$.

The following statement holds.

Remark 2. Let π, π_0 be given by (2.1) and (2.2). Assume **H** and **G** hold.

- a) If $2\delta_1 > 1$, then Theorems 2 (resp. 1) hold in $H_p^{\pi_0}(E)$ (resp. $H_p^{\pi_0}(\mathbf{R}^d)$).
- b) If $2\delta_2 < 1$ and $2\delta_1 > \delta_2$, then Theorems 2 (resp. 1) hold in $H_p^{\pi_0}(E)$ (resp. $H_p^{\pi_0}(\mathbf{R}^d)$).

Proof. We verify that the assumptions of Theorems 2 and 1 hold. Indeed, **H** implies that there are $0 < c \leq C$ so that

$$\begin{aligned} cr^{-d-2\delta_1} &\leq j(r) \leq Cr^{-d-2\delta_2}, r \leq 1, \\ cr^{-d-2\delta_2} &\leq j(r) \leq Cr^{-d-2\delta_1}, r > 1. \end{aligned}$$

Hence $2\delta_1 \leq \sigma \leq 2\delta_2$. In this case $\kappa(R) = j(R)^{-1} R^{-d}$, $R > 0$, is a scaling function: $\kappa(\varepsilon R) \leq l(\varepsilon) \kappa(R)$, $\varepsilon, R > 0$, with

$$l(\varepsilon) = \begin{cases} C_1 \varepsilon^{2\delta_1} & \text{if } \varepsilon \leq 1, \\ C_1 \varepsilon^{2\delta_2} & \text{if } \varepsilon > 1 \end{cases}$$

for some $C_1 > 0$. Hence

$$\gamma(t) = l^{-1}(t) = \begin{cases} C_1^{-1/2\delta_1} t^{1/2\delta_1} & \text{if } t \leq C_1, \\ C_1^{-1/2\delta_2} t^{1/2\delta_2} & \text{if } t > C_1. \end{cases}$$

We see easily that α_1 is any number $> 2\delta_2$ and α_2 is any number $< 2\delta_1$. The measure μ^0 for π is

$$\mu^0(dy) = \mu^{0,\pi}(dy) = c_1 \int \chi_{dy}(rw) \chi_{\{r \leq 1\}} r^{-1-2\delta_1} \rho_0(w) S(dw) dr;$$

and μ^0 for π_0 is

$$\mu^0(dy) = \mu^{0,\pi_0}(dy) = c'_1 \int \chi_{dy}(rw) \chi_{\{r \leq 1\}} r^{-1-2\delta_1} dw dr.$$

Integrability conditions $\mathbf{D}(\kappa, l)(iii)$ easily follow from a) or b). \square

3. AUXILIARY RESULTS

In this section we present some auxiliary results.

3.1. Some L_p estimates. We start with the following observation.

Remark 3. If $\pi \in \mathfrak{A}^\sigma$, then for any $f \in C_0^\infty(\mathbf{R}^d)$,

$$|L^\pi f|_{L_p(\mathbf{R}^d)} \leq \left| \int_{|y| \leq 1} \dots \right|_{L_p(\mathbf{R}^d)} + \left| \int_{|y| > 1} \dots \right|_{L_p(\mathbf{R}^d)} \leq C |f|_{2,p}.$$

Hence $H_p^2(\mathbf{R}^d) \subseteq H_p^\pi(\mathbf{R}^d)$ and the embedding is continuous. The same holds for $H_p^2(E) \subseteq H_p^\pi(E)$.

We will use the following equality for Sobolev norm estimates.

Lemma 1. (Lemma 2.1 in [5]) For $\alpha \in (0, 1)$ and $u \in \mathcal{S}(\mathbf{R}^d)$,

$$(3.1) \quad u(x+y) - u(x) = C \int k^{(\alpha)}(y, z) \partial^\alpha u(x-z) dz,$$

where the constant $C = C(\alpha, d)$ and

$$k^{(\alpha)}(z, y) = |z+y|^{-d+\alpha} - |z|^{-d+\alpha}.$$

Moreover, there is a constant $C = C(\alpha, d)$ such that for each $y \in \mathbf{R}^d$

$$\int |k^{(\alpha)}(z, y)| dz \leq C |y|^\alpha.$$

Corollary 1. Let $\alpha \in (0, 1), p \geq 1$. Then

(i) for $y \in \mathbf{R}^d$,

$$(3.2) \quad |\partial^\alpha u|_{L_p(\mathbf{R}^d)} \leq C |u|_{H_p^1(\mathbf{R}^d)}, u \in \mathcal{S}(\mathbf{R}^d);$$

$$(3.3) \quad |u(\cdot + y) - u|_{L_p(\mathbf{R}^d)} \leq C |\partial^\alpha u|_{L_p(\mathbf{R}^d)} |y|^\alpha,$$

$$(3.4) \quad |u(\cdot + y) - u - y \cdot \nabla u|_{L_p(\mathbf{R}^d)} \leq C |\partial^\alpha \nabla u|_{L_p(\mathbf{R}^d)} |y|^{1+\alpha},$$

$u \in \mathcal{S}(\mathbf{R}^d)$.

(ii) for any $\varepsilon > 0$,

$$\partial^\alpha [u(\varepsilon \cdot)] = \varepsilon^\alpha (\partial^\alpha u)(\varepsilon x), \partial^\alpha \nabla [u(\varepsilon \cdot)] = \varepsilon^{1+\alpha} (\partial^\alpha \nabla u)(\varepsilon x), x \in \mathbf{R}^d,$$

Proof. Let $u \in \mathcal{S}(\mathbf{R}^d)$. Since for $x \in \mathbf{R}^d$,

$$\begin{aligned} |\partial^\alpha u(x)| &\leq \int_{|y| \leq 1} \int_0^1 |y \cdot \nabla u(x + sy)| ds \frac{dy}{|y|^{d+\alpha}} \\ &\quad + \int_{|y| > 1} [|u(x + y)| + u(x)] \frac{dy}{|y|^{d+\alpha}} \end{aligned}$$

(3.2) follows. Applying generalized Minkowski inequality to (3.1), we derive easily (3.3). Similarly, using

$$u(x + y) - u(x) - y \cdot \nabla u(x) = \int_0^1 y \cdot [\nabla u(x + sy) - \nabla u(x)] ds$$

and (3.3) we derive (3.4).

Changing the variable of integration,

$$\partial^\alpha [u(\varepsilon \cdot)](x) = \varepsilon^\alpha \int [u(\varepsilon x + y) - u(\varepsilon x)] \frac{dy}{|y|^{d+\alpha}} = \varepsilon^\alpha (\partial^\alpha u)(\varepsilon x), x \in \mathbf{R}^d.$$

□

Corollary 2. Let $\pi \in \mathfrak{A}^\sigma$ and

$$\int_{|z| \leq 1} |z|^{\alpha_1} \pi(dz) + \int_{|z| > 1} |z|^{\alpha_2} \pi(dz) \leq N,$$

where $\alpha_1, \alpha_2 \in (0, 1]$ if $\sigma \in (0, 1)$; $\alpha_1, \alpha_2 \in (1, 2]$ if $\sigma \in (1, 2)$; $\alpha_1 \in (1, 2]$ and $\alpha_2 \in (0, 1]$ for $\sigma = 1$.

Then there is a constant $C = C(N)$ such that for any $v \in \mathcal{S}(\mathbf{R}^d)$, (assuming $\partial^\gamma = \nabla$ if $\gamma = 1$),

$$\begin{aligned} |L^\pi v|_{L_1} &\leq C (|\partial^{\alpha_1} v|_{L_1} + |v|_{L_1}), \\ |L^\pi v|_{L_1} &\leq C (|\nabla v|_{L_1} + |\partial^{\alpha_2} v|_{L_1}), \end{aligned}$$

if $\sigma \in (0, 1)$;

$$\begin{aligned} |L^\pi v|_{L_1} &\leq C (|\partial^{\alpha_1-1} \nabla v|_{L_1} + |v|_{L_1}), \\ |L^\pi v|_{L_1} &\leq C (|D^2 v|_{L_1} + |\partial^{\alpha_2} v|_{L_1}) \end{aligned}$$

if $\sigma = 1$;

$$\begin{aligned} |L^\pi v|_{L_1} &\leq C (|\partial^{\alpha_1-1} \nabla v|_{L_1} + |\nabla v|_{L_1}), \\ |L^\pi v|_{L_1} &\leq C (|D^2 v|_{L_1} + |\partial^{\alpha_2-1} \nabla v|_{L_1}) \end{aligned}$$

if $\sigma \in (1, 2)$.

Proof. By Corollary 1, for $|y| \leq 1$

$$|v(\cdot + y) - v - \chi_\sigma(y) y \cdot \nabla u|_{L_1} \leq \begin{cases} |\partial^{\alpha_1} v|_{L_1} |y|^{\alpha_1} & \text{if } \sigma \in (0, 1) \\ |\partial^{\alpha_1-1} \nabla v|_{L_1} |y|^{\alpha_1} & \text{if } \sigma \in [1, 2) \end{cases}$$

and for $|y| > 1$,

$$|v(\cdot + y) - v - \chi_\sigma(y) y \cdot \nabla u|_{L_1} \leq \begin{cases} 2|v|_{L_1} & \text{if } \sigma \in (0, 1], \\ 2|\nabla v|_{L_1} |y|^{\alpha_2} & \text{if } \sigma \in (1, 2). \end{cases}$$

On the other hand, for $|y| \leq 1$

$$|v(\cdot + y) - v - \chi_\sigma(y) y \cdot \nabla u|_{L_1} \leq \begin{cases} |\nabla v|_{L_1} |y|^{\alpha_1} & \text{if } \sigma \in (0, 1), \\ |D^2 v|_{L_1} |y|^{\alpha_1} & \text{if } \sigma \in [1, 2), \end{cases}$$

and for $|y| > 1$,

$$|v(\cdot + y) - v - \chi_\sigma(y) y \cdot \nabla u|_{L_1} \leq \begin{cases} |\partial^{\alpha_2} v|_{L_1} & \text{if } \sigma \in (0, 1], \\ 2|\partial^{\alpha_2-1} \nabla v|_{L_1} |y|^{\alpha_2} & \text{if } \sigma \in (1, 2). \end{cases}$$

The statement follows. \square

In addition, the following holds.

Lemma 2. For any $\beta \in [0, 1]$, $a \geq 0$, $|z| \leq 1$ and $u \in \mathcal{S}(\mathbf{R}^d)$,

$$\begin{aligned} & \int_{|x| \geq a} |u(x+z) - u(x)| dx \\ & \leq 2^{1-\beta} \left(\int_{|x| \geq (a-1) \vee 0} |u(x)| dx \right)^{1-\beta} \left(\int_{|x| \geq (a-1) \vee 0} |\nabla u(x)| dx \right)^\beta |z|^\beta, \end{aligned}$$

Proof. Let $u \in \mathcal{S}(\mathbf{R}^d)$. For $\beta \in [0, 1]$, $x, z \in \mathbf{R}^d$,

$$|u(x+z) - u(x)| \leq |u(x+z) - u(x)|^{1-\beta} \left(\int_0^1 |\nabla u(x+sz)| ds \right)^\beta |z|^\beta,$$

and

$$\begin{aligned} & |u(x+z) - u(x) - z \cdot \nabla u(x)| \\ & \leq \int_0^1 |\nabla u(x+sz) - \nabla u(x)| ds |z| \end{aligned}$$

By Hölder inequality, for $|z| \leq 1$,

$$\begin{aligned}
 & \int_{|x| \geq a} |u(x+z) - u(x)| dx \\
 & \leq \int_{|x| \geq a} |u(x+z) - u(x)|^{1-\beta} \left(\int_0^1 |\nabla u(x+sz)| ds \right)^\beta dx |z|^\beta \\
 & \leq \left(\int_{|x| \geq a} |u(x+z) - u(x)| dx \right)^{1-\beta} \left(\int_0^1 \int_{|x| \geq a} |\nabla u(x+sz)| ds dx \right)^\beta |z|^\beta \\
 & \leq \left(2 \int_{|x| \geq (a-1) \vee 0} |u(x)| dx \right)^{1-\beta} \left(\int_{|x| \geq (a-1) \vee 0} |\nabla u(x)| dx \right)^\beta |z|^\beta.
 \end{aligned}$$

□

Corollary 3. For any $\beta \in [0, 1]$, $a \geq 0$, $|z| \leq 1$ and $u \in \mathcal{S}(\mathbf{R}^d)$,

$$\begin{aligned}
 & \int_{|x| \geq a} |u(x+z) - u(x) - z \cdot \nabla u(x)| dx \\
 & \leq 2^{1-\beta} \left(\int_{|x| \geq (a-1) \vee 0} |\nabla u(x)| dx \right)^{1-\beta} \left(\int_{|x| \geq (a-1) \vee 0} |D^2 u(x)| dx \right)^\beta |z|^{1+\beta}.
 \end{aligned}$$

Proof. For $\beta \in [0, 1]$, $x, z \in \mathbf{R}^d$, $|z| \leq 1$,

$$\begin{aligned}
 & |u(x+z) - u(x) - z \cdot \nabla u(x)| \\
 & \leq \int_0^1 |\nabla u(x+sz) - \nabla u(x)| ds |z|,
 \end{aligned}$$

and the claim follows by Lemma 2. □

3.2. Density estimates. We start with the following simple statement about the existence of a probability density function (pdf).

Lemma 3. Let μ^0 be a nonnegative measure on \mathbf{R}_0^d such that $\chi_{|y| \leq 1} \mu^0(dy) = \mu^0(dy)$ and

$$\begin{aligned}
 \int |y| d\mu^0 & \leq K_0 \text{ if } \sigma \in (0, 1), \\
 \int |y|^2 d\mu^0 & \leq K_0 \text{ if } \sigma \in \{1, 2\}.
 \end{aligned}$$

Let η be a r.v. such that

$$(3.5) \quad \mathbf{E} e^{i2\pi\xi \cdot \eta} = \exp\{\psi_0(\xi)\}, \xi \in \mathbf{R}^d,$$

where

$$\psi_0(\xi) = \int \left[e^{-i2\pi\xi \cdot y} - 1 - \chi_\sigma(y) i2\pi\xi \cdot y \right] \mu^0(dy), \xi \in \mathbf{R}^d.$$

Assume $n \geq 0$ and

$$(3.6) \quad \int |\xi|^n [1 + 1_{n \geq 1} \zeta(\xi)]^{d+3} \exp \{\phi_0(\xi)\} d\xi \leq K_0,$$

where $\phi_0(\xi) = \operatorname{Re} \psi_0(\xi)$, $\xi \in \mathbf{R}^d$ and

$$\zeta(\xi) = \int_{|y| \leq 1} \chi_\sigma(y) |y| [(|\xi| |y|) \wedge 1] \mu^0(dy), \xi \in \mathbf{R}^d.$$

Then η has a pdf $p_0(x)$, $x \in \mathbf{R}^d$, such that

$$\sup_x \left| \partial^\beta p_0(x) \right| + \int (1 + |x|^2) \left| \partial^\beta p_0(x) \right| dx \leq C \quad \forall |\beta| \leq n$$

for some $C = C(d, K_0)$.

Proof. By Proposition I.2.5 in [7], η has a continuous bounded density

$$(3.7) \quad p_0(x) = \int e^{-i2\pi x \cdot \xi} \exp \{\psi_0(\xi)\} d\xi$$

if

$$\int \exp \{-\phi_0(\xi)\} d\xi < \infty.$$

The assumption (3.6) implies that for any multiindex $|\beta| \leq n$,

$$\partial^\beta p_0(x) = \int e^{-i2\pi x \cdot \xi} (-i2\pi \xi)^\beta \exp \{\psi_0(\xi)\} d\xi, x \in \mathbf{R}^d,$$

is a bounded continuous function. The function $(1 + |x|^2) \partial^\beta p_0$ is integrable if

$$\begin{aligned} (3.8) \quad & (-i2\pi x_j)^{d+1} (-i2\pi x_k)^2 \partial^\beta p_0(x) \\ &= \int \partial_{\xi_j}^{d+1} \partial_{\xi_k}^2 [e^{-i2\pi x \cdot \xi}] (-i2\pi \xi)^\beta \exp \{\psi_0(\xi)\} d\xi \\ &= (-1)^{d+3} \int e^{-ix \cdot \xi} \partial_{\xi_j}^{d+1} \partial_{\xi_k}^2 [(-i2\pi \xi)^\beta \exp \{\psi_0(\xi)\}] d\xi \end{aligned}$$

is bounded for all j, k . Since $\partial^\mu \psi_0(\xi)$ is bounded for $|\mu| \geq 2$ and

$$|\nabla \psi_0(\xi)| \leq C(1 + \zeta(\xi)), \xi \in \mathbf{R}^d,$$

the boundedness of (3.8) follows from assumption (3.6). Therefore $p_0(x)$ has n bounded continuous derivatives and for any multiindex $|\beta| \leq n$,

$$(3.9) \quad \int (1 + |x|^2) \left| \partial^\beta p_0(x) \right| dx \leq C$$

with $C = C(d, K_0)$. □

We will need the following tail estimate.

Lemma 4. *Let $\pi \in \mathfrak{A}$. Assume*

$$(3.10) \quad \int_{|z| \leq 1} |z|^{\alpha_1} \pi(dz) + \int_{|z| > 1} |z|^{\alpha_2} \pi(dz) \leq N,$$

where $\alpha_1, \alpha_2 \in (0, 1]$ if $\sigma \in (0, 1)$; $\alpha_1, \alpha_2 \in (1, 2]$ if $\sigma \in (1, 2)$; $\alpha_1 \in (1, 2]$ and $\alpha_2 \in [0, 1)$ if $\sigma = 1$. Let ζ_t be the associated Levy process, that is

$$\mathbf{E} e^{i2\pi\xi \cdot \zeta_t} = \exp\{\psi(\xi) t\}, t \geq 0,$$

with

$$\psi(\xi) = \int \left[e^{i2\pi\xi \cdot y} - 1 - i2\pi\chi_\sigma(y) y \cdot \xi \right] d\pi, \xi \in \mathbf{R}^d.$$

Let $t > 0$ and $\mathcal{L}_t(dy)$ be the distribution measure of ζ_t on \mathbf{R}^d . Then for each $\delta > 0$ there is a constant $C = C(\delta, N)$ such that

$$\mathcal{L}_t(\{|y| > \delta\}) \leq Ct.$$

Proof. Recall

$$(3.11) \quad \zeta_t = \int_0^t \int \chi_\sigma(y) y q(ds, dy) + \int_0^t \int (1 - \chi_\sigma(y)) y p(ds, dy), t \geq 0,$$

$p(ds, dy)$ is Poisson point measure with

$$\mathbf{E} p(ds, dy) = \pi(dy) ds, q(ds, dy) = p(ds, dy) - \pi(dy) ds.$$

Now, $\zeta_t = \bar{\zeta}_t + \tilde{\zeta}_t$ with

$$\begin{aligned} \bar{\zeta}_t &= \int_0^t \int_{|y| \leq 1} \chi_\sigma(y) y q(ds, dy) + \int_0^t \int_{|y| \leq 1} (1 - \chi_\sigma(y)) y p(ds, dy), \\ \tilde{\zeta}_t &= \int_0^t \int_{|y| > 1} \chi_\sigma(y) y q(ds, dy) + \int_0^t \int_{|y| > 1} (1 - \chi_\sigma(y)) y p(ds, dy), \end{aligned}$$

$t \geq 0$.

Case 1: $\sigma \in (0, 1)$. In this case (3.10) holds with $\alpha_1, \alpha_2 \in (0, 1]$. Then

$$|\bar{\zeta}_t|^{\alpha_1} = \sum_{s \leq t} [|\bar{\zeta}_{s-} + \Delta \bar{\zeta}_s|^{\alpha_1} - |\bar{\zeta}_{s-}|^{\alpha_1}] \leq \sum_{s \leq t} |\Delta \bar{\zeta}_s|^{\alpha_1},$$

and

$$\mathbf{E} |\bar{\zeta}_t|^{\alpha_1} \leq t \int_{|y| \leq 1} |y|^{\alpha_1} \pi(dy) \leq Nt.$$

Similarly, $\mathbf{E} |\tilde{\zeta}_t|^{\alpha_2} \leq Nt$.

Case 2: $\sigma \in (1, 2)$. In this case, $\alpha_1, \alpha_2 \in (1, 2]$. Then

$$\mathbf{E} [\bar{\zeta}_t^2] = \int_{|y| \leq 1} |y|^2 \pi(dy) t \leq Nt,$$

and

$$\mathbf{E} [\tilde{\zeta}_t] \leq 2t \int_{|y| > 1} |y|^{\alpha_2} \pi(dy) \leq Nt$$

Case 3: $\sigma = 1$. In this case, $\alpha_1 \in (1, 2]$ and $\alpha_2 \in [0, 1)$. Similarly as above, we find that

$$\begin{aligned} \mathbf{E}[\bar{\zeta}_t^2] &= t \int_{|y| \leq 1} |y|^2 \pi(dy) \leq Nt, \\ \mathbf{E}[\tilde{\zeta}_t^{\alpha_2}] &\leq Nt. \end{aligned}$$

The statement is proved. \square

Let $\pi \in \mathfrak{A}^\sigma$ and $p(dt, dy)$ be a Poisson point measure on $[0, \infty) \times \mathbf{R}_0^d$ such that $\mathbf{E}p(dt, dy) = \pi(dy) dt$. Let $q(dt, dy) = p(dt, dy) - \pi(dy) dt$. We associate to L^π the stochastic process with independent increments

$$(3.12) \quad Z_t = Z_t^\pi = \int_0^t \int \chi_\sigma(y) y q(ds, dy) + \int_0^t \int (1 - \chi_\sigma(y)) y p(ds, dy), t \geq 0.$$

By Ito formula,

$$(3.13) \quad \mathbf{E}e^{i2\pi\xi \cdot Z_t^\pi} = \exp\{\psi^\pi(\xi) t\}, t \geq 0, \xi \in \mathbf{R}^d,$$

where

$$\psi^\pi(\xi) := \int [\exp(i2\pi\xi \cdot y) - 1 - i2\pi y \cdot \xi \chi_\sigma(y)] \pi(dy).$$

Let $\kappa(R)$, $R > 0$, be a scaling function, $Z_t = Z_t^R$ be the stochastic process with independent increments associated with $\tilde{\pi}_R = \kappa(R) \pi_R$, i.e.,

$$\mathbf{E}e^{i2\pi\xi \cdot Z_t^R} = \exp\{\psi^{\tilde{\pi}_R}(\xi) t\}$$

with

$$\psi^{\tilde{\pi}_R}(\xi) = \int [e^{i2\pi\xi \cdot y} - 1 - i2\pi\chi_\sigma(y) y \cdot \xi] d\tilde{\pi}_R, \xi \in \mathbf{R}^d.$$

Note Z_t^R and $R^{-1}Z_{\kappa(R)t}^\pi$, $t > 0$, have the same distribution.

Lemma 5. *Let $\pi \in \mathfrak{A}^\sigma$, κ be a scaling function with scaling factor l . Assume*

$$\tilde{\pi}_R(dy) = \kappa(R) \pi_R(dy) \geq 1_{\{|y| \leq 1\}} \mu^0(dy)$$

with μ^0 satisfying the assumptions of Lemma 3 (in particular, (3.6) with $n \geq 0$ and the constant K_0), and Let

$$\psi_0(\xi) = \int [e^{-2\pi\xi \cdot y} - 1 - \chi_\sigma(y) \xi \cdot y] \mu^0(dy), \xi \in \mathbf{R}^d.$$

a) For each $t > 0$, $R > 0$, we have $Z_t^R = \eta_t + \eta'_t$ (in distribution), η_t and $\tilde{\eta}_t$ are independent with

$$(3.14) \quad \mathbf{E}e^{i2\pi\xi \cdot \eta_t} = \exp\{\psi_0(\xi \gamma(t))\}, \xi \in \mathbf{R}^d,$$

and $\mu_{\gamma(t)^{-1}}^0 \leq t\tilde{\pi}_R$, where $\gamma(t) = l^{-1}(t) = \inf\{s : l(s) \geq t\}$. Moreover, $\eta_t = \gamma(t) \eta$ (in distribution), where η is a r.v. in Lemma 3.

b) For every $t > 0$, $R > 0$, the process Z_t^R (equivalently $R^{-1}Z_{\kappa(R)t}^\pi$) has a bounded continuous probability density function

$$p^R(t, x) = \gamma(t)^{-d} \int p_0\left(\frac{x - y}{\gamma(t)}\right) P_{t,R}(dy), x \in \mathbf{R}^d,$$

where $P_{t,R}(dy)$ is the distribution measure of η'_t on \mathbf{R}^d and p_0 is pdf of η . Moreover, $p^R(t, x)$ has n bounded continuous derivatives such that for any multiindex $|\beta| \leq n$,

$$\begin{aligned} \int \left| \partial^\beta p^R(t, x) \right| dx &\leq \gamma(t)^{-|\beta|} \int \left| \partial^\beta p_0(x) \right| dx, \\ \sup_{x \in \mathbf{R}^d} \left| \partial^\beta p^R(t, x) \right| &\leq \gamma(t)^{-d-|\beta|} \sup_x \left| \partial^\beta p_0(x) \right|, \end{aligned}$$

and for any $\alpha \in (0, 1)$ such that $|\beta| + \alpha < n$

$$(3.15) \quad \int \left| \partial^\alpha \partial^\beta p^R(t, x) \right| dx \leq \gamma(t)^{-|\beta|-\alpha} \int \left| \partial^\alpha \partial^\beta p_0(x) \right| dx.$$

c) Assume, in addition, that there exist α_1 and α_2 and a constant $N > 0$ such that

$$\int_{|z| \leq 1} |z|^{\alpha_1} \tilde{\pi}_R(dz) + \int_{|z| > 1} |z|^{\alpha_2} \tilde{\pi}_R(dz) \leq N \quad \forall R > 0,$$

where $\alpha_1, \alpha_2 \in (0, 1]$ if $\sigma \in (0, 1)$; $\alpha_1, \alpha_2 \in (1, 2]$ if $\sigma \in (1, 2)$; $\alpha_1 \in (1, 2]$ and $\alpha_2 \in [0, 1]$ if $\sigma = 1$. Then for each $a > 0$ there is $C = C(d, a, N, K_0, n)$ such that for any multiindex $|\beta| \leq n, R > 0, t > 0$,

$$\int_{|x| > a} \left| \partial^\beta p^R(t, x) \right| dx \leq C \left(\gamma(t)^{2-|\beta|} + t \gamma(t)^{-|\beta|} \right).$$

Proof. a) Let $R > 0, t > 0$. Since $l(l^{-1}(t)) = t$, we have $\kappa(R)t \geq \kappa(Rl^{-1}(t)) = \kappa(R\gamma(t))$. Hence

$$(3.16) \quad \tilde{\pi}_R t \geq \kappa(R\gamma(t)) \pi_{R\gamma(t)/\gamma(t)} \geq \mu_{\gamma(t)^{-1}}^0,$$

and $\mu_{\gamma(t)^{-1}}^0(dy) = \mu^0(\gamma(t)^{-1}dy)$ is the Levy measure of a random variable, denoted η_t , such that (3.14) holds. Let

$$\psi^{t\tilde{\pi}_R}(\xi) = \psi_0(\xi\gamma(t)) + \psi'(\xi), \xi \in \mathbf{R}^d.$$

The inequality (3.16) implies that (see e.g. [2]) $\psi' = \psi^{\Pi_t}$ with $\Pi_t = \tilde{\pi}_R t - \mu_{\gamma(t)^{-1}}^0$ and $\exp\{\psi'(\xi)\}$ is characteristic function of a random variable η'_t independent of η_t . Obviously the distribution of Z_t^R coincides with the distribution of the sum $\eta_t + \eta'_t$. If η is a r.v. with characteristic function (3.5), then $\eta_t = \gamma(t)\eta$ in distribution.

b) First we prove the existence of the probability density function of η whose characteristic function is $\exp\{\psi_0(\xi)\}$. Note that $\phi_0(\xi) = \operatorname{Re} \psi_0(\xi), \xi \in \mathbf{R}^d$. Let $t > 0$. By part a), $Z_t^R = \gamma(t)\eta + \eta'_t$ (in distribution), η and η'_t are independent. The pdf of $\eta\gamma(t)$ is

$$(3.17) \quad p_0(t, x) = \gamma(t)^{-d} p_0(x/\gamma(t)), x \in \mathbf{R}^d.$$

Let $P_{t,R}(dy)$ be the distribution measure of η'_t on \mathbf{R}^d . Since $\eta\gamma(t)$ and η'_t are independent, Z_t^R has a density

$$p^R(t, x) = \int p_0(t, x - y) P_{t,R}(dy), x \in \mathbf{R}^d.$$

According to (3.17) (see (3.9) as well), for any $|\beta| \leq n$,

$$\partial^\beta p^R(t, x) = \int \partial^\beta p_0(t, x - y) P_{t,R}(dy),$$

and, according to Corollary 1(ii),

$$(3.18) \quad \sup_{x,R} \left| \partial^\beta p^R(t, x) \right| \leq \gamma(t)^{-d-|\beta|} \sup_x \left| \partial^\beta p_0(x) \right| < \infty,$$

$$\int \left| \partial^\beta p^R(t, x) \right| dx \leq \gamma(t)^{-|\beta|} \int \left| \partial^\beta p_0(x) \right| dx < \infty.$$

Similarly, see Corollary 1, (3.15) follows.

c) Let $a > 0, |\beta| \leq n$.

Then

$$\begin{aligned} \int_{|x|>a} \left| \partial^\beta p^R(t, x) \right| dx &= \gamma(t)^{-d-|\beta|} \int_{|x|>a} \left| \int (\partial^\beta p_0) \left(\frac{x-y}{\gamma(t)} \right) P_{t,R}(dy) \right| dx \\ &\leq \int \int_{|x-y|>a/2} \dots + \int \int_{|y|>a/2} \dots \\ &\leq C[\gamma(t)^{2-|\beta|} \int |x|^2 \left| \partial^\beta p_0(x) \right| dx + t\gamma(t)^{-|\beta|} \left| \partial^\beta p_0 \right|_{L_1}], \end{aligned}$$

because by Lemma 4 there is $C = C(N, a)$ such that

$$P_{t,R}(|y| > a/2) \leq Ct.$$

□

We will need some estimates involving the operators L^π .

Lemma 6. *Let $\pi_0 \in \mathfrak{A}^\sigma, \kappa$ be a scaling function with scaling factor l , and $\mathbf{D}(\kappa, l)(i)-(ii)$ hold for π_0 . Let $\pi \in \mathfrak{A}^\sigma$ be such that*

$$\int_{|z|\leq 1} |z|^{\alpha_1} \tilde{\pi}_R(dz) + \int_{|z|>1} |z|^{\alpha_2} \tilde{\pi}_R(dz) \leq N \quad \forall R > 0,$$

for some $N_2 > 0$, where $\tilde{\pi}_R(dy) = \kappa(R) \pi_R(dy)$, and α_1, α_2 are exponents in assumption $\mathbf{D}(\kappa, l)$ for π_0 . Let $R > 0$ and $p^R(t, x), x \in \mathbf{R}^d$, be pdf of $R^{-1}Z_{\kappa(R)t}^{\pi_0}$ (see Lemma 5), $\gamma(t) = l^{-1}(t), t > 0$, and $L^{0,R}$ be the operator corresponding to Levy measure $\kappa(R) \pi_0(Rdy)$. Then there is $C = C(d, N_0, N)$ such that

a)

$$(3.19) \quad \int_{|x|>2} \left| L^{\tilde{\pi}_R} p^R(t, x) \right| dx$$

$$\leq C \left[1 + 1_{\sigma \in (1,2)} \gamma(t)^{-1} + \gamma(t)^{-\alpha_1} \left(\gamma(t)^2 + t \right) \right],$$

for all $t > 0$;

b) denoting $I_2 = \{t > 0 : \gamma(t) > 1\}$,

$$(3.20) \quad |L^{\tilde{\pi}_R} \nabla p^R(t, \cdot)|_{L_1} \leq C \gamma(t)^{-(1+\alpha_2)}, t \in I_2$$

c)

$$(3.21) \quad |L^{\tilde{\pi}_R} L^{0,R} p^R(t, \cdot)|_{L_1} \leq C \gamma(t)^{-2\alpha_2}, t \in I_2.$$

Proof. a) For any $t > 0$,

$$\int_{|x|>2} |L^{\tilde{\pi}_R} p^R(t, x)| dx \leq \int \int_{|x|>2} |\nabla_z^\sigma p^R(t, x)| dx \pi(dz).$$

Now, for $|z| \leq 1$, by Lemmas 2 and 5,

$$\begin{aligned} & \int_{|x|>2} |\nabla_z^\sigma p^R(t, x)| dx \\ & \leq C |z|^{\alpha_1} \left(\int_{|x|>1} |p^R(t, x)| dx \right)^{1-\alpha_1} \left(\int_{|x|>1} |\nabla p^R(t, x)| dx \right)^{\alpha_1} \\ & \leq C |z|^{\alpha_1} [\gamma(t)^2 + t] \gamma(t)^{-\alpha_1} \text{ if } \sigma \in (0, 1); \end{aligned}$$

and by Corollary 3 and Lemma 5,

$$\begin{aligned} & \int_{|x|>2} |\nabla_z^\sigma p^R(t, x)| dx \\ & \leq C |z|^{\alpha_1} \left(\int_{|x|>1} |\nabla p^R(t, x)| dx \right)^{2-\alpha_1} \left(\int_{|x|>1} |D^2 p^R(t, x)| dx \right)^{\alpha_1-1} \\ & \leq C |z|^{\alpha_1} [t + \gamma(t)^2] \gamma(t)^{\alpha_1} \text{ if } \sigma \in [1, 2). \end{aligned}$$

For $|z| > 1$,

$$\int_{|x|>2} |\nabla_z^\sigma p^R(t, x)| dx \leq 2 \text{ if } \sigma \in (0, 1],$$

and

$$\int_{|x|>2} |\nabla_z^\sigma p^R(t, x)| dx \leq C(1 + |z| \int_{|x|>2} |\nabla p^R(t, x)| dx) \text{ if } \sigma \in (1, 2).$$

Hence by Lemma 5 c),

$$\int \int_{|x|>2} |\nabla_z^\sigma p^R(t, x)| dx \pi(dz) \leq C [1 + \gamma(t)^{-\alpha_1} (\gamma(t)^2 + t)]$$

if $\sigma \in (0, 1]$, and

$$\begin{aligned} \int \int_{|x|>2} |\nabla_z^\sigma p^{*R}(t, x)| dx \pi(dz) & \leq C \gamma(t)^{-\alpha_1} (\gamma(t)^2 + t) \\ & \quad + C (1 + \gamma(t)^{-1}) \end{aligned}$$

if $\sigma \in (1, 2)$.

b) Let $t \in I_2$. By Corollary 2 and Lemma 5 b),

$$\begin{aligned} |L^{\tilde{\pi}_R} \nabla p^R(t, \cdot)|_{L_1} &\leq C \left(|D^2 p^R(t, \cdot)|_{L_1} + |\partial^{\alpha_2} \nabla p^R(t, \cdot)|_{L_1} \right) \\ &\leq C \left[\gamma(t)^{-2} + \gamma(t)^{-(1+\alpha_2)} \right] \leq C \gamma(t)^{-(1+\alpha_2)} \end{aligned}$$

if $\sigma \in (0, 1)$. Similarly,

$$|L^R \nabla p^R(t, \cdot)|_{L_1} \leq C \left[\gamma(t)^{-3} + \gamma(t)^{-(1+\alpha_2)} \right] \leq C \gamma(t)^{-(1+\alpha_2)}$$

if $\sigma = 1$, and

$$\begin{aligned} |L^R \nabla p^R(t, \cdot)|_{L_1} &\leq C \left(|D^3 p^R(t, \cdot)|_{L_1} + |\partial^{\alpha_2-1} D^2 p^R(t, \cdot)|_{L_1} \right) \\ &\leq C \left[\gamma(t)^{-3} + \gamma(t)^{-(1+\alpha_2)} \right] \leq C \gamma(t)^{-(1+\alpha_2)} \end{aligned}$$

if $\sigma \in (1, 2)$.

c) Let $t \in I_2$, i.e., $\gamma(t) > 1$. By Corollary 2,

$$\begin{aligned} &|L^{\tilde{\pi}_R} L^{0,R} p^R(t, \cdot)|_{L_1} \\ &\leq C \left(|L^{0,R} \nabla p^R(t, \cdot)|_{L_1} + |L^{0,R} \partial^{\alpha_2} p^R(t, \cdot) v|_{L_1} \right) \\ &\leq C \left(|D^2 p^R(t, \cdot)|_{L_1} + |\partial^{\alpha_2} \nabla p^R(t, \cdot)|_{L_1} + |\partial^{\alpha_2} \partial^{\alpha_2} p^R(t, \cdot) v|_{L_1} \right) \end{aligned}$$

if $\sigma \in (0, 1)$;

$$\begin{aligned} &|L^{\tilde{\pi}_R} L^{0,R} p^R(t, \cdot)|_{L_1} \\ &\leq C \left(|L^{0,R} D^2 p^R(t, \cdot)|_{L_1} + |L^{0,R} \partial^{\alpha_2} p^R(t, \cdot) v|_{L_1} \right) \\ &\leq C \left(|D^4 p^R(t, \cdot)|_{L_1} + |\partial^{\alpha_2} D^2 p^R(t, \cdot)|_{L_1} + |\partial^{\alpha_2} \partial^{\alpha_2} p^R(t, \cdot) v|_{L_1} \right) \end{aligned}$$

if $\sigma = 1$;

$$\begin{aligned} &|L^{\tilde{\pi}_R} L^{0,R} p^R(t, \cdot)|_{L_1} \\ &\leq C \left(|L^{0,R} D^2 p^R(t, \cdot)|_{L_1} + |L^{0,R} \partial^{\alpha_2-1} \nabla p^R(t, \cdot) v|_{L_1} \right) \\ &\leq C \left(|D^4 p^R(t, \cdot)|_{L_1} + |\partial^{\alpha_2-1} D^3 p^R(t, \cdot)|_{L_1} + |\partial^{\alpha_2-1} \partial^{\alpha_2-1} D^2 p^R(t, \cdot) v|_{L_1} \right) \end{aligned}$$

if $\sigma \in (1, 2)$. The estimate (3.21) follows by Lemma 5. \square

3.3. Estimates of ψ^π . We present now some properties of the functions $\psi^\pi(\xi)$, $\xi \in \mathbf{R}^d$, with $\pi \in \mathfrak{A}^\sigma$.

Lemma 7. *Let $\pi \in \mathfrak{A}^\sigma$ and $\kappa(R)$, $R > 0$, be a scaling function, and $\tilde{\pi}_R(dy) = \kappa(R) \pi(Rdy)$.*

a) Assume there is $N_1 > 0$ so that

$$(3.22) \quad \begin{aligned} \int (|y| \wedge 1) \tilde{\pi}_R(dy) &\leq N_2 \text{ if } \sigma \in (0, 1), \\ \int (|y|^2 \wedge 1) \tilde{\pi}_R(dy) &\leq N_2 \text{ if } \sigma = 1, \\ \int_{|y| \leq 1} |y|^2 \tilde{\pi}_R(dy) + \int_{|y| > 1} |y| \tilde{\pi}_R(dy) &\leq N_2 \text{ if } \sigma \in (1, 2) \end{aligned}$$

for any $R > 0$. Then there is a constant C_1 so that for all $\xi \in \mathbf{R}^d$,

$$\begin{aligned} \int [1 - \cos(2\pi\xi y)] \pi(dy) &\leq C_1 N_2 \kappa(|\xi|^{-1})^{-1}, \\ \int |\sin(2\pi\xi \cdot y) - 2\pi\chi_\sigma(y) \xi \cdot y| \pi(dy) &\leq C_1 N_2 \kappa(|\xi|^{-1})^{-1}, \end{aligned}$$

assuming $\kappa(|\xi|^{-1})^{-1} = 0$ if $\xi = 0$.

b) Assume there is a $n_1 > 0$ such that

$$(3.23) \quad \int_{|y| \leq 1} |\xi \cdot y|^2 \tilde{\pi}_R(dy) \geq n_1,$$

for all $R > 0$ and $\xi \in S_{d-1} = \{\xi \in \mathbf{R}^d : |\xi| = 1\}$. Then there is a constant $c_2 = c_2(l) > 0$ such that

$$\int [1 - \cos(2\pi\xi y)] \pi(dy) \geq c_2 n_1 \kappa(|\xi|^{-1})^{-1}$$

for all $\xi \in \mathbf{R}^d$, assuming $\kappa(|\xi|^{-1})^{-1} = 0$ if $\xi = 0$.

Proof. The following simple trigonometric estimates hold:

$$(3.24) \quad \begin{aligned} |\sin x - x| &\leq \frac{|x|^3}{6}, 1 - \cos x \leq \frac{1}{2}x^2, x \in \mathbf{R}, \\ 1 - \cos x &\geq \frac{x^2}{\pi} \text{ if } |x| \leq \pi/2. \end{aligned}$$

a) Let $\xi \neq 0$. Denoting $\hat{\xi} = \xi/|\xi|$, and using (3.24),

$$\begin{aligned} \int \left| 1 - \cos(2\pi\hat{\xi}|\xi|y) \right| \pi(dy) &= \kappa(|\xi|^{-1})^{-1} \int \left| 1 - \cos(2\pi\hat{\xi} \cdot y) \right| \tilde{\pi}_{|\xi|^{-1}}(dy) \\ &\leq \kappa(|\xi|^{-1})^{-1} 2\pi^2 \int (|y|^2 \wedge 1) \tilde{\pi}_{|\xi|^{-1}}(dy), \end{aligned}$$

and there is C_1 so that

$$\begin{aligned} &\int |\sin(2\pi\xi \cdot y) - 2\pi\chi_\sigma(y) \xi \cdot y| \pi(dy) \\ &= \kappa(|\xi|^{-1})^{-1} \int \left| \sin(2\pi\hat{\xi} \cdot y) - 2\pi\chi_\sigma(y) \hat{\xi} \cdot y \right| \tilde{\pi}_{|\xi|^{-1}}(dy) \\ &\leq C_1 N_2 \kappa(|\xi|^{-1})^{-1} \end{aligned}$$

for all $\xi \in \mathbf{R}^d$.

b) By (3.24), for all $\xi \in \mathbf{R}^d$,

$$\begin{aligned}
& \int [1 - \cos(2\pi\xi \cdot y)] \pi(dy) \\
&= \int [1 - \cos(2\pi\hat{\xi} \cdot y)] \pi_{|\xi|^{-1}}(dy) \geq \int_{|y| \leq \frac{1}{4}} 4\pi |\hat{\xi} \cdot y|^2 \pi_{|\xi|^{-1}}(dy) \\
&= 4^{-1} \int_{|4y| \leq 1} \pi |\hat{\xi} \cdot 4y|^2 \pi_{|\xi|^{-1}}(dy) = 4^{-1} \kappa(|4\xi|^{-1})^{-1} \int_{|y| \leq 1} \pi |\hat{\xi} \cdot y|^2 \tilde{\pi}_{|4\xi|^{-1}}(dy) \\
&\geq n_1 4^{-1} \pi \kappa(|\xi|^{-1})^{-1} \frac{\kappa(|\xi|^{-1})}{\kappa(|4\xi|^{-1})}.
\end{aligned}$$

Let l be a scaling factor of κ , i.e., $\kappa(\varepsilon R) \leq l(\varepsilon) \kappa(R)$, $\varepsilon, R > 0$. Then

$$\frac{\kappa(|\xi|^{-1})}{\kappa(|4\xi|^{-1})} \geq \frac{1}{l(4^{-1})} > 0 \quad \forall \xi \in \mathbf{R}^d.$$

The claim follows. \square

For $\pi \in \mathfrak{A}^\sigma$, let

$$\begin{aligned}
\tilde{\psi}(\xi) &= \tilde{\psi}^\pi(\xi) = \int [\cos(2\pi\xi y) - 1] \pi(dy) = \operatorname{Re} \psi^\pi(\xi), \\
\varphi(\xi) &= \varphi^\pi(\xi) = \int [\sin(2\pi\xi \cdot y) - 2\pi\chi_\sigma(y) \xi \cdot y] \pi(dy) = \operatorname{Im} \psi^\pi(\xi),
\end{aligned}$$

$\xi \in \mathbf{R}^d$. An obvious consequence of Lemma 7 (note $\psi^{\pi^*}(\xi) = \psi^\pi(-\xi)$, $\xi \in \mathbf{R}^d$) is the following

Corollary 4. *Let κ be a scaling function with scaling factor l and both assumptions, (3.22) and (3.23), of Lemma 7 hold for $\pi \in \mathfrak{A}^\sigma$. Then there is constant $c = c(n_1, N_1, l) > 0$ such that*

$$c |\psi^\pi(\xi)| \leq |\tilde{\psi}^\pi(\xi)| \leq |\psi^\pi(\xi)|, \quad \xi \in \mathbf{R}^d,$$

and $|\varphi^\pi(\xi)| \leq c^{-1} |\tilde{\psi}^\pi(\xi)|$, $\xi \in \mathbf{R}^d$.

Note that it implies

$$c |\psi^{\pi^*}(\xi)| \leq |\tilde{\psi}^\pi(\xi)| = |\tilde{\psi}^\pi(-\xi)| \leq |\psi^{\pi^*}(\xi)|, \quad \xi \in \mathbf{R}^d,$$

where $\pi^*(dy) = \pi(-dy)$.

4. PROOF OF THE MAIN RESULTS

In this section we prove the main results in three steps. First we prove the existence and uniqueness of classical solutions for smooth input functions. Then we derive $|u|_{L^{\pi_0}}$ -norm estimates with constants independent of the regularity of the input function. Finally, continuity estimate of L^π with

respect to $|u|_{L^{\pi_0}}$ -norm allows to pass to the limit and derive the results for the input function $f \in L_p$.

4.1. Existence and uniqueness for smooth input functions. For $E = [0, T] \times \mathbf{R}^d$, we denote by $\tilde{C}^\infty(E)$ the space of all measurable functions f on E such that for any multiindex $\gamma \in \mathbf{N}_0^d$ and for all $1 \leq p < \infty$

$$\sup_{(t,x) \in E} |D^\gamma f(t, x)| + \sup_{t \in [0, T]} |D^\gamma f(t, \cdot)|_{L_p(\mathbf{R}^d)} < \infty.$$

Similarly, let $\tilde{C}^\infty(\mathbf{R}^d)$ be the space of all measurable functions f on \mathbf{R}^d such that for any multiindex $\gamma \in \mathbf{N}_0^d$ and for all $1 \leq p < \infty$

$$\sup_{x \in \mathbf{R}^d} |D^\gamma f(x)| + |D^\gamma f|_{L_p(\mathbf{R}^d)} < \infty.$$

Next we suppose that $f \in \tilde{C}^\infty(E)$ and derive some estimates for the solution.

Lemma 8. *Let $f \in \tilde{C}^\infty(E)$ then there is unique $u \in \tilde{C}^\infty(E)$ solving (1.1). Moreover,*

$$(4.1) \quad u(t, x) = \int_0^t e^{-\lambda(t-s)} \mathbf{E} f(s, x + Z_{t-s}^\pi) ds, (t, x) \in E,$$

and for $p \in [1, \infty]$ and any multiindex $\gamma \in \mathbf{N}_0^d$,

$$(4.2) \quad |D^\gamma u|_{L_p(E)} \leq \rho_\lambda |D^\gamma f|_{L_p(E)},$$

$$(4.3) \quad |D^\gamma u(t)|_{L_p(\mathbf{R}^d)} \leq \int_0^t |D^\gamma f(s)|_{L_p(\mathbf{R}^d)} ds, t \geq 0.$$

where $\rho_\lambda = (1/\lambda) \wedge T$.

Proof. Denote $Z_t = Z_t^\pi, t \geq 0$. *Uniqueness.* Let $u_1, u_2 \in \tilde{C}^\infty(E)$ solve (1.1) and $u = u_1 - u_2$. Then u solve (1.1) with $f = 0$. Let $(t, x) \in E$. By Ito formula for $e^{\lambda(t-s)} u(t-s, x + Z_s)$, $0 \leq s \leq t$, we have

$$-u(t, x) = \mathbf{E} \int_0^t e^{\lambda(t-s)} [-\partial_t + L^\pi - \lambda] u(t-s, x + Z_s) ds = 0.$$

Hence $u(t, x) = 0$ for all $(t, x) \in E$.

Existence. Let $f \in \tilde{C}^\infty(E)$. Set

$$u(t, x) = \int_0^t e^{-\lambda(t-s)} \mathbf{E} f(s, x + Z_{t-s}^\pi) ds, (t, x) \in E.$$

Then

$$D^\gamma u(t, x) = \int_0^t e^{-\lambda(t-s)} \mathbf{E} D^\gamma f(s, x + Z_{t-s}^\pi) ds, (t, x) \in E,$$

and (4.3) follows.

By Hölders inequality for $\lambda > 0$,

$$|D^\gamma u|_{L_p(T)}^p \leq \lambda^{-p} \int_0^T \int_0^t \lambda e^{-\lambda(t-s)} |D^\gamma f(s, \cdot)|_{L_p(\mathbf{R}^d)}^p ds dt \leq \lambda^{-p} |D^\gamma f|_{L_p(T)}^p.$$

By Hölder inequality for $\lambda \geq 0$,

$$|D^\gamma u|_{L_p(T)}^p \leq \int_0^T t^{p-1} \int_0^t |D^\gamma f(s, \cdot)|_{L_p(\mathbf{R}^d)}^p ds dt \leq \frac{T^p}{p} |D^\gamma f|_{L_p(T)}^p.$$

We fix $s \in [0, T]$, $x \in \mathbf{R}^d$, and applying Ito formula with $e^{-\lambda r} f(s, x + \cdot)$ and Z_r , $0 \leq r \leq t - s$, we have

$$\begin{aligned} & e^{-\lambda(t-s)} f(s, x + Z_{t-s}) \\ = & f(s, x) + \int_0^{t-s} \int_{\mathbf{R}_0} e^{-\lambda r} [f(s, x + Z_r + y) - f(s, x + Z_{r-})] q(dr, dy) \\ & + \int_0^{t-s} e^{-\lambda r} (L^\pi - \lambda) f(s, x + Z_r) dr \end{aligned}$$

Taking expectation on both sides, and integrating with respect to s , we obtain by Fubini's theorem for each $(t, x) \in E$,

$$\begin{aligned} & \int_0^t e^{-\lambda(t-s)} \mathbf{E} f(s, x + Z_{t-s}) ds \\ = & \int_0^t f(s, x) ds + \int_0^t \int_0^{t-s} e^{-\lambda r} \mathbf{E} [L^\pi - \lambda] f(s, x + Z_r) dr ds \\ = & \int_0^t f(s, x) ds + \int_0^t \int_s^t e^{-\lambda(r-s)} \mathbf{E} [L^\pi - \lambda] f(s, x + Z_{r-s}) dr ds \\ = & \int_0^t f(s, x) ds + \int_0^t \int_0^r e^{-\lambda(r-s)} \mathbf{E} [L^\pi - \lambda] f(s, x + Z_{r-s}) ds dr. \end{aligned}$$

Since for each $(r, x) \in E$

$$\int_0^r e^{-\lambda(r-s)} \mathbf{E} L^\pi f(s, x + Z_{r-s}) ds = L^\pi u(r, x),$$

it follows that for each $(t, x) \in E$,

$$u(t, x) = \int_0^t f(s, x) ds + \int_0^t [L^\pi u(r, x) - \lambda u(r, x)] dr.$$

The statement follows. \square

Similarly we can handle the problem (1.2).

Lemma 9. *Let $f \in \tilde{C}^\infty(\mathbf{R}^d)$ then there is unique $u \in \tilde{C}^\infty(\mathbf{R}^d)$ solving (1.2). Moreover,*

$$(4.4) \quad u(x) = \int_0^\infty e^{-\lambda t} \mathbf{E} f(x + Z_t^\pi) dt, x \in \mathbf{R}^d,$$

and for $p \in [1, \infty]$ and any multiindex $\gamma \in \mathbf{N}_0^d$,

$$(4.5) \quad |D^\gamma u|_{L_p(\mathbf{R}^d)} \leq (1/\lambda) |D^\gamma f|_{L_p(\mathbf{R}^d)}.$$

Proof. Denote $Z_t = Z_t^\pi, t \geq 0$. *Uniqueness.* Let $u_1, u_2 \in \tilde{C}^\infty(E)$ solve (1.1) and $u = u_1 - u_2$. Then u solve (1.1) with $f = 0$. Let $x \in \mathbf{R}^d$. By Ito formula for $e^{-\lambda t} u(x + Z_t), 0 \leq s \leq t$, we have

$$e^{-\lambda t} \mathbf{E} u(x + Z_t) - u(x) = \mathbf{E} \int_0^t e^{-\lambda s} [L^\pi u(x + Z_s) - \lambda u(x + Z_s)] ds = 0.$$

Passing to the limit as $t \rightarrow \infty$ we obtain that $u(x) = 0$ for all $x \in \mathbf{R}^d$.

Existence. Let $f \in \tilde{C}^\infty(\mathbf{R}^d)$. Set

$$u(x) = \int_0^\infty e^{-\lambda t} \mathbf{E} f(x + Z_t^\pi) dt, x \in \mathbf{R}^d.$$

By direct estimate using Hölder inequality, as in Lemma 8, (4.5) readily follows, i.e. $u \in \tilde{C}^\infty(\mathbf{R}^d)$. We fix $x \in \mathbf{R}^d$, and applying Ito formula with $e^{-\lambda t} f(x + Z_t), 0 \leq t$, we have for all $t > 0, x \in \mathbf{R}^d$,

$$e^{-\lambda t} \mathbf{E} f(x + Z_t) = f(x) + \int_0^t e^{-\lambda s} \mathbf{E} (L^\pi - \lambda) f(s, x + Z_s) ds$$

Passing to the limit as $t \rightarrow \infty$ we have

$$L^\pi u(x) - \lambda u(x) + f(x) = 0, x \in \mathbf{R}^d.$$

The statement follows. \square

An obvious consequence of Lemma 9 is

Corollary 5. *Let $f \in \tilde{C}^\infty(\mathbf{R}^d)$ and*

$$u(x) = \int_0^\infty e^{-\lambda t} \mathbf{E} f(x + Z_t^\pi) dt, x \in \mathbf{R}^d.$$

Then $u \in \tilde{C}^\infty(\mathbf{R}^d)$, and

$$\begin{aligned} L^\pi u(x) &= L^\pi \int_0^\infty e^{-\lambda t} \mathbf{E} f(x + Z_t^\pi) dt = \int_0^\infty e^{-\lambda t} \mathbf{E} L^\pi f(x + Z_t^\pi) dt, \\ (L^\pi - \lambda) u(x) &= (L^\pi - \lambda) \int_0^\infty e^{-\lambda t} \mathbf{E} f(x + Z_t^\pi) dt \\ &= \int_0^\infty e^{-\lambda t} \mathbf{E} (L^\pi - \lambda) f(x + Z_t^\pi) dt = -f(x), x \in \mathbf{R}^d. \end{aligned}$$

4.2. L_2 -estimates. We derive first some L_2 -estimates independent of the regularity of f . Given $\pi \in \mathfrak{A}^\sigma$, let

$$\tilde{\psi} = \tilde{\psi}^\pi = \operatorname{Re} \psi^\pi(\xi) = \int [\cos(2\pi y \cdot \xi) - 1] \pi(dy), \xi \in \mathbf{R}^d.$$

Let for $v \in \tilde{C}^\infty(\mathbf{R}^d)$ or $v \in \tilde{C}^\infty(E)$ we define

$$|v|_{\tilde{\psi}, 2} = \left| \mathcal{F}^{-1} \tilde{\psi}^\pi \hat{v} \right|_{L_2(\mathbf{R}^d)} \quad \text{or} \quad |v|_{\tilde{\psi}, 2; E} = \left| \mathcal{F}^{-1} \tilde{\psi}^\pi \hat{v} \right|_{L_2(E)};$$

in the case $v \in \tilde{C}^\infty(E)$, \hat{v} denotes Fourier transform in x .

Lemma 10. *Let $\pi \in \mathfrak{A}^\sigma$.*

a) Let $f \in \tilde{C}^\infty(E)$ and $u \in \tilde{C}^\infty(E)$ be the unique solution to (??), defined in Lemma 8. Then

$$(4.6) \quad |u|_{\tilde{\psi}, 2; E} \leq |f|_{L_2(T)}.$$

Moreover,

$$(4.7) \quad |u|_{L_2(T)} \leq (\lambda^{-1} \wedge T) |f|_{L_2(T)}, |u(t)|_{L_2(\mathbf{R}^d)} \leq \int_0^t |f(s)|_{L_2(\mathbf{R}^d)} ds, t \geq 0,$$

and for $t \geq 0$,

$$(4.8) \quad \int_0^t \int \int [u(s, x+y) - u(s, x)]^2 \pi(dy) dx ds \leq \left(\int_0^t |f(s)|_{L_2(\mathbf{R}^d)} ds \right)^2.$$

b) Let $f \in \tilde{C}^\infty(\mathbf{R}^d)$ and $v \in \tilde{C}^\infty(\mathbf{R}^d)$ be the unique solution to (1.2), defined in Lemma 9. Then

$$(4.9) \quad |v|_{\tilde{\psi}, 2} \leq |f|_{L_2(\mathbf{R}^d)}.$$

Moreover,

$$(4.10) \quad |v|_{L_2(\mathbf{R}^d)} \leq (1/\lambda) |f|_{L_2(\mathbf{R}^d)},$$

and

$$(4.11) \quad \int \int [v(x+y) - v(x)]^2 \pi(dy) dx \leq \frac{2}{\lambda} |f|_{L_2(\mathbf{R}^d)}^2.$$

Proof. a) Let $f \in \tilde{C}^\infty(E)$. Taking Fourier transform in x in the representation (4.1) we find that

$$\hat{u}(t, \xi) = \int_0^t e^{-\lambda(t-s)} \exp\{\psi^\pi(\xi)(t-s)\} \hat{f}(s, \xi) ds, (t, \xi) \in E.$$

Hence, by Hölder inequality, for any $(t, \xi) \in E$,

$$\left| \tilde{\psi}^\pi(\xi) \hat{u}(t, \xi) \right|^2 \leq \left| \tilde{\psi}^\pi(\xi) \right|^2 \int_0^t \exp\left\{ \tilde{\psi}^\pi(\xi)(t-s) \right\} \left| \hat{f}(s, \xi) \right|^2 ds,$$

and, by Fubini theorem,

$$\left| \tilde{\psi}^\pi \hat{u} \right|_{L_2(T)}^2 \leq \left| \hat{f} \right|_{L_2(T)}^2.$$

The inequality (4.7) is derived in Lemma 8. We derive the remaining inequalities using chain rule and integrating. For any $(t, x) \in E$,

$$(4.12) \quad u(t, x)^2 = 2 \int_0^t u(s, x) [L^\pi u(s, x) - \lambda u(s, x) + f(s, x)] ds.$$

Now, for any $(s, x) \in E, y \in \mathbf{R}^d$,

$$(4.13) \quad \begin{aligned} & 2u(s, x) [u(s, x + y) - u(s, x) - \chi_\sigma(y) y \cdot \nabla u(s, x)] \\ &= -[u(s, x + y) - u(s, x)]^2 \\ & \quad + u(s, x + y)^2 - u(s, x)^2 - \chi_\sigma(y) y \cdot \nabla [u(s, x)^2]. \end{aligned}$$

Using (4.13) and integrating both sides of (4.12) in x , we have for any $t \in [0, T]$,

$$\begin{aligned} & |u(t)|_{L_2(\mathbf{R}^d)}^2 + \int_0^t \int \int [u(s, x + y) - u(s, x)]^2 \pi(dy) dx ds \\ & \leq 2 \int_0^t \int f(s, x) u(s, x) dx ds \leq 2 \int_0^t |f(s)|_{L_2(\mathbf{R}^d)} |u(s)|_{L_2(\mathbf{R}^d)} \\ & \leq \left(\int_0^t |f(s)|_{L_2(\mathbf{R}^d)} ds \right)^2 \end{aligned}$$

and (4.8) follows.

b) Let $f \in \tilde{C}^\infty(\mathbf{R}^d)$. Taking Fourier transform in (4.4) we find that

$$\hat{v}(\xi) = \int_0^\infty e^{-\lambda t} \exp\{\psi^\pi(\xi) t\} \hat{f}(\xi) dt, \xi \in \mathbf{R}^d.$$

Hence for any $\xi \in \mathbf{R}^d$,

$$\left| \tilde{\psi}^\pi(\xi) \hat{v}(\xi) \right|^2 \leq \left| \tilde{\psi}^\pi(\xi) \right| \int_0^\infty \exp\{\tilde{\psi}^\pi(\xi) t\} \left| \hat{f}(\xi) \right|^2 dt \leq \left| \hat{f}(\xi) \right|^2,$$

and

$$\left| \tilde{\psi}^\pi \hat{v} \right|_{L_2(\mathbf{R}^d)}^2 \leq \left| \hat{f} \right|_{L_2(\mathbf{R}^d)}^2.$$

Inequality (4.10) was derived in Lemma 9. Multiplying both sides of (5.1) by $2v$ and integrating as in the part a), we have

$$\begin{aligned} & \int \int [v(x + y) - v(x)]^2 \pi(dy) dx \\ & \leq 2 |v|_{L_2(\mathbf{R}^d)} |f|_{L_2(\mathbf{R}^d)} \leq \frac{2}{\lambda} |f|_{L_2(\mathbf{R}^d)}^2. \end{aligned}$$

□

Remark 4. For $v \in \tilde{C}^\infty(\mathbf{R}^d)$, by Plancherel's equality,

$$\begin{aligned} \int \int [v(x + y) - v(x)]^2 \pi(dy) dx &= \int \int |e^{i2\pi\xi \cdot y} - 1|^2 \pi(dy) |\hat{v}(\xi)|^2 d\xi \\ &= 2 \int \left(-\tilde{\psi}^\pi(\xi) \right) |\hat{v}(\xi)|^2 d\xi \\ &= 2 \left| \sqrt{-\tilde{\psi}} \hat{v} \right|_{L_2(\mathbf{R}^d)}^2 = 2 |v|_{(-\tilde{\psi})^{1/2}, 2}^2, \end{aligned}$$

where $|v|_{(-\tilde{\psi})^{1/2},2} = \left| \mathcal{F}^{-1} \sqrt{-\tilde{\psi}} \hat{v} \right|_{L_2(\mathbf{R}^d)}$.

4.3. Continuity of L^π and proof of Theorem 1. We show first the L_2 continuity as follows.

Lemma 11. *Let $\pi, \pi_0 \in \mathfrak{A}^\sigma$. Assume there is $C_0 > 0$ so that*

$$|\psi^\pi(\xi)| \leq C_0 |\psi^{\pi_0}(\xi)|, \xi \in \mathbf{R}^d.$$

Then

$$(4.14) \quad |L^\pi \varphi|_{L_2(\mathbf{R}^d)} \leq C_0 |L^{\pi_0} \varphi|_{L_2(\mathbf{R}^d)}, \varphi \in \tilde{C}^\infty(\mathbf{R}^d).$$

Proof. Let $\varphi \in \tilde{C}^\infty(\mathbf{R}^d)$. By Plancherel equality,

$$|L^\pi \varphi|_{L_2(\mathbf{R}^d)} = |\psi^\pi \hat{\varphi}|_{L_2(\mathbf{R}^d)} \leq C_0 |\psi^{\pi_0} \hat{\varphi}|_{L_2(\mathbf{R}^d)} = C_0 |L^{\pi_0} \varphi|_{L_2(\mathbf{R}^d)}.$$

□

Let $\pi_0 \in \mathfrak{A}^\sigma$, κ be a scaling function with a scaling factor l and **D**(κ, l)(i)-(ii) hold for π_0 . Let $\gamma(t) = \inf(s > 0 : l(s) > t)$, $t > 0$. According to Lemma 5, for each $t > 0$, the associated Levy process $Z_t^{\pi_0}$ has a bounded probability density function $p^{\pi_0}(t, x)$, $x \in \mathbf{R}^d$. Also, $\int |\nabla p^{\pi_0}(t, x)| dx \leq C \gamma(t)^{-1}$, $t > 0$, if $\sigma \in (1, 2)$.

Define for $v \in \tilde{C}^\infty(\mathbf{R}^d)$, $\lambda > 0$, $\varepsilon \geq 0$,

$$\begin{aligned} K_\lambda^\varepsilon v(x) &= \int_\varepsilon^\infty e^{-\lambda t} \mathbf{E} v(x + Z_t^{\pi_0}) dt = \int_\varepsilon^\infty \int e^{-\lambda t} v(x + y) p^{\pi_0}(t, y) dy dt \\ &= \int v(x - y) \int_\varepsilon^\infty e^{-\lambda t} p^{\pi_0^*}(t, y) dt dy \\ &= \int v(y) \int_\varepsilon^\infty e^{-\lambda t} p^{\pi_0^*}(t, x - y) dt dy, x \in \mathbf{R}^d. \end{aligned}$$

Let

$$\begin{aligned} (4.15) \quad T_\lambda^\varepsilon v(x) &= L^\pi K_\lambda^\varepsilon v(x) = \int L^\pi v(x - y) \int_\varepsilon^\infty e^{-\lambda t} p^{\pi_0^*}(t, y) dt dy \\ &= \int_\varepsilon^\infty e^{-\lambda t} \int L^\pi p^{\pi_0^*}(t, x - y) v(y) dy dt \\ &= \int m_\lambda^\varepsilon(x - y) v(y) dy, v \in \tilde{C}^\infty(\mathbf{R}^d), \end{aligned}$$

with

$$m^\varepsilon(x) = m_\lambda^\varepsilon(x) = \int_\varepsilon^\infty e^{-\lambda t} L^\pi p^{\pi_0^*}(t, x) dt, x \in \mathbf{R}^d.$$

Note that according to Lemma 5b) and Corollaries 2 and 1,

$$\int |m_\lambda^\varepsilon(x)| dx < \infty.$$

Lemma 12. *Let $\pi_0 \in \mathfrak{A}^\sigma$, κ be a scaling function with scaling factor l , and $D(\kappa, l)$ hold for π_0 . Let $\pi \in \mathfrak{A}^\sigma$ be such that*

$$\int_{|z| \leq 1} |z|^{\alpha_1} \tilde{\pi}_R(dz) + \int_{|z| > 1} |z|^{\alpha_2} \tilde{\pi}_R(dz) \leq N \quad \forall R > 0,$$

where $\tilde{\pi}_R(dy) = \kappa(R) \pi_R(dy)$, and α_1, α_2 are exponents in assumption $D(\kappa, l)$ for π_0 .

Then for each $p \in (1, \infty)$ there is a constant $C = C(d, p, \kappa, l, N_0, N, N_1, c_1)$ such that

$$|T_\lambda^\varepsilon v|_{L_p} \leq C |v|_{L_p}, v \in \tilde{C}^\infty(\mathbf{R}^d).$$

Proof. 1. First we prove the statement for $p = 2$. Observe that

$$\widehat{m}_\lambda^\varepsilon(\xi) = \psi^\pi(\xi) \int_\varepsilon^\infty \exp\left\{\psi^{\pi_0^*}(\xi)t - \lambda t\right\} dt, \xi \in \mathbf{R}^d.$$

Hence by Lemma 7 and Corollary 4 there is $C = C(N, c_1)$ such that

$$\left|\widehat{m}_\lambda^\varepsilon(\xi)\right| \leq CN_2 \left|\psi^{\pi_0^*}(\xi)\right| \int_0^\infty \exp\left\{\tilde{\psi}^{\pi_0^*}(\xi)t - \lambda t\right\} dt \leq CN_2$$

for all $\xi \in \mathbf{R}^d$, and

$$|T_\lambda^\varepsilon v|_{L_2} \leq C |v|_{L_2}, v \in \tilde{C}^\infty(\mathbf{R}^d).$$

2. Since we already have an L_2 -estimate, according to Theorem 3 of Chapter I in [8], it suffices to show that

$$(4.16) \quad \int_{|x| \geq 3|s|} |m_\lambda^\varepsilon(x-s) - m_\lambda^\varepsilon(x)| dx \leq C, \quad \forall s \neq 0.$$

Let $s \neq 0, R = |s|$. Changing the variable in (4.16), we see that we have to prove that

$$(4.17) \quad R^d \int_{|x| \geq 3} |m_\lambda^\varepsilon(R(x-\hat{s})) - m_\lambda^\varepsilon(Rx)| dx \leq C, \quad |\hat{s}| = 1, R > 0.$$

Let p^{*R} be the pdf corresponding to the Levy measure $\kappa(R) \pi_0^*(Rdy)$, and let $L^R = L^{\tilde{\pi}_R}$ with $\tilde{\pi}_R = \kappa(R) \pi(Rdy)$. For $R > 0$, changing the variables of integration, we have

$$\begin{aligned} m_\lambda^\varepsilon(Rx) &= \int_\varepsilon^\infty e^{-\lambda t} (\nabla_y^\sigma p^{\pi_0^*})(t, Rx) \pi^\varepsilon(dy) dt \\ &= \int_{\varepsilon/\kappa(R)}^\infty e^{-\lambda \kappa(R)t} (\nabla_{Ry}^\sigma p^{\pi_0^*})(\kappa(R)t, Rx) \tilde{\pi}_R(dy) dt \\ &= \int_{\varepsilon/\kappa(R)}^\infty e^{-\lambda \kappa(R)t} R^{-d} (L^R p^{*R})(t, x) dt, x \in \mathbf{R}^d. \end{aligned}$$

In order to prove (3.11) it is enough to show that

$$(4.18) \quad \begin{aligned} & \int_{|x|>3} \int_0^\infty |L^R p^{*R}(t, x - \hat{s}) - L^R p^{*R}(t, x)| dt dx \\ & \leq C, |\hat{s}| = 1, R > 0, \end{aligned}$$

with $C = C(d, p, \kappa, l, N_0, N, N_1)$.

Let $I_1 = \{t > 0 : \gamma(t) \leq 1\}$, $I_2 = \{t > 0 : \gamma(t) > 1\}$. Now,

$$\begin{aligned} & \int_{|x|>3} \int_0^\infty |L^R p^{*R}(t, x - \hat{s}) - L^R p^{*R}(t, x)| dt dx \\ & \leq \int_{|x|>3} \int_{I_1} |L^R p^{*R}(t, x - \hat{s}) - L^R p^{*R}(t, x)| dt dx \\ & \quad + \int_{|x|>3} \int_{I_2} |L^R p^{*R}(t, x - \hat{s}) - L^R p^{*R}(t, x)| dt dx \\ & = A_1 + A_2. \end{aligned}$$

By Lemma 6,

$$\begin{aligned} A_1 &= \int_{|x|>3} \int_{I_1} |L^R p^{*R}(t, x - \hat{s}) - L^R p^{*R}(t, x)| dt dx \\ &\leq 2 \int_{|x|>2} \int_{I_1} |L^R p^{*R}(t, x)| dt dx \\ &\leq C \int_{I_1} \left(1 + t\gamma(t)^{-\alpha_1} + 1_{\sigma \in (1,2)} \gamma(t)^{-1}\right) dt \leq C. \end{aligned}$$

By Fubini theorem,

$$\begin{aligned} A_2 &= \int_{|x|>3} \int_{I_2} |L^R p^{*R}(t, x - \hat{s}) - L^R p^{*R}(t, x)| dt dx \\ &\leq \int_{|x|>3} \int_{I_2} \int_0^1 |L^R \nabla p^{*R}(t, x - r\hat{s})| dr dt dx \\ &\leq 2 \int_{I_2} \int_{|x|>2} |L^R \nabla p^{*R}(t, x)| dx dt \leq 2 \int_{I_2} \|L^R \nabla p^{*R}(t, \cdot)\|_{L_1} dt. \end{aligned}$$

By Lemma 6,

$$\int_{I_2} \|L^R \nabla p^{*R}(t, \cdot)\|_{L_1} dt \leq C \int_{I_2} \gamma(t)^{-(1+\alpha_2)} dt \leq C.$$

The statement is proved. \square

Corollary 6. *Let $\pi_0 \in \mathfrak{A}^\sigma$, κ be a scaling function with scaling factor l , and $D(\kappa, l)$ hold for π_0 . Let $\pi \in \mathfrak{A}^\sigma$ be such that*

$$\int_{|z|\leq 1} |z|^{\alpha_1} \tilde{\pi}_R(dz) + \int_{|z|>1} |z|^{\alpha_2} \tilde{\pi}_R(dz) \leq N \quad \forall R > 0,$$

where $\tilde{\pi}_R(dy) = \kappa(R)\pi_R(dy)$, and α_1, α_2 are exponents in assumption $\mathbf{D}(\kappa)$ for π_0 . Let $v \in \tilde{C}^\infty(\mathbf{R}^d)$ and $u \in \tilde{C}^\infty(\mathbf{R}^d)$ be the unique solution to

$$(4.19) \quad (L^{\pi_0} - \lambda)u = v \text{ in } \mathbf{R}^d.$$

Then for each $p \in (1, \infty)$ there is a constant $C = C(d, p, \kappa, l, N_0, N, N_1, c_1)$ such that

$$|L^\pi u|_{L_p} \leq C |v|_{L_p}.$$

Proof. Let $v \in \tilde{C}^\infty(\mathbf{R}^d)$, $\lambda > 0$. There is a unique $u \in \tilde{C}^\infty(\mathbf{R}^d)$ solving (4.19). According to Lemma 9,

$$L^\pi u(x) = \int_0^\infty e^{-\lambda t} \mathbf{E} L^\pi v(x + Z_t^{\pi_0}) dt, x \in \mathbf{R}^d.$$

By (4.15),

$$(4.20) \quad \begin{aligned} T_\lambda^\varepsilon v(x) &= \int L^\pi v(x - y) \int_\varepsilon^\infty e^{-\lambda t} p^{\pi_0^*}(t, y) dt dy \\ &= \int_\varepsilon^\infty e^{-\lambda t} \mathbf{E} L^\pi v(x + Z_t^{\pi_0}) dt, x \in \mathbf{R}^d. \end{aligned}$$

By Lemma 12, for each $p \in (1, \infty)$ there is a constant $C = C(d, p, \kappa, l, N_0, N, N_1, c_1)$ such that

$$(4.21) \quad |T_\lambda^\varepsilon v|_{L_p} \leq C |v|_{L_p}.$$

Passing to the limit in (4.21) and (4.20) as $\varepsilon \rightarrow 0$, we have

$$|L^\pi u|_{L_p} \leq C |v|_{L_p}.$$

□

Now we prove the continuity of L^π -norm.

Proposition 1. Let $\pi_0 \in \mathfrak{A}^\sigma$, κ be a scaling function with scaling factor l , and $\mathbf{D}(\kappa, l)$ hold for π_0 . Let $\pi \in \mathfrak{A}^\sigma$ be such that

$$\int_{|z| \leq 1} |z|^{\alpha_1} \tilde{\pi}_R(dz) + \int_{|z| > 1} |z|^{\alpha_2} \tilde{\pi}_R(dz) \leq N \quad \forall R > 0,$$

where $\tilde{\pi}_R(dy) = \kappa(R)\pi_R(dy)$, and α_1, α_2 are exponents in assumption $\mathbf{D}(\kappa)$ for π_0 .

Then for each $p \in (1, \infty)$ there is a constant $C = C(d, p, \kappa, l, N_0, N, N_1, c_1)$ such that

$$|L^\pi f|_{L_p} \leq C |L^{\pi_0} f|_{L_p}, f \in \tilde{C}^\infty(\mathbf{R}^d).$$

Proof. Let $f \in \tilde{C}^\infty(\mathbf{R}^d)$, $\lambda > 0$. Then $v = (L^{\pi_0} - \lambda)f \in \tilde{C}^\infty(\mathbf{R}^d)$, and, by Corollary 6, there is $C = C(d, p, \kappa, \mu^0, N_2, N, C_0)$ such that

$$|L^\pi f|_{L_p} \leq C |v|_{L_p} = C |(L^{\pi_0} - \lambda)f|_{L_p}.$$

Since C does not depend on $\lambda > 0$, the statement follows. □

4.3.1. *Proof of Theorem 1. Existence.* Let $f \in L_p(\mathbf{R}^d)$. There is a sequence $f_n \in C_0^\infty(\mathbf{R}^d)$ such that $f_n \rightarrow f$ in L_p . For each n , there is unique $u_n \in \tilde{C}^\infty(\mathbf{R}^d)$ solving (1.2). Hence

$$(L^\pi - \lambda)(u_n - u_m) = f_n - f_m.$$

By Corollary 6 and Lemma 9,

$$\begin{aligned} |L^{\pi_0}(u_n - u_m)|_{L_p} &\leq C|f_n - f_m|_{L_p} \rightarrow 0, \\ |u_n - u_m|_{L_p} &\leq \frac{1}{\lambda}|f_n - f_m|_{L_p} \rightarrow 0, \end{aligned}$$

as $n, m \rightarrow \infty$. Hence there is $u \in H_p^{\pi_0}$ so that $u_n \rightarrow u$ in $H_p^{\pi_0}$. Using Proposition 1, we can pass to the limit in $(L^\pi - \lambda)u_n = f_n$ as $n \rightarrow \infty$. Obviously, $(L^\pi - \lambda)u = f$ in L_p .

Uniqueness. Assume $u_1, u_2 \in H_p^{\pi_0}(\mathbf{R}^d)$ solve (1.2). Then $u = u_1 - u_2 \in H_p^{\pi_0}$ solves $(L^\pi - \lambda)u = 0$, i.e. $\forall \varphi \in \tilde{C}^\infty(\mathbf{R}^d)$

$$\int \varphi (L^\pi - \lambda)u = \int u (L^{\pi^*} - \lambda) \varphi dx = 0$$

According to Lemma 9, $\int u f dx = 0 \forall f \in \tilde{C}^\infty(\mathbf{R}^d)$. Hence $u = 0$ a.e. The statement is proved.

4.4. **Proof of Theorem 2.** Let $f \in \tilde{C}^\infty(E)$ and $u \in \tilde{C}^\infty(E)$ be the solution to

$$\begin{aligned} (4.22) \quad \partial_t u &= L^\pi u - \lambda u + f \text{ in } E, \\ u(0, \cdot) &= 0. \end{aligned}$$

By Lemma 5, the associated process Z_t^π has a density function $p^\pi(t, x)$, $x \in \mathbf{R}^d$. Then $p^{\pi^*}(t, x) = p^\pi(t, -x)$, $x \in \mathbf{R}^d$, is pdf of $Z_t^{\pi^*}$. By Lemma 8,

$$\begin{aligned} u(t, x) &= \int_0^t e^{-\lambda(t-s)} \mathbf{E} f(s, x + Z_{t-s}^\pi) ds \\ &= \int_0^t \int e^{-\lambda(t-s)} f(s, x - y) p^{\pi^*}(t - s, y) dy ds \\ &= \int_0^t \int e^{-\lambda(t-s)} p^{\pi^*}(t - s, x - y) f(s, y) dy ds, \end{aligned}$$

and

$$\begin{aligned} (4.23) \quad \mathcal{L}^{\pi_0} u(t, x) &= \int_0^t e^{-\lambda(t-s)} \mathbf{E} L^{\pi_0} f(s, x + Z_{t-s}^\pi) ds \\ &= \int_0^t \int e^{-\lambda(t-s)} L^{\pi_0} f(s, x - y) p^{\pi^*}(t - s, y) dy ds \\ &= \int_0^t \int e^{-\lambda(t-s)} (L^{\pi_0} p^{\pi^*})(t - s, x - y) f(s, y) dy ds. \end{aligned}$$

Hence

$$L^{\pi_0} u(t, x) = \int_{-\infty}^{\infty} \int e^{-\lambda(t-s)} K(t-s, x-y) \chi_{[0, \infty)}(t-s) f(s, y) ds dy,$$

where $f(s, y) = \chi_{[0, \infty)}(s) f(s, y)$, $(s, y) \in \mathbf{R}^{d+1}$,

$$K(t, x) = L^{\pi_0} p^{\pi^*}(t, x) \chi_{[0, \infty)}(t), (t, x) \in \mathbf{R}^{d+1}.$$

Let $\varepsilon \in (0, 1)$,

$$K^\varepsilon(t, x) = L^{\pi_0} p^{\pi^*}(t, x) \chi_{[\varepsilon, \infty)}(t), (t, x) \in \mathbf{R}^{d+1},$$

and consider for $h \in C_0^\infty(\mathbf{R}^{d+1})$,

$$\begin{aligned} (4.24) T_\lambda^\varepsilon h(t, x) &= \int_{-\infty}^{t-\varepsilon} \int e^{-\lambda(t-s)} p^{\pi^*}(t-s, x-y) L^{\pi_0} h(s, y) dy ds \\ &= \int_{-\infty}^{\infty} \int K_\lambda^\varepsilon(t-s, x-y) h(s, y) dy ds, (t, x) \in \mathbf{R}^d, \end{aligned}$$

where $K_\lambda^\varepsilon(t, x) = e^{-\lambda t} K^\varepsilon(t, x)$, $(t, x) \in \mathbf{R} \times \mathbf{R}^d = \mathbf{R}^{d+1}$.

Claim 1. For each $p \in (1, \infty)$ there is a constant $C = C(d, p, l, N, N_0, N_1, c_1)$ such that

$$(4.25) \quad |T_\lambda^\varepsilon h|_{L_p} \leq C |h|_{L_p}, h \in C_0^\infty(\mathbf{R}^{d+1}),$$

here $L_p = L_p(\mathbf{R}^{d+1})$.

Proof. We will apply Calderon-Zygmund theorem (see Theorem 5 in Appendix). First we will prove the estimate in L_2 . Then, according to Theorem 5, the proof reduces to verification of Hörmander condition (see below).

1. We start with $p = 2$. Obviously,

$$\left| \widehat{T_\lambda^\varepsilon h}(t, \xi) \right| \leq \int_{-\infty}^t e^{-\lambda(t-s)} |\psi^{\pi_0}(\xi)| \exp \left\{ \operatorname{Re} \psi^{\pi^*}(\xi)(t-s) \right\} |\hat{h}(s, \xi)| ds,$$

$(t, \xi) \in E$. Hence by Hölder inequality, Fubini theorem, Lemma 7 and Corollary 4, we have $|\psi^{\pi_0}(\xi)| \leq C |\operatorname{Re} \psi^{\pi^*}(\xi)|$, $\xi \in \mathbf{R}^d$, for some $C = C(N, c_1, l)$ and

$$\left| \widehat{T_\lambda^\varepsilon h} \right|_{L_2}^2 \leq C |h|_{L_2(T)},$$

i.e., (4.25) follows (cf. the proof of Lemma 10).

2. We prove (4.25) for $p \in (1, 2)$ using a version of Calderon-Zygmund theorem (Theorem 5 in Appendix). Let \mathbb{Q} be the collection of sets $Q_\delta = Q_\delta(t, x) = (t - \kappa(\delta), t + \kappa(\delta)) \times B_\delta(x)$, $(t, x) \in \mathbf{R} \times \mathbf{R}^d = \mathbf{R}^{d+1}$, $\delta > 0$.

Note

(i) $(t, x) \notin (s - \kappa(c\delta), s + \kappa(c\delta)) \times B_{c\delta}(y) \iff (t-s, x-y) \notin Q_{c\delta}(0) \iff |t-s| \geq \kappa(c\delta) \text{ or } |x-y| \geq c\delta;$

(ii) $(\bar{s}, \bar{y}) \in Q_\delta(s, y) \iff (\bar{s} - s, \bar{y} - y) \in Q_\delta(0) \iff |s - \bar{s}| < \kappa(\delta) \text{ and } |y - \bar{y}| < \delta.$

According to Theorem 5, it is enough to show

$$(4.26) \quad \int \chi_{Q_{c\delta}(0)^c}(t-s, x-y) |K_\lambda^\varepsilon(t-\bar{s}, x-\bar{y}) - K_\lambda^\varepsilon(t-s, x-y)| dxdt \leq A$$

for all $(\bar{s}, \bar{y}) \in Q_\delta(s, y) \iff (\bar{s}-s, \bar{y}-y) \in Q_\delta(0)$. Equivalently, we have to prove that

$$\begin{aligned} & \int \chi_{Q_{c\delta}(0)^c}(t, x) |K_\lambda^\varepsilon(t-\tilde{s}, x-\tilde{y}) - K_\lambda^\varepsilon(t, x)| dxdt \\ &= \delta^d \kappa(\delta) \int \chi_{Q_{c\delta}(0)^c}(\kappa(\delta)t, \delta x) \left| K_\lambda^{\varepsilon, \delta}(t-\hat{s}, x-\hat{y}) - K_\lambda^{\varepsilon, \delta}(t, x) \right| dxdt \\ &\leq A \quad \forall |\hat{s}| \leq 1, |\hat{y}| \leq 1, \end{aligned}$$

where

$$K_\lambda^{\varepsilon, \delta}(t, x) = K_\lambda^\varepsilon(\kappa(\delta)t, \delta x), (t, x) \in \mathbf{R}^{d+1}.$$

Fix $c > 4$ such that $(\bar{c}-1, \infty) \subseteq I_2$ with $\bar{c} = l(1/c)^{-1} > 3$. Let $G = (-\bar{c}, \bar{c}) \times B_c(0)$. Since $\chi_G(t, x) \leq \chi_{Q_{c\delta}(0)}(\kappa(\delta)t, \delta x)$, it is enough to prove that

$$(4.27) \quad \delta^d \kappa(\delta) \int_{\mathbf{R}^{d+1}} \chi_{G^c}(t, x) \left| K_\lambda^{\varepsilon, \delta}(t-\hat{s}, x-\hat{y}) - K_\lambda^{\varepsilon, \delta}(t, x) \right| dxdt \leq A$$

for all $|\hat{s}| \leq 1, |\hat{y}| \leq 1$. Since

$$\delta^d \kappa(\delta) K_\lambda^{\varepsilon, \delta}(t, x) = e^{-\lambda t} L^{0, \delta} p^{\tilde{\pi}_\delta^*}(t, x) \chi_{(\varepsilon/\kappa(\delta), \infty)}(t), (t, x) \in \mathbf{R}^{d+1},$$

with $L^{0, \delta} = L^{\tilde{\pi}_{0, \delta}}$, $\tilde{\pi}_{0, \delta}(dy) = \kappa(\delta) \pi_0(\delta dy)$, $\tilde{\pi}_\delta^*(dy) = \kappa(\delta) \pi^*(\delta dy)$, we rewrite (4.27) as

$$\begin{aligned} B &= \int_{\mathbf{R}^{d+1}} \chi_{G^c}(t, x) |e^{-\lambda(t-\hat{s})} L^{0, \delta} p^{\tilde{\pi}_\delta^*}(t-\hat{s}, x-\hat{y}) \chi_{(\varepsilon/\kappa(\delta), \infty)}(t-\hat{s}) \\ (4.28) \quad &- e^{-\lambda t} L^{0, \delta} p^{\tilde{\pi}_\delta^*}(t, x) \chi_{(\varepsilon/\kappa(\delta), \infty)}(t)| dxdt \\ &\leq A \end{aligned}$$

Since $G^c \subseteq \{(t, x) : |t| \leq \bar{c}, |x| \geq 3\} \cup \{(t, x) : |t| \geq \bar{c}\} = G_1 \cup G_2$,

$$B \leq \int_{G_1} \dots + \int_{G_2} \dots = B_1 + B_2.$$

By Lemma 6 a),

$$\begin{aligned} B_1 &\leq 2 \int_0^{\bar{c}+1} \int_{|x|>2} \left| L^{0, \delta} p^{\tilde{\pi}_\delta^*}(t, x) \right| dt dx \\ &\leq C \int_0^{\bar{c}+1} \left(1 + t \gamma(t)^{-\alpha_1} + 1_{\sigma \in (1, 2)} \gamma(t)^{-1} \right) dt. \end{aligned}$$

Estimate of B_2 . We have

$$\begin{aligned}
 & B_2 \\
 & \leq \int_{\bar{c}}^{\infty} \int |L^{0,\delta} p^{\pi^*}(t, x)| \left| \chi_{(\varepsilon/\kappa(\delta), \infty)}(t) - \chi_{(\varepsilon/\kappa(\delta), \infty)}(t - \hat{s}) \right| dt dx \\
 & \quad + \int_{\bar{c}}^{\infty} \int |e^{-\lambda(t-\hat{s})} L^{0,\delta} p^{\pi^*}(t - \hat{s}, x - \hat{y}) - e^{-\lambda t} L^{0,\delta} p^{\pi^*}(t, x)| dt dx \\
 & = B_{21} + B_{22}.
 \end{aligned}$$

Then $B_{21} \leq C$ by Lemma 5 b). Since

$$\begin{aligned}
 & e^{-\lambda(t-\hat{s})} L^{0,\delta} p^{\pi^*}(t - \hat{s}, x - \hat{y}) - e^{-\lambda t} L^{0,\delta} p^{\pi^*}(t, x) \\
 & = \int_0^1 [-\lambda e^{-\lambda(t-r\hat{s})} L^{0,\delta} p^{\pi^*}(t - r\hat{s}, x - r\hat{y}) \\
 & \quad + e^{-\lambda(t-r\hat{s})} L^{0,\delta} \partial_t p^{\pi^*}(t - r\hat{s}, x - r\hat{y})] (-\hat{s}) dr \\
 & \quad + \int_0^1 e^{-\lambda(t-r\hat{s})} \hat{y} \cdot L^{0,\delta} \nabla p^{\pi^*}(t - r\hat{s}, x - r\hat{y}) dr
 \end{aligned}$$

we have

$$\begin{aligned}
 & B_{22} \\
 & \leq \int_{\bar{c}}^{\infty} \int \int_0^1 \lambda e^{-\lambda(t-r\hat{s})} |L^{0,\delta} p^{\pi^*}(t - r\hat{s}, x - r\hat{y})| dr dt dx \\
 & \quad + \int_{\bar{c}}^{\infty} \int \int_0^1 |L^{0,\delta} \partial_t p^{\pi^*}(t - r\hat{s}, x - r\hat{y})| dr dt dx \\
 & \quad + \int_{\bar{c}}^{\infty} \int \int_0^1 |L^{0,\delta} \nabla p^{\pi^*}(t - r\hat{s}, x - r\hat{y})| dr dt dx \\
 & = b_1 + b_2 + b_3.
 \end{aligned}$$

By Lemma 5 b),

$$b_1 \leq \int_{\bar{c}-1}^{\infty} \int \lambda e^{-\lambda t} |L^{0,\delta} p^{\pi^*}(t, x)| dt dx \leq C.$$

Since $\partial_t p^{\pi^*}(t, x) = L^{\pi} p(t, x)$, $t > 0$, we have by Lemma 6 c),

$$\begin{aligned}
 b_2 & = \int_{\bar{c}}^{\infty} \int \int_0^1 |L^{0,\delta} L^{\pi} p^{\pi^*}(t - r\hat{s}, x - r\hat{y})| dr dt dx \\
 & \leq \int_{\bar{c}-1}^{\infty} \int |L^{0,\delta} L^{\pi} p^{\pi^*}(t, x)| dt dx \leq C \int_{\bar{c}-1}^{\infty} \gamma(t)^{-2\alpha_2} dt.
 \end{aligned}$$

Finally, by Lemma 6 b),

$$\begin{aligned}
 b_3 & \leq \int_{\bar{c}-1}^{\infty} \int |L^{0,\delta} \nabla p^{\pi^*}(t, x)| dt dx \\
 & \leq C \int_{\bar{c}-1}^{\infty} \gamma(t)^{-1-\alpha_2} dt.
 \end{aligned}$$

Claim is proved for $p \in (1, 2)$ by Theorem 5.

3. We prove the statement for $p > 2$ in a standard way (by duality argument). First note that

$$L^{\pi_0} p^{\pi^*}(t, -x) = L^{\pi_0^*} p^{\pi}(t, x), (t, x) \in \mathbf{R}^{d+1},$$

and let

$$\tilde{K}^\varepsilon(t, x) = K^\varepsilon(t, -x) = L^{\pi_0^*} p^{\pi}(t, x) \chi_{[\varepsilon, \infty)}(t), (t, x) \in \mathbf{R}^{d+1},$$

and

$$\tilde{T}_\lambda^\varepsilon g(s, y) = \int e^{-\lambda(s-t)} \tilde{K}^\varepsilon(s-t, y-x) g(t, x) dt dx, (s, y) \in \mathbf{R}^{d+1}.$$

Let $1/p + 1/q = 1, h, g \in C_0^\infty(\mathbf{R}^{d+1})$. Then, denoting $\tilde{g}(t, x) = g(-t, x), (t, x) \in \mathbf{R}^{d+1}$, we have (by Fubini theorem and changing the variable of integration)

$$\begin{aligned} & \int T_\lambda^\varepsilon h(t, x) g(t, x) dt dx \\ &= \int \int e^{-\lambda(t-s)} K^\varepsilon(t-s, x-y) h(s, y) ds dy g(t, x) dt dx \\ &= \int \int e^{-\lambda(s-t)} \tilde{K}^\varepsilon(s-t, y-x) g(-t, x) dt dx h(-s, y) ds dy \\ &= \int \tilde{T}_\lambda^\varepsilon \tilde{g}(s, y) h(-s, y) ds dy, \end{aligned}$$

and (4.25) holds for $\tilde{T}_\lambda^\varepsilon$ and $q \in (1, 2)$ (see Corollary 4). Hence by Hölder inequality,

$$\begin{aligned} & \left| \int T_\lambda^\varepsilon h(t, x) g(t, x) dt dx \right| \\ & \leq \left| \tilde{T}_\lambda^\varepsilon \tilde{g} \right|_{L_q} \|h\|_{L_p} \leq C \|g\|_{L_q} \|h\|_{L_p}, \end{aligned}$$

and (4.25) holds for $p > 2$ as well, that is for all $p \in (1, \infty)$. The claim is proved. \square

Now, we see that for $h(t, x) = \chi_{[0, T]}(t) f(t, x)$ with $f \in \tilde{C}(E)$,

$$(4.29) \quad T_\lambda^\varepsilon h(t, x) = \int_0^{t-\varepsilon} e^{-\lambda(t-s)} \mathbf{E} L^{\pi_0} f(s, x + Z_{t-s}^\pi) ds, (t, x) \in E.$$

If $u \in \tilde{C}^\infty(E)$ solves (4.22), then

$$(4.30) \quad L^{\pi_0} u(t, x) = \int_0^t e^{-\lambda(t-s)} \mathbf{E} L^{\pi_0} f(s, x + Z_{t-s}^\pi) ds,$$

and $|T_\lambda^\varepsilon h - L^{\pi_0} u|_{L_p} \rightarrow 0$ as $\varepsilon \rightarrow 0$. Since the constant $C = C(d, p, l, N, N_0, N_1, c_1)$ in the above Claim 1 does not depend on ε , passing to the limit in it we get the estimate

$$(4.31) \quad |L^{\pi_0} u|_{L_p} \leq C \|f\|_{L_p}.$$

We finish the proof of Theorem 2 the same way as the proof of Theorem 1. Let $f \in L_p(\mathbf{R}^d)$. There is a sequence $f_n \in \tilde{C}^\infty(E)$ such that $f_n \rightarrow f$ in $L_p(E)$. For each n , there is unique $u_n \in \tilde{C}^\infty(E)$ solving (1.1). Hence

$$\partial_t(u_n - u_m) = (L^\pi - \lambda)(u_n - u_m) + f_n - f_m.$$

By (4.31) and Lemma 8,

$$\begin{aligned} (4.32) \quad |L^{\pi_0}(u_n - u_m)|_{L_p(E)} &\leq C|f_n - f_m|_{L_p(E)} \rightarrow 0, \\ |u_n - u_m|_{L_p(E)} &\leq \left(\frac{1}{\lambda} \wedge T\right) |f_n - f_m|_{L_p(E)} \rightarrow 0, \\ |u_n(t) - u_m(t)|_{L_p(\mathbf{R}^d)} &\leq |f_n - f_m|_{L_p(E)} \quad \forall t \in [0, T], \end{aligned}$$

as $n, m \rightarrow \infty$. Hence there is $u \in \mathcal{H}_p^{\pi_0}$ so that $u_n \rightarrow u$ in $\mathcal{H}_p^{\pi_0}$. Moreover,

$$(4.33) \quad \sup_{t \leq T} |u_n(t) - u(t)|_{L_p(\mathbf{R}^d)} \rightarrow 0,$$

and, according to Proposition 1,

$$(4.34) \quad |L^\pi f|_{L_p(E)} \leq C|L^{\pi_0} f|_{L_p(E)}, f \in \tilde{C}^\infty(E).$$

Hence (see (4.32)-(4.34)) we can pass to the limit in the equation

$$(4.35) \quad u_n(t) = \int_0^t [L^\pi u_n(s) - \lambda u_n(s) + f_n(s)] ds, 0 \leq t \leq T.$$

Obviously, (4.35) holds for u and f . We proved the existence part of Theorem 2.

Uniqueness. Assume $u_1, u_2 \in \mathcal{H}_p^{\pi_0}$ solve (1.1). Then $u = u_1 - u_2 \in \mathcal{H}_p^{\pi_0}$ solves (4.22) with $f = 0$. Now, let $\varphi \in \tilde{C}^\infty(E)$, and $\tilde{\varphi}(t, x) = \varphi(T - t, x)$, $(t, x) \in E$. By Lemma 8, there is unique $\tilde{v} \in \tilde{C}^\infty(E)$ solving (4.22) with $f = \tilde{\varphi}$ and π^* instead of π . Let $v(t, x) = \tilde{v}(T - t, x)$, $(t, x) \in E$. Then $\partial_t v + L^{\pi^*} v - \lambda v + \varphi = 0$ in E and $v(T) = v(T, \cdot) = 0$. Integrating by parts,

$$\begin{aligned} \int_E \varphi u &= \int_E u \left(-\partial_t v - L^{\pi^*} v + \lambda v \right) \\ &= \int_E v (\partial_t u - L^\pi u + \lambda u) = 0. \end{aligned}$$

Hence $\int_E u \varphi dx = 0 \quad \forall \varphi \in \tilde{C}^\infty(E)$. Hence $u = 0$ a.e. Theorem 2 is proved.

5. APPENDIX

Given a function $\kappa : (0, \infty) \rightarrow (0, \infty)$, consider the collection \mathbb{Q} of sets $Q_\delta = Q_\delta(t, x) = (t - \kappa(\delta), t + \kappa(\delta)) \times B_\delta(x)$, $(t, x) \in \mathbf{R} \times \mathbf{R}^d = \mathbf{R}^{d+1}$, $\delta > 0$. The volume $|Q_\delta(t, x)| = c_0 \kappa(\delta) \delta^d$. We will need the following assumptions.

A1. κ is continuous, $\lim_{\delta \rightarrow 0} \kappa(\delta) = 0$ and $\lim_{\delta \rightarrow \infty} \kappa(\delta) = \infty$.

A2. There is a constant C_1 and a nondecreasing continuous function $l(\varepsilon), \varepsilon > 0$, such that $\lim_{\varepsilon \rightarrow 0} l(\varepsilon) = 0$ and

$$\kappa(\varepsilon r) \leq l(\varepsilon) \kappa(r), r > 0, \varepsilon > 0.$$

Since $Q_\delta(t, x)$ not "exactly" increases in δ , we present the basic estimates involving maximal functions based on the system $\mathbb{Q} = \{Q_\delta\}$.

5.1. Vitali Lemma, maximal functions. We start with engulfing property.

Lemma 13. *Let A2 hold. If $Q_\delta(t, x) \cap Q_{\delta'}(r, z) \neq \emptyset$ with $\delta' \leq \delta$, then there is $K_0 \geq 3$ such that $Q_{K_0\delta}(t, x)$ contains both, $Q_\delta(t, x)$ and $Q_{\delta'}(r, z)$, and*

$$|Q_\delta(t, x)| \leq |Q_{K_0\delta}(t, x)| \leq K_0^d l(K_0) |Q_\delta(t, x)|.$$

Proof. Let $(s, y) \in Q_\delta(t, x) \cap Q_{\delta'}(r, z)$ with $\delta' \leq \delta$. If $(r', z') \in Q_{\delta'}(r, z)$, then $|z' - x| \leq 3\delta$, and using **A2**,

$$|r' - t| \leq |r' - r| + |r - s| + |s - t| \leq 2\kappa(\delta') + \kappa(\delta) \leq [2l(1) + 1]\kappa(\delta).$$

We choose $K_0 \geq 3$ so that $[2l(1) + 1]l(1/K_0) \leq 1$. By **A2**,

$$[2l(1) + 1]\kappa(\delta) \leq [2l(1) + 1]l(1/K_0)\kappa(K_0\delta) \leq \kappa(K_0\delta).$$

Hence $Q_{\delta'}(r, z) \subseteq Q_{K_0\delta}(t, x)$ and, obviously, $Q_\delta(t, x) \subseteq Q_{K_0\delta}(t, x)$. Also,

$$|Q_{K_0\delta}(t, x)| = c_0 K_0^d \delta^d \kappa(K_0\delta) \leq c_0 K_0^d \delta^d l(K_0) \kappa(\delta) = K_0^d l(K_0) |Q_\delta(t, x)|.$$

□

Now, following 3.1.1 in [8], we prove Vitali covering lemma.

Lemma 14. *Let $E \subseteq \mathbf{R} \times \mathbf{R}^d$ be a union of a finite collection $\{Q'\}$ of sets from the system $\{Q_\delta(t, x) : (t, x) \in \mathbf{R}^{d+1}, \delta > 0\}$ and **A2** hold.*

There is a positive $c = \frac{1}{K_0^d l(K_0)}$ and a disjoint subcollection

$\{Q^k = Q_{\delta_k}(t_k, x_k), 1 \leq k \leq m\}$ such that

$$\sum_{k=1}^m |Q^k| \geq c |E|.$$

Proof. Let $Q^1 = Q_{\delta_1}(t_1, x_1)$ be the set of the collection $\{Q'\}$ with maximal δ . Let $Q^2 = Q_{\delta_2}(t_2, x_2)$ be the set with maximal δ among remaining sets in $\{Q'\}$ that do not intersect Q^1 . According to Lemma 13, $Q_{K_0\delta_1}(t_1, x_1)$ contains Q^1 and all Q_δ in $\{Q'\}$ that intersect Q^1 and such that $\delta \leq \delta_1$. Continuing we get $Q_{K_0\delta_k}(t_k, x_k)$ containing $Q^k = Q_{\delta_k}(t_k, x_k)$ and all Q_δ in $\{Q'\}$ that intersect Q^k and such that $\delta \leq \delta_k$. So we obtain a finite disjoint subcollection $\{Q^k = Q_{\delta_k}(t_k, x_k), 1 \leq k \leq m\}$ such that $\cup_{k=1}^m Q_{K_0\delta_k}(t_k, x_k) \supseteq Q_\delta$ for any Q_δ in $\{Q'\}$. Hence $\cup_{k=1}^m Q_{K_0\delta_k}(t_k, x_k) \supseteq E$, and by Lemma 13,

$$|E| \leq \sum_{k=1}^m |Q_{K_0\delta_k}| \leq K_0^d l(K_0) \sum_{k=1}^m |Q^k|.$$

□

Remark 5. *The statement of the Lemma 14 still holds if instead of **A2** we assume that there is a constant C so that $C\kappa(\delta) \geq \kappa(\delta')$ whenever $\delta \geq \delta'$.*

Following [8], for a locally integrable function $f(t, x)$ on \mathbf{R}^{d+1} we define

$$(A_\delta f)(t, x) = \frac{1}{|Q_\delta(t, x)|} \int_{Q_\delta(t, x)} f(s, y) ds dy, (t, x) \in \mathbf{R} \times \mathbf{R}^d, \delta > 0$$

and the maximal function of f by

$$\mathcal{M}f(t, x) = \sup_{\delta > 0} (A_\delta |f|)(t, x), (t, x) \in \mathbf{R}^{d+1}.$$

We use collection \mathbb{Q} to define a larger, noncentered maximal function of f , as

$$\widetilde{\mathcal{M}}f(t, x) = \sup_{(t, x) \in Q} \frac{1}{|Q|} \int_Q |f(s, y)| ds dy, (t, x) \in \mathbf{R}^{d+1},$$

where sup is taken over all $Q \in \mathbb{Q}$ that contain (t, x) .

Remark 6. *Let **A2** hold and K_0 be a constant in Lemma 13. For a locally integrable f on \mathbf{R}^{d+1} ,*

$$\mathcal{M}f \leq \widetilde{\mathcal{M}}f \leq \frac{1}{K_0^{dl}(K_0)} \mathcal{M}f.$$

Indeed, if $(t, x) \in Q' = Q_\delta(t', x')$, then by Lemma 13

$$\frac{1}{|Q'|} \int_{Q'} |f| \leq \frac{K_0^{dl}(K_0)}{|Q_{K_0\delta}(t, x)|} \int_{Q_{K_0\delta}(t, x)} |f|.$$

Note $\widetilde{\mathcal{M}}f$ is lower semicontinuous as a sup of lower semicontinuous functions.

Theorem 1.3.1 in [8] holds for \mathbb{Q} (we sketch its proof).

Theorem 3. *Let **A2** hold and f be measurable function on $\mathbf{R}^{d+1} = \mathbf{R} \times \mathbf{R}^d$.*

- (a) *If $f \in L_p, 1 \leq p \leq \infty$, then $\mathcal{M}f$ is finite a.e.*
- (b) *If $f \in L_1$, then for every $\alpha > 0$,*

$$|\{\mathcal{M}f(t, x) > \alpha\}| \leq \frac{c}{\alpha} \int |f| dm.$$

- (c) *If $f \in L_p, 1 < p \leq \infty$, then $\mathcal{M}f \in L_p$ and*

$$|\mathcal{M}f|_{L_p} \leq N_p |f|_{L_p},$$

where N_p depends only on p, l and K_0 .

Proof. (b) Let $E_\alpha = \{\widetilde{\mathcal{M}}f(t, x) > \alpha\}$ and $E \subseteq E_\alpha$ be any compact subset.

Since $\widetilde{\mathcal{M}}f$ is lower semicontinuous, E_α is open. By definition of $\widetilde{\mathcal{M}}f$ for each $(t, x) \in E$, there is $Q \in \mathbb{Q}$ so that $(t, x) \in Q$ and

$$|Q| \leq \frac{1}{\alpha} \int_Q |f|.$$

Since E is compact there exist a finite number $Q_{\delta_1}(t_1, x_1), \dots, Q_{\delta_n}(t_n, x_n) \in \mathbb{Q}$ so that $E \subseteq \cup_{j=1}^n Q_{\delta_j}(t_j, x_j)$. By Lemma 14, there is a subcovering of disjoint sets Q^1, \dots, Q^m so that

$$|E| \leq c \sum_{k=1}^m |Q^k| \leq \frac{c}{\alpha} \sum_{k=1}^m \int_{Q^k} |f| \leq \frac{c}{\alpha} \int |f|.$$

with $c = K_0^{dl}(K_0)$. Taking sup over all such compacts E we get (b).

c) Let $f_1 = f \chi_{\{|f| > \alpha/2\}}$. Note that $\widetilde{\mathcal{M}}f \leq \widetilde{\mathcal{M}}f_1 + \frac{\alpha}{2}$. Hence by part (b)

$$\left| \left\{ \widetilde{\mathcal{M}}f > \alpha \right\} \right| \leq \left| \left\{ \widetilde{\mathcal{M}}f_1 > \alpha/2 \right\} \right| \leq \frac{2c}{\alpha} \int_{|f| > \alpha/2} |f| dm.$$

On the other hand,

$$\begin{aligned} \int \left(\widetilde{\mathcal{M}}f \right)^p &= p \int_0^\infty \left| \left\{ \widetilde{\mathcal{M}}f > \alpha \right\} \right| \alpha^{p-1} d\alpha \leq p \frac{2c}{\alpha} \int_0^\infty \int_{|f| > \alpha/2} |f| \alpha^{p-1} d\alpha \\ &= 2cp \int \int_0^{2|f|} \alpha^{p-2} d\alpha |f| = c \frac{p}{p-1} 2^p \int |f|^p. \end{aligned}$$

□

Corollary 7. *Let $f \in L_1$. Then*

$$\lim_{\delta \rightarrow 0} A_\delta f(t, x) = f(t, x) \text{ a.e.}$$

and $|f(t, x)| \leq \mathcal{M}f(t, x)$ a.e. Moreover, for every $\alpha > 0$,

$$\left| \left\{ \widetilde{\mathcal{M}}f(t, x) > \alpha \right\} \right| \leq \frac{2c}{\alpha} \int_{\{\widetilde{\mathcal{M}}f(t, x) > \alpha/2\}} |f| dm,$$

where c is a constant in Theorem 3.

Proof. Let $f \in L_1, \varepsilon > 0$. There is $g \in C_c(\mathbf{R}^{d+1})$ so that $|f - g|_{L_1} \leq \varepsilon$. Let $\eta > 0$. Since g is uniformly continuous, for all (t, x)

$$\begin{aligned} &|A_\delta g(t, x) - g(t, x)| \\ &\leq \frac{1}{|Q_\delta(t, x)|} \int_{Q_\delta(t, x)} |g(s, y) - g(t, x)| ds dy \leq \eta \end{aligned}$$

if $\delta \leq \delta_0$ for some $\delta_0 > 0$. Hence $\sup_{t, x} |A_\delta g(t, x) - g(t, x)| \rightarrow 0$ as $\delta \rightarrow 0$.

Now for $(t, x) \in \mathbf{R}^{d+1}$,

$$\begin{aligned} &\limsup_{\delta \rightarrow 0} |A_\delta f(t, x) - f(t, x)| \\ &\leq \limsup_{\delta \rightarrow 0} |A_\delta f(t, x) - A_\delta g(t, x)| + |g(t, x) - f(t, x)| \\ &\leq \mathcal{M}(f - g)(t, x) + |g(t, x) - f(t, x)|. \end{aligned}$$

Hence for any $\alpha > 0$, by Theorem 3,

$$\begin{aligned} & \left| \left\{ \limsup_{\delta \rightarrow 0} |A_\delta f(t, x) - f(t, x)| > \alpha \right\} \right| \\ & \leq |\{\mathcal{M}(f - g) > \alpha/2\}| + |\{|g - f| > \alpha/2\}| \\ & \leq \frac{2c\varepsilon}{\alpha} + \frac{2\varepsilon}{\alpha}. \end{aligned}$$

Since ε and α are arbitrary, it follows that $\limsup_{\delta \rightarrow 0} |A_\delta f(t, x) - f(t, x)| = 0$ a.e. Hence for almost all (t, x) ,

$$\begin{aligned} |f(t, x)| &= \left| \lim_{\delta \rightarrow 0} A_\delta f(t, x) \right| \leq \lim_{\delta \rightarrow 0} \frac{1}{|Q_\delta(t, x)|} \int_{Q_\delta(t, x)} |f(t, y)| dt dy \\ &\leq \sup_{\delta > 0} \frac{1}{|Q_\delta(t, x)|} \int_{Q_\delta(t, x)} |f(t, y)| dt dy = \mathcal{M}f(t, x). \end{aligned}$$

Finally, for $f_1 = f\chi_{\{|f| > \alpha/2\}}$ we have $\tilde{\mathcal{M}}f \leq \tilde{\mathcal{M}}f_1 + \frac{\alpha}{2}$, and by Theorem 3(b),

$$\left| \left\{ \tilde{\mathcal{M}}f > \alpha \right\} \right| \leq \left| \left\{ \tilde{\mathcal{M}}f_1 > \alpha/2 \right\} \right| \leq \frac{2c}{\alpha} \int_{|f| > \alpha/2} |f| dm \leq \frac{2c}{\alpha} \int_{\tilde{\mathcal{M}}f > \alpha/2} |f| dm.$$

□

5.2. Calderon-Zygmund decomposition. Assume **A1**, **A2** hold. Let $F \subseteq \mathbf{R} \times \mathbf{R}^d$ be closed and $O = F^c = \mathbf{R}^{d+1} \setminus F$. For $(t, x) \in O$, let

$$D(t, x) = \inf \{ \delta > 0 : Q_\delta(t, x) \cap F \neq \emptyset \}.$$

For each $(t, x) \in O$, $D(t, x) \in (0, \infty)$. Let K_0 be a constant in Lemma 13. We fix $A > 1$ so that $l(1/A) < 1$ and $\varepsilon > 0$ so that $l(2K_0\varepsilon) < 1$, $\varepsilon \leq \frac{1}{4AK_0^3} <$

1. Then, denoting $D = D(t, x)$, we have

$$\kappa(\varepsilon D) \leq l(2\varepsilon) \kappa(D/2) \leq \kappa\left(\frac{D}{2}\right), \kappa(\varepsilon D) \leq l(\varepsilon) \kappa(D) \leq \kappa(D),$$

$$\kappa(D) \leq l(1/A) \kappa(AD) < \kappa(AD),$$

and

$$\kappa(\varepsilon D) \leq l(2K_0\varepsilon) \kappa(D/2K_0) \leq \kappa(D/2K_0).$$

Consider the covering $Q_{\varepsilon D(t, x)}(t, x)$, $(t, x) \in O$, of O . Let

$$Q^k = Q_{\varepsilon D(t_k, x_k)}(t_k, x_k), k \geq 1,$$

be its maximal disjoint subcollection: for any $Q_{\varepsilon D(t, x)}(t, x)$ there is k so that $Q^\varepsilon(t, x) \cap Q^k \neq \emptyset$. Let

$$Q^{*k} = Q_{D(t_k, x_k)/2}(t_k, x_k), Q^{**k} = Q_{AD(t_k, x_k)}(t_k, x_k).$$

Note that $Q^k \subseteq Q^{*k} \subseteq Q_{D(t_k, x_k)}(t_k, x_k) \subseteq O$, $Q^{**k} \cap F \neq \emptyset$. We will show that $\cup_k Q^{*k} = O$. Let $(t, x) \in O$ and $Q_{\varepsilon D(t_k, x_k)}(t_k, x_k) \cap Q_{\varepsilon D(t, x)}(t, x) \neq \emptyset$ for some k . Since

$$\begin{aligned} Q_{\varepsilon D(t_k, x_k)}(t_k, x_k) &\subseteq Q_{D(t_k, x_k)}(t_k, x_k) \subseteq Q_{AD(t_k, x_k)}(t_k, x_k), \\ Q_{\varepsilon D(t, x)}(t, x) &\subseteq Q_{D(t, x)/(2K_0)}, \end{aligned}$$

it follows that

$$Q_{D(t, x)/(2K_0)}(t, x) \cap Q_{AD(t_k, x_k)}(t_k, x_k) \neq \emptyset.$$

We show by contradiction that $AD(t_k, x_k) \geq D(t, x)/2K_0$. If not so, then $AD(t_k, x_k) < D(t, x)/2K_0$, and, by Lemma 13, $Q_{D(t, x)/2K_0}(t, x)$ and $Q_{AD(t_k, x_k)}(t_k, x_k)$ are contained in $Q_{D(t, x)/2}(t, x) \subseteq O$: a contradiction to $Q_{AD(t_k, x_k)}(t_k, x_k) \cap F \neq \emptyset$. Therefore $AD(t_k, x_k) \geq D(t, x)/2K_0$ and $2AK_0\varepsilon D(t_k, x_k) \geq \varepsilon D(t, x)$. Now, $Q_{\varepsilon D(t_k, x_k)}(t_k, x_k) \subseteq Q_{2AK_0\varepsilon D(t_k, x_k)}(t_k, x_k)$ and $Q_{\varepsilon D(t_k, x_k)}(t_k, x_k) \cap Q_{\varepsilon D(t, x)}(t, x) \neq \emptyset$. Hence by Lemma 13, $Q_{\varepsilon D(t, x)}$ is contained in $Q_{2AK_0^2\varepsilon D(t_k, x_k)}(t_k, x_k)$. Since $2AK_0^2\varepsilon \leq \frac{1}{2K_0}$, it follows by Lemma 1 that

$$Q_{\varepsilon D(t, x)}(t, x) \subseteq Q_{2AK_0^2\varepsilon D(t_k, x_k)}(t_k, x_k) \subseteq Q_{D(t_k, x_k)/2}(t_k, x_k) = Q^{*k}.$$

So we proved the following statement.

Lemma 15. *Assume **A1**, **A2** hold. Given a closed nonempty F , there are sequences Q^k , Q^{*k} and Q^{**k} in \mathbb{Q} having the same center but with radius expanded by the same factor $c_1^{**} > c_1^* > c_1$ so that $Q^k \subseteq Q^{*k} \subseteq Q^{**k}$ (all of them are of the form $Q_{bD(t_k, x_k)}(t_k, x_k)$ with $b = c_1, c_1^*, c_1^{**}$ correspondingly) and*

- (a) *the sets Q^k are disjoint.*
- (b) *$\cup_k Q^{*k} = O = F^c$.*
- (c) *$Q^{**k} \cap F \neq \emptyset$ for each k .*

Remark 7. *Assume **A1**, **A2** hold and $Q^k \subseteq Q^{*k} \subseteq Q^{**k}$ be the sequences in \mathbb{Q} from Lemma 15. It is easy to find a sequence of disjoint measurable sets C^k so that $Q^k \subseteq C^k \subseteq Q^{*k}$ and $\cup_k C^k = O$. For example (see Remark, p. 15, in [8]),*

$$C^k = Q^{*k} \cap (\cup_{j < k} C^j)^c \cap (\cup_{j > k} Q^j)^c.$$

Now we derive Calderon-Zygmund decomposition for \mathbb{Q} .

Theorem 4. *Assume **A1**, **A2** hold. Let $f \in L_1(\mathbf{R} \times \mathbf{R}^d)$, $\alpha > 0$ and $O_\alpha = \{\widetilde{\mathcal{M}}f > \alpha\}$. Consider the sets $Q^k \subseteq C^k \subseteq Q^{*k} \subseteq O$ of Lemma 15 and Remark 7 associated to O_α .*

There is a decomposition $f = g + b$ with

$$(5.1) \quad g(x) = \begin{cases} f(x) & \text{if } x \notin O_\alpha, \\ \frac{1}{|C^k|} \int_{C^k} f & \text{if } x \in C^k, k \geq 1, \end{cases}$$

and with $b = \sum_k b_k$, where

$$(5.2) \quad b_k = \chi_{C^k} \left[f(x) - \frac{1}{|C^k|} \int_{C^k} f \right], k \geq 1,$$

(note C^k are disjoint, $\cup_k C^k = O_\alpha$). Also,

(i) $|g(x)| \leq c\alpha$ for a.e. x .

(ii) $\text{support}(b_k) \subseteq Q^{*k}$,

$$\int b_k = 0 \text{ and } \int |b_k| \leq c\alpha |Q^{*k}|.$$

$$(iii) \sum_k |Q^{*k}| \leq \frac{c}{\alpha} \int |f|.$$

Proof. The set $O_\alpha = \{\widetilde{\mathcal{M}}f > \alpha\}$ is open. We can apply Lemma 15 and Remark 7 to it and consider the sets $Q^k \subseteq C^k \subseteq Q^{*k} \subseteq E_\alpha$ with C^k disjoint and $\cup_k C^k = E_\alpha$.

Define g by (5.2). Hence $f = g + \sum_k b_k$ with b_k given by (5.2). Obviously

$$\sum_k |Q^k| \leq |E_\alpha|.$$

(i) By Corollary 7, $|f(x)| \leq \alpha$ a.e. on $O_\alpha^c = \{\widetilde{\mathcal{M}}f(t, x) \leq \alpha\}$. Hence: so $|g(x)| \leq \alpha$ a.e. on E_α^c . On the other hand, if $Q^{**k} \in \mathbb{Q}$ is the sequence of Lemma 15, then

$$\frac{1}{|Q^{**k}|} \int_{Q^{**k}} |f| \leq \alpha$$

because $Q^{**k} \cap O_\alpha^c \neq \emptyset$ and $\widetilde{\mathcal{M}}f(t, x) \leq \alpha$ on O_α^c (the definition of $\widetilde{\mathcal{M}}$ implies it). Since $|Q^k| \leq |C^k| \leq |Q^{*k}| \leq |Q^{**k}| \leq l \left(\frac{c_1^{**}}{c_1} \right) |Q^k|$ and $C^k \subseteq Q^{**k}$, it follows that

$$|g| \leq \bar{c}\alpha.$$

(ii) Only inequality is not trivial:

$$\int |b_k| \leq 2 \int_{C^k} |f| \leq 2 |Q^{**k}| \frac{1}{|Q^{**k}|} \int_{Q^{**k}} |f| \leq c\alpha |Q^{*k}|.$$

(iii) We have

$$|O_\alpha| = \left| \left\{ \widetilde{\mathcal{M}}f(t, x) > \alpha \right\} \right| \geq \sum_k |Q^k| \geq \bar{c} \sum_k |Q^{*k}|$$

and the inequality follows by Theorem 3. \square

5.3. L_p -estimates. Let now

$$(Tf)(t, x) = \int_{\mathbf{R}^{d+1}} K(t, x, s, y) f(s, y) ds dy, (t, x) \in \mathbf{R}^{d+1}.$$

We assume that T is defined and bounded on L_q :

$$(5.3) \quad |Tf|_{L_q} \leq C |f|_{L_q}, f \in L_q.$$

In addition, we assume that there is a constant $A > 0$ so that (denoting (s_k, y_k) the center of Q_k^* ,

$$(5.4) \quad \sup_{(s, y) \in Q_k^*} \int_{(Q_k^{**})^c} |K(t, x, s, y) - K(t, x, s_k, y_k)| dt dx \leq A.$$

The assumption (5.4) holds if there are constants $c > 1, A > 0$ so that for any $Q_\delta \in \mathcal{Q}$,

$$(5.5) \quad \int_{\mathbf{R}^{d+1} \setminus Q_{c\delta}(s, y)} |K(t, x, \bar{s}, \bar{y}) - K(t, x, s, y)| dx dt \leq A \quad \forall (\bar{s}, \bar{y}) \in Q_\delta(s, y)$$

Theorem 5. *Let **A1**, **A2**, (5.3) and (5.5) hold. Then T is bounded in L_p -norm on $L_p \cap L_q$ if $1 < p < q$. More precisely,*

$$|T(f)|_{L_p} \leq A_p |f|_{L_p}, f \in L_p \cap L_q, 1 < p < q,$$

where A_p depends only on the constant A and p .

Proof. As [8] says, it is enough to prove that

$$m(|Tf| > \alpha) \leq \frac{A'}{\alpha} \int |f| dx, f \in L_1 \cap L_q, \alpha > 0,$$

where A' depends on A .

For a large constant c' (to be determined) we estimate $m(|Tf| > c'\alpha)$. For a fixed $\alpha > 0$ we consider the decomposition $f = g + b$ in Theorem 4. First note that

$$m(\cup_n Q_n^{**}) \leq \sum_n m(Q_n^{**}) \leq c \sum_n m(Q_n^*) \leq \frac{c}{\alpha} \int |f|.$$

It is enough to show that

$$|\{|Tg| > (c'/2)\alpha\}| + |\{|Tb| > (c'/2)\alpha\}| \leq A'/\alpha \int |f| dx.$$

First $g \in L_q$. Indeed,

$$\int |g|^q dx = \int_{(\cup_k Q_k^*)^c} |g|^q + \int_{\cup_k Q_k^*} |g|^q \leq c\alpha^{q-1} \int |f|$$

because

$$\begin{aligned} \int_{(\cup_k Q_k^*)^c} |g|^q &\leq c\alpha^{q-1} \int_{(\cup_k Q_k^*)^c} |f|, \\ \int_{\cup_k Q_k^*} |g|^q &\leq c\alpha^q \sum_k |Q_k^*| \leq c\alpha^{q-1} \int |f|. \end{aligned}$$

By Chebyshev inequality,

$$\begin{aligned} |\{|Tg| > (c'/2)\alpha\}| &\leq \left(\frac{c'\alpha}{2}\right)^{-q} |Tg|_{L_q}^q \leq \left(\frac{c'\alpha}{2}\right)^{-q} A^q |g|_{L_q}^q \\ &\leq c \left(\frac{c'\alpha}{2}\right)^{-q} A^q \alpha^{q-1} \int |f| \leq \frac{A'}{\alpha} |f|_{L_1}. \end{aligned}$$

Now,

$$\int_{(\cup_k Q_k^{**})^c} |Tb| \leq \sum_k \int_{(Q_k^{**})^c} |Tb_k|$$

Since for $x \notin Q_k^*$, denoting by (s_k, y_k) the center of Q_k^* (and Q_k^{**}), we have. denoting $f_k = 1/|C^k| \int_{C_k} f$,

$$\begin{aligned} Tb_k &= \int_{C_k} K(t, x, s, y) [f(s, y) - f_k] ds dy \\ &= \int_{C_k} [K(t, x, s, y) - K(t, x, s_k, y_k)] [f(s, y) - f_k] ds dy \end{aligned}$$

and

$$\begin{aligned} \int_{(Q_k^{**})^c} |Tb_k| dx &\leq \int |b_k| ds dy \sup_{(s,y) \in Q_k^*} \int_{(Q_k^{**})^c} |K(t, x, s, y) - K(t, x, s_k, y_k)| dt dx \\ &\leq cA\alpha |Q_k^*|. \end{aligned}$$

Hence

$$\int_{(\cup_k Q_k^{**})^c} |Tb| \leq cA\alpha \sum_k |Q_k^*| \leq cA \int |f|.$$

□

REFERENCES

- [1] Dong H. and Kim D., On L_p - estimates of non-local elliptic equations, Journal of Functional Analysis 262 (2012), 1166-1199.
- [2] Farkas, W., Jacob, N., Schilling, R., Function spaces related to continuous negative definite functions: ψ -Bessel potential spaces, Diss. Math. 393, 2001.
- [3] Kim, I. and Kim, K.-H., An L_p -theory for a class of non-local elliptic equations related to nonsymmetric measurable kernels, J. Math. Anal. Appl, 434 (2016), pp. 1302-1335.
- [4] Kim, P., Song, R. and Vondraček, Z., Global uniform boundary Harnack principle with explicit decay rate and its application, Stoch. Proc. Appl., 124 (2014), pp 235-267.
- [5] Komatsu, T., On the martingale problem for generators of stable processes with perturbations, Osaka J. Math., 21 (1984), pp 113-132.
- [6] Krylov, N.V., Lectures on Elliptic and Parabolic Equations in Sobolev Spaces, AMS, 2008.

- [7] Sato, K., Levy Processes and Infinitely Divisible Distributions, Cambridge University Press, 1999.
- [8] Stein, E., Harmonic Analysis, Princeton University Press, 1993.
- [9] Zhang Xicheng, L^p -maximal regularity of nonlocal parabolic equations and applications, Ann. I.H. Poincaré-AN 30 (2013), pp 573-614.

UNIVERSITY OF SOUTHERN CALIFORNIA, LOS ANGELES