

NEW RESULT ON CHERN CONJECTURE FOR MINIMAL HYPERSURFACES AND ITS APPLICATION

HONGWEI XU AND ZHIYUAN XU

Dedicated to Professor Shiing-shen Chern on the occasion of his 105th birthday

ABSTRACT. We verify that if M is a compact minimal hypersurface in \mathbb{S}^{n+1} whose squared length of the second fundamental form satisfying $0 \leq |A|^2 - n \leq \frac{n}{22}$, then $|A|^2 \equiv n$ and M is a Clifford torus. Moreover, we prove that if M is a complete self-shrinker with polynomial volume growth in \mathbb{R}^{n+1} whose equation is given by (1.1), and if the squared length of the second fundamental form of M satisfies $0 \leq |A|^2 - 1 \leq \frac{1}{21}$, then $|A|^2 \equiv 1$ and M is a round sphere or a cylinder. Our results improve the rigidity theorems due to Q. Ding and Y. L. Xin [14, 16].

1. INTRODUCTION

The famous Chern Conjecture for minimal hypersurfaces in a sphere was proposed by S. S. Chern [10, 11] in 1968 and 1970, and was listed in the Problem Section by S. T. Yau [46] in 1982, as stated

Chern Conjecture. (A)(Standard version) *Let M be a compact minimal hypersurface with constant scalar curvature in the unit sphere \mathbb{S}^{n+1} . Then the possible values of the scalar curvature of M form a discrete set.*

(B)(Refined version) *Let M be a compact minimal hypersurface with constant scalar curvature in the unit sphere \mathbb{S}^{n+1} . Then M is isoparametric.*

(C)(Stronger version) *Let M be a compact minimal hypersurface in the unit sphere \mathbb{S}^{n+1} . Denote by A the second fundamental form of M . Set $a_k = (k - \text{sgn}(5 - k))n$, for $k \in \{m \in \mathbb{Z}^+; 1 \leq m \leq 5\}$. Then we have*
(i) For any fixed $k \in \{m \in \mathbb{Z}^+; 1 \leq m \leq 4\}$, if $a_k \leq |A|^2 \leq a_{k+1}$, then

2010 *Mathematics Subject Classification.* 53C44; 53C42.

Key words and phrases. Chern conjecture, minimal hypersurface, the second pinching theorem, the second fundamental form, self-shrinker.

Research supported by the National Natural Science Foundation of China, Grant Nos. 11531012, 11371315; and the China Postdoctoral Science Foundation, Grant No. 2016M590530.

M is isoparametric, and $|A|^2 \equiv a_k$ or $|A|^2 \equiv a_{k+1}$.
(ii) If $|A|^2 \geq a_5$, then M is isoparametric, and $|A|^2 \equiv a_5$.

It is well-known that the Chern Conjecture consists of several pinching problems. In the late 1960s, Simons, Lawson, and Chern-do Carmo-Kobayashi [11, 21, 32] solved the first pinching problem for compact minimal hypersurfaces in the unit sphere \mathbb{S}^{n+1} and verified that if $|A|^2 \leq n$, then $|A|^2 \equiv 0$ and M is the great sphere \mathbb{S}^n , or $|A|^2 \equiv n$ and M is one of the Clifford torus $\mathbb{S}^k(\sqrt{\frac{k}{n}}) \times \mathbb{S}^{n-k}(\sqrt{\frac{n-k}{n}})$, $1 \leq k \leq n-1$. More generally, they proved a rigidity theorem for compact minimal submanifolds in a sphere under a sharp pinching condition. Further developments on the first pinching problem have been made by many other authors [7, 13, 18, 23, 25, 31, 36, 37, 45], etc.

As a special part of the Chern Conjecture, the following problem has been open for more than 40 years.

The Second Pinching Problem. *Let M be a compact minimal hypersurface in the unit sphere \mathbb{S}^{n+1} .*

- (i) *If $|A|^2$ is constant, and if $n \leq |A|^2 \leq 2n$, then $|A|^2 = n$, or $|A|^2 = 2n$.*
(ii) *If $n \leq |A|^2 \leq 2n$, then $|A|^2 \equiv n$, or $|A|^2 \equiv 2n$.*

In 1983, Peng and Terng [28, 29] initiated the study of the second pinching problem for minimal hypersurfaces in a unit sphere, and made the following breakthrough on the Chern Conjecture.

Theorem A. *Let M be a compact minimal hypersurface in the unit sphere \mathbb{S}^{n+1} .*

- (i) *If $|A|^2$ is constant, and if $n \leq |A|^2 \leq n + \frac{1}{12n}$, then $|A|^2 = n$.*
(ii) *If $n \leq 5$, and if $n \leq |A|^2 \leq n + \tau_1(n)$, where $\tau_1(n)$ is a positive constant depending only on n , then $|A|^2 \equiv n$.*

During the past three decades, there have been some important progresses on the Chern Conjecture [1, 4, 5, 6, 9, 14, 17, 27, 30, 33, 34, 35, 38, 39, 40, 42, 43, 44, 47], etc. In 1993, Chang [4] proved Chern Conjecture (A) in dimension three. Yang-Cheng [42, 43, 44] improved the pinching constant $\frac{1}{12n}$ in Theorem A(i) to $\frac{2}{3}$. Later, Suh-Yang [33] improved this pinching constant to $\frac{3}{7}n$.

In 2007, Wei and Xu [35] proved that if M is a compact minimal hypersurface in \mathbb{S}^{n+1} , $n = 6, 7$, and if $n \leq |A|^2 \leq n + \tau_2(n)$, where $\tau_2(n)$ is a positive constant depending only on n , then $|A|^2 \equiv n$. Later, Zhang [47] extended the second pinching theorem due to Peng-Terng [29] and Wei-Xu [35] to the case of $n = 8$. In 2011, Ding and Xin [14] verified the following important rigidity theorem, as stated

Theorem B. *Let M be an n -dimensional compact minimal hypersurface in the unit sphere \mathbb{S}^{n+1} . If $n \geq 6$, the squared norm of the second fundamental form satisfies $0 \leq |A|^2 - n \leq \frac{n}{23}$, then $|A|^2 \equiv n$, i.e., M is a Clifford torus.*

In the present paper, we first give a refined version of Ding-Xin's rigidity theorem for minimal hypersurfaces in a sphere.

Theorem 1.1. *Let M be an n -dimensional compact minimal hypersurface in the unit sphere \mathbb{S}^{n+1} . If the squared norm of the second fundamental form satisfies $0 \leq |A|^2 - n \leq \frac{n}{22}$, then $|A|^2 \equiv n$ and M is a Clifford torus.*

Moreover, we consider the isometric immersion $X : M \rightarrow \mathbb{R}^{n+1}$. If the position vector X evolves in the direction of the mean curvature vector \vec{H} , then it gives rise to a solution to the mean curvature flow

$$\begin{cases} \frac{\partial}{\partial t} X(x, t) = \vec{H}(x, t), & x \in M, \\ X(x, 0) = X(x). \end{cases}$$

An important class of solutions to the above mean curvature flow equations are self-shrinkers, which satisfy

$$(1.1) \quad H = -X^N,$$

where X^N is the projection of X on the unit inner normal vector ξ , i.e., $X^N = \langle X, \xi \rangle$.

Rigidity problems of self-shrinkers have been studied extensively [2, 8, 12, 15, 16, 20, 22, 24]. In 2011, Le and Sesum [22] proved that any n -dimensional complete self-shrinker with polynomial volume growth in \mathbb{R}^{n+1} whose squared norm of the second fundamental form satisfies $|A|^2 < 1$ must be a hyperplane. Afterwards, Cao and Li [2] generalized this rigidity result to arbitrary codimensional cases and proved that if M is an n -dimensional complete self-shrinker with polynomial volume growth in \mathbb{R}^{n+q} , and if $|A|^2 \leq 1$, then M must be one of the generalized cylinders. Under the assumption that $|A|$ is constant, Cheng and Wei [8] proved that if M is an n -dimensional complete self-shrinker with polynomial volume growth in \mathbb{R}^{n+1} , and if $0 \leq |A|^2 - 1 \leq \frac{3}{7}$, then $|A|^2 = 1$.

In 2014, Ding and Xin [16] proved the second pinching theorem for self-shrinkers in the Euclidean space.

Theorem C. *Let M be an n -dimensional complete self-shrinker with polynomial volume growth in \mathbb{R}^{n+1} . If the squared norm of the second fundamental form satisfies $0 \leq |A|^2 - 1 \leq 0.022$, then $|A|^2 \equiv 1$ and M is a round sphere or a cylinder.*

In Section 3, we improve Ding-Xin's pinching constant in Theorem C and prove the following rigidity theorem for self-shrinkers in the Euclidean space.

Theorem 1.2. *Let M be an n -dimensional complete self-shrinker with polynomial volume growth in \mathbb{R}^{n+1} . If the squared norm of the second fundamental form satisfies $0 \leq |A|^2 - 1 \leq \frac{1}{21}$, then $|A|^2 \equiv 1$ and M is a round sphere or a cylinder.*

2. MINIMAL HYPERSURFACES IN THE UNIT SPHERE

Let M be an $n(\geq 2)$ -dimensional compact minimal hypersurface in the unit sphere \mathbb{S}^{n+1} . We shall make use of the following convention on the range of indices:

$$1 \leq i, j, k, \dots \leq n.$$

We choose a local orthonormal frame $\{e_1, e_2, \dots, e_{n+1}\}$ near a fixed point $x \in M$ over \mathbb{S}^{n+1} such that $\{e_1, e_2, \dots, e_n\}$ are tangent to M .

Let $\{\omega_1, \omega_2, \dots, \omega_{n+1}\}$ be the dual frame fields of $\{e_1, e_2, \dots, e_{n+1}\}$. Denote by R_{ijkl} and $A = \sum_{i,j} h_{ij} \omega_i \otimes \omega_j$ the Riemannian curvature tensor and the second fundamental form of M , respectively. Then we have $\text{Trace } A = \sum_i h_{ii} = 0$. Put $S = |A|^2 = \sum_{i,j} h_{ij}^2$. We denote the first, the second and the third covariant derivatives of the second fundamental form of M by

$$\begin{aligned} \nabla A &= \sum_{i,j,k} h_{ijk} \omega_i \otimes \omega_j \otimes \omega_k, \\ \nabla^2 A &= \sum_{i,j,k,l} h_{ijkl} \omega_i \otimes \omega_j \otimes \omega_k \otimes \omega_l, \\ \nabla^3 A &= \sum_{i,j,k,l,m} h_{ijklm} \omega_i \otimes \omega_j \otimes \omega_k \otimes \omega_l \otimes \omega_m. \end{aligned}$$

We have the Gauss and Codazzi equations.

$$(2.1) \quad R_{ijkl} = \delta_{ik} \delta_{jl} - \delta_{il} \delta_{jk} + h_{ik} h_{jl} - h_{il} h_{jk},$$

and

$$(2.2) \quad h_{ijk} = h_{ikj}.$$

By the Gauss equations, the scalar curvature of M is given by

$$R = n(n-1) - S.$$

We also have the Ricci identities.

$$(2.3) \quad h_{ijkl} = h_{ijlk} + \sum_m h_{mj} R_{mikl} + \sum_m h_{im} R_{mjkl},$$

$$(2.4) \quad h_{ijklm} = h_{ijkml} + \sum_r h_{rjk} R_{rilm} + \sum_r h_{irk} R_{rjlm} + \sum_r h_{ijr} R_{rklm}.$$

Choose a suitable orthonormal frame $\{e_1, e_2, \dots, e_n\}$ at x such that $h_{ij} = \lambda_i \delta_{ij}$ for all i, j . Then we have

$$(2.5) \quad \frac{1}{2} \Delta S = S(n - S) + |\nabla A|^2.$$

Hence

$$(2.6) \quad |\nabla S|^2 = \frac{1}{2} \Delta S^2 - S \Delta S = \frac{1}{2} \Delta S^2 + 2S^2(S - n) - 2S|\nabla A|^2.$$

We get

$$(2.7) \quad \begin{aligned} \frac{1}{2} \Delta |\nabla A|^2 &= (2n + 3 - S) |\nabla A|^2 - \frac{3}{2} |\nabla S|^2 + |\nabla^2 A|^2 \\ &\quad + 3(2B_2 - B_1), \end{aligned}$$

where $B_1 = \sum_{i,j,k,l,m} h_{ijk} h_{ijl} h_{km} h_{ml}$ and $B_2 = \sum_{i,j,k,l,m} h_{ijk} h_{klm} h_{im} h_{jl}$.

Following [29] (also see [14, 40]), we have

$$(2.8) \quad |\nabla^2 A|^2 - \frac{3S(S-n)^2}{2(n+4)} \geq \frac{3}{4} \sum_{i \neq j} t_{ij}^2 = \frac{3}{4} \sum_{i,j} t_{ij}^2,$$

$$(2.9) \quad 3(B_1 - 2B_2) \leq \sigma S |\nabla A|^2,$$

where $t_{ij} = h_{ijij} - h_{jiji}$ and $\sigma = \frac{\sqrt{17}+1}{2}$.

Moreover, we have

$$(2.10) \quad \int_M (B_1 - 2B_2) dM = \int_M (Sf_4 - f_3^2 - S^2 - \frac{|\nabla S|^2}{4}) dM,$$

where $f_k = \text{Trace } A^k = \sum_i \lambda_i^k$.

Set $G = \sum_{i,j} t_{ij}^2 = \sum_{i,j} (\lambda_i - \lambda_j)^2 (1 + \lambda_i \lambda_j)^2$. Thus, we have

$$(2.11) \quad G = 2Sf_4 - 2f_3^2 - 2S^2 - 2S(S - n).$$

This implies that

$$(2.12) \quad \frac{1}{2} \int_M G dM = \int_M [(B_1 - 2B_2) - |\nabla A|^2 + \frac{1}{4} |\nabla S|^2] dM.$$

Now we are in a position to prove our rigidity theorem for minimal hypersurfaces in a sphere.

Proof of Theorem 1.1. It follows from (2.7), (2.8) and (2.12) that

$$(2.13) \quad \int_M [(S - 2n - \frac{3}{2})|\nabla A|^2 + \frac{3}{2}(B_1 - 2B_2) + \frac{9}{8}|\nabla S|^2]dM \geq 0.$$

Under the pinching condition $n \leq S \leq n + \frac{n}{k}$, (2.13) together with (2.6) implies

$$(2.14) \quad \begin{aligned} 0 &\leq \int_M [(S - 2n - \frac{3}{2})|\nabla A|^2 + \frac{3}{2}(B_1 - 2B_2) \\ &\quad + \frac{9}{4}(S^3 - nS^2 - S|\nabla A|^2)]dM \\ &\leq \int_M [(\frac{k+9}{4k}n - \frac{5}{4}S - \frac{3}{2})|\nabla A|^2 + \frac{3}{2}(B_1 - 2B_2)]dM. \end{aligned}$$

When $n = 2, 3$, combining (2.9) and (2.14), we get

$$(2.15) \quad 0 \leq \int_M \left(\frac{k+9}{4k}n + \frac{\sqrt{17}-4}{4}S - \frac{3}{2} \right) |\nabla A|^2 dM.$$

When $n = 4, 5$, by Lemma 3.4 in [47], we have

$$(2.16) \quad 3(B_1 - 2B_2) \leq \eta_n S |\nabla A|^2,$$

where $\eta_4 = 2.16$ and $\eta_5 = 2.23$. It follows from (2.14) and (2.16) that

$$(2.17) \quad 0 \leq \int_M \left(\frac{1+\frac{9}{k}}{4}n - \frac{5-2\eta_n}{4}S - \frac{3}{2} \right) |\nabla A|^2 dM.$$

Taking $k = 22$, we see that the coefficients of the integrals in (2.15) and (2.17) are both negative. Therefore, $\nabla A = 0$ and $|A|^2 = n$, for $n \leq 5$.

When $n \geq 6$, the following lemma can be found in [14].

Lemma 2.1. *If $n \geq 6$, $n \leq S \leq \frac{16}{15}n$, then $3(B_1 - 2B_2) \leq (S + 4 + C_1(n)G^{1/3})|\nabla A|^2$, where $C_1(n) = \left(\frac{3-\sqrt{6}-4p}{\sqrt{6}-1+13p} (6 - \sqrt{6} - 13p)^2 \right)^{1/3}$ and $p = \frac{1}{13(n-2)}$.*

For $\theta \in (0, 1)$, from (2.7), (2.8) and (2.12), using Lemma 2.1 and Young's inequality, we drive the following inequality.

$$\begin{aligned}
& \frac{3(1-\theta)}{4} \int_M G dM + \int_M \frac{3S(S-n)^2}{2(n+4)} dM \\
\leq & \int_M \left[(S-2n-3)|\nabla A|^2 + \frac{3}{2}|\nabla S|^2 + 3(B_1-2B_2) - \frac{3\theta}{4}G \right] dM \\
= & \int_M (S-2n-3 + \frac{3\theta}{2})|\nabla A|^2 dM + (\frac{3}{2} - \frac{3\theta}{8}) \int_M |\nabla S|^2 dM \\
& + (3 - \frac{3\theta}{2}) \int_M (B_1 - 2B_2) dM \\
\leq & \int_M (S-2n-3 + \frac{3\theta}{2})|\nabla A|^2 dM + (\frac{3}{2} - \frac{3\theta}{8}) \int_M |\nabla S|^2 dM \\
& + (1 - \frac{\theta}{2}) \int_M (S+4 + C_1(n)G^{1/3})|\nabla A|^2 dM \\
\leq & \int_M \left[(2 - \frac{\theta}{2})S - 2n + 1 - \frac{\theta}{2} \right] |\nabla A|^2 dM + (\frac{3}{2} - \frac{3\theta}{8}) \int_M |\nabla S|^2 dM \\
(2.18) \quad & + \frac{3(1-\theta)}{4} \int_M G dM + C_3 \int_M |\nabla A|^3 dM,
\end{aligned}$$

where $C_3 = C_3(n, \theta) = \frac{4}{9}C_1(n)^{3/2}(1 - \frac{\theta}{2})^{3/2}(1 - \theta)^{-1/2}$.

For $\epsilon > 0$, (2.5) together with the divergence theorem and Cauchy-Schwarz's inequality implies

$$\begin{aligned}
\int_M |\nabla A|^3 dM & \leq \int_M S(S-n)|\nabla A| dM + \epsilon \int_M |\nabla^2 A|^2 dM \\
(2.19) \quad & + \frac{1}{16\epsilon} \int_M |\nabla S|^2 dM.
\end{aligned}$$

Under the pinching condition $n \leq S \leq n + \delta(n) (\leq \frac{n}{15})$, for $\epsilon_1 = \theta_1 \epsilon > 0$, we estimate $\int_M S(S-n)|\nabla A| dM$ as follows.

$$\begin{aligned}
\int_M S(S-n)|\nabla A| dM & \leq 2(n+\delta)\epsilon_1 \int_M S(S-n) dM \\
& + \frac{1}{8(n+\delta)\epsilon_1} \int_M S(S-n)|\nabla A|^2 dM \\
(2.20) \quad & \leq \left[2(n+\delta)\epsilon_1 + \frac{S-n}{8\epsilon_1} \right] \int_M |\nabla A|^2 dM.
\end{aligned}$$

From (2.7) and (2.9), we have

$$(2.21) \quad \begin{aligned} \int_M |\nabla^2 A|^2 dM &\leq \frac{3}{2} \int_M |\nabla S|^2 dM \\ &+ \int_M [(\sigma + 1)S - 2n - 3] |\nabla A|^2 dM. \end{aligned}$$

Substituting (2.19), (2.20) and (2.21) into (2.18), we obtain

$$(2.22) \quad \begin{aligned} 0 &\leq \int_M \left[\left(2 - \frac{\theta}{2}\right)S - 2n + 1 - \frac{\theta}{2} \right] |\nabla A|^2 dM \\ &+ \left(\frac{3}{2} - \frac{3\theta}{8}\right) \int_M |\nabla S|^2 dM - \int_M \frac{3S(S-n)^2}{2(n+4)} dM \\ &+ C_3 \left\{ \left[2(n+\delta)\epsilon_1 + \frac{S-n}{8\epsilon_1} \right] \int_M |\nabla A|^2 dM \right. \\ &+ \frac{3\epsilon}{2} \int_M |\nabla S|^2 dM + \frac{1}{16\epsilon} \int_M |\nabla S|^2 dM \\ &\left. + \int_M \epsilon [(\sigma + 1)S - 2n - 3] |\nabla A|^2 dM \right\} \\ &= \int_M \left\{ \left(2 - \frac{\theta}{2}\right)S - 2n + 1 - \frac{\theta}{2} + \epsilon C_3 [(\sigma + 1)S - 2n - 3] \right. \\ &\left. + C_3 \left[2(n+\delta)\epsilon_1 + \frac{S-n}{8\epsilon_1} \right] \right\} |\nabla A|^2 dM - \int_M \frac{3S(S-n)^2}{2(n+4)} dM \\ &+ \left(3 - \frac{3\theta}{4} + 3\epsilon C_3 + \frac{C_3}{8\epsilon}\right) \int_M [S(S-n)^2 - (S-n)|\nabla A|^2] dM \\ &\leq \int_M \left\{ 1 - \frac{\theta}{2}(n+1) + \epsilon C_3(n\sigma - n - 3) + 2\epsilon_1 C_3(n+\delta) \right. \\ &\left. - \left[1 - \frac{\theta}{4} + \frac{C_3}{8\epsilon} - \frac{C_3}{8\epsilon_1} + \epsilon C_3(2-\sigma) \right] (S-n) \right\} |\nabla A|^2 dM \\ &+ \delta \left(3 - \frac{3\theta}{4} + 3\epsilon C_3 + \frac{C_3}{8\epsilon} - \frac{3}{2(n+4)} \right) \int_M |\nabla A|^2 dM \\ &= \int_M \left\{ 1 - \frac{\theta}{2}(n+1) + \epsilon C_3(n\sigma + n - 3 + 5\delta) + \frac{C_3\delta}{8\epsilon} \right. \\ &\left. + \delta \left(3 - \frac{3\theta}{4} - \frac{3}{2(n+4)} \right) + 2\epsilon(\theta_1 - 1)C_3(n+\delta) \right. \\ &\left. - \left[1 - \frac{\theta}{4} + \frac{C_3}{8\epsilon} - \frac{C_3}{8\epsilon_1} + \epsilon C_3(2-\sigma) \right] (S-n) \right\} |\nabla A|^2 dM. \end{aligned}$$

Let $\epsilon = \sqrt{\frac{\delta}{8[\sigma n + n - 3 + 5\delta + 2(n + \delta)(\theta_1 - 1)]}}$, $\theta = 0.866$ and $\theta_1 = 0.83$. Then

$$C_3(n) = \frac{4}{9} \times 0.567^{3/2} \times 0.134^{-1/2} \times \sqrt{\frac{3 - \sqrt{6} - 4p}{\sqrt{6} - 1 + 13p}} (6 - \sqrt{6} - 13p),$$

where $p = \frac{1}{13(n-2)}$.

Thus, (2.22) is reduced to

$$(2.23) \quad \begin{aligned} 0 \leq & \left[-0.433n + C_3 \sqrt{\frac{\delta}{2} [\sigma n + n - 3 + 5\delta - 0.34(n + \delta)]} \right. \\ & \left. + (2.3505 - \frac{3}{2(n+4)})\delta + 0.567 \right] \int_M |\nabla A|^2 dM \\ & + \left[-0.7835 - C_3 \left(2\epsilon - \sigma\epsilon - \frac{0.17}{6.64\epsilon} \right) \right] \int_M (S - n) |\nabla A|^2 dM. \end{aligned}$$

We put

$$\delta = \frac{n}{22},$$

$$\begin{aligned} g_1(x) = & -0.433x + \sqrt{\frac{16 \times 0.567^3}{81 \times 0.134}} \times \sqrt{\frac{3 - \sqrt{6} - \frac{4}{13(x-2)}}{\sqrt{6} - 1 + \frac{1}{x-2}}} \\ & \times (6 - \sqrt{6} - \frac{1}{x-2}) \sqrt{\frac{x}{44} \left(\frac{\sqrt{17} + 3}{2}x - 3 + \frac{5x}{22} - \frac{0.34 \times 23x}{22} \right)} \\ & + \frac{2.3505x}{22} - \frac{3x}{44(x+4)} + 0.567, \end{aligned}$$

$$\begin{aligned} g_2(x) = & -0.7835 - \sqrt{\frac{16 \times 0.567^3}{81 \times 0.134}} \times \sqrt{\frac{3 - \sqrt{6} - \frac{4}{13(x-2)}}{\sqrt{6} - 1 + \frac{1}{x-2}}} \\ & \times (6 - \sqrt{6} - \frac{1}{x-2}) \left[\sqrt{\frac{(1 - \sqrt{17})^2 x}{4 \times 176 \left(\frac{\sqrt{17} + 3}{2}x - 3 + \frac{5x}{22} - \frac{0.34 \times 23x}{22} \right)}} \right. \\ & \left. - \frac{17}{664} \times \sqrt{\frac{176}{x} \left(\frac{\sqrt{17} + 3}{2}x - 3 + \frac{5x}{22} - \frac{0.34 \times 23x}{22} \right)} \right]. \end{aligned}$$

Now we are in a position to prove that both $g_1(x)$ and $g_2(x)$ are negative for $x \geq 6$.

Case I. Noting that

$$0.433x - \frac{2.3505x}{22} + \frac{3x}{44(x+4)} - 0.567 > 0, \text{ for } x \geq 6,$$

we take

$$\begin{aligned} g_1(x) &= \left[\sqrt{\frac{16 \times 0.567^3}{81 \times 0.134}} \times \sqrt{\frac{3 - \sqrt{6} - \frac{4}{13(x-2)}}{\sqrt{6} - 1 + \frac{1}{x-2}}} \left(6 - \sqrt{6} - \frac{1}{x-2}\right) \right. \\ &\quad \times \sqrt{\frac{x}{44} \left(\frac{\sqrt{17} + 3}{2}x - 3 + \frac{5x}{22} - \frac{0.34 \times 23x}{22} \right)} \\ &\quad \left. + 0.433x - \frac{2.3505x}{22} + \frac{3x}{44(x+4)} - 0.567 \right]^{-1} \\ &\quad \times \left(\frac{1}{(x-2)(x+4)} \right)^2 \frac{Z(x) - W(x)}{(\sqrt{6} - 1)(x-2) + 1}, \end{aligned}$$

where

$$\begin{aligned} Z(x) &= \frac{0.567^3 \times 16}{0.134 \times 81 \times 44} [(6 - \sqrt{6})(x-2) - 1]^2 \left[(3 - \sqrt{6})(x-2) - \frac{4}{13} \right] \\ &\quad \times \left[\frac{(11\sqrt{17} + 38 - 0.34 \times 23)x}{22} - 3 \right] (x+4)^2 x, \end{aligned}$$

and

$$\begin{aligned} W(x) &= \left[0.433x(x+4) - \frac{2.3505x(x+4)}{22} + \frac{3x}{44} - 0.567(x+4) \right]^2 \\ &\quad \times [(\sqrt{6} - 1)(x-2) + 1](x-2)^2. \end{aligned}$$

We get

$$(2.24) \quad Z(x) - W(x) \leq Q_1(x), \text{ for } x \geq 0,$$

where $Q_1(x) = -0.00868x^7 + 0.0575x^6 + 0.207x^5 - 3.126x^4 + 2.331x^3 + 30.434x^2 - 69.56x + 40$.

By calculating the determinant of the Sylvester matrix associated to Q_1 and Q_1' , we get

$$(2.25) \quad R(Q_1, Q_1') \approx -32.12.$$

It implies that the discriminant of $Q_1(x)$ is negative, which equals to $\frac{(-1)^{21}}{-0.00868} R(Q_1, Q_1')$. By a classical result for discriminant of polynomials, there exists at least one pair of complex conjugate roots for $Q_1(x)$. By a direct computation, we have

$$\begin{aligned} Q_1(-6) &= 501.124, \quad Q_1(-5) = -166.787, \quad Q_1(0) = 40, \\ Q_1(1.5) &= -1.74319, \quad Q_1(2) = 0.44096, \quad Q_1(3) = -11.8077. \end{aligned}$$

Thus, $Q_1(x)$ possesses five real roots in the interval $(-6, 3)$.

Hence

$$Q_1(x) < 0, \text{ for } x \geq 3.$$

Therefore, we have

$$g_1(x) < 0, \text{ for } x \geq 6.$$

Case II. Put

$$U_1(x) = \sqrt{\frac{3 - \sqrt{6} - \frac{4}{13(x-2)}}{\sqrt{6} - 1 + \frac{1}{x-2}}}, \quad U_2(x) = \left(6 - \sqrt{6} - \frac{1}{x-2}\right)$$

and

$$U_3(x) = \sqrt{\frac{22x}{(11\sqrt{17} + 38 - 0.34 \times 23)x - 66}}.$$

By a direct computation, we have

$$\frac{\partial}{\partial x} U_1(x) = \frac{35 - 9\sqrt{6}}{26U_1[(\sqrt{6} - 1)(x - 2) + 1]^2},$$

$$\frac{\partial}{\partial x} U_2(x) = \frac{1}{(x-2)^2}, \quad \frac{\partial}{\partial x} U_3(x) = -\frac{3U_3^3(x)}{2x^2}.$$

Thus, we have

$$\begin{aligned} \frac{\partial}{\partial x} g_2(x) &= \sqrt{\frac{16 \times 0.567^3}{81 \times 0.134}} \frac{\partial}{\partial x} \left\{ U_1(x) U_2(x) \right. \\ &\quad \left. \times \left[\frac{17\sqrt{11}}{166U_3(x)} + \frac{(\sqrt{17} - 1)\sqrt{11}}{88} U_3(x) \right] \right\} \\ &= \sqrt{\frac{16 \times 0.567^3}{81 \times 0.134}} \left\{ \left[\frac{(35 - 9\sqrt{6})U_2}{26U_1[(\sqrt{6} - 1)(x - 2) + 1]^2} + \frac{U_1}{(x - 2)^2} \right] \right. \\ &\quad \left. \times \left[\frac{17\sqrt{11}}{166U_3} + \frac{(\sqrt{17} - 1)\sqrt{11}}{88} U_3 \right] \right. \\ &\quad \left. - \left[\frac{(\sqrt{17} - 1)\sqrt{11}}{88} - \frac{17\sqrt{11}}{166U_3^2} \right] \frac{3U_1 U_2 U_3^3}{2x^2} \right\} \end{aligned}$$

$$\begin{aligned}
&= \frac{\sqrt{\frac{16 \times 0.567^3}{81 \times 0.134}} [(11\sqrt{17} + 38 - 0.34 \times 23)x - 66]^{-2}}{U_1 U_3 (x-2)^2 x [(\sqrt{6} - 1)(x-2) + 1]^2} \\
&\quad \times \left\{ \left[\frac{(35 - 9\sqrt{6})U_2(x-2)^2}{26} + U_1^2 [(\sqrt{6} - 1)(x-2) + 1]^2 \right] \right. \\
&\quad \times \left[\frac{17\sqrt{11}}{166} + \frac{(\sqrt{17} - 1)\sqrt{11}}{88} U_3^2 \right] [(11\sqrt{17} + 30.18)x - 66]^2 x \\
&\quad \left. - \frac{3U_3^2}{x} [(11\sqrt{17} + 30.18)x - 66]^2 [(\sqrt{6} - 1)(x-2) + 1]^2 \right. \\
&\quad \left. \times U_1^2 \left[\frac{(\sqrt{17} - 1)\sqrt{11}U_3^2}{176} - \frac{17\sqrt{11}}{332} \right] U_2(x-2)^2 \right\} \\
&= \frac{\sqrt{\frac{16 \times 0.567^3}{81 \times 0.134}} [(11\sqrt{17} + 38 - 0.34 \times 23)x - 66]^{-2}}{U_1 U_3 (x-2)^2 x [(\sqrt{6} - 1)(x-2) + 1]^2} R(x),
\end{aligned}$$

where

$$\begin{aligned}
R(x) &= \left\{ \frac{(35 - 9\sqrt{6})[(6 - \sqrt{6})(x-2) - 1](x-2)}{26} \right. \\
&\quad \left. + \left[(3 - \sqrt{6})(x-2) - \frac{4}{13} \right] [(\sqrt{6} - 1)(x-2) + 1] \right\} \\
&\quad \times \left\{ \frac{17\sqrt{11}}{166} [(11\sqrt{17} + 30.18)x - 66]^2 x \right. \\
&\quad \left. + \frac{(\sqrt{17} - 1)\sqrt{11}}{4} [(11\sqrt{17} + 30.18)x - 66]x^2 \right\} \\
&\quad - 66 \left[(3 - \sqrt{6})(x-2) - \frac{4}{13} \right] [(\sqrt{6} - 1)(x-2) + 1] \\
&\quad \times \left\{ \frac{(\sqrt{17} - 1)\sqrt{11}x}{8} - \frac{17\sqrt{11}}{332} [(11\sqrt{17} + 30.18)x - 66] \right\} \\
&\quad \times [(6 - \sqrt{6})(x-2) - 1](x-2).
\end{aligned}$$

We have

$$(2.26) \quad R(x) \geq 10000Q_2(x), \text{ for } x \geq 0,$$

where $Q_2(x) = 0.7633x^5 - 5.1552x^4 + 13.4534x^3 - 17.435x^2 + 11.598x - 3.2064$.

By calculating the determinant of the Sylvester matrix associated to Q_2 and Q_2' , we get

$$(2.27) \quad R(Q_2, Q_2') \approx -0.145.$$

Hence, the discriminant of $Q_2(x)$ is negative. So, there exists at least one pair of complex conjugate roots for $Q_2(x)$.

On the other hand, we get

$$\begin{aligned} Q_2(0) &= -3.2064, & Q_2(1) &= 0.0181, \\ Q_2(2) &= -0.1808, & Q_2(3) &= 5.8251. \end{aligned}$$

Hence

$$Q_2(x) > 0, \text{ for } x \geq 3.$$

It implies that $g_2(x)$ is increasing for $x \geq 3$. Notice that

$$\lim_{x \rightarrow \infty} g_2(x) \approx -0.044.$$

Therefore, we have

$$g_2(x) < 0, \text{ for } x \geq 6.$$

From(2.23), we have $\nabla A = 0$ and $S = n$, i.e., M is a Clifford torus. \square

Remark. In fact, we can enlarge the second pinching interval in Theorem 1.1 to $[n, n + \frac{n}{21.6}]$ by modifying the parameters θ and θ_1 .

3. SELF-SHRINKERS IN THE EUCLIDEAN SPACE

Let M be an n -dimensional complete self-shrinker with polynomial volume growth in \mathbb{R}^{n+1} . We shall make use of the following convention on the range of indices:

$$1 \leq i, j, k, \dots \leq n.$$

We choose a local orthonormal frame $\{e_1, e_2, \dots, e_{n+1}\}$ near a fixed point $x \in M$ over \mathbb{R}^{n+1} such that $\{e_i\}$ are tangent to M and e_{n+1} equals to the unit inner normal vector ξ . Let $\{\omega_1, \omega_2, \dots, \omega_{n+1}\}$ be the dual frame fields of $\{e_1, e_2, \dots, e_{n+1}\}$. Denote by A , H , S and R_{ijkl} the second fundamental form, the mean curvature, the squared length of the second fundamental form and the Riemannian curvature tensor of M , respectively. Then we have

$$(3.1) \quad A = \sum_{i,j} h_{ij} \omega_i \otimes \omega_j, \quad h_{ij} = h_{ji},$$

$$(3.2) \quad S = |A|^2 = \sum_{i,j} h_{ij}^2,$$

$$(3.3) \quad H = \text{Trace } A = \sum_i h_{ii}.$$

The Gauss and Codazzi equations are given by

$$(3.4) \quad R_{ijkl} = h_{ik}h_{jl} - h_{il}h_{jk},$$

and

$$(3.5) \quad h_{ijk} = h_{ikj}.$$

We have the Ricci identities.

$$(3.6) \quad h_{ijkl} - h_{ijlk} = \sum_m h_{im} R_{mjkl} + \sum_m h_{mj} R_{mikl},$$

$$(3.7) \quad h_{ijklm} - h_{ijkml} = \sum_r h_{rjk} R_{rilm} + \sum_r h_{irk} R_{rjlm} + \sum_r h_{ijr} R_{rklm}.$$

To simplify computation, we choose the local orthonormal frame such that its tangential covariant derivatives vanish at $x \in M$, i.e., $\bar{\nabla}_{e_i} e_j = h_{ij} \xi$.

Then we have

$$(3.8) \quad \nabla_{e_i} H = -\nabla_{e_i} \langle X, \xi \rangle = h_{ik} \langle X, e_k \rangle,$$

and

$$(3.9) \quad \begin{aligned} \text{Hess } H(e_i, e_j) &= -\nabla_{e_i} \nabla_{e_j} \langle X, \xi \rangle \\ &= h_{ijk} \langle X, e_k \rangle + h_{ij} - H h_{ik} h_{kj}. \end{aligned}$$

In [12], Colding and Minicozzi introduced the linear operator

$$\mathcal{L} = \Delta - \langle X, \nabla(\cdot) \rangle = e^{\frac{|X|^2}{2}} \text{Div}(e^{-\frac{|X|^2}{2}} \nabla(\cdot)).$$

They showed that \mathcal{L} is self-adjoint respect to the measure ρdM , where $\rho = e^{-\frac{|X|^2}{2}}$.

Thus, we get

$$(3.10) \quad \begin{aligned} \mathcal{L}|A|^2 &= \Delta|A|^2 - \langle X, \nabla|A|^2 \rangle \\ &= 2 \sum_{i,j} h_{ij} \Delta h_{ij} + 2|\nabla A|^2 - 2 \sum_{i,j,k} h_{ij} h_{ijk} \langle X, e_k \rangle \\ &= 2 \left(\sum_{i,j} h_{ij} \nabla_{e_i} \nabla_{e_j} H + H \sum_{i,j,k} h_{ij} h_{jk} h_{ki} - |A|^4 \right) \\ &\quad + 2|\nabla A|^2 - 2 \sum_{i,j,k} h_{ij} h_{ijk} \langle X, e_k \rangle \\ &= 2|A|^2 - 2|A|^4 + 2|\nabla A|^2. \end{aligned}$$

We have the following equalities for $|\nabla S|^2$ and $|\nabla^2 A|^2$.

$$(3.11) \quad |\nabla S|^2 = \frac{1}{2} \mathcal{L} S^2 + 2S^2(S-1) - 2S|\nabla A|^2,$$

$$(3.12) \quad |\nabla^2 A|^2 = \frac{1}{2} \mathcal{L} |\nabla A|^2 + (|A|^2 - 2)|\nabla A|^2 + 3(B_1 - 2B_2) + \frac{3}{2} |\nabla S|^2,$$

where $B_1 = \sum_{i,j,k,l,m} h_{ijk}h_{ijl}h_{km}h_{ml}$ and $B_2 = \sum_{i,j,k,l,m} h_{ijk}h_{klm}h_{im}h_{jl}$.

Following [16, 29], we have

$$(3.13) \quad 3(B_1 - 2B_2) \leq \sigma S |\nabla A|^2,$$

where $\sigma = \frac{\sqrt{17}+1}{2}$.

We choose a local orthonormal frame $\{e_i\}$ such that $h_{ij} = \mu_i \delta_{ij}$ at x . By Ricci identity (3.6), we have

$$(3.14) \quad t_{ij} := h_{ijij} - h_{jiji} = \mu_i \mu_j (\mu_i - \mu_j).$$

By a computation, we obtain

$$(3.15) \quad \int_M (B_1 - 2B_2) \rho dM = \int_M \left(\frac{1}{2}G - \frac{1}{4}|\nabla S|^2 \right) \rho dM,$$

where $G = \sum_{i,j} t_{ij}^2 = 2(Sf_4 - f_3^2)$ and $f_k = \text{Trace } A^k = \sum_i \mu_i^k$.

By Lemma 4.2 in [16], we have

$$(3.16) \quad 3(B_1 - 2B_2) \leq (S + C_2 G^{1/3}) |\nabla A|^2,$$

where $C_2 = \frac{2\sqrt{6}+3}{\sqrt[3]{21\sqrt{6}+103/2}}$.

Set $u_{ijkl} = \frac{1}{4}(h_{ijkl} + h_{lijk} + h_{klij} + h_{jkli})$. Notice that u_{ijkl} is symmetric in i, j, k, l . Then we have

$$\begin{aligned} \sum_{i,j,k,l} (h_{ijkl}^2 - u_{ijkl}^2) &= \frac{1}{16} \sum_{i,j,k,l} [(h_{ijkl} - h_{lijk})^2 + (h_{ijkl} - h_{klij})^2 \\ &\quad + (h_{ijkl} - h_{jkli})^2 + (h_{lijk} - h_{klij})^2 \\ &\quad + (h_{lijk} - h_{jkli})^2 + (h_{klij} - h_{jkli})^2] \\ &\geq \frac{6}{16} \sum_{i \neq j} [(h_{ijij} - h_{jiji})^2 + (h_{jiji} - h_{ijij})^2] \\ (3.17) \quad &= \frac{3}{4}G, \end{aligned}$$

i.e.,

$$(3.18) \quad |\nabla^2 A|^2 \geq \frac{3}{4}G.$$

Now we are in a position to prove our rigidity theorem for self-shrinkers in the Euclidean space.

Proof of Theorem 1.2. Combining (3.12), (3.15), (3.16) and (3.18), using Young's inequality, we drive the following inequality for arbitrary

$\theta \in (0, 1)$.

$$\begin{aligned}
& \frac{3}{4}(1-\theta) \int_M G \rho dM + \frac{3\theta}{8} \int_M |\nabla S|^2 \rho dM \\
\leq & \int_M |\nabla^2 A|^2 \rho dM - \frac{3\theta}{2} \int_M B \rho dM \\
= & \int_M [(S-2)|\nabla A|^2 + \frac{3}{2}|\nabla S|^2] \rho dM \\
& + 3\left(1 - \frac{\theta}{2}\right) \int_M (B_1 - 2B_2) \rho dM \\
\leq & \int_M [(S-2)|\nabla A|^2 + \frac{3}{2}|\nabla S|^2] \rho dM \\
& + \left(1 - \frac{\theta}{2}\right) \int_M (S + C_2 G^{1/3}) |\nabla A|^2 \rho dM \\
\leq & \int_M \left\{ \left[\left(2 - \frac{\theta}{2}\right) S - 2 \right] |\nabla A|^2 + \frac{3}{2} |\nabla S|^2 \right\} \rho dM \\
(3.19) \quad & + C_4 \int_M |\nabla A|^3 \rho dM + \frac{3}{4}(1-\theta) \int_M G \rho dM,
\end{aligned}$$

where $C_4(\theta) = \frac{4}{9} C_2^{3/2} \left(1 - \frac{\theta}{2}\right)^{3/2} (1-\theta)^{-1/2}$. This implies that

$$\begin{aligned}
0 \leq & \int_M \left[\left(2 - \frac{\theta}{2}\right) S - 2 \right] |\nabla A|^2 \rho dM \\
(3.20) \quad & + \left(\frac{3}{2} - \frac{3\theta}{8}\right) \int_M |\nabla S|^2 \rho dM + C_4 \int_M |\nabla A|^3 \rho dM.
\end{aligned}$$

Applying the divergence theorem, we get

$$\begin{aligned}
& - \int_M \nabla |\nabla A| \cdot \nabla S \rho dM \\
= & \int_M |\nabla A| \Delta S \rho dM + \int_M |\nabla A| \nabla S \cdot \nabla \rho dM \\
= & \int_M |\nabla A| \Delta S \rho dM + \int_M |\nabla A| (-\langle X, \nabla S \rangle) \rho dM \\
(3.21) \quad & = \int_M |\nabla A| \mathcal{L} S \rho dM.
\end{aligned}$$

From (3.10), (3.12), (3.13) and (3.21), for arbitrary $\epsilon > 0$, we have

$$\begin{aligned}
\int_M |\nabla A|^3 \rho dM &= \int_M (S^2 - S + \frac{1}{2} \mathcal{L}|A|^2) |\nabla A| \rho dM \\
&= \int_M (S^2 - S) |\nabla A| \rho dM - \frac{1}{2} \int_M \nabla |\nabla A| \cdot \nabla S \rho dM \\
&\leq \int_M (S^2 - S) |\nabla A| \rho dM + \epsilon \int_M |\nabla^2 A|^2 \rho dM \\
&\quad + \frac{1}{16\epsilon} \int_M |\nabla S|^2 \rho dM \\
&\leq \int_M (S^2 - S) |\nabla A| \rho dM + \left(\frac{3\epsilon}{2} + \frac{1}{16\epsilon}\right) \int_M |\nabla S|^2 \rho dM \\
(3.22) \quad &+ \epsilon \int_M (\sigma S + S - 2) |\nabla A|^2 \rho dM.
\end{aligned}$$

From (3.11), we have

$$(3.23) \quad \frac{1}{2} \int_M |\nabla S|^2 \rho dM = \int_M S(S-1)^2 \rho dM - \int_M (S-1) |\nabla A|^2 \rho dM.$$

When $1 \leq S \leq 1 + \delta$, for $\epsilon_1 = \theta_1 \epsilon > 0$, we obtain

$$\begin{aligned}
\int_M S(S-1) |\nabla A| \rho dM &\leq (2 + 2\delta) \epsilon_1 \int_M S(S-1) \rho dM \\
&\quad + \frac{1}{(8 + 8\delta) \epsilon_1} \int_M S(S-1) |\nabla A|^2 \rho dM \\
&\leq (2 + 2\delta) \epsilon_1 \int_M S(S-1) \rho dM \\
(3.24) \quad &+ \frac{1}{8\epsilon_1} \int_M (S-1) |\nabla A|^2 \rho dM.
\end{aligned}$$

Substituting (3.22), (3.23) and (3.24) into (3.20), we have

$$\begin{aligned}
0 &\leq \int_M \left[\left(2 - \frac{\theta}{2}\right) S - 2 \right] |\nabla A|^2 \rho dM + \left(\frac{3}{2} - \frac{3\theta}{8}\right) \int_M |\nabla S|^2 \rho dM \\
&\quad + C_4 \left[(2 + 2\delta) \epsilon_1 \int_M S(S-1) \rho dM + \frac{1}{8\epsilon_1} \int_M (S-1) |\nabla A|^2 \rho dM \right. \\
&\quad \left. + \epsilon \int_M (\sigma S + S - 2) |\nabla A|^2 \rho dM + \left(\frac{3\epsilon}{2} + \frac{1}{16\epsilon}\right) \int_M |\nabla S|^2 \rho dM \right]
\end{aligned}$$

$$\begin{aligned}
&\leq \int_M \left[\left(2 - \frac{\theta}{2}\right)S - 2 + \frac{C_4}{8\epsilon_1}(S-1) + 2C_4\epsilon_1(1+\delta) \right. \\
&\quad \left. + C_4\epsilon(\sigma S + S - 2) \right] |\nabla A|^2 \rho dM + \left[3 - \frac{3\theta}{4} + C_4\left(3\epsilon + \frac{1}{8\epsilon}\right) \right] \\
&\quad \times \left[\int_M \delta S(S-1)\rho dM - \int_M (S-1)|\nabla A|^2 \rho dM \right] \\
&= \int_M \left[-\frac{\theta}{2} + \frac{\sqrt{17}+3}{2}C_4\epsilon + \left(5\epsilon + \frac{1}{8\epsilon}\right)\delta C_4 \right. \\
&\quad \left. + \left(3 - \frac{3\theta}{4}\right)\delta + 2C_4(1+\delta)(\epsilon_1 - \epsilon) \right] |\nabla A|^2 \rho dM \\
(3.25) \quad &- \int_M \left[1 - \frac{\theta}{4} - \frac{\sqrt{17}-3}{2}C_4\epsilon + \frac{C_4}{8\epsilon} - \frac{C_4}{8\epsilon_1} \right] (S-1)|\nabla A|^2 \rho dM.
\end{aligned}$$

Let $\epsilon_1 = \theta_1\epsilon$, $\theta_1 = 0.81$, $\epsilon = \sqrt{\frac{\delta}{4(\sqrt{17}+3)+16(1+\delta)(\theta_1-1)+40\delta}}$, $\theta = 0.836$. Then we have $C_4 \leq 1.066218$. Thus, (3.25) is reduced to

$$\begin{aligned}
0 &\leq \left[\frac{C_4}{4} \sqrt{40\delta^2 - 3.04(1+\delta)\delta + 4(\sqrt{17}+3)\delta} \right. \\
&\quad \left. - 0.418 + 2.373\delta \right] \int_M |\nabla A|^2 \rho dM \\
&\quad + \left[\frac{C_4}{8} \left(4\sqrt{17}\epsilon - 12\epsilon - \frac{1}{\epsilon} + \frac{1}{0.81\epsilon} \right) - 0.791 \right] \\
(3.26) \quad &\times \int_M (|A|^2 - 1)|\nabla A|^2 \rho dM.
\end{aligned}$$

We take $\delta = 1/21$. Then the coefficients of the integrals in (3.26) are both negative. Therefore, we have $|\nabla A| \equiv 0$ and $|A| \equiv 1$, i.e., M is a round sphere or a cylinder. \square

To attack the Chern conjecture, we would like to suggest a program that divides the second optimal pinching problem for minimal hypersurfaces in a sphere into several steps. Precisely, we propose the following unsolved problem.

Problem 3.1. *Let M be a compact minimal hypersurface in \mathbb{S}^{n+1} . Denote by A the second fundamental form of M . Is it possible to prove that there exists a positive integer k with $1 \leq k \leq 21$, such that if $0 \leq |A|^2 - n \leq \frac{n}{k}$, then $|A|^2 \equiv n$ and M is a Clifford torus, or $|A|^2 \equiv 2n$ and $k = 1$?*

For self-shrinkers in the Euclidean space, we would like to propose the following problem.

Problem 3.2. *Let M be an n -dimensional complete self-shrinker with polynomial volume growth in \mathbb{R}^{n+1} . Denote by A the second fundamental form of M . Is it possible to prove that there exists a positive integer k with $1 \leq k \leq 20$, such that if $0 \leq |A|^2 - 1 \leq \frac{1}{k}$, then $|A|^2 \equiv 1$ and M is a round sphere or a cylinder, or $|A|^2 \equiv 2$ and $k = 1$?*

REFERENCES

1. S. C. de Almeida and F. G. B. Brito, Closed 3-dimensional hypersurfaces with constant mean curvature and constant scalar curvature, *Duke Math. J.*, **61**(1990), 195-206.
2. H. D. Cao and H. Z. Li, A gap theorem for self-shrinkers of the mean curvature flow in arbitrary codimension, *Calc. Var. Partial Differ. Equations*, **46**(2013), 879-889.
3. E. Cartan, Sur des familles remarquables d'hypersurfaces isoparamétriques dans les espaces sphériques, *Math. Z.*, **45**(1939), 335-367.
4. S. P. Chang, On minimal hypersurfaces with constant scalar curvatures in S^4 , *J. Differ. Geom.*, **37**(1993), 523-534.
5. S. P. Chang, A closed hypersurface of constant scalar and mean curvatures in S^4 is isoparametric, *Comm. Anal. Geom.*, **1**(1993), 71-100.
6. S. P. Chang, On closed hypersurfaces of constant scalar curvatures and mean curvatures in S^{n+1} , *Pacific J. Math.*, **165**(1994), 67-76.
7. Q. M. Cheng and H. Nakagawa, Totally umbilic hypersurfaces, *Hiroshima Math. J.*, **20**(1990), 1-10.
8. Q. M. Cheng and G. X. Wei, A gap theorem of self-shrinkers, *Trans. Amer. Math. Soc.*, **367**(2015), 4895-4915.
9. S. Y. Cheng, On the Chern conjecture for minimal hypersurface with constant scalar curvatures in the spheres, *Tsing Hua Lectures on Geometry and Analysis*, International Press, Cambridge, MA, 1997, 59-78.
10. S. S. Chern, Minimal submanifolds in a Riemannian manifold. University of Kansas, Department of Mathematics Technical Report 19, Univ. of Kansas, Lawrence, Kan., 1968.
11. S. S. Chern, M. do Carmo and S. Kobayashi, Minimal submanifolds of a sphere with second fundamental form of constant length, *Functional Analysis and Related Fields*, Springer-Verlag, Berlin, 1970, 59-75.
12. T. H. Colding and W. P. Minicozzi II, Generic mean curvature flow I: generic singularities, *Ann. of Math.*, **175**(2012), 755-833.
13. Qing Ding, On spectral characterizations of minimal hypersurfaces in a sphere, *Kodai Math. J.*, **17**(1994), 320-328.
14. Qi Ding and Y. L. Xin, On Chern's problem for rigidity of minimal hypersurfaces in the spheres, *Adv. Math.*, **227**(2011), 131-145.
15. Qi Ding and Y. L. Xin, Volume growth, eigenvalue and compactness for self-shrinkers, *Asian J. Math.*, **17**(2013), 443-456.
16. Qi Ding and Y. L. Xin, The rigidity theorems of self-shrinkers, *Trans. Amer. Math. Soc.*, **366**(2014), 5067-5085.
17. J. Q. Ge and Z. Z. Tang, Chern conjecture and isoparametric hypersurfaces, *Differential geometry*, *Adv. Lect. Math.*, **22**, International Press, Somerville, MA, 2012, 49-60.

18. J. R. Gu, H. W. Xu, Z. Y. Xu and E. T. Zhao, A survey on rigidity problems in geometry and topology of submanifolds, Proceedings of the 6th International Congress of Chinese Mathematicians, *Adv. Lect. Math.*, **37**, Higher Education Press & International Press, Beijing-Boston, 2016, 79-99.
19. G. Huisken, Flow by mean curvature of convex surfaces into spheres, *J. Differ. Geom.*, **22**(1984), 237-266.
20. G. Huisken, Asymptotic behavior for singularities of the mean curvature flow, *J. Differ. Geom.*, **31**(1990), 285-299.
21. B. Lawson, Local rigidity theorems for minimal hypersurfaces, *Ann. of Math.*, **89**(1969), 187-197.
22. Nam Q. Le and N. Sesum, Blow-up rate of the mean curvature during the mean curvature flow and a gap theorem for self-shrinkers, *Comm. Anal. Geom.*, **19**(2011), 1-27.
23. A. M. Li and J. M. Li, An intrinsic rigidity theorem for minimal submanifolds in a sphere, *Arch. Math.*, **58**(1992), 582-594.
24. H. Z. Li and Y. Wei, Classification and rigidity of self-shrinkers in the mean curvature flow, *J. Math. Soc. Japan*, **66**(2014), 709-734.
25. Z. Q. Lu, Normal scalar curvature conjecture and its applications, *J. Funct. Anal.*, **261**(2011), 1284-1308.
26. H. F. Münzner, Isoparametrische hyperflächen in sphären, I, II, *Math. Ann.*, **251**, **256**(1980, 1981), 57-71, 215-232.
27. H. Muto, The first eigenvalue of the Laplacian of an isoparametric minimal hypersurface in a unit sphere, *Math. Z.*, **197**(1988), 531-549.
28. C. K. Peng and C. L. Terng, Minimal hypersurfaces of sphere with constant scalar curvature, *Ann. of Math. Stud.*, **103**, Princeton Univ. Press, Princeton, NJ, 1983, 177-198.
29. C. K. Peng and C. L. Terng, The scalar curvature of minimal hypersurfaces in spheres, *Math. Ann.*, **266**(1983), 105-113.
30. M. Scherfner, S. Weiss and S. T. Yau, A review of the Chern conjecture for isoparametric hypersurfaces in spheres, *Advances in Geometric Analysis*, *Adv. Lect. Math.*, **21**, International Press, Somerville, MA, 2012, 175-187.
31. Y. B. Shen, On intrinsic rigidity for minimal submanifolds in a sphere, *Sci. China Ser. A*, **32**(1989), 769-781.
32. J. Simons, Minimal varieties in Riemannian manifolds, *Ann. of Math.*, **88**(1968), 62-105.
33. Y. J. Suh. and H. Y. Yang, The scalar curvature of minimal hypersurfaces in a unit sphere, *Comm. Contemp. Math.*, **9**(2007), 183-200.
34. L. Verstraelen, Sectional curvature of minimal submanifolds, In: Proceedings Workshop on Differential Geometry, Univ. Southampton, 1986, 48-62.
35. S. M. Wei and H. W. Xu, Scalar curvature of minimal hypersurfaces in a sphere, *Math. Res. Lett.*, **14**(2007), 423-432.
36. H. W. Xu, Pinching theorems, global pinching theorems and eigenvalues for Riemannian submanifolds, Ph.D. dissertation, Fudan University, 1990.
37. H. W. Xu, A rigidity theorem for submanifolds with parallel mean curvature in a sphere, *Arch. Math.*, **61**(1993), 489-496.
38. H. W. Xu and L. Tian, A new pinching theorem for closed hypersurfaces with constant mean curvature in S^{n+1} , *Asian J. Math.*, **15**(2011), 611-630.
39. H. W. Xu and Z. Y. Xu, The second pinching theorem for hypersurfaces with constant mean curvature in a sphere, *Math. Ann.*, **356**(2013), 869-883.

40. H. W. Xu and Z. Y. Xu, A new characterization of the Clifford torus via scalar curvature pinching, *J. Funct. Anal.*, **267**(2014), 3931-3962.
41. H. W. Xu and Z. Y. Xu, A refined version of the second pinching theorem for CMC hypersurfaces in a sphere, in preparation.
42. H. C. Yang and Q. M. Cheng, A note on the pinching constant of minimal hypersurfaces with constant scalar curvature in the unit sphere, *Kexue Tongbao*, **35**(1990), 167-170; *Chinese Sci. Bull.*, **36**(1991), 1-6.
43. H. C. Yang and Q. M. Cheng, An estimate of the pinching constant of minimal hypersurfaces with constant scalar curvature in the unit sphere, *Manuscripta Math.*, **84**(1994), 89-100.
44. H. C. Yang and Q. M. Cheng, Chern's conjecture on minimal hypersurfaces, *Math. Z.*, **227**(1998), 377-390.
45. S. T. Yau, Submanifolds with constant mean curvature, I, II, *Amer. J. Math.*, **96**, **97**(1974, 1975), 346-366, 76-100.
46. S. T. Yau, Problem section, In: Seminar on Differential Geometry, *Ann. of Math. Stud.*, **102**, Princeton Univ. Press, Princeton, NJ, 1982, 669-706.
47. Q. Zhang, The pinching constant of minimal hypersurfaces in the unit spheres, *Proc. Amer. Math. Soc.*, **138**(2010), 1833-1841.

CENTER OF MATHEMATICAL SCIENCES, ZHEJIANG UNIVERSITY, HANGZHOU 310027,
CHINA

E-mail address: xuhw@cms.zju.edu.cn; srxwing@zju.edu.cn