

# The Fadell-Rabinowitz index and multiplicity of non-contractible closed geodesics on Finsler $\mathbb{R}P^n$

Hui Liu<sup>1\*</sup>

<sup>1</sup> Key Laboratory of Wu Wen-Tsun Mathematics, Chinese Academy of Sciences,  
School of Mathematical Sciences, University of Science and Technology of China,  
Hefei, Anhui 230026, People's Republic of China

## Abstract

*In this paper, we prove that for every irreversible Finsler  $n$ -dimensional real projective space  $(\mathbb{R}P^n, F)$  with reversibility  $\lambda$  and flag curvature  $K$  satisfying  $\frac{16}{9} \left( \frac{\lambda}{1+\lambda} \right)^2 < K \leq 1$  with  $\lambda < 3$ , there exist at least  $n-1$  non-contractible closed geodesics. In addition, if the metric  $F$  is bumpy with  $\frac{64}{25} \left( \frac{\lambda}{1+\lambda} \right)^2 < K \leq 1$  and  $\lambda < \frac{5}{3}$ , then there exist at least  $2[\frac{n+1}{2}]$  non-contractible closed geodesics, which is the optical lower bound due to Katok's example. The main ingredients of the proofs are the Fadell-Rabinowitz index theory of non-contractible closed geodesics on  $(\mathbb{R}P^n, F)$  and the  $S^1$ -equivariant Poincaré series of the non-contractible component of the free loop space on  $\mathbb{R}P^n$ .*

**Key words:** Real projective space, non-contractible closed geodesics, Fadell-Rabinowitz index, Poincaré series, multiplicity.

**AMS Subject Classification:** 53C22, 58E05, 58E10.

## 1 Introduction and main results

In this paper, we are interested in the existence of non-contractible closed geodesics on real projective space  $\mathbb{R}P^n$  with a irreversible Finsler metric, which is the typically non-simply connected manifold with the fundamental group  $\pi_1(\mathbb{R}P^n) = \mathbf{Z}_2$ .

A closed curve on a Finsler manifold is a closed geodesic if it is locally the shortest path connecting any two nearby points on this curve. As usual, on any Finsler manifold  $(M, F)$ , a closed geodesic  $c : S^1 = \mathbf{R}/\mathbf{Z} \rightarrow M$  is *prime* if it is not a multiple covering (i.e., iteration) of any other closed geodesics. Here the  $m$ -th iteration  $c^m$  of  $c$  is defined by  $c^m(t) = c(mt)$ . The inverse curve

---

\*Partially supported by NSFC (No.11401555), Anhui Provincial Natural Science Foundation (No. 1608085QA01).  
E-mail: huiliu@ustc.edu.cn.

$c^{-1}$  of  $c$  is defined by  $c^{-1}(t) = c(1-t)$  for  $t \in \mathbf{R}$ . Note that unlike Riemannian manifold, the inverse curve  $c^{-1}$  of a closed geodesic  $c$  on a irreversible Finsler manifold need not be a geodesic. We call two prime closed geodesics  $c$  and  $d$  *distinct* if there is no  $\theta \in (0, 1)$  such that  $c(t) = d(t + \theta)$  for all  $t \in \mathbf{R}$ . We shall omit the word *distinct* when we talk about more than one prime closed geodesic. For a closed geodesic  $c$  on  $(M, F)$ , denote by  $P_c$  the linearized Poincaré map of  $c$ . Recall that a Finsler metric  $F$  is *bumpy* if all the closed geodesics on  $(M, F)$  are non-degenerate, i.e.,  $1 \notin \sigma(P_c)$  for any closed geodesic  $c$ .

There have been a great deal of works on the multiplicity of closed geodesics for simply connected manifolds (cf. [Ano], [Ban], [Bot], [Fra], [Kat], [Kli], [Liu], [LiL], [Lon], [LyF], [Zil]). In particular, Gromoll and Meyer [GM2] of 1969 established the existence of infinitely many distinct closed geodesics on  $M$ , provided that the Betti number sequence  $\{b_p(\Lambda M; \mathbf{Q})\}_{p \in \mathbf{N}}$  of the free loop space  $\Lambda M$  of  $M$  is unbounded is unbounded. Then in [ViS] of 1976, for compact simply connected manifold  $M$ , Vigué-Poirrier and Sullivan further proved this Betti number sequence is bounded if and only if  $M$  satisfies

$$H^*(M; \mathbf{Q}) \cong T_{d,n+1}(x) = \mathbf{Q}[x]/(x^{n+1} = 0)$$

with a generator  $x$  of degree  $d \geq 2$  and height  $n + 1 \geq 2$ , where  $\dim M = dn$ . Due to their works, many mathematicians are concerned with the compact globally symmetric spaces of rank 1 which consist in

$$S^n, \quad \mathbb{R}P^n, \quad \mathbb{C}P^n, \quad \mathbb{H}P^n \quad \text{and} \quad CaP^2,$$

where the Gromoll-Meyer assumption does not hold.

Recently the Maslov-type index theory for symplectic paths has been applied to study the closed geodesic problem on simply connected manifolds. In 2005, Bangert and Long proved the existence of at least two distinct closed geodesics on every Finsler  $(S^2; F)$  (which was published as [BaL] in 2010). Since then in the last ten years, a great number of results on the multiplicity of closed geodesics on simply connected Finsler manifolds has appeared, for which we refer readers to [DuL1]-[DuL2], [LoD], [Rad4]-[Rad6], [Wan1]-[Wan2], [Tai1], [HiR], [DLW1]-[DLW2] and the references therein.

As for the multiplicity of closed geodesics on non-simply connected manifolds whose free loop space possesses bounded Betti number sequence, Ballman et al. [BTZ] proved in 1981 that every Riemannian manifold with the fundamental group being a nontrivial finitely cyclic group and possessing a generic metric has infinitely many distinct closed geodesics. In 1984, Bangert and Hingston [BaH] proved that any Riemannian manifold with fundamental group an infinite cyclic group has infinitely many distinct closed geodesics. Since then, there seem to be very few works on the multiplicity of closed geodesics on non-simply connected manifolds. The main reason is that the topological structures of the free loop spaces on these manifolds are not well known, so that

the classical Morse theory is difficult to be applicable. In [XL] of 2015, Xiao and Long studied the topological structure of the non-contractible component of the free loop space on the real projective spaces with odd dimensions. Furthermore, they obtain the resonance identity for the non-contractible closed geodesics provided that the total number of closed geodesics on  $\mathbb{R}P^{2n+1}$  is finite. Based on these results, Duan, Long and Xiao [DLX] further proved the existence of at least two distinct non-contractible closed geodesics on  $\mathbb{R}P^3$  endowed with a bumpy and irreversible Finsler metric. In a very recent paper [Tai2], Taimanov studied the rational equivariant cohomology of the spaces of non-contractible loops in compact space forms and proved the existence of at least two distinct non-contractible closed geodesics on  $\mathbb{R}P^2$  endowed with a bumpy and irreversible Finsler metric.

In this paper, we continue to study the multiplicity of non-contractible closed geodesics on  $\mathbb{R}P^n$  under some natural conditions and obtain the following theorems:

**Theorem 1.1.** *On every irreversible Finsler  $n$ -dimensional real projective space  $(\mathbb{R}P^n, F)$  with reversibility  $\lambda$  and flag curvature  $K$  satisfying  $\frac{16}{9} \left( \frac{\lambda}{1+\lambda} \right)^2 < K \leq 1$  with  $\lambda < 3$ , there exist at least  $n - 1$  non-contractible closed geodesics.*

**Theorem 1.2.** *On every bumpy irreversible Finsler  $n$ -dimensional real projective space  $(\mathbb{R}P^n, F)$  with reversibility  $\lambda$  and flag curvature  $K$  satisfying  $\frac{64}{25} \left( \frac{\lambda}{1+\lambda} \right)^2 < K \leq 1$  with  $\lambda < \frac{5}{3}$ , there exist at least  $2\lfloor \frac{n+1}{2} \rfloor$  non-contractible closed geodesics.*

**Theorem 1.3.** *On every irreversible Finsler  $n$ -dimensional real projective space  $(\mathbb{R}P^n, F)$  with reversibility  $\lambda$  and flag curvature  $K$  satisfying  $\frac{4m^2}{(n-1)^2} \left( \frac{\lambda}{1+\lambda} \right)^2 < K \leq 1$  with  $\frac{n-1}{2} \leq m < n-1$ , there exist at least  $m + 1$  non-contractible closed geodesics.*

**Remark.** A. Katok [Kat] in 1973 found some non-symmetric Finsler metrics on  $S^n$  with only finitely many prime closed geodesics. The smallest number of closed geodesics on  $S^n$  that one obtains in these examples is  $2\lfloor \frac{n+1}{2} \rfloor$  (cf. [Zil]). Then D. V. Anosov in I.C.M. of 1974 conjectured that the lower bound of the number of closed geodesics on any Finsler sphere  $(S^n, F)$  should be  $2\lfloor \frac{n+1}{2} \rfloor$ , i.e., the number of closed geodesics in Katok's example. Let  $(S^n, N_\alpha)$  denote the sphere  $S^n$  endowed with the Katok metric, then it induces a Finsler metric for  $\mathbb{R}P^n$ , which is still denoted by  $N_\alpha$  for simplicity. Then there exist precisely  $2\lfloor \frac{n+1}{2} \rfloor$  non-contractible closed geodesics on  $(\mathbb{R}P^n, N_\alpha)$  (cf. Proposition 3.1, Remark 3.2 of [XL]). Thus it is natural to conjecture that the lower bound of the number of non-contractible closed geodesics on any Finsler real projective space  $(\mathbb{R}P^n, F)$  should be  $2\lfloor \frac{n+1}{2} \rfloor$ . Our Theorem 1.2 gives a confirmed answer to this conjecture for a generic case. Note that a closed geodesic on  $\mathbb{R}P^n$  is orientable if and only if  $n$  is odd, the iteration formulae of Maslov-type index of orientable and non-orientable closed geodesics are different, cf. [LiL], [Liu]. In the proofs of our theorems, we will not need the iteration formulae of Maslov-type index, but we use Morse index and rational coefficient homology only, so the orientable or non-orientable problem is inessential for our proofs.

The main ingredient of the proofs of Theorems 1.1 and 1.3 is the Fadell-Rabinowitz index

theory for non-contractible component of free loop space on  $(\mathbb{R}P^n, F)$ , which we will study in Section 3. In Section 2, we review the Morse theory of non-contractible closed geodesics, especially the  $S^1$ -equivariant Poincaré series of the non-contractible component of the free loop space on  $\mathbb{R}P^n$  derived by Taimanov [Tai2] which is the main ingredient of the proof of Theorem 1.2. The proofs of Theorems 1.1-1.3 will be given in Section 4.

## 2 Morse theory of non-contractible closed geodesics

Let  $M = (M, F)$  be a compact Finsler manifold, the space  $\Lambda = \Lambda M$  of  $H^1$ -maps  $\gamma : S^1 \rightarrow M$  has a natural structure of Riemannian Hilbert manifolds on which the group  $S^1 = \mathbf{R}/\mathbf{Z}$  acts continuously by isometries (cf. [Kli], [She]). This action is defined by  $(s \cdot \gamma)(t) = \gamma(t + s)$  for all  $\gamma \in \Lambda$  and  $s, t \in S^1$ . For any  $\gamma \in \Lambda$ , the energy functional is defined by

$$E(\gamma) = \frac{1}{2} \int_{S^1} F(\gamma(t), \dot{\gamma}(t))^2 dt. \quad (2.1)$$

It is  $C^{1,1}$  and invariant under the  $S^1$ -action. The critical points of  $E$  of positive energies are precisely the closed geodesics  $\gamma : S^1 \rightarrow M$ . The index form of the functional  $E$  is well defined along any closed geodesic  $c$  on  $M$ , which we denote by  $E''(c)$ . As usual, we denote by  $i(c)$  and  $\nu(c) - 1$  the Morse index and nullity of  $E$  at  $c$ . In the following, we denote by

$$\Lambda^\kappa = \{d \in \Lambda \mid E(d) \leq \kappa\}, \quad \Lambda^{\kappa-} = \{d \in \Lambda \mid E(d) < \kappa\}, \quad \forall \kappa \geq 0. \quad (2.2)$$

For a closed geodesic  $c$  we set  $\Lambda(c) = \{\gamma \in \Lambda \mid E(\gamma) < E(c)\}$ .

For  $m \in \mathbf{N}$  we denote the  $m$ -fold iteration map  $\phi_m : \Lambda \rightarrow \Lambda$  by  $\phi_m(\gamma)(t) = \gamma(mt)$ , for all  $\gamma \in \Lambda, t \in S^1$ , as well as  $\gamma^m = \phi_m(\gamma)$ . If  $\gamma \in \Lambda$  is not constant then the multiplicity  $m(\gamma)$  of  $\gamma$  is the order of the isotropy group  $\{s \in S^1 \mid s \cdot \gamma = \gamma\}$ . For a closed geodesic  $c$ , the mean index  $\hat{i}(c)$  is defined as usual by  $\hat{i}(c) = \lim_{m \rightarrow \infty} i(c^m)/m$ . Using singular homology with rational coefficients we consider the following critical  $\mathbf{Q}$ -module of a closed geodesic  $c \in \Lambda$ :

$$\overline{C}_*(E, c) = H_*((\Lambda(c) \cup S^1 \cdot c)/S^1, \Lambda(c)/S^1; \mathbf{Q}). \quad (2.3)$$

Following [Rad1], Section 6.2, we can use finite-dimensional approximations to  $\Lambda$  to apply the results of D. Gromoll and W. Meyer [GM1] to a given closed geodesic  $c$  which is isolated as a critical orbit. Then we have

**Proposition 2.1.** *Let  $k_j(c) \equiv \dim \overline{C}_j(E, c)$ . Then  $k_j(c)$  equal to 0 when  $j < i(c)$  or  $j > i(c) + \nu(c)$  and if  $\nu(c) = 0$ , then  $k_j(c) = 1$  for  $j = i(c)$ .*

For  $M = \mathbb{R}P^n$ , it is well known that  $\pi_1(\mathbb{R}P^n) = \mathbf{Z}_2 = \{e, g\}$  with  $e$  being the identity and  $g$  being the generator of  $\mathbf{Z}_2$  satisfying  $g^2 = e$ . Then the free loop space  $\Lambda M$  possesses a natural decomposition

$$\Lambda M = \Lambda_e M \bigsqcup \Lambda_g M,$$

where  $\Lambda_e M$  and  $\Lambda_g M$  are the two connected components of  $\Lambda M$  whose elements are homotopic to  $e$  and  $g$  respectively.

**Lemma 2.2.** (cf. Theorem 3 of [Tai2]) *For  $M = \mathbb{R}P^n$ , we have*

(i) *When  $n = 2k + 1$  is odd, the  $S^1$ -cohomology ring of  $\Lambda_g M$  has the form*

$$H^{S^1,*}(\Lambda_g M; \mathbf{Q}) = \mathbf{Q}[w, z] / \{w^{k+1} = 0\}, \quad \deg(w) = 2, \quad \deg(z) = 2k$$

*Then the  $S^1$ -equivariant Poincaré series of  $\Lambda_g M$  is given by*

$$\begin{aligned} P^{S^1}(\Lambda_g M; \mathbf{Q})(t) &= \frac{1 - t^{2k+2}}{(1 - t^2)(1 - t^{2k})} \\ &= \frac{1}{1 - t^2} + \frac{t^{2k}}{1 - t^{2k}} \\ &= (1 + t^2 + t^4 + \cdots + t^{2k} + \cdots) + (t^{2k} + t^{4k} + t^{6k} + \cdots), \end{aligned}$$

*which yields Betti numbers*

$$\begin{aligned} b_q &= \text{rank} H_q^{S^1}(\Lambda_g M; \mathbf{Q}) \\ &= \begin{cases} 2, & \text{if } q \in \{j(n-1) \mid j \in \mathbf{N}\}, \\ 1, & \text{if } q \in 2\mathbf{N}_0 \setminus \{j(n-1) \mid j \in \mathbf{N}\}, \\ 0, & \text{otherwise.} \end{cases} \end{aligned} \quad (2.4)$$

(ii) *When  $n = 2k$  is even, the  $S^1$ -cohomology ring of  $\Lambda_g M$  has the form*

$$H^{S^1,*}(\Lambda_g M; \mathbf{Q}) = \mathbf{Q}[w, z] / \{w^{2k} = 0\}, \quad \deg(w) = 2, \quad \deg(z) = 4k - 2$$

*Then the  $S^1$ -equivariant Poincaré series of  $\Lambda_g M$  is given by*

$$\begin{aligned} P^{S^1}(\Lambda_g M; \mathbf{Q})(t) &= \frac{1 - t^{4k}}{(1 - t^2)(1 - t^{4k-2})} \\ &= \frac{1}{1 - t^2} + \frac{t^{4k-2}}{1 - t^{4k-2}} \\ &= (1 + t^2 + t^4 + \cdots + t^{2k} + \cdots) + (t^{4k-2} + t^{2(4k-2)} + t^{3(4k-2)} + \cdots), \end{aligned}$$

*which yields Betti numbers*

$$\begin{aligned} b_q &= \text{rank} H_q^{S^1}(\Lambda_g M; \mathbf{Q}) \\ &= \begin{cases} 2, & \text{if } q \in \{2j(n-1) \mid j \in \mathbf{N}\}, \\ 1, & \text{if } q \in 2\mathbf{N}_0 \setminus \{2j(n-1) \mid j \in \mathbf{N}\}, \\ 0, & \text{otherwise.} \end{cases} \end{aligned} \quad (2.5)$$

Note that for a non-contractible prime closed geodesic  $c$ ,  $c^m \in \Lambda_g M$  if and only if  $m$  is odd. Thus we have the following Morse inequality for non-contractible closed geodesics:

**Theorem 2.3.** (cf. Theorem I.4.3 of [Cha]) *Assume that  $M = \mathbb{R}P^n$  be a Finsler manifold with finitely many non-contractible prime closed geodesics, denoted by  $\{c_j\}_{1 \leq j \leq k}$ . Set*

$$M_q = \sum_{1 \leq j \leq k, m \geq 1} \dim \overline{C}_q(E, c_j^{2m-1}), \quad q \in \mathbf{Z}.$$

*Then for every integer  $q \geq 0$  there holds*

$$\begin{aligned} M_q - M_{q-1} + \cdots + (-1)^q M_0 &\geq b_q - b_{q-1} + \cdots + (-1)^q b_0, \\ M_q &\geq b_q. \end{aligned} \tag{2.6}$$

### 3 Fadell-Rabinowitz index theory for non-contractible closed geodesics on $(\mathbb{R}P^n, F)$

In 1978, Fadell and Rabinowitz [FaR] introduced a topological index theory by using the equivariant Borel cohomology. Via a classifying map the cohomology ring  $H^*(BG)$  of a classifying space is a subring of the equivariant cohomology  $H_G^*(X)$  of the  $G$ -space  $X$ . For a characteristic class  $\eta \in H^*(BG)$  the Fadell-Rabinowitz index  $index_\eta X$  of  $X$  is the smallest non-negative integer  $k$  with  $\eta^k = 0$  in  $H_G^*(X)$ . Hence the classes  $\eta^j$  for  $j = 0, \dots, index_\eta X - 1$  define a sequence of subordinate cohomology classes. Fadell-Rabinowitz used this index to study the bifurcation of time periodic solutions from an equilibrium solution for Hamiltonian systems of ordinary differential equations. Ekeland and Hofer in [EkH] of 1987 used this index for the group  $S^1$  to derive a close relationship between the set of Maslov-type indices of closed characteristics on convex Hamiltonian energy surfaces in  $\mathbf{R}^{2n}$  and the set of even positive integers, which is the core in studying multiplicity and ellipticity of closed characteristics on compact convex hypersurfaces (cf. [LoZ]). Rademacher in [Rad2] of 1994 developped the relative version of Fadell-Rabinowitz index to study the existence of closed geodesics on compact manifolds, which played an important role in studying multiplicity and stability of closed geodesics on Finsler spheres, cf. [Rad4], [Wan3].

In this section, we use only singular homology modules with  $\mathbf{Q}$ -coefficients and apply the Fadell-Rabinowitz index to the non-contractible component  $\Lambda_g M$  of the free loop space on  $M = \mathbb{R}P^n$ , which are the main ingredients for proving our multiplicity results of non-contractible closed geodesics.

It is well known that there exists a non-contractible closed geodesic  $c$  on  $(\mathbb{R}P^n, F)$ , which is the minimal point of the energy functional  $E$  on  $\Lambda_g$ , set  $E(c) = \frac{\kappa_1^2}{2}$  for  $\kappa_1 > 0$ . For any  $\kappa > 0$ , we denote by

$$\Lambda_g^\kappa = \{\gamma \in \Lambda_g \mid E(\gamma) \leq \frac{1}{2}\kappa^2\}. \tag{3.1}$$

We define the function  $d_g : [\kappa_1, +\infty] \rightarrow \mathbf{N} \cup \{\infty\}$  by

$$d_g(\kappa) = index_\eta \Lambda_g^\kappa,$$

where  $H^*(BS^1) = H^*(CP^\infty) = \mathbf{Q}[\eta]$ .

**Definition 3.1.** The upper index  $\underline{\sigma}$  and lower index  $\overline{\sigma}$  are given by:

$$\underline{\sigma} = \liminf_{\kappa \rightarrow \infty} \frac{d_g(\kappa)}{\kappa}, \quad \overline{\sigma} = \limsup_{\kappa \rightarrow \infty} \frac{d_g(\kappa)}{\kappa},$$

we call  $[\underline{\sigma}, \overline{\sigma}]$  the global index interval.

Similarly to Proposition 5.2 of [Rad2], as a consequence of the Morse-Schoenberg comparison theorem and the proof of Myers' theorem, we have the following estimates for the global index interval in terms of the curvature.

**Lemma 3.2.** Let  $K$  and  $Ric$  be the flag curvature and Ricci curvature of the manifold  $(\mathbb{R}P^n, F)$  respectively, then we have

- (a) If  $K \leq \Delta^2$ , then  $\overline{\sigma} \leq \frac{\Delta(n-1)}{2\pi}$ .
- (b) If  $K \geq \Delta^2$ , then  $\overline{\sigma} \leq \frac{\delta(n-1)}{2\pi}$ .
- (c) If  $Ric \geq \delta^2(n-1)$ , then  $\underline{\sigma} \geq \frac{\delta}{2\pi}$ .

**Proof.** We give the proof of the first estimate; the other proofs are analogous. We can choose a sequence  $(F_i)$  of bumpy metrics converging in the strong  $C^2$  topology to  $F$  using the bumpy metrics theorem. Let  $E_i(\gamma) = \frac{1}{2} \int_{S^1} F_i(\dot{\gamma}(t))^2 dt$  be the energy functional and  $\Lambda_i^\kappa = \{\gamma \in \Lambda_g \mid E_i(\gamma) \leq \frac{1}{2}\kappa^2\}$ .

Choose a sequence  $(a_i) \subset \mathbf{R}$  with  $a = \lim_{i \rightarrow \infty} a_i$  and  $\Lambda_g^a \subset \Lambda_i^{a_i}$  for all  $i$ . We can choose a sequence  $(\Delta_i) \subset \mathbf{R}$  such that the sectional curvature of  $F_i$  is bounded from above by  $\Delta_i^2$  and  $\Delta = \lim_{i \rightarrow \infty} \Delta_i$ . It follows from the Morse-Schoenberg comparison theorem that the index  $i(c)$  of a closed geodesic  $c$  of the metric  $F_i$  with length at most  $a_i$  satisfies

$$i(c) \leq \left( \frac{a_i}{\pi} \Delta_i + 1 \right) (n-1) =: k_i. \quad (3.2)$$

Hence it follows from the Morse inequalities that  $H_{S^1}^k(\Lambda_i^{a_i}) = 0$  for all  $k > k_i$ . Since  $\Lambda_g^a \subset \Lambda_i^{a_i}$  it follows from the composition

$$H^k(BS^1) \rightarrow H_{S^1}^k(\Lambda_i^{a_i}) \rightarrow H_{S^1}^k(\Lambda_g^a).$$

of restriction homomorphisms that  $\eta^k = 0$  in  $H_{S^1}^k(\Lambda_g^a)$  for all  $k > \frac{k_i}{2}$ . Hence we have  $d_g(a) \leq \frac{k_i}{2}$ , which together with (3.2) and Definition 3.1 gives  $\overline{\sigma} \leq \frac{\Delta(n-1)}{2\pi}$ .  $\blacksquare$

**Corollary 3.3.** For  $\mathbb{R}P^n$  endowed with any Finsler metric  $F$ , we have  $\lim_{\kappa \rightarrow +\infty} d_g(\kappa) = +\infty$ .

**Proof.** Let  $(S^n, N_\alpha)$  denote the sphere  $S^n$  endowed with the Katok metric, then it induces a Finsler metric for  $\mathbb{R}P^n$ , which is still denoted by  $N_\alpha$  for simplicity. Therefore the natural projection

$$\pi : (S^n, N_\alpha) \rightarrow (\mathbb{R}P^n, N_\alpha)$$

is locally isometric. Since  $(S^n, N_\alpha)$  and  $(\mathbb{R}P^n, N_\alpha)$  have constant flag curvature 1, then for  $(\mathbb{R}P^n, N_\alpha)$ ,  $\underline{\sigma} > 0$  by Lemma 3.2. Thus for  $\mathbb{R}P^n$  endowed with any Finsler metric  $F$ ,  $\underline{\sigma} > 0$

also holds which implies  $\lim_{\kappa \rightarrow +\infty} d_g(\kappa) = +\infty$ . In fact, Let  $F, F^*$  be two Finsler metrics on  $\mathbb{R}P^n$  with

$$F^*(X)^2/D^2 \leq F(X)^2 \leq D^2 F^*(X)^2$$

for all tangent vector  $X$  and some  $D > 1$ . Let  $[\underline{\sigma}, \bar{\sigma}]$ ,  $[\underline{\sigma}^*, \bar{\sigma}^*]$  be the global index intervals with respect to the metrics  $F, F^*$ . Then by definition we have

$$D^{-1}\underline{\sigma} \leq \underline{\sigma}^* \leq D\underline{\sigma}, \quad D^{-1}\bar{\sigma} \leq \bar{\sigma}^* \leq D\bar{\sigma}$$

which implies our desired result.  $\blacksquare$

**Lemma 3.4.** *The function  $d_g : [\kappa_1, +\infty) \rightarrow \mathbf{N}$  is non-decreasing and  $\lim_{\lambda \searrow \kappa} d_g(\lambda) = d_g(\kappa)$ . Each discontinuous point of  $d_g$  is a critical value of the energy functional  $E$  on  $\Lambda_g$ . In particular, if  $d_g(\kappa) - d_g(\kappa-) \geq 2$ , then there are infinitely many prime non-contractible closed geodesics  $c$  with energy  $\kappa$ , where  $d_g(\kappa-) = \lim_{\epsilon \searrow 0} d_g(\kappa - \epsilon)$ ,  $t \searrow a$  means  $t > a$  and  $t \rightarrow a$ .*

**Proof.** We have that  $d_g(\lambda)$  is finite by the proof of Lemma 3.2. The monotony of  $d_g(\lambda)$  follows from the monotonicity property of Fadell-Rabinowitz index. Let  $V$  be a closed  $S^1$  invariant neighbourhood of  $\Lambda_g^\kappa$  with

$$d_g(\kappa) = \text{index}_\eta \Lambda_g^\kappa = \text{index}_\eta V,$$

which follows from the continuity property of Fadell-Rabinowitz index. Then there exist  $\epsilon, \bar{\epsilon} > 0$  such that for all  $\gamma \in \Lambda_g^{\kappa+\epsilon} - (\Lambda_g^{\kappa-\epsilon} \cup V)$  the gradient  $\nabla E$  of the energy functional  $E$  satisfies  $\|\nabla E(\gamma)\| \geq \bar{\epsilon}$ , cf. [Kli]. Hence there is a  $T > 0$  such that  $\Phi_T(\Lambda_g^{\kappa+\epsilon}) \subset V \cup \Lambda_g^{\kappa-\epsilon}$ , where  $\Phi_t : \Lambda_g \rightarrow \Lambda_g$  is the flow of the vector field  $-\nabla E$  on the non-contractible component  $\Lambda_g$  of the free loop space. It together with the monotonicity and continuity properties of Fadell-Rabinowitz index gives

$$d_g(\kappa) = \text{index}_\eta \Lambda_g^\kappa = \text{index}_\eta V \geq \text{index}_\eta \Lambda_g^{\kappa+\epsilon} = d_g(\kappa + \epsilon),$$

which implies  $d_g(\kappa) = d_g(\kappa + \epsilon)$  and also  $\lim_{\lambda \searrow \kappa} d_g(\lambda) = d_g(\kappa)$ .

In the following, we prove  $\text{index}_\eta Cr(\kappa) \geq d_g(\kappa) - d_g(\kappa-)$ , where  $Cr$  is the critical set of  $E$  on  $\Lambda_g$  with positive energy, that is,  $Cr(\kappa)$  is the set of non-contractible closed geodesics of length  $\kappa$ .

In fact, we assume that  $d_g(\kappa) - d_g(\kappa-) \geq 1$ ,  $U$  is a closed  $S^1$  invariant neighbourhood of  $Cr(\kappa)$  with  $\text{index}_\eta U = \text{index}_\eta Cr(\kappa)$  by the continuity property of Fadell-Rabinowitz index. Then there exists an  $\epsilon > 0$  such that  $U \cup \Lambda_g^{\kappa-\epsilon}$  is a  $S^1$  deformation retract of  $\Lambda_g^{\kappa+\epsilon}$ , that is

$$H_{S^1}^*(U \cup \Lambda_g^{\kappa-\epsilon}) = H_{S^1}^*(\Lambda_g^{\kappa+\epsilon})$$

and  $d_g(\kappa) = d_g(\kappa + \epsilon)$ ,  $d_g(\kappa-) = d_g(\kappa - \epsilon)$ . Then by the subadditivity property of Fadell-Rabinowitz index, we obtain

$$d_g(\kappa) = d_g(\kappa + \epsilon) \leq \text{index}_\eta U + \text{index}_\eta \Lambda_g^{\kappa-\epsilon} = \text{index}_\eta Cr(\kappa) + d_g(\kappa-)$$



which implies that each discontinuous point of  $d_g$  is a critical value of the energy functional  $E$  on  $\Lambda_g$  and for  $d_g(\kappa) - d_g(\kappa-) \geq 2$ , there are infinitely many prime non-contractible closed geodesics  $c$  with energy  $\kappa$ , since the Fadell-Rabinowitz index of a set consisting of finite  $S^1$  orbits is 1. ■

For each  $i \geq 1$ , define

$$\kappa_i = \inf\{\delta \in \mathbf{R} \mid d_g(\delta) \geq i\}. \quad (3.3)$$

Then  $\kappa_i$  is well defined by Corollary 3.3.

Now we can obtain the main results in this section.

**Theorem 3.5.** *Suppose there are only finitely many prime non-contractible closed geodesics on  $(\mathbb{R}P^n, F)$ . Then each  $\kappa_i$  is a critical value of  $E$  on  $\Lambda_g$ . If  $\kappa_i = \kappa_j$  for some  $i < j$ , then there are infinitely many prime non-contractible closed geodesics on  $(\mathbb{R}P^n, F)$ .*

**Proof.** By definition, each  $\kappa_i$  is a discontinuous point of  $d_g$ , then it is a critical value of  $E$  by Lemma 3.4. Now suppose  $\kappa_i = \kappa_j$  for some  $i < j$ . Then by (3.3), we have  $d_g(\kappa_i-) < i$  and  $d_g(\kappa_i) = d_z(\kappa_j) \geq j \geq d_z(\kappa_i-) + 2$ . Hence we have  $d_g(\kappa_i) - d_g(\kappa_i-) \geq 2$ . Thus Lemma 3.4 implies there are infinitely many prime non-contractible closed geodesics  $c$  with energy  $\kappa_i$ . This proves the theorem. ■

**Theorem 3.6.** *Suppose there are only finitely many prime non-contractible closed geodesics on  $(\mathbb{R}P^n, F)$ . Then for every  $i \in \mathbf{N}$ , there exists a non-contractible closed geodesic  $c$  on  $(\mathbb{R}P^n, F)$  such that*

$$E(c) = \kappa_i, \quad \overline{C}_{2i-2}(E, c) \neq 0. \quad (3.4)$$

**Proof.** Choose  $\epsilon$  small enough such that the interval  $(\kappa_i - \epsilon, \kappa_i + \epsilon)$  contains no critical values of  $E$  except  $\kappa_i$ . Since any  $\gamma \in \Lambda_g^{\kappa_i+\epsilon} \setminus \Lambda_g^{\kappa_i-\epsilon}$  is not a fixed point of the  $S^1$ -action, its isotropy group is finite. Hence we can use Lemma 6.11 of [FaR] to obtain

$$H_{S^1}^*(\Lambda_g^{\kappa_i+\epsilon}, \Lambda_g^{\kappa_i-\epsilon}) \cong H^*(\Lambda_g^{\kappa_i+\epsilon}/S^1, \Lambda_g^{\kappa_i-\epsilon}/S^1). \quad (3.5)$$

By the finiteness assumption of the number of prime non-contractible closed geodesics, a small perturbation on the energy functional can be applied to reduce each critical orbit to nearby non-degenerate ones. Thus similar to the proofs of Lemma 2 of [GM1] and Lemma 4 of [GM2], all the homological  $\mathbf{Q}$ -modules of  $(\Lambda_g^{\kappa_i+\epsilon}, \Lambda_g^{\kappa_i-\epsilon})$  is finitely generated. Therefore we can apply Theorem 5.5.3 and Corollary 5.5.4 on pages 243-244 of [Spa1] to obtain

$$H_*(\Lambda_g^{\kappa_i+\epsilon}/S^1, \Lambda_g^{\kappa_i-\epsilon}/S^1) \cong H^*(\Lambda_g^{\kappa_i+\epsilon}/S^1, \Lambda_g^{\kappa_i-\epsilon}/S^1). \quad (3.6)$$

Then we combining (3.5)-(3.6) with Theorem 1.4.2 of [Cha] to obtain

$$H_{S^1}^*(\Lambda_g^{\kappa_i+\epsilon}, \Lambda_g^{\kappa_i-\epsilon}) = \bigoplus_{E(c)=\kappa_i} \overline{C}_*(E, c). \quad (3.7)$$

Similar to P.431 of [EkH], we have

$$H^{2(i-1)}((\Lambda_g^{\kappa_i+\epsilon})_{S^1}, (\Lambda_g^{\kappa_i-\epsilon})_{S^1}) \xrightarrow{q^*} H^{2(i-1)}((\Lambda_g^{\kappa_i+\epsilon})_{S^1}) \xrightarrow{p^*} H^{2(i-1)}((\Lambda_g^{\kappa_i-\epsilon})_{S^1}), \quad (3.8)$$

where  $p$  and  $q$  are natural inclusions. Denote by  $f : (\Lambda_g^{\kappa_i+\epsilon})_{S^1} \rightarrow CP^\infty$  a classifying map and let  $f^\pm = f|_{(\Lambda_g^{\kappa_i\pm\epsilon})_{S^1}}$ . Then clearly each  $f^\pm : (\Lambda_g^{\kappa_i\pm\epsilon})_{S^1} \rightarrow CP^\infty$  is a classifying map on  $(\Lambda_g^{\kappa_i\pm\epsilon})_{S^1}$ . Let  $\eta \in H^2(CP^\infty)$  be the first universal Chern class.

By definition of  $\kappa_i$ , we have  $d_g(\kappa_i - \epsilon) < i$ , hence  $(f^-)^*(\eta^{i-1}) = 0$ . Note that  $p^*(f^+)^*(\eta^{i-1}) = (f^-)^*(\eta^{i-1})$ . Hence the exactness of (3.8) yields a  $\sigma \in H^{2(i-1)}((\Lambda_g^{\kappa_i+\epsilon})_{S^1}, (\Lambda_g^{\kappa_i-\epsilon})_{S^1})$  such that  $q^*(\sigma) = (f^+)^*(\eta^{i-1})$ . Since  $d_g(\kappa_i + \epsilon) \geq i$ , we have  $(f^+)^*(\eta^{i-1}) \neq 0$ . Hence  $\sigma \neq 0$ , and then

$$H_{S^1}^{2i-2}(\Lambda_g^{\kappa_i+\epsilon}, \Lambda_g^{\kappa_i-\epsilon}) = H^{2i-2}((\Lambda_g^{\kappa_i+\epsilon})_{S^1}, (\Lambda_g^{\kappa_i-\epsilon})_{S^1}) \neq 0.$$

Thus (3.4) follows from (3.7). ■

Note that by Proposition 2.1, (3.4) yields

$$E(c) = \kappa_i, \quad i(c) \leq 2i - 2 \leq i(c) + \nu(c). \quad (3.9)$$

Similar to Definition 1.4 of [LoZ], we have

**Definition 3.7.** A prime non-contractible closed geodesic  $c$  is  $(2m-1, i)$ -**variationally visible**, if there exist some  $m, i \in \mathbf{N}$  such that (3.4) holds for  $c^{2m-1}$  and  $\kappa_i$ . Denote by  $\mathcal{V}(\mathbb{R}P^n, F)$  the set of variationally visible non-contractible closed geodesics on  $(\mathbb{R}P^n, F)$ .

**Theorem 3.8.** Assume that  $\#\mathcal{V}(\mathbb{R}P^n, F) < +\infty$ , then we have

$$\sum_{c \in \mathcal{V}(\mathbb{R}P^n, F)} \frac{1}{\hat{i}(c)} \geq 1. \quad (3.10)$$

**Proof.** Applying Theorem 3.6 and (3.9), for every integer  $i \geq 1$ , there exists a prime non-contractible closed geodesic  $c_i \in \mathcal{V}(\mathbb{R}P^n, F)$  and some  $m_i \in \mathbf{N}$ , such that

$$|i(c_i^{2m_i-1}) - 2i| \leq 2n, \quad E(c_i^{2m_i-1}) = \kappa_i. \quad (3.11)$$

Define the density of a sequence  $a_i$  in  $\mathbf{N}$  to be the number(if it exists)

$$\delta := \lim_{N \rightarrow +\infty} \frac{1}{N} \# \{i \mid a_i \leq N\},$$

then relation (3.11) clearly implies that the sequences  $\{i(c_i^{2m_i-1})\}$  and  $2i$  have the same density  $\delta$ , namely  $\frac{1}{2}$ . Now consider the sequence  $k \rightarrow i(c^{2m-1})$  for  $c \in \mathcal{V}(\mathbb{R}P^n, F)$ , they have the same density as the sequences  $(2m-1)\hat{i}(c)$ , namely  $\delta(c) = \frac{1}{2\hat{i}(c)}$ . Since  $\kappa_j \neq \kappa_k$  for  $j \neq k$ , then each point in the sequence  $i(c_i^{2m_i-1})$  belongs to one of the sequences  $i(c^{2m-1})$  and no point in the sequences  $i(c^{2m-1})$  can be used twice. So its density must be no more than the sum of their densities, that is

$$\sum_{c \in \mathcal{V}(\mathbb{R}P^n, F)} \frac{1}{2\hat{i}(c)} = \sum_{c \in \mathcal{V}(\mathbb{R}P^n, F)} \delta(c) \geq \delta = \frac{1}{2},$$

which is precisely (3.10). ■

## 4 Multiplicity of non-contractible closed geodesics on $(\mathbb{R}P^n, F)$

In this section, we give proofs of the multiplicity results of non-contractible closed geodesics on  $(\mathbb{R}P^n, F)$  stated in Section 1 by using Fadell-Rabinowitz theory for non-contractible closed geodesics obtained in Section 3, the  $S^1$ -equivariant Poincaré series of the non-contractible component of the free loop space on  $\mathbb{R}P^n$  and some estimations on the index and mean index of non-contractible closed geodesics.

The following lemmas proved by Rademacher are useful for us to control the index of non-contractible closed geodesics.

**Lemma 4.1.** (cf. Lemma 1 of [Rad4]) *Let  $c$  be a closed geodesic on a Finsler manifold  $(M, F)$  of dimension  $n$  with positive flag curvature  $K \geq \delta$  for some  $\delta > 0$ , if the length  $L(c)$  satisfies  $L(c) > \frac{k\pi}{\sqrt{\delta}}$  for some positive integer  $k$  then  $i(c) \geq k(n-1)$ .*

**Lemma 4.2.** (cf. Theorem 1 and Theorem 4 of [Rad3]) *Let  $M$  be a compact and simply-connected manifold with a Finsler metric  $F$  with reversibility  $\lambda$  and flag curvature  $K$  satisfying  $0 < K \leq 1$  resp.  $\frac{\lambda^2}{(\lambda+1)^2} < K \leq 1$  if the dimension  $n$  is odd. Then the length of a closed geodesic is bounded from below by  $\pi \frac{\lambda+1}{\lambda}$ .*

Now we can estimate the index of non-contractible closed geodesics.

**Proposition 4.3.** *Let  $c$  be a non-contractible closed geodesic on Finsler  $n$ -dimensional real projective space  $(\mathbb{R}P^n, F)$  with reversibility  $\lambda$  and flag curvature  $K$  satisfying  $\frac{16}{9} \left( \frac{\lambda}{1+\lambda} \right)^2 < K \leq 1$  with  $\lambda < 3$ , then  $i(c^m) \geq 2(n-1)$  for  $m \geq 3$ .*

**Proof.** Since  $(S^n, F)$  is the universal double covering of  $(\mathbb{R}P^n, F)$ , then  $c^2$  is a closed geodesic on  $(S^n, F)$ . By Lemma 4.2, for  $m \geq 3$  we have

$$L(c^m) = \frac{m}{2} L(c^2) \geq \frac{m\pi}{2} \frac{\lambda+1}{\lambda} \geq \frac{3\pi}{2} \frac{\lambda+1}{\lambda} = 2 \frac{3\pi(\lambda+1)}{4\lambda}. \quad (4.1)$$

By assumption, we can choose an  $\delta > \frac{16}{9} \left( \frac{\lambda}{1+\lambda} \right)^2$  such that  $K \geq \delta$ , it together with (4.1) gives  $L(c^m) > \frac{2\pi}{\sqrt{\delta}}$  for  $m \geq 3$  which implies  $i(c^m) \geq 2(n-1)$  by Lemma 4.1. ■

**Proposition 4.4.** *Let  $c$  be a non-contractible closed geodesic on Finsler  $n$ -dimensional real projective space  $(\mathbb{R}P^n, F)$  with reversibility  $\lambda$  and flag curvature  $K$  satisfying  $\frac{64}{25} \left( \frac{\lambda}{1+\lambda} \right)^2 < K \leq 1$  with  $\lambda < \frac{5}{3}$ , then  $i(c^m) \geq 4(n-1)$  for  $m \geq 5$ .*

**Proof.** Since  $(S^n, F)$  is the universal double covering of  $(\mathbb{R}P^n, F)$ , then  $c^2$  is a closed geodesic on  $(S^n, F)$ . By Lemma 4.2, for  $m \geq 5$  we have

$$L(c^m) = \frac{m}{2} L(c^2) \geq \frac{m\pi}{2} \frac{\lambda+1}{\lambda} \geq \frac{5\pi}{2} \frac{\lambda+1}{\lambda} = 4 \frac{5\pi(\lambda+1)}{8\lambda}. \quad (4.2)$$

By assumption, we can choose an  $\delta > \frac{64}{25} \left( \frac{\lambda}{1+\lambda} \right)^2$  such that  $K \geq \delta$ , it together with (4.2) gives  $L(c^m) > \frac{4\pi}{\sqrt{\delta}}$  for  $m \geq 5$  which implies  $i(c^m) \geq 4(n-1)$  by Lemma 4.1. ■

**Proof of Theorem 1.1.** We assume that there exist finitely many prime non-contractible closed geodesics. Then by Theorems 3.5-3.6, there exist non-contractible closed geodesics  $c_i$  such that

$$E(c_i) = \kappa_i, \quad \overline{C}_{2i-2}(E, c_i) \neq 0, \quad (4.3)$$

and  $\kappa_i \neq \kappa_j$  for positive integers  $i \neq j$ . Combining Proposition 2.1 with (4.3), we obtain  $i(c_i) \leq 2i - 2 \leq i(c_i) + \nu(c_i)$  for any  $i \in \mathbf{N}$ . Then by Proposition 4.3, note that even iterations of  $c_i$  do not belong to the non-contractible component  $\Lambda_g$  of the free loop space, then for  $1 \leq i \leq n - 1$ ,  $c_i$  is prime. Thus we get  $n - 1$  non-contractible prime closed geodesics  $\{c_i \mid 1 \leq i \leq n - 1\}$ . ■

**Proof of Theorem 1.2.** Denote all the prime non-contractible closed geodesics by  $\{c_i \mid 1 \leq i \leq k\}$ . Then by Proposition 4.4, we have  $i(c_i^m) \geq 4(n - 1)$  for  $m \geq 5$  which implies that if  $c_i^m$  has contribution to  $m_q$  for  $0 \leq q \leq 4(n - 1) - 2$ , then  $m = 1$  or  $3$ . Thus we obtain

$$m_{4n-6} + m_{4n-8} + \cdots + m_2 + m_0 \leq 2k. \quad (4.4)$$

In the following, we prove in two cases:

(i)  $n$  is odd. In this case, by (2.6) of Theorem 2.3 and (2.4) of Lemma 2.2, we have

$$m_{4n-6} + m_{4n-8} + \cdots + m_2 + m_0 \geq b_{4n-6} + b_{4n-8} + \cdots + b_2 + b_0 = 2n + 1. \quad (4.5)$$

Then  $k \geq n + 1$  follows from (4.4) and (4.5).

(ii)  $n$  is even. In this case, by (2.6) of Theorem 2.3 and (2.5) of Lemma 2.2, we have

$$m_{4n-6} + m_{4n-8} + \cdots + m_2 + m_0 \geq b_{4n-6} + b_{4n-8} + \cdots + b_2 + b_0 = 2n - 1. \quad (4.6)$$

Then  $k \geq n$  follows from (4.4) and (4.6). The proof is complete. ■

The following lemma helps us to control the mean index of a closed geodesic:

**Lemma 4.5.** (cf. Lemma 2 of [Rad4]) *Let  $c$  be a closed geodesic on a compact and simply-connected Riemannian manifold of dimension  $n$  with a non-reversible Finsler metric with reversibility  $\lambda$  and flag curvature  $0 < \delta \leq K \leq 1$  where  $\delta > \frac{\lambda^2}{(\lambda+1)^2}$  if  $n$  is odd. Then  $\hat{i}(c) \geq \sqrt{\delta} \frac{\lambda+1}{\lambda} (n-1)$ .*

**Proof of Theorem 1.3.** Denote all the prime non-contractible closed geodesics by  $\{c_i \mid 1 \leq i \leq k\}$ . Since  $(S^n, F)$  is the universal double covering of  $(\mathbb{R}P^n, F)$ , then  $c_i^2$  is a closed geodesic on  $(S^n, F)$ . By Lemma 4.5, we have

$$\hat{i}(c_i^2) > \sqrt{\frac{4m^2}{(n-1)^2}} \left( \frac{\lambda}{1+\lambda} \right)^2 \frac{\lambda+1}{\lambda} (n-1) = 2m,$$

which implies

$$\hat{i}(c_i) > m. \quad (4.7)$$

On the other hand, by Theorem 3.8, we have

$$\sum_{1 \leq i \leq k} \frac{1}{\hat{i}(c)} \geq 1,$$

which together with (4.7) gives  $k \geq m + 1$ . ■

**Acknowledgements.** I would like to sincerely thank my advisor, Professor Yiming Long, for introducing me to the theory of closed geodesics and for his valuable advices and comments on this paper.

## References

- [Ano] D. V. Anosov, Geodesics in Finsler geometry. *Proc. I.C.M.* (Vancouver, B.C. 1974), Vol. 2. 293-297 Montreal (1975) (Russian), *Amer. Math. Soc. Transl.* 109 (1977) 81-85.
- [BTZ] W. Ballmann, G. Thorbergsson, W. Ziller, Closed geodesics and the fundamental group. *Duke Math. J.* 48 (1981), 585-588.
- [Ban] V. Bangert, On the existence of closed geodesics on two-spheres. *Internat. J. Math.* 4 (1993), no. 1, 1-10.
- [BaH] V. Bangert, N. Hingston, Closed geodesics on manifolds with infinite abelian fundamental group. *J. Differ. Geom.* 19 (1984), 277-282.
- [BaL] V. Bangert, Y. Long, The existence of two closed geodesics on every Finsler  $n$ -sphere. *Math. Ann.* 346 (2010), no. 2, 335-366.
- [Bot] Bott, R., On the iteration of closed geodesics and the Sturm intersection theory. *Comm. Pure Appl. Math.* 9 (1956) 171-206.
- [Cha] K. C. Chang, Infinite Dimensional Morse Theory and Multiple Solution Problems. Birkhäuser. Boston. 1993.
- [DuL1] H. Duan and Y. Long, Multiple closed geodesics on bumpy Finsler  $n$ -spheres. *J. Diff. Equa.* 233 (2007), no. 1, 221-240.
- [DuL2] H. Duan and Y. Long, The index growth and multiplicity of closed geodesics. *J. Funct. Anal.* 259 (2010), no. 7, 1850-1913.
- [DLX] H. Duan, Y. Long and Y. Xiao, Two closed geodesics on  $\mathbb{R}P^{2n+1}$  with a bumpy Finsler metric. *Calc. Var. and PDEs.* 54 (2015), 2883-2894.
- [DLW1] H. Duan, Y. Long and W. Wang, Two closed geodesics on compact simply-connected bumpy Finsler manifolds. *J. Differ. Geom.*, to appear.
- [DLW2] H. Duan, Y. Long and W. Wang, The enhanced common index jump theorem for symplectic paths and non-hyperbolic closed geodesics on Finsler manifolds. arXiv:1510.02872. Submitted.

- [EkH] I. Ekeland and H. Hofer, Convex Hamiltonian energy surfaces and their closed trajectories, *Comm. Math. Phys.* 113 (1987), 419-467.
- [FaR] E. Fadell and P. Rabinowitz, Generalized cohomological index theories for Lie group actions with an application to bifurcation questions for Hamiltonian systems. *Invent. Math.* 45 (1978), no. 2, 139–174.
- [Fra] J. Franks, Geodesics on  $S^2$  and periodic points of annulus homeomorphisms. *Invent. Math.* 108 (1992), no. 2, 403-418.
- [GM1] D. Gromoll and W. Meyer, On differentiable functions with isolated critical points. *Topology.* 8 (1969) 361-369.
- [GM2] D. Gromoll, W. Meyer, Periodic geodesics on compact Riemannian manifolds. *J. Differ. Geom.* 3 (1969), 493-510.
- [HiR] N. Hingston and H.-B. Rademacher, Resonance for loop homology of spheres. *J. Differ. Geom.* 93 (2013), 133-174.
- [Kat] A. B. Katok, Ergodic properties of degenerate integrable Hamiltonian systems. *Izv. Akad. Nauk SSSR.* 37 (1973) (Russian), *Math. USSR-Izv.* 7 (1973) 535-571.
- [Kli] W. Klingenberg, Lectures on Closed Geodesics. Springer. Berlin. 1978.
- [Liu] C. Liu, The relation of the Morse index of closed geodesics with the Maslov-type index of symplectic paths. *Acta Math. Sinica.* English Series 21 (2005) 237-248.
- [LiL] C. Liu and Y. Long, Iterated index formulae for closed geodesics with applications. *Science in China.* 45 (2002) 9-28.
- [Lon] Y. Long, Index Theory for Symplectic Paths with Applications. Progress in Math. 207, Birkhäuser. Basel. 2002.
- [LoD] Y. Long and H. Duan, Multiple closed geodesics on 3-spheres. *Adv. Math.* 221 (2009), no. 6, 1757-1803.
- [LoZ] Y. Long and C. Zhu, Closed characteristics on compact convex hypersurfaces in  $\mathbf{R}^{2n}$ . *Ann. of Math.* 155 (2002) 317-368.
- [LyF] L. A. Lyusternik and A. I. Fet, Variational problems on closed manifolds. *Dokl. Akad. Nauk SSSR (N.S.)* 81 (1951) 17-18 (in Russian).
- [Rad1] H.-B. Rademacher, Morse Theorie und geschlossene Geodatische. *Bonner Math. Schriften* Nr. 229 (1992).
- [Rad2] H.-B. Rademacher, The Fadell-Rabinowitz index and closed geodesics. *J. London. Math. Soc.* 50 (1994) 609-624.
- [Rad3] H.-B. Rademacher, A Sphere Theorem for non-reversible Finsler metrics. *Math. Annalen.* 328 (2004) 373-387.

- [Rad4] H.-B. Rademacher, Existence of closed geodesics on positively curved Finsler manifolds. *Ergodic Theory Dynam. Systems.* 27 (2007), no. 3, 957–969.
- [Rad5] H.-B. Rademacher, The second closed geodesic on the complex projective plane. *Front. Math. China.* 3 (2008), 253-258.
- [Rad6] H.-B. Rademacher, The second closed geodesic on Finsler spheres of dimension  $n > 2$ . *Trans. Amer. Math. Soc.* 362 (2010), no. 3, 1413-1421.
- [She] Z. Shen, Lectures on Finsler Geometry. World Scientific. Singapore. 2001.
- [Spa1] E. H. Spanier, Algebraic Topology. McGraw-Hill Book Comp. New York. 1966.
- [Tai1] I.A. Taimanov, The type numbers of closed geodesics (2010). arXiv:0912.5226.
- [Tai2] I.A. Taimanov, The spaces of non-contractible closed curves in compact space forms (2016). arXiv:1604.05237v1.
- [ViS] M. Vigué-Poirrier, D. Sullivan, The homology theory of the closed geodesic problem. *J. Differ. Geom.* 11 (1976), 633-644.
- [Wan1] W. Wang, Closed geodesics on positively curved Finsler spheres. *Adv. Math.* 218 (2008), 1566-1603.
- [Wan2] W. Wang, On a conjecture of Anosov. *Adv. Math.* 230 (2012), 1597-1617.
- [Wan3] W. Wang, On the average indices of closed geodesics on positively curved Finsler spheres. *Math. Ann.* 355 (2013), 1049-1065.
- [XL] Y. Xiao, Y. Long, Topological structure of non-contractible loop space and closed geodesics on real projective spaces with odd dimensions. *Adv. Math.* 279 (2015), 159-200.
- [Zil] W. Ziller, Geometry of the Katok examples. *Ergod. Th. Dyn. Sys.* 3 (1982) 135-157.