

MEAN CURVATURE FLOW OF ARBITRARY CODIMENSION IN COMPLEX PROJECTIVE SPACES

LI LEI AND HONGWEI XU

ABSTRACT. In this paper, we investigate the mean curvature flow of submanifolds of arbitrary codimension in $\mathbb{C}\mathbb{P}^m$. We prove that if the initial submanifold satisfies a pinching condition, then the mean curvature flow converges to a round point in finite time, or converges to a totally geodesic submanifold as $t \rightarrow \infty$. Consequently, we obtain a new differentiable sphere theorem for submanifolds in $\mathbb{C}\mathbb{P}^m$. Our work improves the convergence theorem for mean curvature flow due to Pipoli and Sinestrari [27].

1. INTRODUCTION

Let $F_0 : M^n \rightarrow N^{n+q}$ be an n -dimensional submanifold in the $(n+q)$ -dimensional Riemannian manifold N . The mean curvature flow with initial value F_0 is a smooth family of immersions $F : M \times [0, T) \rightarrow N^{n+q}$ satisfying

$$(1.1) \quad \begin{cases} \frac{\partial}{\partial t} F(x, t) = H(x, t), \\ F(\cdot, 0) = F_0, \end{cases}$$

where $H(x, t)$ is the mean curvature vector of the submanifold $M_t = F_t(M)$, $F_t = F(\cdot, t)$.

In 1984, Huisken [14] proved that uniformly convex hypersurfaces in Euclidean space will converge to a round point along the mean curvature flow. Afterwards, Huisken obtained convergence results for mean curvature flow of convex hypersurfaces in Riemannian manifolds [15] and pinched hypersurfaces in spheres [16]. In [1], Andrews constructed a fully nonlinear parabolic flow of surfaces in the three-sphere, and proved an optimal convergence result for this flow.

For higher codimensional submanifolds, Andrews and Baker [2] proved an optimal convergence theorem for the mean curvature flow and an optimal differentiable sphere theorem for submanifolds in \mathbb{R}^{n+q} . Meanwhile, by using the Ricci flow [4, 5, 6, 12], Xu and Gu [37] proved a general differentiable sphere theorem for submanifolds in space forms independently. Afterwards, Baker [3] and Liu-Xu-Ye-Zhao [22] proved a sharp convergence theorem for the mean curvature flow of submanifolds in the space form $\mathbb{F}^{n+q}(c)$ with $c \neq 0$. Later, Liu, Xu and Zhao [23] obtained a convergence result for mean curvature flow of arbitrary codimension in Riemannian manifolds. Recently, inspired by the rigidity theory of submanifolds [30, 34, 35, 36, 40], and by developing new techniques, Lei

2010 *Mathematics Subject Classification.* 53C44; 53C40; 53C20; 58J35.

Key words and phrases. Mean curvature flow, submanifolds of arbitrary codimension, complex projective space, convergence theorem, differentiable sphere theorem.

Research supported by the National Natural Science Foundation of China, Grant Nos. 11531012, 11371315.

and Xu [17, 18, 19] verified an optimal convergence theorem for the mean curvature flow of submanifolds in hyperbolic spaces and a new convergence theorem for the mean curvature flow of submanifolds in spheres, which improve the convergence theorems due to Baker [3], Huisken [16] and Liu-Xu-Ye-Zhao [22]. For more results on rigidity, sphere and convergence theorems, we refer the readers to [8, 9, 10, 11, 20, 21, 24, 25, 26, 29, 31, 32, 33, 38, 39, 40, 41].

More recently, Pipoli and Sinestrari [27] obtain the following convergence theorem for mean curvature flow of small codimension in the complex projective space.

Theorem A. *Let $F_0 : M^n \rightarrow \mathbb{C}\mathbb{P}^{\frac{n+q}{2}}$ be a closed submanifold of dimension n and codimension q in the complex projective space with Fubini-Study metric. Suppose either $n \geq 5$ and $q = 1$, or and $2 \leq q < \frac{n-3}{4}$. Let $F : M^n \times [0, T) \rightarrow \mathbb{C}\mathbb{P}^{\frac{n+q}{2}}$ be the mean curvature flow with initial value F_0 . If F_0 satisfies*

$$|h|^2 < \begin{cases} \frac{1}{n-1}|H|^2 + 2, & q = 1, \\ \frac{1}{n-1}|H|^2 + \frac{n-3-4q}{n}, & q \geq 2, \end{cases}$$

then F_t converges to a round point in finite time, or converges to a totally geodesic submanifold as $t \rightarrow \infty$. In particular, M is diffeomorphic to either \mathbb{S}^n or $\mathbb{C}\mathbb{P}^{n/2}$.

In this paper, we investigate the mean curvature flow of arbitrary codimensional submanifolds in the complex projective space, and prove the following theorem.

Theorem 1.1. *Let $F_0 : M^n \rightarrow \mathbb{C}\mathbb{P}^{\frac{n+q}{2}}$ be an n -dimensional closed submanifold in $\mathbb{C}\mathbb{P}^{\frac{n+q}{2}}$. Suppose that the dimension and codimension satisfy either (i) $q = 1$ and $n \geq 3$, (ii) $2 \leq q < n - 4$, or (iii) $q \geq n - 4 \geq 2$. Let $F : M^n \times [0, T) \rightarrow \mathbb{C}\mathbb{P}^{\frac{n+q}{2}}$ be the mean curvature flow with initial value F_0 . If F_0 satisfies*

$$|h|^2 < \begin{cases} \varphi(|H|^2), & q = 1, \\ \frac{1}{n-1}|H|^2 + 2 - \frac{3}{n}, & 2 \leq q < n - 4, \\ \psi(|H|^2), & q \geq n - 4, \end{cases}$$

then F_t converges to a round point in finite time, or converges to a totally geodesic submanifold as $t \rightarrow \infty$. In particular, M is diffeomorphic to \mathbb{S}^n or $\mathbb{C}\mathbb{P}^{n/2}$.

Here the functions φ and ψ are defined as (3.2) and (3.13), and $\varphi(|H|^2)$ and $\psi(|H|^2)$ are given by

$$\varphi(|H|^2) = 2 + a_n + \left(b_n + \frac{1}{n-1}\right) |H|^2 - \sqrt{b_n^2 |H|^4 + 2a_n b_n |H|^2},$$

$$\psi(|H|^2) = \frac{9}{n^2 - 3n - 3} + \frac{n^2 - 3n}{n^3 - 4n^2 + 3} |H|^2 - \frac{3\sqrt{|H|^4 + \frac{2}{n}(n-1)(n^2-3)|H|^2 + 9(n-1)^2}}{n^3 - 4n^2 + 3},$$

where $a_n = 2\sqrt{(n^2 - 4n + 3)b_n}$, $b_n = \min \left\{ \frac{n-3}{4n-4}, \frac{2n-5}{n^2+n-2} \right\}$.

Remark 1. *By a computation, if $n > 3$, we have $\varphi(x) > \frac{x}{n-1} + 2$ for $x \geq 0$. Furthermore, if $n \geq 3$, we have $\varphi(x) > \sqrt{2(n-3)}$ for $x \geq 0$. Therefore, Theorem 1.1 substantially improves Theorem A.*

The function ψ satisfies $\psi(x) \geq \frac{x}{n}$ for $x \geq 0$, and the equality holds if and only if $x = 0$. In addition, we have $\lim_{x \rightarrow +\infty} \psi(x)/x = \frac{1}{n-1}$.

Since $\psi(0) = 0$, the pinching condition $|h|^2 < \psi(|H|^2)$ implies that the submanifold has nonzero mean curvature. Thus, if $q \geq n - 4$, it doesn't occur that the mean curvature flow in Theorem 1.1 converges to a totally geodesic submanifold. When $q \geq n - 4 \geq 2$, we obtain a refined result under the weakly pinching condition.

Theorem 1.2. *Let $F_0 : M^n \rightarrow \mathbb{C}\mathbb{P}^{\frac{n+q}{2}}$ be a closed submanifold of dimension n and codimension q in $\mathbb{C}\mathbb{P}^{\frac{n+q}{2}}$, where $q \geq n - 4 \geq 2$. Let $F : M^n \times [0, T) \rightarrow \mathbb{C}\mathbb{P}^{\frac{n+q}{2}}$ be the mean curvature flow with initial value F_0 . If F_0 satisfies*

$$|h|^2 \leq \psi(|H|^2),$$

then one of the following holds:

- (i) F_t is congruent to the totally geodesic $\mathbb{R}\mathbb{P}^n$ or $\mathbb{C}\mathbb{P}^{n/2}$ for each $t \in [0, +\infty)$;
- (ii) F_t converges to a round point in finite time. In particular, M is diffeomorphic to \mathbb{S}^n .

2. NOTATIONS AND FORMULAS

Let $\mathbb{C}\mathbb{P}^m$ be the m -dimensional complex projective space with the Fubini-Study metric g_{FS} . Let J be its complex structure. We denote by ∇ the Levi-Civita connection of $(\mathbb{C}\mathbb{P}^m, g_{\text{FS}})$. Since the Fubini-Study metric is a Kähler metric, we have $\nabla J = 0$. The curvature tensor \bar{R} of $\mathbb{C}\mathbb{P}^m$ can be written as

$$(2.1) \quad \begin{aligned} \bar{R}(X, Y, Z, W) &= \langle X, Z \rangle \langle Y, W \rangle - \langle X, W \rangle \langle Y, Z \rangle \\ &+ \langle X, JZ \rangle \langle Y, JW \rangle - \langle X, JW \rangle \langle Y, JZ \rangle \\ &+ 2\langle X, JY \rangle \langle Z, JW \rangle. \end{aligned}$$

Let (M^n, g) be a real n -dimensional Riemannian submanifold immersed in $(\mathbb{C}\mathbb{P}^m, g_{\text{FS}})$. Let q be its codimension, i.e., $n + q = 2m$. At a point $p \in M$, let T_pM and N_pM be the tangent space and normal space, respectively. For a vector in $T_pM \oplus N_pM$, we denote by $(\cdot)^T$ and $(\cdot)^N$ its projections onto T_pM and N_pM , respectively. We use the same symbol ∇ to represent the connections of tangent bundle TM and normal bundle NM . Denote by $\Gamma(E)$ the spaces of smooth sections of a vector bundles E . For $X, Y \in \Gamma(TM)$, $\xi \in \Gamma(NM)$, the connections ∇ are given by $\nabla_X Y = (\bar{\nabla}_X Y)^T$ and $\nabla_X \xi = (\bar{\nabla}_X \xi)^N$. The second fundamental form of M is defined as $h(X, Y) = (\bar{\nabla}_X Y)^N$.

Throughout this paper, we shall make the following convention on indices:

$$1 \leq A, B, C, \dots \leq n + q, \quad 1 \leq i, j, k, \dots \leq n, \quad n + 1 \leq \alpha, \beta, \gamma, \dots \leq n + q.$$

We choose a local orthonormal frame $\{e_i\}$ for the tangent bundle and a local orthonormal frame $\{e_\alpha\}$ for the normal bundle. With the local frame, the components of h are given by $h_{ij}^\alpha = \langle h(e_i, e_j), e_\alpha \rangle$. The mean curvature vector is defined as $H = \sum_\alpha H^\alpha e_\alpha$, where $H^\alpha = \sum_i h_{ii}^\alpha$. Let $\mathring{h} = h - \frac{1}{n}H \otimes g$ be the traceless second fundamental form. We have the relations $|\mathring{h}|^2 = |h|^2 - \frac{1}{n}|H|^2$ and $|\nabla \mathring{h}|^2 = |\nabla h|^2 - \frac{1}{n}|\nabla H|^2$.

We denote by (J_{AB}) the matrix of J with respect to the frame $\{e_A\}$, i.e.,

$$J_{AB} = \langle e_A, J e_B \rangle.$$

This matrix satisfies $J_{AB} = -J_{BA}$ and $\sum_B J_{AB} J_{BC} = -\delta_{AC}$.

At each point $p \in M$, we define a tensor $P : N_pM \rightarrow T_pM$ by

$$P\xi = (J\xi)^T \quad \text{for } \xi \in N_pM.$$

Then we have

$$|P|^2 = \sum_\alpha |P e_\alpha|^2 \leq \sum_\alpha |e_\alpha|^2 = q,$$

and

$$|P|^2 = \sum_{\alpha,i} (J_{i\alpha})^2 = \sum_{A,i} (J_{iA})^2 - \sum_{i,j} (J_{ij})^2 = n - \sum_{i,j} (J_{ij})^2.$$

We have the following estimates for the gradient of the second fundamental form.

Lemma 2.1. *For an n -dimensional submanifold in $\mathbb{C}\mathbb{P}^{\frac{n+q}{2}}$, we have*

$$|\nabla h|^2 \geq \begin{cases} \frac{3}{n+2} |\nabla H|^2 + 2(n-1), & q = 1, \\ \frac{3}{n+8} |\nabla H|^2 + 2(n-q)|P|^2, & 2 \leq q < n, \\ \frac{3}{n+8} |\nabla H|^2, & q \geq n. \end{cases}$$

Proof. Let S be the symmetric part of ∇h , i.e., $S_{ijk}^\alpha = \frac{1}{3}(\nabla_i h_{jk}^\alpha + \nabla_j h_{ik}^\alpha + \nabla_k h_{ij}^\alpha)$. Using the same argument as in the proof of Lemma 2.2 in [14], we have

$$(2.2) \quad |S|^2 \geq \frac{3}{n+2} \sum_{\alpha,i} \left(\sum_k S_{ikk}^\alpha \right)^2.$$

By the Codazzi equation, we have $\sum_k S_{ikk}^\alpha = \nabla_i H^\alpha + 2 \sum_k J_{\alpha k} J_{ki}$. Then we obtain

$$(2.3) \quad \sum_i \left(\sum_k S_{ikk} \right)^2 = |\nabla H|^2 + 4 \sum_{\alpha,k} \nabla_i H^\alpha J_{\alpha k} J_{ki} + 4 \sum_{\alpha,i} \left(\sum_k J_{\alpha k} J_{ki} \right)^2.$$

Using the Codazzi equation again, we get

$$(2.4) \quad \begin{aligned} |S|^2 &= \frac{1}{3} |\nabla h|^2 + \frac{2}{3} \sum_{\alpha,i,j,k} \nabla_k h_{ij}^\alpha \nabla_j h_{ik}^\alpha \\ &= \frac{1}{3} |\nabla h|^2 + \frac{2}{3} \sum_{\alpha,i,j,k} \nabla_k h_{ij}^\alpha (\nabla_k h_{ij}^\alpha + \bar{R}_{\alpha ijk}) \\ &= |\nabla h|^2 + \frac{2}{3} \sum_{\alpha,i,j,k} \nabla_k h_{ij}^\alpha \bar{R}_{\alpha ijk} \\ &= |\nabla h|^2 + \frac{2}{3} \sum_{\alpha,i,j,k} (\nabla_i h_{jk}^\alpha + \bar{R}_{\alpha jki}) \bar{R}_{\alpha ijk} \\ &= |\nabla h|^2 + \frac{2}{3} \sum_{\alpha,i,j,k} \bar{R}_{\alpha jki} \bar{R}_{\alpha ijk} \\ &= |\nabla h|^2 - 2 \sum_{\alpha,i,j,k} [J_{\alpha j} J_{ji} J_{\alpha k} J_{ki} + (J_{\alpha i})^2 (J_{jk})^2] \\ &= |\nabla h|^2 - 2 \sum_{\alpha,i} \left(\sum_k J_{\alpha k} J_{ki} \right)^2 - 2|P|^2(n - |P|^2). \end{aligned}$$

From (2.2), (2.3) and (2.4), we obtain

$$(2.5) \quad \begin{aligned} |\nabla h|^2 &\geq \frac{1}{n+2} \left[3|\nabla H|^2 + 12 \sum_{\alpha,i,k} \nabla_i H^\alpha J_{\alpha k} J_{ki} + 2(n+8) \sum_{\alpha,i} \left(\sum_k J_{\alpha k} J_{ki} \right)^2 \right] \\ &\quad + 2|P|^2(n - |P|^2). \end{aligned}$$

If $q = 1$, we have $\sum_k J_{\alpha k} J_{ki} = 0$ and $|P|^2 = 1$. Thus (2.5) becomes

$$|\nabla h|^2 \geq \frac{3}{n+2} |\nabla H|^2 + 2(n-1).$$

If $q \geq 2$, from (2.5) we get

$$\begin{aligned} |\nabla h|^2 &\geq \frac{2}{n+2} \sum_{\alpha,i} \left(3\sqrt{\frac{1}{n+8}} \nabla_i H^\alpha + \sqrt{n+8} \sum_k J_{\alpha k} J_{ki} \right)^2 \\ &\quad + \frac{3}{n+8} |\nabla H|^2 + 2|P|^2(n - |P|^2). \end{aligned}$$

□

Let $F : M^n \times [0, T) \rightarrow \mathbb{C}\mathbb{P}^m$ be a mean curvature flow in a complex projective space. For a fixed t , letting $F_t = F(\cdot, t)$, then $F_t : M^n \rightarrow \mathbb{C}\mathbb{P}^m$ is a Riemannian submanifold in $\mathbb{C}\mathbb{P}^m$. We denote by M_t the submanifold at time t . Following [2, 27], we have the evolution equations below.

$$\begin{aligned} \frac{\partial}{\partial t} |h|^2 &= \Delta |h|^2 - 2|\nabla h|^2 \\ &\quad + 2 \sum_{\alpha,\beta} \left(\sum_{i,j} h_{ij}^\alpha h_{ij}^\beta \right)^2 + 2 \sum_{i,j,\alpha,\beta} \left(\sum_k (h_{ik}^\alpha h_{jk}^\beta - h_{ik}^\beta h_{jk}^\alpha) \right)^2 \\ &\quad + 2 \sum_{\alpha,\beta,i,j,k} h_{ij}^\alpha h_{ij}^\beta \bar{R}_{\alpha k \beta k} + 8 \sum_{\alpha,\beta,i,j,k} \dot{h}_{ik}^\alpha \dot{h}_{jk}^\beta \bar{R}_{\alpha \beta ij} \\ &\quad + 4 \sum_{\alpha,i,j,k,l} (h_{ik}^\alpha h_{jl}^\alpha \bar{R}_{ijkl} - h_{ik}^\alpha h_{jk}^\alpha \bar{R}_{iljl}), \\ \frac{\partial}{\partial t} |H|^2 &= \Delta |H|^2 - 2|\nabla H|^2 + 2 \sum_{i,j} \left(\sum_\alpha H^\alpha h_{ij}^\alpha \right)^2 + 2 \sum_{\alpha,\beta,k} H^\alpha H^\beta \bar{R}_{\alpha k \beta k}. \end{aligned}$$

From (2.1), these evolution equations can be written as

Lemma 2.2. *For mean curvature flow $F : M^n \times [0, T) \rightarrow \mathbb{C}\mathbb{P}^m$, we have*

- (i) $\frac{\partial}{\partial t} |h|^2 = \Delta |h|^2 - 2|\nabla h|^2 - 2n|h|^2 + 4|H|^2 + 2R_1 + 2S_1$,
- (ii) $\frac{\partial}{\partial t} |H|^2 = \Delta |H|^2 - 2|\nabla H|^2 + 2n|H|^2 + 2R_2 + 6S_2$,

where

$$\begin{aligned} R_1 &= \sum_{\alpha,\beta} \left(\sum_{i,j} h_{ij}^\alpha h_{ij}^\beta \right)^2 + \sum_{i,j,\alpha,\beta} \left(\sum_k (h_{ik}^\alpha h_{jk}^\beta - h_{ik}^\beta h_{jk}^\alpha) \right)^2, \\ R_2 &= \sum_{i,j} \left(\sum_\alpha H^\alpha h_{ij}^\alpha \right)^2, \\ S_1 &= 3 \sum_{i,j,k} \left(\sum_\alpha h_{ij}^\alpha J_{k\alpha} \right)^2 + 4 \sum_{\alpha,\beta,i,j,k} \dot{h}_{ik}^\alpha \dot{h}_{jk}^\beta (J_{i\alpha} J_{j\beta} - J_{i\beta} J_{j\alpha}) \\ &\quad + 6 \sum_{\alpha,i,j,k,l} (\dot{h}_{ij}^\alpha \dot{h}_{kl}^\alpha J_{il} J_{jk} - \dot{h}_{ik}^\alpha \dot{h}_{jk}^\alpha J_{il} J_{jl}) + 8 \sum_{\alpha,\beta,i,j,k} \dot{h}_{ik}^\alpha \dot{h}_{jk}^\beta J_{\alpha\beta} J_{ij}, \\ S_2 &= \sum_k \left(\sum_\alpha H^\alpha J_{k\alpha} \right)^2. \end{aligned}$$

where $a = 2\sqrt{(n^2 - 4n + 3)b}$, $b = \min \left\{ \frac{n-3}{4n-4}, \frac{2n-5}{n^2+n-2} \right\}$, $c_\varepsilon = b + \frac{1}{n-1+\varepsilon}$, $d_\varepsilon = 2 - 2\varepsilon + a$, $e = \sqrt{\varepsilon}$. We define $\varphi = \varphi_0$.

Let $\hat{\varphi}_\varepsilon(x) = \varphi_\varepsilon(x) - \frac{x}{n}$. The following lemma will be proven in the Appendix.

Lemma 3.1. *For sufficiently small ε , the function $\hat{\varphi}_\varepsilon$ satisfies*

- (i) $2x\hat{\varphi}_\varepsilon''(x) + \hat{\varphi}_\varepsilon'(x) < \frac{2(n-1)}{n(n+2)}$,
- (ii) $\hat{\varphi}_\varepsilon(x)(\varphi_\varepsilon(x) - n + 3) - x\hat{\varphi}_\varepsilon'(x)(\varphi_\varepsilon(x) + n + 3) < 2(n-1)$,
- (iii) $\hat{\varphi}_\varepsilon(x) - x\hat{\varphi}_\varepsilon'(x) > 1$.

Suppose that M_0 is an n -dimensional closed hypersurface in $\mathbb{C}\mathbb{P}^{\frac{n+1}{2}}$ satisfying $|h|^2 < \varphi(|H|^2)$. Let $F : M^n \times [0, T] \rightarrow \mathbb{C}\mathbb{P}^{\frac{n+1}{2}}$ be a mean curvature flow with initial value M_0 . We will show that the pinching condition is preserved along the flow. For convenience, we denote $\hat{\varphi}_\varepsilon(|H|^2)$, $\hat{\varphi}_\varepsilon'(|H|^2)$, $\hat{\varphi}_\varepsilon''(|H|^2)$ by $\hat{\varphi}_\varepsilon$, $\hat{\varphi}_\varepsilon'$, $\hat{\varphi}_\varepsilon''$, respectively.

Theorem 3.2. *If the initial value M_0 satisfies $|h|^2 < \varphi(|H|^2)$, then there exists a small positive number ε , such that for all $t \in [0, T]$, we have $|h|^2 < \varphi_\varepsilon(|H|^2) - \varepsilon|H|^2 - \varepsilon$.*

Proof. Since M_0 is compact, there exists a small positive number ε , such that M_0 satisfies $|\dot{h}|^2 < \hat{\varphi}_\varepsilon$.

From Lemma 3.1 (i), we have

$$(3.3) \quad \begin{aligned} \left(\frac{\partial}{\partial t} - \Delta \right) \hat{\varphi}_\varepsilon &= -2(\hat{\varphi}_\varepsilon' + 2\hat{\varphi}_\varepsilon'' \cdot |H|^2) |\nabla H|^2 + 2\hat{\varphi}_\varepsilon' \cdot |H|^2 (|h|^2 + n + 3) \\ &\geq -\frac{4(n-1)}{n(n+2)} |\nabla H|^2 + 2\hat{\varphi}_\varepsilon' \cdot |H|^2 (|h|^2 + n + 3). \end{aligned}$$

Let $U = |\dot{h}|^2 - \hat{\varphi}_\varepsilon$. We obtain

$$\begin{aligned} \frac{1}{2} \left(\frac{\partial}{\partial t} - \Delta \right) U &\leq -|\nabla \dot{h}|^2 + \frac{2(n-1)}{n(n+2)} |\nabla H|^2 \\ &\quad + |\dot{h}|^2 (|h|^2 - n + 3) - \hat{\varphi}_\varepsilon' \cdot |H|^2 (|h|^2 + n + 3). \end{aligned}$$

By Lemma 2.1, we have

$$-|\nabla \dot{h}|^2 + \frac{2(n-1)}{n(n+2)} |\nabla H|^2 \leq -2(n-1).$$

Thus, at the points where $U = 0$, we get

$$\frac{1}{2} \left(\frac{\partial}{\partial t} - \Delta \right) U \leq -2(n-1) + \hat{\varphi}_\varepsilon(\varphi_\varepsilon - n + 3) - \hat{\varphi}_\varepsilon' \cdot |H|^2 (\varphi_\varepsilon + n + 3) < 0.$$

Applying the maximum principle, we obtain $U < 0$ for all $t \in [0, T]$.

By choosing a suitable small ε , we complete the proof of Theorem 3.2. \square

Let $f_\sigma = |\dot{h}|^2 / \hat{\varphi}_\varepsilon^{1-\sigma}$, where $\sigma \in (0, 1)$ is a positive constant. Then we have

Lemma 3.3. *If M_0 satisfies $|h|^2 < \varphi(|H|^2)$, then there exists a small positive number ε , such that the following inequality holds along the mean curvature flow.*

$$\frac{\partial}{\partial t} f_\sigma \leq \Delta f_\sigma + \frac{2}{\hat{\varphi}_\varepsilon} |\nabla f_\sigma| |\nabla \hat{\varphi}_\varepsilon| - \frac{2\varepsilon f_\sigma}{n|\dot{h}|^2} \left| \nabla \dot{h} \right|^2 + 2\sigma |h|^2 f_\sigma - \frac{\varepsilon}{n} f_\sigma.$$

Proof. By a straightforward calculation, we have

$$\begin{aligned} \left(\frac{\partial}{\partial t} - \Delta\right) f_\sigma &= f_\sigma \left[\frac{1}{|\dot{h}|^2} \left(\frac{\partial}{\partial t} - \Delta\right) |\dot{h}|^2 - \frac{1-\sigma}{\dot{\varphi}_\varepsilon} \left(\frac{\partial}{\partial t} - \Delta\right) \dot{\varphi}_\varepsilon \right] \\ &\quad + 2(1-\sigma) \frac{\langle \nabla f_\sigma, \nabla \dot{\varphi}_\varepsilon \rangle}{\dot{\varphi}_\varepsilon} - \sigma(1-\sigma) f_\sigma \frac{|\nabla \dot{\varphi}_\varepsilon|^2}{|\dot{\varphi}_\varepsilon|^2}. \end{aligned}$$

Using (3.1) and (3.3), we have

$$\begin{aligned} \left(\frac{\partial}{\partial t} - \Delta\right) f_\sigma &\leq 2f_\sigma \left[-\frac{|\nabla \dot{h}|^2}{|\dot{h}|^2} + \frac{2(n-1)}{n(n+2)} \frac{|\nabla H|^2}{\dot{\varphi}_\varepsilon} \right] \\ &\quad + 2f_\sigma \left[|h|^2 - n + 3 - (1-\sigma) \frac{\dot{\varphi}'_\varepsilon \cdot |H|^2}{\dot{\varphi}_\varepsilon} (|h|^2 + n + 3) \right] \\ &\quad + \frac{2}{\dot{\varphi}_\varepsilon} |\nabla f_\sigma| |\nabla \dot{\varphi}_\varepsilon|. \end{aligned}$$

From Lemma 2.1 and Theorem 3.2, we have

$$\begin{aligned} -\frac{|\nabla \dot{h}|^2}{|\dot{h}|^2} + \frac{2(n-1)}{n(n+2)} \frac{|\nabla H|^2}{\dot{\varphi}_\varepsilon} &\leq -\frac{|\nabla \dot{h}|^2}{|\dot{h}|^2} + \frac{|\nabla \dot{h}|^2 - 2(n-1)}{\dot{\varphi}_\varepsilon} \\ &\leq \frac{|\dot{h}|^2 - \dot{\varphi}_\varepsilon}{|\dot{h}|^2 \dot{\varphi}_\varepsilon} |\nabla \dot{h}|^2 - \frac{2(n-1)}{\dot{\varphi}_\varepsilon} \\ &\leq -\varepsilon \frac{|H|^2 + 1}{|\dot{h}|^2 \dot{\varphi}_\varepsilon} |\nabla \dot{h}|^2 - \frac{2(n-1)}{\dot{\varphi}_\varepsilon} \\ &\leq -\frac{\varepsilon}{n|\dot{h}|^2} |\nabla \dot{h}|^2 - \frac{2(n-1)}{\dot{\varphi}_\varepsilon}. \end{aligned}$$

From Lemma 3.1 (ii) and (iii), we have

$$\begin{aligned} &|h|^2 - n + 3 - (1-\sigma) \frac{\dot{\varphi}'_\varepsilon \cdot |H|^2}{\dot{\varphi}_\varepsilon} (|h|^2 + n + 3) \\ &= \frac{1-\sigma}{\dot{\varphi}_\varepsilon} [(\dot{\varphi}_\varepsilon - \dot{\varphi}'_\varepsilon \cdot |H|^2) |h|^2 - \dot{\varphi}'_\varepsilon \cdot |H|^2 (n+3)] - n + 3 + \sigma |h|^2 \\ &\leq \frac{1-\sigma}{\dot{\varphi}_\varepsilon} [(\dot{\varphi}_\varepsilon - \dot{\varphi}'_\varepsilon \cdot |H|^2) (\varphi_\varepsilon - \varepsilon |H|^2 - \varepsilon) - \dot{\varphi}'_\varepsilon \cdot |H|^2 (n+3)] - n + 3 + \sigma |h|^2 \\ &= \frac{1-\sigma}{\dot{\varphi}_\varepsilon} [(\dot{\varphi}_\varepsilon - \dot{\varphi}'_\varepsilon \cdot |H|^2) \varphi_\varepsilon - \dot{\varphi}'_\varepsilon \cdot |H|^2 (n+3)] - n + 3 + \sigma |h|^2 \\ &\quad - \frac{(1-\sigma)\varepsilon}{\dot{\varphi}_\varepsilon} (\dot{\varphi}_\varepsilon - \dot{\varphi}'_\varepsilon \cdot |H|^2) (|H|^2 + 1) \\ &\leq (1-\sigma) \left[n - 3 + \frac{2(n-1)}{\dot{\varphi}_\varepsilon} \right] - n + 3 + \sigma |h|^2 - \frac{(1-\sigma)\varepsilon}{\dot{\varphi}_\varepsilon} (|H|^2 + 1) \\ &\leq \sigma |h|^2 + \frac{2(n-1)}{\dot{\varphi}_\varepsilon} - \frac{\varepsilon}{2n}. \end{aligned}$$

This completes the proof of the lemma. \square

3.2. The case of $2 \leq q < n - 4$.

At a fixed point $p \in M$, we always choose the orthonormal frame $\{e_\alpha\}$ for $N_p M$ such that $H = |H|e_{n+1}$. For fixed α , let $|\mathring{h}^\alpha|^2 = \sum_{i,j} (\mathring{h}_{ij}^\alpha)^2$. Set $\rho_1 = |\mathring{h}^{n+1}|^2$, $\rho_2 = \sum_{\alpha > n+1} |\mathring{h}^\alpha|^2$, $\theta_1 = |Pe_{n+1}|^2$ and $\theta_2 = \sum_{\alpha > n+1} |Pe_\alpha|^2$. Notice that $\theta_1 \leq 1$ and $\theta_2 \leq q - 1$.

Lemma 3.4.

- (i) $R_1 \leq |h|^4 - \frac{2}{n}\rho_2|H|^2 + 2\rho_1\rho_2 + \frac{1}{2}\rho_2^2$,
- (ii) $R_2 = |H|^2(|h|^2 - \rho_2)$,
- (iii) $S_1 \leq \frac{3}{n}S_2 + 3|\mathring{h}|^2 + 8\sqrt{\theta_2\rho_1\rho_2} + 4\theta_2\rho_2$.

Proof. The estimates of R_1 and R_2 are similar to that in [2]. We choose an orthonormal frame $\{e_i\}$ for the tangent space, such that $\mathring{h}_{ij}^{n+1} = \mathring{\lambda}_i\delta_{ij}$. Then we have

$$\begin{aligned}
 R_1 &= \rho_1^2 + \frac{2}{n}\rho_1|H|^2 + \frac{1}{n^2}|H|^4 \\
 (3.4) \quad &+ 2 \sum_{\alpha > n+1} \left(\sum_i \mathring{\lambda}_i \mathring{h}_{ii}^\alpha \right)^2 + 2 \sum_{\substack{\alpha > n+1 \\ i \neq j}} \left((\mathring{\lambda}_i - \mathring{\lambda}_j) \mathring{h}_{ij}^\alpha \right)^2 \\
 &+ \sum_{\alpha, \beta > n+1} \left(\sum_{i,j} \mathring{h}_{ij}^\alpha \mathring{h}_{ij}^\beta \right)^2 + \sum_{\substack{\alpha, \beta > n+1 \\ i, j}} \left(\sum_k (\mathring{h}_{ik}^\alpha \mathring{h}_{jk}^\beta - \mathring{h}_{jk}^\alpha \mathring{h}_{ik}^\beta) \right)^2.
 \end{aligned}$$

Using the Cauchy-Schwarz inequality, we get

$$\begin{aligned}
 \left(\sum_i \mathring{\lambda}_i \mathring{h}_{ii}^\alpha \right)^2 + \sum_{i \neq j} \left((\mathring{\lambda}_i - \mathring{\lambda}_j) \mathring{h}_{ij}^\alpha \right)^2 &\leq \rho_1 \sum_i (\mathring{h}_{ii}^\alpha)^2 + 2 \sum_{i \neq j} (\mathring{\lambda}_i^2 + \mathring{\lambda}_j^2) (\mathring{h}_{ij}^\alpha)^2 \\
 &\leq \rho_1 \sum_i (\mathring{h}_{ii}^\alpha)^2 + 2\rho_1 \sum_{i \neq j} (\mathring{h}_{ij}^\alpha)^2 \\
 &\leq 2\rho_1 |\mathring{h}^\alpha|^2.
 \end{aligned}$$

It follows from Theorem 1 of [20] that

$$\sum_{\alpha, \beta > n+1} \left(\sum_{i,j} \mathring{h}_{ij}^\alpha \mathring{h}_{ij}^\beta \right)^2 + \sum_{\substack{\alpha, \beta > n+1 \\ i, j}} \left(\sum_k (\mathring{h}_{ik}^\alpha \mathring{h}_{jk}^\beta - \mathring{h}_{jk}^\alpha \mathring{h}_{ik}^\beta) \right)^2 \leq \frac{3}{2}\rho_2^2.$$

Thus we obtain

$$\begin{aligned}
 R_1 &\leq \rho_1^2 + \frac{2}{n}\rho_1|H|^2 + \frac{1}{n^2}|H|^4 + 4\rho_1\rho_2 + \frac{3}{2}\rho_2^2 \\
 &= |h|^4 - \frac{2}{n}\rho_2|H|^2 + 2\rho_1\rho_2 + \frac{1}{2}\rho_2^2.
 \end{aligned}$$

R_2 can be written as

$$(3.5) \quad R_2 = \sum_{i,j} (|H|\mathring{h}_{ij}^{n+1})^2 = |H|^2(|h|^2 - \rho_2).$$

Next, we have

$$\begin{aligned}
(3.6) \quad & \frac{1}{n} \sum_k \left(\sum_{\alpha} H^{\alpha} J_{k\alpha} \right)^2 + \sum_{i,j,k} \left(\sum_{\alpha} \mathring{h}_{ij}^{\alpha} J_{k\alpha} \right)^2 \\
&= \frac{S_2}{n} + \sum_{i,j} \left| P\mathring{h}(e_i, e_j) \right|^2 \\
&= \frac{S_2}{n} + \left| P\mathring{h} \right|^2.
\end{aligned}$$

Choose an orthonormal frame $\{e_i\}$ such that the matrix (J_{ij}) takes the form of (2.6). Thus

$$\begin{aligned}
(3.7) \quad & 6 \sum_{\alpha, i, j, k, l} (h_{ij}^{\alpha} h_{kl}^{\alpha} J_{il} J_{jk} - h_{ik}^{\alpha} h_{jl}^{\alpha} J_{il} J_{jk}) + 8 \sum_{\alpha, \beta, i, j, k} h_{ik}^{\alpha} h_{jk}^{\beta} J_{\alpha\beta} J_{ij} \\
&= 6 \sum_{i, k, \alpha} \left(-\mathring{h}_{ik}^{\alpha} \mathring{h}_{k\bar{i}}^{\alpha} J_{i\bar{i}} J_{k\bar{k}} - \left(\mathring{h}_{ik}^{\alpha} J_{i\bar{i}} \right)^2 \right) + 8 \sum_{i, k, \alpha, \beta} \mathring{h}_{ik}^{\alpha} \mathring{h}_{i\bar{k}}^{\beta} J_{\alpha\beta} J_{i\bar{i}} \\
&= -3 \sum_{i, k, \alpha} \left(\mathring{h}_{ik}^{\alpha} J_{k\bar{k}} + \mathring{h}_{ik}^{\alpha} J_{i\bar{i}} \right)^2 - 4 \sum_{i, k, \alpha} \left[\left(\mathring{h}_{ik}^{\alpha} J_{k\bar{k}} + \mathring{h}_{ik}^{\alpha} J_{i\bar{i}} \right) \sum_{\beta} \mathring{h}_{ik}^{\beta} J_{\alpha\beta} \right] \\
&\leq \frac{4}{3} \sum_{i, k, \alpha} \left(\sum_{\beta} \mathring{h}_{ik}^{\beta} J_{\alpha\beta} \right)^2 \\
&= \frac{4}{3} \sum_{i, k} \left| \left(J\mathring{h}(e_i, e_k) \right)^N \right|^2 \\
&= \frac{4}{3} \sum_{i, k} \left(\left| J\mathring{h}(e_i, e_k) \right|^2 - \left| P\mathring{h}(e_i, e_k) \right|^2 \right) \\
&= \frac{4}{3} \left(\left| \mathring{h} \right|^2 - \left| P\mathring{h} \right|^2 \right).
\end{aligned}$$

For fixed α, β , we choose an orthonormal frame $\{e_i\}$, such that the $n \times n$ matrix $(J_{i\alpha} J_{j\beta} - J_{j\alpha} J_{i\beta})$ takes the form of (2.6). Thus we get

$$\begin{aligned}
& \sum_{i, j, k} \mathring{h}_{ik}^{\alpha} \mathring{h}_{jk}^{\beta} (J_{i\alpha} J_{j\beta} - J_{j\alpha} J_{i\beta}) \\
&= \sum_{i, k} \mathring{h}_{ik}^{\alpha} \mathring{h}_{i\bar{k}}^{\beta} (J_{i\alpha} J_{i\bar{\beta}} - J_{i\bar{\alpha}} J_{i\beta}) \\
&\leq \sum_{i, k} \left| \mathring{h}_{ik}^{\alpha} \mathring{h}_{i\bar{k}}^{\beta} \right| \sqrt{(J_{i\alpha})^2 + (J_{i\bar{\alpha}})^2} \sqrt{(J_{i\beta})^2 + (J_{i\bar{\beta}})^2} \\
&\leq \sum_{i, k} \left| \mathring{h}_{ik}^{\alpha} \mathring{h}_{i\bar{k}}^{\beta} \right| |Pe_{\alpha}| |Pe_{\beta}| \\
&\leq \sqrt{\sum_{i, k} \left(\mathring{h}_{ik}^{\alpha} \right)^2} \sqrt{\sum_{i, k} \left(\mathring{h}_{i\bar{k}}^{\beta} \right)^2} |Pe_{\alpha}| |Pe_{\beta}| \\
&= \left| \mathring{h}^{\alpha} \right| \left| \mathring{h}^{\beta} \right| |Pe_{\alpha}| |Pe_{\beta}|.
\end{aligned}$$

Then

$$\begin{aligned}
& \sum_{\alpha, \beta, i, j, k} \dot{h}_{ik}^\alpha \dot{h}_{jk}^\beta (J_{i\alpha} J_{j\beta} - J_{j\alpha} J_{i\beta}) \\
& \leq \sum_{\alpha \neq \beta} |Pe_\alpha| |Pe_\beta| \left| \dot{h}^\alpha \right| \left| \dot{h}^\beta \right| \\
& = 2|Pe_{n+1}| \left| \dot{h}^{n+1} \right| \sum_{\alpha > n+1} |Pe_\alpha| \left| \dot{h}^\alpha \right| + \sum_{\substack{\alpha, \beta > n+1 \\ \alpha \neq \beta}} |Pe_\alpha| |Pe_\beta| \left| \dot{h}^\alpha \right| \left| \dot{h}^\beta \right| \\
& \leq 2\sqrt{\rho_1} \sum_{\alpha > n+1} |Pe_\alpha| \left| \dot{h}^\alpha \right| + \left(\sum_{\alpha > n+1} |Pe_\alpha| \left| \dot{h}^\alpha \right| \right)^2.
\end{aligned}$$

Using the Cauchy inequality, we have

$$\left(\sum_{\alpha > n+1} |Pe_\alpha| \left| \dot{h}^\alpha \right| \right)^2 \leq \left(\sum_{\alpha > n+1} |Pe_\alpha|^2 \right) \left(\sum_{\alpha > n+1} \left| \dot{h}^\alpha \right|^2 \right) = \theta_2 \rho_2.$$

So

$$\sum_{\alpha, \beta, i, j, k} \dot{h}_{ik}^\alpha \dot{h}_{jk}^\beta (J_{i\alpha} J_{j\beta} - J_{j\alpha} J_{i\beta}) \leq 2\sqrt{\theta_2 \rho_1 \rho_2} + \theta_2 \rho_2.$$

Thus we obtain $S_1 \leq \frac{3}{n} S_2 + 3|\dot{h}|^2 + 8\sqrt{\theta_2 \rho_1 \rho_2} + 4\theta_2 \rho_2$. \square

Let $F : M \times [0, T) \rightarrow \mathbb{C}P^{\frac{n+q}{2}}$ be a mean curvature flow with $2 \leq q \leq n-6$. Suppose that the initial submanifold M_0 satisfies the pinching condition $|\dot{h}|^2 < k|H|^2 + l$, where $k = \frac{1}{n(n-1)}$, $l = 2 - \frac{3}{n}$. Since M_0 is compact, there exists a small positive number ε , such that M_0 satisfies $|\dot{h}|^2 < (k|H|^2 + l)(1 - \varepsilon)$.

Now we prove that the pinching condition is preserved.

Theorem 3.5. *If M_0 satisfies $|\dot{h}|^2 < (k|H|^2 + l)(1 - \varepsilon)$, then this condition holds for all time $t \in [0, T)$.*

Proof. Let $k_\varepsilon = k(1 - \varepsilon)$, $l_\varepsilon = l(1 - \varepsilon)$. We set $U = |\dot{h}|^2 - k_\varepsilon |H|^2 - l_\varepsilon$. From the evolution equations we have

$$\begin{aligned}
\frac{1}{2} \left(\frac{\partial}{\partial t} - \Delta \right) U &= k_\varepsilon |\nabla H|^2 - \left| \nabla \dot{h} \right|^2 - n \left(|\dot{h}|^2 + k_\varepsilon |H|^2 \right) \\
&\quad + R_1 - \left(k_\varepsilon + \frac{1}{n} \right) R_2 + S_1 - 3 \left(k_\varepsilon + \frac{1}{n} \right) S_2.
\end{aligned}$$

Now we show that $\left(\frac{\partial}{\partial t} - \Delta \right) U$ is negative at all points where $|\dot{h}|^2 = k_\varepsilon |H|^2 + l_\varepsilon$.

From Lemma 3.4, if $|\dot{h}|^2 = k_\varepsilon |H|^2 + l_\varepsilon$, we get the following estimates.

$$\begin{aligned}
R_1 - \left(k_\varepsilon + \frac{1}{n} \right) R_2 &\leq \left(|\dot{h}|^2 - k_\varepsilon |H|^2 \right) |h|^2 - \left(\frac{1}{n} - k_\varepsilon \right) |H|^2 \rho_2 + 2\rho_1 \rho_2 + \frac{1}{2} \rho_2^2 \\
&\leq l_\varepsilon |h|^2 - (n-2)k_\varepsilon |H|^2 \rho_2 + 2\rho_1 \rho_2 + \frac{1}{2} \rho_2^2 \\
&= \left(k_\varepsilon + \frac{1}{n} \right) l_\varepsilon |H|^2 + l_\varepsilon^2 - (n-2)(\rho_1 + \rho_2 - l_\varepsilon) \rho_2 + 2\rho_1 \rho_2 + \frac{1}{2} \rho_2^2 \\
&= \left(k_\varepsilon + \frac{1}{n} \right) l_\varepsilon |H|^2 + l_\varepsilon^2 + (n-2)l_\varepsilon \rho_2 - (n-4)\rho_1 \rho_2 - \left(n - \frac{5}{2} \right) \rho_2^2,
\end{aligned}$$

and

$$S_1 - 3 \left(k_\varepsilon + \frac{1}{n}\right) S_2 \leq 3|\dot{h}|^2 + 8\sqrt{\theta_2 \rho_1 \rho_2} + 4\theta_2 \rho_2 \leq 3k_\varepsilon |H|^2 + 3l_\varepsilon + 8\sqrt{\theta_2 \rho_1 \rho_2} + 4\theta_2 \rho_2.$$

Thus, at a point where $U = 0$, we get

$$\begin{aligned} \frac{1}{2} \left(\frac{\partial}{\partial t} - \Delta \right) U &\leq k_\varepsilon |\nabla H|^2 - \left| \nabla \dot{h} \right|^2 - n(2k_\varepsilon |H|^2 + l_\varepsilon) \\ &\quad + \left(k_\varepsilon + \frac{1}{n}\right) l_\varepsilon |H|^2 + l_\varepsilon^2 + (n-2)l_\varepsilon \rho_2 - (n-4)\rho_1 \rho_2 - \left(n - \frac{5}{2}\right) \rho_2^2 \\ (3.8) \quad &\quad + 3k_\varepsilon |H|^2 + 3l_\varepsilon + 8\sqrt{\theta_2 \rho_1 \rho_2} + 4\theta_2 \rho_2 \\ &= k_\varepsilon |\nabla H|^2 - \left| \nabla \dot{h} \right|^2 + [(3-2n)k_\varepsilon + \left(k_\varepsilon + \frac{1}{n}\right) l_\varepsilon] |H|^2 \\ &\quad + (3-n+l_\varepsilon)l_\varepsilon + (n-2)l_\varepsilon \rho_2 - \left(n - \frac{5}{2}\right) \rho_2^2 \\ &\quad - (n-4)\rho_1 \rho_2 + 8\sqrt{\theta_2 \rho_1 \rho_2} + 4\theta_2 \rho_2. \end{aligned}$$

By the definition of $k_\varepsilon, l_\varepsilon$, we have

$$(3-2n)k_\varepsilon + \left(k_\varepsilon + \frac{1}{n}\right) l_\varepsilon \leq 0.$$

On the other hand, we have

$$(3.9) \quad -(n-4)\rho_1 \rho_2 + 8\sqrt{\theta_2 \rho_1 \rho_2} \leq \frac{16\theta_2}{n-4}$$

and

$$(3.10) \quad (3-n+l_\varepsilon)l_\varepsilon + (n-2)l_\varepsilon \rho_2 \leq \frac{(n-2)^2 l_\varepsilon}{4(n-3-l_\varepsilon)} \rho_2^2 \leq \frac{(n-2)^2 l}{4(n-3-l)} \rho_2^2.$$

Hence, the RHS of (3.8) is not greater than

$$k_\varepsilon |\nabla H|^2 - \left| \nabla \dot{h} \right|^2 + \left[\frac{(n-2)^2 l}{4(n-3-l)} - \left(n - \frac{5}{2}\right) \right] \rho_2^2 + 4\theta_2 \rho_2 + \frac{16\theta_2}{n-4}.$$

Since $n \geq q + 6 \geq 8$, we have $\frac{(n-2)^2 l}{4(n-3-l)} - \left(n - \frac{5}{2}\right) < 0$. Then we get

$$\begin{aligned} &\left[\frac{(n-2)^2 l}{4(n-3-l)} - \left(n - \frac{5}{2}\right) \right] \rho_2^2 + 4\theta_2 \rho_2 + \frac{16\theta_2}{n-4} \\ &\leq \frac{4}{n - \frac{5}{2} - \frac{(n-2)^2 l}{4(n-3-l)}} \theta_2^2 + \frac{16\theta_2}{n-4} \\ (3.11) \quad &\leq \left[\frac{4(q-1)}{n - \frac{5}{2} - \frac{(n-2)^2 l}{4(n-3-l)}} + \frac{16}{n-4} \right] \theta_2 \\ &\leq \left[\frac{4(n-7)}{n - \frac{5}{2} - \frac{(n-2)^2 l}{4(n-3-l)}} + \frac{16}{n-4} \right] \theta_2 \\ &= 8 \left[1 - \frac{-60 + 76n - 16n^2 + n^3}{(n-4)(-18 + 42n - 19n^2 + 2n^3)} \right] \theta_2 \leq 8\theta_2. \end{aligned}$$

By Lemma 2.1, we get

$$k_\varepsilon |\nabla H|^2 - \left| \nabla \dot{h} \right|^2 + 8\theta_2 \leq 0.$$

Therefore, the RHS of (3.8) is nonpositive for small ε . Then the assertion follows from the maximum principle. \square

Let $f_\sigma = |\dot{h}|^2/(k|H|^2 + l)^{1-\sigma}$, where $\sigma \in (0, 1)$ is a positive constant. Then we have

Lemma 3.6. *If M_0 satisfies $|\dot{h}|^2 < (k|H|^2 + l)(1 - \varepsilon)$, then the following inequality holds along the mean curvature flow.*

$$\frac{\partial}{\partial t} f_\sigma \leq \Delta f_\sigma + \frac{2k|\nabla f_\sigma||\nabla|H|^2|}{k|H|^2 + l} - \frac{2f_\sigma}{3|\dot{h}|^2} \left| \nabla \dot{h} \right|^2 + 2\sigma f_\sigma(|h|^2 + n) - 2\varepsilon l f_\sigma.$$

Proof. By a direct calculation, we have

$$\begin{aligned} \left(\frac{\partial}{\partial t} - \Delta \right) f_\sigma &= f_\sigma \left[\frac{1}{|\dot{h}|^2} \left(\frac{\partial}{\partial t} - \Delta \right) |\dot{h}|^2 - \frac{1-\sigma}{|H|^2 + l/k} \left(\frac{\partial}{\partial t} - \Delta \right) |H|^2 \right] \\ &\quad + 2(1-\sigma) \frac{\langle \nabla f_\sigma, \nabla |H|^2 \rangle}{|H|^2 + l/k} - \sigma(1-\sigma) f_\sigma \frac{|\nabla |H|^2|^2}{|H|^2 + l/k}. \end{aligned}$$

Now we estimate $(\partial_t - \Delta)|\dot{h}|^2$ and $(\partial_t - \Delta)|H|^2$. From (3.9), (3.11) and Lemma 3.4, we get

$$\begin{aligned} \left(\frac{\partial}{\partial t} - \Delta \right) |\dot{h}|^2 &\leq -2 \left| \nabla \dot{h} \right|^2 + 2|\dot{h}|^2(|h|^2 - n + 3) - \frac{2}{n} \rho_2 |H|^2 \\ &\quad + 4\rho_1 \rho_2 + \rho_2^2 + 16\sqrt{\theta_2 \rho_1 \rho_2} + 8\theta_2 \rho_2 \\ &\leq -2 \left| \nabla \dot{h} \right|^2 + 2|\dot{h}|^2(|h|^2 - n + 3) - \frac{2}{n} \rho_2 |H|^2 \\ &\quad + 2(n-2)\rho_1 \rho_2 + \rho_2^2 + \frac{32\theta_2}{n-4} + 8\theta_2 \rho_2 \\ (3.12) \quad &= -2 \left| \nabla \dot{h} \right|^2 + 2|\dot{h}|^2(|h|^2 - n + 3 + (n-2)\rho_2) - \frac{2}{n} \rho_2 |H|^2 \\ &\quad + (5-2n)\rho_2^2 + \frac{32\theta_2}{n-4} + 8\theta_2 \rho_2 \\ &\leq -2 \left| \nabla \dot{h} \right|^2 + 2|\dot{h}|^2(|h|^2 - n + 3 + (n-2)\rho_2) - \frac{2}{n} \rho_2 |H|^2 \\ &\quad + 16\theta_2 - \frac{(n-2)^2 l}{2(n-3-l)} \rho_2^2. \end{aligned}$$

We also have

$$\begin{aligned}
& -\frac{1-\sigma}{|H|^2+l/k} \left(\frac{\partial}{\partial t} - \Delta \right) |H|^2 \\
\leq & \frac{2|\nabla H|^2}{|H|^2+l/k} - \frac{2(1-\sigma)|H|^2}{|H|^2+l/k} (|h|^2 - \rho_2 + n) \\
\leq & \frac{2k|\nabla H|^2}{|\dot{h}|^2} - \left(2 - \frac{2l}{k|H|^2+l} \right) (|h|^2 - \rho_2 + n) + 2\sigma(|h|^2 + n) \\
= & \frac{2k|\nabla H|^2}{|\dot{h}|^2} + \frac{2l(|h|^2 - \rho_2 + n)}{k|H|^2+l} - 2(|h|^2 - \rho_2 + n) + 2\sigma(|h|^2 + n) \\
\leq & \frac{2k|\nabla H|^2}{|\dot{h}|^2} + \frac{2l[(1-\varepsilon)(k|H|^2+l) + \frac{1}{n}|H|^2 - \rho_2 + n]}{k|H|^2+l} \\
& - 2(|h|^2 - \rho_2 + n) + 2\sigma(|h|^2 + n) \\
\leq & \frac{2k|\nabla H|^2}{|\dot{h}|^2} + \frac{2l(\frac{1}{n}|H|^2 - \rho_2 + n)}{k|H|^2+l} + 2l(1-\varepsilon) - 2(|h|^2 - \rho_2 + n) + 2\sigma(|h|^2 + n).
\end{aligned}$$

Then we get

$$\begin{aligned}
& \left(\frac{\partial}{\partial t} - \Delta \right) f_\sigma \\
\leq & f_\sigma \left[\frac{1}{|\dot{h}|^2} \left(\frac{\partial}{\partial t} - \Delta \right) |h|^2 - \frac{1-\sigma}{|H|^2+l/k} \left(\frac{\partial}{\partial t} - \Delta \right) |H|^2 \right] + \frac{2|\nabla f_\sigma||\nabla|H|^2|}{|H|^2+l/k} \\
\leq & \frac{2f_\sigma}{|\dot{h}|^2} \left[-|\nabla \dot{h}|^2 + 8\theta_2 + k|\nabla H|^2 \right] + \frac{2|\nabla f_\sigma||\nabla|H|^2|}{|H|^2+l/k} \\
& + 2f_\sigma[-2n+3+(n-1)\rho_2+l(1-\varepsilon)+\sigma(|h|^2+n)] \\
& + 2f_\sigma \left[-\frac{1}{|\dot{h}|} \left[\frac{1}{n}\rho_2|H|^2 - \frac{(n-2)^2l}{4(n-3-l)}\rho_2^2 \right] + \frac{l(\frac{1}{n}|H|^2 - \rho_2 + n)}{k|H|^2+l} \right] \\
\leq & \frac{2f_\sigma}{|\dot{h}|^2} \left[-|\nabla \dot{h}|^2 + 8\theta_2 + k|\nabla H|^2 \right] + \frac{2|\nabla f_\sigma||\nabla|H|^2|}{|H|^2+l/k} \\
& + 2f_\sigma[(n-1)(\rho_2-l) + \sigma(|h|^2+n) - \varepsilon l] \\
& + \frac{2f_\sigma}{k|H|^2+l} \left[-\frac{1}{n}\rho_2|H|^2 - \frac{(n-2)^2l}{4(n-3-l)}\rho_2^2 + l(\frac{1}{n}|H|^2 - \rho_2 + n) \right].
\end{aligned}$$

By Lemma 2.1 and the condition $n \geq q+6 \geq 8$, we have

$$-|\nabla \dot{h}|^2 + 8\theta_2 + k|\nabla H|^2 \leq -\frac{1}{3}|\nabla \dot{h}|^2.$$

Using (3.10), we get

$$\begin{aligned}
& -\frac{1}{n}\rho_2|H|^2 - \frac{(n-2)^2l}{4(n-3-l)}\rho_2^2 + l(\frac{1}{n}|H|^2 - \rho_2 + n) \\
\leq & -\frac{1}{n}\rho_2|H|^2 - (3-n+l)l - (n-2)l\rho_2 + l(\frac{1}{n}|H|^2 - \rho_2 + n) \\
= & (n-1)(l-\rho_2)(k|H|^2+l).
\end{aligned}$$

Therefore, we complete the proof of this lemma. \square

3.3. The case of $q \geq n - 4$.

We choose an orthonormal frame $\{e_\alpha\}$ for the normal space such that $H = |H|e_{n+1}$. Let $\rho_1 = |\mathring{h}^{n+1}|^2$, $\rho_2 = \sum_{\alpha > n+1} |\mathring{h}^\alpha|^2$. We have

Lemma 3.7.

- (i) $R_1 \leq |h|^4 - \frac{2}{n}\rho_2|H|^2 + 2|\mathring{h}|^2\rho_2 - \frac{3}{2}\rho_2^2$,
- (ii) $R_2 = |H|^2(|h|^2 - \rho_2)$,
- (iii) $S_1 \leq \frac{3}{n}S_2 + (2n + 3)|\mathring{h}|^2$.

Proof. Using the same proof as Lemma 3.4, we obtain (i) and (ii).

Now we re-estimate S_1 . From (3.6) and (3.7), we have

$$S_1 \leq \frac{3}{n}S_2 + 3|\mathring{h}|^2 + 4 \sum_{\alpha, \beta, i, j, k} \mathring{h}_{ik}^\alpha \mathring{h}_{jk}^\beta (J_{i\alpha} J_{j\beta} - J_{i\beta} J_{j\alpha}).$$

With a local orthonormal frame, let v be a vector given by $v_k = \mathring{h}_{ik}^\alpha J_{i\alpha}$. We define two tensors D and E by

$$D_{ijk} = \mathring{h}_{ij}^\alpha J_{k\alpha} + \frac{\mathring{h}_{ik}^\alpha J_{j\alpha} + \mathring{h}_{jk}^\alpha J_{i\alpha}}{n + \eta}, \quad E_{ijk} = -\frac{2\delta_{ij}v_k}{(n + \eta)\eta} + \frac{\delta_{ik}v_j + \delta_{jk}v_i}{\eta},$$

where $\eta = \sqrt{n^2 + n - 2}$. Then we have $\langle D - E, E \rangle = 0$. This implies $|D|^2 \geq |E|^2$. By the definitions of D and E , we get

$$|D|^2 = \frac{5n - 2 - 4\eta}{(n - 2)^2} \left(n\mathring{h}_{ij}^\alpha \mathring{h}_{ij}^\beta J_{k\alpha} J_{k\beta} + 2\mathring{h}_{ik}^\alpha \mathring{h}_{jk}^\beta J_{i\beta} J_{j\alpha} \right),$$

$$|E|^2 = \frac{2(5n - 2 - 4\eta)}{(n - 2)^2} \mathring{h}_{ik}^\alpha \mathring{h}_{jk}^\beta J_{i\alpha} J_{j\beta}.$$

Thus we obtain

$$\sum_{\alpha, \beta, i, j, k} \mathring{h}_{ik}^\alpha \mathring{h}_{jk}^\beta (J_{i\alpha} J_{j\beta} - J_{i\beta} J_{j\alpha}) \leq \frac{n}{2} \sum_{i, j, k} \left(\sum_{\alpha} \mathring{h}_{ij}^\alpha J_{k\alpha} \right)^2 \leq \frac{n}{2} |\mathring{h}|^2.$$

□

Define a function $\psi : [0, +\infty) \rightarrow \mathbb{R}$ by

$$(3.13) \quad \psi(x) := \frac{9}{n^2 - 3n - 3} + \frac{n(n-3)}{n^3 - 4n^2 + 3}x - \frac{3\sqrt{x^2 + \frac{2}{n}(n-1)(n^2-3)x + 9(n-1)^2}}{n^3 - 4n^2 + 3}.$$

Let $\mathring{\psi}(x) = \psi(x) - \frac{x}{n}$. From Lemma 7.3 in the Appendix, the function $\mathring{\psi}$ has the following properties.

Lemma 3.8. *The function $\mathring{\psi}$ satisfies*

- (i) $0 \leq \mathring{\psi}'(x) < \frac{1}{n(n-1)}$, $0 \leq \mathring{\psi}(x) \leq \frac{x}{n(n-1)}$,
- (ii) $\max_{x \geq 0} \left(2x\mathring{\psi}''(x) + \mathring{\psi}'(x) \right) < \frac{2(n-4)}{n(n+8)}$,
- (iii) $3\mathring{\psi}(x) + \left(\mathring{\psi}(x) - x\mathring{\psi}'(x) \right) \left(\mathring{\psi}(x) + \frac{x}{n} + n \right) \leq 0$, and the equalities hold if and only $x = 0$,
- (iv) $0 \leq x\mathring{\psi}'(x) - \mathring{\psi}(x) < 2$.

For convenience, we denote $\dot{\psi}(|H|^2)$, $\dot{\psi}'(|H|^2)$, $\dot{\psi}''(|H|^2)$ by $\dot{\psi}$, $\dot{\psi}'$, $\dot{\psi}''$, respectively. Let $F : M^n \times [0, T) \rightarrow \mathbb{C}\mathbb{P}^{\frac{n+q}{2}}$ ($q \geq n - 4 \geq 2$) be a mean curvature flow whose initial value M_0 satisfies $|h|^2 < \dot{\psi}(|H|^2)$. Since M_0 is compact, there exists a small positive number ε , such that M_0 satisfies $|\dot{h}|^2 < \dot{\psi} - \varepsilon|H|^2 - \varepsilon$.

Theorem 3.9. *If the initial value M_0 satisfies $|\dot{h}|^2 < \dot{\psi} - \varepsilon|H|^2 - \varepsilon$, then this pinching condition holds for all $t \in [0, T)$.*

Proof. From Lemma 2.1 and Lemma 3.7, we get

$$(3.14) \quad \begin{aligned} \left(\frac{\partial}{\partial t} - \Delta\right) |\dot{h}|^2 &= -2 \left| \nabla \dot{h} \right|^2 - 2n |\dot{h}|^2 + 2R_1 - \frac{2}{n} R_2 + 2S_1 - \frac{6}{n} S_2 \\ &\leq -2 \left| \nabla \dot{h} \right|^2 + 2|\dot{h}|^2 |h|^2 - \frac{2}{n} \rho_2 |H|^2 + 4|\dot{h}|^2 \rho_2 + 2(n+3)|\dot{h}|^2. \end{aligned}$$

We have the following evolution equation of $\dot{\psi}$.

$$(3.15) \quad \begin{aligned} \left(\frac{\partial}{\partial t} - \Delta\right) \dot{\psi} &= -2\dot{\psi}' \cdot |\nabla H|^2 - \dot{\psi}'' \cdot |\nabla |H|^2|^2 + 2\dot{\psi}'(n|H|^2 + R_2 + 3S_2) \\ &\geq -2 \left(\dot{\psi}' + 2\dot{\psi}'' \cdot |H|^2 \right) |\nabla H|^2 + 2\dot{\psi}' \cdot |H|^2 (|h|^2 + n - \rho_2). \end{aligned}$$

Let $U = |\dot{h}|^2 - \dot{\psi} + \varepsilon|H|^2 + \varepsilon$. We obtain

$$\begin{aligned} \frac{1}{2} \left(\frac{\partial}{\partial t} - \Delta\right) U &\leq \left(\dot{\psi}' - \varepsilon + 2\dot{\psi}'' \cdot |H|^2 \right) |\nabla H|^2 - \left| \nabla \dot{h} \right|^2 \\ &\quad + |\dot{h}|^2 \left(|\dot{h}|^2 + \frac{1}{n} |H|^2 + n + 3 \right) - \left(\dot{\psi}' - \varepsilon \right) |H|^2 \left(|\dot{h}|^2 + \frac{1}{n} |H|^2 + n \right) \\ &\quad + \rho_2 \left(-\frac{1}{n} |H|^2 + 2|\dot{h}|^2 + \left(\dot{\psi}' - \varepsilon \right) |H|^2 \right). \end{aligned}$$

By Lemma 2.1 and Lemma 3.8 (ii), the first line of the RHS of the formula above is nonpositive. From $|\dot{h}|^2 = U + \dot{\psi} - \varepsilon|H|^2 - \varepsilon$, we obtain

$$(3.16) \quad \begin{aligned} \frac{1}{2} \left(\frac{\partial}{\partial t} - \Delta\right) U &\leq U \left(U + 2\dot{\psi} + n + 3 - 2\varepsilon + \left(\frac{1}{n} - \dot{\psi}' - \varepsilon \right) |H|^2 + 2\rho_2 \right) \\ &\quad + \dot{\psi} \left(\dot{\psi} + \frac{1}{n} |H|^2 + n + 3 \right) - \dot{\psi}' \cdot |H|^2 \left(\dot{\psi} + \frac{1}{n} |H|^2 + n \right) \\ &\quad + \varepsilon \left[-n - 3 + \varepsilon + \dot{\psi}' \cdot |H|^2 - 2\dot{\psi} \right. \\ &\quad \quad \left. + |H|^2 \left(\dot{\psi}' \cdot |H|^2 - \dot{\psi} - 3 - \frac{1}{n} + \varepsilon \right) \right] \\ &\quad + \rho_2 \left(-\frac{1}{n} |H|^2 + 2\dot{\psi} + \dot{\psi}' |H|^2 - \varepsilon(3|H|^2 + 2) \right). \end{aligned}$$

This together with Lemma 3.8 implies

$$(3.17) \quad \frac{1}{2} \left(\frac{\partial}{\partial t} - \Delta\right) U < U \left(U + 2\dot{\psi} + n + 3 - 2\varepsilon + \left(\frac{1}{n} - \dot{\psi}' - \varepsilon \right) |H|^2 \right).$$

Then the assertion follows from the maximum principle. \square

Now we prove that the mean curvature flow has finite maximal existence time in this case.

Lemma 3.10. *If the initial value M_0 satisfies $|\dot{h}|^2 < \dot{\psi}$, then T is finite.*

Proof. Let $U = |\dot{h}|^2 - \dot{\psi}$. Then $U < 0$ holds for all $t \in [0, T)$. From (3.17), we have

$$\begin{aligned} \left(\frac{\partial}{\partial t} - \Delta \right) U &\leq 2U \left(U + 2\dot{\psi} + n + 3 + \left(\frac{1}{n} - \dot{\psi}' \right) |H|^2 \right) \\ &\leq 2U \left(U + 2\dot{\psi} \right) \\ &= 2U \left(2|\dot{h}|^2 - U \right) \leq -2U^2. \end{aligned}$$

From the maximum principle, U will blow up in finite time. Therefore, T must be finite. \square

Let $f_\sigma = |\dot{h}|^2 / \dot{\psi}^{1-\sigma}$, where $\sigma \in (0, 1)$ is a positive constant. Then we have

Lemma 3.11. *If M_0 satisfies $|\dot{h}|^2 < \dot{\psi}$, then there exists a small positive constant ε , such that the following inequality holds along the mean curvature flow.*

$$\frac{\partial}{\partial t} f_\sigma \leq \Delta f_\sigma + \frac{2}{\dot{\psi}} |\nabla f_\sigma| |\nabla \dot{\psi}| - \frac{2\varepsilon f_\sigma}{|\dot{h}|^2} |\nabla \dot{h}|^2 + 2\sigma |h|^2 f_\sigma + 4n f_\sigma.$$

Proof. By a straightforward calculation, we have

$$\begin{aligned} \left(\frac{\partial}{\partial t} - \Delta \right) f_\sigma &= f_\sigma \left[\frac{1}{|\dot{h}|^2} \left(\frac{\partial}{\partial t} - \Delta \right) |\dot{h}|^2 - \frac{1-\sigma}{\dot{\psi}} \left(\frac{\partial}{\partial t} - \Delta \right) \dot{\psi} \right] \\ &\quad + 2(1-\sigma) \frac{\langle \nabla f_\sigma, \nabla \dot{\psi} \rangle}{\dot{\psi}} - \sigma(1-\sigma) f_\sigma \frac{|\nabla \dot{\psi}|^2}{|\dot{\psi}|^2}. \end{aligned}$$

Using (3.14) and (3.15), we have

$$\begin{aligned} \left(\frac{\partial}{\partial t} - \Delta \right) f_\sigma &\leq 2f_\sigma \left[-\frac{|\nabla \dot{h}|^2}{|\dot{h}|^2} + (1-\sigma) \frac{\dot{\psi}' + 2H^2 \dot{\psi}''}{\dot{\psi}} |\nabla H|^2 \right] \\ &\quad + 2f_\sigma \left[|h|^2 + n + 3 - (1-\sigma) \frac{\dot{\psi}' \cdot |H|^2}{\dot{\psi}} (|h|^2 + n) \right] \\ &\quad + 2f_\sigma \rho_2 \left[-\frac{|H|^2}{n|\dot{h}|^2} + 2 + (1-\sigma) \frac{\dot{\psi}' \cdot |H|^2}{\dot{\psi}} \right] + \frac{2}{\dot{\psi}} |\nabla f_\sigma| |\nabla \dot{\psi}|. \end{aligned}$$

From Lemma 2.1 and Lemma 3.8 (ii), we have

$$\begin{aligned} &-\frac{|\nabla \dot{h}|^2}{|\dot{h}|^2} + (1-\sigma) \frac{\dot{\psi}' + 2H^2 \dot{\psi}''}{\dot{\psi}} |\nabla H|^2 \\ &\leq -\frac{|\nabla \dot{h}|^2}{|\dot{h}|^2} + (1-\sigma) \frac{(1-\varepsilon) \frac{2(n-4)}{n(n+8)}}{\dot{\psi}} |\nabla H|^2 \\ &\leq -\frac{|\nabla \dot{h}|^2}{|\dot{h}|^2} + \frac{(1-\varepsilon) |\nabla \dot{h}|^2}{|\dot{h}|^2} = -\frac{\varepsilon |\nabla \dot{h}|^2}{|\dot{h}|^2}. \end{aligned}$$

By Lemma 3.8 (iii), we get

$$\begin{aligned}
& |h|^2 + n + 3 - (1 - \sigma) \frac{\dot{\psi}' \cdot |H|^2}{\dot{\psi}} (|h|^2 + n) \\
& \leq (1 - \sigma) \frac{\dot{\psi} - \dot{\psi}' \cdot |H|^2}{\dot{\psi}} |h|^2 + n + 3 + \sigma |h|^2 \\
& \leq 2n + \sigma |h|^2.
\end{aligned}$$

By Lemma 3.8 (i), we have

$$-\frac{|H|^2}{n|\dot{h}|^2} + 2 + (1 - \sigma) \frac{\dot{\psi}' \cdot |H|^2}{\dot{\psi}} \leq \frac{-\frac{1}{n}|H|^2 + 2\dot{\psi} + \dot{\psi}' \cdot |H|^2}{\dot{\psi}} \leq 0.$$

This completes the proof of the lemma. \square

4. AN ESTIMATE FOR TRACELESS SECOND FUNDAMENTAL FORM

Let $F : M^n \times [0, T) \rightarrow \mathbb{C}\mathbb{P}^{\frac{n+q}{2}}$ be a mean curvature flow. Suppose that the initial value M_0 satisfies the condition in Theorem 1.1. We put

$$W = \begin{cases} \dot{\varphi}_\varepsilon, & q = 1, \\ \frac{1}{n(n-1)} |H|^2 + 2 - \frac{3}{n}, & 2 \leq q < n - 4, \\ \dot{\psi}, & q \geq n - 4. \end{cases}$$

By the conclusions of the previous section, there exists a sufficiently small positive number ε , such that for all $t \in [0, T)$, the following pinching condition holds.

$$|\dot{h}|^2 < W - \varepsilon |H|^2 - \varepsilon.$$

From this inequality and the definition of W , we have $\varepsilon |H|^2 + \varepsilon < W < \frac{|H|^2}{n(n-1)} + n$.

We introduce an auxiliary function:

$$f_\sigma = \frac{|\dot{h}|^2}{W^{1-\sigma}},$$

In this section, we will show that f_σ decays exponentially.

Lemma 4.1. *There exist positive constants ε and C_1 depending on M_0 , such that*

$$\frac{\partial}{\partial t} f_\sigma \leq \Delta f_\sigma + \frac{2C_1}{|\dot{h}|} |\nabla f_\sigma| |\nabla \dot{h}| - \frac{\varepsilon f_\sigma}{|\dot{h}|^2} |\nabla \dot{h}|^2 + \frac{4n\sigma}{\varepsilon} f_\sigma W + (\chi - 2\varepsilon) f_\sigma,$$

where $\chi = 5n \frac{\max\{q-n+5, 0\}}{q-n+5}$.

Proof. Combining the conclusions of Lemmas 4.1, 3.6 and 3.11, we have the following inequality with some suitable small ε .

$$\frac{\partial}{\partial t} f_\sigma \leq \Delta f_\sigma + \frac{2}{W} |\nabla f_\sigma| |\nabla W| - \frac{\varepsilon f_\sigma}{|\dot{h}|^2} |\nabla \dot{h}|^2 + 2\sigma f_\sigma (|h|^2 + n) + (\chi - 2\varepsilon) f_\sigma.$$

By the definition of W , there exists a constant B_1 such that $|\nabla W| < B_1 |\nabla |H|^2|$. Let C_1 be a constant such that $2B_1 |H| / \sqrt{W} < \frac{1}{n} C_1$. From Lemma 2.1, we have

$$(4.1) \quad \frac{|\nabla W|}{W} \leq \frac{2B_1 |H| |\nabla H|}{\sqrt{W} |\dot{h}|} \leq \frac{\frac{1}{n} C_1 |\nabla H|}{|\dot{h}|} \leq \frac{C_1 |\nabla \dot{h}|}{|\dot{h}|}.$$

We also have

$$|h|^2 + n < W + \frac{1}{n}|H|^2 + n < W + n(|H|^2 + 1) < \frac{2n}{\varepsilon}W.$$

Thus we complete the proof of this lemma. \square

We need the following estimate for the the Laplacian of $|\mathring{h}|^2$.

Lemma 4.2. *There exists a positive constant $C_2 > 1$, such that*

$$\Delta|\mathring{h}|^2 \geq 2\langle \mathring{h}, \nabla^2 H \rangle + 2\left(\varepsilon|\mathring{h}|^2 - C_2\right)W.$$

Proof. We have the following identity

$$\Delta|\mathring{h}|^2 = 2\langle \mathring{h}, \nabla^2 H \rangle + 2|\nabla\mathring{h}|^2 - 2R_1 + 2R_3 + 2n|\mathring{h}|^2 - 2S_1 + \frac{\varepsilon}{n}S_2 + 6S_3,$$

where $R_3 = H^\alpha h_{ik}^\alpha h_{ij}^\beta h_{jk}^\beta$, $S_3 = \mathring{h}_{ij}^\alpha H^\beta J_{i\alpha} J_{j\beta}$. Then there exists a positive constant B_2 only depending on n , such that

$$\Delta|\mathring{h}|^2 \geq 2\langle \mathring{h}, \nabla^2 H \rangle + 2(R_3 - R_1) - 2B_2|h|^2.$$

We choose a local orthonormal frame, such that $H = |H|e_{n+1}$ and $\mathring{h}^1 = \text{diag}(\mathring{\lambda}_1, \dots, \mathring{\lambda}_n)$.

Let $\rho_1 = |\mathring{h}^{n+1}|^2$, $\rho_2 = \sum_{\alpha > n+1} |\mathring{h}^\alpha|^2$. Expanding R_3 , we get

$$R_3 = \frac{1}{n^2}|H|^4 + \frac{3\rho_1 + \rho_2}{n}|H|^2 + |H| \sum_{\alpha, i} \mathring{\lambda}_i \left(\mathring{h}_{ii}^\alpha\right)^2 + \frac{|H|}{2} \sum_{\substack{\alpha > 1 \\ i \neq j}} \left(\mathring{\lambda}_i + \mathring{\lambda}_j\right) \left(\mathring{h}_{ij}^\alpha\right)^2.$$

By Lemma 2.6 of [28] or Proposition 1.6 of [31], we have

$$\begin{aligned} \sum_{\alpha, i} \mathring{\lambda}_i \left(\mathring{h}_{ii}^\alpha\right)^2 &\geq -\frac{n-2}{\sqrt{n(n-1)}}\sqrt{\rho_1} \left(|\mathring{h}|^2 - \sum_{\substack{\alpha > 1 \\ i \neq j}} \left(\mathring{h}_{ij}^\alpha\right)^2\right) \\ &\geq -\frac{n-2}{\sqrt{n(n-1)}} \left(\frac{1}{2} \left(P_1 + |\mathring{h}|^2\right) |\mathring{h}| - \sqrt{\rho_1} \sum_{\substack{\alpha > 1 \\ i \neq j}} \left(\mathring{h}_{ij}^\alpha\right)^2\right) \\ &\geq -\frac{n-2}{\sqrt{n(n-1)}} \left(|\mathring{h}|^3 - \frac{1}{2}|\mathring{h}|\rho_2 - \sqrt{\rho_1} \sum_{\substack{\alpha > 1 \\ i \neq j}} \left(\mathring{h}_{ij}^\alpha\right)^2\right). \end{aligned}$$

We also have

$$\sum_{\substack{\alpha > 1 \\ i \neq j}} \left(\mathring{\lambda}_i + \mathring{\lambda}_j\right) \left(\mathring{h}_{ij}^\alpha\right)^2 \geq \sum_{\substack{\alpha > 1 \\ i \neq j}} -\sqrt{2\left(\mathring{\lambda}_i^2 + \mathring{\lambda}_j^2\right)} \left(\mathring{h}_{ij}^\alpha\right)^2 \geq -\sqrt{2\rho_1} \sum_{\substack{\alpha > 1 \\ i \neq j}} \left(\mathring{h}_{ij}^\alpha\right)^2.$$

Note that $\frac{n-2}{\sqrt{n(n-1)}} > \frac{\sqrt{2}}{2}$ if $n \geq 6$, and $\sum_{\alpha > 1, i \neq j} \left(\mathring{h}_{ij}^\alpha\right)^2 = 0$ if $q = 1$. We get

$$(4.2) \quad R_3 \geq \frac{1}{n^2}|H|^4 + \frac{3\rho_1 + \rho_2}{n}|H|^2 - \frac{n-2}{\sqrt{n(n-1)}}|H| \left(|\mathring{h}|^3 - \frac{1}{2}|\mathring{h}|\rho_2\right).$$

From (4.2) and Lemma 3.4 (i), we get

$$\begin{aligned}
R_3 - R_1 &\geq |\dot{h}|^2 \left(-|\dot{h}|^2 + \frac{1}{n}|H|^2 - \frac{n-2}{\sqrt{n(n-1)}}|H||\dot{h}| \right) \\
&\quad + \rho_2 \left(-2|\dot{h}|^2 + \frac{n-2}{2\sqrt{n(n-1)}}|H||\dot{h}| + \frac{3}{2}\rho_2 \right) \\
&\geq |\dot{h}|^2 \left[\varepsilon(|H|^2 + 1) - W + \frac{1}{n}|H|^2 - \frac{n-2}{\sqrt{n(n-1)}}|H|\sqrt{W} \right] \\
&\quad - \rho_2 \left[|\dot{h}| \left(2\sqrt{W} - \frac{n-2}{2\sqrt{n(n-1)}}|H| \right) - \frac{3}{2}\rho_2 \right] \\
&\geq |\dot{h}|^2 \left[\varepsilon(|H|^2 + 1) + \frac{(n-2)|H|^2}{n(n-1)} - n - \frac{n-2}{\sqrt{n(n-1)}}|H|\sqrt{\frac{|H|^2}{n(n-1)} + n} \right] \\
&\quad - \rho_2 \left[|\dot{h}| \left(2\sqrt{\frac{|H|^2}{n(n-1)} + n} - \frac{n-2}{2\sqrt{n(n-1)}}|H| \right) - \frac{3}{2}\rho_2 \right] \\
&\geq |\dot{h}|^2 [\varepsilon(|H|^2 + 1) - (n-1)n] - \rho_2 \left[|\dot{h}| \cdot 2\sqrt{n} - \frac{3}{2}\rho_2 \right] \\
&\geq |\dot{h}|^2 [\varepsilon(|H|^2 + 1) - (n-1)n] - \frac{2n}{3}|\dot{h}|^2.
\end{aligned}$$

By choosing suitable large C_2 , we complete the proof of this lemma. \square

From (4.1) and Lemma 4.2, we have

$$\begin{aligned}
\Delta f_\sigma &= f_\sigma \left(\frac{\Delta|\dot{h}|^2}{|\dot{h}|^2} - (1-\sigma)\frac{\Delta W}{W} \right) - 2(1-\sigma)\frac{\langle \nabla f_\sigma, \nabla W \rangle}{W} + \sigma(1-\sigma)f_\sigma\frac{|\nabla W|^2}{W^2} \\
&\geq \frac{f_\sigma\Delta|\dot{h}|^2}{|\dot{h}|^2} - (1-\sigma)\frac{f_\sigma\Delta W}{W} - \frac{2C_1|\nabla f_\sigma||\nabla\dot{h}|}{|\dot{h}|} \\
&\geq \frac{2\langle \dot{h}, \nabla^2 H \rangle}{W^{1-\sigma}} + 2\varepsilon f_\sigma W - 2C_2 W^\sigma - (1-\sigma)\frac{f_\sigma\Delta W}{W} - \frac{2C_1|\nabla f_\sigma||\nabla\dot{h}|}{|\dot{h}|}.
\end{aligned}$$

Multiplying both sides of the above inequality by f_σ^{p-1} , we get

$$2\varepsilon f_\sigma^p W \leq f_\sigma^{p-1} \Delta f_\sigma + (1-\sigma)\frac{f_\sigma^p \Delta W}{W} - \frac{2f_\sigma^{p-1} \langle \dot{h}, \nabla^2 H \rangle}{W^{1-\sigma}} + \frac{2C_1 f_\sigma^{p-1} |\nabla f_\sigma| |\nabla \dot{h}|}{|\dot{h}|} + 2C_2 f_\sigma^{p-1} W^\sigma.$$

The Young's inequality yields

$$\begin{aligned}
2C_2 f_\sigma^{p-1} W^\sigma &\leq 2C_2 W \left[\frac{1}{p} \left(\frac{2C_2}{\varepsilon W^{(1-\sigma)}} \right)^p + \frac{p-1}{p} \left(\frac{\varepsilon}{2C_2} \right)^{\frac{p}{p-1}} f_\sigma^p \right] \\
&\leq 2C_2 \left(\frac{1}{p} \left(\frac{2C_2}{\varepsilon^{(2-\sigma)}} \right)^p + \frac{\varepsilon}{2C_2} f_\sigma^p W \right) \\
&\leq C_3^p + \varepsilon f_\sigma^p W,
\end{aligned}$$

where C_3 is a positive constant depending on M_0 . Thus

$$(4.3) \quad \varepsilon f_\sigma^p W \leq f_\sigma^{p-1} \Delta f_\sigma + (1-\sigma) \frac{f_\sigma^p \Delta W}{W} - \frac{2f_\sigma^{p-1} \langle \dot{h}, \nabla^2 H \rangle}{W^{1-\sigma}} + \frac{2C_1 f_\sigma^{p-1} |\nabla f_\sigma| |\nabla \dot{h}|}{|\dot{h}|} + C_3^p.$$

Then integrate both side of (4.3) over M_t . By the divergence theorem, we have

$$(4.4) \quad \int_{M_t} f_\sigma^{p-1} \Delta f_\sigma d\mu_t = -(p-1) \int_{M_t} f_\sigma^{p-2} |\nabla f_\sigma|^2 d\mu_t.$$

From (4.1), we have

$$(4.5) \quad \begin{aligned} \int_{M_t} \frac{f_\sigma^p}{W} \Delta W d\mu_t &= - \int_{M_t} \left\langle \nabla \left(\frac{f_\sigma^p}{W} \right), \nabla W \right\rangle d\mu_t \\ &= \int_{M_t} \left(-\frac{p f_\sigma^{p-1}}{W} \langle \nabla f_\sigma, \nabla W \rangle + \frac{f_\sigma^p}{W^2} |\nabla W|^2 \right) d\mu_t \\ &\leq \int_{M_t} \left(\frac{C_1 p f_\sigma^{p-1}}{|\dot{h}|} |\nabla f_\sigma| |\nabla \dot{h}| + \frac{C_1^2 f_\sigma^p}{|\dot{h}|^2} |\nabla \dot{h}|^2 \right) d\mu_t. \end{aligned}$$

We also have

$$(4.6) \quad \begin{aligned} &- \int_{M_t} \frac{f_\sigma^{p-1} \langle \dot{h}, \nabla^2 H \rangle}{W^{1-\sigma}} d\mu_t \\ &= \int_{M_t} \nabla_i \left(\frac{f_\sigma^{p-1}}{W^{1-\sigma}} \dot{h}_{ij}^\alpha \right) \nabla_j H^\alpha d\mu_t \\ &= \int_{M_t} \left[\frac{(p-1) f_\sigma^{p-2}}{W^{1-\sigma}} \dot{h}_{ij}^\alpha \nabla_i f_\sigma - \frac{(1-\sigma) f_\sigma^{p-1}}{W^{2-\sigma}} \dot{h}_{ij}^\alpha \nabla_i W + \frac{f_\sigma^{p-1}}{W^{1-\sigma}} \nabla_i \dot{h}_{ij}^\alpha \right] \nabla_j H^\alpha d\mu_t \\ &\leq \int_{M_t} \left[\frac{(p-1) f_\sigma^{p-1}}{|\dot{h}|} |\nabla f_\sigma| + \frac{f_\sigma^{p-1}}{W^{2-\sigma}} |\dot{h}| |\nabla W| + \frac{f_\sigma^{p-1}}{W^{1-\sigma}} n |\nabla \dot{h}| \right] |\nabla H| d\mu_t \\ &\leq \int_{M_t} \left[\frac{(p-1) f_\sigma^{p-1}}{|\dot{h}|} |\nabla f_\sigma| + \frac{C_1 f_\sigma^{p-1}}{W^{1-\sigma}} |\dot{h}| |\nabla \dot{h}| + \frac{f_\sigma^{p-1}}{W^{1-\sigma}} n |\nabla \dot{h}| \right] n |\nabla \dot{h}| d\mu_t \\ &\leq \int_{M_t} \left[\frac{n(p-1) f_\sigma^{p-1}}{|\dot{h}|} |\nabla f_\sigma| |\nabla \dot{h}| + \frac{(C_1 + n^2) f_\sigma^p}{|\dot{h}|^2} |\nabla \dot{h}|^2 \right] d\mu_t. \end{aligned}$$

Putting (4.3)-(4.6) together, we get

$$(4.7) \quad \int_{M_t} W f_\sigma^p d\mu_t \leq \frac{C_4}{\varepsilon} \int_{M_t} \left[\frac{p f_\sigma^{p-1}}{|\dot{h}|} |\nabla f_\sigma| |\nabla \dot{h}| + \frac{f_\sigma^p}{|\dot{h}|^2} |\nabla \dot{h}|^2 + C_3^p \right] d\mu_t,$$

where C_4 is a positive constant independent of t .

Combining Lemma 4.1 and (4.7), we obtain

$$\begin{aligned}
\frac{d}{dt} \int_{M_t} f_\sigma^p d\mu_t &= p \int_{M_t} f_\sigma^{p-1} \frac{\partial f_\sigma}{\partial t} d\mu_t - \int_{M_t} f_\sigma^p |H|^2 d\mu_t \\
&\leq p \int_{M_t} f_\sigma^{p-2} \left[-(p-1) |\nabla f_\sigma|^2 + \left(2C_1 + \frac{4n\sigma C_4 p}{\varepsilon^2} \right) \frac{f_\sigma}{|\dot{h}|} |\nabla f_\sigma| |\nabla \dot{h}| \right. \\
(4.8) \quad &\quad \left. - \left(\varepsilon - \frac{4n\sigma C_4}{\varepsilon^2} \right) \frac{f_\sigma^2}{|\dot{h}|^2} |\nabla \dot{h}|^2 \right] d\mu_t \\
&\quad + p \int_{M_t} f_\sigma^p \left(\frac{4n\sigma C_4 C_3^p}{\varepsilon^2} + \chi - 2\varepsilon \right) d\mu_t.
\end{aligned}$$

Now we show that the L^p -norm of f_σ decays exponentially.

Lemma 4.3. *There exist positive constants C_5, p_0, σ_0 depending on M_0 , such that for all $p \geq p_0$ and $\sigma \leq \sigma_0/\sqrt{p}$, we have*

$$\left(\int_{M_t} f_\sigma^p d\mu_t \right)^{\frac{1}{p}} < C_5 e^{-\varepsilon t}.$$

Proof. The expression in the square bracket of the right hand side of (4.8) is a quadratic polynomial. With p_0 large enough and σ_0 small enough, its discriminant satisfies $\left(2C_1 + \frac{4n\sigma C_4 p}{\varepsilon^2} \right)^2 - 4(p-1) \left(\varepsilon - \frac{4n\sigma C_4}{\varepsilon^2} \right) < 0$. We also have $\frac{4n\sigma C_4 C_3^p}{\varepsilon^2} < \varepsilon$. Then we get

$$\frac{d}{dt} \int_{M_t} f_\sigma^p d\mu_t \leq p(\chi - \varepsilon) \int_{M_t} f_\sigma^p d\mu_t.$$

Since $\chi = 5n \frac{\max\{q-n+5, 0\}}{q-n+5}$, we have $\int_{M_t} f_\sigma^p d\mu_t \leq e^{-p\varepsilon t} \int_{M_0} f_\sigma^p d\mu_0$ for $q < n-4$, and $\int_{M_t} f_\sigma^p d\mu_t \leq e^{5npT} \int_{M_0} f_\sigma^p d\mu_0$ for $q \geq n-4$. Noting that T is finite if $q \geq n-4$, we obtain the conclusion. \square

Let $g_\sigma = f_\sigma e^{\varepsilon t/2}$. By the Sobolev inequality on submanifolds [13] and a Stampacchia iteration procedure, we obtain that g_σ is uniformly bounded for all t (see [14] or [18] for the details). Then we obtain the following theorem.

Theorem 4.4. *There exist positive constants σ and C_0 depending only on M_0 , such that for all $t \in [0, T)$ we have*

$$|\dot{h}|^2 \leq C_0 (|H|^2 + 1)^{1-\sigma} e^{-\varepsilon t/2}.$$

5. A GRADIENT ESTIMATE

In the following, we derive an estimate for $|\nabla H|^2$ along the mean curvature flow. Firstly, the same as Proposition 4.3 in [27], we have the following result.

Lemma 5.1. *There exists a constant $C_6 > 1$ depending only on n , such that*

$$\frac{\partial}{\partial t} |\nabla H|^2 \leq \Delta |\nabla H|^2 + C_6 (|H|^2 + 1) |\nabla h|^2.$$

Secondly, we need the following estimates.

Lemma 5.2. *Along the mean curvature flow, we have*

- (i) $\frac{\partial}{\partial t}|H|^4 \geq \Delta|H|^4 - 24n|H|^2|\nabla h|^2 + \frac{4}{n}|H|^6$,
 - (ii) $\frac{\partial}{\partial t}|\dot{h}|^2 \leq \Delta|\dot{h}|^2 - \frac{1}{3}|\nabla h|^2 + C_7(|H|^2 + 1)|\dot{h}|^2$,
 - (iii) $\frac{\partial}{\partial t}(|H|^2|\dot{h}|^2) \leq \Delta(|H|^2|\dot{h}|^2) - \frac{1}{6}|H|^2|\nabla h|^2 + C_9|\nabla h|^2 + C_8(|H|^2 + 1)^2|\dot{h}|^2$,
- where C_7, C_8, C_9 are sufficiently large constants.

Proof. (i) From Lemma 2.2 (ii) we derive that

$$\frac{\partial}{\partial t}|H|^4 = \Delta|H|^4 - 4|H|^2|\nabla H|^2 - 2|\nabla|H|^2|^2 + 4|H|^2(R_2 + 3S_2) + 4n|H|^4.$$

From Lemma 3.7, we get $R_2 + 3S_2 \geq R_2 \geq \frac{1}{n}|H|^4$. Then from Lemma 2.1, we have $4|H|^2|\nabla H|^2 + 2|\nabla|H|^2|^2 \leq 24n|H|^2|\nabla h|^2$.

(ii) We have

$$\frac{\partial}{\partial t}|\dot{h}|^2 = \Delta|\dot{h}|^2 - 2|\nabla\dot{h}|^2 - 2n|\dot{h}|^2 + 2R_1 - \frac{2}{n}R_2 + 2S_1 - \frac{6}{n}S_2.$$

From Lemma 2.1, we get $2|\nabla\dot{h}|^2 \geq \frac{1}{3}|\nabla h|^2$. Choosing a large constant C_7 , we obtain inequality (ii).

(iii) It follows from the evolution equations that

$$\begin{aligned} \frac{\partial}{\partial t}(|H|^2|\dot{h}|^2) &= \Delta(|H|^2|\dot{h}|^2) + 2|H|^2\left(R_1 - \frac{R_2}{n} + S_1 - \frac{3S_2}{n}\right) + 2|\dot{h}|^2(R_2 + 3S_2) \\ &\quad - 2|H|^2|\nabla\dot{h}|^2 - 2|\dot{h}|^2|\nabla H|^2 - 2\langle\nabla|H|^2, \nabla|\dot{h}|^2\rangle. \end{aligned}$$

From Lemma 2.1, we get $-2|H|^2|\nabla\dot{h}|^2 \leq -\frac{1}{3}|H|^2|\nabla h|^2$.

From the preserved pinching condition $|\dot{h}|^2 < W$, we have

$$2|H|^2\left(R_1 - \frac{R_2}{n} + S_1 - \frac{3S_2}{n}\right) + 2|\dot{h}|^2(R_2 + 3S_2) < C_8(|H|^2 + 1)^2|\dot{h}|^2.$$

Using Theorem 4.4, we have

$$-2\langle\nabla|H|^2, \nabla|\dot{h}|^2\rangle \leq 8|H||\nabla H||\dot{h}||\nabla h| \leq 8n\sqrt{C_0}|H|(|H|^2 + 1)^{\frac{1-\sigma}{2}}|\nabla h|^2.$$

By Young's inequality, there exists a positive constant C_9 , such that $-2\langle\nabla|H|^2, \nabla|\dot{h}|^2\rangle \leq (C_9 + \frac{1}{6}|H|^2)|\nabla h|^2$. \square

Now we prove a gradient estimate for mean curvature.

Theorem 5.3. *For all $\eta \in (0, \frac{\varepsilon^{1/2}}{8n\pi})$, there exists a number $\Psi(\eta)$ depending on η and M_0 , such that*

$$|\nabla H|^2 < [(\eta|H|)^4 + \Psi(\eta)^2]e^{-\varepsilon t/4}.$$

Proof. Define a scalar

$$f = \left(|\nabla H|^2 + B_1|\dot{h}|^2 + B_2|H|^2|\dot{h}|^2\right)e^{\varepsilon t/4} - (\eta|H|)^4,$$

where B_1, B_2 are two positive constants.

From Lemmas 5.1 and 5.2, we obtain

$$\begin{aligned} \left(\frac{\partial}{\partial t} - \Delta\right) f &\leq \left[e^{\varepsilon t/4} \left(C_6 - \frac{B_2}{6} \right) + 24n\eta^4 \right] |H|^2 |\nabla h|^2 \\ &\quad + e^{\varepsilon t/4} \left[\left(C_6 - \frac{B_1}{3} + B_2 C_9 \right) |\nabla h|^2 + \sigma |\nabla H|^2 \right] \\ &\quad + e^{\varepsilon t/4} [B_1 C_7 (|H|^2 + 1) + B_2 C_8 (|H|^2 + 1)^2 + \sigma (B_1 + B_2 |H|^2)] |\dot{h}|^2 \\ &\quad - \frac{4\eta^4}{n} |H|^6. \end{aligned}$$

We choose the constants B_1 and B_2 , such that $C_6 - \frac{B_2}{6} < -1$ and $C_6 - \frac{B_1}{3} + B_2 C_9 < -1$. Then applying Theorem 4.4, we get

$$(5.1) \quad \left(\frac{\partial}{\partial t} - \Delta\right) f \leq e^{-\varepsilon t/4} \left[B_3 (|H|^2 + 1)^2 (|H|^2 + 1)^{1-\sigma} - \frac{4\eta^4}{n} |H|^6 \right].$$

Consider the expression in the bracket of (5.1). Since the coefficient of $|H|^6$ is negative, it has an upper bound $\Psi_2(\eta)$. Then we have $(\frac{\partial}{\partial t} - \Delta) f \leq \Psi_2(\eta) e^{-\varepsilon t/4}$. It follows from the maximum principle that f is bounded. This completes the proof of Theorem 5.3. \square

6. CONVERGENCE

In order to estimate the diameter of M_t , we need the well-known Myers theorem.

Theorem 6.1. (Myers) *Let Γ be a geodesic of length l on M . If the Ricci curvature satisfies $\text{Ric}(X) \geq (n-1)\frac{\pi^2}{l^2}$, for each unit vector $X \in T_x M$, at any point $x \in \Gamma$, then Γ has conjugate points.*

Then we obtain the following lemma.

Lemma 6.2. *Let M be an n -dimensional submanifold in $\mathbb{C}\mathbb{P}^{\frac{n+q}{2}}$. If there exist positive constants L and ε , $\varepsilon < \frac{1}{n(n-1)}$, such that M satisfies $|h|^2 < \left(\frac{1}{n-1} - \varepsilon\right) |H|^2 + L$, $\max_M |H|^2 > 2nL/\varepsilon$ and $|\nabla H| < 2\eta^2 \max_M |H|^2$ for all $0 < \eta < \frac{\varepsilon^{1/2}}{8n\pi}$, then we have*

$$\frac{\min_M |H|^2}{\max_M |H|^2} > 1 - \eta \quad \text{and} \quad \text{diam } M \leq (2\eta |H|_{\max})^{-1}.$$

Proof. By Proposition 2 in [30], the Ricci curvature of M satisfies

$$\begin{aligned} \text{Ric} &\geq \frac{n-1}{n} \left(n + \frac{2}{n} |H|^2 - |h|^2 - \frac{n-2}{\sqrt{n(n-1)}} |H| |\dot{h}| \right) \\ &\geq \frac{n-1}{n} \left[n + \frac{2}{n} |H|^2 - \left(\frac{1}{n-1} - \varepsilon \right) |H|^2 - L - \frac{n-2}{\sqrt{n(n-1)}} |H| \sqrt{\frac{|H|^2}{n(n-1)} + L} \right] \\ &\geq \frac{n-1}{n} \left[n + \left(\frac{n-2}{n(n-1)} + \varepsilon \right) |H|^2 - L - (n-2) \left(\frac{|H|^2}{n(n-1)} + L \right) \right] \\ &\geq \frac{n-1}{n} [\varepsilon |H|^2 - nL]. \end{aligned}$$

Let x be a point on M where $|H|$ achieves its maximum. Consider all the geodesics of length $l = (4\eta \max_M |H|)^{-1}$ starting from x . Since $|\nabla |H|^2| < 4\eta^2 \max_M |H|^3$, we have $|H|^2 > \max_M |H|^2 - 4\eta^2 \max_M |H|^3 \cdot l = (1 - \eta) \max_M |H|^2$ along such a geodesic. Thus we have $\text{Ric} > \frac{n-1}{n} [\varepsilon |H|^2 - nL] > \frac{n-1}{n} \varepsilon \left(\frac{1}{2} - \eta \right) \max_M |H|^2 >$

$(n-1)\pi^2/l^2$ on such a geodesic. Then from Myers' theorem, these geodesics can reach any point of M .

Therefore, we obtain $\min_M |H|^2 > (1-\eta)\max_M |H|^2$ and $\text{diam } M \leq 2l$. \square

Now we show that under the assumption of Theorem 1.1, the mean curvature flow converges to a point or a totally geodesic submanifold.

Theorem 6.3. *If M_0 satisfies $|\mathring{h}|^2 < W$ and T is finite, then F_t converges to a round point as $t \rightarrow T$.*

Proof. If T is finite, we have $\max_{M_t} |h|^2 \rightarrow \infty$ as $t \rightarrow T$. Let $\underline{H} = \min_{M_t} |H|$, $\overline{H} = \max_{M_t} |H|$. From the preserved pinching condition, we get $\overline{H} \rightarrow \infty$ as $t \rightarrow T$.

By Theorem 5.3, for any $\eta \in \left(0, \frac{\varepsilon^{1/2}}{8n\pi}\right)$, we have $|\nabla H| < (\eta|H|)^2 + \Psi(\eta)$. Since $\overline{H} \rightarrow \infty$ as $t \rightarrow T$, there exists a time τ depending on η , such that for $t > \tau$, $\overline{H}^2 > \Psi(\eta)/\eta^2$. Then we have $|\nabla H| < 2\eta^2\overline{H}^2$. Using Lemma 6.2, we obtain $\text{diam } M_t \rightarrow 0$ and $\underline{H}/\overline{H} \rightarrow 1$ as $t \rightarrow T$.

Now we dilate the metric of the ambient space such that the submanifold maintains its volume along the flow. Using the same method as in [23], we can prove that the rescaled mean curvature flow converges to a round sphere as the reparameterized time tends to infinity. \square

Theorem 6.4. *If M_0 satisfies $|\mathring{h}|^2 < W$ and $T = \infty$, then F_t converges to a totally geodesic submanifold $\mathbb{C}\mathbb{P}^{n/2}$ as $t \rightarrow \infty$.*

Proof. Firstly, we prove $|H|^2 < Ce^{-\varepsilon t/8}$ by contradiction. Suppose that $\overline{H}^2 \cdot e^{\varepsilon t/8}$ is unbounded. Then for any small positive number η , there exists a time τ , such that $\overline{H}^2(\tau) \cdot e^{\varepsilon\tau/8} > \Psi(\eta)/\eta^2$. By Theorem 5.3, we have $|\nabla H| < 2\eta^2\overline{H}^2$ on M_τ . By Lemma 6.2, we have $\underline{H}^2(\tau) > (1-\eta)\overline{H}^2(\tau) > \frac{1-\eta}{\eta^2}\Psi(\eta)e^{-\varepsilon\tau/8}$. This together with Theorem 5.3 yields $|\nabla H|^2 < (\eta|H|)^4 + \frac{\eta^4}{(1-\eta)^2}\underline{H}^4(\tau) \cdot e^{\varepsilon(\tau-t)/4}$. From the evolution equation of $|H|^2$, we have

$$(6.1) \quad \frac{\partial}{\partial t}|H|^2 > \Delta|H|^2 + \frac{1}{n}|H|^4 - \frac{1}{2n}\underline{H}^4(\tau) \cdot e^{\varepsilon(\tau-t)/4}.$$

Using the maximum principle, we get $|H|^2 \geq \underline{H}^2(\tau)$ if $t \geq \tau$. Then (6.1) yields $\frac{\partial}{\partial t}|H|^2 > \Delta|H|^2 + \frac{1}{2n}|H|^4$ for $t \geq \tau$. Hence, $|H|^2$ blows up in finite time. This contradicts the infinity of T . Therefore, we obtain $|H|^2 < Ce^{-\varepsilon t/8}$.

From Theorem 4.4, we have $|h|^2 = |\mathring{h}|^2 + \frac{1}{n}|H|^2 \leq Ce^{-\varepsilon t/8}$. Since $|h| \rightarrow 0$ as $t \rightarrow \infty$, M_t converges to a totally geodesic submanifold M_∞ as $t \rightarrow \infty$.

In the case of $q \geq n-4$, since $|h|^2 < \psi(|H|^2)$ is preserved, thus the flow can't converge to a totally geodesic submanifold. So the dimension and codimension of M_∞ satisfies $q < n-4$. From the fact that the totally geodesic submanifolds of $\mathbb{C}\mathbb{P}^m$ are totally real submanifolds $\mathbb{R}\mathbb{P}^n$ and Kähler submanifolds $\mathbb{C}\mathbb{P}^{n/2}$ [7], we see that M_∞ must be $\mathbb{C}\mathbb{P}^{n/2}$. \square

Combining the results of Theorems 6.3 and 6.4, we complete the proof of Theorem 1.1.

At last, we prove the convergence result under the weakly pinching condition for the mean curvature flow with $q \geq n-4 \geq 2$.

Proof of Theorem 1.2. From (3.16), we have

$$\begin{aligned} & \frac{1}{2} \left(\frac{\partial}{\partial t} - \Delta \right) \left(|\dot{h}|^2 - \dot{\psi} \right) \\ & \leq \left(|\dot{h}|^2 - \dot{\psi} \right) \left(|\dot{h}| + \dot{\psi} + n + 3 - 2\varepsilon + \left(\frac{1}{n} - \dot{\psi}' - \varepsilon \right) |H|^2 + 2\rho_2 \right) \\ & \quad + \dot{\psi} \left(\dot{\psi} + \frac{1}{n} |H|^2 + n + 3 \right) - \dot{\psi}' \cdot |H|^2 \left(\dot{\psi} + \frac{1}{n} |H|^2 + n \right). \end{aligned}$$

Using the strong maximum principle, we obtain either $|\dot{h}|^2 < \dot{\psi}$ for some $t > 0$, or $|\dot{h}|^2 = \dot{\psi}$ holds for all $t \in [0, t_0)$.

If $|\dot{h}|^2 < \dot{\psi}$ for some $t > 0$, then it follows from Theorem 1.1 that F_t converges to a round point.

If $|\dot{h}|^2 = \dot{\psi}$ holds for all $t \in [0, t_0)$, we have

$$\dot{\psi} \left(\dot{\psi} + \frac{1}{n} |H|^2 + n + 3 \right) - \dot{\psi}' \cdot |H|^2 \left(\dot{\psi} + \frac{1}{n} |H|^2 + n \right) = 0.$$

From Lemma 3.8 (iii), we get $|H| = 0$. Thus $|h|^2 = \psi(0) = 0$. Therefore, $F_t(M)$ is a totally geodesic submanifold for each $t \in [0, +\infty)$. \square

7. APPENDIX

For an odd integer $n \geq 3$, and a real number $\varepsilon \in [0, 1]$, we define a function $\varphi_\varepsilon : [0, +\infty) \rightarrow \mathbb{R}$ by

$$\varphi_\varepsilon(x) := d_\varepsilon + c_\varepsilon x - \sqrt{b^2 x^2 + 2abx + e},$$

where $a = 2\sqrt{(n^2 - 4n + 3)b}$, $b = \min \left\{ \frac{n-3}{4n-4}, \frac{2n-5}{n^2+n-2} \right\}$, $c_\varepsilon = b + \frac{1}{n-1+\varepsilon}$, $d_\varepsilon = 2 - 2\varepsilon + a$, $e = \sqrt{\varepsilon}$. We set $\varphi = \varphi_0$.

Lemma 7.1. *The function φ satisfies*

- (i) $\frac{x}{n-1} + 2 < \varphi(x) < \frac{x}{n-1} + n$,
- (ii) $\varphi(x) > \sqrt{2(n-3)}$.

Proof. If $n = 3$, then $a = b = 0$. We have $\varphi(x) = 2 + \frac{x}{n-1}$. It's easy to verify the inequalities above.

If $n \geq 5$, by direct computations, we get

$$\begin{aligned} \varphi'(x) &= c_0 - \frac{bx + a}{\sqrt{x^2 + 2ax/b}}, \\ \varphi''(x) &= \frac{a^2}{b(x^2 + 2ax/b)^{3/2}}. \end{aligned}$$

Since $\left(\varphi(x) - \frac{x}{n-1} \right)'' = \varphi''(x) > 0$ and $\lim_{x \rightarrow \infty} \varphi'(x) = \frac{1}{n-1}$, we have $\varphi'(x) < \frac{1}{n-1}$. Hence we get

$$2 = \lim_{x \rightarrow \infty} \left(\varphi(x) - \frac{x}{n-1} \right) < \varphi(x) - \frac{x}{n-1} \leq \varphi(0) = 2 + a < n.$$

If $n \geq 5$, we figure out that

$$\min_{x \geq 0} \varphi(x) = \varphi \left(\frac{ac_0}{b\sqrt{c_0^2 - b^2}} - \frac{a}{b} \right) = d_0 - \frac{ac_0}{b} + \frac{a}{b} \sqrt{c_0^2 - b^2}.$$

If $n = 5$, we have $\min_{x \geq 0} \varphi(x) = 4\sqrt{2} - 2$. If $n \geq 7$, we have $\min_{x \geq 0} \varphi(x) = 2 + 2\sqrt{\frac{n-3}{2n-5}} (\sqrt{5n-8} - \sqrt{n+2}) > \sqrt{2(n-3)}$. \square

Letting $\check{\varphi}_\varepsilon(x) = \varphi_\varepsilon(x) - \frac{x}{n}$, we have

Lemma 7.2. *There exists a positive constant ε_1 depending on n , such that for all $\varepsilon \in (0, \varepsilon_1)$, the function φ_ε satisfies the following inequalities*

- (i) $2x\check{\varphi}_\varepsilon''(x) + \check{\varphi}_\varepsilon'(x) < \frac{2(n-1)}{n(n+2)}$,
- (ii) $\check{\varphi}_\varepsilon(x)(\varphi_\varepsilon(x) - n + 3) - x\check{\varphi}_\varepsilon'(x)(\varphi_\varepsilon(x) + n + 3) < 2(n-1)$,
- (iii) $\check{\varphi}_\varepsilon(x) - x\check{\varphi}_\varepsilon'(x) > 1$.

Proof. If $n = 3$, then $a = b = 0$. We have $\varphi_\varepsilon(x) = 2 - 2\varepsilon + \frac{x}{2+\varepsilon} - \sqrt[4]{\varepsilon}$. It's easy to verify these inequalities.

If $n \geq 5$, by direct computations, we get

$$\begin{aligned}\check{\varphi}_\varepsilon'(x) &= c_\varepsilon - \frac{1}{n} - \frac{b^2x + ab}{\sqrt{b^2x^2 + 2abx + e}}, \\ \check{\varphi}_\varepsilon''(x) &= \frac{b^2(a^2 - e)}{(b^2x^2 + 2abx + e)^{3/2}}, \\ \check{\varphi}_\varepsilon'''(x) &= -\frac{3b^3(a^2 - e)(bx + a)}{(b^2x^2 + 2abx + e)^{5/2}}.\end{aligned}$$

Then we have

$$\begin{aligned}2x\check{\varphi}_\varepsilon''(x) + \check{\varphi}_\varepsilon'(x) &= c_\varepsilon - \frac{1}{n} - \frac{b^3x^2(bx + 3a) + eb(3bx + a)}{(b^2x^2 + 2abx + e)^{3/2}} \\ &< c_\varepsilon - \frac{1}{n} \\ &< \frac{2(n-1)}{n(n+2)}.\end{aligned}$$

This proves inequality (i).

Setting $f(x) = \check{\varphi}_\varepsilon(x)(\varphi_\varepsilon(x) - n + 3) - x\check{\varphi}_\varepsilon'(x)(\varphi_\varepsilon(x) + n + 3)$, we get

$$\begin{aligned}f(x) &= e + d_\varepsilon(d_\varepsilon + 3 - n) + (2 + ab + c_\varepsilon(d_\varepsilon - 2n))x \\ &\quad - (b^2x^2 + 2abx + e)^{-\frac{1}{2}} \times \\ &\quad [(3ab(d_\varepsilon + 1 - n) + c_\varepsilon e)x + b(ac_\varepsilon + b(d_\varepsilon - 2n))x^2 + e(2d_\varepsilon + 3 - n)].\end{aligned}$$

Then we figure out

$$\begin{aligned}\lim_{x \rightarrow +\infty} f(x) &= \frac{a^2c_\varepsilon}{b} + d_\varepsilon(d_\varepsilon + 3 - n) + a(n - 3 - 2d_\varepsilon) + e \left(1 - \frac{c_\varepsilon}{b}\right) \\ &= 2(n-1) + \frac{2\varepsilon(n^2 - 10n + 13 + 3\varepsilon(n-3) + 2\varepsilon^2)}{n-1+\varepsilon} + e \left(1 - \frac{c_\varepsilon}{b}\right) \\ &< 2(n-1),\end{aligned}$$

and

$$\begin{aligned}f'(x) &= 2 + ab + c_\varepsilon d_\varepsilon - 2c_\varepsilon n - (b^2x^2 + 2abx + e)^{-3/2} \times \\ &\quad [(3a^2b^2(d_\varepsilon + 1 - n) - 3be(b - ac_\varepsilon + bn))x \\ &\quad + b^2(ac_\varepsilon + bd_\varepsilon - 2nb)x^2(bx + 3a) \\ &\quad + c_\varepsilon e^2 + abe(d_\varepsilon - 2n)].\end{aligned}$$

Then we have

$$\lim_{x \rightarrow +\infty} f'(x) = 2 + ab + c_\varepsilon d_\varepsilon - 2c_\varepsilon n - ac_\varepsilon - bd_\varepsilon + 2bn = 0.$$

Furthermore, we get

$$f''(x) = [b(2b + bd_\varepsilon - ac_\varepsilon)x^2 + (ab(1 - n + d_\varepsilon) - c_\varepsilon e)x - (n + 1)e] \frac{3b^2(a^2 - e)}{(b^2x^2 + 2abx + e)^{5/2}}.$$

From $b \leq \frac{n-3}{4n-4}$, we obtain

$$2b + bd_\varepsilon - ac_\varepsilon = (4 - 2\varepsilon)b - \frac{2\sqrt{(n^2 - 4n + 3)b}}{n - 1 + \varepsilon} < 0,$$

and

$$1 - n + d_\varepsilon = 3 - n + 2\sqrt{(n^2 - 4n + 3)b} - 2\varepsilon < 0.$$

So, we get $f''(x) < 0$. Then we have $f'(x) > 0$. From this we deduce that

$$f(x) < \lim_{x \rightarrow +\infty} f(x) < 2(n - 1).$$

Thus, inequality (ii) is proved.

We have

$$\begin{aligned} \hat{\varphi}_\varepsilon(x) - x\hat{\varphi}'_\varepsilon(x) &= d_\varepsilon - \frac{abx + e}{\sqrt{b^2x^2 + 2abx + e}} \\ &> d_\varepsilon - \frac{abx}{\sqrt{b^2x^2}} - \frac{e}{\sqrt{e}} \\ &= 2 - 2\varepsilon - \sqrt[4]{\varepsilon}. \end{aligned}$$

This implies inequality (iii). \square

For each integer $n \geq 6$, we define the function $\psi : [0, +\infty) \rightarrow \mathbb{R}$ by

$$\psi(x) := \nu + \kappa x - \sqrt{\lambda^2 x^2 + 2\lambda\mu x + \nu^2},$$

where $\kappa = \lambda + \frac{1}{n-1}$, $\lambda = \frac{3}{n^3 - 4n^2 + 3}$, $\mu = \nu + \frac{3}{n}$, $\nu = \frac{9}{n^2 - 3n - 3}$.

Lemma 7.3. *The function ψ satisfies the following inequalities.*

- (i) $\frac{1}{n} \leq \psi'(x) < \frac{1}{n-1}$, $\frac{x}{n} \leq \psi(x) \leq \frac{x}{n-1}$, and the equalities hold if and only $x = 0$,
- (ii) $\max_{x \geq 0} (2x\psi''(x) + \psi'(x)) < \frac{3}{n+8}$,
- (iii) $3\psi(x) - \frac{3x}{n} + (\psi(x) - x\psi'(x))(\psi(x) + n) \leq 0$, and the equalities hold if and only $x = 0$,
- (iv) $0 \leq x\psi'(x) - \psi(x) < 2$.

Proof. Taking derivatives, we get

$$\begin{aligned} \psi'(x) &= \kappa - \frac{\lambda^2 x + \lambda\mu}{\sqrt{\lambda^2 x^2 + 2\lambda\mu x + \nu^2}}, \\ \psi''(x) &= \frac{\lambda^2(\mu^2 - \nu^2)}{(\lambda^2 x^2 + 2\lambda\mu x + \nu^2)^{3/2}}, \\ \psi'''(x) &= -\frac{3\lambda^3(\mu^2 - \nu^2)(\lambda x + \mu)}{(\lambda^2 x^2 + 2\lambda\mu x + \nu^2)^{5/2}}. \end{aligned}$$

(i) We have $\psi'(0) = \kappa - \frac{\lambda\mu}{\nu} = \frac{1}{n}$ and $\lim_{x \rightarrow +\infty} \psi'(x) = \kappa - \lambda = \frac{1}{n-1}$. Since $\psi''(x) > 0$, we get $\frac{1}{n} < \psi'(x) < \frac{1}{n-1}$ for $x > 0$. Thus we obtain $\frac{x}{n} < \psi(x) < \frac{x}{n-1}$ for $x > 0$.

(ii) Letting $g(x) = 2x\psi''(x) + \psi'(x)$, we have

$$g(x) = \kappa - \frac{\lambda^4 x^3 + 3\lambda^3 \mu x^2 + 3\lambda^2 \nu^2 x + \lambda \mu \nu^2}{(\lambda^2 x^2 + 2\lambda \mu x + \nu^2)^{3/2}},$$

and

$$g'(x) = 2x\psi'''(x) + 3\psi''(x) = \frac{3\lambda^2(\mu^2 - \nu^2)(\nu^2 - \lambda^2 x^2)}{(\lambda^2 x^2 + 2\lambda \mu x + \nu^2)^{5/2}}.$$

Thus we get

$$\begin{aligned} \max_{x \geq 0} g(x) &= g\left(\frac{\nu}{\lambda}\right) = \kappa - \lambda \sqrt{\frac{2\nu}{\mu + \nu}} \\ &= \frac{n(n-3) - 3\sqrt{\frac{6n}{n^2+3n-3}}}{n^3 - 4n^2 + 3} \\ &< \frac{3}{n+8}. \end{aligned}$$

(iii) We have

$$\begin{aligned} &3\psi(x) - \frac{3x}{n} + (\psi(x) - x\psi'(x))(\psi(x) + n) \\ &= \left[\nu(n+3+2\nu) + \left(\lambda\mu + 3\kappa + \kappa\nu - \frac{3}{n}\right)x \right] \\ &\quad - \frac{\lambda(\kappa\mu + 3\lambda + \lambda\nu)x^2 + (\kappa\nu^2 + \lambda\mu(n+6+3\nu))x + \nu^2(n+3+2\nu)}{\sqrt{\lambda^2 x^2 + 2\lambda\mu x + \nu^2}}. \end{aligned}$$

Set

$$\begin{aligned} h(x) &:= \left[\nu(n+3+2\nu) + \left(\lambda\mu + 3\kappa + \kappa\nu - \frac{3}{n}\right)x \right]^2 (\lambda^2 x^2 + 2\lambda\mu x + \nu^2) \\ &\quad - [\lambda(\kappa\mu + 3\lambda + \lambda\nu)x^2 + (\kappa\nu^2 + \lambda\mu(n+6+3\nu))x + \nu^2(n+3+2\nu)]^2. \end{aligned}$$

Now we need to prove $h(x) \leq 0$. Since $\kappa = \lambda + \frac{1}{n-1}$, $\mu = \nu + \frac{3}{n}$, we have

$$\lambda\mu + 3\kappa + \kappa\nu - \frac{3}{n} = \kappa\mu + 3\lambda + \lambda\nu.$$

Putting $A = \kappa\mu + 3\lambda + \lambda\nu$, $B = n+3+2\nu$, $C = \kappa\nu^2 + \lambda\mu(n+6+3\nu)$, we get

$$\begin{aligned} h(x) &= (\nu B + Ax)^2 (\lambda^2 x^2 + 2\lambda\mu x + \nu^2) - (\lambda Ax^2 + Cx + \nu^2 B)^2 \\ &= 2\lambda A(\mu A + \lambda\nu B - C)x^3 \\ &\quad + [\nu^2(A - \lambda B)^2 + 4\lambda\mu\nu AB - C^2]x^2 \\ &\quad + 2\nu^2 B(\nu A + \lambda\mu B - C)x. \end{aligned}$$

By the definitions of $\kappa, \lambda, \mu, \nu$, we have

$$\mu A + \lambda\nu B = \nu A + \lambda\mu B = C,$$

and

$$\nu^2(A - \lambda B)^2 + 4\lambda\mu\nu AB - C^2 = -\frac{81(n^3 - 12n + 9)^2}{n^2(n-1)^2(n^2 - 3n - 3)^4} < 0.$$

Thus, inequality (iii) is proved.

(iv) It follows from inequalities (i) and (iii) that $\psi(x) - x\psi'(x) \leq 0$. By direct computations, we get

$$\begin{aligned} x\psi'(x) - \psi(x) &= -\nu + \frac{\lambda\mu x + \nu^2}{\sqrt{\lambda^2 x^2 + 2\lambda\mu x + \nu^2}} \\ &\leq -\nu + \frac{\lambda\mu x}{\sqrt{\lambda^2 x^2}} + \frac{\nu^2}{\sqrt{\nu^2}} \\ &= \mu < 2. \end{aligned}$$

□

REFERENCES

- [1] B. ANDREWS, *Positively curved surfaces in the three-sphere*, Proc. ICM 2002, Vol.2, Higher Education Press, Beijing, 2002, 221-230.
- [2] B. ANDREWS and C. BAKER, *Mean curvature flow of pinched submanifolds to spheres*, J. Differential Geom. **85**(2010), 357-395.
- [3] C. BAKER, *The mean curvature flow of submanifolds of high codimension*, arXiv:1104.4409, 2011.
- [4] C. BÖHM and B. WILKING, *Manifolds with positive curvature are space forms*, Ann. of Math. **167**(2008), 1079-1097.
- [5] S. BRENDLE, *A general convergence result for the Ricci flow in higher dimensions*, Duke Math. J. **145**(2008), 585-601.
- [6] S. BRENDLE and R. SCHOEN, *Manifolds with 1/4-pinched curvature are space forms*, J. Amer. Math. Soc. **22**(2009), 287-307.
- [7] B. Y. CHEN and K. OGIUE, *Two theorems on Kaehler manifolds*, Michigan Math. J. **21**(1974), 225-229.
- [8] B. Y. CHEN and M. OKUMURA, *Scalar curvature, inequality and submanifold*, Proc. Amer. Math. Soc. **38**(1973), 605-608.
- [9] Q. M. CHENG and K. NONAKA, *Complete submanifolds in Euclidean spaces with parallel mean curvature vector*, Manuscripta Math. **105**(2001), 353-366.
- [10] S. S. CHERN, M. DO CARMO and S. KOBAYASHI, *Minimal submanifolds of a sphere with second fundamental form of constant length*, In: Functional Analysis and Related Fields, F. E. Browder (ed.), Springer, New York, 1970, 59-75.
- [11] J. R. GU and H. W. XU, *The sphere theorems for manifolds with positive scalar curvature*, J. Differential Geom. **92**(2012), 507-545.
- [12] R. HAMILTON, *Three manifolds with positive Ricci curvature*, J. Differential Geom. **17**(1982), 255-306.
- [13] D. HOFFMAN and J. SPRUCK, *Sobolev and isoperimetric inequalities for Riemannian submanifolds*, Comm. Pure Appl. Math. **27**(1974), 715-727; **28**(1975), 765-766.
- [14] G. HUISKEN, *Flow by mean curvature of convex surfaces into spheres*, J. Differential Geom. **20**(1984), 237-266.
- [15] G. HUISKEN, *Contracting convex hypersurfaces in Riemannian manifolds by their mean curvature*, Invent. Math. **84**(1986), 463-480.
- [16] G. HUISKEN, *Deforming hypersurfaces of the sphere by their mean curvature*, Math. Z. **195**(1987), 205-219.
- [17] L. LEI and H. W. XU, *An optimal convergence theorem for mean curvature flow of arbitrary codimension in hyperbolic spaces*, arXiv:1503.06747, 2015.
- [18] L. LEI and H. W. XU, *A new version of Huisken's convergence theorem for mean curvature flow in spheres*, arXiv:1505.07217, 2015.
- [19] L. LEI and H. W. XU, *Mean curvature flow of arbitrary codimension in spheres and sharp differentiable sphere theorem*, arXiv:1506.06371v2, 2015.
- [20] A. M. LI and J. M. LI, *An intrinsic rigidity theorem for minimal submanifolds in a sphere*, Arch. Math. **58**(1992), 582-594.
- [21] H. Z. LI and X. F. WANG, *A differentiable sphere theorem for compact Lagrangian submanifolds in complex Euclidean space and complex projective space*, Comm. Anal. Geom. **22**(2014), 269-288.

- [22] K. F. LIU, H. W. XU, F. YE and E. T. ZHAO, *Mean curvature flow of higher codimension in hyperbolic spaces*, Comm. Anal. Geom. **21**(2013), 651-669.
- [23] K. F. LIU, H. W. XU and E. T. ZHAO, *Mean curvature flow of higher codimension in Riemannian manifolds*, arXiv:1204.0107, 2012.
- [24] K. F. LIU, H. W. XU and E. T. ZHAO, *Some recent progress on mean curvature flow of arbitrary codimension*, to appear in Proceedings of the Sixth International Congress of Chinese Mathematicians, Taipei, July 14-19, 2013, AMS/IP, Studies in Advanced Math.
- [25] M. OKUMURA, *Submanifolds and a pinching problem on the second fundamental tensors*, Trans. Amer. Math. Soc. **178** (1973), 285-291.
- [26] M. OKUMURA, *Hypersurfaces and a pinching problem on the second fundamental tensor*, Amer. J. Math. **96** (1974), 207-213.
- [27] G. PIPOLI and C. SINISTRARI, *Mean curvature flow of pinched submanifolds of $\mathbb{C}\mathbb{P}^n$* , arXiv:1502.00519v3, 2015, to appear in Comm. Anal. Geom.
- [28] W. SANTOS, *Submanifolds with parallel mean curvature vector in spheres*, Tohoku Math. J. **46**(1994), 403-415.
- [29] K. SHIOHAMA, *Sphere theorems*, Handbook of Differential Geometry, Vol. **1**, F.J.E. Dillen and L.C.A. Verstraelen (eds.), Elsevier Science B.V. Amsterdam, 2000.
- [30] K. SHIOHAMA and H. W. XU, *The topological sphere theorem for complete submanifolds*, Compositio Math. **107**(1997), 221-232.
- [31] K. SHIOHAMA and H. W. XU, *A general rigidity theorem for complete submanifolds*, Nagoya Math. J. **150**(1998), 105-134.
- [32] J. SIMONS, *Minimal varieties in riemannian manifolds*, Ann. of Math. **88**(1968), 62-105.
- [33] M. T. WANG, *Lectures on mean curvature flows in higher codimensions*, Handbook of Geometric Analysis, Adv. Lect. Math. (ALM), Vol. **1**, No. 1, International Press, Somerville, MA, 2008, 525-543.
- [34] H. W. XU, *Pinching theorems, global pinching theorems, and eigenvalues for Riemannian submanifolds*, Ph.D. dissertation, Fudan University, 1990.
- [35] H. W. XU, *A rigidity theorem for submanifolds with parallel mean curvature in a sphere*, Arch. Math. **61**(1993), 489-496.
- [36] H. W. XU, *Rigidity of submanifolds with parallel mean curvature in space forms*, preprint, 1993.
- [37] H. W. XU and J. R. GU, *An optimal differentiable sphere theorem for complete manifolds*, Math. Res. Lett. **17**(2010), 1111-1124.
- [38] H. W. XU and J. R. GU, *Geometric, topological and differentiable rigidity of submanifolds in space forms*, Geom. Funct. Anal. **23**(2013), 1684-1703.
- [39] H. W. XU and E. T. ZHAO, *Topological and differentiable sphere theorems for complete submanifolds*, Comm. Anal. Geom. **17**(2009), 565-585.
- [40] S. T. YAU, *Submanifolds with constant mean curvature I, II*, Amer. J. Math. **96** (1974), 346-366; **97** (1975), 76-100.
- [41] X. P. ZHU, *Lectures on mean curvature flows*, Studies in Advanced Mathematics Vol. **32**, International Press, Somerville, MA, 2002.

CENTER OF MATHEMATICAL SCIENCES, ZHEJIANG UNIVERSITY, HANGZHOU 310027, CHINA
E-mail address: lei-li@zju.edu.cn; xuhw@cms.zju.edu.cn