Necessary and sufficient criterion for k-separability of N-qubit noisy GHZ states

Xiao-yu Chen^a, Li-zhen Jiang^a Zhu-an Xu^b

^aCollege of Information and Electronic Engineering, Zhejiang Gongshang University, Hangzhou, Zhejiang 310018, China ^bDepartment of Physics, Zhejiang University, Hangzhou, Zhejiang 310027, China

Abstract

A Multipartite entangled state has many different kinds of entanglement specified by the number of partitions. The most essential example of multipartite entanglement is the entanglement of multi-qubit Greenberger-Horne-Zeilinger (GHZ) state in white noise. We explicitly construct the entanglement witnesses for these states with stabilizer generators of the GHZ states. For a N qubit GHZ state in white noise, we demonstrate the necessary and sufficient criterion of separability when it is divided into k parties with $N \leq 2k-1$ for arbitrary N and k. The criterion covers more than a half of all kinds of partial entanglement for N-qubit GHZ states in white noise. For the rest of multipartite entanglement problems, we present a method to obtain the sufficient conditions of separability. As an application, we consider N qubit GHZ state as a codeword of the degenerate quantum code passing through depolarizing channel. We find that the output state is neither genuinely entangled nor fully separable when the quantum channel capacity reduces from positive to zero.

PACS number(s): 03.67.Mn; 03.65.Ud

1 Introduction

Entanglement in multipartite systems is a key resource for quantum information and communication protocols[1, 2]. In experiments, various multipartite entangled states have been generated[3, 4, 5, 6, 7, 8]. The question whether or not an experimentally generated multipartite state is (partially) entangled has become a highly relevant topic for quantum information theory. Although an enormous amount of work has been devoted to detect multipartite entanglement [9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20], we are still very far away from the characterization of multipartite entanglement. The experimental detection of entanglement is typically done via the construction of proper entanglement witnesses [2, 19, 21], and multipartite witnesses have been considered [24, 25, 26, 27].

Among all the multipartite entangled states, the

Greenberger-Horne-Zeilinger (GHZ) states are the simplest ones. The GHZ entanglement, originally introduced to explore the extreme violation of local realism against quantum mechanics, is an important resource for multipartite quantum communication tasks such as quantum cryptographic conferencing (QCC) [28, 29, 30], quantum secret sharing (QSS) [31, 32, 33, 34] and third-man quantum cryptography [35]. Typically, the experimental preparations of multipartite entangled states [3, 4, 5] are the N qubit GHZ states. The imperfection and noise in the preparations are usually described by white noise. Most probably, an experimental prepared multipartite state is the mixture of N qubit GHZ state with white noise (noisy GHZ state, also known as generalized Werner state [36] or Werner-Popescu state [37, 38]). Meanwhile the noisy GHZ state describes the output state of the GHZ state passing through a depolarizing channel.

Suppose there is a composed Hilbert space $\mathcal{H} = \mathcal{H}_1 \otimes \cdots \otimes \mathcal{H}_n$. Consider a partition $\mathcal{I} = {\mathcal{I}_1, ..., \mathcal{I}_k}$ of the index set $\mathcal{J} = {1, ..., n}$. A quantum state $\sigma_{\mathcal{I}}$ is called separable for the given partition \mathcal{I} , if it can be written as a classical mixture of product states:

$$\sigma_{\mathcal{I}} = \sum_{i} q_{i} |\psi_{\mathcal{I}_{1}}^{(i)}\rangle \langle \psi_{\mathcal{I}_{1}}^{(i)}| \otimes \cdots \otimes |\psi_{\mathcal{I}_{k}}^{(i)}\rangle \langle \psi_{\mathcal{I}_{k}}^{(i)}|, \qquad (1)$$

with q_i being a classical probability distribution, $|\psi_{\mathcal{I}_j}^{(i)}\rangle$ is a pure state of subset \mathcal{I}_j . A state σ is called *k*-separable if it can be written to be

$$\sigma = \sum_{\mathcal{I}:|\mathcal{I}|=k} q'_{\mathcal{I}} \sigma_{\mathcal{I}}.$$
 (2)

Where $|\mathcal{I}|$ is the number of elements in the set \mathcal{I} , $q'_{\mathcal{I}}$ is a classical probability distribution. The summation is over all possible k partite partitions. If a quantum state cannot be written in the form of Eq.(2), it is referred to as k-inseparable. A 2-inseparable (not biseparable) state is also called genuinely entangled. A N-inseparable (not fully separable) state is an entangled state.

Although the biseparable (k = 2) condition and the fully separable (k = N) condition for a N qubit noisy GHZ state are known [11, 39, 40], the conditions for the k-separability

^{*}Email:xychen@zjgsu.edu.cn

with 2 < k < N almost remain to be uncovered for the state.

The partition of an N qubit noisy GHZ state is greatly simplified by the symmetry of the state. In fact we just count the number of qubits in a party without considering which qubits are in the party. Let the number of qubits in subset \mathcal{I}_j be n_j . We may alternatively denote the partition as $n_1|n_2|n_3| \dots |n_j| \dots |n_k$. Then $\sum_{j=1}^k n_j = N$.

We will show that a properly chosen entanglement witness constructed with stabilizer generators of GHZ state can detect the k-separability of a noisy GHZ state. The paper is organized as follows: In section 2, we describe the entanglement witnesses for noisy GHZ states. In section 3 and section 4, the necessary and sufficient conditions are demonstrated for the k separability of N qubit noisy GHZ states when $N \leq 2k-1$. We also give the witnesses explicitly in section 3. In section 5, the sufficient conditions of all the other separabilities for $N \leq 12$ are shown. In section 6, we discuss the application of our findings to GHZ state in depolarizing channel. The conclusions are drown in section 7. Several lemmas or theorem are proven in the appendix.

2 Entanglement Witnesses of Noisy GHZ states

A witness of multipartite entanglement (more strictly kinseparability) is a Hermitian operator \hat{W} with $tr(\rho_s \hat{W}) \geq$ 0 for all k-separable state ρ_s , and $tr(\rho \hat{W}) < 0$ for at least one state ρ . A witness \hat{W} is optimal if $tr(\rho_s \hat{W}) = 0$ is achieved for some k-separable state ρ_s . Let \hat{M} be a Hermitian operator, the optimal witness can be produced as $\hat{W} = \Lambda \mathbb{I} - \hat{M}$, where \mathbb{I} is the identity operator of the quantum system, and

$$\Lambda = \max_{\rho_s} tr(\rho_s \hat{M}),\tag{3}$$

The method leading to a k-separable criterion is to first choose an Hermitian operator \hat{M} , then find the Λ with respect to all the k-separable state ρ_s . If there is a state ρ with $tr(\rho \hat{M}) > \Lambda$, then the state ρ is called k-inseparable. Let $\mathcal{L} = \frac{\Lambda}{tr(\rho \hat{M})}$, the necessary criterion of k-separability is

$$\mathcal{L} \le 1. \tag{4}$$

The criterion may not be efficient by the ad hoc nature of choosing \hat{M} . For a give state ρ , we may modify \hat{M} to decrease \mathcal{L} such that the criterion is more efficient. For a noisy N qubit GHZ state, we propose a proper \hat{M} then prove the necessary criterion to be also sufficient.

A noisy N qubit GHZ state is

$$\rho_{ghzN} = p \left| GHZ_N \right\rangle \left\langle GHZ_N \right| + \frac{(1-p)}{2^N} \mathbb{I}.$$
 (5)

Where $|GHZ_N\rangle = \frac{1}{\sqrt{2}}(|0\rangle^{\otimes N} + |1\rangle^{\otimes N})$. In the following, we will denote $|\chi\rangle \langle \chi|$ as $|\chi\rangle \langle \cdot|$ for short. The stabilizer of $|GHZ_N\rangle$ is an operator Abel group created by generators $\{\hat{K}_1, \ldots, \hat{K}_N\}$ such that $\hat{K} |GHZ_N\rangle = |GHZ_N\rangle$ for any stabilizer operator $\hat{K} = \hat{K}_1^{j_1} \cdots \hat{K}_N^{j_N}$ with $j_i = 0, 1$. The generators are

$$\begin{array}{rcl}
\dot{K}_{1} &=& \sigma_{1} \otimes \sigma_{1} \otimes \cdots \sigma_{1}, \\
\dot{K}_{2} &=& \sigma_{3} \otimes \sigma_{3} \otimes I \otimes \cdots I, \\
\dot{K}_{3} &=& \sigma_{3} \otimes I \otimes \sigma_{3} \otimes I \otimes \cdots I, \\
&& & & & \\
\dot{K}_{N} &=& \sigma_{3} \otimes I \otimes \cdots I \otimes \sigma_{3}, \\
\end{array}$$
(6)

with $\sigma_1, \sigma_2, \sigma_3$ being Pauli matrices and $I = \sigma_0$ is the 2×2 identity matrix. The projector onto the GHZ state has a direct representation in terms of the corresponding stabilizer, namely,

$$|GHZ_N\rangle \langle GHZ_N| = \frac{1}{2^N} \sum_{\hat{K} \in stabilizer} \hat{K}$$
(7)

We have

$$tr(\rho_{ghzN}\hat{K}) = \begin{cases} 1, \text{ for } \hat{K} = \mathbb{I} \\ p, \text{ otherwise.} \end{cases}$$
(8)

for a stabilizer operator \hat{K} .

Since ρ_{ghzN} is a linear mixture of stabilizer operators, we may suppose the Hermitian operator \hat{M} to be a linear combination of stabilizer operators instead of all elements of Pauli group. The most general form of a Hermitian operator \hat{M} is

$$\hat{M} = \sum_{j_1,\dots,j_N=0}^{1} M_{j_1 j_2 \dots j_N} \hat{K}_1^{j_1} \cdots \hat{K}_N^{j_N}.$$
 (9)

An observation of (6) shows that the generators $K_2, ..., K_n$ constitute a subgroup of the stabilizer. The subgroup is responsible for the diagonal entries of ρ_{ghzN} . The coset (the product of K_1 with the subgroup) is responsible for the anti-diagonal entries of ρ_{ghzN} . Due to the symmetry of ρ_{ghzN} , it is reasonable that the Hermitian operator (thus the witness) also possesses the symmetry. We then assume $M_{j_1j_2...j_N}$ to be a function of $|\mathbf{j}|$ instead of \mathbf{j} , where $\mathbf{j} = (j_2, ..., j_N)$. Furthermore, we assume $\hat{M} = \hat{M}_D + \hat{M}_A$, with

$$\hat{M}_D = \sum_{\mathbf{j}} M_{\lceil |\mathbf{j}|/2\rceil} \hat{K}_2^{j_2} \cdots \hat{K}_N^{j_N}, \qquad (10)$$

$$\hat{M}_A = \hat{K}_1 \sum_{\mathbf{j}} \hat{K}_2^{j_2} \cdots \hat{K}_N^{j_N}, \qquad (11)$$

The reasons are as follows. All coefficients $M_{1j_2...j_N}$ in \hat{M}_A are set to be equal (without loss of generality, they are set to be 1), since in computational basis we have

 $\hat{M}_A = 2^{N-1} (|0\rangle^{\otimes N} \langle 1|^{\otimes N} + |1\rangle^{\otimes N} \langle 0|^{\otimes N})$ due to (6) and (7). Such a choice of M_A is reasonable since it is proportional to the anti-diagonal part of the state ρ_{ghzN} . All the stabilizer operators in \hat{M}_D contribute to diagonal part of the Hermitian operator \hat{M} (thus the witness operator) in computational basis. The coefficients $M_{\lceil |\mathbf{j}|/2 \rceil}$ in M_D are so chosen such that the stabilizer operators with equal number of σ_3 have the same coefficient. For example, all the generators K_i (i = 2, ..., N) have two σ_3 in their tensor product expressions, the coefficient for these stabilizer operators is M_1 . The stabilizer operators with two generators $K_i K_j$ $(i, j = 2, ..., N; i \neq j)$ also have two σ_3 in their tensor product expressions, the coefficient for them is M_1 too. The coefficients in M_D are so chosen that we keep the qubit exchange symmetry for the Hermitian operator. We also assume $M_0 = 0$ since a nonzero M_0 only leads to a displacement of Λ .

Notice that $tr(\rho_{ghzN}\hat{M}_A) = 2^{N-1}p$. There are $C_N^{2i} = \frac{N!}{(N-2i)!(2i)!}$ different tensor products $\sigma_3^{j_1} \otimes \ldots \otimes \sigma_3^{j_N}$ subject to $\sum_l j_l = 2i, j_l \in \{0, 1\}$, each of the tensor products is a stabilizer operator. We then have

$$tr(\rho_{ghzN}\hat{M}) = p(\sum_{i=1}^{\lfloor N/2 \rfloor} M_i C_N^{2i} + 2^{N-1}), \qquad (12)$$

From (4), the necessary condition for the k-separability of ρ_{ghzN} then is

$$p \le \frac{\Lambda}{\left(\sum_{i=1}^{\lfloor N/2 \rfloor} M_i C_N^{2i} + 2^{N-1}\right)}.$$
(13)

In convention, Λ is with respect to k-partition.

3 Necessary condition of $k \ (k \ge \frac{N+1}{2})$ separability

A k-partite partition $n_1|n_2|n_3| \dots |n_l| \dots |n_k$ splits the N qubit system into k parties. There are many k-partite partitions since the number of qubits in each party varies from 1 to at most N - k + 1. The notation of the partition can be shorten as $n_1^2|n_3| \dots |n_l| \dots |n_k$ if $n_2 = n_1$. So alternatively, we may denote the k-partite partition as $1^{N_1}|2^{N_2}|3^{N_3}|\dots|m^{N_m}$ with $k = \sum_{i=1}^m N_i$, $N = \sum_{i=1}^m iN_i$. The number of i qubit parties is N_i in the partition. One of the main findings of this paper is that the coefficients M_i of the Hermitian operator M should form an arithmetic progression.

$$M_i = \frac{4i - N}{N_1},\tag{14}$$

We stress that N_1 is the number of single qubit parties in the partition. We start with a special k-partite partition $1^{k-1}|(N-k+1)$ for the necessary condition of kseparability.

3.1 Partition 1|1|...|1|L

To obtain Λ , the maximal mean of the Hermitian operator \hat{M} with respect to all k-separable states, we only need to consider pure k-separable states. For the partition $1^{N_1}|L$, we have $N_1 = k - 1$, L = N - k + 1, there is a k-separable pure state

$$\rho_s = \bigotimes_{i=1}^{N_1} \varrho_i \otimes \varrho_{N_1+1,\dots,N},\tag{15}$$

where $\rho_i = \frac{1}{2}(I + x_i\sigma_1 + y_i\sigma_2 + z_i\sigma_3)$, $(i = 1, \ldots, N_1)$ is the pure state of the *ith* qubit, with $x_i^2 + y_i^2 + z_i^2 = 1$; $\rho_{N_1+1,\ldots,N} \equiv |\psi\rangle \langle \psi|$ is the pure state for the last *L* qubits. Then

$$tr(\rho_s \hat{M}_D) = \sum_{\mathbf{j}} M_{\lceil |\mathbf{j}|/2\rceil} z_1^{j'} \prod_{i=2}^{N_1} z_i^{j_i} \\ \times \langle \psi | \bigotimes_{l=N_1+1}^N \sigma_3^{j_l} |\psi \rangle, \qquad (16)$$

where $j' = \mod(|\mathbf{j}|, 2)$. We also have

$$tr(\rho_{s}\hat{M}_{A}) = \sum_{\mathbf{j}} (-i)^{|\mathbf{j}|+j'} x_{1}^{1-j'} y_{1}^{j'} \prod_{i=2}^{N_{1}} x_{i}^{1-j_{i}} y_{i}^{j_{i}}$$
$$\langle \psi | \bigotimes_{l=N_{1}+1}^{N} \sigma_{1}^{1-j_{l}} \sigma_{2}^{j_{l}} | \psi \rangle .$$
(17)

We may write $tr(\rho_s \hat{M}) = \langle \psi | \mathcal{M} | \psi \rangle$, where \mathcal{M} is a $2^L \times 2^L$ matrix. The maximum of $tr(\rho_s \hat{M})$ then is equal to the maximal eigenvalue of \mathcal{M} . Denote the diagonal elements of matrix \mathcal{M} as $m_{\mathbf{i},\mathbf{i}}$, where $\mathbf{i} = (i_1, i_2, \ldots, i_L)$ is a binary string and we denote the weight of \mathbf{i} as $|\mathbf{i}|$. Then $m_{\mathbf{i},\mathbf{i}} =$ $\Gamma_{|\mathbf{i}|}$ with $\Gamma_{|\mathbf{i}|}$ defined in (48) of the appendix. Let $\mathcal{M}_{\mathbf{i}} =$ $m_{\mathbf{i},\mathbf{i}}|\mathbf{i}\rangle\langle\mathbf{i}|$, then we have

Lemma 1 All the non-diagonal entries of matrix $|0^{\otimes L}\rangle\langle 1^{\otimes L}|$ \mathcal{M} is nullified except the entries of $|1^{\otimes L}\rangle\langle 0^{\otimes L}|,$ and namely, \mathcal{M} isblock di- $\mathcal{M} = \mathcal{M}_0 \bigoplus_{i \neq 0, i \neq 1} \mathcal{M}_i,$ agonalized aswhere $\mathbf{0} = (0, \dots, 0), \mathbf{1} = (1, \dots, 1).$ The submatrix \mathcal{M}_0 in the computational basis $|0^{\otimes L}\rangle$ and $|1^{\otimes L}\rangle$ is

$$\mathcal{M}_0 = \left[\begin{array}{cc} \Gamma_0 & 2^{L-1}(c-id) \\ 2^{L-1}(c+id) & \Gamma_L \end{array} \right]$$
(18)

where $c = \prod_{i=1}^{N_1} \sin \theta_i \cos \varphi$, $d = \prod_{i=1}^{N_1} \sin \theta_i \sin \varphi$, with $\varphi = \sum_{j=1}^{N_1} \varphi_j$.

Moreover, we have

Lemma 2 The maximal eigenvalue of \mathcal{M}_0 is

$$\Lambda_0 = \frac{N}{N_1}.\tag{19}$$

Using (14), the necessary criterion (13) is simplified to

$$p \le \frac{\Lambda}{\frac{N}{N_1} + 2^{N-1}}.$$
(20)

Theorem 1 The maximal eigenvalue of \mathcal{M} is $\Lambda = \frac{N}{N_1}$. The necessary condition of separability for ρ_{ahzN} is

$$p \le [1 + 2^{N-1}N_1/N]^{-1} \tag{21}$$

with respect to the partition $1^{N_1}|L$.

The proofs of Lemma 1, Lemma 2 and Theorem 1 are shown in the appendix.

3.2 Further splits of the last L qubits

We consider further splits of the last L qubits into two parties. Denote the partition after split as $1^{N_1}|\overline{L} = 1^{N_1}|l|L-l$ with $N = N_1 + L$, $k = N_1 + 2$. Keep in mind that we have exhausted all the single qubit parties into 1^{N_1} , the split of the last L qubits do not produce new single qubit partite. The smallest piece from the further split of the last L qubits is a two qubit partite, namely $l \ge 2, L - l \ge 2$. On the other hand, we have noticed that the maximum of $tr(\rho_s M)$ do not increase by the split of the last L qubits. This is due to the fact that when we split the pure state of the last L qubits $|\psi\rangle$ into a product of pure states, the matrix \mathcal{M} does not change since we only change the last part of ρ_s and do not change the state of the first N_1 qubits and the operator \hat{M} . The maximal eigenvalue of \mathcal{M} is achieved when $|\psi\rangle$ is the corresponding eigenvector, namely $\langle \psi | \mathcal{M} | \psi \rangle \geq \langle \psi' | \mathcal{M} | \psi' \rangle$ for all the other state $| \psi' \rangle$ including the product state produced by the split of the last L qubits. We conclude that the maximum of $tr(\rho_s \hat{M})$ for partition $1^{N_1} | \overline{L}$ can not exceed $\frac{N}{N_1}$.

Next we should prove that the maximal eigenvalue $\frac{N}{N_1}$ of \mathcal{M} is achievable by the state with partition $1^{N_1}|\overline{L}$. Let $|\psi'\rangle = |\psi_l\rangle|\psi_{L-l}\rangle$ be the product state of the partition \overline{L} , with rather generic states $|\psi_l\rangle = \alpha_1|0^{\otimes l}\rangle + \beta_1|1^{\otimes l}\rangle$ and $|\psi_{L-l}\rangle = \alpha_2|0^{\otimes L-l}\rangle + \beta_2|1^{\otimes L-l}\rangle$ subject to the constrain that $\alpha_1\alpha_2|0^{\otimes L}\rangle + \beta_1\beta_2|1^{\otimes L}\rangle \equiv |\psi_a\rangle \sim |\psi\rangle$ is the unnormalized eigenvector corresponding to the largest eigenvalue of \mathcal{M} . Let $|\psi_b\rangle = \alpha_1\beta_2|0^{\otimes l}\rangle|1^{\otimes l-l}\rangle + \beta_1\alpha_2|1^{\otimes l}\rangle|0^{\otimes L-l}\rangle$, then $|\psi'\rangle = |\psi_a\rangle + |\psi_b\rangle$. Then we have

$$\langle \psi' | \mathcal{M} | \psi' \rangle = \langle \psi' | \mathcal{M}_0 \oplus \mathcal{M}_{0^l 1^{L-l}} \oplus \mathcal{M}_{1^l 0^{L-l}} | \psi' \rangle$$

$$= \langle \psi_a | \mathcal{M}_0 | \psi_a \rangle + \langle \psi_b | \mathcal{M}_{0^l 1^{L-l}} \oplus \mathcal{M}_{1^l 0^{L-l}} | \psi_b \rangle$$

$$= (|\alpha_1 \alpha_2|^2 + |\beta_1 \beta_2|^2) \Lambda_0 + |\alpha_1 \beta_2|^2 \Gamma_{L-l} + |\alpha_2 \beta_1|^2 \Gamma_l$$

$$= \frac{N}{N_l}.$$

$$(22)$$

The last equality comes from Lemma 2 and the fact that $\Gamma_l = \frac{N}{N_1}$ for 1 < l < L proven in the appendix.

Based on the proof that the largest eigenvalue of \mathcal{M} is achievable for \overline{L} part of the partition $1^{N_1}|\overline{L}$, it is straightforward to show that the largest eigenvalue of \mathcal{M} is also achievable for the $2^{N_2}|3^{N_3}|...|m^{N_m}$ part of a partition $1^{N_1}|2^{N_2}|3^{N_3}|...|m^{N_m}$. Theorem 1 then is true for any partition $1^{N_1}|2^{N_2}|3^{N_3}|...|m^{N_m}$.

3.3 k-partite separability

We then consider the k $(k \geq \frac{N+1}{2})$ partite separability. The maximum of $tr(\rho_s \hat{M})$ for a k-partite partition $1^{N_1}|2^{N_2}|3^{N_3}|...|m^{N_m}$ (where $k = \sum_{i=1}^m N_i$, $N = \sum_{i=1}^m iN_i$) is equal to $\frac{N}{N_1}$ as shown in the last subsection. We will analyze the possible largest $\frac{N}{N_1}$ for k $(k \geq \frac{N+1}{2})$ partite separability.

We first distribute each partite with one qubit and we remain N-k qubits. Then we have many strategies to distribute the remained qubits. The best way to decrease the number of single qubit parties (in order to increase $\frac{N}{N_1}$) is to distribute the remained N-k qubits to N-k parties, then we have $N_1 = k - (N-k)$ single qubit parties and (N-k) two qubit parties, the resultant partition is $1^{(2k-N)}|2^{(N-k)}$. The condition $k \geq \frac{N+1}{2}$ guarantees $N_1 \geq 1$, thus there is at least one single qubit party. The maximum of $tr(\rho_s \hat{M})$ for the partition $1^{(2k-N)}|2^{(N-k)}$ is equal to the maximum of $tr(\rho_s \hat{M})$ for the partition $1^{(2k-N)}|2^{(N-k)}|2^{(N-k)}$.

$$\Lambda \le \frac{N}{k - 2N} \tag{23}$$

for all $k \ (k \ge \frac{N+1}{2})$ partite separable noisy GHZ states. The achievable upper bound $\frac{N}{k-2N}$ of Λ leads to the following theorem.

Theorem 2 The necessary condition of the k-separability $(k \ge \frac{N+1}{2})$ of the noisy N qubit GHZ state ρ_{qhzN} is

$$p \le [1 + 2^{N-1}(2k - N)/N]^{-1}.$$
 (24)

4 Sufficient conditions for $k \ (k \ge \frac{N+1}{2})$ separability

4.1 Partition $1^{N_1}|L$

The way of $tr(\rho_s \hat{M})$ achieving its maximum $\Lambda = \frac{N}{N_1}$ hints the sufficient condition. We consider the case that the maximum value of $tr(\rho_s \hat{M})$ is achieved by the separable state with $z_i = 0$ for $i = 1, \ldots, N_1$, hence the pure state of the *i*th qubit is $\rho_i = \frac{1}{2}(I + x_i\sigma_1 + y_i\sigma_2)$ with $x_i^2 + y_i^2 = 1$. We may assume $x_i = \cos \varphi_i$, $y_i = \sin \varphi_i$, then $\rho_i = |\beta_i\rangle \langle \beta_i|$, where $|\beta_i\rangle = \frac{1}{\sqrt{2}}(|0\rangle + e^{i\varphi_i} |1\rangle)$. The state of the last L qubits is the eigenvector of \mathcal{M}_0 corresponding to eigenvalue $\Lambda_0 = \frac{N}{N_1}$. The eigenvector is

$$|\psi\rangle = \frac{1}{\sqrt{2}} (|0^{\otimes L}\rangle + e^{-i\varphi}|1^{\otimes L}\rangle).$$
 (25)

Where $\varphi = \sum_{i=1}^{N_1} \varphi_i$ as defined in Lemma 1. The $k(k = N_1 + 1)$ particle separable pure state is

$$|\Omega\rangle = \frac{1}{\sqrt{2^{N_1}}} \prod_{i=1}^{N_1} (|0\rangle + e^{i\varphi_i} |1\rangle) |\psi\rangle.$$
 (26)

We have the k-separable state $\rho_{s0} = \int |\Omega\rangle \langle \Omega| \prod_{i=1}^{N_1} \frac{d\varphi_i}{2\pi}$, which is

$$\rho_{s0} = \frac{1}{2^{N_1+1}} [2 | GHZ_N \rangle \langle \cdot |$$

$$+ \sum_{\mathbf{j} \neq \mathbf{0}; j_1, \cdots, j_{N_1} = 0,}^{1} (| j_1 \cdots j_{N_1} 0^{\otimes L} \rangle \langle \cdot |$$

$$+ | \overline{j_1} \cdots \overline{j_{N_1}} 1^{\otimes L} \rangle \langle \cdot |)], \qquad (27)$$

where the binary string $\mathbf{j} = (j_1, j_2, \dots, j_{N_1}), \ \overline{j_i} = 1 - j_i$. Averaging over all the cases of qubit permutations we arrive at a k-separable state

$$\rho_{s1} = [2 | GHZ_N \rangle \langle \cdot | + N^{-1} N_1 (T_1 + T_{N-1}) \\ + \sum_{i=2}^{N-2} (C_N^i)^{-1} (C_{N_1}^i + C_{N_1}^{i-L}) T_i] / 2^{N_1 + 1}.$$
(28)

where $T_i = \sum_{j_1+j_2+\dots+j_N=i} |j_1j_2\dots j_N\rangle \langle \cdot|$. In convention, we have $C_{N_1}^i = 0$ if $i > N_1$ and $C_{N_1}^{i-L} = 0$ if i < L.

Lemma 3 The inequality

$$(C_{N_1}^i + C_{N_1}^{i-L})/C_N^i < N_1/N$$
(29)

is true for all N and N_1 (0 < N_1 < N - 1).

The proof is shown in appendix.

Denote

$$\overline{\rho_{s2}} = \sum_{i=2}^{N-2} [N_1/N - (C_{N_1}^i + C_{N_1}^{i-N+N_1})/C_N^i]T_i/2^{N_1+1} + N^{-1}N_1(T_0 + T_N)/2^{N_1+1},$$
(30)

where $T_0 = |0^{\otimes N}\rangle \langle \cdot|$, $T_N = |1^{\otimes N}\rangle \langle \cdot|$. Then $\overline{\rho_{s2}}$ is an unnormalized full separable state. We have $\overline{\rho_s} = \rho_{s1} + \overline{\rho_{s2}}$ being an unnormalized k-separable $(k = N_1 + 1)$ state in the sense that the system is divided into N_1 single qubit parties and one L qubit party. The normalized k-separable state is

$$\rho_s = \frac{|GHZ_N\rangle \langle \cdot|}{1 + 2^{N-1}N_1/N} + \frac{N_1/(2N)}{1 + 2^{N-1}N_1/N} I_{2^N}.$$
 (31)

Hence, for the partition $1^{N_1}|L$ (up to qubit permutation) the noisy N qubit GHZ state is separable iff

$$p \le [1 + 2^{N-1}N_1/N]^{-1}.$$
 (32)

4.2 Further splits of the last L qubits

We consider the partition $1^{N_1}|\overline{L}$, where $\overline{L} = L_1|L_2|\cdots|L_m$ is a split of L with each L_i containing at least two qubits. Hence the number of parties is $k = N_1 + m$. Suppose with respect to the partition $1^{N_1}|\overline{L}$ there is a k-separable pure state

$$|\Omega\rangle = \frac{1}{\sqrt{2^{N_1+m}}} \prod_{i=1}^{N_1} (|0\rangle + e^{i\varphi_i} |1\rangle) \prod_{j=1}^m (|0^{\otimes L_j}\rangle + e^{i\varphi'_j} |1^{\otimes L_j}\rangle),$$
(33)

with $\varphi'_m = -\sum_{i=1}^{N_1} \varphi_i - \sum_{j=1}^{m-1} \varphi_j$. We have the k-separable state $\rho_{s3} = \int |\Omega\rangle \langle \Omega| \prod_{i=1}^{N_1} \frac{d\varphi_i}{2\pi} \prod_{j=1}^{m-1} \frac{d\varphi'_j}{2\pi}$, which is

$$\rho_{s3} = \frac{1}{2^{N_1+m}} [2 |GHZ_N\rangle \langle \cdot |$$

$$+ \sum_{\mathbf{j}\neq\mathbf{0}} (\left| j_1 \cdots j_{N_1} j_{N_1+1}^{\otimes L_1} \cdots j_{N_1+m-1}^{\otimes L_{m-1}} \mathbf{0}^{\otimes L_m} \right\rangle \langle \cdot |$$

$$+ \left| \overline{j_1} \cdots \overline{j_{N_1}} \overline{j_{N_1+1}^{\otimes L_1}} \cdots \overline{j_{N_1+m-1}^{\otimes L_{m-1}}} \mathbf{1}^{\otimes L_m} \right\rangle \langle \cdot |)], \quad (34)$$

Where the summation is over all $\mathbf{j} \in \{0, 1\}^{\otimes (N_1+m)}$ except $\mathbf{j} = \mathbf{0}$, and $\overline{j^{\otimes i}} = (1-j)^{\otimes i}$ for binary j.

Lemma 4 The averaging of ρ_{s3} in (34) over all the qubit permutations can be written as follows:

$$\rho_{s4} = \frac{1}{2^{N_1 + m}} [2 | GHZ_N \rangle \langle \cdot | + \sum_{i=1}^{N-1} (C_N^i)^{-1} f_m(i) T_i], \quad (35)$$

with

$$\sum_{i=1}^{N-1} f_m(i) = 2^{N_1+m} - 2, f_m(i) = f_m(N-i), f_m(1) \le N (36)$$
$$0 \le f_m(i) \le f_m(1)C_N^i/N \quad \text{for all } i \in [1, N-1]. (37)$$

We will use mathematical induction to prove this lemma in the appendix.

We then produce the noisy GHZ state by mixing some fully separable state with ρ_{s4} due to conditions (36) and (37). So that the obtained N qubit noisy GHZ state is k-separable with $k = N_1 + m$. We have the following sufficient criterion:

Theorem 3 The sufficient condition of k-separability $(k = N_1 + m)$ for a N qubit noisy GHZ state is

$$p \le [1 + 2^{N-1} N_1 / N]^{-1} \tag{38}$$

where $N_1 > 0$ is the number of parties with single qubit, the partition is $1^{N_1} | \overline{L}$ with \overline{L} being $L_1 | L_2 | \cdots | L_m$, each partite L_j has two or more qubits.

From (38), we can see that the less the number N_1 , the larger the right hand side of (38), the better the sufficient condition. If the number of parties is fixed to be $k \ (k \geq \frac{N+1}{2})$, the smallest number of N_1 is 2k - N, we have the sufficient criterion:




Figure 1: (Color online)The entanglement properties of N-qubit GHZ states mixed with white noise, $\rho_{ghzN} = p |GHZ_N\rangle \langle GHZ_N | + \frac{1-p}{2^N} I_{2^N}$. It was known before [39],[40] that the states are fully separable iff $p \leq \frac{1}{1+2^{N-1}}$ (the downmost solid line), The states biseparable iff $p \leq \frac{2^{N-1}-1}{2^N-1}$ [11](the uppermost dashed line). Our results show that the states are N-1 separable iff $p \leq \frac{1}{1+\frac{N-2}{2^N-1}}$, k separable iff $p \leq \frac{1}{1+\frac{2k-N}{2^N-1}}$ when $k \geq \frac{N+1}{2}$. They are shown by solid lines for k-separability with k = N, N - 1, N - 2, N - 3, N - 4, N - 5 from left to right. The dashed lines are for k-separability with k = 2, 3, 4, 5 from top down.

Theorem 4 The sufficient condition of k-separability $(k \ge \frac{N+1}{2})$ for noisy GHZ state is

$$p \le [1 + 2^{N-1}(2k - N)/N]^{-1},$$
 (39)

The sufficient condition is realized by a proper state of the partition $1^{(2k-N)}|2^{(N-k)}$.

In figure 1, we display the k partite separable conditions for N - k = 1, 2, 3, 4, 5.

5 Other *k*-separability

There are many k-separability noisy GHZ states that can not be fit into the former regime of $k \ (k \ge \frac{N+1}{2})$ separability or biseparability. The first case is the tri-separability of noisy GHZ states for $N \ge 6$. The partitions are 1|2|3 and 2^3 for N = 6. These partitions give rise to the tripartite states:

$$\rho_{1|2|3}, \rho_{2^3} = \frac{1}{2^3} (2 |GHZ_6\rangle \langle \cdot | + \sum_{i=1}^5 v_i \frac{T_i}{C_6^i}), \qquad (40)$$

where $\mathbf{v} = (1, 1, 2, 1, 1)$ for $\rho_{1|2|3}$ and $\mathbf{v} = (0, 3, 0, 3, 0)$ for ρ_{2^3} , respectively. Consider the mixture of this two kinds

of partitions, the mixed state is $\rho_{s5} = q\rho_{1|2|3} + (1-q)\rho_{2^3}$. We then have

$$\rho_{s5} = \frac{1}{2^3} (2 |GHZ_6\rangle \langle \cdot | + \sum_{i=1}^5 u_i \frac{T_i}{C_6^i}), \qquad (41)$$

where $\mathbf{u} = (q, q+3(1-q), 2q, q+3(1-q), q)$. Let $u_1/C_6^1 = u_2/C_6^2 \ge u_3/C_6^3$, then $q = \frac{2}{3}$. Denote the coefficient of T_1 as $\frac{1}{\tau}$, then $\tau = 9$. The tripartite separable sufficient condition is

$$p \le \frac{1}{1+2^5/\tau} = \frac{9}{41}.$$
(42)

To find the mixed state that has maximal τ , we need to solve the linear programming problem. This is because the mixture probability is positive and we let $u_1/C_N^1 = u_2/C_N^2 = \cdots = u_i/C_N^i \geq u_{i+j}/C_N^{i+j}$ to optimize τ , for some $i \leq \lceil \frac{N}{2} \rceil$ and all positive $j \leq \lceil \frac{N}{2} \rceil - i$. For $N \leq 12$, we list all these mixed optimal states in

For $N \leq 12$, we list all these mixed optimal states in Table I.

Table I. The k-separabilities of $N = 6, ..., 12; \tau \cdot fractions$ are τ times the fractions of partitions. For the tri-separable of 6 qubit noisy GHZ state, partitions $2^3, 1|2|3$ contribute $q = \frac{1}{3}$ and $1 - q = \frac{2}{3}$ fractions in the optimization of τ . Hence $\tau \cdot fractions = (3, 6)$. p_s is the critical value such that the noisy GHZ state is k-separable if $p < p_s$.

t $p \leq p$	p_s .				
N	k	partitions	$\tau \cdot fractions$	au	p_s
6	3	$2^3, 1 2 3$	3, 6	9	$\frac{9}{41}$
7	3	$2^2 3,1 3^2$	10.5, 7	17.5	$\frac{35}{163}$
8	3	$1 3 4,2 3^2,2^2 4$	8, 24, 2	34	$\frac{17}{81}$
8	4	$1 2^2 3, 2^4$	8,3	11	$\frac{11}{139}$
9	3	$1 4^2, 2 3 4, 3^3$	16, 36, 9	61	$\frac{61}{327}$
9	4	$1 2 3^2, 2^3 3$	9,9	18	$\frac{39}{137}$
10	3	$1 4 5,2 4^2,3^2 4$	10, 45, 60	115	$\frac{115}{627}$
10	4	$1 3^3, 2^2 3^2$	10, 22.5	32.5	$\frac{65}{1089}$ $\frac{13}{13}$
10	5	$1 2^3 3, 2^5$	10, 3	13	$\frac{13}{525}$
11	3	$3 4^2, 2 4 5, 3^2 5, 1 5^2$	137.5, 55, 13.75, 11	$\frac{869}{4}$	$\frac{869}{4965}$
11	4	$2^{2} 3 4,2 3^{3},$ $1 3^{2} 4$	4.4, 46.2, 11	61.6	$\frac{77}{1375}$
11	5	$2^4 3, 1^2 3^3$	11, 11	22	$\frac{11}{523}$
12	3	$\begin{array}{c} 4^3, 3 4 5, \\ 2 5^2, 1 5 6 \end{array}$	$\frac{275}{3}, 220, 66, 12$	$\frac{1169}{3}$	$\frac{1169}{7313}$
12	4	$3^4, 2^2 4^2,$ $1 3 4^2$	55, 33, 12	100	$\frac{25}{537}$
12	5	$1 2 3^3, 2^3 3^2$	12, 18	30	$\frac{15}{1039}$
12	6	$1 2^4 3, 2^6$	12, 3	15	$\frac{15}{2063}$

6 Discussion

The N qubit GHZ state as a codeword of the 'tree' code was utilized to explore the nonadditivity of channel capacity of depolarizing channel [41]. The 'tree' code and the repetition code [42] (or 'cat' code [43]) have the same Hilbert subspace but with different bases, they are the same code. A depolarizing channel \mathcal{E} is a completely positive trace preserving map, it maps an input qubit state ρ to an output state $\mathcal{E}(\rho)$, with

$$\mathcal{E}(\rho) = (1 - 3q)\rho + q\sigma_1\rho\sigma_1 + q\sigma_2\rho\sigma_2 + q\sigma_3\rho\sigma_3.$$
(43)

Where the noise q characterizes the depolarizing chan-When the N qubit GHZ state passing through nel. N parallel depolarizing channels, the output state is $\mathcal{E}^{\otimes N}(|GHZ_N\rangle\langle GHZ_N|).$ The output state is a mixture of N qubit GHZ state and a color noise (a diagonal state in the computational basis), the fraction of $|GHZ_N\rangle\langle GHZ_N|$ in the output state is $(1-4q)^N$. We wonder if the output state is entangled, partially separable or fully separable. We may use ρ_{ghzN} to approximate $\mathcal{E}^{\otimes N}(|GHZ_N\rangle \langle GHZ_N|)$. In table II, we use q_{th} to denote the threshold noise (derived from Fidelity = 1 - 3q [43]) of depolarizing channel. Quantum coherent information is positive for the input of 'tree' code when the channel noise is less than the threshold noise, otherwise it is define to be zero. The codewords of the code are $|GHZ_N\rangle$ and $|GHZ_{N-}\rangle = \frac{1}{\sqrt{2}}(|0^{\otimes N}\rangle - |1^{\otimes N}\rangle)$ and they have equal prior probabilities. We compare q_{th} with the various critical values of q for different separabilities of ρ_{qhzN} . The critical values of q are denoted as q_{N-k} for the partition $1^{(2k-N)}|2^{(N-k)}$. When $q > q_{N-k}$, the state ρ_{ghzN} is separable for the partition $1^{(2k-N)}|2^{(N-k)}$. When $q > q_0$, the state ρ_{qhzN} is fully separable.

Notice that the biseparabilities of both ρ_{ghzN} and $\mathcal{E}^{\otimes N}(|GHZ_N\rangle \langle GHZ_N|)$ are detected by the same witness $\frac{\mathbb{I}}{2} - |GHZ_N\rangle \langle GHZ_N|$, they have the same biseparable critical value q_{bi} when they have the same fractions of $|GHZ_N\rangle$. Table II shows us that $q_{th} > q_{bi}$ for all $N \geq 3$. This means that the state $\mathcal{E}^{\otimes N}(|GHZ_N\rangle \langle GHZ_N|)$ is entangled for N = 2 and not genuinely entangled for all $N \geq 3$ when the noise of channel arrives the threshold.

On the other hand, $q_{th} < q_0$. The noise of the output state $\mathcal{E}^{\otimes N}(|GHZ_N\rangle \langle GHZ_N|)$ is a color noise. Thus some of the diagonal elements should be smaller than the average noise level, which is the noise level of ρ_{ghzN} if the same fractions of $|GHZ_N\rangle$ in the two states are assumed. The smaller the diagonal element, the easier the entanglement takes place. Hence, $\mathcal{E}^{\otimes N}(|GHZ_N\rangle \langle GHZ_N|)$ is easier to be entangled than ρ_{ghzN} , and tolerant a higher level of noise than the later to keep to be entangled. Thus $q_{th} < q_0$ holds for output state $\mathcal{E}^{\otimes N}(|GHZ_N\rangle \langle GHZ_N|)$. So that when the state $|GHZ_N\rangle$ is transmitted over depolarizing channel, the output state can not be fully separable if the channel noise is less than its threshold. The output state should maintain some kinds of entanglement.

Table II. Comparison of the threshold noise q_{th} to various critical values of q for different separabilities of ρ_{ghzN} .

N	q_0	q_1	q_2	q_3	q_{th}	q_{bi}
2	0.1057				0.0628	0.1057
3	0.1038	0.0615			0.0634	0.0615
4	0.1057	0.0828			0.0633	0.0434
5	0.1081	0.0941	0.0624		0.0635	0.0338
6	0.1104	0.1010	0.0840		0.0634	0.0279
7	0.1123	0.1056	0.0950	0.0704	0.0634	0.0238

In summary, the output state of depolarizing channel with threshold noise is neither genuinely entangled nor fully separable when we input GHZ state. It is entangled for N = 2 and partially entangled for $N \ge 3$. Thus the k-separability makes sense.

7 Conclusion

We find the entanglement witnesses for almost a half of all kinds of entanglement of an N qubit GHZ state in white noise. The witnesses are linear combinations of the stabilizer group elements of GHZ states. The combinational coefficient of a stabilizer element relies on the number of generators in the stabilizer element. The necessary and sufficient condition for k partite separability has been given for arbitrary N when $k \ge \frac{N+1}{2}$. The necessary and sufficient condition is achieved by a partition (up to qubit permutations) with N - k double qubit parties and 2k - N single qubit parties. The k separability condition is $p < [1 + 2^{N-1}(2k - N)/N]^{-1}$, where p is the fraction of pure GHZ state in the noisy N qubit GHZ state. This gives rise to the necessary and sufficient conditions for more than a half of the k-separability of multi-qubit noisy GHZ states. For all the other k-separable problems of noisy multi-qubit GHZ states, we find that the sufficient separable conditions are achieved by the mixtures of the partitions with the same k (number of parties) but different qubit number distributions. We display all the separabilities of N qubit noisy GHZ states for $N \leq 12$. As an application, we have approximated the output of GHZ state passing through depolarizing channel with the GHZ state in white noise. It has been shown the output state is neither genuinely entangled nor fully separable when the depolarizing channel has threshold noise.

GHZ states are the simplest graph states. The method of constructing entanglement witness operators with linear combinations of stabilizer operators in this paper is also useful for entanglement detection of noisy graph states. A stabilizer codeword passing through certain quantum channel may become a noisy graph state. Thus our method may find more applications in quantum information transmission.

Acknowledgment

Supported by the National Natural Science Foundation of China (Grant Nos. 11375152) and (partially) supported by National Basic Research Program of China (Grant No. 2014CB921203) are gratefully acknowledged.

Appendix

Proof of Lemma 1

Proof: In computational basis, we use $|\psi\rangle = \sum_{\mathbf{i}} \alpha_{\mathbf{i}} |\mathbf{i}\rangle$ to denote the pure state of the last *L* qubits in ρ_s with the partition $1^{N_1}|L$. Where $\mathbf{i} = (i_1, i_2, \dots, i_L)$ is a binary string and we denote the weight of \mathbf{i} as $|\mathbf{i}|$. We rewrite (16) as

$$tr(\rho_s \hat{M}_D) = \sum_{i=1}^{\lfloor N/2 \rfloor} M_i \sum_{n=\max\{0,2i-N_1\}}^{\min\{L,2i\}} S_{2i-n}q_n.$$
(44)

Where $S_0 = 1, S_i = \sum_{1 \le j_1 < j_2 < \dots < j_i \le N_1} \prod_{l=1}^i z_{j_l}$, and

$$q_{n} = \sum_{|\overline{\mathbf{j}}|=n} \langle \psi | \bigotimes_{l=N_{1}+1}^{N} \sigma_{3}^{j_{l}} | \psi \rangle$$
$$= \sum_{|\overline{\mathbf{j}}|=n} \sum_{\mathbf{i}} |\alpha_{\mathbf{i}}|^{2} (-1)^{\mathbf{i} \cdot \overline{\mathbf{j}}}, \qquad (45)$$

where $\mathbf{\overline{j}} = (j_{N_1+1}, \cdots, j_N)$ is a binary string and $|\mathbf{\overline{j}}|$ is the weights of $\mathbf{\overline{j}}$. We may write $q_n = \sum_{l=0}^{L} \sum_{|\mathbf{i}|=l} |\alpha_{\mathbf{i}}|^2 w_{n,l}$ with

$$w_{n,l} = \sum_{j=\max\{0,n+l-L\}}^{\min\{n,l\}} (-1)^j C_{L-l}^{n-j} C_l^j.$$
(46)

Then

$$tr(\rho_s \hat{M}_D) = \sum_{l=0}^{L} \sum_{|\mathbf{i}|=l} |\alpha_{\mathbf{i}}|^2 \Gamma_l, \qquad (47)$$

with

$$\Gamma_{l} = \sum_{m=1}^{\left\lceil \frac{N_{1}}{2} \right\rceil} S_{2m-1} \sum_{i=1}^{\left\lceil \frac{L}{2} \right\rceil} M_{i+m-1} w_{2i-1,l} \\ + \sum_{m=0}^{\left\lfloor \frac{N_{1}}{2} \right\rfloor} S_{2m} \sum_{i=0}^{\left\lfloor \frac{L}{2} \right\rfloor} M_{i+m} w_{2i,l}.$$

If we denote

$$\mathcal{X} = \sum_{|\mathbf{\overline{j}}||even} (-1)^{|\mathbf{\overline{j}}|/2} \bigotimes_{l=N_1+1}^N \sigma_1^{1-j_l} \sigma_2^{j_l}, \qquad (49)$$

....

$$\mathcal{Y} = \sum_{|\mathbf{\bar{j}}||odd} (-1)^{(|\mathbf{\bar{j}}|-1)/2} \bigotimes_{l=N_1+1}^N \sigma_1^{1-j_l} \sigma_2^{j_l}, \quad (50)$$

Equation (17) reads

$$tr(\rho_s \hat{M}_A) = \langle \psi | c \mathcal{X} - d \mathcal{Y} | \psi \rangle, \qquad (51)$$

Then we have

$$\langle \psi | \mathcal{X} | \psi \rangle = 2^{L-1} (\alpha_{\mathbf{0}} \alpha_{\mathbf{1}}^* + \alpha_{\mathbf{0}}^* \alpha_{\mathbf{1}}), \qquad (52)$$

$$\langle \psi | \mathcal{Y} | \psi \rangle = i 2^{L-1} (\alpha_0 \alpha_1^* - \alpha_0^* \alpha_1), \qquad (53)$$

We have used $\langle 0^{\otimes (L-l)} 1^{\otimes l} | \mathcal{X} | 1^{\otimes (L-l)} 0^{\otimes l} \rangle = 0$ if $l \neq 0$ and $l \neq L$. This is due to the fact that for each term in \mathcal{X} if there are odd number of σ_2 in the last l qubits, then there are odd number of σ_2 in the first L - l qubits too, so that an extra -1 factor emerges due to σ_2^2 for such a term. The probabilities of odd and even number of σ_2 in the last l qubits are equal. The terms of \mathcal{X} are concealed with each other in the evaluation of the matrix element $\langle 0^{\otimes (L-l)} 1^{\otimes l} | \mathcal{X} | 1^{\otimes (L-l)} 0^{\otimes l} \rangle$. The null result is also true for \mathcal{Y} when $l \neq 0$ and $l \neq L$.

From (46), we have $w_{i,0} = C_L^i, w_{i,L} = (-1)^i C_L^i$. Thus

$$\Gamma_0 = N/N_1 + 2^{L-1}(a+b), \tag{54}$$

$$\Gamma_L = N/N_1 + 2^{L-1}(a-b), \qquad (55)$$

where

$$a = -1 + N_1^{-1} \sum_{m=1}^{\lfloor N_1/2 \rfloor} [4m - N_1] S_{2m}, \qquad (56)$$

$$b = N_1^{-1} \sum_{m=1}^{\lceil N_1/2 \rceil} [4m - 2 - N_1] S_{2m-1}.$$
 (57)

Proof of Lemma 2

Proof: This is equivalent to $a + \sqrt{b^2 + c^2 + d^2} \le 0$. We should show that (i) $a \le 0$, (ii) $a^2 - b^2 \ge c^2 + d^2$. Let

$$u = N_1^{-1} \sum_{j=1}^{N_1} \prod_{n=1}^{N_1} [1 + (-1)^{\delta_{j,n}} z_n], \qquad (58)$$

$$v = N_1^{-1} \sum_{j=1}^{N_1} \prod_{n=1}^{N_1} [1 - (-1)^{\delta_{j,n}} z_n].$$
 (59)

(48) A direct calculation shows that

$$u = 1 + \sum_{i=1}^{N_1} [1 - 2i/N_1] S_i, \tag{60}$$

$$v = 1 + \sum_{i=1}^{N_1} [1 - 2i/N_1] (-1)^i S_i.$$
 (61)

Hence $\frac{1}{2}(u+v) = -a$ and $\frac{1}{2}(u-v) = -b$. We arrive at (i) $a \leq 0$, since $u \geq 0$ and $v \geq 0$ due to $z_n \in [-1, 1]$ for all n. We further have $a^2 - b^2 = uv$, they are

$$N_1^{-2} \sum_{i=1}^{N_1} \sum_{j=1}^{N_1} (1-z_i)^2 (1+z_j)^2 \prod_{n=1, n \neq i, j}^{N_1} (1-z_n^2), \quad (62)$$

meanwhile $c^2 + d^2 = \prod_{i=1}^{N_1} (1 - z_i^2)$, thus we arrive at (ii)

$$a^{2} - b^{2} - c^{2} - d^{2} = 8N_{1}^{-2} \sum_{1 \le i < j \le N_{1}} (z_{i} - z_{j})^{2} (63)$$
$$\times \prod_{n=1, n \ne i, j}^{N_{1}} (1 - z_{n}^{2}) \ge 0.$$
(64)

Proof of Theorem 1

Proof: The matrix $\mathcal{M} = \bigoplus_{i \neq 0,1} \mathcal{M}_{i,i} \oplus \mathcal{M}_0$. Except for the submatrix \mathcal{M}_0 , the other eigenvalues of \mathcal{M} are just the other diagonal elements $\mathcal{M}_{i,i} = \Gamma_{|i|}$. From the definition of $w_{n,l}$ in (46), it is not difficult to show that

$$\sum_{i=0}^{\lfloor \frac{L}{2} \rfloor} w_{2i,l} = 0, \sum_{i=1}^{\lfloor \frac{L}{2} \rfloor} w_{2i-1,l} = 0,$$
(65)

$$\sum_{i=0}^{\lfloor \frac{L}{2} \rfloor} i w_{2i,l} = \begin{cases} 0, \text{ for } 1 < l < L-1 \\ -2^{L-3}, \text{ for } l = 1, L-1 \end{cases}$$
(66)

$$\sum_{i=1}^{\left\lceil \frac{L}{2} \right\rceil} i w_{2i-1,l} = \begin{cases} 0, \text{ for } 1 < l < L-1 \\ -2^{L-3}, \text{ for } l = 1 \\ 2^{L-3}, \text{ for } l = L-1 \end{cases}$$
(67)

Using (14), we have

$$\sum_{i=1}^{\left\lfloor \frac{L}{2} \right\rfloor} M_{i+m-1} w_{2i-1,l} = \frac{2^{L-1}}{N_1} (-\delta_{l,1} + \delta_{l,L-1}), \quad (68)$$

$$\sum_{i=0}^{\lfloor \frac{L}{2} \rfloor} M_{i+m} w_{2i,l} = \frac{N}{N_1} \delta_{m,0} - \frac{2^{L-1}}{N_1} (\delta_{l,1} + \delta_{l,L-1}).$$
(69)

From (67), for 1 < l < L - 1, we have

$$\Gamma_l = \frac{N}{N_1};\tag{70}$$

For l = 1, we arrive at

$$\Gamma_{1} = \frac{N}{N_{1}} - \frac{2^{L-1}}{N_{1}} \sum_{m=0}^{N_{1}} S_{m}
= \frac{N}{N_{1}} - \frac{2^{L-1}}{N_{1}} \prod_{m=1}^{N_{1}} (1+z_{m})
\leq \frac{N}{N_{1}},$$
(71)

and

$$\Gamma_{L-1} = \frac{N}{N_1} - \frac{2^{L-1}}{N_1} \prod_{m=1}^{N_1} (1 - z_m)$$

$$\leq \frac{N}{N_1}.$$
 (72)

Thus we have proved that all the eigenvalues of \mathcal{M} are up bounded by $\frac{N}{N_1}$. The up bound is achievable. Hence for our choice of the witness in (14), the largest eigenvalue of the matrix \mathcal{M} , thus the maximum of $tr(\rho_s \hat{M})$ is $\frac{N}{N_1}$ for the partition $1^{N_1}|L$. The M_i in (14) yields $\sum_{i=1}^{\lfloor N/2 \rfloor} M_i C_N^{2i} = \frac{N}{N_1}$.

Proof of Lemma 3

Proof: For $i > N_1$ or i < L, the inequality (29) is apparently true. So we consider $L \le i \le N_1$. Since L > 1 is assumed, we have 1 < i < N - 1. The inequality can be rewritten as

$$(N-i)(N-i-1)\cdots(N-i-L+1) + i(i-1)\cdots(i-L+1) < (N-1)(N-2)\cdots N_1.$$
(73)

For L = 2, 3, the inequality becomes N > i + 1, which is true. Let the inequality be true for L = l, we will show that the inequality will be true for L = l + 1 too. Notice that N > i + 1, we have ^{i-l}/_{N-l-1} < 1. Meanwhile i > 1 leads to N-l-1/N-l-1 < 1. Thus we have

$$(N-i)(N-i-1)\cdots(N-i-l+1)\frac{N-i-l}{N-l-1} + i(i-1)\cdots(i-l+1)\frac{i-l}{N-l-1} < (N-1)(N-2)\cdots(N-l),$$
(74)

which is the inequality for L = l + 1.

Proof of Lemma 4

When m = 1, the partition reduces to $1^{N_1}|L$, the state ρ_{s4} reduces to ρ_{s1} . We can verify that (36) and (37) are true for m = 1 case.

Let equation (36) be true for m = m' case. For the case of m = m' + 1, let us consider a partition $1^{N_1} |\overline{L}| l$ of N + lqubit system with $l \ge 2$. The state for this partition can be described by

$$f_{m'+1}(i) = \begin{cases} f_{m'}(i) + f_{m'}(i-l) \text{ for } i \in [l,N]; \\ f_{m'}(i) \text{ for } i < l; \\ f_{m'}(i-l) \text{ for } i > N. \end{cases}$$
(75)

We want show that $f_{m'+1}(i)/C_{N+l}^i \leq f_{m'+1}(1)/(N+l)$ for $i \in [1, N+l-1]$. Apparently, we only need to show it for $i \in [l, N]$. We have $f_{m'+1}(i) = f_{m'}(i) + f_{m'}(i-l) \leq$ $f_{m'}(1)(C_N^i + C_N^{i-l})/N$. We have $f_{m'+1}(1) = f_{m'}(1)$ for $l \geq 2$ (in fact $f_{m'}(1) = N_1$), thus what we need to show is

$$(C_N^i + C_N^{i-l})/N \le C_{N+l}^i/(N+l).$$
 (76)

It is the inequality proved in Lemma 3. Thus for the partition $1^{N_1}|\overline{L}|l$ of the N+l qubit noisy GHZ state with $l \geq 2$, we have $0 \leq f_{m'+1}(i) \leq f_{m'+1}(1)C_{N+l}^i/(N+l)$ for all $i \in [1, N+l-1]$. Hence the assumption (35) is true.

References

- R. Horodecki, P. Horodecki, M. Horodecki, K. Horodecki, Rev. Mod. Phys. 81, 865(2009).
- [2] O. Gühne, G. Tóth, Phys. Rep. **474**, 1(2009).
- [3] T. Monz, P. Schindler, J. T. Barreiro, M. Chwalla, D. Nigg, W. A. Coish, M. Harlander, W. Hänsel, M. Hennrich, and R. Blatt, Phys. Rev. Lett. 106, 130506(2011).
- [4] X.C. Yao, T.X. Wang, P. Xu, H. Lu, G.S. Pan, X.H. Bao, C.Z. Peng, C.Y. Lu, Y.A. Chen, and J.W. Pan, Nature Photonics 6, 225(2012).
- [5] P. Neumann, N. Mizuochi, F. Rempp, P. Hemmer, et al., Science, **320**, 1326(2008).
- [6] C. A. Sackett, D. Kielpinski, B. E. King, C. Langer, V. Meyer, C. J. Myatt, M. Rowe, Q. A. Turchette, W. M. Itano, D. J. Wineland, and C. Monroe, Nature 404, 256 (2000).
- [7] Z. Zhao, Y.A. Chen, A.N. Zhang, T. Yang, H. J. Briegel, and J.W. Pan, Nature 430, 54 (2004).
- [8] W.B. Gao, C.Y. Lu, X.C. Yao, P. Xu, O. Gühne, A. Goebel, Y.A. Chen, C. Z. Peng, Z.B. Chen, and J.W. Pan, Nature Physics 6, 331 (2010).
- [9] M. Seevinck, J. Uffink, Phys. Rev. A 78, 032101 (2008)
- [10] M. Huber, F. Mintert, A. Gabriel, B. C. Hiesmayr, Phys. Rev. Lett. **104**, 210501 (2010).
- [11] O. Guhne, M. Seevinck, New J. Phys. 12, 053002 (2010).
- [12] P. Badziag, C. Brukner, W. Laskowski, T. Paterek, M. Zukowski, Phys. Rev. Lett **100**, 140403 (2008)
- [13] A. S. M. Hasan, P. S. Joag, Quantum Inf. Comput. 8, 773 (2008).
- [14] W. Laskowski, M. Markiewicz, T. Paterek, M. Zukowski, Phys. Rev. A 84, 062305 (2011).
- [15] J. I. de Vicente, M. Huber, Phys. Rev. A 84, 062306 (2011).
- [16] N. Ananth, V. K. Chandrasekar, M. Senthilvelan, Eur. Phys. J. D 69, 56 (2015).
- [17] O. Gühne Phys. Lett. A, **375**, 406-410 (2011).
- [18] X.Y. Chen, L.Z. Jiang, P. Yu, M. Tian, Quant. Inf. Process. 14,2463-2476(2015).
- [19] X.Y. Chen, L.Z. Jiang, Z. A. Xu, Quant. Inf. Process. 16, 95,(2017).

- [20] B. Jungnitsch, T. Moroder, and O. Gühne, Phys. Rev. Lett. 106, 190502 (2011).
- [21] B. M. Terhal, Phys. Lett. A **271**, 319 (2000).
- [22] J. Sperling and W. Vogel, Phys. Rev. Lett. 111, 110503 (2013).
- [23] A. Kay, Phys. Rev. A, 83, 020303(R), (2011).
- [24] P. van Loock and A. Furusawa, Phys. Rev. A 67, 052315 (2003).
- [25] M. Bourennane et al., Phys. Rev. Lett. 92, 087902 (2004).
- [26] J. Eisert, F. G. S. L. Brandão, and K. M. R. Audenaert, New J. Phys. 9, 46 (2007).
- [27] D. Chruściński and A. Kossakowski, J. Phys. A: Math. Theor. 41, 145301 (2008).
- [28] H. Shibata, T. Honjo, and K. Shimizu, Opt. Lett. 39, 5078 (2014).
- [29] J. Yin et al., Nature (London) 488, 185 (2012).
- [30] X. S. Ma et al., Nature (London) **489**, 269 (2012).
- [31] S. Bose, V. Vedral, and P. L. Knight, Phys. Rev. A 57,822 (1998).
- [32] R. Cleve, D. Gottesman, and H.-K. Lo, Phys. Rev. Lett. 83, 648 (1999).
- [33] W. Tittel et al., Phys. Rev. A 63, 042301 (2001); Y.A. Chen et al., Phys. Rev. Lett. 95, 200502 (2005); C. Schmid et al., Phys. Rev. Lett. 95, 230505 (2005); S. Gaertner et al., Phys. Rev. Lett. 98, 020503 (2007).
- [34] B. Bell et al., Nat. Commun. 5, 5480 (2014).
- [35] M. Żukowski et al., Acta Phys. Pol. **93**, 187 (1998).
- [36] A.O. Pittenger and M.H. Rubin, Optics Comm. 179, 447(2000).
- [37] S. Abe, Phys. Rev. A 65, 052323 (2002).
- [38] A. S Nayak, Sudha, A. R. U. Devi, and A. K. Rajagopal, arXiv.org: 1712.00746
- [39] W. Dür and I. Cirac, Phys. Rev. A **61**, 042314(2000).
- [40] R. Schack and C. M. Caves, J. Mod. Opt. 47, 387 (2000).
- [41] X. Y. Chen and L. Z. Jiang, Phys. Rev. A 83, 052316 (2011).
- [42] G. Smith and J. A. Smolin, Phys. Rev. Lett. 98, 030501 (2007).
- [43] D. P. DiVincenzo, P. W. Shor, and J. A. Smolin, Phys. Rev. A 57, 830 (1998).