# The Star-formation History and Accretion-Disk Fraction Among the K-Type Members of the Scorpius-Centaurus OB Association

Mark J. Pecaut,<sup>1\*</sup> Eric E. Mamajek<sup>2</sup>

<sup>1</sup>Rockhurst University, Department of Physics, 1100 Rockhurst Rd, Kansas City, MO 64110-2508, USA <sup>2</sup>University of Rochester, Department of Physics and Astronomy, Rochester, NY 14627-0171, USA

Accepted XXX. Received YYY; in original form ZZZ

### ABSTRACT

We present results of a spectroscopic survey for new K- and M-type members of Scorpius-Centaurus (Sco-Cen), the nearest OB Association (~100-200 pc). Using an X-ray, proper motion and color-magnitude selected sample, we obtained spectra for 361 stars, for which we report spectral classifications and Li and H $\alpha$  equivalent widths. We identified 156 new members of Sco-Cen, and recovered 51 previously published members. We have combined these with previously known members to form a sample of 493 solar-mass ( $\sim$ 0.7–1.3 M $_{\odot}$ ) members of Sco-Cen. We investigated the star-formation history of this sample, and re-assessed the ages of the massive main-sequence turn-off and the G-type members in all three subgroups. We performed a census for circumstellar disks in our sample using WISE infrared data and find a protoplanetary disk fraction for K-type stars of  $4.4^{+1.6}_{-0.9}$ % for Upper Centaurus-Lupus and Lower Centaurus-Crux at ~16 Myr and  $9.0^{+4.0}_{-2.2}$ % for Upper Scorpius at ~10 Myr. These data are consistent with a protoplanetary disk e-folding timescale of  $\sim$ 4–5 Myr for  $\sim$ 1  $M_{\odot}$ stars, twice that previously quoted (Mamajek 2009), but consistent with the Bell et al. revised age scale of young clusters. Finally, we construct an age map of Scorpius-Centaurus which clearly reveals substructure consisting of concentrations of younger and older stars. We find evidence for strong age gradients within all three subgroups. None of the subgroups are consistent with being simple, coeval populations which formed in single bursts, but likely represents a multitude of smaller star formation episodes of hundreds to tens of stars each.

## Key words:

**Key words:** open clusters and associations: individual(Scorpius-Centaurus, Sco OB2, Upper Scorpius, Upper Centaurus-Lupus, Lower Centaurus-Crux) – stars: pre-main-sequence – (stars): circumstellar matter

#### 1 INTRODUCTION AND BACKGROUND

Because of their utility for studying star and planet formation and evolution, a great deal of effort has been directed towards identifying and characterizing samples of nearby young stars. These samples have been indispensible for direct imaging of brown dwarf and planetary-mass companions (e.g., Kraus & Ireland 2012), imaging debris disks in scattered light (e.g., Kalas et al. 2015), exploring the star-disk interaction in young stars (e.g., Nguyen et al. 2009), and studying the evolution of gas-rich and dusty debris disks as a function of age (e.g., Mamajek 2009; Chen et al. 2011).

The nearest of these samples are small, diffuse, nearby associations of kinematically related stars, (e.g., the  $\beta$  Pictoris moving group, TW Hya Association; Zuckerman & Song 2004; Torres et al. 2008). These typically have tens of known members, with masses ~0.02-2 M<sub>☉</sub>. While these are extremely useful due to their proximity, studies based on these samples may be subject to small **CONSECTIONAL TRANSFO** fortunately, there exist large collections of young stars, only slightly more distant, in nearby OB associations. OB

associations contain members across the mass spectrum, including the hot and massive O- and B-type stars for which they are named, intermediate-mass A/F stars, lower-mass G/K/M stars, and very low-mass free-floating substellar objects. Though the lowermass (< 1.5 M<sub>☉</sub>) stars blend in with the Galactic field population and are therefore much more difficult to identify than the OB stars, they comprise the dominant stellar component of OB associations, typically present in the thousands (Briceño et al. 2007).

Scorpius-Centaurus (Sco-Cen), the nearest OB association, has been the subject of several efforts to identify its lower-mass population (e.g., de Zeeuw et al. 1999; Preibisch & Zinnecker 1999; Mamajek et al. 2002; Rizzuto et al. 2011, 2015). Sco-Cen harbors a barely explored population of thousands of low-mass K and Mtype stars (Preibisch & Mamajek 2008). The association consists of three classic subregions first defined by Blaauw (1946), and later

refined by de Zeeuw et al. (1999): Upper Scorpius (US), Upper Centaurus-Lupus (UCL), and Lower Centaurus-Crux (LCC). The subgroups have mean distances of 145 pc, 140 pc, and 118 pc, respectively (de Zeeuw et al. 1999). Though many K/M type members of US have been identified in surveys by Walter et al. (1994), Preibisch et al. (1998), Preibisch & Zinnecker (1999), Preibisch et al. (2001), Preibisch et al. (2002), and Rizzuto et al. (2015), UCL and LCC occupy larger regions of the sky, and have received less attention, with only  $\sim$ 90 K/M stars identified in both subgroups, mostly in Mamajek et al. (2002), Preibisch & Mamajek (2008), Song et al. (2012), and a few new M-type members discovered by Rodriguez et al. (2011) and Murphy et al. (2015) using Galaxy Evolution Explorer (GALEX) UV observations. A deep DECam imaging survey of UCL and LCC is underway which is yielding dozens of members down to the deuterium-burning limit (Moolekamp et al., in prep).

In this study, we identify and characterize a new sample of K/M-type (~0.7-1.3  $M_{\odot}$ ) members of Sco-Cen selected through their X-ray emission and proper motions. We describe the results of a low-resolution spectroscopic survey to identify new low-mass K-and M-type members of Sco-Cen. We combine our newly identified members with members discovered by previous surveys to estimate the accretion disk fraction, and probe the star-formation history of each subgroup. Finally, we place these results in context with the results from other stellar populations.

## 2 SAMPLE SELECTION

We aim to find new K- and M- type members of all three subgroups of Sco-Cen. Low-mass pre-main sequence stars are X-ray luminous, with  $\log(L_x/L_{bol}) \simeq -3$ , due to their strong magnetic dynamos and coronal X-ray emission (Feigelson & Montmerle 1999). To build our candidate star list, we cross-referenced the PPMX propermotion catalog (Röser et al. 2008) with X-ray sources within 40" in the ROSAT Bright Source (Voges et al. 1999) and Faint Source (Voges et al. 2000) catalogs. We restricted our search to the de Zeeuw et al. (1999) boundaries for Sco-Cen and adopted their proper motion limits with an extra 10 mas  $yr^{-1}$  to allow for larger proper motion uncertainties and possible kinematic sub-structure. For objects at a distance of  $\sim 140$  pc, this extra 10 mas yr<sup>-1</sup> translates to  $\simeq$  7 km s<sup>-1</sup>. For US, UCL, and LCC this gives limits of  $\mu$  < 47 mas yr<sup>-1</sup>, 12 mas yr<sup>-1</sup> <  $\mu$  < 55 mas yr<sup>-1</sup>, and 15 mas yr<sup>-1</sup> <  $\mu$  < 55 mas yr^{-1}, respectively, with  $\mu_{lpha}$  < 10 mas yr^{-1} and  $\mu_{\delta} < 30$  mas yr<sup>-1</sup> for all three subgroups. We also required that our proper motions were less than 50% uncertain,  $\sigma_{\mu} < 0.5\mu$ , to avoid candidates with poorly constrained proper motions. The mean proper motion magnitude varies across the association due to its large extent on the sky and will also vary for stars of different distances, this being the principle behind kinematic parallaxes. The intrinsic J-K<sub>S</sub> color of an unreddened K0V dwarf is 0.48 mag (Pecaut & Mamajek 2013), and we aimed to find low-mass K- or M-type members of Sco-Cen so we further made color-magnitude cuts and required our candidates to have color  $J-K_S > 0.5$  mag and magnitude 7.0 < J < 11.0 from the 2MASS Point Source Catalog (Skrutskie et al. 2006). To avoid very large integration times on the SMARTS 1.5m telescope, we chose J < 11.0 mag, which corresponds to  $V \sim 12.5$  mag for an unreddened K0V dwarf. Stars brighter than  $J \sim 7.0$  mag would have been covered by previous surveys (e.g., Mamajek et al. 2002), so we chose J > 7.0 mag. At the time we performed our sample selection, accurate V-band magnitudes were not available for most stars in our parent sample, so

we used J-band magnitudes to filter the brighter and fainter ends of our parent sample. We removed candidates located in UCL just below US, in 343 < l < 350 and 0 < b < 10 in galactic coordinates. This region overlaps with "Lower Sco" (Mamajek 2013; Nguyen et al. 2013; Mamajek et al. 2013), and will be discussed in forthcoming papers (Mamajek et al., in prep.; Nguyen et al., in prep). This left us with 677 candidates, listed in Table 1. Before assembling a list of targets for observation, we searched the literature to record those stars which had been studied in previous surveys from which spectral type, Li or H $\alpha$  measurements were available in sufficient detail to make a membership determination. We then omitted stars which were studied in Mamajek et al. (2002), Preibisch et al. (1998), Preibisch et al. (2001), Preibisch et al. (2002), Ardila et al. (2000), Köhler et al. (2000), Slesnick et al. (2006), Krautter et al. (1997), Wichmann et al. (1997), Riaz et al. (2006), and Torres et al. (2006). This left us with 365 candidates in our spectroscopic target list. Objects PPMX J121431.8-511015, PPMX J134751.4-490148, PPMX J143751.3-545649, and PPMX J154348.8-392737 were not observed because they were within 60" of known Sco-Cen members. The first two may constitute new companions to Sco-Cen members MML 9 and MML 38, respectively (listed in Table 2). The status of the third (PPMX J143751.3-545649) with respect to MML 47 is unclear. The fourth, PPMX J154348.8-392737, is a previously catalogued companion to HD 140197 (listed as SEE 247 AB in the Washington Double Star catalog; Mason et al. 2001). This finally leaves 361 candidates which were spectroscopically observed.

2MASS	$\mu_{\alpha}$ (mas yr <sup>-1</sup> )	$\mu_{\delta}$ (mas yr <sup>-1</sup> )	Ref.	V (mag)	Ref.	<i>B–V</i> (mag)	Ref.	J (mag)	H (mag)	K <sub>S</sub> (mag)	Note
10004365-6522155	$-11.7 \pm 1.9$	$12.4\pm1.9$	РХ	$10.972\pm0.046$	A7	$0.650\pm0.077$	A7	$9.614 \pm 0.026$	$9.193 \pm 0.026$	$9.018 \pm 0.019$	
10065573-6352086	$\textbf{-19.6} \pm \textbf{1.5}$	$10.3\pm1.8$	U4	$10.950\pm0.012$	A7	$0.862\pm0.018$	A7	$9.262\pm0.028$	$8.744 \pm 0.061$	$8.580 \pm 0.024$	
10092184-6736381	$\textbf{-14.6} \pm \textbf{1.8}$	$19.6\pm1.0$	U4	$11.512 \pm 0.017$	A7	$0.849\pm0.051$	A7	$10.027\pm0.026$	$9.497\pm0.024$	$9.382\pm0.021$	
10111521-6620282	$\textbf{-26.5}\pm1.6$	$11.3\pm1.6$	U4	$12.242\pm0.028$	A7	$0.825\pm0.035$	A7	$10.680\pm0.024$	$10.223 \pm 0.026$	$10.073 \pm 0.023$	
10293275-6349156	$\textbf{-11.3}\pm1.4$	$14.7\pm1.5$	U4	$11.373 \pm 0.014$	A7	$0.822\pm0.027$	A7	$9.818\pm0.022$	$9.349\pm0.022$	$9.290\pm0.019$	

Table 1. Photometry and proper motion data for candidates in the Sco-Cen region

Adopted 2MASS JHK<sub>S</sub> magnitudes are PSF-fit photometry unless otherwise specified; (a) 2MASS JHK<sub>S</sub> aperture photometry; (b) 2MASS JHK<sub>S</sub> 6x catalog (Cutri et al. 2012); (c) 2MASS H psf photometry /  $JK_S$  aperture photometry;

References – (PX) PPMX, Röser et al. (2008); (U4) UCAC4, Zacharias et al. (2013); (T2) Tycho-2, Høg et al. (2000); (A6) APASS DR6 (Henden et al. 2012); (A7) APASS DR7 (Henden et al. 2012); (HP) Hipparcos, ESA (1997); (W97) Wichmann et al. (1997); (T06) Torres et al. (2006); Only the first five rows are shown; this table is available in its entirety in the electronic version of the journal.

Table 2. New Candidate Double Stars in Sco-Cen

Primary	Primary	Secondary	J	PA	sep.	epoch	Notes
Name	PPMX	PPMX	mags	deg	″	yr	
MML 9	121434.0-511012	121431.8-511015	8.7,10.0	261.34	21.17	2010.5589	1,2
MML 38	134750.5-490205	134751.4-490148	9.3,10.1	27.36	19.01	2010.5589	2

Notes: (1) The RAVE 4th data release (Kordopatis et al. 2013) reports radial velocities of  $12.6 \pm 3.5$  km s<sup>-1</sup> and  $11.3 \pm 3.6$  km s<sup>-1</sup> for the primary and secondary, respectively, supporting the notion that they constitute a physical pair. (2) Astrometry from AllWISE positions (Cutri & et al. 2014).

## **3 OBSERVATIONS AND DATA**

## 3.1 Spectra

Low-resolution red (~5600Å-6900Å) optical spectra were obtained from the SMARTS 1.5m telescope at Cerro Tololo Inter-American Observatory (CTIO), Chile. Observations were made in queue mode with the RC spectrograph between July 2009 and October 2010, and in classical mode during the nights of UT 7-17 April 2010. The spectra were taken with the "47/Ib" setup, which consists of a grating with groove density of 831 grooves  $mm^{-1}$ , blaze wavelength 7100Å, a GG495 filter and a slit width of 110.5 $\mu$ m, giving a spectral resolution of ~3.1 Å in the red optical. One comparison arc of Ne was taken immediately before three consecutive exposures of each target. The data were reduced using the SMARTS RC Spectrograph IDL pipeline written by Fred Walter<sup>1</sup>. The three object images are median filtered, bias-trimmed, overscan- and bias-subtracted, and flat-fielded. The spectrum is wavelength-calibrated using the Ne comparison frames. Finally, we normalize the spectra to the continuum with a low order spline in preparation for spectral classification.

In addition, we reanalyzed the low resolution spectra from the Mamajek et al. (2002) study. These blue and red spectra were taken at Siding Springs Observatory in UT 20-24 April 2000. The blue spectra have spectral resolution of ~2.8Å with spectral coverage of ~3850-5400Å. The red spectra have spectral resolution of ~1.3Å with spectral coverage of ~6200-7150Å. Further details regarding these spectra are described in Mamajek et al. (2002).

#### 3.2 Photometry

Our compiled photometry is listed in Table 1. Six of our candidates have BV photometry available from the Hipparcos catalog, and we adopted it where possible. For  $\sim 130$  candidates, we adopted BV photometry from the Tycho-2 Catalog (Høg et al. 2000), converted to the Johnson system using the conversions in Mamajek et al. (2002, 2006). For  $\sim$ 350 candidates, we adopted BV photometry from the AAVSO Photometric All-Sky Survey<sup>2</sup> (APASS) Data Release 6 and Data Release 7 (Henden et al. 2012). and the Search for Associations Containing Young stars catalog (SACY; Torres et al. 2006) for  $\sim 100$  candidates. For stars with V  $\leq 12$  mag, conservative estimates for SACY photometric uncertainties are 0.01 mag (C.A.O. Torres, 2012 private communication). We adopt  $JHK_S$ photometry from the Two Micron All-Sky Survey Point Source Catalog (2MASS; Skrutskie et al. 2006). We adopt 2MASS aperture photometry when the PSF is poorly fit as indicated by the quality flags ("qflg" other than 'A') in the 2MASS catalog. These data are indicated in the notes of Table 1.

For candidate members, we use mid-IR photometry from the Wide-field Infrared Survey Explorer (*WISE*; Wright et al. 2010; Cutri & et al. 2012). We visually examined all images of members which exhibit infrared excesses (identified in section 4.9.2) in one or more WISE bands. We flagged photometry which could be affected by blends with adjacent objects, unresolved binaries, or extended emission from nearby bright stars. We exclude photometry with unreliable detections or uncertainties exceeding 0.25 mag. We followed the scheme for evaluating the photometry in Luhman

& Mamajek (2012). Our mid-IR WISE photometry is listed in Table 3.

A few stars in our sample had WISE photometry contaminated by nearby objects. PDS 415 and CD-23 12840 are a  $\sim 4''$  binary, but only CD-23 12840 is in the PPMX catalog and thus only CD-23 12840 is in our sample. Both components of the pair are listed in the WISE All-Sky catalog but CD-23 12840 has  $K_S - W1 < 0.0$ . Examination of the WISE images shows the two are blended. PDS 415 is detected as an infrared source in the AKARI catalog (Ishihara et al. 2010) but CD-23 12840 is not detected. Thus we conclude that PDS 415 is the source of the infrared excess and the CD-23 12840 Ks band excess may be spurious. We exclude the CD-23 12840 WISE photometry. HD 326277 is blended with a nearby object in W1 and W2 and had no reliable detection in W3 and W4. HD 326277 has Spitzer 3.6µm and 4.5µm IRAC photometry in GLIMPSE ([3.6]=8.165±0.040 mag, [4.5]=8.178±0.044 mag; Churchwell et al. 2009) which differs from the WISE W1 and W2 by  $\sim 0.5$  mag. This suggests that the WISE photometry for this star is contaminated by blends and therefore we exclude the WISE photometry for HD 326277.

#### 3.3 Astrometry

Although our original selection scheme utilized proper motions from the PPMX catalog, we adopt proper motions for our analysis from several catalogs, including the Hipparcos catalog (van Leeuwen 2007), the Tycho-2 catalog (Høg et al. 2000), the PPMX catalog (Röser et al. 2008) and the UCAC4 catalog (Zacharias et al. 2013). At the time we performed our sample selection, the PPMX catalog the most complete homogeneous proper motion catalog with absolute proper motions (i.e., on the International Celestial Reference System, or ICRS; Röser et al. 2008). However, for our analysis we make use of proper motions for kinematic distance estimates, and therefore desire the most precise and well-constrained proper motions available. Therefore, for our calculations, we select the source catalog on a case-by-case basis, adopting the proper motions which have the smallest uncertainties. The majority of our adopted proper motions listed in Table 1 are from the UCAC4 catalog.

## 4 ANALYSIS

## 4.1 Spectral Classification

The optical spectra were visually classified against a dense grid of carefully chosen spectral standards<sup>3</sup> with spectral coverage from  $\sim$ 5600Å–6900Å. We use the same spectral standards and classification criteria as described in Pecaut & Mamajek (2013). The G/K stars are on the classification system of Keenan & McNeil (1989) and the M-type stars are on the classification system of Kirkpatrick et al. (1991).

While estimating temperature types for our sample we ignored the Na I doublet at  $\sim$ 5889/5896Å because it increases in strength with surface gravity, and is thus useful in discriminating between dwarfs and giants. For those pre-main sequence (pre-MS) members of Sco–Cen, we expected these to have a Na I doublet line similar to but weaker than dwarfs (Spinrad 1962; Lawson et al. 2009; Schlieder et al. 2012). Once a temperature type had been established, we compared the Na I doublet to that of a dwarf and a

<sup>&</sup>lt;sup>1</sup> http://www.astro.sunysb.edu/fwalter/SMARTS/smarts\_

<sup>15</sup>msched.html#RCpipeline

<sup>&</sup>lt;sup>2</sup> http://www.aavso.org/apass

<sup>3</sup> see http://www.pas.rochester.edu/~emamajek/spt/

giant of the same temperature subclass, assigning an appropriate luminosity class. In a few cases the star had a Na I doublet feature which closely resembled a giant (luminosity class III) but the relative strength of the Ca I at  $\lambda$  6102,  $\lambda$ 6122 and  $\lambda$ 6162 lines relative to the Fe I line at  $\lambda$ 6137 resembled that of a dwarf (luminosity class V). In these cases we assigned the intermediate luminosity class of a subgiant (luminosity class IV). For the early to mid G-type stars, the Na I doublet for subgiants is nearly indistinguishable from that of dwarfs (see Figure 2), so the Na I doublet was only used to assign luminosity classes from spectral type ~G5 to ~M3.

2MASS	WISE	W1	W2	W3	W4	$E(K_S - W1)$	$E(K_S - W2)$	$E(K_S - W3)$	$E(K_S - W4)$
J	J	(mag)	(mag)	(mag)	(mag)	(mag)	(mag)	(mag)	(mag)
10065573-6352086	100655.70-635208.6	$8.513 \pm 0.023$	$8.531 \pm 0.021$	$8.446 \pm 0.018$	$8.301\pm0.131$	$-0.025 \pm 0.033$	$-0.006 \pm 0.032$	$0.041\pm0.030$	$0.115\pm0.133$
10092184-6736381	100921.83-673637.9	$9.056\pm0.019$	$9.177\pm0.018$	$9.201\pm0.020$	$8.916\pm0.190$	$0.233\pm0.028$	$0.142\pm0.028$	$0.078\pm0.029$	$0.291\pm0.191$
10313710-6901587	103137.09-690158.7	$9.414\pm0.023$	$9.431\pm0.021$	$9.360\pm0.025$	$9.622\pm0.435$	$\textbf{-0.013} \pm 0.034$	$\textbf{-0.011} \pm 0.033$	$0.013\pm0.035$	$\textbf{-0.325} \pm 0.436$
10334180-6413457	103341.79-641345.6	$9.236\pm0.025$	$9.218\pm0.021$	$9.167\pm0.030$		$\textbf{-0.038} \pm 0.035$	$\textbf{-0.001} \pm 0.032$	$0.003\pm0.038$	
10412300-6940431	104122.96-694042.8	$8.304\pm0.022$	$8.330\pm0.020$	$8.216\pm0.015$	$7.858\pm0.082$	$\textbf{-0.008} \pm 0.030$	$0.012\pm0.028$	$0.093\pm0.025$	$0.382\pm0.084$

Table 3. Infrared photometry and Infrared Excesses for Members of Sco-Cen

All infrared photometry (*W*1, *W*2, *W*3 and *W*4) above is adopted from the *WISE* catalog (Cutri & et al. 2012). Only the first five rows are shown; this table is available in its entirety in the electronic version of the journal.

We also revised the spectral classifications for Sco-Cen members studied in Mamajek et al. (2002). The spectral types from Mamajek et al. (2002) were tied very closely to those of the Michigan Spectral Survey (Houk 1978, 1982). The Mamajek et al. (2002) spectra were re-classified by M. Pecaut during the survey of Pecaut & Mamajek (2013) using the standards of Keenan & McNeil (1989), and systematic differences were noted. In Table 4 we adopt the revised classifications for the Mamajek et al. (2002) stars. There are clearly systematic differences between the Michigan types and those on the modern MK system (as noted in Appendix C.1 of Pecaut & Mamajek 2013). Put simply, a G2V star classified in the Michigan survey corresponds more closely to a G0.5V on the modern grid of G-dwarf standards (Keenan & Mc-Neil 1989), as classified by Gray and collaborators (e.g. Gray et al. 2003, 2006). This is verified by comparison of the colors and effective temperatures of stars classified both by Houk and Gray et al. While the Michigan survey mostly relied on the MK system of Johnson & Morgan (1953) (including changes up through early 1970s), there were minor shifts to the MK system for stars hotter than G0 by Morgan (Morgan & Keenan 1973; Morgan et al. 1978), and for GK-type stars by Keenan & McNeil (1989) after the Michigan survey was initiated in the late 1960's. The modern M dwarf classifications rely on the standard sequence of Kirkpatrick et al. (1991) and its later additions (e.g. Henry et al. 2002). The spectral classification surveys of bright stars undertaken by Gray and collaborators (Gray et al. 2001, 2003, 2006) are based on the last generations of Morgan's and Keenan's hot and cool star spectral sequences (Morgan et al. 1978, Keenan & McNeil 1989) which are in common modern use. We compared stars of given spectral type in the Michigan survey to those of Gray et al. (2001, 2003, 2006), and in Table 5 we provide an estimate of the modern dwarf spectral type for given Michigan spectral types. Very few of the Michigan spectral classes appear to correspond closely to the same type of star on the modern MK grid (e.g. F6V, K0V, K3V being rare exceptions). Unfortunately, the differences are most pronounced in the early G-type dwarfs, where Michigan G2V corresponds to Gray's G0.5V, Michigan G3V corresponds to G1.5V, etc. This is partially due to Houk's choice of G dwarf standards (e.g. using  $\beta$  Com as their G2V standard, despite it being considered a G0V or F9.5V standard elsewhere), but also due to Keenan's minor adjustments to the G dwarf standards throughout the 1980's. As the Keenan & McNeil (1989) grid defines the modern GK dwarf sequence and is common use, classifications that used this grid are to be considered on the modern MK system following Table 5. For intercomparison of samples of stars classified in the Michigan survey with samples classified on the modern MK system, we recommend converting the Michigan spectral types to the modern MK system using the conversions in Table 5. None map exactly to G2V.

2MASS	SpT	Ref.	EW(Li)	Ref.			Members	Young Star	Sco-Cen		
			(Å)		Li?	μ?	$\log(g)$ ?	HRD?	Final	Ref.	Ref.
10004365-6522155	G3V	PM	0.24	PM	Y	Ν	Y?		Ν		
10065573-6352086	K0Ve	T06	0.35	T06	Y	Y		Y	Y	T06	PM
10092184-6736381	K1IV(e)	PM	0.32	PM	Y	Y	Y?	Y	Y	PM	PM
10111521-6620282	K0IV	PM	0.33	PM	Y	Y	Y?	Ν	Ν		
10293275-6349156	K1IV	PM	0.23	PM	Y?	Ν	Y?		Ν		

#### Table 4. Membership properties for Sco-Cen candidates

References in the final two columns refer to the first work to identify the star as young ("Young Star Ref.") and the first work to identify it as a member of Sco-Cen ("Sco-Cen Ref.").

References – (PM) - This work; (S49) - Struve & Rudkjøbing (1949); (H54) - Henize (1954); (D59) - Dolidze & Arakelyan (1959); (T62) - The (1962); (T64) - The (1964); (G67) - Garrison (1967); (S72) - Satyvaldiev (1972); (F75) - Filin & Satyvoldiev (1975); (P75) - Petrov & Satyvoldiev (1975); (H76) - Henize (1976); (S77) - Schwartz (1977); (S82) - Satyvoldiev (1982); (S86) - Stephenson (1986); (F87) - Finkenzeller & Basri (1987); (D88) - Downes & Keyes (1988); (He88) - Herbig & Bell (1988); (Ho88) - Houk & Smith-Moore (1988); (I89) - Ichikawa & Nishida (1989); (M89) - Mathieu et al. (1989); (S91) - Stocke et al. (1991); (V91) - Vazquez & Feinstein (1991); (B92) - Bouvier & Appenzeller (1992); (C92) - Carballo et al. (1992); (G92) - Gregorio-Hetem et al. (1992); (H94) - Hughes et al. (1994); (W94) - Walter et al. (1994); (R95) - Randich et al. (1995); (B96) - Brandner et al. (1996); (P96) - Park & Finley (1996); (F97) - Feigelson & Lawson (1997); (K97) - Krautter et al. (1997); (W97) - Wichmann et al. (1997); (M98) - Martin et al. (1998); (P98) - Preibisch et al. (1998); (S98) - Sciortino et al. (1998); (D99) - de Zeeuw et al. (1999); (P99) - Preibisch & Zinnecker (1999); (W99) - Wichmann et al. (1999); (H00) - Hoogerwerf et al. (2000); (K00) - Köhler et al. (2000); (W00) - Webb (2000); (P01) - Preibisch & Zinnecker (1999); (W99) - Wichmann et al. (2001); (M02) - Mamajek et al. (2002); (P02) - Preibisch et al. (2002); (R03) - Reid (2003); (M05) - Mamajek (2005); (T06) - Torres et al. (2006); (R06) - Riaz et al. (2001); (R11) - Rodriguez et al. (2011); (L12) - Luhman & Mamajek (2012); (M12) - Mamajek et al. (2012); (S12) - Song et al. (2012); (R15) - Rizzuto et al. (2015); (A) - measured from spectrum downloaded from the ESO archive, program 083.A-9003(A); (B) - measured from spectrum downloaded from the ESO archive, program 083.A-9003(A); (B) - measured from spectrum downloaded from the ESO archive, program 083.A-9003(A); (B) - mea

Only the first five rows are shown; this table is available in its entirety in the electronic version of the journal.

Table 5. Comparing Michigan Spec	ctral Types to Gray MK Spectral Types
----------------------------------	---------------------------------------

(1) SpT Houk		(2) SpT Gray	(3) SpT Houk		(4) SpT Gray	(5) SpT Houk		(6) SpT Gray
F0V F1V	$\rightarrow$ $\rightarrow$	F1V F2V	G0V G0/1V	$\rightarrow$	F9.5V F9.5V	K0V K0/1V	$\rightarrow$ $\rightarrow$	K0V K1V
F2V	$\rightarrow$	F3.5V	G0/2V	$\rightarrow$	G3:V	K1V	$\rightarrow$	K1.5V
F2/3V	$\rightarrow$	F3.5V	G1V	$\rightarrow$	G0+V	K1/2V	$\rightarrow$	K2V
F3V	$\rightarrow$	F4V	G1/2V	$\rightarrow$	G0V	K2V	$\rightarrow$	K2.5V
F3/5V	$\rightarrow$	F5V	G2V	$\rightarrow$	G0.5V	K2/3V	$\rightarrow$	K3V
F5V	$\rightarrow$	F5.5V	G2/3V	$\rightarrow$	G1V	K3V	$\rightarrow$	K3V
F5/6V	$\rightarrow$	F6V	G3V	$\rightarrow$	G1.5V	K3/4V	$\rightarrow$	K4V
F6V	$\rightarrow$	F6V	G3/5V	$\rightarrow$	G4V	K4V	$\rightarrow$	K4+V
F6/7V	$\rightarrow$	F6.5V	G5V	$\rightarrow$	G4.5V	K4/5V	$\rightarrow$	K5V
F7V	$\rightarrow$	F7.5V	G5/6V	$\rightarrow$	G6V	K5V	$\rightarrow$	K6-V
F7/8V	$\rightarrow$	F8V	G8V	$\rightarrow$	G8.5V	K5/M0V	$\rightarrow$	K7V
F8V	$\rightarrow$	F8.5V				M0V	$\rightarrow$	K6.5V
F8/G0V	$\rightarrow$	F9V						

## 4.2 Distance Calculation

Very few of our candidate members have measured trignometric parallaxes, so we estimate distances to each candidate by calculating a "kinematic" or "cluster" parallax (e.g., de Bruijne 1999a). This method uses the centroid space motion of the group with the proper motion of the candidate member and the angular separation to the convergent point of the group to estimate the distance to the individual member, with the assumption that they are co-moving. We emphasize that kinematic parallaxes are only meaningful for true members and meaningless for non-members. Kinematic parallaxes have been shown to reduce the scatter in the H-R diagram for cluster members over trignometric parallaxes (de Bruijne 1999b). We adopt the formalism and methods of Mamajek (2005) and adopt the updated Sco-Cen subgroup space motions listed in Chen et al. (2011). In addition to providing improved distance estimates over simply adopting the mean subgroup distances, the kinematic parallaxes allow us identify non-members and assess the likelihood that the candidate member is a bona-fide member.

#### 4.3 Membership Criteria

In order to identify likely members from our sample, we demand that all available data paint a consistent picture of association membership, and therefore consider several indicators to discriminate against interlopers. Based on previous surveys (see Preibisch & Mamajek 2008, and references therein), we expect the K and M-type stars to be pre-MS, and therefore have Li absorption stronger, on average, than that of a  $\sim$ 30-50 Myr-old population. We also expect that they will exhibit surface gravities intermediate between dwarfs and giants. We identify those stars which have the following:

(i) Appropriate levels of Li absorption in their spectra, using the Li 6708Å line,

(ii) Dwarf or subgiant surface gravities, using the Na I doublet at 5889/5896Å,

(iii) Kinematic distances consistent with Sco-Cen,

(iv) H-R diagram positions broadly consistent with membership in Sco-Cen (e.g., neither below the main sequence nor above a 1 Myr isochrone).

We discuss these criteria in detail in the following sections. Borderline cases were examined closely.

## 4.3.1 Lithium

Pre-MS stars transition from fully convective to having deep convective envelopes with radiative interiors during their contraction. These stars can mix Li from the stellar photosphere throughout the convection zone down to interior regions of the star where it is destroyed at  $T\gtrsim 2.6$  MK (Bodenheimer 1965; Strom 1994). Thus, for ~0.7-1.3 M<sub>☉</sub> stars like those targeted in this study, a high photospheric Li abundance is only present when the star is very young. Based on the published ages (de Geus et al. 1989; Preibisch & Zinnecker 1999; Mamajek et al. 2002; Pecaut et al. 2012), we expect members of Sco-Cen to exhibit stronger Li absorption lines at a given  $T_{eff}$ , on average and within some acceptable scatter, than a ~30-50 Myr sample.

We measured the equivalent width of the 6708Å Li feature for all stars in our spectroscopic sample. The values were measured Table 6. Polynomial fits to Li vs  $T_{eff}$  data

Polynomial Coefficient	IC 2602	Pleiades	Sco-Cen
$ \begin{array}{c} a_0\\ a_1\\ a_2\\ a_3 \end{array} $	$\begin{array}{c} -1.869522 \times 10^6 \\ 1.472855 \times 10^6 \\ -3.862308 \times 10^5 \\ 3.371432 \times 10^4 \end{array}$	$\begin{array}{c} -1.639425 \times 10^6 \\ 1.290057 \times 10^6 \\ -3.379855 \times 10^5 \\ 2.948343 \times 10^4 \end{array}$	$\begin{array}{c} -3.576491 \times 10^{6} \\ 2.890342 \times 10^{6} \\ -7.778982 \times 10^{5} \\ 6.972998 \times 10^{4} \end{array}$

Note:  $EW(Li)[mÅ] = a_0 + a_1(\log(T_{eff})) + a_2(\log(T_{eff}))^2 + a_3(\log(T_{eff}))^3$ Polynomial fit is valid from  $\log(T_{eff})=3.800$  dex to 3.580 dex.

with IRAF<sup>4</sup> with Vogt profiles after the spectrum was normalized to the continuum. Given our spectral resolution and the repeatability of our measurements, we estimate that our reported EW(Li) values are accurate to  $\sim 0.05$ Å. The 6708Å feature is a blend with a nearby Fe I line at 6707.44Å, unresolved at our resolution, and thus our EW(Li) values are probably overestimated by  $\sim 0.02$ Å. However, this is smaller than our uncertainties and we do not attempt to correct for this blend. Many stars in our parent sample were not observed because they had EW(Li) values available in the literature; for these stars we simply adopted the previously published values.

We compare our EW(Li) measurements at a given Teff to those in nearby open clusters. In Figure 1 we plot  $T_{\rm eff}\,$  versus EW(Li) for our sample along with data for the young open clusters IC 2602 (Randich et al. 1997) and the Pleiades (Soderblom et al. 1993; Jones et al. 1996). Most stars in our sample exhibit much larger EW(Li) at a given T<sub>eff</sub> than either IC 2602 (~45 Myr; Dobbie et al. 2010) or the Pleiades (~125 Myr; Stauffer et al. 1998). These data suggest that Li is largely undepleted for the  $\sim 10$ -16 Myr-old stars in Sco-Cen hotter than spectral type  $\sim$ K3, but the Li depletion becomes very strong for stars cooler than  $\sim$ K3  $({\sim}1.1~M_{\odot}).$  We fit low-order polynomials to the IC 2602 and Pleiades T<sub>eff</sub> versus EW(Li) data; the polynomial coefficents are listed in Table 6. For our Li-based membership criterion, if the EW(Li) was above the polynomial fit to IC 2602 we marked that star as 'Y' in Table 4; if EW(Li) was below the IC 2602 polynomial fit but above the Pleiades polynomial fit, we marked it as 'Y?', and if the EW(Li) is below the Pleiades polynomial fit we marked it as 'N'. Stars studied in Preibisch & Zinnecker (1999), Köhler et al. (2000) and Krautter et al. (1997) were confirmed as Li-rich but the measurements were not reported. We marked these candidates as 'Y' in Table 4. In total, we marked 482 candidates as 'Y', 153 as 'N', and 38 as 'Y?'.

## 4.3.2 Surface Gravity

Based on previous nuclear and pre-MS age determinations of the subgroups of Sco–Cen (Mamajek et al. 2002; Pecaut et al. 2012), the low-mass stars in Sco-Cen are expected to be pre-main sequence with surface gravities intermediate between dwarfs and giants. For candidates in our spectroscopic sample we can examine spectral features sensitive to surface gravity, primarily the Na I doublet at 5889Å/5896Å. Details on using Na I as a surface gravity indicator for identifying young stars is discussed in detail in Schlieder

<sup>4</sup> IRAF is distributed by the National Optical Astronomy Observatory, which is operated by the Association of Universities for Research in Astronomy (AURA) under cooperative agreement with the National Science Foundation.



**Figure 1.** Measured EW(Li) from the 6708Å line plotted against  $\log(T_{eff})$  for our X-ray sample. Plotted dashed lines are polynomial fits to data from surveys in IC 2602 (Randich et al. 1997) and the Pleiades (Soderblom et al. 1993; Jones et al. 1996). The location of the EW(Li) for our Sco-Cen candidate members is used to determine the membership indicator in Table 4. Those above the IC 2602 curve are marked "Y", those above the Pleiades curve are marked "Y", those above the Pleiades marked "Y"; the polynomial fit to the Sco-Cen candidates marked "Y"; the polynomial coefficients are listed in Table 6.

et al. (2012). The Na I doublet decreases in strength at a given  $T_{eff}$  as surface gravity decreases. However, the observed strength of the Na I doublet in subgiants among G-type stars is difficult to distinguish from dwarfs. The differences in the Na I doublet strength between dwarfs and subgiants becomes more useful in mid-K-type spectra, as shown in Figure 2. For this reason we are fairly conservative in our surface gravity membership criterion, marking G-type stars with luminosity classes of 'IV' with 'Y', 'V' with 'Y?' and 'III' with 'N'. Note that we only use this criterion for stars which we have observed spectroscopically; luminosity classifications from other authors for stars in our parent sample are not used for this criterion. We assigned 345 stars into these two categories, with 300 marked as 'Y?' and 45 marked as 'N'.

## 4.3.3 Distances

As mentioned previously, the majority of our sample lack trigonometric distance estimates. However, if we calculate a kinematic parallax estimate using the best available proper motion, we can then estimate distances for each star in our sample and verify that this lies within the range of expected distances for the association. Stars with discrepant kinematic distances are identified as non-members. We use the results of the membership study of Rizzuto et al. (2011), which makes use of the revised Hipparcos trignometric parallaxes (van Leeuwen 2007), to establish our distance criteria. Based on the spatial distribution of their candidate members, Rizzuto et al. (2011) concluded that strict boundaries between the subgroups are somewhat arbitrary, so we do not use distinct kinematic distance criteria for each subgroup. To establish parallax ( $\overline{\omega}$ ) limits based on galactic longitude and latitude (l, b), we fit a plane to the B-type members with membership probabilities >50% from the Rizzuto et al. (2011) study, obtaining:



**Figure 2.** Measured EW(Na) from the Na I doublet at 5889Å/5896Å line plotted against  $\log(T_{\rm eff})$  for our spectroscopic sample. Plotted are polynomial fits to dwarf and giant spectral standard stars used for classification.

$$\varpi = (-0.0208 \pm 0.0061) \times (l - < l >) + (-0.0299 \pm 0.0158) \times (b - < b >) + (7.7761 \pm 0.0969)$$

with mean galactic coordinates  $\langle l \rangle = 328.579^{\circ}$ ,  $\langle b \rangle = 13.261^{\circ}$ . The 1 $\sigma$  dispersion in the fit is  $\sigma_{\sigma} \simeq 1.25$  mas. Therefore we model the shape of Sco-Cen with a continuous plane and depth characterized by 1 $\sigma$  dispersion of ~1.25 mas. For our kinematic distance criteria, we mark stars with 'Y' in Table 4 if their kinematic parallaxes are within the 2 $\sigma$  (95% C.L.) dispersion from the plane described above, and 'Y?' if they are between 2 $\sigma$  and 3 $\sigma$ . We mark stars with 'N' if they are beyond 3 $\sigma$  of the dispersion from the plane. This criteria effectively provides different distance limits as a function of galactic longitude and latitude, giving typical 2 $\sigma$  distance limits of  $141^{+77}_{-37}$  pc,  $130^{+62}_{-32}$  pc, and  $117^{+49}_{-42}$  pc for US, UCL and LCC, respectively. 677 stars in our sample have been placed in these categories, with 495 marked as 'Y', 80 as 'Y?', and 102 as 'N'.

#### 4.3.4 HRD position

As a final membership check, we place these stars on the H-R diagram using their kinematic distances, which, for this purpose, *assumes* they are co-moving with other members in the association. Objects that have H-R diagram positions inconsistent with membership, e.g., below the main sequence, are rejected. Below we discuss the few stars which were Li-rich and fell inside our kinematic distance criteria but were rejected because they had discrepant H-R diagram positions. Two stars were rejected since they fell below the main sequence and two stars were rejected since they were well above the 1 Myr isochrone. Unresolved binarity is insufficient to account for these discrepant positions. Further analysis can be found in section 4.5.

## 4.3.5 Interlopers

*US:* HD 144610 is a Li-rich K1IV-III which has a kinematic parallax of  $\pi_{kin}$ =3.41±0.69 mas which is consistent with a H-R diagram position of log(T<sub>eff</sub>)=3.70±0.01, log(L/L<sub>☉</sub>)=1.42±0.18, well above the 1 Myr isochrone, so we reject it.

*UCL*: 2MASS J14301461-4520277 is a Li-rich K3IV(e) which, using the kinematic parallax of  $\pi_{\rm kin} = 10.92 \pm 0.92$  mas, has an H-R diagram position well below the main sequence  $(\log(T_{\rm eff})=3.66\pm0.02, \log(L/L_{\odot})=-0.91\pm0.08)$ , so we reject its membership to UCL. 2MASS J16100321-5026121 is a Li-rich K0III (Torres et al. 2006) which lies well above the 1 Myr isochrone  $(\log(T_{\rm eff})=3.70\pm0.01, \log(L/L_{\odot})=1.16\pm0.08)$ , calculated using the kinematic parallax of  $\pi_{\rm kin}=4.19\pm0.38$  mas, so we consider a UCL interloper.

*LCC:* 2MASS J10111521-6620282, lies well below the main sequence, with  $\log(T_{eff})=3.70\pm0.01$ ,  $\log(L/L_{\odot})=-0.86\pm0.09$ , calculated using the predicted kinematic parallax of  $\pi_{kin}=9.63\pm0.94$  mas. This is puzzling because it is a Li-rich KOIV, with a proper motion in excellent agreement with membership in LCC. It does not exhibit an infrared excess which could be a signature of an edge-on disk, and the B - V and  $V - K_S$  colors are consistent with a negligibly reddened young K0. However, it is  $\sim 0.4$  dex underluminous for the main sequence, so we consider it a likely LCC interloper.

## 4.3.6 Final Membership Assessment

Though we require the Li, surface gravity indicators, kinematic distance criterion, and H-R diagram criterion to indicate membership, the Li and kinematic distance criteria are the most restrictive and are responsible for identifying most non-members. Stars with 'Y' or 'Y?' in each of the four membership categories have been identified members of Sco-Cen. Though we have made every effort to remove interlopers, our list of candidate members may still contain a few non-members, though it will be dominated by true members. From our current sample, we identify 493 stars as likely members, listed in Table 7. The 180 stars rejected as Sco-Cen members are listed in Table 8. 156 are newly identified young stars.

Table 7. Stellar properties for Sco-Cen members

ID	2MASS	Grp.	SpT	Ref.	EW(Hα) (Å)	Ref.	A <sub>V</sub> (mag)	$\pi_{kin}$ (mas)	$\begin{array}{c} log(T_{eff}) \\ (dex) \end{array}$	$\begin{array}{c} log(L/L_{\odot}) \\ (dex) \end{array}$	Age (Myr)	$\begin{array}{c} Mass \\ (M_{\odot}) \end{array}$	Disk Type	Other Name
1	10065573-6352086	LCC	K0Ve	T06	0.0	T06	0.16±0.04	$7.54{\pm}0.83$	3.702±0.009	$-0.059 \pm 0.097$	14	1.1		TYC 8951-289-1
2	10092184-6736381	LCC	K1IV(e)	PM	-0.2	PM	$0.00{\pm}0.10$	$7.03{\pm}0.76$	$3.692 {\pm} 0.012$	$\text{-}0.334 {\pm} 0.096$	29	0.9		TYC 9210-1484-1
3	10313710-6901587	LCC	K2.5IV(e)	PM	-0.1	PM	$0.00{\pm}0.13$	$6.91 {\pm} 0.72$	$3.668 {\pm} 0.019$	$\text{-}0.380{\pm}0.095$	21	0.9		
4	10334180-6413457	LCC	K2.5IV(e)	PM	0.0	PM	$0.00{\pm}0.15$	$5.29{\pm}0.85$	$3.668 {\pm} 0.019$	$\text{-}0.077 {\pm} 0.143$	7	1.2		V542 Car
5	10412300-6940431	LCC	G8Ve	T06	0.0	T06	$0.00{\pm}0.02$	$11.05{\pm}0.97$	$3.717 {\pm} 0.007$	$-0.250{\pm}0.078$	33	0.9		TYC 9215-1181-1

 $EW(H\alpha) < 0$  denotes emission,  $EW(H\alpha) > 0$  denotes absorption. We report the Houk type for the star 16102888-2213477 = HD 145208 (G5V), but internally use the T<sub>eff</sub> and BC for the equivalent Gray type (G4.5V) in our calculations. See Table 5 for more information.

References – (PM) - This work; (G67) - Garrison (1967); (F87) - Finkenzeller & Basri (1987); (Ho88) - Houk & Smith-Moore (1988); (W94) - Walter et al. (1994); (H94) - Hughes et al. (1994); (W97) - Wichmann et al. (1997); (K97) - Krautter et al. (1997); (P98) - Preibisch et al. (1998); (P99) - Preibisch & Zinnecker (1999); (W99) - Wichmann et al. (1999); (K00) - Köhler et al. (2000); (P01) - Preibisch et al. (2001); (M02) - Mamajek et al. (2002); (P02) - Preibisch et al. (2002); (Z01) - Zuckerman et al. (2001); (T06) - Torres et al. (2006); (R06) - Riaz et al. (2006); (C06) - Carpenter et al. (2006); (W07) - White et al. (2007); (M11) - Müller et al. (2011); (D12) - Dahm et al. (2012); (A) - measured from spectrum downloaded from the ESO archive, program 083.A-9003(A); (B) - measured from spectrum downloaded from the ESO archive, program 081.C-0779(A);

Only the first five rows are shown; this table is available in its entirety in the electronic version of the journal.

Table 8. Objects rejected as Sco-Cen members

2MASS	SpT	Ref.	Rejection Reason	Other Names
10004365-6522155 10111521-6620282 10293275-6349156 10342989-6235572 10441393-6446273	G3V K0IV K1IV G7Ve K1V(e)	PM PM PM PM PM	μ HRD position μ μ	TYC 8964-165-1 HD 307772 HD 307960

References- (PM) - This work; (K97) - Krautter et al. (1997); (P98) - Preibisch et al. (1998); (K00) - Köhler et al. (2000); (M02) - Mamajek et al. (2002); (T06) - Torres et al. (2006);

Only the first five rows are shown; this table is available in its entirety in the electronic version of the journal.

## 4.4 Extinction

We estimate the reddening and extinction for Sco-Cen members using the spectral type-intrinsic color sequence for pre-MS stars described in Pecaut & Mamajek (2013). These intrinsic colors are calibrated using low-mass members of nearby moving groups which have negliglible extinction, and bracket the age range of the Sco-Cen subgroups (5-30 Myr). Many of the stars in our sample have very low reddening, and where we obtained a non-physical negative extinction we set the extinction to zero. We used a total-to-selective extinction of R<sub>V</sub>=3.1 and calculated E(B-V), E(V-J), E(V-H), and E(V-K<sub>S</sub>) with A<sub>J</sub>/A<sub>V</sub>=0.27, A<sub>H</sub>/A<sub>V</sub>=0.17, A<sub>K<sub>S</sub></sub>/A<sub>V</sub>=0.11 (Fiorucci & Munari 2003) which allowed us to estimate extinctions to each star individually using as many as four different colors. We adopted the median  $A_V$ , and adopted the standard deviation of the  $A_V$  values as a conservative estimate of their uncertainty. For the stars which lack reliable V band photometry, we estimate their extinctions using  $E(J-K_S)$ . Our extinction estimates are listed in Table 7.

#### 4.5 H-R Diagram for Sco-Cen Members

We place our Sco-Cen members on a theoretical H-R diagram in order to compare with theoretical models and obtain individual isochronal age estimates. We adopt the effective temperature scale  $(T_{eff})$  and bolometric correction  $(BC_V)$  scale from the Pecaut & Mamajek (2013) study<sup>5</sup>, which was constructed specifically for 5-30 Myr old pre-MS stars. This Teff and BC<sub>V</sub> scale was derived by fitting the spectral energy distributions of members of young, nearby moving groups to the BT-Settl theoretical atmospheric models (Allard et al. 2012). Though this Teff and BC scale is dependent on model atmospheres, the method used to develop the resultant Teff scale is in good agreement with Teff values derived from angular diameter measurements (e.g., Boyajian et al. 2012). We combine our individual extinction estimates together with our kinematic parallaxes and 2MASS J-band magnitudes to estimate the bolometric luminosities of our stars and place them on a theoretical H-R diagram. We compare our data to the pre-MS models of Dotter et al. (2008) in Figure 3.

There is considerable scatter in the H-R diagram so we construct empirical isochrones by plotting the median luminosity along the H-R diagram. Though there is some scatter, the relative age rank of the three groups is consistent with that found in Mamajek et al. (2002), Preibisch & Mamajek (2008), Pecaut et al. (2012); from oldest to youngest: LCC, UCL and US, with UCL and LCC approximately coeval. The other striking feature of the empirical isochrones is the mass-dependent age trend. The lower mass stars appear younger against the theoretical isochrones than the higher mass stars. This is the same mass-dependent age trend seen in other studies (e.g., Hillenbrand 1997; Hillenbrand et al. 2008; Bell et al. 2012, 2013, 2014). The likely origin of the mass-dependent age trend is difficulties in handling convection with magnetic fields in young, low-mass stars in the evolutionary models, perhaps due to missing physics (see e.g., Feiden 2016). At an age of 10 Myr and 15 Myr, our observational uncertainties yield individual age uncertainties of  $\pm 3$  Myr and  $\pm 4$  Myr, respectively.

<sup>5</sup> Pecaut & Mamajek (2013) adopted an absolute bolmetric magnitude for the Sun of  $M_{\odot}$  = 4.755 mag, ~ 0.015 mag higher than the recently drafted IAU bolometric magnitude scale (2015 IAU resolution B2). Thus the BC scale in Pecaut & Mamajek (2013) should be shifted down by ~0.02 mag to conform to the IAU scale. The luminosity estimates in this present work are on the 2015 IAU scale.



**Figure 3.** H-R diagram for X-ray selected Sco-Cen members with  $J-K_S > 0.50$  as described in the text. Circles, triangles and star symbols are US, UCL and LCC candidate members, respectively. Plotted for comparison are 5, 10, 15, 20 and 30 Myr isochrones from Dotter et al. (2008). Artificial scatter of  $\delta T_{\text{eff}}$ =±0.001 dex has been introduced for plotting purposes.



Figure 4. H-R diagram with empirical isochrones for US (solid), UCL (dashed) and LCC (dotted). All three subgroups exhibit a mass-dependent age trend.

## 4.6 Ages

The presence of massive stars in OB associations also allows for the opportunity for a comparison of ages obtained through different parts of the H-R diagram. As massive stars burn through their nuclear fuel, they will expand and leave the main sequence. At the same time, the pre-main sequence members of the OB association will be contracting towards the main sequence. The evolutionary models will predict ages for each of these segments of the H-R diagram, but different aspects of stellar physics are important in each of these segments. In this section, we examine the ages for both



Figure 5. Distribution of raw isochronal ages obtained for the Sco-Cen members of our sample using the Dotter et al. (2008) evolutionary models.

the low-mass pre-MS stars as well as the massive main sequence turn-off stars.

## 4.6.1 Pre-MS Ages

We estimate individual ages for our pre-MS Sco-Cen members by linearly interpolating between theoretical model isochrones. We use the Dartmouth models (Dotter et al. 2008), the Pisa models (Tognelli et al. 2011), the PARSEC models (Chen et al. 2014), and the Exeter/Lyon models (Baraffe et al. 2015). These models each assume slightly different stellar compositions, but are each calibrated to reproduce the H-R diagram position of the Sun at its current age. The Dartmouth evolutionary models adopt the Grevesse & Sauval (1998) solar composition with a protosolar He fraction of Y=0.2740, while the Pisa evolutionary models adopt the Asplund et al. (2005) solar composition with a protosolar He fraction of Y=0.2533. The PARSEC models adopt the Caffau et al. (2008, 2009) solar abundances with a initial He fraction of Y=0.284, whereas the Exeter/Lyon models use the Caffau et al. (2011) solar abundances, Y=0.271. The distributions of ages we obtain with the Dartmouth evolutionary models is shown in Figure 5. We obtain median ages of 5±2 Myr, 9±1 Myr and 8±1 Myr (standard error of the median) for US, UCL and LCC, respectively. Surprisingly, these ages are half the mean ages obtained from the F-type members of Sco-Cen (Pecaut et al. 2012). Given the large Teffdependent trend with age (Figure 4), we decide against adopting mean subgroup ages based on the K- and M-type members of Sco-Cen. A detailed discussion of our reasoning is available in section 5.1.

In order to extract useful age estimates for the subgroups, we re-examine the G-type pre-MS ages as well as the main-sequence turn-off ages. We do this for two reasons, other than the obvious desire to quote reliable mean ages. The first reason is that estimating realistic intrinsic age spreads requires a reliable age as free of systematic uncertainties as possible. We use Monte Carlo simulations to constrain our intrinsic age spreads, taking into account realistic binarity statistics and observational uncertainties, the details

of which are described in section 4.7. Because isochrones evenly spaced in age will not generally be evenly spaced in luminosity, a large intrinsic spread in luminosity at a young age could still be consistent with a small age spread, whereas the same intrinsic spread in luminosity at an older age would imply a larger intrinsic age spread. The second reason we reevalute the G-type ages and nuclear ages is that updated Teff and bolometric corrections (BCs) have been made available since the ages were last estimated in Mamajek et al. (2002). For the hot stars, Nieva (2013) used modern non-local thermodynamic equilibrium (NLTE) spectral synthesis models to re-evaluate the  $T_{\rm eff}\,$  and BC scale for the massive OB stars (34,000 K >  $T_{eff}$ > 15,800 K). Their results for dwarfs and subgiants are 1500-6000 K hotter than the scale of Napiwotzki et al. (1993) that was adopted in the study of Mamajek et al. (2002). As previously mentioned, Pecaut & Mamajek (2013) have constructed an updated T<sub>eff</sub> and BC scale applicable to the Sco-Cen G-type stars. Both of these studies should allow us to place Sco-Cen members on the H-R diagram with reduced systematic uncertainties.

#### 4.6.2 Isochronal Ages for G-type Pre-MS Stars

To estimate the ages of the G-type pre-MS stars, we collect the Gtype members from this study together with those in Walter et al. (1994), Preibisch & Zinnecker (1999), Mamajek et al. (2002), and Preibisch & Mamajek (2008). We supplement this with G-type stars studied in Torres et al. (2006) in the Sco-Cen field which are Lirich and have proper motions consistent with membership; G-type stars identified as members here conform to the membership criteria discussed in section 4.3. The stellar properties for these stars are listed in Table 7 and Table 9, together with the median mass and age obtained through a comparison of their H-R diagram positions with the evolutionary models of Dotter et al. (2008), Tognelli et al. (2011), Chen et al. (2014), and Baraffe et al. (2015).

Table 9: Stellar parameters for Sco-Cen G-Type stars

Object	2MASS	Spectral	Ref.	Region	$A_V$	$\log(T_{\text{eff}})$	$\log(L/L_{\odot})$	Age	Mass
Name		Туре			(mag)	(dex)	(dex)	(Myr)	$(M_{\odot})$
TYC 9216-0524-1	10495605-6951216	G8V	T06	LCC	$0.00 {\pm} 0.05$	$3.717 \pm 0.007$	$-0.382 \pm 0.069$	48	0.9
TYC 9212-1782-1	10594094-6917037	G6V	T06	LCC	$0.00 \pm 0.02$	$3.732 \pm 0.008$	$0.428 {\pm} 0.085$	6	1.6
CD-48 6632	11350376-4850219	G7V	T06	LCC	$0.00 \pm 0.15$	$3.709 \pm 0.008$	$0.048 \pm 0.075$	11	1.2
HIP 57524	11472454-4953029	G0V	PM	LCC	$0.18 \pm 0.04$	$3.782 \pm 0.004$	$0.413 \pm 0.061$	17	1.3
CPD-63 2126	12041439-6418516	G8V	T06	LCC	$0.00 \pm 0.04$	$3.717 \pm 0.007$	$0.012 \pm 0.071$	15	1.2
MML 3	12044888-6409555	GIV	PM	LCC	$0.38 \pm 0.06$	3.776±0.007	$0.484 \pm 0.084$	14	1.3
HIP 58996	12054748-5100121	GIV	PM	LCC	$0.15 \pm 0.04$	$3.776 \pm 0.007$	$0.407 \pm 0.057$	16	1.3
MML 7	12113815-7110360	GSV	PM	LCC	$0.23 \pm 0.02$	$3.740\pm0.009$	$0.338 \pm 0.067$	10	1.4
HIP 59854	12162/83-5008356	GIV	PM	LCC	$0.31 \pm 0.05$	3.776±0.007	$0.531 \pm 0.061$	13	1.4
MML 14	12211648-5317450	GIV	PM	LCC	$0.39 \pm 0.05$	3.776±0.007	$0.433 \pm 0.067$	15	1.3
MML 17	12223322-5353489	GUV	PM TOC		$0.24 \pm 0.03$	$3.782 \pm 0.004$	$0.448 \pm 0.000$	10	1.5
CD-54 4705	12242005-5445540	COV	100 DM		$1.00\pm0.39$	$3.740 \pm 0.009$	$0.398 \pm 0.085$	) 15	1.8
HIP 00883	12204003-3327195	COV	PM DM		$0.20\pm0.04$	$3.782 \pm 0.004$	$0.482 \pm 0.037$	13	1.5
HIP 00915 MML 20	12290224-0455000	G2V G0V	PM DM		$0.44 \pm 0.04$	$3.769 \pm 0.009$	$0.498 \pm 0.060$	12	1.4
MINIL 29 LID 62947	13023732-3439370	G4V	F M DM		$0.44 \pm 0.10$ 0.10 $\pm 0.02$	$3.782 \pm 0.004$ $3.750 \pm 0.000$	$0.237 \pm 0.072$ 0.220 $\pm 0.058$	12	1.2
CD 50 4620	13030330-0413332	G2V	T06		$0.19\pm0.03$	$3.750\pm0.009$	$0.320\pm0.038$	12	1.5
CD-39 4029 MML 32	13132810-0000443	GOV	DM		$0.10\pm0.04$ 0.75±0.14	$3.739 \pm 0.009$ $3.782 \pm 0.004$	$0.289 \pm 0.008$ 0.616 $\pm 0.102$	10	1.2
MML 32	13220446 4503231	GOV	DM		$0.75\pm0.14$	$3.782\pm0.004$ $3.782\pm0.004$	$0.010\pm0.102$ 0.220±0.057	25	1.5
HIP 65/23	132/3512-55572/2	G3V	T06		$0.10\pm0.03$	$3.752\pm0.004$ $3.759\pm0.009$	$0.220\pm0.057$ 0.332 $\pm0.057$	14	1.2
HIP 65517	13254783-4814577	GIV	PM		$0.00\pm0.02$ 0.23 $\pm0.05$	$3.759 \pm 0.007$ $3.776 \pm 0.007$	$0.332 \pm 0.037$ 0.104 \pm 0.53	28	1.5
MMI 35	13342026-5240360	GOV	PM		$0.23 \pm 0.03$ 0.38 $\pm 0.07$	$3.770\pm0.007$ $3.782\pm0.004$	$0.104\pm0.053$ $0.478\pm0.058$	15	1.1
MML 37	13432853-5436434	GOV	PM	LCC	$0.33\pm0.04$	$3.782\pm0.004$ $3.782\pm0.004$	$0.478\pm0.053$ 0.213 $\pm0.053$	25	1.5
HIP 67522	13500627-4050090	GOV	PM	UCI	$0.33\pm0.04$ 0.23 $\pm0.04$	$3.782\pm0.004$ $3.782\pm0.004$	$0.213 \pm 0.053$ 0.233 $\pm 0.063$	23	1.2
MML 42	14160567-6917359	G1V	PM	LCC	$0.17\pm0.02$	$3.776\pm0.007$	$0.253 \pm 0.003$ $0.262 \pm 0.059$	27	1.2
HIP 71178	14332578-3432376	G9IV	PM	UCL	$0.12\pm0.02$	$3.709\pm0.008$	$0.031\pm0.057$	12	1.2
MML 49	14473176-4800056	G7IV	PM	UCL	$0.00\pm0.04$	$3.723 \pm 0.007$	$-0.109\pm0.067$	24	1.0
MML 52	14571962-3612274	G3V	PM	UCL	$0.43 \pm 0.03$	$3.759 \pm 0.009$	$0.305\pm0.062$	15	1.3
MML 56	15011155-4120406	G0V	PM	UCL	$0.32 \pm 0.04$	$3.782 \pm 0.004$	$0.613 \pm 0.068$	12	1.5
MML 57	15015882-4755464	G0V	PM	UCL	$0.24{\pm}0.04$	$3.782 \pm 0.004$	$0.333 {\pm} 0.071$	20	1.2
MML 61	15125018-4508044	G2V	PM	UCL	$0.31 \pm 0.04$	$3.769 {\pm} 0.009$	$0.165 {\pm} 0.082$	24	1.1
MML 62	15180174-5317287	G6V	PM	UCL	$0.31 {\pm} 0.01$	$3.732{\pm}0.008$	$0.041 {\pm} 0.063$	18	1.1
TYC 8298-1675-1	15193702-4759341	G9IV	T06	UCL	$0.00{\pm}0.02$	$3.709 {\pm} 0.008$	$-0.336 \pm 0.069$	38	0.9
HIP 75483	15251169-4659132	G9V	T06	UCL	$0.00{\pm}0.16$	$3.709 {\pm} 0.008$	$-0.220 \pm 0.067$	27	1.0
HIP 75924	15302626-3218122	G3V	PM	UCL	$0.33 {\pm} 0.11$	$3.759 {\pm} 0.009$	$0.572{\pm}0.063$	8	1.6
HIP 76472	15370466-4009221	G0V	PM	UCL	$0.52{\pm}0.03$	$3.782{\pm}0.004$	$0.560{\pm}0.060$	13	1.4
MML 69	15392440-2710218	G5V	PM	UCL	$0.34{\pm}0.11$	$3.740{\pm}0.009$	$0.451 {\pm} 0.070$	7	1.6
HIP 77135	15445769-3411535	G3V	PM	UCL	$0.43 {\pm} 0.04$	$3.759{\pm}0.009$	$0.066 {\pm} 0.064$	27	1.1
HIP 77144	15450184-4050310	G0V	PM	UCL	$0.19{\pm}0.10$	$3.782{\pm}0.004$	$0.407 {\pm} 0.060$	18	1.3
HIP 77190	15454266-4632334	G5IV	T06	UCL	$0.00{\pm}0.03$	$3.740{\pm}0.009$	$-0.200 \pm 0.059$	39	0.9
MML 72	15465179-4919048	G7V	PM	UCL	$0.13 {\pm} 0.04$	$3.723 {\pm} 0.007$	$0.136{\pm}0.064$	12	1.3
HIP 77656	15511373-4218513	G7V	PM	UCL	$0.00 {\pm} 0.01$	$3.723 \pm 0.007$	$0.339 {\pm} 0.062$	7	1.6
TYC 7846-1538-1	15532729-4216007	G2	T06	UCL	$0.88 {\pm} 0.35$	$3.769 \pm 0.009$	$0.382 {\pm} 0.067$	15	1.3
HD 142361	15545986-2347181	G2IV	W94	US	$0.46 \pm 0.12$	$3.769 \pm 0.009$	$0.314 \pm 0.070$	18	1.2
HD 142506	15554883-2512240	G3	P99	US	$0.40 \pm 0.03$	$3.759 \pm 0.009$	$0.432 \pm 0.076$	11	1.4
HIP 78133	15571468-4130205	G3V	T06	UCL	$0.00 \pm 0.07$	$3.759 \pm 0.009$	$0.063 \pm 0.066$	27	1.1
HD 143099	15595826-3824317	G0V	T66	UCL	$0.00 \pm 0.05$	$3.782 \pm 0.004$	$0.449 \pm 0.066$	16	1.3
CD-24 12445	16000078-2509423	G0	P99	US	$0.37 \pm 0.17$	$3.782 \pm 0.004$	$0.256 \pm 0.074$	23	1.2
TYC 7333-1260-1	16010792-3254526	GIV	T06	UCL	$0.21 \pm 0.06$	3.776±0.007	$0.371 \pm 0.061$	17	1.2
MML 75	16010896-3320141	G5IV	PM	UCL	$0.81 \pm 0.05$	$3.740 \pm 0.009$	$0.480 \pm 0.066$	6	1.6
HIP 78483	16011842-2652212	G2IV	106	US	$0.33 \pm 0.07$	$3.769 \pm 0.009$	$0.710\pm0.077$	8	1.7
HIP 78581	16024415-3040013	GIV	H82	US	$0.08 \pm 0.05$	$3.7/6\pm0.007$	$0.353 \pm 0.069$	18	1.2
ПГ /9402 ММ 92	10120035-2319406	G2V COV	H88	05	$0.33 \pm 0.13$	5.709±0.009	$0.88/\pm0.077$	5	2.0
MML 82	10211219-4030204	GV	PM	UCL	$0.00\pm0.07$	$3./1/\pm0.007$	$0.204 \pm 0.063$	8	1.4
MML 85	16232955-3958008		PM	UCL	$0.48 \pm 0.12$	$3.7750 \pm 0.007$	$0.212 \pm 0.084$	24	1.1
111P 00520	10233383-2940401	COV	100	05	$0.03 \pm 0.02$	$3.739 \pm 0.009$	$0.439 \pm 0.071$	11	1.4
HIP 80636	10202991-2/41203	GOV	П62 DM	US	$0.00\pm0.03$	$3.782 \pm 0.004$ $3.782 \pm 0.004$	$0.121\pm0.011$ 0.563 $\pm0.060$	9 12	1.0
HID 81380	16271286 2000201	GOV G2V	DM	UCL	0.30±0.03	$3.762 \pm 0.004$ 3.760 \pm 0.00	$0.303 \pm 0.000$ 0.708 $\pm 0.060$	15	1.4 1.9
HIP 81447	16380553-3401106	GOV	PM	UCL	$0.12 \pm 0.16$	$3.782 \pm 0.004$	$0.814 \pm 0.065$	8	1.7

G-type stars not included in Table 7. Ages and masses were estimated using the median of Baraffe et al. (2015), Chen et al. (2014), Tognelli et al. (2011) and Dotter et al. (2008) evolutionary models. We report the Houk types for the stars HIP 78581 (G1V), HIP 79462 (G2V) and HIP 80535 (G0V), but adopt the  $T_{eff}$  and BC for the equivalent Gray types in our calculations. See Table 5 for the conversions and section 4.1 for discussion.

Spectral Type references: (PM) This work; (T06) Torres et al. (2006); (W94) Walter et al. (1994); (P99) Preibisch & Zinnecker (1999); (T66) Thackeray (1966); (H82) Houk (1982); (H088) Houk & Smith-Moore (1988)

Adopted median subgroup ages (Table 11) are preferable to uncertain individual ages listed above.

## 4.6.3 Isochronal Turn-off Ages for Massive Stars

To estimate nuclear ages, we collect membership lists for the most massive stars from de Zeeuw et al. (1999), Mamajek et al. (2002), Rizzuto et al. (2011) and Pecaut et al. (2012). We collect Stromgren  $ubvy\beta$  and Johnson UBV photometry from Hauck & Mermilliod (1998) and Mermilliod & Mermilliod (1994). We estimate the extinction towards each early B-type member with both the Q-method (Johnson & Morgan 1953; updated in Pecaut & Mamajek 2013) using Johnson UBV photometry, and the prescription of Shobbrook (1983), using Stromgren  $ubvy\beta$  photometry. For all the massive stars these two methods give similar  $A_V$  within the uncertainties, so we adopt the mean value. We calculate the  $T_{eff}\,$  and  $BC_{\it V}$  for each star using the calibrations with Q, [u-b], [c1] and  $\beta$  in Nieva (2013); we adopt the median  $T_{eff}$  when available. We note that the calibrations derived by Nieva (2013) give systematically hotter T<sub>eff</sub> than the calibrations of Napiwotzki et al. (1993) and Balona (1994), which were used in Mamajek et al. (2002) and Pecaut et al. (2012), respectively, in the most recent estimation of Sco-Cen nuclear ages. Because of this we expect to obtain younger turn-off ages than previously estimated.

Our stellar parameters for the main-sequence turn-off in Sco-Cen are listed in Table 10, along with an individual age estimate of each object from a comparison the rotating tracks of Ekström et al. (2012), rotating at  $v_{eq} = 0.4 v_{crit}$ , with the H-R diagram position of the stars. A plot of the main-sequence turnoff for all three subgroups is shown in Figure 6. To estimate the turnoff age of Upper Sco we use  $\tau$  Sco,  $\omega$  Sco,  $\sigma$  Sco,  $\beta^1$  Sco,  $\pi$  Sco, and  $\delta$  Sco. We obtain a median turn-off age for US of  $\sim$ 7 Myr. We concur with de Zeeuw et al. (1999) that o Sco is unlikely to be a US member.<sup>6</sup> For Upper Centaurus-Lupus, we use  $\mu$  Cen,  $\delta$  Lup,  $\alpha$  Lup,  $\mu^1$  Sco,  $\mu^2$  Sco,  $\beta$  Lup,  $\gamma$  Lup,  $\nu$  Cen,  $\eta$  Cen,  $\eta$  Lup,  $\phi$  Cen,  $\varepsilon$  Lup,  $\kappa$  Cen, and HR 6143. We obtain a median age for UCL of  $\sim$ 19 Myr. For Lower Centaurus-Crux, we use  $\alpha^1$  Cru,  $\beta$  Cru,  $\beta$  Cen,  $\delta$  Cru and  $\alpha$  Mus. We obtain a median age of  $\sim 11$  Myr. Note that these are all in the southern part of LCC, for which Preibisch & Mamajek (2008) estimated  $\sim$ 12 Myr. There are no turnoff stars in the northern part of LCC, the hottest of which appear to be older ( $\sim 20$  Myr). Our nuclear median subgroup ages are summarized in Table 11.



**Figure 6.** Theoretical H-R diagram for the upper main sequence members of Sco-Cen with isochrones from the rotating evolutionary models of Ekström et al. (2012) with solar metallicity (Z = 0.014).

 $^6$  o Sco is an A5II (Gray & Garrison 1989) with a parallax of 3.71±0.54 mas (van Leeuwen 2007). Its evolutionary status, proper motion, and distance (~270 pc) are inconsistent with membership in Upper Sco.

Object	Spectral	Ref.	Region	$A_V$	$\log(T_{eff})$	$\log(L/L_{\odot})$	Age
Name	Туре			(mag)	(dex)	(dex)	Myr
$\tau$ Sco	B0V	1	US	$0.109{\pm}0.029$	$4.505 {\pm} 0.004$	4.351±0.125	1
$\delta$ Sco	B0.5IV	1	US	$0.432{\pm}0.025$	$4.479 {\pm} 0.030$	$4.664{\pm}0.133$	7
$\beta^1$ Sco	B0.5V	1	US	$0.547{\pm}0.031$	$4.464{\pm}0.021$	$4.394{\pm}0.096$	7
$\sigma$ Sco	B1III	1	US	$1.203 {\pm} 0.009$	$4.466 {\pm} 0.028$	$5.019 {\pm} 0.126$	8
ωSco	B1V	1	US	$0.647 {\pm} 0.013$	$4.454{\pm}0.016$	$4.013 {\pm} 0.048$	1
$\pi$ Sco	B1V+B2:V:	1	US	$0.204{\pm}0.005$	$4.437 {\pm} 0.030$	$4.411 \pm 0.119$	10
1 Sco	B1.5Vn	1	US	$0.462{\pm}0.008$	$4.355 {\pm} 0.016$	$3.490{\pm}0.049$	5
v Sco	B2IV	1	US	$0.756 {\pm} 0.021$	$4.341 {\pm} 0.012$	$3.800 {\pm} 0.100$	23
$\beta^2$ Sco	B2IV-V	1	US	$0.538{\pm}0.025$	$4.340{\pm}0.018$	$3.190{\pm}0.130$	<1
$\rho$ Sco	B2IV-V	1	US	$0.081{\pm}0.033$	$4.335 {\pm} 0.032$	$3.561{\pm}0.074$	18
13 Sco	B2V	1	US	$0.112 {\pm} 0.005$	$4.300 {\pm} 0.022$	$3.234{\pm}0.052$	15
$\alpha$ Lup	B1.5III	2	UCL	$0.094{\pm}0.009$	$4.360{\pm}0.014$	$4.237 {\pm} 0.037$	19
$\delta$ Lup	B1.5IV	2	UCL	$0.030{\pm}0.017$	$4.384{\pm}0.033$	$4.455{\pm}0.145$	15
$\mu^1$ Sco	B1.5IV	2	UCL	$0.060 {\pm} 0.010$	$4.363 {\pm} 0.030$	$4.011 {\pm} 0.142$	19
$\eta$ Cen	B1.5Vn	2	UCL	$0.021{\pm}0.021$	$4.349 {\pm} 0.021$	$3.796 {\pm} 0.024$	21
β Lup	B2III	2	UCL	$0.045 {\pm} 0.008$	$4.354{\pm}0.018$	$3.891{\pm}0.043$	20
HR 6143	B2III	2	UCL	$0.161 {\pm} 0.016$	$4.326 {\pm} 0.019$	$3.580{\pm}0.050$	23
v Cen	B2IV	2	UCL	$0.025{\pm}0.003$	$4.386{\pm}0.031$	$3.772 {\pm} 0.070$	8
$\mu^2$ Sco	B2IV	2	UCL	$0.051{\pm}0.014$	$4.356 {\pm} 0.033$	$3.724 {\pm} 0.072$	17
$\phi$ Cen	B2IV	2	UCL	$0.000 {\pm} 0.019$	$4.345 {\pm} 0.030$	$3.666 {\pm} 0.069$	19
γLup	B2IV	2	UCL	$0.018 {\pm} 0.003$	$4.337 {\pm} 0.030$	$3.880{\pm}0.085$	24
κCen	B2IV	2	UCL	$0.015 {\pm} 0.008$	$4.306 {\pm} 0.029$	$3.591{\pm}0.082$	30
$v^1$ Cen	B2IV-V	2	UCL	$0.009 {\pm} 0.020$	$4.318 {\pm} 0.027$	$3.410 {\pm} 0.062$	18
εLup	B2IV-V	2	UCL	$0.035 {\pm} 0.022$	$4.277 {\pm} 0.029$	$3.692{\pm}0.114$	> 30
μ Cen	B2IV-Ve	2	UCL	$0.257 {\pm} 0.030$	$4.456 {\pm} 0.040$	$4.163 {\pm} 0.092$	4
χ Cen	B2V	2	UCL	$0.018 {\pm} 0.012$	$4.302 {\pm} 0.025$	$3.342{\pm}0.061$	23
$\eta$ Lup	B2.5IV	2	UCL	$0.003 {\pm} 0.003$	$4.352 {\pm} 0.031$	$3.685 {\pm} 0.070$	17
τLib	B2.5V	2	UCL	$0.028 {\pm} 0.006$	$4.242 {\pm} 0.031$	$3.218 {\pm} 0.071$	> 30
θ Lup	B2.5Vn	2	UCL	$0.031 {\pm} 0.020$	$4.248 {\pm} 0.035$	$3.106 {\pm} 0.092$	> 30
HR 5471	B3V	2	UCL	$0.045 {\pm} 0.002$	4.251±0.029	$3.034{\pm}0.064$	> 30
HR 5378	B7IIIp	2	UCL	$0.038 {\pm} 0.012$	$4.273 \pm 0.019$	$3.152 \pm 0.049$	28
β Cru	B0.5III	2	LCC	$0.077 {\pm} 0.006$	$4.458 {\pm} 0.014$	$4.414 {\pm} 0.079$	8
$\alpha^1$ Cru	B0.5IV	2	LCC	$0.018 {\pm} 0.041$	$4.436 {\pm} 0.038$	$4.664 \pm 0.094$	11
β Cen	B1III	2	LCC	$0.075 {\pm} 0.031$	$4.428 {\pm} 0.022$	$4.905 {\pm} 0.072$	11
$\xi^2$ Cen	B1.5V	2	LCC	$0.061 {\pm} 0.005$	$4.308 {\pm} 0.029$	$3.329 \pm 0.069$	18
HR 4618	B2IIIne	2	LCC	$0.043 \pm 0.009$	$4.247 \pm 0.010$	$3.058 {\pm} 0.075$	> 30
$\delta$ Cru	B2IV	2	LCC	$0.000 {\pm} 0.007$	$4.403 {\pm} 0.025$	$3.838 {\pm} 0.056$	6
$\alpha$ Mus	B2IV-V	2	LCC	$0.015 {\pm} 0.015$	4.351±0.024	$3.696 \pm 0.054$	18
$\mu^1$ Cru	B2IV-V	2	LCC	$0.067 \pm 0.009$	$4.297 {\pm} 0.025$	$3.309 \pm 0.057$	24
$\sigma$ Cen	B2V	2	LCC	$0.060 \pm 0.022$	4.303±0.032	$3.358 {\pm} 0.073$	23
ζ Cru	B2.5V	2	LCC	$0.021 \pm 0.021$	4.246±0.025	$3.048 {\pm} 0.070$	> 30
-							

Table 10. Stellar parameters for Sco-Cen main sequence turnoff stars

Ages were estimated using the Ekström et al. (2012) rotating evolutionary models with solar abundances. Spectral Type references: (1) Hiltner et al. (1969); (2) Garrison (1967);

 Table 11. Adopted subgroup ages

Method	US	UCL	LCC
	(Myr)	(Myr)	(Myr)
MS Turnoff	$7\pm 2$	$19\pm 2$	$11\pm 2$
G-Type Pre-MS	10 $\pm 1$	$15\pm 1$	$15\pm 1$
F-Type Pre-MS <sup>a</sup>	13 $\pm 1$	$16\pm 1$	$17\pm 1$
Adopted	10±3	16±2	15±3

Uncertainties reported above are the standard error of the mean which represents the uncertainty in how well the mean value is characterized; these numbers do not represent the spread in ages. Median ages are derived considering the B-type main-sequence turn-off, the F-type pre-MS turn-on and the pre-MS G-type stars in each subgroup.

<sup>a</sup> Adopted F-type Pre-MS ages from Pecaut et al. (2012).

We summarize our derived nuclear, F-type pre-MS and G-type pre-MS ages in Table 11 with our adopted values for each subregion.

#### 4.7 Intrinsic Age Spreads

The H-R diagram positions of Sco-Cen members have a large degree of scatter, and hence a large apparent scatter in inferred ages. Some of this scatter is due to observational uncertainties and unresolved multiplicity, but some scatter may be due to true age spreads within the subgroups. Previous studies have found very small intrinsic age spreads in US, but larger age spreads in UCL and LCC (Preibisch et al. 2002; Slesnick et al. 2006; Mamajek et al. 2002; Preibisch & Mamajek 2008; Pecaut & Mamajek 2013). We perform Monte Carlo simulations in order to model the scatter caused by the observational uncertainties, the effects of unresolved binarity, and an intrinsic age spread. We create many populations of 10<sup>4</sup> simulated members with a gaussian distribution of ages, assuming a Kroupa (2001) initial mass function (IMF), a spot filling factor from 0% to 50%, a multiplicity fraction of 0.44 and companion mass ratio distribution given by a flat power law distribution with  $\gamma=0.3$  (using multiplicity properties for  $0.7 \,\mathrm{M_{\odot}}$  $M_* \lesssim 1.3 M_{\odot}$  population I main sequence stars as summarized in Duchêne & Kraus 2013 and Raghavan et al. 2010). To create the simulated population, each star is assigned a mass and an age, determined to be either binary or single, assigned a companion mass if binary, and assigned a spot coverage ratio from a random uniform distribution ranging from 0% to 50%, which alters its H-R diagram position according to the Somers & Pinsonneault (2015) correction factors. We then introduce dispersion in their H-R diagram positions with the median observational uncertainties from our sample. This is designed to simulate a population with a given mean age, an intrinsic age spread, the effects of unresolved binarity, cool spots, and observational uncertainties. Following Hillenbrand et al. (2008), we compare the the observed luminosity spreads around the empirical isochrones in Sco-Cen members to the simulated luminosity spreads. We emphasize that in this comparison, we only compare the distributions of luminosities around the median. We do not compare the median ages obtained since we observe a massdependent age trend. However, unlike Hillenbrand et al. (2008), we only model star formation with a given mean age and intrinsic (gaussian) spread, rather than consider accelerating star formation or other distributions of ages.



**Figure 7.** Observed luminosity spreads around the empirical isochrone (solid line) compared with the same for a simulated population (dashed line). The simulations used mean ages of 10 Myr, 16 Myr and 15 Myr for US, UCL, and LCC, respectively. We obtained best-fit intrinsic age spreads of  $\pm$ 7 Myr,  $\pm$ 7 Myr,  $\pm$ 6 Myr for US, UCL and LCC, shown above with simulations made with the Baraffe et al. (2015) evolutionary models.

Table 12. Subregion intrinsic age spreads

Evolutionary Models	US at 10 Myr (Myr)	UCL at 16 Myr (Myr)	LCC at 15 Myr (Myr)
Exeter/Lyon	$\pm 8$	$\pm 8$	$\pm 6$
Parsec	$\pm 6$	±7	$\pm 6$
Pisa	$\pm 8$	$\pm 8$	$\pm 6$
Dartmouth	$\pm 6$	±7	$\pm 5$
Adopted Intrinsic Age Spreads	±7	±7	±6

Age spreads were estimated comparing observed luminosity spreads to simulated populations with a given intrinsic age spread, taking into account multiplicity, spots and observational uncertainties. Adopted median ages are listed in Table 11.

We compare our observed luminosity spread with the luminosity spread from the simulations using a Anderson-Darling goodness-of-fit test (e.g., Hou et al. 2009), and adopt the age spread which best matches the simulated luminosity spreads for each Sco-Cen subgroup. We adopted the median subgroup ages listed in Table 11 as the mean age of our simulated populations to compare the luminosity spreads in our simulations to those from our observations. We adopt 1 $\sigma$  gaussian age spreads of  $\pm 7$  Myr,  $\pm 7$  Myr and  $\pm 6$  Myr for US, UCL and LCC, respectively. A plot showing the best matches to the observed luminosity spreads is shown in Figure 7. Our results are summarized in Table 12. These age spreads indicate that 68% of star formation in each subgroup occured over a period of ~14 Myr for US and UCL, and over a period of ~12 Myr for LCC.

## 4.8 Spatial Variation of Ages

Blaauw (1964) divided Sco-Cen into the subgroups Upper Scor-

pius, Upper Centaurus-Lupus and Lower Centaurus Crux. However, Rizzuto et al. (2011) notes that, based on their updated membership study of Sco-Cen, the distribution of probable members indicates that the subgroups cannot be defined in a non-arbitrary manner. US has consistently been shown to be younger than the other two subgroups, and UCL and LCC have typically been assigned similar ages (Preibisch & Mamajek 2008; Mamajek et al. 2002; Pecaut et al. 2012). Given the large spatial extent of UCL and LCC, however, it is too simplistic to place them into two groups, each characterized by a single mean age. For example, the  $\sim 50$  pc size of LCC and adopted mean age of  $\sim 17$  Myr leads to an expected individual age uncertainty of  $\sim$ 50%, or  $\sim$ 8 Myr, based on a star formation timescale-size relation (see discussion in Soderblom et al. 2014). Here we attempt to examine the age structure of the entire association to discern if we see evidence for spatial substructure based on systematic differences in age as a function of position on the sky.

One possible method to investigate spatial variations of ages would be to simply evaluate the age of each member against pre-MS evolutionary tracks and spatially average their ages to create an age map. However, given the strong mass-dependent age trend we find with all the evolutionary tracks, this would tend to show regions with large numbers of low-mass stars as younger regions, when in fact this may be a systematic effect due to the evolutionary tracks. Therefore we do not use this method.

Systematically younger regions will tend to be more luminous than the mean association luminosity as a function of  $T_{eff}$ . This is comparison does not depend on *any* theoretical models. To probe for statistically significant spatial variations of ages, *independent* of any evolutionary tracks, we look for concentrations of stars which lie above or below the average luminosity as a function of  $T_{eff}$ . We use all the pre-MS members which have been studied spectroscopically. We place all stars from all three subgroups together on the H-R diagram and construct an empirical isochrone for the entire association, by fitting a line to  $log(L/L_{\odot})$  as a function of  $T_{eff}$ . This is plotted in Figure 8. Each star lies on the H-R diagram above or below this empirical isochrone. For each star we calculate this offset in units of the luminosity spread  $\sigma$  above or below the linear fit:

offset =  $(\log(L/L_{\odot}) - \langle \log(L/L_{\odot}) \rangle) / \sigma_{\log(L/L_{\odot})}$ 

We then spatially average their offset, above or below the empirical isochrone, in galactic coordinates, shown in Figure 9. Regions on the sky with stars which are systematically more luminious than the average of the association will lie systematically above the empirical isochrone and will appear on the age map as younger. Likewise, stars in older regions will tend to be less luminous on average than the average of the association. In Figure 9, we plot a spatial intensity map of the median offset from the empirical isochrone. The colored regions in Figure 9 correspond to the same colored regions above or below the empirical isochrone in Figure 8. Spatially averaging their offset from the empirical isochrone allows us to determine which regions have concentrations of systematically younger or older stars, relative to the age of the entire association, and independent of any theoretical models. This method avoids biases due to any mass-dependent age trends, and can identify younger or older regions without reference to any evolutionary tracks. The median distance of the F- and G-type stars is within  $\sim$ 4 pc of the median distance of the K-type stars in each subgroup, and is not systematically biased nearer or farther. Furthermore, the variable extinction across the association ranges from  $\sim 0.0 - 0.5$  mag (25% to 75% interquartile range), which would not significantly bias the spatial distribution of ages. Thus it is likely



**Figure 8.** Empirical isochrone for 657 F/G/K-type members of all subgroups of Sco-Cen. We fit a line to create the empirical isochrone. We use individual stars' offsets above or below this empirical isochrone to create a relative age map, shown in Figure 9. The colored regions shown here correspond to the ages on the map in Figure 9.

that our selection methods do not exhibit any systematic spatial biases for younger or older ages across the association that would bias the age map.

We wish to assign mean isochronal ages of these regions using evolutionary models. In order to assign ages to the regions shown on the map, we use the F5 through G9 stars to anchor the regions to an age. For example, the F5 through G9 stars that lie  $0.33\sigma$  to  $0.66\sigma$  below the median isochrone, irrespective of their location in the association, have a median age of 18 Myr. Therefore, any region with a median offset of  $0.33\sigma$  to  $0.66\sigma$  below the empirical isochrone is assigned an age of ~18 Myr. The age map makes use of 657 F-M pre-MS stars to establish *relative* ages from the mean Sco-Cen empirical isochrone, but the ages are adopted from the F5-G9 stars. We discuss our motivations for adopting the F- and G-type ages in section 5.1.



q







Figure 10. Representative spectra of new Sco-Cen members with strong  $H\alpha$  emission consistent with accretion.

#### 4.9 Circumstellar Disks

## 4.9.1 Spectroscopic Accretion Disk Fraction

Giant planet formation is fundamentally limited by the lifetimes of gas-rich protoplanetary disks surrounding the host star (Pollack et al. 1996). The gas disk dissipation timescale therefore provides an upper limit to the giant planet formation timescale. Differences in disk dissipation timescales for stars in different mass bins can provides critical data for inferring how the planet formation process differs around stars of various masses. Additionally, a census for gas-rich circumstellar disks allows for follow–up studies of the gas disk itself (e.g., Zhang et al. 2013) or the star–disk interaction (e.g., Bouvier et al. 2007; Nguyen et al. 2009).

Here we perform a census of accretion disks for our sample using H $\alpha$  emission as an accretion diagnostic. Various criterion have been proposed using H $\alpha$  as an accretion indicator, and we adopt the spectral type dependent empirical criterion of Barrado y Navascués & Martín (2003). If our measured or adopted EW(H $\alpha$ ) (see Table 7) exceeds the Barrado y Navascués & Martín (2003) criterion, we count the object as an accretor. Our accretion disk fraction excludes the 28 Sco-Cen members in our sample which lack H $\alpha$  measurements. Sample spectra for Sco-Cen members with H $\alpha$  in emission consistent with accretion are shown in Figure 10.

Using the Barrado y Navascués & Martín (2003) criteria, we identified 10/108 accretors in US, or a spectroscopic accretion disk fraction of  $9.3^{+3.6}_{-2.1}\%$ . In UCL and LCC we have 5/154 and 4/127 accretors, or spectroscopic accretion disk fractions of  $3.2^{+2.1}_{-0.9}\%$  and  $3.1^{+2.4}_{-0.9}\%$ , respectively. These disk fractions include all K and M-type members in our sample with EW(H $\alpha$ ) measurements, and therefore have masses predominantly ~0.7-1.3 M<sub> $\odot$ </sub>, with a few M-type stars as low as ~0.5 M<sub> $\odot$ </sub>; for statistics on the K-type members *only*, refer to Table 13.

## 4.9.2 Infrared Excess Disk Fraction

Infrared photometry can be used to identify the presence of circumstellar matter with different wavelengths used to probe matter at



**Figure 11.** H– $K_S$  versus J–H colors for new candidate Sco-Cen members. The dwarf and pre-MS stellar locus from Pecaut & Mamajek (2013) are included shown as the dashed and solid lines, respectively.



**Figure 12.** Spectral type versus dereddened  $H-K_S$  color for Sco-Cen members. The solid line is the pre-MS photospheric colors from Pecaut & Mamajek (2013). The dashed line is the  $3\sigma$  dispersion in the photospheric colors. Members with color excesses above the  $3\sigma$  dispersion in the photospheric colors in this and all bands at longer wavelength are identified as having a  $K_S$  band excess (gray stars).

different temperatures. To probe cooler, dusty debris around young stars,  $20\mu$ m and longer wavelengths are useful. We examine Sco-Cen members for excesses in  $H-K_S$ ,  $K_S-W1$ ,  $K_S-W2$ ,  $K_S-W3$ , and  $K_S-W4$  colors, shown in Figure 12 and Figure 14, in order to identify Sco-Cen members exhibiting IR excesses which may indicate the presence of a disk and allow us to infer its evolutionary state. These stars are also plotted against H $\alpha$  equivalent width (EW(H $\alpha$ )) in Figure 13, demonstrating that accreting stars identified with the Barrado y Navascués & Martín (2003) EW(H $\alpha$ ) criteria also typically exhibit a W1 band excess due to the presence of a warm cir-

Band/Criteria/Disk Type	US	UCL	LCC
EW(Hα)	6/84 (7.1 <sup>+3.9</sup> %)	$5/145 (3.4^{+2.2}_{-1.0}\%)$	4/119 (3.4 <sup>+2.5</sup> %)
$K_S$	$6/89 (6.7^{+3.7}_{-1.8}\%)$	$6/158 (3.8^{+2.2}_{-1.0}\%)$	$2/119(1.7^{+2.1}_{-0.5}\%)$
W1	$11/89(12.4^{+4.3}_{-2.7}\%)$	$9/157 (5.7^{+2.4}_{-1.3}\%)$	$4/119(3.4^{+2.5}_{-1.0}\%)$
W2	$14/89(15.7^{+\overline{4.6}}_{-3.1}\%)$	$9/157 (5.7^{+2.4}_{-1.3}\%)$	$4/119(3.4^{+2.5}_{-1.0}\%)$
W3	$24/89(27.0^{+5.2}_{-4.1}\%)$	$17/157 (10.8^{+3.0}_{-2.0}\%)$	$8/118 (6.8^{+3.1}_{-1.6}\%)$
W4	$25/36(69.4^{+6.5}_{-8.5}\%)$	$33/72 \ (45.8^{+\overline{5.9}}_{-\overline{5.6}}\%)$	8/86 (9.3 <sup>+4.1</sup> <sub>-2.2</sub> %)
Full	$8/89~(9.0^{+4.0}_{-2.2}\%)$	8/157 (5.1 <sup>+2.4</sup> %)	$4/118(3.4^{+2.5}_{-1.0}\%)$
Transitional	$2/89 (2.2^{+\overline{2.8}}_{-0.7}\%)$	0/157 (<2.3%; 95% C.L.)	0/118 (<3.0%; 95% C.L.)
Evolved	$2/89 (2.2^{+2.8}_{-0.7}\%)$	$1/157 (0.6^{+1.4}_{-0.2}\%)$	0/118 (<3.0%; 95% C.L.)

Table 13. Infrared excess and spectroscopic accretion disk fractions for K-Type ( $\sim$ 0.7-1.3 M $_{\odot}$ ) members in Sco-Cen

Spectroscopic Accretion and IR Excess fractions shown above for K-type members of Sco-Cen only, with masses  $\sim$ 0.7-1.3 M $_{\odot}$ .



**Figure 13.**  $K_S$ –W2 color versus EW(H $\alpha$ ) for Sco-Cen members. Members spectroscopically identified as accretors with the Barrado y Navascués & Martín (2003) criteria are shown as stars. Members exhibiting a  $K_S$ –W2 color excess are shown as large circle.

cumstellar gas disk. We identify stars above the  $3\sigma$  dispersion in the young stellar locus, as defined in Pecaut & Mamajek (2013), as having an excess in that band. We use  $3\sigma$  as a conservative criterion, to avoid reporting excesses with a small confidence level. To avoid reporting false excesses due to scatter in the photometry, we only report an infrared excess if that object is also above the  $3\sigma$ dispersion in the young stellar locus for that band *and* all bands at longer wavelengths.

Disks in young populations such as Sco-Cen may be found in a variety of stages of evolution. Here we attempt to classify the disks into the disk classification scheme described by Espaillat et al. (2012), using the observational criteria described by Luhman & Mamajek (2012). Based on the boundary between the full disks and transitional, evolved and debris disks for the stars classified in Luhman & Mamajek (2012), we identify "full disks" as those with  $E(K_S-W3) > 1.5$  and  $E(K_S-W4) > 3.2$ , "transitional disks" as those with  $E(K_S-W4) > 3.2$  but have  $E(K_S-W3) < 1.5$ . We then identify "evolved disks" as those with  $E(K_S-W4) > 3.2$  and  $E(K_S-W3) > 0.5$  and "debris disks" as those with  $E(K_S-W4) < 3.2$  and  $E(K_S-W3) < 0.5$ . Color excesses and classifications are shown in Figure 15. Four stars had excesses in W1, W2 or W3 but no reliable W4. We classify them as evolved or debris disks based on their lack of spectroscopic signatures of accretion and their  $E(K_S-W2)$  and  $E(K_S-W3)$ .

Disks classified using this method are listed in Table 7. Our sample contains a small number of G- and M-type members of Sco-Cen. Therefore, in Table 13 we summarize infrared excess fractions for only K-type ( $\sim 0.7$ -1.3 M<sub> $\odot$ </sub>) Sco-Cen members in the 2MASS  $K_S$  band and the four WISE bands.

## 5 DISCUSSION

#### 5.1 Which Ages are Reliable?

Depending on which isochronal ages we examine, we obtain different mean subgroup ages. However, given the Teff-dependent age trend present in Figure 4, we see several reasons to distrust ages from M-type stars. The most obvious reason is that evolutionary models have difficulty predicting the radii of main-sequence M-type stars, systematically underestimating their radii by  $\sim$ 5-20% (Torres & Ribas 2002; Boyajian et al. 2012; see also Kraus et al. 2011). Hillenbrand & White (2004) have compared dynamically constrained masses with predictions from pre-MS evolutionary models and found that the models systematically underpredict the masses of stars 5%-20% for masses under 0.5  $M_{\odot}.$  On the other hand, they found that for masses above 1.2  $M_{\odot}$ , dynamical masses and predicted masses from all models are consistent. As stated in Hillenbrand et al. (2008), the likely source for the poor performance of models in the low-mass regime is incomplete knowledge of opacity sources and the difficulty in modeling convection. One of the large sources of uncertainty particularly applicable to modeling low-mass stars is the role magnetic fields play in convection, as discussed in Feiden (2016). Strong magnetic fields can give rise to large star spots on the surface of the star, which will may cause the star to exhibit more than one surface temperature (Stauffer et al. 2003). Somers & Pinsonneault (2015) have studied the effect of star spots on inferred ages and masses in pre-main sequence stars, and find that spots will tend to inflate the radii of young stars and shift them to a cooler T<sub>eff</sub>. The effect is that spots will make older stars appear to be younger and less massive than implied by the evolutionary models. Somers & Pinsonneault (2015) offer age and mass correction factors which can be used to estimate less biased, more accurate masses and ages from published evolutionary tracks. Given the problems with radii and mass discrepancies from models



**Figure 14.** Spectral type versus infrared colors  $K_S$ –W1,  $K_S$ –W2,  $K_S$ –W3, and  $K_S$ –W4 for Sco-Cen members. The solid lines are the pre-MS photospheric colors from Pecaut & Mamajek (2013). Members with color excesses above the  $3\sigma$  dispersion of photospheric colors in that band and all bands at longer wavelength are identified as having excess emission from circumstellar disks. Objects with color uncertainties greater than 0.25 mag are not shown.

as well as the potential influence of spots, and that there are highermass F- and G-type isochronal ages available, it is preferable to avoid the adopting ages from the K- and M-type members.

What about kinematic ages? Kinematic "expansion" or "traceback" ages are not dependent on stellar interior models and therefore offer the prospect of nearly model-free ages. Song et al. (2012) argues that kinematic expansion ages can function as modelindependent age benchmarks which can then be used to establish a model-indepedent age scale. Age indicators such as Li can then be used to establish relative ages among different stellar populations. This is an interesting idea in principle, however there are a few major issues with kinematic traceback ages.

One major problem is that one must assume that the association was in a physically smaller configuration at some time in the past. The data presented in section 4.8 indicates substantial substructure in Sco-Cen. This is consistent with work in other OB associations, for example, Cygnus OB2 also exhibits substantial substructure (Wright et al. 2014, 2016), and was never a compact, bound cluster. The presences of substantial substructure invalidates the assumption that the members of these associations were in more compact configurations in the past, which prevents the determination of a meaningful kinematic age.

Another major issue is that the results from kinematic ages are very sensitive to the implementation. A recent example is that in the TW Hydra Association (TWA). A commonly quoted age for TWA is 8 Myr based on the kinematic traceback of de la Reza et al. (2006). However, their kinematic traceback was based on a sample of only four stars, which was contaminated by TWA 19, a member of LCC (Mamajek et al. 2002). Mamajek (2005) calculated a kinematic expansion age for TWA using kinematic parallaxes and a vetted list of members and obtained a lower limit of  $\sim 10$  Myr on the expansion age at 95% confidence, though the data were only weakly consistent with expansion. More recently, Weinberger et al. (2012) performed a kinematic traceback of TWA using a vetted list of members and trignometric parallaxes. The Weinberger et al. (2012) traceback result indicated the members were never in a significantly more compact configuration. Another TWA study, Ducourant et al. (2014), independently obtained trignonmetric parallaxes for 13 stars, identify 31 as a co-moving association, 25 of which had radial velocity and trignometric parallax data. This study obtained a traceback age for TWA of 7.5±0.7 Myr. However, this result was based on a sample of 16 stars with converging motions, obtained after removing 9 stars which systematically drifted from the center of the association when traced back in time. Similarly,



Figure 15. Color excesses above the photosphere for stars in Sco-Cen exhibiting an infrared excess.

an often-quoted age for Upper Sco is the 5 Myr expansion age derived from proper motion data by Blaauw (1978). However, it was demonstrated by Brown et al. (1997), using simulations of expanding OB associations, that expansion ages inferred from proper motions alone all converged to  $\sim$ 4 Myr, no matter the actual kinematic age. A recent examination of the expansion age in Upper Sco by Pecaut et al. (2012), using radial velocity data, gave a lower limit of  $\sim 10$  Myr at 95% confidence, though the data were consistent with no expansion. Finally, we mention that the adopted kinematic expansion age of the  $\beta$  Pictoris moving group (BPMG) of 12 Myr, estimated by Song et al. (2003), has been re-evaluated by Mamajek & Bell (2014). The Mamajek & Bell (2014) study found the modern BPMG kinematic data was only weakly indicitave of expansion, and that the age is only weakly constrained by the kinematic data, giving a 95 confidence limit of 13-58 Myr. We conclude that the there is no well-constrained kinematic traceback age for either Sco-Cen or the groups used to bracket its age (e.g., TWA, BPMG) that has withstood the scrutiny of improved data, and that they simply do not yield useful age constraints given the current precision of the available data.

Another relevant chronometric technique is the use of the lithium depletion boundary (LDB) to determine the age of a stellar population. By detecting the stellar T<sub>eff</sub> or luminosity above which all the stars have exhibit Li depletion and comparing this with evolutionary model predictions, one can obtain an age which is independent of distance. LDB ages have been calculated for several of the nearby, young moving groups (Mentuch et al. 2008; Binks & Jeffries 2014) but the subgroups of Sco-Cen do not yet have a reliable LDB age. The results of Cargile et al. (2010) suggests that lithium depletion boundary ages and modern nuclear main sequence turn-off ages are in agreement when convective core overshooting is included in the models of high-mass stars (e.g., Ekström et al. 2012). However, LDB ages are typically much older than pre-MS contraction ages, and it has been suggested that this problem is related to the radii discrepancy in M-type stars (Yee & Jensen 2010; Somers & Pinsonneault 2015).

Recent discoveries of eclipsing binaries, particularly in Upper Scorpius using the data obtained using the Kepler K2 mission, should help evolutionary models significantly by providing well-constrained radii and masses at these young ages. Particularly, the recently published discoveries by David et al. (2015), Alonso et al. (2015), Lodieu et al. (2015), and Kraus et al. (2015), will add a significant number of benchmark eclipsing binaries with tightly constrained, nearly model-independent parameters for objects in Upper Scorpius. Masses and radii for the eclipsing binary UScoC-TIO 5 from Kraus et al. (2015), when compared to the Baraffe et al. (2015) evolutionary models, already provide some concordance with the older ~10 Myr ages from the F- and G-type stars, though more theoretical work remains to be completed.

### 5.2 How "Coeval" are the Sco-Cen Subgroups?

Previous studies of Sco-Cen have attempted to quantify the observed age spread in the subgroups. In Upper Sco, Preibisch et al. (2002), adopting an age of 5 Myr, concluded that the age spread was  $\leq$ 1-2 Myr. Their results account for the effects of binarity, a distance spread, and observational uncertainties. This is consistent with the results of Slesnick et al. (2006), who similarly constrained the age spread in the northern part of Upper Sco to be less than  $\pm$ 3 Myr (uniform distribution) using similar assumptions. In their study of UCL and LCC, Mamajek et al. (2002) have examined age spreads in the older subgroups and have constrained the star formation to have occurred over a time period of  $\pm$ 3 Myr and  $\pm$ 2 Myr (1 $\sigma$ ) for UCL and LCC, respectively.

Our age spreads are larger than those previously reported, with  $1\sigma$  age spreads of  $\pm 7$  Myr,  $\pm 7$  Myr, and  $\pm 6$  Myr for US, UCL and LCC, respectively. Our age spread of  $\pm 7$  Myr for Upper Sco is much larger than the age spreads detected by Preibisch et al. (2002) and Slesnick et al. (2008). However, the age map in Figure 9 indicates that there is an age gradient from the southeastern part of US to the northwest, with the northwestern part being younger. The Slesnick et al. (2008) and Preibisch et al. (2002) lowmass samples were drawn from smaller regions ( $\sim 150 \text{ deg}^2$  and  $\sim 160 \text{ deg}^2$ , respectively) than our sample in Upper Sco (drawn from the entire  $\sim 320 \text{ deg}^2$ ), which could be responsible for the smaller detected age spreads. However, the likely reason our inferred age spreads in US are larger than previous results is that we adopt a mean age of 10 Myr, twice as old as the Slesnick et al. (2008) or Preibisch et al. (2002) studies. A given luminosity spread at a younger age corresponds to a smaller inferred dispersion in ages than the same luminosity spread at an older age, simply because the younger isochrones are spaced farther apart in luminosity than those at older ages.

Slesnick et al. (2008) suggests that spreads in H-R diagram positions may not be an accurate proxy for spreads in age. They demonstrate this by comparing two nearly identical spectra for members in US, with spectroscopic surface gravity indicators which are indicitive of nearly identical surface gravity. However, their H-R diagram positions suggests their ages differ by more than 10 Myr! Jeffries et al. (2011) use constraints on the disk lifetime to show that the age spread in the Orion Nebula Cluster must be less than 0.14 dex in log(Age), though the age spreads inferred from the H-R diagram show a 0.4 dex dispersion in log(Age). These results suggest that scatter in the H-R diagram may not be a reliable indicator of age spreads. However, the H-R diagram remains the best observational indicator available at the present time for revealing any intrinsic age spreads. See Soderblom et al. (2014) for a more detailed and complete discussion regarding age spreads.

As stated previously, our age map does indicate there is a clear age gradient in US as it merges into UCL. The southern part of LCC is also noticably younger than other parts of LCC, confirming the suggestion first raised in Preibisch & Mamajek (2008). We note that there are no main-sequence turnoff stars in northern LCC, which, considering the younger age of southern LCC, accounts for the turn-off age of LCC being much younger than UCL. The star formation history of Sco-Cen, as inferred from the H-R diagram positions, appears to be more complex than previously treated. The spatial distribution of ages is suggestive that the current division of three distinct, coeval subgroups is overly simplistic and a separation into smaller units may be warranted. We avoid speculating further on scenarios of triggered star formation here, leaving that discussion to a future study.

#### 5.3 Circumstellar Disk Census

Observations of young clusters and associations of ages from  $\sim 1$ to >100 Myr have given strong indication that the protoplanetary disk dispersal timescale is very short, with an e-folding time of  $\sim$ 2.5 Myr (Mamajek 2009; Fedele et al. 2010). This is qualitatively consistent with what we seen in Sco-Cen: 9% of Upper Sco K-type stars host an optically thick protoplanetary disk ("Full Disk" in the Espaillat et al. 2012 nomenclature) at an age of  $\sim 10$  Myr, whereas  $\simeq 4\%$  of the K-type stars in the older subgroups UCL and LCC host a full disk (Table 13). However, the e-folding time of 2.5 Myr was estimated adopting an Upper Sco age of 5 Myr, along with many other young clusters. Naylor (2009) has argued that pre-MS ages systematically underestimate cluster ages, and that ages based on high-mass stars, typically double the ages estimated from the lowmass stars, are more likely to be correct (Bell et al. 2013). Using the ages we adopt for the Sco-Cen subgroups, what disk dispersal timescale does this imply? We plot the primordial disk fractions ("Full Disk" from Table 13) as a function of age in Figure 16 and, following Mamajek (2009), fit an exponential decay curve to the data ( $f_{disk} = e^{-t/\tau_{disk}}$ ). We obtain a mean protoplanetary disk efolding timescale of 4.7 Myr for K-type stars ( $\sim 0.7$ -1.3 M $_{\odot}$ ). This is  $\sim 2$  Myr longer than the timescale estimated in Mamajek (2009), and would imply a longer timescale available for planet formation, but consistent with the findings of Bell et al. (2013).

Finally, we note that our K-type disk fraction is larger than what is observed for the higher mass-stars in the same subgroups (Mamajek et al. 2002; Pecaut et al. 2012; Carpenter et al. 2009; Luhman & Mamajek 2012). This is consistent with the massdependent trend first identified by Carpenter et al. (2006) and fur-



**Figure 16.** Mean subgroup age versus disk fraction ("Full Disk"; see Table 13) for K-type stars, ~0.7-1.3 M<sub>☉</sub>, from the three subgroups of Sco-Cen. The best-fit exponential decay curvey  $f_{disk} = e^{-t/\tau_{disk}}$  has  $\tau_{disk}$ =4.7 Myr.

ther confirmed by Luhman & Mamajek (2012) in Upper Sco, as well as the results of Hernández et al. (2007) from the younger  $\sigma$  Ori association at ~6 Myr (Bell et al. 2013). Ribas et al. (2015) have summarized these mass-dependent disk fraction trends using the nearby young stellar associations within ~500 pc, and have come to a similar conclusion – that the disk dispersal time depends on stellar mass, with the low-mass stars retaining their disks longer. For Sco-Cen, however, we still have very poor statistics for M-type stars in the older subgroups, which highlights the need for future surveys to push the membership census to cooler spectral types in UCL and LCC (e.g., Murphy et al. 2015).

### 6 CONCLUSIONS

We can summarize the findings from our survey as follows:

(i) We have performed a survey for new, low-mass K- and Mtype members of all three subgroups of Sco-Cen. Using Li, X-ray, and proper motion data, we identify 156 new pre-MS members of Sco-Cen.

(ii) Using our newly identified members together with previously known members of Sco-Cen, we utilize H $\alpha$  as an accretion diagnostic and identify stars with H $\alpha$  emission levels consistent with accretion. We estimate a spectroscopic accretion disk fraction of  $7.1^{+3.9}_{-1.9}\%$ ,  $3.4^{+2.2}_{-1.0}\%$  and  $3.4^{+2.5}_{-1.0}\%$  for solar analog pre-MS K-type stars (~0.7-1.3 M<sub>☉</sub>) in US, UCL and LCC, respectively, consistent with a protoplanetary disk decay e-folding timescale of ~4.7 Myr, or half-life of ~3.3 Myr.

(iii) Similar to previous results in other star-forming regions (e.g., Hillenbrand 1997; Hillenbrand et al. 2008; Bell et al. 2013), we observe a  $T_{eff}$ -dependent age trend in all three subgroups of Sco-Cen, for all sets of evolutionary tracks.

(iv) We adopt median ages of  $10\pm3$  Myr,  $16\pm2$  Myr and  $15\pm3$  Myr for US, UCL and LCC, respectively, when considering the revised nuclear ages as well as the pre-MS contraction ages from the F- and G-type stars.

(v) We obtain estimates for the intrinsic age spread in each subgroup through a grid of Monte Carlo simulations which take into account binarity, spots, and observational uncertainties. Assuming the median ages obtained above, and modeling the age distribution as a gaussian, we find that 68% of the star formation in US, UCL, and LCC occured over timescales of  $\pm 7$  Myr,  $\pm 7$  Myr, and  $\pm 6$  Myr, respectively. Thus when adopting an age of  $\sim 10$  Myr for Upper Sco, we detect an intrinsic age spread of  $\pm 7$  Myr (1 $\sigma$ ).

(vi) Using members from our X-ray sample as well as F- and G-type members of Sco-Cen, we create an age map of the Sco-Cen complex. We find that the star-formation histories of the UCL and LCC, the older subgroups, are indicitative of substructure and are not consistent with a simple triggered star-formation scenario. The groups are not each monolithic episodes of star formation, but likely an ensemble of small subgroups, exhibiting significant substructure.

#### ACKNOWLEDGEMENTS

We would like to thank the anonymous referee, whose comments and suggestions improved the quality of this work. This work has been supported by funds from the School of Arts and Sciences at the University of Rochester and NSF grants AST-1008908 and AST-1313029. EEM acknowledges support from the NASA Nexus for Exoplanet System Science program (NExSS). We thank Fred Walter for the use of his SMARTS RC-spectrograph pipeline for reducing the data obtained at the SMARTS 1.5m telescope at Cerro Tololo, Chile, as well as Jose Velasquez and Manuel Hernandez for their help and advice at the telescope. This research was made possible through the use of the AAVSO Photometric All-Sky Survey (APASS), funded by the Robert Martin Ayers Sciences Fund. This publication makes use of data products from the Two Micron All Sky Survey, which is a joint project of the University of Massachusetts and the Infrared Processing and Analysis Center/California Institute of Technology, funded by the National Aeronautics and Space Administration and the National Science Foundation. This publication makes use of data products from the Wide-field Infrared Survey Explorer, which is a joint project of the University of California, Los Angeles, and the Jet Propulsion Laboratory/California Institute of Technology, funded by the National Aeronautics and Space Administration.

#### REFERENCES

- Allard F., Homeier D., Freytag B., 2012, Royal Society of London Philosophical Transactions Series A, 370, 2765
- Alonso R., Deeg H. J., Hoyer S., Lodieu N., Palle E., Sanchis-Ojeda R., 2015, A&A, 584, L8
- Ardila D., Martín E., Basri G., 2000, AJ, 120, 479
- Asplund M., Grevesse N., Sauval A. J., 2005, in Barnes III T. G., Bash F. N., eds, Astronomical Society of the Pacific Conference Series Vol. 336, Cosmic Abundances as Records of Stellar Evolution and Nucleosynthesis. p. 25
- Balona L. A., 1994, MNRAS, 268, 119
- Baraffe I., Homeier D., Allard F., Chabrier G., 2015, A&A, 577, A42
- Barrado y Navascués D., Martín E. L., 2003, AJ, 126, 2997
- Bell C. P. M., Naylor T., Mayne N. J., Jeffries R. D., Littlefair S. P., 2012, MNRAS, 424, 3178
- Bell C. P. M., Naylor T., Mayne N. J., Jeffries R. D., Littlefair S. P., 2013, MNRAS, 434, 806
- Bell C. P. M., Rees J. M., Naylor T., Mayne N. J., Jeffries R. D., Mamajek E. E., Rowe J., 2014, MNRAS, 445, 3496

- Binks A. S., Jeffries R. D., 2014, MNRAS, 438, L11
- Blaauw A., 1946, Publications of the Kapteyn Astronomical Laboratory Groningen, 52, 1
- Blaauw A., 1964, ARA&A, 2, 213
- Blaauw A., 1978, Internal Motions and Age of the Sub-Association Upper Scorpio. p. 101
- Bodenheimer P., 1965, ApJ, 142, 451
- Bouvier J., Appenzeller I., 1992, A&AS, 92, 481
- Bouvier J., Alencar S. H. P., Harries T. J., Johns-Krull C. M., Romanova M. M., 2007, Protostars and Planets V, pp 479–494
- Boyajian T. S., et al., 2012, ApJ, 757, 112
- Brandner W., Alcala J. M., Kunkel M., Moneti A., Zinnecker H., 1996, A&A, 307, 121
- Briceño C., Preibisch T., Sherry W. H., Mamajek E. A., Mathieu R. D., Walter F. M., Zinnecker H., 2007, Protostars and Planets V, pp 345– 360
- Brown A. G. A., Dekker G., de Zeeuw P. T., 1997, MNRAS, 285, 479
- Caffau E., Ludwig H.-G., Steffen M., Ayres T. R., Bonifacio P., Cayrel R., Freytag B., Plez B., 2008, A&A, 488, 1031
- Caffau E., Maiorca E., Bonifacio P., Faraggiana R., Steffen M., Ludwig H.-G., Kamp I., Busso M., 2009, A&A, 498, 877
- Caffau E., Ludwig H.-G., Steffen M., Freytag B., Bonifacio P., 2011, Sol. Phys., 268, 255
- Carballo R., Wesselius P. R., Whittet D. C. B., 1992, A&A, 262, 106
- Cargile P. A., James D. J., Jeffries R. D., 2010, ApJ, 725, L111
- Carpenter J. M., Mamajek E. E., Hillenbrand L. A., Meyer M. R., 2006, ApJ, 651, L49
- Carpenter J. M., Mamajek E. E., Hillenbrand L. A., Meyer M. R., 2009, ApJ, 705, 1646
- Chen C. H., Mamajek E. E., Bitner M. A., Pecaut M., Su K. Y. L., Weinberger A. J., 2011, ApJ, 738, 122
- Chen Y., Girardi L., Bressan A., Marigo P., Barbieri M., Kong X., 2014, MNRAS, 444, 2525
- Churchwell E., et al., 2009, PASP, 121, 213
- Cutri R. M., et al. 2012, VizieR Online Data Catalog, 2311, 0
- Cutri R. M., et al. 2014, VizieR Online Data Catalog, 2328, 0
- Cutri R. M., et al., 2012, VizieR Online Data Catalog, 2281, 0
- Dahm S. E., Slesnick C. L., White R. J., 2012, ApJ, 745, 56
- David T. J., Hillenbrand L. A., Cody A. M., Carpenter J. M., Howard A. W., 2015, preprint, (arXiv:1510.08087)
- Dobbie P. D., Lodieu N., Sharp R. G., 2010, MNRAS, 409, 1002
- Dolidze M. V., Arakelyan M. A., 1959, Azh, 36, 444
- Dotter A., Chaboyer B., Jevremović D., Kostov V., Baron E., Ferguson J. W., 2008, ApJS, 178, 89
- Downes R. A., Keyes C. D., 1988, AJ, 96, 777
- Duchêne G., Kraus A., 2013, preprint, (arXiv:1303.3028)
- Ducourant C., Teixeira R., Galli P. A. B., Le Campion J. F., Krone-Martins A., Zuckerman B., Chauvin G., Song I., 2014, A&A, 563, A121
- ESA 1997, VizieR Online Data Catalog, 1239, 0
- Ekström S., et al., 2012, A&A, 537, A146
- Espaillat C., et al., 2012, ApJ, 747, 103
- Fedele D., van den Ancker M. E., Henning T., Jayawardhana R., Oliveira J. M., 2010, A&A, 510, A72
- Feiden G. A., 2016, in Kastner J. H., Stelzer B., Metchev S. A., eds, IAU Symposium Vol. 314, IAU Symposium. pp 79–84 (arXiv:1507.06460), doi:10.1017/S1743921315006067
- Feigelson E. D., Lawson W. A., 1997, AJ, 113, 2130
- Feigelson E. D., Montmerle T., 1999, ARA&A, 37, 363
- Filin A. Y., Satyvoldiev V., 1975, Peremennye Zvezdy, 20, 161
- Finkenzeller U., Basri G., 1987, ApJ, 318, 823
- Fiorucci M., Munari U., 2003, A&A, 401, 781
- Garrison R. F., 1967, ApJ, 147, 1003
- Gray R. O., Garrison R. F., 1989, ApJS, 69, 301
- Gray R. O., Napier M. G., Winkler L. I., 2001, AJ, 121, 2148
- Gray R. O., Corbally C. J., Garrison R. F., McFadden M. T., Robinson P. E., 2003, AJ, 126, 2048
- Gray R. O., Corbally C. J., Garrison R. F., McFadden M. T., Bubar E. J., McGahee C. E., O'Donoghue A. A., Knox E. R., 2006, AJ, 132, 161

- Gregorio-Hetem J., Lepine J. R. D., Quast G. R., Torres C. A. O., de La Reza R., 1992, AJ, 103, 549
- Grevesse N., Sauval A. J., 1998, Space Sci. Rev., 85, 161
- Hauck B., Mermilliod M., 1998, A&AS, 129, 431
- Henden A. A., Levine S. E., Terrell D., Smith T. C., Welch D., 2012, Journal of the American Association of Variable Star Observers (JAAVSO), 40, 430
- Henize K. G., 1954, ApJ, 119, 459
- Henize K. G., 1976, ApJS, 30, 491
- Henry T. J., Walkowicz L. M., Barto T. C., Golimowski D. A., 2002, AJ, 123, 2002
- Herbig G. H., Bell K. R., 1988, Third Catalog of Emission-Line Stars of the Orion Population : 3 : 1988
- Hernández J., et al., 2007, ApJ, 671, 1784
- Hillenbrand L. A., 1997, AJ, 113, 1733
- Hillenbrand L. A., White R. J., 2004, ApJ, 604, 741
- Hillenbrand L. A., Bauermeister A., White R. J., 2008, in van Belle G., ed., Astronomical Society of the Pacific Conference Series Vol. 384, 14th Cambridge Workshop on Cool Stars, Stellar Systems, and the Sun. p. 200 (arXiv:astro-ph/0703642)
- Hiltner W. A., Garrison R. F., Schild R. E., 1969, ApJ, 157, 313
- Høg E., et al., 2000, A&A, 355, L27
- Hoogerwerf R., de Bruijne J. H. J., de Zeeuw P. T., 2000, ApJ, 544, L133
- Hou A., Parker L. C., Harris W. E., Wilman D. J., 2009, ApJ, 702, 1199
- Houk N., 1978, Michigan catalogue of two-dimensional spectral types for the HD stars
- Houk N., 1982, Michigan Catalogue of Two-dimensional Spectral Types for the HD stars. Volume\_3.
- Houk N., Smith-Moore M., 1988, Michigan Catalogue of Two-dimensional Spectral Types for the HD Stars. Volume 4
- Hughes J., Hartigan P., Krautter J., Kelemen J., 1994, AJ, 108, 1071
- Ichikawa T., Nishida M., 1989, AJ, 97, 1074
- Ishihara D., et al., 2010, A&A, 514, A1
- Jeffries R. D., Littlefair S. P., Naylor T., Mayne N. J., 2011, MNRAS, 418, 1948
- Johnson H. L., Morgan W. W., 1953, ApJ, 117, 313
- Jones B. F., Fischer D. A., Stauffer J. R., 1996, AJ, 112, 1562
- Kalas P. G., et al., 2015, ApJ, 814, 32
- Keenan P. C., McNeil R. C., 1989, ApJS, 71, 245
- Kirkpatrick J. D., Henry T. J., McCarthy Jr. D. W., 1991, ApJS, 77, 417
- Kiss L. L., et al., 2011, MNRAS, 411, 117
- Köhler R., Kunkel M., Leinert C., Zinnecker H., 2000, A&A, 356, 541
- Kordopatis G., et al., 2013, AJ, 146, 134
- Kraus A. L., Ireland M. J., 2012, ApJ, 745, 5
- Kraus A. L., Tucker R. A., Thompson M. I., Craine E. R., Hillenbrand L. A., 2011, ApJ, 728, 48
- Kraus A. L., Cody A. M., Covey K. R., Rizzuto A. C., Mann A. W., Ireland M. J., 2015, ApJ, 807, 3
- Krautter J., Wichmann R., Schmitt J. H. M. M., Alcala J. M., Neuhauser R., Terranegra L., 1997, A&AS, 123, 329
- Kroupa P., 2001, MNRAS, 322, 231
- Lawson W. A., Lyo A.-R., Bessell M. S., 2009, MNRAS, 400, L29
- Lodieu N., et al., 2015, preprint, (arXiv:1511.03083)
- Luhman K. L., Mamajek E. E., 2012, ApJ, 758, 31
- Mamajek E. E., 2005, ApJ, 634, 1385
- Mamajek E. E., 2009, in T. Usuda, M. Tamura, & M. Ishii ed., American Institute of Physics Conference Series Vol. 1158, American Institute of Physics Conference Series. pp 3–10 (arXiv:0906.5011), doi:10.1063/1.3215910
- Mamajek E. E., 2013, in American Astronomical Society Meeting Abstracts #221. p. 331.06
- Mamajek E. E., Bell C. P. M., 2014, MNRAS, 445, 2169
- Mamajek E. E., Meyer M. R., Liebert J., 2002, AJ, 124, 1670
- Mamajek E. E., Meyer M. R., Liebert J., 2006, AJ, 131, 2360
- Mamajek E. E., Quillen A. C., Pecaut M. J., Moolekamp F., Scott E. L., Kenworthy M. A., Collier Cameron A., Parley N. R., 2012, AJ, 143, 72
- Mamajek E. E., Pecaut M. J., Nguyen D. C., Bubar E. J., 2013, in Protostars and Planets VI Posters.

- Martin E. L., Montmerle T., Gregorio-Hetem J., Casanova S., 1998, MN-RAS, 300, 733
- Mason B. D., Wycoff G. L., Hartkopf W. I., Douglass G. G., Worley C. E., 2001, AJ, 122, 3466
- Mathieu R. D., Walter F. M., Myers P. C., 1989, AJ, 98, 987
- Mentuch E., Brandeker A., van Kerkwijk M. H., Jayawardhana R., Hauschildt P. H., 2008, ApJ, 689, 1127
- Mermilliod J.-C., Mermilliod M., 1994, Catalogue of Mean UBV Data on Stars
- Morgan W. W., Keenan P. C., 1973, ARA&A, 11, 29
- Morgan W. W., Abt H. A., Tapscott J. W., 1978, Revised MK Spectral Atlas for stars earlier than the sun
- Müller A., Carmona A., van den Ancker M. E., van Boekel R., Henning T., Launhardt R., 2011, A&A, 535, L3
- Murphy S. J., Lawson W. A., Bento J., 2015, MNRAS, 453, 2220
- Napiwotzki R., Schoenberner D., Wenske V., 1993, A&A, 268, 653

Naylor T., 2009, MNRAS, 399, 432

- Nguyen D. C., Jayawardhana R., van Kerkwijk M. H., Brandeker A., Scholz A., Damjanov I., 2009, ApJ, 695, 1648
- Nguyen D. C., Mamajek E., Pecaut M., 2013, in Protostars and Planets VI Posters.
- Nieva M.-F., 2013, A&A, 550, A26
- Park S., Finley J. P., 1996, AJ, 112, 693
- Pecaut M. J., Mamajek E. E., 2013, ApJS, 208, 9
- Pecaut M. J., Mamajek E. E., Bubar E. J., 2012, ApJ, 746, 154
- Petrov P. P., Satyvoldiev V., 1975, Peremennye Zvezdy Prilozhenie, 2, 221
- Pollack J. B., Hubickyj O., Bodenheimer P., Lissauer J. J., Podolak M., Greenzweig Y., 1996, Icarus, 124, 62
- Preibisch T., Mamajek E., 2008, The Nearest OB Association: Scorpius-Centaurus (Sco OB2). pp 235-+
- Preibisch T., Zinnecker H., 1999, AJ, 117, 2381
- Preibisch T., Guenther E., Zinnecker H., Sterzik M., Frink S., Roeser S., 1998, A&A, 333, 619
- Preibisch T., Guenther E., Zinnecker H., 2001, AJ, 121, 1040
- Preibisch T., Brown A. G. A., Bridges T., Guenther E., Zinnecker H., 2002, AJ, 124, 404
- Raghavan D., et al., 2010, ApJS, 190, 1
- Randich S., Schmitt J. H. M. M., Prosser C. F., Stauffer J. R., 1995, A&A, 300, 134
- Randich S., Aharpour N., Pallavicini R., Prosser C. F., Stauffer J. R., 1997, A&A, 323, 86
- Reid N., 2003, MNRAS, 342, 837
- Riaud P., Mawet D., Absil O., Boccaletti A., Baudoz P., Herwats E., Surdej J., 2006, A&A, 458, 317
- Riaz B., Gizis J. E., Harvin J., 2006, AJ, 132, 866
- Ribas Á., Bouy H., Merín B., 2015, A&A, 576, A52
- Rizzuto A. C., Ireland M. J., Robertson J. G., 2011, MNRAS, 416, 3108
- Rizzuto A. C., Ireland M. J., Kraus A. L., 2015, MNRAS, 448, 2737
- Rodriguez D. R., Bessell M. S., Zuckerman B., Kastner J. H., 2011, ApJ, 727, 62
- Röser S., Schilbach E., Schwan H., Kharchenko N. V., Piskunov A. E., Scholz R.-D., 2008, A&A, 488, 401
- Satyvaldiev V., 1972, Astronomicheskij Tsirkulyar, 728, 5
- Satyvoldiev V., 1982, Peremennye Zvezdy Prilozhenie, 4, 109
- Schlieder J. E., Lépine S., Rice E., Simon M., Fielding D., Tomasino R., 2012, AJ, 143, 114
- Schwartz R. D., 1977, ApJS, 35, 161
- Sciortino S., Damiani F., Favata F., Micela G., 1998, A&A, 332, 825
- Shobbrook R. R., 1983, MNRAS, 205, 1215
- Skrutskie M. F., et al., 2006, AJ, 131, 1163
- Slesnick C. L., Carpenter J. M., Hillenbrand L. A., 2006, AJ, 131, 3016
- Slesnick C. L., Hillenbrand L. A., Carpenter J. M., 2008, ApJ, 688, 377
- Soderblom D. R., Jones B. F., Balachandran S., Stauffer J. R., Duncan D. K., Fedele S. B., Hudon J. D., 1993, AJ, 106, 1059
- Soderblom D. R., Hillenbrand L. A., Jeffries R. D., Mamajek E. E., Naylor T., 2014, Protostars and Planets VI, pp 219–241
- Somers G., Pinsonneault M. H., 2015, ApJ, 807, 174
- Song I., Zuckerman B., Bessell M. S., 2003, ApJ, 599, 342

- Song I., Zuckerman B., Bessell M. S., 2012, AJ, 144, 8
- Spinrad H., 1962, ApJ, 135, 715
- Stauffer J. R., Schultz G., Kirkpatrick J. D., 1998, ApJ, 499, L199
- Stauffer J. R., Jones B. F., Backman D., Hartmann L. W., Barrado y Navascués D., Pinsonneault M. H., Terndrup D. M., Muench A. A., 2003, AJ, 126, 833
- Stephenson C. B., 1986, ApJ, 300, 779
- Stocke J. T., Morris S. L., Gioia I. M., Maccacaro T., Schild R., Wolter A., Fleming T. A., Henry J. P., 1991, ApJS, 76, 813
- Strom S. E., 1994, in Caillault J.-P., ed., Astronomical Society of the Pacific Conference Series Vol. 64, Cool Stars, Stellar Systems, and the Sun. p. 211
- Struve O., Rudkjøbing M., 1949, ApJ, 109, 92
- Thackeray A. D., 1966, Mem. RAS, 70, 33
- The P.-S., 1962, Contributions from the Bosscha Observervatory, 15, 0
- The P.-S., 1964, PASP, 76, 293
- Tognelli E., Prada Moroni P. G., Degl'Innocenti S., 2011, A&A, 533, A109 Torres G., Ribas I., 2002, ApJ, 567, 1140
- Torres C. A. O., Quast G. R., da Silva L., de La Reza R., Melo C. H. F., Sterzik M., 2006, A&A, 460, 695
- Torres C. A. O., Quast G. R., Melo C. H. F., Sterzik M. F., 2008, Young Nearby Loose Associations. p. 757
- Vazquez R. A., Feinstein A., 1991, A&AS, 87, 383
- Viana Almeida P., Santos N. C., Melo C., Ammler-von Eiff M., Torres C. A. O., Quast G. R., Gameiro J. F., Sterzik M., 2009, A&A, 501, 965
- Voges W., et al., 1999, A&A, 349, 389
- Voges W., et al., 2000, IAU Circ., 7432, 3
- Walter F. M., Vrba F. J., Mathieu R. D., Brown A., Myers P. C., 1994, AJ, 107, 692
- Webb R. A., 2000, PhD thesis, University of California, Los Angeles
- Weinberger A. J., Anglada-Escudé G., Boss A. P., 2012, preprint,
- (arXiv:1211.2233) White R. J., Gabor J. M., Hillenbrand L. A., 2007, AJ, 133, 2524
- Wichmann R., Sterzik M., Krautter J., Metanomski A., Voges W., 1997, A&A, 326, 211
- Wichmann R., Covino E., Alcalá J. M., Krautter J., Allain S., Hauschildt P. H., 1999, MNRAS, 307, 909
- Wright E. L., et al., 2010, AJ, 140, 1868
- Wright N. J., Parker R. J., Goodwin S. P., Drake J. J., 2014, MNRAS, 438, 639
- Wright N. J., Bouy H., Drew J. E., Sarro L. M., Bertin E., Cuillandre J.-C., Barrado D., 2016, preprint, (arXiv:1605.03583)
- Yee J. C., Jensen E. L. N., 2010, ApJ, 711, 303
- Zacharias N., Finch C. T., Girard T. M., Henden A., Bartlett J. L., Monet D. G., Zacharias M. I., 2013, AJ, 145, 44
- Zhang K., Pontoppidan K. M., Salyk C., Blake G. A., 2013, ApJ, 766, 82
- Zuckerman B., Song I., 2004, ARA&A, 42, 685
- Zuckerman B., Webb R. A., Schwartz M., Becklin E. E., 2001, ApJ, 549, L233
- de Bruijne J. H. J., 1999a, MNRAS, 306, 381
- de Bruijne J. H. J., 1999b, MNRAS, 310, 585
- de Geus E. J., de Zeeuw P. T., Lub J., 1989, A&A, 216, 44
- de Zeeuw P. T., Hoogerwerf R., de Bruijne J. H. J., Brown A. G. A., Blaauw A., 1999, AJ, 117, 354
- de la Reza R., Jilinski E., Ortega V. G., 2006, AJ, 131, 2609 van Leeuwen F., 2007, A&A, 474, 653
- This paper has been typeset from a TEX/LATEX file prepared by the author.