

LOGIC OF LOCAL INFERENCE FOR CONTEXTUALITY IN QUANTUM PHYSICS AND BEYOND

KOHEI KISHIDA

University of Oxford

ABSTRACT. Contextuality in quantum physics provides a key resource for quantum information and computation. The topological approach in [3, 2] characterizes contextuality as “global inconsistency” coupled with “local consistency”, revealing it to be a phenomenon also found in many other fields. This has yielded a logical method of detecting and proving the “global inconsistency” part of contextuality. Our goal is to capture the other, “local consistency” part, which requires a novel approach to logic that is sensitive to the topology of contexts. To achieve this, we formulate a logic of local inference by using context-sensitive theories and models in regular categories. This provides a uniform framework for local consistency, and lays a foundation for high-level methods of detecting, proving, and moreover using contextuality as computational resource.

1. INTRODUCTION

Quantum physics provides quantum computing with immense advantage over classical computing. Among its non-classical properties, non-locality is known to be a basis for quantum communication (see [19]). It is in fact a special case of a more general property called *contextuality*, which recent studies [22, 10] suggest is an essential source of the computational power of quantum computers. This motivates the search for structural, higher-level expressions of contextuality that are independent of the formalism of quantum mechanics.

One conception of contextuality that originated in [11] exploits the structure of *presheaf*: As was shown in [11], a certain type of contextuality of a quantum system amounts to the absence of global sections from presheaves modelling the behaviors of the system. The recent, “sheaf-theoretic” approach [3] expands this insight by viewing contextuality in more general terms, as a matter of *topology* in data of measurements and outcomes: A wider range of contextuality is then characterized as the “global inconsistency” of the “locally consistent”. This has on the one hand shown that contextual phenomena can be found in various other fields such as relational database theory (see [1]), and on the other hand made it possible to apply various tools—sheaf theory, cohomology, linear algebra, for instance—to contextuality. One idea that has emerged is to formulate contextuality argument in logical terms [2]: One describes a presheaf model using logical formulas, and proves its contextuality by deriving contradiction from the formulas.

This method, however, only shows the global inconsistency of a given set of formulas; we know them to be locally consistent only because they describe a locally consistent model that is given. Nonetheless, when designing ways of exploiting contextuality, we may well first obtain a set of

E-mail address: Kohei.Kishida@cs.ox.ac.uk.

This work has been supported by the grant FA9550-12-1-0136 of the U.S. AFOSR.

formulas or a specification, and then check if there is a model satisfying it. This requires a logic in which consistency means local consistency. The chief goal of this paper is to deliver such a new logic of local inference. The two logics—one for global inconsistency and the other for local consistency—together lay a foundation for high-level logical methods of not only showing but also using contextuality as resource.

Section 2 reviews the sheaf approach to contextuality, which takes presheaves valued in **Sets**. Then Section 3 defines what we call “inchworm logic”, a novel logic of local inference for contextual models. We formulate this on the basis of regular logic, since its vocabulary captures the essence of local inference. Semantics is provided for this logic in Section 4, where we generalize **Sets**-valued presheaf models to ones valued in any regular category **S**. This encompasses cases that prove useful and powerful in applications: E.g., presheaves of abelian groups, R -modules, etc. serve the purpose of cohomology; indeed, Čech cohomology is used to detect the contextuality of **Sets**-valued presheaves [2]. This paper gives a uniform way of using **S**-valued presheaves directly as models of contextual logic.

2. CONTEXTUAL MODELS

We first review the idea behind the formalism of [2], stressing that it applies to more settings than just quantum ones. The idea captures contextuality as a matter of topological nature, which we illustrate with a simplicial formulation equivalent to the presheaf formulation of [2]. We also present a modification of the latter that can readily be generalized in Section 4.

2.1. Topological Models for Contextuality. The formalism of [3] concerns variables and values in general. Their bare-bones structure consists of a set X of variables and, for each $x \in X$, a set A_x of possible values of x . So we have an X -indexed family of sets A_x . This can model various settings in which we make queries against a system and it answers, as observed in [1, 2]; e.g.,

- We measure properties $x \in X$ of a physical system and it gives back outcomes $a \in A_x$.
- A relational database has attributes $x \in X$, and $a \in A_x$ are possible data values for x .
- $x \in X$ are sentences of propositional logic and a set of models assign to them boolean values $a \in A_x = \mathbf{2}$. Or $x \in X$ may simply be boolean variables.

We often make a query regarding several variables in combination; a set $U \subseteq X$ of variables the query concerns forms a *context* in which the system gives back a result. Contexts play essential rôles in the following two kinds of constraints, (a) on answers and (b) on queries.

(a) When we make a query in a context U , the system returns (one or a set of) tuples $s \in \prod_{x \in U} A_x$ of values. It then has the subset $A_U \subseteq \prod_{x \in U} A_x$ of “admissible” tuples that can be part of query results, and it is often the information on A_U that we want. E.g.,

- From a relational database we retrieve data with an attribute list U , and the database returns the relation A_U on sets A_x ($x \in U$) as a table.
- Given a set of models and a set U of sentences, A_U is the set of combinations of values that U can take; e.g., a pair $\varphi, \neg\varphi \in U$ only take values $(1, 0)$ or $(0, 1)$.
- We may measure a physical system in various states and find that some set U of quantities always satisfies a certain equation that characterizes A_U .

(A tuple $s \in \prod_{x \in U} A_x$ is a dependent function, so it is formally a set of the form $\{(x, s(x)) \mid x \in U \text{ and } s(x) \in A_x\}$; but we may refer to it as “ $(s(x), s(y), \dots)$ over (x, y, \dots) ”.)

(b) We have the family $C \subseteq \mathcal{P}X$ of contexts in which queries can be made and answered. We may not be able to make a query in a context $V \subseteq X$ (i.e. $V \notin C$) for reasons such as:

- V may have too many variables to deal with feasibly.
- A database schema may have no table encompassing all the attributes in V .
- Quantum mechanics may deem it impossible to measure all the properties in V at once.

In these examples, if queries can be made in a context U , they can be in any $V \subseteq U$; we also assume that queries can be made in $\{x\}$ for any $x \in X$, but only in finite U . So C is an (abstract) *simplicial complex* on X , i.e. a \subseteq -downward closed subfamily of $\mathcal{P}_{\text{fin}}X$ with $\bigcup_{U \in C} U = X$. Also, if a tuple s of values is admissible, so is any $t \subseteq s$. Hence, whenever $V \subseteq U \in C$, the projection of tuples

$$-|_V : \prod_{x \in U} A_x \rightarrow \prod_{x \in V} A_x :: s \mapsto s|_V$$

restricts to $A_{V \subseteq U} : A_U \rightarrow A_V$. Thus $A : C^{\text{op}} \rightarrow \mathbf{Sets}$ forms a presheaf on the poset C .

In fact, A is a *separated* presheaf. Generally, for any subfamily C of $\mathcal{P}X$ closed under binary intersection, whenever $\bigcup_i U_i = U \in C$ for $U_i \in C$, a presheaf P on C has the map

$$\langle P_{U_i \subseteq U} \rangle_i : P_U \rightarrow \prod_i P_{U_i} :: s \mapsto (s|_{U_i})_i$$

land in the set of *matching families* for $(U_i)_i$,

$$\text{Match}_{(U_i)_i, P} = \{ (t_i)_i \in \prod_i P_{U_i} \mid t_j|_{U_j \cap U_k} = t_k|_{U_j \cap U_k} \text{ for every pair } j, k \}.$$

Then P is called *separated* if each of these $\langle P_{U_i \subseteq U} \rangle_i$ is injective, and a *sheaf* if each of those $\langle P_{U_i \subseteq U} \rangle_i : P_U \rightarrow \text{Match}_{(U_i)_i, P}$ is bijective (see [17]). Yet, on a simplicial complex C , separated presheaves and sheaves have simpler descriptions:

Fact 1. A presheaf P on a simplicial complex C is a sheaf iff $P_U = \prod_{x \in U} P_x$ for all $U \in C$. And P is separated iff it is a subpresheaf of a sheaf, i.e. iff $P_U \subseteq \prod_{x \in U} P_x$ for all $U \in C$.

This shows that our A above is a separated presheaf, but not generally a sheaf. So let us write $\mathbf{sPsh}(C)$ for the full subcategory of $\mathbf{Sets}^{C^{\text{op}}}$ of separated presheaves. Note that every sheaf F has $F_\emptyset = 1$, a singleton. A separated presheaf P has $P_\emptyset = 1$, too, unless it is the empty presheaf $U \mapsto \emptyset$, i.e. the model is inconsistent in every context (hence modelling, e.g., a physical system that never produces outcomes in any context of measurements).

2.2. Presheaves and Bundles. The constraints (a) and (b) above are, indeed, matters of topology; this idea will be useful in Subsection 2.3. Given a separated presheaf A as in Subsection 2.1, its underlying family of X -indexed sets $(A_x)_{x \in X}$ is equivalent to a set over X , viz. $\pi : \sum_{x \in X} A_x \rightarrow X :: (x, a) \mapsto x$, by $\mathbf{Sets}^X \simeq \mathbf{Sets}/X$. The base X comes with a simplicial complex C , but so does $\sum_{x \in X} A_x$, taking tuples $s \in A_U$ as simplices, i.e. $\mathcal{A} = \bigcup_{U \in C} A_U$. And π is a *simplicial map*, or a *bundle* of simplicial complexes, since $s \in A_U \subseteq \mathcal{A}$ implies $\pi[s] = U \in C$. On the other hand, any given bundle $\pi : \mathcal{A} \rightarrow C$ has a family of $A_U = \{s \in \mathcal{A} \mid \pi[s] = U\}$ and $A_{V \subseteq U} : s \mapsto s|_V$.

A simplicial map $\pi : \mathcal{A} \rightarrow C$ is called *non-degenerate* if $\pi|_s$ is injective for every $s \in \mathcal{A}$. Our π above is non-degenerate, because every $s \in A_U$ is a *local section* of the bundle π , meaning $s : U \rightarrow \sum_{x \in X} A_x$ such that $\pi \circ s = 1_U$. Let us write **Simp** and **ndSimp** for the categories of simplicial maps and of non-degenerate ones, respectively. It is easy to check that for every simplicial complex C , the slice category **ndSimp**/ C is a full subcategory of **Simp**/ C ; i.e., it is the category of non-degenerate bundles




FIGURE 1. Bundles for (c) the Hardy model and (d) the PR-box

and simplicial maps over C . Then, extending $\mathbf{Sets}^X \simeq \mathbf{Sets}/X$, the correspondence described above gives

Fact 2. $\mathbf{sPsh}(C) \simeq \mathbf{ndSimp}/C$ for any simplicial complex C .

So here is a topological reading of (a) and (b). Each $U \in C$ is a local, small enough region of the space X of variables. The topology on the space \mathcal{A} of values then distinguishes those tuples $s \in \prod_{x \in U} A_x$ in A_U from the others and deems the former to be continuous sections. We refer to objects of $\mathbf{sPsh}(C)$ and \mathbf{ndSimp}/C interchangeably as *topological models*.

2.3. Contextuality in Physics, Databases, and More. Since queries can only be made locally, i.e. in contexts $U \in C$, answers to queries can only be observed locally in those contexts. One might think this is just a matter of convenience or efficiency, since we can simply make multiple queries in multiple contexts U_i to cover $X = \bigcup_i U_i$ and to recover the global information. In certain topologies C , however, this may be prevented due to contextuality.

Given a non-degenerate bundle $\pi : \mathcal{A} \rightarrow C$ over a simplicial complex C on X , consider a *global section* of it, i.e. $g \in \prod_{x \in X} A_x$ such that $g|_U \in A_U$ for all $U \in C$; it corresponds to a matching family $g \in \text{Match}_{C,p}$, since $U \mapsto \prod_{x \in U} A_x$ ($U \subseteq X$) is a sheaf. It is an assignment of values to all the variables that satisfies every constraint on combinations of values. E.g., in classical logic, the models are exactly the global sections; so the consistency of a sentence x means that $(x, 1)$ is part of a global section. Then, in the physical setting, it may seem natural to similarly think of global sections g as states of the system, assigning values to all the quantities—so, although we can only make a query locally in a context $U \in C$, the system in a state g actually has a value $g(x)$ assigned to every quantity x , and the answer we receive in the context U is simply $g|_U$. This assumption, that any section we observe is part of a context-independent global section, holds not just in classical logic but also in classical physics—but breaks down in quantum physics, precisely when contextuality arises.

Figure 1 shows “Bell-type” scenarios in which Alice and Bob measure properties of a system, perhaps a quantum one. The base C expresses constraints of type (b) above: Alice can make at most one of two measurements a_1 and a_2 at a time, so she chooses one; similarly Bob chooses from b_1 and b_2 —so there are four possible combinations of measurements, indicated by the four edges of C . Alice and Bob repeat measurements in different contexts, and learn that each $x \in X = \{a_1, a_2, b_1, b_2\}$ has two possible outcomes 0 and 1, but that some combinations of outcomes are never obtained. \mathcal{A} expresses these constraints, of type (a), with edges indicating possible combinations. E.g., \mathcal{A} of (c) deems every joint outcome of (a_1, b_1) possible, with $A_{\{a_1, b_1\}} = 2 \times 2$; but $(0, 0)$ is not a possible joint outcome of (a_2, b_2) .

The models in Figure 1 all violate the classical assumption above, and are examples of

Definition 3. A topological model is said to be *logically contextual* if not all of its local sections extend to global ones, and *strongly contextual* if it has no global section at all (see [2]).

(c) of Figure 1 represents an example of logical contextuality due to [9] that is realizable in quantum physics. It has several global sections, e.g. the one marked in green; call it g . So, when Alice and Bob measure (a_1, b_1) and observe $(0, 0)$, the classical explanation is possible that the system was in the state g and had outcomes $g(x)$ assigned to all the measurements $x \in X$, and that Alice and Bob have simply retrieved that information on U . On the other hand, the local section in red, $(1, 1)$ over (a_1, b_1) , does not extend to any global section. This means that the classical explanation is simply impossible for this joint outcome. Furthermore, the classical explanation is never possible in the strongly contextual (d). This model, called the *PR box* [21], is not quantum-realizable (though it plays an important rôle in the quantum information literature), but quantum physics exhibits many other instances of strong contextuality.

The upshot is that contextuality consists in *global inconsistency coupled with local consistency*: A section $s \in A_U$ is consistent locally, in the sense of satisfying the constraint on query results in the context U , but it may be inconsistent globally, in the sense of contradicting all the other constraints and thereby failing to extend to a global section.

The general definition of contextuality in terms of global sections can also be applied to relational databases: Contextuality then corresponds exactly to the absence of a universal relation [1]. In fact, the natural join

$$\bowtie_{U \in C} A_U = \{g \in \prod_{x \in X} A_x \mid g|_U \in A_U \text{ for all } U \in C\}$$

of relations A_U (which is the largest of universal relations if there are any) is, simply by definition, the set of global sections. Hence, e.g. in Figure 1, the sections in red are lost from $\bowtie_{U \in C} A_U$.

2.4. No-Signalling Principle. Local consistency means, partly, that a local section may exist without extending to global. Yet it involves more—viz. a constraint that is called the *no-signalling* principle in the physical setting [7]. For a topological model A , it amounts to the condition that every $A_{U \subseteq V} : A_V \rightarrow A_U :: s \mapsto s|_U$ (i.e. the projection of admitted tuples, or the restriction of sections) is a surjection.

An example violating no-signalling is (e) of Figure 1: $A_{\{b_1\} \subseteq \{a_2, b_1\}} : A_{\{a_2, b_1\}} \rightarrow A_{\{b_1\}}$ is not surjective. Suppose Alice and Bob make measurements, Bob chooses to measure b_1 , and he observes 1, which is not in the image of $A_{\{b_1\} \subseteq \{a_2, b_1\}}$. This means that Bob has received the signal from Alice (no matter how far away she may be!) that she has chosen a_1 and not a_2 .

To see why no-signalling should be part of local consistency, regard \mathcal{A} in (e) as representing a relational database. It has tables $A_{\{a_1, b_1\}}$ and $A_{\{a_2, b_1\}}$; but, when queried about the attribute b_1 , they yield different results of projection, differing in whether 1 is in or not. Thus, no-signalling means the consistency of projections (see [1]). Indeed, as we will see in Subsection 4.3, no-signalling means a sort of coherence of A as a semantic model of logic.

Definition 4. We say that a separated presheaf $A : C^{\text{op}} \rightarrow \mathbf{Sets}$ is no-signalling if it satisfies (1), and that a non-degenerate bundle $\pi : \mathcal{A} \rightarrow C$ is no-signalling if it satisfies (2):

- (1) Every $A_{U \subseteq V} : A_V \rightarrow A_U$ is a surjection.
- (2) If $\pi[s] \subseteq U$ for $s \in \mathcal{A}$ and $U \in C$, then there is some $t \in \mathcal{A}$ such that $s \subseteq t$ and $\pi[t] = U$.

Clearly, (1) and (2) coincide via $\mathbf{sPsh}(C) \simeq \mathbf{ndSimp}/C$ (Fact 2). Hence their full subcategories of no-signalling models are equivalent. Note that (1) or (2) implies $A_U \neq \emptyset$ for all $U \in C$, if $A_\emptyset \neq \emptyset$. So, while the empty model is no-signalling, all the other, nonempty no-signalling models (which are, essentially, the “empirical models” of [2]) are locally consistent.

3. CONTEXTUAL LOGICS

After reviewing a kind of contextuality argument for topological models, we explain why the logic of such argument is *not* supposed to be sound with respect to those models, and why we need another logic, viz. the logic of topological models. We then introduce our candidate for such a logic.

3.1. Contextuality Argument: Logic of Global Inconsistency. Viewing A_U as representing a constraint on assignments of values to variables $x \in U$, we can describe a topological model A using formulas in contexts $U \in C$ of variables. E.g., the assignments of $(0, 0)$ and $(1, 1)$ to (x, y) satisfy the equation $x \oplus y = 0$, where \oplus is for XOR, i.e. addition modulo 2; the assignments $(0, 1)$ and $(1, 0)$ satisfy $x \oplus y = 1$. Therefore the PR box, (d) of Figure 1, satisfies the following set of equations:

$$(3) \quad a_1 \oplus b_1 = 0, \quad a_1 \oplus b_2 = 0, \quad a_2 \oplus b_1 = 0, \quad a_2 \oplus b_2 = 1$$

These are in fact inconsistent: Their right-hand sides sum to 1, but the left to 0 regardless of the values of variables (since each variable occurs twice). This is to say that no global assignment of values satisfies all the constraints of A_U , i.e., that A is strongly contextual.

A family of arguments of this sort, using XOR (or parity) equations, has been given to show the strong contextuality of a range of quantum examples; the first instance in literature was in [18] for the GHZ state [8]. This sort of so-called “all-vs-nothing argument” was formalized and generalized in [2]. On the other hand, one may also adopt more expressive languages, such as Boolean formulas, to express a wider range of constraints.

Formulas can also be used to show logical (and not strong) contextuality. E.g., the Hardy model, (c) of Figure 1, satisfies the antecedents of

$$(4) \quad \neg a_1 \vee \neg b_2, \quad \neg a_2 \vee \neg b_1, \quad a_2 \vee b_2 \quad \vdash \quad \neg a_1 \vee \neg b_1$$

but not the consequent, due to the contextual section $(1, 1)$ over (a_1, b_1) . This shows that this local section can be part of no global assignment satisfying all the constraints.

Yet this kind of contextuality argument needs some reflection. The inconsistency of a set Γ of formulas, $\Gamma \vdash \perp$, does not mean that Γ has no model; in fact, the PR box, (d) of Figure 1, satisfies all the equations in (3). In the same vein, the derivability $\Gamma \vdash \varphi$ does not mean that every model of Γ satisfies φ ; the Hardy model (c) satisfies Γ but not φ of (4). So the logic of \vdash here is *not* sound with respect to contextual models—indeed, that is the whole point of the argument. Invalidating \vdash precisely means contextuality: $\Gamma \vdash \perp$ really means that no global section satisfies Γ ; it is why any model of Γ has no global section. $\Gamma \vdash \varphi$ means that every global section satisfying Γ satisfies φ ; it is why any model satisfying Γ but not φ must have local sections (viz. ones not satisfying φ) that fail to extend to global sections.

In this sense, the logic of \vdash here is a “global logic” of global sections. We should then note that this logic, by itself, says very little about local consistency. To see this, consider:

$$(5) \quad a_1 \oplus b_1 = 0, \quad a_1 \oplus b_1 = 1$$




FIGURE 2. Charlie and an inchworm

This set of equations is, like (3), inconsistent. It is, however, inconsistent not just globally but also locally: Not only does no global section satisfy both equations, no local section over the context $\{a_1, b_1\}$ does; a model A satisfies (5) only if $A_{\{a_1, b_1\}} = \emptyset$ (the physical system can give no outcomes to the measurements a_1, b_1 ; the sentences a_1, b_1 are not just inconsistent but can have no truth values). Yet the global logic does not tell us why (3) is locally consistent whereas (5) is not. Thus the kind of argument above is really a “global-inconsistency argument”: It shows contextuality only because we already know the formulas to be locally consistent, having obtained them as descriptions of some model.

3.2. “Inchworm Logic” of Local Inference. In contrast to (3), $\Gamma \vdash \perp$ of (5) means local inconsistency over $\{a_1, b_1\}$ since both equations in Γ are in the context $\{a_1, b_1\}$. Turning $\Gamma, \varphi \vdash \perp$ into the form of inference, if Γ, φ are in a context U , then $\Gamma \vdash \neg\varphi$ gives local entailment over U . E.g., the antecedents of

$$a_1 = 0, \quad b_1 = 0 \quad \vdash \quad a_1 \oplus b_1 = 0$$

rule out all the sections over (a_1, b_1) except $(0, 0)$, which satisfies the consequent.

Indeed, local inference can be carried out across different contexts, validly in no-signalling models, subject to one constraint. To see this, expand the base C in Figure 1 from (f) of Figure 2 to (g), where the four triangles are in C —so a new experimenter, Charlie, can make his measurement c along with Alice and Bob. Now rewrite the locally consistent (3) in the inference form (6) (we replace $x \oplus y = 0$ with simpler $x = y$) and compare it to (7):

$$(6) \quad a_1 = b_1, \quad a_1 = b_2, \quad a_2 = b_1 \quad \vdash \quad a_2 = b_2$$

$$(7) \quad a_1 = b_1, \quad a_1 = c, \quad a_2 = b_1 \quad \vdash \quad a_2 = c$$

(7) is valid in no-signalling models, whereas (6) is not (the PR box is a countermodel, as it is a model for (3)). The only difference is c replacing b_2 —this tiny difference, however, enables us to obtain (7) in the following two steps:

$$(8) \quad \frac{\frac{a_1 = b_1 \quad a_1 = c}{b_1 = c} U \quad a_2 = b_1}{a_2 = c} V$$

The first step is within the context $U = \{a_1, b_1, c\}$, hence valid locally: Every section over U satisfying the antecedents satisfies the consequent. Similarly, the second step is valid within $V = \{a_2, b_1, c\}$. The key aspect is that the formula in the middle, $b_1 = c$, can be in the context $U \cap V$ and so in U or in V . The upshot is that information gets passed on from a larger context U to a smaller $U \cap V$ and then to another larger V —just like the locomotion of an inchworm, if (h) of Figure 2 helps to visualize it. Crucially, the no-signalling property is essential when the inchworm moves from a larger context to a smaller: E.g., the first step of (8) concludes that every section over $\{a_1, b_1, c\}$ satisfies $b_1 = c$; but

then, in the absence of no-signalling, there may be a section over $\{b_1, c\}$ violating $b_1 = c$ without extending to $\{a_1, b_1, c\}$. We will discuss the semantic rôle of no-signalling further in Subsection 4.3.

3.3. Formalizing the Inchworm. We formalize and generalize the idea of “inchworm inference”. As in the example in Subsection 3.2, an inchworm logic is obtained by constraining a global logic. We assume this logic to be (at least) regular, i.e. to have \top , \wedge , and \exists , for the reasons explained shortly.

Definition 5. Let \mathcal{L} be a language of regular logic (or richer) whose variables include X . For each $x \in X$, write T_x for the type of x , and then, for each $\bar{x} \subseteq X$, write $\Phi_{\bar{x}}$ for the set of formulas in the context $\bar{x} : T_{\bar{x}}$. Given a simplicial complex C on X , the C -contextual fragment of \mathcal{L} is $\Phi_C = \bigcup_{U \in C} \Phi_U$. By a C -contextual language \mathcal{L}_C , we simply mean a pair of such \mathcal{L} and Φ_C . Now let \mathbb{T} be a regular theory in \mathcal{L} given by an entailment relation \vdash (which is *not* required to be binary). Then the inchworm fragment of \vdash in \mathcal{L}_C is the entailment relation \vdash_C on Φ_C defined inductively by the following. We write $\Gamma_U = \Gamma \cap \Phi_U$.

- (9) $\Gamma \vdash_C \varphi$ if there is $U \in C$ such that $\varphi \in \Phi_U$ and $\Gamma_U \vdash \varphi$.
- (10) If $\Gamma \vdash_C \varphi$ and $\Delta, \varphi \vdash_C \psi$ then $\Gamma, \Delta \vdash_C \psi$.

(9) expresses the idea that \vdash within a single context is valid locally as well. In (10), note that the first two instances of \vdash_C may be witnessed by different contexts. Observe also that $\Gamma \vdash_C \varphi$ entails $\Gamma \vdash \varphi$; thus \vdash_C is a fragment of \vdash .

Example 6. Let \mathcal{L} have T as a basic type; $0, 1$ be constants of type T ; and \oplus be a function symbol of type $\oplus : T \times T \rightarrow T$. Let $T_x = T$ for all $x \in X$. So, e.g., $x : T, y : T \mid x \oplus y = 1$ makes sense, and is in $\Phi_{\{x, y\}}$. This gives equations of the kind relevant to the examples in Subsection 3.1. Note that Φ_C is a union. E.g., for (f) of Figure 2, $a_i = 0$ is in $\Phi_{\{a_i\}} \subseteq \Phi_C$ for both $i = 1, 2$, but $a_1 = 0 \wedge a_2 = 0$ is *not* in Φ_C since $\{a_1, a_2\} \notin C$.

We assume \mathcal{L} to have \top and \wedge , so that pieces of information can be combined within the same context. The inchworm moves from a smaller context U to a larger V via the order embedding $i : (\Phi_U, \vdash_U) \hookrightarrow (\Phi_V, \vdash_V)$, and from V to U via the left adjoint $\exists_{V \setminus U}$ of i .

$$\Phi_V \begin{array}{c} \xrightarrow{\exists_{V \setminus U}} \\ \xleftarrow[\begin{array}{c} \perp \\ i \end{array}]{\phantom{\exists_{V \setminus U}}} \end{array} \Phi_U$$

Then $\exists_{V \setminus U} \dashv i$ means that, for any $\varphi \in \Phi_V$, $\exists_{V \setminus U}. \varphi \in \Phi_U$ encapsulates all and only the information that φ entails on U . We also have $\exists_{V \setminus U} \circ i \cong 1$, so a piece of information that can be both about U and about V undergoes no change when carried across U and V .

4. CONTEXTUAL SEMANTICS IN REGULAR CATEGORIES

Our definition of model using a sheaf generalizes by replacing **Sets** with any category **S** with finite limits, since the base C is a simplicial complex. Yet, for the sake of no-signalling, we moreover need **S** to be regular. References on regular categories and their categorical logic include [20, 5, 12]. We then lay out how to model the inchworm logic in **S**.

4.1. Topological Models in Regular Categories. Let C be a simplicial complex on a set X , and \mathbf{S} be a category with finite limits. By a presheaf on C valued in \mathbf{S} , we mean any contravariant functor $P : C^{\text{op}} \rightarrow \mathbf{S}$. Then the definitions of separated presheaf and sheaf generalize straightforwardly to

Definition 7. We say that an \mathbf{S} -valued presheaf P on C is separated if the arrow

$$\langle P_{U_i \subseteq U} \rangle_i : P_U \rightarrow \prod_i P_{U_i}$$

is monic whenever $\bigcup_i U_i = U$, and a sheaf if, whenever $\bigcup_i U_i = U$, $\langle P_{U_i \subseteq U} \rangle_i$ is an equalizer as follows, where $p_j : \prod_i P_{U_i} \rightarrow P_{U_j}$ and $p_k : \prod_i P_{U_i} \rightarrow P_{U_k}$ are the projections.

$$P_U \xrightarrow{\langle P_{U_i \subseteq U} \rangle_i} \prod_i P_{U_i} \xrightarrow[\langle P_{U_j \cap U_k \subseteq U_k} \circ p_k \rangle_{j,k}]{\langle P_{U_j \cap U_k \subseteq U_j} \circ p_j \rangle_{j,k}} \prod_{j,k} P_{U_j \cap U_k}$$

Again, every sheaf F has $F_\emptyset = 1$, the terminal object of \mathbf{S} , and every separated presheaf P has $P_\emptyset \rightarrow 1$. Also, for simpler descriptions, Fact 1 generalizes to

Fact 8. An \mathbf{S} -valued presheaf P on a simplicial complex C is a sheaf iff $P_U = \prod_{x \in U} P_x$ for all $U \in C$. And P is separated iff it is a subpresheaf of a sheaf, i.e. iff each $\langle P_{\{x\} \subseteq U} \rangle_{x \in U} : P_U \rightarrow \prod_{x \in U} P_x$ is monic, i.e. iff each P_U is a relation in \mathbf{S} on $(P_x)_{x \in U}$.

Next we define the no-signalling property for \mathbf{S} -valued separated presheaves. In doing so, we need to choose from several generalizations of the notion of surjection in (1) of Definition 4; the one that serves our purpose is the one that provides semantics for $\exists : \Phi_V \rightrightarrows \Phi_U : i$ in the inchworm logic. This is the principal reason we need \mathbf{S} to be regular; then, in \mathbf{S} , every arrow $f : C \rightarrow D$ gives rise to the adjoint pair

$$\text{Sub}_{\mathbf{S}}(C) \xrightleftharpoons[\begin{smallmatrix} \perp \\ f^{-1} \end{smallmatrix}]{\exists_f} \text{Sub}_{\mathbf{S}}(D)$$

(e.g. [5, Lemma 2.5]), and moreover $\exists_f \circ f^{-1} = 1_{\text{Sub}(D)}$ (and so f^{-1} is an order embedding) if f is a regular epi (essentially, [12, Corollary D1.2.8]). Therefore the right generalization of Definition 4 is the following Definition 9, with an alternative description in Fact 10.

Definition 9. A separated presheaf A on a simplicial complex C valued in a regular category \mathbf{S} is said to be no-signalling if every $A_{U \subseteq V} : A_V \rightarrow A_U$ is a regular epi.

Fact 10. Let F be a sheaf on a simplicial complex C . Then a family $(i_U : A_U \rightarrow F_U)_{U \in C}$ of subobjects forms a subpresheaf of F , and hence a separated presheaf, iff $A_V \leq F_{U \subseteq V}^{-1}(A_U)$, or equivalently $\exists_{F_{U \subseteq V}}(A_V) \leq A_U$, whenever $U \subseteq V \in C$. Moreover, a separated presheaf $i : A \rightarrow F$ is no-signalling iff $\exists_{F_{U \subseteq V}}(A_V) = A_U$ whenever $U \subseteq V \in C$.

4.2. Global Inconsistency in Regular Categories. Definition 3 of contextuality for **Sets**-valued presheaves can now extend to ones valued in any regular category \mathbf{S} . Let A be an \mathbf{S} -valued separated presheaf on a simplicial complex C on X . It is a subpresheaf of a sheaf F on C . In fact, let us assume, just in this subsection, that X is finite (or that \mathbf{S} is complete); then, by Fact 8 (or a straightforward generalization), F extends uniquely to a sheaf on $\mathcal{P}X$, viz. $F : U \mapsto \prod_{x \in U} F_x$. Then the set of global sections of A —i.e. the natural join

$$\bigvee_{U \in C} A_U = \{g \in \prod_{x \in X} A_x \mid g|_U \in A_U \text{ for all } U \in C\}$$

of the relations $A_U \subseteq F_U$ —generalizes to the \mathbf{S} -valued case:

Fact 11. Given any \mathbf{S} -valued separated presheaf A , let F be a sheaf such that $i : A \rightarrowtail F$ and, using A_U as predicates in the internal language of \mathbf{S} , define

$$\bowtie A = \llbracket \bar{x} : F_X \mid \bigwedge_{U \in C} A_U(F_{U \subseteq X} \bar{x}) \rrbracket = \bigwedge_{U \in C} F_{U \subseteq X}^{-1}(A_U) \rightarrowtail F_X.$$

Then $\bowtie A$ is the limit of A as a diagram in \mathbf{S} .

For each $U \in C$, write $\rho_U : \bowtie A \rightarrow A_U$ for the restriction of $F_{U \subseteq X}$ to $\bowtie A$; it generalizes the restriction of global sections to local sections over U . Definition 3 then extends to

Definition 12. An \mathbf{S} -valued separated presheaf A is said to be logically contextual if not every $\rho_U : \bowtie A \rightarrow A_U$ is a regular epi. A is moreover said to be strongly contextual if $\bowtie A$ is not well-supported, i.e. if the unique arrow $!_{\bowtie A} : \bowtie A \rightarrow 1$ is not a regular epi.

Rewriting this in the internal language of \mathbf{S} , the strong contextuality of A means that \mathbf{S} fails $\exists \bar{x} : F_X. \bowtie A(\bar{x})$, i.e., that no global section \bar{x} satisfies all the constraints A_U . The logical contextuality means that

$$\bar{x} : F_V \mid A_V(\bar{x}) \vdash \exists \bar{y} : F_{X \setminus V}. \bowtie A(\langle \bar{x}, \bar{y} \rangle)$$

fails in \mathbf{S} for some $V \in C$, i.e., that not every local section \bar{x} over V satisfying A_V extends to a global section $\langle \bar{x}, \bar{y} \rangle$ satisfying all A_U . (We should stress that the definition makes sense for any separated presheaves A and not just no-signalling ones.)

4.3. Contextual Interpretation. In Definition 5 we defined a contextual language \mathcal{L}_C and logic \vdash_C simply as a global language \mathcal{L} and logic \vdash paired with their contextual fragments. Our definition of an interpretation of them in regular categories goes in parallel.

Definition 13. Given a contextual language $\mathcal{L}_C = (\mathcal{L}, \Phi_C)$, an interpretation of it in a regular category \mathbf{S} is simply an interpretation $\llbracket - \rrbracket$ of \mathcal{L} in \mathbf{S} . The images of T_x and Φ_U then play special rôles: For each $\bar{x} \in C$, we have

- $\llbracket T_{\bar{x}} \rrbracket = \prod_{x \in \bar{x}} \llbracket T_x \rrbracket$; therefore $\llbracket T_- \rrbracket : C^{\text{op}} \rightarrow \mathbf{S}$ forms a sheaf by Fact 8.
- Moreover, $\llbracket \bar{x} : T_{\bar{x}} \mid \varphi \rrbracket \rightarrowtail \llbracket T_{\bar{x}} \rrbracket$ for each $\varphi \in \Phi_{\bar{x}}$.

So we may write $(\llbracket - \rrbracket, F)$ for the interpretation $\llbracket - \rrbracket$, where F is the sheaf $F : \bar{x} \mapsto \llbracket T_{\bar{x}} \rrbracket$. We may also write $\llbracket \varphi \rrbracket_{\bar{x}} \rightarrowtail F_{\bar{x}}$ for $\llbracket \bar{x} : T_{\bar{x}} \mid \varphi \rrbracket$.

Example 14. Expanding Example 6, take $\llbracket - \rrbracket$ in **Sets** with $\llbracket T \rrbracket = \mathbf{2}$ and the obvious $\llbracket 0 \rrbracket$, $\llbracket 1 \rrbracket$, and $\llbracket \oplus \rrbracket$. Then we have a sheaf $\llbracket T_- \rrbracket : U \mapsto \mathbf{2}^U$ and, e.g.,

$$\llbracket x : T, y : T \mid x \oplus y = 0 \rrbracket \rightarrowtail \llbracket T_{\{x,y\}} \rrbracket = \mathbf{2} \times \mathbf{2} \xrightarrow[\quad]{\llbracket \oplus \rrbracket} \mathbf{1} \xrightarrow[\quad]{\llbracket 0 \rrbracket} \mathbf{2}$$

is an equalizer of $\llbracket \oplus \rrbracket$ and $\llbracket 0 \rrbracket \circ !$.

An interpretation $\llbracket - \rrbracket$ of \mathcal{L} is said to model a sequent $\Gamma \vdash \varphi$ if some finite $\Delta \subseteq \Gamma$ has $\bigwedge_{\psi \in \Delta} \llbracket \psi \rrbracket_U \leq \llbracket \varphi \rrbracket_U$. This makes sense whether $U \in C$ or not. Nevertheless, if $U \notin C$, then $\bigwedge_{\psi \in \Delta} \llbracket \psi \rrbracket_U \leq \llbracket \varphi \rrbracket_U$ only means the global entailment and not the local one. Take, e.g.,

Example 15. Expanding Example 14, the presheaf model A of the PR box, (d) of Figure 1, is a subpresheaf of $\llbracket T_- \rrbracket$ described by (3): E.g. $A_{\{a_i, b_1\}} = \llbracket a_1 \oplus b_1 = 0 \rrbracket_{\{a_1, b_1\}} \rightarrowtail \mathbf{2} \times \mathbf{2}$. Then the global inconsistency, and strong contextuality $\Gamma \vdash \perp$ in particular, of the equations Γ in (3) means $\bowtie A = \bigcap_{\varphi \in \Gamma} \llbracket \varphi \rrbracket_X \subseteq \llbracket \perp \rrbracket_X = \emptyset$. Yet Γ is locally consistent, modelled by the PR box.

This is why, to model the inchworm logic of local inference, we need a presheaf on different contexts $U \in C$, as opposed to an intersection in a single context $V \notin C$, to the left of \leq .

Definition 16. Suppose $(\llbracket - \rrbracket, F)$ is an interpretation of a contextual language \mathcal{L}_C . Then let us say that a subpresheaf $A \rightarrowtail F$ is a *pre-model* in $(\llbracket - \rrbracket, F)$ of a formula $\varphi \in \Phi_U$ in a context $U \in C$, and write $A \models_U \varphi$, to mean that $A_U \leq \llbracket \varphi \rrbracket_U$.

Fact 17. If $A \leq B$ for subpresheaves A and B of F , then $B \models_U \varphi$ implies $A \models_U \varphi$.

Note, however, that this notion of pre-model is context-dependent and concerns formulas in contexts as opposed to formulas *per se*. When $U \subseteq V \in C$ and $\varphi \in \Phi_U$, Fact 10 yields

(11) $A \models_U \varphi$ entails $A \models_V \varphi$, because

$$\frac{\exists_{F_{U \subseteq V}}(A_V) \leq A_U \leq \llbracket \varphi \rrbracket_U}{A_V \leq F_{U \subseteq V}^{-1} \llbracket \varphi \rrbracket_U = \llbracket \varphi \rrbracket_V}$$

(12) Suppose A is no-signalling. Then $A \models_V \varphi$ entails $A \models_U \varphi$. This is because

$$\frac{A_V \leq \llbracket \varphi \rrbracket_V = F_{U \subseteq V}^{-1} \llbracket \varphi \rrbracket_U}{A_U = \exists_{F_{U \subseteq V}}(A_V) \leq \llbracket \varphi \rrbracket_U}$$

If A is not no-signalling, (12) may fail, and then inchworm inference fails. E.g., in (8), the first step purports to show that, if $A \models_U a_1 = b_1$ and $A \models_U a_1 = c$, then $A \models_U b_1 = c$ and so $A \models_{U \cap V} b_1 = c$; but the “and so” step here requires (12). In this sense, as mentioned in Subsection 2.4, no-signalling means the context-independent coherence of a presheaf as a model of formulas. Therefore

Definition 18. A pre-model A is called a (no-signalling) *model* if it is no-signalling. Then we say that A is a model of a formula $\varphi \in \Phi_C$, and write $A \models \varphi$, to mean that A is a pre-model of φ in any suitable context, i.e., that $A_U \leq \llbracket \varphi \rrbracket_U$ for every $U \in C$ such that $\varphi \in \Phi_U$.

Theorem 19. Let $\llbracket - \rrbracket$ be an interpretation of \mathcal{L}_C that models a theory \vdash in \mathcal{L} . Then the inchworm logic \vdash_C of \vdash is sound with respect to the no-signalling models in $\llbracket - \rrbracket$: If $\Gamma \vdash_C \varphi$, then $A \models \varphi$ for every no-signalling model A of Γ in $\llbracket - \rrbracket$.

4.4. The Inchworm and No-Signalling. Subsection 4.3 primarily concerned how given presheaves modelled formulas. We showed in particular that no-signalling validated inchworm inference. Let us discuss, on the other hand, how the description by given formulas yields a model. This shows the other direction of the connection between no-signalling and the inchworm, from the latter to the former.

We say a set $\Gamma \subseteq \Phi_C$ of formulas of \mathcal{L}_C is *C-finite* if Γ_U is finite for each $U \in C$.

Definition 20. Let $(\llbracket - \rrbracket, F)$ be an interpretation of \mathcal{L}_C . Given any C-finite $\Gamma \subseteq \Phi_C$, define $\mathbb{M}_F(\Gamma)$ as a family $(\mathbb{M}_F(\Gamma)_U = \bigwedge_{\varphi \in \Gamma_U} \llbracket \varphi \rrbracket_U \rightarrowtail F_U)_{U \in C}$ of subobjects of F_U .

Fact 21. $\mathbb{M}_F(\Gamma)$ is the largest subpresheaf A of F such that $A \models_U \Gamma_U$ for each $U \in C$.

$\mathbb{M}_F(\Gamma)$ generally fails to be no-signalling. E.g., A in (e) of Figure 1 is $\mathbb{M}_F(\Gamma)$ given by $\Gamma = \{\varphi\}$ for $\varphi = (a_2 \wedge \neg b_1) \vee (\neg a_2 \wedge \neg b_1)$; since φ cannot be in the context $\{b_1\}$, $\Gamma_{\{b_1\}} = \emptyset$ and $A_{\{b_1\}} = \mathbf{2}$. Yet the description by Γ sometimes manages to give a no-signalling $\mathbb{M}_F(\Gamma)$.

Fact 22. Let $(\llbracket - \rrbracket, F)$ be an interpretation of \mathcal{L}_C that models a theory \vdash in \mathcal{L} . We say $\Gamma \subseteq \Phi_C$ is *inchworm-saturated* if $\Gamma_V \vdash \varphi$ implies $\Gamma_U \vdash \exists_{V \setminus U} \varphi$ whenever $U \subseteq V \in C$ and $\varphi \in \Phi_V$. Now, if a C-finite Γ is inchworm-saturated, then $\mathbb{M}_F(\Gamma)$ is no-signalling.

When Γ is inchworm-saturated, it may not be deductively closed, but the inchworm cannot bring a new piece of information ψ to a context U from another V , since ψ follows from the information Γ_U that U already has. Fact 22 means that, if Γ is inchworm-saturated and if each Γ_U finite and consistent (and has $\mathbb{M}_F(\Gamma)_U$ nonempty or well-supported), then Γ is locally consistent, modelled by a no-signalling model $\mathbb{M}_F(\Gamma)$. E.g., (3) is inchworm-saturated, with each context consistent, so it gives the PR box, (d) of Figure 1, as $\mathbb{M}_F(\Gamma)$.

On the other hand, even when a description Γ is not inchworm-saturated and $\mathbb{M}_F(\Gamma)$ fails to be no-signalling, the inchworm can carve out the “no-signalling interior” of $\mathbb{M}_F(\Gamma)$, if Γ can be saturated in finite (or C -finite) steps.

Theorem 23. *Let $(\llbracket - \rrbracket, F)$ be an interpretation of \mathcal{L}_C that models a theory \vdash in \mathcal{L} . Given $\Gamma \subseteq \Phi_C$, suppose there is a C -finite and inchworm-saturated $\Delta \subseteq \Phi_C$ such that $\Gamma \subseteq \Delta$ and $\Gamma \vdash_C \varphi$ for all $\varphi \in \Delta$. Then $\mathbb{M}_F(\Delta)$ is the largest no-signalling subpresheaf of $\mathbb{M}_F(\Gamma)$.*

Take again the example from right after Fact 21: $\mathbb{M}_F(\Gamma)$ of $\Gamma = \{\varphi\}$ for $\varphi = (a_2 \wedge \neg b_1) \vee (\neg a_2 \wedge \neg b_1)$ is A in (e) of Figure 1 and fails to be no-signalling. Yet $\varphi \vdash \neg b_1$, so $\Gamma \vdash_C \neg b_1$, and $\Delta = \Gamma \cup \{\neg b_1\}$ is inchworm-saturated, with $\neg b_1 \in \Delta_{[b_1]}$. Hence, by Theorem 23, the inchworm carves out a no-signalling $\mathbb{M}_F(\Delta)$ by removing the red sections from (e). Indeed, in many applications (e.g. all the examples in Sections 2 and 3), the theory \vdash satisfies

- (13) Given any $\Gamma \subseteq \Phi_C$ (that may not be C -finite), for each $U \in C$ there is a finite $\Delta_U \subseteq \Gamma_U$ such that $\Delta_U \vdash \varphi$ for all $\varphi \in \Gamma_U$.

This guarantees the supposition of Theorem 23: Given any C -finite $\Gamma \subseteq \Phi_C$, take its \vdash_C -deductive closure $\Gamma^* = \{\varphi \mid \Gamma \vdash_C \varphi\}$ as Γ in (13) and obtain Δ_U ; then $\Delta = \Gamma \cup \bigcup_{U \in C} \Delta_U$ is such as in Theorem 23. Therefore Theorem 23 applies and leads to a family of completeness results organized by Lemma 24, which transfers a completeness theorem of a global theory to its inchworm fragment. It yields, e.g., Theorem 25, since any (global) regular theory has a “conservative model” in a “classifying category” (e.g. [5, Proposition 6.4]).

Lemma 24. *Suppose that a theory \vdash in \mathcal{L} satisfies (13), and that $\llbracket - \rrbracket$ is a conservative model of \vdash , meaning that, for any $\Gamma \subseteq \Phi_C$, $\bigwedge_{\psi \in \Delta} \llbracket \psi \rrbracket_U \leq \llbracket \varphi \rrbracket_U$ for some $\Delta \subseteq \Gamma$ if but also only if $\Gamma \vdash \varphi$. Then $\Gamma \vdash_C \varphi$ iff $A \models \varphi$ for every no-signalling model A of Γ in $\llbracket - \rrbracket$.*

Theorem 25. *Let \vdash be a regular theory satisfying (13). Then, for any $\Gamma \subseteq \Phi_C$, $\Gamma \vdash_C \varphi$ iff $A \models \varphi$ for every no-signalling model A of Γ in every model $\llbracket - \rrbracket$ of \vdash in any regular category.*

4.5. Completion for Completeness. Generally, the property (13) may fail and the inchworm saturation may not be attained in finite steps.

Example 26. In Figure 1, replace each $A_x = \mathbf{2}$ with \mathbb{Z} , and let Γ be the set of formulas

$$a_1 = b_2, \quad b_1 = a_1, \quad a_2 = b_1, \quad b_2 = a_2 + 1, \quad b_2 > 0$$

in the obvious \mathcal{L} and \vdash . Then $\Gamma \vdash_C a_1 > 0, b_1 > 0, a_2 > 0, b_2 > 1, \dots, x > n$ for every $x \in X$ and $n \in \mathbb{N}$, whereas $\Gamma \not\vdash_C \perp$ (although the empty presheaf is the only no-signalling model of Γ). So there cannot be any such Δ as in Theorem 23. (Note that the topology of C is essential: E.g., if we take $C = \mathcal{P}X$ instead, then $\Gamma \vdash_C \perp$ by $\Gamma \vdash \perp$.)

Thus, even if Γ is finite, the set $\{\llbracket \varphi \rrbracket_U \mid \varphi \in \Gamma^*_U\}$ may have no minimum (though it is lower-bounded by $\exists_{F_{U \subseteq X}} (\bigwedge_{\psi \in \Gamma} \llbracket \psi \rrbracket_X)$ if X is finite); then, in a regular category in general, $\bigwedge_{\varphi \in \Gamma^*_U} \llbracket \varphi \rrbracket_U$ may not exist. So, instead of the semilattice $\text{Sub}_S(F_U)$ of subobjects, let us use a completion of it, viz. the semilattice $\text{Filt}(\text{Sub}_S(F_U))$ of filters in $\text{Sub}_S(F_U)$, and assign a filter of subobjects, instead of a subobject, to each $U \in C$.

Definition 27. Suppose $(\llbracket - \rrbracket, F)$ is an interpretation of a contextual language \mathcal{L}_C . Then, by a *filter model* in $(\llbracket - \rrbracket, F)$, we mean a presheaf $G : C^{\text{op}} \rightarrow \mathbf{Sets}$ such that

- $G_U \in \text{Filt}(\text{Sub}_S(F_U))$ for every $U \in C$.
- For $U \subseteq V \in C$,

$$G_U = \{ S \multimap F_U \mid F_{U \subseteq V}^{-1}(S) \in G_V \} = \{ \exists_{F_{U \subseteq V}}(S) \multimap F_U \mid S \in G_V \},$$

so $G_{U \subseteq V} : G_V \rightarrow G_U :: S \mapsto \exists_{F_{U \subseteq V}}(S)$ is a surjection.

We say G models φ , and write $G \models \varphi$, to mean that $\llbracket \varphi \rrbracket_U \in G_U$ whenever $\varphi \in \Phi_U$.

Then we have the filter versions of Theorem 19, Fact 22, and completeness results organized by Lemma 24. Observe that every (no-signalling) model A is a “principal” filter model, $U \mapsto \uparrow A_U = \{ S \multimap F \mid A_U \leq S \}$; so Theorem 28 is stronger than Theorem 19.

Theorem 28. Let $\llbracket - \rrbracket$ be a model of a theory \vdash in \mathcal{L} . Then the inchworm logic \vdash_C of \vdash is sound with respect to the filter models in $\llbracket - \rrbracket$: If $\Gamma \vdash_C \varphi$, then $G \models \varphi$ for every filter model G of Γ in $\llbracket - \rrbracket$.

Fact 29. Let $(\llbracket - \rrbracket, F)$ be a model of a theory \vdash in \mathcal{L} . Given any $\Gamma \subseteq \Phi_C$, the family $\text{FM}_F(\Gamma) = (\{ S \multimap F_U \mid \llbracket \varphi \rrbracket_U \leq S \text{ for some } \varphi \in \Gamma^*_U \})_{U \in C}$ is a filter model of Γ in $(\llbracket - \rrbracket, F)$. Moreover, for any filter model G of Γ in $(\llbracket - \rrbracket, F)$, $\text{FM}_F(\Gamma)_U \subseteq G_U$ for each $U \in C$.

Lemma 30. Suppose $\llbracket - \rrbracket$ is a conservative model of a theory \vdash in \mathcal{L} . Then $\Gamma \vdash_C \varphi$ iff $G \models \varphi$ for every filter model G of Γ in $\llbracket - \rrbracket$.

5. CONCLUSION

We have formulated contextual models as presheaves valued in regular categories, as well as providing “inchworm logic” for local inference in those contextual models. We have also proven “completeness-transfer lemmas” (Lemmas 24 and 30), so that completeness theorems in categorical logic transfer straightforwardly to inchworm logic.

Let us conclude the paper by discussing connections and applications between the framework of this paper and other approaches or other fields as future work. First of all, categorical logic has a long tradition (since [16]) of viewing local truth as a modal operator. Indeed, the logic of local information in this paper is closely related to the dynamic-logical characterization of contextuality in [13]. There is also a connection to model theory. For instance, the similarity between inchworm inference and Craig interpolation should be obvious; indeed, by defining Φ_U more generally as a “language in the vocabulary U ”, we can prove a stronger version of Robinson’s joint consistency theorem (see [6, Subsection 4.1.1]) that is sensitive to the topology of C .

As explained in Section 2, presheaf models, whether no-signalling or not, can model Boolean valuations. This enables us to transfer and apply techniques from satisfiability problems to quantum contextuality as computational resource. Another connection is to the structure of valuation algebra, which is used for local computation [14]. In fact, our presheaf models can also be formulated in

terms of valuation algebras, as a C -indexed family of valuations satisfying certain conditions. We can expect these connections to help extend local computation to situations in classical computing where contextual phenomena arise.

The generality of taking presheaves in regular categories is also expected to facilitate applications. In cohomology, it is typical to use presheaves valued in regular categories, such as presheaves of abelian groups, R -modules, etc. Therefore the framework of this paper applies to the logic of local inference within such presheaves. One can also take regular categories of structures that are used for other purposes such as modelling processes in quantum physics. In addition, the connection to logical paradoxes [2] is also relevant. As shown in [15, 4], regular categories provide background for self-referential and other fixpoint paradoxes; so our formalism will unify the two perspectives on paradoxes.

REFERENCES

- [1] S. Abramsky. Relational databases and Bell’s theorem. In V. Tannen, et al., editors, *In Search of Elegance in the Theory and Practice of Computation: Essays Dedicated to Peter Buneman*, pages 13–35. Springer, 2013.
- [2] S. Abramsky, R. Soares Barbosa, K. Kishida, R. Lal, and S. Mansfield. Contextuality, cohomology and paradox. In S. Kreutzer, editor, *CSL 2015*, pages 211–228. 2015.
- [3] S. Abramsky and A. Brandenburger. The sheaf-theoretic structure of non-locality and contextuality. *New J. Phys.*, 13:113036–113075, 2011.
- [4] S. Abramsky and J. A. Zvesper. From Lawvere to Brandenburger-Keisler: Interactive forms of diagonalization and self-reference. *J. Comput. Syst. Sci.*, 81(5):799–812, 2015.
- [5] C. Butz. Regular categories and regular logic. BRICS Lecture Series LS-98-2, BRICS, 1998.
- [6] D. M. Gabbay and L. Maksimova. *Interpolation and Definability*. Clarendon Press, 2005.
- [7] G.-C. Ghirardi, A. Rimini, and T. Weber. A general argument against superluminal transmission through the quantum mechanical measurement process. *Lettere al Nuovo Cimento*, 27:293–298, 1980.
- [8] D. M. Greenberger, M. A. Horne, and A. Zeilinger. Going beyond Bell’s theorem. In M. Kafatos, editor, *Bell’s Theorem, Quantum Theory, and Conceptions of the Universe*, pages 69–72. Kluwer, 1989.
- [9] L. Hardy. Nonlocality for two particles without inequalities for almost all entangled states. *Phys. Rev. Lett.*, 71:1665–1668, 1993.
- [10] M. Howard, J. Wallman, V. Veitch, and J. Emerson. Contextuality supplies the ‘magic’ for quantum computation. *Nature*, 510:351–355, 2014.
- [11] C. J. Isham and J. Butterfield. Topos perspective on the Kochen–Specker theorem: I. Quantum states as generalized valuations. *Int. J. Theor. Phys.*, 37(11):2669–2733, 1998.
- [12] P. T. Johnstone. *Sketches of an Elephant: A Topos Theory Compendium*, volume 2. Clarendon Press, Oxford, 2002.
- [13] K. Kishida. Stochastic relational presheaves and dynamic logic for contextuality. In B. Coecke, et al., editors, *QPL 2014*, pages 115–132. 2014.
- [14] J. Kohlas, M. Pouly, and C. Schneuwly. Generic local computation. *J. Comput. Syst. Sci.*, 78(1):348–369, 2012.
- [15] F. W. Lawvere. Diagonal arguments and cartesian closed categories. *Lecture Notes in Mathematics*, 92:134–145, 1969.
- [16] F. W. Lawvere. Quantifiers and sheaves. *Actes, Congrès intern. math.*, 1:329–334, 1970.
- [17] S. Mac Lane and I. Moerdijk. *Sheaves in Geometry and Logic: A First Introduction to Topos Theory*. Springer, 1992.
- [18] N. D. Mermin. Extreme quantum entanglement in a superposition of macroscopically distinct states. *Phys. Rev. Lett.*, 65:1838–1840, 1990.
- [19] M. A. Nielsen and I. L. Chuang. *Quantum Computation and Quantum Information: 10th Anniversary Edition*. Cambridge University Press, 2011.
- [20] J. van Oosten. Basic category theory. BRICS Lecture Series LS-95-1, BRICS, 1995.
- [21] S. Popescu and D. Rohrlich. Quantum nonlocality as an axiom. *Found. Phys.*, 24:379–385, 1994.
- [22] R. Raussendorf. Contextuality in measurement-based quantum computation. *Phys. Rev. A*, 88:022322, 2013.

APPENDIX A. PROOFS

Fact 1. A presheaf P on a simplicial complex C is a sheaf iff $P_U = \prod_{x \in U} P_x$ for all $U \in C$. P is separated iff it is a subpresheaf of a sheaf, i.e. iff $P_U \subseteq \prod_{x \in U} P_x$ for all $U \in C$.

Proof. This is just an instance of Fact 8 with $\mathbf{S} = \mathbf{Sets}$. □

Fact 2. $\mathbf{sPsh}(C) \simeq \mathbf{ndSimp}/C$ for any simplicial complex C .

Proof. Let C be a simplicial complex on a set X . Define functors

$$\mathbf{sPsh}(C) \begin{array}{c} \xleftarrow{F} \\ \xrightarrow{G} \end{array} \mathbf{ndSimp}/C$$

as follows. First we define F . Let P be a separated presheaf on C . By Fact 1, $P_U \subseteq \prod_{x \in U} P_x$ for all $U \in C$. Then let FP be a map $\pi : A \rightarrow X$ from a simplicial complex \mathcal{A} on A such that

- $A = \sum_{x \in X} P_x$ and π is the projection $\pi : A \rightarrow X :: (x, a) \mapsto x$.
- $\mathcal{A} = \bigcup_{U \in C} P_U$. That is, for each $S \subseteq A$, we have $S \in \mathcal{A}$ iff there is some $U \in C$ such that $S \in P_U$, i.e. such that $S = \{(x, s(x)) \mid x \in U\}$ for some $s \in P_U \subseteq \prod_{x \in U} P_x$.

Then $FP = \pi : \mathcal{A} \rightarrow C$ is clearly a non-degenerate bundle over C .

Let $\vartheta : P \rightarrow Q$ be a natural transformation between separated presheaves P and Q on C ; write $FP = \pi_1 : \mathcal{A}_1 \rightarrow C$ and $FQ = \pi_2 : \mathcal{A}_2 \rightarrow C$, with \mathcal{A}_1 and \mathcal{A}_2 on the sets $A_1 = \sum_{x \in X} P_x$ and $A_2 = \sum_{x \in X} Q_x$. Then let $F\vartheta : A_1 \rightarrow A_2$ be the map such that

$$F\vartheta : \sum_{x \in X} P_x \rightarrow \sum_{x \in X} Q_x :: (x, a) \mapsto (x, \vartheta_x(a)).$$

$F\vartheta$ is a non-degenerate map of bundles from FP to FQ over C :

- If $s \in \mathcal{A}_1$, then $s \in P_U \subseteq \prod_{x \in U} P_x$ for some $U \in C$, and so

$$F\vartheta[s] = \{(x, \vartheta_x(s(x)) \mid x \in U\} = \{(x, (\vartheta_U(s))(x)) \mid x \in U\} = \vartheta_U(s) \in \mathcal{A}_2$$

by the naturality of ϑ .

$$\begin{array}{ccc} P_U & \xrightarrow{\vartheta_U} & Q_U \\ \downarrow & \cong & \downarrow \\ P_x & \xrightarrow{\vartheta_x} & Q_x \end{array} \qquad \begin{array}{ccc} s & \xrightarrow{\quad} & \vartheta_U(s) \\ \downarrow & & \downarrow \\ s(x) & \xrightarrow{\quad} & \vartheta_x(s(x)) = (\vartheta_U(s))(x) \end{array}$$

- $\pi_2 \circ F\vartheta = \pi_1$ because $\pi_2 \circ F\vartheta(x, a) = \pi_2(x, \vartheta_x(a)) = x = \pi_1(x, a)$.
- Because π_1 is non-degenerate and $\pi_2 \circ F\vartheta = \pi_1$, $F\vartheta$ is non-degenerate as well.

Next we define G . Given a non-degenerate bundle $\pi : \mathcal{A} \rightarrow C$, for each $U \in C$ let

$$(G\pi)_U = P_U = \{s \in \mathcal{A} \mid \pi[s] = U\}.$$

Then $G\pi = P : U \mapsto P_U$ is a presheaf, with $P_{V \subseteq U} : P_U \rightarrow P_V :: s \mapsto s \cap \pi^{-1}[V]$. Moreover $\{P_{\{x\} \subseteq U}\}_{x \in U}$ is jointly monic; so $G\pi = P$ is a separated presheaf by Fact 1.

Given a non-degenerate map of bundles $f : \pi_1 \rightarrow \pi_2$ between $\pi_i : \mathcal{A}_i \rightarrow C$ ($i = 1, 2$), let $Gf : G\pi_1 \rightarrow G\pi_2$ be the family $\{(Gf)_U\}_{U \in C}$ of functions

$$(Gf)_U : (G\pi_1)_U \rightarrow (G\pi_2)_U :: s \mapsto f[s],$$

which lands in $(G\pi_2)_U$ because f is a map over C , i.e. $\pi_2 \circ f[s] = \pi_1[s]$. To show Gf to be a natural transformation, fix any $s \in (G\pi_1)_U$. Then the non-degeneracy of π_1 , π_2 , and f , together with $\pi_2 \circ f = \pi_1$, implies that the triangle on the left below commutes and that all the three arrows are bijections. Therefore the diagram on the right, with $h = \pi_2^{-1}[-] \cap f[s]$, commutes as well:

$$\begin{array}{ccc}
 s & \xrightarrow{f|_s} & f[s] \\
 \pi_1|_s \searrow & \cong & \swarrow \pi_2|_{f[s]} \\
 & U &
 \end{array}
 \qquad
 \begin{array}{ccc}
 \mathcal{P}(s) & \xleftarrow{f[-]} & \mathcal{P}(f[s]) \\
 & \xrightarrow{f^{-1}[-] \cap s} & \\
 \pi_1^{-1}[-] \cap s \searrow & & \swarrow \pi_1[-] \\
 & \mathcal{P}(U) & \\
 h \nearrow & & \nwarrow \pi_2[-]
 \end{array}$$

In particular, $h = f[\pi_1^{-1}[-] \cap s]$, which means that the square below commutes.

$$\begin{array}{ccc}
 (G\pi_1)_U & \xrightarrow{(Gf)_U} & (G\pi_2)_U \\
 \downarrow & \cong & \downarrow \\
 (G\pi_1)_V & \xrightarrow{(Gf)_V} & (G\pi_2)_V
 \end{array}
 \qquad
 \begin{array}{ccc}
 s \vdash & \xrightarrow{\quad} & f[s] \\
 \downarrow & & \downarrow \\
 s \cap \pi_1^{-1}[V] \vdash & \xrightarrow{\quad} & f[s \cap \pi_1^{-1}[V]] = f[s] \cap \pi_2^{-1}[V]
 \end{array}$$

Thus Gf is a natural transformation.

It is obvious that $G \circ F \cong 1_{\mathbf{S}\mathbf{Psh}(C)}$ and $F \circ G \cong 1_{\mathbf{ndSimp}/C}$ naturally. \square

Fact 8. An \mathbf{S} -valued presheaf P on a simplicial complex C is a sheaf iff $P_U = \prod_{x \in U} P_x$ for all $U \in C$. P is separated iff it is a subpresheaf of a sheaf, i.e. iff each $\langle P_{\{x\} \subseteq U} \rangle_{x \in U} : P_U \rightarrow \prod_{x \in U} P_x$ is monic, i.e. iff each P_U is a relation in \mathbf{S} on $(P_x)_{x \in U}$.

Proof. For the sheaf part, suppose P is a sheaf. First, $\bigcup_{i \in \emptyset} U_i = \emptyset$ gives an equalizer

$$P_\emptyset \xrightarrow{\quad} 1 \xrightleftharpoons[1]{1} 1$$

that is an iso; so $P_\emptyset = 1$. Then for every $U \in C$, $\bigcup_{x \in U} \{x\} = U$ gives an equalizer

$$P_U \xrightarrow{\langle P_{\{x\} \subseteq U} \rangle_{x \in U}} \prod_{x \in U} P_x \xrightleftharpoons[\langle P_{\{y\} \cap \{z\} \subseteq \{z\}} \circ p_z \rangle_{y,z \in U}]{\langle P_{\{y\} \cap \{z\} \subseteq \{y\}} \circ p_y \rangle_{y,z \in U}} \prod_{y,z \in U} P_{\{y\} \cap \{z\}}$$

but then the parallel arrows are both isos, and therefore the equalizer is an iso, i.e. $P_U = \prod_x P_x$. For the other direction, suppose $P_U = \prod_{x \in U} P_x$ for all $U \in C$ and fix $\bigcup_i U_i = U$. Then the equalizer we want is

$$\prod_{x \in U} P_x \xrightarrow{\langle \langle p_x \rangle_{x \in U_i} \rangle_i} \prod_i \prod_{x \in U_i} P_x \xrightleftharpoons[h = \langle \langle p_x \rangle_{x \in U_j \cap U_k} \circ p_k \rangle_{j,k}]{g = \langle \langle p_x \rangle_{x \in U_j \cap U_k} \circ p_j \rangle_{j,k}} \prod_{j,k} \prod_{x \in U_j \cap U_k} P_x$$

But it is easy to check that any f such that $g \circ f = h \circ f$ has the form $\langle \langle f_x \rangle_{x \in U_i} \rangle_i$, and hence factors uniquely as $\langle \langle p_x \rangle_{x \in U_i} \rangle_i \circ \langle f_x \rangle_{x \in U}$. Therefore $\langle \langle p_x \rangle_{x \in U_i} \rangle_i$ is an equalizer as desired.

For the presheaf part, fix P and number the conditions as follows:

- (i) P is separated, i.e. $\langle P_{\{x\} \subseteq U} \rangle_{x \in U} : P_U \rightarrow \prod_i P_{U_i}$ is monic whenever $\bigcup_i U_i = U$.
- (ii) P is a subpresheaf of a sheaf.
- (iii) Each $\langle P_{\{x\} \subseteq U} \rangle_{x \in U} : P_U \rightarrow \prod_{x \in U} P_x$ is monic.
- (iv) Each P_U is a relation in \mathbf{S} on $(P_x)_{x \in U}$.

(iv) is defined by (iii). (i) implies (iii) as an instance. (iii) implies (ii), since $U \mapsto \prod_{x \in U} P_x$ is a sheaf by the sheaf part above. So suppose (ii), that $m : P \rightarrowtail F$ for some sheaf F . This means that the following square commutes, in which $\langle F_{U_i \subseteq U} \rangle_i$ is an equalizer and so monic.

$$\begin{array}{ccc} P_U & \xrightarrow{\langle P_{U_i \subseteq U} \rangle_i} & \prod_i P_{U_i} \\ m_U \downarrow & \cong & \downarrow \langle m_{U_i} \rangle_i \\ F_U & \xrightarrow{\langle F_{U_i \subseteq U} \rangle_i} & \prod_i F_{U_i} \end{array}$$

Therefore $\langle P_{U_i \subseteq U} \rangle_i$ is monic, too. Thus (i). \square

Fact 10. Let F be a sheaf on a simplicial complex C . Then a family $(i_U : A_U \rightarrowtail F_U)_{U \in C}$ of subobjects forms a subpresheaf of F , and hence a separated presheaf, iff $A_V \leq F_{U \subseteq V}^{-1}(A_U)$, or equivalently $\exists_{F_{U \subseteq V}}(A_V) \leq A_U$, whenever $U \subseteq V \in C$. Moreover, a separated presheaf $i : A \rightarrowtail F$ is no-signalling iff $\exists_{F_{U \subseteq V}}(A_V) = A_U$ whenever $U \subseteq V \in C$.

Proof. Fix $U \subseteq V \in C$ and write $f = F_{U \subseteq V}$. Then, in the following diagram, the longer dotted arrow making the outer square commute, i.e. $A_{U \subseteq V}$, exists iff the shorter one exists, i.e. iff $A_V \leq f^{-1}(A_U)$. Hence the first “iff” in the fact.

$$\begin{array}{ccc} A_V & \xrightarrow{\quad A_{U \subseteq V} \quad} & A_U \\ \downarrow i_V & \lrcorner & \downarrow i_U \\ F_V & \xrightarrow{f} & F_U \end{array}$$

$f^{-1}(A_U)$ is the object of the top square, and $f^{-1}(A_U) \rightarrowtail A_U$ is the arrow from the top-left to the top-right.

For the second “iff”, factorize $A_{U \subseteq V}$ into a regular epi e followed by a mono m as follows; it gives $\exists_f(A_V)$ because $(i_U \circ m) \circ e$ is a regular epi-mono factorization of $f \circ i_V$.

$$\begin{array}{ccccc} A_V & \xrightarrow{e} & \exists_f(A_V) & \xrightarrow{m} & A_U \\ i_V \downarrow & & \downarrow & \lrcorner & \downarrow i_U \\ F_V & \xrightarrow{f} & F_U & & \end{array}$$

Then, because a regular epi-mono factorization is unique, $A_{U \subseteq V} = m \circ e$ is a regular epi iff m is an iso. \square

Fact 11. (Assume that X is finite or that \mathbf{S} is complete.) Given any \mathbf{S} -valued separated presheaf A , let F be a sheaf such that $i : A \rightarrowtail F$ and, using A_U as predicates in the internal language of \mathbf{S} , define

$$\bowtie A = \llbracket \bar{x} : F_X \mid \bigwedge_{U \in C} A_U(F_{U \subseteq X} \bar{x}) \rrbracket = \bigwedge_{U \in C} F_{U \subseteq X}^{-1}(A_U) \rightarrowtail F_X.$$

Then $\bowtie A$ is the limit of A as a diagram in \mathbf{S} .

Proof. Recall that the sheaf F extends to $\mathcal{P}X$. So, for every $U \in \mathcal{C}$, write $p_U = F_{U \subseteq X} : F_X \rightarrow F_U$, and define an arrow $\rho_U : \bowtie A \rightarrow A_U$ as the composition of the two top arrows in the following.

$$\begin{array}{ccccc}
 \bowtie A & \xrightarrow{\quad} & p_U^{-1}(A_U) & \xrightarrow{\quad} & A_U \\
 & \searrow & \downarrow & \lrcorner & \downarrow \\
 & & F_X & \xrightarrow{p_U} & F_U
 \end{array}$$


Then $\bowtie A$ is a cone of the diagram A because, for every $U \subseteq V \in \mathcal{C}$, we have $A_{U \subseteq V} \circ \rho_V = \rho_U$ by chasing the following commutative diagram, where $f = F_{U \subseteq V}$.

$$\begin{array}{ccccc}
 \bowtie A & \xrightarrow{\quad} & p_V^{-1}(A_V) & \xrightarrow{\quad} & A_V \\
 & \searrow & \downarrow & \searrow & \downarrow \\
 & & p_U^{-1}(A_U) & \xrightarrow{\quad} & A_U \\
 & & \downarrow & \searrow & \downarrow \\
 & & F_X & \xrightarrow{\quad} & F_U \\
 & & \downarrow & \searrow & \downarrow \\
 & & F_V & \xrightarrow{\quad} & F_U
 \end{array}$$

Now we claim that $\bowtie A$ is the terminal among the cones of A . Fix any cone C of A with $g_U : C \rightarrow A_U$ for all $U \in \mathcal{C}$. Then, by Fact 8, for every $U \subseteq X$ we have $\langle i_x \circ g_x \rangle_{x \in U} : C \rightarrow F_U$. Therefore, for every $U \in \mathcal{C}$, the outer square below commutes and hence the arrow making the triangles commute, u_U , exists uniquely:

$$\begin{array}{ccccc}
 C & \xrightarrow{\quad} & p_U^{-1}(A_U) & \xrightarrow{\quad} & A_U \\
 \downarrow \scriptstyle \langle i_x \circ g_x \rangle_{x \in X} & \searrow \scriptstyle u_U & \downarrow & \lrcorner & \downarrow \scriptstyle i_U \\
 & & F_X & \xrightarrow{p_U} & F_U
 \end{array}$$

Then the definition $\bowtie A = \bigwedge_{U \in C} p_U^{-1}(A_U)$ means that there is a unique $v : C \rightarrow \bowtie A$ that makes the following diagram commute for each $U \subseteq V \in C$.



□

Fact 17. If $A \leq B$ for subpresheaves A and B of F , then $B \models_U \varphi$ implies $A \models_U \varphi$.

Proof. Immediate by Definition 16 of \models_U . □

Theorem 19. Let $\llbracket - \rrbracket$ be an interpretation of \mathcal{L}_C that models a theory \vdash in \mathcal{L} . Then the inchworm logic \vdash_C of \vdash is sound with respect to the no-signalling models in $\llbracket - \rrbracket$: If $\Gamma \vdash_C \varphi$, then $A \models \varphi$ for every no-signalling model A of Γ in $\llbracket - \rrbracket$.

Proof. Immediate from Theorem 28, since every model is a filter model. □

Fact 21. $\mathbb{M}_F(\Gamma)$ is the largest subpresheaf A of F such that $A \models_U \Gamma_U$ for each $U \in C$.

Proof. Let $U \subseteq V \in C$. Then, because $\Gamma_U \subseteq \Gamma_V$, because $\llbracket \varphi \rrbracket_V = F_{U \subseteq V}^{-1} \llbracket \varphi \rrbracket_U$ for each $\varphi \in \Phi_U \subseteq \Phi_V$, and because the right adjoint $F_{U \subseteq V}^{-1}$ preserves meets, we have

$$\begin{aligned} \mathbb{M}_F(\Gamma)_V &= \bigwedge_{\varphi \in \Gamma_V} \llbracket \varphi \rrbracket_V \leq \bigwedge_{\varphi \in \Gamma_U} \llbracket \varphi \rrbracket_V = \bigwedge_{\varphi \in \Gamma_U} F_{U \subseteq V}^{-1} \llbracket \varphi \rrbracket_U \\ &= F_{U \subseteq V}^{-1} \left(\bigwedge_{\varphi \in \Gamma_U} \llbracket \varphi \rrbracket_U \right) = F_{U \subseteq V}^{-1} (\mathbb{M}_F(\Gamma)_U). \end{aligned}$$

Thus $\mathbb{M}_F(\Gamma)$ is a subpresheaf of F by Fact 10, and $\varphi \in \Gamma_U$ implies $\mathbb{M}_F(\Gamma)_U = \bigwedge_{\psi \in \Gamma_U} \llbracket \psi \rrbracket_U \leq \llbracket \varphi \rrbracket_U$, i.e. $\mathbb{M}_F(\Gamma) \models_U \varphi$. Now, let A be a subpresheaf of F such that $A \models_U \Gamma_U$ for every $U \in C$. Then, for every $U \in C$ it has $A_U \leq \bigwedge_{\varphi \in \Gamma_U} \llbracket \varphi \rrbracket_U = \mathbb{M}_F(\Gamma)_U$. Thus $A \leq \mathbb{M}_F(\Gamma)$. □

Fact 22. Let $(\llbracket - \rrbracket, F)$ be an interpretation of \mathcal{L}_C that models a theory \vdash in \mathcal{L} . We say $\Gamma \subseteq \Phi_C$ is *inchworm-saturated* if $\Gamma_V \vdash \varphi$ implies $\Gamma_U \vdash \exists_{V \setminus U} \varphi$ whenever $U \subseteq V \in C$ and $\varphi \in \Phi_V$. Now, if a C -finite Γ is inchworm-saturated, then $\mathbb{M}_F(\Gamma)$ is no-signalling.

Proof. Fix any $U \subseteq V \in C$. Then the inchworm saturation of C -finite Γ implies that, since $\Gamma_V \vdash \bigwedge \Gamma_V$, we have $\Gamma_U \vdash \exists_{V \setminus U} \bigwedge \Gamma_V$. Therefore a model $\llbracket - \rrbracket$ of \vdash has

$$\mathbb{M}_F(\Gamma)_U = \bigwedge_{\varphi \in \Gamma_U} \llbracket \varphi \rrbracket_U \leq \llbracket \exists_{V \setminus U} \bigwedge \Gamma_V \rrbracket_U = \exists_{F_{U \subseteq V}} (\mathbb{M}_F(\Gamma)_V).$$

Thus $\mathbb{M}_F(\Gamma)$ is no-signalling by Fact 10. \square

Theorem 23. *Let $(\llbracket - \rrbracket, F)$ be an interpretation of \mathcal{L}_C that models a theory \vdash in \mathcal{L} . Given $\Gamma \subseteq \Phi_C$, suppose there is a C -finite and inchworm-saturated $\Delta \subseteq \Phi_C$ such that $\Gamma \subseteq \Delta$ and $\Gamma \vdash_C \varphi$ for all $\varphi \in \Delta$. Then $\mathbb{M}_F(\Delta)$ is the largest no-signalling subpresheaf of $\mathbb{M}_F(\Gamma)$.*

Proof. $\mathbb{M}_F(\Delta)$ is no-signalling by Fact 22. Hence Fact 21 implies $\mathbb{M}_F(\Delta) \models \Gamma \subseteq \Delta$ and hence that $\mathbb{M}_F(\Delta)$ is a subpresheaf of $\mathbb{M}_F(\Gamma)$. Now fix any no-signalling subpresheaf A of $\mathbb{M}_F(\Gamma)$. Then $A \models \Gamma$ by Facts 17 and 21. This implies $A \models \Delta$ by soundness (Theorem 19), since $\Gamma \vdash_C \varphi$ for all $\varphi \in \Delta$. Therefore $A \leq \mathbb{M}_F(\Delta)$ by Fact 21. \square

Lemma 24. *Suppose that a theory \vdash in \mathcal{L} satisfies (13), and that $\llbracket - \rrbracket$ is a conservative model of \vdash , meaning that, for any $\Gamma \subseteq \Phi_C$, $\bigwedge_{\psi \in \Delta} \llbracket \psi \rrbracket_U \leq \llbracket \varphi \rrbracket_U$ for some $\Delta \subseteq \Gamma$ if but also only if $\Gamma \vdash \varphi$. Then $\Gamma \vdash_C \varphi$ iff $A \models \varphi$ for every no-signalling model A of Γ in $\llbracket - \rrbracket$.*

Proof. The “only if” part is soundness (Theorem 19). So, for the “if” part, suppose $A \models \varphi$ for every no-signalling model A of Γ in $\llbracket - \rrbracket$. By (13), Theorem 23 applies and yields $\mathbb{M}_F(\Delta)$, which models $\Gamma \subseteq \Delta$. Hence $\mathbb{F}\mathbb{M}_F(\Delta) \models \varphi$. This means that, for $U \in C$ such that $\varphi \in \Phi_U$, $\bigwedge_{\psi \in \Delta_U} \llbracket \psi \rrbracket_U = \mathbb{F}\mathbb{M}_F(\Delta)_U \leq \llbracket \varphi \rrbracket_U$. Therefore $\Delta_U \vdash \varphi$ by the conservativity of $\llbracket - \rrbracket$ and so $\Delta_U \vdash_C \varphi$. Then, since $\Gamma \vdash_C \psi$ for all $\psi \in \Delta_U$, we have $\Gamma \vdash_C \varphi$. \square

Theorem 25. *Let \vdash be a regular theory satisfying (13). Then, for any $\Gamma \subseteq \Phi_C$, $\Gamma \vdash_C \varphi$ iff $A \models \varphi$ for every no-signalling model A of Γ in every model $\llbracket - \rrbracket$ of \vdash in any regular category.*

Proof. The “only if” part is soundness (Theorem 19). The “if” part is by Lemma 24, since \vdash has a conservative model in some regular category (e.g. [5, Proposition 6.4]). \square

Theorem 28. *Let $\llbracket - \rrbracket$ be a model of a theory \vdash in \mathcal{L} . Then the inchworm logic \vdash_C of \vdash is sound with respect to the filter models in $\llbracket - \rrbracket$: If $\Gamma \vdash_C \varphi$, then $G \models \varphi$ for every filter model G of Γ in $\llbracket - \rrbracket$.*

Proof. Assume the antecedent of (9), that there is $U \in C$ such that $\varphi \in \Phi_U$ and $\Gamma_U \vdash \varphi$. Fix any filter model G of Γ in $(\llbracket - \rrbracket, F)$. Then $\llbracket \psi \rrbracket_U \in G_U$ for all $\psi \in \Gamma_U$, which implies $\bigwedge_{\psi \in \Gamma_U} \llbracket \psi \rrbracket_U \in G_U$ because G_U is a filter. Yet $\Gamma_U \vdash \varphi$ means $\bigwedge_{\psi \in \Gamma_U} \llbracket \psi \rrbracket_U \leq \llbracket \varphi \rrbracket_U$. Therefore $\llbracket \varphi \rrbracket_U \in G_U$ since G_U is closed upward. Thus $G \models \varphi$.

Suppose that $G \models \varphi$ for every filter model G of Γ in $(\llbracket - \rrbracket, F)$, and that $G \models \psi$ for every filter model G of Δ in $(\llbracket - \rrbracket, F)$. Then every filter model G of $\Gamma \cup \Delta$ in $(\llbracket - \rrbracket, F)$ has $G \models \varphi$ and so $G \models \psi$. Therefore, the induction with (9) and (10) shows the theorem. \square

Fact 29. *Let $(\llbracket - \rrbracket, F)$ be a model of a theory \vdash in \mathcal{L} . Given any $\Gamma \subseteq \Phi_C$, the family $\mathbb{F}\mathbb{M}_F(\Gamma) = (\{S \mapsto F_U \mid \llbracket \varphi \rrbracket_U \leq S \text{ for some } \varphi \in \Gamma^*_U\})_{U \in C}$ is a filter model of Γ in $(\llbracket - \rrbracket, F)$. Moreover, for any filter model G of Γ in $(\llbracket - \rrbracket, F)$, $\mathbb{F}\mathbb{M}_F(\Gamma)_U \subseteq G_U$ for each $U \in C$.*

Proof. Fix $U \in C$. $\mathbb{F}\mathbb{M}_F(\Gamma)_U$ is upward closed by definition. $\top \in \mathbb{F}\mathbb{M}_F(\Gamma)_U$ since $\Gamma \vdash_C \top$. Now suppose $S, S' \in \mathbb{F}\mathbb{M}_F(\Gamma)_U$, i.e. $\llbracket \varphi \rrbracket_U \leq S$ and $\llbracket \psi \rrbracket_U \leq S'$ for some $\varphi, \psi \in \Gamma^*_U$. Then $\llbracket \varphi \wedge \psi \rrbracket_U \leq S \wedge S'$, where $\varphi \wedge \psi \in \Gamma^*_U$; so $S \wedge S' \in \mathbb{F}\mathbb{M}_F(\Gamma)_U$. Thus $\mathbb{F}\mathbb{M}_F(\Gamma)_U$ is a filter.

Fixing $U \subseteq V \in C$, we claim $\mathbb{F}\mathbb{M}_F(\Gamma)_U = \{S \mapsto F_U \mid F_{U \subseteq V}^{-1}(S) \in \mathbb{F}\mathbb{M}_F(\Gamma)_V\}$. Suppose $S \in \mathbb{F}\mathbb{M}_F(\Gamma)_U$, i.e. $\llbracket \varphi \rrbracket_U \leq S$ for some $\varphi \in \Gamma^*_U$. Then $\llbracket \varphi \rrbracket_V = F_{U \subseteq V}^{-1} \llbracket \varphi \rrbracket_U \leq F_{U \subseteq V}^{-1}(S)$ for $\varphi \in \Gamma^*_U \subseteq \Gamma^*_V$. Thus $F_{U \subseteq V}^{-1}(S) \in \mathbb{F}\mathbb{M}_F(\Gamma)_V$. On the other hand, suppose $F_{U \subseteq V}^{-1}(S) \in \mathbb{F}\mathbb{M}_F(\Gamma)_V$, i.e. $\llbracket \psi \rrbracket_V \leq F_{U \subseteq V}^{-1}(S)$ for some $\psi \in \Gamma^*_V$. This implies by $\exists_{F_{U \subseteq V}} \dashv F_{U \subseteq V}^{-1}$ that $\llbracket \exists_{V \setminus U} \psi \rrbracket_U \leq S$ for $\exists_{V \setminus U} \psi \in \Gamma^*_U$, i.e. $S \in \mathbb{F}\mathbb{M}_F(\Gamma)_U$. \square

Lemma 30. *Suppose $\llbracket - \rrbracket$ is a conservative model of a theory \vdash in \mathcal{L} . Then $\Gamma \vdash_C \varphi$ iff $G \models \varphi$ for every filter model G of Γ in $\llbracket - \rrbracket$.*

Proof. Since the “only if” part is soundness (Theorem 28), suppose $G \models \varphi$ for every filter model G of Γ in $\llbracket - \rrbracket$. Then $\mathbb{FM}_F(\Gamma) \models \varphi$ in particular. This means that, for $U \in C$ such that $\varphi \in \Phi_U$, $\llbracket \varphi \rrbracket_U \in \mathbb{FM}_F(\Gamma)_U$, i.e. $\llbracket \psi \rrbracket_U \leq \llbracket \varphi \rrbracket_U$ for some $\psi \in \Gamma^*_U$. Therefore $\Gamma \vdash_C \psi$ and $\psi \vdash \varphi$ by the conservativity of $\llbracket - \rrbracket$. Thus $\Gamma \vdash_C \varphi$. \square