

A Simpler Self-reduction Algorithm for Matroid Path-width*

Petr Hliněný

Faculty of Informatics, Masaryk University, Brno, Czech Republic

hlineny@fi.muni.cz

February 6, 2022

Abstract

Path-width of matroids naturally generalizes the better known parameter of path-width for graphs, and is NP-hard by a reduction from the graph case. While the term *matroid path-width* was formally introduced by Geelen–Gerards–Whittle [JCTB 2006] in pure matroid theory, it was soon recognized by Kashyap [SIDMA 2008] that it is the same concept as long-studied so called trellis complexity in coding theory, later named *trellis-width*, and hence it is an interesting notion also from the algorithmic perspective. It follows from a result of Hliněný [JCTB 2006] that the decision problem, whether a given matroid over a finite field has path-width at most t , is fixed-parameter tractable (FPT) in t , but this result does not give any clue about constructing a path-decomposition. The first constructive and rather complicated FPT algorithm for path-width of matroids over a finite field was given by Jeong–Kim–Oum [SODA 2016]. Here we propose a simpler “self-reduction” FPT algorithm for a path-decomposition. Precisely, we design an efficient routine that constructs an optimal path-decomposition of a matroid by calling any subroutine for testing whether the path-width of a matroid is at most t (such as the aforementioned decision algorithm for matroid path-width).

Keywords: matroid; path-width; trellis-width; fixed-parameter tractability

1 Introduction

An ordinary path-decomposition of a graph G , see [16], is a sequence of sets $(X_i \subseteq V(G) : i = 1, \dots, p)$, such that; (i) $\bigcup_{i=1}^p X_i = V(G)$ and for every $1 \leq i < j < k \leq p$, we have $X_j \supseteq X_i \cap X_k$, and (ii) for every $e = uv \in E(G)$ there is $1 \leq i \leq p$ such that $u, v \in X_i$. The width of this decomposition equals $\max_{1 \leq i \leq p} |X_i| - 1$, and the *path-width* of G is the minimum width over all path-decompositions of G . This notion, together with related tree-width, has received great attention in the Graph Minors project of Robertson and Seymour.

*Supported by the project 17-00837S of the Czech Science Foundation.

There is another, more recent view of path-width; the *matroid path-width* defined first by Geelen, Gerards and Whittle [4] in matroid research. We refer to Section 2 for the definition. While the two variants of path-width are indeed tightly related, there is no simple explicit formula between the ordinary path-width and the matroid path-width of the same graph. Matroid path-width of graphs has been recently studied in some papers, e.g. [13]. Our interest in matroid path-width, however, lies beyond the graph case.

A similar notion to path-width has been considered for quite some time also in the area of coding theory, under various names such as the “trellis complexity” of a code, e.g. [17, 9]. In 2008, Kashyap [12] observed that this is the same parameter as the aforementioned path-width [4] of a vector matroid represented by the generator matrix of a linear code. He introduced for it the new name *trellis-width* of a linear code, and proved that computing trellis-width is NP-hard by a reduction from graph path-width. Kashyap also asked, as one of the main open problems in [12], how difficult it is to decide whether the trellis-width of a linear code over a fixed finite field is at most t , and to construct the corresponding optimal decomposition in the YES case, where $t \in \mathbb{N}$ is a fixed parameter.

Concerning the first half of Kashyap’s question, the decision problem is in FPT (*fixed-parameter tractable*) which follows already from the author’s papers [6, 7]. Recall that a parameterized problem is in FPT if it admits an algorithm with runtime of order $\mathcal{O}(f(t) \cdot n^c)$ where t is the parameter, n the input size and c a constant. We briefly sketch two key ideas on which an FPT algorithm for deciding ‘trellis-width $\leq t$ ’ is based (see Section 4 for full details):

- The branch-width of the underlying vector matroid of a linear code is upper-bounded in terms of t , the assumed trellis-width bound. Hence there are only finitely many “minimal obstructions” for the property ‘trellis-width $\leq t$ ’ for each $t \in \mathbb{N}$ and each finite field \mathbb{F} , which follows from [3]. (A similar observation occurs also in Kashyap [12].)
- For bounded branch-width of a vector matroid, over any finite field \mathbb{F} , we can construct an approximate branch-decomposition of it in FPT, see [6]. Then we can, again in FPT, check presence of each one of these finitely many obstructions, see [5, 7].

A careful reader may immediately notice a problem of the suggested scheme—in what way can we get a corresponding trellis- or path-decomposition from it? The sad truth is that in *no way*. To get a corresponding decomposition, a new approach is needed.

Speaking in general, situations in which we get an algorithm which efficiently computes the value of a solution to a certain problem, but not the witnessing solution, are not common in algorithm theory, however, they are also not rare. In such situations, the so-called *self-reduction* routine helps, that is, repeated calls to the algorithms for a solution value (on various inputs derived from the given one) are used to find an admissible solution of the given instance. For a brief example, imagine having an oracle for testing 3-colourability of any graph; how could we then find an actual 3-colouring of a particular graph G using it? The corresponding self-reduction algorithm is quite simple: trying to add new edges to G as long as the oracle certifies that a 3-colouring still exists, the final outcome will be a complete tri-partite graph exhibiting three valid colour classes of G .

Our situation is analogous to that of constructing an optimal matroid branch-decomposition, for which the aforementioned paper [6] provided an approximate construction and an exact decision (the value) in FPT. Building upon that, Oum and the author [8] later designed a self-reduction routine which constructs an optimal branch-decomposition of a matroid over a finite field, by calling the decision subroutine for exact branch-width. It appears very natural to try to extend the self-reduction approach of [8] also for path-decompositions but this, unfortunately, does not easily work. Instead, Jeong, Kim and Oum [10, 11] designed a rather complicated standalone algorithm for the construction of an optimal path-decomposition of a matroid over a finite field, which runs in FPT time for the parameter path-width. In their algorithm, they refer back to the ideas and techniques of Bodlaender and Kloks [1] from graphs.

In this paper we complete the whole picture by providing a new *self-reduction routine* for constructing an optimal matroid path-decomposition, partially inspired by [8]. That is, our routine uses recursive calls to (any) decision subroutine for exact path-width to efficiently construct the output path-decomposition. As the decision subroutine we may use, e.g., the above mentioned FPT algorithm for matroid path-width over a finite field based on [6, 7].

In a nutshell, we contribute the following:

1. A nonuniform FPT algorithm that, for fixed parameters t and $|\mathbb{F}|$, inputs an n -element matroid M represented by a matrix over a finite field \mathbb{F} , and in $\mathcal{O}(n^3)$ time constructs a path-decomposition of M of width $\leq t$ or concludes that the path-width of M is $> t$. This is *not* a better or faster algorithm than in aforementioned Jeong, Kim and Oum [10, 11], but the advantage of our approach is in much simpler design and proof of the algorithm. (*Section 3 – Theorem 3.2 for a generic algorithm, and Theorem 4.3 for improved runtime*)
2. An FPT algorithm that, for a fixed parameter t , a given *oracle function* \mathcal{P} testing if the path-width of a matroid is $\leq t$, and an input n -element abstract matroid M , constructs a path-decomposition of M of width $\leq t$ or concludes that the path-width of M is $> t$. This part is not achieved by [10, 11]. (*Section 4 – Theorem 4.4*)

Regarding (1.), a ‘*nonuniform FPT algorithm*’ means that there is a sequence of algorithms for each values of the parameters $t, |\mathbb{F}|$, rather than one universal algorithm. This weakness is only due to the used decision subroutine for matroid path-width in which we do not know explicitly the finite list of obstructions. If, on the other hand, a different decision algorithm for matroid path-width is found in the future (which may be easier than such a constructive algorithm), or some explicit bound on the path-width obstructions is proved (as in the case of branch-width [2]), then our results immediately give corresponding uniform FPT algorithms.

2 Preliminaries

We refer to the textbook of Oxley [15] for standard matroid material and terminology.

Matroids; rank and connectivity A *matroid* is a pair $M = (E, \mathcal{B})$ where $E = E(M)$ is the ground set of M (elements of M), and $\mathcal{B} \subseteq 2^E$ is a nonempty collection

of *bases* of M , no two of which are in an inclusion. Moreover, matroid bases satisfy the “exchange axiom”: if $B_1, B_2 \in \mathcal{B}$ and $x \in B_1 \setminus B_2$, then there is $y \in B_2 \setminus B_1$ such that $(B_1 \setminus \{x\}) \cup \{y\} \in \mathcal{B}$. We consider only finite matroids.

All matroid bases have the same cardinality called the *rank* $r(M)$ of the matroid. Subsets of bases are called *independent*, and sets that are not independent are *dependent*. Minimal dependent sets are called *circuits*. The *rank function* $r_M(X)$ of M maps subsets of $E(M)$ to non-negative integers; $r_M(X)$ equals the maximum cardinality of an independent subset of a set $X \subseteq E(M)$. The rank function is *submodular*, meaning that $r_M(X) + r_M(Y) \geq r_M(X \cup Y) + r_M(X \cap Y)$ for any $X, Y \subseteq E(M)$, and it fully defines a matroid on its ground set. A matroid M is *uniform* if all subsets of $E(M)$ of size equal to $r(M)$ are bases, and it is also denoted by $U_{r,n}$ where $r = r(M)$ and $n = |E(M)|$.

For $X \subseteq E$, *deletion of X* results in the matroid $M \setminus X$ which is defined by the restriction of the rank function r_M to $E \setminus X$. On the other hand, *contraction of X* results in the matroid M/X which is defined by the rank function $r_{M/X}(Y) := r_M(X \cup Y) - r_M(X)$ for all $Y \subseteq E \setminus X$. Matroids of the form $M/X \setminus Y$ are called *minors* of M .

The *closure of a set $X \subseteq E$ in M* , denoted by $cl_M(X)$, is defined by

$$cl_M(X) := \{e \in E : r_M(X \cup \{e\}) = r_M(X)\}.$$

The closure of X , hence, includes all elements dependent on (or spanned by) X . Sets X such that $X = cl_M(X)$ are closed, or *flats*.

We, moreover, define the (symmetric and submodular) *connectivity function* of M by

$$\lambda_M(X) := r_M(X) + r_M(E \setminus X) - r(M)$$

for all subsets $X \subseteq E$. Any bipartition (X, Y) of E (where $Y = E \setminus X$) is called a *separation in M* of connectivity value $\lambda_M(X) = \lambda_M(Y)$, or shortly a *k -separation*, if $\lambda_M(X) = k - 1$ and both $|X|, |Y| \geq k$. Informally, λ_M measures how much the two sides of a separation “share together” in terms of rank. A matroid is *connected* if and only if it has no 1-separation. It is well-known that in a connected matroid, every two elements belong to a common circuit (this is analogous to graph 2-connectivity).

We also define the following extension of the connectivity function which will be useful in our context

$$\mu_M(X, A) := r_M(X \cup A) + r_M((E \setminus X) \cup A) - r(M).$$

For example, $\mu_M(X, A) = \lambda_M(X)$ if and only if $A \subseteq cl_M(X) \cap cl_M(E \setminus X)$ or, in other words, if A is spanned by both X and $E \setminus X$. If $e \in cl_M(X) \cap cl_M(E \setminus X)$, then we say that e is *in the guts of the bipartition $(X, E \setminus X)$* . As another example we mention that, if $\mu_M(X, A) = r_M(A)$ then every element in the guts of $(X, E \setminus X)$ belongs to the closure of A .

Matroid path-width [4] Let M be an n -element matroid. Any permutation $Y = (e_1, e_2, \dots, e_n)$ of the elements $E(M)$ is called a *path-decomposition* of M . The *width* of (e_1, e_2, \dots, e_n) is defined

$$w_M(Y) = w_M(e_1, e_2, \dots, e_n) := \max_{i=1, \dots, n} \lambda_M(\{e_1, \dots, e_i\}),$$

and the *path-width* $\text{pw}(M)$ of M is the least width over all path-decompositions of M , i.e.

$$\text{pw}(M) := \min_{\text{permut. } \pi \in S_n} w_M(e_{\pi(1)}, e_{\pi(2)}, \dots, e_{\pi(n)}).$$

We say, for any $1 \leq i < n$, that the bipartition $(\{e_1, \dots, e_i\}, \{e_{i+1}, \dots, e_n\})$ is *displayed* by the path-decomposition Y , and we refer to $(\{e_1, \dots, e_i\}, \{e_{i+1}, \dots, e_n\})$ as to the *bipartition at position i* .

The notion of matroid path-width is related to the better known parameter of branch-width. A tree T is *cubic* if its vertex degrees are 3 or 1. A *branch-decomposition* of a matroid M is a pair (T, τ) where T is a cubic tree and $\tau : E(M) \rightarrow \ell(T)$ is a bijection of the elements of M to the leaves of T . Every edge $e \in E(T)$ partitions the leaves of T into two sets L_1, L_2 , and we say the bipartition $(\tau^{-1}(L_1), \tau^{-1}(L_2))$ is *displayed by* (T, τ) . We define the width of e as $\lambda_M(\tau^{-1}(L_1)) + 1$ and the width of (T, τ) as the maximum of widths over all edges of T . The *branch-width* $\text{bw}(M)$ of M is the minimum width over all branch-decompositions of M .

A cubic tree is a caterpillar if it is obtained by connecting leaves to a path. *Linear branch-width* of a matroid M is defined as ordinary branch-width with a restriction that the cubic tree T must be a caterpillar. One can easily observe that this notion coincides with that of matroid path-width (except the artificial ‘+1’ term above); the path-width of M is always one less than its linear branch-width. Consequently, we have:

Lemma 2.1. *For any matroid M , we have $\text{bw}(M) \leq \text{pw}(M) + 1$.* □

Assorted matroid claims We list some elementary and intuitive technical claims about matroids which will be used in the proof of our algorithm.

Lemma 2.2. *Let M be a matroid and $C_1, C_2 \subseteq E(M)$ be two circuits of M such that $|C_1 \cap C_2| = 1$ and $r_M(C_1) + r_M(C_2) = r_M(C_1 \cup C_2) + 1$. Then $C_1 \Delta C_2$ (the symmetric difference) is also a circuit of M .*

Proof. Let $C_1 \cap C_2 = \{f\}$. By the standard circuit exchange axiom there exists a circuit of M contained in the set $C_3 := (C_1 \cup C_2) \setminus \{f\} = C_1 \Delta C_2$. Consider any $e \in C_3$ where, up to symmetry, $e \in C_2 \setminus C_1$. We have $|C_3 \setminus \{e\}| = |C_1| - 1 + |C_2| - 1 - 1 = r_M(C_1) + r_M(C_2) - 1 = r_M(C_1 \cup C_2)$. At the same time, since C_1, C_2 are circuits and $f \in C_1 \cap C_2$, we have $r_M(C_1 \cup C_2) = r_M(C_1 \cup (C_2 \setminus \{e\})) = r_M((C_1 \setminus \{f\}) \cup (C_2 \setminus \{e, f\})) = r_M(C_3 \setminus \{e\})$ and so $C_3 \setminus \{e\}$ is independent. Therefore, C_3 itself is the circuit. □

Lemma 2.3. *Let M be a matroid and $X \subseteq E = E(M)$. If $e, f \in E$ such that $\mu_M(X, \{e\}) = \mu_M(X, \{f\}) = \mu_M(X, \{e, f\}) = \lambda_M(X) + 1$, then either $e, f \in X$ or $e, f \notin X$.*

Proof. Let $Y = E \setminus X$. Assume the contrary, i.e. up to symmetry, $e \in X$ and $f \in Y$. From $r_M(X) + r_M(Y) - r(M) + 1 = \lambda_M(X) + 1 = \mu_M(X, \{f\}) = r_M(X \cup \{f\}) + r_M(Y) - r(M)$ we immediately get $r_M(X \cup \{f\}) = r_M(X) + 1$ and, by symmetry, $r_M(Y \cup \{e\}) = r_M(Y) + 1$. This leads to

$$\begin{aligned} \lambda_M(X) + 1 &= \mu_M(X, \{e, f\}) = r_M(X \cup \{f\}) + r_M(Y \cup \{e\}) - r(M) \\ &= r_M(X) + r_M(Y) + 2 - r(M) = \lambda_M(X) + 2, \end{aligned}$$

a contradiction. □

Lemma 2.4. *Let M be a matroid and N a minor of M . Then $\text{pw}(N) \leq \text{pw}(M)$.*

Proof. Consider $X \subseteq E(M)$ and $e \in E(M)$. It is well-known that $\lambda_{M \setminus e}(X \setminus \{e\}) \leq \lambda_M(X)$ and $\lambda_{M/e}(X \setminus \{e\}) \leq \lambda_M(X)$. Hence, by induction on $|E(M)| - |E(N)|$, the restriction of any path-decomposition of M is a path-decomposition of N of at most the same width. \square

Lemma 2.5. *Let M be an n -element matroid and (e_1, \dots, e_n) be a path-decomposition of M of width $t = w_M(e_1, \dots, e_n)$. For an index i let $X = \{e_1, \dots, e_i\}$ and $Y = \{e_{i+1}, \dots, e_n\} = E(M) \setminus X$ such that $\lambda_M(X) = t$. Assume that there exists a circuit $C \subseteq E(M)$ such that no element of C is in the guts of (X, Y) and $\mu_M(X, C) = r_M(C)$. Then $X \cap C \neq \emptyset \neq Y \cap C$.*

Proof. We proceed by means of contradiction, aiming to show that $w_M(e_1, \dots, e_n) > t$. Up to symmetry, let $Y \cap C = \emptyset$, meaning that $C \subseteq X$. Let $j \leq i$ be the largest index such that $e_j \in C$, and $X' = \{e_1, \dots, e_{j-1}\}$, $Y' = \{e_j, \dots, e_n\}$. From the assumptions $\lambda_M(X) = r_M(X) + r_M(Y) - r(M) = t$ and $\mu_M(X, C) = r_M(X) + r_M(Y \cup C) - r(M) = r_M(C)$, we derive

$$r_M(C) + r_M(Y) - r_M(Y \cup C) = t. \quad (1)$$

We have $r_M(C \setminus \{e_j\}) = r_M(C)$ since C is a circuit, and $r_M(Y \cup \{e_j\}) = r_M(Y) + 1$ since e_j is not in the guts of (X, Y) . Hence we can rewrite (1) as

$$r_M(C \setminus \{e_j\}) + r_M(Y \cup \{e_j\}) - r_M(Y \cup C) = t + 1. \quad (2)$$

Note that $Y \cup \{e_j\} \subseteq Y'$ and $C \setminus \{e_j\} \subseteq X'$. We conclude the proof by showing

$$\begin{aligned} \lambda_M(X') &= r_M(X') + r_M(Y') - r(M) \\ &= r_M(X') + (r_M(Y') - r_M(X' \cup Y')) \\ &\geq r_M(X') + (r_M(Y \cup \{e_j\}) - r_M(X' \cup Y \cup \{e_j\})) \\ &= r_M(Y \cup \{e_j\}) + (r_M(X') - r_M(X' \cup Y \cup \{e_j\})) \\ &\geq r_M(Y \cup \{e_j\}) + (r_M(C \setminus \{e_j\}) - r_M((C \setminus \{e_j\}) \cup Y \cup \{e_j\})) \\ &= r_M(C \setminus \{e_j\}) + r_M(Y \cup \{e_j\}) - r_M(C \cup Y) = t + 1, \end{aligned}$$

using submodularity and (2). This however contradicts $w_M(e_1, \dots, e_n) = t$. \square

Lemma 2.6. *Let M be a matroid and $Y \subseteq E(M)$, $Y' = E(M) \setminus Y$. Assume that $Q \subseteq E(M)$ is such that all elements of Q are in the guts of (Y, Y') and $r_M(Q) = \lambda_M(Y)$. If C is a circuit of M and $e \in C \setminus Y$, then there exists a circuit C' of M such that $e \in C' \subseteq (C \setminus Y) \cup Q$ and $C' \subseteq \text{cl}_M(Y')$.*

Proof. If $C \subseteq \text{cl}_M(Y')$, we are done. Otherwise, both the sets $C \cap Y, C \cap Y'$ are nonempty and independent, and $r_M(Q) > 0$. Let $Q_1 \subseteq Q$ be a basis of Q , and M' be the restriction of M onto $Q_1 \cup C$. We aim to show that the set $D := (C \setminus Y) \cup Q_1$ is dependent. On the contrary, assume that D is independent and choose $D_1 \subseteq C \setminus D$ such that $D \cup D_1$ is a basis of M' . Let $M_1 := M'/D_1$ be obtained by contracting D_1 (hence M_1 has a basis D). Then $C \setminus D_1$ is a circuit of M_1 and $\emptyset \neq (C \setminus D_1) \cap Y \subseteq \text{cl}_{M_1}(Q_1)$. Consequently, $[(C \setminus D_1) \setminus ((C \setminus D_1) \cap Y)] \cup Q_1 = (C \setminus Y) \cup Q_1 = D$ is dependent in M_1 , and so it is in M since $r_{M_1}(D) = r_M(D)$.

Hence, dependent D contains a circuit C' of M , and since Q_1 is independent, we may choose C' such that $e \in C'$. Finally, $C' \subseteq (C \setminus Y) \cup Q$ and $C' \subseteq \text{cl}_M(Y')$ are true by the definition of D . \square

Matroid representation and extensions A standard example of a matroid is given by a set of vectors (forming the columns of a matrix \mathbf{A}) with usual linear independence. The matrix \mathbf{A} is then called a *(vector) representation* of the matroid. We will consider only representations \mathbf{A} over finite fields. Since non-zero scaling of vectors does not change linear dependencies, vector representations can also be seen as point configurations in the projective space over \mathbb{F} , which will be the view followed throughout this paper. (Note that parallel vectors are represented by the same points.)

We now briefly illustrate the “geometric” meaning of matroid terms.

- The matroid closure of a set X corresponds to the affine closure or *span* $\langle X \rangle$ of the points representing X (note that considering the points of X in a projective space, $\langle X \rangle$ does *not* contain the origin $\mathbf{0}$). The rank of X is the dimension or rank of the span of X .
- For a bipartition (X, Y) of M , the *guts* of (X, Y) consists exactly of the points in the intersection of the spans of X and Y , that is $\langle X \rangle \cap \langle Y \rangle$, and $\lambda_M(X)$ is the rank of this guts. The value of $\mu_M(X, A)$ equals the rank of the space spanned by $(\langle X \rangle \cap \langle Y \rangle) \cup A$.
- All the previous entities can be straightforwardly computed by means of standard linear algebra over the matrix \mathbf{A} .

There is one particular operation we need to discuss in close detail. For a matroid M we say that a matroid M_1 is a *free extension of M by element e* if $e \in E(M_1)$ and $M = M_1 \setminus e$, $r(M_1) = r(M)$, and for every $X \subseteq E(M)$ we have $r_{M_1}(X \cup \{e\}) = r_M(X) + 1$ unless $r_M(X) = r(M)$. This is equivalent to claiming that every circuit of M_1 containing e has full rank $r(M_1)$. Informally saying, e is added to M without any unforced dependency – geometrically, in a general position. We will also say that e is *freely placed* in M (see also (M2) in Section 4). We will use the following:

Lemma 2.7. *Let M be a matroid of rank r represented by a matrix \mathbf{A} over a finite field \mathbb{F} . Let α be a root of an irreducible polynomial of degree r in \mathbb{F} , and denote by $\mathbf{b} = (1, \alpha, \dots, \alpha^{r-1})^T$. Let $\mathbb{F}(\alpha)$ be the extension field of \mathbb{F} obtained by adjoining α to \mathbb{F} . Then the matrix $[\mathbf{A} | \mathbf{b}]$ over $\mathbb{F}(\alpha)$ represents a free extension of M by an element b .*

Proof. Assume the contrary, that \mathbf{b} is a linear combination over $\mathbb{F}(\alpha)$ of the columns of a column-submatrix $\mathbf{A}' \subseteq \mathbf{A}$ of rank less than r . Since \mathbf{A}' has r rows denoted by $\mathbf{a}'_1, \mathbf{a}'_2, \dots, \mathbf{a}'_r$, they are linearly dependent as vectors, and so for some $\lambda_1, \dots, \lambda_r \in \mathbb{F}$ (not all 0) it holds $\lambda_1 \mathbf{a}'_1 + \lambda_2 \mathbf{a}'_2 + \dots + \lambda_r \mathbf{a}'_r = \mathbf{0}$. However, since \mathbf{b} is a linear combination of the columns of \mathbf{A}' , we have also $\lambda_1 + \lambda_2 \alpha + \dots + \lambda_r \alpha^{r-1} = 0$. This contradicts the assumption that α is a root of an irreducible polynomial of degree r over \mathbb{F} . \square

The next two lemmas cover some simple properties of path-decompositions of represented matroids.

Lemma 2.8. *Let \mathbb{F} be a finite field, $|\mathbb{F}| \geq 3$, and $t \geq 2$ be an integer. Denote by P the point set of some rank- t projective space Σ over \mathbb{F} . Then, for any permutation (p_1, \dots, p_k) of P there exists i such that $\langle p_1, \dots, p_i \rangle = \Sigma = \langle p_{i+1}, \dots, p_k \rangle$. In other words, $w_M(p_1, \dots, p_k) = t$ where M is the matroid represented by P .*

Proof. We have $|P| = k = \frac{q^t-1}{q-1}$ points where $q = |\mathbb{F}|$ ([15]), and every proper subspace of Σ has at most $k' = \frac{q^{t-1}-1}{q-1}$ points. Since, by simple calculus, $k' < \lfloor k/2 \rfloor$ when $q \geq 3$, we are done by choosing $i = \lfloor k/2 \rfloor$. \square

Lemma 2.9. *For $i = 1, 2$, let M_i be a matroid represented over a finite field \mathbb{F} , and Y_i be a path decomposition of M_i of width at most t . Assume there exist prefixes Z_i of Y_i , $i = 1, 2$, such that $\langle Z_1 \rangle \cap \langle E(M_1) \setminus Z_1 \rangle = \langle E(M_1) \rangle \cap \langle E(M_2) \rangle \subseteq \langle E(M_2) \setminus Z_2 \rangle$ and $r_{M_2}(E(M_2) \setminus Z_2) \leq t$. Then the matroid M' represented by $(E(M_1) \setminus Z_1) \cup E(M_2)$ has path-width at most t .*

Proof. We form a path-decomposition Y of M' by appending $Y_1 \setminus (Z_1 \cup E(M_2))$ after Y_2 . Let the considered subspaces (of the projective space over \mathbb{F}) be $\Pi := \langle E(M_1) \rangle \cap \langle E(M_2) \rangle$ and $\Sigma := \langle E(M_2) \setminus Z_2 \rangle \supseteq \Pi$. Let (X, X') be a bipartition of M' displayed by Y . If $X \subseteq Z_2 \subseteq E(M_2)$, then $\langle X \rangle \cap \langle E(M_1) \rangle \subseteq \Pi$, and since $\langle E(M') \setminus X \rangle \supseteq \Sigma \supseteq \Pi$, we have $\langle X \rangle \cap \langle E(M') \setminus X \rangle = \langle X \rangle \cap \langle E(M_2) \setminus X \rangle$ which is of rank $\leq t$ by the assumption $w_{M_2}(Y_2) \leq t$.

If $Z_2 \subseteq X \subseteq Y_2$, then $\langle X \rangle \cap \langle E(M') \setminus X \rangle \subseteq \langle \Sigma \cup \Pi \rangle = \Sigma$ is easily of rank $\leq t$. In the remaining case of $Y_2 \subseteq X$ we get, similarly as in the first case, $\langle X \rangle \cap \langle E(M') \setminus X \rangle \subseteq \langle \Pi \cup (X \setminus E(M_2)) \rangle \cap \langle E(M') \setminus X \rangle \subseteq \langle Z_1 \cup (X \cap E(M_1)) \rangle \cap \langle E(M_1) \setminus (X \cup Z_1) \rangle$ which is of rank $\leq t$ by the assumption $w_{M_1}(Y_1) \leq t$. \square

3 Self-reduction Algorithm

In this section we give our core result—a self-reduction routine that, for a fixed parameter t , constructs an optimal path-decomposition of a given represented matroid of path-width t , using an oracle which can decide whether the path-width of a given matroid is at most t . We stress that our routine can work with *any* oracle (subroutine) for deciding the path-width value, and that it is not restricted to only representable matroids as we will see in the next Section 4.

Motivation For easier understanding of the problem we are dealing with, we start this section with a brief overview of the algorithm for constructing an optimal branch-decomposition of a given (represented) matroid of branch-width t from [8]; it is based on the following decision step:

- [8] Assume $X \subseteq E = E(M)$ is such that $\lambda_M(X) \leq t$ and that $M[X]$ (the restriction of M to X) has branch-width $\leq t$. The task is to decide whether M has a branch-decomposition of width t such that “ X forms one branch” of the decomposition.

The way this decision task is implemented in [8] is based on extending $M \setminus X$ with a bounded number of elements so that every optimal branch-decomposition of it displays a separation whose guts is geometrically identical with that of $(X, E \setminus X)$ of M (then a branch formed by X can be simply added to this place). Besides implementing this key decision task, the rest of the algorithm of [8] is an easy recursive composition routine (merging branches until the whole tree is constructed). On a very high level, our new algorithm will do the same thing tailored to path-width – see next. Though, the underlying details will be very different and more complicated due to the fact that one *cannot* “add a branch” to a path-decomposition as to a branch-decomposition.

Algorithm outline We give a high-level description of our new path-decomposition algorithm. We now treat a given matroid M represented over a finite field \mathbb{F} as a point configuration in a projective geometry over \mathbb{F} (recall Section 2): Let M be the input matroid and $E = E(M)$, $n = |E|$, where the points of E are given as vectors over \mathbb{F} . For a simplification of the arguments, we assume that $|\mathbb{F}| \geq 3$, that is, if M is given with a representation over $GF(2)$ then we equivalently view it over $\mathbb{F} = GF(4)$.

Assume that $\text{pw}(M) = t$.

- (I) For $i = 1, 2, \dots, n$, suppose that we have got a sequence $X = (e_1, \dots, e_{i-1}) \in E^{i-1}$ such that there exists a path-decomposition of M of width t which starts with the prefix X (note that initially $X = \emptyset$ and our assumption is trivial).
- (II) For each $f \in E \setminus X$, we set $X_f = (e_1, \dots, e_{i-1}, f)$. If $\lambda_M(X_f) \leq t$, we test whether there exists a path-decomposition of M of width t which starts with the prefix X_f .
- (III) If the test of (II) succeeds for some (any) f —which has to happen for at least one value by the assumption—we let $X := X_f$ and continue with (I).

Clearly, this scheme results in the construction of a path-decomposition (e_1, \dots, e_n) of M of width t . Hence it remains to explain implementation of crucial Step (II).

For convenience, we refer by X_f also to the underlying set of the sequence X_f from the above outline. Unlike in the easier case of [8], it is now not sufficient to test $M \setminus X_f$ for path-width $\leq t$ under the condition that the guts of the bipartition $(X_f, E \setminus X_f)$ is geometrically identical to the guts of some bipartition displayed by the corresponding optimal path-decomposition. We actually need that the corresponding optimal path-decomposition of $M \setminus X_f$ can be “prefixed” with this guts without increasing the width (which could be impossible if the displaying bipartition is somewhere in the middle of the decomposition). This goal we achieve by adding to $M \setminus X_f$ a special set D of points of rank $t + 1$ and path-width t (in fact, D is represented over an extension field of \mathbb{F}). Denoting by M' the new matroid on $(E(M) \setminus X_f) \cup D$, it is then easy to see that the path-width of M' is $\leq t$ if the answer to (II) is YES (see Lemma 2.9 with $M_1 = M$, $Z_1 = X_f$ and $M_2 \sim D$). Proving the converse of this claim constitutes the core of the proof below.

The formal details are given below, in Algorithm 3.1 and its proof.

Algorithm 3.1. Let \mathbb{F} be a fixed finite field and $t \in \mathbb{N}$ a fixed parameter. Let \mathcal{P} be an oracle which, given any matroid N represented over \mathbb{F} , correctly decides whether $\text{pw}(N) \leq t$. Let M be an input connected n -element matroid of rank r , given as an $r \times n$ matrix \mathbf{A} over \mathbb{F} , and assume $\text{pw}(M) = t$.

1. We pad \mathbf{A} with 0’s to make an $(r + t + 1) \times n$ matrix (informally, adding “extra dimensions” useful in the computation). For simplicity, we will refer to the columns of the matrix as to the elements of M , with understanding that all computations will be carried out by means of linear algebra (i.e., in the matrix) in a natural way.
2. Let initially $X := \emptyset$. For $i = 1, 2, \dots, n$, we repeat the following instructions:
 - (a) We have got $X = (e_1, \dots, e_{i-1}) \in E(M)^{i-1}$ where the elements of the sequence are distinct, and we use the symbol X to refer both to the sequence and the underlying set of elements of M .

- (b) We choose $f \in E(M) \setminus X$ such that $\lambda_M(X \cup \{f\}) \leq t$, and set $X_f := (e_1, \dots, e_{i-1}, f)$.
- (c) We compute the guts $\Gamma := \langle X_f \rangle \cap \langle E(M) \setminus X_f \rangle$ and choose a subspace $\Sigma \supseteq \Gamma$ of rank exactly t and an element $d_0 \notin \Sigma$, such that $\langle \Sigma \cup \{d_0\} \rangle \cap \langle E(M) \rangle = \Gamma$. (Note that the rank of Γ , by (2c), may be smaller than t , and we use some of the “extra dimensions” from Step (1) for placing d_0 and $\Sigma \supseteq \Gamma$ of rank exactly t .) Let P denote the set of all points of Σ in the finite projective geometry over \mathbb{F} . Specially, for $t = 1$, we form P by two parallel points.
- (d) Let N_0 denote the matroid of rank $t + 1$ induced by the points of $P \cup \{d_0\}$, and $\mathbb{F}_0 = \mathbb{F}$. For $j = 1, 2, \dots, t$, let N_j be the matroid constructed as a free extension of N_{j-1} by an element d_j . By Lemma 2.7, N_j is represented over the extension field \mathbb{F}_j obtained from \mathbb{F}_{j-1} by adjoining a root of degree $r(N_0) = t + 1$. At the end, let $D_0 := \{d_0, d_1, \dots, d_t\}$, $D := P \cup D_0$ and $\mathbb{F}' = \mathbb{F}_t$.
- (e) For the matroid M' induced on the point set $(E(M) \setminus X_f) \cup D$ in the projective geometry over \mathbb{F}' , we ask the oracle \mathcal{P} whether $\text{pw}(M') \leq t$.
 - If the answer is NO, then we repeat the Steps from (2b) for another choice of f .
 - If the answer is YES, then we update $X := X_f$ and continue the cycle in Step (2) with the next value of i until $i = n$.

3. We output the path-decomposition $X = (e_1, \dots, e_n)$ of M of width t .

Note that, in Step (2e), some element e of M may be in the guts of $(X_f, E \setminus X_f)$ and then e is represented by the same point as some element of P in M' . It actually does not matter whether we consider these two elements as identical or a parallel pair.

Theorem 3.2. *Let \mathbb{F} , t and \mathcal{P} be as in Algorithm 3.1. For any connected n -element input matroid M represented by a matrix over \mathbb{F} , such that $\text{pw}(M) = t$, Algorithm 3.1 correctly outputs a path-decomposition of M of width t . With fixed parameters \mathbb{F} and t , the algorithm computes in FPT time $\mathcal{O}(n^4)$ and, in addition, makes $\mathcal{O}(n^2)$ calls to the oracle \mathcal{P} .*

Proof. We start with justifying correctness of the algorithm. Thanks to the condition $\lambda_M(X \cup \{f\}) \leq t$ in Step (2b) of Algorithm 3.1, we know that the (eventual) output of the algorithm must be a path-decomposition of M of width t . Consequently, it is enough to prove that for every iteration of Step (2) there is a choice of $f \in E(M) \setminus X$ which correctly succeeds in the test of Step (2e). Assuming, for this moment, the following

Claim 3.3. *in Step (2e), $\text{pw}(M') \leq t$ if and only if there exists a path-decomposition of M of width t which starts with the prefix X_f ,*

the rest of the proof follows by a straightforward induction on i .

It is hence enough to prove Claim 3.3. In one direction (\Leftarrow), assume that there exists a path-decomposition $Y = (e_1, \dots, e_n)$ of M of width t which starts with the prefix X_f . We give a path-decomposition $Y' = (e'_1, e'_2, \dots)$ of the matroid N_t induced by the point set D , where $e'_1 = d_0, e'_2 = d_1, \dots, e'_{t+1} = d_t$ and this is followed by the elements of P in any order. The bipartition at position $j + 1$ in Y' , for $j < t$, has the guts $\langle \{d_0, \dots, d_j\} \rangle$ of

rank $\leq t$. At positions $j + 1$ for $j \geq t$, on the other hand, the guts is always Σ of rank t (or its subspace). Therefore, we can set $M_1 = M$, $Z_1 = X_f$ and $M_2 = N_t$, $Z_2 = D_0$ and apply Lemma 2.9, to conclude that $\text{pw}(M') \leq t$.

In the opposite direction (\Rightarrow) of Claim 3.3, we assume that $\text{pw}(M') \leq t$. Recall the set P of the points of Σ over \mathbb{F} from Step (2c), and the matroid M' on the point set $(E(M) \setminus X_f) \cup P \cup D_0$ from Step (2e). Let $Y' = (e'_1, \dots, e'_p)$ be an optimal path-decomposition of M' where $p = |E(M')|$. We first aim to show that there exists an index $1 \leq j \leq p$ such that the guts at the position j in Y' contains Σ (and so it equals Σ and $\text{pw}(M') = t$). If $t = 1$, then P consists of two parallel points (parallel to single-point Σ) and we simply choose a position between those points. For $t > 1$ this conclusion follows from Lemma 2.8 applied to the restriction of Y' onto P .

Let $Y'_j = \{e'_1, \dots, e'_j\}$ where $\lambda_{M'}(Y'_j) = t$ by the previous paragraph. Recall the point set $D_0 = \{d_0, \dots, d_t\}$ from Step (2d). We first claim that, up to possible reversal of the sequence Y' , we have $D_0 \subseteq Y'_j$. This easily follows from the conclusion of Lemma 2.3 since, for any $0 \leq a < b \leq t$, we have $\langle \Sigma \cup \{d_a\} \rangle = \langle \Sigma \cup \{d_b\} \rangle = \langle \Sigma \cup D_0 \rangle$ of rank $t + 1$, and so the condition of the lemma $\mu_{M'}(Y'_j, \{d_a\}) = \mu_{M'}(Y'_j, \{d_b\}) = \mu_{M'}(Y'_j, \{d_a, d_b\}) = t + 1$ holds true.

Second, we claim that $(E(M) \setminus X_f) \cap Y'_j \subseteq \langle P \rangle = \Sigma$. Suppose the contrary, that $Z := ((E(M) \setminus X_f) \cap Y'_j) \setminus \Sigma \neq \emptyset$ (here we view $E(M) \setminus X_f$ as points in a projective space). Note that $Z \cap \Sigma = \emptyset$. We first consider the subcase that $\langle Z \rangle \cap \Sigma \neq \emptyset$. Informally, we are going to argue that the spans of Z and D_0 “freely overlap” in Σ and so, for any $g \in Z \cup D_0$, the span of $(Z \cup D_0) \setminus \{g\}$ still contains $\langle D_0 \rangle$. Then the path-decomposition Y' , at some position before j , must contain $\langle D_0 \rangle$ in the guts, but this is impossible since the rank of D_0 is $t + 1$. The corresponding formal argument follows.

Let $M'' = M' \setminus D_0$. We choose $Z_0 \subseteq Z$ minimal by inclusion such that $\langle Z_0 \rangle \cap \Sigma \neq \emptyset$, and so the rank of $\langle Z_0 \rangle \cap \Sigma$ is one (in matroid terms this reads $r_{M''}(Z_0) + r_{M''}(P) = r_{M''}(Z_0 \cup P) + 1$). Since P contains all the points of Σ in the projective geometry over \mathbb{F} (in matroid terms, P is a modular flat in M'' which is represented over \mathbb{F}), we have that $\langle Z_0 \rangle \cap P \neq \emptyset$, and by minimality of Z_0 we have $\langle Z_0 \rangle \cap P = \{p_0\}$. Consequently, $C_0 = Z_0 \cup \{p_0\}$ is a circuit in M'' and so also in M' . Now we look at the set $C_1 := D_0 \cup \{p_0\}$ in M' which is of rank $t + 1$ and cardinality $t + 2$, and hence is dependent. Since d_1, \dots, d_t have been chosen as free extensions in Step (2d), there cannot be any smaller circuits in C_1 and so C_1 itself is a circuit. We apply Lemma 2.2 to C_0 and C_1 , obtaining a circuit $C_2 := C_0 \Delta C_1$ of M' , where no element of C_2 belongs to Σ (our guts at position j). Since $C_2 \supseteq D_0$, the span of C_2 contains Σ and so $\mu_{M'}(Y'_j, C_2) = r_{M'}(C_2)$ and the conditions of Lemma 2.5 are fulfilled for C_2 . However, the conclusion of the lemma contradicts our assumption $C_2 \subseteq Y'_j$.

Next, still under the assumption $Z \neq \emptyset$, we consider the subcase that $\langle Z \rangle \cap \Sigma = \emptyset$. Recall that $\Gamma = \langle X_f \rangle \cap \langle E(M) \setminus X_f \rangle = \langle \Sigma \cup D_0 \rangle \cap \langle E(M) \rangle$. Let $Z' := (E(M) \setminus X_f) \setminus Y'_j$ and note that $E(M) \setminus X_f \supseteq Z \cup Z' \supseteq (E(M) \setminus X_f) \setminus \Sigma = (E(M) \setminus X_f) \setminus \Gamma$. Hence, $\langle X_f \rangle \cap \langle \Gamma \cup Z \cup Z' \rangle = \Gamma$ and so $\langle Z \rangle \cap \langle X_f \cup \Gamma \cup Z' \rangle = \langle Z \rangle \cap \langle \Gamma \cup Z' \rangle$. Since M is connected, we in particular have $\emptyset \neq \langle Z \rangle \cap \langle \Gamma \cup X_f \cup Z' \rangle = \langle Z \rangle \cap \langle \Gamma \cup Z' \rangle$. The latter in turn means, again from the path-decomposition Y' of M' with the guts $\Sigma \supseteq \Gamma$ at position j , that $\langle Z \rangle \cap \Sigma \neq \emptyset$ – the case already being considered above.

To recapitulate, the assumed path-decomposition Y' of M' has the (geometric) guts

$\Sigma = \langle Y'_j \rangle \cap \langle E(M') \setminus Y'_j \rangle$ at position j . We have also shown that $Y'_j \cap (E(M) \setminus X_f) \subseteq \Sigma$. Hence, if we form Y'' by restricting Y' to the elements of $E(M) \setminus X_f$, the concatenated sequence (X_f, Y'') will be a path-decomposition of M of width t . The proof of Claim 3.3 is finished.

The last point is to address runtime complexity of Algorithm 3.1. Note that the finite field \mathbb{F} and the value of t are fixed parameters. In particular, arithmetic operations over \mathbb{F} and \mathbb{F}' (which depends only on \mathbb{F} and t) take constant time each. Also note that $r \leq n$. We n times iterate at Step (2), and each iteration costs the following. We are choosing at most n values of f in Step (2b), and for each we compute the subspace Γ . Knowing $\langle X \rangle \cap \langle E(M) \setminus X \rangle$ already from the previous level, the computation of $\Gamma = \langle X_f \rangle \cap \langle E(M) \setminus X_f \rangle$ takes $\mathcal{O}(n^2)$ in Step 2d by standard linear algebra (the rank of Γ is at most constant t). The rank of $\Sigma \supseteq \Gamma$ and cardinality of the set P are constants depending on \mathbb{F} and t . Step (2d) takes $\mathcal{O}(1)$ time since it depends only on \mathbb{F} and t and not on the input M . In fact, the point set D_0 needs to be computed only once during the whole algorithm and then linearly transformed to match actual Σ . This amounts to $\mathcal{O}(n^4)$ total time and $\mathcal{O}(n^2)$ calls to the oracle \mathcal{P} in Step (2e). \square

4 Algorithmic Consequences

So far, in Section 3 we have restricted attention to connected matroids, but this is not any problem since we may easily concatenate path-decompositions of connected components of a general matroid. To make use of Algorithm 3.1, we also need to provide an implementation of the oracle \mathcal{P} (which tests the value of path-width $\leq t$, as sketched in the Introduction). This will be done by Theorem 4.1. A class \mathcal{N} of matroids is *minor-closed* if, for every matroid $M \in \mathcal{N}$, all minors of M also belong to \mathcal{N} . A matroid $M \notin \mathcal{N}$ is an *obstruction for membership in \mathcal{N}* if all proper minors of M belong to \mathcal{N} .

Theorem 4.1 (Geelen–Gerards–Whittle [3],¹ and Hliněný [7]). *Let \mathbb{F} be a fixed finite field and $k \in \mathbb{N}$ a fixed parameter. For any minor-closed class \mathcal{N} of matroids, there are finitely many obstructions for membership in \mathcal{N} which are representable over \mathbb{F} and have branch-width at most k . Consequently, there is an FPT algorithm which, given an n -element matroid M represented by a matrix over \mathbb{F} , in time $\mathcal{O}(n^3)$ correctly decides whether $M \in \mathcal{N}$ or outputs that the branch-width of M is more than k .*

Direct implementation The way we use Theorem 4.1 in an implementation of the oracle \mathcal{P} combines Lemma 2.1 with Lemma 2.4; the matroids of path-width at most t have branch-width at most $t + 1$ and form a minor-closed class \mathcal{P}_t for which we can test membership in FPT time $\mathcal{O}(n^3)$. Note, though, that this approach results in a nonuniform FPT algorithm since we do not explicitly know the finite lists of obstructions for the classes \mathcal{P}_t , $t \in \mathbb{N}$. In combination with Theorem 3.2 we immediately get:

Corollary 4.2. *Let \mathbb{F} be a fixed finite field and $t \in \mathbb{N}$ a fixed parameter. There is a nonuniform FPT algorithm parameterized by t and $|\mathbb{F}|$ which, given an n -element matroid*

¹We remark that Geelen, Gerards and Whittle have announced a “matroid minors” theorem which does not require a bound on branch-width to claim finite number of \mathbb{F} -representable obstructions for \mathcal{N} , but that is not fully published yet. For our purpose, the version of [3] is sufficient.

M represented by a matrix over \mathbb{F} , in time $\mathcal{O}(n^3)$ decides whether $\text{pw}(M) \leq t$. Consequently, if $\text{pw}(M) \leq t$, there is a nonuniform FPT algorithm parameterized by t and $|\mathbb{F}|$, which in time $\mathcal{O}(n^5)$ outputs a path-decomposition of M of width t . \square

We remark that, in the setting of nonuniform algorithms, Algorithm 3.1 as used in Corollary 4.2 can be further simplified by the following observation. The point configuration D constructed in Step (2d) is unique, up to a linear transformation, for given parameters t, \mathbb{F} , and hence it can be hard-coded into the (anyway nonuniform) algorithm with the smallest possible extension field \mathbb{F}' which can represent D (this would quite likely be a much smaller field than the one computed by brute force in Step (2d)).

Improving runtime Runtime dependence on n of the algorithm of Corollary 4.2 can be improved to $\mathcal{O}(n^3)$ by using the same implementation tricks as in [8], based on earlier [6].

Theorem 4.3. *Let \mathbb{F} be a fixed finite field and $t \in \mathbb{N}$ a fixed parameter. There is a nonuniform FPT algorithm parameterized by t and $|\mathbb{F}|$ which, given an n -element matroid M represented by a matrix over \mathbb{F} , in time $\mathcal{O}(n^3)$ outputs a path-decomposition of M of width t or certifies that $\text{pw}(M) > t$.*

Proof sketch. In the improved algorithm, we follow the general scheme of [8, Section 6] but in a simplified way. This is possible thanks to the fact that Algorithm 3.1, at each iteration, works with only one “active guts” of a bipartition $(X, E \setminus X)$, unlike the algorithm of [8] which builds many branches of the desired branch-decomposition concurrently.

We modify the main steps of Algorithm 3.1 as follows:

1. For the input matroid M represented by the matrix \mathbf{A} , we use [6] to compute a branch-decomposition of M of width at most $3t + 3$ —actually, a so-called *3t-boundaried parse tree* \mathcal{T} for M —or to confirm that $\text{bw}(M) > t + 1$ and so $\text{pw}(M) \geq \text{bw}(M) - 1 > t$. This step takes $\mathcal{O}(n^3)$ time for fixed t, \mathbb{F} .
2. Each task performed in Steps 2c and 2d can be done in time $\mathcal{O}(n)$ within the parse tree \mathcal{T} (we refer to [8, Section 6] for corresponding details). It is important to compute within \mathcal{T} (and not on whole \mathbf{A}), for which purpose we each time “enlarge” every node of \mathcal{T} by the constant-rank subspace Σ . Subsequently, Step 2e can test $\text{pw}(M') \leq t$ by checking (non-)presence of the finitely many obstructions for ‘path-width $\leq t$ ’. This test can also be done in time $\mathcal{O}(n)$ by [7] since minor obstructions are MSO-definable.

Altogether, runtime is $\mathcal{O}(n^3 + n^2 \cdot n) = \mathcal{O}(n^3)$ for fixed t, \mathbb{F} . \square

Abstract matroids Besides its simplicity, our Algorithm 3.1 has another theoretical advantage over the constructive algorithm of [11]. While the authors of [11] directly compute with points and subspaces in a finite projective geometry, and it does not seem possible to extend their approach to infinite projective geometries or abstract matroids, we can easily adapt our algorithm to work even with *abstract matroids* given by a rank oracle (although our algorithm also directly worked with the points of a subspace Σ , that was only for convenience and clarity, and could be rather easily replaced by an abstract handling).

In this respect we mention the algorithm of Nagamochi [14] which computes an optimal path-decomposition for an arbitrary submodular function (and hence including the case of a matroid given by a rank oracle). Though, its runtime is of order $\mathcal{O}(n^{f(t)})$ where t is the path-width (complexity class XP) while we aim for an FPT algorithm.

We say that an abstract matroid M is given by a *rank oracle* if the input consists of the ground set $E = E(M)$ and an oracle function $\mathcal{R} : 2^E \rightarrow \mathbb{N}$ such that $\mathcal{R}(X) = r_M(X)$ for all $X \subseteq E$. Algorithms then handle M by asking \mathcal{R} so called *rank queries*. In this setting we have got the following algorithm.

Theorem 4.4. *Let $t \in \mathbb{N}$ and \mathcal{P} be an oracle function which, for any matroid N given by a rank oracle, correctly decides whether $\text{pw}(N) \leq t$. There is an algorithm that, for an input n -element matroid M given by a rank oracle \mathcal{R} , outputs a path-decomposition of M of width t or correctly answers that $\text{pw}(M) > t$. The algorithm makes $\mathcal{O}(n^2)$ calls to the oracle function \mathcal{P} and, neglecting the fixed parameter t , asks $\mathcal{O}(n^2)$ rank queries.*

Before moving onto the proof, we need one more technical concept. We are going to modify the matroid M (which we do not completely know—we cannot read all the ranks of sets in M in polynomial time!). Instead, we will modify the rank oracle \mathcal{R} by prescribing its (efficient) answers to rank queries involving elements which we add to M . In this respect we define the following three elementary operations:

(M1) *Adding a coloop a to M* defines, for every $X \subseteq E(M)$, that $\mathcal{R}(X \cup \{a\}) := \mathcal{R}(X) + 1$.

(M2) *Placing b freely into the closure of $Z \subseteq E(M)$* defines, for every $X \subseteq E(M)$,

- $\mathcal{R}(X \cup \{b\}) := \mathcal{R}(X)$ if $r_M(Z \cup X) = r_M(X)$, and
- $\mathcal{R}(X \cup \{b\}) := \mathcal{R}(X) + 1$ otherwise.

(M3) *Placing c freely into the guts of (the bipartition of) $Z \subseteq E(M)$* means, for $X \subseteq E(M)$,

- $\mathcal{R}(X \cup \{c\}) := \mathcal{R}(X)$ if $\mu_M(Z, X) = r_M(X)$, and
- $\mathcal{R}(X \cup \{c\}) := \mathcal{R}(X) + 1$ otherwise.

An informal geometric explanation of these operations follows. (M1) simply “adds another dimension” with a . (M2) puts the new point b in general position (i.e., without unforced linear dependencies) into the span $\langle Z \rangle$. (M3) similarly puts the new point c in general position into the guts $\langle Z \rangle \cap \langle E(M) \setminus Z \rangle$. It is a routine exercise to prove that the rank oracle defined by each one of (M1), (M2), (M3) is the rank function of a matroid.

Proof of Theorem 4.4. Again, we may restrict our attention to connected input matroids M . We modify some steps of Algorithm 3.1 as follows:

- Step (1) is not needed.
- In step (2c), let $k = \lambda_M(X_f)$. First, we $(t - k)$ -times (if $k < t$) repeat the operation (M1) of adding a coloop. Let P_0 denote the set of coloops added to M this way. We then $(t + k)$ -times repeat the operation (M3) of placing a new element freely into the guts of $(X_f \cup P_0, (E(M) \setminus X_f) \cup P_0)$ —to be formally precise, we consider for this operation the elements of P_0 duplicated. Let $P \supseteq P_0$ denote the set of all

the $2t$ added elements, which is of rank t (one may observe that P actually induces a uniform matroid $U_{t,2t}$).

- In Step (2d), we add a new coloop d_0 by (M1). Then, for $j = 1, \dots, t$, we iteratively do the operation (M2) of freely placing a new element d_j into the closure of $P \cup \{d_0\}$. Again, let $D_0 := \{d_0, d_1, \dots, d_t\}$ and $D := P \cup D_0$.
- In Step (2e), we let M' be the matroid defined on the ground set $(E(M) \setminus X_f) \cup D$ by the rank oracle \mathcal{R}' constructed from \mathcal{R} by the above modifications.

In the proof of the modified algorithm, we can essentially repeat the setup and most of the arguments of the proof of Theorem 3.2, translated into the abstract setting of the rank functions of M and M' . Such as, geometric span of points representing M' is translated as the closure operation in M' and, in particular, the subspace $\Sigma = \langle P \rangle$ is now written as $cl_{M'}(P)$. Though, the following two steps in the proof of the forward direction of Claim 3.3 need separate formal arguments:

- Assuming a path-decomposition $Y' = (e'_1, \dots, e'_p)$ of M' of width t , we, instead of invoking Lemma 2.8, argue simply as follows: We define index j as the minimum $1 \leq j \leq t$ such that $|P \cap Y'_j| = t$. Since the elements of P have been each freely placed into a rank- t flat, the t -element set $P \cap Y'_j$ is independent, and so is the complement $P \setminus Y'_j$. Consequently, all elements of P belong to the guts of the bipartition at the position j of Y' , a situation analogous to the former proof.
- Second, we differently argue that $(E(M) \setminus X_f) \cap Y'_j \subseteq cl_{M'}(P)$. Assuming $Z := ((E(M) \setminus X_f) \cap Y'_j) \setminus cl_{M'}(P) \neq \emptyset$, we again aim to find a circuit $C_2 \subseteq Y'_j$ contradicting the conclusion of Lemma 2.5. A full proof of the existence of such C_2 is left for coming Lemma 4.5 (in which $X = X_f$ and $C_2 = D_0 \cup Z_0$).

Assuming now Lemma 4.5, the proof is finished. \square

Lemma 4.5. *Let M be a connected matroid, $X \subseteq E = E(M)$ and $\emptyset \neq Z \subseteq E \setminus X$. Assume that $P \subseteq E \setminus X$ is such that $r_M(P) = t \geq \lambda_M(X)$, $\mu_M(X, P) = t$, and $Z \cap cl_M(P) = \emptyset$. Furthermore, assume that M_0 is a matroid on the ground set $E \cup D_0$ where $D_0 = \{d_0, d_1, \dots, d_t\}$, the restriction of M_0 to E is M , and d_0 is a coloop w.r.t. M and each d_i is freely placed (M2) in the closure of $P \cup \{d_0\}$ w.r.t. $E \cup \{d_0, \dots, d_{i-1}\}$ for $i = 1, \dots, t$. Let M' be M_0 restricted to $E(M_0) \setminus X$. If $\lambda_{M'}(D_0 \cup Z) \leq t$, then there exists $Z_0 \subseteq Z$ such that $D_0 \cup Z_0$ is a circuit of M' .*

Proof. Let $E' = E \setminus Z$. Observe that $|D_0| = r_{M'}(D_0) = t+1$ (since D_0 is independent both in M_0 and M'), $\lambda_{M'}(D_0) = t$ and $cl_{M'}(D_0) \supseteq P$, but $D_0 \cap cl_{M'}(P) = \emptyset = D_0 \cap cl_{M'}(E')$. From the assumptions $\lambda_{M'}(D_0 \cup Z) \leq t$ and $Z \cap P = \emptyset$ we get that actually $\lambda_{M'}(D_0 \cup Z) = t = r_{M'}(P)$ and all elements of P are in the guts of $(D_0 \cup Z, E' \setminus Z)$ in M' .

Let $e_1 \in X$ and $e_2 \in Z$ be arbitrary. Since M is connected, there exists a circuit $C \subseteq M$, $C \ni e_1, e_2$. We apply Lemma 2.6 to M , C and $Y := X$, $Q := P$, $e := e_2$. The obtained circuit C' satisfies: $e_2 \in C' \subseteq (C \setminus X) \cup P \subseteq E(M')$ and $C' \subseteq cl_{M'}(E')$.

In the matroid M' , we let $Y := cl_{M'}(E' \setminus Z) \supseteq P$. From previous $\lambda_{M'}(D_0 \cup Z) = r_{M'}(P)$ where P is in the guts, we have $Z \cap Y \subseteq cl_{M'}(P)$; and since $Z \cap cl_{M'}(P) = \emptyset$ by the assumptions, we then get $Z \cap Y = \emptyset$. In the matroid $M'' = M'/d_0$ obtained by contracting

d_0 , we have $D_1 = \{d_1, \dots, d_t\} \subseteq cl_{M''}(P)$ and $cl_{M''}(D_1) = cl_{M''}(P)$, since $cl_{M'}(D_0) \supseteq P$ and $r_{M''}(D_1) = t = r_{M''}(P)$. Note that C' is a circuit of M'' , too, since M'' restricted to E' equals $M \setminus X$. Denoting $Y' = E(M'') \setminus Y$, we have $e_2 \in Z \cap C' \subseteq Y' \cap C'$. In this setup, we apply Lemma 2.6 to M'' , Y and $C := C'$, $Q := D_1$, $e := e_2$. The circuit C'' that we obtain, satisfies $C'' \subseteq (C' \setminus Y) \cup D_1$, and so $C'' \subseteq Z_0 \cup D_1$ where $Z_0 = Z \cap C''$ (since $Y \supseteq E' \setminus Z$).

Back in the matroid M' (uncontracting d_0), $C'' \cup \{d_0\}$ is a circuit of M' , and $C'' \supseteq D_1$ since the elements of D_1 have been freely placed—they do not have unforced dependencies in M' . Hence this circuit is $D_0 \cup Z_0 = C'' \cup \{d_0\}$. \square

5 Conclusions

We have shown a relatively simple oracle algorithm which can construct an optimal path-decomposition of a given matroid if it is provided with a subroutine testing the value of matroid path-width. This completes the picture of width decompositions of (\mathbb{F} -represented) matroids in the following sense: While for the matroid branch-width, a non-constructive FPT decision algorithm has been known since [6], followed by a natural self-reduction constructive algorithm in [8], no such FPT self-reduction approach to constructing an optimal matroid path-decomposition seemed possible along similar lines before.

Specifically for matroids represented over a finite field \mathbb{F} , this result provides an alternative to the recent algorithm of Jeong, Kim and Oum [10, 11] which uses a direct and complicated construction based on ideas originally developed for graphs by Bodlaender and Kloks [1]. Though, there is price we have to pay for simplicity of our algorithm; our approach provides a nonuniform FPT algorithm, caused by the fact that we have yet no explicit bound on the size of the minor-minimal obstructions for path-width $\leq t$ (unlike the case of branch-width in which an explicit bound [2] readily provides a uniform FPT algorithm [8]).

Moreover, our self-reduction oracle algorithm readily generalizes to abstract matroids given by rank oracles, as proved in Theorem 4.4. Although we are currently not aware of an FPT algorithm which could test path-width $\leq t$ for matroids given by rank oracles, such algorithms could probably emerge in the future (cf. [14]) for other matroid classes, and then Theorem 4.4 will be readily applicable also to these new classes. Along the same line, it is likely that in the future an explicit bound on the obstructions for path-width $\leq t$ will be found and then Algorithm 3.1 will immediately turn uniform.

Acknowledgments

We would like to thank the anonymous referees for careful reading and checking all the proofs, and for many suggestions which helped to improve this paper a lot.

References

- [1] H. L. BODLAENDER AND T. KLOKS, *Efficient and constructive algorithms for the path-width and treewidth of graphs*, J. Algorithms, 21 (1996), pp. 358–402.

- [2] J. GEELLEN, A. GERARDS, N. ROBERTSON, AND G. WHITTLE, *On the excluded minors for the matroids of branch-width k* , J. Comb. Theory, Ser. B, 88 (2003), pp. 261–265, [https://doi.org/10.1016/S0095-8956\(02\)00046-1](https://doi.org/10.1016/S0095-8956(02)00046-1).
- [3] J. GEELLEN, A. GERARDS, AND G. WHITTLE, *Branch-width and well-quasi-ordering in matroids and graphs*, J. Comb. Theory, Ser. B, 84 (2002), pp. 270–290, <https://doi.org/10.1006/jctb.2001.2082>.
- [4] J. GEELLEN, B. GERARDS, AND G. WHITTLE, *On Rota’s conjecture and excluded minors containing large projective geometries*, J. Comb. Theory, Ser. B, 96 (2006), pp. 405–425, <https://doi.org/10.1016/j.jctb.2005.09.005>.
- [5] P. HLINĚNÝ, *On matroid properties definable in the MSO logic*, in Mathematical Foundations of Computer Science, MFCS 2003, Proceedings, vol. 2747 of Lecture Notes in Computer Science, Springer, 2003, pp. 470–479, https://doi.org/10.1007/978-3-540-45138-9_41.
- [6] P. HLINĚNÝ, *A parametrized algorithm for matroid branch-width*, SIAM J. Comput., 35 (2005), pp. 259–277, <https://doi.org/10.1137/S0097539702418589>.
- [7] P. HLINĚNÝ, *Branch-width, parse trees, and monadic second-order logic for matroids*, J. Combin. Theory Ser. B, 96 (2006), pp. 325–351, <https://doi.org/10.1016/j.jctb.2005.08.005>.
- [8] P. HLINĚNÝ AND S. OUM, *Finding branch-decomposition and rank-decomposition*, SIAM J. Comput., 38 (2008), pp. 1012–1032, <https://doi.org/10.1137/070685920>.
- [9] K. JAIN, I. MANDOIU, AND V. VAZIRANI, *The ‘Art of Trellis Decoding’ is computationally hard – for large fields*, IEEE Trans. Information Theory, 44 (1998), pp. 1211–1214, <https://doi.org/10.1109/18.669287>.
- [10] J. JEONG, E. KIM, AND S. OUM, *Constructive algorithm for path-width of matroids*, in Proceedings of the Twenty-Seventh Annual ACM-SIAM Symposium on Discrete Algorithms, SODA 2016, SIAM, 2016, pp. 1695–1704, <https://doi.org/10.1137/1.9781611974331.ch116>.
- [11] J. JEONG, E. J. KIM, AND S. OUM, *The “art of trellis decoding” is fixed-parameter tractable*, IEEE Trans. Information Theory, 63 (2017), pp. 7178–7205, <https://doi.org/10.1109/TIT.2017.2740283>.
- [12] N. KASHYAP, *Matroid pathwidth and code trellis complexity*, SIAM J. Discrete Math., 22 (2008), pp. 256–272, <https://doi.org/10.1137/070691152>.
- [13] A. KOUTSONAS, D. THILIKOS, AND K. YAMAZAKI, *Outerplanar obstructions for matroid pathwidth*, Discrete Mathematics, 315–316 (2014), pp. 95–101, <https://doi.org/10.1016/j.disc.2013.10.007>.
- [14] H. NAGAMOCHI, *Linear layouts in submodular systems*, in Algorithms and Computation - 23rd International Symposium, ISAAC 2012, Proceedings, vol. 7676 of Lecture Notes in Computer Science, Springer, 2012, pp. 475–484, https://doi.org/10.1007/978-3-642-35261-4_50.
- [15] J. G. OXLEY, *Matroid Theory (Oxford Graduate Texts in Mathematics)*, Oxford University Press, Inc., New York, NY, USA, 2006.

- [16] N. ROBERTSON AND P. SEYMOUR, *Graph minors. I. Excluding a forest*, J. Comb. Theory, Ser. B, 35 (1983), pp. 39–61, [https://doi.org/10.1016/0095-8956\(83\)90079-5](https://doi.org/10.1016/0095-8956(83)90079-5).
- [17] A. VARDY, *Trellis structure of codes*, in Handbook of coding theory. Vol. 1. Part 1: Algebraic coding. Vol. 2. Part 2: Connections, Part 3: Applications, Amsterdam: Elsevier, 1998, pp. 1989–2117.