arXiv:1606.04160v2 [cs.LG] 7 Mar 2017

The Crossover Process: Learnability and Data Protection from Inference Attacks

Richard Nock
Data61, The Australian National University & The University of Sydney
richard.nock@data61.csiro.au

Giorgio Patrini
Data61 & The Australian National University
giorgio.patrini@anu.edu.au

Finnian Lattimore
The Australian National University & Data61
finnian.lattimore@nicta.com.au

Tiberio Caetano
Ambiata & The University of Sydney
tiberio.caetano@gmail.com

Abstract

It is usual to consider data protection and learnability as conflicting objectives. This is not always the case: we show how to jointly control inference — seen as the attack — *and* learnability by a noise-free process that mixes training examples, the Crossover Process (CP). One key point is that the CP is typically able to alter joint distributions *without* touching on marginals, nor altering the sufficient statistic for the class. In other words, it saves (and sometimes improves) generalization for supervised learning, but can alter the relationship between covariates — and therefore fool measures of nonlinear independence and causal inference into misleading *ad-hoc* conclusions. For example, a CP can increase / decrease odds ratios, bring fairness or break fairness, tamper with disparate impact, strengthen, weaken or reverse causal directions, change observed statistical measures of dependence. For each of these, we quantify changes brought by a CP, as well as its statistical impact on generalization abilities via a new complexity measure that we call the Rademacher CP complexity. Experiments on a dozen readily available domains validate the theory.

Keywords: Supervised learning, Privacy, Fairness, Statistical inference, Causality.

1 Introduction

We study the problem of sensitive data sharing under two conflicting objective: protection of the data from unwanted inference and guarantees that supervised learning can be effective after the data protection mechanism has been applied. The two goals are inherently in tension, yet not necessarily in contradiction. Any credible solution to the combined issues would have a considerable impact on open data policies for research and commercial enterprises.

We motivate our goal by an example. A medical laboratory aims to release to the research community a newly collected dataset about its genetic, behavioural, habits and infection history of the patients affected by cancer, with the intent of letting other institutions to test their predictive models on it so as to improve diagnosis methodologies. Even assuming that we perfectly anonymize it, this data is still extremely sensitive. Anyone possessing it could directly make statistical queries and causal inference between specific patient traits, or combine it with his/her own private data using powerful causal inference techniques [Pearl and Bareinboim, 2014].

Ideally, researchers should be able to release the data and hide discriminatory and sensitive relationships, such as smoking tendency by ethnicity and gender-prone infections [Rizvi and Haritsa, 2002], *while* making sure that the utility of the dataset for predicting the sickness state remains unaltered. Can we design a procedure that would transform the data in a form apt to publication, that is, erasing any trace of statistical or causal relationship between particular pairs of attributes *and*, at the same time, leaving prediction performance virtually untouched? Several streams of research may share similarities with this open question thus we start by covering the current background.

Privacy is a growing concern in the public sphere [Barocas and Nissenbaum, 2014a, Enserink and Chin, 2015, Graham et al., 2015, Machanavajjhala and Kifer, 2015] (and references therein). Two leading mechanisms for the private release of data are differential privacy and k-anonymity [Dwork and Roth, 2014, Enserink and Chin, 2015, Sweeney, 2002]. They guarantee *individual level* protection, *i.e.* identifiability. We depart from this view on the problem because we are concerned with inference at global level over the present data, for which for example differential privacy does not provide any sort of guarantee.

In fact, as pointed out in Barocas and Nissenbaum [2014a], "even when individuals are not *identifiable* they may still be *reachable* [...] and subject to *consequential inferences* and predictions taken on that basis". The reference is to the possibility of performing *inference attacks* by a malicious agent willing to uncover *causal relationships*, or even just measure *statistical independence*, between sensitive covariates. Even when true causality is sometimes considered "a research field in its infancy" [Graham et al., 2015], it is hard to exaggerate the recent burst in causal inference techniques [Cornia and Mooij, 2014, Doran et al., 2014, Gretton et al., 2005, 2007, Hoyer et al., 2008, Janzing et al., 2012, Li et al., 2016, Mooij et al., 2016], as well as the threats this may pose on privacy [Barocas and Nissenbaum, 2014a,b, Enserink and Chin, 2015, Kusner et al., 2016, Machanavajjhala and Kifer, 2015].

In simple terms with a quite general example, the attacker estimates from the data/outputs some $\Pr[U|\mathcal{V}]$, where U is typically a sensitive attribute and $\mathcal{V}$ is built from one (or a set of) protected attribute(s) [Hardt et al., 2016]. Protection against such attacks involve in general controlling similar estimates or odds ratios. Since the advent of differential privacy [Dwork and Roth, 2014], these questions have received a steadily increasing treatment, with a further surge over the past two years over fairness considerations [Feldman et al., 2015, Hardt et al., 2016, Kleinberg et al., 2016, Machanavajjhala and Kifer, 2015].

When it comes to supervised learning, there is often a single sensitive attribute U to protect, in general a score or prediction [Feldman et al., 2015, Hardt et al., 2016, Kleinberg et al., 2016]. This is quite restrictive for our purpose if we consider that the total number of observation variables is blowing up in hundreds, thousands or more in mainstream datasets. This is not to say that previous techniques do not or cannot apply, but there could be at least a serious combinatorial overload to treating a lot of sensitive attributes with techniques fit for one. Finally, data protection is not the sole constraint — otherwise, communicating noise would just solve the problem. Guaranteed protection has to come with provable utility, *i.e.* learnability [Dwork and Roth, 2014, Hardt et al., 2016]. In the context of differential privacy, the trade-off does not play favourably for learning [Duchi et al., 2013].

In the design a solution to the problem, we keep in mind an additional requirement. At the age where protection is shifting towards statistical information — in constrast with computation, *e.g.* for public key encryption —, a good protection mechanism that targets specific utility is one that, knowing all the public part of the protection mechanism[1], gives the *least* information about any other sensible content. This is not trivial to satisfy. In fact, knowing for example that a dataset was protected with a specific technique *for fairness* (say, odds ratios = 1, Hardt et al. [2016]) leaks information: if the attacker sees that some attributes that are important for him do not display fairness in data, then they were probably not treated and he can use this data for his own analyses [Pearl and Bareinboim, 2014]. If, however, a "suspicious" amount of fairness is detected, then the attributes were probably treated, and if those attributes relate smoking and cancer, then it is not hard to imagine the most likely imbalance in original odds that justified protection (smoking causes cancer). In sum, we should seek a data protection mechanism able to bring, *or break*, fairness, and thereby be able to just *fool* inference into misleading *ad-hoc* conclusions.

In short, we carry out protection at the *upstream level*, *i.e.* the training sample's[2] and we target two goals: the dataset's utility for the black-box supervised prediction task remains *within control*[3], but it is surgically altered against fined-grained specific inference attacks among description features. Alteration can work in *all directions*: increasing / decreasing odds, being fair or breaking fairness, tampering with disparate impact, strengthening, weakening or reversing causal directions, changing observed statistical measures of dependence, and so on.

We show that this task is within reach **and it involves the same protection process for all**. It can also be very surgical — for example, marginals can remain untouched and therefore may not raise suspicions about protection. Coping with the desired level of protection to statistical independence and causal inference attacks may require wrangling the complete data, but this may be done with a tight *explicit* control of its utility for supervised learning, and, as we show, it may even yield *better* models for prediction. Although counterintuitive, this last fact should not come with great surprise considering the success of sophisticated noisification methods, *e.g.* dropout [Srivastava et al., 2014], to enhance learning.

[1] This is Kerckhoff's principle, Machanavajjhala and Kifer [2015].

[2] Neither the algorithm [Chaudhuri et al., 2011], nor its output Hardt et al. [2016] but on the input data as in *local privacy* [Duchi et al., 2013].

[3] Like [Chaudhuri et al., 2011], we investigate the *generalization* abilities impacted by data protection, and do not remain within the realm of the empirical risk.

Our **main contribution** is the introduction of the *Crossover Process*, CP. An analogy may be done with the biological crossover: a population of DNA strands gets mixed with a crossover, but there is a single zone for chiasma (*i.e.* contact point) for the whole population. In the same way as DNA strands exchange genetic material during recombination, feature values get mixed between observations during a CP , although in a more general way than in genetic recombination. The key to learning and generalization is that the CP may be done without changing the sufficient statistic for the class [Patrini et al., 2014, 2016], nor touching class-based marginals. The key to interfering with measures of (un)fairness, independence and causal calculus is that the CP is able to surgically alter joint distributions. Our contribution is therefore twofolds: (i) we introduce the CP and show how it drives the generalisation abilities of linear and some non-linear classifiers by the introduction of a new statistical complexity measure, the Rademacher CP complexity (RCP). We show that the RCP can be very significantly smaller than the standard empirical Rademacher complexity, thereby being a lightweight player — and a tractable knob — for generalisation. Then, (ii) on the data protection standpoint, we show

(a) how the components of a CP may be chosen to alter odds ratios and measures of (un)fairness, equality of opportunity, equalized odds or disparate impact [Feldman et al., 2015, Hardt et al., 2016, Kleinberg et al., 2016],

(b) how it can be built to alter the powerful Hilbert-Schmidt independence criterion [Gretton et al., 2007],

(c) how it may be devised to blow-up causal estimation errors [Cornia and Mooij, 2014], and finally

(d) how it can interfere with identifiable causal queries on a causal graph in the *do* calculus framework [Pearl, 2000, Shpitser and Pearl, 2006].

Targeting all these different models of dependence *exhaustively* would require far more than the paper's current size and technical content. Yet, all of them are important and forgetting one would reduce *de facto* the scope of the CP from the protection standpoint. This is why we deliberately chose to make a very specific treatment of some, in particular for [Cornia and Mooij, 2014].
Organisation of the paper — Section §2 gives general definitions. §3 presents the Crossover Process, §4 relates the CP to measures of (un)fairness and §5 presents its relationships with learnability. §6 shows the impact of the CP on measures of independence and §7 shows the impact of the CP on causal queries. A last Section discusses and concludes. An Appendix provides all proofs, additional results and some extensive experiments performed to assess the theory. A movie[4], presented in Subsection 10.2, shows the effects of the CP on a popular domain for causal discovery [Hoyer et al., 2008]. Figure 1 provides a high-level overview of the papers topics for the main technical Sections.

2 General notations and definitions

Learning setting — We let $[m] \doteq \{1, 2, ..., m\}$ and $\Sigma_m \doteq \{\boldsymbol{\sigma} \in \{-1, 1\}^m\}$. $\mathcal{X} \subseteq \mathbb{R}^d$ is a domain of observations. Examples are couples (observation, label) $\in \mathcal{X} \times \Sigma_1$, sampled i.i.d. according to

[4]Available anonymously at `https://youtu.be/4d5Z23cwEyY`




Figure 1: Organisation and dependencies of the main Sections of the paper according to learning and data protection. Sections are independent within each rectangle.

some unknown but fixed distribution $\mathcal{D}$. We denote $\mathcal{F} \doteq [d]$ the set of observation attributes (or features). $\mathcal{S} \doteq \{(\boldsymbol{x}_i, y_i), i \in [m]\} \sim \mathcal{D}_m$ is a training sample of $|\mathcal{S}| = m$ examples. For any vector $\boldsymbol{z} \in \mathbb{R}^d$, z_j denotes its coordinate j. Finally, notation $x \sim X$ for X a set denotes uniform sampling in X, and the mean operator is $\boldsymbol{\mu}_{\mathcal{S}} \doteq \mathbb{E}_{(\boldsymbol{x},y)\sim\mathcal{S}}[y \cdot \boldsymbol{x}]$ [Patrini et al., 2014, 2016].

In supervised learning, the task is to learn a classifier $\mathcal{H} \ni h : \mathcal{X} \to \mathbb{R}$ from $\mathcal{S}$ with good generalisation properties, that is, having a small *true risk* $\mathbb{E}_{(\boldsymbol{x},y)\sim\mathcal{D}}[L_{0/1}(y, h(\boldsymbol{x}))]$, with $L_{0/1}(z, z') \doteq 1_{zz'\leq 0}$ the 0/1 loss ($1_{\cdot}$ is the indicator variable). In general, this is achieved by minimising over $\mathcal{S}$ a *φ-risk* $\mathbb{E}_{(\boldsymbol{x},y)\sim\mathcal{S}}[\varphi(yh(\boldsymbol{x}))] = (1/m) \cdot \sum_i \varphi(y_i h(\boldsymbol{x}_i))$, where $\varphi(z) \geq 1_{z\leq 0}$ is a *surrogate* of the 0/1 loss. In this paper, φ is any differentiable proper symmetric (PS) loss [Nock and Nielsen, 2009, Patrini et al., 2014] (symmetric meaning that there is no class-dependent misclassification cost). The logistic, square and Matsushita losses are examples of PS losses. Set $\mathcal{H}$ is a predefined set of classifiers, such as linear separators, decision trees, etc. .

Matrix quantities — The set of unnormalised column stochastic matrices, $\mathcal{M}_n \subset \mathbb{R}^{n\times n}$, is the superset of column stochastic matrices for which we drop the non-negativity constraint, thus keeping the sole constraint of unit per-*column* sums. We let $S_n \subset \mathcal{M}_n$ denote the symmetric group of order n. For any $\mathrm{A}, \mathrm{B} \in \mathbb{R}^{n\times n}$ and $\mathrm{M} \in \mathcal{M}_n$, we let

$$\langle \mathrm{A}, \mathrm{B} \rangle_{\mathrm{M}} \doteq \mathrm{tr}\left((\mathrm{I}_n - \mathrm{M})^\top \mathrm{A} (\mathrm{I}_n - \mathrm{M}) \mathrm{B}\right)$$

denote the *centered inner product* of A and B with respect to M. It is a generalisation of the centered inner product used in kernel statistical tests of independence [Gretton et al., 2005], for which $\mathrm{M} = (1/n)\mathbf{1}\mathbf{1}^\top$.

Without loss of generality, we shall assume that indices in $\mathcal{S}$ cover first the positive class: $(y_i = +1 \wedge y_{i'} = -1) \Rightarrow i < i'$. A key subset of matrices of $\mathbb{R}^{m\times m}$ consists of block matrices whose coordinates on indices corresponding to different classes in $\mathcal{S}$ are zero: block-class matrices.

Definition 1 $\mathrm{A} \in \mathbb{R}^{m\times m}$ *is a* ***block-class matrix*** *iff* $(y_i \cdot y_{i'} = -1) \Rightarrow \mathrm{A}_{ii'} = 0, \forall i, i'$.

An asterisk exponent in a subset of matrices indicates the intersection of the set with block class matrices, such as for $\mathcal{M}_n^* \subset \mathcal{M}_n$ and $S_n^* \subset S_n$. Finally, matrix entries are noted with double indices like $\mathrm{M}_{ii'}$; replacing an index by a dot, ".", indicates a sum over the index, like $\mathrm{M}_{i\cdot} \doteq \sum_{i'} \mathrm{M}_{ii'}$.

3 The Crossover Process

The Crossover process (CP) transforms $\mathcal{S}$ in two steps: the split and the shuffle step. In the split step, a bi-partition of the features set $\mathcal{F}$ is computed: $\mathcal{F} = \mathcal{F}_{\rm a} \cup \mathcal{F}_{\rm s}$. $\mathcal{F}_{\rm a}$ is the *a*nchor set and $\mathcal{F}_{\rm s}$ is the *s*huffle set. To perform the shuffle step, we need additional notations. Without loss of generality, we assume $\mathcal{F}_{\rm a} \doteq [d_{\rm a}]$ and $\mathcal{F}_{\rm s} \doteq \{d_{\rm a} + j, j \in [d_{\rm s}]\}$, $d_{\rm a} > 0, d_{\rm s} > 0, d_{\rm a} + d_{\rm s} = d$. So, $\mathcal{F}_{\rm a}$ contains the first $d_{\rm a}$ features and $\mathcal{F}_{\rm s}$ contains the last $d_{\rm s}$ features. Let I_d be the identity matrix, and $[\mathrm{F}^{\rm a}|\mathrm{F}^{\rm s}] = \mathrm{I}_d$ a vertical block partition where $\mathrm{F}^{\rm a} \in \mathbb{R}^{d \times d_{\rm a}}$ ($\mathrm{F}^{\rm s} \in \mathbb{R}^{d \times d_{\rm s}}$) has columns representing the features of $\mathcal{F}_{\rm a}$ ($\mathcal{F}_{\rm s}$) — we use notation $[.]$ both for integer sets and block matrices without ambiguity. Finally, we define the (*row*-wise) observation matrix $\mathrm{S} \in \mathbb{R}^{m \times d}$ with $(\mathrm{S})_{ij} \doteq x_{ij}$. Let $\mathbf{1}_i$ be the i^{th} canonical basis vector.

Definition 2 *For any block partition $[\mathrm{F}^{\rm a}|\mathrm{F}^{\rm s}] = \mathrm{I}_d$ and any **shuffle matrix** $\mathrm{M} \in \mathcal{M}_n$, the Crossover process $\mathscr{T} \doteq \textsc{cp}(\mathcal{S}; \mathrm{F}^{\rm a}, \mathrm{F}^{\rm s}, \mathrm{M})$ returns m-sample $\mathcal{S}^{\mathscr{T}}$ such that its observation matrix is $\mathrm{S}^{\rm M} \doteq [\mathrm{SF}^{\rm a}|\mathrm{MSF}^{\rm s}]$, and each example $\mathcal{S}^{\mathscr{T}} \ni (\boldsymbol{x}_i^{\rm M}, y_i) \doteq ((\mathrm{S}^{\rm M})^\top \mathbf{1}_i, y_i)$.*

We consider M fixed beforehand. Figure 2 (top) presents the CP on a toy data with M a permutation matrix (invertible). Figure 2 (bottom) presents another example with M block-uniform (non invertible).

4 The Crossover Process and measures of (un)fairness

Before drilling into the technical impact on learnability of a CP, it is good to make a small incursion in how the CP can be used a simple model of data protection that has received a surge of treatment over the last years [Machanavajjhala and Kifer, 2015, Feldman et al., 2015, Hardt et al., 2016, Kleinberg et al., 2016]. It is sometimes related to as fairness, equality of opportunity, equalized odds or disparate impact. It essentially builds on odds ratios.

Let x_C and x_A be two binary attributes and $\boldsymbol{\pi}$ a predicate defined on other description variables, like for example $\boldsymbol{\pi} \equiv \boldsymbol{x}_{\mathcal{V}} = \boldsymbol{v}$, where $\boldsymbol{x}_{\mathcal{V}} \subseteq \mathcal{F} \backslash \{x_C, x_A\}$ and $\boldsymbol{v}$ is an instantiation of $\boldsymbol{x}_{\mathcal{V}}$. Define the *odds ratio*

$$\rho(x_C, x_A, \boldsymbol{\pi}|\mathcal{S}) \doteq \frac{\Pr_{\mathcal{S}}[x_C = 1 | x_A = 0, \boldsymbol{\pi}]}{\Pr_{\mathcal{S}}[x_C = 1 | x_A = 1, \boldsymbol{\pi}]} \ . \tag{1}$$

In this definition, x_C is the sensitive feature, x_A is a protected attribute [Hardt et al., 2016] and the eventual additional features in $\boldsymbol{x}_{\mathcal{V}}$ are a *private* subset of attributes. For example, private attributes can contain additional features on which we want to constrain fairness measures, like qualification in Hardt et al. [2016]. If $\boldsymbol{\pi} = \top$ (the predicate that is always true), we just write $\rho(x_C, x_A|\mathcal{S})$.

Definition 3 *Let x_A, x_C be two binary attributes and $\boldsymbol{\pi}$ a predicate defined on other description variables. For any $\rho \in \mathbb{R}_+$, we say that sample $\mathcal{S}$ has ρ-odds ratio for the triple $(x_A, x_C, \boldsymbol{\pi})$ iff $\rho(x_C, x_A, \boldsymbol{\pi}|\mathcal{S}) = \rho$.*

We can also replace real ρ by a subset $\mathcal{R} \subseteq \mathbb{R}_+$, in which case we must have $\rho(x_C, x_A, \boldsymbol{\pi}|\mathcal{S}) \in \mathcal{R}$. Here are some examples of how this definition aligns with previous works. If one takes $\boldsymbol{\pi} \equiv x_{\hat{C}} = 1$ where $x_{\hat{C}}$ is a proxy for x_C, like an estimate for x_C obtained using a specific procedure, then requiring $\rho = 1$ brings the condition for balance on the positive class from Kleinberg et al. [2016];




Figure 2: *Top*: example of CP with M a block-class permutation matrix (the two blocks are in bold). *Bottom*: toy domain where $d = 3$, but all examples have zero z-coordinate (not shown). The CP uniformly mixes examples by class. The domain consists of two spirals (red for positive, green for negative examples) with $\mathcal{D}$ = uniform distribution. Arrows depict respectively the optimal direction (black), and the directions learned by minimizing φ = square loss over $\mathcal{S}$ (light green, "ex") and $\mathcal{S}^{\mathcal{T}}$ (blue, "CP"). The right plot displays test errors (y-scale) on uniform sampling of datasets of different sizes (x-scale). The effect of the CP is to produce in $\mathcal{S}^{\mathcal{T}}$ *two* distinct examples that *average* the positive / negative examples, and yield a better approximation of the optimum.

if on the other hand $\boldsymbol{\pi} \equiv x_{\hat{C}} = 0$, then requiring $\rho = 1$ brings the condition for balance on the negative class from Kleinberg et al. [2016]; if finally $\boldsymbol{\pi} = \top$, requiring $\rho = 1$ brings the condition for balance within groups from Kleinberg et al. [2016]. Replacing $\rho = 1$ by $\mathcal{R} \doteq [1 - \epsilon, 1 + \epsilon]$ brings the corresponding approximated fairness conditions of Kleinberg et al. [2016]. Permuting, in the balance for positive class, the role of x_C and $x_{\hat{C}}$, still with $\alpha = 1$, brings the condition for equal opportunity in Hardt et al. [2016], adding a second 1-odds ratio condition for $(x_A, x_{\hat{C}}, x_{\hat{C}} = 0)$ brings equalized odds in Hardt et al. [2016]. Finally, replacing, $\rho = 1$ by $\mathcal{R} = (0.8, +\infty)$ in the condition for balance within groups above yields the no-disparate impact condition of Feldman et al. [2015]. Let us see now what a simple CP can do to alter odds like in eq. (1), via the following Definition.

Definition 4 *We say that a* CP $\mathcal{T}$ *shifts the odds ratio for the triple* $(x_A, x_C, \boldsymbol{\pi})$ *by* Δ *on dataset* $\mathcal{S}$ *iff* $\rho(x_C, x_A, \boldsymbol{\pi}|\mathcal{S}^{\mathcal{T}}) = \rho(x_C, x_A, \boldsymbol{\pi}|\mathcal{S}) + \Delta$.

Lemma 5 *Suppose Table 1 describes the observed joint distribution for attributes* x_A *and* x_C *in sample* $\mathcal{S}$*, conditioned on* $\boldsymbol{\pi}$ *being true (hence,* $a + b + c + d \leq m$*). Let*

$$\Delta(i) \doteq \frac{b+d}{d-i} \cdot \frac{i}{d} \ , \forall i \in \mathbb{Z} \ . \tag{2}$$

	$x_C = 0$	$x_C = 1$
$x_A = 0$	a	b
$x_A = 1$	c	d

Table 1: Contingency table (conditioned on $\boldsymbol{\pi}$ being true) for two binary attributes A and C between which a CP is going to change the dependency relationships and odds ratios ($a+b+c+d \leq m$, see text for details).

Then, for any sample $\mathcal{S}$ and any $i \in \{-\min\{b,c\}, -\min\{b,c\}+1, ..., \min\{a,d\}\}$, there exists a CP *$\mathscr{T}_i$ that shifts the odds ratio for the triple $(x_A, x_C, \boldsymbol{\pi})$ by $\Delta(i)$ on $\mathcal{S}$.*

(proof in Subsection 9.1) The proof of the Lemma involves very simple CPs, for which the shuffle matrix M is a permutation matrix. As a consequence, the Lemma implies that such a simple CP can produce fairness ($\rho(x_C, x_A, \boldsymbol{\pi}|\mathcal{S}^{\mathscr{T}}) = 1$) as long as $b \leq d + 2\min\{a,d\}$ and $d \leq b + 2\min\{b,c\}$, *i.e.* as long as the joint distribution is not too unbalanced; since i can take on both positive and negative values, a CP can also shifts the odds ratio in $\mathcal{S}$ to smaller or larger values. If $b = d$, it can therefore also break fairness with shifts $\Delta(i) = 2i/(d-i)$.

5 The Crossover Process and learnability

Generalization — We now explore the effect of the CP on generalisation. We need two assumptions on $\mathcal{H}$ and φ. The first is a weak linearity condition on $\mathcal{H}$:

(i) $\forall h \in \mathcal{H}$, $\exists$ classifiers $h_{\mathrm{a}}, h_{\mathrm{s}}$ over $\mathcal{F}_{\mathrm{a}}, \mathcal{F}_{\mathrm{s}}$ s. t. $h(\boldsymbol{x}) = h_{\mathrm{a}}((\mathrm{F}^{\mathrm{a}})^\top \boldsymbol{x}) + h_{\mathrm{s}}((\mathrm{F}^{\mathrm{s}})^\top \boldsymbol{x})$.

$(\mathrm{F}^{\mathrm{s}})^\top \boldsymbol{x}$ picks the features of $\boldsymbol{x}$ in $\mathcal{F}_{\mathrm{s}}$. Such an assumption is also made in the feature bagging model [Sutton et al., 2006]. Any linear classifier satisfies **(i)**, but also any linear combination of arbitrary classifiers, each learnt over one of $\mathcal{F}_{\mathrm{a}}$ and $\mathcal{F}_{\mathrm{s}}$. We let $\mathcal{H}_{\mathrm{s}}$ denote the set of all h_{s}. The second assumption postulates that key quantities are bounded [Bartlett and Mendelson, 2002]:

(ii) $0 \leq \varphi(z) \leq K_\varphi, \forall z$ and $|h_{\mathrm{s}}((\mathrm{F}^{\mathrm{s}})^\top \boldsymbol{x})| \leq K_{\mathrm{s}}, \forall \boldsymbol{x} \in \mathcal{X}, \forall h_{\mathrm{s}} \in \mathcal{H}_{\mathrm{s}}$.

Let $\mathrm{R}_{\mathcal{S}}(\mathcal{H}) \doteq \mathbb{E}_{\boldsymbol{\sigma}\sim\Sigma_m}\left[\sup_{h\in\mathcal{H}} |(1/m)\cdot\sum_i \sigma_i h(\boldsymbol{x}_i)|\right]$ be the empirical Rademacher complexity of $\mathcal{H}$. Additionally, we coin the Rademacher CP complexity, RCP.

Definition 6 *The Rademacher* CP *complexity (*RCP*) of $\mathcal{H}$ with respect to $\mathscr{T} \doteq \mathrm{CP}(\mathcal{S}; \mathrm{F}^{\mathrm{a}}, \mathrm{F}^{\mathrm{s}}, \mathrm{M})$ is:*

$$\mathrm{RCP}_{\mathscr{T}}(\mathcal{H}) \doteq \mathbb{E}_{\boldsymbol{\sigma}\sim\Sigma_m}\left[\sup_{h\in\mathcal{H}_{\mathrm{s}}} \left| \frac{1}{m}\sum_i \sigma_i \left(h((\mathrm{SF}^{\mathrm{s}})^\top \mathbf{1}_i) - h((\mathrm{MSF}^{\mathrm{s}})^\top \mathbf{1}_i)\right)\right|\right] . \quad (3)$$

Notice that the RCP is computed over the shuffle set of features only, and $(\mathrm{SF}^{\mathrm{s}})^\top \mathbf{1}_i = (\mathrm{F}^{\mathrm{s}})^\top \boldsymbol{x}_i$. The next Theorem expresses a generalisation bound *wrt* the CP.

Theorem 7 *Consider any $\mathcal{H}$, φ and split $\mathcal{F} = \mathcal{F}_{\rm a} \cup \mathcal{F}_{\rm s}$ such that **(i)** and **(ii)** hold. For any m and any $\delta > 0$, with probability $\geq 1 - \delta$ over i.i.d. m-sample $\mathcal{S}$, we have:*

$$\mathbb{E}_{\mathcal{D}}\left[L_{0/1}(y, h(\boldsymbol{x}))\right] \;\leq\; \mathbb{E}_{\mathcal{S}\mathscr{T}}\left[\varphi(yh(\boldsymbol{x}))\right] + \textsc{rcp}_{\mathscr{T}}(\mathcal{H}) + \frac{4}{b_\varphi} \cdot \textsc{r}_{\mathcal{S}}(\mathcal{H}) + (2K_\varphi + K_{\rm s}) \cdot \sqrt{\frac{2}{m}\log\frac{3}{\delta}}\ ,$$

for every classifier h and every $\mathscr{T} \doteq \textsc{cp}(\mathcal{S}; \mathrm{F}^{\rm a}, \mathrm{F}^{\rm s}, \mathrm{M})$ such that $\mathrm{M} \in \mathcal{M}_m^*$. Here, $b_\varphi > 0$ is a constant depending on φ.

(proof in Subsection 9.2) Notice that Theorem 7 requires that M is a block-class matrix. A key to the proof is the invariance of the mean operator: $\boldsymbol{\mu}_{\mathcal{S}} = \boldsymbol{\mu}_{\mathcal{S}\mathscr{T}}$. Theorem 7 says that a key to good generalisation is the control of $\textsc{rcp}_{\mathscr{T}}(\mathcal{H})$. We would typically want it to be small compared to the Rademacher complexity penalty. The rest of this Section shows that (and when) this is indeed achievable.

Upperbounds on $\textsc{rcp}_{\mathscr{T}}(\mathcal{H})$ — We consider different configurations of $\mathcal{H}$ and / or $\mathscr{T}$:

Setting (A): Classifiers $h^{\rm s}$ and $h^{\rm a}$ in **(i)** above are linear;

Setting (B): $\mathrm{M} \in S_m^*$.

The following Lemma establishes a first bound on $\textsc{rcp}_{\mathscr{T}}(\mathcal{H})$.

Lemma 8 *if $\mathscr{T}$ satisfies the conditions of Theorem 7, then $\textsc{rcp}_{\mathscr{T}}(\mathcal{H}) \leq 2 \cdot \textsc{r}_{\mathcal{S}'}(\mathcal{H}_{\rm s})$, for $\mathrm{S}' \doteq (\mathrm{I}_m - \mathrm{M})\mathrm{SF}^{\rm s}$ in Setting (A), and $\mathrm{S}' \doteq \mathrm{SF}^{\rm s}$ in Setting (B). S' is the row-wise observation matrix of $\mathcal{S}'$.*

Proof (Sketch) Consider for example Setting (B). In this case, recalling that $(\mathrm{SF}^{\rm s})^\top \mathbf{1}_i = (\mathrm{F}^{\rm s})^\top \boldsymbol{x}_i$ and letting $\varsigma : [m] \to [m]$ denote the permutation that M represents, we have because of the triangle inequality:

$$\begin{aligned}
\textsc{rcp}_{\mathscr{T}}(\mathcal{H}) \;&=\; \mathbb{E}_{\boldsymbol{\sigma}\sim\Sigma_m}\left[\sup_{h\in\mathcal{H}_{\rm s}}\left|\frac{1}{m}\sum_i \sigma_i\left(h((\mathrm{F}^{\rm s})^\top\boldsymbol{x}_i) - h((\mathrm{F}^{\rm s})^\top\boldsymbol{x}_{\varsigma(i)})\right)\right|\right] \qquad (4)\\
&\leq\; \mathbb{E}_{\boldsymbol{\sigma}\sim\Sigma_m}\left[\sup_{h\in\mathcal{H}_{\rm s}}\left|\frac{1}{m}\sum_i \sigma_i h((\mathrm{F}^{\rm s})^\top\boldsymbol{x}_i)\right|\right] + \mathbb{E}_{\boldsymbol{\sigma}\sim\Sigma_m}\left[\sup_{h\in\mathcal{H}_{\rm s}}\left|\frac{1}{m}\sum_i \sigma_i h((\mathrm{F}^{\rm s})^\top\boldsymbol{x}_{\varsigma(i)})\right|\right]\\
&= 2 \cdot \textsc{r}_{\mathcal{S}'}(\mathcal{H}_{\rm s})\ ,
\end{aligned}$$

as claimed. The case of Setting (A) follows the same path. ■

Lemma 8 says that $\textsc{rcp}_{\mathscr{T}}(\mathcal{H})$ is at most twice a Rademacher complexity over the *shuffle set*. This bound is however loose since many terms can cancel in the sum of eq. (4), and the inequality does not take this into account. In particular,

Theorem 9 *Under Setting (A), suppose any $h_{\rm s}$ is of the form $h_{\rm s}(\boldsymbol{x}) = \boldsymbol{\theta}^\top\boldsymbol{x}$ with $\|\boldsymbol{\theta}\|_2 \leq r_{\rm s}$, for some $r_{\rm s} > 0$. Let $\mathrm{K}^{\rm s} \doteq \mathrm{SF}^{\rm s}(\mathrm{SF}^{\rm s})^\top$. Then $\exists u \in (0,1)$ depending only in $\mathcal{S}$ such that for any $\mathrm{M} \in \mathcal{M}_m$,*

$$\textsc{rcp}_{\mathscr{T}}(\mathcal{H}) \;\leq\; (ur_{\rm s}/m) \cdot \sqrt{\langle \mathrm{I}_m, \mathrm{K}^{\rm s}\rangle_{\mathrm{M}}}\ . \qquad (5)$$

Notice that K^{s} is a Gram matrix in the shuffle feature space. The proof technique (Subsection 9.4) relies on a data-dependent expression for u which depends on the cosines of angles between the observations in $\mathcal{S}$. It can be used to refine and improve a popular bound on the empirical Rademacher complexity of linear classifiers [Kakade et al., 2008] (we give the proof in Theorem 20 in the Appendix). We now investigate an upperbound on Setting (B) in which classifiers in $\mathcal{H}^{\mathrm{s}}$ are (rooted) directed acyclic graph (DAG), like decision trees, with bounded real valued predictions (say, $K_{\mathrm{s}} > 0$) at the leaves. Each classifier h_{s} defines a partition over $\mathcal{X}$. We let $\mathcal{H}^{\mathrm{s}}_{+}$ be the subset of $\mathcal{H}^{\mathrm{s}}$ in which all leaves have in absolute value the largest magnitude, *i.e.*, K_{s}. Remark that we may have $|\mathcal{H}^{\mathrm{s}}_{+}| \ll \infty$ while $|\mathcal{H}^{\mathrm{s}}| = \infty$ in general.

Theorem 10 *Under Setting (B), suppose $\mathcal{H}^{\mathrm{s}}$ is* DAG *and assumption **(ii)** is satisfied. Suppose that $\log|\mathcal{H}^{\mathrm{s}}_{+}| \geq (4\varepsilon/3)\cdot m$ for some $\varepsilon > 0$. Then, letting $\mathrm{odd_cycle}(\mathrm{M})$ denote the set of odd cycles (excluding fixed points) of* M*, we have:*

$$\textsc{rcp}_{\mathscr{T}}(\mathcal{H}) \quad \leq \quad K_{\mathrm{s}} \cdot \sqrt{\frac{2}{m} \cdot \log \frac{|\mathcal{H}^{\mathrm{s}}_{+}|}{(1+\varepsilon)^{|\mathrm{odd_cycle}(\mathrm{M})|}}} \quad . \tag{6}$$

(proof in Subsection 9.5) The assumption on $\mathcal{H}^{\mathrm{s}}_{+}$ is not restrictive and would be met by decision trees, branching programs, etc. (and subsets). Usual bounds on the Rademacher complexity of decision trees would roughly be the right-hand side of (6) *without* the denominator in the $\log$ (see for example [Schapire and Freund, 2012, Chapter 5]). Hence, the RCP may be significantly smaller than the Rademacher complexity for more "involved" CPs. The number of cycles is not the only relevant parameter of the CP on which relies non-trivial bounds on $\textsc{rcp}_{\mathscr{T}}(\mathcal{H})$: the Appendix presents, for the interested reader, a proof that the number of fixed points is another parameter which can decrease significantly the *expected* RCP (by a factor $\sqrt{1-|\mathrm{fixed_points}|/m}$), when CPs are picked at random (see Theorem 22 and discussion in Subsection 9.5).

At last, we notice that Theorem 7 gives a perhaps counterintuitive rationale for the CP that goes beyond our framework to machine learning at large: *learning over a* CP*'ed $\mathcal{S}$ may improve generalisation over $\mathcal{D}$ as well*. By means of words, learning over transformed data may improve generalisation over the initial domain. Figure 2 (bottom) gives a toy example for which this holds. It is also not hard to exhibit domains for which we even have:

$$\min_{h} \mathbb{E}_{\mathcal{S}\mathscr{T}}\left[\varphi(yh(\boldsymbol{x}))\right] + \textsc{rcp}_{\mathscr{T}}(\mathcal{H}) \quad < \quad \min_{h} \mathbb{E}_{\mathcal{S}}\left[\varphi(yh(\boldsymbol{x}))\right] \quad . \tag{7}$$

In order not to load the paper's body, we present such an example in the Appendix (Subsection 9.3). Having discussed learning guarantees, we are ready to dive into more applications of the Crossover Process for data protection. In addition to the measures of (un)fairness developed in Section 4, we develop the CP in two other frameworks: Hilbert-Schmidt independence and *do* calculus. In the former one, our results exploit the design of the shuffle matrix M to alter independence; in the latter one, our results exploit the split step of the CP to interfere with causal inference.

6 The Crossover Process and statistical independence

Here, we assume that $\mathcal{S}$ is subject to quantitative tests of independence, that is, assessing $\mathcal{U} \perp\!\!\!\perp \mathcal{V}$ for some $\mathcal{U}, \mathcal{V} \subset \mathcal{X}$. We compute CPs such that $\mathcal{U} \subseteq \mathcal{F}_{\mathrm{a}}$ and $\mathcal{V} \subseteq \mathcal{F}_{\mathrm{s}}$, so that the CP alters

the measure of independence. One popular criterion to determine (conditional) (in)dependence is Hilbert-Schmidt Independence Criterion [Doran et al., 2014, Gretton et al., 2005, 2007].

Definition 11 *Let $\mathcal{U} \subset [d]$ and $\mathcal{V} \subset [d]$ be non-empty and disjoint. Let K^u and K^v be two kernel functions over $\mathcal{U}$ and $\mathcal{V}$ computed using $\mathcal{S}$. The (unnormalised) Hilbert-Schmidt Independence Criterion (*HSIC*) between $\mathcal{U}$ and $\mathcal{V}$ is defined as* $\text{HSIC}(\mathrm{K}^u, \mathrm{K}^v) \doteq \langle \mathrm{K}^u, \mathrm{K}^v \rangle_{(1/m)\mathbf{1}\mathbf{1}^\top}$.

(We choose not to normalise the HSIC: various exist but they mainly rely on a multiplicative factor depending on m only, so they do not affect the results to come.) The choice of M in the CP directly influences the value the result of HSIC; therefore, we can design a search strategy aimed to alter it. In the same way as we did for Section 4, we provide a Definition for the alteration in the HSIC criterion caused by a CP, and then a Theorem that quantifies it precisely for any CP.

Definition 12 *We say that a* CP $\mathscr{T}$ *shifts the* HSIC *criterion between $\mathcal{U}$ and $\mathcal{V}$ by Δ (on dataset $\mathcal{S}$) iff* $\text{HSIC}_{\mathscr{T}} - \text{HSIC} = \Delta$*, where* HSIC *and* $\text{HSIC}_{\mathscr{T}}$ *respectively denote the* HSIC *before and after applying the* CP.

Theorem 13 *For any dataset $\mathcal{S}$, let $\tilde{\boldsymbol{u}} \doteq (1/m)\sum_i \lambda_i(\mathbf{1}^\top \boldsymbol{u}_i)\boldsymbol{u}_i$, $\tilde{\boldsymbol{v}} \doteq (1/m)\sum_i \mu_i(\mathbf{1}^\top \boldsymbol{v}_i)\boldsymbol{v}_i$, where $\{\lambda_i, \boldsymbol{u}_i\}_{i\in[d]}$, $\{\mu_i, \boldsymbol{v}_i\}_{i\in[d]}$ are respective eigensystems of K^u and K^v. Then for* ***any*** CP $\mathscr{T}$*, $\mathscr{T}$ shifts the* HSIC *criterion between $\mathcal{U}$ and $\mathcal{V}$ by Δ (on $\mathcal{S}$) with*

$$\Delta \;=\; 2m \cdot \tilde{\boldsymbol{u}}^\top (\mathrm{I}_m - \mathrm{M})\tilde{\boldsymbol{v}} \;, \tag{8}$$

where M *is the shuffling matrix of the* CP.

(proof in Subsection 9.6) This result shows that altering the HSIC criterion with a CP is also algorithmic friendly: (a) while storing kernels requires $O(m^2)$ space, controlling the evolution of the HSIC requires only *linear*-space information about kernels, and (b) this information can be computed beforehand, and can be efficiently approximated from low-rank approximations of the kernels [Bach, 2013]. Theorem 13 also shows that the sign of the shift Δ is determined by the eigenspace of the shuffling matrix (when it can be diagonalized).

We now go one step further in showing the algorithmic friendliness of computing a CP that shifts HSIC. In the following Theorem, we *compose* CP processes with T different elementary permutation shuffling matrices. Notice that since the composition of permutation matrices is a permutation matrix, when the matrix of the final process $\text{HSIC}_{\mathscr{T}_T}$ is block class, Theorem 7 can be applied *directly* to $\text{HSIC}_{\mathscr{T}_T}$, so we get, at each iteration, a quantification of the protection achieved (shift Δ) and the impact on learnability. We let $\mathscr{R}^{u,v} \doteq m\,(1 - (\mathrm{K}^u_{\cdot\cdot} + \mathrm{K}^v_{\cdot\cdot})/(2m^2))$.

Theorem 14 *Suppose $\mathscr{T}$ is built by a sequence of $T = \epsilon m$ elementary permutation ($\epsilon > 0$) and the kernels K^u and K^v have unit diagonal. Suppose that the initial* $\text{HSIC} > \mathscr{R}^{u,v}$ *(before applying the* CP*). Then there exists such a sequence of elementary permutations such that $\mathscr{T}_T$ shifts the* HSIC *criterion between $\mathcal{U}$ and $\mathcal{V}$ by Δ (on $\mathcal{S}$) with*

$$\Delta \;\leq\; -(1-\alpha)\cdot(\text{HSIC} - \mathscr{R}^{u,v}) \;,$$

where $\alpha \doteq \exp(-8\epsilon) < 1$.

The proof (Subsection 9.7) states a more general result, not restricted to unit diagonal kernels. Theorem 14 is a worst-case result that shows sufficient conditions for negative shift, and therefore decrease the HSIC criterion. Note that some sequences of permutations may be much more efficient in decreasing HSIC. If we compare this bound to Theorem 3 in [Gretton et al., 2007], then $\mathscr{R}^{u,v}$ may be *below* the expectation of the HSIC, so indeed we can obtain $\Delta < 0$ and somehow trick statistical tests in keeping independence after the CP, while they would eventually reject it before.

Remark: it is in fact possible to kill two birds with one stone, namely trick statistical tests into keeping independence *and* then incur arbitrarily large errors in estimating causal effects. Our basis is the Cornia-Mooij (CM) model [Cornia and Mooij, 2014] which, for space considerations, we defer to the Appendix (Subsection 9.8). One interesting feature of this particular causal graph is that it is so simple that it may be found as subgraph of real-world domains, thus for which the results we give would directly transfer.

7 The Crossover Process and causality

We assume here basic knowledge of the causality and *do*-calculus frameworks Pearl [2000]. We consider the case where the causal directed acyclic graph is known, and the goal is to interfere with the inference of causal effects between covariates. The key challenge for causal inference is the existence of confounding variables that are causes of both the exposure and outcome variables. For example, suppose impact of hormone replacement therapy on women's health was captured by the causal DAG $I \to T \to H, I \to H$, where I represents income, T represents taking the treatment and H is the health outcome of interest. If wealthier women are more likely to see a doctor for treatment and also have generally better health, then $\Pr[H|T]$ will be more positive than the true causal effect $\Pr[H|do(T)]$.

Adjusting for such nuisance or confounding variables, either by matching [Greenwood, 1945, Rubin, 1973] or regression [Fisher, 1935], is a central tool in economics and social sciences [Morgan and Winship, 2014]. The back-door criterion [Pearl, 2000] clarifies which variables it is appropriate to condition on in order to achieve unbiased estimates of causal effects. The CP can be designed to interfere with obtaining causal estimates via such adjustments.

Let $G = (\mathcal{U} \cup \mathcal{F}, \mathcal{A})$ be a causal directed acyclic graph over observable vertices $\mathcal{F}$, latent variables $\mathcal{U}$ and arcs $\mathcal{A}$ [Pearl, 2000]. We are given a set $\mathcal{Q} \doteq \{(x_i, x_i'), i \in [q]\}$ of q causal queries, each of which represents the estimation of $\Pr[x_i|\mathrm{do}(x_i')]$.

A covariate adjustment (adjustment for short) for a query (x_i, x_i') is a set $\mathcal{Z}_i \subset \mathcal{F}$ such that $x_i', x_i \notin \mathcal{Z}_i$ and

$$\Pr[x_i|\mathrm{do}(x_i')] \;=\; \sum_{z \sim \mathcal{Z}_i} \Pr[x_i|x_i', z] \Pr[z] \;, \tag{9}$$

An adjustment is not guaranteed to exist. In our example, for the query (H, T), there is no adjustment if $I \in \mathcal{U}$. An adjustment is minimal iff it does not contain any other adjustment as proper subset. Note that $\mathcal{Z}_i$ can be the empty set.

In the same way as we did for Sections 4 and 6, we provide a Definition for the alteration in the *do*-calculus caused by a CP, and then a Lemma that qualifies it more precisely for any CP. It is a weaker result than the former ones because we do not quantify a shift, but rather just state changes in estimation.

Definition 15 *We say that a* CP $\mathscr{T}$ *interferes with causal inference via adjustment for a query* (x_i, x_i') *if the solution to (9) differs between the datasets* $\mathcal{S}$ *and* $\mathcal{S}^{\mathscr{T}}$.

We put no constraint on the magnitude of the change, so interfering with causal queries is essentially a matter of biasing the distributions involved in the right-hand side of (9). Let Z_i denote the set of minimal adjustments for query (x_i, x_i').

Lemma 16 *Let* $\mathcal{V}_i = x_i' \cup x_i \cup \mathcal{Z}_i$. *A* CP *interferes with causal inference via adjustment for the query* (x_i, x_i') *iff* $\forall \mathcal{Z}_i \in \mathsf{Z}_i$, $\exists$ *variables* $v_{\mathrm{a}}, v_{\mathrm{s}} \in \mathcal{V}_i$ *such that* $v_{\mathrm{a}} \in \mathcal{F}_{\mathrm{a}}$ *and* $v_{\mathrm{s}} \in \mathcal{F}_{\mathrm{s}}$, *where* $\mathcal{F}_{\mathrm{a}}$ *and* $\mathcal{F}_{\mathrm{s}}$ *are the shuffle and anchor set of features of the* CP.

The proof is a direct consequence of eq. (9) and the fact that the shuffle matrix M alters joint distributions between variables that do not belong to the same split set, without touching marginals. We can always interfere with a single query (x_i, x_i') by ensuring x_i and x_i' are in different splits. To simultaneously interfere with the set of queries $\mathcal{Q}$, we must first find the set of minimal adjustments $\mathsf{Z}_i, \forall i \in [q]$, then select a split that satisfies Lemma 16 for every query. This involves heavy combinatorics. Enumerating the adjustments can be done with cubic *delay* per adjustment [Textor and Liskiewicz, 2012](the set of minimal adjustments for a given query can grow exponentially with d). The second step subsumes the infamous Set Splitting problem [Garey and Johnson, 1979], which is NP-Complete. In practice, causal graphs must often be constructed by humans so this approach can still be computationally feasible for a small set of queries. Exploring the addition of constraints on the graph (such as sparsity) to develop more efficient algorithms is an interesting avenue for future research.

A causal query is *identifiable* if we can obtain an expression for it purely in terms of distributions over the observable variables $\mathcal{F}$. The existence of an adjustment is sufficient but not necessary for identifiability. In theory, a causal query for which we have interfered with any adjustments, could still be identified via another approach. The *do*-calculus [Pearl, 2000] and Identify Algorithm [Shpitser and Pearl, 2006] provide a complete framework for determining if a query is identifiable and computing an expression for it. In principle, we could utilize the CP to interfere with all routes to identifiability. This would require an algorithm that could enumerate the expressions for a causal query. We are not aware of such an algorithm in the literature. In practice, expressions that are not of the form of 9 are rarely used.

8 Discussion and conclusion

This paper introduces the Crossover Process (CP), a mechanism that cross-modifies data using a generalisation of stochastic matrices. This process can be used to cope with data optimisation for supervised learning, as well as for the problem of handling a process-level protection on data such as causal inference attacks on a supervised learning dataset. In this case, the CP allows to release data with spotless low-level description (variable names, observed values, marginals), substantial utility (learnability), but disclosing dependences and causal effects under control, and thus that could even be crafted to be conflicting with a ground truth to protect[5]. We have chosen to focus here

[5]Note that the initial data may not be lost, as opposed to differential privacy: knowing the noise parameters does not allow to revert differential privacy protection, while a CP protection is reversible when the shuffling matrix M is invertible.

on three major components of the actual trends, namely (un)fairness measures, statistical measures of causal inference and causal queries. In these directions, there are some very interesting and non-trivial avenues for future research, like for example the control of combinatorial blow-up in the worst case for causal queries, ideally as a function of the causal graph structure. There are also more applications of the CP in the field of causal discovery. Suppose for example that description features denote transactions. Since we modify joint distributions without touching on marginals, our technique has direct applications in causal rule mining, with the potential to fool any level-wise association rule mining algorithms, that is, any spawn of Apriori [Li et al., 2016].

The theory we develop fo CP introduces a new complexity measure of the process, the Rademacher CP complexity. We do believe that the CP is also a good contender in the pool of methods optimising data for learning, and it may provide new metrics, algorithms and tools to devise improved solutions that fit to challenging domains not restricted to optimizing learning or data privacy.

References

F. Bach. Sharp analysis of low-rank kernel matrix approximations. In *26 th COLT*, pages 185–209, 2013.

S. Barocas and H. Nissenbaum. Big data's end run around anonymity and consent. In J. Lane, V. Stodden, S. Bender, and H. Nissenbaum, editors, *Privacy, Big Data, and the Public Good*, pages 44–75. Cambridge University Press, 2014a.

S. Barocas and H. Nissenbaum. Big data's end run around procedural privacy protection. *Communications of the ACM*, 57:31–33, 2014b.

P.-L. Bartlett and S. Mendelson. Rademacher and gaussian complexities: Risk bounds and structural results. *JMLR*, 3:463–482, 2002.

S. Chatterjee. Stein's method for concentration inequalities. *Probability Theory and Related Fields*, 138:305–321, 2007.

K. Chaudhuri, C. Monteleoni, and A.-D. Sarwate. Differentially private empirical risk minimization. *JMLR*, 12:1069–1109, 2011.

N. Cornia and J.-M. Mooij. Type-II errors of independence tests can lead to arbitrarily large errors in estimated causal effects: An illustrative example. In *30th UAI Workshops*, pages 35–42, 2014.

G. Doran, K. Muandet, K. Zhang, and B. Schölkopf. A permutation-based kernel conditional independence test. In *30 th UAI*, 2014.

J. C. Duchi, M. I. Jordan, and M. J. Wainwright. Local privacy and statistical minimax rates. In *Proc. of the 53 rd FOCS*, pages 429–438, 2013.

C. Dwork and A. Roth. The algorithmic foudations of differential privacy. *Foundations and Trends in Theoretical Computer Science*, 9:211–407, 2014.

M. Enserink and G. Chin. The end of privacy. *Science*, 347:490–491, 2015.

M. Feldman, S.-A. Friedler, J. Moeller, C. Scheidegger, and S. Venkatasubramanian. Certifying and removing disparate impact. In *21st KDD*, pages 259–268, 2015.

R.-A Fisher. *The design of experiments*. Oliver and Boyd, Edinburgh, 1935.

M.R. Garey and D.S. Johnson. *Computers and Intractability, a guide to the theory of NP-Completeness*. Bell Telephone Laboratories, 1979.

S. Graham, W. Press, S.-J. Gates, M. Gorenberg, J.-P. Holden, E. Lander, C. Mundie, M. Savitz, and E. Schmidt. Big data and privacy: a technological perspective. CreateSpace Independent Publishing Platform, 2015. — President's Council of Advisors on Science and Technology.

E. Greenwood. *Experimental sociology: A study in method*. King's crown Press, 1945.

A. Gretton, O. Bousquet, A.-J. Smola, and B. Schölkopf. Measuring statistical dependence with Hilbert-Schmidt norms. In *16th ALT*, pages 63–77, 2005.

A. Gretton, K. Fukumizu, C.-H. Teo, L. Song, B. Schölkopf, and A. Smola. A kernel statistical test of independence. In *NIPS*20*, pages 585–592, 2007.

M. Hardt, E. Price, and N. Srebro. Equality of opportunity in supervised learning. In *NIPS*29*, pages 3315–3323, 2016.

P.-O. Hoyer, D. Janzing, J.-M. Mooij, J. Peters, and B. Schölkopf. Nonlinear causal discovery with additive noise models. In *NIPS*21*, pages 689–696, 2008.

D. Janzing, J.-M. Mooij, K. Zhang, J. Lemeire, J. Zscheischler, P. Daniusis, B. Steudel, and B. Schölkopf. Information-geometric approach to inferring causal directions. *Artificial Intelligence*, 182-183:1–31, 2012.

S. Kakade, K. Sridharan, and A. Tewari. On the complexity of linear prediction: Risk bounds, margin bounds, and regularization. In *NIPS*21*, pages 793–800, 2008.

J.-M. Kleinberg, S. Mullainathan, and M. Raghavan. Inherent trade-offs in the fair determination of risk scores. *CoRR*, abs/1609.05807, 2016.

M.-J. Kusner, Y. Sun, K. Sridharan, and K.-Q. Weinberger. Inferring the causal direction privately. In *19th AISTATS*, 2016.

J. Li, T.-D. Le, L. Liu, J. Liu, Z. Jin, B. Sun, and S. Ma. From observational studies to causal rule mining. *ACM Trans. IST*, 7:1–27, 2016.

A Machanavajjhala and D. Kifer. Designing statistical privacy for your data. *Communications of the ACM*, 58:58–67, 2015.

C. McDiarmid. Concentration. In M. Habib, C. McDiarmid, J. Ramirez-Alfonsin, and B. Reed, editors, *Probabilistic Methods for Algorithmic Discrete Mathematics*, pages 1–54. Springer Verlag, 1998.

J.-M. Mooij, J. Peters, D. Janzing, J. Zscheischler, and B. Schölkopf. Distinguishing cause from effect using observational data: methods and benchmarks. *JMLR*, 2016.

S.-L. Morgan and C. Winship. *Counterfactuals and causal inference*. Cambridge University Press, 2014.

R. Nock and F. Nielsen. On the efficient minimization of classification-calibrated surrogates. In *NIPS*21*, pages 1201–1208, 2008.

R. Nock and F. Nielsen. Bregman divergences and surrogates for learning. *IEEE Trans.PAMI*, 31: 2048–2059, 2009.

G. Patrini, R. Nock, P. Rivera, and T. Caetano. (Almost) no label no cry. In *NIPS*27*, 2014.

G. Patrini, F. Nielsen, R. Nock, and M. Carioni. Loss factorization, weakly supervised learning and label noise robustness. In *33 rd ICML*, pages 708–717, 2016.

J. Pearl. *Causality: models, reasoning, and inference*. Cambridge University Press, 2000.

J. Pearl and E. Bareinboim. External validity: from do-calculus to transportability across populations. *Statistical Science*, 29:579–595, 2014.

F. Pedregosa, G. Varoquaux, A. Gramfort, V. Michel, B. Thirion, O. Grisel, M. Blondel, P. Prettenhofer, R. Weiss, V. Dubourg, J. Vanderplas, A. Passos, D. Cournapeau, M. Brucher, M. Perrot, and E. Duchesnay. Scikit-learn: Machine learning in python. *JMLR*, 12:2825–2830, 2011.

S.-J. Rizvi and J.-R. Haritsa. Maintaining data privacy in association rule mining. In *Proc. of the 28 th VLDB*, pages 682–693, 2002.

D.-B. Rubin. Matching to remove bias in observational studies. *Biometrics*, pages 159–183, 1973.

R.-E. Schapire and Y. Freund. *Boosting: foundations and algorithms*. MIT press, 2012.

I. Shpitser and J. Pearl. Identification of joint interventional distributions in recursive semi-markovian causal models. In *Proceedings of the National Conference on Artificial Intelligence*, volume 21, page 1219. Menlo Park, CA; Cambridge, MA; London; AAAI Press; MIT Press; 1999, 2006.

L. Song, A. Smola, A. Gretton, J. Bedo, and K.-M. Borgwardt. Feature selection via dependence maximization. *JMLR*, 13:1393–1434, 2012.

N. Srivastava, G. Hinton, A. Krizhevsky, I. Sutskever, and R. Salakhutdinov. Dropout: a simple way to prevent neural networks from overfitting. *JMLR*, 15:1929–1958, 2014.

C.-A. Sutton, M. Sindelar, and A. McCallum. Reducing weight undertraining in structured discriminative learning. In *10th HLT-NAACL*, 2006.

L. Sweeney. Achieving k-anonymity privacy protection using generalization and suppression. *Int. J. Uncertainty, Fuzziness and Knowledge-based systems*, 10:571–588, 2002.

J. Textor and M. Liskiewicz. Adjustment criteria in causal diagrams: An algorithmic perspective. *CoRR*, abs/1202.3764, 2012.

9 Proofs

We shall use the following notations and shorthands. We shall sometimes replace notation $\boldsymbol{x}_i$ by $\boldsymbol{x}_i^b$ for $b \in \{-,+\}$, indicating explicitly an observation from class $b1$. We also let m_b denote the number of examples in class $b1$ for $b \in \{-,+\}$ ($m = m_+ + m_-$). Furthermore, $[m]_{m'} \doteq \{m' + i : i \in [m]\}$ ($m \in \mathbb{N}_*, m' \in \mathbb{N}$). Matrix $\mathrm{U}_m \doteq (1/m)\mathbf{1}\mathbf{1}^\top$ denotes the uniform Markov chain.

9.1 Proof of Lemma 5

Consider CP $\mathscr{T}$ in which the anchor contains x_C and the variables of $\boldsymbol{\pi}$, and the shuffle set contains x_A. Other features can be split arbitrarily. Let the shuffle matrix $\mathrm{M} \in S_n$ be any permutation matrix that transfers ω observations for which $x_C = 1$ and $x_A = 1$ to observations for which $x_C = 1$ and $x_A = 0$, thereby also moving ω observations for which $x_C = 0$ and $x_A = 0$ to observations for which $x_C = 0$ and $x_A = 1$. The resulting contingency table is therefore:

	$x_C = 0$	$x_C = 1$
$x_A = 0$	$a - \omega$	$b + \omega$
$x_A = 1$	$c + \omega$	$d - \omega$

Plus, we get the additional constraints that $\omega \leq d$ and $\omega \leq a$. But we can also transfer ω' in the opposite direction, in which case the resulting contingency table is therefore:

	$x_C = 0$	$x_C = 1$
$x_A = 0$	$a + \omega'$	$b - \omega'$
$x_A = 1$	$c - \omega'$	$d + \omega'$

This time, we get the additional constraints that $\omega' \leq b$ and $\omega' \leq c$. The corresponding odds ratio satisfy:

$$\rho(x_C, x_A, \boldsymbol{\pi}|\mathcal{S}^{\mathscr{T}}) = \frac{b+\omega}{d-\omega} \in \left[\frac{b}{d}, \frac{b+\min\{a,d\}}{d-\min\{a,d\}}\right] , \tag{10}$$

$$\rho(x_C, x_A, \boldsymbol{\pi}|\mathcal{S}^{\mathscr{T}}) = \frac{b-\omega'}{d+\omega'} \in \left[\frac{b-\min\{b,c\}}{d+\min\{b,c\}}, \frac{b}{d}\right] . \tag{11}$$

Therefore, there always exists a CP $\mathscr{T}$ bringing ρ-odds for the triple $(x_A, x_C, \boldsymbol{\pi})$ in *modified* sample $\mathcal{S}^{\mathscr{T}}$, for

$$\rho = \frac{b+i}{d-i} \ , \forall i \in \{-\min\{b,c\}, -\min\{b,c\}+1, ..., \min\{a,d\}\} \ . \tag{12}$$

There just remain to compute the difference in odds ratios,

$$\begin{aligned} \Delta(i) &\doteq \rho(x_C, x_A, \boldsymbol{\pi}|\mathcal{S}^{\mathscr{T}}) - \rho(x_C, x_A, \boldsymbol{\pi}|\mathcal{S}) \\ &= \frac{b+i}{d-i} - \frac{b}{d} \\ &= \frac{(b+d)}{d-i} \cdot \frac{i}{d} \ , \end{aligned} \tag{13}$$

as claimed.

9.2 Proof of Theorem 7

The first steps of the proof are the same as [Bartlett and Mendelson, 2002] (Theorems 5, 8). We sketch them. First, for any CP $\mathscr{T}$,

$$\begin{aligned}\mathbb{E}_{\mathcal{D}}\left[L_{0/1}(y,h(\boldsymbol{x}))\right] &\leq \mathbb{E}_{\mathcal{D}}\left[\varphi(yh(\boldsymbol{x}))\right] \\ &\leq \mathbb{E}_{\mathcal{S}\mathscr{T}}\left[\varphi(yh(\boldsymbol{x}))\right] + \sup_{h\in\mathcal{H}}\left\{\mathbb{E}_{\mathcal{D}}\left[\varphi(yh(\boldsymbol{x}))\right] - \mathbb{E}_{\mathcal{S}\mathscr{T}}\left[\varphi(yh(\boldsymbol{x}))\right]\right\} \quad . \quad (14)\end{aligned}$$

Then, since $\varphi(z) \in [0, K_\varphi]$ (assumption **(ii)**), the use of the independent bounded differences inequality [McDiarmid, 1998] yield for any sample $\mathcal{S}$ sampled from $\mathcal{D}$, and any $\varsigma \in S_m$, and any δ_1, we have with probability $\geq 1-\delta_1$:

$$\begin{aligned}&\sup_{h\in\mathcal{H}}\left\{\mathbb{E}_{\mathcal{D}}\left[\varphi(yh(\boldsymbol{x}))\right] - \mathbb{E}_{\mathcal{S}\mathscr{T}}\left[\varphi(yh(\boldsymbol{x}))\right]\right\} \\ &\quad\leq \mathbb{E}_{\mathcal{S}\sim\mathcal{D}}\left[\sup_{h\in\mathcal{H}}\left\{\mathbb{E}_{\mathcal{D}}\left[\varphi(yh(\boldsymbol{x}))\right] - \mathbb{E}_{\mathcal{S}\mathscr{T}}\left[\varphi(yh(\boldsymbol{x}))\right]\right\}\right] + K_\varphi\cdot\sqrt{\frac{2}{m}\log\frac{1}{\delta_1}} \quad . \quad (15)\end{aligned}$$

We also have, because of the convexity of $\sup$ (see [Bartlett and Mendelson, 2002]),

$$\begin{aligned}&\mathbb{E}_{\mathcal{S}\sim\mathcal{D}}\left[\sup_{h\in\mathcal{H}}\left\{\mathbb{E}_{\mathcal{D}}\left[\varphi(yh(\boldsymbol{x}))\right] - \mathbb{E}_{\mathcal{S}\mathscr{T}}\left[\varphi(yh(\boldsymbol{x}))\right]\right\}\right] \\ &\quad\leq \mathbb{E}_{\mathcal{S},\mathcal{S}'\sim\mathcal{D}}\left[\sup_{h\in\mathcal{H}}\left\{\mathbb{E}_{\mathcal{S}'}\left[\varphi(yh(\boldsymbol{x}))\right] - \mathbb{E}_{\mathcal{S}\mathscr{T}}\left[\varphi(yh(\boldsymbol{x}))\right]\right\}\right] \quad .\end{aligned}$$

The proof now takes a fork compared to [Bartlett and Mendelson, 2002], as we integrate new steps to upperbound the right-hand side. We split the right supremum in two, one which involves different datasets of size m not being subject to CP, and one which involves the same dataset with and without CP:

$$\begin{aligned}&\mathbb{E}_{\mathcal{S},\mathcal{S}'\sim\mathcal{D}}\left[\sup_{h\in\mathcal{H}}\left\{\mathbb{E}_{\mathcal{S}'}\left[\varphi(yh(\boldsymbol{x}))\right] - \mathbb{E}_{\mathcal{S}\mathscr{T}}\left[\varphi(yh(\boldsymbol{x}))\right]\right\}\right] \\ &\quad= \mathbb{E}_{\mathcal{S},\mathcal{S}'\sim\mathcal{D}}\left[\sup_{h\in\mathcal{H}}\left\{\left(\mathbb{E}_{\mathcal{S}'}\left[\varphi(yh(\boldsymbol{x}))\right] - \mathbb{E}_{\mathcal{S}}\left[\varphi(yh(\boldsymbol{x}))\right]\right) + \left(\mathbb{E}_{\mathcal{S}}\left[\varphi(yh(\boldsymbol{x}))\right] - \mathbb{E}_{\mathcal{S}\mathscr{T}}\left[\varphi(yh(\boldsymbol{x}))\right]\right)\right\}\right] \\ &\quad\leq \underbrace{\mathbb{E}_{\mathcal{S},\mathcal{S}'\sim\mathcal{D}}\left[\sup_{h\in\mathcal{H}}\left\{\mathbb{E}_{\mathcal{S}'}\left[\varphi(yh(\boldsymbol{x}))\right] - \mathbb{E}_{\mathcal{S}}\left[\varphi(yh(\boldsymbol{x}))\right]\right\}\right]}_{\doteq A} \\ &\qquad+ \underbrace{\mathbb{E}_{\mathcal{S}\sim\mathcal{D}}\left[\sup_{h\in\mathcal{H}}\left\{\mathbb{E}_{\mathcal{S}}\left[\varphi(yh(\boldsymbol{x}))\right] - \mathbb{E}_{\mathcal{S}\mathscr{T}}\left[\varphi(yh(\boldsymbol{x}))\right]\right\}\right]}_{\doteq B} \quad . \quad (16)\end{aligned}$$

We handle A and B separately.

Upperbound on A. Handling A is achieved in the usual way [Bartlett and Mendelson, 2002]. Following the usual symmetrisation trick [Bartlett and Mendelson, 2002] (Theorem 8), and the

fact [Bartlett and Mendelson, 2002] (Theorem 12.4) that φ is $1/b_\varphi$-Lipschitz [Nock and Nielsen, 2008] for some $b_\varphi > 0$, we obtain that with probability $\geq 1 - \delta_1$, we have:

$$\begin{aligned}
&\mathbb{E}_{\mathcal{S},\mathcal{S}'\sim\mathcal{D}}\left[\sup_{h\in\mathcal{H}}\left\{\mathbb{E}_{\mathcal{S}'}\left[\varphi(yh(\boldsymbol{x}))\right] - \mathbb{E}_{\mathcal{S}}\left[\varphi(yh(\boldsymbol{x}))\right]\right\}\right] \\
&\leq \frac{4}{b_\varphi}\mathbb{E}_{\boldsymbol{\sigma}\sim\Sigma_m}\left[\sup_{h\in\mathcal{H}}\left|\frac{1}{m}\sum_i \sigma_i h\left(\boldsymbol{x}_i\right)\right|\right] + K_\varphi \cdot \sqrt{\frac{2}{m}\log\frac{1}{\delta_1}} \ , \qquad (17)
\end{aligned}$$

$\forall \delta_1 > 0$.

Upperbound on B. This penalty appears when $\mathrm{M} \neq \mathrm{I}_m$. The trick is because φ is proper symmetric, there is a simple way to make appear Rademacher variables and a particular Rademacher complexity, which follows from the fact that [Patrini et al., 2014]:

$$\mathbb{E}_{\mathcal{S}}\left[\varphi(yh(\boldsymbol{x}))\right] = \frac{b_\varphi}{2m}\sum_{\sigma\in\Sigma_1}\sum_i \varphi(\sigma h(\boldsymbol{x}_i)) - \frac{\overline{h}\left(\mathcal{S}\right)}{2} \ , \qquad (18)$$

where

$$\overline{h}\left(\mathcal{S}\right) \doteq \frac{1}{m}\cdot\sum_i y_i h(\boldsymbol{x}_i) \qquad (19)$$

is the *h-mean-operator*, a statistics which can be proven to be minimally sufficient for classes given h [Patrini et al., 2014]. Now, assumption **(i)** yields the invariance of $\overline{h}\left(\mathcal{S}\right)$ under permutation operation. This is proved in the following Lemma.

Lemma 17 *(mean-operator consistency of $\mathscr{T}$) Under the conditions of Theorem 7,*

$$\overline{h}\left(\mathcal{S}\right) = \overline{h}\left(\mathcal{S}^{\mathscr{T}}\right) \ , \forall h, \forall \mathcal{S}, \forall \mathscr{T} \ . \qquad (20)$$

Proof To prove it, we let $\oplus$ denote the vector concatenation operation over the features of $\mathcal{F}_{\mathrm{a}}$ and $\mathcal{F}_{\mathrm{s}}$. We now first write using Assumption **(i)**:

$$\begin{aligned}
m\cdot\overline{h}\left(\mathcal{S}\right) &= \sum_i y_i h(\boldsymbol{x}_i) \\
&= \sum_i y_i \cdot h((\mathrm{SF}^{\mathrm{a}})^\top \mathbf{1}_i \oplus (\mathrm{SF}^{\mathrm{s}})^\top \mathbf{1}_i) \\
&= \sum_i y_i \cdot h_{\mathrm{a}}((\mathrm{SF}^{\mathrm{a}})^\top \mathbf{1}_i) + \sum_i y_i \cdot h_{\mathrm{s}}((\mathrm{SF}^{\mathrm{s}})^\top \mathbf{1}_i) \ , \qquad (21)
\end{aligned}$$

and also, for the same reason,

$$m\cdot\overline{h}\left(\mathcal{S}^{\mathscr{T}}\right) = \sum_i y_i \cdot h_{\mathrm{a}}((\mathrm{SF}^{\mathrm{a}})^\top \mathbf{1}_i) + \sum_i y_i \cdot h_{\mathrm{s}}((\mathrm{MSF}^{\mathrm{s}})^\top \mathbf{1}_i) \ . \qquad (22)$$

Therefore, we need to prove an equality that depends only upon the features of $\mathcal{F}_{\mathrm{s}}$:

$$\sum_i y_i \cdot h_{\mathrm{s}}((\mathrm{SF}^{\mathrm{s}})^\top \mathbf{1}_i) = \sum_i y_i \cdot h_{\mathrm{s}}((\mathrm{MSF}^{\mathrm{s}})^\top \mathbf{1}_i) \ . \qquad (23)$$

We have two cases to consider to prove eq. (23). Notice that since it is a block-class matrix, M admits the following block matrix decomposition:

$$\mathrm{M} \;=\; \left[\begin{array}{c|c} \mathrm{M}_+ & 0 \\ 0 & \mathrm{M}_- \end{array}\right] , \tag{24}$$

with $\mathrm{M}_b \in \mathbb{R}^{m_b \times m_b}$. We distinguish two cases.
Case 1 — Setting (A). In this case,

$$\begin{aligned}
\sum_{i\in[m]} y_i \cdot h_{\mathrm{s}}((\mathrm{MSF}^{\mathrm{s}})^\top \mathbf{1}_i) &= \sum_{i\in[m]} y_i \cdot h_{\mathrm{s}}\left(\bigoplus_{k\in[d_{\mathrm{s}}]_{d_{\mathrm{a}}}} \sum_{l\in[m]} \mathrm{M}_{il}\mathrm{S}_{lk}\right) \\
&= \sum_{i\in[m]} y_i \cdot h_{\mathrm{s}}\left(\sum_{l\in[m]} \mathrm{M}_{il} \bigoplus_{k\in[d_{\mathrm{s}}]_{d_{\mathrm{a}}}} \mathrm{S}_{lk}\right) \\
&= \sum_{i\in[m]} y_i \cdot h_{\mathrm{s}}\left(\sum_{l\in[m]} \mathrm{M}_{il}(\mathrm{SF}^{\mathrm{s}})^\top \mathbf{1}_l\right) \\
&= \sum_{i\in[m]} y_i \sum_{l\in[m]} \mathrm{M}_{il} \cdot h_{\mathrm{s}}\left((\mathrm{SF}^{\mathrm{s}})^\top \mathbf{1}_l\right) \qquad (25) \\
&= \sum_{l\in[m]} \left(\sum_{i\in[m]} \mathrm{M}_{il}\right) y_l \cdot h_{\mathrm{s}}\left((\mathrm{SF}^{\mathrm{s}})^\top \mathbf{1}_l\right) \qquad (26) \\
&= \sum_{l\in[m]} y_l \cdot h_{\mathrm{s}}\left((\mathrm{SF}^{\mathrm{s}})^\top \mathbf{1}_l\right) \; . \qquad (27)
\end{aligned}$$

Here, $\oplus$ denotes the concatenation operator. Eq. (25) holds because of Setting (A), eq. (26) holds because of the class-consistency assumption (eq. (24)), and eq. (27) holds because $\mathrm{M} \in \mathcal{M}_m$.

Case 2 — Setting (B). We have $\mathrm{M} \in S_m^*$ (and no further assumption on h_{s}). In this case, letting $\varsigma : [m] \to [m]$ represent the (block-class) permutation, we have:

$$\begin{aligned}
\sum_{i\in[m]} y_i \cdot h_{\mathrm{s}}((\mathrm{MSF}^{\mathrm{s}})^\top \mathbf{1}_i) &= \sum_{i\in[m]} y_i \cdot h_{\mathrm{s}}\left(\bigoplus_{k\in[d_{\mathrm{s}}]_{d_{\mathrm{a}}}} \sum_{l\in[m]} \mathrm{M}_{il}\mathrm{S}_{lk}\right) \\
&= \sum_{i\in[m]} y_i \cdot h_{\mathrm{s}}\left(\bigoplus_{k\in[d_{\mathrm{s}}]_{d_{\mathrm{a}}}} \mathrm{S}_{\varsigma(i)k}\right) \\
&= \sum_{i\in[m]} y_i \cdot h_{\mathrm{s}}\left((\mathrm{SF}^{\mathrm{s}})^\top \mathbf{1}_{\varsigma(i)}\right) \\
&= \sum_{i\in[m]} y_{\varsigma(i)} \cdot h_{\mathrm{s}}\left((\mathrm{SF}^{\mathrm{s}})^\top \mathbf{1}_{\varsigma(i)}\right) \qquad (28) \\
&= \sum_{l\in[m]} y_l \cdot h_{\mathrm{s}}\left((\mathrm{SF}^{\mathrm{s}})^\top \mathbf{1}_l\right) \; . \qquad (29)
\end{aligned}$$

Eq. (28) holds because M is a block-class matrix (eq. (24), and eq. (29) holds because ς is a permutation. This ends the proof of Lemma 17 ■

As a remark, when $h_{\mathrm{s}}(\boldsymbol{x}) = \boldsymbol{\theta}_{\mathrm{s}}^{\top}\boldsymbol{x}_{\mathrm{s}}$, with $\boldsymbol{\theta}_{\mathrm{s}} \in \mathbb{R}^{d_{\mathrm{s}}}$, eq. (20) shows the invariance of the mean operator

$$\mu_{\mathcal{S}\mathscr{T}} = \mu_{\mathcal{S}} \doteq \frac{1}{m} \cdot \sum_i y_i \boldsymbol{x}_i \ (\forall \mathscr{T}) \ , \tag{30}$$

as minimal sufficient statistic for the classes [Patrini et al., 2014]. Using eqs. (18) and (20) yield the first following identity ($\forall \mathcal{S}, \mathscr{T}, h$):

$$\begin{aligned}
&\mathbb{E}_{\mathcal{S}}\left[\varphi(yh(\boldsymbol{x}))\right] - \mathbb{E}_{\mathcal{S}\mathscr{T}}\left[\varphi(yh(\boldsymbol{x}))\right] \\
&= \mathbb{E}_{\mathcal{S}}\left[\varphi(yh(\boldsymbol{x}))\right] - \mathbb{E}_{\mathcal{S}\mathscr{T}}\left[\varphi(yh(\boldsymbol{x}))\right] \\
&= \frac{b_\varphi}{2m} \left\{ \begin{array}{c} \sum_{\sigma\in\Sigma_1} \sum_i \varphi(\sigma h((\mathrm{SF}^{\mathrm{a}})^{\top}\mathbf{1}_i \oplus (\mathrm{SF}^{\mathrm{s}})^{\top}\mathbf{1}_i)) \\ - \\ \sum_{\sigma\in\Sigma_1} \sum_i \varphi(\sigma h((\mathrm{SF}^{\mathrm{a}})^{\top}\mathbf{1}_i \oplus (\mathrm{MSF}^{\mathrm{s}})^{\top}\mathbf{1}_i)) \end{array} \right\} \\
&= \frac{b_\varphi}{2m} \cdot \mathbb{E}_{\boldsymbol{\sigma}\sim\Sigma_m} \left[\begin{array}{c} \sum_i \varphi(\sigma_i h((\mathrm{SF}^{\mathrm{a}})^{\top}\mathbf{1}_i \oplus (\mathrm{SF}^{\mathrm{s}})^{\top}\mathbf{1}_i)) \\ - \\ \sum_i \varphi(\sigma_i h((\mathrm{SF}^{\mathrm{a}})^{\top}\mathbf{1}_i \oplus (\mathrm{MSF}^{\mathrm{s}})^{\top}\mathbf{1}_i)) \end{array} \right] \qquad (31) \\
&\leq \frac{1}{2m} \cdot \mathbb{E}_{\boldsymbol{\sigma}\sim\Sigma_m} \left[\left| \sum_i \sigma_i h((\mathrm{SF}^{\mathrm{a}})^{\top}\mathbf{1}_i \oplus (\mathrm{SF}^{\mathrm{s}})^{\top}\mathbf{1}_i) - \sigma_i h((\mathrm{SF}^{\mathrm{a}})^{\top}\mathbf{1}_i \oplus (\mathrm{MSF}^{\mathrm{s}})^{\top}\mathbf{1}_i) \right| \right] \qquad (32) \\
&= \frac{1}{2m} \cdot \mathbb{E}_{\boldsymbol{\sigma}\sim\Sigma_m} \left[\left| \sum_i \sigma_i \left\{ \begin{array}{c} (h_{\mathrm{a}}((\mathrm{SF}^{\mathrm{a}})^{\top}\mathbf{1}_i) - h_{\mathrm{a}}((\mathrm{SF}^{\mathrm{a}})^{\top}\mathbf{1}_i)) \\ + \\ (h_{\mathrm{s}}((\mathrm{SF}^{\mathrm{s}})^{\top}\mathbf{1}_i) - h_{\mathrm{s}}((\mathrm{MSF}^{\mathrm{s}})^{\top}\mathbf{1}_i)) \end{array} \right\} \right| \right] \qquad (33) \\
&= \frac{1}{2m} \cdot \mathbb{E}_{\boldsymbol{\sigma}\sim\Sigma_m} \left[\left| \sum_i \sigma_i (h_{\mathrm{s}}((\mathrm{SF}^{\mathrm{s}})^{\top}\mathbf{1}_i) - h_{\mathrm{s}}((\mathrm{MSF}^{\mathrm{s}})^{\top}\mathbf{1}_i)) \right| \right] \ . \qquad (34)
\end{aligned}$$

Eq. (31) holds because $\boldsymbol{\sigma}$ is Rademacher. Ineq. (32) holds because F_φ is $(1/b_\varphi)$-Lipschitz [Nock and Nielsen, 2008]. Eq. (33) holds because of assumption **(i)**. We thus get the following upperbound for B in ineq. (16):

$$\begin{aligned}
&\mathbb{E}_{\mathcal{S}\sim\mathcal{D}} \left[\sup_{h\in\mathcal{H}} \left\{ \mathbb{E}_{\mathcal{S}}\left[\varphi(yh(\boldsymbol{x}))\right] - \mathbb{E}_{\mathcal{S}\mathscr{T}}\left[\varphi(yh(\boldsymbol{x}))\right] \right\} \right] \\
&\leq \mathbb{E}_{\mathcal{S}\sim\mathcal{D}} \left[\sup_{h_{\mathrm{s}}} \mathbb{E}_{\boldsymbol{\sigma}\sim\Sigma_m} \left[\left| \frac{1}{m} \sum_i \sigma_i (h_{\mathrm{s}}((\mathrm{SF}^{\mathrm{s}})^{\top}\mathbf{1}_i) - h_{\mathrm{s}}((\mathrm{MSF}^{\mathrm{s}})^{\top}\mathbf{1}_i)) \right| \right] \right] \\
&\leq \mathbb{E}_{\mathcal{S}\sim\mathcal{D},\boldsymbol{\sigma}\sim\Sigma_m} \left[\sup_{h_{\mathrm{s}}} \left| \frac{1}{m} \sum_i \sigma_i (h_{\mathrm{s}}((\mathrm{SF}^{\mathrm{s}})^{\top}\mathbf{1}_i) - h_{\mathrm{s}}((\mathrm{MSF}^{\mathrm{s}})^{\top}\mathbf{1}_i)) \right| \right] \ , \qquad (35)
\end{aligned}$$

where ineq. (35) holds because of the convexity of $\sup$. Using assumption **(ii)** ($h_{\mathrm{s}}(.) \in [0, K_{\mathrm{s}}]$), another use of the independent bounded differences inequality [McDiarmid, 1998] yield with prob-

ability $\geq 1-\delta_2$:

$$\mathbb{E}_{\mathcal{S}\sim\mathcal{D},\boldsymbol{\sigma}\sim\Sigma_m}\left[\sup_{h_{\mathrm{s}}}\left|\frac{1}{m}\sum_i \sigma_i(h_{\mathrm{s}}((\mathrm{SF}^{\mathrm{s}})^\top \mathbf{1}_i) - h_{\mathrm{s}}((\mathrm{MSF}^{\mathrm{s}})^\top \mathbf{1}_i))\right|\right]$$
$$\leq \mathbb{E}_{\boldsymbol{\sigma}\sim\Sigma_m}\left[\sup_{h_{\mathrm{s}}}\left|\frac{1}{m}\sum_i \sigma_i(h_{\mathrm{s}}((\mathrm{SF}^{\mathrm{s}})^\top \mathbf{1}_i) - h_{\mathrm{s}}((\mathrm{MSF}^{\mathrm{s}})^\top \mathbf{1}_i))\right|\right] + K_{\mathrm{s}}\cdot\sqrt{\frac{2}{m}\log\frac{1}{\delta_2}}\ . \quad (36)$$

We now put altogether ineqs. (15), (17) and (36) and obtain that with probability $\geq 1-(2\delta_1+\delta_2)$, we shall have

$$\begin{aligned}\mathbb{E}_{\mathcal{D}}\left[L_{0/1}(y,h(\boldsymbol{x}))\right] \leq\ & \mathbb{E}_{\mathcal{S}\mathscr{T}}\left[\varphi(yh(\boldsymbol{x}))\right] \\ & + \frac{4}{b_\varphi}\cdot\mathbb{E}_{\boldsymbol{\sigma}\sim\Sigma_m}\left[\sup_{h\in\mathcal{H}}\left|\frac{1}{m}\sum_i \sigma_i h\left(\boldsymbol{x}_i\right)\right|\right] \\ & + \mathbb{E}_{\boldsymbol{\sigma}\sim\Sigma_m}\left[\sup_{h_{\mathrm{s}}}\left|\frac{1}{m}\sum_i \sigma_i(h_{\mathrm{s}}((\mathrm{SF}^{\mathrm{s}})^\top \mathbf{1}_i) - h_{\mathrm{s}}((\mathrm{MSF}^{\mathrm{s}})^\top \mathbf{1}_i))\right|\right] \\ & + 2K_\varphi\cdot\sqrt{\frac{2}{m}\log\frac{1}{\delta_1}} + K_{\mathrm{s}}\cdot\sqrt{\frac{2}{m}\log\frac{1}{\delta_2}}\ . \end{aligned} \quad (37)$$

To simplify this expression, we fix $\delta_1=\delta_2=\delta/3$ and get with probability $\geq 1-\delta$,

$$\begin{aligned}\mathbb{E}_{\mathcal{D}}\left[L_{0/1}(y,h(\boldsymbol{x}))\right] \leq\ & \mathbb{E}_{\mathcal{S}\mathscr{T}}\left[\varphi(yh(\boldsymbol{x}))\right] \\ & + \mathbb{E}_{\boldsymbol{\sigma}\sim\Sigma_m}\left[\sup_{h_{\mathrm{s}}}\left|\frac{1}{m}\sum_i \sigma_i(h_{\mathrm{s}}((\mathrm{SF}^{\mathrm{s}})^\top \mathbf{1}_i) - h_{\mathrm{s}}((\mathrm{MSF}^{\mathrm{s}})^\top \mathbf{1}_i))\right|\right] \\ & + \frac{4}{b_\varphi}\cdot\mathbb{E}_{\boldsymbol{\sigma}\sim\Sigma_m}\left[\sup_{h\in\mathcal{H}}\left|\frac{1}{m}\sum_i \sigma_i h\left(\boldsymbol{x}_i\right)\right|\right] + (2K_\varphi+K_{\mathrm{s}})\cdot\sqrt{\frac{2}{m}\log\frac{3}{\delta}} \\ & = \mathbb{E}_{\mathcal{S}\mathscr{T}}\left[\varphi(yh(\boldsymbol{x}))\right] + \mathrm{RCP}_{\mathscr{T}}(\mathcal{H}) + \frac{4}{b_\varphi}\cdot\mathbb{E}_{\boldsymbol{\sigma}\sim\Sigma_m}\left[\sup_{h\in\mathcal{H}}\left|\frac{1}{m}\sum_i \sigma_i h\left(\boldsymbol{x}_i\right)\right|\right] \\ & + (2K_\varphi+K_{\mathrm{s}})\cdot\sqrt{\frac{2}{m}\log\frac{3}{\delta}}\ , \end{aligned}$$

from which we obtain the statement of Theorem 7.

9.3 Example of domain and CP for which (min risk over $\mathcal{S}^{\mathscr{T}}$ + Rademacher CP complexity) is strictly smaller than (min risk over $\mathcal{S}$)

We exhibit a toy domain which shows that

$$\min_h \mathbb{E}_{\mathcal{S}\mathscr{T}}\left[\varphi(yh(\boldsymbol{x}))\right] + \mathrm{RCP}_{\mathscr{T}}(\mathcal{H}) \ <\ \min_h \mathbb{E}_{\mathcal{S}}\left[\varphi(yh(\boldsymbol{x}))\right]\ , \quad (38)$$

for φ = square loss. Let $\mathcal{X} = \mathbb{R}^2$, with $\mathcal{S}$ consisting of 2 copies of observation $(0,0)$ (positive), 2 copies of observation $(1,1)$ (positive), and 1 copy of observation $(-1,-1)$ (negative). We enumerate the examples in $\mathcal{S}$ in this order. The CP satisfies $\mathrm{F}^{\mathrm{a}} \doteq \{x\}$, $\mathrm{F}^{\mathrm{s}} \doteq \{y\}$ and

$$\mathbf{M} \doteq \begin{bmatrix} 0 & 0 & 1 & 0 & 0 \\ 0 & 0 & 0 & 1 & 0 \\ 1 & 0 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 1 \end{bmatrix} ,$$

so that $\mathcal{S}^{\mathscr{T}}$ consists of two copies of observation $(1,0)$ (positive), two copies of $(0,1)$ (positive) and one copy of $(-1,-1)$ (the same observation as in $\mathcal{S}$). Let $h \doteq \boldsymbol{\theta}$, with its coordinates denoted x and y. The square loss L over $\mathcal{S}$ equals:

$$L = \frac{1}{5} \cdot \left(2 + 2(1-x-y)^2 + (1-x-y)^2\right) ,$$

which is minimized for $x = y = 1/2$ and yields $L = 2/5$. The square loss $L^{\mathscr{T}}$ over $\mathcal{S}^{\mathscr{T}}$ equals:

$$L^{\mathscr{T}} = \frac{1}{5} \cdot \left(2(1-x)^2 + 2(1-y)^2 + (1-x-y)^2\right) , \tag{39}$$

which is minimized for $x = y = 3/4$ and yields $L^{\mathscr{T}} = 1/10$. Assuming all linear separators have ℓ_∞ norm bounded by $3/4$ (which allows to have both solutions above), the RCP is

$$\begin{aligned} \mathrm{RCP}_{\mathscr{T}}(\mathcal{H}) &\doteq \mathbb{E}_{\boldsymbol{\sigma}\sim\Sigma_m}\left[\sup_{h\in\mathcal{H}_{\mathrm{s}}}\left|\frac{1}{m}\sum_i \sigma_i \left(h((\mathrm{SF}^{\mathrm{s}})^\top \mathbf{1}_i) - h((\mathrm{MSF}^{\mathrm{s}})^\top \mathbf{1}_i)\right)\right|\right] \\ &= \frac{1}{5}\cdot\frac{3}{4}\cdot\frac{1}{16}\cdot\sum_{\boldsymbol{\sigma}\in\Sigma_4}|-\sigma_1-\sigma_2+\sigma_3+\sigma_4| \\ &= \frac{3}{20}\cdot\frac{1}{16}\cdot\sum_{\boldsymbol{\sigma}\in\Sigma_4}\left|\mathbf{1}^\top\boldsymbol{\sigma}\right| \\ &= \frac{3}{20}\cdot\frac{1}{16}\cdot(4\cdot 2 + 2\cdot 8 + 0\cdot 6) \qquad (40) \\ &= \frac{3}{20}\cdot\frac{24}{16} = \frac{9}{40} . \qquad (41) \end{aligned}$$

We then check that

$$L^{\mathscr{T}} + \mathrm{RCP}_{\mathscr{T}}(\mathcal{H}) = \frac{13}{40} < \frac{2}{5} = L , \tag{42}$$

as claimed.

9.4 Proof of Theorem 9

The proof of Theorem 9 follows from the proof of a more general Theorem that we prove here. We say that $(\mathrm{M}, \mathrm{K}^{\mathrm{s}})$ satisfies the (γ,δ)-correlation assumption for some $0 < \delta, \gamma \leq 1$ iff the following two assumptions hold:

(a) $((\mathrm{I}_m - \mathrm{M})\mathrm{K}^{\mathrm{s}}(\mathrm{I}_m - \mathrm{M})^\top)_{ii} \geq (1-\delta)\cdot(1/m)\cdot \mathrm{tr}\left((\mathrm{I}_m - \mathrm{M})\mathrm{K}^{\mathrm{s}}(\mathrm{I}_m - \mathrm{M})^\top\right), \forall i \in [m]$;

(b) $|((\mathrm{I}_m - \mathrm{M})\mathrm{K}^{\mathrm{s}}(\mathrm{I}_m - \mathrm{M})^\top)_{ii'}/\sqrt{((\mathrm{I}_m - \mathrm{M})\mathrm{K}^{\mathrm{s}}(\mathrm{I}_m - \mathrm{M})^\top)_{ii}((\mathrm{I}_m - \mathrm{M})\mathrm{K}^{\mathrm{s}}(\mathrm{I}_m - \mathrm{M})^\top)_{i'i'}}| \geq 1-\gamma, \forall i, i' \in [m]$.

Regardless of $\mathcal{S}$, there always exist $0 < \delta, \gamma \leq 1$ for which this holds, but the bound may be quantitatively better when at least one is small.

Theorem 18 *Using notations of Theorem 9, there exists $\epsilon > 0$ such that for any $\mathcal{S}$ and $\mathscr{T}$ for which $(\mathrm{M}, \mathrm{K}^{\mathrm{s}})$ satisfies the (γ, δ)-correlation assumption, we have*

$$\mathrm{RCP}_{\mathscr{T}}(\mathcal{H}) \quad \leq \quad u \cdot \frac{r_{\mathrm{s}}}{\sqrt{m}} \cdot \sqrt{\frac{1}{m}\cdot \langle \mathrm{I}_m, \mathrm{K}^{\mathrm{s}}\rangle_{\mathrm{M}} \mathrm{I}_m} \ . \tag{43}$$

with

$$u \quad \doteq \quad \frac{1}{m} + \kappa(\epsilon)\left(1 - \frac{1}{m}\right) \ , \tag{44}$$

and $\kappa(\epsilon) = 1 - ((1-\delta)(1-\epsilon)(1-\gamma))^2 \in (0,1)$.

Furthermore,

$$\frac{1}{m}\cdot \langle \mathrm{I}_m, \mathrm{K}^{\mathrm{s}}\rangle_{\mathrm{M}} \quad = \quad 2 \cdot \sum_{j \in [d_{\mathrm{s}}]} \mathbb{V}(\mathrm{SF}^{\mathrm{s}}\mathbf{1}_j)(1 - \rho(\mathrm{SF}^{\mathrm{s}}\mathbf{1}_j, \mathrm{MSF}^{\mathrm{s}}\mathbf{1}_j)) \ . \tag{45}$$

Proof We observe that $\forall \boldsymbol{\sigma} \in \Sigma_m$,

$$\begin{aligned} &\arg \sup_{\boldsymbol{\theta} \in \mathbb{R}^{d_{\mathrm{s}}} : \|\boldsymbol{\theta}\|_2 \leq r_{\mathrm{s}}} \left| \frac{1}{m} \sum_i \sigma_i \boldsymbol{\theta}^\top \left(((\mathrm{I}_m - \mathrm{M})\mathrm{SF}^{\mathrm{s}})^\top \mathbf{1}_i\right)\right| \\ &= \quad \frac{r_{\mathrm{s}}}{\|\sum_i \sigma_i ((\mathrm{I}_m - \mathrm{M})\mathrm{SF}^{\mathrm{s}})^\top \mathbf{1}_i\|_2} \sum_i \sigma_i ((\mathrm{I}_m - \mathrm{M})\mathrm{SF}^{\mathrm{s}})^\top \mathbf{1}_i \ , \end{aligned}$$

and so:

$$\begin{aligned} \mathrm{RCP}_{\mathscr{T}}(\mathcal{H}) \quad &= \quad \mathbb{E}_{\boldsymbol{\sigma}\sim\Sigma_m}\left[\sup_{h \in \mathcal{H}_{\mathrm{s}}} \left| \frac{1}{m}\sum_i \sigma_i \left(h((\mathrm{SF}^{\mathrm{s}})^\top \mathbf{1}_i) - h((\mathrm{MSF}^{\mathrm{s}})^\top \mathbf{1}_i)\right)\right|\right] \qquad (46) \\ &= \quad \frac{r_{\mathrm{s}}}{m}\cdot \mathbb{E}_{\boldsymbol{\sigma}\sim\Sigma_m}\left[\left\|\sum_i \sigma_i ((\mathrm{I}_m - \mathrm{M})\mathrm{SF}^{\mathrm{s}})^\top \mathbf{1}_i\right\|_2\right] \\ &= \quad \frac{r_{\mathrm{s}}}{m}\cdot \sqrt{\sum_i \|((\mathrm{I}_m - \mathrm{M})\mathrm{SF}^{\mathrm{s}})^\top \mathbf{1}_i\|_2^2} \\ &\quad \cdot \mathbb{E}_{\boldsymbol{\sigma}\sim\Sigma_m}\left[\sqrt{1 + \frac{\sum_{i\neq i'} \sigma_i\sigma_{i'} \mathbf{1}_i^\top (\mathrm{I}_m - \mathrm{M})\mathrm{SF}^{\mathrm{s}}(\mathrm{SF}^{\mathrm{s}})^\top (\mathrm{I}_m - \mathrm{M})^\top \mathbf{1}_{i'}}{\sum_i \|((\mathrm{I}_m - \mathrm{M})\mathrm{SF}^{\mathrm{s}})^\top \mathbf{1}_i\|_2^2}}\right] \\ &\doteq \quad \frac{r_{\mathrm{s}}}{m}\cdot \sqrt{\sum_i \|((\mathrm{I}_m - \mathrm{M})\mathrm{SF}^{\mathrm{s}})^\top \mathbf{1}_i\|_2^2}\cdot \mathbb{E}_{\boldsymbol{\sigma}\sim\Sigma_m}\left[\sqrt{1 + u(\boldsymbol{\sigma})}\right] \ , \qquad (47) \end{aligned}$$

with

$$u(\boldsymbol{\sigma}) \doteq \frac{\sum_{i\neq i'}\sigma_i\sigma_{i'}\mathbf{1}_i^\top(\mathrm{I}_m-\mathrm{M})\mathrm{SF}^{\mathrm{s}}(\mathrm{SF}^{\mathrm{s}})^\top(\mathrm{I}_m-\mathrm{M})^\top\mathbf{1}_{i'}}{\sum_i\|((\mathrm{I}_m-\mathrm{M})\mathrm{SF}^{\mathrm{s}})^\top\mathbf{1}_i\|_2^2} \ . \tag{48}$$

Let us call for short $\boldsymbol{\delta}_i \doteq ((\mathrm{I}_m-\mathrm{M})\mathrm{SF}^{\mathrm{s}})^\top\mathbf{1}_i$, so that eq. (48) can be simplified to $u(\boldsymbol{\sigma}) = (\sum_i\|\boldsymbol{\delta}_i\|_2^2)^{-1}\sum_{i\neq i'}\sigma_i\sigma_{i'}\boldsymbol{\delta}_i^\top\boldsymbol{\delta}_{i'}$. $\forall n\in\mathbb{N}_*$, we have

$$\begin{aligned}
&\mathbb{E}_{\boldsymbol{\sigma}\sim\Sigma_m}\left[u^n(\boldsymbol{\sigma})\right] \\
&= \frac{1}{\left(\sum_i\|\boldsymbol{\delta}_i\|_2^2\right)^n}\cdot\mathbb{E}_{\boldsymbol{\sigma}\sim\Sigma_m}\left[\sum_{i_1\neq i'_1}\sum_{i_2\neq i'_2}\cdots\sum_{i_n\neq i'_n}\prod_{k=1}^n\sigma_{i_k}\sigma_{i'_k}\boldsymbol{\delta}_{i_k}^\top\boldsymbol{\delta}_{i'_k}\right] \\
&= \frac{1}{\left(\sum_i\|\boldsymbol{\delta}_i\|_2^2\right)^n}\cdot\sum_{i_1\neq i'_1}\sum_{i_2\neq i'_2}\cdots\sum_{i_n\neq i'_n}\mathbb{E}_{\boldsymbol{\sigma}\sim\Sigma_m}\left[\prod_{k=1}^n\sigma_{i_k}\sigma_{i'_k}\boldsymbol{\delta}_{i_k}^\top\boldsymbol{\delta}_{i'_k}\right] \\
&= \frac{1}{\left(\sum_i\|\boldsymbol{\delta}_i\|_2^2\right)^n}\cdot\sum_{i_1\neq i'_1}\sum_{i_2\neq i'_2}\cdots\sum_{i_n\neq i'_n}\prod_{(i,i')\in\{(i_k,i'_k)\}_{k=1}^n}\mathbb{E}_{\boldsymbol{\sigma}\sim\Sigma_m}\left[(\sigma_i\sigma_{i'})^{n(i,i')}\right]\left(\boldsymbol{\delta}_i^\top\boldsymbol{\delta}_{i'}\right)^{n(i,i')} \ ,
\end{aligned} \tag{49}$$

with $n(i,i') \doteq |\{k:(i,i')=(i_k,i'_k)\}|$ satisfying $\sum n(i,i') = n$. Whenever $n(i,i')$ is odd, $\mathbb{E}_{\boldsymbol{\sigma}\sim\Sigma_m}\left[(\sigma_i\sigma_{i'})^{n(i,i')}\right] = 0$ (because $\boldsymbol{\sigma}$ is Rademacher), and it is 1 otherwise. We get, if n is even:

$$\begin{aligned}
&\mathbb{E}_{\boldsymbol{\sigma}\sim\Sigma_m}\left[u^n(\boldsymbol{\sigma})\right] \\
&= \frac{1}{\left(\sum_i\|\boldsymbol{\delta}_i\|_2^2\right)^n}\cdot\underbrace{\sum_{0<\ell\leq n}\ \sum_{\substack{\{n_k\}_{k=1}^\ell\subset\mathbb{N}_* \\ \text{s.t. } 2\sum n_k = n}}\ \sum_{\substack{\{(i_k,i'_k)\}_{k=1}^\ell \\ \text{s.t. } i_k\neq i'_k,\forall k}}\prod_{k=1}^\ell\left(\boldsymbol{\delta}_{i_k}^\top\boldsymbol{\delta}_{i'_k}\right)^{2n_k}}_{\doteq\zeta(n)} \ ,
\end{aligned} \tag{50}$$

and $\mathbb{E}_{\boldsymbol{\sigma}\sim\Sigma_m}\left[u^n(\boldsymbol{\sigma})\right] = 0$ if n is odd. Since

$$\sqrt{1+x} = 1+\sum_{n\in\mathbb{N}_*}\frac{1}{2^n n!}\cdot\prod_{k=0}^{n-1}(1-2k)x^n \ , \tag{51}$$

we get after combining with eqs (47) and (50) and using the definition of $\zeta(.)$ in eq. (50):

$$\begin{aligned}
\text{RCP}_{\mathscr{T}}(\mathcal{H}) &= \frac{r_{\mathrm{s}}}{m}\cdot\sqrt{\sum_i\|\boldsymbol{\delta}_i\|_2^2}\cdot\left(1-\sum_{n\in\mathbb{N}_*}\frac{\prod_{k=1}^{2n-1}(2k-1)\cdot\zeta(2n)}{(2n)!\left(2\sum_i\|\boldsymbol{\delta}_i\|_2^2\right)^{2n}}\right) \\
&= \frac{r_{\mathrm{s}}}{m}\cdot\sqrt{\sum_i\|\boldsymbol{\delta}_i\|_2^2}\cdot\left(1-\sum_{n\in\mathbb{N}_*}\frac{\prod_{k=1}^{2n-1}(2k-1)}{(2m)^{2n}(2n)!}\cdot\tilde{\zeta}(2n)\right) \ ,
\end{aligned} \tag{52}$$

with:

$$\begin{aligned}
&\tilde{\zeta}(2n) \\
&\doteq \sum_{0<\ell\leq 2n}\ \sum_{\substack{\{n_k\}_{k=1}^\ell\subset\mathbb{N}_* \\ \text{s.t. } \sum n_k = n}}\ \sum_{\substack{\{(i_k,i'_k)\}_{k=1}^\ell \\ \text{s.t. } i_k\neq i'_k,\forall k}}\prod_{k=1}^\ell\left(\begin{array}{c}\frac{\left\|\boldsymbol{\delta}_{i_k}\right\|_2^2}{\frac{1}{m}\cdot\sum_p\|\boldsymbol{\delta}_p\|_2^2} \\ \cdot\frac{\left\|\boldsymbol{\delta}_{i'_k}\right\|_2^2}{\frac{1}{m}\cdot\sum_p\|\boldsymbol{\delta}_p\|_2^2}\cdot\gamma_{i_k,i'_k}\end{array}\right)^{2n_k} \ ,
\end{aligned} \tag{53}$$

and $\gamma_{i,i'} \doteq \cos(\boldsymbol{\delta}_i, \boldsymbol{\delta}_{i'})$. Remark that eq. (52) is an equality. We now use assumption **(a)** and obtain

$$\tilde{\zeta}(2n) \;\geq\; (1-\delta)^{2n} \sum_{0<\ell\leq 2n} \sum_{\substack{\{n_k\}_{k=1}^{\ell} \subset \mathbb{N}_* \\ \text{s.t. } \sum n_k = n}} \sum_{\substack{\{(i_k, i'_k)\}_{k=1}^{\ell} \\ \text{s.t. } i_k \neq i'_k, \forall k}} \prod_{k=1}^{\ell} \left(\gamma_{i_k,i'_k}\right)^{2n_k} \;. \tag{54}$$

Denote for short $\mathcal{U}(n)$ the set of eligible triples (n_k, i_k, i'_k) in the summation. We get because of assumption **(b)** $\tilde{\zeta}(2n) \geq |\mathcal{U}(n)|((1-\delta)(1-\gamma))^{2n}$, and so, using the shorthand

$$\mathscr{L} \;\doteq\; \frac{1}{m} \cdot \sum_i \|\boldsymbol{\delta}_i\|_2^2 \;, \tag{55}$$

we obtain our first upperbound,

$$\text{RCP}_{\mathscr{T}}(\mathcal{H}) \;\leq\; \frac{r_{\text{s}}}{\sqrt{m}} \cdot \sqrt{\mathscr{L}} \cdot \left(1 - \sum_{n\in\mathbb{N}_*} |\mathcal{U}(n)| \left(\frac{(1-\delta)(1-\gamma)}{m}\right)^{2n} \cdot \frac{\prod_{k=1}^{2n-1}(2k-1)}{2^{2n}(2n)!}\right) \tag{56}$$

Lemma 19 *There exists a constant $\epsilon > 0$ such that $\forall n \in \mathbb{N}_*$,*

$$\frac{\prod_{k=1}^{2n-1}(2k-1)}{(2n)!} \;\geq\; (2(1-\epsilon))^{2n} \;. \tag{57}$$

Proof We proceed by induction, letting $g_\epsilon(n) \doteq (2(1-\epsilon))^{2n}$ and

$$f(n) \;\doteq\; \frac{\prod_{k=1}^{2n-1}(2k-1)}{(2n)!} \;, \tag{58}$$

and for $\epsilon = \epsilon_* \doteq 1 - 1/(2\sqrt{2})$. We remark that $g_{\epsilon_*}(1) = f(1)$. Furthermore,

$$\begin{aligned} f(n+1) = \frac{\prod_{k=1}^{2(n+1)-1}(2k-1)}{(2(n+1))!} \;&=\; \frac{(2n-1)(2n+1)}{(n+1)(n+2)} \cdot \frac{\prod_{k=1}^{2n-1}(2k-1)}{(2n)!} \\ &\geq\; \frac{(2n-1)(2n+1)}{(n+1)(n+2)} \cdot (2(1-\epsilon))^{2n} \qquad (59) \\ &=\; \frac{(2n-1)(2n+1)}{(n+1)(n+2)(2(1-\epsilon))^2} \cdot g_\epsilon(n+1) \;, \end{aligned}$$

where ineq. (59) uses the induction hypothesis. We prove the Lemma once we prove that the factor on the right is at least 1, that is, for $\epsilon = \epsilon_*$, we need to prove

$$\frac{(2n-1)(2n+1)}{(n+1)(n+2)} \;\geq\; \frac{1}{2} \;, \tag{60}$$

which is indeed the case since the left function is strictly increasing over $\mathbb{N}$ and equals the right-hand side for $n = 1$. ■

So we get from ineq. (56):

$$\begin{aligned}
\mathrm{RCP}_{\mathscr{T}}(\mathcal{H}) &\leq \frac{r_{\mathrm{s}}}{\sqrt{m}} \cdot \sqrt{\mathscr{L}} \cdot \left(1 - \sum_{n \in \mathbb{N}_*} |\mathcal{U}(n)| \left(\frac{(1-\delta)(1-\gamma)(1-\epsilon)}{m}\right)^{2n}\right) \\
&\leq \frac{r_{\mathrm{s}}}{\sqrt{m}} \cdot \sqrt{\mathscr{L}} \cdot \left(1 - u_*(m) \sum_{n \in \mathbb{N}_*} \left(\frac{(1-\delta)(1-\gamma)(1-\epsilon)}{m}\right)^{2n}\right) \quad , \qquad (61)
\end{aligned}$$

where $u_*(m)$ satisfies $u_*(m) \leq \min_n |\mathcal{U}(n)|$. We finally get:

$$\begin{aligned}
\mathrm{RCP}_{\mathscr{T}}(\mathcal{H}) &\leq \frac{r_{\mathrm{s}}}{\sqrt{m}} \cdot \sqrt{\mathscr{L}} \cdot \left(1 - (1-\kappa(\epsilon)) \cdot \frac{u_*(m)}{m^2 - (1-\kappa(\epsilon))}\right) \\
&\leq \frac{r_{\mathrm{s}}}{\sqrt{m}} \cdot \sqrt{\mathscr{L}} \cdot \left(1 - (1-\kappa(\epsilon)) \cdot \frac{u_*(m)}{m^2}\right) \\
&\leq \frac{r_{\mathrm{s}}}{\sqrt{m}} \cdot \sqrt{\mathscr{L}} \cdot \left(\frac{1}{m} + \kappa(\epsilon)\left(1 - \frac{1}{m}\right)\right) \quad , \qquad (62)
\end{aligned}$$

with $\kappa(\epsilon) = 1 - ((1-\delta)(1-\epsilon)(1-\gamma))^2 > 0$, since $u_*(m) \geq m(m-1)$ (obtained for $\ell = 2n$ in eq. (54)). We finish the proof by remarking that $\mathscr{L}$ in eq. (55) satisfies

$$\begin{aligned}
\mathscr{L} &= \frac{1}{m} \cdot \sum_i \mathbf{1}_i^\top (\mathrm{I}_m - \mathrm{M}) \mathrm{SF}^{\mathrm{s}} (\mathrm{SF}^{\mathrm{s}})^\top (\mathrm{I}_m - \mathrm{M})^\top \mathbf{1}_i \\
&= \frac{1}{m} \cdot \mathrm{tr}\left((\mathrm{I}_m - \mathrm{M}) \mathrm{SF}^{\mathrm{s}} (\mathrm{SF}^{\mathrm{s}})^\top (\mathrm{I}_m - \mathrm{M})^\top\right) \\
&= \frac{1}{m} \cdot \mathrm{tr}\left((\mathrm{I}_m - \mathrm{M}) \mathrm{K}^{\mathrm{s}} (\mathrm{I}_m - \mathrm{M})^\top\right) \\
&= \frac{1}{m} \cdot \mathrm{tr}\left((\mathrm{I}_m - \mathrm{M})^\top \mathrm{I}_m (\mathrm{I}_m - \mathrm{M}) \mathrm{K}^{\mathrm{s}}\right) \\
&= \frac{1}{m} \cdot \langle \mathrm{I}_m, \mathrm{K}^{\mathrm{s}} \rangle_{\mathbf{M}} \quad . \qquad (63)
\end{aligned}$$

Ineq. (62) and eq. (63) allow to conclude the proof of Theorem 18 with

$$u \doteq \frac{1}{m} + \kappa(\epsilon)\left(1 - \frac{1}{m}\right) \quad , \qquad (64)$$

which, since $\kappa(\epsilon) < 1$, satisfies indeed $u \in (0,1)$. This achieves the main part of the proof of Theorem 18. To prove eq. (45), we just have to write (letting $\varsigma : [m] \to [m]$ represent the

corresponding permutation),

$$
\begin{aligned}
&\frac{1}{m}\cdot\langle \mathrm{I}_m, \mathrm{K}^{\mathrm{s}}\rangle_{\mathbf{M}} \\
&= \frac{1}{m}\sum_i \left\|\boldsymbol{x}_i^{\mathrm{s}} - \boldsymbol{x}_{\varsigma(i)}^{\mathrm{s}}\right\|_2^2 \\
&= \sum_{j\in[d_{\mathrm{s}}]}\frac{1}{m}\cdot\sum_i (x_{ij}^{\mathrm{s}} - x_{\varsigma(i)j}^{\mathrm{s}})^2 \\
&= 2\cdot\sum_{j\in[d_{\mathrm{s}}]}\left\{\frac{1}{m}\cdot\sum_i (x_{ij}^{\mathrm{s}})^2 - \frac{1}{m}\cdot\sum_i x_{ij}^{\mathrm{s}}x_{\varsigma(i)j}^{\mathrm{s}}\right\} \qquad (65)\\
&= 2\cdot\sum_{j\in[d_{\mathrm{s}}]}\left\{\frac{1}{m}\cdot\sum_i (x_{ij}^{\mathrm{s}})^2 - \left(\frac{1}{m}\cdot\sum_i x_{ij}^{\mathrm{s}}\right)^2 + \left(\frac{1}{m}\cdot\sum_i x_{ij}^{\mathrm{s}}\right)^2 - \frac{1}{m}\cdot\sum_i x_{ij}^{\mathrm{s}}x_{\varsigma(i)j}^{\mathrm{s}}\right\} \\
&= 2\cdot\sum_{j\in[d_{\mathrm{s}}]}\left\{\mathbb{V}(\mathrm{SF}^{\mathrm{s}}\mathbf{1}_j) - \mathbb{C}\mathrm{ov}(\mathrm{SF}^{\mathrm{s}}\mathbf{1}_j, \mathrm{MSF}^{\mathrm{s}}\mathbf{1}_j)\right\} \\
&= 2\cdot\sum_{j\in[d_{\mathrm{s}}]}\mathbb{V}(\mathrm{SF}^{\mathrm{s}}\mathbf{1}_j)\left\{1 - \frac{\mathbb{C}\mathrm{ov}(\mathrm{SF}^{\mathrm{s}}\mathbf{1}_j, \mathrm{MSF}^{\mathrm{s}}\mathbf{1}_j)}{\sqrt{\mathbb{V}(\mathrm{SF}^{\mathrm{s}}\mathbf{1}_j)}\sqrt{\mathbb{V}(\mathrm{MSF}^{\mathrm{s}}\mathbf{1}_j)}}\right\} \qquad (66)\\
&= 2\cdot\sum_{j\in[d_{\mathrm{s}}]}\mathbb{V}(\mathrm{SF}^{\mathrm{s}}\mathbf{1}_j)(1-\rho(\mathrm{SF}^{\mathrm{s}}\mathbf{1}_j, \mathrm{MSF}^{\mathrm{s}}\mathbf{1}_j))\ ,
\end{aligned}
$$

where eq. (66) follows from the fact that $\mathbb{V}(\mathrm{SF}^{\mathrm{s}}\mathbf{1}_j) = \mathbb{V}(\mathrm{MSF}^{\mathrm{s}}\mathbf{1}_j)$. ■

Remark: it is worthwhile remarking that the proof of Theorem 9 can also be applied to upperbound the empirical Rademacher complexity of linear functions, without modifications, except for the handling of $\mathscr{L}$. In this case, the proof improves the upperbound known [Kakade et al., 2008] (Theorem 1) by factor u in eq. (64). This is due to the fact that the proof in [Kakade et al., 2008] takes into account only the maximum norm in the observations of $\mathcal{S}$, and not the angles between the observations. We now state the corresponding Theorem.

Theorem 20 *Following [Gretton et al., 2007], we let $\mathbf{i}_r^m$ denote the set of r-tuples drawn without replacement drawn from $[m]$. Suppose $\mathcal{S}$ satisfies the following for some $\delta, \gamma > 0$ and $r_x > 0$:*

(a) $\|\boldsymbol{x}_i\|_2^2 \geq (1-\delta)\cdot(1/m)\sum_{i'}\|\boldsymbol{x}_{i'}\|_2^2$, $\forall i \in [m]$;

(b) $\mathbb{E}_{(i,i')\sim\mathbf{i}_2^m}[|\cos(\boldsymbol{x}_i, \boldsymbol{x}_{i'})|] \geq 1-\gamma$;

(c) $\|\boldsymbol{x}_i\|_2 \leq r_x$, $\forall i \in [m]$.

Then, assuming that $\mathcal{H}$ contains linear classifiers of the form $\boldsymbol{\theta}^\top\boldsymbol{x}$ with $\|\boldsymbol{\theta}\|_2 \leq r_\theta$, there exists $\epsilon > 0$ such that the empirical Rademacher complexity of $\mathcal{H}$ satisfies:

$$
\mathrm{R}_{\mathcal{S}}(\mathcal{H}) \ \leq\ \left(\frac{1}{m} + \kappa(\epsilon)\left(1-\frac{1}{m}\right)\right)\cdot\frac{r_x r_\theta}{\sqrt{m}}\ , \qquad (67)
$$

with $\kappa(\epsilon) = 1 - ((1-\delta)(1-\epsilon)(1-\gamma))^2 \in (0,1)$.




Figure 3: A permutation ς defines an oriented graph whose vertices are examples and permutation ς defines arcs (here, $\varsigma(1) = 2$ for example; black dots are positive examples, red dots are negative examples). Each term in the $\sup$ of eq. (68) is a weighted cut (one weighted cut for each $\boldsymbol{\sigma} \in \{-1, 1\}^m$): the squares depict the examples for which $w(.) \neq 0$ in Lemma 21 for the $\boldsymbol{\sigma}$ displayed. In this example of $\boldsymbol{\sigma}$, only two examples out of the six would bring a non-zero weight $w(.)$.

[Kakade et al., 2008]'s proof relies on **(c)**. Since **(a)** and **(b)** can always be satisfied for some $\delta, \gamma > 0$, ineq. (43) holds under their setting as well; however, it becomes better than theirs as both δ, γ are small, so in particular in the case where observations start to be heavily correlated and be of approximately the same norm. Indeed, in this case, $\sum_i \sigma_i \boldsymbol{x}_i$ will often have *small* magnitude, because $\Sigma_m \ni \boldsymbol{\sigma} \sim \{-1, 1\}$ and thus many vectors will approximately cancel through the sum in many draws of $\boldsymbol{\sigma}$.

9.5 Proof of Theorem 10

We first start by a Lemma which shows that indeed $\text{RCP}_{\mathscr{T}}(\mathcal{H})$ can be significantly smaller than a Rademacher complexity.

Lemma 21 *Suppose setting (B) holds in Theorem 7. Then*

$$\text{RCP}_{\mathscr{T}}(\mathcal{H}) \quad = \quad 2 \cdot \mathbb{E}_{\boldsymbol{\sigma} \sim \Sigma_m} \left[\sup_{h_{\text{s}} \in \mathcal{H}_{\text{s}}} \left| \frac{1}{m} \sum_i w(i) h_{\text{s}}((\text{SF}^{\text{s}})^\top \mathbf{1}_i) \right| \right] \quad , \tag{68}$$

where $w(i) \doteq (1/2) \cdot (\sigma_i - \sigma_{\varsigma^{-1}(i)}) \in \{-1, 0, 1\}$.

The proof of this Lemma is straightforward. What is interesting is, since $w(.)$ can take on zero values, to what extent $\text{RCP}_{\mathscr{T}}(\mathcal{H})$ can be smaller than the corresponding Rademacher complexity in which $w(.)$ would be replaced by $\boldsymbol{\sigma} \in \{-1, 1\}^m$, and what drives this reduction. Figure 3 displays qualitatively this intuition on a simple example. We now investigate a quantitative derivation of the reduction for DAG classifiers.

We let $\mathrm{cut}_\varsigma(\boldsymbol{\sigma}) \doteq \{i : \sigma_i \neq \sigma_{\varsigma^{-1}(i)}\}$. We simplify notations in the proof and drop notation b so that notation $h^i_{\mathrm{s}} \doteq h_{\mathrm{s}}((\mathbf{SF}^{\mathrm{s}})^\top \mathbf{1}_i)$ where $i \in [m]$. We let

$$\mu \doteq \mathbb{E}_{\boldsymbol{\sigma}\sim\Sigma_m}\left[\sup_{h_{\mathrm{s}}\in\mathcal{H}^{\mathrm{s}}} \sum_{i\in\mathrm{cut}_\varsigma(\boldsymbol{\sigma})} \sigma_i h^i_{\mathrm{s}}\right] \ , \tag{69}$$

which is a generalisation of $m \cdot \textsc{rcp}_{\mathscr{T}}(\mathcal{H})$ to any permutation $\varsigma \in S_m$, and not just a block-class permutation in S^*_m as assumed in Setting (B). We assume basic knowledge of Massart's finite class Lemma's proof. Using Jensen's inequality, we arrive, after the same chain of derivations, for any $t > 0$, to:

$$\exp(t\mu) \ \leq \ \frac{1}{2^m}\sum_{\boldsymbol{\sigma}\in\{-1,1\}^m} \sup_{h_{\mathrm{s}}\in\mathcal{H}^{\mathrm{s}}} \exp\left(t \cdot \sum_{i\in\mathrm{cut}_\varsigma(\boldsymbol{\sigma})} \sigma_i h^i_{\mathrm{s}}\right) \ . \tag{70}$$

The proof (of Massart's Lemma) now involves replacing the $\sup$ by a sum. Remark that when h is DAG, the $\sup$ implies that each h^j_{s} is in fact $\in \{\pm K_{\mathrm{s}}\}$. So let us use $\mathcal{H}^{\mathrm{s}}_+ \subseteq \mathcal{H}^{\mathrm{s}}$, the set of classifiers whose output is in $\{\pm K_{\mathrm{s}}\}$. We get:

$$\exp(t\mu) \ \leq \ \sum_{h_{\mathrm{s}}\in\mathcal{H}^{\mathrm{s}}_+} \frac{1}{2^m}\sum_{\boldsymbol{\sigma}\in\{-1,1\}^m} \exp\left(t \cdot \sum_{i\in\mathrm{cut}_\varsigma(\boldsymbol{\sigma})} \sigma_i h^i_{\mathrm{s}}\right) \ . \tag{71}$$

To identify better permutations, we name in this proof $\varsigma \in S^*_m$ the permutation represented by M, so that we also have

$$\mathrm{odd_cycle}(\mathbf{M}) \ \doteq \ \mathrm{odd_cycle}(\varsigma) \ .$$

Since each coordinate of $\boldsymbol{\sigma}$ is chosen uniformly at random, cycles in a permutation are disjoint and $\exp(a+b) = \exp(a)\exp(b)$, the inner sigma in ineq. (71) factors over the cycles of permutation ς:

$$\begin{aligned}
&\sum_{h_{\mathrm{s}}\in\mathcal{H}^{\mathrm{s}}_+} \frac{1}{2^m}\sum_{\boldsymbol{\sigma}\in\{-1,1\}^m} \exp\left(t \cdot \sum_{i\in\mathrm{cut}_\varsigma(\boldsymbol{\sigma})} \sigma_i h^i_{\mathrm{s}}\right) \\
&= \ \frac{1}{2^m}\sum_{h_{\mathrm{s}}\in\mathcal{H}^{\mathrm{s}}_+} \prod_{U\in\mathrm{cycle}(\varsigma)} \left(\sum_{\boldsymbol{\sigma}\in\{-1,1\}^{|U|}} \exp\left(t \cdot \sum_{i\in\mathrm{cut}_\varsigma(\boldsymbol{\sigma})\cap U} \sigma_i h^U_i\right)\right) \ ,
\end{aligned} \tag{72}$$

where h^U indicates coordinates of h in U and $\mathrm{cycle}(\varsigma)$ is the set of cycles *without 1-cycles* (*i.e.* fixed points) — we have let m_ς denote the number of fixed points of ς. We have used the fact that cycles define a partition of $[m]$.

Now, whenever U contains an *odd* number of indexes, whatever $\boldsymbol{\sigma}$, the sum over $\mathrm{cut}_\varsigma(\boldsymbol{\sigma}) \cap U$ cannot cover the sum over U: there always remains at least one vertex which does not belong to the sum (In Figure 3, the red dot cycle displays this fact on an example). Let us denote $i_{\boldsymbol{\sigma}} \in [|U|]$

this vertex. Since $\exp(x) + \exp(-x) \geq 2 + x^2$ and $h_{\mathrm{s}} \in \mathcal{H}^{\mathrm{s}}_+$, we get:

$$\begin{aligned}
&\sum_{\boldsymbol{\sigma}\in\{-1,1\}^{|U|}} \exp\left(t \cdot \sum_{i\in \mathrm{cut}_\varsigma(\boldsymbol{\sigma})\cap U} \sigma_i h_i^U\right) \\
&\leq \frac{1}{2+t^2K_{\mathrm{s}}^2} \cdot \sum_{\boldsymbol{\sigma}\in\{-1,1\}^{|U|}} \left\{\exp\left(t \cdot \sum_{i\in \mathrm{cut}_\varsigma(\boldsymbol{\sigma})\cap U} \sigma_i h_i^U\right) \cdot \left(\exp(th_{i_{\boldsymbol{\sigma}}}^U) + \exp(-th_{i_{\boldsymbol{\sigma}}}^U)\right)\right\} \quad (73) \\
&\leq \frac{2}{2+t^2K_{\mathrm{s}}^2} \cdot \sum_{\boldsymbol{\sigma}\in\{-1,1\}^{|U|}} \exp\left(t \cdot \sum_{i\in U} \sigma_i h_i^U\right) \ , \quad (74)
\end{aligned}$$

since the multiplication in ineq. (73) duplicates part of the terms in (74). We now plug this bound in eq. (71 — 72) and finish the derivation following Massart's finite class Lemma:

$$\begin{aligned}
\exp(t\mu) &\leq \left(\frac{2}{2+t^2K_{\mathrm{s}}^2}\right)^{|\mathrm{odd_cycle}(\varsigma)|} \cdot \frac{1}{2^m} \sum_{h_{\mathrm{s}}\in\mathcal{H}^{\mathrm{s}}_+} \sum_{\boldsymbol{\sigma}\in\{-1,1\}^m} \exp\left(t \cdot \sum_i \sigma_i h_i\right) \\
&= \left(\frac{2}{2+t^2K_{\mathrm{s}}^2}\right)^{|\mathrm{odd_cycle}(\varsigma)|} \cdot \sum_{h_{\mathrm{s}}\in\mathcal{H}^{\mathrm{s}}_+} \prod_i \left(\frac{\exp(th_{\mathrm{s}}^i) + \exp(-th_{\mathrm{s}}^i)}{2}\right) \\
&\leq \left(\frac{2}{2+t^2K_{\mathrm{s}}^2}\right)^{|\mathrm{odd_cycle}(\varsigma)|} \cdot |\mathcal{H}^{\mathrm{s}}_+| \exp\left(\frac{t^2}{2} \cdot \sum_i \left(\max_{h\in\mathcal{H}^{\mathrm{s}}_+} h_{\mathrm{s}}^i\right)^2\right) \quad (75) \\
&\leq \left(\frac{2}{2+t^2K_{\mathrm{s}}^2}\right)^{|\mathrm{odd_cycle}(\varsigma)|} \cdot |\mathcal{H}^{\mathrm{s}}_+| \exp\left(\frac{t^2 m K_{\mathrm{s}}^2}{2}\right) \ .
\end{aligned}$$

Ineq. (75) holds because $(\exp(x)+\exp(-x))/2 \leq \exp(x^2/2)$. Taking logs and rearranging yields:

$$\mu \leq \frac{1}{t} \cdot \log \frac{|\mathcal{H}^{\mathrm{s}}_+|}{\left(1 + \frac{t^2K_{\mathrm{s}}^2}{2}\right)^{|\mathrm{odd_cycle}(\varsigma)|}} + \frac{tmK_{\mathrm{s}}^2}{2} \ . \quad (76)$$

Now, suppose that we can choose t such that

$$\frac{t^2K_{\mathrm{s}}^2}{2} \geq \varepsilon \ , \quad (77)$$

for some $\varepsilon > 0$. In this case, ineq. (76) implies

$$\mu \leq \frac{1}{t} \cdot \log \frac{|\mathcal{H}^{\mathrm{s}}_+|}{(1+\varepsilon)^{|\mathrm{odd_cycle}(\varsigma)|}} + \frac{tmK_{\mathrm{s}}^2}{2} \ , \quad (78)$$

which is of the form $\mu \leq A/t + Bt$ with $B > 0$. Taking $t = \sqrt{A/B}$ yields, using the fact that each classifier in $\mathcal{H}^{\mathrm{s}}_+$:

$$\mu \leq K_{\mathrm{s}} \cdot \sqrt{2m \log \frac{|\mathcal{H}^{\mathrm{s}}_+|}{(1+\varepsilon)^{|\mathrm{odd_cycle}(\varsigma)|}}} \ . \quad (79)$$

Dividing by m gives the statement of Theorem 10. We need however to check that ineq. (77) holds, which, since $t = \sqrt{A/B}$, yields that we must have, after simplification:

$$\log |\mathcal{H}^{\mathrm{s}}_{+}| \;\; \geq \;\; \varepsilon m + |\mathrm{odd_cycle}(\varsigma)| \log(1+\varepsilon) \;\; . \tag{80}$$

Since we have excluded fixed points, $|\mathrm{odd_cycle}(\varsigma)| \leq m/3$, and since $\log(1+x) \leq x$, a sufficient condition is $\log |\mathcal{H}^{\mathrm{s}}_{+}| \geq 4\varepsilon m/3$, which is the Theorem's assumption.

We now show a bound on the expected RCP when M in $\mathscr{T}$ is picked uniformly at random, with or without the class consistency requirement, as a function of the non-fixed points in the permutations. Following [Gretton et al., 2007], we let $\mathbf{i}_r^m$ denote the set of r-tuples drawn without replacement drawn from $[m]$. We also let $\mathbf{i}_r^{m,b}$ denote the set of r-tuples drawn without replacement drawn from $[m] \cap \{i : y_i = b1\}$, for $b \in \{-,+\}$.

Theorem 22 *Under the joint Settings of Theorem 9 and Setting (B), let S_m^k denotes the set of permutations with exactly k non-fixed points, and S_m^{kb} its subset of block-class permutations with non-fixed points in class $b \in \{-,+\}$. Then for $S \in \{S_m^k, S_m^{k-}, S_m^{k+}\}$ the following holds over the uniform sampling of permutations:*

$$\mathbb{E}_{\mathbf{M}\sim S}[\mathrm{RCP}_{\mathscr{T}}(\mathcal{H})] \;\; \leq \;\; u \cdot \frac{r_{\mathrm{s}}}{\sqrt{m}} \cdot \sqrt{\frac{k}{m} \cdot \mathscr{Q}} \;\; , \tag{81}$$

where $\mathscr{Q} \doteq \mathbb{E}_{(i,i')\sim \mathbf{i}_2^m}[\|\boldsymbol{x}_i^{\mathrm{s}} - \boldsymbol{x}_{i'}^{\mathrm{s}}\|_2^2]$ if $S = S_m^k$, and $\mathscr{Q} \doteq \mathbb{E}_{(i,i')\sim \mathbf{i}_2^{m,b}}[\|\boldsymbol{x}_i^{\mathrm{s}} - \boldsymbol{x}_{i'}^{\mathrm{s}}\|_2^2]$ if $S = S_m^{kb}$ ($b \in \{-,+\}$).

Proof We make the proof for $S = S_m^k$. The two other cases follow in the same way. We have from Theorem 9, because of Jensen inequality:

$$\begin{aligned}
\mathbb{E}_{\mathbf{M}\sim S_m^k}\left[\mathrm{RCP}_{\mathscr{T}}(\mathcal{H})\right] \;\; &\leq \;\; u \cdot \frac{r_{\mathrm{s}}}{\sqrt{m}} \cdot \mathbb{E}_{\mathbf{M}\sim S_m^k}\left[\sqrt{\frac{1}{m} \cdot \langle \mathrm{I}_m, \mathrm{K}^{\mathrm{s}} \rangle_{\mathbf{M}}}\right] \\
&\leq \;\; u \cdot \frac{r_{\mathrm{s}}}{\sqrt{m}} \cdot \sqrt{\frac{1}{m} \cdot \mathbb{E}_{\mathbf{M}\sim S_m^k}\left[\langle \mathrm{I}_m, \mathrm{K}^{\mathrm{s}} \rangle_{\mathbf{M}}\right]} \;\; .
\end{aligned} \tag{82}$$

We now decompose the expectation inside and first condition on the set of permutations whose set of non fixed points are the same set of k examples, say for $i \in [k]$. Let us call S_m^{k*} this subset of

S_m^k. In this case, we obtain:

$$\begin{aligned}
&\mathbb{E}_{\mathrm{M}\sim S_m^{k*}}\left[\langle \mathrm{I}_m, \mathrm{K}^{\mathrm{s}}\rangle_{\mathrm{M}}\right] \\
&= \mathbb{E}_{\mathrm{M}\sim S_m^{k*}}\left[\mathrm{tr}\left((\mathrm{I}_m-\mathrm{M})\mathrm{K}^{\mathrm{s}}(\mathrm{I}_m-\mathrm{M})^\top\right)\right] \\
&= \mathrm{tr}\left(\mathrm{K}^{\mathrm{s}}\right)+\mathbb{E}_{\mathrm{M}\sim S_m^{k*}}\left[\mathrm{tr}\left(\mathrm{M}\mathrm{K}^{\mathrm{s}}\mathrm{M}^\top\right)\right]-2\cdot\mathbb{E}_{\mathrm{M}\sim S_m^{k*}}\left[\mathrm{tr}\left(\mathrm{M}\mathrm{K}^{\mathrm{s}}\right)\right] \qquad (83)\\
&= 2\cdot\left(\mathrm{tr}\left(\mathrm{K}^{\mathrm{s}}\right)-\mathrm{tr}\left(\mathbb{E}_{\mathrm{M}\sim S_m^{k*}}\left[\mathrm{M}\right]\mathrm{K}^{\mathrm{s}}\right)\right) \\
&= 2\cdot\left(\sum_{i\in[m]}\mathrm{K}^{\mathrm{s}}_{ii}-\sum_{i\in[k]}\frac{1}{k}\cdot\sum_{i'\in[k]}\mathrm{K}^{\mathrm{s}}_{ii'}-\sum_{i\in[m]\setminus[k]}\mathrm{K}^{\mathrm{s}}_{ii}\right) \qquad (84)\\
&= 2\cdot\left(\frac{k-1}{k}\cdot\sum_{i\in[k]}\mathrm{K}^{\mathrm{s}}_{ii}-\frac{1}{k}\cdot\sum_{(i,i')\in\mathbf{i}_2^k}\mathrm{K}^{\mathrm{s}}_{ii'}\right) \\
&= \frac{2}{k}\cdot\left(\sum_{(i,i')\in\mathbf{i}_2^k}\frac{\mathrm{K}^{\mathrm{s}}_{ii}+\mathrm{K}^{\mathrm{s}}_{i'i'}}{2}-\mathrm{K}^{\mathrm{s}}_{ii'}\right) \\
&= \frac{1}{k}\cdot\sum_{(i,i')\in\mathbf{i}_2^k}\|\boldsymbol{x}_i^{\mathrm{s}}-\boldsymbol{x}_{i'}^{\mathrm{s}}\|_2^2\ . \qquad (85)
\end{aligned}$$

In eq. (83) we use the fact that K^{s} is symmetric. Eq. (84) uses the fact that

$$\mathbb{E}_{\mathrm{M}\sim S_m^{k*}}\left[\mathrm{M}\right] = \left[\begin{array}{c|c} \mathrm{U}_k & 0 \\ 0 & \mathrm{I}_{m-k}\end{array}\right],$$

where we recall that $\mathrm{U}_k \doteq \frac{1}{k}\cdot\mathbf{1}\mathbf{1}^\top$ (main file, Definition 11). There remains to average eq. (85) over the set of all permutations whose set of fixed points is a different $(m-k)$-subset of $[m]$ and the statement of Theorem 22 is proven for $S=S_m^k$. ■

The key point in the bound is factor k/m, which implies that when permutations have lots of fixed points, say $(1-\Omega(1))m$, then the RCP may just vanish (as m increases) wrt the Rademacher complexity, whose dependency on m is $\Omega(1/\sqrt{m})$ [Kakade et al., 2008].

9.6 Proof of Theorem 13

The proof stems from the following Theorem, which just assumes that K^u and K^v can be diagonalized (hence, it is applies to a more general setting than kernel functions).

Theorem 23 *Let K^u and K^v be two diagonalisable matrices with respective eigendecomposition $\{\lambda_i,\boldsymbol{u}_i\}_{i\in[d]}$ and $\{\mu_i,\boldsymbol{v}_i\}_{i\in[d]}$, with eigenvalues eventually duplicated up to their algebraic multiplicity. Letting $\overline{a}\doteq(1/m)\mathbf{1}^\top\boldsymbol{a}$ denote the average coordinate in $\boldsymbol{a}$, the difference in Hilbert-Schmidt Independence Criterion with respect to shuffling M satisfies:*

$$\mathrm{HSIC}(\mathrm{K}^u,\mathrm{K}^v)-\mathrm{HSIC}(\mathrm{K}^u,\mathrm{M}\mathrm{K}^v\mathrm{M}^\top) = -2m\cdot\left(\sum_i\lambda_i\overline{u}_i\boldsymbol{u}_i\right)^\top(\mathrm{I}_m-\mathrm{M})\left(\sum_i\mu_i\overline{v}_i\boldsymbol{v}_i\right) \qquad (86)$$

Hence, if $\mathrm{M} \in S_m^e$ *permutes* ℓ *and* ℓ' *in* $[m]$*, then* $\textsc{hsic}(\mathrm{K}^u, \mathrm{M}\mathrm{K}^v\mathrm{M}^\top) > \textsc{hsic}(\mathrm{K}^u, \mathrm{K}^v)$ *iff:*

$$\left(\sum_i \lambda_i \overline{u}_i (u_{i\ell} - u_{i\ell'})\right)\left(\sum_i \mu_i \overline{v}_i (v_{i\ell} - v_{i\ell'})\right) \;>\; 0 \;. \tag{87}$$

Proof Being symmetric, K^u and K^v can be diagonalized as $\mathrm{K}^u = \sum_i \lambda_i \boldsymbol{u}_i \boldsymbol{u}_i^\top$ and $\mathrm{K}^v = \sum_i \mu_i \boldsymbol{v}_i \boldsymbol{v}_i^\top$. We use definition $\mathrm{U}_m \doteq (1/m)\mathbf{1}\mathbf{1}^\top$ for short. The following is folklore or can be can be checked after analytic derivations that we omit:

$$\begin{aligned}
\mathrm{tr}\left(\mathrm{U}_m \mathrm{K}^u \mathrm{K}^v\right) &= m\left(\sum_i \lambda_i \overline{u}_i \boldsymbol{u}_i\right)\left(\sum_i \mu_i \overline{v}_i \boldsymbol{v}_i\right) \\
&= \mathrm{tr}\left(\mathrm{K}^u \mathrm{U}_m \mathrm{K}^v\right) \\
\mathrm{tr}\left(\mathrm{U}_m \mathrm{K}^u \mathrm{U}_m \mathrm{K}^v\right) &= m^2\left(\sum_i \lambda_i (\overline{u}_i)^2\right)\left(\sum_i \mu_i (\overline{v}_i)^2\right) \\
\mathrm{tr}\left(\mathrm{K}^u \mathrm{K}^v\right) &= \sum_i \lambda_i \mu_i \;.
\end{aligned} \tag{88}$$

We thus get

$$\begin{aligned}
\textsc{hsic}(\mathrm{K}^u, \mathrm{K}^v) &= \langle \mathrm{K}^u, \mathrm{K}^v \rangle_{\mathrm{U}_m} \\
&= \mathrm{tr}\left((\mathrm{I}_m - \mathrm{U}_m)\mathrm{K}^u(\mathrm{I}_m - \mathrm{U}_m)\mathrm{K}^v\right) \\
&= \mathrm{tr}\left(\mathrm{K}^u\mathrm{K}^v\right) - \mathrm{tr}\left(\mathrm{U}_m\mathrm{K}^u\mathrm{K}^v\right) - \mathrm{tr}\left(\mathrm{K}^u\mathrm{U}_m\mathrm{K}^v\right) + \mathrm{tr}\left(\mathrm{U}_m\mathrm{K}^u\mathrm{U}_m\mathrm{K}^v\right) \\
&= \sum_i \lambda_i\mu_i - 2m \cdot \left(\sum_i \lambda_i \overline{u}_i \boldsymbol{u}_i\right)^\top \left(\sum_i \mu_i \overline{v}_i \boldsymbol{v}_i\right) \\
&\quad + m^2 \cdot \left(\sum_i \lambda_i (\overline{u}_i)^2\right) \cdot \left(\sum_i \mu_i (\overline{v}_i)^2\right) \\
&= m^2\left(\frac{1}{m^2} \cdot \sum_i \lambda_i\mu_i - \frac{2}{m}\sum_{i,j} \lambda_i \overline{u}_i \boldsymbol{u}_i^\top \mu_j \overline{v}_j \boldsymbol{v}_j + \sum_{i,j} \lambda_i\mu_j (\overline{u}_i)^2 (\overline{v}_j)^2\right) \\
&= m^2\left(\frac{1}{m^2} \cdot \sum_i \lambda_i\mu_i - \frac{1}{m^2} \cdot \sum_{i,j} \lambda_i\mu_j (\boldsymbol{u}_i^\top \boldsymbol{v}_j)^2 + \sum_{i,j} \lambda_i\mu_j \left(\frac{1}{m}\boldsymbol{u}_i^\top \boldsymbol{v}_j - \overline{u}_i\overline{v}_j\right)^2\right) \\
&= m^2\left(\frac{1}{m^2} \cdot \boldsymbol{\lambda}^\top (\mathrm{I} - \mathrm{C})\,\boldsymbol{\mu} + \boldsymbol{\lambda}^\top \mathrm{J} \boldsymbol{\mu}\right) \qquad (89) \\
&= \boldsymbol{\lambda}^\top \left((\mathrm{I} - \mathrm{C}) + m^2 \cdot \mathrm{J}\right)\boldsymbol{\mu} \;. \qquad (90)
\end{aligned}$$

We have used here the square cosine matrix C with $\mathrm{C}_{ij} \doteq \cos^2(\boldsymbol{u}_i, \boldsymbol{u}_j)$, and the square correlation matrix J with $\mathrm{J}_{ij} \doteq ((1/m)\boldsymbol{u}_i^\top \boldsymbol{v}_j - \overline{u}_i\overline{v}_j)^2$. Now, suppose we perform CP $\mathscr{T}$ with shuffling matrix M. K^v and its eigendecomposition become after shuffling

$$\mathrm{M}\mathrm{K}^v\mathrm{M}^\top \;=\; \sum_i \mu_i (\mathrm{M}\boldsymbol{v}_i)(\mathrm{M}\boldsymbol{v}_i)^\top \;. \tag{91}$$

Remark that shuffling affects the order in the coordinate of *all* eigenvectors. So the difference between the two Hilbert-Schmidt Independence Criteria (before - after shuffling) is:

$$\begin{aligned}
&\textsc{hsic}(\mathrm{K}^u, \mathrm{K}^v) - \textsc{hsic}(\mathrm{K}^u, \mathrm{M}\mathrm{K}^v\mathrm{M}^\top) \\
&= \sum_{i,j} \lambda_i\mu_j((\boldsymbol{u}_i^\top \mathrm{M}\boldsymbol{v}_j)^2 - (\boldsymbol{u}_i^\top \boldsymbol{v}_j)^2) \\
&\quad - \sum_{i,j} \lambda_i\mu_j \left\{ \left(\boldsymbol{u}_i^\top \mathrm{M}\boldsymbol{v}_j - m\overline{u}_i\overline{v}_j\right)^2 - \left(\boldsymbol{u}_i^\top \boldsymbol{v}_j - m\overline{u}_i\overline{v}_j\right)^2 \right\} \\
&= \sum_{i,j} \lambda_i\mu_j \cdot \boldsymbol{u}_i^\top(\mathrm{M} - \mathrm{I}_m)\boldsymbol{v}_j \cdot \boldsymbol{u}_i^\top(\mathrm{M} + \mathrm{I}_m)\boldsymbol{v}_j \\
&\quad - \sum_{i,j} \lambda_i\mu_j \left\{ \boldsymbol{u}_i^\top(\mathrm{M} - \mathrm{I}_m)\boldsymbol{v}_j \cdot (\boldsymbol{u}_i^\top(\mathrm{M} + \mathrm{I}_m)\boldsymbol{v}_j - 2m\overline{u}_i\overline{v}_j) \right\} \\
&= 2m \cdot \sum_{i,j} (\lambda_i\overline{u}_i)\boldsymbol{u}_i^\top(\mathrm{M} - \mathrm{I}_m)(\mu_j\overline{v}_j) \\
&= 2m \cdot \left(\sum_i \lambda_i\overline{u}_i\boldsymbol{u}_i\right)^\top (\mathrm{M} - \mathrm{I}_m) \left(\sum_i \mu_i\overline{v}_i\boldsymbol{v}_i\right) \quad . \qquad (92)
\end{aligned}$$

We now remark that whenever $\mathrm{M} \in S_m^{\mathrm{e}}$, if it permutes ℓ and ℓ' in $[m]$, then $\boldsymbol{a}(\mathrm{M} - \mathrm{I}_m)\boldsymbol{b} = a_\ell(b_{\ell'} - b_\ell) + a_{\ell'}(b_\ell - b_{\ell'}) = -(a_\ell - a_{\ell'})(b_\ell - b_{\ell'})$, so we get:

$$\begin{aligned}
&\textsc{hsic}(\mathrm{K}^u, \mathrm{K}^v) - \textsc{hsic}(\mathrm{K}^u, \mathrm{M}\mathrm{K}^v\mathrm{M}^\top) \\
&= -2m \left(\sum_i \lambda_i\overline{u}_i(u_{i\ell} - u_{i\ell'})\right)\left(\sum_i \mu_i\overline{v}_i(v_{i\ell} - v_{i\ell'})\right) \quad , \qquad (93)
\end{aligned}$$

and we get ineq. (87). ■

This ends the proof of Theorem 13.

9.7 Proof of Theorem 14

The Theorem is a direct consequence of the following Theorem.

Theorem 24 *Let* K^u *and* K^v *be two kernel functions over* $\mathcal{S}$*. Then for any elementary permutation* $\mathrm{M} \in S_m^e$ *that permutes* ℓ *and* ℓ' *in* $[m]$*,*

$$\textsc{hsic}(\mathrm{K}^u, \mathrm{M}\mathrm{K}^v\mathrm{M}^\top) - \textsc{hsic}(\mathrm{K}^u, \mathrm{K}^v) \quad = \quad -2m \cdot \mathbb{C}\mathrm{ov}(\boldsymbol{\delta}^u_{\ell\ell'}, \boldsymbol{\delta}^v_{\ell\ell'}) + \mathscr{R}^{u,v}_{\ell\ell'} \quad , \qquad (94)$$

with

$$\mathscr{R}^{u,v}_{\ell\ell'} \quad \doteq \quad (\mathrm{K}^u_{\ell\ell} - \mathrm{K}^u_{\ell'\ell})(\mathrm{K}^v_{\ell\ell} - \mathrm{K}^v_{\ell'\ell}) + (\mathrm{K}^u_{\ell'\ell'} - \mathrm{K}^u_{\ell'\ell})(\mathrm{K}^v_{\ell'\ell'} - \mathrm{K}^v_{\ell'\ell}) \quad . \qquad (95)$$

Furthermore, the uniform sampling of elementary permutations in S_m^e *satisfies:*

$$\mathbb{E}_{\mathrm{M}\sim S_m^e}\left[\textsc{hsic}(\mathrm{K}^u, \mathrm{M}\mathrm{K}^v\mathrm{M}^\top)\right] \quad = \quad \left(1 - \frac{8}{m-1}\right) \cdot \textsc{hsic}(\mathrm{K}^u, \mathrm{K}^v) + \frac{8}{m-1} \cdot \mathscr{R}^{u,v} \quad , \qquad (96)$$

with

$$\mathscr{R}^{u,v} \doteq \sum_i \mathrm{K}^u_{ii}\mathrm{K}^v_{ii} - \frac{1}{m}\cdot\left(\frac{\sum_i \mathrm{K}^u_{ii}\mathrm{K}^v_{.i} + \sum_i \mathrm{K}^u_{.i}\mathrm{K}^v_{ii}}{2}\right) . \tag{97}$$

Here, when replacing an index notation by a point, ".", we denote a sum over all possible values of this index.

Proof We first decompose $\mathrm{HSIC}(\mathrm{K}^u, \mathrm{K}^v)$:

$$\mathrm{HSIC}(\mathrm{K}^u, \mathrm{K}^v) = \sum_{i,i'} \mathrm{K}^u_{ii'}\mathrm{K}^v_{ii'} - \frac{2}{m}\cdot\sum_i \mathrm{K}^u_{i.}\mathrm{K}^v_{i.} + \frac{1}{m^2}\mathrm{K}^u_{..}\mathrm{K}^u_{..} . \tag{98}$$

for any $(\ell, \ell') \in \mathbf{i}_2^m$. For any $\mathrm{M} \in S_m^{\mathrm{e}}$ denoting an elementary permutation ς of the features in $\mathcal{F}^{\mathrm{s}}$ such that $\mathcal{V} \subseteq \mathcal{F}^{\mathrm{s}}$ and $\varsigma(\ell) = \ell'$, $\varsigma(\ell') = \ell$, we obtain:

$$\begin{aligned}
&\mathrm{HSIC}(\mathrm{K}^u, \mathrm{M}\mathrm{K}^v\mathrm{M}^\top) - \mathrm{HSIC}(\mathrm{K}^u, \mathrm{K}^v) \\
&= 2\cdot\left(\sum_{i\neq\ell,\ell'} \mathrm{K}^u_{\ell i}\mathrm{K}^v_{\ell' i} + \sum_{i\neq\ell,\ell'} \mathrm{K}^u_{\ell' i}\mathrm{K}^v_{\ell i} - \sum_{i\neq\ell,\ell'} \mathrm{K}^u_{\ell i}\mathrm{K}^v_{\ell i} - \sum_{i\neq\ell,\ell'} \mathrm{K}^u_{\ell' i}\mathrm{K}^v_{\ell' i}\right) \\
&\quad -\frac{2}{m}\cdot\mathrm{K}^u_{\ell.}\mathrm{K}^v_{\ell'.} - \frac{2}{m}\cdot\mathrm{K}^u_{\ell'.}\mathrm{K}^v_{\ell.} + \frac{2}{m}\cdot\mathrm{K}^u_{\ell.}\mathrm{K}^v_{\ell.} + \frac{2}{m}\cdot\mathrm{K}^u_{\ell'.}\mathrm{K}^v_{\ell'.} \\
&= -2\left(\sum_i (\mathrm{K}^u_{\ell i} - \mathrm{K}^u_{\ell' i})(\mathrm{K}^v_{\ell i} - \mathrm{K}^v_{\ell' i}) - \frac{1}{m}\cdot(\mathrm{K}^u_{\ell.} - \mathrm{K}^u_{\ell'.})(\mathrm{K}^v_{\ell.} - \mathrm{K}^v_{\ell'.})\right) \\
&\quad +(\mathrm{K}^u_{\ell\ell} - \mathrm{K}^u_{\ell'\ell})(\mathrm{K}^v_{\ell\ell} - \mathrm{K}^v_{\ell'\ell}) + (\mathrm{K}^u_{\ell'\ell'} - \mathrm{K}^u_{\ell'\ell})(\mathrm{K}^v_{\ell'\ell'} - \mathrm{K}^v_{\ell'\ell}) \\
&= -2m\cdot\mathbb{C}\mathrm{ov}(\boldsymbol{\delta}^u_{\ell\ell'}, \boldsymbol{\delta}^v_{\ell\ell'}) + \mathscr{R}^{u,v}_{\ell\ell'} ,
\end{aligned}$$

which is eq. (94). We also have:

$$\begin{aligned}
\mathbb{E}_{\mathrm{M}\sim S_m^{\mathrm{e}}}\left[\sum_i (\mathrm{K}^u_{\ell i} - \mathrm{K}^u_{\ell' i})(\mathrm{K}^v_{\ell i} - \mathrm{K}^v_{\ell' i})\right] &= \frac{4}{m}\cdot\sum_{i,i'} \mathrm{K}^u_{ii'}\mathrm{K}^v_{ii'} - \frac{4}{m(m-1)}\cdot\sum_i \mathrm{K}^u_{i.}\mathrm{K}^v_{i.} \\
&\quad +\frac{4}{m(m-1)}\cdot\sum_{i,i'} \mathrm{K}^u_{ii'}\mathrm{K}^v_{ii'} \\
&= \frac{4}{m-1}\cdot\sum_{i,i'} \mathrm{K}^u_{ii'}\mathrm{K}^v_{ii'} - \frac{4}{m(m-1)}\cdot\sum_i \mathrm{K}^u_{i.}\mathrm{K}^v_{i.} ,
\end{aligned}$$

$$\begin{aligned}
\mathbb{E}_{\mathrm{M}\sim S_m^{\mathrm{e}}}\left[(\mathrm{K}^u_{\ell.} - \mathrm{K}^u_{\ell'.})(\mathrm{K}^v_{\ell.} - \mathrm{K}^v_{\ell'.})\right] &= \frac{4}{m}\cdot\sum_i \mathrm{K}^u_{i.}\mathrm{K}^v_{i.} - \frac{4}{m(m-1)}\cdot\sum_{i\in[m]} \mathrm{K}^u_{i.}\sum_{i'\in[m]\setminus\{i\}} \mathrm{K}^v_{i'.} \\
&= \frac{4}{m}\cdot\sum_i \mathrm{K}^u_{i.}\mathrm{K}^v_{i.} - \frac{4}{m(m-1)}\cdot\sum_i \mathrm{K}^u_{i.}\cdot(\mathrm{K}^v_{..} - \mathrm{K}^v_{i.}) \\
&= \frac{4}{m}\cdot\sum_i \mathrm{K}^u_{i.}\mathrm{K}^v_{i.} - \frac{4}{m(m-1)}\mathrm{K}^u_{..}\mathrm{K}^v_{..} + \frac{4}{m(m-1)}\cdot\sum_i \mathrm{K}^u_{i.}\mathrm{K}^v_{i.} \\
&= \frac{4}{m-1}\cdot\sum_i \mathrm{K}^u_{i.}\mathrm{K}^v_{i.} - \frac{4}{m(m-1)}\mathrm{K}^u_{..}\mathrm{K}^v_{..} ,
\end{aligned}$$

and finally

$$\begin{aligned}\mathbb{E}_{\mathbf{M}\sim S_m^{\mathrm{e}}}[\mathscr{R}_{\ell\ell'}^{u,v}] &= \frac{8}{m-1}\cdot\left[\sum_i \mathrm{K}_{ii}^u\mathrm{K}_{ii}^v - \frac{1}{m}\cdot\left(\frac{\sum_i \mathrm{K}_{ii}^u\mathrm{K}_{\cdot i}^v + \sum_i \mathrm{K}_{\cdot i}^u\mathrm{K}_{ii}^v}{2}\right)\right] \\ &\doteq \frac{8}{m-1}\cdot\mathscr{R}^{u,v}\ , \end{aligned} \tag{99}$$

since

$$\begin{aligned}&\mathbb{E}_{\mathbf{M}\sim S_m^{\mathrm{e}}}[(\mathrm{K}_{\ell\ell}^u - \mathrm{K}_{\ell'\ell}^u)(\mathrm{K}_{\ell\ell}^v - \mathrm{K}_{\ell'\ell}^v)] \\ &= \frac{4}{m(m-1)}\sum_i\sum_{i'\neq i}\mathrm{K}_{ii}^u\mathrm{K}_{ii}^v - \frac{2}{m(m-1)}\sum_i\sum_{i'\neq i}\mathrm{K}_{ii}^u\mathrm{K}_{i'i}^v - \frac{2}{m(m-1)}\sum_i\sum_{i'\neq i}\mathrm{K}_{i'i}^u\mathrm{K}_{ii}^v \\ &= \frac{4}{m}\sum_i\mathrm{K}_{ii}^u\mathrm{K}_{ii}^v - \frac{2}{m(m-1)}\sum_i\mathrm{K}_{ii}^u(\mathrm{K}_{\cdot i}^v - \mathrm{K}_{ii}^v) - \frac{2}{m(m-1)}\sum_i(\mathrm{K}_{i\cdot}^u - \mathrm{K}_{ii}^u)\mathrm{K}_{ii}^v \\ &= \frac{4}{m}\sum_i\mathrm{K}_{ii}^u\mathrm{K}_{ii}^v + \frac{4}{m(m-1)}\sum_i\mathrm{K}_{ii}^u\mathrm{K}_{ii}^v - \frac{2}{m(m-1)}\sum_i\mathrm{K}_{ii}^u\mathrm{K}_{\cdot i}^v - \frac{2}{m(m-1)}\sum_i\mathrm{K}_{i\cdot}^u\mathrm{K}_{ii}^v \\ &= \frac{4}{m-1}\sum_i\mathrm{K}_{ii}^u\mathrm{K}_{ii}^v - \frac{2}{m(m-1)}\sum_i\mathrm{K}_{ii}^u\mathrm{K}_{\cdot i}^v - \frac{2}{m(m-1)}\sum_i\mathrm{K}_{i\cdot}^u\mathrm{K}_{ii}^v \\ &= \mathbb{E}_{\mathbf{M}\sim S_m^{\mathrm{e}}}[(\mathrm{K}_{\ell'\ell'}^u - \mathrm{K}_{\ell'\ell}^u)(\mathrm{K}_{\ell'\ell'}^v - \mathrm{K}_{\ell'\ell}^v)]\ . \end{aligned}$$

So we obtain

$$\begin{aligned}&\mathbb{E}_{\mathbf{M}\sim S_m^{\mathrm{e}}}\left[\mathrm{HSIC}(\mathrm{K}^u, \mathrm{M}\mathrm{K}^v\mathrm{M}^\top) - \mathrm{HSIC}(\mathrm{K}^u, \mathrm{K}^v)\right] \\ &= -\frac{8}{m-1}\cdot\sum_{i,i'}\mathrm{K}_{ii'}^u\mathrm{K}_{ii'}^v + \frac{8}{m(m-1)}\cdot\sum_i\mathrm{K}_{i\cdot}^u\mathrm{K}_{i\cdot}^v \\ &\quad + \frac{8}{m(m-1)}\cdot\sum_i\mathrm{K}_{i\cdot}^u\mathrm{K}_{i\cdot}^v - \frac{8}{m^2(m-1)}\mathrm{K}_{\cdot\cdot}^u\mathrm{K}_{\cdot\cdot}^v + \frac{8}{m-1}\cdot\mathscr{R}^{u,v} \\ &= -\frac{8}{m-1}\cdot\mathrm{HSIC}(\mathrm{K}^u, \mathrm{K}^v) + \frac{8}{m-1}\cdot\mathscr{R}^{u,v}\ . \end{aligned}$$

This ends the proof of Theorem 24. ■

When kernel functions have unit diagonal (such as for the Gaussian kernel), eq. (97) simplifies to:

$$\mathscr{R}^{u,v} \doteq m\left(1 - \frac{1}{m^2}\cdot\left(\frac{\mathrm{K}_{\cdot\cdot}^u + \mathrm{K}_{\cdot\cdot}^v}{2}\right)\right)\ . \tag{100}$$

Hence, provided we perform $T \doteq \epsilon m$ elementary permutations, there exists a sequence of such permutations such that the composition $\mathrm{M}_* \doteq \mathrm{M}_T\mathrm{M}_{T-1}\cdots\mathrm{M}_1$ satisfies:

$$\begin{aligned}\mathrm{HSIC}(\mathrm{K}^u, \mathrm{M}_*\mathrm{K}^v\mathrm{M}_*^\top) &\leq \left(1 - \frac{8}{m-1}\right)^{\epsilon m}\cdot\mathrm{HSIC}(\mathrm{K}^u, \mathrm{K}^v) + \left[1 - \left(1 - \frac{8}{m-1}\right)^{\epsilon m}\right]\cdot\mathscr{R}^{u,v} \\ &\leq \alpha(\epsilon)\cdot\mathrm{HSIC}(\mathrm{K}^u, \mathrm{K}^v) + (1-\alpha(\epsilon))\cdot\mathscr{R}^{u,v}\ , \end{aligned} \tag{101}$$




Figure 4: The Cornia-Mooij model [Cornia and Mooij, 2014]. Left: belief, right: true model.

with

$$\alpha(\epsilon) \doteq \exp(-8\epsilon) \ , \tag{102}$$

as long as $\text{HSIC}(\text{K}^u, \text{K}^v) \geq \mathscr{R}^{u,v}$. We have used the fact that

$$\begin{aligned} \left(1 - \frac{8}{m-1}\right)^{\epsilon m} &\leq \exp\left(-\frac{8\epsilon m}{m-1}\right) \\ &\leq \exp(-8\epsilon) \ . \end{aligned} \tag{103}$$

This achieves the proof of Theorem 14.

9.8 The Cornia-Mooij model and results

We now show how to trick statistical tests into keeping independence *and* then incur arbitrarily large errors in estimating causal effects, *via* the use of CPs. The model we refer to is the Cornia-Mooij (CM) model [Cornia and Mooij, 2014], shown in Figure 4. In the CM model, there are $d = 3$ observation variables, and a true model which relies on a weak conditional dependence $x_1 \not\perp\!\!\!\perp x_3 | x_2$. [Cornia and Mooij, 2014] show that *if* one keeps the independence assumption H_0 that $x_1 \perp\!\!\!\perp x_3 | x_2$, this can lead to very high causal estimation errors, as measured by $|\mathbb{E}[x_3|x_2] - \mathbb{E}[x_3|\text{do}(x_2)]|/|x_2|$ [Cornia and Mooij, 2014]. We show that the CP is precisely able to trick statistics into keeping H_0.

There is a hidden confounder x_4, which is assumed to be independent from x_1. The true model makes the following statistical dependence assumptions:

- $x_1 \not\perp\!\!\!\perp x_2$,
- $x_2 \not\perp\!\!\!\perp x_3$,
- and the most important one, which we scramble through the CP, $x_1 \not\perp\!\!\!\perp x_3 | x_2$.

We chose this simple model because (a) it belongs to the few worst-case models for causality analysis, and (b) it shows, in addition to jamming (non)linear correlations, how CP can also jam partial correlations. In the CM model, there are $d = 3$ observation variables, and a true model which relies on a weak conditional dependence $x_1 \not\perp\!\!\!\perp x_3 | x_2$. [Cornia and Mooij, 2014] show that *if* one keeps the independence assumption H_0 that $x_1 \perp\!\!\!\perp x_3 | x_2$, this can lead to very high causal

estimation errors[6]. We show that it is possible, through a CP, to trick statistics into *keeping H_0 as well*. In the following, $\mathcal{F}_{\rm s} = \{x_3\}$. It is shown in [Cornia and Mooij, 2014] that $x_1 \perp\!\!\!\perp x_3 | x_2$ iff the partial correlation $\rho_{(13)\cdot 2} \doteq (\rho_{13} - \rho_{12}\rho_{23})/\sqrt{(1-\rho_{12}^2)(1-\rho_{23}^2)}$ vanishes. Assuming $\rho_{(13)\cdot 2}$ is large enough in the dataset we have (so that we would reject H_0 from observing $\mathcal{S}$), we show how to reduce it through a sequence of CPs, using a similar strategy as in Theorem 14, the main difference being that we rely on block-class permutations. For any $\varsigma \in S_m$, notation 3^{ς} indicates column variable 3 shuffled. We assume $\rho_{(13)\cdot 2} > 0$ (the same analysis can be done if $\rho_{(13)\cdot 2} < 0$).

Theorem 25 *Suppose that there exists $\epsilon > 0$ such that $\rho_{12}^2 \leq 1-\epsilon$ and $\rho_{23^{\varsigma}}^2 \leq 1-\epsilon$ for any $\varsigma \in S_m^*$. Then there exists $T > 0$ and a sequence of T elementary permutations in S_m^* such that $\rho_{(13^{\varsigma})\cdot 2}$ is strictly decreasing in the sequence and meets at the end $\rho_{(13^{\varsigma})\cdot 2} \leq \mathscr{R}$ with*

$$\mathscr{R} \;\doteq\; (1-\epsilon)^{-1} \cdot p_+(1-p_+) \cdot (\tilde{\mu}_1 - \rho_{12} \cdot \tilde{\mu}_2) \cdot \tilde{\mu}_3 \;\;,$$

and $\tilde{\mu}_j \doteq (1/(2\sqrt{v_j})) \cdot \sum_{y'} y' \mathbb{E}_{(\boldsymbol{x},y)\sim \mathcal{S}}[x_j | y = y']$.

Proof

The Theorem is a direct consequence of the following Lemma.

Lemma 26 *Let c_{jk} denote the covariance between columns j and k, $\mu_l^b \doteq (1/m_b) \sum_{i:y_i=b} x_{il}$ and $p \doteq m_+/m$. Suppose that $j \in \mathcal{F}_{\rm a}$ and $k \in \mathcal{F}_{\rm s}$. Then as long as*

$$c_{jk} \;>\; p(1-p) \cdot (\mu_j^+ - \mu_j^-) \cdot (\mu_k^+ - \mu_k^-) \;\;, \tag{104}$$

there always exist $\varsigma \in S_m^*$ such that $\rho_{jk^{\varsigma}} < \rho_{jk}$, where k^{ς} denote column variable k shuffled according to ς in the corresponding CP.

Proof Let us denote for short $\boldsymbol{c}_j \in \mathbb{R}^m$ the j^{th} feature column. We have because of the fact that $\mu_{k^{\varsigma}} = \mu_k$ and $v_{k^{\varsigma}} = v_k$ ($v_{\cdot}$ being the variance):

$$\rho_{jk^{\varsigma}} - \rho_{jk} \;=\; \frac{1}{m\sqrt{v_j v_k}} \cdot \left(\boldsymbol{c}_j^{\top} \left(\mathrm{M}_{\varsigma} - \mathrm{I}_m \right) \boldsymbol{c}_k \right) \;\;, \tag{105}$$

where M_{ς} is the shuffling matrix of permutation ς. Matrix $\mathrm{M}_{\varsigma} - \mathrm{I}_m$ has only four non-zero coordinates: in (ℓ,ℓ) and (ℓ',ℓ') (both -1), and in (ℓ,ℓ') and (ℓ',ℓ) (both 1), so we get:

$$\begin{aligned} \rho_{jk^{\varsigma}} - \rho_{jk} \;&=\; \frac{1}{m\sqrt{v_j v_k}} \cdot (x_{\ell j}(x_{\ell' k} - x_{\ell k}) + x_{\ell' j}(x_{\ell k} - x_{\ell' k})) \\ &=\; -\frac{1}{m\sqrt{v_j v_k}} \cdot ((x_{\ell j} - x_{\ell' j})(x_{\ell k} - x_{\ell' k})) \;\;. \end{aligned} \tag{106}$$

Hence, $\rho_{jk^{\varsigma}} < \rho_{jk}$ iff the sign of $x_{\ell j} - x_{\ell' j}$ is the same as the sign of $x_{\ell k} - x_{\ell' k}$. Let $\pi^b(\ell)$ the predicate $\rho_{jk^{\varsigma}} \geq \rho_{jk}$, for any elementary permutation $\varsigma \in S_m^*$ that changes ℓ to index ℓ' of the same

[6] As measured by $|\mathbb{E}[x_3|x_2] - \mathbb{E}[x_3|\mathrm{do}(x_2)]|/|x_2|$ [Cornia and Mooij, 2014].

class b ($\varsigma(\ell)=\ell'$, $\varsigma(\ell')=\ell$). If $\pi^b(\ell)$ is true, then, averaging over all such permutations, we obtain:

$$\begin{aligned} 0 \;\geq\; & -\frac{1}{m\sqrt{v_j v_k}}\cdot\frac{1}{m_b}\sum_{\ell'}\left((x^b_{\ell j}-x^b_{\ell' j})(x^b_{\ell k}-x^b_{\ell' k})\right) \\ & = -\frac{1}{m\sqrt{v_j v_k}}\cdot\left(x^b_{\ell j}x^b_{\ell k}-x^b_{\ell j}\mu^b_k-x^b_{\ell k}\mu^b_j+\mu^b_{jk}\right) \\ & = -\frac{1}{m\sqrt{v_j v_k}}\cdot\left(c^b_{jk}+(x^b_{\ell j}-\mu^b_j)(x^b_{\ell k}-\mu^b_k)\right) \;\;, \end{aligned}$$

i.e. we have:

$$(x^b_{\ell j}-\mu^b_j)(x^b_{\ell k}-\mu^b_k) \;\geq\; -c^b_{jk}\;. \tag{107}$$

Assume now that $\pi^b(\ell)$ holds over any $\ell\in[m_b]$. As long as c^b_{jk} is strictly positive, we thus obtain, averaging ineq. (107) over all $\ell\in[m_b]$,

$$\begin{aligned} c^b_{jk} \;\doteq\; & \frac{1}{m_b}\sum_{\ell}(x^b_{\ell j}-\mu^b_j)(x^b_{\ell k}-\mu^b_k) \\ \leq\; & -c^b_{jk} \\ <\; & 0\;, \end{aligned} \tag{108}$$

a contradiction. Hence, as long as $c^b_{jk}>0$, there must exist $(\ell,\ell')\in\mathbf{i}_2^{m_b}$ such that the elementary permutation $\varsigma(\ell)=\ell'$, $\varsigma(\ell')=\ell$ satisfies

$$c^b_{jk^\varsigma} \;<\; c^b_{jk}\;, \tag{109}$$

and this holds for $b\in\{-,+\}$. Now remark that

$$\begin{aligned} c_{jk} \;=\; & p\mu^+_{jk}+(1-p)\mu^-_{jk}-(p\mu^+_j+(1-p)\mu^-_j)(p\mu^+_k+(1-p)\mu^-_k) \\ =\; & pc^+_{jk}+(1-p)c^-_{jk}+p(1-p)(\mu^+_j-\mu^-_j)(\mu^+_k-\mu^-_k)\;, \end{aligned} \tag{110}$$

and so as long as whichever $c^+_{jk}>0$ or $c^-_{jk}>0$, we can always find an elementary permutation that decreases the one chosen. When no more elementary permutations achieve that, $c_{jk}\leq p(1-p)(\mu^+_j-\mu^-_j)(\mu^+_k-\mu^-_k)$, which yields the statement of the Lemma. ■

To prove the Theorem, remark that

$$\begin{aligned} \rho_{13^\varsigma}-\rho_{12}\rho_{23^\varsigma} \;=\; & \frac{1}{\sqrt{v_1 v_3}}\left(\frac{1}{m}\cdot\sum_i\left(x_{i1}-\frac{c_{12}}{v_2}\cdot x_{i2}\right)x_{\varsigma(i)3}\right)-\left(\frac{\mu_1\mu_3}{\sqrt{v_1 v_3}}-c_{12}\frac{\mu_2\mu_3}{v_2\sqrt{v_1 v_3}}\right) \\ =\; & \frac{1}{\sqrt{v_1 v_3}}\left(\frac{1}{m}\cdot\sum_i\left(x_{i1}-\frac{c_{12}}{v_2}\cdot x_{i2}\right)x_{\varsigma(i)3}-\left(\mu_1-\frac{c_{12}\mu_2}{v_2}\right)\mu_3\right) \end{aligned} \tag{111}$$

We apply Lemma 26 to linearly transformed column $\boldsymbol{c}'\doteq\boldsymbol{c}_1-\frac{c_{12}}{v_2}\boldsymbol{c}_2$ and column $\boldsymbol{c}_3$ and obtain that as long as

$$\rho_{13}-\rho_{12}\rho_{23} \;>\; \frac{p(1-p)}{\sqrt{v_1 v_3}}\left((\mu^+_1-\mu^-_1)-\frac{c_{12}}{v_2}(\mu^+_2-\mu^-_2)\right)(\mu^+_3-\mu^-_3)\;, \tag{112}$$

there always exist a block-class elementary permutation ς that is going to make $\rho_{13^\varsigma} - \rho_{12}\rho_{23^\varsigma} < \rho_{13} - \rho_{12}\rho_{23}$. When no such permutation exist anymore, we have, letting ς_T denote the composition of all elementary permutations performed so far and $\varsigma_* \doteq \arg\max_{\varsigma \in S_m^*} \rho^2_{23^\varsigma}$,

$$\begin{aligned}
\frac{\rho_{13} - \rho_{12}\rho_{23}}{\sqrt{1-\rho_{12}^2}\sqrt{1-\rho^2_{23^{\varsigma_T}}}} &\leq \frac{\rho_{13} - \rho_{12}\rho_{23}}{\sqrt{1-\rho_{12}^2}\sqrt{1-\rho^2_{23^{\varsigma_*}}}} \\
&= \frac{p(1-p)}{\sqrt{v_1 v_2 - c_{12}^2}\sqrt{v_2 v_3 - c^2_{23^{\varsigma_*}}}} \\
&\cdot \left(v_2(\mu_1^+ - \mu_1^-) - c_{12}(\mu_2^+ - \mu_2^-)\right)(\mu_3^+ - \mu_3^-) \\
&= \frac{p(1-p)}{1-\epsilon}\left(\frac{\mu_1^+ - \mu_1^-}{\sqrt{v_1}} - \rho_{12}\cdot\frac{\mu_2^+ - \mu_2^-}{\sqrt{v_2}}\right)\cdot\frac{\mu_3^+ - \mu_3^-}{\sqrt{v}_3} \qquad (113)
\end{aligned}$$

as long as $c_{12}^2 \leq (1-\epsilon)v_1 v_2$ and $c^2_{23^{\varsigma_*}} \leq (1-\epsilon)v_2 v_3$. We just have to use the fact that

$$\tilde{\mu}_j = \frac{\mu_j^+ - \mu_j^-}{\sqrt{v_j}} \qquad (114)$$

using the main file notation to conclude (End of the proof of Theorem 25). ■

Remark that the proof also shows that the conditions to blow up the type-II error (Corollary 2.1 in [Cornia and Mooij, 2014]) are not affected by the CP. To see that it is possible to still make the Type II error blow, up, in the CM model, the Type II error can be made at least K/v_2 [Cornia and Mooij, 2014] where the coefficient K does not depend on ς ([Cornia and Mooij, 2014], Corollary 2.1). Since v_2 is also not altered by the permutations, the Type II error can still be blown up following [Cornia and Mooij, 2014]'s construction.

We now show that the iterative process is actually not necessary if one has enough data: sampling $\varsigma \sim S_m^*$ jams $\rho_{(13^\varsigma)\cdot 2}$ up to bounds competitive with Theorem 25 with high probability. Such good concentration results also hold for HSIC [Song et al., 2012].

Theorem 27 *For any $\delta > 0$, provided $m = \Omega((1/\delta)\log(1/\delta))$, the uniform sampling of ς in S_m^* satisfies*

$$\mathbb{P}_{\varsigma \sim S_m^*}[\rho_{(13^\varsigma)\cdot 2} \leq \mathscr{R} + \delta] \geq 1 - \delta \ ,$$

where $\mathscr{R}$ is defined in Theorem 25.

Proof We detail first the sampling process of ς. It relies on the fundamental property that a permutation uniquely factors as a product of disjoint cycles, and so a block-class permutation ς factors uniquely as two permutations ς_+ and ς_-, each of which acts in one of the two classes. Therefore, sampling uniformly each of ς_+ and ς_- results in an uniform sampling of a block-class ς.

The Theorem stems from the following Lemma, whose notations follow Lemma 26.

Lemma 28 *For any $q > 0$, as long as*

$$m = \Omega\left(\frac{1}{q}\log\frac{1}{q}\right) \ , \qquad (115)$$

there is probability $\geq 1-q$ *that a randomly chosen block-class permutation* ς *shall bring*

$$c_{jk^\varsigma} \quad \in \quad \left[p(1-p)(\mu_j^+ - \mu_j^-)(\mu_k^+ - \mu_k^-) - q, p(1-p)(\mu_j^+ - \mu_j^-)(\mu_k^+ - \mu_k^-) + q\right] \quad . \tag{116}$$

Proof Let $S_m^{*b} \subset S_m^*$ denote the set of block-class permutations whose set of fixed points contains all examples from class $\neq b1$, $\forall b \in \{-,+\}$. We have

$$\mathbb{E}_{\varsigma \sim S_m^{*b}}\left[\sum_{l:y_l=b} x_{lj}^b x_{\varsigma(l)k}^b\right] = m_b \mu_j^b \mu_k^b \quad ,$$

and since $(1/m_b)\sum_{l:y_l=b} x_{lj}^b x_{\varsigma(l)k}^b - \mu_j^b \mu_k^b = c_{jk^\varsigma}^b$, if we sample uniformly at random $\varsigma \sim S_m^{*b}$, then we get from [Chatterjee, 2007] (Proposition 1.1):

$$\begin{aligned}
\mathbb{P}_{\varsigma \sim S_m^{*b}}[|c_{jk^\varsigma}^b| \geq t] \quad &= \quad \mathbb{P}_{\varsigma \sim S_m^{*b}}\left[\left|\sum_{l:y_l=b} x_{lj}^b x_{\varsigma(l)k}^b - m_b \mu_j^b \mu_k^b\right| \geq m_b t\right] \\
&\leq \quad 2\exp\left(-\frac{m_b t^2}{4\mu_j^b \mu_k^b + 2t}\right) \quad .
\end{aligned} \tag{117}$$

We want the right hand side to be no more than some δ_b; equivalently, we want

$$t^2 - \left(\frac{2}{m_b}\log\frac{2}{\delta_b}\right)t - \frac{2}{m_b}\log\frac{2}{\delta_b} \quad \geq \quad 0 \quad , \tag{118}$$

which holds provided

$$t \quad \geq \quad \frac{2(1+o(1))}{m_b}\log\frac{2}{\delta_b} \quad , \tag{119}$$

where the little-oh is measured wrt m_b. Since a block-class permutation factors as two fully determined permutations from S_m^{*+} and S_m^{*-}, if we fix $\delta_+ = \delta_- = \delta/2$, we get that if we sample uniformly at random these two permutations $\varsigma_+ \sim S_m^{*+}$ and $\varsigma_- \sim S_m^{*-}$, then we shall have simultaneously

$$|c_{jk^\varsigma}^b| \quad \leq \quad \frac{2(1+o(1))}{m_b}\log\frac{4}{\delta} \quad , \forall b \in \{-,+\} \quad , \tag{120}$$

which implies for the factored permutation ς,

$$\begin{aligned}
|c_{jk^\varsigma} - p(1-p)(\mu_j^+ - \mu_j^-)(\mu_k^+ - \mu_k^-)| \quad &\leq \quad \sum_b \frac{m_b}{m} \cdot \frac{2(1+o(1))}{m_b}\log\frac{4}{\delta} \\
&= \quad \frac{2(1+o(1))}{m}\log\frac{4}{\delta}
\end{aligned} \tag{121}$$

from eq. (110). Hence, if

$$m \quad = \quad \Omega\left(\frac{1}{\delta}\log\frac{1}{\delta}\right) \quad , \tag{122}$$

there will be probability $\geq 1-\delta$ that c_{jk^ς} is within additive δ from $p(1-p)(\mu_j^+ - \mu_j^-)(\mu_k^+ - \mu_k^-)$. ■

We get that with probability $\geq 1-\delta$, a randomly chosen block-class permutation ς shall make

$$\frac{\rho_{13^\varsigma} - \rho_{12}\rho_{23^\varsigma}}{\sqrt{1-\rho_{12}^2}\sqrt{1-\rho_{23^\varsigma}^2}} \leq \frac{p(1-p)}{1-\epsilon}\left(\frac{\mu_1^+ - \mu_1^-}{\sqrt{v_1}} - \rho_{12}\cdot\frac{\mu_2^+ - \mu_2^-}{\sqrt{v_2}}\right)\cdot\frac{\mu_3^+ - \mu_3^-}{\sqrt{v_3}} + \delta \ , \quad (123)$$

and the Theorem is proven (End of the proof of Theorem 27). ■

10 Experiments

10.1 Domains and setup

Domain characteristics are described in Table 2. In particular, the process for train/test split is there given in detail for each dataset. Some domains deserve more comments.

Similarly to [Hoyer et al., 2008], we consider only two features of the *abalone* datasets; those are *rings* (the age) and *length*, which are provably causally linked –*age* causes *length*–, and hence correlated. We predict the attribute *diameter* (reasonably caused by *age* as well); to turn this into a binary classification problem, we classify if the *diameter* is above or below the average one. (We also exclude abalone examples which have missing *sex* attribute.) For the experiments, we train with 200 examples and held out 567, both picked at random. The *digoxin* domain [Doran et al., 2014] is already defined by only two features, *digoxin* and *urine*, which are conditionally independent given *creatine*. From those, we predict if the level of *creatine* is above or below average. Domains Liver disorder, Auto+MPG, Arrhythmia and Diabete are part of the benchmark of domains of [Mooij et al., 2016].

The *synthetic* dataset is generated by the function `datasets.make_classification` of the *scikit-learn* python library [Pedregosa et al., 2011], with 6 features, 3 informative for the class prediction, and 3 more that are linear combinations of the formers. The rational of this toy domain is to craft two feature subspaces highly correlated.

All training sets are standardized, and the same transformation is then applied to the respective test sets. The partition of the feature space is defined by the first split $F = \lfloor d/2 \rfloor$, and its complement; features are taken in the order defined by the datasets.

Unless stated differently, models are trained with L_2 regularisation by *scikit-learn*'s `linear_model.LogisticRegression`. The hyper-parameter λ is optimized by 5-folds cross validation on the grid $\{10^{-5}, 10^{-4}, \ldots, 10^4\}$.

10.2 Explanation of the movie

Along with this draft comes a movie displaying the impact on the p-value of DT, in the context of the decrease of the HSIC. The movie shows 200 iterations of DT on the Abalone2D domain, along with the modification of the point cloud (classes are red / blue). The p-value is indicated. Notice that it begins at value 0 up to 13 digits before DT starts. The two lines indicate the classifier learnt over the current data (plain gold line) and compare with the initial classifier learnt over the data




Figure 5: Crop of the movie (see text for explanation).

before running DT (dashed gold line). The big number (167 in Figure 5) is the iteration number. Finally, the polygon displayed is the convex envelope of the initial data.

10.3 Main experiments

Our applications use the same meta-level algorithm (Algorithm 1) which operates in Setting (A) ∩ Setting (B) (M in S_m^*, linear classifiers) (Section 5), iteratively composing block-class elementary permutations. Here, $S_m^{e*} \subset S_m^*$ is the set of block-class elementary permutations. The iteration step minimises a criterion $\mathscr{G}_1$ over S_m^{e*} and potentially, after the update of the CP matrix, a criterion $\mathscr{G}_2$ over $\mathcal{H}$. The optimization of $\mathscr{G}_1$ is performed by a simple greedy search in the space of S_m^{e*}. The experimental setup (a dozen readily available domains) and results are provided *in extenso* in Section 10.4; Table 3 summarises them. The split step and the choice of $\mathcal{F}_a$ are highly domain and task dependent: to keep experiments of reasonable length, unless otherwise stated, we put in $\mathcal{F}_a$ the first half of features. In Abalone2D and Digoxin, this jams a particular ground truth (see below). Also, φ=logistic loss.

10.3.1 Disrupting dependence and causality

General experiments — We run Algorithm 1 without step 1.3, and let $\mathscr{G}_1$ be HSIC. As a proof of concept, we show that we can destroy the significance of statistical tests for independence, commonly as base for causal inference; in particular, we measure the change in the p-value computed on top of HSIC as in [Gretton et al., 2007]. We use two Gaussian kernels for K^u and K^v, each

Algorithm 1 Crossover Learning $(\mathcal{S}, T, \boldsymbol{\theta}_0, \mathrm{M}_0, \mathrm{F}^{\mathrm{a}}, \mathrm{F}^{\mathrm{s}}; \mathscr{G}_1, [\mathscr{G}_2])$

Input Sample $\mathcal{S}$, iterations T, classifier $\boldsymbol{\theta}_0$, initial CP $\mathscr{T}$ (matrices $\mathrm{M}_0 \in S_m^*$, $[\mathrm{F}^{\mathrm{a}}|\mathrm{F}^{\mathrm{s}}] = \mathrm{I}_m$);
Step 1 : **for** $t = 1, 2, ..., T$
  Step 1.1 : $\mathrm{M} \leftarrow \arg\min_{\mathrm{M}' \in S_m^{\mathrm{e}*}} \mathscr{G}_1(\mathrm{M}' \circ \mathrm{M}_{t-1}[|\boldsymbol{\theta}_{t-1}])$;
    // finds update of shuffle matrix
  Step 1.2 : $\mathrm{M}_t \leftarrow \mathrm{M} \circ \mathrm{M}_{t-1}$;
    // updates shuffle matrix
  [Step 1.3 : $\boldsymbol{\theta}_t \leftarrow \arg\min_{\boldsymbol{\theta} \in \mathbb{R}^d} \mathscr{G}_2(\boldsymbol{\theta}|\mathrm{M}_t)$;]
    // (optionally) updates classifier
Return classifier $\boldsymbol{\theta}_T$ and / or CP'ed dataset $\mathcal{S}^{\mathscr{T}(\mathrm{M}_T)}$

computed over its full subset of features (Subsection 10.1) — hence, we do not seek to alter specifically the dependence between two features, but between the two sets of features defined by the anchor and shuffle sets. On most domains (Table 3, top row), the p-value of the independence test starts close to zero at the beginning of the CP, which implies that in general both anchor and shuffle sets are (predictably) dependent. In general, we achieve a good control of the RCP and manage in several cases to decrease the true error as well through the process.

Specific dependences — The general experiments revealed that we manage to blow-up the p-value, even for domains for which the ground truth clearly *implies the alternative hypothesis* H_1. To dive into this phenomenon, we have considered two sets of experiments on which HSIC is computed over two specific features that are known to have a causal relationship. To disrupt the dependence, we thus put one of the features in the anchor set and one in the shuffle set of features (*i.e.*, after we have split the feature set in two, if both features belong to the same set, we switch one with a randomly chosen feature of the other set).

In the first set of experiments, we consider datasets with $d = 2$ features, so that this surgical disruption embeds kernels measured over the complete set of features. Experiments are reported in Table 3 for domains Abalone2D and Digoxin (Subsection 10.1). In Abalone2D for example, a gold standard for dependence [Hoyer et al., 2008], the final p-values is more than *ten billion* times the initial value. For Digoxin domain, another popular domain [Doran et al., 2014] with ground truth, p is very small at the beginning (which corresponds to the ground truth $D \not\perp\!\!\!\perp U$; D = digoxin clearance, U = urin flow). After shuffling, we obtain $p > 0.4$, which easily brings $D \perp\!\!\!\perp U$, while ground truth is $D \perp\!\!\!\perp U|C$ (C = creatinine clearance). A rather surprising fact is that in both cases, the effect on test error is minimal, considering that Abalone2D and Digoxin have $d = 2$ attributes only.

In the second set of experiments, we consider several domains with a larger d, between 7 and 279. These domains belong to the benchmarks of [Mooij et al., 2016] in which specific pairs are known to have specific causal relationships, referred to as "causal tasks", indicated in Subsection 10.1. We have targeted one causal task for each domain. Table 4 summarizes the results obtained. In all domains, the p-value is blown up at almost no expense in test error. Quite remarkably, the initial value is indeed $p = 0$ (up to *sixteen* digits), while we manage at the end of the process to get p that exceeds 1‰, which would be quite sufficient to raise doubts about the causal relationships for sensitive domains. For example, in `pair0016`, the final $p > 2$‰ might lead us to keep the independence assumption between horsepower and acceleration. In Arrhythmia, we would keep the (obviously fake) independence between age and weight. In the Liver disorder do-

main, our experiment has the following interesting consequence. Causal task `pair0034` is the causality relationship between alcohol consumption and the measure of alkaline phosphatase (ALP, [Mooij et al., 2016]). It is known that ALP elevation may be caused by heavy alcohol consumption. By keeping the independence assumption for such a value of p, one may just discourage specific blood tests related to alcohol consumption if they were to be designed from this domain. Again, the variation of the test error is minimal (if any) for all these domains.

10.3.2 Data optimisation for efficient learning

In the previous subsection, we showed how a CP may be carried out to target directly the disruption of causal relationships. In this subsection, we analyse how (and when) it can be devised to improve the test performances of classifiers. We perform Algorithm 1 with $\mathrm{M}_0 = \mathrm{I}_m$, $\mathscr{G}_1(\mathrm{M}|\boldsymbol{\theta}) = \mathbb{E}_{\mathcal{S}^{\mathscr{T}}(\mathbf{M})}\left[\varphi(y\boldsymbol{\theta}^\top \boldsymbol{x})\right]$ (Theorem 7) and $\mathscr{G}_2(\boldsymbol{\theta}|\mathrm{M}) = \mathbb{E}_{\mathcal{S}^{\mathscr{T}}}\left[\varphi(y\boldsymbol{\theta}^\top \boldsymbol{x})\right] + \lambda\|\boldsymbol{\theta}\|_2^2$ where λ is learnt through cross-validation. The algorithm returns classifier $\boldsymbol{\theta}_T$. The bottom row in Table 3, and Subsection 10.4 shows how we almost always find some permutations that reduce the test error compared to the initial data, even when a specific data optimisation should care for a risk of overfitting, which seems to occur for problems with a very small number of features (see Ionosphere and Abalone2D, Subsection 10.4). It appears also that when the (bound on the) RCP flattens, it may indicate a regime where substantial reductions can be obtained on test error as witnessed by *e.g.* Synthetic, Heart, Glass (see also BreastWisc in Table 11). We also remarked that the regime where the (bound on the) RCP flattens can be associated with a peak or decrease of the number of odd cycles as seen in Table 14, which, interestingly, can be used to provide upperbounds on the RCP as well (Theorem 10). Finally, the results on Glass display that some domains (predictably) make it possible to kill to birds in one shots at little effort: in this case, the CP both reduces the test error and increases the p-value for HSIC computed as in the general experiments of Subsection 10.3.1.

10.4 Complete experimental results

name	$m^{(*)}$	d	source	notes
digoxin	35	2	[Doran et al., 2014]	features are cond. independent given label
glass	146	9	UCI	
abalone-2D	200, 567	2	UCI, [Hoyer et al., 2008]	subsample, {rings, length} predict diameter
synthetic	200	13	scikit-learn	
heart	270	13	UCI	
liver disorders	345	7	UCI, [Mooij et al., 2016]	predict mcv $< 30^{th}$ percentile, task `pair0034`
ionosphere	351	34	UCI	
auto+mpg	398	8	UCI, [Mooij et al., 2016]	predict mpg $<$ mean, task `pair0016` (feature vectors with missing values removed)
arrhythmia	452	279	UCI, [Mooij et al., 2016]	task `pair0023`, missing values replaced by 0
breastw	683	10	UCI	
australian	690	14	UCI	
diabete(-2)	768	8	UCI	Pima domain (-2 = task `pair0038`, [Mooij et al., 2016], half of the dataset used for training)

Table 2: Domains considered. $^{(*)}$ When only one number appears in the column, 1/5 of m was hold out at random for test; when two numbers are present, the first is training set size, while the second is test size, that is fixed by the dataset description.




Table 3: Experiments performed with CP. Top row: reduction in HSIC task; bottom row: data optimisation task. References to domain names are provided in Subsection 10.1. "test-error*" is test error over initial, non shuffled data. "discrepancy" is an upperbound on the RCP provided by Theorem 9. The rightmost column aggregates the information of all curves for the task in the row, for two different domains.




Table 4: HSIC reduction on specific causal tasks [Mooij et al., 2016] (referred to as `pair00XX`); datasets indicated on pictures; "p_{init}" is the initial p-value, 0_N indicating zero up to N^{th} digit (see Table 3 for additional notations, and text for details).




Table 5: Results on domain Digoxin. Left: Data optimisation; right: HSIC reduction. Color codes are the same on all plots. See text for details.




Table 6: Results on domain Glass. Left: Data optimisation; right: HSIC reduction. Color codes are the same on all plots. Color codes are the same on all plots. See text for details.




Table 7: Results on domain Synthetic. Left: Data optimisation; right: HSIC reduction. Color codes are the same on all plots. Color codes are the same on all plots. See text for details.




Table 8: Results on domain Heart. Left: Data optimisation; right: HSIC reduction. Color codes are the same on all plots. See text for details.




Table 9: Results on domain Ionosphere. Left: Data optimisation; right: HSIC reduction. Color codes are the same on all plots. See text for details.




Table 10: Results on domain Abalone. Left: Data optimisation; right: HSIC reduction. Color codes are the same on all plots. The scale of the p-value curve is not the same as in the main file: here, its scale is the same as for the HSIC curve, which explains why it seems to be flat while the value for the first iterations is the zero-machine and the values for the last exceed one per thousand. See text for details.




Table 11: Results on domain BreastWisc. Left: Data optimisation; right: HSIC reduction. Color codes are the same on all plots. See text for details.




Table 12: Results on domain Australian. Left: Data optimisation; right: HSIC reduction. Color codes are the same on all plots. See text for details.




Table 13: Results on domain Diabete_scale. Left: Data optimisation; right: HSIC reduction. Color codes are the same on all plots. See text for details.




Table 14: Number of odd cycles (excluding fixed points, normalized by m) for the data optimization experiments in Table 3.

10.5 Comparisons of block-class vs arbitrary permutations

We now compare DT as in Algorithm 1 to the one where we relax the constraint that permutations must be block-class (implying the invariance of the mean operator). See Tables 15, 16. The results are a clear advocacy for the constraint, as relaxing it brings poor results, from both the φ-risk and test error standpoints.




Table 15: Comparison, for data optimisation, of algorithm DT in which elementary permutation matrices are constrained to be block-class (left), and *not* constrained to be block-class (right).




Table 16: Comparison (cont'd), for data optimisation, of algorithm DT in which elementary permutation matrices are constrained to be block-class (left), and *not* constrained to be block-class (right).