

Running Head: Autocorrelation in ecological data

The basis function approach for modeling autocorrelation in ecological data

Trevor J. Hefley¹

Department of Fish, Wildlife, and Conservation Biology, Colorado State University

Department of Statistics, Colorado State University

Kristin M. Broms

Department of Fish, Wildlife, and Conservation Biology, Colorado State University

Brian M. Brost

Department of Fish, Wildlife, and Conservation Biology, Colorado State University

Frances E. Buderman

Department of Fish, Wildlife, and Conservation Biology, Colorado State University

Shannon L. Kay

Department of Statistics, Colorado State University

Henry R. Scharf

Department of Statistics, Colorado State University

John R. Tipton

Department of Statistics, Colorado State University

Perry J. Williams

Department of Fish, Wildlife, and Conservation Biology, Colorado State University

Department of Statistics, Colorado State University

Mevin B. Hooten

U.S. Geological Survey, Colorado Cooperative Fish and Wildlife Research Unit

¹Corresponding author; E-mail: Trevor.Hefley@colostate.edu.

Department of Fish, Wildlife, and Conservation Biology, Colorado State University

Department of Statistics, Colorado State University

ABSTRACT

Analyzing ecological data often requires modeling the autocorrelation created by spatial and temporal processes. Many of the statistical methods used to account for autocorrelation can be viewed as regression models that include basis functions. Understanding the concept of basis functions enables ecologists to modify commonly used ecological models to account for autocorrelation, which can improve inference and predictive accuracy. Understanding the properties of basis functions is essential for evaluating the fit of spatial or time-series models, detecting a hidden form of multicollinearity, and analyzing large data sets. We present important concepts and properties related to basis functions and illustrate several tools and techniques ecologists can use when modeling autocorrelation in ecological data.

Key words: autocorrelation; Bayesian model; collinearity; dimension reduction; semiparametric regression; spatial statistics; time series

INTRODUCTION

Ecological processes interact at multiple temporal and spatial scales, generating complex spatio-temporal patterns (Levin 1992). The science of ecology is concerned with understanding, describing, and predicting components of these spatio-temporal patterns using limited and noisy observations. An important consideration when developing an ecological model is how to include the spatial, temporal, or spatio-temporal aspects of the process (Legendre 1993). For example, species distribution models are used to understand

and predict how the abundance of plants and animals varies across space and time (Elith and Leathwick 2009). The abundance of a species within a patch of habitat might depend on environmental covariates (e.g., minimum annual temperature), but might also depend on the abundance in surrounding patches. Dispersal of organisms among habitat patches can make abundance in nearby patches more similar than could be explained by environmental covariates alone (Legendre and Fortin 1989). Correlation among observations that depends on the proximity (in space, time, or both) of neighboring observations is known as autocorrelation (Table 1). Disentangling autocorrelation from environmental covariates is critical to understanding endogenous and exogenous factors that influence populations and ecosystems (Borcard et al. 1992). Moreover, properly accounting for autocorrelation is necessary for obtaining reliable statistical inference (e.g., Fieberg and Ditmer 2012).

Isolating the effect of autocorrelation in an ecological model can be accomplished by including a function that captures the dependence between observations that are close in space or time. The mathematical form of the function that best describes the autocorrelation is always unknown and may be complex, but can be approximated by a combination of simple basis functions. Most ecologists have encountered basis functions (e.g., polynomial regression), but may not be aware of the breadth of situations in which they can be used to model autocorrelation in ecological data. For example, basis functions are used in semiparametric models, such as generalized additive models, but are also implicitly used in spatial or time-series models. Understanding how basis functions can be used to model autocorrelation is essential for evaluating the fit of spatial, time-series, or spatio-temporal models, detecting a hidden form of multicollinearity (Hodges and Reich 2010), and facilitating the analysis of large data sets (Wikle 2010). More importantly,

understanding the basis function approach will enable ecologists to tailor many commonly used models to account for autocorrelation, which can improve both inference and predictive accuracy (e.g., Hooten et al. 2003; Conn et al. 2015; Buderman et al. 2016).

We have three goals in this paper: 1) introduce concepts and terminology related to basis functions and autocorrelation; 2) demonstrate the connections between commonly used methods to model autocorrelation; and 3) develop a working knowledge of the basis function approach so ecologists can devise ways to model autocorrelation in commonly used ecological models. We first introduce basis functions and then the concepts of first-order and second-order model specifications. To illustrate these concepts, we present three examples: a standard regression model, a time-series model, and a spatial model, each applied to different types of ecological data. We include supplementary material comprised of a tutorial that contains additional descriptions, data, and example computer code.

BASIS FUNCTIONS

Consider a simple linear regression model of pelagic bioluminescence density (sources) as a function of water depth (Fig. 1a; Gillibrand et al. 2007)

$$\mathbf{y} = \alpha_0 \mathbf{z}_0 + \alpha_1 \mathbf{z}_1 + \boldsymbol{\varepsilon} , \quad (1)$$

where \mathbf{y} is an $n \times 1$ vector of bioluminescence density, \mathbf{z}_0 is an $n \times 1$ vector of ones, \mathbf{z}_1 is an $n \times 1$ vector that contains the depth in meters of the observed bioluminescence sources, α_0 is the intercept, α_1 is a regression coefficient, and $\boldsymbol{\varepsilon}$ is an $n \times 1$ vector that contains independent and normally distributed error terms with variance σ_{ε}^2 (i.e., $\boldsymbol{\varepsilon} \sim N(\mathbf{0}, \sigma_{\varepsilon}^2 \mathbf{I})$, where \mathbf{I} is the $n \times n$ identity matrix). In this simple linear regression model, the basis coefficients are α_0 and α_1 and the basis vectors \mathbf{z}_0 and \mathbf{z}_1 are the depths raised to the

power 0 and 1 (Table 1). It may not be common to refer to transformations of a covariate as basis vectors; however, the transformations form a “basis” of possible values in covariate space. The function that transforms a covariate into a basis vector is referred to as a basis function. Although the terms basis function and basis vector tend to be used interchangeably in the literature, basis functions typically exist in continuous space, whereas basis vectors are the output from a function and exist in discrete space (e.g., where depth was measured; Table 1; see Appendix A for further discussion). All transformations of a covariate are known collectively as a basis expansion (e.g., \mathbf{z}_0 and \mathbf{z}_1). Finally, as in simple linear regression, the expected density of bioluminescence at the measured depths is the linear combination of basis vectors $\alpha_0\mathbf{z}_0 + \alpha_1\mathbf{z}_1$ (Fig. 1a).

It is clear from Fig. 1a that the simple linear regression model does not adequately capture the relationship between bioluminescence and depth. A more flexible basis expansion that better captures the relationship is the polynomial regression model that includes the quadratic effect of depth

$$\mathbf{y} = \alpha_0\mathbf{z}_0 + \alpha_1\mathbf{z}_1 + \alpha_2\mathbf{z}_2 + \boldsymbol{\varepsilon} , \quad (2)$$

where α_2 is the basis coefficient for the squared effect of depth (\mathbf{z}_2 ; Fig. 1b). Some models that use basis functions can be respecified, which can aid in interpretation and increase computational efficiency and stability. For example, we can respecify Eq. 2 using a different basis expansion, but in a way that yields the exact same model

$$\mathbf{y} = \alpha_1(\mathbf{z}_1 - k_1)^2 + \alpha_2(\mathbf{z}_1 - k_2)^2 + \alpha_3(\mathbf{z}_1 - k_3)^2 + \boldsymbol{\varepsilon} , \quad (3)$$

where \mathbf{z}_1 is an $n \times 1$ vector of the observed depths in meters, k_j is the j^{th} depth of interest ($j = 1, 2, 3$), and α_j is the basis coefficient. The two basis expansions in Eq. 2 and Eq. 3 have different basis vectors and will yield different estimates of α_j , but result in the exact same polynomial curve when fit to the data. For example, let $k_1 = 1140$ m, $k_2 = 2620$ m, and $k_3 = 3420$ m, and compare the basis vectors and predicted bioluminescence (cf., Fig. 1b and 1c). An interactive figure of this example can be found in Appendix B.

Even if the specifications result in the exact same model, there are many reasons why one basis expansion might be preferred over others. For example, the number of parameters in the model can be reduced if a basis expansion results in some basis coefficients that can be assumed to be zero (see confidence intervals for α_j in Appendix B). In addition, some basis expansions might have scientific interpretations. For example, the model in Eq. 3 states that the density of bioluminescence is a function of the distance between the observed depth and three locations in the water column, that we might believe are biologically important. Finally, some basis expansions may reduce the correlation among basis vectors. For example, the coefficient of determination (R^2) for the basis vectors \mathbf{z}_1 and \mathbf{z}_2 in Eq. 2 is 0.96, whereas, the maximum R^2 among the three basis vectors in Eq. 3 is 0.25.

Model assumptions

A critical assumption is that a linear combination of basis vectors adequately approximates the unknown relationship between the observed response and basis vectors. In a regression context, this assumption is analogous to assuming the model includes all covariates that linearly influence the response. For example, it is assumed in Eq. 2 and Eq. 3 that a linear combination of the basis vectors (e.g., $\alpha_0\mathbf{z}_0 + \alpha_1\mathbf{z}_1 + \alpha_2\mathbf{z}_2$) adequately represents the unknown functional relationship between depth and the density of bioluminescence. A

result of this assumption is that the basis vectors must span the space of interest, otherwise the model is inadequate. In the bioluminescence example, the space of interest is depth and the bases in Eq. 2 and Eq. 3 span the set of all second-degree polynomial functions. That is, we assume the true underlying relationship between the density of bioluminescence and depth can be appropriately modeled as a second-degree polynomial. Because Eq. 2 and Eq. 3 are both second-degree polynomials and the two basis expansions span the same space, the estimated curves are exactly the same (cf., Fig. 1b and 1c). If one or more of the basis vectors were removed from the polynomial regression model, the basis vectors would not span the same space of all second-degree polynomials and therefore could not create the exact curve shown in Fig. 1b and 1c.

Generalizations

Now consider an unknown function $\eta(x)$ that describes a pattern or process in nature that generates autocorrelation. This function can be continuous or discrete, but the true form is always unknown. For example, $\eta(x)$ might describe the similarity in abundance between habitat patches in geographic space, population regulation influenced by endogenous factors in temporal space (e.g., density dependence), or how net primary productivity changes with temperature in covariate space. Although the true form of the function $\eta(x)$ is unknown, we can approximate it with a combination of simple basis functions. We can combine the basis functions in a linear equation such that $\eta(x) \approx \sum_{j=1}^m \alpha_j f_j(x)$; this is a general notation that contains m basis functions $f_j(x)$ ($j = 1, \dots, m$). For example, in the polynomial regression model (Eq. 2), $f_1(x) = x^0$, $f_2(x) = x^1$, and $f_3(x) = x^2$. In what follows, we use matrix notation and write $\boldsymbol{\eta} \equiv \mathbf{Z}\boldsymbol{\alpha}$, where $\boldsymbol{\eta}$ is an $n \times 1$ vector representing an approximation of the unknown function $\eta(\mathbf{x})$ at the n locations \mathbf{x} in the space of interest, \mathbf{Z} is an $n \times m$ matrix containing the basis vectors, and $\boldsymbol{\alpha}$ is an $m \times 1$ vector of

basis coefficients. We also use the matrix notation $\mathbf{X}\boldsymbol{\beta}$, where \mathbf{X} is an $n \times p$ matrix of traditional covariates and $\boldsymbol{\beta}$ is a $p \times 1$ vector of traditional regression coefficients. Our notation is used to identify when a basis vector should be treated as a traditional covariate (and included in \mathbf{X}) or treated as a formal basis expansion (and included in \mathbf{Z}). In some applications, there is no practical difference between including basis vectors in \mathbf{X} or \mathbf{Z} , whereas, in other applications, the choice of notation is used to designate whether the coefficients are treated as fixed (\mathbf{X}) or random (\mathbf{Z}) effects.

MODEL SPECIFICATIONS

Specifying a statistical model involves combining probability distributions and deterministic mathematical equations, both with unknown parameters, in a way that properly describes the ecological process (Hobbs and Hooten 2015). As we demonstrated in the polynomial regression example, there are multiple ways to specify the exact same model (e.g., Eq. 2 and Eq. 3). An important concept related to specifying a model is first-order and second-order model specifications (Table 1; Cressie and Wikle 2011; Hobbs and Hooten 2015), which is also known in the mixed-model literature as G-side and R-side specifications (Littell et al. 2006; Stroup 2012). Specifically, this concept relates to placing a function that describes the autocorrelation in either the mean (expected value) or the covariance of a probability distribution. The concept of first-order and second-order model specification is important for understanding the equivalence of certain spatial and time-series models, efficient implementation of Bayesian models that account for autocorrelation, and for detecting a hidden form of multicollinearity. A general understanding of hierarchical models or mixed models is necessary for what follows. Introductions to hierarchical models or mixed models include: Hobbs and Hooten (2015), Hodges (2013), Littell et al. (2006),

Ruppert et al. (2003), Stroup (2012), Wood (2006), and Zuur et al. (2009).

First-order specification

A first-order specification defines a function that describes the autocorrelation in the mean structure of a distribution (Table 1). For example, consider the linear regression model $\mathbf{y} = \mathbf{X}\boldsymbol{\beta} + \boldsymbol{\varepsilon}$, where $\mathbf{y} \equiv (y_1, \dots, y_n)'$ are the observed data. Assuming independent and normally distributed errors ($\boldsymbol{\varepsilon} \sim N(\mathbf{0}, \sigma_\varepsilon^2 \mathbf{I})$), we can write this model as

$$\mathbf{y} \sim N(\mathbf{X}\boldsymbol{\beta}, \sigma_\varepsilon^2 \mathbf{I}) . \quad (4)$$

An assumption of Eq. 4 is that a linear combination of the covariates serves as a good approximation to the underlying relationship. If there is evidence that the model in Eq. 4 is inadequate (e.g., correlation among the residual errors), a linear combination of basis vectors may be added to the mean to improve model fit and satisfy model assumptions, such that

$$\mathbf{y} \sim N(\mathbf{X}\boldsymbol{\beta} + \mathbf{Z}\boldsymbol{\alpha}, \sigma_\varepsilon^2 \mathbf{I}) . \quad (5)$$

The basis expansion (\mathbf{Z}) account for additional complexity in the mean structure that is not explained by the covariates in \mathbf{X} . For example, \mathbf{Z} could account for the lack of fit (spatial autocorrelation) in the linear or polynomial model in the bioluminescence example (Fig. 1; Table 1).

Second-order specification

A second-order specification defines a function that describes the autocorrelation in the covariance of a probability distribution. For example, consider the linear model

$\mathbf{y} = \mathbf{X}\boldsymbol{\beta} + \boldsymbol{\eta} + \boldsymbol{\varepsilon}$ where $\boldsymbol{\varepsilon} \sim N(\mathbf{0}, \sigma_\varepsilon^2 \mathbf{I})$, $\boldsymbol{\eta} \sim N(\mathbf{0}, \sigma_\alpha^2 \mathbf{R})$, and the random effect $\boldsymbol{\eta}$ results in

correlated errors. We can write the model as:

$$\mathbf{y} \sim N(\mathbf{X}\boldsymbol{\beta}, \sigma_\varepsilon^2\mathbf{I} + \sigma_\alpha^2\mathbf{R}), \quad (6)$$

where \mathbf{R} is a correlation matrix that accounts for autocorrelation among observations. The correlation matrix \mathbf{R} is often specified using a correlation function that depends on a distance measure between two observations in the space of interest (Table 1).

Equivalent specifications

In some situations, the first-order and second-order specifications result in the same model.

To demonstrate this for a specific case, we make the additional assumption that the basis coefficients in Eq. 5 are normally-distributed random effects (i.e., $\boldsymbol{\alpha} \sim N(\mathbf{0}, \sigma_\alpha^2\mathbf{I})$).

Equivalent probability density functions can be obtained by integrating out the random effect $\boldsymbol{\alpha}$ from the first-order specification in Eq. 5 to yield

$$\begin{aligned} \mathbf{y} &\sim \int N(\mathbf{X}\boldsymbol{\beta} + \mathbf{Z}\boldsymbol{\alpha}, \sigma_\varepsilon^2\mathbf{I})N(\mathbf{0}, \sigma_\alpha^2\mathbf{I})d\boldsymbol{\alpha} \\ &= N(\mathbf{X}\boldsymbol{\beta}, \sigma_\varepsilon^2\mathbf{I} + \sigma_\alpha^2\mathbf{Z}\mathbf{Z}'), \end{aligned} \quad (7)$$

where equivalence between the first- and second-order specifications holds if the correlation matrix \mathbf{R} is the outer product of the basis expansion \mathbf{Z} (i.e., $\mathbf{R} \equiv \mathbf{Z}\mathbf{Z}'$; see example below).

The integration in Eq. 7 effectively “moves” the autocorrelation modeled by the basis vectors in the mean structure to the covariance structure. For example, consider a mixed-effects model where \mathbf{Z} is used to represent autocorrelation due to a site or grouping effect among observations:

$$\mathbf{Z} = \begin{bmatrix} 1 & 0 & 0 \\ 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \\ 0 & 0 & 1 \end{bmatrix}, \quad (8)$$

where, y_1 and y_2 were observed at the first site, y_3 and y_4 were observed at the second site, etc. If we assume the basis coefficients are normally-distributed random effects, then

$$\mathbf{R} = \mathbf{Z}\mathbf{Z}' = \begin{bmatrix} 1 & 0 & 0 \\ 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \\ 0 & 0 & 1 \end{bmatrix} \begin{bmatrix} 1 & 1 & 0 & 0 & 0 & 0 \\ 0 & 0 & 1 & 1 & 0 & 0 \\ 0 & 0 & 0 & 0 & 1 & 1 \end{bmatrix} = \begin{bmatrix} 1 & 1 & 0 & 0 & 0 & 0 \\ 1 & 1 & 0 & 0 & 0 & 0 \\ 0 & 0 & 1 & 1 & 0 & 0 \\ 0 & 0 & 1 & 1 & 0 & 0 \\ 0 & 0 & 0 & 0 & 1 & 1 \\ 0 & 0 & 0 & 0 & 1 & 1 \end{bmatrix}, \quad (9)$$

where \mathbf{R} is called the compound symmetry correlation matrix (Littell et al. 2006; Zuur et al. 2009; Stroup 2012). The model that is obtained by using the first-order specification in Eq. 8 is the exact same model that would be obtained by using the correlation matrix in Eq. 9.

Although one may start with a basis expansion \mathbf{Z} , many methods developed to model autocorrelation start by choosing a correlation matrix \mathbf{R} . When starting with a correlation matrix \mathbf{R} , a basis expansion \mathbf{Z} can be obtained using matrix decomposition methods (e.g., spectral decomposition; Lorenz 1956; Cressie and Wikle 2011; pg. 156). For example,

consider the regression model in Eq. 6, and let $\mathbf{R}(\phi)$ be an order-one autoregressive correlation matrix (AR(1))

$$\mathbf{R}(\phi) = \begin{bmatrix} 1 & \phi^1 & \phi^2 & \dots & \phi^{t-1} \\ \phi^1 & 1 & \phi^1 & \dots & \phi^{t-2} \\ \phi^2 & \phi^1 & 1 & \vdots & \phi^{t-3} \\ \vdots & \vdots & \vdots & \ddots & \vdots \\ \phi^{t-1} & \phi^{t-2} & \phi^{t-3} & \ddots & 1 \end{bmatrix}, \quad (10)$$

where $-1 < \phi < 1$. The AR(1) correlation matrix in Eq. 10 is commonly used in time-series analysis to model temporal correlation that diminishes exponentially with a rate of decay that depends on ϕ . When a correlation matrix, or basis expansion depends on parameters, we include the parameters in parentheses (e.g., $\mathbf{R}(\phi)$ and $\mathbf{Z}(\phi)$).

A correlation matrix can be decomposed to produce basis vectors that are useful in the first-order specification. One approach to obtain basis vectors from $\mathbf{R}(\phi)$ is the spectral decomposition: $\mathbf{R}(\phi) = \mathbf{Q}\mathbf{\Lambda}\mathbf{Q}'$, where \mathbf{Q} are the eigenvectors and $\mathbf{\Lambda}$ is a diagonal matrix with elements that contain the eigenvalue associated with each eigenvector (Cressie and Wikle 2011; pgs. 156-157). Using the spectral decomposition, the basis expansion can be written as $\mathbf{Z}(\phi) = \mathbf{Q}\mathbf{\Lambda}^{\frac{1}{2}}$. As an example, for three observations $\mathbf{y} \equiv (y_1, y_2, y_3)'$ collected at times $t = 1, 2, 3$, the AR(1) correlation matrix using $\phi = 0.5$ is

$$\mathbf{R}(\phi) = \begin{bmatrix} 1 & 0.5 & 0.25 \\ 0.5 & 1 & 0.5 \\ 0.25 & 0.5 & 1 \end{bmatrix}. \quad (11)$$

The spectral decomposition of $\mathbf{R}(\phi)$ is

$$\mathbf{R}(\phi) = \mathbf{Q}\mathbf{\Lambda}\mathbf{Q}' = \begin{bmatrix} -0.54 & -0.71 & 0.45 \\ -0.64 & 0 & -0.77 \\ -0.54 & 0.71 & 0.45 \end{bmatrix} \begin{bmatrix} 1.84 & 0 & 0 \\ 0 & 0.75 & 0 \\ 0 & 0 & 0.41 \end{bmatrix} \begin{bmatrix} -0.54 & -0.64 & -0.54 \\ -0.71 & 0 & 0.71 \\ 0.45 & -0.77 & 0.45 \end{bmatrix}. \quad (12)$$

The matrices of eigenvectors (\mathbf{Q}) and eigenvalues ($\mathbf{\Lambda}$) in Eq. 12 can be used to construct the basis expansion

$$\mathbf{Z}(\phi) = \mathbf{Q}\mathbf{\Lambda}^{\frac{1}{2}} = \begin{bmatrix} -0.74 & -0.61 & 0.29 \\ -0.87 & 0 & -0.49 \\ -0.74 & 0.61 & 0.29 \end{bmatrix}. \quad (13)$$

Alternatively, one might use the eigenvectors \mathbf{Q} as basis vectors (i.e., $\mathbf{Z}(\phi) \equiv \mathbf{Q}(\phi)$; Griffith and Peres-Neto 2006), which would require specifying a non-constant variance for the basis coefficients such that $\boldsymbol{\alpha} \sim \mathbf{N}(\mathbf{0}, \sigma_{\alpha}^2 \mathbf{\Lambda})$. Converting models between first- and second-order specifications using the techniques we have presented is critical for harnessing the full power of basis functions for modeling autocorrelation in ecological data.

Generalized models

Basis functions can be used with any response distribution (e.g., Poisson, binomial). For example, the generalized linear mixed model can include random effects for the coefficients associated with the basis vectors that account for autocorrelation (Bolker et al. 2009).

Similarly, basis functions can also be embedded within Bayesian hierarchical models to account for autocorrelation (see Example 3). In all of the examples presented below, we assume that the basis coefficients are normally-distributed random effects; however, this is not a necessary assumption. As with the generalized linear mixed model or Bayesian

hierarchical models, distributions for random effects other than the normal could be used (e.g., gamma, t-distribution; Higdon 2002; Lee et al. 2006; Johnson et al. 2013; Gelman et al. 2013; Hobbs and Hooten 2015).

EXAMPLE 1: PELAGIC BIOLUMINESCENCE VERSUS DEPTH GRADIENT

In the Basis Functions section, we initially modeled the density of bioluminescence using depth or transformations of depth as covariates with coefficients that were fixed effects. Depth can also be thought of as the spatial location in the water column, so it is natural to model the density of bioluminescence using a spatial model instead of regressing on depth directly. In the three following model specifications that we use to capture spatial autocorrelation, \mathbf{X} is an $n \times 1$ matrix of ones and β is a constant intercept term (i.e., $\mathbf{X}\beta = \beta_0$). As a result, the influence of depth will be modeled in either the basis expansion \mathbf{Z} or the correlation matrix \mathbf{R} , in accordance with the first-order or second-order specification, respectively.

Spatial regression model: a second-order specification

Consider the model in Eq. 6 where the correlation matrix $\mathbf{R}(\phi)$ is specified using a parametric correlation function that depends on a range parameter ϕ (Cressie and Wikle 2011; Banerjee et al. 2014). The range parameter controls how correlation diminishes as the distance between locations increases. For this example, we use a Gaussian correlation function

$$r_{ij}(\phi) = e^{-\frac{d_{ij}^2}{\phi}}, \quad (14)$$

where d_{ij} is the distance between locations i and j (note that $d_{ij} = 0$ for $i = j$) and $r_{ij}(\phi)$ is the element in the i^{th} row and j^{th} column of $\mathbf{R}(\phi)$. For the bioluminescence example, d_{ij} is the difference in depth between observations i and j . Given the second-order

specification, it is not immediately clear how to estimate the influence of depth on bioluminescence (i.e., $\beta_0 + \boldsymbol{\eta}$), which requires the spatial random effect $\boldsymbol{\eta}$. To predict bioluminescence at the observed (and unobserved) depths using the second-order specification, we used best linear unbiased prediction (BLUP; Kriging in the spatial context; see Robinson 1991 for derivation). The predicted spatial random effect ($\boldsymbol{\eta}$) for the observed depths, given estimates of all other parameters, can be obtained using

$$\hat{\boldsymbol{\eta}} = \hat{\sigma}_\alpha^2 \mathbf{R}(\hat{\phi}) \left(\hat{\sigma}_\varepsilon^2 \mathbf{I} + \hat{\sigma}_\alpha^2 \mathbf{R}(\hat{\phi}) \right)^{-1} \left(\mathbf{y} - \mathbf{X}\hat{\boldsymbol{\beta}} \right). \quad (15)$$

The fitted spatial model (Fig. 2a; $\hat{\beta}_0 + \hat{\boldsymbol{\eta}}$) captures fine scale (local) variability better than the polynomial regression model (Fig. 1b and 1c; Appendix C).

Spatial regression model: a first-order specification

The spatial regression model can also be implemented using basis vectors. Consider the first-order specification from Eq. 5 where $\boldsymbol{\alpha} \sim N(\mathbf{0}, \sigma_\alpha^2 \mathbf{I})$ and $\mathbf{Z}(\phi)$ is obtained from a spectral decomposition of $\mathbf{R}(\phi)$. Using the BLUP from Eq. 15, basis coefficients are equivalent to

$$\hat{\boldsymbol{\alpha}} = \hat{\sigma}_\alpha^2 \mathbf{Z}(\hat{\phi})' \left(\hat{\sigma}_\varepsilon^2 \mathbf{I} + \hat{\sigma}_\alpha^2 \mathbf{Z}(\hat{\phi}) \mathbf{Z}(\hat{\phi})' \right)^{-1} \left(\mathbf{y} - \mathbf{X}\hat{\boldsymbol{\beta}} \right). \quad (16)$$

Note that because $\mathbf{R}(\hat{\phi}) \equiv \mathbf{Z}(\hat{\phi}) \mathbf{Z}(\hat{\phi})'$, by definition $\mathbf{Z}(\hat{\phi}) \hat{\boldsymbol{\alpha}}$ is the same as $\hat{\boldsymbol{\eta}}$ in Eq. 15. The bioluminescence for any given depth (i.e., $\hat{\beta}_0 + \mathbf{Z}(\hat{\phi}) \hat{\boldsymbol{\alpha}}$) from the first-order specification is shown in Fig. 2b along with three of the eigen basis vectors multiplied by the corresponding basis coefficients. The fitted values from the first- and second-order specifications are exactly the same (cf., Fig. 2a to 2b) because both specifications result in an equivalent model.

Even if the initial model formulation is a second-order specification that uses a correlation function, the equivalent first-order specification is often useful. Three important uses for the first-order specification are: 1) it allows for assessment of collinearity between covariates in \mathbf{X} and basis vectors in $\mathbf{Z}(\phi)$ (see Example 2); 2) basis vectors can be visualized and certain types of basis expansions have useful ecological interpretation (e.g., Griffith and Peres-Neto 2006); and 3) certain types of basis expansions are useful for dimension reduction required to fit models to large data sets (Wikle, 2010). We will demonstrate the utility of the first-order specification in the following examples.

Modeling spatial autocorrelation using kernel basis functions

Another method that can be used to model autocorrelation is kernel regression. Kernel regression is a semiparametric regression technique widely used by statisticians and the machine learning community (Bishop, 2006; Hastie et al., 2009; James et al., 2013). Regression models that employ kernel basis functions are written using the first-order specification. The commonly used Gaussian kernel basis function is defined as:

$$z_{ij}(\phi) = e^{-\frac{2d_{ij}^2}{\phi}}, \quad (17)$$

where $z_{ij}(\phi)$ is the element in the i^{th} row and j^{th} column of $\mathbf{Z}(\phi)$, and d_{ij} is the distance between the i^{th} data point and the j^{th} knot ($j = 1, \dots, m$ where m is the number of basis vectors). Knots are locations in the space of interest where each basis vector is anchored (e.g., k_j in the bioluminescence example; Table 1). In Fig. 2c, we show the density of bioluminescence predicted from the kernel regression model (i.e., $\hat{\beta}_0 + \mathbf{Z}(\hat{\phi})\hat{\alpha}$) and three of its basis vectors (from a total $m = 17$) multiplied by the corresponding basis coefficients estimated using Eq. 16. Comparison of the eigen and kernel basis vectors reveals that the

two types of basis functions look different, but the fitted curves are nearly equivalent (cf., Fig. 2b and 2c). Importantly, as the number of basis vectors and knots increases to infinity (on a grid), the first-order model specification that uses a Gaussian kernel basis function (Eq. 17) converges to the second-order specification that uses a Gaussian correlation function (Eq. 14; Higdon 2002). An interactive figure that allows users to experiment with basis functions for this example can be found in Appendix B.

Regression models that use kernel basis functions are useful because they allow for more complex correlation structures compared to models that rely on correlation functions (Higdon 2002; Sampson 2010; Table 2). Further, the number of basis vectors and coefficients can be controlled by the user depending on the level of computational efficiency required. Choosing the dimension of the basis expansion (m) to be less than n is known as dimension reduction. For example, the spatial model that relies on a second-order specification and uses a correlation function can be converted to a first-order model that requires 51 eigen basis vectors, but the kernel regression uses a pre-selected number of kernel basis vectors (17 for this example). Dimension reduction usually relies on first-order model specification and is helpful for modeling autocorrelation in large data sets, allowing for statistical inference that would otherwise be computationally infeasible.

EXAMPLE 2: POPULATION TREND

A common task in applied ecology is to infer if the abundance of a species is declining over time. This task often involves fitting a trend to a time series of abundance indices and inferring if the associated regression coefficient is negative. For this example, we use the simple linear model

$$\mathbf{y} \sim N(\beta_0 + \beta_1 \mathbf{t}, \sigma_\epsilon^2 \mathbf{I}), \quad (18)$$

where \mathbf{y} is a $t \times 1$ vector containing the count or index of population size from each time period, and \mathbf{t} is the corresponding vector of times.

Bobwhite quail (*Colinus virginianus*) are a common species that occurs throughout a large portion of the United States, but are declining in abundance in many regions (Veech 2006). We present an index of bobwhite quail population size obtained from annual whistle-count surveys in Nemaha County, Nebraska (Fig. 3a; see Hefley et al. 2013 for a detailed description of the data). We fit a simple linear regression model to these data using maximum likelihood which results in $\hat{\beta}_1 = -1.16$ and a 95% confidence interval of $[-1.88, -0.44]$. The fitted regression model suggests a decline in abundance; however, a clear pattern in the residuals is present, possibly due to underlying population dynamics of bobwhite quail (Fig. D1; Appendix D). If autocorrelation is ignored, the uncertainty associated with regression coefficients will be underestimated and may cause the researcher to overstate the statistical significance of the decline (Cressie 1993; Hoeting 2009). The underlying population dynamics that generated the autocorrelation are likely complex and building mechanistic models that account for the process is challenging (Hefley et al. 2013). A simpler approach for modeling the endogenous population dynamics is to assume that the population size (or some transformation thereof) can be modeled as

$\mathbf{y} \sim N(\beta_0 + \beta_1 \mathbf{t}, \sigma_\alpha^2 \mathbf{R}(\phi) + \sigma_\epsilon^2 \mathbf{I})$, where $\mathbf{R}(\phi)$ can be any appropriate correlation matrix.

When we account for the autocorrelation using the AR(1) correlation matrix (Eq. 10), we obtain $\hat{\beta}_1 = -1.10$ and a 95% confidence interval of $[-2.61, 0.41]$. The 95% confidence interval now covers zero and is approximately twice as wide compared to the model that does not account for autocorrelation. The fitted trend lines for the two models ($\hat{\beta}_0 + \hat{\beta}_1 \mathbf{t}$) appear nearly identical, but when the temporal process $\boldsymbol{\eta}$ is included ($\hat{\beta}_0 + \hat{\beta}_1 \mathbf{t} + \hat{\boldsymbol{\eta}}$; where $\boldsymbol{\eta}$ is estimated using Eq. 15) the fit is much better because the residuals appear to be

uncorrelated (see Fig. D1-D4 in Appendix D).

To demonstrate two different first-order model specifications that account for temporal autocorrelation, we use eigen basis vectors obtained from a spectral decomposition of the the AR(1) correlation matrix (Fig. 3b), as well as compactly supported uniform kernel basis function with knots placed at each year (Fig. 3c; Appendix D). In contrast to the spatial model used in the bioluminescence example, the AR(1) correlation matrix does not have a corresponding kernel that can be used as an approximation. Consequently, the first-order model that uses a uniform kernel basis function results in a different fit to the data when compared to the models that used an AR(1) correlation matrix (cf., Fig. 3b to 3c). Both models, however, appear to capture the temporal autocorrelation and result in similar estimates ($\hat{\beta}_1 = -1.28$, 95% confidence interval $[-2.59, 0.03]$ for the uniform kernel basis function model).

This example also demonstrates that it is important to check for multicollinearity between basis vectors and covariates when modeling autocorrelation. As with traditional regression models, severe multicollinearity can negatively influence inference (Dormann et al. 2013). The potential for multicollinearity is evident in the first-order specification, however, the collinearity is effectively “hidden” in the correlation matrix of the second-order specification (Hodges and Reich 2010; Hanks et al. 2015). For example, the coefficient of determination between the covariate year (\mathbf{t} in Eq. 18) and second eigen basis vector is $R^2 = 0.80$.

EXAMPLE 3: PREDICTING THE DISTRIBUTION OF A SPECIES

In this example, we fit three different models that account for spatial autocorrelation to illustrate concepts presented in previous sections. Many ecological studies aim to predict

the presence or abundance of a species at unsampled locations using species distribution models applied to count, presence-absence, and presence-only data (Elith and Leathwick 2009). Generalized linear mixed models with a spatial random effect are well-suited to model a species distribution using count or presence-absence data (Bolker et al. 2009). For example, Hooten et al. (2003) used a binary spatial regression model to predict the probability of pointed-leaved tick trefoil (*Desmodium glutinosum*) occurring in 10×10 m plots across a 328 ha area from presence-absence data collected at 216 plots (Fig. 4a). A common problem when predicting the distribution of a species is that data are sparse relative to the prediction domain. For this example, only 0.66% of the plots within the prediction domain were sampled (Fig. 4a). In this application, Hooten et al. (2003) specified a second-order spatial random effect to increase the predictive ability of a binary regression model and to account for spatial autocorrelation generated by a complex ecological process. A suitable second-order spatial binary model for presence-absence data is

$$\begin{aligned} \mathbf{y} &\sim \text{Bernoulli}(g(\mathbf{X}\boldsymbol{\beta} + \boldsymbol{\eta})) \\ \boldsymbol{\eta} &\sim \text{N}(\mathbf{0}, \sigma_{\alpha}^2 \mathbf{R}(\phi)) , \end{aligned} \tag{19}$$

where \mathbf{y} is an $n \times 1$ vector with elements equal to 1 if the species is present and 0 if the species is absent at a sampled location, $g(\cdot)$ is the probit link function, and $\boldsymbol{\eta}$ is the spatial random effect. As in Hooten et al. (2003), we specified the correlation matrix in Eq. 19 using an exponential correlation function

$$r_{ij}(\phi) = e^{-\frac{d_{ij}}{\phi}} , \tag{20}$$

where d_{ij} is the distance between locations i and j . Although there are numerous ways to

implement the binary regression model, we adopt a Bayesian approach. The associated prior distributions and covariates are described in Hooten et al. (2003). After fitting the model, we predicted the probability of occurrence at all 32,768 plots within the prediction domain. The predicted probability of occurrence depends on several covariates and has a strong spatial component (Fig. 4b). Unlike the Gaussian kernel basis function which approximates a Gaussian correlation function (see bioluminescence example), there is no kernel basis function that can approximate an exponential covariance function (see Fig. 2 in Higdon, 2002 for details).

Evaluating the likelihood for any second-order model requires inverting the correlation matrix $\mathbf{R}(\phi)$. For the geostatistical (continuous space) spatial model, inverting the correlation matrix has a computational cost that increases according to the cube of the sample size. For this example, when $n = 216$, fitting the Bayesian spatial model requires approximately 45 seconds per 1,000 Markov chain Monte Carlo (MCMC) samples obtained on a laptop computer with a 2.8 GHz quad-core processor, 16 GB of RAM, and optimized basic linear algebra subprograms, but would require about an hour per 1,000 MCMC samples obtained from the same model if the sample size was $n = 1,000$. For large spatial datasets, a variety of dimension reduction and computationally efficient implementations can be used to model the spatial autocorrelation. The majority of efficient methods involve modeling the spatial autocorrelation using basis functions and a first-order model specification. To illustrate dimension reduction, we model the spatial autocorrelation using two different types of basis functions: the predictive process and thin plate regression splines. The predictive process is similar to kernel regression methods, except the basis expansion is slightly different and the basis coefficients are correlated in geographic space (Banerjee et al. 2008, 2014). The predictive process approach models the spatial process by

smoothing over a finite number of representative locations as follows

$$\begin{aligned}\mathbf{Z}(\phi) &\equiv \mathbf{C}(\phi)\mathbf{R}^*(\phi)^{-1} \\ \boldsymbol{\alpha} &\sim \text{N}(\mathbf{0}, \sigma_{\alpha}^2 \mathbf{R}^*(\phi))\end{aligned}\tag{21}$$

where $\mathbf{R}^*(\phi)$ is the $m \times m$ correlation matrix for preselected knots (Fig. 4c) and $\mathbf{C}(\phi)$ is the $n \times m$ correlation matrix between the observed data and knots. Using the predictive process method with $m = 50$ knots, the Bayesian model requires approximately three seconds per 1,000 MCMC samples obtained and the predicted probability of occurrence looks similar when compared to the second-order spatial model (cf. Fig. 4b to Fig. 4c; Appendix E). Furthermore, the predictive process method can be implemented using readily available software (Finley et al. 2015).

Generalized additive models (GAMs) are similar to models that use spatial random effects, but rely on techniques and basis functions commonly used in semiparametric regression (Ruppert et al. 2003). Specifying GAM typically require choosing a type of basis function, the number of basis vectors, and the location and number of knots. The most significant difference between the previous methods we have demonstrated and GAMs is the type of basis functions used. Many different basis functions are used to specify GAMs and introductions can be found in Hastie et al. (2009), James et al. (2013), Ruppert et al. (2003), and Wood (2006). Although GAMs are implemented under a Bayesian paradigm (Crainiceanu et al. 2005; Gelman et al. 2013; Ch. 20), penalized maximum likelihood methods are commonly used (Wood 2006). For illustrative purposes, we implement a GAM using thin plate regression splines to model the spatial autocorrelation. This implementation is available in standard software and may be particularly useful for very

large data sets (e.g., $n \approx 10^6$), requiring approximately two seconds to fit the model to our data using 50 basis coefficients (Wood et al. 2015). The predicted probability of occurrence is shown in Fig. 4d and is comparable to both specifications of the Bayesian spatial model (Fig. 4). We expected similarity between the GAM and the spatial model because there is a connection between first-order models that use spline basis functions and second-order spatial models (Nychka 2000).

DISCUSSION

Autocorrelation: the two cultures

“What is one person’s covariance structure is another persons’ mean structure (Cressie 1993; pg. 25).” Within subfields of statistics that focus on dependent data (e.g., spatial statistics), there is no general consensus on whether the influence of autocorrelation should be specified in the mean or covariance structure of a probability distribution. The notion that first-order specified models that use basis functions and second-order specified spatial and time-series models are both useful for dependent data has been a topic of discussion for several decades among statisticians (e.g., Cressie 1989 and comments by Wahba 1990; Laslett 1994 and comments by Handcock et al. 1994 and Mächler 1994). As we have demonstrated, the two approaches can result in the same model or an approximation thereof. With regard to which method to use, there are entire books written about correlation functions from a single perspective (e.g., Kriging in a spatial context; Stein 1999) and about certain classes of basis functions (Nason 2010; Table 2). Given the diversity of approaches, it is difficult to make specific recommendations. Our goal is to encourage researchers to consider both perspectives, rather than one or the other.

First-order or second-order?

Models that use second-order specifications can be converted to the equivalent first-order specification to assess collinearity among basis vectors and covariates of interest (e.g., Hodges and Reich 2010). Modeling the autocorrelation using a first-order specification can be beneficial when the autocorrelation does not follow a standard correlation function, such as the case with data collected from streams and rivers (Peterson and Ver Hoef 2010; Sampson 2010; Isaak et al. 2014) or moving animals (Buderman et al. 2016). Although we have not demonstrated this in our examples, the first-order specification might be more appealing when specifying theory-based ecological models (e.g., using partial differential equations), because the first-order model specification is naturally hierarchical (Wikle and Hooten 2010). Using the first-order specification, the conditional distribution of the data (or unobserved latent process) can be selected and knowledge of the process can be incorporated into the mean structure (e.g., Wikle 2003; Hooten and Wikle 2008; Wikle and Hooten 2010). Although scientifically meaningful, second-order model specifications may be more challenging to understand when compared to first-order models specifications (Hanks *under review*). Second-order specifications, however, can facilitate more computationally efficient algorithms. Thus, many contemporary models for autocorrelation are specified in terms of first-order structure and then converted to second-order structure for implementation (e.g., Finley et al. 2015).

Choosing basis functions

Choosing basis functions requires an understanding of both the underlying ecological process and the properties of the basis functions. For example, a property of the polynomial basis function is that it has a global support, thus an observation at one

location influences the fit of the model at another location, no matter how far apart the two locations are. This is why polynomial basis expansions often fails to model fine scale structure (cf., Fig. 1b to 2b). From an ecological perspective, the global support of polynomial basis functions implies that the underlying ecological process is connected across the entire space of interest. If the ecological process is thought to have discontinues, then basis functions that capture discontinuous structure and have compact support are a better choice (e.g., the uniform kernel used in Example 2; Table 2).

When selecting a basis function to model autocorrelation, standard model checking procedures are critical to ensure that model assumptions are met (e.g., checking for correlated residuals, multicollinearity, lack of fit, overfitting). Formal model selection may also be useful for selecting the optimal basis functions (Gelfand et al. 2012; Gelman et al. 2013 Ch. 20; Hooten and Hobbs 2015). Computational considerations may also be important when choosing a basis function. For example, orthogonal basis functions often result in more stable computational algorithms because the basis vectors are independent, obviating collinearity between basis vectors. We illustrated only a small fraction of the basis functions that could be used, thus we recommend that practitioners become familiar with the variety of options to ensure that the chosen basis function matches the goals of the study. To facilitate this, we have provided a brief summary of common basis functions, their properties, and useful references in Table 2.

Implementation

Typically, only a small number of covariates are included in a regression model, but one may want to include as many or more basis vectors as there are observations. For example, there are as many eigen basis vectors as there are unique locations in the dataset when the correlation matrix is specified using a correlation function. When many basis vectors are

used to model autocorrelation, the model can overfit the data. Adding constraints to high-dimensional estimation problems is a common technique to prevent overfitting. Such methods include regularization, penalized maximum likelihood estimation (e.g., ridge regression), treating the basis coefficients as random effects, or using a prior distribution that induces shrinkage (regularization) in a Bayesian model. There are important connections between methods that impose constraints to prevent overfitting that we have not presented here, but are important to understand when implementing models that use basis functions (Hooten and Hobbs 2015).

When fitting models data sets where dimension reduction is required, there is a trade-off between the reduction in dimension and the fit of the model. The fit of the model is influenced by dimension reduction because choosing the number of basis vectors to include in a model is an implicit form of regularization (Gelman et al. 2013, Ch. 20; Hooten and Hobbs 2015). Determining which basis functions are optimal for approximating correlation functions, how many basis vectors are needed, and the locations of knots are active areas of research (Gelfand et al., 2012). A general rule of thumb is to choose fewer basis vectors than the number of unique locations in the dataset, but as large as possible given the computational restrictions so that the predictions are accurate (e.g., on out-of-sample data). A detailed summary of dimension reduction approaches is beyond the scope of our work, but accessible introductions can be found in Paciorek (2007b), Wikle (2010), Cressie and Wikle (2011), and Banerjee et al. (2014).

Basis function model specifications have also become popular in spatio-temporal modeling, both in environmental (e.g., Wikle 2002) and ecological applications (e.g., Hooten and Wikle 2007). In practice, we find that understanding the properties of basis functions is critical to implementing computationally efficient Bayesian hierarchical models

that account for spatial, temporal, or spatio-temporal autocorrelation. In addition, using basis functions as part of a Bayesian hierarchical model makes many spatio-temporal models accessible to users of JAGS, NIMBLE, Stan, and WinBugs (Crainiceanu et al. 2005). Using the tools and techniques we have presented in this paper, basis function components can be added to existing hierarchical models to account for autocorrelation.

Inference and collinearity

For some applications, collinearity among covariates and basis vectors might occur and the development of remedial methods is a current topic of research in spatial statistics (Reich et al. 2006; Paciorek 2010; Hodges and Reich 2010; Hodges 2013; Hughes and Haran 2013; Hanks et al. 2015). The effects of collinearity among covariates and basis vectors have been noted in the ecological literature as well, particularly in a spatial context (e.g., Bini et al. 2009; Kühn 2007). In our experience, collinearity among covariates and basis vectors is a difficult challenge in applied problems. In some cases, the conventional wisdom that applies to collinearity among covariates can also be applied to basis vectors, but new intuition is needed when basis coefficients are treated as random effects (Hodges and Reich 2010; Paciorek 2010; Hodges 2013; Hanks et al. 2015; Murakami and Griffith 2015). As with collinearity among covariates in linear regression models, there is no clear remedy for extreme cases.

Conclusion

Ecologists face many choices when specifying models. One important choice is how to model autocorrelation. Autocorrelation, however, is not limited to specific domains and can occur in any space (e.g., covariate space, time, three-dimensional Euclidian space). Many methods used to model autocorrelation are general and can be understood as

generalized linear mixed models that employ basis expansions and treat basis coefficients as random effects. Using the basis function approach, we find that many of the commonly used ecological models can be modified to incorporate autocorrelation.

ACKNOWLEDGEMENTS

We thank Evan Cooch, Perry de Valpine, Devin Johnson, and three anonymous reviewers for valuable insight and early discussions about this work. The authors acknowledge support for this research from USGS G14AC00366. Any use of trade, firm, or product names is for descriptive purposes only and does not imply endorsement by the U.S. Government.

LITERATURE CITED

- Banerjee, S., Carlin, B. P., and , A. E. (2014). *Hierarchical Modeling and Analysis for Spatial Data*. CRC Press.
- Banerjee, S., Gelfand, A. E., Finley, A. O., and Sang, H. (2008). Gaussian predictive process models for large spatial data sets. *Journal of the Royal Statistical Society: Series B (Statistical Methodology)*, 70(4):825–848.
- Bini, M., Diniz-Filho, J. A. F., Rangel, T. F., Akre, T. S., Albaladejo, R. G., Albuquerque, F. S., Aparicio, A., Araujo, M. B., Baselga, A., Beck, J., et al. (2009). Coefficient shifts in geographical ecology: an empirical evaluation of spatial and non-spatial regression. *Ecography*, 32(2):193–204.
- Bishop, C. (2006). *Pattern Recognition and Machine Learning*. Springer.
- Bolker, B. M., Brooks, M. E., Clark, C. J., Geange, S. W., Poulsen, J. R., Stevens, M. H. H., and White, J.-S. S. (2009). Generalized linear mixed models: a practical guide for

- ecology and evolution. *Trends in Ecology & Evolution*, 24(3):127–135.
- Borcard, D., Legendre, P., and Drapeau, P. (1992). Partialling out the spatial component of ecological variation. *Ecology*, 73(3):1045–1055.
- Buderman, F. E., Hooten, M. B., Ivan, J. S., and Shenk, T. M. (2016). A functional model for characterizing long-distance movement behaviour. *Methods in Ecology and Evolution*, 7(3):264–273.
- Conn, P. B., Johnson, D. S., Hoef, J. M. V., Hooten, M. B., London, J. M., and Boveng, P. L. (2015). Using spatiotemporal statistical models to estimate animal abundance and infer ecological dynamics from survey counts. *Ecological Monographs*, 85(2):235–252.
- Crainiceanu, C., Ruppert, D., and Wand, M. P. (2005). Bayesian analysis for penalized spline regression using WinBUGS. *Journal of Statistical Software*, 14.
- Cressie, N. (1989). Geostatistics. *The American Statistician*, 43(4):197–202.
- Cressie, N. (1993). *Statistics for Spatial Data*. John Wiley & Sons.
- Cressie, N. and Wikle, C. (2011). *Statistics for Spatio-Temporal Data*. Wiley.
- Dormann, C. F., Elith, J., Bacher, S., Buchmann, C., Carl, G., Carré, G., Marquéz, J. R. G., Gruber, B., Lafourcade, B., Leitão, P. J., et al. (2013). Collinearity: a review of methods to deal with it and a simulation study evaluating their performance. *Ecography*, 36(1):27–46.
- Elith, J. and Leathwick, J. R. (2009). Species distribution models: ecological explanation and prediction across space and time. *Annual Review of Ecology, Evolution, and Systematics*, 40(1):677.
- Fieberg, J. and Ditmer, M. (2012). Understanding the causes and consequences of animal movement: a cautionary note on fitting and interpreting regression models with time-dependent covariates. *Methods in Ecology and Evolution*, 3(6):983–991.

- Finley, A. O., Banerjee, S., and Gelfand, A. E. (2015). spBayes for large univariate and multivariate point-referenced spatio-temporal data models. *Journal of Statistical Software*, 63:1–28.
- Gelfand, A. E., Banerjee, S., and Finley, A. O. (2012). Spatial design for knot selection in knot-based dimension reduction models. In *Spatio-temporal Design: Advances in Efficient Data Acquisition*, pages 142–169. Wiley.
- Gelman, A., Carlin, J., Stern, H., Dunson, D., Vehtari, A., and Rubin, D. (2013). *Bayesian Data Analysis*. Chapman & Hall/CRC. Taylor & Francis.
- Gillibrand, E., Jamieson, A., Bagley, P., Zuur, A., and Priede, I. (2007). Seasonal development of a deep pelagic bioluminescent layer in the temperate NE atlantic ocean. *Marine Ecology Progress Series*, 341:37–44.
- Griffith, D. A. and Peres-Neto, P. R. (2006). Spatial modeling in ecology: the flexibility of eigenfunction spatial analyses. *Ecology*, 87(10):2603–2613.
- Handcock, M. S., Meier, K., and Nychka, D. (1994). Comment on Kriging and splines: an empirical comparison of their predictive performance in some applications. *Journal of the American Statistical Association*, 89(426):401–403.
- Hanks, E. M. (under review). A constructive spatio-temporal approach to modeling spatial covariance. *Journal of the American Statistical Association*, <http://arxiv.org/abs/1506.03824>.
- Hanks, E. M., Schliep, E. M., Hooten, M. B., and Hoeting, J. A. (2015). Restricted spatial regression in practice: geostatistical models, confounding, and robustness under model misspecification. *Environmetrics*, 26(4):243–254.
- Hastie, T., Tibshirani, R., and Friedman, J. (2009). *The Elements of Statistical Learning: Data Mining, Inference, and Prediction, Second Edition*. Springer Series in Statistics.

Springer.

- Hefley, T. J., Tyre, A. J., and Blankenship, E. E. (2013). Statistical indicators and state-space population models predict extinction in a population of bobwhite quail. *Theoretical Ecology*, 6(3):319–331.
- Higdon, D. (2002). Space and space-time modeling using process convolutions. *Quantitative Methods for Current Environmental Issues*, 3754.
- Hobbs, N. T. and Hooten, M. B. (2015). *Bayesian Models: A Statistical Primer for Ecologists*. Princeton University Press.
- Hodges, J. S. (2013). *Richly Parameterized Linear Models: Additive, Time Series, and Spatial Models Using Random Effects*. CRC Press.
- Hodges, J. S. and Reich, B. J. (2010). Adding spatially-correlated errors can mess up the fixed effect you love. *The American Statistician*, 64(4):325–334.
- Hoeting, J. A. (2009). The importance of accounting for spatial and temporal correlation in analyses of ecological data. *Ecological Applications*, 19(3):574–577.
- Hooten, M. B. and Hobbs, N. T. (2015). A guide to Bayesian model selection for ecologists. *Ecological Monographs*, 85(1):3–28.
- Hooten, M. B., Larsen, D. R., and Wikle, C. K. (2003). Predicting the spatial distribution of ground flora on large domains using a hierarchical Bayesian model. *Landscape Ecology*, 18(5):487–502.
- Hooten, M. B. and Wikle, C. K. (2007). Shifts in the spatio-temporal growth dynamics of shortleaf pine. *Environmental and Ecological Statistics*, 14(3):207–227.
- Hooten, M. B. and Wikle, C. K. (2008). A hierarchical Bayesian non-linear spatio-temporal model for the spread of invasive species with application to the Eurasian collared-dove. *Environmental and Ecological Statistics*, 15(1):59–70.

- Hughes, J. and Haran, M. (2013). Dimension reduction and alleviation of confounding for spatial generalized linear mixed models. *Journal of the Royal Statistical Society: Series B (Statistical Methodology)*, 75(1):139–159.
- Isaak, D. J., Peterson, E. E., Ver Hoef, J. M., Wenger, S. J., Falke, J. A., Torgersen, C. E., Sowder, C., Steel, E. A., Fortin, M., Jordan, C. E., et al. (2014). Applications of spatial statistical network models to stream data. *Wiley Interdisciplinary Reviews: Water*, 1(3):277–294.
- James, G., Witten, D., Hastie, T., and Tibshirani, R. (2013). *An Introduction to Statistical Learning*. Springer.
- Johnson, D. S., Ream, R. R., Towell, R. G., Williams, M. T., and Guerrero, J. D. L. (2013). Bayesian clustering of animal abundance trends for inference and dimension reduction. *Journal of Agricultural, Biological, and Environmental Statistics*, 18(3):299–313.
- Krainski, E., Lindgren, F., Simpson, D., and Rue, H. (2016). The r-inla tutorial on spde models. *Journal of Geographical Systems*, <http://www.math.ntnu.no/inla/r-inla.org/tutorials/spde/spde-tutorial.pdf>.
- Kühn, I. (2007). Incorporating spatial autocorrelation may invert observed patterns. *Diversity and Distributions*, 13(1):66–69.
- Laslett, G. M. (1994). Kriging and splines: an empirical comparison of their predictive performance in some applications. *Journal of the American Statistical Association*, 89(426):391–400.
- Lee, Y., Nelder, J. A., and Pawitan, Y. (2006). *Generalized Linear Models with Random Effects: Unified Analysis via H-likelihood*. CRC Press.
- Legendre, P. (1993). Spatial autocorrelation: trouble or new paradigm? *Ecology*, 74(6):1659–1673.

- Legendre, P. and Fortin, M. J. (1989). Spatial pattern and ecological analysis. *Vegetatio*, 80(2):107–138.
- Levin, S. A. (1992). The problem of pattern and scale in ecology. *Ecology*, 73(6):1943–1967.
- Lindgren, F., Rue, H., and Lindström, J. (2011). An explicit link between Gaussian fields and Gaussian Markov random fields: the stochastic partial differential equation approach. *Journal of the Royal Statistical Society: Series B (Statistical Methodology)*, 73(4):423–498.
- Littell, R. C., Stroup, W. W., Milliken, G. A., Wolfinger, R. D., and Schabenberger, O. (2006). *SAS for Mixed Models*. SAS institute.
- Lorenz, E. N. (1956). Empirical orthogonal functions and statistical weather prediction. <http://www.o3d.org/abracco/Atlantic/Lorenz1956.pdf>.
- Mächler, M. (1994). Comment on Kriging and splines: an empirical comparison of their predictive performance in some applications. *Journal of the American Statistical Association*, 89(426):403–405.
- Murakami, D. and Griffith, D. A. (2015). Random effects specifications in eigenvector spatial filtering: a simulation study. *Journal of Geographical Systems*, 17(4):311–331.
- Nason, G. (2010). *Wavelet Methods in Statistics with R*. Springer Science & Business Media.
- Nychka, D. W. (2000). Spatial-process estimates as smoothers. In *Smoothing and Regression: Approaches, Computation, and Applications*, pages 393–424. Wiley.
- Paciorek, C. (2010). The importance of scale for spatial-confounding bias and precision of spatial regression estimators. *Statistical Science*, 25(1):107–125.
- Paciorek, C. J. (2007a). Bayesian smoothing with Gaussian processes using Fourier basis functions in the spectralgp package. *Journal of Statistical Software*, 19(2).

- Paciorek, C. J. (2007b). Computational techniques for spatial logistic regression with large data sets. *Computational Statistics and Data Analysis*, 8(51):3631–3653.
- Peterson, E. E. and Ver Hoef, J. (2010). A mixed-model moving-average approach to geostatistical modeling in stream networks. *Ecology*, 91(3):644–651.
- Reich, B. J., Hodges, J. S., and Zadnik, V. (2006). Effects of residual smoothing on the posterior of the fixed effects in disease-mapping models. *Biometrics*, 62(4):1197–1206.
- Robinson, G. K. (1991). That BLUP is a good thing: the estimation of random effects. *Statistical Science*, 6(1):15–32.
- Ruppert, D., Wand, P., and Carroll, R. (2003). *Semiparametric Regression*. Cambridge Series in Statistical and Probabilistic Mathematics. Cambridge University Press.
- Sampson, P. D. (2010). Constructions for nonstationary spatial processes. In *Handbook of Spatial Statistics*, pages 119–130. CRC Press.
- Stein, M. L. (1999). *Interpolation of Spatial Data: Some Theory for Kriging*. Springer Science & Business Media.
- Stroup, W. W. (2012). *Generalized Linear Mixed Models: Modern Concepts, Methods and Applications*. CRC press.
- Veech, J. A. (2006). Increasing and declining populations of northern bobwhites inhabit different types of landscapes. *Journal of Wildlife Management*, 70(4):922–930.
- Wahba, G. (1990). Comment on Cressie. *The American Statistician*, 44(3):255–258.
- Wikle, C. K. (2002). A kernel-based spectral model for non-Gaussian spatio-temporal processes. *Statistical Modelling: An International Journal*, 2(4):299–314.
- Wikle, C. K. (2003). Hierarchical Bayesian models for predicting the spread of ecological processes. *Ecology*, 84(6):1382–1394.
- Wikle, C. K. (2010). Low rank representations for spatial processes. In *Handbook of Spatial*

Statistics, pages 107–118. CRC Press.

Wikle, C. K. and Hooten, M. B. (2010). A general science-based framework for dynamical spatio-temporal models. *Test*, 19(3):417–451.

Wood, S. (2006). *Generalized Additive Models: An Introduction with R*. Chapman & Hall/CRC Texts in Statistical Science. Taylor & Francis.

Wood, S. N., Goude, Y., and Shaw, S. (2015). Generalized additive models for large data sets. *Journal of the Royal Statistical Society: Series C (Applied Statistics)*, 64(1):139–155.

Zuur, A., Ieno, E. N., Walker, N., Saveliev, A. A., and Smith, G. M. (2009). *Mixed Effects Models and Extensions in Ecology with R*. Springer Science & Business Media.

SUPPLEMENTAL MATERIAL

Appendix A

Additional comments about basis functions.

Appendix B

Interactive illustration of Fig. 1 & Fig. 2

https://hootenlab.shinyapps.io/hooten_lab_manuscript_figures.

Appendix C

Tutorial with data and R code for the bioluminescence example.

Appendix D

Tutorial with data and R code for the time series example.

Appendix E

Tutorial with data and R code for the spatial example.

Table 1. Glossary of terms and definitions.

Term	Definition
Autocorrelation	Correlation between observations based on some measure of distance or time that exists after the influence of all covariates is accounted for
Basis expansion	A collection of basis vectors from a single covariate
Basis vector	Any transformation of a covariate
Basis function	Any mathematical function that transforms a covariate
Compact support	A support that does not includes all possible locations or time points
Correlation function	A function that describes the autocorrelation between observations
Correlation matrix	A positive semi-definite matrix whose elements are the correlation between observations
Covariate	Any quantity that can be measured and is associated with an observation (e.g., the time or spatial location of the observation)
Dependence	Correlation between observations defined in a general space (spatial or temporal dependence is equivalent to autocorrelation)
First-order specification	When a function that models the dependence is specified in the mean (expected value) of a probability distribution
Global support	A support that includes all possible locations or time points
Second-order specification	When a function that models dependence is specified in the covariance of a probability distribution
Support	The set of locations or time points where the basis function results in non-zero values

Table 2. Common types of basis functions, important properties and references.

Basis function	Orthogonal	Support	Notable use	Reference
Eigen	yes	global	Dimension reduction and detecting multicollinearity between basis vectors and covariates in second-order models	Hodges and Reich (2010), Hodges (2013; Ch. 10), Cressie and Wikle (2011; Ch. 5)
Fourier	yes	global	Large data sets with a smooth effect of autocorrelation	Paciorek (2007a), Cressie and Wikle (2011; Ch. 3)
Kernel	no	global or compact	Large data sets and a directional effect of autocorrelation	Higdon (2002), Sampson (2010), Peterson and Ver Hoef (2010)
Piecewise linear	no	local	Implementing numerical solutions to stochastic partial differential equations	Lindgren et al. (2011), Krainski et al. (2016)
Polynomial	no	global	Modeling simple nonlinear effects of autocorrelation	Carroll et al. (2003; Ch. 2), James et al. (2013; Ch. 7)
Splines	no	global or compact	Large data sets with smooth effects of autocorrelation	Carroll et al. (2003; Ch. 3), Wood et al. (2006), Hastie et al. (2009; Ch. 5), James et al. (2013; Ch. 7)
Wavelets	yes	global and compact	Modeling discontinuous effects of autocorrelation	Nason (2008)

Figure 1. Scatterplots showing the density of pelagic bioluminescence (sources) versus water depth (points). The top panels show fitted regression models (black lines). The corresponding basis vectors multiplied by the estimated coefficients (colored curves) are shown in the bottom panels. Simple linear regression model (a; Eq. 1) with corresponding constant (red) and linear basis vectors (blue). Polynomial regression model (b; Eq. 2) with corresponding constant (red), linear (blue), and quadratic basis vectors (green). Polynomial regression model (c) similar to that shown in panel (b), except with basis vectors calculated relative to three water depths (k_j in Eq. 3; vertical colored lines). Note that the basis vectors are multiplied by the estimated coefficients and summed to produce the fitted curves (black lines). See Appendix B for an interactive version of this figure.

Figure 2. Scatterplots showing the density of pelagic bioluminescence (sources) versus water depth (points). The top panels shows the fitted curve (black lines) obtained from a second-order specification that uses a Gaussian correlation function (a), the equivalent first-order specification that uses eigen basis vectors (b), and a first-order specification that uses a Gaussian kernel basis function (c). The bottom panels show the intercept term (red), eigen basis vectors (b), and Gaussian kernel basis vectors (c). For illustrative purposes, only the product of three basis vectors and coefficients are shown (with knots located at the vertical lines in panel (c)). There are 51 eigen basis vectors and coefficients that sum to produce the fitted curve (black line) in panel (b), and 17 kernel basis vectors that sum to produce the fitted curve (black line) in panel (c). See Appendix B for an interactive version of this figure.

Figure 3. Scatterplots of bobwhite quail population indices over time (points). The top panels show fitted regression models (black lines) obtained from a second-order specification that uses an AR(1) correlation matrix (a), the equivalent first-order specification that uses

eigen basis vectors (b), and a first-order specification that uses a compactly supported uniform kernel basis function (c). The bottom panels show the fixed effects term (red), three eigen basis vectors (b), and three compactly supported kernel basis vectors with knots located at the vertical lines (c). All basis vectors are multiplied by basis coefficients.

Figure 4. Prediction domain from the Missouri Ozark Forest Ecosystem Project presented in Hooten et al. (2003). Red and black points (a) represent the 10×10 m plot locations that were sampled ($n = 216$) and whether pointed-leaved tick trefoil was present (red) or absent (black). The heat maps (b,c,d) show the predicted probability of occurrence in 32,768 plots from a binary spatial regression model (b; Eq. 19), a reduced dimension binary spatial regression model using predictive process basis functions (Eq. 21) with knots located within the prediction domain represented by + (c), and a generalized additive model that uses thin plate regression splines (d).




Figure 1:




Figure 2:




Figure 3:




Figure 4: