

Supports of Implicit Dependence Copulas

Songkiat Sumetkijakan

Department of Mathematics and Computer Science
Faculty of Science, Chulalongkorn University
Phayathai Road, Bangkok, Thailand

June 24, 2016

Abstract

A copula of continuous random variables X and Y is called an *implicit dependence copula* if there exist functions α and β such that $\alpha(X) = \beta(Y)$ almost surely, which is equivalent to C being factorizable as the $*$ -product of a left invertible copula and a right invertible copula. Every implicit dependence copula is supported on the graph of $f(x) = g(y)$ for some measure-preserving functions f and g but the converse is not true in general.

We obtain a characterization of copulas with implicit dependence supports in terms of the non-atomicity of two newly defined associated σ -algebras. As an application, we give a broad sufficient condition under which a self-similar copula has an implicit dependence support. Under certain extra conditions, we explicitly compute the left invertible and right invertible factors of the self-similar copula.

Keywords: Markov operators; non-atomic; bivariate copulas; implicit dependence; self-similar copulas; fractal support

MSC2010: 28A20; 28A35; 46B20; 60A10; 60B10

1 Introduction

It is well-known that the bivariate copula of two random variables completely captures their dependence structure. Notable examples are the *independence copula* $\Pi(u, v) = uv$, which corresponds to independent random variables, and the copulas of completely dependent random variables, called *complete dependence copulas*. Since it was discovered [10, 9, 8] that there are complete dependence copulas arbitrarily closed to Π in the uniform norm, many norms have been introduced and investigated in the literature [3] giving rise to measures of dependence such as ω in [17] and ζ_1 in [18]. These dependence measures defined in terms of the copula's first partial derivatives attain the maximum value 1 at least for complete dependence copulas and the minimum value 0 when and only when the copula is Π . However, with respect to the measure of mutual complete

dependence (MCD) ω , the independence copula can still be approximated by *implicit dependence copulas* [1], defined as copulas of random variables X and Y which are implicitly dependent in the sense that $f(X) = g(Y)$ a.s. for some Borel measurable functions f and g . For Rényi-type measures of dependence [13] such as ω_* in [15] and ν_* in [7], with respect to which all complete dependence copulas have measure 1, it can be proved that all implicit dependence copulas also have maximum measure 1. It is then evident that implicit dependence copulas plays a crucial role in understanding as well as comparing and contrasting measures of MCD and Rényi-type dependence measures. Closely related and constituting a much larger class than the implicit dependence copulas are the copulas whose supports are that of an implicit dependence copula.

We shall investigate copulas with implicit dependence supports via their corresponding Markov operators and associated pairs of σ -algebras. Our approach is motivated by the characterization of idempotent copulas via their σ -algebras of invariant sets in [4]. In particular, they proved that non-atomic idempotent copulas must be of the form $L * L^t$ for some left invertible copula L (where C^t denotes the transpose of C , i.e. $C^t(x, y) = C(y, x)$, and $*$ is the product of copulas first defined and studied in [2, 12]). We define σ_C and σ_C^* associated with a copula C for which the corresponding Markov operator T_C maps indicator functions of sets in σ_C to indicator functions of sets in σ_C^* . We prove that copulas with implicit dependence supports are exactly copulas whose both σ -algebras are non-atomic. Our main result finds an application in copulas with fractal supports introduced by Fredricks, Nelsen and Rodríguez-Lallena [6]. Given a transformation matrix A , there is a unique *self-similar copula* C_A such that $[A](C_A) = C_A$, where $[A]$ maps the class of bivariate copulas into itself according to the weights given by the entries in A . As a consequence, we obtain a broad sufficient condition on a transformation matrix A under which the self-similar copula C_A is non-atomic and hence has an implicit dependence support. Working directly with the transformation matrix A , a sufficient condition under which C_A is an implicit dependence copula is also given. It would be even more interesting if we had a characterization of implicit dependence copulas via behaviors of their σ -algebras. We are hopeful that our future attempts will not be futile as such a characterization would be beneficial in studies involving products of implicit dependence copulas.

The manuscript is organized as follows. Section 2 lays the necessary background on copulas and Markov operators for the rest of the paper. We then define the associated σ -algebras of a copula and prove their basic properties in Section 3. Section 4 gives a definition of non-atomic copulas and some of their fundamental properties summarizing in a characterization of non-atomic copulas. In the final section, the characterization is used in an investigation of copulas with fractal support. We also give a sufficient condition on a transformation matrix under which the induced invariant copula can be written as the product of a left invertible copula and a right invertible copula.

2 Background on copulas and Markov operators

Let λ denote the Lebesgue measure on \mathbb{R} , $I \equiv [0, 1]$ and $\mathcal{B} \equiv \mathcal{B}(I)$ the Borel σ -algebra on I . Since we always consider λ -integrable functions on I that are measurable with respect to various sub- σ -algebras \mathcal{M} of \mathcal{B} , we will denote by $L^1(\mathcal{M})$ the class of λ -integrable \mathcal{M} -measurable functions on I . \mathcal{B} -measurable functions are called *Borel functions*. For $A \in \mathcal{B}$, $\mathbb{1}_A$ denotes the indicator function of A and $\mathbb{1} \equiv \mathbb{1}_I$.

A (bivariate) *copula* C is a function from I^2 to I which is the joint distribution function of two random variables uniformly distributed on $[0, 1]$. For random variables X, Y with joint distribution $F_{X,Y}$ and continuous marginal distributions F_X and F_Y , there exists, by the Sklar's theorem, a unique copula C , called the *copula of X and Y* , for which $F_{X,Y}(x, y) = C(F_X(x), F_Y(y))$ for all x, y . The *independence copula* is the product copula $\Pi(u, v) = uv$. *Complete dependence copulas* are either the copulas $C_{ef} \equiv C_{e,f}$ or $C_{fe} \equiv C_{f,e}$ where $e(x) = x$ and f is a measure-preserving function on I in the sense that $\lambda(f^{-1}(B)) = \lambda(B)$ for all $B \in \mathcal{B}$. Here, $C_{f,g}(u, v) = \lambda(f^{-1}([0, u]) \cap g^{-1}([0, v]))$ for $u, v \in I$. The *comonotonic* and *countermonotonic* copulas are $M = C_{e,e}$ and $W = C_{e,1-e}$, respectively. Each copula C induces a *doubly stochastic measure* μ_C by $\mu_C((a, b] \times (c, d]) = C(b, d) - C(b, c) - C(a, d) + C(a, c)$. The *support* of C is then defined as the support of the induced measure μ_C , i.e. the complement of the union of all open sets having zero μ_C -measure. One can construct a new copula by taking any convex combinations of two copulas. Any two copulas C, D also give rise to a new copula via the **-product*: $(C * D)(u, v) = \int_0^1 \partial_2 C(u, t) \partial_1 D(t, v) dt$. The binary operation $*$ makes the class of copulas a monoid with null element Π and identity M . If $C * D = M$ then C is a left inverse of D and D is a right inverse of C . The left invertible copulas are exactly the complete dependence copulas C_{ef} , while the right invertible copulas are exactly the complete dependence copulas C_{fe} . See [11, 5] for comprehensive introductions to many aspects of copulas.

A linear operator T on $L^1(\mathcal{B})$ is called a *Markov operator* if

- i. $T\mathbb{1} = \mathbb{1}$,
- ii. $\int_0^1 T\psi d\lambda = \int_0^1 \psi d\lambda$ for all $\psi \in L^1$, which is equivalent to $T^*\mathbb{1} = \mathbb{1}$, and
- iii. $T\psi \geq 0$ for all $\psi \geq 0$, which means T is positive.

So a Markov operator must be a bounded linear operator on L^1 (and L^∞). From [12], for each copula C , there corresponds a Markov operator T_C defined by

$$(T_C\psi)(x) = \frac{d}{dx} \int_0^1 \partial_2 C(x, t) \psi(t) dt.$$

In fact, the mapping $\Phi: C \mapsto T_C$ is an isomorphism from the set of copulas endowed with the $*$ -product onto the set of Markov operators under the composition. In particular, $T_{C*D} = T_C \circ T_D$. Define also $(T_C^*\varphi)(y) = \frac{d}{dy} \int_0^1 \partial_1 C(t, y) \varphi(t) dt$.

The copula C and the Markov operators T_C and T_C^* are also related by the identities

$$C(x, y) = \int_0^x T_C \mathbf{1}_{[0, y]}(t) dt \quad \text{and} \quad C(x, y) = \int_0^y T_C^* \mathbf{1}_{[0, x]}(t) dt. \quad (1)$$

In fact, it is a good exercise in functional analysis to verify that the Markov operator T_C^* coincides with the extension of the adjoint of T_C to a Markov operator on L^1 , i.e. $T_C^* = (T_C)^*$.

Let us quote a very useful result from [4, Theorem 2.11] where, for brevity, we denote $T_{C_{f,g}} = T_{f,g}$.

Theorem 2.1. *Let f be a measure-preserving Borel function and $\psi \in L^1(\mathcal{B})$. Then $[T_{ef}\psi](x) = \psi \circ f(x)$ and $[T_{fe}(\psi \circ f)](x) = \psi(x)$ for almost all $x \in [0, 1]$.*

3 Associated σ -algebras

Unless stated otherwise, all equalities of two functions hold λ -almost everywhere and all equalities of two sets mean that their symmetric difference has Lebesgue measure zero. The integral on the whole unit interval I is denoted simply by \int .

Let C be the copula of random variables X and Y which are uniformly distributed on $[0, 1]$. Then for any Borel sets $R, S \subseteq [0, 1]$, we have

$$T_C \mathbf{1}_S(x) = P(Y \in S | X = x) \quad \text{and} \quad T_C^* \mathbf{1}_R(y) = P(X \in R | Y = y).$$

They are called *transition probabilities*. See [2] We also have $T_C^* = T_{C^t}$ and $(T_C^*)^* = T_C$. Roughly speaking, if $T_C \mathbf{1}_S = \mathbf{1}_R$, then it happens with probability one that if $X \in R$ then $Y \in S$. For each copula C or Markov operator $T = T_C$, let us define

$$\begin{aligned} \sigma_C &= \sigma_T \equiv \{S \in \mathcal{B} : T \mathbf{1}_S = \mathbf{1}_R \text{ for some } R\} \quad \text{and} \\ \sigma_C^* &= \sigma_T^* \equiv \{R \in \mathcal{B} : T \mathbf{1}_S = \mathbf{1}_R \text{ for some } S\}. \end{aligned}$$

Example 1. Let us explicitly compute T_C for $C = \Pi$, $C =$ some complete dependence copulas and $C = \frac{M+W}{2}$.

1. $T_\Pi \mathbf{1}_S = \mathbf{1}_R$ is equivalent to $\mathbf{1}_R(x) = \frac{d}{dx} \int_S \partial_2 \Pi(x, t) dt = \lambda(S)$ for a.e. $x \in [0, 1]$. So $\lambda(S) = 0$ or 1 and hence $\lambda(R) = \lambda(S) = 0$ or 1 . Thus, $\sigma_\Pi = \sigma_\Pi^* = \{S \in \mathcal{B} : \lambda(S) = 0 \text{ or } 1\}$.
2. With essentially the same arguments as that in [4], σ_M , σ_M^* , σ_W and σ_W^* are the Borel sets.
3. For $C = \frac{M+W}{2}$, $T_C \mathbf{1}_S(x) = \frac{1}{2} \mathbf{1}_S(x) + \frac{1}{2} \mathbf{1}_S(1-x)$ a.e. x . If S is symmetric with respect to $\frac{1}{2}$, i.e. $x \in S$ if and only if $1-x \in S$, then $T_C \mathbf{1}_S = \mathbf{1}_S$. Conversely, $T_C \mathbf{1}_S(x) = \mathbf{1}_R(x)$ implies that $x \in S$ if and only if $1-x \in S$. Moreover, for such a symmetric set S , $(\mathbf{1}_R - \mathbf{1}_S)(x) = (T_C \mathbf{1}_S - \mathbf{1}_S)(x) = \frac{1}{2}(\mathbf{1}_S(1-x) - \mathbf{1}_S(x)) = 0$ a.e. x . That is, $R = S$. Hence, $\sigma_{\frac{M+W}{2}} = \sigma_{\frac{M+W}{2}}^* = \{S \subseteq [0, 1] : x \in S \Leftrightarrow 1-x \in S\}$.

4. For $0 < \alpha < 1$, let L_α denote the complete dependence copula whose support consists of the line segments $y = \frac{x}{\alpha}$, $0 \leq x \leq \alpha$, and $y = \frac{x-\alpha}{1-\alpha}$, $\alpha < x \leq 1$. Then a direct computation yields $T_{L_\alpha} \mathbf{1}_S = \mathbf{1}_{(\alpha S) \cup (\alpha + (1-\alpha)S)}$ and hence $\sigma_{L_\alpha} = \mathcal{B}$ and $\sigma_{L_\alpha}^* = \{\alpha S \cup (\alpha + (1-\alpha)S) : S \in \mathcal{B}\}$.

Listed below are basic properties of sets S and R linked by T .

Proposition 3.1. *Let T be a Markov operator and $S_1, S_2, R_1, R_2 \in \mathcal{B}$. Then*

1. *if $T\mathbf{1}_{S_1} = \mathbf{1}_{R_1}$ then $T^*\mathbf{1}_{R_1} = \mathbf{1}_{S_1}$;*
2. *if $T\mathbf{1}_{S_1} = \mathbf{1}_{R_1}$ then $T\mathbf{1}_{S_1^c} = \mathbf{1}_{R_1^c}$;*
3. *if $T\mathbf{1}_{S_i} = \mathbf{1}_{R_i}$ for $i = 1, 2$, then $T\mathbf{1}_{S_1 \cap S_2} = \mathbf{1}_{R_1 \cap R_2}$;*
4. *the classes σ_T and σ_T^* are σ -algebras; and*
5. *$\sigma_T^* = \sigma_{T^*}$, that is $\sigma_C^* = \sigma_{C^*}$.*

Proof. 1: By the definition of $T^*\mathbf{1}_R$ applied to $\mathbf{1}_S$ with the canonical identification between $(L^1)^*$ and L^∞ , $\int_S T^*\mathbf{1}_R d\lambda = \int \mathbf{1}_S T^*\mathbf{1}_R d\lambda = \int \mathbf{1}_S \mathbf{1}_R d\lambda = \lambda(R) = \int_S \mathbf{1}_S d\lambda$. Since $0 \leq T^*\mathbf{1}_R \leq 1$ (a.e.), $T^*\mathbf{1}_R = \mathbf{1}_S$ as desired. 2: If $T\mathbf{1}_S = \mathbf{1}_R$ then it follows from $T\mathbf{1}_S + T\mathbf{1}_{S^c} = T(\mathbf{1}_S + \mathbf{1}_{S^c}) = T\mathbf{1} = \mathbf{1} = \mathbf{1}_R + \mathbf{1}_{R^c}$ that $T\mathbf{1}_{S^c} = \mathbf{1}_{R^c}$. 3: Suppose $T\mathbf{1}_{S_i} = \mathbf{1}_{R_i}$ for $i = 1, 2$ and let $S = S_1 \cap S_2$, $R = R_1 \cap R_2$. Since $S \subseteq S_i$, $\mathbf{1}_{S_i} - \mathbf{1}_S \geq 0$ and so $\mathbf{1}_{R_i} - T\mathbf{1}_S = T(\mathbf{1}_{S_i} - \mathbf{1}_S) \geq 0$. Therefore $T\mathbf{1}_S \leq \min(\mathbf{1}_{R_1}, \mathbf{1}_{R_2}) = \mathbf{1}_{R_1 \cap R_2} = \mathbf{1}_R$. But $\int (\mathbf{1}_R - T\mathbf{1}_S) d\lambda = \int \mathbf{1}_R d\lambda - \int \mathbf{1}_S d\lambda = \lambda(R) - \lambda(S) = 0$, where the last equality follows from considering $T^*\mathbf{1}_{R_2}(\mathbf{1}_{S_1}) = \mathbf{1}_{R_2}(T\mathbf{1}_{S_1})$. Hence, $T\mathbf{1}_S = \mathbf{1}_R$. 4: Suppose $T\mathbf{1}_{S_i} = \mathbf{1}_{R_i}$ for all $i \in \mathbb{N}$. By 3., if S_1, S_2, \dots are mutually disjoint then so is the sequence R_1, R_2, \dots because $\sum_{i=1}^\infty \mathbf{1}_{R_i} = \sum_{i=1}^\infty T\mathbf{1}_{S_i} = T(\sum_{i=1}^\infty \mathbf{1}_{S_i}) = T(\mathbf{1}_{\cup_i S_i}) \leq \mathbf{1}$. Thus, $T(\mathbf{1}_{\cup_i S_i}) = \mathbf{1}_{\cup_i R_i}$. Generally, we write $\cup_i S_i$ as the disjoint union $\cup_i \tilde{S}_i$, where $\tilde{S}_i = S_i \setminus \cup_{j < i} \tilde{S}_j$. Letting $T\mathbf{1}_{\tilde{S}_i} = \mathbf{1}_{\tilde{R}_i}$, it follows from 2., 3. and the disjoint union case above that $T\mathbf{1}_{S_i \setminus \cup_{j < i} \tilde{S}_j} = \mathbf{1}_{R_i \setminus \cup_{j < i} \tilde{R}_j}$. Hence, $\tilde{R}_i = R_i \setminus \cup_{j < i} \tilde{R}_j$ by induction. Consequently, $T(\mathbf{1}_{\cup_i S_i}) = \mathbf{1}_{\cup_i R_i}$. 5: This clearly follows from 1. and the fact that $T^{**} = T$. \square

Remark. By Theorem 2.1, $T_L^* \circ T_L = I$, the identity map, on $L^1(\mathcal{B})$ if the copula L is left invertible. However, it follows from the above Proposition that for all copula C , $T_C^* \circ T_C = I$ on $\{\mathbf{1}_S : S \in \sigma(C)\}$, or equivalently on the class of $\sigma(C)$ -measurable functions.

Theorem 3.2. *Let T be a Markov operator with associated σ -algebras σ_T and σ_T^* . If ψ is σ_T -measurable then $T\psi$ is σ_T^* -measurable.*

Proof. By the linearity of T and the definition of σ_T and σ_T^* , if ψ is a simple σ_T -measurable function then $T\psi$ is simple and σ_T^* -measurable. The case when ψ is σ_T -measurable follows from the continuity of T . \square

4 Non-atomic copulas

Definition 1. Let \mathcal{S} be a sub- σ -algebra of \mathcal{B} . A set S in \mathcal{S} is called an *atom* in \mathcal{S} if $\lambda(S) > 0$ and for all $E \in \mathcal{S}$, either $\lambda(S \cap E) = \lambda(S)$ or $\lambda(S \cap E) = 0$. The σ -algebra \mathcal{S} is said to be *non-atomic* if there are no atoms in \mathcal{S} ; otherwise, it is called *atomic*. \mathcal{S} is *totally atomic* if there is a (countable) collection of essentially disjoint atoms E_1, E_2, \dots in \mathcal{S} such that $\sum_i \lambda(E_i) = 1$. We say that a bivariate copula C is *non-atomic* if both σ_C and σ_C^* are non-atomic. And C is called *totally atomic* if both σ_C and σ_C^* are totally atomic.

Note that the non-atomicity is a generalization of the notion of the same name in [4], which is defined only for idempotent copulas C via their invariant sets defined as Borel sets S for which $T_C \mathbf{1}_S = \mathbf{1}_S$. However, the two notions agree for idempotent copulas.

Proposition 4.1. *If a copula C is non-atomic and idempotent then $\sigma_C = \sigma_C^*$ is the σ -algebra of invariant sets.*

Proof. Since idempotent copula is symmetric, $T_C^* = T_{C^t} = T_C$. Consequently, $T_C^* \circ T_C = T_C \circ T_C = T_{C * C} = T_C$. Now, if $T_C \mathbf{1}_S = \mathbf{1}_R$ then $T_C^* \mathbf{1}_R = \mathbf{1}_S$ and so $T_C^* \circ T_C \mathbf{1}_S = \mathbf{1}_S$. Hence S is an invariant set of C and so is R as $\mathbf{1}_R = T_C \mathbf{1}_S = \mathbf{1}_S$. That is, σ_C and σ_C^* are subsets of the class of invariant sets. The converse inclusions are clear. \square

Example 2. It is evident from the computations of σ_C and σ_C^* in Example 1 that Π is totally atomic but $M, W, \frac{M+W}{2}$ and L_α are non-atomic. In fact, every complete dependence copula is non-atomic.

Proposition 4.2. *Let C be a copula.*

1. $\sigma_C = \mathcal{B}$ if and only if C is left invertible.
2. $\sigma_C^* = \mathcal{B}$ if and only if C is right invertible.

Proof. We shall prove only 1. as 2. can be proved in a similar manner. If $C = C_{eg}$ then $T_C = T_{eg}$ maps ψ to $\psi \circ g$. So for every $B \in \mathcal{B}$, $T_C \mathbf{1}_B = \mathbf{1}_B \circ g = \mathbf{1}_{g^{-1}(B)}$. Hence, $\sigma_C = \mathcal{B}$. Conversely, Theorem 5 in [16] implies in particular that T is multiplicative, meaning $T(\psi \cdot \phi) = T\psi \cdot T\phi$ for all $\psi, \phi \in L^\infty$, if and only if $T = T_{eg}$ for some measure-preserving function g . By the linearity and continuity of $T = T_C$, it then suffices to show that $T(\mathbf{1}_B \cdot \mathbf{1}_E) = T\mathbf{1}_B \cdot T\mathbf{1}_E$ for all $B, E \in \mathcal{B}$. In fact, if $T\mathbf{1}_B = \mathbf{1}_{B'}$ and $T\mathbf{1}_E = \mathbf{1}_{E'}$ then $T\mathbf{1}_{B \cap E} = \mathbf{1}_{B' \cap E'}$ (by Proposition 3.1(3)) and the desired equality follows. \square

Lemma 4.3. *Let C be a copula with associated Markov operator T_C and doubly stochastic measure μ_C and $R, S \subseteq \mathcal{B}$. $T_C \mathbf{1}_S = \mathbf{1}_R$ if and only if $\mu_C(R \times S) = \lambda(R) = \lambda(S)$.*

Proof. Note the fact that $\mu_C(R \times S) = \int_R T_C \mathbf{1}_S d\lambda$. If $T_C \mathbf{1}_S = \mathbf{1}_R$ then $\mu_C(R \times S) = \lambda(R)$ and, by the property of T_C , $\lambda(R) = \int \mathbf{1}_R d\lambda = \int T_C \mathbf{1}_S d\lambda =$

$\int \mathbb{1}_S d\lambda = \lambda(S)$. Conversely, if $\mu_C(R \times S) = \lambda(R) = \lambda(S)$ then $\int_R T_C \mathbb{1}_S d\lambda = \mu_C(R \times S) = \lambda(S) = \int \mathbb{1}_S d\lambda = \int T_C \mathbb{1}_S d\lambda$ and $\int_R T_C \mathbb{1}_S d\lambda = \mu_C(R \times S) = \lambda(R) = \int_R \mathbb{1} d\lambda$. Since $T_C \mathbb{1}_S(t) \in [0, 1]$, $T_C \mathbb{1}_S(t) = 0$ for all $t \notin R$ and $T_C \mathbb{1}_S(t) = 1$ for all $t \in R$, i.e. $T_C \mathbb{1}_S = \mathbb{1}_R$. \square

Proposition 4.4. *Let T be a Markov operator and f and g be measure-preserving functions on $[0, 1]$. Then the following are equivalent.*

1. $T\mathbb{1}_{g^{-1}(B)} = \mathbb{1}_{f^{-1}(B)}$ for all $B \in \mathcal{B}$.
2. $T(\theta \circ g) = \theta \circ f$ for all Borel functions $\theta \in L^1$.
3. $T\psi = T_{ef} \circ T_{ge} \psi$ for all $g^{-1}(\mathcal{B})$ -measurable functions ψ .

Proof. 1 \Rightarrow 2: For every Borel set $B \subseteq I$, $T(\mathbb{1}_B \circ g) = T\mathbb{1}_{g^{-1}(B)} = \mathbb{1}_{f^{-1}(B)} = \mathbb{1}_B \circ f$. So, $T(\psi \circ g) = \psi \circ f$ for all simple Borel functions ψ . By the standard measure-theoretic argument, $T(\theta \circ g) = \theta \circ f$ for every Borel function θ .

2 \Rightarrow 3: Using Theorem 2.1, 2. implies that $T_{fe} \circ T \circ T_{eg} \theta = T_{fe}(T(\theta \circ g)) = T_{fe}(\theta \circ f) = \theta$ for every Borel function θ , and hence $T_{fe} \circ T \circ T_{eg} = I = T_M$, that is $C_{fe} * C * C_{eg} = M$. This means that $C_{fe} * C$ is the unique left inverse of C_{eg} , or $C_{fe} * C = C_{ge}$. Left multiplying by C_{ef} yields $C_{ef} * C_{fe} * C = C_{ef} * C_{ge}$, i.e. $T_{ef} \circ T_{fe} \circ T = T_{ef} \circ T_{ge}$. On the other hand, by Theorem 2.1, for any Borel function θ , $T_{ef} \circ T_{fe}(\theta \circ f) = (\theta \circ f)$ which gives $T_{ef} \circ T_{fe} \circ T(\theta \circ g) = T(\theta \circ g)$. Since $\{\theta \circ g: \theta \text{ is a Borel function}\}$ coincides with the class of $g^{-1}(\mathcal{B})$ -measurable functions, $T_{ef} \circ T_{fe} \circ T = T$ on the class of L^1 -functions which are $g^{-1}(\mathcal{B})$ -measurable. Therefore, $T\psi = (T_{ef} \circ T_{ge})\psi$ for all $(g^{-1}(\mathcal{B}) = \sigma_C)$ -measurable $\psi \in L^1$.

3 \Rightarrow 1: This is clear from taking $\psi = \mathbb{1}_{g^{-1}(B)}$. \square

The equivalence relation \approx on \mathcal{B} is defined as follows: $E \approx F$ if and only if the symmetric difference $E \Delta F$ has Lebesgue measure zero. Of course, \approx is still an equivalence relation on any $\mathcal{S} \subseteq \mathcal{B}$. The equivalence class of S in \mathcal{S} is denoted by $[S]_{\mathcal{S}}$ or just $[S]$ if no confusion can arise. The collection of equivalence classes in \mathcal{S} is denoted by $[\mathcal{S}]$. $[\mathcal{S}]$ is in fact a measure algebra induced by the Lebesgue measure λ . That is, $[\mathcal{S}]$ is a Boolean σ -algebra with respect to the operations $[S] \vee [R] = [S \cup R]$ and $[S] \wedge [R] = [S \cap R]$ together with $\lambda: [\mathcal{S}] \rightarrow [0, 1]$ defined overloadedly by $\lambda([S]) = \lambda(S)$ and satisfying $\lambda(\bigvee_{i=1}^{\infty} [A_i]) = \sum_{i=1}^{\infty} \lambda([A_i])$ if $[A_i] \wedge [A_j] = [\emptyset]$ for $i \neq j$. See [14, p.398].

T_C induces a well-defined equivalence class function $\Upsilon_C: [\sigma_C] \rightarrow [\sigma_C^*]$ mapping $[S]$ to $[R]$ if and only if $T_C \mathbb{1}_S = \mathbb{1}_R$. It follows from the defining property ii of T that $\lambda(R) = \lambda(S)$ and hence Υ_C is measure-preserving. The following lemma summarizes results from Theorem 3.2 and the proof of Theorem 3.3 in [4].

Lemma 4.5 ([4]). *Let $\mathcal{S} \subseteq \mathcal{B}$ be a non-atomic σ -algebra. Then*

1. *there exists a surjective isomorphism $\Psi: [\mathcal{S}] \rightarrow [\mathcal{B}]$ which means $\Psi([S]^c) = \Psi([S])^c$, $\Psi([S_1] \vee [S_2]) = \Psi([S_1]) \vee \Psi([S_2])$, $\lambda([S]) = \lambda(\Psi([S]))$ and Ψ is onto (it follows that Ψ is one-to-one and an isometry with respect to the*

metric $\rho([S], [R]) = \lambda([S] \triangle [R])$ and preserves countable unions and intersections); and

2. for any surjective isomorphism $\Psi: [\mathcal{S}] \rightarrow [\mathcal{B}]$, there exists a unique (a.e.) measure-preserving Borel function $h: [0, 1] \rightarrow [0, 1]$ such that $h^{-1}(\mathcal{B}) \subseteq \mathcal{S}$ (in fact, they are essentially equivalent,) and that $h^{-1}(B) \in \Psi^{-1}([B])$ for all $B \in \mathcal{B}$.

Proof. Enumerate $\mathbb{Q} \cap [0, 1] = \{r_n\}_{n \in \mathbb{N}}$ and set $I_n = [0, r_n]$. For each $n \in \mathbb{N}$, choose S_n in the equivalence class $\Psi^{-1}([I_n])$ so that $S_k = [0, 1]$ if $r_k = 1$ and $S_k \subset S_l$ whenever $r_k < r_l$. Define a Borel measure-preserving function $h(x) = \inf \{r_k : x \in S_k\}$. For $S \in \mathcal{S}$, choose $B_0 \in \Psi([S])$ and set $S_0 = h^{-1}(B_0)$. Since Ψ is 1-1, $[S] = \Psi^{-1}([B_0])$. We prove that $h^{-1}(B_0) \in \Psi^{-1}([B_0])$ by considering $\mathcal{M} = \{B \in \mathcal{B} : h^{-1}(B) \in \Psi^{-1}([B])\}$. Since \mathcal{M} is a monotone class containing all $[0, r_k]$, it contains all Borel sets and we have the claim. To prove uniqueness, if $k: [0, 1] \rightarrow [0, 1]$ is a measure-preserving Borel function such that $k^{-1}(B) \in \Psi^{-1}([B])$ for all $B \in \mathcal{B}$, then $k^{-1}([0, r_n]) \in \Psi^{-1}([I_n])$. So $\lambda(h^{-1}([0, r_n]) \triangle k^{-1}([0, r_n])) = 0$ for all n . By Proposition 10 in [14, Chapter 11 (p. 261)], $h = k$ a.e. \square

Theorem 4.6. *Let C be a copula with associated Markov operator T_C and doubly stochastic measure μ_C . For measure-preserving functions f and g on $[0, 1]$, the following are equivalent.*

1. $\mu_C(\text{graph}\{f = g\}) = 1$, where $\text{graph}\{f = g\} = \{(x, y) : f(x) = g(y)\}$.
2. $T_C = T_{ef} \circ T_{ge}$ on the class of $g^{-1}(\mathcal{B})$ -measurable functions.
3. C is non-atomic with $\sigma_C \supseteq g^{-1}(\mathcal{B})$ and $\sigma_C^* \supseteq f^{-1}(\mathcal{B})$.

Proof. $1 \Rightarrow 2$: By Proposition 4.4, Lemma 4.3 and the measure-preserving property of f and g , it suffices to show that $\mu_C(f^{-1}(B) \times g^{-1}(B)) = \lambda(f^{-1}(B))$ for all $B \in \mathcal{B}$. This follows from $\mu_C(f^{-1}(B) \times g^{-1}(B)) \leq \mu_C(f^{-1}(B) \times I) = \lambda(f^{-1}(B))$ and

$$\begin{aligned} \mu_C(f^{-1}(B) \times g^{-1}(B)) &\geq \mu_C(\{(x, y) : f(x) = g(y) \in B\}) \\ &= \mu_C(\{(x, y) : f(x) = g(y)\} \cap \{(x, y) : f(x) \in B\}) \\ &= \lambda(f^{-1}(B)). \end{aligned}$$

$2 \Rightarrow 1$: For each $n \in \mathbb{N}$ and $i = 1, 2, \dots, 2^n$, put $I_{i,n} = [\frac{i-1}{2^n}, \frac{i}{2^n}]$ and $B_n = \bigcup_i f^{-1}(I_{i,n}) \times g^{-1}(I_{i,n})$. Then $B_1 \supseteq B_2 \supseteq \dots \supseteq B_n \supseteq \dots$ and

$$\mu_C(B_n) = \sum_{i=1}^{2^n} \mu_C(f^{-1}(I_{i,n}) \times g^{-1}(I_{i,n})) = \sum_{i=1}^{2^n} \lambda(f^{-1}(I_{i,n})) = \sum_{i=1}^{2^n} \lambda(I_{i,n}) = 1,$$

where we have used Lemma 4.3 in the second equality and the measure-preserving property of f in the third. Set $B = \bigcap_{n=1}^{\infty} B_n$. We have $\mu_C(B) = \lim_{n \rightarrow \infty} \mu_C(B_n) =$

1. It then suffices to show that $B = \{(x, y) : f(x) = g(y)\}$. First, if $f(x) \neq g(y)$ then there exists $n \in \mathbb{N}$ such that $\frac{1}{2^n} < |f(x) - g(y)|$ and hence $(x, y) \notin B_n$. Conversely, it is clear that $\{(x, y) : f(x) = g(y)\} = \bigcup_i \{(x, y) : f(x) = g(y) \in I_{i,n}\} \subseteq B_n$ for all n .

2 \Rightarrow 3: By Proposition 4.4, $T_C \mathbb{1}_{g^{-1}(B)} = \mathbb{1}_{f^{-1}(B)}$ for all $B \in \mathcal{B}$, which implies that $g^{-1}(\mathcal{B}) \subseteq \sigma_C$ and $f^{-1}(\mathcal{B}) \subseteq \sigma_C^*$. But both $g^{-1}(\mathcal{B})$ and $f^{-1}(\mathcal{B})$ are non-atomic because g and f are measure-preserving. Therefore the finer σ -algebras σ_C and σ_C^* are non-atomic as well.

3 \Rightarrow 2: Our proof is in three steps. i) By applying Lemma 4.5(1) to σ_C and σ_C^* , it follows that $\Upsilon_C : [\sigma_C] \rightarrow [\sigma_C^*]$ defined earlier is one-to-one, onto, measure-preserving and preserves order, complementation and the lattice operation on equivalence classes corresponding to countable unions of monotonic sequence of sets, i.e. Υ_C is a surjective isomorphism. ii) By Lemma 4.5(2), there exists $\Xi_C : [\sigma_C^*] \rightarrow [\mathcal{B}]$ and a unique (a.e.) Borel function $f : I \rightarrow I$ such that $f^{-1}(B) \in \Xi_C^{-1}([B])$ for all $B \in \mathcal{B}$. So the composition $\Xi_C \circ \Upsilon_C : [\sigma_C] \rightarrow [\mathcal{B}]$ satisfies the same properties and induces a unique Borel function $g : I \rightarrow I$ such that

$$g^{-1}(B) \in (\Xi_C \circ \Upsilon_C)^{-1}([B]) \quad \text{for all } B \in \mathcal{B}. \quad (2)$$

iii) From (2), $\Upsilon_C([g^{-1}(B)]) = \Xi_C^{-1}([B])$ and, by the definition of Υ_C , $T_C \mathbb{1}_{g^{-1}(B)} = \mathbb{1}_{f^{-1}(B)}$ for all $B \in \mathcal{B}$. And the proof is done by Proposition 4.4. \square

In particular, for measure-preserving functions f, g on I , the support of $C_{ef} * C_{ge}$ “is” the graph of $f(x) = g(y)$, that is $\mu_{C_{ef} * C_{ge}}(\text{graph}\{f = g\}) = 1$, and its mass is distributed uniformly in the sense that $T_{C_{ef} * C_{ge}} \mathbb{1}_B = T_{ef}(T_{ge} \mathbb{1}_B)$ is $(\sigma_C^* = f^{-1}(\mathcal{B}))$ -measurable for all $B \in \mathcal{B}$.

Example 3. For a fixed $\alpha \in (0, 1)$, consider $C_\alpha = \alpha M + (1 - \alpha)W$ with Markov operator T_α . It is readily verified that $\sigma_{C_\alpha} = \sigma_{C_\alpha}^* = \{S \in \mathcal{B} : S = 1 - S\}$ and $T_\alpha \mathbb{1}_S = \mathbb{1}_S$ for all $S \in \sigma_{C_\alpha}$. But only $T_{\frac{1}{2}}$ has the property that $T_{\frac{1}{2}} \mathbb{1}_B$ is $\sigma_{C_{\frac{1}{2}}}$ -measurable for all $B \in \mathcal{B}$. Note also that $C_{\frac{1}{2}} = L_{\frac{1}{2}} * L_{\frac{1}{2}}^t$.

5 Copulas with fractal support

Let us recall the construction of copulas with self-similar supports in Fredricks et al. [6] put in the context of patched copulas [21, 1].

Definition 2. A *transformation matrix* is a matrix A with nonnegative entries, for which the sum of all entries is 1 and every row and column has at least one non-zero entry.

Given a transformation matrix $A = [a_{ij}]_{k \times \ell}$ where the first index (i) is the column number from left ($i = 1$) to right ($i = k$) and the second index (j) is the row number from bottom ($j = 1$) to top ($j = \ell$), the matrix multiplication of A and $B = [b_{jm}]_{\ell \times n}$ is defined by $[a_{ij}][b_{jm}] = \left[\sum_{j=1}^{\ell} a_{ij} b_{jm} \right]$. $A^t = [a_{ji}]_{\ell \times k}$.

This unconventional entry arrangement syncs well with the product of copulas defined [2] as

$$C * D(u, v) = \int_0^1 \partial_2 C(u, t) \partial_1 D(t, v) dt.$$

Let p_i denote the sum of the entries in the first i columns of A and let q_j denote the sum of the entries in the first j rows of A where $p_0 = 0$ and $q_0 = 0$. The partitions $\{p_i\}_{i=0}^k$ and $\{q_j\}_{j=0}^\ell$ of $[0, 1]$ yield a rectangular partition of $[0, 1]^2$ consisting of $k \cdot \ell$ rectangles with vertices (p_i, q_j) . Given a copula C , we construct a new copula denoted by $[A](C)$ by placing a scaled copy of C in each of the $k \cdot \ell$ rectangles weighted according to the mass given by the corresponding entry in A . It is defined for $(u, v) \in R_{ij} = [p_{i-1}, p_i] \times [q_{j-1}, q_j]$ by

$$\begin{aligned} [A](C)(u, v) &= \sum_{i' < i, j' < j} a_{i'j'} + \frac{u - p_{i-1}}{p_i - p_{i-1}} \sum_{j' < j} a_{ij'} + \frac{v - q_{j-1}}{q_j - q_{j-1}} \sum_{i' < i} a_{i'j} \\ &\quad + a_{ij} C\left(\frac{u - p_{i-1}}{p_i - p_{i-1}}, \frac{v - q_{j-1}}{q_j - q_{j-1}}\right). \end{aligned}$$

Here, empty sums are zero by convention. See [6] for more details. It will be more convenient to view $[A](C)$ as the so-called *patched copula* [21, 1] defined as

$$[A](C)(u, v) = \sum_{i=1}^k \sum_{j=1}^\ell a_{ij} C(F_i(u), G_j(v))$$

where $F_i(u) = \min\left(1, \frac{u - p_{i-1}}{p_i - p_{i-1}}\right) \mathbb{1}_{(p_{i-1}, \infty)}(u)$ is the uniform distribution on $[p_{i-1}, p_i]$ and $G_j(v) = \min\left(1, \frac{v - q_{j-1}}{q_j - q_{j-1}}\right) \mathbb{1}_{(q_{j-1}, \infty)}(v)$ is the uniform distribution on $[q_{j-1}, q_j]$. Denote $\Delta p_i = p_i - p_{i-1}$ and $\Delta q_j = q_j - q_{j-1}$.

We then investigate how this rectangular patching of a copula C according to the transformation matrix A affects C in terms of their Markov operators.

$$\begin{aligned} T_{[A](C)} f(x) &= \sum_{i=1}^k \sum_{j=1}^\ell \frac{a_{ij}}{\Delta q_j} \frac{d}{dx} \int_{q_{j-1}}^{q_j} \partial_2 C(F_i(x), G_j(t)) f(t) dt \\ &= \sum_{i=1}^k \sum_{j=1}^\ell a_{ij} \frac{d}{dx} \int_0^1 \partial_2 C(F_i(x), s) f \circ G_j^{-1}(s) ds \\ &= \sum_{i=1}^k \sum_{j=1}^\ell \frac{a_{ij}}{\Delta p_i} T_C(f \circ G_j^{-1})(F_i(x)). \end{aligned} \tag{3}$$

Consequently, $T_{[A](C)}(f)(F_i^{-1}(x)) = \sum_{j=1}^\ell \frac{a_{ij}}{\Delta p_i} T_C(f \circ G_j^{-1})(x)$, which can be written in matrix form as

$$\left[T_{[A](C)}(f)(F_i^{-1}(x)) \right]_{k \times 1} = \left[\frac{a_{ij}}{\Delta p_i} \right]_{k \times \ell} \left[T_C(f \circ G_j^{-1})(x) \right]_{\ell \times 1}.$$

See [19, 20] for essentially the same identity in terms of Markov kernels.

Define inductively $[A]^n(C) = [A]([A]^{n-1}(C))$ for $n \geq 2$. Fredricks et al. [6] showed that for any transformation matrix $A \neq [1]$ and copula C , $[A]^n(C)$ converges (pointwise and hence uniformly) to a unique copula C_A , as $n \rightarrow \infty$. Moreover, C_A is the fixed point of $[A]$, i.e. $[A](C_A) = C_A$. However, since uniform convergence will not suffice for our purposes, we shall investigate the convergence of $[A]^k(D)$ with respect to a stronger norm with respect to which the $*$ -product is jointly continuous. We choose the modified Sobolev norm defined as $\|C\|_S^2 = \|C\|_1^2 + \|C\|_2^2$ where $\|C\|_i^2 = \int_0^1 \int_0^1 |\partial_i C(u, v)|^2 du dv$. See [3].

Proposition 5.1. *Let A be a transformation matrix whose dimension is at least 2×2 and let C and D be copulas. Then*

$$\|[A](C) - [A](D)\|_S \leq r \|C - D\|_S$$

where $r^2 = \max \left(\sum_{j=1}^{\ell} \sum_{i=1}^k a_{ij}^2 \frac{q_j - q_{j-1}}{p_i - p_{i-1}}, \sum_{j=1}^{\ell} \sum_{i=1}^k a_{ij}^2 \frac{p_i - p_{i-1}}{q_j - q_{j-1}} \right) < 1$ and $\|\cdot\|_S$ is the modified Sobolev norm.

Proof. Denote $A = [a_{ij}]_{k \times \ell}$ with $k, \ell \geq 2$ and let $\{p_i\}_{i=0}^k, \{q_j\}_{j=0}^{\ell}$ be the induced partitions of $[0, 1]$ on the x -axis and y -axis, respectively. For $u \in (p_{i-1}, p_i)$ and $v \in (q_{j-1}, q_j)$, $\partial_1[A](C)(u, v) = a_{ij} F'_i(u) \partial_1 C(F_i(u), G_j(v)) + \sum_{k=1}^{j-1} a_{ik} F'_i(u)$ and so

$$|\partial_1[A](C)(u, v) - \partial_1[A](D)(u, v)| = \frac{a_{ij}}{p_i - p_{i-1}} |\partial_1 C(F_i(u), G_j(v)) - \partial_1 D(F_i(u), G_j(v))|.$$

Hence, $\|[A](C) - [A](D)\|_1^2$ is equal to

$$\begin{aligned} & \sum_{j=1}^{\ell} \sum_{i=1}^k \frac{a_{ij}^2}{(p_i - p_{i-1})^2} \int_{q_{j-1}}^{q_j} \int_{p_{i-1}}^{p_i} |\partial_1 C(F_i(u), G_j(v)) - \partial_1 D(F_i(u), G_j(v))|^2 du dv \\ &= \left(\sum_{j=1}^{\ell} \sum_{i=1}^k a_{ij}^2 \frac{q_j - q_{j-1}}{p_i - p_{i-1}} \right) \|C - D\|_1^2. \end{aligned}$$

A similar proof yields $\|[A](C) - [A](D)\|_2^2 = \left(\sum_{j=1}^{\ell} \sum_{i=1}^k a_{ij}^2 \frac{p_i - p_{i-1}}{q_j - q_{j-1}} \right) \|C - D\|_2^2$.

Now, since $a_{ij} \leq p_i - p_{i-1}$ and $\sum_i a_{ij} = q_j - q_{j-1}$,

$$\sum_{j=1}^{\ell} \sum_{i=1}^k a_{ij}^2 \frac{q_j - q_{j-1}}{p_i - p_{i-1}} \leq \sum_{j=1}^{\ell} \sum_{i=1}^k a_{ij} (q_j - q_{j-1}) = \sum_{j=1}^{\ell} (q_j - q_{j-1})^2 < \sum_{j=1}^{\ell} (q_j - q_{j-1}) = 1,$$

where we have used the assumption that $\ell \geq 2$ in the last inequality. In fact, the same assumption implies that $a_{ij} < p_i - p_{i-1}$ for some j . The same line of proof using $k \geq 2$ shows $\sum_{j=1}^{\ell} \sum_{i=1}^k a_{ij}^2 \frac{p_i - p_{i-1}}{q_j - q_{j-1}} < 1$ and the desired result follows. \square

We now have that the mapping $[A]$ is a contraction on the class of copulas \mathcal{C}_2 with respect to the modified Sobolev norm. Using the fact [3] that \mathcal{C}_2 is complete with respect to the modified Sobolev norm, it follows from the Contraction-Mapping Theorem that:

Theorem 5.2. *For each transformation matrix A of dimension at least 2×2 , and for any initial copula D , the sequence $\{[A]^r(D)\}_r$ converges to the copula C_A in the modified Sobolev norm.*

Definition 3. Let $A = [a_{ij}]_{k \times \ell}$ be a transformation matrix, $\emptyset \neq I \subseteq \{1, 2, \dots, k\}$ and $\emptyset \neq J \subseteq \{1, 2, \dots, \ell\}$. The pair (I, J) is called an *invariant pair* of A if $a_{ij} = 0$ for all $(i, j) \in (I \times J^c) \cup (I^c \times J)$ and $a_{ij} > 0$ for some $(i, j) \in I \times J$. Two invariant pairs (I_1, J_1) and (I_2, J_2) of A are called *disjoint* if $I_1 \cap I_2 = \emptyset$ and $J_1 \cap J_2 = \emptyset$. It follows that if they are not disjoint then both $I_1 \cap I_2$ and $J_1 \cap J_2$ are not empty.

We say that A is *disjointly decomposable* if A has a finite number of pairwise disjoint invariant pairs $(I_1, J_1), (I_2, J_2), \dots, (I_N, J_N)$ such that $\bigcup_{n=1}^N I_n = \{1, 2, \dots, k\}$ and $\bigcup_{n=1}^N J_n = \{1, 2, \dots, \ell\}$. For each $n = 1, 2, \dots, N$, let us denote by A_n the $k \times \ell$ matrix whose (i, j) th entry is a_{ij} if $i \in I_n$ and $j \in J_n$ and is equal to zero otherwise. Observe that $A = \sum_{n=1}^N A_n$ and in particular, every non-zero entry in A appears in exactly one A_n . We also say that A is *disjointly decomposable* as the sum $\sum_{n=1}^N A_n$ or *disjointly decomposable* by N invariant pairs.

Such a disjoint decomposition of A gives rise to two partitions of $[0, 1]$: $\{Q_n\}_{n=1}^N$ and $\{P_n\}_{n=1}^N$ defined by $Q_n = \bigcup_{j \in J_n} (q_{j-1}, q_j)$ and $P_n = \bigcup_{i \in I_n} (p_{i-1}, p_i)$. Then $\lambda(Q_n) > 0$ and $\lambda(P_n) > 0$ for all $n = 1, \dots, N$. Each pair (I_n, J_n) induces set functions \mathcal{G}_n and \mathcal{F}_n mapping $B \in \mathcal{B}$ to

$$\mathcal{G}_n(B) \equiv \bigcup_{j \in J_n} G_j^{-1}(B) \subseteq Q_n \quad \text{and} \quad \mathcal{F}_n(B) \equiv \bigcup_{i \in I_n} F_i^{-1}(B) \subseteq P_n.$$

Note that $G_j^{-1}((0, 1)) = (q_{j-1}, q_j)$ and $F_i^{-1}((0, 1)) = (p_{i-1}, p_i)$.

Remark. If A has N invariant pairs $(I_1, J_1), (I_2, J_2), \dots, (I_N, J_N)$, not necessarily disjoint, such that $\bigcup_{n=1}^N I_n = \{1, 2, \dots, k\}$ and $\bigcup_{n=1}^N J_n = \{1, 2, \dots, \ell\}$ then A is disjointly decomposable by $N' \geq N$ invariant pairs. In fact, without loss of generality, if (I_1, J_1) and (I_2, J_2) are not disjoint then $I' \equiv I_1 \cap I_2 \neq \emptyset$ and $J' \equiv J_1 \cap J_2 \neq \emptyset$ and it can be shown that $(I_1 \setminus I_2, J_1 \setminus J_2), (I_2 \setminus I_1, J_2 \setminus J_1), (I', J')$ are pairwise disjoint invariant pairs. Replacing $(I_1, J_1), (I_2, J_2)$ with these three pairs gives us a finer list of invariant pairs. Then repeat the process until all invariant pairs are pairwise disjoint.

Lemma 5.3. *If (I, J) is an invariant pair of a transformation matrix A and $S, R \in \mathcal{B}$ are such that $T_C \mathbb{1}_S = \mathbb{1}_R$, then $T_{[A](C)} \mathbb{1}_{\bigcup_{j \in J} G_j^{-1}(S)} = \mathbb{1}_{\bigcup_{i \in I} F_i^{-1}(R)}$.*

Proof. For $j \in J$, we have $\mathbb{1}_{\bigcup_{j' \in J} G_{j'}^{-1}(S)} \circ G_j^{-1}(x) = \mathbb{1}_S(x)$; otherwise, $\mathbb{1}_{\bigcup_{j' \in J} G_{j'}^{-1}(S)} \circ G_j^{-1}(x) = 0$. So, by (3),

$$\begin{aligned} T_{[A](C)} \mathbb{1}_{\bigcup_{j \in J} G_j^{-1}(S)} &= \sum_{i=1}^k \sum_{j=1}^{\ell} \frac{a_{ij}}{\Delta p_i} T_C \left(\mathbb{1}_{\bigcup_{j' \in J} G_{j'}^{-1}(S)} \circ G_j^{-1} \right) (F_i(x)) \\ &= \sum_{i=1}^k \sum_{j \in J} \frac{a_{ij}}{\Delta p_i} T_C(\mathbb{1}_S)(F_i(x)) \\ &= \sum_{i \in I} \frac{\sum_{j \in J} a_{ij}}{\Delta p_i} T_C(\mathbb{1}_S)(F_i(x)), \end{aligned}$$

where we have used the assumption that $a_{ij} = 0$ if $i \notin I$ and $j \in J$. Since for each $i \in I$, $a_{ij} = 0$ for all $j \notin J$, the sum $\sum_{j \in J} a_{ij}$ is equal to Δp_i and

$$T_{[A](C)} \mathbb{1}_{\bigcup_{j \in J} G_j^{-1}(S)} = \sum_{i \in I} T_C(\mathbb{1}_S)(F_i(x)) = \sum_{i \in I} \mathbb{1}_R(F_i(x)) = \mathbb{1}_{\bigcup_{i \in I} F_i^{-1}(R)}(x).$$

□

Lemma 5.4. *Let C be a copula. Suppose a transformation matrix A is disjointly decomposable by N invariant pairs. If $T_{[A](C)} \mathbb{1}_S = \mathbb{1}_R$ then for $n = 1, \dots, N$, $T_{[A](C)} \mathbb{1}_{S_n} = \mathbb{1}_{R_n}$ where $S_n \equiv S \cap Q_n$ and $R_n \equiv R \cap P_n$.*

Proof. For each $n = 1, \dots, N$, the positivity of $T_{[A](C)}$ implies that $T_{[A](C)} \mathbb{1}_{S_n} \leq T_{[A](C)} \mathbb{1}_S = \mathbb{1}_R$. If it holds that $T_{[A](C)} \mathbb{1}_{S_n} = 0$ for a.e. x not in P_n , then $T_{[A](C)} \mathbb{1}_{S_n} \leq \mathbb{1}_{R \cap P_n} = \mathbb{1}_{R_n}$. Summing over all n gives

$$\mathbb{1}_R = T_{[A](C)} \mathbb{1}_S = \sum_n T_{[A](C)} \mathbb{1}_{S_n} \leq \sum_n \mathbb{1}_{R_n} = \mathbb{1}_R$$

and hence $T_{[A](C)} \mathbb{1}_{S_n} = \mathbb{1}_{R_n}$ for all n . We then prove the claim. If $x \notin P_n \equiv \bigcup_{i' \in I_n} (p_{i'-1}, p_{i'})$, then $F_i(x) = 0$ or 1 for all $i \in I_n$ and

$$\begin{aligned} T_{[A](C)} \mathbb{1}_{S_n}(x) &= \sum_{i \in I_n} \sum_{j \in J_n} \frac{a_{ij}}{\Delta p_i} T_C \left(\sum_{j' \in J_n} \mathbb{1}_{S \cap (q_{j'-1}, q_{j'})} \circ G_j^{-1} \right) (F_i(x)) \\ &= \sum_{i \in I_n} \sum_{j \in J_n} \frac{a_{ij}}{\Delta p_i} T_C \left(\mathbb{1}_{S \cap (q_{j-1}, q_j)} \circ G_j^{-1} \right) (F_i(x)) = 0. \end{aligned}$$

We use the convention that 0 and 1 are not in the support of all functions. □

Theorem 5.5. *Suppose a transformation matrix A is disjointly decomposable by $N \geq 2$ invariant pairs. Then the invariant copula C_A is non-atomic.*

Proof. Let $S, R \in \mathcal{B}$ be such that $T_{C_A} \mathbb{1}_S = \mathbb{1}_R$ and $\lambda(S) = \lambda(R) > 0$. Since $C_A = [A](C_A)$, by Lemma 5.4, $T_{C_A} \mathbb{1}_{S_n} = \mathbb{1}_{R_n}$ for all $n = 1, 2, \dots, N$ where $S_n \equiv \mathcal{G}_n(S) \subseteq Q_n$ and $R_n \equiv \mathcal{F}_n(R) \subseteq P_n$ are such that $S = \bigcup_n S_n$ and




Figure 1: Supports of $[A_1]^r(\Pi)$, $r = 1, 2, \dots, 5$

$R = \bigcup_n R_n$. If one of the $\lambda(S_n)$'s is strictly between 0 and $\lambda(S)$ then S and R are not atoms of σ_{C_A} and $\sigma_{C_A}^*$, respectively. Otherwise, there is an n_1 such that $\lambda(S_{n_1}) = \lambda(S)$ and we repeat the process by applying Lemma 5.4 to $T_{C_A} \mathbf{1}_{S_{n_1}} = \mathbf{1}_{R_{n_1}}$ and obtain $S_{n_1,n} \equiv \mathcal{G}_n(S_{n_1}) \subseteq \mathcal{G}_n(Q_{n_1})$ and $R_{n_1,n} \equiv \mathcal{F}_n(R_{n_1}) \subseteq \mathcal{F}_n(P_{n_1})$ are such that $S_{n_1} = \bigcup_n S_{n_1,n}$, $R_{n_1} = \bigcup_n R_{n_1,n}$ and $T_{C_A} \mathbf{1}_{S_{n_1,n}} = \mathbf{1}_{R_{n_1,n}}$ for all $n = 1, 2, \dots, N$. If some $\lambda(S_{n_1,n})$ lies between 0 and $\lambda(S)$ then we are done. Otherwise, there must be an n_2 such that $\lambda(S_{n_1,n_2}) = \lambda(S)$. This process will certainly stop because $\lambda(\mathcal{G}_{n_k}(\dots(\mathcal{G}_{n_2}(Q_{n_1})))) = \lambda(Q_{n_k}) \cdots \lambda(Q_{n_2})\lambda(Q_{n_1}) \rightarrow 0$ as $k \rightarrow \infty$ and $\lambda(S) > 0$. \square

Example 4. Let $A_1 = K_0 + K_1$ and $A_2 = K_0 + K_2$ where

$$K_0 = \begin{bmatrix} 0 & 0 & 0 \\ 0 & 1/3 & 0 \\ 0 & 0 & 0 \end{bmatrix}, \quad K_1 = \begin{bmatrix} 1/12 & 0 & 1/4 \\ 0 & 0 & 0 \\ 1/4 & 0 & 1/12 \end{bmatrix} \quad \text{and} \quad K_2 = \begin{bmatrix} 1/6 & 0 & 1/6 \\ 0 & 0 & 0 \\ 1/6 & 0 & 1/6 \end{bmatrix}.$$

Then both transformation matrices A_1 and A_2 are disjointly decomposable by 2 invariant pairs $(\{1, 3\}, \{1, 3\})$ and $(\{2\}, \{2\})$. By Theorem 5.5, the invariant copulas C_{A_1} and C_{A_2} are non-atomic and hence, by Theorem 4.6, it is supported in the support of an implicit dependence copula. In fact, they share the same support shown in Figure 1. Note also that both copulas are symmetric as the corresponding matrices are. We will see later that (only) C_{A_2} can be factored as $L_1 * L_2^t$ for some left invertible copulas L_1, L_2 .

A transformation matrix $L = [\lambda_{in}]_{k \times N}$ is said to be a *left complete dependence matrix* if there is exactly one nonzero entry in each column, i.e. for each i , there exists a positive integer $n_i \leq N$ such that $\lambda_{in} = 0$ for $n \neq n_i$. *Right complete dependence matrices* are defined similarly.

Lemma 5.6. *Let L be a left complete dependence matrix with dimension at least 2×2 and C be a left invertible copula. Then $[L](C), [L]^2(C), \dots, [L]^r(C), \dots$ are left invertible copulas converging to a left invertible copula C_L . The same statement also holds if all occurrences of “left” are replaced by “right.”*

Proof. Since the general case can be proved by induction on r , it suffices to show that $[L](C)$ is left invertible. By definition, the x -partition of $[L](C)$ is determined by the only non-zero entry in each column: $p_0 = 0$ and $p_i = \sum_{i' \leq i} \lambda_{i'n_{i'}}$, $i = 1, \dots, k$. So $[L](C)(u, v) = \sum_{i=1}^k \lambda_{in_i} C(F_i(u), G_{n_i}(v))$ and

$$\begin{aligned} [L](C)^t * [L](C)(u, v) &= \sum_{i'=1}^k \sum_{i=1}^k \lambda_{in_i} \lambda_{i'n_{i'}} \int_0^1 \frac{\partial}{\partial t} C(F_{i'}(t), G_{n_{i'}}(u)) \frac{\partial}{\partial t} C(F_i(t), G_{n_i}(v)) dt \\ &= \sum_{i=1}^k \lambda_{in_i}^2 \int_0^1 \partial_1 C(F_i(t), G_{n_i}(u)) \partial_1 C(F_i(t), G_{n_i}(v)) F_i'(t)^2 dt \\ &= \sum_{i=1}^k \lambda_{in_i} C^t * C(G_{n_i}(u), G_{n_i}(v)) \\ &= \sum_{i=1}^k \lambda_{in_i} G_{n_i}(\min(u, v)) = M(u, v). \end{aligned}$$

Therefore, $[L](C)$ is left invertible. In general, we have $[L]^r(C)^t * [L]^r(C) = M$ and so, by taking the limit as $r \rightarrow \infty$ with respect to the modified Sobolev norm (see Theorem 5.2), $C_L^t * C_L = M$. \square

Theorem 5.7. *Let A be a $k \times \ell$ transformation matrix, with $k, \ell \geq 2$, which is disjointly decomposable by $N \geq 2$ invariant pairs. If all A_1, A_2, \dots, A_N have rank one, then $C_A = L * R$ for some left invertible copula L and right invertible copula R .*

Proof. If A_n has rank one, then there exist a row matrix $L_n = [\lambda_{in}]_{k \times 1}$ and a column matrix $R_n = [\rho_{nj}]_{1 \times \ell}$ such that $|L_n| = |A_n| = |R_n|$ and

$$A_n = \frac{1}{|A_n|} L_n R_n, \quad (4)$$

where $|A|$ denotes the sum of the absolute values of all entries in a matrix A . Stacking up L_n 's vertically and R_n 's horizontally, we obtain transformation matrices

$$L = [\lambda_{in}]_{k \times N} = \begin{bmatrix} L_N \\ \vdots \\ L_1 \end{bmatrix} \quad \text{and} \quad R = [\rho_{nj}]_{N \times \ell} = \begin{bmatrix} R_1 & \cdots & R_N \end{bmatrix}.$$

Since I_n 's are disjoint, each column of L has exactly one non-zero entry. Similarly, each row of R has exactly one non-zero entry.

We then show that $[L](C_1) * [R](C_2) = [A](C_1 * C_2)$ for any copulas C_1 and C_2 . For $m = 1, 2, \dots, N$, denote by H_m the uniform distribution on $\left[\sum_{n=1}^{m-1} |A_n|, \sum_{n=1}^m |A_n|\right]$. So $H'_m = \frac{1}{|A_m|}$ on its support. For $u, v \in [0, 1]$,

$$\begin{aligned} & [L](C_1) * [R](C_2)(u, v) \\ &= \int_0^1 \left(\sum_{i=1}^k \sum_{n=1}^N \lambda_{in} \partial_2 C_1(F_i(u), H_n(t)) \right) \left(\sum_{j=1}^\ell \sum_{m=1}^N \rho_{mj} \partial_1 C_2(H_m(t), G_j(v)) \right) dt \\ &= \sum_{i=1}^k \sum_{j=1}^\ell \left(\sum_{n=1}^N \frac{\lambda_{in} \rho_{nj}}{|A_n|} \right) C_1 * C_2(F_i(u), G_j(v)) \end{aligned} \quad (5)$$

$$= [A](C_1 * C_2)(u, v). \quad (6)$$

It is left to verify that the sum over n in (5) is equal to the (i, j) th-element in A . Since $\sum_{n=1}^N A_n$ is a disjoint decomposition of $A = [a_{ij}]_{k \times \ell}$, the equation (4) implies that if $(i, j) \in I_m \times J_m$ for some (unique) m then

$$a_{ij} = \frac{\text{the } (i, j)\text{th element of } L_m R_m}{|A_m|} = \frac{\lambda_{im} \rho_{mj}}{|A_m|} = \sum_{n=1}^N \frac{\lambda_{in} \rho_{nj}}{|A_n|},$$

where the last equality follows from the fact that $\lambda_{in} \rho_{nj} = 0$ if $(i, j) \notin I_n \times J_n$. It also clearly follows from this fact that if $(i, j) \notin I_m \times J_m$ for all m then $\sum_{n=1}^N \frac{\lambda_{in} \rho_{nj}}{|A_n|} = 0 = a_{ij}$.

By (6), a proof by induction on $r = 1, 2, \dots$ yields $[L]^r(C_1) * [R]^r(C_2) = [A]([L]^{r-1}(C_1) * [R]^{r-1}(C_2)) = \dots = [A]^r(C_1 * C_2)$. So $[L]^r(E) * [R]^r(E) = [A]^r(E)$ for all $r \geq 1$ if E is an idempotent copula, e.g. $E = M$ or Π . By Lemma 5.6, $[L]^r(M)$ are left invertible copulas and $[R]^r(M)$ are right invertible copulas.

By Theorem 5.2, with respect to the modified Sobolev norm, $[L]^r(M)$, $[R]^r(M)$ and $[A]^r(M)$ converge respectively to a left invertible copula C_L , a right invertible copula C_R and a copula C_A satisfying $[L](C_L) = C_L$, $[R](C_R) = C_R$ and $[A](C_A) = C_A$. As a consequence of the joint continuity of the $*$ -product with respect to $\|\cdot\|_S$, we obtain $C_L * C_R = C_A$ as desired. \square

Example 5. Fix $r \in (0, 1/2)$ and consider the transformation matrix

$$\begin{bmatrix} r/2 & 0 & r/2 \\ 0 & 1-2r & 0 \\ r/2 & 0 & r/2 \end{bmatrix} = \begin{bmatrix} 0 & 0 & 0 \\ 0 & 1-2r & 0 \\ 0 & 0 & 0 \end{bmatrix} + \begin{bmatrix} r/2 & 0 & r/2 \\ 0 & 0 & 0 \\ r/2 & 0 & r/2 \end{bmatrix}.$$

Both matrices on the right hand side have rank one which implies by Theorem 5.7 that the invariant copula can be factored as the product of a left invertible copula and a right invertible copula. When $r = 1/3$, the transformation matrix is A_2 in Example 4. The copula factors, C_{L_2} and C_{R_2} , of C_{A_2} are shown (approximately) in Figure 2, where $L_2 = \begin{bmatrix} 0 & 1/3 & 0 \\ 1/3 & 0 & 1/3 \end{bmatrix} = R_2^t$.




Figure 2: Supports of $[A_2]^5(\Pi)$, $[L_2]^5(\Pi)$ and $[R_2]^5(\Pi)$ in Example 5




Figure 3: Supports of $[A_3]^5(\Pi)$, $[L_3]^5(\Pi)$ and $[R_3]^5(\Pi)$ in Example 6

Example 6. Let us also consider the factorization of a non-symmetric invariant copula by first setting

$$K_3 = \begin{bmatrix} 3/28 & 0 & 27/140 \\ 0 & 0 & 0 \\ 1/7 & 0 & 9/35 \end{bmatrix}, \quad L_3 = \begin{bmatrix} 0 & 3/10 & 0 \\ 1/4 & 0 & 9/20 \end{bmatrix} \quad \text{and} \quad R_3 = \begin{bmatrix} 3/10 & 0 \\ 0 & 3/10 \\ 4/10 & 0 \end{bmatrix}.$$

Then the non-symmetric transformation matrix $A_3 = K_0 + K_3$ is disjointly decomposable by 2 invariant pairs $(\{1, 3\}, \{1, 3\})$ and $(\{2\}, \{2\})$. Since K_0 and K_3 have rank one, it follows from Theorem 5.7 and its proof that $C_{A_3} = C_{L_3} * C_{R_3}$. Approximations of their supports are illustrated in Figure 3.

6 Acknowledgements

The author is grateful for the financial support from the Commission on Higher Education and the Thailand Research Fund through grant no. RSA5680037. We would also like to thank Wolfgang Trutschnig and Anusorn Junchuay for sharing and helping with the R codes used to produce copulas with fractal supports and their copula factors.

References

- [1] Chaidee, N., Santiwipanont, T., Sumetkijakan S.: Patched approximations and their convergence. *Comm. Statist. Theory Methods.* 45, 2654–2664 (2016)
- [2] Darsow, W.F., Nguyen, B., Olsen, E.T.: Copulas and Markov processes. *Illinois J. Math.* 36, 600–642 (1992)
- [3] Darsow, W.F., Olsen, E.T.: Norms for copulas. *Int. J. Math. Math. Sci.* 18, 417–436 (1995)
- [4] Darsow, W.F., Olsen, E.T.: Characterization of idempotent 2-copulas. *Note Mat.* 30, 147–177 (2010)
- [5] Durante, F., Sempi, C.: Principles of copula theory. CRC Press (2015)
- [6] Fredricks, G.A., Nelsen, R.B., Rodríguez-Lallena, J.A.: Copulas with fractal supports. *Insurance Math. Econom.* 37, 42–48 (2005)
- [7] Kamnitui, N., Santiwipanont, T., Sumetkijakan, S.: Dependence measuring from conditional variances. *Depend. Model.* 3, 98–112 (2015)
- [8] Kimeldorf, G., Sampson, A.R.: Monotone dependence. *Ann. of Statist.* 6, 895–903 (1978)
- [9] Mikusiński, P., Sherwood, H., Taylor, M.D.: Probabilistic interpretations of copulas and their convex sums. In: Dall’Aglia, G., Kotz, S., Salinetti, G. (eds.) *Advances in Probability Distributions with Given Marginals: Beyond the Copulas*, pp. 111–128. Kluwer Academic Publishers, Dordrecht (1991)
- [10] Mikusiński, P., Sherwood, H., Taylor, M.D.: Shuffles of min. *Stochastica* 13, 61–74 (1992)
- [11] Nelsen, R.B.: *An Introduction to Copulas*. 2nd ed. Springer-Verlag, New York (2006)
- [12] Olsen, E.T., Darsow, W.F., Nguyen, B.: Copulas and Markov operators. *Lecture Notes-Monograph Series* 28, 244–259 (1996)
- [13] Rényi, A.: On measures of dependence. *Acta. Math. Acad. Sci. Hungar.* 10, 441–451 (1959)
- [14] Royden, H.L.: *Real analysis*, 3rd ed. Macmillan, New York (1988).
- [15] Ruankong, P., Santiwipanont, T., Sumetkijakan, S.: Shuffles of copulas and a new measure of dependence. *J. Math. Anal. Appl.* 398, 392–402 (2013)
- [16] Sakai, Y., Shimogaki, T.: Equimeasurability of functions and doubly stochastic operators. *Kodai Math. Sem. Rep.* 24, 203–211 (1972)

- [17] Siburg, K.F., Stoimenov, P.A.: A measure of mutual complete dependence. *Metrika* 71, 239–251 (2009)
- [18] Trutschnig, W.: On a strong metric on the space of copulas and its induced dependence measure. *J. Math. Anal. Appl.* 384, 690–705 (2011)
- [19] Trutschnig, W.: Idempotent copulas with fractal support. In: Greco, S., Bouchon-Meunier, B., Coletti, G., Fedrizzi, M., Matarazzo, B., Yager, R.R. (eds.) *Advances in Computational Intelligence*, Vol. 298, pp. 161–170. Springer, Berlin Heidelberg (2012)
- [20] Trutschnig, W., Sánchez, J.F.: Idempotent and multivariate copulas with fractal support. *J. Statist. Plann. Inference*, 142, 3086–3096 (2012)
- [21] Yanting, Z., Jingping, Y., Jianhua, Z.H.: Approximation of bivariate copulas by patched bivariate Fréchet copulas. *Insurance Math. Econom.* 48, 246–256 (2011)