

Buhmann covariance functions, their compact supports, and their smoothness

E. PORCU*, V.P. ZASTAVNYI[†], and M. BEVILACQUA[‡]

March 2, 2022

Abstract

We consider the Buhmann class of compactly supported radial basis functions, which includes a wealth of special cases that have been studied in both numerical analysis and spatial statistics literatures. In particular, the celebrated Wu, Wendland and Missing Wendland functions are notable special cases of this class. We propose a very simple difference operator and show the conditions for which the application of it to Buhmann functions preserves positive definiteness on m -dimensional Euclidean spaces. We also show that the application of the difference operator increases smoothness at the origin, whilst keeping positive definiteness in the same m -dimensional Euclidean space, as well as compact support. Thus, our operator is a competitor of the celebrated Montée operator, which allows to increase the smoothness at the origin, at the expense of losing positive definiteness in the space where the radial basis function is originally defined. The proofs of our results highlight surprising connections with past literatures on celebrated class of functions. Amongst them, absolute and completely monotone functions. *Keywords:* Buhmann functions; Compact Support; Completely Monotonic; Fourier transforms; Laplace transforms.

*Departamento de Matemática, Universidad Federico Santa Maria,

[†]Department of Mathematics, Donetsk National University

[‡]Department of Statistics, University of Valparaiso.

1 Introduction

Interpolation of data has a notable importance in both numerical analysis and geostatistical communities: in geostatistics, the underlying structure of the data is assumed to be a stochastic process, with the interpolation procedure known as kriging [23]. The method is mathematically equivalent to kernel interpolation, a method used in numerical analysis for the same problem, but derived under completely different modelling assumptions.

Radial basis functions are well-known and successful tools for the interpolation of data in many dimensions [5]. Several radial basis functions of compact support that give rise to non-singular interpolation problems have been proposed, and we cite [19, 12, 5, 30, 20, 32, 33] and the impressive reviews in [8] and [21], amongst others. In particular, the motivation behind Buhmann's [5] tour de force is to propose radial functions of compact support that also give positive definite matrices and have genuinely banded interpolation matrices (similarly to the multiquadric and Gaussian kernels). Of such nature are those ones we will discuss in the present paper. Early examples of radial functions with compact support that have a simple piecewise polynomial structure are due to [32]. Then Schaback and Wu [22], Wendland [30] and finally Schaback [20] established several of their special properties, such as certain optimality facts about their degree and smoothness. The functions proposed by Buhmann are closely related to the so-called multiply monotone radial basis functions as discussed in [15].

Radial basis functions are known under the name of covariance (correlation) functions in the geostatistical community: the use of compactly supported covariance functions has been advocated in a number of papers, and we refer the reader to [13], with the references therein, and to [7] for a recent effort under the framework on multivariate Gaussian fields. Covariance functions with compact support represent the building block for the construction of methods allowing to overcome the big data problem ([11]). The recent work of [4] brought even more attention on the role of some classes of compactly supported covariances for asymptotically optimal prediction on a bounded set of \mathbb{R}^d . From the cited works it has become apparent that the smoothness at the origin (intended as even extension) of a compactly supported and isotropic covariance function plays a crucial role for both estimation and prediction. Wendland functions [30] have been especially popular, being compactly supported over balls of \mathbb{R}^d with arbitrary radii, and additionally allowing for a continuous parameterization of

differentiability at the origin, in a similar way to the Matérn family ([25]).

The tour de force in [30], [13] and [39] puts emphasis on linear operators, called *Montée*, that allow to increase the smoothness of a given radial function, being positive definite on \mathbb{R}^d . This is done at the expense of losing positive definiteness, which is only achieved on \mathbb{R}^{d-2} , for $d \geq 3$.

Figure 1 depicts the following situation: Dashed lines report Wendland functions [30] with unit compact support, for $k = 0, 1, 2$ (from left to right) being the parameter that allows to govern differentiability at the origin. Dashed-dotted lines report the respective Wendland functions for a compact support equal to 0.75. The functions depicted with continuous lines report their weighted differences (see Equation (17)). We can clearly appreciate that the level of differentiability changes for these last.

This gives a substantial motivation for the present work: we consider the Buhmann functions as defined in Zastavnyi [36] on the basis of [5]. This class includes as special case many other classes of compactly supported covariance functions, such as Askey [1] Wendland [30] and Missing Wendland [20] functions, as well as the Zastavnyi [34, 35, 36] and Trigub [26, 27] classes. Finally, also Wu functions [32] and the celebrated spherical model [29] are included as special cases. We show that some parameterized differences of Buhmann functions preserve positive definiteness in m -dimensional Euclidean spaces, and we then determine the exact level of smoothness induced by such operation.

We focus throughout on the class Φ_m of continuous functions $\varphi : [0, \infty) \rightarrow \mathbb{R}$ such that $\varphi(\|\mathbf{x}\|)$ is positive definite on \mathbb{R}^m . Thus, $\varphi(\|\cdot\|)$ with $\varphi(0) = 1$ is the correlation function of some Gaussian field.

Apparently, the functions $C(\cdot) := \varphi(\|\cdot\|)$ are radially symmetric and Schoenberg's theorem (1938, see [6] for a more recent discussion) uniquely identifies them as scale mixtures of the type

$$\varphi(t) = \int_{[0, \infty)} \Omega_m(rt) F(dr), t \geq 0, \quad (1)$$

with F a uniquely determined probability measure, and $\Omega_m(\cdot)$ being the characteristic function of a random vector that is uniformly distributed on the spherical shell of \mathbb{R}^m . Daryl and Porcu [6] put emphasis on the measure F , termed Schoenberg measure there. It is well known [13] that any random vector \mathbf{X} of \mathbb{R}^m with characteristic function φ can be written as $\mathbf{X} = \boldsymbol{\eta}R$, with $\boldsymbol{\eta}$ having Ω_m as characteristic function, R a positive random variable

distributed according to F , and η and R are stochastically independent. The identity above is intended as equality in distribution.

The class Φ_m is nested, with the following inclusion relation

$$\Phi_1 \supset \Phi_2 \supset \dots \supset \Phi_\infty := \bigcap_{m \geq 1} \Phi_m, \quad m \in \mathbb{N},$$

being strict (see, for example, [6, 13]). A function $f : (0, \infty) \rightarrow \mathbb{R}$ is called completely monotone if it is infinitely often differentiable and $(-1)^n f^{(n)}(x) \geq 0$, for all $n \in \mathbb{Z}_+$ and for all $x > 0$. The set of completely monotone functions on $(0, \infty)$ is denoted \mathcal{CM} . By Schoenberg's theorem, $\varphi(t) \in \Phi_\infty$ if and only if $\varphi(\sqrt{t}) \in \mathcal{CM}$ with $\varphi(0) = \varphi(+0) < \infty$.

The plan of the paper is the following. Section 2 introduces the Buhmann class and offers a formal statement of the problem. Section 3 exposes the structure of the solution. We also characterize the smoothness of differences of Buhmann functions in a neighborhood of the origin. Section 4 illustrates connections with previous literature.

2 Buhmann functions. Statement of the problem

2.1 Buhmann's class and relations with previous literature

We denote $C(\mathbb{R}^m)$ the set of continuous functions on \mathbb{R}^m , for $m = 1, 2, \dots$. Let $\delta, \mu, \nu \in \mathbb{C}_+ := \{z \in \mathbb{C} : \operatorname{Re} z > 0\}$ and $\alpha \in \mathbb{C}$. Zastavnyi [36] (2006) proposed the following even functions given on \mathbb{R} :

$$\varphi_{\delta, \mu, \nu, \alpha}(x) := \begin{cases} \int_{|x|}^1 (s^2 - x^2)^{\nu-1} (1 - s^\delta)^{\mu-1} s^{\alpha-2\nu+1} ds, & |x| < 1 \\ 0, & |x| \geq 1. \end{cases} \quad (2)$$

If $\delta, \mu, \nu \in \mathbb{C}_+$, then arguments in Proposition 1 and Theorem 1 in [36] show, respectively, that $\varphi_{\delta, \mu, \nu, \alpha} \in C(-1, 1)$ if and only if $\alpha \in \mathbb{C}_+$ and that $\varphi_{\delta, \mu, \nu, \alpha} \in C(\mathbb{R})$ if and only if $\alpha, \mu + \nu - 1 \in \mathbb{C}_+$. If $\delta, \mu, \nu, \alpha \in \mathbb{C}_+$, then $\varphi_{\delta, \mu, \nu, \alpha}(0) = B(\alpha/\delta, \mu)/\delta$, with B denoting the Beta function.

The functions $\varphi_{\delta, \mu, \nu, \alpha}$ coincide (modulo some positive factors) with the functions

$$\phi_{\delta, \varrho, \lambda, \alpha}(x) \equiv 2\varphi_{2\delta, \varrho+1, \lambda+1, 2\alpha+2}(x), \quad x \in \mathbb{R},$$

introduced by Martin Buhmann [5]. We thus term them Buhmann functions throughout. The class $\varphi_{\delta,\mu,\nu,\alpha}$ includes a wealth of interesting special cases. For instance, $\mu\delta\varphi_{\delta,\mu,1,\delta}(x) = (1 - |x|^\delta)_+^\mu$, which implies that $\mu\varphi_{1,\mu,1,1}$ coincides with the Askey functions [1]. Also, we have that

$$\varphi_{1,\mu,\nu,2\nu-1}(x) \equiv h_{\mu,\nu}(x) \equiv \frac{2^{\nu-1}\Gamma(\nu)}{\mu}\psi_{\mu,\nu-1}(x), \quad x \in \mathbb{R}, \quad (3)$$

with the functions $h_{\mu,\nu}$ being introduced by Zastavnyi (2002) [35, 34] and defined as follows: $h_{\mu,\nu}(x) := 0$ for $|x| \geq 1$ and

$$h_{\mu,\nu}(x) := \int_{|x|}^1 (2u - |x|)g_{\mu,\nu}(u)g_{\mu,\nu}(u - |x|) du, \quad |x| < 1, \quad (4)$$

where $g_{\mu,\nu}(u) := u^{\mu-1}(1 - u^2)^{\nu-1}$, $u \in (0, 1)$, $\mu, \nu \in \mathbb{C}_+$.

Functions of the form (4) arise in the study of exponential type entire functions without zeros in the lower half-plane [37, Proposition 5.1]. The functions $\psi_{\mu,\nu-1}$, with $\mu > 0$, $\nu \in \mathbb{N}$, have been introduced in 1995 by Wendland [30], and they have been termed Wendland functions in both numerical analysis and geostatistical literatures: for $\mu > 0$, $k \in \mathbb{Z}_+$, we have

$$\psi_{\mu,0}(x) := \psi_\mu(x) := (1 - |x|)_+^\mu, \quad \psi_{\mu,k} := I^k\psi_\mu \quad (k \in \mathbb{N}),$$

where $I(f)(x) := \int_{|x|}^{+\infty} sf(s) ds$ is the Matheron's [14] Montée operator (provided the integral is well defined), and where I^k is the k -fold application of the operator I . Arguments in [30] and subsequently [13] show that $I\varphi$ belongs to the class Φ_{m-2} whenever $\varphi \in \Phi_m$, for $m \geq 3$. For $k < 2m$, the k -fold application of the Montée operators shows that $I^k\varphi \in \Phi_{m-2k}$, $k \in \mathbb{N}$.

Gneiting [13, Equation (17)] has proposed a generalization of Wendland functions on the basis of the fractional Montée operator, which coincides with the normalized Buhmann functions $\varphi_{1,\mu+1,\nu,2\nu}(x)/\varphi_{1,\mu+1,\nu,2\nu}(0)$, $\mu, \nu > 0$, as well as with the functions $h_{\mu,\nu+1}(x)/h_{\mu,\nu+1}(0) \equiv \psi_{\mu,\nu}(x)/\psi_{\mu,\nu}(0)$ (see Equation (6)). Arguments in [5] show that Wu functions [32] and consequently the spherical model are special cases of the Buhmann class.

For $r \in \mathbb{Z}_+$ and $k \in \mathbb{N}$, we have

$$h_{r+k,r+1}(x) \equiv B(r+k, 2r+1)A_{r,2k-1}(x),$$

with the splines $A_{r,2k-1}$ introduced by Trigub (1987), and we refer to [26], [27, § 6.2.13, 6.2.16, 6.3.12] for their analytical expression which is not reported here. Equation above

in turn highlights the explicit connection between Trigub splines and Wendland functions:

$A_{r,2k-1}(x) \equiv \psi_{r+k,r}(x)/\psi_{r+k,r}(0)$, for $r \in \mathbb{Z}_+$ and $k \in \mathbb{N}$.

For a proof of the identities above, the reader is referred to Zastavnyi and Trigub [35, Remarks 10 and 11], to [34, Theorems 12 and 13], [36] and [38, § 4.7].

Arguments in Proposition 4 of [36] show that, for $\delta, \mu, \nu \in \mathbb{C}_+$ and $x \in \mathbb{R}$,

$$\begin{aligned} \varphi_{2, \frac{\mu}{2}, \frac{\mu}{2} + \nu, 2\nu-1}(x) &\equiv \frac{2^{\mu-1} \Gamma\left(\frac{\mu}{2}\right) \Gamma\left(\frac{\mu}{2} + \nu\right)}{\Gamma(\mu)} \varphi_{1, \mu, \nu, 2\nu-1}(x), \\ 2\nu \varphi_{\delta, \mu+1, \nu, 2\nu}(x) &\equiv \delta \mu \varphi_{\delta, \mu, \nu+1, 2\nu+\delta}(x), \end{aligned} \quad (5)$$

and, for $\mu, \nu \in \mathbb{C}_+$ and $x \in \mathbb{R}$, we also have the obvious identities:

$$\varphi_{1, \mu+1, \nu, 2\nu}(x) \equiv \frac{\mu}{2\nu} \varphi_{1, \mu, \nu+1, 2\nu+1}(x) \equiv \frac{\mu}{2\nu} h_{\mu, \nu+1}(x) \equiv 2^{\nu-1} \Gamma(\nu) \psi_{\mu, \nu}(x). \quad (6)$$

2.2 Buhmann class and its Fourier and Laplace Transforms

After the illustration of the relation between Buhmann and other celebrated classes of radial basis function, we need some preliminary material in order to provide a better description of the results coming subsequently. For a function h defined on $(0, \infty)$ and $m \in \mathbb{C}$, we define the Hankel transform \mathfrak{F}_m as follows:

$$\mathfrak{F}_m(h)(t) := t^{1-\frac{m}{2}} \int_0^\infty h(u) u^{\frac{m}{2}} J_{\frac{m}{2}-1}(tu) du = \int_0^\infty h(u) u^{m-1} j_{\frac{m}{2}-1}(tu) du, \quad t > 0, \quad (7)$$

where J_λ is the Bessel function of the first kind (see [28, Sec. 3.1]) and

$$j_\lambda(x) := \frac{J_\lambda(x)}{x^\lambda} = \frac{1}{2^\lambda} \sum_{k=0}^\infty \frac{1}{\Gamma(k + \lambda + 1)} \cdot \frac{\left(-\frac{x^2}{4}\right)^k}{k!}, \quad x \in \mathbb{C}, \quad \lambda \in \mathbb{C}. \quad (8)$$

Remark 1 For $m \in \mathbb{N}$ the transform \mathfrak{F}_m is connected with the Fourier transforms F_m of radial functions through the identity

$$F_m(h(\|\cdot\|))(\mathbf{x}) = (2\pi)^{\frac{m}{2}} \mathfrak{F}_m(h)(\|\mathbf{x}\|), \quad \mathbf{x} \in \mathbb{R}^m.$$

These facts and Bochner-Khintchine theorem (see, for example [18, 24, 27]) imply that, if h is a continuous functions on $[0, \infty)$ and $\int_0^\infty t^{m-1} |h(t)| dt < \infty$, then $h \in \Phi_m$ if and only if $\mathfrak{F}_m(h)$ is nonnegative on the positive real line.

For $\delta, \mu, \alpha + 1 \in \mathbb{C}_+$ and $\nu \in \mathbb{C}$, we define the function $I_{\delta, \mu, \nu, \alpha} : \mathbb{R}_+ \rightarrow \mathbb{C}$ through

$$\begin{aligned} I_{\delta, \mu, \nu, \alpha}(t) &:= t^{-\alpha-1-\delta(\mu-1)} \int_0^t (t^\delta - u^\delta)^{\mu-1} u^{\alpha-\nu+\frac{1}{2}} J_{\nu-\frac{1}{2}}(u) \, du \\ &= \int_0^1 (1-x^\delta)^{\mu-1} x^\alpha j_{\nu-\frac{1}{2}}(tx) \, dx, \quad t > 0. \end{aligned} \quad (9)$$

The following result reports succinctly a collection of useful results from [36] (Theorems 2, 3 and Proposition 4 (Assertions 1,3)).

Theorem 1 (Zastavnyi [36]) *Let the functions I and \mathfrak{F}_m as being defined through Equations (9) and (7), respectively. Denote with I' the first derivative of I . Then, the following assertions are true:*

1. *Let $\delta, \mu, \nu, m, \alpha + m \in \mathbb{C}_+$. Then $\mathfrak{F}_m(\varphi_{\delta, \mu, \nu, \alpha})(t) = 2^{\nu-1} \Gamma(\nu) I_{\delta, \mu, \frac{m-1}{2} + \nu, m-1+\alpha}(t)$.*

Moreover, if $n, m - n + 2\nu \in \mathbb{C}_+$, then

$$\mathfrak{F}_m(\varphi_{\delta, \mu, \nu, \alpha})(t) = \frac{2^{\frac{n-m}{2}} \Gamma(\nu)}{\Gamma(\frac{m-n}{2} + \nu)} \mathfrak{F}_n(\varphi_{\delta, \mu, \frac{m-n}{2} + \nu, m-n+\alpha})(t), \quad t \geq 0.$$

2. *Let $\delta, \mu, \alpha + 1 \in \mathbb{C}_+$ and $\nu \in \mathbb{C}$. Then*

$$I'_{\delta, \mu, \nu, \alpha}(t) = -t I_{\delta, \mu, \nu+1, \alpha+2}(t), \quad t > 0. \quad (10)$$

3. *If $\mu, \nu \in \mathbb{C}_+$, $t > 0$, then*

$$I_{1, \mu, \nu, 2\nu-1}(t) = \frac{2^{\frac{1}{2}-\nu} \Gamma(\mu) \Gamma(2\nu)}{\Gamma(\nu + \frac{1}{2}) \Gamma(\mu + 2\nu)} {}_1F_2\left(\nu; \frac{\mu + 2\nu}{2}, \frac{\mu + 2\nu + 1}{2}; -\frac{t^2}{4}\right). \quad (11)$$

4. *Let $\delta, \mu, \nu \in \mathbb{C}_+$ and $\alpha \in \mathbb{C}$. Then*

$$\varphi_{\delta, \mu, \nu+1, \alpha+2}(t) = 2\nu \int_{|t|}^{\infty} u \varphi_{\delta, \mu, \nu, \alpha}(u) \, du, \quad t \neq 0. \quad (12)$$

A relevant remark is that Equation (11) describes the spectral density of the Wendland functions. Another remarkable consequence of Theorem 1 is that, for $\mu, \nu, m, 2\nu - 1 + m \in \mathbb{C}_+$,

$$\mathfrak{F}_m(h_{\mu, \nu})(t) = \mathfrak{F}_m(\varphi_{1, \mu, \nu, 2\nu-1})(t) = \frac{2^{\frac{1-m}{2}} \Gamma(\nu)}{\Gamma(\frac{m-1}{2} + \nu)} \mathfrak{F}_1(h_{\mu, \frac{m-1}{2} + \nu})(t), \quad t \geq 0, \quad (13)$$

which in turn shows, in concert with [35, Lemma 12], that in some cases the Hankel transforms above can be written in closed form. Specifically, we have

$$\begin{aligned} \mathfrak{F}_m(h_{\mu,\nu})(t) &= \mathfrak{F}_m(\varphi_{1,\mu,\nu,2\nu-1})(t) = 2^{\nu-1}\Gamma(\nu)I_{1,\mu,\frac{m-1}{2}+\nu,m-1+2\nu-1}(t) = \\ &D(m,\mu,\nu) \cdot {}_1F_2\left(\frac{m-1}{2} + \nu; \frac{m-1}{2} + \nu + \frac{\mu}{2}, \frac{m-1}{2} + \nu + \frac{\mu+1}{2}; -\frac{t^2}{4}\right), \quad (14) \\ \text{with } D(m,\mu,\nu) &:= \frac{2^{-\frac{m}{2}}\Gamma(\nu)\Gamma(\mu)\Gamma(m-1+2\nu)}{\Gamma(\frac{m}{2} + \nu)\Gamma(\mu+m-1+2\nu)}, \quad \mu, \nu, m, 2\nu-1+m \in \mathbb{C}_+. \end{aligned}$$

Let us use the abuse of notation $\widehat{h}_{\mu,\nu}$ for the one dimensional Fourier transform of the function $h_{\mu,\nu}$. We also denote with L the Laplace transform operator. For $\mu, \nu \in \mathbb{C}_+$, arguments in Zastavnyi and Trigub [35, Equation (44)] show that

$$L\left(t^{2\nu+\mu-1}\widehat{h}_{\mu,\nu}(t)\right)(x) := \int_0^\infty e^{-tx} t^{2\nu+\mu-1}\widehat{h}_{\mu,\nu}(t) dt = \frac{\Gamma^2(\nu)\Gamma(\mu)2^{2\nu-1}}{x^\mu(1+x^2)^\nu}, \quad x > 0. \quad (15)$$

Thus, for $\mu, \nu, m, 2\nu-1+m \in \mathbb{C}_+$ and $x > 0$, we have

$$\begin{aligned} &L\left(t^{m-1+2\nu+\mu-1}\mathfrak{F}_m(h_{\mu,\nu})(t)\right)(x) \\ &= \frac{2^{\frac{1-m}{2}}\Gamma(\nu)}{\Gamma(\frac{m-1}{2} + \nu)} L\left(t^{m-1+2\nu+\mu-1}\mathfrak{F}_1(h_{\mu,\frac{m-1}{2}+\nu})(t)\right)(x) \\ &= \frac{2^{\frac{1-m}{2}}\Gamma(\nu)}{\Gamma(\frac{m-1}{2} + \nu)} \cdot \frac{\Gamma^2(\frac{m-1}{2} + \nu)\Gamma(\mu)2^{m-1+2\nu-1}}{(2\pi)^{\frac{1}{2}}} \cdot \frac{1}{x^\mu(1+x^2)^{\frac{m-1}{2}+\nu}} \\ &= \frac{C(m,\mu,\nu)}{x^\mu(1+x^2)^{\frac{m-1}{2}+\nu}}, \\ \text{with } C(m,\mu,\nu) &:= \frac{2^{-\frac{m}{2}}\Gamma(\nu)\Gamma(\mu)\Gamma(m-1+2\nu)}{\Gamma(\frac{m}{2} + \nu)} = D(m,\mu,\nu)\Gamma(\mu+m-1+2\nu). \end{aligned} \quad (16)$$

Remark 2 Equation (16) is the crux of the proof of the main part of Theorem 11 in [34]:

(i) If $\nu > \frac{1}{2}$ and $\mu \geq \max\{\nu, 1\}$, then $h_{\mu,\nu} \in \Phi_1$. If, additionally, $(\mu; \nu) \neq (1; 1)$, then there exist constants $c_i > 0$, $i = 1, 2$, depending on μ and ν only, such that

$$c_1 \leq (1+t^2)^\nu \cdot \widehat{h}_{\mu,\nu}(t) \leq c_2, \quad t \in \mathbb{R}.$$

(ii) If $\nu \geq 1$, then $h_{\mu,\nu} \in \Phi_1 \iff \mu \geq \nu$.

(iii) If $m \geq 2$, then $h_{\mu,\nu} \in \Phi_m \iff \nu > \frac{1}{2}$ and $\mu \geq \frac{m-1}{2} + \nu$. In this case, there exist two constants $c_i > 0$, $i = 1, 2$, depending on μ, ν and m , and such that

$$c_1 \leq (1+t^2)^{\frac{m-1}{2}+\nu} \cdot \mathfrak{F}_m(h_{\mu,\nu})(t) \leq c_2, \quad t \geq 0.$$

This theorem is related to the positiveness of the function $I_{1,\mu,\nu,2\nu-1}(t)$ for all $t > 0$. Theorems on positiveness of the functions $I_{\delta,\mu,\nu,\alpha}(t)$ are obtained in [36, Theorems 4,5,6] (the well-known cases given before Theorem 4 from [36]).

2.3 Statement of the Problem

We are now able to state our main

Problem 1 Let $\mu > 0$, $\nu > \frac{1}{2}$, $\mu + \nu > 1$. Then $h_{\mu,\nu} \in C(\mathbb{R})$ (see [35, 34]). Let $\varepsilon > 0$ and

$$f_{\mu,\nu,\varepsilon,\beta_1,\beta_2}(x) := \beta_2^\varepsilon h_{\mu,\nu}\left(\frac{x}{\beta_2}\right) - \beta_1^\varepsilon h_{\mu,\nu}\left(\frac{x}{\beta_1}\right), \quad x \in \mathbb{R}. \quad (17)$$

Let $m \in \mathbb{N}$. Show the conditions on (ε, μ, ν) such that, for any $\beta_2 > \beta_1 > 0$, we have

$$f_{\mu,\nu,\varepsilon,\beta_1,\beta_2} \in \Phi_m. \quad (18)$$

Next section details the structure of the solution and offers the exact smoothness of the new covariance resulting from the differences of Buhmann functions.

3 Structure of the solution

Let us start with a general assertion regarding the structure of Problem 1.

Proposition 1 *The following conditions are equivalent:*

1. *Condition (18) is satisfied.*
2. *For any $\beta_2 > \beta_1 > 0$, the function $t \mapsto \beta_2^{\varepsilon+m} \mathfrak{F}_m(h_{\mu,\nu})(\beta_2 t) - \beta_1^{\varepsilon+m} \mathfrak{F}_m(h_{\mu,\nu})(\beta_1 t)$ is nonnegative in interval $(0, \infty)$.*
3. *The function $t^{\varepsilon+m} \mathfrak{F}_m(h_{\mu,\nu})(t) = 2^{\nu-1} \Gamma(\nu) t^{\varepsilon+m} I_{1,\mu, \frac{m-1}{2} + \nu, m-1+2\nu-1}(t)$ increases in the interval $(0, \infty)$.*
4. *The following inequality is true:*

$$\begin{aligned} & (\varepsilon + m) I_{1,\mu, \frac{m-1}{2} + \nu, m-1+2\nu-1}(t) + t I'_{1,\mu, \frac{m-1}{2} + \nu, m-1+2\nu-1}(t) = \\ & (\varepsilon + m) I_{1,\mu, \frac{m-1}{2} + \nu, m-1+2\nu-1}(t) - t^2 I_{1,\mu, \frac{m-1}{2} + \nu+1, m-1+2\nu+1}(t) \geq 0, \quad \forall t > 0 \end{aligned}$$

5. Let $n := \frac{m-1}{2} + \nu$. Then,

$$(\varepsilon + m)\mathfrak{F}_1(h_{\mu,n})(t) - \frac{t^2}{2n}\mathfrak{F}_1(h_{\mu,n+1})(t) \geq 0, \quad \forall t > 0.$$

6. We have

$$L \left(t^{2n+\mu-1} \left((\varepsilon + m)\widehat{h}_{\mu,n}(t) - \frac{t^2}{2n}\widehat{h}_{\mu,n+1}(t) \right) \right) (x) = \\ \Gamma^2(n)\Gamma(\mu)2^{2n-1} \left(\frac{\varepsilon + m}{x^\mu(1+x^2)^n} - \frac{2n}{x^\mu(1+x^2)^{n+1}} \right) \in \mathcal{CM}.$$

7.

$$\frac{\varepsilon - 2\nu + 1 + (\varepsilon + m)x^2}{x^\mu(1+x^2)^{\frac{m-1}{2}+\nu+1}} \in \mathcal{CM} \quad (19)$$

Remark 3 It follows from Hausdorff-Bernstein-Widder theorem (see, for example, [9, 18, 24, 31]) that if $g \in C[0, +\infty)$ and its Laplace transform

$$Lg(x) := \int_0^{+\infty} e^{-xs}g(s) ds$$

converges for all $x > 0$, then $g(s) \geq 0$ for $s \geq 0$ if and only if $Lg \in \mathcal{CM}$.

The proof Proposition 1 is an easy consequence of Remark 1 in concert with the Hausdorff-Bernstein-Widder theorem (see Remark 3), Theorem 1 (statements 1 and 2), and equalities (13) and (15).

Note that, if $\mu, \nu > 0$, then $x^{-\mu}(1+x^2)^{-\nu} \in \mathcal{CM}$ if and only if $\widehat{h}_{\mu,\nu}(t) \geq 0$ for all $t > 0$, if and only if $I_{1,\mu,\nu,2\nu-1}(t) \geq 0$ for all $t > 0$ (see Equations (15) and (14)).

Proposition 1 will now be combined with the following facts:

1. If $\nu \geq 1$, then $x^{-\mu}(1+x^2)^{-\nu} \in \mathcal{CM}$ if and only if $\mu \geq \nu$. The sufficiency of this result can be found in [10], and the necessity has been proved in [16], [33, Lemma 8]).
2. If $0 < \nu < 1$, $\mu \geq 1$, then $x^{-\mu}(1+x^2)^{-\nu} \in \mathcal{CM}$ [16] and [37, Example 5.4], [38, § 4.7, Example 4.7.7].
3. If $\nu > 0$, $\mu \geq 2\nu$, then $x^{-\mu}(1+x^2)^{-\nu} \in \mathcal{CM}$ [2].
4. If $n = 1, 2, 3$, then $(a+x^2)/(x^n(1+x^2)^n) \in \mathcal{CM}$ if and only if $a \geq 1/(2^{n-1}+1)$ [35, § 2].

We can now combine the first three sufficient conditions above to obtain the following assertion: if $\nu > 0$ and $\mu \geq \min\{2\nu; \max\{1, \nu\}\}$, then $x^{-\mu}(1+x^2)^{-\nu} \in \mathcal{CM}$.

The combination of these facts with Proposition 1 has just offered the proof of the following

Theorem 2 *The following assertions are true:*

1. *If (18) is true, then $\varepsilon \geq 2\nu - 1$.*
2. *If $m \in \mathbb{N}$, $\nu > \frac{1}{2}$, $\varepsilon \geq 2\nu - 1$ and $\mu \geq (m-1)/2 + \nu + 3$, then condition (18) is true. If, in addition, $\varepsilon = 2\nu - 1$, then (18) is true if and only if $\mu \geq (m-1)/2 + \nu + 3$.*
3. *Suppose that for some $n = 1, 2, 3$, we have $\varepsilon \geq 2^{1-n}(m + (2\nu - 1)(2^{n-1} + 1))$, $(m-1)/2 + \nu + 1 - n > 0$ and*

$$\mu - n \geq \min \left\{ m - 1 + 2\nu + 2 - 2n; \max \left\{ 1; \frac{m-1}{2} + \nu + 1 - n \right\} \right\}.$$

Then, condition (18) is true.

We now provide a characterization result for the following problem: whether the condition (18) is satisfied for fixed $\beta_2, \beta_1 > 0$ (and not for any $\beta_2 > \beta_1 > 0$ as in the problem 1).

Theorem 3 *Let $\mu > 0$, $\nu > \frac{1}{2}$, $\mu + \nu > 1$ and $m \in \mathbb{N}$. Let $\varepsilon \in \mathbb{R}$, $\beta_2, \beta_1 > 0$ and $a := \beta_2/\beta_1$.*

Then $f_{\mu, \nu, \varepsilon, \beta_1, \beta_2} \in \Phi_m$ if and only if

$$\frac{1}{x^\mu(1+x^2)^{\frac{m-1}{2}+\nu}} - \frac{a^{2\nu-1-\varepsilon}}{x^\mu(1+a^2x^2)^{\frac{m-1}{2}+\nu}} \in \mathcal{CM}. \quad (20)$$

If $f_{\mu, \nu, \varepsilon, \beta_1, \beta_2} \in \Phi_m$ and $\beta_2 > \beta_1 > 0$, then $\varepsilon \geq 2\nu - 1$.

Proof. From Remarks 1 and 3, it follows that the following conditions are equivalent:

1. $f_{\mu, \nu, \varepsilon, \beta_1, \beta_2} \in \Phi_m$.
2. $\beta_2^{\varepsilon+m} \mathfrak{F}_m(h_{\mu, \nu})(\beta_2 t) - \beta_1^{\varepsilon+m} \mathfrak{F}_m(h_{\mu, \nu})(\beta_1 t) \geq 0$ for all $t \in (0, \infty)$.
3. $L \left(t^{m-1+2\nu+\mu-1} \left(\beta_2^{\varepsilon+m} \mathfrak{F}_m(h_{\mu, \nu})(\beta_2 t) - \beta_1^{\varepsilon+m} \mathfrak{F}_m(h_{\mu, \nu})(\beta_1 t) \right) \right) (x) \in \mathcal{CM}$.

Furthermore, we have (see (16)):

$$\beta^{\varepsilon+m} L \left(t^{m-1+2\nu+\mu-1} \mathfrak{F}_m(h_{\mu, \nu})(\beta t) \right) (x) = \frac{\beta^{\varepsilon-2\nu+1}}{x^\mu(1+x^2/\beta^2)^{\frac{m-1}{2}+\nu}}, \quad \beta > 0, x > 0.$$

Suppose that condition (20) is satisfied. Since completely monotone functions are non-negative on $(0, +\infty)$, we have that $1 - a^{2\nu-1-\varepsilon} \geq 0$. If, additionally, $a = \beta_2/\beta_1 > 1$, then $\varepsilon \geq 2\nu - 1$. The proof is completed. \blacksquare

Direct inspection of the proof of Proposition 1 as well as the proof of Theorem 3 shows that the following proposition is true.

Proposition 2 *Let $\mu, \nu, m, 2\nu - 1 + m > 0$ and $\varepsilon \in \mathbb{R}$. Then following conditions are equivalent:*

1. $\frac{\varepsilon - 2\nu + 1 + (\varepsilon + m)x^2}{x^\mu(1 + x^2)^{\frac{m-1}{2}+\nu+1}} \in \mathcal{CM}$.
2. $\frac{1}{x^\mu(1 + x^2)^{\frac{m-1}{2}+\nu}} - \frac{a^{2\nu-1-\varepsilon}}{x^\mu(1 + a^2x^2)^{\frac{m-1}{2}+\nu}} \in \mathcal{CM}, \forall a > 1$.
3. $\frac{1}{x^\mu(1 + x^2)^{\frac{m-1}{2}+\nu}} - \frac{a_n^{2\nu-1-\varepsilon}}{x^\mu(1 + a_n^2x^2)^{\frac{m-1}{2}+\nu}} \in \mathcal{CM}$ for some sequence $a_n > 1, a_n \rightarrow 1$.

If the first condition is satisfied, or if the second condition is satisfied for some $a > 1$, then $\varepsilon \geq 2\nu - 1$.

Proof The crux of the proof is in the following equality, which holds for $a > 1$:

$$\int_1^a \frac{(\varepsilon - 2\nu + 1 + (\varepsilon + m)x^2t^2)t^{2\nu-\varepsilon-2}}{x^\mu(1 + x^2t^2)^{\frac{m-1}{2}+\nu+1}} dt = \frac{1}{x^\mu(1 + x^2)^{\frac{m-1}{2}+\nu}} - \frac{a^{2\nu-1-\varepsilon}}{x^\mu(1 + a^2x^2)^{\frac{m-1}{2}+\nu}}.$$

Since completely monotone functions are closed under scale mixtures, the equality above provides the implication $1 \Rightarrow 2$. Assertion $2 \Rightarrow 3$ is obvious. To prove assertion $3 \Rightarrow 1$, it is necessary to take in the last equation $a = a_n$, divide both sides by $a_n - 1$ and take the limit as $n \rightarrow \infty$. Finally, we make use of the well-known fact: if a sequence of completely monotone functions converges pointwise on $(0, +\infty)$, then the limit function is also completely monotone. \blacksquare

We conclude this section detailing the exact smoothness of the differences of Buhmann functions.

Theorem 4 *Let $\nu \in \mathbb{N}$, $\mu > 0$, $\varepsilon \in \mathbb{R}$, $\beta_2, \beta_1 > 0$, and $\beta_2 \neq \beta_1$. Let $q := \min(\beta_1, \beta_2)$. Then,*

1. *If $\varepsilon \neq 2\nu - 1$, then $f_{\mu,\nu,\varepsilon,\beta_1,\beta_2} \in C^{2\nu-2}(-q, q)$, and $f_{\mu,\nu,\varepsilon,\beta_1,\beta_2} \notin C^{2\nu-1}(-q, q)$.*
2. *If $\varepsilon = 2\nu - 1$, $\mu \notin \{1, 2\}$, then $f_{\mu,\nu,\varepsilon,\beta_1,\beta_2} \in C^{2\nu}(-q, q)$, and $f_{\mu,\nu,\varepsilon,\beta_1,\beta_2} \notin C^{2\nu+1}(-q, q)$.*

3. If $\varepsilon = 2\nu - 1$, $\mu = 1$ or $\mu = 2$, then $f_{\mu,\nu,\varepsilon,\beta_1,\beta_2}$ is a even polynomial of degree at most $\mu + 2\nu - 2$ on $[-q, q]$, and therefore $f_{\mu,\nu,\varepsilon,\beta_1,\beta_2} \in C^\infty(-q, q)$.

Proof. If $\mu, \nu > 0$, then arguments in [35, Equality (40)] show that

$$h_{\mu,\nu}(x) = (1-x)^{\mu+\nu-1} \int_0^1 t^{\mu-1} (1-t)^{\nu-1} (1-t+(1+t)x)^{\nu-1} dt, \quad x \in (0, 1). \quad (21)$$

Let $\nu \in \mathbb{N}$. Then from Proposition 1 in [36] we have that $h_{\mu,\nu} \in C^{2\nu-2}(-1, 1)$ and $h_{\mu,\nu} \notin C^{2\nu-1}(-1, 1)$. Thus, from (21) it follows that

$$\begin{aligned} h_{\mu,\nu}(x) &= \sum_{k=0}^{\infty} a_k(\mu, \nu) x^{2k} + |x|^{2\nu-1} \sum_{k=0}^{\infty} b_k(\mu, \nu) x^{2k}, \quad |x| < 1, \\ b_0(\mu, \nu) &\neq 0, \quad b_1(\mu, \nu) = \frac{h_{\mu,\nu}^{(2\nu+1)}(+0)}{(2\nu+1)!}. \end{aligned} \quad (22)$$

From (12) we have that $h'_{\mu,\nu}(x) = -2(\nu-1)x h_{\mu,\nu-1}(x)$ for $\nu \geq 2$, $x > 0$, and $h_{\mu,\nu}^{(k+1)}(x) = -2(\nu-1)(x h_{\mu,\nu-1}^{(k)}(x) + k h_{\mu,\nu-1}^{(k-1)}(x))$ for $k \geq 1$, $0 < x < 1$. From the last equation it follows that $h_{\mu,\nu}^{(k+1)}(+0) = -2(\nu-1)k h_{\mu,\nu-1}^{(k-1)}(+0)$ for $\nu \geq 2$, $k \geq 1$, and for $\nu \geq 2$, $k \geq 2\nu-3$, we have (for convenience, we consider that $0!! := 1$ and $(-1)!! := 1$)

$$h_{\mu,\nu}^{(k+1)}(+0) = (-2)^{\nu-1} (\nu-1)! \cdot \frac{k!!}{(k-2\nu+2)!!} h_{\mu,1}^{(k-2\nu+3)}(+0). \quad (23)$$

The equality (23) is true for $\nu = 1$, $k \geq -1$. It's obvious that $h_{\mu,1}(x) = (1-|x|)_+^\mu / \mu$ and $h_{\mu,1}^{(p)}(+0) = (-1)^p \Gamma(\mu) / \Gamma(\mu-p+1)$, $p \in \mathbb{Z}_+$. Therefore, for $k \geq \nu-1$, with $\nu \in \mathbb{N}$, we have

$$\begin{aligned} h_{\mu,\nu}^{(2k+1)}(+0) &= (-4)^{\nu-1} (\nu-1)! \cdot \frac{k!}{(k-\nu+1)!} \cdot h_{\mu,1}^{(2k-2\nu+3)}(+0) \\ &= -(-4)^{\nu-1} (\nu-1)! \cdot \frac{k!}{(k-\nu+1)!} \cdot \frac{\Gamma(\mu)}{\Gamma(\mu-2k+2\nu-2)}; \\ h_{\mu,\nu}^{(2\nu+1)}(+0) &= -(-4)^{\nu-1} (\nu-1)! \nu! (\mu-1)(\mu-2). \end{aligned}$$

On the other hand, (22) shows that, for $|x| < q = \min(\beta_1, \beta_2)$,

$$\begin{aligned} f_{\mu,\nu,\varepsilon,\beta_1,\beta_2}(x) &= \sum_{k=0}^{\infty} a_k(\mu, \nu) (\beta_2^{\varepsilon-2k} - \beta_1^{\varepsilon-2k}) x^{2k} + \\ &+ |x|^{2\nu-1} \sum_{k=0}^{\infty} b_k(\mu, \nu) (\beta_2^{\varepsilon-2\nu+1-2k} - \beta_1^{\varepsilon-2\nu+1-2k}) x^{2k}. \end{aligned}$$

Assertions 1. and 2. are thus proved.

If, in addition, $\mu \in \mathbb{N}$, then $h_{\mu,\nu}$ is a polynomial of degree $\mu + 2\nu - 2$ on the compact interval $[0, 1]$. If $\mu = 1$ or $\mu = 2$, then in the second sum of (22), the terms with $k \geq 1$ vanish. Therefore, if $\varepsilon = 2\nu - 1$, $\mu = 1$ or $\mu = 2$, then $f_{\mu,\nu,\varepsilon,\beta_1,\beta_2}$ on the interval $[-q, q]$ is a even polynomial of degree $\leq \mu + 2\nu - 2$. Assertion 3. is proved. The proof is completed. ■

4 Connections with previous literature

The k -fold application of the Montée operator to the Askey function $\mathcal{A}_\mu(t) = (1 - t)_+^\mu$, for $\mu \geq (m + 1)/2$, results in Wendland functions $\psi_{\mu,k}$ as described in previous section. Table 1 depicts the role of the mapping $f_{\mu,\nu,\varepsilon,\beta_1,\beta_2}$ as defined through Equation (17). In particular, we consider $f_{\mu,k+1,2k+1,\beta_1,\beta_2}$, for $k = 0, 1, 2$. Expressions of the corresponding Wendland functions are reported in the second column of the same table. According to Theorem 2, $f_{\mu,k+1,2k+1,\beta_1,\beta_2} \in \Phi_m$ if and only if $\mu \geq (m + 7)/2 + k$. The third and fourth column allow to describe the action of the mapping $f_{\mu,k+1,2k+1,\beta_1,\beta_2}$. When $\mu \notin \{1, 2\}$, one can clearly appreciate the increase in terms of differentiability at the origin.

k	$\psi_{\mu,k}$	D_{before}	D_{after}
0	$(1 - x)_+^\mu$	$C(\{0\})$	$C^2(\{0\})$
1	$(1 - x)_+^{\mu+1}(1 + (\mu + 1)x)$	$C^2(\{0\})$	$C^4(\{0\})$
2	$(1 - x)_+^{\mu+2}(1 + (\mu + 2)x + \frac{1}{3}((\mu + 2)^2 - 1))$	$C^4(\{0\})$	$C^6(\{0\})$

Table 1: Examples of the Wendland functions $\psi_{\mu,k}$ for $k = 0, 1, 2$ and $\mu \notin \{1, 2\}$. For all cases, $x \geq 0$. The first column reports the values of k , and the corresponding expression in the second column reports the analytic expression of $\psi_{\mu,k}$. In the third column, D_{before} depicts the differentiability of $\psi_{\mu,k}$. In the last column, D_{after} stays for the differentiability at the origin of $f_{\mu,k+1,2k+1,\beta_1,\beta_2}$. Observe that when $\mu = 1$ or $\mu = 2$, then $D_{\text{after}} = \infty$.




Figure 1: Continuous Lines: $f_{(d+1)/2+k+3,k+1,2k+1,0.75,1}$, defined according to Equation (17) for $k = 0, 1, 2$ (from left to right). Dashed and Dashed-dot Lines report Wendland functions with $k = 0, 1, 2$ and compact supports $\beta_1 = 0.75$ and $\beta_2 = 1$. All the functions are normalized with their value at the origin.

The Wendland radial basis functions are piecewise polynomial compactly supported reproducing kernels in Hilbert spaces which are normequivalent to Sobolev spaces. But they only cover the Sobolev spaces $H^{d/2+k+1/2}(\mathbb{R}^d)$, when $k \in \mathbb{N}$. Motivated by this fact, Robert Schaback [20] covered the case of the integer order spaces in even dimensions. Namely, he derived the missing Wendland functions working for half-integer k and even dimensions, reproducing integer-order Sobolev spaces in even dimensions, and showing that they turn out to have two additional non-polynomial terms: a logarithm and a square root.

Other walks through dimensions have been recently proposed by [17] through the Generalized Askey functions $\varphi_{n,k,m} : [0, \infty) \rightarrow \mathbb{R}$ defined through

$$\varphi_{n,k,m}(t) = t^{k-n}(1-t)_+^{n+m+1}F(n-k, n+1, n+m+2, 1-1/t), \quad t \geq 0,$$

with F being the Gauss hypergeometric function. The parameters (n, m, k) are then shown to be crucial in order to determine when Φ_m (see their Proposition 2.3).

To our knowledge, the only case of compactly supported correlation functions which is not covered by this work is the case of the Euclid's hat ([19, 12]), which is the self-convolution of an indicator function supported on the unit ball in \mathbb{R}^d . As noted by [20], while Euclid's hat is not differentiable and Wus functions have zeros in their Fourier transform, Wendlands functions have no such drawbacks. They are polynomials on $[0, 1]$ and yield positive definite $2k$ -times differentiable radial basis functions on \mathbb{R}^d . Given these properties, their polynomial degree $\lfloor d/2 \rfloor + 3k + 1$ is minimal.

The proof of Theorem 2 highlights explicit connections with previous literature devoted to (sub) classes of completely monotone functions. A function $f : [0, \infty) \rightarrow \mathbb{R}$ is called Logarithmically completely monotonic on $(0, \infty)$, and denoted $f \in \mathcal{L}(0, \infty)$ if and only if it is infinitely often differentiable on $(0, \infty)$ and

$$(-1)^n [\log f(x)]^{(n)} \geq 0, \quad x \geq 0.$$

By well known results (see [3], with the references therein) $f \in \mathcal{L} \Leftrightarrow f^\alpha \in \mathcal{CM} \Leftrightarrow (f)^{1/n} \in \mathcal{CM}$, for all $\alpha > 0$ and $n \in \mathbb{N}$.

Berg, Porcu and Mateu [3] introduced the so called *Auxiliary* family

$$f_{\alpha,\beta}(x) = \frac{1}{x^\alpha(1+x^\beta)}, \quad x > 0,$$

with α, β positive parameters. They show that $f_{\alpha,\beta} \in \mathcal{L}$ for $\alpha \geq 0$ and $0 \leq \beta \leq 1$. Additionally, $f_{\alpha,2} \in \mathcal{L}$ if and only if $\alpha \geq 2$. This result is one of the crux for describing the complete monotonicity of the Dagum family. Known cases, when $x^{-\mu}(1+x^2)^{-\nu} \notin \mathcal{CM}$ (if $\mu, \nu > 0$, then this is equivalent to the inequality $I_{1,\mu,\nu,2\nu-1} \geq 0$ is not true for all $t > 0$): 1) $\mu < 0$ or $\mu = 0, \nu \neq 0$ (it is obvious); 2) $\mu < \nu$; 3) $0 < \mu = \nu < 1$. Proof of the latter two cases, see, for example, [16], [33, Lemma 8] and [36, Theorem 5].

Acknowledgement

We are grateful to Professor R. Furrer for helpful remarks through the preparation of the manuscript. Research work of Moreno Bevilacqua was partially supported by grant FONDECYT 11121408 from Chilean government. Research work of Emilio Porcu was partially supported by grant FONDECYT 1130647 from Chilean government.

References

- [1] Askey, R. Radial characteristic functions, Technical Report, University of Wisconsin, 1973.
- [2] Askey, R. and Pollard, H. Some absolutely monotonic and completely monotonic functions. SIAM J. Math. Anal. , 1974, 5, 58–63.

- [3] Berg, C., Porcu, E. and Mateu, J. The Dagum family of completely monotonic functions. *Bernoulli*, 2008, 14 (4), 1134–1149.
- [4] Bevilacqua, M., Faouzi, T., Furrer, R. and Porcu, E. Estimation and Prediction using generalized Wendland Covariance Functions under fixed domain asymptotics, Technical Report, University of Valparaiso. To be submitted for publication.
- [5] Buhmann, M.D. A new class of radial basis functions with compact support. *Mathematics of Computation*, 2000, 70 (233), 307–318.
- [6] Daley, D.J. and Porcu, E. Dimension walks through Schoenberg spectral measures. *Proceedings of the American Mathematical Society*, 2014, 142 (5), 1813–1824.
- [7] Daley, D.J., Porcu, E. and Bevilacqua, M. Classes of compactly supported covariance functions for multivariate random fields. *Stochastic Environmental Research Risk Assessment*, 2015, 29 (4), 1249–1263.
- [8] Fassahuer, G. Positive Definite Kernels: Past, Present and Future. *Dolomites Research Notes on Approximation*, 2011, 4, 21–63.
- [9] Feller, W., Completely monotone functions and sequences. *Trans. Amer. Math. Soc.*, 1939, **5**, 662–774.
- [10] Fields, J.L. and Ismail, M.E.H. On the positivity of some ${}_1F_2$'s. *SIAM J. Math. Anal.*, 1975, 6, 551–559.
- [11] Furrer, R., Genton, M. and Nychka, D. Covariance Tapering for Interpolation of Large Spatial Datasets, *Journal of Computational and Graphical Statistics*, 2006, 15 (3), 502–523.
- [12] Gneiting, T. Radial positive definite functions generated by Euclid's hat. *J. Multivar. Anal.*, 1999, 69(1), 88–119.
- [13] Gneiting, T. Compactly supported correlation functions. *Journal of Multivariate Analysis*, 2002, 83, 493–508.
- [14] Matheron, G.. *Les variables régionalisées et leur estimation*, 1965, Masson Paris.

- [15] Micchelli, C.A. Interpolation of scattered data: distance matrices and conditionally positive definite functions. *Constructive Approximation*, 1986, 2, 11–22.
- [16] Moak, D. Completely monotonic functions of the form $s^{-b}(s^2 + 1)^{-a}$. *Rocky Mountain J. Mathematics*, 1987, 17, 719–725.
- [17] Porcu, E. and Zastavnyi, V.P. Generalized Askey Functions and their walks through dimensions. *Exposithiones Mathematicae*, 2014, 32 (2), 190–198.
- [18] Sasvari, Z. *Multivariate Characteristic and Correlation Functions*, De Gruyter, Berlin, 2013.
- [19] Schaback, R.: Creating surfaces from scattered data using radial basis functions. In: Lyche, T., Daehlen, M., Schumaker, L.L. (eds.) *Mathematical Methods for Curves and Surfaces*, 1996, 477–496. Vanderbilt University Press, Nashville.
- [20] Schaback, R. The Missing Wendland Functions. *Advances in Computational Mathematics*, 2010, 43, 67–81.
- [21] Schaback, R. and Wendland, H. Kernel Techniques: From Machine Learning to Meshless Methods, *Acta Numerica*, 2006, 1–97.
- [22] Schaback, R. and Wu, Z. Operators on radial functions, *Journal of Computational and Applied Mathematics*, 1996, 73, 257–270.
- [23] Scheuerer, M., Schaback, R. and Schlather, M. Interpolation of spatial data – A stochastic or a deterministic problem? *European Journal of Applied Mathematics*, 2013, 24 (4), 601–629.
- [24] Schilling, R.L., Song, R. Vondraček, Z.. *Bernstein functions*, Berlin, Boston, 2010. De Gruyter.
- [25] Stein, M.L. *Interpolation of Spatial Data. Some Theory of Kriging*, 1999, Springer Verlag, New York.
- [26] Trigub, R. M., Positive-definite compactly supported radial functions of polynomial form and maximal smoothness. *Mat. Fiz. Anal. Geom.* (2002), 9 (3), 394–400. (<http://www.mathnet.ru/links/7b3900405c11b813f29a918b05dc26a5/jmag301.pdf>)

- [27] Trigub R.M, Belinsky, E.S. Fourier Analysis and Approximation of Functions, Boston, Dordrecht, London: Kluwer-Springer, 2004.
- [28] G. N. Watson, A Treatise on the Theory of Bessel Functions, Cambridge University, Cambridge (1944).
- [29] Wackernagel, H. Multivariate Geostatistics: An Introduction with Applications, Springer Science & Business Media, 2003.
- [30] Wendland, H. Piecewise polynomial, positive definite and compactly supported radial functions of minimal degree. *Advances in Comp. Math.* 1995, 4, 389–396.
- [31] Widder, D. V., The Laplace Transform. Princeton: Princeton University Press, 1946.
- [32] Wu, Z. Compactly supported positive definite radial functions, *Advances in Computational Mathematics*, 1995, 4, 283–292.
- [33] Zastavnyi, V. P. On positive definiteness of some functions. *Journal of Multivariate Analysis*, 2000, 73, 55–81.
- [34] Zastavnyi, V.P. Positive-definite radial functions and splines. *Doklady Mathematics*, 2002, 66 (2), 213-216.
- [35] Zastavnyi, V.P., and Trigub, R.M. Positive-definite splines of a special form. *Sbornik: Mathematics*, 2002, 193 (12), 1771–1800.
- [36] Zastavnyi, V.P. On some properties of Buhmann functions. *Ukrainian Mathematical Journal*, 2006, 58 (8), 1184–1208.
- [37] Zastavnyi V. P. On Exponential Type Entire Functions without Zeros in the Open Lower Half-plane. *Ukranian Math. Bull*, 2006, 3, 395–422. (<http://arxiv.org/abs/1606.08255>)
- [38] Zastavnyi, V.P. Problems Related to Positive Definite Functions. In *Positive Definite Functions: from Schoenberg to Space-Time Challenges*. J. Mateu and E. Porcu Eds. – Editorial Universitat Jaume I, 2008, 63–114.
- [39] zu Castell, W. Recurrence relations for radial positive definite functions. *J. Math. Anal. Appl.*, 2002, 271, 108–123.