

# ON STABLE CMC HYPERSURFACES WITH FREE-BOUNDARY IN A EUCLIDEAN BALL

EZEQUIEL BARBOSA

ABSTRACT. In this note, we observe that if  $B$  is a ball in a Euclidean space with dimension  $n$ ,  $n \geq 3$ , then a stable CMC hypersurface  $\Sigma$  with free boundary in  $B$  satisfies

$$nA \leq L \leq nA \left( \frac{1 + \sqrt{1 + 4(n+1)H^2}}{2} \right),$$

where  $L$ ,  $A$  and  $H$  denote the length of  $\partial\Sigma$ , the area of  $\Sigma$  and the mean curvature of  $\Sigma$ , respectively. Consequently, if the boundary  $\partial\Sigma$  is embedded then  $\Sigma$  must be totally geodesic or starshaped with respect to the center of the ball. This result is an improvement of a theorem proved by A. Ros and E. Vergasta [R-V]. In particular, if  $n = 3$ , the only stable CMC surfaces with free boundary in  $B$  are the totally geodesic disks or the spherical caps. This last result was proved very recently by I. Nunes [N] using an extended stability result and a modified Hersch type balancing argument to get a better control on the genus. We don't use that modified Hersch type argument. However, we use a Nunes type Stability Lemma and a crucial result due to A. Ros and E. Vergasta.

## 1. INTRODUCTION

Given a smooth compact and convex domain  $B$  in  $\mathbb{R}^{n+1}$ , denote by  $\partial B$  and  $\text{int } B$  the boundary and the interior of  $B$ , respectively. A CMC free-boundary hypersurface in  $B$  is a constant mean curvature hypersurface  $\Sigma \subset B$  meeting  $\partial B$  orthogonally along  $\partial\Sigma$ . That kind of hypersurfaces are solutions for the problem of finding critical points of the area functional among all compact hypersurfaces  $\Sigma \subset B$  with  $\partial\Sigma \subset \partial B$  which divides  $B$  into two subsets of prescribed volumes. If a CMC free-boundary hypersurface  $\Sigma \subset B$  has nonnegative second variation of area for all preserving volume variations we name it as a CMC free-boundary stable hypersurface. For more details about CMC free-boundary hypersurfaces, see the following references and references therein: [N], [R], [R-V], [So].

In [R-V], Ros and Vergasta studied stable CMC hypersurfaces with free boundary when  $B$  is a ball and proved the following result. Denote by  $L$  the length of the boundary  $\partial\Sigma$  and by  $A$  the area of  $\Sigma$ .

**Theorem 1.1** (Ros-Vergasta [R-V]). *Let  $B \subset \mathbb{R}^n$ ,  $n \geq 3$ , be a closed ball. Let  $\Sigma \subset B$  be a CMC free-boundary stable hypersurface with embedded boundary in  $B$ . If  $L \geq nA$  then  $\Sigma$  is totally geodesic or starshaped with respect to the center of the ball.*

In order to improve the result above, we use a Nunes type Stability Lemma (see Lemma 2.1) to prove that always  $L \geq nA$ . More precisely, we obtain the following result.

---

The author partially supported by CNPq-Brazil.

**Theorem 1.2.** *Let  $B \subset \mathbb{R}^n$ ,  $n \geq 3$ , be a closed ball. If  $\Sigma \subset B$  is a CMC free-boundary stable hypersurface in  $B$ , then*

$$nA \leq L \leq nA \left( \frac{1 + \sqrt{1 + 4(n+1)H^2}}{2} \right).$$

*In particular, if  $\partial\Sigma$  is embedded, then  $\Sigma$  is totally geodesic or starshaped with respect to the center of the ball.*

As a direct consequence, we obtain the following corollary.

**Corollary 1.1.** *Let  $B \subset \mathbb{R}^n$ ,  $n \geq 3$ , be a closed ball. If  $\Sigma \subset B$  is a CMC stable hypersurface with embedded free-boundary in  $B$  and  $0 \in \Sigma$  then  $\Sigma$  is totally geodesic.*

Now, noting that free-boundary surfaces must have embedded boundary (see Theorem 11 in [R-V]), we obtain a complete topological classification.

**Corollary 1.2.** *Let  $B \subset \mathbb{R}^3$  be a closed ball. If  $\Sigma \subset B$  is a CMC stable surface with free-boundary then  $\Sigma$  is a totally geodesic disk or a spherical cap.*

It's worthy to mention that the Corollary 1.2 was proved recently by I. Nunes [N] using a powerful stability result and a modified Hersch type balancing argument to get a better control on the genus and on the number of connected components of the boundary of the surface. In fact, I. Nunes proved a more general result which, joint with Theorem 11 in [R-V], gives us the result above as a corollary.

**Theorem 1.3** (I. Nunes [N]). *Let  $\Omega \subset \mathbb{R}^3$  be a smooth compact convex domain. Suppose that the second fundamental form  $\Pi^{\partial\Omega}$  of  $\partial\Omega$  satisfies the pinching condition*

$$k h \leq \Pi^{\partial\Omega} \leq (3/2)k h.$$

*for some constant  $k > 0$ , where  $h$  denotes the induced metric on  $\partial\Omega$ . If  $\Sigma \subset \Omega$  is an immersed orientable compact stable CMC surface with free boundary, then  $\Sigma$  has genus zero and  $\Sigma$  has at most two connected components.*

In order to prove Theorem 1.2, we apply the same idea as that applied by I. Nunes in the proof of the main result for the free-boundary surfaces case in [N]. I. Nunes showed that the stability of a free-boundary CMC surface implies that the quadratic form given by the second variation of area is nonnegative for all functions  $f$  such that  $f = 0$  on  $\partial\Sigma$  regardless of whether it satisfies  $\int_{\Sigma} f dvol_{\Sigma} = 0$  or not. That is what we are calling Nunes Stability Lemma. Then I. Nunes was able to apply a modified Hersch type balancing argument to obtain a better control on the genus of  $\Sigma$ . We use that idea for high dimension combined with some Ros-Vergasta results.

## 2. NUNES TYPE STABILITY LEMMA

Let  $B$  be a compact convex domain in  $\mathbb{R}^{n+1}$ . Let  $\varphi : \Sigma^n \rightarrow B$  be an immersion of a smooth orientable manifold  $\Sigma$  with boundary  $\partial\Sigma$  such that  $\varphi(\partial\Sigma) = \varphi(\Sigma) \cap \partial B$ . Let's denote the unit normal vector of the hypersurface  $\Sigma$  by  $N$ . The immersion  $\varphi$  is called free boundary if  $\varphi(\Sigma)$  meets  $\partial B$  orthogonally. The second fundamental form  $A$  of  $\Sigma$  is the endomorphism  $A(X) = -\nabla_X N$ , where  $X \in T\Sigma$ . The mean curvature of  $\Sigma$  is then given by  $H = \frac{1}{n} \text{Tr}A$ .

If we consider a smooth variation  $\phi : \Sigma \times [0, \varepsilon) \rightarrow B$  that preserves  $\partial B$  and such that  $\phi(\cdot, 0) = \varphi(\cdot)$  then it natural to consider the following two functions:

$$A(t) = \int_{\Sigma} dvol_{\Sigma_t} \quad \text{and} \quad V(t) = \int_{\Sigma \times [0, t]} \phi^* dvol_{\mathbb{R}^{n+1}}.$$

The variation  $\phi$  is called volume preserving if  $V(t) \equiv 0$ . Let  $f$  be the function defined by  $f = \langle \frac{\partial}{\partial t} \phi(x, 0), N(x) \rangle$  where  $x \in \Sigma$  then the first variation formula yields:

$$A'(0) = -n \int_{\Sigma} H f dvol_{\Sigma} + \int_{\partial \Sigma} f \langle N, \nu \rangle ds \quad \text{and} \quad V'(0) = \int_{\Sigma} f dvol_{\Sigma}.$$

It follows that CMC hypersurfaces with free boundary are critical points for the area functional  $A(t)$  when restricted to volume preserving variations. The converse is also true, see [R-V]. If  $A''(0) \geq 0$  is nonnegative for every volume preserving variation then the immersion  $\phi$  is called Stable CMC. It can be shown that this is equivalent to have for every  $f \in C^\infty(\Sigma)$  with  $\int_{\Sigma} f dvol_{\Sigma} = 0$  that

$$I(f, f) = \int_{\Sigma} |\nabla f|^2 - |A_{\Sigma}|^2 f^2 dvol_{\Sigma} - \int_{\partial \Sigma} \Pi(N, N) f^2 ds \geq 0$$

**Lemma 2.1** (Nunes Type Stability Lemma). *Let  $\Sigma$  be an immersed stable hypersurface with constant mean curvature with free boundary in  $B$ . If  $f \in C^\infty(\Sigma)$  is such that  $f(x) = 0$  for every  $x \in \partial \Sigma$  then*

$$I(f, f) = \int_{\Sigma} |\nabla f|^2 - |A_{\Sigma}|^2 f^2 dvol_{\Sigma} \geq \frac{1}{n+1} \left( \frac{\int_{\Sigma} f dvol_{\Sigma}}{A} \right)^2 \int_{\partial \Sigma} \Pi(N, N) ds.$$

In particular, If  $f \in C^\infty(\Sigma)$  is such that  $f(x) = 0$  for every  $x \in \partial \Sigma$  then

$$I(f, f) = \int_{\Sigma} |\nabla f|^2 - |A_{\Sigma}|^2 f^2 dvol_{\Sigma} \geq 0.$$

*Proof.* Let  $f_i$  be the function  $f_i = \langle e_i, N \rangle$  where  $\{e_i\}$  is the canonical orthonormal basis of  $\mathbb{R}^{n+1}$ . A simple computation yields:

$$(2.1) \quad \Delta f_i + |A_{\Sigma}|^2 f_i = 0.$$

Plugging these functions on the quadratic form  $I$  we have:

$$\begin{aligned} \sum_{i=1}^{n+1} I(f_i, f_i) &= \int_{\Sigma} \sum_{i=1}^{n+1} |\nabla f_i|^2 - |A_{\Sigma}|^2 f_i^2 dvol_{\Sigma} - \int_{\partial \Sigma} \sum_{i=1}^{n+1} \Pi(N, N) f_i^2 ds \\ &= - \sum_{i=1}^{n+1} \int_{\Sigma} f_i \Delta f_i + |A_{\Sigma}|^2 f_i dvol_{\Sigma} + \int_{\partial \Sigma} \sum_{i=1}^{n+1} f_i \frac{\partial f_i}{\partial \nu} - \Pi(N, N) ds \\ &= \int_{\partial \Sigma} \frac{1}{2} \frac{\partial}{\partial \nu} \left( \sum_{i=1}^{n+1} f_i^2 \right) ds - \int_{\partial \Sigma} \Pi(N, N) ds = - \int_{\partial \Sigma} \Pi(N, N) ds. \end{aligned}$$

We have used that  $\sum_{i=1}^{n+1} f_i^2 = |N|^2 = 1$ . It follows that, given a function  $f$  such that  $f = 0$  on  $\partial \Sigma$ , at least one of the  $f_i$  have the property that

$$I(f_i, f_i) \leq -\frac{1}{n+1} \int_{\partial \Sigma} \Pi(N, N) ds < 0 \quad \text{and} \quad f_i \neq f.$$

In fact, if for each  $f_i$  we have  $I(f_i, f_i) > -\frac{1}{n+1} \int_{\partial\Sigma} \Pi(N, N) ds$  or  $f_i = f$  then we obtain that

$$\sum_{i=1}^{n+1} I(f_i, f_i) > -\frac{m}{n+1} \int_{\partial\Sigma} \Pi(N, N) ds$$

for some positive integer  $m \leq n+1$ , since when  $f_i = f$  we obtain  $I(f_i, f_i) = 0$ . This gives us the contradiction

$$-\int_{\partial\Sigma} \Pi(N, N) ds = \sum_{i=1}^{n+1} I(f_i, f_i) > -\frac{m}{n+1} \int_{\partial\Sigma} \Pi(N, N) ds.$$

Hence, let  $f_i$  be the function satisfying that condition. Note that, because of the stability of  $\Sigma$ , we have that  $\int_{\Sigma} f_i dvol_{\Sigma} \neq 0$ . Assume that  $\int_{\Sigma} f dvol_{\Sigma} \neq 0$ . Now, consider the function  $\bar{f} = cf$ , where

$$c = \frac{\int_{\Sigma} f_i dvol_{\Sigma}}{\int_{\Sigma} f dvol_{\Sigma}}.$$

We have  $\int_{\Sigma} (\bar{f} - f_i) dvol_{\Sigma} = 0$ . Using (2.1) and that  $\bar{f} = 0$  at  $\partial\Sigma$  we have

$$0 \leq I(\bar{f} - f_i, \bar{f} - f_i) = I(\bar{f}, \bar{f}) - 2I(\bar{f}, f_i) + I(f_i, f_i) \leq I(\bar{f}, \bar{f}) - \frac{1}{n+1} \int_{\partial\Sigma} \Pi(N, N) ds.$$

This implies that

$$I(f, f) \geq \left( \frac{\int_{\Sigma} f dvol_{\Sigma}}{\int_{\Sigma} f_i dvol_{\Sigma}} \right)^2 \frac{1}{n+1} \int_{\partial\Sigma} \Pi(N, N) ds.$$

It follows from Holder's inequality and  $\sum_{i=1}^{n+1} f_i^2 = |N|^2 = 1$  that

$$\left( \int_{\Sigma} f_i dvol_{\Sigma} \right)^2 = \left| \int_{\Sigma} f_i dvol_{\Sigma} \right|^2 \leq \left( \int_{\Sigma} |f_i| dvol_{\Sigma} \right)^2 \leq A^2.$$

This finishes the proof.  $\square$

### 3. PROOF OF THEOREM 1.2

*Proof.* Assume that  $\Sigma$  is a stable free-boundary hypersurface in  $B$ . Consider then the support function  $u = \langle \psi, N \rangle$  of  $\Sigma$ , where  $\psi$  is the immersion of  $\Sigma$  in  $B$ . It satisfies the following

$$\begin{cases} \Delta u + |\sigma|^2 u = -nH & \text{on } \Sigma \\ u = 0 & \text{on } \partial\Sigma \end{cases}$$

Moreover, taking the diverge of the tangent component  $\psi - uN$  of  $\psi$  is given by

$$\operatorname{div}(\psi - uN) = n + nHu.$$

It follows from the Divergence Theorem that

$$(3.2) \quad L = n \left( A + \int_{\Sigma} Hudvol_{\Sigma} \right).$$

Since  $u = 0$  on  $\partial\Sigma$ , it follows from the stability of  $\Sigma$  and Nunes Stability Lemma that

$$nH \int_{\Sigma} u dvol_{\Sigma} = \int_{\Sigma} |\nabla_{\Sigma} u|^2 - |A_{\Sigma}|^2 u^2 dvol_{\Sigma} \geq 0.$$

Note that, if  $H = 0$ , then  $L = nA$ . Assume that  $H \neq 0$ . First, as was done by Ros-Vergasta in [R-V], we will first prove that either  $u \geq 0$  or  $u \leq 0$  on  $\Sigma$ . Suppose, by

contradiction, that  $u$  changes sign. Consider  $\Sigma^+$  (resp.  $\Sigma^-$ ) the subset of  $\Sigma$  where  $u$  is positive (resp. negative) and define  $u^+, u^- \in H^1(\Sigma)$  by

$$u^+(p) = \begin{cases} u(p) & \text{if } p \in \Sigma^+ \\ 0 & \text{if } p \in \Sigma \setminus \Sigma^+ \end{cases} \quad \text{and} \quad u^-(p) = \begin{cases} u(p) & \text{if } p \in \Sigma^- \\ 0 & \text{if } p \in \Sigma \setminus \Sigma^- \end{cases}$$

A direct computation gives

$$I(u^-, u^-) = nH \int_{\Sigma} u^- dvol_{\Sigma}$$

and

$$I(u^+, u^+) = nH \int_{\Sigma} u^+ dvol_{\Sigma}.$$

Now we define  $\tilde{u} = u^+ + au^-$ , where  $a$  is a positive constant such that  $\int_{\Sigma} \tilde{u} dvol_{\Sigma} = 0$ . It follows that  $\tilde{u}$  is not identically null and

$$I(\tilde{u}, \tilde{u}) = -naH \int_{\Sigma} u dvol_{\Sigma}.$$

As in Ros-Vergasta [R-V], pag. 29, we obtain that either  $u \geq 0$  or  $u \leq 0$  on  $\Sigma$ . We can choose the orientation on  $\Sigma$  such that  $u \geq 0$ . Since  $H \neq 0$  and  $\int_{\Sigma} HudA \geq 0$ , we get that  $H > 0$ . Therefore,  $u$  satisfies:  $u \geq 0$ ,  $u = 0$  on  $\partial\Sigma$  and  $\Delta u = |\sigma|^2 u - nH < 0$ . By the maximum principle for subharmonic functions we obtain that  $u$  is strictly positive on  $int\Sigma$ . This gives us that  $\int_{\Sigma} u dvol_{\Sigma} \neq 0$ . It follows from the Nunes Stability Lemma that

$$\begin{aligned} \int_{\Sigma} |\nabla_{\Sigma} u|^2 - |A_{\Sigma}|^2 u^2 dvol_{\Sigma} &\geq \frac{1}{n+1} \left( \frac{\int_{\Sigma} u dvol_{\Sigma}}{A} \right)^2 \int_{\partial\Sigma} \Pi(N, N) ds \\ &= \frac{L}{n+1} \left( \frac{\int_{\Sigma} u dvol_{\Sigma}}{A} \right)^2. \end{aligned}$$

Hence, we obtain

$$(3.3) \quad n \int_{\Sigma} Hu dvol_{\Sigma} \geq \frac{L}{n+1} \left( \frac{\int_{\Sigma} u dvol_{\Sigma}}{A} \right)^2,$$

since

$$n \int_{\Sigma} Hu dvol_{\Sigma} = \int_{\Sigma} |\nabla_{\Sigma} u|^2 - |A_{\Sigma}|^2 u^2 dvol_{\Sigma}.$$

From (3.2) we have that

$$\frac{\int_{\Sigma} u dvol_{\Sigma}}{A} = \frac{L - nA}{nHA}.$$

Then, from (3.2) and (3.3), we conclude that

$$L = nA + nH \int_{\Sigma} u dvol_{\Sigma} \geq nA + \frac{L}{n+1} \left( \frac{\int_{\Sigma} u dvol_{\Sigma}}{A} \right)^2 = nA + \frac{L}{n+1} \left( \frac{L - nA}{nHA} \right)^2.$$

This implies that

$$L - nA \geq \frac{L}{n+1} \left( \frac{L - nA}{nHA} \right)^2.$$

Therefore,

$$L^2 - nAL - n^2 A^2 (n+1) H^2 \leq 0.$$

This implies that

$$L \leq nA \left( \frac{1 + \sqrt{1 + 4(n+1)H^2}}{2} \right).$$

□

#### 4. PROOF OF COROLLARY 1.2

*Proof.* As in the proof of Theorem 11 in [R-V], we obtain that  $\partial\Sigma$  is embedded. Now, applying the Theorem 1.2, we obtain that  $\Sigma$  is totally geodesic or starshaped with respect to the center of the ball. Since starshaped surfaces must have genus 0, we obtain that  $\Sigma$  is totally geodesic or a spherical cap. □

#### REFERENCES

- [R-V] A. Ros and E. Vergasta, Stability for Hypersurfaces of Constant Mean Curvature with Free Boundary, *Geometriae Dedicata* 56 (1995), 19-33.
- [N] I. Nunes, On stable constant mean curvature surfaces with free-boundary. 2016 (arXiv: 1605.09625v1).
- [R] A. Ros, Stability of minimal and constant mean curvature surfaces with free boundary, *Mat. Contemp.* 35 (2008), 221-240.
- [So] R. Souam, On stability of stationary hypersurfaces for the partitioning problem for balls in spce forms, *Math. Z.* 224 (1997), 195-208.

UNIVERSIDADE FEDERAL DE MINAS GERAIS (UFMG), DEPARTAMENTO DE MATEMÁTICA, CAIXA POSTAL 702, 30123-970, BELO HORIZONTE, MG, BRAZIL  
*E-mail address:* ezequiel@mat.ufmg.br