

The Steinhaus-Weil property: its converse, subcontinuity and Solecki amenability

by

N. H. Bingham and A. J. Ostaszewski

Abstract.

The Steinhaus-Weil theorem that concerns us here is the simple, or classical, ‘interior-points’ property – that in a Polish topological group a non-negligible set B has the identity as an interior point of $B^{-1}B$. There are various converses; the one that mainly concerns us is due to Simmons and Mospan. Here the group is locally compact, so we have a *Haar* reference measure η . The Simmons-Mospan theorem states that a (regular Borel) measure has such a Steinhaus-Weil property if and only if it is absolutely continuous with respect to the Haar measure. In Part I (Propositions 1-7, Theorems 1-4) we exploit the connection between the interior-points property and a selective form of infinitesimal invariance afforded by a certain family of *selective* reference measures σ , drawing on Solecki’s amenability at 1 (and using Fuller’s notion of subcontinuity). In Part II (Propositions 8, 9, Theorems 5, 6) we develop a number of relatives of the Simmons-Mospan theorem. In Part III (Theorems 7, 8) we link this with topologies of Weil type. We close in Part IV with Propositions 12-13 and Theorem B – concerning the ‘composite interior-point’ property (of AB^{-1}) and the Borell ‘relative interior-point’ property (relative to the Cameron-Martin space) – and complements.

Keywords. Steinhaus-Weil property, amenability at 1, measure subcontinuity, Simmons-Mospan theorem, Weil topology, interior-points property, Haar measure, Lebesgue decomposition, left Haar null, Cameron-Martin space.

Classification: Primary 22A10, 43A05; Secondary 28C10.

Table of Contents

- 0. Introduction
- I. Subcontinuity and Solecki’s amenability
 - 1. Measure under translation – preliminaries
 - 2. Subcontinuity of measures
- II. Around the Simmons-Mospan Theorem
 - 3. A Lebesgue Decomposition

- 4. Discontinuity: the Simmons-Mospan Theorem
- III. Weil-like topologies
 - 5. Weil-like topologies: preliminaries
 - 6. Weil-like topologies: theorems
- IV. Other interior-point properties
 - 7.1. The Steinhaus property AA^{-1} versus the Steinhaus property AB^{-1}
 - 7.2. Borell's interior-point property
 - 8. Complements
- References

0 Introduction

We begin by stating the Steinhaus-Weil Theorem in its simplest form (Steinhaus [Ste] for the line, Weil [Wei, §11, p. 50] for a Polish locally compact group, Grosse-Erdmann [GroE]):

Theorem SW. *In a locally compact Polish group G with (left) Haar measure η_G , for non-null Borel B , $B^{-1}B$ (and likewise BB^{-1}) contains a neighbourhood of the identity.*

The context we work in here and throughout, unless otherwise stated, is that groups and spaces are assumed separable. This both simplifies the exposition and emphasizes that we need only the axiom of Dependant Choices (DC – ‘what is needed to make induction work’), rather than the Axiom of Choice (AC); cf. [BinO7]. For comments concerning non-separable settings, see §8.1.

The interior-point property of the measure-theoretically ‘non-negligible’ set B of the theorem is referred to as the *Steinhaus-Weil property*, which encompasses the category variant due to Piccard [Pic] and Pettis [Pet], cf. Cor. 2' and Th. 7B (by reference, when appropriate, to the *quasi-interior* of a set – the largest open set equivalent to it modulo a meagre set). This important result has many ramifications; for example, it is basic to the theory of regular variation – see e.g. [BinGT, Th. 1.1.1].

We are also concerned here with its converse, the Simmons-Mospan theorem ([Sim, Th. 1], [Mos, Th. 7], recently rediscovered in the abelian case [BarFF, Th. 10]), stated in its simplest form.

Theorem SM. *In a locally compact Polish group, a Borel measure has the Steinhaus-Weil property if and only if it is absolutely continuous with respect to Haar measure.*

The proof of this and related (converse) results emerges in §4 after due preparatory work; regarding the reduction of the non-separable to the separable case, see the closing remark of §4. The results below hinge on work of Solecki [Sol2] on amenability at 1 and the concept of subcontinuity (see §2 and below). These are aimed at freeing up the classical dependency on local compactness and the corresponding standard (Haar) reference measure. To the best of our knowledge such aims, in respect of topological groups, were last undertaken by Xia in 1972 in Chapter 3 of [Xia], where the emphasis is on (relative) quasi-invariance (cf. §7.2), a topic we pursue in the companion paper [Bin08] with tools developed here. (The additive subgroup context of the more recent [Bana] sits in a topological vector space.)

For G a topological group with (admissible) metric d (briefly: metric group), denote by $\mathcal{M}(G)$ the family of Borel regular σ -finite measures on G , with $\mathcal{P}(G) \subseteq \mathcal{M}(G)$ the probability measures ([Kec, §17E], [Par]), by $\mathcal{P}_{\text{fin}}(G)$ the larger family of finitely-additive regular probability measures (cf. [Bin]), and by $\mathcal{M}_{\text{sub}}(G)$ submeasures (monotone, finitely subadditive set functions μ with $\mu(\emptyset) = 0$). Here *regular* is taken to imply both *inner* regularity (inner approximation by compact subsets, also called the *Radon* property, as in [Bog2, II §7.1] and [Sch]), and *outer* regularity (outer approximation by open sets). We recall that a σ -finite Borel measure on a metric space is necessarily outer regular ([Bog2, II. Th. 7.1.7], [Kal, Lemma 1.34], cf. [Par, Th. II.1.2] albeit for a probability measure) and, when the metric space is complete, inner regular ([Bog2, II. Th. 7.1.7], cf. [Par, Ths. II.3.1 and 3.2]). When G is locally compact we denote Haar measure by η_G or just η (η denoting capital eta in Greek). For X a metric space, we denote by $\mathcal{K} = \mathcal{K}(X)$ the family of compact subsets of X (the *hyperspace* of X in §1, where we view it as a topological space under the Hausdorff metric, or the Vietoris topology). For $\mu \in \mathcal{M}(G)$ we write ${}_g\mu(\cdot) := \mu(g\cdot)$ and $\mu_g(\cdot) := \mu(\cdot g)$; $\mathcal{M}(\mu)$ denotes the μ -measurable sets of G and $\mathcal{M}_+(\mu)$ those of finite positive measure, and $\mathcal{K}_+(\mu) := \mathcal{K}(G) \cap \mathcal{M}_+(\mu)$. For G a Polish group, recall that $E \subseteq G$ is *universally measurable* ($E \in \mathcal{U}(G)$) if E is measurable with respect to every measure $\mu \in \mathcal{P}(G)$ – for background, see e.g. [Kec, §21D], cf. [Fre, 434D, 432], [Sho]; these form a σ -algebra. Examples are analytic subsets (see e.g. [Rog, Part 1 §2.9], or [Kec, Th. 21.10], [Fre, 434Dc]) and the σ -algebra that

they generate. Beyond these are the *provably* Δ_2^1 sets of [FenN] – cf. [BinO7].

Recall that E is *left Haar null*, $E \in \mathcal{HN}$, as in Solecki [Sol1,2,3] (following [Chr1,2]) if there are $B \in \mathcal{U}(G)$ covering E and $\mu \in \mathcal{P}(G)$ with

$$\mu(gB) = 0 \quad (g \in G).$$

(The terminal brackets here and below indicate *universal quantification* over the free variable.) So if $B \in \mathcal{U}(G)$ is not left Haar null, then for each $\mu \in \mathcal{P}(G)$ there is compact $K = K_\mu \subseteq B$ and $g \in G$ with

$${}_g\mu(K) > 0.$$

The question then arises whether there is also $\delta > 0$ with ${}_g\mu(Kt) > 0$ for *all* $t \in B_\delta$, for $B_\delta = B_\delta(1_G)$ the open δ -ball centered at 1_G : a *right-sided property complementing* the earlier *left-sided* property (of nullity, or otherwise). If this is the case for some μ , then (see Corollary 2' in §2) $1_G \in \text{int}(K^{-1}K) \subseteq \text{int}(E^{-1}E)$; indeed, one has

$$K \cap Kt \in \mathcal{M}_+({}_g\mu) \quad (t \in B_\delta), \quad (*M)$$

and this implies (see Lemma 1, §2):

$$B_\delta \subseteq \text{int}(K^{-1}K) \subseteq \text{int}(E^{-1}E).$$

As this clearly forces local-compactness of G (see Lemma 1 below), for the more general context we weaken the ‘complementing right-sided property’ to hold only *selectively*: on a subset of B_δ of the form

$$\{z \in B_\delta : |\mu(Kz) - \mu(K)| < \varepsilon\}$$

(cf. $B_\delta^\Delta(\mu)$ in §2). This turns attention to refining the topology \mathcal{T}_d of G via the metrics

$$d_K(x, y) := d(x, y) + |\mu(Kx) - \mu(Ky)|$$

(for a family of sets $K \in \mathcal{K}_+(\mu)$ – cf. the Struble sampler of §5), determined in Theorem 5 below, aiming at a relative-interior-points property in the finer topology, the theme of the companion paper [BinO8], cf. §8.2. See (in addition to Theorem SW above) Kemperman [Kem] (cf. [Kuc, Lemma 3.7.2], [BinO1, Th. K], [BinO5, Th. 1(iv)]).

We are guided by the close relation between the measure-theoretic *Steinhaus-Weil-like* property (*M) and its category version

$$K \cap Kt \in \mathcal{B}_+(\tau), \quad (*B)$$

where the latter term refers to non-meagre *Baire* sets (= with the Baire property) of τ , a refinement of the ambient topology $\mathcal{T}_G = \mathcal{T}_d$ of G , the latter conveniently taken to be generated by a *left*-invariant metric $d = d_L^G$ with associated group-norm (§5) $\|x\| := d(x, 1_G) = d(tx, t)$ (so that $B_\delta(t) = tB_\delta$ – see Prop. 1). We refer to the (left) invariance of $\mathcal{B}_+(\tau)$ (under translation) as the (left) *Nikodym property* of τ .

In Part I below (§1,2, Props. 1-7, Th. 1-4: Subcontinuity Theorem, Aggregation Theorem, Shift-compactness Theorem, Strong Subcontinuity Theorem), in the context of a metric or Polish group G , we study continuity properties of the maps $m_K : t \mapsto \mu(Kt)$ in the light of theorems of Solecki [Sol2] and of Theorem SM above and related results. The key here is Fuller’s notion of subcontinuity, as applied to the function $m_K(t)$ at $t = 1_G$. This yields a fruitful interpretation of Solecki’s notion of *amenability at 1_G* via *selective subcontinuity* and linkage to *shift-compactness* (see Th. 3 below; the term is borrowed from [Par, III.2]). Since commutative Polish groups are amenable at 1 [Sol2, Th. 1(ii)], this widens the field of applicability of shift-compactness to non-Haar-null subsets of these, as in [BinO4], and leads to a conjecture (see remarks preceding Theorem 3) as to whether \mathcal{HN} comprises the negligible sets of some refinement topology of \mathcal{T}_d . In Part II (§3,4, Props. 8-9, Theorems 5,6: Disaggregation Theorem, Generalized Simmons Theorem) we study measure discontinuity and extend results of Simmons beyond his locally-compact context by reference to measures exhibiting selective subcontinuity. This draws on results in the companion paper [BinO8], which is concerned with the relative-interior-point property inspired by the *Cameron-Martin* theory of Gaussian measures (§7.2 and §8.2). In Part III (§6, 7, Props. 10, 11, Theorems 7, 7M, 7B, 8) we study the converse problem of the *Weil topologies* generated by the *Fréchet-Nikodym* pseudo-norms, defined for $E \in \mathcal{M}_+(G)$ by:

$$\|t\|_\mu^E := \mu(tE\Delta E) \quad (t \in G).$$

For ease of reference – duly identified, as here – we study EE^{-1} (for $\|t\|_\mu^E > 0$) when following Simmons and Weil, and $E^{-1}E$ when following Solecki. In Part IV (§7, Props. 12-13, Th. K, Th. B) we study the relation of the AA^{-1} interior-point property to that for AB^{-1} , and discuss the *Borell interior-point property* relative to the Cameron-Martin ‘embedded subspace’. We close with a section of complements (§8).

Część I

Subcontinuity and Solecki's amenability

1 Measure under translation – preliminaries

We begin with a form of the ‘telescope’ or ‘tube’ lemma (cf. [Mun, Lemma 5.8]), applied in §2. Our usage of upper semicontinuity in relation to set-valued maps follows [Rog], cf. [Bor].

Proposition 1 (cf. [Hey, 1.2.8]). *For a metric group G and compact $K \subseteq G$, the map $t \mapsto Kt$ is upper semicontinuous; in particular, for $\mu \in \mathcal{M}(G)$,*

$$m_K : t \mapsto \mu(Kt)$$

is upper semicontinuous, hence μ -measurable. In particular, if $m_K(t) = 0$, then m is continuous at t .

Proof. For K compact and $V \supseteq K$ open, pick for each $k \in K$ a $\delta(k) > 0$ with $kB_{2\delta(k)} \subseteq V$. By compactness, there are k_1, \dots, k_n with $K \subseteq \bigcup_j k_j B_{\delta(k_j)} \subseteq V$; then for $\delta := \min_j r(k_j) > 0$

$$Kt \subseteq \bigcup_j k_j B_{\delta(k_j)} t \subseteq \bigcup_j k_j B_{2\delta(k_j)} \subseteq V \quad (\|t\| < \delta).$$

To prove upper semicontinuity of m_K , fix $t \in G$. For $\varepsilon > 0$, as Kt is compact, choose by outer regularity an open $U \supseteq Kt$ with $\mu(U) < \mu(Kt) + \varepsilon$; as before, there is an open ball B_δ at 1_G with $KtB_\delta \subseteq U$, and then $\mu(KtB_\delta) \leq \mu(U) < \mu(Kt) + \varepsilon$. The final assertion follows from positivity of m_K . \square

We continue with an analogue. The result is folklore, cf. [BeeV, Th. 3.2(i)]; it comes close to matters touched on in [Ost1, §3]. Here and below the vertical section of a set A is denoted $A_x := \{y : (x, y) \in A\}$.

Proposition 2 (Sectional upper semicontinuity). *For a metric group G , compact $F \subseteq G$ and compact $K \subseteq G^2$, the map*

$$x \mapsto K_x \quad (x \in F)$$

is upper semicontinuous.

Proof. For $V \subseteq G$ open with $K_x \subseteq V$, suppose for $x_n \in F$ with $x_n \rightarrow x$ that $(x_n, y_n) \in K \setminus (G \times V)$. By compactness of K , we may suppose w.l.o.g. that $y_n \rightarrow y$. Then $(x, y) \in K \setminus (G \times V)$, and so $(x, y) \in \{x\} \times K_x$ and $y \notin V$; but $y \in K_x \subseteq V$, a contradiction. \square

From Prop. 2 on *upper* semicontinuity, we obtain information about $m_K : t \mapsto \mu(Kt)$ below. This links with *lower* semicontinuity. By a theorem of Fort, the ε -continuity points (defined in terms of the Hausdorff metric: see [For]) of an upper semicontinuous compact-valued mapping of a metric space into a totally bounded metric space form a dense open set, implying in a real-valued context such as here *continuity* on a co-meagre set. We return to this shortly in Theorem LB below.

Proposition 3 (Sectional upper semicontinuity under a measure).

For a metric group G , compact $F \subseteq G$ and compact $K \subseteq G^2$, and $\mu \in \mathcal{M}(G)$, the map

$$m : x \mapsto \mu(K_x) \quad (x \in F = \text{proj}_1 K)$$

is upper semicontinuous, and so Borel.

Proof. Fix $x \in F$. Let $\varepsilon > 0$. By outer regularity, take V open in G with $K_x \subseteq V$ and $\mu(V) < \mu(K_x) + \varepsilon$. By Prop. 2 $x \mapsto \{x\} \times K_x$ is upper semicontinuous on F ; so for some open neighbourhood U of x

$$K \cap (U \times G) \subseteq K \cap (U \times V).$$

So for $y \in F \cap U$

$$K_y \subseteq V,$$

and so $\mu(K_y) \leq \mu(V) < \mu(K_x) + \varepsilon$, proving the first assertion. The second assertion follows since

$$m^{-1}(a, b) = \bigcap_{n \in \mathbb{N}} m^{-1}[0, b) \setminus m^{-1}[0, a + 1/n). \quad \square$$

For further results on Borel-measurability of regular Borel measures see [BeeV, Th. 2.2] (there termed ‘Radon measures’).

We will need the following result in Parts II and III (see Lemma 2, §4, and Th. 7, §6), preferable to the usual Fubini Theorem as using qualitative rather than quantitative measure theory (like the Kuratowski-Ulam Theorem [FreNR]). Interestingly, it may be proved by mimicking the proof of Prop. 1 above, yielding a simplification to that by Eric van Downen [vDo], itself a simplification of that in [Oxt2, Ch. 14]: for the proof (omitted here), see §8.12.

Theorem FN (Fubini theorem for null sets). *For a metric group G and $A \subseteq G^2$ measurable under $\mu \times \nu$, with $\mu, \nu \in \mathcal{M}(G)$: if the ‘exceptional set’ of points x for which the vertical section A_x is ν -non-null is itself μ -null, then A is $\mu \times \nu$ -null.*

We close this section with a study of the continuity properties of the map $m_K : t \mapsto \mu(Kt)$ for compact K , extending Prop. 3.

Corollary 1 (Fort [For]). *In Proposition 2, $t \mapsto \mu(Kt)$ is lower semi-continuous (so also continuous) on a co-meagre set.*

We can improve on the preceding result by recourse to a natural generalization, for our compact sectional context, of the classical continuity theorems of Luzin [Hal, §55] and Baire [Oxt2, Th. 8.1] – see also [Sch, Ch. 1, §5]. Below for a compact metric space X , we denote by $\mathcal{K}(X)$ the *hyperspace of X* , the space of compact subsets of X under the Hausdorff metric, or Vietoris topology; here this is also a compact space ([Eng, 2.7.28], [Kec, Th. 4.25], [Mic]). Then (LB for ‘Luzin-Baire’):

Theorem LB. *For G a metric group and compact $K \subseteq G^2$, the map $\kappa : G \rightarrow \mathcal{K}(G) : x \mapsto K_x$ is Borel-measurable, and so*

(i) *κ is continuous relative to a co-meagre set.*

For $\mu \in \mathcal{P}(G)$:

(ii) *for each $\varepsilon > 0$ there is a Borel set S_ε with $\mu(G \setminus S_\varepsilon) < \varepsilon$ such that $x \mapsto K_x$ is continuous on S_ε ; equivalently:*

(ii)' *there is an increasing sequence of Borel sets S_n with union μ -almost all of G such that $x \mapsto K_x$ is continuous on each S_n .*

Proof. For U, V open, the set $\{x : K_x \subseteq U \text{ and } K_x \cap V \neq \emptyset\}$ is σ -compact; indeed, by Prop. 2, $\{x : K_x \subseteq U\}$ is open, whereas for $F := \text{proj}_1(K)$, a compact set

$$\{x : K_x \cap U \neq \emptyset\} = \text{proj}_1[(F \times U) \cap K],$$

which is the projection of a σ -compact set. Then (ii) follows from Luzin's theorem ([Hal, §55]), and for (ii)' see [BinO1, p. 142]. Its extension to a regular (i.a. \mathcal{G} -outer regular) σ -finite measure may be made via Egoroff's Theorem [Hal, §21 Th. A] – see [Zak]. \square

A first corollary is the following result on the continuity of the map

$$x \mapsto \|x\|_E^\mu = \mu(xE\Delta E),$$

for measurable E , by compact approximation. Below, the sets C_x associated with points x should be interpreted as neighbourhoods of x in the spirit of a Hashimoto ideal topology for the ideal of μ -null sets, for which see [LukMZ], or [BinO5]. This mimicks Weil's *proof* of the 'fragmentation lemma' (§8, Lemma 3) in [Hal, Ch. XII §62 Th. A] (cf. [Wei, Ch. VII, §31]).

Proposition 4 (Almost everywhere continuity). *For a metric group G , $\delta > 0$, $\mu \in \mathcal{P}(G)$, $E \in \mathcal{M}_+(\mu)$, and $F \in \mathcal{K}_+(\mu)$:*

there is a compact $C \subseteq F$ with $\mu(F \setminus C) < \delta$ such that for any $\varepsilon > 0$ and each $x \in C$ there is a μ -non-null measurable $C_x \subseteq C$ containing x with

$$|\mu(xE\Delta E) - \mu(yE\Delta E)| < \varepsilon \quad (y \in C_x).$$

In particular, there is an increasing family of compact sets C_n with union μ -almost all of G satisfying the above with C_n for C .

Proof. Fix $E \subseteq G$ measurable and for F compact with $\mu(F) > 0$, put

$$\begin{aligned} H &:= \bigcup_{x \in F} \{x\} \times (E\Delta xE) = \bigcup_{x \in F} \{x\} \times ((xE \setminus E) \cup (E \setminus xE)) \\ &= \bigcup_{x \in F} (\{x\} \times xE) \setminus \bigcup_{x \in F} (\{x\} \times E) \cup \bigcup_{x \in F} (\{x\} \times E) \setminus \bigcup_{x \in F} \{x\} \times (xE), \end{aligned}$$

which is measurable. So $F \subseteq \text{proj}_1(H)$ and for $x \in F$

$$H_x := xE\Delta E.$$

We will work inductively, taking successively smaller values of ε . Fix $\varepsilon > 0$ with $\varepsilon < \mu(F)$ and choose K a finite union of compact rectangles with

$$(\mu \times \mu)(H\Delta K) < \varepsilon^2.$$

So

$$K = \bigcup_{j \leq n} (F^j \times K^j),$$

say. Let $S = S_\varepsilon := \{x \in F : \mu((H\Delta K)_x) \geq \varepsilon\}$. Then $\mu(S) \leq \varepsilon$; otherwise

$$\varepsilon^2 > (\mu \times \mu)(H\Delta K) \geq \varepsilon\mu(S) \geq \varepsilon^2,$$

a contradiction. So

$$\mu(F \setminus S) > \mu(F) - \varepsilon.$$

So we may choose a compact set $C = C_\varepsilon \subseteq F \setminus S_\varepsilon$ with $\mu(F \setminus C) < \varepsilon$ and

$$\mu(H_x \Delta K_x) \leq \varepsilon \quad (x \in C_\varepsilon).$$

But $K_x \in \{K^1, \dots, K^n\}$, so

$$C_\varepsilon = \bigcup_{j \leq n} C_\varepsilon^j, \quad \text{where } C_\varepsilon^j := \{x \in C_\varepsilon : \mu(H_x \Delta K^j) \leq \varepsilon\}.$$

Now for $x, y \in C_\varepsilon^j$

$$\mu(H_x \Delta H_y) \leq \mu(H_x \Delta K^j) + \mu(K^j \Delta H_y) \leq 2\varepsilon,$$

and w.l.o.g. the sets C_ε^j may be assumed compact.

Now fix $\delta > 0$ and repeat the construction inductively, taking in turn $\varepsilon = \varepsilon_n = 2^{-n}\mu(F)\delta$, to obtain a sequence of compact sets $F = C^0 \supseteq C = C^1 \supseteq C^2 \supseteq C^3 \dots$ with $C^n = C_{\varepsilon_n}$ as above, and $\mu(C^n \setminus C^{n+1}) < \varepsilon_n$. These have non-null intersection C_0 :

$$\mu(F \setminus C_0) < \sum_{n \geq 1} 2^{-n}\mu(F)\delta = \mu(F)\delta.$$

Now for each $\varepsilon > 0$ there is n with $\varepsilon_n < \varepsilon$. So for $x \in C_0 \subseteq C^n = C_{\varepsilon_n} = \bigcup_{j \leq n} C_{\varepsilon_n}^j$

$$\mu(H_x \Delta H_y) < \mu(H_x \Delta H_y) + 2\varepsilon_n < \mu(H_x) + 2\varepsilon \quad (x, y \in C_{\varepsilon_n}^j \cap C_0).$$

So

$$|\mu(H_x) - \mu(H_y)| < 4\varepsilon \quad (x, y \in C_{\varepsilon_n}^j \cap C_0). \quad \square$$

A proof similar to but simpler than that above (omitted here – see §8.13) improves Prop. 1:

Proposition 5 (Almost everywhere upper semicontinuity). *For a metric group G , $\delta > 0$, $\mu \in \mathcal{P}(G)$, $E \in \mathcal{M}_+(\mu)$, and $F \in \mathcal{K}_+(\mu)$:*

there is a compact $C \subseteq F$ with $\mu(F \setminus C) < \delta$ such that for any $\varepsilon > 0$ each $x \in C$ has a neighbourhood U_x with

$$\mu(yE) < \mu(xE) + \varepsilon \quad (y \in C \cap U_x).$$

In particular, there are disjoint compact sets C with union μ -almost all of G for which this holds.

2 Subcontinuity of measures

Proposition 1 above, on upper semicontinuity, motivates the following definitions, the key one being an adaptation of *subcontinuity* (of functions) due to Fuller [Ful] (for which see Remark 4 below) to the context of measures. We focus on the *right-sided* version of the concept. Subcontinuity is a natural auxiliary in the quest for fuller forms of continuity: as one instance, see [Bou] for the step from separate to joint continuity; as another, classic instance, note that a subcontinuous set-valued map with closed graph (yet another relative of upper semicontinuity) is continuous – see [HoIN] for an extensive bibliography. Here its relevance to the Steinhaus-Weil Theorem (which seems to be new here) yields Theorems 1 and 3, linking *amenability at 1* with *shift-compactness*, for which see Theorem 3 below (the latter term is borrowed from [Par, III.2]).

Definition ([BinO5]). For $\mu \in \mathcal{P}_{\text{fin}}(G)$, and (compact) $K \in \mathcal{K}(G)$, noting that $\mu_\delta(K) := \inf\{\mu(Kt) : t \in B_\delta\}$ is weakly decreasing in δ , put

$$\mu_-(K) := \sup_{\delta > 0} \inf\{\mu(Kt) : t \in B_\delta\},$$

and, for $\mathbf{t} = \{t_n\}$ a *null sequence*, i.e. with $t_n \rightarrow 1_G$,

$$\mu_-^{\mathbf{t}}(K) := \liminf_{n \rightarrow \infty} \mu(Kt_n).$$

Then

$$0 \leq \mu_-(K) \leq \mu(K) = \inf_{\delta > 0} \sup\{\mu(Kt) : t \in B_\delta\},$$

by Proposition 1. We say that a null sequence \mathbf{t} is *non-trivial* if $t_n \neq 1_G$ infinitely often. Define (with the quantification convention of the Introduction) as follows:

- (i) μ is *translation-continuous* (‘*continuous*’ or ‘*mobile*’) if $\mu(K) = \mu_-(K)$ ($K \in \mathcal{K}(G)$);
- (ii) μ is *maximally discontinuous* at $K \in \mathcal{K}(G)$ if $0 = \mu_-(K) < \mu(K)$;
- (iii) μ is *subcontinuous* if $0 < \mu_-(K) \leq \mu(K)$ ($K \in \mathcal{K}_+(\mu)$);
- (iv) μ is (*selectively*) *subcontinuous at* $K \in \mathcal{K}_+(\mu)$ *along* \mathbf{t} if $\mu_-^{\mathbf{t}}(K) > 0$.

Remarks. 1. $m_K(\cdot)$ is *continuous* if μ is continuous, since $m_K(st) = m_{Ks}(t)$ and Ks is compact whenever K is compact; for directional continuity of measures in linear spaces see [Bog3, §3.1]. In [LiuR] (cf. [LiuRW], [Gow1,2])

- a Radon measure μ on a space X , on which a group G acts homeomorphically, is called *mobile* if $t \mapsto \mu(Kt)$ is continuous for all $K \in \mathcal{K}(X)$.
2. For G locally compact (i) holds for μ the left Haar measure η_G , and also for $\mu \ll \eta_G$ (absolutely continuous w.r.t. to η_G).
 3. A measure μ singular w.r.t. Haar measure is maximally discontinuous for its support: this is at the heart of the analysis offered by Simmons (and independently, much later by Mospan) – see Corollary 2' below.
 4. *Subcontinuity*, in the sense of [Ful], of a map $f : G \rightarrow (0, \infty)$ requires that, for every $t_n \rightarrow t \in G$, there is a subsequence $t_{m(n)}$ with $f(t_{m(n)})$ convergent in the range (i.e. to a positive value). The distinguished role of null sequences emerges below in the *Subcontinuity Theorem* (Theorem 1). Null sequences should be viewed here as selecting stepwise (or even pathwise, under local connectedness, as suggested by Tomasz Natkaniec) ‘asymptotic directions’ justifying the phrase ‘*along t*’ in (iv) above, and allowing (iv) to be interpreted as a *selective subcontinuity* in ‘direction’ \mathbf{t} . The analogous selective concept in a linear space is ‘along a vector’ as in [Bog3, §3.1].
 5. *Selective versus uniform subcontinuity*. Definition (iii) is equivalent to demanding for $K \in \mathcal{K}_+(\mu)$ that *any* null sequence $\mathbf{t} = \{t_n\}$ have a subsequence $\mu(Kt_{m(n)})$ bounded away from 0; then (iii) may be viewed as demanding ‘uniform subcontinuity’: selective subcontinuity along *each* \mathbf{t} for all $K \in \mathcal{K}_+(\mu)$.
 6. *Left- versus right-sided versions*. Writing $\tilde{\mu}(E) := \mu(E^{-1})$ for the *inverse measure* captures versions associated with right-sided translation such as $\tilde{\mu}_-$ and

$$\tilde{\mu}_-^{\mathbf{t}}(K) := \liminf_{n \rightarrow \infty} \mu(t_n K).$$

Definition. We will say that μ is *symmetric* if $\mu = \tilde{\mu}$; then B is null iff B^{-1} is null.

In Lemma 1 below it suffices for μ to be a bounded, regular submeasure which is *supermodular*:

$$\mu(E \cup F) \geq \mu(E) + \mu(F) - \mu(E \cap F);$$

recall, however, from [Bog2, 1.12.37] the opportunity to replace, for any $K \in \mathcal{K}(G)$, a supermodular submeasure μ by a dominating $\mu' \in \mathcal{M}_{\text{fin}}(G)$, i.e. with $\mu'(K) \geq \mu(K)$.

For $K \in \mathcal{K}_+(\mu)$ and $\delta, \Delta > 0$, put

$$B_\delta^\Delta = B_\delta^{K, \Delta}(\mu) := \{z \in B_\delta : \mu(Kz) > \Delta\},$$

which is monotonic in $\Delta : B_\delta^\Delta \subseteq B_\delta^{\Delta'}$ for $0 < \Delta' \leq \Delta$. Note that $1_G \in B_\delta^\Delta$ for $0 < \Delta < \mu(K)$.

The specialization below to a mobile measure (see above) may be found in [Gow1,2].

Lemma 1 (cf. [BinO5, Th. 2.5]). *Let $\mu \in \mathcal{P}_{\text{fin}}(G)$ for G a metric group. For $K \in \mathcal{K}_+(\mu)$, if $\mu_-^{\mathbf{t}}(K) > 0$ for some non-trivial null sequence \mathbf{t} , then there are $\delta > 0$ and $0 < \Delta < \mu_-^{\mathbf{t}}(K)$ with $t_n \in B_\delta^\Delta$ for all large enough n and*

$$\Delta \leq \mu(K \cap Kt) \quad (t \in B_\delta^\Delta),$$

so that

$$K \cap Kt \in \mathcal{M}_+(\mu) \quad (t \in B_\delta^\Delta). \quad (*)$$

In particular,

$$K \cap Kt \neq \emptyset \quad (t \in B_\delta^\Delta),$$

or, equivalently,

$$B_\delta^\Delta \subseteq K^{-1}K, \quad (**)$$

so that B_δ^Δ has compact closure.

A fortiori, if $\mu_-(K) > 0$, then $\delta, \Delta > 0$ may be chosen with $\Delta < \mu_-(K)$ and $B_\delta \subseteq B_\delta^\Delta$ so that () and (**) hold with B_δ replacing B_δ^Δ , and in particular G is locally compact.*

Proof. For the first part take $\Delta := \mu_-^{\mathbf{t}}(K)/4$. Then, for $t \in B_\delta^\Delta$ with $\delta > 0$ arbitrary, $\mu(Kt) > 2\Delta$, and so, since $t_n \in B_\delta$ for all large enough n , also $t_n \in B_\delta^\Delta$ for all large enough n , so $B_\delta^\Delta(K) \setminus \{1_G\}$ is non-empty for \mathbf{t} non-trivial.

Put $H_t := K \cap Kt \subseteq K$. By outer regularity of μ , choose U open with $K \subseteq U$ and $\mu(U) < \mu(K) + \Delta$. By upper semicontinuity of $t \mapsto Kt$, w.l.o.g. $KB_\delta \subseteq U$ for some $\delta > 0$. For $t \in B_\delta^\Delta$, by finite additivity of μ , since $2\Delta < \mu(Kt)$

$$\begin{aligned} 2\Delta + \mu(K) - \mu(H_t) &\leq \mu(Kt) + \mu(K) - \mu(H_t) = \mu(Kt \cup K) \\ &\leq \mu(U) \leq \mu(K) + \Delta. \end{aligned}$$

Comparing extreme ends of this chain of inequalities gives

$$0 < \Delta \leq \mu(H_t) \quad (t \in B_\delta^\Delta).$$

For $t \in B_\delta^\Delta$, as $K \cap Kt \in \mathcal{M}_+(\mu)$, take $s \in K \cap Kt \neq \emptyset$; then $s = kt$ for some $k \in K$, so $t = k^{-1}s \in K^{-1}K$. Conversely, $t \in B_\delta^\Delta \subseteq K^{-1}K$ yields $t = k^{-1}k'$ for some $k, k' \in K$; then $k' = kt \in K \cap Kt$.

By the compactness of $K^{-1}K$, B_δ^Δ has compact closure.

As for the final assertions, if $\mu_-(K) > 0$, take $\Delta := \mu_-(K)/2$. Then $\inf\{\mu(Kt) : t \in B_\delta\} > \Delta$ for all small enough $\delta > 0$, and so in particular $\mu(Kt) > \Delta$ for $t \in B_\delta$, i.e. $B_\delta \subseteq B_\delta^\Delta$. So the argument above applies for such $\delta > 0$ with B_δ in lieu of B_δ^Δ , just as before. Here the compactness of $K^{-1}K$ now implies local compactness of G itself. \square

As an immediate and useful corollary, we have

Lemma 1'. *For $\mu \in \mathcal{P}_{\text{fin}}(G)$, with G a metric group, any null sequence \mathbf{t} and any $K \in \mathcal{K}(G)$: if $\mu_-^{\mathbf{t}}(K) > 0$, then there is $m \in \mathbb{N}$ with*

$$0 < \mu_-^{\mathbf{t}}(K)/4 < \mu(K \cap Kt_n) \quad (n > m). \quad (*')$$

In particular,

$$t_n \in K^{-1}K \quad (n > m). \quad (**')$$

Proof. Apply Lemma 1 to obtain $\Delta, \delta > 0$; for $t \in B_\delta^\Delta$, $\mu(Kt) > \Delta$, so as above $t_n \in B_\delta^\Delta$ for all large enough n . \square

This permits a connection with left Haar null sets; recall that a group G is *amenable at 1* [Sol2] (see below for the origin of this term) if, given $\boldsymbol{\mu} := \{\mu_n\}_{n \in \mathbb{N}} \subseteq \mathcal{P}(G)$ with $1_G \in \text{supp}(\mu_n)$, for $n \in \mathbb{N}$ there are σ and σ_n in $\mathcal{P}(G)$ with $\sigma_n \ll \mu_n$ satisfying:

$$\sigma_n * \sigma(K) \rightarrow \sigma(K) \quad (K \in \mathcal{K}(G)).$$

We refer to σ (or $\sigma(\boldsymbol{\mu})$ if context requires) as a *Solecki measure* and to the measures σ_n , if needed, as *associated measures* (corresponding to the sequence $\{\mu_n\}_{n \in \mathbb{N}}$).

Solecki explains ([Sol2, end of §2]) the use of the term ‘amenability at 1’ as a localization (via the restriction that supports contain 1_G) of a *Reiter-like condition* [Pat, Prop. 0.4] which characterizes amenability: for $\mu \in \mathcal{P}(G)$ and $\varepsilon > 0$, there is $\nu \in \mathcal{P}(G)$ with

$$|\nu * \mu(K) - \nu(K)| < \varepsilon \quad (K \in \mathcal{K}(G)).$$

Lemma 1 and the next several results disaggregate Solecki's Interior-point Theorem [Sol2, Th 1(ii)] (Corollary 2 below), shedding more light on it and in particular connecting it to shift-compactness (Theorem 3 below). Indeed, we see that interior-point theorem itself as an 'aggregation' phenomenon. Theorem 5 of Part II below identifies subgroups with a 'disaggregation' topology, refining \mathcal{T}_G by using sets of the form $B_\delta^{K\Delta}(\sigma)$, the measures σ being provided in our first result:

Theorem 1 (Subcontinuity Theorem, after Solecki [Sol2, Th. 1(ii)]). *For G Polish and amenable at 1_G and \mathbf{t} a null sequence, there is $\sigma = \sigma(\mathbf{t}) \in \mathcal{P}(G)$ such that for each $K \in \mathcal{K}_+(\sigma)$ there is a subsequence $\mathbf{s} = \mathbf{s}(K) := \{t_{m(n)}\}$ with*

$$\lim_n \sigma(Kt_{m(n)}) = \nu(K) \quad (n \in \mathbb{N}), \text{ so } \quad \sigma_-^{\mathbf{s}}(K) > 0.$$

Proof. For $\mathbf{t} = \{t_n\}$ null, put $\mu_n := 2^{n-1} \sum_{m \geq n} 2^{-m} \delta_{t_m^{-1}} \in \mathcal{P}(G)$; then $1_G \in \text{supp}(\mu_n) \supseteq \{t_m^{-1} : m > n\}$. By definition of amenability at 1_G , in $\mathcal{P}(G)$ there are σ and $\sigma_n \ll \mu_n$, with $\sigma_n * \sigma(K) \rightarrow \sigma(K)$ for all $K \in \mathcal{K}(G)$. For $n \in \mathbb{N}$ choose $\alpha_{mn} \geq 0$ with $\sum_{m \geq n} \alpha_{mn} = 1$ ($n \in \mathbb{N}$) and with $\sigma_n := \sum_{m \geq n} \alpha_{mn} \delta_{t_m^{-1}}$.

Fix $K \in \mathcal{K}_+(\sigma)$ and θ with $0 < \theta < 1$. As K is compact, $\sigma_n * \sigma(K) \rightarrow \sigma(K)$; then w.l.o.g.

$$\sigma_n * \sigma(K) > \theta \sigma(K) \quad (n \in \mathbb{N}).$$

Then for each n

$$\sup\{\sigma(Kt_m) : m \geq n\} \cdot \sum_{m \geq n} \alpha_{mn} \geq \sum_{m \geq n} \alpha_{mn} \sigma(Kt_m) > \theta \sigma(K).$$

So for each n there is $m = m(\theta) \geq n$ with

$$\sigma(Kt_m) > \theta \sigma(K).$$

Now choose $m(n) \geq n$ inductively so that $\sigma(Kt_{m(n)}) > (1 - 2^{-n})\sigma(K)$; then, by Proposition 1, $\lim_n \sigma(Kt_{m(n)}) = \sigma(K)$: σ is subcontinuous along $\mathbf{s} := \{t_{m(n)}\}$ on K . \square

Remark. The selection above of the subsequence \mathbf{s} mirrors the role of 'admissible directions' which we encounter later in Cameron-Martin theory (§7.2 and below §8.2).

We are now able to deduce Solecki's interior-point theorem in a slightly stronger form, which asserts that the sets B_δ^Δ reconstruct the open sets of G using the compact subsets of a 'non-negligible set', as follows.

Theorem 2 (Aggregation Theorem). *For G Polish and amenable at 1_G , if $E \in \mathcal{U}(G)$ is not left Haar null – then, setting*

$$\hat{E} := \bigcup_{\delta, \Delta > 0, g \in G, \mathbf{t}} \{B_\delta^{gK, \Delta}(\nu(\mathbf{t})) : K \subseteq E, K \in \mathcal{K}_+(\nu(\mathbf{t})), \Delta < \nu(\mathbf{t})(gK)\},$$

$$1_G \in \text{int}(\hat{E}) \subseteq \hat{E} \subseteq E^{-1}E.$$

In particular, for E open, $1_G \in \text{int}(\hat{E})$.

Proof. Suppose otherwise; then for each n there is

$$t_n \in B_{1/n} \setminus \hat{E}.$$

Consider $\sigma = \sigma(\mathbf{t})$. As E is not left Haar null, there is g with $\sigma(gE) > 0$. Choose compact $K \subseteq gE$ with $\sigma(K) > 0$. Then with $h := g^{-1}$ and $H := hK \subseteq E$, $\sigma(K) = \sigma(gH) = \sigma^s(gH) > 0$ for some subsequence $\mathbf{s} = \{t_{m(m)}\}$. So, as in Lemma 1, with $\Delta := \sigma(gH)/4$ for some $\delta > 0$

$$B_\delta^{gH\Delta}(\sigma(\mathbf{t})) \subseteq (gH)^{-1}gH = H^{-1}H \subseteq E^{-1}E.$$

Choose n with $n > 1/\delta$. Then $t_n \in B_\delta$ for all $m > n$; so for infinitely many k

$$t_{m(k)} \in B_\delta^{gH\Delta}(\sigma(\mathbf{t})) \subseteq \hat{E},$$

a contradiction. As for the last assertion, for E open, D countable and dense, $G \subseteq \bigcup_{d \in D} dE$, so for any $\mu \in \mathcal{P}(G)$ (in particular for σ) $\mu(dE) > 0$ for some $d \in D$, and so E is not left Haar null. \square

The immediate consequence is

Corollary 2 (Solecki's Interior-Point Theorem [Sol2, Th 1(ii)]). *For G Polish and amenable at 1_G , if $E \in \mathcal{U}(G)$ is not left Haar null, then $1_G \in \text{int}(E^{-1}E)$.*

Corollary 2'. *For G a Polish group, if $B \in \mathcal{U}(G)$ is not left Haar null and is in $\mathcal{M}_+(\mu)$ for some subcontinuous $\mu \in \mathcal{P}_{\text{fin}}(G)$, then (**) holds for some $\delta > 0$.*

In particular, (**) holds in a locally compact group G , for any Baire non-meagre set E .

Proof. The first assertion is immediate from Lemma 1. As for the second, for a non-meagre Baire set E , if \tilde{E} is the quasi-interior and $K \subseteq \tilde{E}$ is compact with non-empty interior, then $\eta_G(K) > 0$. Since η is subcontinuous, there is $\delta > 0$ with

$$Kt \cap K \neq \emptyset \quad (||t|| < \delta).$$

A fortiori,

$$\tilde{E}t \cap \tilde{E} \neq \emptyset \quad (||t|| < \delta);$$

then $U := (Et) \cap \tilde{E} \neq \emptyset$, since $(Et) \cap \tilde{E} = \tilde{E}t$ (the Nikodym property of the usual topology of G). So since U is open and non-meagre, also $Et \cap \tilde{E} \neq \emptyset$, and so again (**). \square

The next result establishes the embeddability by (left-sided) translation of an appropriate subsequence of a given *null* sequence into a given target set that (like-sidedly) is non-left-Haar null. This property of embedding into a *non-negligible* set, first studied in respect of category and measure negligibility on \mathbb{R} by Kestelman and much later independently by Borwein and Ditor and thereafter also by other authors, mostly for combinatorial challenges, has emerged as an important general unifying principle, termed *shift-compactness*, applicable in a much wider context embracing metric groups G under various topologies refining \mathcal{T}_G and so defining various notions of negligibility: for the background here see [BinO2,3], [MilO]. Its consequences include various uniform-boundedness theorems as well as the Effros and the Open Mapping Theorems. Here we establish the said property, announced in [MilO], in relation to the ideal \mathcal{HN} of left Haar null sets. (It is a σ -ideal for Polish G in the presence of amenability at 1 [Sol2, Th 1(i)].) This leaves open the ‘converse question’ of a refinement topology for which \mathcal{HN} is the associated notion of negligibility; this seems plausible under the continuum hypothesis, CH, if one restricts attention only to Borel sets in \mathcal{HN} and their subsets by lifting a result concerning \mathbb{R} in [CieJ, Cor. 4.2] to G – see also the Remark following our next result.

Theorem 3 (Shift-compactness Theorem for \mathcal{HN}). *For G Polish and amenable at 1_G , if $E \in \mathcal{U}(G)$ is not left Haar null and z_n is null, then there are $s \in E$ and an infinite $\mathbb{M} \subseteq \mathbb{N}$ with*

$$\{sz_m : m \in \mathbb{M}\} \subseteq E.$$

Indeed, this holds for quasi all $s \in E$, i.e. off a left Haar null set.

Proof. Put $t_n := z_n^{-1}$, which is null. With $\sigma = \sigma(\mathbf{t})$ as in the Subcontinuity Theorem, since E is not left Haar null, there is g with $\sigma(gE) > 0$. For this g , put $\mu := {}_g\sigma$. Fix a compact $K_0 \subseteq E$ with $\mu(K_0) > 0$ and then, passing to a subsequence of \mathbf{t} as necessary (by Th. 1), we may assume that $\mu_-^{\mathbf{t}}(K_0) > 0$. Proceed to choose inductively a sequence $m(n) \in \mathbb{N}$, and decreasing compact sets $K_n \subseteq K_0 \subseteq E$ with $\mu(K_n) > 0$ such that

$$\mu(K_n \cap K_n t_{m(n)}) > 0.$$

To check the inductive step, suppose K_n already defined. As $\mu(K_n) > 0$, by the Subcontinuity Theorem, there is a subsequence $\mathbf{s} = \mathbf{s}(K_n)$ of \mathbf{t} with $\mu_-^{\mathbf{s}}(K_n) > 0$. By Lemma 1', there is $k(n) > n$ such that $\mu(K_n \cap K_n s_{k(n)}) > 0$. Putting $t_{m(n)} = s_{k(n)}$ and $K_{n+1} := K_n \cap K_n t_{m(n)} \subseteq K_n$ completes the inductive step.

By compactness, select s with

$$s \in \bigcap_{m \in \mathbb{N}} K_m \subseteq K_{n+1} = K_n \cap K_n t_{m(n)} \quad (n \in \mathbb{N});$$

choosing $k_n \in K_n \subseteq K$ with $s = k_n t_{m(n)}$ gives $s \in K_0 \subseteq E$, and

$$s z_{m(n)} = s t_{m(n)}^{-1} = k_n \in K_n \subseteq K_0 \subseteq E.$$

Finally take $\mathbb{M} := \{m(n) : n \in \mathbb{N}\}$.

As for the final assertion, we follow the idea of the Generic Completeness Principle [BinO1, Th. 3.4] (but with $\mathcal{U}(G)$ for $\mathcal{B}a$ there): define

$$F(H) := \bigcap_{n \in \mathbb{N}} \bigcup_{m > n} H \cap H t_m \quad (H \in \mathcal{U}(G));$$

then $F : \mathcal{U}(G) \rightarrow \mathcal{U}(G)$ and F is monotone ($F(S) \subseteq F(T)$ for $S \subseteq T$); moreover, $s \in F(H)$ iff $s \in H$ and $s z_m \in H$ for infinitely many m . We are to show that $E_0 := E \setminus F(E)$ is left Haar null. Suppose otherwise. Then renaming g and K_0 as necessary, w.l.o.g. both $\mu(E_0) > 0$ and $K_0 \subseteq E_0$ (and $\mu(K_0) > 0$). But then, as above, $\emptyset \neq F(K_0) \cap K_0 \subseteq F(E) \cap E_0$, a contradiction, since $F(E) \cap E_0 = \emptyset$. \square

Remark. In the setting of Th. 3 any non-empty open set U is not left Haar null (as $\{dU : D \in D\}$ with D countable dense covers G), hence neither is

$U \setminus H$ for $H \in \mathcal{HN}$. So the (Hashimoto ideal) topology generated by such sets includes \mathcal{HN} among its negligible sets.

Corollary 3. *For G Polish and amenable at 1_G and z_n null, there is $\mu \in \mathcal{P}(G)$ such that for $K \in \mathcal{K}_+(\mu)$*

$$K \cap Kz_m^{-1} \in \mathcal{M}_+(\mu) \quad \text{for infinitely many } m \in \mathbb{N},$$

iff for μ -quasi all $s \in K$ there is an infinite $\mathbb{M} \subseteq \mathbb{N}$ with

$$\{sz_m : m \in \mathbb{M}\} \subseteq K.$$

Proof. We will refer to the function F of the preceding proof. First proceed as in the proof of Th. 3 above, taking $t_n := z_n^{-1}$ and $g = 1_G$ (so that $\mu = \sigma$). Fix K with $\mu(K) > 0$. For the forward direction, continue as in the proof of Th. 3 with $K_0 = K$ and observe that the proof above needs only that $s_{k(n)} \in K_n^{-1}K_n$ occurs infinitely often whenever $\mu(K_n) > 0$. This yields the desired conclusion that $\mu(K \setminus F(K)) = 0$. For the converse direction, suppose that $\mu(F(K)) > 0$. Since for each $n \in \mathbb{N}$

$$F(K) \subseteq \bigcup_{m>n} K \cap Kt_m,$$

we have $\mu(K \cap Kt_m) > 0$ for some $m > n$; so

$$K \cap Kt_m \in \mathcal{M}_+(\mu) \quad \text{for infinitely many } m. \quad \square$$

Remark. With E as in the Shift-compactness Theorem, if $z_n \in B_{1/n} \setminus E^{-1}E$, then z_n is null; so, for some $s \in E$, $sz_m \in E$ for infinitely many m . Then, for any such m ,

$$z_m \in E^{-1}E,$$

contradicting the choice of z_m . So $1_G \in \text{int}(E^{-1}E)$, i.e. E has the Steinhaus-Weil property, as before.

The following sharpens a result due (for Lebesgue measure on \mathbb{R}) to Mospan [Mos] by providing the converse below; it is antithetical to Lemma 1 (and so to Theorem 3).

Proposition 6 (Mospan property). *For G a metric group and compact K , if $1_G \notin \text{int}(K^{-1}K)$, then $\mu_-(K) = 0$, i.e. μ is maximally discontinuous; equivalently, there is a ‘null sequence’ $t_n \rightarrow 1_G$ with $\lim_n \mu(Kt_n) = 0$.*

Conversely, if $\mu(K) > \mu_-(K) = 0$, then there is a null sequence $t_n \rightarrow 1_G$ with $\lim_n \mu(Kt_n) = 0$, and there is a compact $C \subseteq K$ with $\mu(K \setminus C) = 0$ with $1_G \notin \text{int}(C^{-1}C)$.

Proof. The first assertion follows from Lemma 1. For the converse, as in [Mos]: suppose that $\mu(Kt_n) = 0$, for some sequence $t_n \rightarrow 1_G$. By passing to a subsequence, we may assume that $\mu(Kt_n) < 2^{-n-1}$. Put $D_m := K \setminus \bigcap_{n \geq m} Kt_n \subseteq K$; then $\mu(K \setminus D_m) \leq \sum_{n \geq m} \mu(Kt_n) < 2^{-m}$, so $\mu(D_m) > 0$ provided $2^{-m} < \mu(K)$. Now choose compact $C_m \subseteq D_m$, with $\mu(D_m \setminus C_m) < 2^{-m}$. So $\mu(K \setminus C_m) < 2^{1-m}$. Also $C_m \cap C_m t_n = \emptyset$, for each $n \geq m$, as $C_m \subseteq K$; but $t_n \rightarrow 1_G$, so the compact set $C_m^{-1}C_m$ contains no interior points. Hence, by Baire's theorem, neither does $C^{-1}C$, since $C = \bigcup_m C_m$, which differs from K by a null set. \square

Proposition 7. *A (regular) Borel measure μ on a locally compact metric topological group G has the Steinhaus-Weil property iff*

- (i) *for each non-null compact set K , the map $m_K : t \rightarrow \mu(Kt)$ is subcontinuous at 1_G ;*
- (ii) *for each non-null compact set K there is no 'null' sequence $t_n \rightarrow 1_G$ with $\mu(Kt_n) \rightarrow 0$.*

Remark. This is immediate from Prop. 6 (cf. [Mos]).

We now prove a strengthening of the Subcontinuity Theorem obtained by assuming a 'concentration property'. That this property holds in an abelian Polish group emerges from an inspection of Solecki's proof of his theorem that an abelian Polish group is amenable at 1.

Definitions. Say that a null sequence \mathbf{t} is *regular* if \mathbf{t} is non-trivial, $\|t_k\|$ is non-increasing, and

$$\|t_k\| \leq r(k) := 1 / [2^k(k+1)] \quad (k \in \mathbb{N}).$$

For regular \mathbf{t} , put $\mu_k = \mu_k(\mathbf{t}) := 2^{k-1} \sum_{m \geq k} 2^{-m} (\delta_{t_m^{-1}} + \delta_{t_m}) = \frac{1}{2} \delta_{t_k^{-1}} + \frac{1}{4} \delta_{t_{k+1}^{-1}} + \dots$. Then $\mu_k(B_{r(k)}) = 1$ for $k \in \mathbb{N}$. Merging \mathbf{t}^{-1} with \mathbf{t} by alternation of terms if necessary, it is now convenient to assume that \mathbf{t} contains as successive pairs inverses of its terms. So, if $\nu_k \ll \mu_k$, then

$$\nu_k := \sum_{m \geq k} a_{km} \delta_{t_m^{-1}},$$

for some non-negative sequence $\mathbf{a}_k := \{a_{kk}, a_{k,k+1}, a_{k,k+2}, \dots\}$ of unit ℓ_1 -norm. Say that $\{\mathbf{a}_k\}$ has the *concentration property* if for some index j and some $\alpha > 0$

$$a_{k,k+j} \geq \alpha > 0 \quad \text{for all large } k;$$

then say that the sequence $\{\nu_k\}$ has the *concentration property*. (This will fail if \mathbf{a}_k has $a_{k,k+k} = 1$, which concentrates measure in an unbounded fashion.)

Definition. Say that a group G is *strongly amenable at 1* if G is amenable at 1, and for each *regular* \mathbf{t} a Solecki measure $\sigma(\mathbf{t})$ has associated measures $\sigma_k(\mathbf{t}) \ll \mu_k(\mathbf{t})$ with the concentration property.

Theorem 4 (Strong amenability at 1, after [Sol2, Prop. 3.3(i)]). *Any abelian Polish group G is strongly amenable at 1.*

Proof. This follows the construction in [Sol2] of the reference measure in the case of $\mu_k(\mathbf{t})$ above. First define the normalized restriction

$$\sigma_k := \mu_k|_{B_{r(k)}} / \mu_k(B_{r(k)})$$

and then set

$$\sigma := *_{k=1}^{\infty} \rho_k \quad \text{for } \rho_k := \frac{1}{k+1} \sum_{i=0}^k \sigma_k^i.$$

(Convolution powers intended here.) Then the argument in [Sol2] shows that $\sigma_k * \sigma(K) \rightarrow \sigma(K)$ (for K compact). However, as \mathbf{t} is regular, $\mu_k \equiv \sigma_k$. But $a_{kk} = 1/2$ (all k), so the measures σ_k here have the concentration property. \square

Definitions (Sequence and measure symmetrization):

1. Merging \mathbf{t}^{-1} with \mathbf{t} by alternation of terms yields the regular sequence $\mathbf{s} = (s_1, s_2, \dots) := (t_1, t_1^{-1}, t_2, t_2^{-1}, \dots)$; we term this the **symmetrized sequence** of \mathbf{t} . (It is ‘symmetric’ in the sense only that $\|s_{2k-1}\| = \|s_{2k}\|$.)
2. For odd k , as $(\mu_k(\mathbf{t}) + \mu_k(\mathbf{t}^{-1}))/2$ is symmetric as a measure, taking $\sigma_{2k}(\mathbf{s}) = \sigma_{2k-1}(\mathbf{s}) := (\mu_k(\mathbf{t}) + \mu_k(\mathbf{t}^{-1}))/2$ in lieu of $\mu_k(\mathbf{t})$ above yields each ρ_k symmetric. So, in the abelian context of Theorem 4 above, the limiting convolution σ is a *symmetric Solecki measure* $\sigma(\mathbf{t})$.

Remark. Performing the symmetrization of the Definition above gives in the proof of Theorem 4 above that $a_{2k-1,2k-1} = a_{2k-1,2k} = a_{2k,2k} = a_{2k,2k+1} = 1/4$, which presents *simultaneous concentration* along \mathbf{t} and \mathbf{t}^{-1} .

We now re-run the proof of Theorem 1 with improved estimates to yield:

Theorem 1_S (Strong Subcontinuity Theorem). *For G a Polish group that is strongly amenable at 1, if \mathbf{t} is regular and $\sigma = \sigma(\mathbf{t})$ is a Solecki measure – then for $K \in \mathcal{K}_+(\sigma)$*

$$\sigma(K) = \lim_n \sigma(Kt_n) = \sigma_-^{\mathbf{t}}(K).$$

Likewise, passing to the symmetrized sequence of \mathbf{t} as above and to a symmetric Solecki measure $\sigma(\mathbf{t})$ with the simultaneous concentration property (for \mathbf{t} and \mathbf{t}^{-1}) corresponding to an abelian context:

$$\sigma(K) = \lim_n \sigma(t_n K).$$

Proof. Fix \mathbf{t} and a corresponding Solecki reference measure $\nu(\mathbf{t})$ and its associated sequence ν_k , which as in Th. 4 has the concentration property. Write $\sigma_k := \sum_{m \geq k} a_{km} \delta_{t_m^{-1}}$; as σ_k has the concentration property, there are n_0, j and $\alpha > 0$ with

$$a_{kk+j} \geq \alpha > 0 \quad (k \geq n_0).$$

Now fix K compact with $\sigma(K) > 0$ and $\varepsilon > 0$. Put

$$\delta := \varepsilon / \left(\frac{2}{\alpha} - 1 \right) > 0,$$

as $\alpha \leq 1$. Then, by upper semicontinuity and by ‘amenability at 1’ (i.e. $\sigma_k * \sigma(K) \rightarrow \sigma(K)$), there is $n_1 = n_1(\varepsilon, K) > n_0$ with

$$\sigma(Kt_k) \leq \sigma(K) + \delta \text{ and } \sigma_k * \sigma(K) \geq \sigma(K) - \delta \quad (k \geq n_1).$$

So (by upper semicontinuity) for $k \geq n_1$

$$\sum_{m \geq k, m \neq k+j} a_{km} \sigma(Kt_m) \leq \sum_{m \geq k, m \neq k+j} a_{km} (\sigma(K) + \delta) = (\sigma(K) + \delta)(1 - a_{kk+j}).$$

Also (by ‘amenability at 1’) for $k \geq n_1$

$$\begin{aligned} a_{kk+j} \sigma(Kt_{k+j}) &\geq \sigma(K) - \delta - \sum_{m \geq k, m \neq k+j} a_{km} \sigma(Kt_m) \\ &\geq \sigma(K) + \delta - 2\delta - (\sigma(K) + \delta)(1 - a_{kk+j}) \\ &= a_{kk+j} \sigma(K) - \delta(2 - a_{kk+j}). \end{aligned}$$

So for $m = k + j > n_1 + j$

$$\sigma(Kt_m) = \sigma(Kt_{k+j}) \geq \sigma(K) - \delta \left(\frac{2}{a_{kk+j}} - 1 \right) \geq \sigma(K) - \delta \left(\frac{2}{\alpha} - 1 \right) = \sigma(K) - \varepsilon.$$

As for the final assertion concerning symmetrization, note that $\sigma(t_n K) = \sigma(K^{-1}t_n^{-1}) \rightarrow \sigma(K^{-1}) = \sigma(K)$, by symmetry of σ . \square

We note an immediate corollary, needed in §4.

Corollary 4. *For G, \mathbf{t} and σ as in Th. 1_S above, and $K, H \in \mathcal{K}_+(\sigma), \delta > 0$: if $0 < \Delta < \sigma(K)$ and $0 < D < \sigma(H)$, then there is n with*

$$B_\delta^{K\Delta} \cap B_\delta^{HD} \supseteq \{t_m : m \geq n\}.$$

Proof. Take $\varepsilon := \min\{\sigma(K) - \Delta, \sigma(H) - D\} > 0$. As $K, H \in \mathcal{K}_+(\sigma)$, there is n such that $\|t_m\| < \delta$ for $m \geq n$ and

$$\sigma(Kt_m) \geq \sigma(K) - \varepsilon \geq \Delta, \quad \sigma(Ht_m) \geq \sigma(H) - \varepsilon \geq D \quad (m \geq n). \quad \square$$

To accommodate varying sided-ness conventions, we close with the left-handed version of Theorem 1, in which $\tilde{\nu}_-^s(K)$ (defined at the start of the section) replaces $\nu_-^s(K)$.

Theorem 1_L (Left Subcontinuity Theorem, after Solecki [Sol2, Th. 1(ii)]). *For G amenable at 1_G , and \mathbf{t} a null sequence, there is $\sigma = \sigma_L(\mathbf{t}) \in \mathcal{P}(G)$ such that for each $K \in \mathcal{K}_+(\sigma)$ there is a subsequence $\mathbf{s} = \mathbf{s}(K) := \{t_{m(n)}\}$ with*

$$\lim_n \sigma(t_{m(n)}K) = \sigma(K) \quad (n \in \mathbb{N}), \text{ so } \quad \tilde{\sigma}_-^s(K) > 0.$$

Proof. From Theorem 1 with $\sigma = \sigma(\mathbf{t})$, recalling that $\tilde{\sigma}(E) := \sigma(E^{-1})$ so that $\sigma(E) = \tilde{\sigma}(E^{-1})$, there is $\{t_{m(n)}\}$ with

$$\tilde{\sigma}(t_{m(n)}^{-1}K^{-1}) > (1 - 2^{-n})\tilde{\sigma}(K^{-1}) \quad (n \in \mathbb{N}), \text{ so } \quad \lim_n \tilde{\sigma}(t_{m(n)}^{-1}K^{-1}) > 0.$$

Since K^{-1} is compact and $t_{m(n)}^{-1}$ is null, replace K^{-1} by K , $t_{m(n)}^{-1}$ by $t_{m(n)}$; then with μ for $\tilde{\sigma}$

$$\mu(t_{m(n)}K) > (1 - 2^{-n})\mu(K) \quad (n \in \mathbb{N}), \text{ so } \quad \lim_n \mu(t_{m(n)}K) > 0. \quad \square$$

Część II

Around the Simmons-Mospan Theorem

We saw in Part I that measure may exhibit selective continuity under translation in the presence of amenability at 1. Here we consider obstacles/obstructions to continuity.

3 A Lebesgue Decomposition

We begin with definitions isolating left-handed components in Christensen's notion of Haar null sets [Chr1], and Solecki's left Haar null sets [Sol2]; right-handed versions have analogous properties. As far as we are aware the component notions in parts (ii)-(iv) below have not been previously studied. Below G is a Polish group.

Definition. (i) **Left μ -null:** For $\mu \in \mathcal{M}(G)$, say that N is *left μ -null* ($N \in \mathcal{M}_0^L(\mu)$) if it is contained in a universally measurable set B such that

$$\mu(gB) = 0 \quad (g \in G).$$

Thus a set S is *left Haar null* ([Sol3] after [Chr1]) if it is contained in a universally measurable set B that is left μ -null for some $\mu \in \mathcal{M}(G)$.

(ii) **Left μ -inversion:** For $\mu \in \mathcal{M}(G)$, say that $N \in \mathcal{M}_0^L(\mu)$ is *left invertibly μ -null* ($N \in \mathcal{M}_0^{L\text{-inv}}(\mu)$) if

$$N^{-1} \in \mathcal{M}_0^L(\mu),$$

so that N^{-1} is contained in a universally measurable set B^{-1} such that

$$\mu(gB^{-1}) = 0 \quad (g \in G).$$

(iii) **Left μ -absolute continuity:** For $\mu, \nu \in \mathcal{M}(G)$, ν is *left absolutely continuous* w.r.t. μ ($\nu <^L \mu$) if $\nu(N) = 0$ for each $N \in \mathcal{M}_0^L(\mu)$, and likewise for the invertibility version: $\nu <^{L\text{-inv}} \mu$.

(iv) **Left μ -singularity:** For $\mu, \nu \in \mathcal{M}(G)$, ν is *left singular* w.r.t. μ (on B) ($\nu \perp^L \mu$ (on B)) if B is a support of ν and B is in $\mathcal{M}_0^L(\mu)$, and likewise $\nu \perp^{L\text{-inv}} \mu$.

Remark. For μ symmetric, since

$${}_{g^{-1}}\mu(B) = \mu_g(B^{-1})$$

if B is left μ -null we may conclude only that B^{-1} is *right* μ -null. The ‘inversion property’, property (ii) above, is thus quite strong (though obvious in the abelian case).

Notice that each of $\mathcal{M}_0^L(\mu)$ and $\mathcal{M}_0^{L\text{-inv}}(\mu)$ forms a σ -algebra (since $g \cup_n B = \cup_n gB$ and $g(\cup_n B)^{-1} = \cup_n gB^{-1}$). This implies the following left versions of the Lebesgue Decomposition Theorem (we need the second one below). The ‘pedestrian’ proof below demonstrates that the Principle of Dependent Choice (DC) suffices, a further example that ‘positive’ results in measure theory follow from DC (as Solovay points out in [Solo, p. 31]).

Theorem LD. *For G a Polish group, $\mu, \nu \in \mathcal{M}(G)$, there are $\nu_a, \nu_s \in \mathcal{M}(G)$ with*

$$\nu = \nu_a + \nu_s \text{ with } \nu_a <^L \mu \text{ and } \nu_s \perp^L \mu,$$

and likewise, there are $\nu'_a, \nu'_s \in \mathcal{M}(G)$ with

$$\nu = \nu'_a + \nu'_s \text{ with } \nu_a <^{L\text{-inv}} \mu \text{ and } \nu_s \perp^{L\text{-inv}} \mu.$$

Proof. As the proof depends on σ -additivity, it will suffice to check the ‘L’ case. Write $G = \cup_n G_n$ with the G_n disjoint, universally measurable, and with each $\nu(G_n)$ finite (say, with all but one term σ -compact, and their complement ν -null). Put $s_n = \sup\{\nu(E) : E \subseteq G_n, E \in \mathcal{M}_0^L(\mu)\}$. In $\mathcal{M}_0^L(\mu)$, for each n with $s_n > 0$, choose $E_{n,m} \subseteq G_n$ with $\nu(E_{n,m}) > s_n - 1/m$, and put $B_n := \cup_m E_{n,m} \subseteq G_n$. Then the sets B_n are disjoint and lie in $\mathcal{M}_0^L(\mu)$, as does also $B := \cup_n B_n$; moreover $\nu(G_n \setminus B_n) = 0$ for each n . Put $A := G \setminus B$. Then $\nu(M) = 0$ for $M \in \mathcal{M}_0^L(\mu)$ with $M \subseteq A$, since $A = \cup_n (G_n \setminus B_n)$. So $\nu_a := \nu|_A <^L \mu$, and $\nu_s := \nu|_B \perp^L \mu$, since $B \in \mathcal{M}_0^L(\mu)$. \square

Remark. The above rests on DC; a simpler argument rests on maximality: choose a maximal disjoint family \mathcal{B} of universally measurable sets $M \in \mathcal{M}_0^L(\mu)$ with finite positive $\nu(M)$; then, their union $B \in \mathcal{M}_0^L(\mu)$ (as \mathcal{B} would be countable, by the σ -finiteness of ν).

4 Discontinuity: the Simmons-Mospan Theorem

It is convenient to begin by repeating the gist of the Simmons-Mospan argument here, as it is short, despite its ‘near perfect disguise’, to paraphrase a phrase from Loomis [Loo, p. 85]. The result follows from their use of Fubini’s Theorem and the Lebesgue decomposition theorem of §3, but here we stress the dependence on Theorem FN and on left μ -inversion. We revert to the Weil left-sided convention and associated KK^{-1} usage.

Proposition 8 (Local almost nullity). *For G a Polish group, $\mu \in \mathcal{M}(G)$, V open and K compact with $K \in \mathcal{M}_0^{\text{L-inv}}(\mu)$, i.e. $K, K^{-1} \in \mathcal{M}_0^{\text{L}}(\mu)$:
– for any $\nu \in \mathcal{M}(G)$, $\nu(tK) = 0$ for μ -almost all $t \in V$, and likewise $\nu(Kt) = 0$.*

Proof. For ν invertibly μ -absolutely continuous (as in §3 above), the conclusion is immediate; for general ν this will follow from Theorem LD (§3), once we have proved the corresponding singular version of the assertion: that is the nub of the proof.

Thus, suppose that $\nu \perp^{\text{L-inv}} \mu$ on K . For $t \in V$ let $t = uw$ be any expression for t as a group product of $u, v \in G$, and note that $\mu(uK^{-1}) = 0$, as $K^{-1} \in \mathcal{M}_0^{\text{L}}(\mu)$. Let H be the set

$$\bigcup_{t \in V} (\{t\} \times tK),$$

here viewed as a union of vertical t -sections. We next express it as a union of u -horizontal sections and apply the Fubini Null Theorem (Th. FN, §1).

Since $u = tk = uwk$ is equivalent to $w = k^{-1}$, the u -horizontal sections of H may now be rewritten, eliminating t , as

$$\{(t, u) : uw = t \in V, u \in tK = uwK\} = \{(uw, u) : uw \in V, uw \in uK^{-1}\}.$$

So H may now be viewed as a union of u -horizontal sections as

$$\bigcup_{u \in G} (V \cap (uK^{-1})) \times \{u\},$$

all of these u -horizontal sections being μ -null. By Th. FN, μ -almost all vertical t -sections of H for $t \in V$ are ν -null. As the assumptions on K are symmetric the right-sided version follows. \square

The result here brings to mind the Dodos Dichotomy Theorem [Dod1, Th. A] for *abelian* Polish groups G : if an analytic set A is witnessed as Haar-null under one measure $\mu \in \mathcal{P}(G)$, then either A is Haar-null for quasi all $\nu \in \mathcal{P}(G)$ or else it is not Haar-null for quasi all such ν , i.e. if $A \in \mathcal{M}_0(\mu)$ (omitting the unnecessary superscript L), then either $A \in \mathcal{M}_0(\nu)$ for quasi all such ν , or $A \notin \mathcal{M}_0(\nu)$ for quasi all such ν (w.r.t. the Prokhorov-Lévy metric in $\mathcal{P}(G)$ [Dud, 11.3, cf. 9.2]). Indeed, [Dod2, Prop. 5] when A is σ -compact A is Haar-null for quasi all $\nu \in \mathcal{P}(G)$. The result is also reminiscent of [Amb, Lemma 1.1].

Before stating the Simmons-Mospan specialization to the Haar context and also to motivate one of the conditions in its subsequent generalizations, we cite (and give a direct proof) of the following known result (equivalence of Haar measure η and its inverse $\tilde{\eta}$), encapsulated in the formula

$$\eta(K^{-1}) = \int_K d\eta(t)/\Delta(t),$$

exhibiting the direct connection between η and $\tilde{\eta}$ via the modular function [HewR, 15.14], or [Hal, §60.5f]; this equivalence result holds more generally between any two probability measures when one is left and the other right quasi-invariant – see [Xia, Cor. 3.1.4]; this is related to a theorem of Mackey’s [Mac] cf. §9.16. As will be seen from the proof, in Lemma H below there is no need to assume the group is separable, a compact metrizable subspace (being totally bounded) is separable.

Lemma H (cf. [Hal, §50(ff); §59 Th. D]). *In a locally compact metrizable group G , for K compact, if $\eta(K) = 0$, then $\eta(K^{-1}) = 0$, and, by regularity, so also for K measurable.*

Proof. Fix a compact K . As K is compact, Δ (the modular function of G) is bounded away from 0 on K , say by $M > 0$; furthermore, K is separable, so pick $\{d_n : n \in \mathbb{N}\}$ dense in K . Then for any $\varepsilon > 0$ there are two (finite) sequences $m(1), \dots, m(n) \in \mathbb{N}$ and $\delta(1), \dots, \delta(n) > 0$ such that $\{B_{\delta(i)}d_{m(i)} : i \leq n\}$ covers K and

$$M \sum_{i \leq n} \eta(B_{\delta(i)}) \leq \sum_{i \leq n} \eta(B_{\delta(i)}) \Delta(d_{m(i)}) = \sum_{i \leq n} \eta(B_{\delta(n)}d_{m(n)}) < \varepsilon.$$

Then

$$\sum_{i \leq n} \eta(d_{m(i)}^{-1}B_{\delta(i)}) = \sum_{i \leq n} \eta(B_{\delta(i)}) \leq \varepsilon/M.$$

But $\{d_{m(i)}^{-1}B_{\delta(i)} : i \leq n\}$ covers K^{-1} by the symmetry of the balls B_{δ} (by the symmetry of the norm); so, as $\varepsilon > 0$ is arbitrary, $\eta(K^{-1}) = 0$.

As for the final assertion, if $\eta(E^{-1}) > 0$ for some measurable E , then $\eta(K^{-1}) > 0$ for some compact $K^{-1} \subseteq E^{-1}$, by regularity; then $\eta(K) > 0$, and so $\eta(E) > 0$. \square

Proposition 8 and Lemma H immediately give:

Theorem S (cf. [Sak, III.11], [Mos], [BarFF, Th. 7]). *For G locally compact with left Haar measure η and ν a Borel measure on G , if the set S is η -null, then for η -almost all t*

$$\nu(tS) = 0.$$

In particular, this is so for S the support of a measure ν singular with respect to η .

This in turn allows us to prove the locally compact (separable) case of the Simmons-Mospan Theorem, as stated in the Introduction. We then pursue a non-locally compact variant.

Proof of Theorem SM. If μ is absolutely continuous w.r.t. Haar measure η , then μ , being invariant, is subcontinuous, and Lemma 1 (in §2) gives the Steinhaus-Weil property. Otherwise, decomposing μ into its singular and absolutely continuous parts w.r.t. η , choose K a compact subset of the support of the singular part of μ ; then $\mu(K) > \mu_-(K) = 0$, by Prop. SM above, and so Prop. 6 (Converse part – see §2) applies. \square

Proposition 9 (after Simmons, cf. [Sim, Lemma] and [BarFF, Th. 8]). *For G a Polish group, $\mu, \nu \in \mathcal{M}(G)$ and $\nu \perp^{\text{L-inv}} \mu$ concentrated on a compact invertibly μ -null set K , there is $B \subseteq K$ such that $K \setminus B$ is ν -null and both BB^{-1} and $B^{-1}B$ have empty interior.*

Proof. As we are concerned only with the subspace $KK^{-1} \cup K^{-1}K$, w.l.o.g. the group G is separable. By Prop. 8, $Z := \{x : \nu(xK) = 0\}$ is dense and so also

$$Z_1 := \{x : \nu(K \cap xK) = 0\},$$

since $\nu(K \cap xK) \leq \nu(xK) = 0$, so that $Z \subseteq Z_1$. Take a denumerable dense set $D \subseteq Z_1$ and put

$$S := \bigcup_{d \in D} K \cap dK.$$

Then $\nu(S) = 0$. Take $B = K \setminus S$. If $\emptyset \neq V \subseteq BB^{-1}$ and $d \in D \cap V$, then for some $b_1, b_2 \in B \subseteq K$

$$d = b_1 b_2^{-1} : \quad b_1 = db_2 \in K \cap dK \subseteq S,$$

a contradiction, since $B \cap S = \emptyset$. So $(K \setminus S)(K \setminus S)^{-1}$ has empty interior. A similar argument based on

$$T := \bigcup_{d \in D} Kd \cap K$$

ensures that also $(K \setminus S \setminus T)^{-1}(K \setminus S \setminus T)$ has empty interior. \square

In order to generalize the Simmons Theorem from its locally compact context we will need to cite the following result. Here $\mathbb{Q}_+ := \mathbb{Q} \cap (0, \infty)$ denotes the positive rationals, and $B_\delta^{K, \Delta}(\sigma) := \{z \in B_\delta : \sigma(Kz) > \Delta\}$ as in §2.

Theorem 5 (Disaggregation Theorem, [BinO8, Th.6, Prop. 4]). *Let G be a Polish group that is strongly amenable at 1, and let \mathbf{t} be a regular null sequence. For $\sigma = \sigma(\mathbf{t})$ there are a countable family \mathcal{H} with $\mathcal{H} \subseteq \mathcal{K}_+(\sigma)$, a countable set $D = D(\mathcal{H}) \subseteq G$ dense in G , and a dense subset $G(\sigma)$ on which the sets below are the sub-basic sets of a metrizable topology:*

$$B_\delta^{K, \Delta}(\sigma) \quad (K \in \mathcal{H}, \delta, \Delta \in \mathbb{Q}_+, \Delta < \sigma(K)).$$

In particular, the space $G(\sigma)$ is continuously and compactly embedded in G . Moreover, each such open set contains a cofinal subsequence of \mathbf{t} .

For a proof we refer the reader to the companion paper [BinO8]; the result relies on Corollary 4 above. The subspace $G(\sigma)$ here is a topological analogue of the Cameron-Martin subspace $H(\gamma)$ of a locally convex topological vector space equipped with a Radon Gaussian measure γ – see §8.2-3.

We are now ready for the promised generalization. This requires equivalence of the Solecki measure and its inverse – valid at least in Polish abelian groups (see Th. 4 of §2 on strong amenability at 1).

Theorem 6 (Generalized Simmons Theorem, cf. [Sim, Th. 2]). *Let G be a Polish group that is strongly amenable at 1 (e.g. if G is abelian), let $\sigma = \sigma(\mathbf{t})$ be a Solecki measure corresponding to a regular null sequence \mathbf{t} , which we assume is equivalent to its inverse $\tilde{\sigma}$ (e.g. if G is abelian), and let*

$G(\sigma)$ be the dense subgroup of the preceding theorem. Then:
 $\nu \in \mathcal{M}(G)$ is left invertibly-singular w.r.t. σ iff ν has a support that is a σ -compact union of compact sets K_n with each of the compact sets $K_n K_n^{-1}$ and $K_n^{-1} K_n$ nowhere dense (equivalently: having empty interior) in the topology of the subgroup $G(\sigma)$, as above.

Proof. Suppose that $\mu \in \mathcal{M}(G)$ is subcontinuous. If $\nu \in \mathcal{M}(G)$, by Theorem LD write

$$\nu = \nu_a + \nu_s \text{ with } \nu_a <^{\text{L-inv}} \mu \text{ and } \nu_s \perp^{\text{L-inv}} \mu.$$

If ν is concentrated as in the statement of the theorem on a σ -compact set B with $B^{-1}B$ having empty interior in $G(\sigma)$, then $\nu_a = 0$, and so ν is left invertibly-singular w.r.t. μ . Indeed, as ν is concentrated on B , so is ν_a . We claim that $\nu_a(B) = 0$. Otherwise, $\nu_a(K_n) > 0$ for some compact $K_n \subseteq B$. So $K = K_n \notin \mathcal{M}_0^{\text{L-inv}}(\sigma)$, as $\nu_a <^{\text{L-inv}} \sigma$. The argument now splits into two cases, according as $K \notin \mathcal{M}_0^{\text{L}}(\sigma)$ or $K^{-1} \notin \mathcal{M}_0^{\text{L}}(\sigma)$

First, suppose that $\sigma(gK) > 0$ for some $g \in G$; then, by Lemma 1, there are $\delta > 0$ and $0 < \Delta < \sigma_-^{\text{t}}(gK)$ with

$$B_\delta^{gK\Delta}(\sigma) \subseteq (gK)^{-1}gK = K^{-1}K \subseteq B^{-1}B,$$

contradicting the above property of B .

Next, suppose that $\sigma(Kg) = \sigma^{-1}(g^{-1}K^{-1}) > 0$ for some $g \in G$; so $\sigma(g^{-1}K^{-1}) > 0$, as $\tilde{\sigma}$ is equivalent to σ . Then, again by Lemma 1, there are $\delta > 0$ and $0 < \Delta < \sigma_-^{\text{t}}(g^{-1}K^{-1})$ with

$$B_\delta^{g^{-1}K^{-1}\Delta}(\sigma) \subseteq (g^{-1}K^{-1})^{-1}g^{-1}K^{-1} = KK^{-1} \subseteq BB^{-1},$$

again contradicting the above property of B . So $\nu = \nu_a$ is invertibly singular w.r.t. μ .

The rest of the proof is as in Simmons [Sim, Th. 2], using Prop. 6: the Baire-category argument still holds, since compactness implies closure under the $G(\sigma(\mathbf{t}))$ -topology, the latter being a finer topology; avoidance of interior points requires second countability, assured by Th. 5. \square

Corollary 5 (Simmons Theorem: [Sim, Th. 2]). *For G separable and locally compact and η left Haar measure:*

$\nu \in \mathcal{M}(G)$ is singular w.r.t. η iff ν has a support that is a σ -compact union of compact sets K_n with each of the compact sets $K_n K_n^{-1}$ nowhere dense (equivalently: having empty interior).

For the non-separable version of the above, see §8.1.

Część III

Weil topologies

5 Weil-like topologies: preliminaries

We turn now to relatives of the *Weil topology*. For background, we refer to Weil's book [Wei, Ch. VII] and Halmos's book [Hal, Ch. XII] (see also §8.4). Weil regarded his result as a *Converse Haar Theorem*, in retrieving the topological-group structure from the measure-algebra structure [Fre] as encoded by the Haar-measurable subsets – cf. [Kod]. (Here one may work either, following Weil, to within a dense embedding in a locally compact group, as in the Remark to Theorem 7M below, or, following Mackey, uniquely up to homeomorphism, granted the further assumption of an analytic Borel structure [Mac, Th. 7.1]; for further information see §8.16.) The alternative view below throws light on this result in that the measure structure is already encoded by the density topology \mathcal{D} via the Haar density theorem, for which see [Mue], [Hal, §61(5), p. 268], cf. [BinO2, §7; Th. 6.10], [BinO5]; this view is partially implicit in [Amb], where refinement of one invariant measure μ_1 by another μ_2 holds when sets in $\mathcal{M}_+(\mu_2)$ contain sets in $\mathcal{M}_+(\mu_1)$ (as in the refinement of one topology by another). This falls within the broader aim of retrieving a *topological* group structure from a given (one-sidedly) invariant topology τ on a group G , when τ arises from refinement of a topological group structure (i.e. starting from a *semitopological* group structure (G, τ)). Also relevant here are *Converse Steinhaus-Weil* results, as in Prop. 7 of §3 above (see also §8.4). For background on group-norms see the textbook treatment in [ArhT, §3.3] (who trace this notion back to Markov) or [BinO2], but note their use of ‘pre-norm’ for what we call (following Pettis [Pet]) a *pseudo-norm*; for quasi-interiors and regular open sets see §8.6. Thus a norm $\|\cdot\| : G \rightarrow [0, \infty)$ satisfies all the three conditions 1-3 below and generates a right-invariant metric $d(x, y) = \|xy^{-1}\|$ and so a topology $\mathcal{T} = \mathcal{T}_d$, just as a right-invariant metric d derives from a separable topology \mathcal{T}_G and generates, via the Birkhoff-Kakutani Theorem ([HewR, Th. 8.3], [Gao, Th. 2.1.1]), the norm $\|x\| = d(x, 1_G)$. A pseudo-norm differs in possibly lacking condition

1.i. (so generates a *pseudo-metric*).

1.i (positivity): $\|g\| > 0$ for $g \neq 1_G$, and 1.ii: $\|1_G\| = 0$;

2 (subadditivity): $\|gh\| \leq \|g\| + \|h\|$,

3 (symmetry): $\|g^{-1}\| = \|g\|$.

Recall that a set function λ defined on $\mathcal{U}(G)$ is a *submeasure* if it is monotone and subadditive with $\lambda(\emptyset) = 0$ (Introduction, [Fre, Ch. 39, §392], [Tal]); by analogy with the term *finitely additive measure* (for background see [Bin], [TomW, Ch. 12]; cf. [Pat]), this is a *finitely subadditive outer measure*, similarly as in Maharam [Mah], albeit in the context of Boolean algebras, but without her positivity condition. Recall from Halmos [Hal, Ch. II §10] that a submeasure is an *outer measure* if in addition it is *countably subadditive*. The set function λ is *left invariant* if $\lambda(gE) = \lambda(E)$ for all $g \in G$ and $E \in \mathcal{U}(G)$.

Propositions 10 and 11 below are motivated by [Hal, Ch. XII §62, cf. Ch. II §9 (2-4)], where G is a locally compact group with λ its left Haar measure, but here the context is broader, allowing in *amenable* groups G (cf. [TomW, Ch. 12], [Pat]). The two results enable the introduction in §6 of Weil-like topologies generated from families of left-invariant *pseudo-metrics* derived from invariant submeasures. The latter rely on the natural *measure-metric*, also known as the *Fréchet-Nikodym metric* ([Fre, §323Ad], [Hal, §40 Th. A], [Bog2, p. 53, 102-3, 418]); see [Drew1,2] (cf. [Web]) for the related literature of Fréchet-Nikodym topologies and their relation to the Vitali-Hahn-Saks Theorem. Maharam [Mah] studies sequential continuity of the order relation (of inclusion, here in the measure algebra), and requires positivity to obtain a (measure-) *metric*; see Talagrand [Tal] (cf. [Fre, §394] and the literature cited there) for a discussion of pathological submeasures (the only measures they dominates under \ll being trivial), and [ChrH] for corresponding exotic abelian Polish groups.

In the setting of a locally compact group G , these pseudo-metrics are implicit in work of Struble: initially, in 1953 [Str1], he used a ('sampler') family of pre-compact open sets $\{E_t : t > 0\}$ to construct a mean on G , thereby referring to a one-parameter family of pseudo-metrics corresponding to the sets E_t ; some twenty years later in 1974 [Str2] (cf. [DieS, Ch. 8]) identifies a left-invariant (proper) metric on G by taking the supremum of pseudo-metrics, each generated from some open set in a countable open base at 1_G . The pseudo-metric makes a very brief appearance in Yamasaki's textbook treatment [Yam, Ch. 1] of Weil's theorem.

Proposition 10 (Weil pseudo-norm, cf. [Fre, §392H], [Yam, Ch. 1, Proof of Th. 4.1]). For G a Polish group, $\lambda \in \mathcal{M}_{\text{sub}}(G)$, a left-invariant submeasure on $\mathcal{U}(G)$, and $E \in \mathcal{U}(G)$ with $\lambda(E) > 0$, put

$$\|g\|_E^\lambda := \lambda(gE\Delta E) \quad (g \in G).$$

Then $\|\cdot\|_E$ defines a group pseudo-norm with associated right-invariant pseudo-metric

$$d_E^\lambda(g, h) = \|gh^{-1}\|_E^\lambda \quad (g, h \in G).$$

Likewise, for λ right-invariant, a pseudo-norm is defined by

$$\|g\|_E^\lambda := \lambda(E\Delta Eg) \quad (g \in G).$$

Proof. Since $\lambda(\emptyset) = 0$, $\|1_G\|_E^\lambda = 0$. By left invariance under a ,

$$\|a^{-1}\|_E^\lambda = \lambda(a^{-1}E\Delta E) = \lambda(a(a^{-1}E\Delta E)) = \lambda(E\Delta aE) = \|a\|_E^\lambda.$$

Also,

$$\|ab\|_E^\lambda \leq \|a\|_E^\lambda + \|b\|_E^\lambda$$

follows from monotonicity, subadditivity and $\lambda(abE\Delta aE) = \lambda(bE\Delta E)$:

$$\begin{aligned} \lambda(abE \setminus E \cup E \setminus abE) &\leq \lambda(abE \setminus aE) \cup (aE \setminus E) \cup (E \setminus aE) \cup (aE \setminus abE) \\ &= \lambda(abE \setminus aE) \cup (aE \setminus abE) \cup (aE \setminus E) \cup (E \setminus aE) \\ &\leq \lambda(abE\Delta aE) + \lambda(E\Delta aE) = \lambda(bE\Delta E) + \lambda(E\Delta aE). \quad \square \end{aligned}$$

Corollary 6 (Kneser for Haar measure, [Kne, Hilfs. 4]). For G a Polish group, $\lambda \in \mathcal{M}_{\text{sub}}(G)$, a left-invariant submeasure on $\mathcal{U}(G)$, and $E \in \mathcal{U}(G)$ with $\lambda(E) > 0$, the set

$$H := \{g \in G : \lambda(gE\Delta E) = 0\}$$

is a subgroup of G closed under the norm $\|g\|_E^\lambda$.

Proof. Indeed $H = \{g \in G : \|g\|_E^\lambda = 0\}$, and so H is a subgroup, since for $g, h \in H$, $\|gh^{-1}\|_E^\lambda \leq \|g\|_E^\lambda + \|h\|_E^\lambda = 0$. \square

Recall now (from the Introduction) that a subset of a Polish group G is *left Haar null* if it is contained in a universally measurable set B such that for some $\mu \in \mathcal{P}(G)$

$$\mu(gB) = 0 \quad (g \in G).$$

It is *Haar null* [Sol1] (cf. [HofT, p. 374]) if it is contained in a universally measurable set B such that for some $\mu \in \mathcal{P}(G)$

$$\mu(gBh) = 0 \quad (g, h \in G).$$

This motivates the following application of Proposition 10 beyond Haar measure. Extending the notation of §3, below $\mathcal{M}_0^L(G)$ (resp. $\mathcal{M}_0(G)$) denotes the family of left-Haar-null (resp. Haar-null) sets of G , and we write

$$\mathcal{U}_+^L(G) := \mathcal{U}(G) \setminus \mathcal{M}_0^L(G), \quad \mathcal{U}_+(G) := \mathcal{U}(G) \setminus \mathcal{M}_0(G).$$

Prop. 10 may be applied to the following measures; those constructed from μ a normalized counting measure (of finite support) are studied in [Sol1].

Proposition 11. *In a Polish group G , for $\mu \in \mathcal{P}(G)$ put*

$$\begin{aligned} \mu_L^*(E) &: = \sup\{\mu(gE) : g \in G\} & (E \in \mathcal{U}(G)), \\ \hat{\mu}(E) &: = \sup\{\mu(gEh) : g, h \in G\} & (E \in \mathcal{U}(G)). \end{aligned}$$

Then μ_L^* (resp. $\hat{\mu}$) is a left invariant (resp. bi-invariant) submeasure, which is positive for $E \in \mathcal{U}_+^L(G)$ (resp. for $E \in \mathcal{U}_+(G)$), i.e. for universally measurable, non-left-Haar null (resp. non-Haar null) sets.

Proof. We consider only $\hat{\mu}$, as the case μ_L^* is similar and simpler (through the omission of h and b below). The set function $\hat{\mu}$ is well defined, with

$$\mu(E) \leq \hat{\mu}(E) \leq 1 \quad (E \in \mathcal{U}(G)),$$

since μ is a probability measure; it is bi-invariant, since

$$\hat{\mu}(aEb) := \sup\{\mu(gaEbh) : g, h \in G\} = \sup\{\mu(gEh) : g, h \in G\},$$

and G is a group. Furthermore, for $B \in \mathcal{U}(G)$

$$\mu(gBh) \leq \hat{\mu}(B) \leq 1, \quad (g, h \in G).$$

So, for $\mu \in \mathcal{P}(G)$

$$0 < \hat{\mu}(B) \leq 1 \quad (B \in \mathcal{U}_+(G)),$$

since there are $g, h \in G$ with $\mu(gBh) > 0$. Countable subadditivity follows (on taking suprema of the leftmost term over g, h) from

$$\mu(g(\bigcup_n A_n)h) \leq \sum_n \mu(gA_nh) \leq \sum_n \hat{\mu}(gA_nh) = \sum_n \hat{\mu}(A_n),$$

for any sequence of sets $A_n \in \mathcal{U}(G)$. \square

Definition. For $\mu \in \mathcal{P}(G)$, $E \in \mathcal{U}(G)$, put

$$B_\varepsilon^E(\mu) := \{x \in G : \|x\|_E^\mu < \varepsilon\}.$$

Our next step uses Prop. 11 to inscribe these balls into EE^{-1} for all small enough $\varepsilon > 0$.

Lemma 2 (Self-intersection Lemma). *In a Polish group G for $E \in \mathcal{U}_+(G)$, and respectively for $E \in \mathcal{U}_+^L(G)$, and $\mu \in \mathcal{P}(G)$,*

$$\left. \begin{array}{l} 1_G \in B_\varepsilon^E(\hat{\mu}) \subseteq EE^{-1} \quad (0 < \varepsilon < \hat{\mu}(E)), \\ 1_G \in B_\varepsilon^E(\mu_L^*) \subseteq EE^{-1} \quad (0 < \varepsilon < \mu_L^*(E)). \end{array} \right\}$$

Equivalently, for $0 < \varepsilon < \hat{\mu}(E)$, and respectively for $0 < \varepsilon < \mu_L^(E)$,*

$$E \cap xE \neq \emptyset \quad (x \in B_\varepsilon^E(\hat{\mu})); \quad E \cap xE \neq \emptyset \quad (x \in B_\varepsilon^E(\mu_L^*)).$$

Proof. We check only the $\hat{\mu}$ case; the other is similar and simpler (through the omission of h below). For $E \in \mathcal{U}_+(G)$, since $\hat{\mu}(E) > 0$ by Prop. 11, we may pick $g, h \in G$ such that $\varepsilon_E := \mu(gEh) > 0$. Consider x and $\varepsilon > 0$ with $\|x\|_E^\mu < \varepsilon \leq \varepsilon_E$. If E and xE are disjoint, then

$$\begin{aligned} \varepsilon_E &= \mu(gEh) \leq \mu(g(E \cup xE)h) \leq \hat{\mu}(g(E \cup xE)h) = \hat{\mu}(E \cup xE) \\ &= \hat{\mu}(xE \triangle E) = \|x\|_E^\mu < \varepsilon \leq \varepsilon_E, \end{aligned}$$

a contradiction. So E and xE do meet. Now first pick $t \in xE \cap E$ and next $s \in E$ so that $t = xs$; then $x = ts^{-1} \in EE^{-1}$. The argument is valid when $\varepsilon_E = \mu(gEh)$ assumes any value in $(0, \hat{\mu}(E)]$, as in Prop. 1M. For the converse, if $B_\varepsilon^E(\hat{\mu}) \subseteq EE^{-1}$, proceed as in Prop. 1M. \square

We need a simple analogue of a result due to Weil ([Wei, Ch. VII, §31], cf. [Hal, Ch. XII §62]). Below τ_1 denotes the τ -open *neighbourhoods* of 1_G . For G locally compact with $\lambda = \eta_G$, the identity

$$2\eta(E) - 2\eta(E \cap xE) = \eta(E \triangle xE) = 1 - 2 \int 1_E(t)1_{E^{-1}}(t^{-1}x)d\eta(t) \quad (\dagger)$$

connects the continuity of the (pseudo-) norm to \mathcal{T}_d -continuity of translation in the topological group structure (G, \mathcal{T}_d) of the locally compact group, and

to continuity of the convolution function here (for E of finite η -measure) – see [HewR, Th. 20.16]; see also [HewR, Th. 20.17] for the well-known connection between the Steinhaus-Weil Theorem and convolution. Such a continuity condition result guarantees that $B_\varepsilon^E(\eta)$ contains points other than 1_G .

Lemma 3 (Fragmentation Lemma; cf. [Hal, Ch. XII §62 Th. A]). *For $\lambda \in \mathcal{M}_{\text{sub}}(G)$ a left-invariant submeasure on a Polish group G equipped with a finer right-invariant topology τ with 1_G -open-nhd family $\tau_1 \subseteq \mathcal{U}_+^L(G)$:*

if the map

$$x \mapsto \|x\|_E^\lambda$$

is continuous under τ at $x = 1_G$ for each $E \in \mathcal{U}_+^L(G)$

– then, for each $\emptyset \neq E, F \in \tau$ and $\varepsilon > 0$ with $\varepsilon < \lambda(E)$, there exists $H \in \tau_1$ with $HH^{-1} \subseteq FF^{-1}$ and

$$\|h'h^{-1}\|_E^\lambda < \varepsilon \quad (h, h' \in H) : \quad HH^{-1} \subseteq B_\varepsilon^E,$$

so that $\text{diam}_E^\lambda(H) \leq \varepsilon$.

Proof. Pick any $f \in F$, and $D \in \tau_1$ satisfying $\|x\|_E^\lambda < \varepsilon/2$ for all $x \in D$. As τ is right-invariant and $1_G \in D \cap Ff^{-1} \in \tau$, pick $H \in \tau_1$ with $H \subseteq D \cap Ff^{-1}$; then

$$HH^{-1} = Hff^{-1}H^{-1} \subseteq FF^{-1}.$$

For $h, h' \in H$, as $h, h' \in D$

$$\|h'f(hf)^{-1}\|_E^\lambda = \|h'h^{-1}\|_E^\lambda \leq \|h'\|_E^\lambda + \|h^{-1}\|_E^\lambda = \|h'\|_E^\lambda + \|h\|_E^\lambda < \varepsilon. \quad \square$$

In the presence of a refinement topology τ on the group G , the lemma motivates further notation: write $\mathcal{P}_{\text{cont}}(G, \tau)$, or just

$$\mathcal{P}_{\text{cont}}(\tau) := \{\mu \in \mathcal{P}(G, \mathcal{I}_d) : g \mapsto \|g\|_E^{\hat{\mu}} := \hat{\mu}(gE\Delta E) \text{ is } \tau\text{-continuous at } 1_G\}.$$

Of necessity attention here focuses on continuity. The characterization question as to which topologies τ yield a non-empty $\mathcal{P}(\tau)$ is in part answered by Theorem 7M below. Indeed, for Haar measure η in the locally compact case,

$$\mu \in \mathcal{P}_{\text{cont}}(\tau) \quad (\mu \ll \eta, \tau \supseteq \mathcal{I}_d),$$

by (†) in the presence of $d\mu/d\eta$ as a kernel:

$$\|x\|_E^\mu = 1 - 2 \int 1_E(t)1_{E^{-1}}(t^{-1}x) \frac{d\mu}{d\eta}(t). \quad (\dagger\dagger)$$

However, $\mathcal{P}(G)$ will contain measures μ singular with respect to η : for such μ , by Th. SM there will be Borel subsets B of positive μ -measure such that BB^{-1} has void \mathcal{T}_d -interior.

6 Weil-like topologies: theorems

Prop. 11 now yields the following result, which embraces known Hashimoto topologies [BinO5] in both the Polish abelian setting, where the left Haar null sets form a σ -ideal (Christensen [Chr1]), and likewise in (the not necessarily abelian) Polish groups that are *amenable at 1* (Solecki [Sol1]); this includes, as additive groups, F - (hence also Banach) spaces – cf. [BinO5,6], where use is made of Hashimoto topologies.

Theorem 7. *Let G be a Polish group and τ both a left- and a right-invariant refinement topology with 1_G -open-nhd family $\tau_1 \subseteq \mathcal{U}_+(G)$.*

Then both the families $\{AA^{-1} : A \in \tau_1\}$ and $\{B_\varepsilon^E(\hat{\mu}) : \emptyset \neq E \in \tau, \mu \in \mathcal{P}(\tau)$ and $0 < \varepsilon \leq \hat{\mu}(E)\}$ generate neighbourhoods of the identity under which G is a topological group. Moreover, the pseudo-norms

$$\{\|\cdot\|_E^\mu : \emptyset \neq E \in \tau, \mu \in \mathcal{P}_{\text{cont}}(\tau)\}$$

are downward directed by refinement as follows: for $\emptyset \neq E, F \in \tau_1$, $\lambda, \mu \in \mathcal{P}(\tau)$ and $\varepsilon < \min\{\hat{\lambda}(E), \hat{\mu}(F)\}$, there is $H \in \tau_1$ such that for $0 < \delta < \min\{\tilde{\lambda}(H), \hat{\mu}(H)\}$

$$B_\delta^H(\lambda) \cap B_\delta^H(\mu) \subseteq B_\varepsilon^E(\lambda) \cap B_\varepsilon^F(\mu).$$

Proof. The proof is similar to but simpler than that of [Hal, Ch. XII §62 Th. A]. Given two (non-left-Haar-null) sets $E, F \in \tau_1$ and $\varepsilon < \min\{\hat{\lambda}(E), \hat{\mu}(F)\}$, by the Fragmentation Lemma (Lemma 3 of §5) applied separately to $\hat{\lambda}$ and to $\hat{\mu}$, there are $A, B \in \tau_1$ with

$$AA^{-1} \subseteq B_\varepsilon^E(\hat{\lambda}), \quad BB^{-1} \subseteq B_\varepsilon^F(\hat{\mu}).$$

Take any $H \in \tau_1$ with $H \subseteq A \cap B$; then

$$HH^{-1} \subseteq AA^{-1} \cap BB^{-1}.$$

Since $H \in \mathcal{U}_+(G)$ (as $\tau_1 \subseteq \mathcal{U}_+(G)$), take δ with $0 < \delta < \min\{\hat{\lambda}(H), \hat{\mu}(H)\}$; then by (*)

$$B_\delta^H(\hat{\lambda}) \cap B_\delta^H(\hat{\mu}) \subseteq HH^{-1} \subseteq AA^{-1} \cap BB^{-1} \subseteq B_\varepsilon^E(\hat{\lambda}) \cap B_\varepsilon^F(\hat{\mu}).$$

(So ‘mutual refinement’ holds between the sets of the form AA^{-1} and those of the form B_ε^E .) As $\|\cdot\|_E^\mu$ is a pre-norm,

$$B_{\varepsilon/2}^E(\hat{\mu})B_{\varepsilon/2}^E(\hat{\mu})^{-1} = B_{\varepsilon/2}^E(\hat{\mu})B_{\varepsilon/2}^E(\hat{\mu}) \subseteq B_\varepsilon^E(\hat{\mu}).$$

By the Fragmentation Lemma again, given any $x \in G$ and $\varepsilon > 0$, choose $H \in \tau_1$ with $HH^{-1} \subseteq B_\varepsilon^E(\hat{\mu})$. Then with $F := xH \in \tau$,

$$B_\varepsilon^F(\hat{\mu}) = \{z : \|z\|_F^\mu < \varepsilon\} \subseteq (xH)(xH)^{-1} = xHH^{-1}x^{-1} \subseteq xB_\varepsilon^E(\hat{\mu})x^{-1}.$$

Finally, for any x_0 with $\|x_0\|_E^\mu < \varepsilon$, put $\delta := \varepsilon - \|x_0\|_E^\mu$. Then for $\|y\|_E^\mu < \delta$,

$$\|x_0 \cdot y\|_E^\mu \leq \|x_0\|_E^\mu + \|y\|_E^\mu < \|x_0\|_E^\mu + \varepsilon - \|x_0\|_E^\mu < \varepsilon,$$

i.e.

$$x_0B_\delta^E(\hat{\mu}) \subseteq B_\varepsilon^E(\hat{\mu}). \quad \square$$

Specializing to locally compact groups yields as a corollary, on writing $B_\varepsilon^E := B_\varepsilon^E(\eta)$:

Theorem 7M. *For G a locally compact group with left Haar measure η , if:*
(i) τ is both a left- and a right-invariant refinement topology with $\tau_1 \subseteq \mathcal{M}_+$,
(ii) for every non-empty $E \in \tau$, the pseudo-norm

$$g \mapsto \|g\|_E := \eta(gE\Delta E) \quad (g \in G)$$

is continuous under τ at $g = 1_G$

– then both the families $\{AA^{-1} : A \in \tau_1\}$ and $\{B_\varepsilon^E : \emptyset \neq E \in \tau \text{ and } 0 < \varepsilon \leq 2\eta(E)\}$ generate neighbourhoods of the identity under which G is a topological group. Moreover, the pseudo-norms

$$\{\|\cdot\|_E : \emptyset \neq E \in \tau\}$$

are downward directed by refinement; indeed, for $\emptyset \neq E, F \in \tau$ and $\varepsilon < 2 \min\{\eta(E), \eta(F)\}$, there is $H \in \tau_1$ such that for $0 < \delta < \eta(H)$

$$B_\delta^H \subseteq B_\varepsilon^E \cap B_\varepsilon^F.$$

Proof. It is enough to replace $\mathcal{P}(G)$ by $\{\eta\}$ (so that λ and μ both refer to η), and to note that if xE and E are disjoint, then $\eta(xE \Delta E) = 2\eta(E)$, so that in Lemma 2 the bound $\eta^*(E)$ in the restriction governing inclusion may be replaced by $2\eta(E)$. \square

Remark. As in [Hal, Ch. XII §62 Th. F], but by the Fragmentation Lemma (and by the countable additivity of η), the Weil-like topology on a locally compact G in Theorem 7M is locally bounded (norm-totally-bounded in some ball). Then G with the Weil-like topology may be densely embedded in its completion \hat{G} , which is in turn locally compact, being locally complete and (totally) bounded. However, the corresponding argument in the case of the preceding more general Theorem 7 fails, since $\hat{\mu}$ there is not necessarily countably additive.

Finally, we give a category version of Theorem 7M, as an easy corollary; indeed, our main task is merely to define what is meant by ‘mutatis mutandis’ in the present context. Given the assumption $\tau_1 \subseteq \mathcal{B}_+$, we are entitled to refer to the usual quasi-interior of any $E \in \mathcal{B}_+$, denoted below by \tilde{E} , as in Cor. 2’; we also write \tilde{B}_ε^E for $B_\varepsilon^{\tilde{E}}(\eta)$.

Theorem 7B. *For G a locally compact group with left Haar measure η , if:*
(i) τ is both a left- and a right-invariant refinement topology with $\tau_1 \subseteq \mathcal{B}_+$ and with the left Nikodym property (preservation of category under left shifts),
(ii) for every non-empty $E \in \tau$ the pseudo-norm

$$g \mapsto \|g\|_{\tilde{E}} := \eta(g\tilde{E} \Delta \tilde{E}) \quad (g \in G)$$

is continuous under τ at $g = 1_G$
– then both the families $\{AA^{-1} : A \in \tau_1\}$ and $\{\tilde{B}_\varepsilon^E : \emptyset \neq E \in \tau \text{ and } 0 < \varepsilon \leq 2\eta(\tilde{E})\}$ generate neighbourhoods of the identity under which G is a topological group. Moreover, the pseudo-norms

$$\{\|\cdot\|_{\tilde{E}} : \emptyset \neq E \in \tau\}$$

are downward directed by refinement; indeed, for $\emptyset \neq E, F \in \tau$ and $\varepsilon < 2 \min\{\eta(\tilde{E}), \eta(\tilde{F})\}$, there is $H \in \tau_1$ such that for $0 < \delta < 2\eta(\tilde{H})$

$$\tilde{B}_\delta^H \subseteq \tilde{B}_\varepsilon^E \cap \tilde{B}_\varepsilon^F.$$

Proof. In place of the inclusion of Lemma 2 we note a result stronger than that valid for \tilde{E} (i.e. inclusion only in $\tilde{E}\tilde{E}^{-1}$): since meagreness is translation-invariant (the ‘Nikodym property’ of [BinO5]), $(xE)^\sim = x\tilde{E}$ for non-meagre Baire E , so $x\tilde{E} \cap \tilde{E} \neq \emptyset$ implies $xE \cap E \neq \emptyset$, and so again

$$\tilde{B}_\varepsilon^E = B_\varepsilon^{\tilde{E}} \subseteq EE^{-1};$$

here again in Lemma 2 the bound $\eta^*(E)$ in the restriction governing inclusion may be replaced by $2\eta(E)$. The proof of Theorem 7 may now be followed verbatim, but for the replacement of $\mathcal{P}(G)$ by $\{\eta\}$, using the stronger inclusion just observed, and $B_\varepsilon(\eta)$ by \tilde{B}_ε . \square

Remark. The last result follows more directly from Th. 7M in a context where there exists on G a *Marczewski measure* (see [TomW, Ch. 13, cf. Ch. 11]), i.e. a finitely additive invariant measure on \mathcal{B} vanishing on bounded members of \mathcal{B}_0 ; this includes $\mathbb{R}, \mathbb{R}^2, \mathbb{S}^1$, albeit under AC [TomW, Cor. 13.3]; cf. [Myc], but not \mathbb{R}^d for $d \geq 3$ [DohF].

With the groundwork of Part I on translation-continuity for *compacts* completed, we close by establishing the promised dichotomy associated with the map

$$x \mapsto \|x\|_E^\mu = \mu(xE\Delta E),$$

for measurable E : Theorem FN (§1) creates a duality between the vanishing of the F -based pseudo-norm and a *dichotomy* for x -translates of E^{-1} in relation to F according as $x \in E$ or $x \notin E$, which are thus unable in each case to distinguish between the points of F . Below we write \forall^μ for the generalized quantifier ‘for μ -a.a.’ (cf. [Kec, 8.J]).

Theorem 8 (Almost Inclusion-Exclusion). *For G a Polish group $\mu \in \mathcal{P}(G)$ and non-null μ -measurable E, F , the vanishing μ -a.e. on F of the E -norm under μ :*

$$\|x\|_F^\mu = \mu(xE\Delta E) = 0 \quad (x \in F),$$

is equivalent to the following Almost Inclusion-Exclusion for translates of E^{-1} :

(i) Inclusion: F is μ -almost covered by μ -almost every translate xE^{-1} for $x \in E$:

$$\mu(F \setminus xE^{-1}) = 0 \quad (\forall^\mu x \in E),$$

(ii) Exclusion: F is μ -almost disjoint from μ -almost every translate xE^{-1} for $x \notin E$:

$$\mu(F \cap xE^{-1}) = 0 \quad (\forall^\mu x \notin E).$$

Proof. By Theorem FN (§1), applied to the set H of Prop. 3, i.e.

$$H := \bigcup_{x \in F} \{x\} \times (xE \Delta E),$$

H has vertical sections H_x almost all μ -null iff μ -almost all of its horizontal sections H^y are μ -null. But, since $y \in xE$ iff $x \in yE^{-1}$, $H^y = F \setminus yE^{-1}$ for $y \in E$ and $H^y := F \cap yE^{-1}$ for $y \in G \setminus E$. \square

Remark. If the inclusion side of the dichotomy of Th. 8 holds for all $x \in E$, then $F \subseteq EE^{-1}$. The converse direction may fail: consider $E = (1, 2) \subseteq \mathbb{R}$ and $F = (-1, 1)$, so that $E - E = F$, but no translate of $-E$ may cover F .

7 Other interior-point properties

7.1 The Steinhaus property AA^{-1} versus the Steinhaus property AB^{-1}

We clarify the relation between two versions of the Steinhaus interior points property: the simple (sometimes called ‘classical’) version concerning sets AA^{-1} and the composite, more embracing one, concerning sets AB^{-1} , for sets from a given family \mathcal{H} . The latter is connected to a strong form of metric transitivity: Kominek [Kom] shows, for a general separable Baire topological group G equipped with an inner regular measure μ defined on some σ -algebra \mathcal{M} , that AB^{-1} has non-empty interior for all $A, B \in \mathcal{M}_+$ iff for each countable dense set D and each $E \in \mathcal{M}_+$ the set $X \setminus DE \in \mathcal{M}_0$; this is recalled in Theorem K below. The composite property is thus related to the Smital property, for which see [BarFN]. Care is required when moving to the alternative property for AB , since the family \mathcal{H} need not be preserved under inversion.

In general the simple property does not imply the composite: Matořsková and Zelený [MatZ] show that in any non-locally compact abelian Polish group there are closed non-(left) Haar null sets A, B such that $A + B$ has empty interior. Recently, Jabłońska [Jab2] has shown that likewise in any non-locally compact abelian Polish group there are closed non-Haar meager sets A, B such that $A + B$ has empty interior. Bartoszewicz and M. and T. Filipczak [BarFF, Ths. 1, 4] analyze the Bernoulli product measure on $\{0, 1\}^{\mathbb{N}}$ with p the probability of the digit 1; see §9.15. The product space may be regarded as comprising canonical binary digit expansions of the additive reals modulo 1 (in which case the measure is not invariant). Here the (Borel) set A of binary expansions with asymptotic frequency p of the digit ‘1’ has $[0, 1)$ as its difference set iff $\frac{1}{4} \leq p \leq \frac{3}{4}$; however $A + A$ has empty interior unless $p = \frac{1}{2}$ (the base 2 simple-normal-numbers case).

Below we identify some conditions on a family of sets A with the simple AA^{-1} property which do imply the AB^{-1} property. What follows is a generalization to a group context of relevant observations from [BinO9] from the classical context of \mathbb{R} .

The motivation for the definition below is that its subject, the space H , is a subgroup of a topological group G from which it inherits a (necessarily) translation-invariant (i.e. either-sidedly) topology τ . Various notions of ‘density at a point’ give rise to ‘density topologies’ [BinO5], which are translation-invariant since they may be obtained via translation to a fixed reference point: early examples, which originate in spirit with Denjoy as interpreted by Haupt and Pauc [HauP], were studied intensively in [GofW], [GofNN], soon followed by [Mar] and [Mue]; more recent examples include [FilW] and others investigated by the Wilczyński school, cf. [Wil].

Proposition 12 below embraces as an immediate corollary the case $H = G$ with G locally compact and σ the Haar density topology (see [BinO8]). Proposition 13 proves that Proposition 12 applies also to the ideal topology (in the sense of [LukMZ]) generated from the ideal of Haar null sets of an abelian Polish group.

We recall that a group H carries a *left semi-topological* structure τ if the topology τ is left invariant [ArhT] ($hU \in \tau$ iff $U \in \tau$); the structure is *semi-topological* if it is also right invariant, i.e. briefly: τ is translation invariant. H is a *quasi-topological group* under τ if τ is both left and right invariant and inversion is τ -continuous.

Definition. For H a group with a translation-invariant topology τ , call a

topology $\sigma \supseteq \tau$ a *Steinhaus refinement* if:

- i) $\text{int}_\tau(AA^{-1}) \neq \emptyset$ for each non-empty $A \in \sigma$, and
- ii) σ is involutive-translation invariant: $hA^{-1} \in \sigma$ for all $A \in \sigma$ and all $h \in H$.

Property (ii) above (called simply ‘invariance’ in [BarFN]) calls apparently for only left invariance, but in fact, via double inversion, delivers translation invariance, since $Uh = (h^{-1}U^{-1})^{-1}$; then H under σ is a semi-topological group with a continuous inverse, so a *quasi-topological group*.

Proposition 12. *If τ is translation-invariant, and $\sigma \supseteq \tau$ is a Steinhaus refinement topology, then $\text{int}_\tau(AB^{-1}) \neq \emptyset$ for non-empty $A, B \in \sigma$. In particular, as σ is preserved under inversion, also $\text{int}_\tau(AB) \neq \emptyset$ for $A, B \in \sigma$.*

Proof. Suppose $A, B \in \sigma$ are non-empty; as $B^{-1} \in \sigma$, choose $a \in A$ and $b \in B$; then by (ii)

$$1_H \in C := a^{-1}A \cap b^{-1}B^{-1} \in \sigma.$$

By (i), for some non-empty $W \in \tau$,

$$W \subseteq CC^{-1} = (a^{-1}A \cap b^{-1}B) \cdot (A^{-1}a \cap B^{-1}b) \subseteq (a^{-1}A) \cdot (B^{-1}b).$$

As τ is translation invariant, $aWb^{-1} \in \tau$ and

$$aWb^{-1} \subseteq AB^{-1},$$

the latter since for each $w \in W$ there are $x \in A, y \in B^{-1}$ with

$$w = a^{-1}x.yb : \quad awb^{-1} = xy \in AB^{-1}.$$

So $\text{int}_\tau(AB^{-1}) \neq \emptyset$. \square

Corollary 7. *In a locally compact group the Haar density topology is a Steinhaus refinement.*

Proof. Property (i) follows from Weil’s theorem since density-open sets are non-null measurable; left translation invariance in (ii) follows from left invariance of Haar measure, while involutive invariance holds, as any measurable set of positive Haar measure has non-null inverse ([HewR, 15.14], cf. §4 Lemma H). \square

A weaker version, inspired by metric transitivity, comes from applying the following concept.

Definition. Say H acts transitively on \mathcal{H} for each $A, B \in \mathcal{H}$ if there is $h \in H$ with $A \cap hB \in \mathcal{H}$.

Thus a locally compact topological group acts transitively on the non-null Haar measurable sets (in fact, either-sidedly); this follows from Fubini's Theorem [Hal, 36C], via the average theorem [Hal, 59.F]:

$$\int_G |g^{-1}A \cap B| dg = |A| \cdot |B^{-1}| \quad (A, B \in \mathcal{M}),$$

($g = ab^{-1}$ iff $g^{-1}a = b$) – cf. [TomW, §11.3 after Th. 11.17].

[MatZ] show that in any non-locally compact abelian Polish group G there exist two non-Haar null sets, $A, B \notin \mathcal{HN}$, such that $A \cap hB \in \mathcal{HN}$ for all h ; that is, G here does *not* act transitively on the non-Haar null sets.

Definition. In a quasi-topological group (H, τ) say that a proper σ -ideal \mathcal{H} has the *Simple Steinhaus Property* AA^{-1} if AA^{-1} has interior points for universally measurable subsets $A \notin \mathcal{H}$. This follows [BarFN].

Proposition 12' (cf. [Kha, Th. 1]). *In a group (H, τ) with τ translation-invariant, if H acts transitively on a family of subsets \mathcal{H} with the simple Steinhaus property, then \mathcal{H} has the composite Steinhaus property: $\text{int}_\tau(AB^{-1}) \neq \emptyset$ for $A, B \in \mathcal{H}$. Furthermore, if \mathcal{H} is preserved under inversion, then also $\text{int}_\tau(AB) \neq \emptyset$ for $A, B \in \mathcal{H}$.*

Proof. For $A, B \in \mathcal{H}$ choose h with $C := A \cap hB \in \mathcal{H}$; then

$$CC^{-1}h = (A \cap hB)(A^{-1} \cap B^{-1}h^{-1}) \subseteq AB^{-1}. \quad \square$$

Proposition 13. *If (H, τ) is a quasi-topological group (i.e. τ is invariant with continuous inversion) carrying a left invariant σ -ideal \mathcal{H} with the Steinhaus property and $\tau \cap \mathcal{H} = \{\emptyset\}$, then the ideal-topology σ with basis*

$$\mathcal{B} := \{U \setminus N : U \in \tau, N \in \mathcal{H}\}$$

is a Steinhaus refinement of τ .

In particular, for (H, τ) an abelian Polish group, the ideal topology generated by the σ -ideal of Haar null subsets is a Steinhaus refinement.

Proof. If $U, V \in \mathcal{B}$ and $w \in U \cap V$, choose $M, N \in \mathcal{H}$ and $W_M, W_N \in \tau$ such that $x \in (W_M \setminus M) \subseteq U$ and $x \in (W_N \setminus N) \subseteq V$. Then as $M \cup N \in \mathcal{H}$,

$$x \in (W_M \cap W_N) \setminus (M \cup N) \in \mathcal{B}.$$

So \mathcal{B} generates a topology σ refining τ . With the same notation, $hU = hW_M \setminus hM \in \sigma$, as $hM \in H$, and $U^{-1} = W_M^{-1} \setminus M^{-1}$. Finally, UU^{-1} has non-empty τ -interior, as $U \notin \mathcal{H}$ and is non-empty.

As for the final assertion concerned with an abelian Polish group context, note that if N is Haar null ($N \in \mathcal{HN}$), then $\mu(hN) = 0$ for some $\mu \in \mathcal{P}(G)$ and all $h \in H$, so $hN \in \mathcal{HN}$ for all $h \in H$. Furthermore, if $A \notin \mathcal{HN}$ then $A^{-1} \notin \mathcal{HN}$, otherwise $\mu(hA^{-1}) = 0$ for some $\mu \in \mathcal{P}(G)$ and all $h \in H$; then, taking $\tilde{\mu}(B) = \mu(B^{-1})$ for Borel B , we have $\tilde{\mu}(A) = 0$ and $\tilde{\mu}(hA) = \mu(A^{-1}h^{-1}) = 0$ for all $h \in H$, a contradiction. \square

Remark. A left Haar null set need not be right Haar null: for one example see [ShiT], and for more general non-coincidence see Solecki [Sol1, Cor. 6]. So the argument in Prop. B does not extend to the family of left Haar null sets \mathcal{LHN} of a *non-commutative* Polish group. Indeed, Solecki [Sol2, Th. 1.4] shows in the context of a countable product of countable groups that the simpler Steinhaus property holds for \mathcal{HN} iff $\mathcal{HN} = \mathcal{LHN}$.

Next, a result from [Kom]. Recall that μ is *quasi-invariant* if μ -nullity is translation invariant. The transitivity assumption (of co-nullity) is motivated by *Smítal's lemma*, which refers to a countable dense set – see [KucS].

Theorem K. ([Kom, Th. 5]). *If $\mu \in P(G)$ is quasi-invariant and there exists a countable subset $H \subseteq G$ with HM co-null for all $M \in M_+(\mu)$, then $\text{int}(AB^{-1}) \neq \emptyset$ for all $A, B \in \mathcal{M}_+(\mu)$.*

Proof. By regularity, we may assume $A, B \in \mathcal{M}_+(\mu)$ are compact, so AB^{-1} is compact. Fix $g \in G$; then by quasi-invariance $\mu(gB) > 0$. So by the transitivity assumption, both $G \setminus HgB$ and $G \setminus HA$ are null, and so $HA \cap HgB \neq \emptyset$. Say $h_1a = h_2gb$, for some $a \in A, b \in B, h_1, h_2 \in H$; then $g = h_2^{-1}h_1ab^{-1}$. As g was arbitrary,

$$G = \bigcup_{h \in H} h_2^{-1}h_1AB^{-1}.$$

By Baire's Theorem, as H is countable, $\text{int}(AB^{-1}) \neq \emptyset$. \square

7.2 Borell's interior-point property

For completeness of this overview of the Steinhaus-Weil interior-point property, we offer in brief here the context and statement of a (by now) classical Steinhaus-like result in probability theory; this differs in that the Polish group now specializes to an infinite-dimensional topological vector space and the reference measure is Gaussian, so no longer invariant. We refer to the companion paper [Bin08] for further details and background literature, and to our generalizations to Polish groups and to other reference measures.

For X a locally convex topological vector space, γ a probability measure on the σ -algebra of the cylinder sets generated by the dual space X^* (equivalently, for X separable Fréchet, e.g. separable Banach, the Borel sets), with $X^* \subseteq L^2(\gamma)$: then γ is called *Gaussian* on X (gamma for Gaussian, following [Bog1]) iff $\gamma \circ \ell^{-1}$ defined by

$$\gamma \circ \ell^{-1}(B) = \gamma(\ell^{-1}(B)) \quad (\text{Borel } B \subseteq \mathbb{R})$$

is Gaussian (normal) on \mathbb{R} for every $\ell \in X^* \subseteq L^2(\gamma)$. For a monograph treatment of Gaussianity in a Hilbert-space setting, see Janson [Jan]. Write $\gamma_h(K) := \gamma(K + h)$ for the translate by h . *Relative quasi-invariance* of γ_h and γ , that for all compact K

$$\gamma_h(K) > 0 \text{ iff } \gamma(K) > 0,$$

holds relative to a set of vectors $h \in X$ (the *admissible directions*) forming a vector subspace known as the *Cameron-Martin space*, $H(\gamma)$. Then γ_h and γ are equivalent, $\gamma \sim \gamma_h$, iff $h \in H(\gamma)$. Indeed, if $\gamma \sim \gamma_h$ fails, then the two measures are mutually singular, $\gamma_h \perp \gamma$ (the Hajek-Feldman Theorem – cf. [Bog1, Th. 2.4.5, 2.7.2]).

Continuing with the assumption above on X^* , as $X \subseteq X^{**} \subseteq L^2(\gamma)$, one can equip $H = H(\gamma)$ with a norm derived from that on $L^2(\gamma)$. In brief, this is done with reference to a natural covariance under γ obtained by regarding $f \in X^*$ as a random variable and working with its zero-mean version $f - \gamma(f)$; then, for $h \in H$, δ_h^γ , the (shifted) evaluation map defined by $\delta_h^\gamma(f) := f(h) - \gamma(f)$ for $f \in X^*$, is represented as $\langle f - \gamma(f), \hat{h} \rangle_{L^2(\gamma)}$ for some $\hat{h} \in L^2(\gamma)$. (Here for γ symmetric $\gamma(f) = 0$, so $\delta_h^\gamma = \delta_h$ is the Dirac measure at h .) This is followed by identifying h with \hat{h} (for $h \in H$), and $|h|_H := \|\hat{h}\|_{L^2(\gamma)}$ is a norm on H arising from the inner product

$$(h, k)_H := \int_X \hat{h}(x)\hat{k}(x)d\gamma(x).$$

Formally, the construction first requires an extension of the domain of δ_h^γ to X_γ^* , the closed span of $\{x^* - \gamma(x^*) : x^* \in X^*\}$ in $L^2(\gamma)$, a Hilbert subspace in which to apply the Riesz Representation Theorem.

We may now state the Steinhaus-like property due, essentially in this form, to Christer Borell. ([LeP, Prop. 1] offers a weaker, ‘one-dimensional-section’ form, which would now be called, as in e.g. [Brz], a ‘radial’ form.)

Theorem B (Borell’s Interior-point Theorem, [Bor, Cor. 4.1] – see [Bog1, p. 64]). *For γ a Gaussian measure on a locally convex topological space X with $X^* \subseteq L^2(\gamma)$, and A any non-null γ -measurable subset A of X , the difference set $A - A$ contains a $|\cdot|_H$ -open nhd (neighbourhood) of 0 in the Cameron Martin space $H = H(\gamma)$, i.e. $(A - A) \cap H$ contains a H -open nhd of 0.*

This flows from the continuity in h of the density of γ_h wrt γ ([Bog1, Cor. 2.4.3]), as given in the *Cameron-Martin-Girsanov formula*:

$$\exp\left(\hat{h}(x) - \frac{1}{2}\|\hat{h}\|_{L^2(\gamma)}^2\right) \quad (CM)$$

(where \hat{h} ‘Riesz-represents’ h , i.e. $x^*(h) = \langle x^*, \hat{h} \rangle$, for $x^* \in X^*$, as above). Thus here a modified Steinhaus Theorem holds: the *relative-interior-point theorem*.

8 Complements

1. Non-separability. Simmons establishes his theorems without separability by using the Kakutani-Kodaira ‘separable quotient’ approximation theorem for metric groups [MonZ, I.2.6], cf. [HewR, Th. 8.7] and the recent generalization [Hu]. (Starting with a fixed compact support permits a reduction to the case of a compactly generated group, which is followed by working in a *separable* quotient group.)

The links between the Effros theorem (§8.7 below), the Baire theorem and the Steinhaus-Weil theorem are pursued at length in [Ost2]. There, any separability assumption is avoided. Instead *sequential* methods are used, for example shift-compactness arguments (§8.4 below).

2. Cameron-Martin theory. In the context of Gaussian measures in probability theory, Bogachev [Bog1] gives a thorough treatment of Cameron-Martin

theory (on translation of Wiener and other Gaussian measures; cf. abstract Wiener spaces, reproducing-kernel Hilbert spaces, etc.). Here results of Steinhaus-Weil type occur, in the context of topological vector spaces [Bog1, p.64], as we have seen in §7.2, albeit only relative to a specific *subspace*. In the companion paper to this, [BinO8], we develop an analogous theory in the context of topological groups.

3. *Smallness.* When the (classical) Cameron-Martin subspace of a topological vector space X , alluded to in §8.1 (and mentioned below Th. 5 in §6), is a separable Hilbert space (as when the Gaussian measure γ is a Radon measure, see [Bog1, 3.2.7], cf. [Bog1, p. 62]), so complete under its norm and hence analytic [Rog], being γ -null it has empty interior. So by a result of Dodos [Dod3, Cor. 9], it is (generically) left-Haar null – left-Haar null for quasi all $\mu \in \mathcal{P}(X)$, in the sense of the Lévy metric on $\mathcal{P}(X)$.

4. *Inclusion-Exclusion dichotomy.* Our Weil-like analysis in Part III focuses on inclusions amongst sets of the form EE^{-1} , the exception being the Inclusion-Exclusion of a set F by an E - or non- E -translate of E^{-1} in Theorem 8 (a dichotomy as between E and its complement). This places most of our study on one side of a related inclusion-exclusion dichotomy, for subsets H, B of a group G one has either inclusion, or ‘near-disjointness’:

$$HH^{-1} \subseteq BB^{-1}, \quad \text{or} \quad HH^{-1} \cap BB^{-1} = \{1_G\}.$$

Inclusion may be equivalently re-phrased to the meeting of distinct pairs of H^{-1} -translates of B :

$$kB \cap k'B \neq \emptyset \quad (k, k' \in H^{-1}), \quad (\text{I})$$

whereas exclusion to their disjointness:

$$kB \cap k'B = \emptyset \quad (\text{distinct } k, k' \in H^{-1}). \quad (\text{E})$$

The duality of the relation of (E) to the results in Th. 8 is clarified by observing that $\mu(F \cap xE^{-1}) = 0$, for a.a. $x \in C$, is equivalent to $\mu(C \cap yE) = 0$, for a.a. $y \in F$. Indeed,

$$0 = \int \int 1_C(x) 1_F(y) 1_{xE^{-1}}(y) d(\mu \times \mu) = \int \int 1_F(y) 1_C(x) 1_{yE}(x) d(\mu \times \mu).$$

The condition (E) gives rise to \mathcal{I}_0 , the σ -ideal introduced in Balcerzak et al. [BalRS], generated by Borel sets B having perfectly many disjoint

translates, as in (E) above with H^{-1} a perfect compact set (i.e. compact and dense-in-itself); continuum-many disjoint translates of a compactum also emerge in a theorem of Ulam concerning a non-locally compact Polish group: see [Oxt1, Th. 1]. Such *perfect exclusions* offer a combinatorial tool, akin to *shift-compactness* (as in Th. 3, subsequence embedding under translation of a null sequence into a non-negligible set – cf. [BinO2,3] [MilO]), and play a key role in the context of groups with *ample generics*; see for instance the small-index property of [HodHLS].

Solecki [Sol3] proves a ‘Fubini for negligibles’-type theorem (cf. Theorem FN in §1 above): the non-negligible vertical sections (relative to a uniformly Steinhaus ideal) of a planar \mathcal{I}_0 -negligible set form a horizontal \mathcal{I}_0 -negligible set. The ideal \mathcal{I}_0 is of interest, as it violates the countable (anti)-chain condition, [BalRS].

5. Steinhaus-Weil property of a Borel measure. In a locally compact group G , the family $\mathcal{M}_+(\mu)$ of finite non-null measurable sets of a Borel measure μ on G fails to have the Steinhaus-Weil property iff there are a null sequence $z_n \rightarrow 1_G$ and a non-null compact set K with $\lim_n \mu(t_n K) = 0$. Equivalently, this is so iff the measure μ is not absolutely continuous with respect to Haar measure: these observations motivated Th. SM.

Alternative characterizations of absolute continuity and singularity of Radon measures on a locally compact group include the following.

First for absolute continuity:

- (i) factorization under convolution: $\mu = f * \nu$ with $f \in L^1(G, \eta)$ [LiuR];
- (ii) *continuity* of the map $g \mapsto \mu_g$, using the total variation norm on the space of measures [LiuR];
- (iii) *separability of the orbit* $\{\mu_g : g \in G\}$ in the space of measures, under the same norm as in (ii) [Lar], [LiuRW], [Tam];
- (iv) ‘Glicksberg separability’: for a fixed sequence $\{g_n\}$, each range set $\mathcal{R}(K) := \{\mu_{g_n}(K) : g_n \in K\}$, for μ -null compact sets K , contains $\{\mu(g_n K) : n \in \mathbb{N}\}$ as a dense subset [Gli];
- (v) *outer-equiregularity* of sequences of translates of the μ -null compact sets K : for every $\varepsilon > 0$, compact μ -null K , and sequence $\{g_n\}$ there is an open $W \supseteq K$ with $\mu(g_n(W \setminus K)) < \varepsilon$ for all n [Gli]

Then for singularity: ‘self-singularity’, e.g. $\mu \perp \mu_g$ for η -almost all g , pursued in [LiuR], [LiuRW], [Pro], cf. [BinO8].

6. Regular open sets. Recall that U is regular open if $U = \text{int}(\text{cl}U)$, and that $\text{int}(\text{cl}U)$ is itself regular open; for background see e.g. [GivH, Ch. 10]. For $\mathcal{D} = \mathcal{D}_{\mathcal{B}}$ the Baire-density topology of a normed topological group, let $\mathcal{D}_{\mathcal{B}}^{RO}$

denote the regular open sets. For $D \in \mathcal{D}_{\mathcal{B}}^{RO}$, put

$$N_D := \{t \in G : tD \cap D \neq \emptyset\} = DD^{-1}, \quad \mathcal{N}_1 := \{N_D : 1_G \in D \in \mathcal{D}_{RO}\};$$

then \mathcal{N}_1 is a base at 1_G (since $1_G \in C \in \mathcal{D}_{RO}$ and $1_G \in D \in \mathcal{D}_{RO}$ yield $1_G \in C \cap D \in \mathcal{D}_{RO}$) comprising \mathcal{T} -neighbourhoods that are $\mathcal{D}_{\mathcal{B}}$ -open (since $DD^{-1} = \bigcup\{Dd^{-1} : d \in D\}$). We raise the (metrizability) question, by analogy with the Weil topology of a measurable group (see §5 and §8.4 above): with $\mathcal{D}_{\mathcal{B}}$ above replaced by a general density topology \mathcal{D} on a group G , when is the topology generated by \mathcal{N}_1 on G a norm topology? Some indications of an answer may be found in [ArhT, §3.3]. We note the following answer in the context of Theorem 7B; compare Struble's Theorem [Str2], or [DieS, Ch. 8]. If there exists a separating sequence D_n , i.e. such that for each $g \neq 1_G$ there is n with $\|g\|_{D_n} = 1$, then

$$\|g\| := \sum_n 2^{-n} \|g\|_{D_n}$$

is a norm, since it is separating and, by the Nikodym property, $(D \cap g^{-1}D) = g^{-1}(gD \cap D) \in \mathcal{B}_0$.

7. The Effros Theorem asserts that a transitive continuous action of a Polish group G on a space X of second category in itself is necessarily 'open', or more accurately is microtransitive (the (continuous) evaluation map $e_x : g \mapsto g(x)$ takes open neighbourhoods E of 1_G to open neighbourhoods that are the orbit sets $E(x)$ of x). It emerges that this assertion is very close to the shift-compactness property: see [Ost2]. The Effros Theorem reduces to the Open Mapping Theorem when G, X are Banach spaces regarded as additive groups, and G acts on X by a linear surjection $L : G \rightarrow X$ via $g(x) = L(g) + x$. Indeed, here $e_0(E) = L(E)$. For a neat proof, choose an open neighbourhood U of 0 in G with $E \supseteq U - U$; then $L(U)$ is Baire (being analytic) and non-meagre (since $\{L(nU) : n \in \mathbb{N}\}$ covers X), and so $L(U) - L(U) \subseteq L(E)$ is an open neighbourhood of 0 in X .

8. Haar null and left Haar null. The two families, which are both left and right translation-invariant (cf. [Sol2, p. 696] – if $\mu \in \mathcal{P}(G)$ witnesses that E is left Haar null, then $\mu_{g^{-1}}$ witnesses that Eg is left Haar null), coincide in Polish abelian groups, and in locally compact second countable groups (where they also coincide with the sets of Haar measure zero – by an application of the Fubini theorem). The former family, however, is in general smaller; indeed, non-Haar null sets need not have the Steinhaus-Weil property – see [Sol2].

9. Beyond local compactness: Haar category-measure duality. In the absence of Haar measure, the definition of left Haar null subsets of a topological group G requires $\mathcal{U}(G)$, the universally measurable sets – by dint of the role of the totality of (probability) measures on G . The natural dual of $\mathcal{U}(G)$ is the class $\mathcal{U}_{\mathcal{B}}(G)$ of *universally Baire sets*, defined for G with a Baire topology as those sets B whose preimages $f^{-1}(B)$ are Baire in any compact Hausdorff space K for any continuous $f : K \rightarrow G$. Initially considered in [FenMW] for $G = \mathbb{R}$, these have attracted continued attention for their role in the investigation of axioms of determinacy and large cardinals – see especially [Woo]; cf. [MarS].

Analogously to the left Haar null sets, define a *left Haar meagre* set as any set M coverable by a universally Baire set B for which there are a compact Hausdorff space K and a continuous $f : K \rightarrow G$ with $f^{-1}(gB)$ meagre in K for all $g \in G$. These were introduced, in the abelian Polish group setting with K metrizable, by Darji [Dar], cf. [Jab1], and shown there to form a σ -ideal of meagre sets (co-extensive with the meagre sets for G locally compact).

10. Metrizability and Christensen's Theorem. An analytic topological group is metrizable; so if also it is a Baire space, then it is a Polish group – [HofT, Th. 2.3.6].

11. Strong Kemperman property: qualitative versus quantitative measure theory. We note that property (*) of the Introduction and Lemma 1 corresponds to the following quantitative property on the line, stated in terms of Haar (i.e. Lebesgue) measure $\eta(\cdot) = |\cdot|$ for sets open in the Lebesgue density topology $\mathcal{D}_{\mathcal{L}}$:

(iv)* strong Kemperman property (see [Kem], [Kuc, Lemma 3.7.2]): for $0 \in U \in \mathcal{D}_{\mathcal{L}}$ there is $\delta > 0$ so that for all $|t| < \delta$

$$|U \cap (t + U)| \geq \varepsilon.$$

This is connected with continuity of the Haar norm. Indeed, since

$$|U \cap (t + U)| = |U| - |U \Delta (t + U)|/2,$$

the inequality above is equivalent to

$$\|t\|_U^\eta := |U \Delta (t + U)| \leq 2(|U| - \varepsilon).$$

The latter holds for any $0 < \varepsilon < |U|$ and for sufficiently small t , by the continuity of the norm $\|t\|_U^\eta$.

12. Proof of Theorem FN of §1. For μ -null $N \subseteq G$ the set $N \times G$ is $\mu \times \nu$ -null, so (by passing to the complement of the null exceptional set of the theorem)

we may assume w.l.o.g. that the exceptional set of A is empty. By inner regularity, it suffices to show that $(\mu \times \nu)(K) = 0$ for all compact $K \subseteq A$.

For K compact, denote by F the (compact) projection of K on the first axis. Let $\varepsilon > 0$. By Prop. 5, for any $x \in F$ there is an open neighbourhood U_x of x and open V_x with $\nu(V_x) < \varepsilon$ and

$$K \cap (U_x \times G) \subseteq R_x := U_x \times V_x.$$

By compactness of F , there are $U^j \times V^j$ for $i = 1, \dots, n$, with U^j, V^j open and $\nu(V^j) < \varepsilon$ such that

$$F \subseteq \bigcup_j U^j : \quad K \subseteq \bigcup_j U^j \times V^j.$$

To disjoint the sets U^j , put

$$S^j := U^j \setminus \bigcup_{j < i} U^j : \quad \bigcup_j U^j = \bigcup_j S^j.$$

Then

$$F = \bigcup_j F \cap S^j : \quad K \subseteq \bigcup_j S^j \times V^j.$$

So

$$\mu(K) \leq \sum_j (\mu \times \nu)(S^j \times V^j) = \sum_j \mu(S^j) \nu(V^j) \leq \sum_j \mu(S^j) \cdot \varepsilon = \varepsilon \mu(F).$$

As $\varepsilon > 0$ was arbitrary, $\mu(K) = 0$. \square

13. *Proof of Proposition 5 (Almost everywhere upper semicontinuity).* As this is much as in Prop. 2, suffice it to indicate the necessary adjustments. For measurable $E \subseteq G$ measurable and compact F with $\mu(F) > 0$, refer to

$$H := \Phi_E(F) = \bigcup_{x \in F} \{x\} \times xE.$$

Instead of the finite union of rectangles K , choose a compact set $K = K_\varepsilon \subseteq H$ with

$$(\mu \times \mu)(H \setminus K) < \varepsilon^2.$$

Then obtain a compact set $C = C_\varepsilon \subseteq F \setminus S_\varepsilon$ with $\mu(F \setminus C) < \varepsilon$ and

$$\mu(H_x \setminus K_x) \leq \varepsilon \quad (x \in C_\varepsilon).$$

For $x \in C_\varepsilon$, by upper semicontinuity of $t \mapsto \mu(K_t)$, there is a neighbourhood U_x^ε of x with

$$\mu(K_y) < \mu(K_x) + \varepsilon \quad (y \in U_x^\varepsilon \cap C).$$

So

$$\mu(H_y) < \mu(K_y) + \varepsilon < \mu(K_x) + 2\varepsilon < \mu(H_x) + 3\varepsilon,$$

i.e.

$$\mu(H_y) < \mu(H_x) + 3\varepsilon \quad (x \in C, y \in U_x^\varepsilon \cap C).$$

Thereafter the inductive argument of Prop. 4 follows almost verbatim. \square

14. Carlson–Simpson Theorem. The shift-compactness property in Th. 3 §3 concerned with embeddings appears to be related to the following van der Waerden-like theorem in Ramsey Theory (cf. [BinO1, §6]). This is due to Carlson and Simpson [CarS] and is a generalization of the famous Hales-Jewett Theorem; for a proof and related literature see [DodKT]:

For every integer $k \geq 2$ and every finite coloring of the set of all (finite) words over $[k] := \{1, \dots, k\}$ there exist a word c over $[k]$ and a sequence of left-variable words $w_n(v)$ over $[k] \cup \{v\}$ such that

$$\{c\} \cup \{c \cdot w_1(a_1) \cdot \dots \cdot w_n(a_n) : a_1, \dots, a_n \in [k] \text{ and } n \in \mathbb{N}\}$$

is monochromatic.

Here \cdot denotes concatenation and $w(a)$ is obtained by substituting a for v . A *word* over S is a string of symbols from the (finite) collection S of symbols (the vocabulary) and a *left-variable word* $w(v)$ over $[k] \cup \{v\}$ means a word over $[k] \cup \{v\}$ beginning with the symbol v , where $v \notin [k]$.

15. Infinite Bernoulli convolutions. The measures above in [BarFF] are related to infinite Bernoulli convolutions with parameter (probability) $p \in (0, 1)$. These have been studied since the work of Erdős in 1939 and 1940; for background and references, see e.g. Cooper [Coo], Solomyak [Solom]. By the law of pure types from probability theory, they are either absolutely continuous (the generic case), or continuous singular. This second case occurs if p is a PV-number – an algebraic number of Pisot-Vijayaraghavan type.

16. Quasi-invariance and the Mackey topology of analytic Borel groups. We stop to comment on the force of full quasi-invariance of a measure in connection with a Steinhaus triple (H, G, μ) with H and G completely metrizable. Both groups, being absolutely Borel, are analytic spaces. So both carry a ‘standard’ Borel structures with H a Borel substructure of G . Mackey [Mac] investigates such Borel groups, defining also a (Borel) measure μ to be ‘standard’ if it has a Borel support. It emerges that every σ -finite Borel measure in an analytic Borel space is standard [Mac, Th. 6.1]. Of interest to us is Mackey’s notion of a ‘measure class’ C_μ , comprising all Borel measures ν

with the same null sets as $\mu : \mathcal{M}_0(\nu) = \mathcal{M}_0(\mu)$. Such a measure class may be closed under translation, and may be right or left invariant; then the common null sets are themselves invariant, and so may be viewed as witnessing quasi-invariance of the measure μ . Mackey shows that a Borel group with a one-sided invariant measure class has a both-sidedly invariant measure class [Mac, Lemma 7.2]; furthermore, if the class is countably generated, then the class contains a left-invariant and a right-invariant measure [Mac, Lemma 7.3]. This enables Mackey to improve on Weil's theorem in showing that an analytic Borel group G with a one-sidedly invariant measure class, in particular one generated by a quasi-invariant measure, has a unique locally compact topology both making G a topological group and generating the given Borel structure.

9 References

- [Amb] W. Ambrose, Measures on locally compact topological groups. *Trans. Amer. Math. Soc.* **61** (1947), 106-121.
- [ArhT] A. Arhangel'skii, M. Tkachenko, *Topological groups and related structures*. World Scientific, 2008.
- [BeeV] G. Beer, L. Villar, Borel measures and Hausdorff distance. *Trans. Amer. Math. Soc.* **307** (1988), 763–772.
- [BalRS] M. Balcerzak, A. Rosłanowski, S. Shelah, Ideals without ccc. *J. Symbolic Logic* **63** (1998), 128-148.
- [Bana] W. Banaszczyk, *Additive subgroups of topological vector spaces*. Lecture Notes in Mathematics **1466**. Springer, 1991.
- [BarFF] A. Bartoszewicz, M. Filipczak, T. Filipczak, On supports of probability Bernoulli-like measures. *J. Math. Anal. Appl.* **462** (2018), 26–35.
- [BarFN] A. Bartoszewicz, M. Filipczak, T. Natkaniec, On Smítal properties. *Topology Appl.* **158** (2011), 2066–2075.
- [Bin] N. H. Bingham, Finite additivity versus countable additivity. *Electronic J. History of Probability and Statistics*, **6** (2010), 35p.
- [BinGT] N. H. Bingham, C. M. Goldie and J. L. Teugels, *Regular variation*, 2nd ed., Cambridge University Press, 1989 (1st ed. 1987).
- [BinO1] N. H. Bingham and A. J. Ostaszewski, Kingman, category and combinatorics. *Probability and mathematical genetics* (Sir John Kingman Festschrift, ed. N. H. Bingham and C. M. Goldie), 135-168, *London Math. Soc. Lecture Notes in Mathematics* **378**, CUP, 2010.

- [BinO2] N. H. Bingham and A. J. Ostaszewski, Normed groups: Dichotomy and duality. *Dissert. Math.* **472** (2010), 138p.
- [BinO3] N. H. Bingham and A. J. Ostaszewski, Dichotomy and infinite combinatorics: the theorems of Steinhaus and Ostrowski. *Math. Proc. Camb. Phil. Soc.* **150** (2011), 1-22.
- [BinO4] N. H. Bingham and A. J. Ostaszewski, Category-measure duality: convexity, mid-point convexity and Berz sublinearity, *Aequationes Mathematicae*, **91** (2017), 801-836 (arXiv:1607.05750).
- [BinO5] N. H. Bingham and A. J. Ostaszewski, Beyond Lebesgue and Baire IV: Density topologies and a converse Steinhaus-Weil theorem. *Topology and its Applications* **239** (2018), 274-292 (arXiv:1607.00031).
- [BinO6] N. H. Bingham and A. J. Ostaszewski, Additivity, subadditivity and linearity: Automatic continuity and quantifier weakening. *Indag. Math. (N.S.)* **29** (2018), 687–713. (arXiv 1405.3948v3).
- [BinO7] N. H. Bingham and A. J. Ostaszewski, Set theory and the analyst, *European J. Math.*, to appear, arXiv:1801.09149v2.
- [BinO8] N. H. Bingham and A. J. Ostaszewski, Beyond Haar and Cameron-Martin: the Steinhaus support. arXiv1805.02325v2.
- [Bog1] V. I. Bogachev, *Gaussian Measures*, Math. Surveys & Monographs **62**, Amer Math Soc., 1998.
- [Bog2] V. I. Bogachev, *Measure theory*. Vol. I, II. Springer-Verlag, Berlin, 2007.
- [Bog3] V. I. Bogachev, *Differentiable measures and the Malliavin calculus*. Math. Surv. Mon. **164**, American Math. Soc, 2010.
- [Bor] K. C. Border, *Fixed point theorems with applications to economics and game theory*. Cambridge University Press, 2nd ed. 1989 (1st ed.1985).
- [Bor] C. Borell, Gaussian Radon measures on locally convex spaces, *Math. Scand.* **36** (1976), 265-284.
- [Bou] A. Bouziad, Continuity of separately continuous group actions in p -spaces. *Topology Appl.* **71** (1996), 119-124.
- [Brz] J. Brzdęk, Subgroups of the group \mathbb{Z}_n and a generalization of the Gołąb-Schinzel functional equation. *Aequat. Math.* **43** (1992), 59–71.
- [CarS] T. J. Carlson and S. G. Simpson, A dual form of Ramsey’s theorem, *Adv. Math.* **53** (1984), 265-290.
- [Chr1] J. P. R. Christensen, On sets of Haar measure zero in abelian Polish groups. Proceedings of the International Symposium on Partial Differential Equations and the Geometry of Normed Linear Spaces (Jerusalem, 1972). *Israel J. Math.* **13** (1972), 255–260 (1973).

- [Chr2] J. P. R. Christensen, *Topology and Borel structure. Descriptive topology and set theory with applications to functional analysis and measure theory*. North-Holland Mathematics Studies **10**, 1974.
- [ChrH] J. P. R. Christensen, W. Herer, On the existence of pathological submeasures and the construction of exotic topological groups. *Math. Ann.* **213** (1975), 203–210.
- [CieJ] K. Ciesielski, J. Jasiński, Topologies making a given ideal nowhere dense or meager. *Topology Appl.* **63** (1995), 277–298.
- [Coo] M. J. P. Cooper, Dimension, measure and infinite Bernoulli convolutions. *Math. Proc. Cambridge Phil. Soc.* **124** (1998), 135–149.
- [DieS] J. Diestel, A. Spalsbury, *The joys of Haar measure*. Grad. Studies in Math. **150**. Amer. Math. Soc., 2014.
- [Dar] U. B. Darji, On Haar meager sets. *Topology Appl.* **160** (2013), 2396–2400.
- [Dod1] P. Dodos, Dichotomies of the set of test measures of a Haar-null set. *Israel J. Math.* **144** (2004), 15–28.
- [Dod2] P. Dodos, On certain regularity properties of Haar-null sets. *Fund. Math.* **191** (2004), 97–109.
- [Dod3] P. Dodos, The Steinhaus property and Haar-null sets. *Bull. Lond. Math. Soc.* **41** (2009), 377–384.
- [DodKT] P. Dodos, V. Kanellopoulos, K. Tyros, A density version of the Carlson and Simpson theorem, *J. Eur. Math. Soc.* **16** (2014), 2097–2164.
- [Drew1] L. Drewnowski, Topological rings of sets, continuous set functions, integration. I, II, III. *Bull. Acad. Polon. Sci. Sér. Sci. Math. Astronom. Phys.* **20** (1972), 269–276; **20** (1972), 277–286; **20** (1972), 439–445.
- [Drew2] L. Drewnowski, On control submeasures and measures. *Studia Math.* **50** (1974), 203–224.
- [Dud] R. M. Dudley. *Real analysis and probability*. Revised ed., Cambridge Studies in Advanced Mathematics **74**. Cambridge University Press, 2002 (1st ed. 1989).
- [Eng] R. Engelking, *General topology*. 2nd ed., Heldermann, 1989 (1st ed. PWN, 1977).
- [Erd1] P. Erdős, On a family of symmetric Bernoulli convolutions. *Amer. J. Math.* **61** (1939), 974–975.
- [Erd2] P. Erdős, On the smoothness properties of Bernoulli convolutions. *Amer. J. Math.* **62** (1940), 180–186.
- [FenMW] Q. Feng, M. Magidor, H. Woodin, Universally Baire sets of reals, in H. Judah, W. Just, H. Woodin (eds.), *Set theory of the continuum*, 203–

- 242, Math. Sci. Res. Inst. Publ. **26**, Springer, 1992.
- [FenN] J. E. Fenstad, D. Normann, On absolutely measurable sets. *Fund. Math.* **81** (1973/74), 91–98.
- [For] M. K. Fort, Jr., A unified theory of semi-continuity. *Duke Math. J.* **16** (1949), 237–246.
- [Fre] D. Fremlin, *Measure theory* Vol. 3: *Measure algebras*. Corrected 2nd printing of the 2002 original. Torres Fremlin, Colchester, 2004.
- [FreNR] D. Fremlin, T. Natkaniec, I. Reclaw, Universally Kuratowski-Ulam spaces. *Fund. Math.* **165** (2000), 239–247.
- [Ful] R. V. Fuller, Relations among continuous and various non-continuous functions. *Pacific J. Math.* **25** (1968), 495–509.
- [Gao] Su Gao, *Invariant descriptive set theory*. Pure and Applied Mathematics **293**. CRC Press, 2009.
- [GivH] S. Givant and P. Halmos, *Introduction to Boolean algebras*, Springer 2009.
- [Gli] I. Glicksberg, Some remarks on absolute continuity on groups. *Proc. Amer. Math. Soc.* **40** (1973), 135–139.
- [Gow1] C. Gowrisankaran, Radon measures on groups. *Proc. Amer. Math. Soc.* **25** (1970), 381–384.
- [Gow2] C. Gowrisankaran, Quasi-invariant Radon measures on groups. *Proc. Amer. Math. Soc.* **35** (1972), 503–506.
- [GroE] K.-G. Grosse-Erdmann, An extension of the Steinhaus-Weil theorem. *Colloq. Math.* **57** (1989), 307–317.
- [Hal] P. R. Halmos, *Measure theory*, Grad. Texts in Math. **18**, Springer 1974 (1st ed. Van Nostrand, 1950).
- [HewR] E. Hewitt, K. A. Ross, *Abstract harmonic analysis*, Vol. I, Grundlehren der math. Wiss. **115**, Springer 1963 [Vol. II, Grundlehren **152**, 1970].
- [Hey] H. Heyer, *Probability measures on locally compact groups*. Ergebnisse Math. Grenzgebiete **94**, Springer, 1977.
- [HodHLS] W. Hodges, I. Hodkinson, D. Lascar, S. Shelah, The small index property for ω -stable ω -categorical structures and for the random graph. *J. London Math. Soc.* **48** (1993), 204–218.
- [HofT] J. Hoffmann-Jørgensen, F. Topsøe, Analytic spaces and their application, in [Rog, Part 3].
- [HolN] L. Holá, B. Novotný, Subcontinuity. *Math. Slovaca* **62** (2012), 345–362.
- [Hu] Z. Hu, A generalized Kakutani-Kodaira theorem. *Proc. Amer. Math. Soc.* **133** (2005), 3437–3440.

- [Jab1] E. Jabłońska, Some analogies between Haar meager sets and Haar null sets in abelian Polish groups. *J. Math. Anal. Appl.* **421** (2015), 1479–1486.
- [Jab2] E. Jabłońska, A theorem of Piccard’s type in abelian Polish groups. *Anal. Math.* **42** (2016), 159–164.
- [Jan] S. Janson, *Gaussian Hilbert spaces*. Cambridge Tracts in Mathematics **129**. Cambridge University Press, 1997.
- [Kal] O. Kallenberg, *Foundations of modern probability*. 2nd ed., Springer, 2002 (1st ed. 1997).
- [Kec] A. S. Kechris, *Classical descriptive set theory*, Graduate Texts in Mathematics **156**, Springer, 1995.
- [Kem] J. H. B. Kemperman, A general functional equation. *Trans. Amer. Math. Soc.* **86** (1957), 28–56.
- [Kha] A. Kharazishvili, Some remarks on the Steinhaus property for invariant extensions of the Lebesgue measure, *Eur. J. Math.*, Online, 2018.
- [Kne] M. Kneser, Summenmengen in lokalkompakten abelschen Gruppen, *Math. Z.* **66** (1956), 88–110.
- [Kod] K. Kodaira, Über die Beziehung zwischen den Massen und den Topologien in einer Gruppe. *Proc. Phys.-Math. Soc. Japan* (3) **23** (1941), 67–119.
- [Kom] Z. Kominek, On an equivalent form of a Steinhaus theorem, *Math. (Cluj)* **30** (53)(1988), 25–27.
- [Kuc] M. Kuczma, *An introduction to the theory of functional equations and inequalities. Cauchy’s equation and Jensen’s inequality*. 2nd ed., Birkhäuser, 2009 (1st ed. PWN, Warszawa, 1985).
- [KucS] M. Kuczma, J. Smítal, On measures connected with the Cauchy equation. *Aequationes Math.* **14** (1976), no. 3, 421–428.
- [Lar] Larsen, R. Measures with separable orbits. *Proc. Amer. Math. Soc.* **19** (1968), 569–572.
- [LeP] R. D. LePage, Subgroups of paths and reproducing kernels, *Ann. Prob.* **1** (1973), 345–347.
- [Loo] L. H. Loomis, *An introduction to abstract harmonic analysis*. Van Nostrand, 1953.
- [LukMZ] J. Lukeš, Jaroslav, J. Malý, L. Zajíček, *Fine topology methods in real analysis and potential theory*. Lecture Notes in Math. **1189**, Springer, 1986.
- [LiuR] T. S. Liu, A. van Rooij, Transformation groups and absolutely continuous measures, *Indag. Math.* **71** (1968), 225–231.
- [LiuRW] T. S. Liu, A. van Rooij, J-K Wang, Transformation groups and ab-

- solutely continuous measures II, *Indag. Math.* **73** (1970), 57–61.
- [Mac] G. W. Mackey, Borel structure in groups and their duals. *Trans. Amer. Math. Soc.* **85** (1957), 134–165.
- [Mah] D. Maharam, An algebraic characterization of measure algebras. *Ann. of Math. (2)* **48** (1947). 154–167.
- [MarS] D. A. Martin, J. R. Steel, Projective determinacy. *Proc. Nat. Acad. Sci. U.S.A.* **85** (1988), 6582–6586.
- [Mic] E. Michael, Topologies on the spaces of subsets. *Trans. Amer. Math. Soc.* **71** (1951), 152–182.
- [MilO] H. I. Miller and A. J. Ostaszewski, Group action and shift-compactness. *J. Math. Anal. App.* **392** (2012), 23–39.
- [MonZ] D. Montgomery, L. Zippin, *Topological transformation groups*. Krieger, 1974 (Interscience, 1955).
- [Mos] Y. V. Mospan, A converse to a theorem of Steinhaus-Weil. *Real An. Exch.* **31** (2005), 291–294.
- [Mue] B. J. Mueller, Three results for locally compact groups connected with the Haar measure density theorem. *Proc. Amer. Math. Soc.* **16** (6) (1965), 1414–1416.
- [Mun] J. R. Munkres, *Topology, a first course*, Prentice-Hall, 1975.
- [Myc] J. Mycielski, Finitely additive measures. *Coll. Math.*, **42** (1979), 309–318.
- [Ost1] A. J. Ostaszewski, Families of compact sets and their universals. *Mathematika* **21** (1974), 116–127.
- [Ost2] A. J. Ostaszewski, Effros, Baire, Steinhaus-Weil and non-separability. *Topology and its Applications* (Mary Ellen Rudin Memorial Volume) **195** (2015), 265–274.
- [Oxt1] Oxtoby, John C. Invariant measures in groups which are not locally compact. *Trans. Amer. Math. Soc.* **60** (1946), 215–237.
- [Oxt2] J. C. Oxtoby, *Measure and category*, 2nd ed. Graduate Texts in Math. 2, Springer, 1980 (1st ed. 1972).
- [PacS] J. Pachl, J. Steprāns, Continuity of convolution and SIN groups. *Canad. Math. Bull.* **60** (2017), 845–854.
- [Par] K. R. Parthasarathy, *Probability measures on metric spaces*. Academic Press, 1967.
- [Pat] A. L. T. Paterson, *Amenability*. Math. Surveys and Mon. **29**, Amer. Math. Soc., 1988
- [Pet] B. J. Pettis, On continuity and openness of homomorphisms in topological groups. *Ann. of Math. (2)* **52** (1950), 293–308.

- [Pic] S. Piccard, Sur les ensembles de distances des ensembles de points d'un espace Euclidien. *Mém. Univ. Neuchâtel* **13**, 212 pp. 1939.
- [Pro] V. Prokaj, A characterization of singular measures, *Real Anal. Exchange* **29** (2003/2004), 805–812.
- [Rog] C. A. Rogers, J. Jayne, C. Dellacherie, F. Topsøe, J. Hoffmann-Jørgensen, D. A. Martin, A. S. Kechris, A. H. Stone, *Analytic sets*, Academic Press, 1980.
- [Rud] W. Rudin, *Functional analysis*, 2nd ed. McGraw-Hill, 1991 (2nd ed.).
- [Sak] S. Saks, *Theory of the integral*. 2nd ed. Dover, 1964 (1st ed. 1937, Monografie Mat. **VII**).
- [Sch] L. Schwartz, *Radon measures on arbitrary topological spaces and cylindrical measures*. Tata Institute of Fundamental Research Studies in Mathematics **6**, Oxford University Press, 1973.
- [Sho] R.M. Shortt, Universally measurable spaces: an invariance theorem and diverse characterizations, *Fundamenta Math.* **121** (1984), 169-176.
- [ShiT] H. Shi, B. S. Thomson, Haar null sets in the space of automorphisms on $[0,1]$. *Real Anal. Exchange* **24** (1998/99), 337–350.
- [Sim] S. M. Simmons, A converse Steinhaus-Weil theorem for locally compact groups. *Proc. Amer. Math. Soc.* **49** (1975), 383-386.
- [Sol1] S. Solecki, Size of subsets of groups and Haar null sets. *Geom. Funct. Anal.* **15** (2005), 246–273.
- [Sol2] S. Solecki, Amenability, free subgroups, and Haar null sets in non-locally compact groups. *Proc. London Math. Soc.* (3) **93** (2006), 693–722.
- [Sol3] S. Solecki, A Fubini theorem. *Topology Appl.* **154** (2007), 2462–2464.
- [Solo] R. M. Solovay, A model of set-theory in which every set of reals is Lebesgue measurable. *Ann. of Math.* (2) **92** (1970), 1–56.
- [Solom] B. Solomyak, Notes on Bernoulli convolutions, in: Fractal geometry and applications: a jubilee of Benoît Mandelbrot. *Proc. Sympos. Pure Math.* **72**, Part 1, 207–230, Amer. Math. Soc., 2004.
- [Ste] H. Steinhaus, Sur les distances des points de mesure positive. *Fund. Math.* **1** (1920), 83-104.
- [Str1] R. Struble, Almost periodic functions on locally compact groups. *Proc. Nat. Acad. Sci. U. S. A.* **39** (1953). 122–126.
- [Str2] R. Struble, Metrics in locally compact groups. *Compositio Math.* **28** (1974), 217–222.
- [Tal] M. Talagrand, Maharam's problem. *Ann. of Math.* (2) **168** (2008), 981–1009.
- [Tam] K.W. Tam, On measures with separable orbit. *Proc. Amer. Math. Soc.*

23 (1969) 409–411.

[TomW] G. Tomkiewicz, S. Wagon, *The Banach-Tarski paradox*. Cambridge University Press, 2016 (1st ed. 1985).

[vDo] E. van Downen, Fubini’s theorem for null sets. *Amer. Math. Monthly* **96** (1989), 718–721.

[Web] H. Weber, FN-topologies and group-valued measures. in E. Pap (ed) *Handbook of measure theory*, Vol. I, 703–743, North-Holland, 2002.

[Wei] A. Weil, *L’intégration dans les groupes topologiques*, Actualités Scientifiques et Industrielles **1145**, Hermann, 1965 (1st ed. 1940).

[Woo] W. H. Woodin, *The axiom of determinacy, forcing axioms, and the nonstationary ideal*. 2nd ed., De Gruyter Series in Logic and its Applications **1**. De Gruyter, 2010.

[Xia] D. X. Xia, *Measure and integration theory on infinite-dimensional spaces*. *Abstract harmonic analysis*. Pure and App. Math. **48**. Academic Press, 1972.

[Yam] Y. Yamasaki, *Measures on infinite-dimensional spaces*. World Scientific, 1985.

[Zak] E. Zakon, A remark on the theorems of Lusin and Egoroff. *Canad. Math. Bull.* **7** (1964), 291–295.

Mathematics Department, Imperial College, London SW7 2AZ; n.bingham@ic.ac.uk
Mathematics Department, London School of Economics, Houghton Street,
London WC2A 2AE; A.J.Ostaszewski@lse.ac.uk