

A Characterization of Wishart Processes and Wishart Distributions

Piotr Graczyk^{*}, Jacek Małecki[†] and Eberhard Mayerhofer[‡]

December 21, 2021

Abstract

A characterization of the existence of non-central Wishart distributions (with shape and non-centrality parameter) as well as the existence of solutions to Wishart stochastic differential equations (with initial data and drift parameter) in terms of their exact parameter domains is given. These two families are the natural extensions of the non-central chi-square distributions and the squared Bessel processes to the positive semidefinite matrices.

^{*}LAREMA, Université d'Angers, France. piotr.graczyk@univ-angers.fr

[†]Faculty of Pure and Applied Mathematics, Wrocław University of Science and Technology, Poland, jacek.malecki@pwr.edu.pl. Supported by the National Science Centre Poland (2013/11/D/ST1/02622).

[‡]University of Limerick, Castletroy, County Limerick, Ireland, eberhard.mayerhofer@ul.ie. Supported by ERC (278295) and SFI (08/SRC/FMC1389).

1 Introduction and Preliminaries

The aim of this paper is to characterize the parameter domain of non-central Wishart distributions (with shape, scale and non-centrality parameters) and that of Wishart processes, a class of positive semi-definite diffusion processes (with drift parameter).

Denote by \mathcal{S}_p the space of symmetric $p \times p$ matrices and let \mathcal{S}_p^+ be the open cone of positive definite matrices, with topological closure $\bar{\mathcal{S}}_p^+$, the positive semi-definite matrices. The classical Gindikin¹ set W_0 is defined as the set of admissible $\beta \in \mathbb{R}$ such that there exists a random matrix X with values in $\bar{\mathcal{S}}_p^+$ (equivalently a measure with support in $\bar{\mathcal{S}}_p^+$) such that its Laplace transform is of the form

$$\mathbb{E}e^{-\text{tr}(uX)} = (\det(I + \Sigma u))^{-\beta}, \quad u \in \bar{\mathcal{S}}_p^+,$$

where $\Sigma \in \mathcal{S}_p^+$. It is well-known (cf. [9], pp. 137, 349) that

$$W_0 = \frac{1}{2}B \cup \left[\frac{p-1}{2}, \infty \right),$$

where $B = \{0, 1, \dots, p-2\}$.

A more intricate question concerns the existence of non-central Wishart distributions, which in addition involves a parameter of non-centrality:

Definition 1.1. *The general non-central Wishart distribution $\Gamma_p(\beta, \omega; \Sigma)$ on $\bar{\mathcal{S}}_p^+$ is defined (whenever it exists) by its Laplace transform*

$$\mathcal{L}(\Gamma_p(\beta, \omega; \Sigma))(u) = (\det(I + \Sigma u))^{-\beta} e^{-\text{tr}(u(I + \Sigma u)^{-1}\omega)}, \quad u \in \bar{\mathcal{S}}_p^+ \quad (1.1)$$

where $\beta > 0$ denotes its shape parameter, $\Sigma \in \mathcal{S}_p^+$ is the scale parameter and the parameter of non-centrality equals $\omega \in \bar{\mathcal{S}}_p^+$.

Random matrices X verifying (1.1) arise in statistics as estimators of the covariance matrix parameter Σ of a normal population. In fact, for the random matrix

$$X = \xi_1 \xi_1^T + \dots + \xi_n \xi_n^T =: q(\xi_1, \dots, \xi_n),$$

where for $i = 1, \dots, n$, $\xi_i \sim \mathcal{N}_p(m_i, \Sigma/2)$ are independent, normally distributed column vectors in \mathbb{R}^p , the Laplace transform of X is given by the right-hand side of (1.1) with $\beta = n/2$ and $\omega = q(m_1, \dots, m_n)$, see Johnson and Kotz [15, Chap.38 (47), p.175].

Accordingly, the pair (ω, β) is said to belong to the non-central Gindikin set W if there exists a random matrix X with values in $\bar{\mathcal{S}}_p^+$ having the Laplace transform (1.1) for a matrix $\Sigma \in \mathcal{S}_p^{+ \ 2}$.

¹The name of this set originates from Gindikin's [12] work in a general multivariate setting.

²If Σ is of maximal rank, this definition is indeed independent of Σ , see Lemma 3.5.

Note the following:

- If $(\omega, \beta) \in W$ then $\beta \geq 0$, otherwise $\mathbb{E}[e^{-\text{tr}(uX)}]$ would be unbounded; and clearly, $(0, \beta) \in W$ if and only if $\beta \in W_0$.
- In the case, where $\text{rank}(\omega) = 1$ and $\beta \neq 0$, the characterization of the non-central Gindikin set W is given in [21]: then $(\omega, \beta) \in W$ if and only if $\beta \in W_0$.
- For $\beta > \frac{p-1}{2}$, Bru [2] shows that Wishart processes have Laplace transform given by (1.1).

The general problem of existence and non-existence of non-central Wishart distributions is studied by Letac and Massam [17]³. In a more recent work Mayerhofer [20] reveals that there is an interplay between the rank of the non-centrality parameter ω and the magnitude of β in the discrete part of the classical Gindikin ensemble: if $(\omega, \beta) \in W$ and $2\beta \in B$, then $\text{rank}(\omega) \leq 2\beta + 1$.

The Laplace transform formulas in Johnson and Kotz [15] and Bru [2] and the results in [20] allow to conjecture⁴ the following:

NCGS Conjecture. *The non-central Gindikin set is characterized by*

$$(\omega, \beta) \in W \Leftrightarrow (2\beta \in [p-1, \infty), \omega \in \bar{\mathcal{S}}_p^+) \text{ or } (2\beta \in B, \text{rank}(\omega) \leq 2\beta).$$

A proof of the NCGS Conjecture has been put forward by the preprint [18]. The proof of [18] is technical⁵ and does not provide an intuitive explanation for the particular parametric restrictions of shape and non-centrality parameter.

The present paper gives a first complete proof of the NCGS conjecture, which reveals and builds on the intimate connection between non-central Wishart distributions and Wishart processes ([2], see also [7, Theorem 1.1]). The latter constitute positive semi-definite solutions $(X_t)_{t \geq 0}$ of stochastic differential equations of the form

$$dX_t = \sqrt{X_t} dW_t + dW_t^T \sqrt{X_t} + \alpha Idt, \quad X_t \in \bar{\mathcal{S}}_p^+, \quad X_0 = x_0 \in \bar{\mathcal{S}}_p^+, \quad (1.2)$$

where $\sqrt{X_t}$ is the unique positive square root of X_t , W is a $p \times p$ matrix of standard Brownian motions, and $\alpha \geq 0$ is the drift parameter.

Wishart processes are natural generalizations of squared Bessel Processes [24]. It is demonstrated in the present paper that the existence of Wishart processes depends crucially on the drift parameter.

The paper proves a necessary and sufficient condition for the existence of Wishart processes, and how this existence issue is related to the one of Wishart distributions. Already Bru [2], who introduces Wishart processes for the first time, realizes the explicit formula for the Laplace transform of X_t :

³However, the statement and proof in [17] are incomplete, as pointed out by [19] and [20].

⁴In [18] and a previous version of this paper, the name *Mayerhofer Conjecture* is used. The conjecture was first presented at the CIMPA Workshop in Hammamet in 2011.

⁵It requires a detailed analysis of the singular and continuous part of certain non-central distributions. Besides, the present version of [18] does not prove that $(w, p) \in W$ implies $(0, p) \in W$.

Proposition 1.2. *Bru([2, Theorem 3]) If the stochastic differential equation (1.2) with $x_0 \in \bar{\mathcal{S}}_p^+$ has a global weak solution in $\bar{\mathcal{S}}_p^+$, then X_t is Wishart distributed for each $t \geq 0$. In particular,*

$$\mathbb{E}^{x_0}[\exp(-\operatorname{tr}(uX_t))] = (\det(I + 2tu))^{-\alpha/2} \exp[-\operatorname{tr}(x_0(I + 2tu)^{-1}u)], \quad u \in \bar{\mathcal{S}}_p^+. \quad (1.3)$$

In the present paper, it is also shown how to construct full-fledged Wishart processes from individual Wishart distributions. The main result is thus a three-fold characterization:

Theorem 1.3. *Let $x_0 \in \bar{\mathcal{S}}_p^+$ and $\alpha \geq 0$. The following are equivalent:*

- (i) *The SDE (1.2) has a global weak solution with $X_0 = x_0$.*
- (ii) *Either $\alpha \geq p - 1$, or $\alpha \in B$ and $\operatorname{rank}(x_0) \leq \alpha$.*
- (iii) *$(x_0, \alpha/2) \in W$.*

Our proof of the NCGS Conjecture (that is, Theorem 1.3 (ii) \Leftrightarrow (iii)) is based on an analysis of affine Wishart semigroups. As a new tool, the action of a class of polynomials on Wishart processes is used, which arise as coefficients of the characteristic polynomial of a symmetric matrix. A full characterization of Wishart processes is provided by (Theorem 1.3 (i) \Leftrightarrow (ii)).

For convenience of the reader, but at the expense of proving an additional implication, Theorem 1.3 is split into two independent theorems in the following two chapters. They require different mathematical tools and therefore can be read independently. Chapter 2 is concerned with the existence of solutions to Wishart stochastic differential equations using elementary stochastic analysis with symmetric polynomials (Theorem 2.4 comprises the equivalence (i) \Leftrightarrow (ii) of Theorem 1.3). Chapter 3 concerns the existence of Wishart distributions (the NCGS conjecture, which comprises (ii) \Leftrightarrow (iii) of Theorem 1.3). Here the Markovian viewpoint is used, in particular the fact that Wishart semigroups are affine Feller semigroups. Finally, in Section 3.3 a conjecture by Damir Filipović [10] on the existence of such semigroups on the cones of lower rank matrices is proved.

2 Gindikin sets for Wishart Processes

This section studies the question of solutions in $\bar{\mathcal{S}}_p^+$ of the Wishart SDE (1.2), using the dynamics of some polynomial functionals of these solutions.

For a symmetric $p \times p$ matrix X , define the elementary symmetric polynomials

$$e_n(X) = \sum_{i_1 < \dots < i_n} \lambda_{i_1}(X) \lambda_{i_2}(X) \dots \lambda_{i_n}(X), \quad n = 1, \dots, p, \quad (2.1)$$

in the eigenvalues $\lambda_1(X) \leq \dots \leq \lambda_p(X)$ of X . Moreover, the convention $e_0(X) \equiv 1$ is used. Up to the sign change, the polynomials e_n are the coefficients of the characteristic polynomial of X , i.e.

$$\det(X - uI) = (-1)^p u^p + (-1)^{p-1} e_1(X) u^{p-1} + \dots - e_{p-1}(X) u + e_p(X)$$

and are polynomial functions of the entries of the matrix X . In particular, $e_p(X) = \det X$.

In [14], symmetric polynomials related to general class of non-colliding particle systems were studied in details. Here similar results are presented, adapted to the matrix SDE

$$dX_t = g(X_t)dW_t h(X_t) + h(X_t)dW_t^T g(X_t) + b(X_t)dt, \quad (2.2)$$

where the continuous functions g, h, b act spectrally⁶ on \mathcal{S}_p and W_t is a Brownian $p \times p$ matrix. Henceforth, abbreviate $\sigma = 2gh$ and $G(x, y) = g^2(x)h^2(y) + g^2(y)h^2(x)$. Furthermore, the natural bijection (2.1) between the eigenvalues $\Lambda = (\lambda_1 \dots \lambda_p)$ and the polynomials $e = (e_1, \dots, e_p)$ is used, extended to the closed Weyl chamber $\bar{C}_+ = \{(x_1, \dots, x_p) \in \mathbb{R}^p : x_1 \leq x_2 \leq \dots \leq x_p\}$, see [14, p.6]. Furthermore, write $\Lambda = \Lambda(e)$ for the inverse bijection on the set $e(\bar{C}_+)$. The notation $e_n^{\bar{i}}$ for the incomplete polynomial of order n , not containing the variable $\lambda_i(e)$, is used; the notation $e_n^{\bar{i}, \bar{j}}$ is analogous. Moreover, set $e_0^{\bar{i}} \equiv 1$ and $e_{-1}^{\bar{i}, \bar{j}} \equiv 0$.

Proposition 2.1. *Let $X = (X_t)_{t \geq 0}$ be a weak solution of (2.2) (with possible finite time blow-up). Then the symmetric polynomials $e_n = e_n(X)$, $n = 1, \dots, p$, are continuous semimartingales described by the system of SDEs ($n = 1, \dots, p$)*

$$de_n = \left(\sum_{i=1}^p \sigma^2(\lambda_i(e))(e_{n-1}^{\bar{i}})^2 \right)^{\frac{1}{2}} dV_n + \left(\sum_{i=1}^p b(\lambda_i(e))e_{n-1}^{\bar{i}} - \sum_{i < j} G(\lambda_i(e), \lambda_j(e))e_{n-2}^{\bar{i}, \bar{j}} \right) dt, \quad (2.3)$$

where V_n are Brownian motions on \mathbb{R} such that $d\langle e_n, e_m \rangle = \sum_{i=1}^p \sigma^2(\lambda_i(e))e_{n-1}^{\bar{i}}e_{m-1}^{\bar{i}}dt$.

Proof. Note that here the equation is considered on \mathcal{S}_p which does not require solutions to live in $\bar{\mathcal{S}}_p^+$ (as is required in reference to Wishart processes). Since the coefficients of the equation (2.2) are continuous, a local weak solution exists. This solution, before its possible blow-up, is considered.

The symmetric polynomials (e_1, \dots, e_p) are given by an analytic function

$$F : \mathcal{S}_p \rightarrow \mathbb{R}^p, \quad X \rightarrow (e_1(X), \dots, e_p(X)),$$

since each elementary symmetric polynomial is given in terms of the coefficients of the matrix X . Thus Itô's formula implies that (e_1, \dots, e_n) are continuous semimartingales (for every starting point x_0 and even when the eigenvalues collide).

The SDEs describing (e_1, \dots, e_p) can be determined similarly as in Propositions 3.1 and 3.2 in [14], which generalize the proof of (4.1) in [2]. One uses the SDEs for the eigenvalues

$$d\lambda_i = 2g(\lambda_i)h(\lambda_i)dB_i + \left(b(\lambda_i) + \sum_{j \neq i} \frac{G(\lambda_i, \lambda_j)}{\lambda_i - \lambda_j} \right) dt, \quad i = 1, \dots, p, \quad (2.4)$$

which are available, according to Theorem 3 from [13], when eigenvalues $\lambda_i(0)$ of x_0 are all distinct and before their eventual collision. However, the Itô formula states that the

⁶ Recall that if $g : \mathbb{R} \mapsto \mathbb{R}$ then $g(X)$ is defined spectrally, i.e. $g(U \operatorname{diag}(\lambda_i)U^T) = U \operatorname{diag}(g(\lambda_i))U^T$, where $U \in SO(p)$.

martingale part and the bounded variation part of (e_1, \dots, e_p) are given in terms of derivatives of the smooth function F and those derivatives have just be determined on the open set $U = \{X \in \mathcal{S}_p : \lambda_i(X) \neq \lambda_j(X) \text{ for all } i \neq j, \text{ with } 1 \leq i, j \leq p\}$. Since the derivatives of F are continuous on \mathcal{S}_p as well as the coefficients in (2.3) (the singular expressions $(\lambda_i - \lambda_j)^{-1}$ appearing in (2.4) are no longer present in (2.3)), one can conclude, by continuity, that the equalities hold on $\bar{U} = \mathcal{S}_p$, i.e. one can drop the conditions that eigenvalues of the initial point are all different and that they are non-colliding for $t > 0$. \square

Using Proposition 2.1 the following characterization of the symmetric polynomials related to Wishart processes is obtained:

Proposition 2.2. *Let X_t be a Wishart process, i.e. a solution of the matrix SDE (1.2). Then the symmetric polynomials $e_n = e_n(X)$, $n = 1, \dots, p$ are semimartingales satisfying the following system of SDEs*

$$de_1 = 2\sqrt{e_1}dV_1 + p\alpha dt, \quad (2.5)$$

$$de_n = M_n(e_1, \dots, e_p)dV_n + (p - n + 1)(\alpha - n + 1)e_{n-1}dt, \quad n = 2, \dots, p - 1, \quad (2.6)$$

$$de_p = 2\sqrt{e_{p-1}e_p}dV_p + (\alpha - p + 1)e_{p-1}dt, \quad (2.7)$$

where V_n , $n = 1, \dots, p$ are one-dimensional Brownian motions and the functions M_n are continuous on \mathbb{R}^p . Furthermore, for $n = 1, \dots, p$, the processes $\mathcal{M}_n(t) := \int_0^t M_n dV_n$ are martingales satisfying

$$\mathbb{E}\left[\int_0^t \langle \mathcal{M}_n, \mathcal{M}_n \rangle_s ds\right] < \infty, \quad \text{for each } t > 0 \text{ and } n = 1, \dots, p. \quad (2.8)$$

Remark 2.3. *Note that by Proposition 2.1, the explicit forms of the martingale parts $d\mathcal{M}_n = M_n(e_1, \dots, e_p)dV_n$ as well as their brackets $d\langle e_n, e_m \rangle$ are known for every $n, m = 1, \dots, p$. Equation (2.5) is given by Bru [2] and is used in the proof of (2.8). Equation (2.7) is just kept for informative reasons. They are both covered by (2.6), by setting $n = 1$ and $n = p$.*

Proof. Applying Proposition 2.1 to the SDE (1.2), one finds that

$$M_n = 2 \left(\sum_{i=1}^p \lambda_i (e_{n-1}^i)^2 \right)^{1/2}. \quad (2.9)$$

Moreover, the drift coefficients of de_n satisfy

$$\sum_{i=1}^p \alpha e_{n-1}^i - \sum_{i < j} (\lambda_i + \lambda_j) e_{n-2}^{\bar{i}, \bar{j}} = (p - n + 1)(\alpha - n + 1)e_{n-1}.$$

It remains to show (2.8), for each $n = 1, \dots, p$. For $n = 1$, by (2.5), $e_1(t)$ is a squared Bessel process. Furthermore, since $e_1(t)$ is non-centrally chi-squared distributed, for each $t > 0$, and for each $m \geq 1$

$$\int_0^t \mathbb{E}[|e_1(s)|^m ds] < \infty, \quad (2.10)$$

hence by Fubini

$$\mathbb{E}\left[\int_0^t |e_1(s)|^m ds\right] < \infty.$$

For $m = 1$, this estimate implies

$$\mathbb{E}\left[\int_0^t \langle \mathcal{M}_1, \mathcal{M}_1 \rangle_s ds\right] < \infty, \quad (2.11)$$

for each $t > 0$. For $1 < n \leq p$ one can use (2.9) to obtain the estimate

$$\langle \mathcal{M}_n, \mathcal{M}_n \rangle = 4 \sum_{i=1}^p \lambda_i(t) (e_{n-1}^i)^2 \leq 4e_{n-1}(t) \leq 4e_1^{2n-2}(t),$$

and thus, by (2.10), one obtains (2.8). \square

Since a Wishart process is $\bar{\mathcal{S}}_p^+$ valued by definition, so $e_n \geq 0$, for all $n = 1, \dots, p$. The idea of the proof of the next Theorem is to show that for $(x_0, \beta) \notin W$, some of the symmetric polynomials e_n become strictly negative.

2.1 Solving the Wishart stochastic differential equations

This section gives a full characterization of the existence of solutions to Wishart SDEs (1.2).

Theorem 2.4. *Let $\alpha \geq 0$, and $x_0 \in \bar{\mathcal{S}}_p^+$. The following are equivalent.*

- (i) *The SDE (1.2) has a global weak solution with $X_0 = x_0$.*
- (ii) *$\alpha \geq p - 1$, or $\alpha \in \{0, 1, \dots, p - 2\}$ and $\text{rank}(x_0) \leq \alpha$.*

Proof. Assume first (i). If $\alpha \geq p - 1$, nothing has to be shown. Suppose, therefore, $\alpha < p - 1$. Recall formulas (2.5)–(2.8) from Proposition 2.2. One can compute explicitly the expected value of the polynomials starting from the first one,

$$\mathbb{E}e_1(t) = e_1(0) + p\alpha \int_0^t ds = e_1(0) + p\alpha t.$$

Therefore

$$\begin{aligned} \mathbb{E}e_2(t) &= e_2(0) + (p-1)(\alpha-1) \int_0^t \mathbb{E}e_1(s) ds \\ &= e_2(0) + (p-1)(\alpha-1)e_1(0)t + p(p-1)\alpha(\alpha-1)\frac{t^2}{2}, \end{aligned}$$

and so on. Consequently $\mathbb{E}e_n(t)$ is a polynomial of degree not greater than n . In particular, the coefficient of t^n is

$$\frac{p(p-1) \cdot \dots \cdot (p-n+1) \cdot \alpha(\alpha-1) \cdot \dots \cdot (\alpha-n+1)}{n!}.$$

If $\alpha \notin B$ and n is the first integer greater than or equal to $\alpha + 1$, then $\mathbb{E}e_n(t)$ is a polynomial of degree n such that the leading coefficient is negative. Consequently, it cannot stay positive for every $t > 0$, which is an impossibility.

If $\alpha = m \in B$, consider $\mathbb{E}e_n(t)$ where $n = m + 1$. Then

$$\mathbb{E}e_n(t) = e_n(0) + (p-n+1)(\alpha-n+1) \int_0^t \mathbb{E}e_{n-1}(s)ds = e_n(0).$$

If $e_n(0) > 0$, then

$$\mathbb{E}e_{n+1}(t) = e_{n+1}(0) + (p-n)(\alpha-n)e_n(0)t,$$

i.e. the leading term is negative and thus $\mathbb{E}e_{n+1}(t) < 0$ for large t . It implies $e_n(0) = 0$, i.e. $\text{rank}(x_0) \leq n-1 = m = \alpha$.

Proof of (ii) \Rightarrow (i):

The existence of global weak solutions for $\alpha \geq p-1$ is proved by Bru [2] (Bru's proof for $\alpha > p-1$ can be easily extended to $\alpha \geq p-1$) and in [3, Theorem 2.6]. Therefore, only the cases $\alpha \in \{0, 1, \dots, p-2\}$ need to be considered. If $\alpha = 0$, then $X = 0$ is the global weak solution of (1.2), for initial value $x_0 = 0$. Let therefore $1 \leq \alpha \leq p-2$, and $\text{rank}(x_0) \leq \alpha$. Let $B_1, B_2, \dots, B_\alpha$ be a sequence of independent, p -dimensional standard Brownian motions, and let $y_1, \dots, y_\alpha \in \mathbb{R}^p$ be such that $x_0 = y_1 y_1^\top + \dots y_\alpha y_\alpha^\top$. Then the process

$$X_t := \sum_{i=1}^{\alpha} (y_i + B_i)(y_i + B_i)^\top$$

is a continuous semimartingale, by construction, and $X_0 = x_0$. Furthermore $dX_t = dM_t + \alpha I dt$, where I is the $p \times p$ unit matrix, and $(M_t)_t$ is a continuous martingale having quadratic variation (A.2). Therefore, by Proposition A.1, the Wishart SDE (1.2) has a global weak solution. \square

Remark 2.5. *Necessity of (ii) can be also proved, if the validity of the NCGS Conjecture is assumed (a fact that is proven in Section 3, and which has not been used above to keep the section self-contained). Suppose the existence of a weak solution. Then by Proposition 1.2, the solution is Wishart distributed, that is, for each $t \geq 0$, $X_t \sim \Gamma_p(\alpha/2, x_0; 2tI)$. By the NCGS Conjecture, $\alpha/2 \in W_0$ and, in addition, if $\alpha < p-1$ then $\text{rank}(x_0) \leq \alpha$.*

3 The NCGS Conjecture and Wishart Semigroups

In this section Wishart semigroups are introduced, which are the main tool for the proof of the NCGS Conjecture in Section 3.2 below. In Section 3.3 all Wishart semigroups on lower rank matrices are characterized.

3.1 Wishart semigroups

For $p \geq 1$, let $D_p(k) \subset \bar{\mathcal{S}}_p^+$ be the sub-cones of rank $\leq k$ matrices, $0 \leq k \leq p$, where clearly $D_p(0) = \{0\}$ and $D_p(p) = \bar{\mathcal{S}}_p^+$. Denote by $f_u(x) = \exp(\text{tr}(-ux))$, where $u, x \in \bar{\mathcal{S}}_p^+$.

Definition 3.1. Let $D \subset \bar{\mathcal{S}}_p^+$ be a closed set. A Wishart semigroup $(P_t)_{t \geq 0}$ on D is a positive, strongly continuous $C_0(D)$ contraction semigroup which for any $u \in \mathcal{S}_p^+$ acts on $f_u|_D$ as

$$P_t f_u(x) = \det(I + 2tu)^{-\alpha/2} e^{-\text{tr}(x(u^{-1} + 2tI)^{-1})}, \quad x \in D. \quad (3.1)$$

Here $\alpha \geq 0$ is called the drift parameter of $(P_t)_{t \geq 0}$.

Note: A Wishart semigroup may or may not exist, depending on the choice of α and D . In Theorem 3.10 below, the existence of Wishart semigroups for $D = D_p(k)$ is characterized.

The following remark summarizes several essential properties of Wishart semigroups:

Remark 3.2. Let $(P_t)_{t \geq 0}$ be a Wishart semigroup with drift parameter α .

- (i) (Markovian representation) In view of the Riesz representation theorem for positive functionals [23, Chapter 2.14], for each $t > 0$, $x \in D$ there exists a positive measure $p_t(x, d\xi)$ on D such that

$$P_t f(x) = \int_D f(\xi) p_t(x, d\xi). \quad (3.2)$$

Furthermore, the semigroup property of $(P_t)_{t \geq 0}$ implies, that $p_t(x, d\xi)$ satisfies the Chapman-Kolmogorov equations, thus $p_t(x, d\xi)$ is a Markov transition function. Hence, the semigroup has a stochastic representation as a Markov process $(\mathbb{P}^x)_{x \in D}$, where for each $x \in D$, \mathbb{P}^x denotes the resulting probability on the canonical path space $D^{\mathbb{R}^+}$ with initial law δ_x , and $X_t(\omega) := \omega(t)$, where $\omega \in D^{\mathbb{R}^+}$. The Markov process (X, \mathbb{P}^x) is called the canonical representation of the semigroup $(P_t)_{t \geq 0}$.

- (ii) (Càdlàg Paths) It is a well-established fact, that any Feller process (that is, a Markov process with strongly continuous C_0 semigroup), has a càdlàg version.
- (iii) (Affine Property) By definition, Wishart semigroups are affine semigroups (see [3]), that is, the Laplace transform of their transition function is of the form

$$\mathbb{E}[e^{-\text{tr}(uX_t)} \mid X_0 = x] = e^{-\phi(t,u) - \text{tr}(\psi(t,u)x)}, \quad (3.3)$$

where

$$\phi(t, u) = \frac{\alpha}{2} \log(\det(I + 2tu)), \quad \psi(t, u) = (u^{-1} + 2tI)^{-1}.$$

- (iv) (Wishart transition function) By definition, the Markovian transition function of a Wishart semigroup $p_t(x, d\xi)$ is $\Gamma_p(\alpha/2, x; 2tI)$ distributed, for each $t \geq 0$ and for all $x \in D$. Furthermore, the support of $\Gamma_p(\alpha/2, x; 2tI)$ is contained in D .

- (v) *(Non-Explosion)* $(P_t)_{t \geq 0}$ is conservative: Let $u_n \in \mathcal{S}_p^+$ such that $u_n \rightarrow 0$ as $n \rightarrow \infty$. By (3.2) and Lebesgue's dominated convergence theorem one thus has

$$P_t 1 = \lim_{n \rightarrow \infty} P_t f_{u_n}(x) = 1.$$

- (vi) *(Semimartingales and Continuity)* If, in addition, one assumes that the linear span of D has non-empty interior, (X, \mathbb{P}_x) for each x is an affine semimartingale, that is, a semimartingale with differential characteristics which are affine functions in the state variable. The continuity of the sample paths of X follows. For more details, see Appendix A.
- (vii) *(Strong Maximum Principle)* For a strongly continuous C_0 semigroup $(P_t)_{t \geq 0}$ with infinitesimal generator \mathcal{A} , the following are equivalent

- (a) \mathcal{A} satisfies the strong maximum principle, that is, $\mathcal{A}f(x_0) \geq 0$, for any $f \in C_0$ that satisfies $f(x) \geq f(x_0)$.
- (b) $(P_t)_{t \geq 0}$ is positive (hence a Feller semigroup).

The proof of (a) \Rightarrow (b) is simple. A proof of the non-trivial implication (b) \Rightarrow (a) employs the positivity of the Yoshida approximations of \mathcal{A} ([8, Corollary 2.8]).

Wishart semigroups on $D = \bar{\mathcal{S}}_p^+$ are well understood; they are the semigroups associated with affine diffusion processes on D . By [3, Theorem 2.4] the following are equivalent:

- The Wishart semigroup with drift parameter α exists with state space $D = \bar{\mathcal{S}}_p^+$.
- $\alpha \geq p - 1$.

However, for strict subsets $D \subset \bar{\mathcal{S}}_p^+$, less is known about Wishart semigroups. In Theorem 3.10 below a new result for the sets of rank $k \leq p - 1$ matrices is given.

Let \mathcal{S}_p^* be the space of rapidly decreasing smooth functions on \mathcal{S}_p , and for a subset $D \subseteq \bar{\mathcal{S}}_p^+$, let $\mathcal{S}_p^*(D) = \{f \mid_D \mid f \in \mathcal{S}_p^*\}$.

For any $f \in \mathcal{S}_p^*(D)$, the action of the following differential operator is well-defined,

$$\mathcal{A}^\sharp f(x) = 2 \operatorname{tr}(x \nabla^2) f(x) + \alpha \operatorname{tr}(\nabla f(x)), \quad (3.4)$$

where the notation of Bru [2]

$$x \nabla^2 := x \cdot \nabla \cdot \nabla$$

is used, with \cdot denoting the matrix multiplication, and ∇ being the matrix of partial differential operators $\nabla = (\nabla_{ij})_{ij}$, where $\nabla_{ij} = \frac{\partial}{\partial x_{ij}}$. This expression reads in canonical coordinates, (cf. the notation of [3, Theorem 2.4])

$$2 \operatorname{tr}(x \nabla^2) = \sum_{i,j,k,l} A(x)_{i,j,k,l} \frac{\partial^2}{\partial x_{ij} \partial x_{kl}},$$

where $A(x)$ is a quadratic form on $(\bar{\mathcal{S}}_p^+)^2$, defined in coordinates as

$$A(x)_{i,j,k,l} = x_{ik} \delta_{jl} + x_{il} \delta_{jk} + x_{jk} \delta_{il} + x_{jl} \delta_{ik}.$$

Proposition 3.3. *Suppose $D_p(1) \subseteq D$, and let $(P_t)_{t \geq 0}$ be a Wishart semigroup on D with infinitesimal generator \mathcal{A} . Then $S_p^*(D) \subset \mathcal{D}(\mathcal{A})$ and $\mathcal{A}f = \mathcal{A}^\# f$ in (3.4) for any $f \in S_p^*(D)$.*

Proof. It is first proved that

$$\mathcal{A}f_u^D = (\mathcal{A}^\# f_u) |_D, \quad (3.5)$$

for any exponential $f_u^D(\cdot) := e^{-\text{tr}(u \cdot)} |_D$. Here the right hand side involves differentiation on the open domain \mathcal{S}_p , and later restriction to D , whereas on the left hand side \mathcal{A} acts directly on f_u^D .

By the definition of the affine property (3.3),

$$\mathcal{A}f_u(x) = (F(u) + \text{tr}(R(u)x)f_u(x), \quad x \in D, \quad (3.6)$$

for $f_u(x) = \exp(-\text{tr}(ux))$ and $u \in \bar{\mathcal{S}}_p^+$, and thus $f_u^D \in \mathcal{D}(\mathcal{A})$. Here

$$F(u) = \frac{\partial \phi(t, u)}{\partial t} \Big|_{t=0} = \alpha \text{tr}(u) \quad (3.7)$$

and

$$R(u) = \frac{\partial \psi(t, u)}{\partial t} \Big|_{t=0} = -2u^2, \quad (3.8)$$

where the differentiation rules for inverse map and determinant ([9, Proposition III.4.2 (ii) and Proposition II.3.3 (i)]) have been used. The assumption that D contains rank one matrices implies that the convex hull of D equals $\bar{\mathcal{S}}_p^+$, and thus F and R are uniquely determined, as the coefficients of the affine (in the state variable x) function

$$x \mapsto F(u) + \text{tr}(xR(u)).$$

A straightforward computation reveals that the action of $\mathcal{A}^\#$ on f_u^D coincides with (3.6), hence (3.5) holds.

According to the density argument [3, Theorem B.3], the linear hull of such exponentials for strictly positive definite u is dense in the space of rapidly decreasing functions on $\bar{\mathcal{S}}_p^+$ and thus equality in (3.5) extends, by convergence properties in the Schwarz class and the closedness of \mathcal{A} , to rapidly decreasing functions. \square

Recall that a time-homogenous Markov process is polynomial if the action of its semigroup can be extended to polynomials of any order ([4, Definition 2.1]).

Proposition 3.4. *Suppose $(P_t)_{t \geq 0}$ is a Wishart semigroup supported on $D \subset \bar{\mathcal{S}}_p^+$ with drift $\alpha \geq 0$. $(P_t)_{t \geq 0}$ is polynomial and its infinitesimal generator acts on symmetric polynomials as follows*

$$\mathcal{A}e_n(x) = (p - n + 1)(\alpha - n + 1)e_{n-1}(x), \quad x \in D, \quad 1 \leq n \leq p. \quad (3.9)$$

Proof. By Proposition A.2, there is a version $(\tilde{X}_t)_{t \geq 0}$ of $(X_t)_{t \geq 0}$ which is a Wishart semimartingale, and thus by Proposition A.1 there exists a $d \times d$ dimensional Brownian motion W such that the pair $((\tilde{X}_t)_{t \geq 0}, W)$ constitutes a global weak solution of the Wishart SDE.

Hence Proposition 2.2 may be applied, that yields the SDE dynamics (2.5)–(2.7). By (2.8), $\int_0^t M_n dV_n$ are true martingales, hence

$$\mathbb{E}^x[e_n(t)] = e_n(x) + (p - n + 1)(\alpha - p + 1)\mathbb{E}^x\left[\int_0^t e_{n-1}(s)ds\right],$$

thus by Lebesgue's dominated convergence theorem,

$$\mathcal{A}e_n(x) = \lim_{t \downarrow 0} \frac{P_t e_n(x) - e_n(x)}{t} = (p - n + 1)(\alpha - p + 1)e_{n-1}(x).$$

□

An equivalence relation \simeq on the space of random variables with values in $\bar{\mathcal{S}}_p^+$ is introduced by defining $X \simeq Y$ if and only if for all $0 \leq r \leq p$

$$\mathbb{P}[\text{rank}(X) = r] > 0 \text{ if and only if } \mathbb{P}[\text{rank}(Y) = r] > 0.$$

Three technical lemmas are useful:

Lemma 3.5. *Let $\beta \geq 0, \omega, \in \bar{\mathcal{S}}_p^+$ and $\Sigma \in \mathcal{S}_p^+$.*

- (i) *(linear automorphism) Let $\Sigma = qq^\top$, where q is a real $p \times p$ matrix. If $X \sim \Gamma_p(\beta, \omega; I)$, then $Y = qXq^\top \sim \Gamma_p(\beta, q\omega q^\top; \Sigma)$ and $Y \simeq X$. Conversely, $Y \sim \Gamma_p(\beta, q\omega q^\top; \Sigma)$ implies $X = q^{-1}Y(q^{-1})^\top \sim \Gamma_p(\beta, \omega; I)$.*
- (ii) *(exponential family) If $X \sim \mu(d\xi) \sim \Gamma_p(\beta, \omega; I)$, then for $v := \Sigma^{-1} - I$ there exists a random variable Y distributed as*

$$Y \sim \frac{\exp(\text{tr}(v\xi))\mu(d\xi)}{\mathbb{E}[\exp(\text{tr}(vX))]} \sim \Gamma_p(\beta, \Sigma\omega\Sigma; \Sigma)$$

and $Y \simeq X$. Conversely, $Y \sim \Gamma_p(\beta, \Sigma\omega\Sigma; \Sigma)$ implies that $X \sim \Gamma_p(\beta, \omega; I)$

- (iii) *If $X \sim \Gamma_p(\beta, \omega; \Sigma)$ then $\Gamma_p(\beta, \tilde{\omega}; \tilde{\Sigma})$ exists for any $\tilde{\omega}$ satisfying $\text{rank}(\tilde{\omega}) \leq \text{rank}(\omega)$ and for any $\tilde{\Sigma} \in \bar{\mathcal{S}}_p^+$.*

Proof. The equivalence relation in (i) holds, since any linear automorphism maintains the rank of matrices. The remaining claims in (i) follow from the following chain of identities, using the very definition of the Wishart distribution in terms of its Laplace transform (using multiplicativity of the determinant and the cyclic property of the trace):

$$\begin{aligned} \mathbb{E}[e^{-\text{tr}(uY)}] &= \mathbb{E}[e^{-\text{tr}(uqXq^\top)}] = \mathbb{E}[e^{-\text{tr}((q^\top uq)X)}] = (\det(I + q^\top uq))^{-\beta} e^{-\text{tr}(q^\top uq(I + q^\top uq)^{-1}\omega)} \\ &= (\det(I + \Sigma u))^{-\beta} e^{u(I + \Sigma u)^{-1}q\omega q^\top}, \end{aligned}$$

i.e. $Y \sim \Gamma_p(\beta, q\omega q^\top; \Sigma)$.

Proof of (ii): Note that due to Proposition B.1, $v = -I + \Sigma^{-1} \in D(\mu)$ and (1.1) holds for v . Hence the first part of the proof of (ii) follows the lines of the proof of [20, Proposition

3.1 (ii)]. Conversely, let $Y \sim \mu_1 = \Gamma_p(\beta, \Sigma\omega\Sigma; \Sigma)$. Then $v_1 = -\Sigma^{-1} + I \in D(\mu_1)$ and, after a few computations, one obtains

$$\int e^{-\text{tr}((u+v_1)\xi)} \mu_1(d\xi) = ((\det(\Sigma))^{-\beta} e^{-\text{tr}((\Sigma-I)\omega)}) (\det(I+u))^{-\beta} e^{-\text{tr}(u(I+u)^{-1}\omega)},$$

where the pre-factor is recognized as

$$(\det(\Sigma))^{-\beta} e^{-\text{tr}((\Sigma-I)\omega)} = \mathbb{E}[e^{-\text{tr}(v_1 Y)}],$$

and the second factor equals

$$(\det(I+u))^{-\beta} e^{-\text{tr}(u(I+u)^{-1}\omega)} = \mathbb{E}[e^{-\text{tr}(uX)}]$$

for $X \sim \Gamma_p(\beta, \omega; I)$.

Finally, for any $u \in -\Sigma^{-1} + \mathcal{S}_p^+$, let

$$\nu(d\xi) := \frac{\exp(-\text{tr}(u\xi))\mu(d\xi)}{\mathbb{E}[\exp(-\text{tr}(uX))]}.$$

Then $\nu(B) > 0$ if and only if $\mu(B) > 0$, for any Borel set $B \subset \bar{\mathcal{S}}_p^+$. Hence $Y \simeq X$ in (ii).

Proof of (iii): Let $\text{rank}(\omega) = r$ with $0 \leq r \leq p$. The following outlines the transformations that map $\Gamma_p(\beta, \omega; \Sigma)$ onto $\Gamma_p(\beta, \omega_1; \Sigma_1)$.

Suppose first $\text{rank}(\omega_1) = r$ and that $\Sigma_1 = q_1 q_1^\top$ is of full rank. By properties of the Natural Exponential Family (ii), one obtains $\Gamma_p(\beta, \Sigma^{-1}\omega\Sigma^{-1}; I)$. By (i) the transformation $\xi \mapsto q_a \xi q_a^\top$, where q_a is an invertible but not necessarily symmetric matrix, yields $\Gamma_p(\beta, q_a \Sigma^{-1}\omega\Sigma^{-1} q_a^\top; \Sigma_a)$, where $\Sigma_a := q_a q_a^\top$. Again using (ii) yields

$$\Gamma_p(\beta, \Sigma_a^{-1} q_a \Sigma^{-1} \omega \Sigma^{-1} q_a^\top \Sigma_a^{-1}; I) = \Gamma_p(\beta, (q_a^{-1})^\top \Sigma^{-1} \omega \Sigma^{-1} q_a^{-1}; I)$$

Finally, by (i), the linear transformation $\xi \mapsto q_1 \xi q_1^\top$ yields

$$\Gamma_p(\beta, q_1 (q_a^{-1})^\top \Sigma^{-1} \omega \Sigma^{-1} q_a^{-1} q_1^\top; \Sigma_1)$$

Note that q_a has not been specified yet. Since the linear automorphism group acts transitively on $\bar{\mathcal{S}}_p^+$ and maintains ranks, there exists q_a such that

$$q_1 (q_a^{-1})^\top \Sigma^{-1} \omega \Sigma^{-1} q_a^{-1} q_1^\top = \omega_1,$$

and thus one obtains the existence of $\Gamma_p(\beta, \omega_1; \Sigma_1)$ for any invertible Σ_1 and any ω_1 with $\text{rank}(\omega_1) = r$.

Finally, let $\text{rank}(\tilde{\omega}) \leq \text{rank}(\omega) = r$ and let $\tilde{\Sigma}$ be not necessarily invertible. Let $(\omega_n)_n$ be a sequence of non-centrality parameters ω_n such that $\lim_{n \rightarrow \infty} \omega_n = \tilde{\omega}$, where $\text{rank}(\omega_n) = r$ for each n , and let $(\Sigma_n)_n$ be a sequence of non-singular matrices Σ_n such that $\lim_{n \rightarrow \infty} \Sigma_n = \tilde{\Sigma}$.

By the previous arguments,

$$\Gamma_p(\beta, \omega_n; \Sigma_n)$$

exists for any $n \in \mathbb{N}$. By Proposition B.1, for each n , the characteristic functions are of the same form, and converge for any $u \in i\mathcal{S}_p$ as $n \rightarrow \infty$ to

$$\left(\det(I + \tilde{\Sigma}u)\right)^{-\beta} e^{-\text{tr}(u(I + \tilde{\Sigma}u)^{-1}\tilde{\omega})}$$

Hence, by Lévy's continuity theorem, the limit is the characteristic function of a positive measure on $\bar{\mathcal{S}}_p^+$, namely $\Gamma_p(\beta, \tilde{\omega}; \tilde{\Sigma})$. \square

Lemma 3.6. *Let Ξ be a positive semi-definite random matrix supported on $D_p(r-1)$ and $\text{rank}(\Xi) = r-1$ with nonzero probability, where $1 \leq r \leq p$. Let further $\eta \sim \mathcal{N}(\mu, \Sigma)$ with $\mu \in \mathbb{R}^p$ and with covariance matrix $\Sigma \in \mathcal{S}_p^+$. If Ξ and η are independent, then $\text{rank}(\Xi + \eta\eta^\top) = r$ with nonzero probability.*

Proof. Assume first the constant case $\Xi = \Xi_0 \in \bar{\mathcal{S}}_p^+$. Without loss of generality, one may assume $\Xi_0 = \text{diag}(I_{r-1}, 0)$, where I_k is the $k \times k$ unit matrix. Define

$$V = \begin{pmatrix} I_{r-1} & -\Omega \\ 0 & I_{p-r+1} \end{pmatrix}$$

with a $(r-1) \times (p-r+1)$ matrix $\Omega_{ij} = \delta_{ij} \frac{\eta_i}{\eta_{r-1+j}}$. Then

$$V(\Xi_0 + \eta\eta^\top)V^\top = \text{diag}(I_{r-1}, (\eta\eta^\top)_{r \leq i, j \leq p})$$

and since $(\eta_k)_{r \leq k \leq p} \sim \mathcal{N}((\mu_k)_{r \leq k \leq p}, (\Sigma_{ij})_{r \leq i, j \leq p})$, it follows that $\eta\eta^\top$ has rank 1 almost surely. Thus $\text{rank}(V(\Xi_0 + \eta\eta^\top)V^\top) = r-1 + 1 = r$ almost surely.

Now consider a random matrix Ξ . Clearly, $\text{rank}(\Xi + \eta\eta^\top) \leq r$. The set $A_\Xi := \{\text{rank}(\Xi(\omega)) = r-1\}$ is Borel, since for $r=1$ it is precisely the set $\{\text{tr}(\Xi) = 0\}$, and for $r > 1$ one has $A_\Xi = \{e_{r-1}(\Xi) = 0\}^c \cap \{e_r(\Xi) = 0\}$. By assumption $\mathbb{P}[A_\Xi] > 0$, thus the first part of the proof implies

$$\mathbb{E}[\text{rank}(\Xi + \eta\eta^\top) \mid \text{rank}(\Xi) = r-1] = r$$

and thus $\text{rank}(\Xi + \eta\eta^\top) = r$ almost surely on A_Ξ . \square

Lemma 3.7. *Suppose $\Xi_0 \in \bar{\mathcal{S}}_p^+$ with $\text{rank}(\Xi_0) = p-1$, and let $\Xi \sim \Gamma_p((p-1)/2, \Xi_0; \Sigma)$, where Σ is non-degenerate. Then $\text{rank}(\Xi) = p-1$ almost surely.*

Proof. By Lemma 3.5 (i), the automorphism $\xi \rightarrow q^{-1}\xi q^{-1}$ with $q = \sqrt{\Sigma}$ yields $q^{-1}\Xi q^{-1} \sim \Gamma_p((p-1)/2, q^{-1}\Xi_0 q^{-1}; I)$, and since $\text{rank}(\Xi_0) = \text{rank}(q^{-1}\Xi_0 q^{-1})$, and $\Xi \simeq q^{-1}\Xi q^{-1}$, one may without loss of generality assume $\Sigma = 2I$.

Let $\mu_i \in \mathbb{R}^p$ for $i = 1, \dots, p-1$ such that $\mu_1\mu_1^\top + \dots + \mu_{p-1}\mu_{p-1}^\top = \Xi_0$. Let x_{ij} , $1 \leq i \leq p$, $1 \leq j \leq p-1$ be a sequence of independent standard normally distributed random variables, and set $x_j = (x_{ij})_{1 \leq i \leq p}$ and $y_j = x_j + \mu_j$. Then (see [19, Section 1]) the random variable

$$X = \sum_{j=1}^{p-1} y_j y_j^\top$$

is $\Gamma_p(\frac{p-1}{2}, \Xi_0; 2I)$ distributed. Furthermore, $x := (x_{ij})_{ij}$ has rank $p-1$ almost surely, hence X has rank $p-1$ almost surely, and thus also Ξ . \square

The following statement concerns the support of Wishart distributions with general shape parameter.

Proposition 3.8. *Suppose $\beta \in \{0, 1/2, \dots, (p-2)/2\}$ and $\Sigma \in \mathcal{S}_p^+$. Suppose $\text{rank}(\omega) = 2\beta + k$, where $1 \leq k \leq p - (2\beta + 1)$. Then $\Gamma_p(\beta, \omega; \Sigma)$, if exists, is supported in $D_p(2\beta)$. In other words, almost surely,*

$$\text{rank}(\Xi) \leq 2\beta \quad (3.10)$$

for any $\Xi \sim \Gamma_p(\beta, \omega; \Sigma)$.

Proof. Suppose first $\beta = 0$ and $\text{rank}(\omega) \geq 1$. Then, also $\Gamma_p(0, \tilde{\omega}; 2tI)$ exists, with $\text{rank}(\tilde{\omega}) = 1$, see Lemma 3.5 (iii). Let $x \in \tilde{\mathcal{S}}_p^+$, then one can write

$$x = \sum_{i=1}^p \mu_i \mu_i^\top, \quad \mu_i \in \mathbb{R}^p$$

Let $t > 0$ be fixed. By Lemma 3.5 (iii), there exist independent random variables $\Xi_i \sim \Gamma_p(\beta = 0, \mu_i \mu_i^\top; 2tI)$, for $i = 1, \dots, p$, and therefore

$$\Xi = \Xi_1 + \dots + \Xi_p \sim \Gamma_p(0, x; 2tI),$$

and thus a transition function of a Wishart semigroup with zero drift is constructed, violating the drift condition for affine Markov processes on $\tilde{\mathcal{S}}_p^+$ [3, Theorem 2.4 and Definition 2.3, equation (2.4)] (which rules out drifts strictly below $(p-1)/2$). Thus $\Gamma_p(\beta, \omega; \Sigma)$ does not exist.

Let now $\beta \in \{1/2, \dots, (p-2)/2\}$, then, since $2\beta + k \geq 2\beta + 1 \geq 2$, there is nothing to show when $p \leq 2$. Set therefore $p \geq 3$. Then,

- $\beta' := (p-1)/2 - \beta$ satisfies $1/2 \leq \beta' \leq (p-2)/2$.
- Since

$$2 \leq \text{rank}(\omega) = 2\beta + k \leq 2\beta + (p - (2\beta + 1)) = p - 1$$

there exists $\omega' \in \tilde{\mathcal{S}}_p^+$ with $\text{rank}(\omega') = (p-1) - \text{rank}(\omega) = (p-1) - (2\beta + k)$ and such that $\omega_* := \omega + \omega'$ satisfies $\text{rank}(\omega_*) = p - 1$. Furthermore, since

$$\text{rank}(\omega') = p - 1 - (2\beta + k) = 2\beta' - k \leq 2\beta'$$

a random variable $Y \sim \Gamma_p(\beta', \omega'; \Sigma)$ exists, independent of Ξ : Let $m_i \in \mathbb{R}^p$ ($i = 1, \dots, n := 2\beta'$) such that

$$m_1 m_1^\top + \dots + m_n m_n^\top = \omega'$$

and ξ_j ($j = 1, \dots, n$) be a sequence of independent, normally distributed random variables with mean m_j , and variance $\Sigma/2$, and independent of Ξ . Then $Y := \xi_1 \xi_1^\top + \dots + \xi_n \xi_n^\top \sim \Gamma_p(\beta', \omega'; \Sigma)$, see the remark following Definition 1.1.

The sum $\Xi' = \Xi + Y$ is $\Gamma_p((p-1)/2, \omega_*, \Sigma)$ distributed. Since $\text{rank}(\omega_*) = p - 1$, Lemma 3.7 applies and yields $\text{rank}(\Xi') = p - 1$ almost surely. Thus, by Lemma 3.6 (applied exactly $2\beta'$ times, since Y is constructed by a sum of $2\beta'$ squares of independent, normally distributed vectors) one must have $\text{rank}(\Xi) \leq 2\beta$ almost surely, as otherwise $\text{rank}(\Xi') > p - 1$ with non-zero probability. \square

3.2 Proof of the NCGS Conjecture.

Proof of \Leftarrow : Sufficiency of conditions in NCGS Conjecture was shown for $2\beta \in B$ in [15, Chap.38 (47), p.175] and for $2\beta > p - 1$ in [2]. The case $2\beta = p - 1$ follows from the case $2\beta > p - 1$ by Lévy continuity theorem arguments [19, 20].

Proof of \Rightarrow : Conversely, suppose the existence of a single distribution $\Gamma_p(\beta, \omega; I)$. Then by Lemma 3.5 (iii), also $\Gamma_p(\beta, 0; I)$ exists. Since the latter is a classical Wishart distribution with non-degenerate scale parameter, $\beta \in W_0$, the classical Gindikin set.

Let $\beta \in \{0, 1/2, \dots, (p-2)/2\}$ and assume, for a contradiction, $\text{rank}(\omega) = 2\beta + l$, where $1 \leq l \leq p - 2\beta$. By Lemma 3.5 (iii) one can obtain non-central Wishart distributions for $\Gamma_p(\beta, \omega'; \Sigma)$ with any $\text{rank}(\omega') \leq 2\beta + l$ and any invertible Σ .

Using, in addition, the support information of Proposition 3.8, one thus obtains a Wishart semigroup $(P_t)_{t \geq 0}$ with state space $D_p(2\beta + l)$ and with drift 2β , by creating $\Gamma_p(\beta, x; 2tI)$, for each $t > 0$, and for each x with $\text{rank}(x) \leq 2\beta + l$. Denote by \mathcal{A} the infinitesimal generator of $(P_t)_{t \geq 0}$.

Distinguish the following two cases.

- (i) $l < p - 2\beta$. Since for all $x \in D_p(2\beta + l)$, $e_{2\beta+l+1}(x) = 0$,

$$\begin{aligned} 0 &= \lim_{t \rightarrow 0} \frac{P_t e_{2\beta+l+1}(x) - e_{2\beta+l+1}(x)}{t} = \mathcal{A} e_{2\beta+l+1}(x) = \\ &= (p - (2\beta + l))(-\beta - l) e_{2\beta+l}(x) \neq 0, \quad \text{for all } x \text{ with } \text{rank}(x) = 2\beta + l, \end{aligned}$$

which is a contradiction. Here, for the last identity Proposition 3.4 has been used.

- (ii) $l = p - 2\beta$. Then $\text{rank}(\omega) = p$ and the semigroup $(P_t)_{t \geq 0}$ acts on $C_0(\bar{S}_p^+)$. The positivity of the Feller semigroup implies that its infinitesimal generator \mathcal{A} satisfies the positive maximum principle. Applied to $e_p(x) = \det(x)$ this implies that

$$\mathcal{A} \det(x_0) \geq 0$$

for any x_0 with $\text{rank}(x_0) < p$. Choose x_0 with $\text{rank}(x_0) = p - 1$, then $e_{p-1}(x_0) > 0$, and therefore by Proposition 3.4 (setting $n = p$ and recalling that $e_p = \det$)

$$\mathcal{A} \det(x_0) = (2\beta - p + 1) e_{p-1}(x_0) < 0$$

because $\beta \in \{0, 1, \dots, \frac{p-2}{2}\}$, by assumption. This violates the positive maximum principle.

These two contradictions imply that indeed $\text{rank}(\omega) \leq 2\beta$, whenever $\beta \in \{0, \dots, \frac{p-2}{2}\}$, and thus the proof of the NCGS conjecture is finished.

Remark 3.9. *Let us mention another proof of the necessity in the NCGS. As above, the existence of a single distribution $\Gamma_p(\beta, \omega; I)$ implies the existence of a Wishart semigroup $(P_t)_{t \geq 0}$ with state space $D_p(2\beta + l)$ and with drift 2β . By Proposition A.2(ii), the Wishart SDE (1.2) has a global weak solution with $X_0 = \omega$. The proof is completed by using Theorem 2.4.*

3.3 A Characterization of Wishart Semigroups

The paper concludes with the following characterization of Wishart semigroups with state spaces $D_p(k)$, the $p \times p$ symmetric positive semi-definite matrices of rank $\leq k$.⁷ The statement has been conjectured by Damir Filipović [10] in 2009.

Theorem 3.10. *Let $k \in \{1, \dots, p\}$ and let $\alpha \geq 0$. The following are equivalent:*

- (i) *The Wishart semigroup with state-space $D = D_p(k)$ exists.*
- (ii) *If $k \in \{1, \dots, p-1\}$, then $\alpha = k$, and if $k = p$, then $\alpha \geq p-1$.*

Proof. If $k = p$, that is $D = \bar{\mathcal{S}}_p^+$, then $\alpha \geq p-1$ due to [3], which also includes a proof of existence. Therefore, only the cases $k < p$ require a proof:

Proof of (ii) \Rightarrow (i): The existence is shown by construction, using squares. See, for instance, the proof of Theorem 2.4, or [19, Examples III.1 and III.2].

Proof of (i) \Rightarrow (ii): Assume the existence of a Wishart semigroup on $D_p(k)$.⁸ Since e_{k+1} vanishes on $D_p(k)$, one obtains by using Proposition 3.4 that

$$0 = (\mathcal{A}e_{k+1})(x) = (p-k)(\alpha-k)e_k(x).$$

Since $k < p$, and $e_k(x) > 0$ for $\text{rank}(x) = k$ matrices, α must be equal to k . \square

A Wishart Semimartingales

Proposition A.1. *Let $(\Omega, \mathcal{F}, (\mathcal{F}_t)_{t \geq 0}, \mathbb{P})$ be a standard filtered probability space. Let $(X_t)_{t \geq 0}$ be a continuous, $\bar{\mathcal{S}}_p^+$ valued semimartingale of the form*

$$dX_t = dM_t + \alpha Idt, \tag{A.1}$$

where $\alpha \geq 0$, and the continuous martingale M_t has quadratic variation

$$d\langle M_{t,ij}, M_{t,kl} \rangle = ((X_t)_{ik}\delta_{jl} + (X_t)_{il}\delta_{jk} + (X_t)_{jk}\delta_{il} + (X_t)_{jl}\delta_{ik}) dt. \tag{A.2}$$

Then there exists an extension $(\tilde{\Omega}, \tilde{\mathcal{F}}, (\tilde{\mathcal{F}}_t)_{t \geq 0}, \tilde{\mathbb{P}})$ of $(\Omega, \mathcal{F}, (\mathcal{F}_t)_{t \geq 0}, \mathbb{P})$ which supports a $d \times d$ standard Brownian motion W such that

$$dX_t = \sqrt{X_t}dW_t + dW_t^\top \sqrt{X_t} + \alpha Idt. \tag{A.3}$$

Proof. This is an application of [22, Theorem V.20.1], where one interprets the SDE (A.3) in vector form, and thus W as a vector of p^2 independent, standard Brownian motions. The details of the proof are the same as those found in [3, p. 53, Proof of Theorem 2.6]. \square

⁷Note that $D_p(k)$ are non-convex domains for $k < p$, but, by Theorem 1.3, Wishart semigroups on $D_p(k)$ cannot be extended to their convex hull $\bar{\mathcal{S}}_p^+$.

⁸Using the NCGS conjecture, the following, weaker, conclusion can be made. Assume the existence of a Wishart semigroup on $D_p(k)$. Then $\Gamma_p(\alpha, x_0, I)$ exists with $\text{rank}(x_0) = k$. By the NCGS Conjecture, $\alpha/2 \in W_0$ and, if $\alpha < p-1$, then $\text{rank}(x_0) \leq \alpha$. This implies $\alpha \geq k$.

Proposition A.2. *Let $D \subset \mathcal{S}_p^+$ such that $D_p(1) \subset D$, and let $(P_t)_{t \geq 0}$ be a Wishart semigroup on D with parameter α . The following hold:*

- (i) *For $x \in D$ let (X, \mathbb{P}^x) be the canonical representation of the Markov semigroup with the initial law δ_x (cf. Remark 3.2(i)). There exists a version \tilde{X} of X that is a continuous semimartingale of the form (A.1) with quadratic variation (A.2).*
- (ii) *For any $x \in D$, the Wishart SDE (1.2) has a global weak solution with $X_0 = x$.*

Proof. Proof of (i): The canonical representation $(X, (\mathbb{P}_x)_{x \in D})$ constitutes a time homogeneous Markov process in the sense of [5, Definition 1] and an affine processes in the sense of [5, Definition 2]. Since $D_p(1) \subset D$, D contains $p \times (p+1)/2 + 1$ affinely independent elements, and thus D satisfies [5, Assumption 1].

Let $\mathcal{F}_t^0 = \sigma(X_s, s \leq t)$ be the filtration generated by the canonical process $X_t(\omega) := \omega(t)$, and let $\mathcal{F}^0 := \vee_{t \geq 0} \mathcal{F}_t$. Then by [5, Theorem 2], there exists a version \tilde{X} of X which is càdlàg. Since $(P_t)_{t \geq 0}$ is conservative, [5, Theorem 6] implies that \tilde{X} is a semimartingale with characteristics (B, C, ν) , where

$$\begin{aligned} B_{t,i} &= \int_0^t b_i(\tilde{X}_{s-}) ds, \\ C_{t,ij} &= \int_0^t c_{ij}(\tilde{X}_{s-}) ds, \\ \nu(\omega; dt, d\xi) &= K(\tilde{X}_t, d\xi) dt. \end{aligned}$$

Here $b : D \rightarrow \mathcal{S}_p$ and $c : D \rightarrow \text{Sym}_+(\mathcal{S}_p)$ are measurable functions, and $K(x, d\xi)$ is a positive kernel ($\text{Sym}_+(V)$ denotes positive semidefinite matrices on a vector space V). From the computations in the proof of Proposition 3.3 it follows that (X, \mathbb{P}_x) is regular in the sense of [5, Definition 7], that is, the coefficients ϕ, ψ are differentiable at $t = 0$, with derivatives $F(u), R(u)$ given by (3.7) and (3.8). On the other hand, by [5, Theorem 7], the functions $F(u), R(u)$ uniquely determine the differential characteristics $b_i(x), c_{ij}(x)$ and $K(x, d\xi)$. A comparison of (3.7)–(3.8) with the expressions of F and R in [5, Theorem 7] finally reveals that $\nu = 0$, i.e., the process \tilde{X} is continuous \mathbb{P}_x -almost surely, because by the semimartingale decomposition

$$X_t = X_0 + \int_0^t dM_s + \int_0^t b(X_s) ds,$$

where M is the continuous martingale part of X .

Proof of (ii): Follows from (i) by applying Proposition A.1. □

B Fourier-Laplace Transform of Wishart distributions

This section shows that the Laplace transform (1.1) can be extended to its maximal domain, which is dictated by the blow up of the right side.

The right side of (1.1) is a real analytic function, which is finite on the domain

$$D(\mu) := -\Sigma^{-1} + \mathcal{S}_p^+$$

but blows up as the argument u approaches the boundary $\partial D(\mu)$, since then the determinant vanishes.

Furthermore, the right side of (1.1) can be extended to a complex analytic function on the complex strip $D(\mu) + i\mathcal{S}_p$ (by just replacing u by $u + iv$, where $v \in \mathcal{S}_p$) and it agrees, by definition, with the left side of (1.1), on a set of uniqueness, namely the open domain \mathcal{S}_p^+ . Hence, by [6, (9.4.4)], equality holds in (1.1) for $u \in \mathcal{S}_p^+ + i\mathcal{S}_p$.

The following extends the validity of (1.1) to its maximal domain $D(\mu) + i\mathcal{S}_p$:

Proposition B.1. *Let $\mu = \Gamma_p(\beta, \omega; \Sigma)$. Then its Fourier-Laplace transform can be extended to the complex strip $D(\mu) + i\mathcal{S}_p$, and (1.1) holds for any $u \in D(\mu) + i\mathcal{S}_p$.*

For the proof, the following fundamental technical statement concerning extension of the Laplace transform of a measure on the non-negative real line is used. It is a refinement of [11, Lemma A.4]:

Lemma B.2. *Let μ be a probability measure on \mathbb{R}_+ , and h an analytic function on $(-\infty, s_1)$, where $s_1 > s_0 \geq 0$ such that*

$$\int_{\mathbb{R}_+} e^{sx} \mu(dx) = h(s) \tag{B.1}$$

for $s \in (-\infty, s_0)$. Then (B.1) also holds for $s \in (-\infty, s_1)$.

Proof. If $s_0 > 0$, the statement follows from [11, Lemma A.4]. Let therefore $s_0 = 0$.

Denote, for $s \leq 0$, $f(s) = \int_{\mathbb{R}_+} e^{sx} \mu(dx)$.

Since $h(s)$ is real analytic at 0, there exists $0 < \varepsilon < s_1$ such that for any $s \in (-\varepsilon, \varepsilon)$

$$h(s) = \sum_{k \geq 0} \frac{c_k}{k!} s^k.$$

Furthermore, by dominated convergence, one obtains iteratively for the left derivatives

$$\int_{\mathbb{R}_+} x^k e^{sx} \mu(dx) = \lim_{t \uparrow 0} \int_{\mathbb{R}_+} x^{k-1} e^{sx} \frac{e^{-tx} - 1}{-t} \mu(dx) = f^{(k)}(s) = h^{(k)}(s), \quad s \leq 0,$$

hence

$$c_k = \int_{\mathbb{R}_+} x^k \mu(dx).$$

Hence, by monotone convergence, for any $s \in (0, \varepsilon)$

$$h(s) = \sum_{k \geq 0} \int_{\mathbb{R}_+} \frac{s^k x^k}{k!} \mu(dx) = \int_{\mathbb{R}_+} \sum_{k \geq 0} \frac{s^k x^k}{k!} \mu(dx) = \int_{\mathbb{R}_+} e^{sx} \mu(dx).$$

Thus $h(s)$ verifies (B.1) on all of $(-\infty, \varepsilon)$. Now the assumptions of [11, Lemma A.4] are verified (setting $s_0 = \varepsilon$), that shows the extension to the maximal domain $(-\infty, s_1)$. \square

Proof of Proposition B.1. For $u = \Sigma^{-1}$, define μ^* as the pushforward of $\mu = \Gamma_p(\beta, \omega; \Sigma)$ under $\xi \mapsto \text{tr}(u\xi) = \text{tr}(\Sigma^{-1}\xi)$. Then μ^* is a probability measure on \mathbb{R}_+ with Laplace transform

$$\begin{aligned} f(t) &:= \int e^{tx} \mu^*(dx) = \int e^{-\text{tr}((-tu)\xi)} \mu(d\xi) \\ &= (\det \Sigma)^{-\beta} \det(\Sigma^{-1}(1-t))^{-\beta} e^{t(1-t)^{-1} \text{tr}(\Sigma^{-1}\omega)}, \quad t \leq 0, \end{aligned} \quad (\text{B.2})$$

and the right side is real analytic for $t < 1$. Hence, by Lemma B.2 the left side is also finite for $t < 1$ and equality holds in (B.2).

Therefore, it is shown that the formula (1.1) can be extended to $u = -t\Sigma^{-1}$, for any $t < 1$. Since $u > -\Sigma^{-1}$ implies $u > -t\Sigma^{-1}$ for some $t < 1$, also for any $u > -\Sigma^{-1}$

$$\int e^{-\text{tr}(u\xi)} \mu(d\xi) \leq \int e^{t \text{tr}(\Sigma^{-1}\xi)} \mu(d\xi) < \infty$$

and therefore the left side of (1.1) exists for any $u > -\Sigma^{-1}$, and thus also the Fourier-Laplace transform exists for any $u + iv$, where $u > -\Sigma^{-1}$ and $v \in \mathcal{S}_p$. Since the Fourier-Laplace transform is complex analytic on the strip $-\Sigma^{-1} + \mathcal{S}_p^+ + i\mathcal{S}_p$, and agrees with the right side of (1.1) on the domain \mathcal{S}_p^+ (which is a set of uniqueness), equality in (1.1) holds by [6, (9.4.4)]. This concludes the proof of Proposition B.1.

References

- [1] A. Ahdida and A. Alfonsi, *Exact and high-order discretization schemes for Wishart processes and their affine extension*. The Annals of Applied Probability 23 (2013), No. 3, pp. 1025–1073.
- [2] M. F. Bru, *Wishart processes*. Journal of Theoretical Probability 4 (1991), No. 4, pp. 725–751.
- [3] C. Cuchiero, D. Filipović, E. Mayerhofer and J. Teichmann, *Affine processes on positive semidefinite matrices*. The Annals of Applied Probability 21 (2011), No. 2, pp. 397–463.
- [4] C. Cuchiero, M. Keller-Ressel and J. Teichmann, *Polynomial processes and their applications to mathematical finance*. Finance and Stochastics 16 (2012), No. 4, pp. 711–740.
- [5] C. Cuchiero and J. Teichmann, *Path properties and regularity of affine processes on general state spaces*. Séminaire de Probabilités XLV Vol. 2078 (2013), pp. 201–244.
- [6] J. Dieudonné, *Foundations of Modern Analysis*. Pure and Applied Mathematics, P. Smith and S. Eilenberg, Eds. New York: Academic Press 10 (1969).
- [7] C. Donati-Martin, Y. Doumerc, H. Matsumoto and M. Yor, *Some properties of the Wishart processes and a matrix extension of the Hartman-Watson laws*. Publications of the Research Institute for Mathematical Sciences (Kyoto University) 40 (2004), No. 4, pp. 1385–1412.

- [8] S.N. Ethier and T.G. Kurtz, *Markov processes: characterization and convergence*. Vol. 282. John Wiley & Sons, 2009.
- [9] J. Faraut and A. Koranyi, *Analysis on symmetric cones*. Oxford Mathematical Monographs. Oxford Science Publications. The Clarendon Press, Oxford University Press, New York, 1994.
- [10] D. Filipović and E. Mayerhofer, Vienna Institute of Finance (2009), private communication.
- [11] D. Filipović and E. Mayerhofer, *Affine Diffusion Processes: Theory and Applications*. Advanced financial Modeling, Walter de Gruyter, Radon Series Comp. Appl. Math 8, 125–164, 2009.
- [12] S. G. Gindikin, *Invariant generalized functions in homogeneous domains*. Functional Analysis and Its Applications 9 (1975), No.1, pp. 50–52.
- [13] P. Graczyk and J. Małeck, *Multidimensional Yamada-Watanabe theorem and its applications to particle systems*. Journal of Mathematical Physics 54 (2013), No.2, pp. 021503-1–021503-15.
- [14] P. Graczyk and J. Małeck, *Strong solutions of non-colliding particle systems*. Electronic Journal of Probability 19 (2014), No. 119, pp. 1–21 .
- [15] N.L. Johnson, S. Kotz, *Distributions in Statistics: Continuous Multivariate Distributions*, John Wiley and Sons, New York, 1972.
- [16] M. Keller-Ressel and E. Mayerhofer, *Exponential Moments of Affine Processes*, Annals of Applied probability 25 (2015), No. 2, pp. 714–752.
- [17] G. Letac and H. Massam, *The noncentral Wishart as an exponential family, and its moments*. Journal of Multivariate Analysis 99 (2008), no. 7, pp. 1393–1417.
- [18] G. Letac and H. Massam, *Existence and non-existence of the non-central Wishart distributions*. ArXiv preprint: 1108.2849 (2011).
- [19] E. Mayerhofer, *Stochastic Analysis Methods in Wishart Theory II*. Modern Methods of Multivariate Statistics, P. Graczyk, A. Hassairi Eds., Travaux en Cours 82, Hermann, Paris, 2014.
- [20] E. Mayerhofer, *On the existence of non-central Wishart distributions*. Journal of Multivariate Analysis 114 (2013), pp. 448–456.
- [21] S. D. Peddada and D. St. P. Richards, *Proof of a conjecture of M. L. Eaton on the characteristic function of the Wishart distribution*. The Annals of Probability 19 (1991), No. 2, pp. 868–874.

- [22] L.C.G. Rogers and D. Williams, *Diffusions, Markov Processes, and Martingales.*, Vol. 2, Cambridge Univ. Press, Cambridge, 2000.
- [23] W. Rudin, *Real and complex analysis (3rd)*. New York: McGraw-Hill Inc, 1986.
- [24] M. J. Yor, *Squared Bessel Processes*. Encyclopedia of Quantitative Finance (2010).