

QUANTILE GRAPHICAL MODELS: PREDICTION AND CONDITIONAL INDEPENDENCE WITH APPLICATIONS TO SYSTEMIC RISK

ALEXANDRE BELLONI*, MINGLI CHEN[‡], AND VICTOR CHERNOZHUKOV[†]

ABSTRACT. We propose two types of Quantile Graphical Models (QGMs) — Conditional Independence Quantile Graphical Models (CIQGMs) and Prediction Quantile Graphical Models (PQGMs). CIQGMs characterize the conditional independence of distributions by evaluating the distributional dependence structure at each quantile index. As such, CIQGMs can be used for validation of the graph structure in the causal graphical models ([60, 89, 94]). One main advantage of these models is that we can apply them to large collections of variables driven by non-Gaussian and non-separable shocks. PQGMs characterize the statistical dependencies through the graphs of the best linear predictors under asymmetric loss functions. PQGMs make weaker assumptions than CIQGMs as they allow for misspecification. Because of QGMs ability to handle large collections of variables and focus on specific parts of the distributions, we could apply them to quantify tail interdependence. The resulting tail risk network can be used for measuring systemic risk contributions that help make inroads in understanding international financial contagion and dependence structures of returns under downside market movements.

We develop estimation and inference methods for QGMs focusing on the high-dimensional case, where the number of variables in the graph is large compared to the number of observations. For CIQGMs, these methods and results include valid simultaneous choices of penalty functions, uniform rates of convergence, and confidence regions that are simultaneously valid. We also derive analogous results for PQGMs, which include new results for penalized quantile regressions in high-dimensional settings to handle misspecification, many controls, and a continuum of additional conditioning events.

Key Words: High-dimensional graphs, conditional independence, prediction, inference, nonlinear correlation, tail risk network, systemic risk, downside movement

Date: First version: November 2012, this version October 28, 2019.

We would like to thank Don Andrews, Debopam Bhattacharya, Peter Bickel, Marianne Bitler, Peter Bühlmann, Colin Cameron, Karim Chalak, Xu Cheng, Valentina Corradi, Alan Crawford, Francis Diebold, Peng Ding, Mirko Draca, Iván Fernández-Val, Bulat Gafarov, Jean-Jacques Forneron, Kenji Fukumizu, Dalia Ghanem, Bryan Graham, Jiaying Gu, Ran Gu, Han Hong, Jungbin Hwang, Cheng Hsiao, Hidehiko Ichimura, Michael Jansson, Oscar Jorda, Chihwa Kao, Kengo Kato, Shakeeb Khan, Roger Koenker, Brad Larsen, Chenlei Leng, Arthur Lewbel, Michael Leung, Song Liu, Francesca Molinari, Whitney Newey, Dong Hwan Oh, David Pacini, Andrew Patton, Aureo de Paula, Victor de la Pena, Elisabeth Perlman, Stephen Portnoy, Demian Pouzo, James Powell, Geert Ridder, Joe Romano, Stephen Ross, Shu Shen, Senay Sokullu, Sami Stouli, Aleksey Tetenov, Takuya Ura, Tiemen Woutersen, Zhijie Xiao, Wenkai Xu, Chaoran Yu, and Yanos Zylberberg for comments and discussions. We would also like to thank the seminar and workshop participants from Aarhus University, Boston College, 2015 Warwick Summer Workshop, 11th World Congress of the Econometric Society, 2017 UCL Workshop on the Theory of Big Data, 2017 California Econometrics Conference, 2018 International Symposium on Financial Engineering and Risk Management, 2018 Shanghai Econometrics Workshop, 2018 York Econometrics Symposium, Bank of England Modelling with Big Data and Machine Learning, London School of Economics, University of Bristol, University of Connecticut, Humboldt University Berlin, the Institute of Statistical Mathematics, USC, UC Berkeley, UC Davis, Stanford University, Warwick Statistics Department and University of Tokyo.

*Duke University, e-mail:abn5@duke.edu.

[‡]University of Warwick, e-mail:m.chen.3@warwick.ac.uk.

[†]Massachusetts Institute of Technology, e-mail:vchern@mit.edu.

1. INTRODUCTION

Co-movements, dependence and influence between variables are fundamental in economics and finance for decision and policy making as well as prediction. To this end, we propose Quantile Graphical Models (QGMs) as a modeling framework and consider their usefulness in three main applications. First, empirical auction models often rely on independent private values or affiliated private values, detecting collusion in these auctions is a form of testing conditional independence. Examples of studying entry, market power, or collusion can be found in [9, 56, 91]. Second, QGMs can be used to identify and measure systemic tail risk. The recent bank and sovereign crisis in the US and Europe have also boosted the interest in the important role of network spill-over effects in contagion and shaping systemic risk ([2, 3, 47, 54]). Many measures of systemic risk focus on spill-overs fit naturally in our QGM setting ([4, 5, 55, 57, 58]). We apply these insights to re-evaluate international financial contagion in volatilities ([35]).¹ Third, QGMs can measure dependence between stock returns for hedging strategies. In financial management settings, risk quantification is crucial, and advanced hedging decisions are typically focused on the tail of the distribution of stock returns rather than the mean. Moreover, such strategies aiming to reduce risk are critical precisely during the market downside movement. Empirical evidence ([6, 7, 87]) points to the non-Gaussianity of the distribution of stock returns, especially during market downturns. Therefore, it is also instructive to understand how dependence (and policy impact) would change as the downside movement of the market becomes more extreme. The proposed QGMs are flexible enough to cover all these cases.

QGMs can be viewed as part of graphical models which have been successively applied to estimate and visualize relationship ([60, 81]). Graphical models are widely used in machine learning, statistical learning, and social science to model the statistical dependence among the components a d -dimensional random vector X_V , in the form of a graph or network $G = (V, E)$. Here V is the node set contains the labels of the components and E is the edge set represents *unknown* statistical relationships that need to be estimated, thus poses novel problems of statistical inference. In the case of Gaussian Graphical Models (GGMs), which assuming X_V are jointly Gaussian distributed, the conditional independence structure is completely characterized by the support of the inverse of the covariance matrix of X_V . Notably in this case, the same graph will characterize conditional independence and the best linear prediction. However, in *non-Gaussian* settings, not only it is harder to characterize conditional independence, there are no reasons for the same graph to characterize both conditional independence and best linear prediction.

QGMs provide an alternative route to learn conditional independence and prediction under asymmetric loss functions which is appealing in non-Gaussian settings. As in non-Gaussian cases these notions do not coincide and there are needs for different estimation approaches. We propose two different QGMs to handle different types of applications. First, we propose Conditional Independence Quantile Graphical Models (CIQGMs) to characterize the conditional independence of distributions through evaluating the distributional dependence structure at each quantile index. Second, we propose Prediction Quantile Graphical Models (PQGMs) in which predictive relationship is the main focus. Note, QGMs also enable us to focus on *specific parts of the distributions* of variables, which play an important role in applications like financial contagion and measuring systemic risk contributions where extreme events are the main interests for practitioners.

¹Works on economic and financial networks include [23, 24, 64]. We refer to [38] for an excellent review on the econometrics literature on networks.

CIQGMs can be used for validation of the graph structure in the causal graphical models ([60, 89, 94]). Conditional independence has a long history in statistical models with consequences towards parameter identification, causal inference, prediction sufficiency, and many others, see [37]. CIQGMs aim to characterize conditional independence via the conditional quantile functions. In such models, we consider a flexible specification that can approximate well the conditional quantile functions (up to a vanishing approximation error). In turn, this allows detecting which variables have a strong or near zero impact on others which can then be used to provide guidance on conditional independence.

PQGMs focus on the prediction of a variable based on linear combinations of other variables (a reduced form relation) under asymmetric losses. An important motivation for proposing PQGMs is to allow for misspecification as the conditional quantile function is typically non-linear in non-Gaussian settings. The linear specification is widely used in practice despite possible misspecification which motivates an analysis for accomodating these issues. We characterize the uniform prediction properties under a family of asymmetric loss functions, which this family enables practitioners to investigate different parts of the tail distribution. Other papers investigated the impact of misspecification on quantile functions are [1, 8, 67, 73]. Our analysis also contributes to the high-dimensional quantile regression by allowing non-vanishing misspecification.

Broadly speaking, QGMs enhance our understanding of statistical dependence among X_V . For example, for each quantile index τ , they provide visualization of the dependence via graphs whose edges represent conditional (quantile) relationships. Given that for each specific quantile index τ we will obtain one such graph, we could have a *graph process* indexed by $\tau \in (0, 1)$. The structure represented by a τ -quantile graph represents a local relation and can be valuable in cases where tail interdependence might be of special interest.² The graph process induced by QGMs has several important features. First, a τ -quantile graph enables different values of edge strength in different directions. This is important because for undirected networks, the distinction is unclear. Second, QGMs can capture the tail interdependence through estimating at a high or low quantile index.³ Third, QGMs can capture the asymmetric dependence structure at different quantiles, which can be particularly useful in empirical applications (e.g., stock market dependence, exchange rate dependence). By considering all the quantiles at once we can characterize conditional independence structure for a set of variables that are not jointly Gaussian.

We also provide and study the estimation procedures that allow us to learn QGMs from the observed data. Our techniques are geared for covering high-dimensional settings where the size of the model is potentially larger than the sample size. These techniques are based on ℓ_1 -penalized quantile regression and Neyman orthogonal equations. For CIQGMs, under mild regularity conditions, we provide rates of convergence and edge properties of the estimated graph that hold *uniformly* over a large class of data generating processes. We provide *simultaneously valid* confidence regions (post-selection) for the coefficients of the CIQGM that are uniformly valid, despite of possible model selection mistakes. Based on proper thresholding, recovery of CIQGMs patterns is possible when coefficients are well separated from zero which parallel the results

²This is similar to the contrast between quantile regression and linear regression, where the latter provides information only on the conditional mean, while the former can provide a more complete description of the distribution of the outcome.

³Examples of high or low quantile index can be $\tau = 0.95$ or $\tau = 0.05$ respectively. The analysis extends to the case of $s^3 \log^5 p = o(n\tau(1 - \tau))$. The case of $n\tau = C$, even in the fixed dimension case, leads to a substantially different analysis and limiting distributions, as shown in [27]. Here s is the sparsity parameter, p is the dimension of conditional variables, n is the sample size, C is a constant.

for graph recovery in the Gaussian case.⁴ For PQGMs, we provide an estimator that achieves an adaptive rate of convergence, which might differ under different conditioning events. Therefore we contribute to the recent active literature on simultaneous valid confidence regions post-model selection, [11, 17, 18, 26, 33, 50] [19, 20, 62, 85, 99, 102, 109]; in particular, the penalty choices and theoretical results are uniformly valid and adaptive to the relevant conditioning events.

Although we build upon the quantile regression literature ([19, 68]), we derive new results for penalized quantile regression in high dimensional settings that are uniformly valid, robust to small coefficients (e.g. allowing for model selection mistakes), allow possibly non-vanishing misspecification, many controls and a continuum of additional conditioning events. These results contribute to a growing literature that relies on quantile based models to characterize the data generating process. [110] considers a globally adaptive quantile regression model, establishes oracle properties, and improved rates of convergence for the high-dimensional case. Screening procedures based on moment conditions motivated by the quantile models have been proposed and analyzed in [59, 103] in the high-dimensional case. [63] considers tail dependence defined via conditional probabilities in a low dimensional setting.

Finally, we view QGMs as complementary to a large body of works on GGMs ([10, 25, 34, 36, 39, 43, 44, 45, 46, 53, 71, 77, 78, 82, 97, 107, 108]). Our work is also complementary to other works trying to relax the joint Gaussian assumption. [75, 76, 104] work with so-called nonparanormal models or semiparametric Gaussian copula models, i.e., the variables follow a joint Gaussian distribution after monotone transformations. [90, 93] work with Sub-Gaussian data, which restricted fatness of the tails. [79, 92, 105] work with discrete-valued random variable, and few types of exponential families. [106] provides results for M-estimators for a subclass of exponential family graphical models. QGMs allow for different sets of distributions.

The rest of the paper is organized as follows. Section 2 provides main motivating examples. Section 3 lays out the foundation of the conceptual framework of QGMs. Section 4 contains estimators for QGMs while Section 5 contains the theoretical guarantees of the estimators. Section 6 provides an empirical application of QGMs to measure systemic risk contribution. Finally, the appendix contains proofs, simulations, and implementation details of the estimators.

Notation. For an integer k , we let $[k] := \{1, \dots, k\}$ denote the set of integers from 1 to k . For a random variable X we denote by \mathcal{X} its support. We use the notation $a \vee b = \max\{a, b\}$ and $a \wedge b = \min\{a, b\}$. We use $\|v\|_p$ to denote the p -norm of a vector v . In particular, the ℓ_2 -norm is denoted by $\|\cdot\|$; the ℓ_0 -“norm” $\|\cdot\|_0$ denotes the number of non-zero components. Given a vector $\delta \in \mathbb{R}^p$, and a set of indices $T \subset \{1, \dots, d\}$, we denote by δ_T the vector in which $\delta_{Tj} = \delta_j$ if $j \in T$, $\delta_{Tj} = 0$ if $j \notin T$. We use \mathbb{E}_n to abbreviate the notation $n^{-1} \sum_{i=1}^n$; for example, $\mathbb{E}_n[f] := \mathbb{E}_n[f(\omega_i)] := n^{-1} \sum_{i=1}^n f(\omega_i)$.

2. MOTIVATING EXAMPLES

2.1. Screening for Collusion. One application of CIQGM is in the empirical auction literature on the examination of entry or bidding prices patterns to detect coordinating groups. As shown in [9], firms bids, after controlling for all information about costs, are *jointly independent* under the competitive model and lack

⁴Similar to graph recovery in the Gaussian case such exact recovery is subject to the lack of uniformity validity critiques of Leeb and Pötscher [74].

of independence is taken as evidence consistent with collusion. Although, collusion is only one alternative explanation, testing conditional independence can be viewed as a screening device to determine whether further investigation is warranted.

Mathematically, denote $Y_{a,t}$ as the amount bid by firm a on project t , and $Z_{a,t}$ as covariates observed in dataset. We define $X_{a,t}$ to be residuals after projecting out $Z_{a,t}$. [9] test whether X_a is independent of X_b , using Fisher's Z-transformation of the coefficient of correlation between X_a and X_b . For two Gaussian random variables, this is equivalent to test pairwise independence. In our terminology, this corresponds to an edge (a, b) not being contained in the graph if and only if

$$X_a \perp X_b. \quad (2.1)$$

Pairwise independence, however, needs not imply *joint independence*. CIQGM can also handle joint independence in the *non-Gaussian* setting. This is because CIQGM works with the following case

$$X_a \perp X_b \mid X_{V \setminus \{a,b\}}, \quad (2.2)$$

namely an edge (a, b) is not contained in the graph if and only if X_b and X_a are independent conditional on all remaining variables $X_{V \setminus \{a,b\}} = \{X_k; k \in V \setminus \{a,b\}\}$, via the equivalence between conditional probabilities and conditional quantiles (details can be found in Section 3.1).

2.2. Systemic Tail Risk. Measuring systemic risk taking into account tail risk network spillover effects is another application of QGMs, e.g. our framework complements to the systemic risk measure CoVaR [4] which ignore tail risk dependencies induced by the underlying financial network structure. Our framework also allows for large scale networks.

Traditional tail risk measures, such as Value of Risk (VaR), focus on the loss of an individual institution. CoVaR attempts to measure the VaR of the whole financial system or a particular financial institution by conditioning on another being in distress. Formally, [4] define institution b 's CoVaR at level τ conditional on a particular outcome of institution a , as the value of $CoVaR_\tau^{b|a}$ that solves

$$P(X_b \leq CoVaR_\tau^{b|a} | \mathbb{C}(X_a)) = \tau, \quad (2.3)$$

for some event $\mathbb{C}(X_a)$ based on X_a . A special case is $\mathbb{C}(X_a) = \{X_a = VaR_\tau^a\}$ which, as interpreted by [4], means with probability τ institution b is in trouble given that institution a is in trouble.

QGMs can work with the case

$$P(X_b \leq CoVaR_\tau^{b|a, V \setminus \{a,b\}} | \mathbb{C}(X_a, X_{V \setminus \{a,b\}})) = \tau, \quad (2.4)$$

with the main difference here is the conditioning events (or variables), i.e. from $\mathbb{C}(X_a)$ to $\mathbb{C}(X_a, X_{V \setminus \{a,b\}})$ with the latter could be high dimensional. Hence, our QGMs take into account risk spillovers from other institutions driving the CoVaR. The identified risk spillovers between all financial institutions constitute a financial tail risk network which, as shown later, can be used for measuring institutions' systemic tail risk contributions. In summary, QGMs can take into account the system-wide network spillover effects via incorporating tail network spillover effects into risk measuring, thus relate systemic risk to tail spillover effects from individual institutions to the whole system.

Another related definitions of tail risk, $\Delta CoVaR$, is defined as the change in the VaR of the whole financial system conditional on a institution being under distress relative to its median state. In terms of

estimation, replacing covariate X_a by the difference between its τ -th quantile (denoted as VaR_τ^a), and its median (denoted as $VaR_{50\%}^a$), yields

$$\Delta CoVaR_\tau^{b|a} = \widehat{\beta}_a^b(\tau)(VaR_\tau^a - VaR_{50\%}^a). \quad (2.5)$$

where $\widehat{\beta}_a^b(\tau)$ comes from pairwise quantile regression of X_b on X_a , and

$$\Delta CoVaR_\tau^{b|a, V \setminus \{a, b\}} = \check{\beta}_a^b(\tau)(VaR_\tau^a - VaR_{50\%}^a), \quad (2.6)$$

where $\check{\beta}_a^b(\tau)$ is estimated via Algorithm 4.1 or 4.2, inference procedures are based on Corollary 1.

After learning a tail risk network, we can use our new network-cooperated $\Delta CoVaR$ to measure the systemic risk contribution of each institution. The systemic risk contribution of institution a can be measured by it "to" or "from" degree, see [5]; or by other network centrality measures, see [61, 70, 84]. To-degrees measure contributions of individual institutions to the overall risk of systemic network events, for institution a is defined as $\delta_a^{to} = \sum_k \Delta CoVaR_\tau^{k|a, V \setminus \{a, k\}}$. From-degrees measure exposure of individual institutions to systemic shocks from the network, for institution a it is defined as $\delta_a^{from} = \sum_k \Delta CoVaR_\tau^{a|k, V \setminus \{a, k\}}$. The net contribution of institution a is defined as $net\text{-}\Delta CoVaR^a = \delta_a^{to} - \delta_a^{from}$.

In Section 6, we revisit the analysis of international financial contagion through the volatility spillovers perspective. We visualize the tail risk interdependence via PQGMs as they allow for heteroskedasticity and asymmetric responses, can be used to model nonlinear tail interdependence, and to visualize potential asymmetric changes in conditional correlations. The estimated contagion network taking into account global interconnectedness is important for Eurozone financial regulators who want to identify globally systemically important EU countries, or for global financial portfolio diversification. Our the systemic tail risk analysis tools mentioned in previous paragraphs, can help with achieve those goals.

2.3. Stock Returns Under Market Downside Movements. Hedging decisions rely on the dependence of various stocks returns. Moreover, hedging is even more relevant during market downside movements, which motivates us to understand interdependence conditional on those events. Stock returns are in general non-Gaussian in those settings, as shown in the empirical finance literature, e.g. [6, 80, 88]. We can parameterize the downside movements by using a random variable W , which could be the market index, and conditional on the event $\Omega_\varpi = \{W \leq \varpi\}$. This allows us to define a ϖ -conditional-CIQGM as $G^I(\tau, \varpi) = (V, E^I(\tau, \varpi))$ and a ϖ -conditional-PQGM as $G^P(\tau, \varpi) = (V, E^P(\tau, \varpi))$, for each $\varpi \in \mathcal{W}$. We might be interest in a fixed ϖ or on a family of values $\varpi \in (-\bar{\varpi}, 0]$. The latter induces $\mathcal{W} = \{W \leq \varpi : \varpi \in (-\bar{\varpi}, 0]\}$.

Figure 1 provides an example using \mathcal{W} -Conditional QGM with $W = \{\text{Market Index Returns}\}$, ϖ as the τ_m -th quantile of the market index returns, and $\tau_m = \{0.15, 0.5, 0.75, 0.9\}$. We obtain daily stock returns from CRSP and use S&P 500 as the market index. The full sample consists of 2769 observations for 86 stocks from Jan 2, 2003 to December 31, 2013. The total number of stocks is 86 due to data availability. We define market movement as when the market index returns are below a pre-specified level (e.g. τ_m -th quantile), hence conditioning on a particular ϖ corresponds to consider the subsample based on whether the corresponding date's market return is less equal to the τ_m -th quantile of the market index returns. The results show higher interdependence under market downside moments, pose different hedging decisions.




FIGURE 1. Stock Returns Interdependence under Different Market Conditions. Note: Presence of edges between nodes indicate that these two nodes (or stocks) are conditionally dependent.

3. QUANTILE GRAPHICAL MODELS

In this section we describe quantile graphical models associated with a d -dimensional random vector X_V where the set $V = [d] = \{1, \dots, d\}$ denotes the labels of the components. These models aim to provide a description of the dependence between the random variables in X_V . In particular, these models induce graphs that allow for visualizing dependence structures. Nonetheless, because of the non-Gaussianity, we consider two fundamentally distinct models (one geared towards conditional independence and one geared towards prediction).

3.1. Conditional Independence Quantile Graphical Models. Conditional independence graphs have been used to provide visualization and insight on the dependence structure between random variables. Each node of the graph is associated with a component of X_V . We denote the conditional independence graph as $G^I = (V, E^I)$ where G^I is an undirected graph with vertex set V and edge set E^I which is represented by an adjacency matrix ($E_{a,b}^I = 1$ if the edge $(a, b) \in G^I$, and $E_{a,b}^I = 0$ otherwise). An edge (a, b) is not contained in the graph if and only if

$$X_a \perp X_b \mid X_{V \setminus \{a,b\}}, \quad (3.7)$$

namely X_b and X_a are independent conditional on all remaining variables $X_{V \setminus \{a,b\}} = \{X_k; k \in V \setminus \{a,b\}\}$.

Comment 3.1 (Conditional Independence Under Gaussianity). In the case that X_V is jointly Gaussian distributed, $X_V \sim N(0, \Sigma)$ with Σ as the covariance matrix of X_V , the conditional independence structure between two components is determined by the inverse of the covariance matrix, i.e. the precision matrix $\Theta = \Sigma^{-1}$. It follows that the non-zero elements in the precision matrix corresponds to the non-zero coefficients of the associated (high dimensional) mean regression. The family of Gaussian distributions with this property is known as a Gauss-Markov random field with respect to the graph G . This observation has motivated a large literature [71] and interesting extensions that allow for transformations of Gaussian variables [75, 76].

In order to achieve a tractable concept for non-Gaussian settings, we use that (3.7) occurs if and only if

$$F_{X_a}(\cdot | X_{V \setminus \{a\}}) = F_{X_a}(\cdot | X_{V \setminus \{a,b\}}) \text{ for all } X_{V \setminus \{a\}} \in \mathcal{X}_{V \setminus \{a\}}. \quad (3.8)$$

In turn, by the equivalence between conditional probabilities and conditional quantiles to characterize a random variable, we have that (3.7) occurs if and only if

$$Q_{X_a}(\tau | X_{V \setminus \{a\}}) = Q_{X_a}(\tau | X_{V \setminus \{a,b\}}) \text{ for all } \tau \in (0, 1), \text{ and } X_{V \setminus \{a\}} \in \mathcal{X}_{V \setminus \{a\}}. \quad (3.9)$$

For a quantile index $\tau \in (0, 1)$, the τ -quantile conditional independence graph is a directed graph $G^I(\tau) = (V, E^I(\tau))$ with vertex set V and edge set $E^I(\tau)$. An edge (a, b) is not contained in the edge set $E^I(\tau)$ if and only if

$$Q_{X_a}(\tau | X_{V \setminus \{a\}}) = Q_{X_a}(\tau | X_{V \setminus \{a,b\}}) \text{ for all } X_{V \setminus \{a\}} \in \mathcal{X}_{V \setminus \{a\}}. \quad (3.10)$$

By the equivalence between (3.8) and (3.9), the union of τ -quantile graphs over $\tau \in (0, 1)$ represents the conditional independence structure of X , namely $E^I = \cup_{\tau \in (0,1)} E^I(\tau)$. We also consider a relaxation of (3.7). For a set of quantile indices $\mathcal{T} \subset (0, 1)$, we say that

$$X_a \perp_{\mathcal{T}} X_b \mid X_{V \setminus \{a,b\}}, \quad (3.11)$$

X_a and X_b are \mathcal{T} -conditionally independent given $X_{V \setminus \{a,b\}}$, if (3.10) holds for all $\tau \in \mathcal{T}$. Thus, we have that (3.7) implies (3.11). We define the \mathcal{T} -quantile graph as $G^I(\mathcal{T}) = (V, E^I(\mathcal{T}))$ where

$$E^I(\mathcal{T}) = \cup_{\tau \in \mathcal{T}} E^I(\tau).$$

Although the conditional independence concept relates to all quantile indices, the quantile characterization described above also lends itself to quantile specific impacts which can be of independent interest.⁵

⁵For example, we might be interested in some extreme events which typically correspond to crises in financial systems.

3.2. Prediction Quantile Graphical Models. Prediction Quantile Graphical Models (PQGMs) are motivated by prediction accuracy under an asymmetric loss function (instead of conditional independence as in Section 3.1). More precisely, for each $a \in V$, we are interested in predicting X_a based on linear combinations of the remaining variables, $X_{V \setminus \{a\}}$, where accuracy is measured with respect to an asymmetric loss function. Formally, PQGMs measure accuracy as

$$\mathcal{L}_a(\tau \mid V \setminus \{a\}) = \min_{\beta} \mathbb{E}[\rho_{\tau}(X_a - X'_{-a}\beta)] \quad (3.12)$$

where $X_{-a} = (1, X'_{V \setminus \{a\}})'$, and the asymmetric loss function $\rho_{\tau}(t) = (\tau - 1\{t \leq 0\})t$ is the check function used in [69].

Importantly, PQGMs are concerned with the best linear predictor under the asymmetric loss function ρ_{τ} which is a specification that is widely used in practice. This is a fundamental distinction with respect to CIQGMs discussed in Section 3.1 where the specification of the conditional quantile was approximately a linear function of transformations of $X_{V \setminus \{a\}}$.⁶ Indeed, we note that under suitable conditions the linear predictor that solves the minimization problem in (3.12) approximates the conditional quantile regression as shown in [16]. (In fact, the conditional quantile function would be linear if X_V was jointly Gaussian distributed.) However, PQGMs do not assume that the conditional quantile function of X_a is well approximated by a linear function and instead it focuses on the best linear predictor.

We define that X_b is predictively uninformative for X_a given $X_{V \setminus \{a,b\}}$ if

$$\mathcal{L}_a(\tau \mid V \setminus \{a\}) = \mathcal{L}_a(\tau \mid V \setminus \{a,b\}) \quad \text{for all } \tau \in (0, 1),$$

i.e., considering a linear function of X_b will not improve our performance of predicting X_a with respect to the asymmetric loss function ρ_{τ} .

Again we can visualize the predictive relationship through a graph process indexed by $\tau \in (0, 1)$. That is, for each $\tau \in (0, 1)$ we have a directed graph $G^P(\tau) = (V, E^P(\tau))$, where an edge $(a, b) \in G^P(\tau)$ only if X_b is predictively informative for X_a given $X_{V \setminus \{a,b\}}$ at the quantile τ . Finally, it is also convenient to define the PQGM associated with a subset $\mathcal{T} \subset (0, 1)$ as $G^P(\mathcal{T}) = (V, E^P(\mathcal{T}))$ where

$$E^P(\mathcal{T}) = \cup_{\tau \in \mathcal{T}} E^P(\tau).$$

3.3. \mathcal{W} -Conditional Quantile Graphical Models. In what follows, we discuss an extension of the QGMs discussed in Sections 3.1 and 3.2 to allows for conditioning on a (possible infinity) family of events $\varpi \in \mathcal{W}$.⁷ Such extension is motivated by several applications in which the interdependence between the random variables in X_V maybe substantially impacted by additional observable events (e.g. downside movements of the market). This general framework allows different forms of conditioning. The main implication of this extension is that QGMs are now graph processes indexed by $\tau \in \mathcal{T} \subset (0, 1)$ and $\varpi \in \mathcal{W}$.

We define X_a and X_b are (\mathcal{T}, ϖ) -conditionally independent,

$$X_a \perp_{\mathcal{T}} X_b \mid X_{V \setminus \{a,b\}}, \varpi \quad (3.13)$$

⁶In Section 3.1 the vector Z^a in equation (4.16) collects the functions of the vector $X_{V \setminus \{a\}}$.

⁷With a slight abuse of notation, we let ϖ to denote the event and also the index of such event. For example, we write $P(\varpi)$ as a shorthand for $P(W \in \Omega_{\varpi})$.

if for all $\tau \in \mathcal{T}$ we have

$$Q_{X_a}(\tau|X_{V \setminus \{a\}}, \varpi) = Q_{X_a}(\tau|X_{V \setminus \{a,b\}}, \varpi). \quad (3.14)$$

The conditional independence edge set associated with (τ, ϖ) is defined analogously as before. We denote them by $E^I(\tau, \varpi)$ and $E^I(\mathcal{T}, \varpi) = \cup_{\tau \in \mathcal{T}} E^I(\tau, \varpi)$ for each $\varpi \in \mathcal{W}$.

The extension of PQGMs proceeds by defining the accuracy under the asymmetric loss function conditionally on ϖ . More precisely, we define

$$\mathcal{L}_a(\tau|V \setminus \{a\}, \varpi) = \min_{\beta} E[\rho_{\tau}(X_a - X'_{-a}\beta) | \varpi]. \quad (3.15)$$

The prediction edge set associated with (τ, ϖ) is also defined analogously as before. We denote them by $E^P(\tau, \varpi)$ and $E^P(\mathcal{T}, \varpi) = \cup_{\tau \in \mathcal{T}} E^P(\tau, \varpi)$, for each $\varpi \in \mathcal{W}$.

4. ESTIMATORS FOR HIGH-DIMENSIONAL QUANTILE GRAPHICAL MODELS

In this section, we propose and discuss estimators for QGMs introduced in Section 3. Throughout it is assumed that we observe a d -dimensional i.i.d. random vector X_V , namely $\{X_{iV} : i = 1, \dots, n\}$. Based on the data observed, unless additional assumptions are imposed we cannot estimate the quantities of interest for all $\tau \in (0, 1)$. Instead, in what follows we will consider a (compact) set of quantile index $\mathcal{T} \subset (0, 1)$. The estimators are intended to handle high dimensional models and a continuum of conditioning events in \mathcal{W} .

4.1. Estimators for CIQGMs. We discuss the specification and propose an estimator for CIQGMs. Although in general it is potentially hard to correctly specify coherent models, the following are simple examples.

Example 1 (Multivariate Gaussian Distribution). Consider the Gaussian case, $X_V \sim N(\mu, \Sigma)$. It follows that for each $a \in V$, the conditional distribution $X_a | X_{V \setminus \{a\}}$ satisfies

$$X_a | X_{V \setminus \{a\}} \sim N \left(\mu_a - \sum_{j \in V \setminus \{a\}} \frac{(\Sigma^{-1})_{aj}}{(\Sigma^{-1})_{aa}} (X_j - \mu_j), \frac{1}{(\Sigma^{-1})_{aa}} \right).$$

Therefore the conditional quantile function of X_a is linear in $X_{V \setminus \{a\}}$ and is given by

$$Q_{X_a}(\tau|X_{V \setminus \{a\}}) = \frac{\Phi^{-1}(\tau)}{(\Sigma^{-1})_{aa}^{1/2}} + \mu_a - \sum_{j \in V \setminus \{a\}} \frac{(\Sigma^{-1})_{aj}}{(\Sigma^{-1})_{aa}} (X_j - \mu_j).$$

Example 2 (Multivariate t -Distribution). Consider the multivariate t distribution case, $X_V \sim t_p(\mu, \Sigma, v)$, with location μ , scale matrix Σ , and degrees of freedom v , as in [42]. It follows that for each $a \in V$, the conditional distribution $X_a | X_{V \setminus \{a\}}$ satisfies

$$X_a | X_{V \setminus \{a\}} \sim t_{p-1} \left(\mu_a - \sum_{j \in V \setminus \{a\}} \frac{(\Sigma^{-1})_{aj}}{(\Sigma^{-1})_{aa}} (X_j - \mu_j), \frac{v + d_1}{v + 1} \frac{1}{(\Sigma^{-1})_{aa}}, v + 1 \right).$$

Therefore the conditional quantile function of X_a is given by

$$Q_{X_a}(\tau|X_{V \setminus \{a\}}) = \sqrt{\frac{v + d_1}{v + 1}} \frac{F_{t_{v+1}}^{-1}(\tau)}{(\Sigma^{-1})_{aa}^{1/2}} + \mu_a - \sum_{j \in V \setminus \{a\}} \frac{(\Sigma^{-1})_{aj}}{(\Sigma^{-1})_{aa}} (X_j - \mu_j),$$

here $d_1 = (X_{V \setminus \{a\}} - \mu_{V \setminus \{a\}})^T \Sigma_{aa}^{-1} (X_{V \setminus \{a\}} - \mu_{V \setminus \{a\}})$.

Example 3 (Multiplicative Error Model). Consider $d = 2$ so that $V = \{1, 2\}$. Assume that X_2 and ε are independent positive random variables. Assume further that they relate to X_1 as

$$X_1 = \alpha + \varepsilon X_2.$$

In this case, we have that the conditional quantile functions are linear and given by

$$Q_{X_1}(\tau|X_2) = \alpha + F_\varepsilon^{-1}(\tau)X_2 \quad \text{and} \quad Q_{X_2}(\tau|X_1) = (X_1 - \alpha)/F_\varepsilon^{-1}(1 - \tau).$$

Example 4 (Additive Error Model). Consider $d = 2$ so that $V = \{1, 2\}$. Let $X_2 \sim U(0, 1)$ and $\varepsilon \sim U(0, 1)$ be independent random variables. Also define the random variable X_1 as

$$X_1 = \alpha + \beta X_2 + \varepsilon.$$

It follows that $Q_{X_1}(\tau|X_2) = \alpha + \beta X_2 + \tau$. However, if $\beta = 0$, we have $Q_{X_2}(\tau|X_1) = \tau$, and for $\beta > 0$, direct calculations yield that

$$Q_{X_2}(\tau|X_1) = \begin{cases} \frac{\tau}{\beta}(X_1 - \alpha), & \text{if } X_1 \leq \alpha + \beta \\ \tau + (1 - \tau)(X_1 - \alpha - \beta), & \text{if } X_1 \geq \alpha + \beta \end{cases}$$

where we note that $X_1 \in [\alpha, 1 + \alpha + \beta]$.

Example 5 (Mixture of Gaussians). Similar to the prior example, consider the case $X_V | \varpi \sim N(\mu_\varpi, \Sigma_\varpi)$ for each $\varpi \in \mathcal{W}$. It follows that for $a \in V$, the conditional distribution satisfies

$$X_a | X_{V \setminus \{a\}}, \varpi \sim N \left(\mu_{\varpi a} - \sum_{j \in V \setminus \{a\}} \frac{(\Sigma^{-1})_{\varpi a j}}{(\Sigma^{-1})_{\varpi a a}} (X_j - \mu_{\varpi j}), \frac{1}{(\Sigma^{-1})_{\varpi a a}} \right).$$

Again the conditional quantile function of X_a is linear in $X_{V \setminus \{a\}}$ and is given by

$$Q_{X_a}(\tau|X_{V \setminus \{a\}}, \varpi) = \frac{\Phi^{-1}(\tau)}{(\Sigma^{-1})_{\varpi a a}^{1/2}} + \mu_{\varpi a} - \sum_{j \in V \setminus \{a\}} \frac{(\Sigma^{-1})_{\varpi a j}}{(\Sigma^{-1})_{\varpi a a}} (X_j - \mu_{\varpi j}).$$

Example 6 (Monotone Transformations). Consider the Gaussian case, for each $a \in V$, $X_a = h_a(Y_a)$ and $Y_V \sim N(\mu, \Sigma)$. It follows that for each $a \in V$, the conditional quantile function satisfies

$$Q_{X_a}(\tau|X_{V \setminus \{a\}}) = h_a \left(\frac{\Phi^{-1}(\tau)}{(\Sigma^{-1})_{aa}^{1/2}} + \mu_a - \sum_{j \in V \setminus \{a\}} \frac{(\Sigma^{-1})_{aj}}{(\Sigma^{-1})_{aa}} (h_j^{-1}(X_j) - \mu_j) \right).$$

In particular, if $(h_a : a \in V)$ are monotone polynomials, the expression above is a sum of monomials with fractional and integer exponents.

Although a linear specification is correct for Examples 1 and 3, Example 2 and 4 illustrate that we need to consider a more general transformation of the covariates X_V in the specification for each conditional quantile function. Nonetheless, specifications with additional non-linear terms can approximate non-drastic departures from normality.

We will consider a conditional quantile representation for each $a \in V$. It is based on transformations of the original covariates $X_{V \setminus \{a\}}$ that create a p -dimensional random vector $Z^a = Z^a(X_{V \setminus \{a\}})$ such that

$$Q_{X_a}(\tau|X_{V \setminus \{a\}}) = Z^a \beta_{a\tau} + r_{a\tau}, \quad \beta_{a\tau} \in \mathbb{R}^p, \quad \text{for all } \tau \in \mathcal{T}, \quad (4.16)$$

where $r_{a\tau}$ denotes a small approximation error. For $b \in V \setminus \{a\}$ we let $I_a(b) := \{j : Z_j^a \text{ depends on } X_b\}$. That is, $I_a(b)$ contains the components of Z^a that are functions of X_b . Under correct specification, if X_a and X_b are conditionally independent, we have $\beta_{a\tau j} = 0$ for all $j \in I_a(b)$, $\tau \in (0, 1)$.

This allows us to connect the conditional independence quantile graph estimation problem with model selection with quantile regression. Indeed, the representation (4.16) has been used in several quantile regression models, see [68]. Under mild conditions this model allows us to identify the process $(\beta_{a\tau})_{\tau \in \mathcal{T}}$ as the solution of the following moment equation

$$\mathbb{E}[(\tau - 1\{X_a \leq Z^a \beta_{a\tau} + r_{a\tau}\})Z^a] = 0. \quad (4.17)$$

In order to allow for a flexible specification, so that the approximation errors are negligible, it is attractive to consider a high-dimensional Z^a where its dimension p is possibly larger than the sample size n . In turn, having a large number of technical controls creates an estimation challenge if the number of coefficients p is not negligible with respect to the sample size n . In such a high dimensional setting, a widely applicable condition that makes estimation possible is approximate sparsity [11, 18, 49]. Formally we require

$$\max_{a \in V} \sup_{\tau \in \mathcal{T}} \|\beta_{a\tau}\|_0 \leq s, \quad \max_{a \in V} \sup_{\tau \in \mathcal{T}} \{\mathbb{E}[r_{a\tau}^2]\}^{1/2} \lesssim \sqrt{s/n}, \quad \text{and} \quad \max_{a \in V} \sup_{\tau \in \mathcal{T}} |\mathbb{E}[f_{a\tau} r_{a\tau} Z^a]| = o(n^{-1/2}), \quad (4.18)$$

where the sparsity parameter s of the model is allowed to grow (at a slower rate) as n grows, and $f_{a\tau} = f_{X_a|X_{V \setminus \{a\}}}(Q_{X_a}(\tau|X_{V \setminus \{a\}})|X_{V \setminus \{a\}})$ denotes the conditional density function evaluated at the corresponding conditional quantile value. This sparsity also has implications on the maximum degree of the associated quantile graph.

Algorithm 4.1 below contains our proposal to estimate $\beta_{a\tau}$, $a \in V$, $\tau \in \mathcal{T}$. It is based on three procedures in order to overcome high-dimensionality. In the first step, we apply a (post-) ℓ_1 -penalized quantile regression. The second step applies (post-)Lasso where the data is weighted by the conditional density function at the conditional quantile.⁸ Finally, the third step relies on constructing (orthogonal) score function that provides immunity to (unavoidable) model selection mistakes.

There are several parameters that need to be specified for Algorithm 4.1. The penalty parameter $\lambda_{V\mathcal{T}}$ is chosen to be larger than the ℓ_∞ -norm of the (rescaled) score at the true quantile function. The work in [12] exploits the fact that this quantity is pivotal in their setting. Here, additional correlation structure would have an impact and the distribution is pivotal only for each $a \in V$. The penalty is based on the maximum of the quantiles of the following random variables (each with pivotal distribution), for $a \in V$

$$\Lambda_{a\mathcal{T}} = \sup_{\tau \in \mathcal{T}} \max_{j \in [p]} \frac{|\mathbb{E}_n[(1\{U \leq \tau\} - \tau)Z_j^a]|}{\sqrt{\tau(1-\tau)}\hat{\sigma}_{aj}^Z} \quad (4.19)$$

where $\{U_i : i = 1, \dots, n\}$ are i.i.d. uniform $(0, 1)$ random variables, and $\hat{\sigma}_{aj}^Z = \{\mathbb{E}_n[(Z_j^a)^2]\}^{1/2}$ for $j \in [p]$. The penalty parameter $\lambda_{V\mathcal{T}}$ is defined as

$$\lambda_{V\mathcal{T}} := \max_{a \in V} \Lambda_{a\mathcal{T}}(1 - \xi/|V| \mid Z^a),$$

that is, the maximum of the $1 - \xi/|V|$ conditional quantile of $\Lambda_{a\mathcal{T}}$ given in (4.19). Regarding the penalty term for the weighted Lasso in Step 2, we recommend a (theoretically valid) iterative choice. We refer to Appendix

⁸We note that an estimate for $f_{a\tau}$ is available from ℓ_1 -penalized quantile regression estimators for $\tau + h$ and $\tau - h$ where h is a bandwidth parameter, see [19, 68] and Comment 4.2.

A for the implementation details of the algorithm. We denote $\|\beta\|_{1,\hat{\sigma}^Z} := \sum_j \hat{\sigma}_{aj}^Z |\beta_j|$ the standardized version of the ℓ_1 -norm.

Algorithm 4.1. (CIQGM Estimator.) *For each $a \in V$, $\tau \in \mathcal{T}$, and $j \in [p]$*

Step 1. *Compute $\hat{\beta}_{a\tau}$ from $\|\cdot\|_{1,\hat{\sigma}^Z}$ -penalized τ -quantile regression of X_a on Z^a with penalty $\lambda_{V\mathcal{T}}\sqrt{\tau(1-\tau)}$.*

Compute $\check{\beta}_{a\tau}$ from τ -quantile regression of X_a on $\{Z_k^a : |\hat{\beta}_{a\tau k}| \geq \lambda_{V\mathcal{T}}\sqrt{\tau(1-\tau)}/\hat{\sigma}_{ak}^Z\}$.

Step 2. *Compute $\tilde{\gamma}_{a\tau}^j$ from the post-Lasso estimator of $f_{a\tau}Z_{-j}^a$ on $f_{a\tau}Z_{-j}^a$.*

Step 3. *Construct the score function $\hat{\psi}_i(\alpha) = (\tau - 1\{X_{ia} \leq Z_{ij}^a\alpha + Z_{i,-j}^a\check{\beta}_{a\tau,-j}\})f_{ia\tau}(Z_{ij}^a - Z_{i,-j}^a\tilde{\gamma}_{a\tau}^j)$ and for $L_{a\tau j}(\alpha) = |\mathbb{E}_n[\hat{\psi}_i(\alpha)]|^2/\mathbb{E}_n[\hat{\psi}_i^2(\alpha)]$, set $\check{\beta}_{a\tau j} \in \arg \min_{\alpha \in \mathcal{A}_{a\tau j}} L_{a\tau j}(\alpha)$.*

Algorithm 4.1 above has been studied in [19] where it is applied to a single triple (a, τ, j) , and we have used the following parameter space for α , $\mathcal{A}_{a\tau j} = \{\alpha \in \mathbb{R} : |\alpha - \check{\beta}_{a\tau j}| \leq 10/\{\hat{\sigma}_{aj}^Z \log n\}\}$. Under similar conditions, results that hold uniformly over $(a, \tau, j) \in V \times \mathcal{T} \times [p]$ are achievable (as shown in the next sections) building upon the tools developed in [12] and [29]. Algorithm 4.1 is tailored to achieve good rates of convergence in the ℓ_∞ -norm. In particular, under standard regularity conditions, with probability approaching 1 we have

$$\sup_{\tau \in \mathcal{T}} \|\beta_{a\tau} - \check{\beta}_{a\tau}\|_\infty \lesssim \sqrt{\frac{\log(p|V|n)}{n}}.$$

In order to create an estimate of $E^I(\tau) = \{(a, b) \in V \times V : \max_{j \in I_a(b)} |\beta_{a\tau j}| > 0\}$, we define

$$\hat{E}^I(\tau) = \left\{ (a, b) \in V \times V : \max_{j \in I_a(b)} \frac{|\check{\beta}_{a\tau j}|}{\text{se}(\check{\beta}_{a\tau j})} > \overline{\text{cv}} \right\}$$

where $\text{se}(\check{\beta}_{a\tau j}) = \{\tau(1-\tau)\mathbb{E}_n[\tilde{v}_{ia\tau j}^2]\}^{-1/2}$ with $\tilde{v}_{ia\tau j} = \hat{f}_{ia\tau}\{Z_{ij}^a - Z_{i,-j}^a\tilde{\gamma}_{a\tau}^j\}$, is an estimate of the standard deviation of the estimator, and the critical value $\overline{\text{cv}}$ is set to account for the uniformity over $a \in V$, $\tau \in \mathcal{T}$, and $j \in [p]$. We discuss in the following sections a data driven procedure based on multiplier bootstrap that is theoretically valid in this high dimensional setting.

Comment 4.1 (Stepdown Procedure for $\overline{\text{cv}}$). Setting a critical value $\overline{\text{cv}}$ that accounts for the multiple hypotheses being tested plays an important role to estimate the graph $\hat{E}^I(\tau)$. Further improvements can be obtained by considering the stepdown procedure of [95] for multiple hypothesis testing that was studied for the high-dimensional case in [28]. The procedure iteratively creates a suitable sequence of decreasing critical values. In each step only null hypotheses that were not rejected are considered to determine the critical value. Thus, as long as any hypothesis is rejected at a step, the critical value decreases and we continue to the next iteration. The procedure stops when no hypothesis in the current active set is rejected.

Comment 4.2 (Estimation of Conditional Density Function). The algorithm above requires the conditional density function $f_{a\tau}$ which typically needs to be estimated in practice. It turns out that estimation of conditional quantiles yields a natural estimator for the conditional density function as

$$f_{a\tau} = \frac{1}{\partial Q_{X_a}(\tau|Z^a)/\partial \tau}.$$

Therefore, based on ℓ_1 -penalized quantile regression estimates at the $\tau + h_n$ and $\tau - h_n$ quantile, where $h = h_n \rightarrow 0$ denotes a bandwidth parameter, we have

$$\hat{f}_{a\tau} = \frac{2h}{\widehat{Q}_{X_a}(\tau + h|Z^a) - \widehat{Q}_{X_a}(\tau - h|Z^a)} \quad (4.20)$$

as an estimator of $f_{a\tau}$. Under smoothness conditions, it has a bias of order h^2 . See [19] and the references therein for additional comments and estimators.

4.2. Estimators for PQGMs. In this section we propose an estimator for PQGMs in which case we are interested in the prediction of X_a , $a \in V$, using a linear combination of $X_{V \setminus \{a\}}$ under the asymmetric loss discussed in (3.12). We will add an intercept as one of the variables for the sake of notation so that $X_{-a} = (1, X'_{V \setminus \{a\}})'$. Given the loss function ρ_τ , the target d -dimensional vector of parameters $\beta_{a\tau}$ is defined as (part of) the solution of the following optimization problem

$$\beta_{a\tau} \in \arg \min_{\beta} \mathbb{E}[\rho_\tau(X_a - X'_{-a}\beta)]. \quad (4.21)$$

As we are interested in the case that d is large, the use of high-dimensional tools to achieve consistent estimators is needed. The estimation procedure we proposed is based on ℓ_1 -penalized quantile regression but additional issues need to be considered to cope with the (non-vanishing) difference between the best linear predictor and the conditional quantile function. Again we consider models that satisfy an approximately sparse condition. Formally, we require the existence of sparse coefficients $\{\bar{\beta}_{a\tau} : a \in V, \tau \in \mathcal{T}\}$ such that

$$\max_{a \in V} \sup_{\tau \in \mathcal{T}} \|\bar{\beta}_{a\tau}\|_0 \leq s \quad \text{and} \quad \max_{a \in V} \sup_{\tau \in \mathcal{T}} \{\mathbb{E}[\{X'_{-a}(\beta_{a\tau} - \bar{\beta}_{a\tau})\}^2]\}^{1/2} \lesssim \sqrt{s/n}, \quad (4.22)$$

where (again) the sparsity parameter s of the model is allowed to grow as n grows. The high-dimensionality prevents us from using (standard) quantile regression methods and regularization methods are needed to achieve good prediction properties.

A key issue is to set the penalty parameter properly so that it bounds from above

$$\max_{a \in V} \sup_{\tau \in \mathcal{T}} \max_{j \in [d]} |\mathbb{E}_n[(1\{X_a \leq X'_{-a}\beta_{a\tau}\} - \tau)X_{-a,j}]|. \quad (4.23)$$

However, it is important to note that we do not assume that the conditional quantile of X_a is a linear function of X_{-a} . Under correct linear specification of the conditional quantile function, ℓ_1 -penalized quantile regression estimator has been studied in [12]. The case that the conditional quantile function differs from a linear specification by vanishing approximation errors has been considered in [65] and [19]. The analysis proposed here aims to allow for non-vanishing misspecification of the quantile function relative to a linear specification while still guarantees good rates of convergence in the ℓ_2 -norm to the best linear specification. Thus the penalty parameter in the penalized quantile regression needs to account for such misspecification and is no longer pivotal as in [12].

In order to handle this issue we propose a two step estimation procedure. In the first step, the penalty parameter λ_0 is conservative and is set via bounds constructed based on symmetrization arguments, similar in spirit to [13, 98]. This leads to $\lambda_0 = 2(1 + 1/16)\sqrt{2\log(8|V|^2/\xi)/n}$. Although this is conservative, under mild conditions this would lead to estimates that can be leverage to fine tune the penalty choice. The second step uses the preliminary estimator to bootstrap (4.23) based on the tools in [28] as follows. Specifically, for estimates $\hat{\varepsilon}_{ia\tau}$ of the “noise” $\varepsilon_{ia\tau} = 1\{X_{ia} \leq X'_{i,-a}\beta_{a\tau}\} - \tau$ for $i \in [n]$, for $a \in V$ define

$$\bar{\Lambda}_{a\mathcal{T}} := 1.1 \sup_{\tau \in \mathcal{T}} \max_{j \in [d]} \frac{|\mathbb{E}_n[g_i \hat{\varepsilon}_{ia\tau} X_{i,-aj}]|}{\{\mathbb{E}_n[\hat{\varepsilon}_{ia\tau}^2 X_{i,-aj}^2]\}^{1/2}} \quad (4.24)$$

where $(g_i)_{i=1}^n$ is a sequence of i.i.d. standard Gaussian random variables. The new penalty parameter $\bar{\lambda}_{V\mathcal{T}}$ is defined as

$$\bar{\lambda}_{V\mathcal{T}} := \max_{a \in V} \bar{\Lambda}_{a\mathcal{T}}(1 - \xi |X_{-a}) \quad (4.25)$$

that is, the maximum of the $(1 - \xi)$ conditional quantile of $\bar{\Lambda}_{a\mathcal{T}}$. The penalty choice above adapts to the unknown correlation structure across components and quantile indices. The following algorithm states the procedure where we denote weighted ℓ_1 -norms by $\|\beta\|_{1,\hat{\sigma}^X} := \sum_j \hat{\sigma}_{aj}^X |\beta_j|$ with $\hat{\sigma}_{aj}^X = \{\mathbb{E}_n[X_j^2]\}^{1/2}$ the standardized version of the ℓ_1 -norm and $\|\beta\|_{1,\hat{\varepsilon}} := \sum_j \hat{\varepsilon}_{a\tau j}^X |\beta_j|$ with $\hat{\varepsilon}_{a\tau j}^X = \{\mathbb{E}_n[\hat{\varepsilon}_{a\tau}^2 X_{-a,j}^2]\}^{1/2}$ a norm based on the estimated residuals.

Algorithm 4.2. (PQGM Estimator.) *For each $a \in V$, and $\tau \in \mathcal{T}$*

Step 1. *Compute $\hat{\beta}_{a\tau}$ from $\|\cdot\|_{1,\hat{\sigma}^X}$ -penalized τ -quantile regression of X_a on X_{-a} with penalty λ_0 .*

Compute $\tilde{\beta}_{a\tau}$ from τ -quantile regression of X_a on $\{X_k : |\hat{\beta}_{a\tau k}| \geq \lambda_0 / \hat{\sigma}_{ak}^X\}$.

Step 2. *For $\hat{\varepsilon}_{ia\tau} = 1\{X_{ia} \leq X'_{i,-a} \tilde{\beta}_{a\tau}\} - \tau$ for $i \in [n]$, and $\xi = 1/n$, compute $\bar{\lambda}_{V\mathcal{T}}$ via (4.25).*

Step 3. *Recompute $\hat{\beta}_{a\tau}$ from $\|\cdot\|_{1,\hat{\varepsilon}}$ -penalized τ -quantile regression of X_a on X_{-a} with penalty $\bar{\lambda}_{V\mathcal{T}}$.*

Compute $\check{\beta}_{a\tau}$ from τ -quantile regression of X_a on $\{X_k : |\hat{\beta}_{a\tau k}| \geq \bar{\lambda}_{V\mathcal{T}} / \hat{\varepsilon}_{a\tau k}^X\}$.

Under regularity conditions stated in Section 5, with probability approaching 1, we have

$$\max_{a \in V} \sup_{\tau \in \mathcal{T}} \|\beta_{a\tau} - \check{\beta}_{a\tau}\| \lesssim \sqrt{\frac{s \log(|V|n)}{n}}.$$

The estimate of the prediction quantile graph is given by the support of $(\check{\beta}_{a\tau})_{a \in V, \tau \in \mathcal{T}}$, namely

$$\hat{E}^P(\tau) = \left\{ (a, b) \in V \times V : |\hat{\beta}_{a\tau b}| > \bar{\lambda}_{V\mathcal{T}} / \hat{\varepsilon}_{a\tau b}^X \right\}.$$

That is, it is induced by covariates selected by the ℓ_1 -penalized estimator. Those thresholded estimators not only have the same rates of convergence as of the original penalized estimators but also possess additional sparsity guarantees.

4.3. Estimators for \mathcal{W} -Conditional Quantile Graphical Models. In order to handle the additional conditioning events Ω_ϖ , $\varpi \in \mathcal{W}$, we propose to modify Algorithms 4.1 and 4.2 based on kernel smoothing. To that extent, we assume the observed data is of the form $\{(X_{iV}, W_i) : i = 1, \dots, n\}$, where W_i might be defined through additional variables. Furthermore, we assume for each conditioning event $\varpi \in \mathcal{W}$ we have access to a kernel function K_ϖ that is applied to W , to represent the relevant observations associated with ϖ (recall that we denote $P(W \in \Omega_\varpi)$ as $P(\varpi)$). We assume that $K_\varpi(W) = 1\{W \in \Omega_\varpi\}$.

Example 7 (Stock Returns Under Market Downside Movements, continued). In Example 2.3, we have W as the market return and the conditioning event as $\Omega_\varpi = \{W \leq \varpi\}$ which is parameterized by $\varpi \in \mathcal{W}$, a closed interval in \mathbb{R} . We might be interest in a fixed ϖ or on a family of values $\varpi \in (-\bar{\varpi}, 0]$. The latter induces $\mathcal{W} = \{\Omega_\varpi = \{W \leq \varpi\} : \varpi \in (-\bar{\varpi}, 0]\}$. The kernel function is simply $K_\varpi(t) = 1\{t \leq \varpi\}$.

This framework encompasses the previous framework by having $K_\varpi(W) = 1$ for all W . Moreover, it allows for a richer class of estimands which require estimators whose properties should hold uniformly over $\varpi \in \mathcal{W}$ as well. Next we propose estimators for this setting, i.e. we generalize the previous methods to account for the additional conditioning on $\varpi \in \mathcal{W}$. In what follows, with a slight abuse of notation we use ϖ to denote not only the index but also the event Ω_ϖ . For further notational convenience, we

denote $u = (a, \tau, \varpi) \in \mathcal{U} := V \times \mathcal{T} \times \mathcal{W}$ so that the set \mathcal{U} collects all the three relevant indices. With $\hat{\sigma}_{a\varpi j}^Z = \{\mathbb{E}_n[K_\varpi(W)(Z_j^a)^2]\}^{1/2}$, we define the following weighted ℓ_1 -norm $\|\beta\|_{1,\varpi} = \sum_{j \in [p]} \hat{\sigma}_{a\varpi j}^Z |\beta_j|$. This norm is ϖ dependent and provides the proper adjustments as we condition on different events associated with different ϖ 's.

We first consider estimators of CIQGMs conditional on the events in \mathcal{W} . In this setting, the model is correctly specified up to small approximation errors. The definition of the penalty parameter will be based on the random variable

$$\Lambda_{a\mathcal{T}\mathcal{W}} = \sup_{\tau \in \mathcal{T}} \max_{\varpi \in \mathcal{W}} \max_{j \in [p]} \left| \frac{\mathbb{E}_n[K_\varpi(W)(1\{U \leq \tau\} - \tau)Z_j^a]}{\sqrt{\tau(1-\tau)}\hat{\sigma}_{a\varpi j}^Z} \right|$$

where U_i are independent uniform $(0, 1)$ random variables, and set the penalty

$$\lambda_{V\mathcal{T}\mathcal{W}} = \max_{a \in V} \Lambda_{a\mathcal{T}\mathcal{W}}(1 - \xi/\{|V|n^{1+2d_W}\}|Z^a, W),$$

that is, the maximum of the $(1 - \xi/\{|V|n^{1+2d_W}\})$ conditional quantile of $\Lambda_{a\mathcal{T}\mathcal{W}}$. Algorithm 4.3 provides the definition of the estimator. Here $\mathcal{A}_{uj} = \{\alpha \in \mathbb{R} : |\alpha - \tilde{\beta}_{uj}| \leq 10/\{\hat{\sigma}_{a\varpi j}^Z \log n\}\}$, and denote $\lambda_u := \lambda_{V\mathcal{T}\mathcal{W}}\sqrt{\tau(1-\tau)}$.

Algorithm 4.3. (\mathcal{W} -Conditional CIQGM Estimator.) For $(a, \tau, \varpi) \in V \times \mathcal{T} \times \mathcal{W}$ and $j \in [p]$

Step 1. Compute $\hat{\beta}_u$ from $\|\cdot\|_{1,\varpi}$ -penalized τ -quantile regression of $K_\varpi(W)(X_a; Z^a)$ with penalty λ_u .

Compute $\tilde{\beta}_u$ from τ -quantile regression of $K_\varpi(W)(X_a; \{Z_k^a : |\hat{\beta}_{uk}| \geq \lambda_u/\hat{\sigma}_{a\varpi j}^Z\})$.

Step 2. Compute $\tilde{\gamma}_{ij}^j$ from the post-Lasso estimator of $K_\varpi(W)f_u Z_j^a$ on $K_\varpi(W)f_u Z_{-j}^a$.

Step 3. Construct the score function $\hat{\psi}_i(\alpha) = K_\varpi(W_i)(\tau - 1\{X_{ia} \leq Z_{ij}^a \alpha + Z_{i,-j}^a \tilde{\beta}_{u,-j}\})f_{iu}(Z_{ij}^a - Z_{i,-j}^a \tilde{\gamma}_{ij}^j)$ and for $L_{uj}(\alpha) = |\mathbb{E}_n[\hat{\psi}_i(\alpha)]|^2/\mathbb{E}_n[\hat{\psi}_i^2(\alpha)]$, set $\tilde{\beta}_{uj} \in \arg \min_{\alpha \in \mathcal{A}_{uj}} L_{uj}(\alpha)$.

Next we consider estimators of PQGMs conditional on the events in \mathcal{W} . Similar to the previous case, for $a \in V$ define

$$\bar{\Lambda}_{a\mathcal{T}\mathcal{W}} := 1.1 \sup_{\tau \in \mathcal{T}} \max_{\varpi \in \mathcal{W}} \max_{j \in [d]} \frac{|\mathbb{E}_n[K_\varpi(W)g\hat{\varepsilon}_{a\tau\varpi}X_{-a,j}]|}{\{\mathbb{E}_n[K_\varpi(W)\hat{\varepsilon}_{a\tau\varpi}^2 X_{-a,j}^2]\}^{1/2}} \quad (4.26)$$

where $(g_i)_{i=1}^n$ is a sequence of i.i.d. standard Gaussian random variables. The new penalty parameter $\bar{\lambda}_{V\mathcal{T}}$ is defined as

$$\bar{\lambda}_{V\mathcal{T}\mathcal{W}} := \max_{a \in V} \bar{\Lambda}_{a\mathcal{T}\mathcal{W}}(1 - \xi|X_{-a}|) \quad (4.27)$$

that is, the maximum of the $(1 - \xi)$ conditional quantile of $\bar{\Lambda}_{a\mathcal{T}\mathcal{W}}$. It will also be useful to define another weighted ℓ_1 -norm, $\|\beta\|_{1,\varpi\hat{\varepsilon}} := \sum_j \hat{\sigma}_{a\tau\varpi j}^{\varepsilon X} |\beta_j|$ with $\hat{\sigma}_{a\tau\varpi j}^{\varepsilon X} = \{\mathbb{E}_n[K_\varpi(W)\hat{\varepsilon}_{a\tau\varpi}^2 X_{-a,j}^2]\}^{1/2}$. We also denote $\hat{\sigma}_{a\varpi j}^X = \{\mathbb{E}_n[K_\varpi(W)X_{-a,j}^2]\}^{1/2}$. The penalty choice and weighted ℓ_1 -norm adapt to the unknown correlation structure across components and quantile indices. The following algorithm states the procedure, with $\lambda_{0\mathcal{W}} = 2(1 + 1/16)\sqrt{2 \log(8|V|^2\{ne/d_W\}^{2d_W}/\xi)}/n$.

Algorithm 4.4. (\mathcal{W} -Conditional PQGM Estimator.) For $(a, \tau, \varpi) \in V \times \mathcal{T} \times \mathcal{W}$

Step 1. Compute $\hat{\beta}_u$ from $\|\cdot\|_{1,\varpi}$ -penalized τ -quantile regression of X_a on X_{-a} with penalty $\lambda_{0\mathcal{W}}$.

Compute $\tilde{\beta}_u$ from τ -quantile regression of $K_\varpi(W)(X_a; \{X_{-a,k} : |\hat{\beta}_{uk}| \geq \lambda_{0\mathcal{W}}/\hat{\sigma}_{a\varpi k}^X\})$.

Step 2. For $\hat{\varepsilon}_{iu} = 1\{X_{ia} \leq X'_{i,-a}\tilde{\beta}_u\} - \tau$ for $i \in [n]$, and $\xi = 1/n$, compute $\bar{\lambda}_{V\mathcal{T}\mathcal{W}}$ via (4.27).

Step 3. Recompute $\tilde{\beta}_u$ from $\|\cdot\|_{1,\varpi\hat{\varepsilon}}$ -penalized τ -quantile regression of $K_\varpi(W)(X_a; X_{-a})$ with penalty $\bar{\lambda}_{V\mathcal{T}\mathcal{W}}$.

Compute $\check{\beta}_u$ from τ -quantile regression of $K_\varpi(W)(X_a; \{X_{-a,k} : |\hat{\beta}_{uk}| \geq \bar{\lambda}_{V\mathcal{T}\mathcal{W}}/\hat{\sigma}_{uk}^\varepsilon\})$.

Comment 4.3 (Computation of Penalty Parameter over \mathcal{W}). The penalty choices require one to maximize over $a \in V$, $\tau \in \mathcal{T}$ and $\varpi \in \mathcal{W}$. The set V is discrete and does not pose a significant challenge. However both other sets are continuous and additional care is needed. In most applications we are concerned with the case that \mathcal{W} is a low dimensional VC class of sets and it impacts the calculation only through indicator functions, which is precisely the case of \mathcal{T} . It follows that only a polynomial number (in n) of different values of τ and ϖ would need to be considered.⁹

5. MAIN THEORETICAL RESULTS

This section is devoted to theoretical guarantees associated with the proposed estimators. We will establish rates of convergence results for the proposed estimators as well as the (uniform) validity of confidence regions. These results build upon and contribute to an increasing literature on the estimation of many processes of interest with (high-dimensional) nuisance parameters.

Throughout, we will provide results for the estimators of the \mathcal{W} -conditional quantile graphical models as those can be generalized the other models by setting $K_\varpi(W) = 1$. Although some of the tools are similar, CIQGMs and PQGMs require different estimators and are subject to different assumptions. Thus, substantial different analyses are required.

5.1. \mathcal{W} -Conditional CIQGM. For $u = (a, \tau, \varpi) \in \mathcal{U}$, define the τ -conditional quantile function of X_a given $X_{V \setminus \{a\}}$ and ϖ as

$$Q_{X_a}(\tau|X_{V \setminus \{a\}}, \varpi) = Z^a \beta_u + r_u, \quad (5.28)$$

where Z^a is a p -dimensional vector of (known) transformations of $X_{V \setminus \{a\}}$, and r_u is an approximation error. The event $\varpi \in \mathcal{W}$ will be used for further conditioning through the function $K_\varpi(W) = 1\{W \in \varpi\}$.

We let $f_{X_a|X_{V \setminus \{a\}}, \varpi}(\cdot|X_{V \setminus \{a\}}, \varpi)$ denote the conditional density function of X_a given $X_{V \setminus \{a\}}$ and $\varpi \in \mathcal{W}$. We define $f_u := f_{X_a|X_{V \setminus \{a\}}, \varpi}(Q_{X_a}(\tau|X_{V \setminus \{a\}}, \varpi)|X_{V \setminus \{a\}}, \varpi)$ as the value of the conditional density function evaluated at the τ -conditional quantile. In our analysis we will consider for $u \in \mathcal{U}$

$$\underline{f}_u = \inf_{\|\delta\|=1} \frac{\mathbb{E}[f_u \{Z^a \delta\}^2 | \varpi]}{\mathbb{E}[\{Z^a \delta\}^2 | \varpi]} \quad \text{and} \quad \underline{f}_{\mathcal{U}} = \min_{u \in \mathcal{U}} \underline{f}_u. \quad (5.29)$$

Moreover, for each $u \in \mathcal{U}$ and $j \in [p]$ we define

$$\gamma_u^j = \arg \min_{\gamma} \mathbb{E}[f_u^2 K_\varpi(W)(Z_j^a - Z_{-j}^a \gamma)^2]. \quad (5.30)$$

This provides a weighted projection to construct the residuals

$$v_{uj} = f_u(Z_j^a - Z_{-j}^a \gamma_u^j)$$

that satisfy $\mathbb{E}[f_u Z_{-j}^a v_{uj} | \varpi] = 0$ for each $(u, j) \in \mathcal{U} \times [p]$.

⁹ A class of sets is said to be a VC class, if the VC dimension is finite. In what follows we use that the VC dimension provides a way to control how much we can overfit the data and it will also lead to (theoretically valid) recommendations for the penalty parameters. For the formal definition of VC class, see [100].

The estimands of interest are $\beta_u \in \mathbb{R}^p$, $u \in \mathcal{U}$, and can be written as the solution of (a continuum of) moment equations. Letting β_{uj} denote the j th component of β_u so that $\beta_{uj} \in \mathbb{R}$ solves

$$\mathbb{E}[\psi_{uj}(X, W, \beta, \eta_{uj})] = 0,$$

where the function ψ_{uj} is given by

$$\psi_{uj}(X, W, \beta, \eta_{uj}) = K_\varpi(W)(\tau - 1\{X_a \leq Z_j^a \beta + Z_{-j}^a \eta_{uj}^{(1)} + \eta_{uj}^{(3)}\})f_u(Z_j^a - Z_{-j}^a \eta_{uj}^{(2)}),$$

and the true value of the nuisance parameter is given by $\eta_{uj} = (\eta_{uj}^{(1)}, \eta_{uj}^{(2)}, \eta_{uj}^{(3)})$ with $\eta_{uj}^{(1)} = \beta_{u,-j}$, $\eta_{uj}^{(2)} = \gamma_u^j$, and $\eta_{uj}^{(3)} = r_u$. In what follows c, C denote some fixed constant, δ_n and Δ_n denote sequences go to zero with $\delta_n = n^{-\mu}$ for some sufficiently small μ . Denote $\mu_{\mathcal{W}} = \inf_{\varpi \in \mathcal{W}} \mathbb{P}(\varpi)$.

Condition CI. Let $u = (a, \tau, \varpi) \in \mathcal{U} := V \times \mathcal{T} \times \mathcal{W}$ and $(X_i, W_i)_{i=1}^n$ denote a sequence of independent and identically distributed random vectors generated accordingly to models (5.28) and (5.30):

(i) Suppose $\sup_{u \in \mathcal{U}, j \in [p]} \{\|\beta_u\| + \|\gamma_u^j\|\} \leq C$ and \mathcal{T} is a fixed compact set: (a) there exists $s = s_n$ such that $\sup_{u \in \mathcal{U}, j \in [p]} \{\|\beta_u\|_0 + \|\bar{\gamma}_u^j\|_0\} \leq s$, $\sup_{u \in \mathcal{U}, j \in [p]} \|\bar{\gamma}_u^j - \gamma_u^j\| + s^{-1/2} \|\bar{\gamma}_u^j - \gamma_u^j\|_1 \leq C\{n^{-1}s \log(|V|pn)\}^{1/2}$, where $\bar{\gamma}_u^j$ is approximately sparse; (b) the conditional distribution function of X_a given $X_{V \setminus \{a\}}$ and ϖ is absolutely continuous with continuously differentiable density $f_{X_a|X_{V \setminus \{a\}}, \varpi}(t|X_{V \setminus \{a\}}, \varpi)$ bounded by \bar{f} and its derivative bounded by \bar{f}' uniformly over $u \in \mathcal{U}$; (c) $|f_u - f_{u'}| \leq L_f \|u - u'\|$, $\|\beta_u - \beta_{u'}\| \leq L_\beta \|u - u'\|^\kappa$ with $\kappa \in [1/2, 1]$, and $\mathbb{E}[|K_\varpi(W) - K_{\varpi'}(W)|] \leq L_K \|\varpi - \varpi'\|$; (d) the VC dimension d_W of the set \mathcal{W} is fixed, $\{Q_{X_a}(\tau|X_{V \setminus \{a\}}, \varpi) : (\tau, \varpi) \in \mathcal{T} \times \mathcal{W}\}$ is a VC-subgraph with VC-dimension $1 + Cd_W$ for every $a \in V$;

(ii) The following moment conditions hold uniformly over $u \in \mathcal{U}$ and $j \in [p]$: $\mathbb{E}[|f_u v_{uj} Z_k^a|^2 | \varpi]^{1/2} \leq C \underline{f}_u$, $\min_{a \in V} \inf_{\|\delta\|=1} \mathbb{E}[\{(X_a, Z^a)\delta\}^2 | \varpi] \geq c$, $\max_{a \in V} \sup_{\|\delta\|=1} \mathbb{E}[\{(X_a, Z^a)\delta\}^4 | \varpi] \leq C$, $\mathbb{E}[f_u^2 (Z^a \delta)^2 | \varpi] \leq C \underline{f}_u^2 \mathbb{E}[(Z^a \delta)^2 | \varpi]$, $\max_{j,k} \frac{\mathbb{E}[|f_u v_{uj} Z_k^a|^3 | \varpi]^{1/3}}{\mathbb{E}[|f_u v_{uj} Z_k^a|^2 | \varpi]^{1/2}} \log^{1/2}(pn|V|) \leq \delta_n \{n\mathbb{P}(\varpi)\}^{1/6}$;

(iii) Furthermore, for some fixed $q \geq 4 \vee (1 + 2d_W)$, $\sup_{u \in \mathcal{U}, \|\delta\|=1} \mathbb{E}[|(X_a, Z^a)\delta|^2 r_u^2 | \varpi] \leq C \mathbb{E}[r_u^2 | \varpi] \leq Cs/n$, $\max_{u \in \mathcal{U}, j \in [p]} |\mathbb{E}[f_u r_u v_{uj} | \varpi]| \leq \delta_n n^{-1/2}$, $\mathbb{E}[\max_{i \leq n} \sup_{u \in \mathcal{U}} |K_\varpi(W) r_{iu}|^q] \leq C$, and with probability $1 - \Delta_n$, uniformly over $u \in \mathcal{U}, j \in [p]$: $\mathbb{E}_n[r_u^2 v_{uj}^2 | \varpi] + \mathbb{E}_n[r_u^2 | \varpi] \lesssim n^{-1} s \log(p|V|n)$, $\mathbb{E}_n[K_\varpi(W)\{r_u + r_u^2\}(Z^a \delta)^2] \leq \delta_n \mathbb{E}_n[K_\varpi(W) f_u (Z^a \delta)^2]$;

(iv) For a fixed $q \geq 4 \vee (1 + d_W)$, $\text{diam}(\mathcal{W}) \leq n^{1/2q}$, $\mathbb{E}[\max_{i \leq n} \|X_{iV}\|_\infty^q \vee \max_{a \in V} \|Z_i^a\|_\infty^q]^{1/q} / \mu_{\mathcal{W}} \leq M_n$, $\mathbb{E}[\max_{i \leq n} \sup_{u \in \mathcal{U}, j \in [p]} |v_{iuj}|^q]^{1/q} \leq L_n$, $(L_f + L_K)^2 M_n^2 \log^2(p|V|n) \leq \delta_n n \mu_{\mathcal{W}}^3 \underline{f}_{\mathcal{U}}^6$, $M_n^4 \log(p|V|n) \log n \leq \delta_n^2 n \mu_{\mathcal{W}}^2 \underline{f}_{\mathcal{U}}^2$, $s^2 \log^2(p|V|n) \leq \delta_n^2 n \underline{f}_{\mathcal{U}}^4 \mu_{\mathcal{W}}^6$, $s^3 \log^3(p|V|n) \leq \delta_n^4 n \underline{f}_{\mathcal{U}}^2 \mu_{\mathcal{W}}^3$, $L_n^2 s \log^{3/2}(p|V|n) \leq \delta_n \underline{f}_{\mathcal{U}} (n \mu_{\mathcal{W}})^{1/2}$, $M_n s \sqrt{\log(p|V|n)} \leq \delta_n n^{1/2} \mu_{\mathcal{W}} \underline{f}_{\mathcal{U}}$.

Condition CI assumes various conditional moment conditions to allow for the estimation to be conditional on $\varpi \in \mathcal{W}$. Those are analogous to the (unconditional) conditions in the high-dimensional literature in quantile regression models, [19]. In particular, condition CI(i) assumes smoothness of the density function, and of coefficients. Condition CI(ii) assumes conditions on the (conditional) population design matrices such as the ratio between eigenvalues. Condition CI(iii) pertains to the approximations errors and assumes mild moment conditions. Finally Condition CI(iv) provides sufficient conditions on the allowed growth of the model via p and $|V|$ relative to the available sample size n . Note, Condition CI(iii) also assume d_W is bounded by fixed q , and the proof can easily be extended to other cases.

Condition CI is a high level condition intended to allow approximate sparse models, approximation errors, tail events in \mathcal{W} , and to require only q moments (going beyond sub-Gaussian variables). When applied to the special case of sub-Gaussian, exactly sparse, and singleton \mathcal{W} , it becomes a relatively standard assumption. For example, without approximation error, e.g. the multivariate Gaussian case, Condition CI(iii) can be removed entirely. By allowing for a large number of variables (and transformations) the approximation errors can be controlled when the quantile functions belong to some smooth function class (e.g. Sobolev space). Although it is outside the scope of the current work, the ideas and results can be generalized to dependent data, using results from [32].

Based on Condition CI, we derive our main results regarding the proposed estimator. Moreover, we also establish new results for ℓ_1 -penalized quantile regression methods that hold uniformly over the indices $u \in \mathcal{U}$. The following theorems summarize these results.

Theorem 1 (Uniform Rates of Convergence for \mathcal{W} -Conditional Penalized Quantile Regression). *Under Condition CI, we have that with probability at least $1 - o(1)$*

$$\|\hat{\beta}_u - \beta_u\| \lesssim \sqrt{\frac{s(1 + d_W) \log(p|V|n)}{nf_u P(\varpi)}}, \quad \text{uniformly over } u = (a, \tau, \varpi) \in \mathcal{U}$$

Moreover, the thresholded estimator $\hat{\beta}^{\bar{\lambda}}$, with $\bar{\lambda} = \sqrt{(1 + d_W) \log(p|V|n)/n}$ and $\hat{\beta}_{uj}^{\bar{\lambda}} = \hat{\beta}_{uj} 1\{|\hat{\beta}_{uj}| > \bar{\lambda} \hat{\sigma}_{a\varpi j}^Z\}$, satisfies the same rate and $\|\hat{\beta}^{\bar{\lambda}}\|_0 \lesssim s$.

Theorem 1 builds upon ideas in [12] however the proof strategy is designed to derive rates that are adaptive to each $u \in \mathcal{U}$. Indeed the rates of convergence are u -dependent and they show a slower rate for rare events $\varpi \in \mathcal{W}$.

Theorem 2 (Uniform Rates of Convergence for \mathcal{W} -Conditional Weighted Lasso). *Under Condition CI, we have that with probability at least $1 - o(1)$*

$$\|\hat{\gamma}_u^j - \gamma_u^j\| \lesssim \frac{1}{f_u} \sqrt{\frac{s(1 + d_W) \log(p|V|n)}{nP(\varpi)}} \quad \text{and} \quad \|\hat{\gamma}_u^j\|_0 \lesssim s, \quad \text{uniformly over } u = (a, \tau, \varpi) \in \mathcal{U}, j \in [p].$$

The following result establishes a uniform Bahadur representation for the final estimators.

Theorem 3 (Uniform Bahadur Representation for \mathcal{W} -Conditional CIQGM). *Under Condition CI, the estimator $(\hat{\beta}_{uj})_{u \in \mathcal{U}, j \in [p]}$ satisfies*

$$\sigma_{uj}^{-1} \sqrt{n}(\hat{\beta}_{uj} - \beta_{uj}) = \mathbb{U}_n(u, j) + O_P(\delta_n) \text{ in } \ell^\infty(\mathcal{U} \times [p]),$$

where $\sigma_{uj}^2 = \tau(1 - \tau)E[K_\varpi(W)v_{uj}^2]^{-1}$ and

$$\mathbb{U}_n(u, j) := \frac{\{\tau(1 - \tau)E[K_\varpi(W)v_{uj}^2]\}^{-1/2}}{\sqrt{n}} \sum_{i=1}^n (\tau - 1\{U_i(a, \varpi) \leq \tau\}) K_\varpi(W_i) v_{i,uj},$$

where $U_1(a, \varpi), \dots, U_n(a, \varpi)$ are i.i.d. uniform $(0, 1)$ random variables, independent of $v_{1,uj}, \dots, v_{n,uj}$.

Theorem 3 plays a key role. However, it is important to note that the marginal distribution of $\mathbb{U}_n(u, j)$ is pivotal. Nonetheless, there is a non-trivial correlation structure between $U(a, \varpi)$ and $U(\tilde{a}, \tilde{\varpi})$. In order to construct confidence regions with non-conservative guarantees, we rely on a multiplier bootstrap

method. We will approximate the process $\mathcal{N} = (\mathcal{N}_{uj})_{u \in \mathcal{U}, j \in [p]}$ by the Gaussian multiplier bootstrap based on estimates $\hat{\psi}_{uj} := \{\tau(1 - \tau)\mathbb{E}_n[K_\varpi(W)\hat{v}_{uj}^2]\}^{-1/2}(\tau - 1\{X_a \leq Z^a\hat{\beta}_u\})K_\varpi(W_i)\hat{v}_{uj}$ of $\bar{\psi}_{uj}(U, W) = \{\tau(1 - \tau)\mathbb{E}_n[K_\varpi(W)v_{uj}^2]\}^{-1/2}(\tau - 1\{U(a, \varpi) \leq \tau\})K_\varpi(W)v_{uj}$, namely

$$\hat{\mathcal{G}} = (\hat{\mathcal{G}}_{uj})_{u \in \mathcal{U}, j \in [p]} = \left\{ \frac{1}{\sqrt{n}} \sum_{i=1}^n g_i \hat{\psi}_{uj}(X_i, W_i) \right\}_{u \in \mathcal{U}, j \in [p]}$$

where $(g_i)_{i=1}^n$ are independent standard normal random variables which are independent from the data $(W_i)_{i=1}^n$. Based on Theorem 5.2 of [28], the following result shows that the multiplier bootstrap provides a valid approximation to the large sample probability law of $\sqrt{n}(\check{\beta}_{uj} - \beta_{uj})_{u \in \mathcal{U}, j \in [p]}$ which is suitable for the construction of uniform confidence bands over the set of indices associated with $I_a(b)$ for all $a, b \in V$. We let \mathcal{P}_n denote the collection of distributions P for the data such that Condition CI is satisfied for given n . This is the collection of all approximately sparse models where the above sparsity conditions, moment conditions, and growth conditions are satisfied.

Corollary 1 (Gaussian Multiplier Bootstrap for \mathcal{W} -Conditional CIQGM). *Under Condition CI with $\delta_n = o(\{(1 + d_W) \log(p|V|n)\}^{-1/2})$, and $(1 + d_W) \log(p|V|n) = o(\{(n/L_n^2)^{1/7} \wedge (n^{1-2/q}/L_n^2)^{1/3}\})$, we have that*

$$\sup_{P \in \mathcal{P}_n} \sup_{t, t' \in \mathbb{R}, u \in \mathcal{U}, b \in V} \left| \mathbb{P}_P \left(\max_{j \in I_a(b)} \frac{|\check{\beta}_{uj} - \beta_{uj}|}{n^{-1/2}\sigma_{uj}} \in [t, t'] \right) - \mathbb{P}_P \left(\max_{j \in I_a(b)} |\hat{\mathcal{G}}_{uj}| \in [t, t'] \mid (X_i, W_i)_{i=1}^n \right) \right| = o(1)$$

Corollary 1 allows the construction of simultaneous confidence regions for the coefficients that are uniformly valid over the set of data generating processes induced by Condition CI. Based on the coefficients whose intervals do not overlap zero, we can construct a conditional independence graph process $\hat{E}^I(\tau, \varpi), \tau \in \mathcal{T}, \varpi \in \mathcal{W}$ that contains the true conditional independence quantile graph with a specified probability.

5.2. \mathcal{W} -Conditional PQGM. In this section, we derive theoretical guarantees for the \mathcal{W} -conditional predictive quantile estimators uniformly over $u = (a, \tau, \varpi) \in \mathcal{U}$. For each $u \in \mathcal{U}$ the estimand of interest is $\beta_u \in \mathbb{R}^p$ that corresponds to the best linear predictor under asymmetric loss function, namely

$$\beta_u \in \arg \min_{\beta} \mathbb{E}[\rho_\tau(X_a - X'_{-a}\beta) \mid \varpi] \quad (5.31)$$

where the event $\varpi \in \mathcal{W}$ is used for further conditioning. In the analysis below, the conditioning is implemented through the function $K_\varpi(W) = 1\{W \in \varpi\}$.

In the analysis of this case, the main issue is to handle the inherent misspecification of the linear form $X'_{-a}\beta_u$ with respect to the true conditional quantile. The first consequence is to handle the identification condition. Given X_{-a} and $\varpi \in \mathcal{W}$, we let $f_u := f_{X_a|X_{-a}, \varpi}(X'_{-a}\beta_u|X_{-a}, \varpi)$ denote the value of the conditional density function evaluated at $X'_{-a}\beta_u$. In our analysis, we will consider

$$\underline{f}_u = \inf_{\|\delta\|=1} \frac{\mathbb{E}[f_u\{X'_{-a}\delta\}^2 \mid \varpi]}{\mathbb{E}[(X'_{-a}\delta)^2 \mid \varpi]} \quad \text{and} \quad \underline{f}_{\mathcal{U}} = \min_{u \in \mathcal{U}} \underline{f}_u. \quad (5.32)$$

We remark that \underline{f}_u defined in (5.32) differs from (5.29) which is the standard conditional density at the true quantile value. It turns out that Knight's identity can be used by exploiting the first order condition associated with the optimization problem (5.31) which yields zero mean condition similar to the conditional quantile condition.

A second consequence of the misspecification is the lack of pivotality of the score. Such pivotal property was convenient in the previous section to define penalty parameters and to conduct inference. We will exploit bounds on the VC-dimension of the relevant classes of sets formally stated below.

Condition P. Let $\mathcal{U} = V \times \mathcal{T} \times \mathcal{W}$ and $(X_i, W_i)_{i=1}^n$ denote a sequence of independent and identically distributed random vectors generated accordingly to models (5.31):

(i) Suppose that $\sup_{u \in \mathcal{U}} \|\beta_u\| \leq C$ and \mathcal{T} is a fixed compact set: (a) there exists $s = s_n$ and $\bar{\beta}_u$ such that $\sup_{u \in \mathcal{U}} \|\bar{\beta}_u\|_0 \leq s$, $\sup_{u \in \mathcal{U}} \|\bar{\beta}_u - \beta_u\| + s^{-1/2} \|\bar{\beta}_u - \beta_u\|_1 \leq \sqrt{s/n}$; (b) the conditional distribution function of X_a given X_{-a} and ϖ is absolutely continuous with continuously differentiable density $f_{X_a|X_{-a}, \varpi}(t | X_{-a}, \varpi)$ such that its values are bounded by \bar{f} and its derivative is bounded by \bar{f}' uniformly over $u \in \mathcal{U}$; (c) $|f_u - f_{u'}| \leq L_f \|u - u'\|$, $\|\beta_u - \beta_{u'}\| \leq L_\beta \|u - u'\|^\kappa$ with $\kappa \in [1/2, 1]$, and $\mathbb{E}[|K_\varpi(W) - K_{\varpi'}(W)|] \leq L_K \|\varpi - \varpi'\|$; (d) the VC dimension d_W of the set \mathcal{W} is fixed, $\{1\{X_a \leq X'_{-a}\beta_u\} : (\tau, \varpi) \in \mathcal{T} \times \mathcal{W}\}$ is a VC-class with VC-dimension $1 + d_W$ for every $a \in V$;

(ii) The following moment conditions hold uniformly over $u \in \mathcal{U}$: $\min_{a \in V} \inf_{\|\delta\|=1} \mathbb{E}[\{X'_{-a}\delta\}^2 | \varpi] \geq c$, $\max_{a \in V} \sup_{\|\delta\|=1} \mathbb{E}[\{X'_{-a}\delta\}^4 | \varpi] \leq C$;

((iii) With probability $1 - \Delta_n$, uniformly over $u \in \mathcal{U}$ and $a \in V$: $\mathbb{E}_n[K_\varpi(W)\{|X'_{-a}(\bar{\beta}_u - \beta_u)| + |X'_{-a}(\bar{\beta}_u - \beta_u)|^2\}(Z^a\delta)^2] \leq \delta_n \mathbb{E}_n[K_\varpi(W)f_u(X'_{-a}\delta)^2]$;

(iv) For a fixed $q \geq 4 \vee (1 + d_W)$, we have that: $\text{diam}(\mathcal{W}) \leq n^{1/2q}$, $\mathbb{E}[\max_{i \leq n} \|X_{iV}\|_\infty^q]^{1/q} / \mu_{\mathcal{W}} \leq M_n$, $M_n^2 \log^7(n|V|) \leq \delta_n n \underline{f}_{\mathcal{U}}^2 \mu_{\mathcal{W}}^2$, $M_n^4 \log(n|V|) \log n \leq \delta_n n \mu_{\mathcal{W}}$, $(L_f + L_K)^2 M_n^2 \log^2(|V|n) \leq \delta_n n \mu_{\mathcal{W}}^3 \underline{f}_{\mathcal{U}}^6$, $M_n^2 s \log^{3/2}(n|V|) \leq \delta_n \underline{f}_{\mathcal{U}} (n \mu_{\mathcal{W}})^{1/2}$, $M_n s \sqrt{\log(n|V|)} \leq \delta_n (n \mu_{\mathcal{W}})^{1/2}$, and $s^3 \log^5(n|V|) \leq \delta_n n \underline{f}_{\mathcal{U}}^2 \mu_{\mathcal{W}}^2$.

Condition P is a high-level condition. It allows to cover conditioning events $\varpi \in \mathcal{W}$ whose probability can decrease to zero (although slower than $n^{-1/4}$).

Next we derive our main results regarding the proposed estimator for the best linear predictor. These results are also new ℓ_1 -penalized quantile regression methods as it holds under possible misspecification of the conditional quantile function and hold uniformly over the indices $u \in \mathcal{U}$. The following theorem summarizes the result.

Theorem 4 (Uniform Rates of Convergence for \mathcal{W} -Conditional Penalized Quantile Regression under Misspecification). *Under Condition P, we have that with probability at least $1 - o(1)$, uniformly over $u = (a, \tau, \varpi) \in \mathcal{U}$,*

$$\|\hat{\beta}_u - \beta_u\| \lesssim \sqrt{\frac{s(1 + d_W) \log(|V|n)}{n \underline{f}_u \mathbb{P}(\varpi)}}.$$

The data-driven choice of penalty parameter helps diminish the regularization bias and also allow to obtain sparse estimators with provably rates of convergence (through thresholding). Moreover, the u specific penalty parameter combined with the new analysis yields an adaptive rate of convergence to each $u \in \mathcal{U}$ unlike previous works.

Comment 5.1 (Simultaneous Confidence Bands for Coefficients in PQGMs). We note that in some applications we might be interested in constructing (simultaneous) confidence bands for the coefficients in PQGMs. In particular, this would include the cases practitioners are using a misspecified linear specification in a

quantile regression model. Provided the conditional density function at $X'_{-u}\beta_u$ can be estimated, a version of Algorithm 4.3 using the penalty parameters in Algorithm 4.4 for the initial step can deliver such confidence regions via a multiplier bootstrap.

6. APPLICATION: INTERNATIONAL FINANCIAL CONTAGION AND SYSTEMIC RISK

There is widespread disagreement about what financial contagion entails, e.g. [41, 51]. The existing measures of contagion are mainly based on linear correlation and can only account for certain types of risk network structure or do not provide inference procedures for the large scale networks estimated. This paper defines contagion occurs whenever the quantile partial correlation from one country to another country is nonzero, i.e. the presence of edges in PQGM. This definition takes into account network spillover effects when identifying contagion and measuring systemic risk. Here, the weight of an edge indicates the strength of contagion effects. The estimated contagion network taking into account global interconnectedness is important for Eurozone financial regulators to identify globally systemically important EU countries, or for global financial portfolio diversification.

We revisit the analysis of international financial contagion, [35]. We provide an alternative approach to the literature by visualizing tail interdependence via PQGM. As shown in Section 2.2, our framework naturally extends the systemic risk measure CoVaR taking into account tail network spillover effects, hence after learning PQGM from data, we identify systemically important countries using our new systemic risk measures. We can also provide inference on networks estimated. To simplify the visualization, we provide graphical visualization for the confidence intervals of $\Delta CoVaRs$ in Figure 3 of Section 6.2.

We focus on examining financial contagion through the volatility spillovers perspective, i.e. recovering volatility interconnectedness. [48] reported that international stock markets are related through their volatilities instead of returns. [40] studied the return and volatility spillovers of 19 countries and found differences in return and volatility spillovers.¹⁰

We use average two-day equity index returns,¹¹ September 2009 to September 2013, from Morgan Stanley Capital International (MSCI). The returns are all translated into dollar-equivalents as of September 6th 2013.¹² We use absolute returns as a proxy for volatility.¹³ We have a total of 45 countries in our sample, there are 21 developed markets (Australia, Austria, Belgium, Canada, Denmark, France, Germany, Hong Kong, Ireland, Italy, Japan, Netherlands, New Zealand, Norway, Portugal, Singapore, Spain, Sweden, Switzerland, the United Kingdom, the United States), 21 emerging markets (Brazil, Chile, Mexico, Greece, Israel, China, Colombia, Czech Republic, Egypt, Hungary, India, Indonesia, Korea, Malaysia, Peru, Philippines, Poland, Russia, Taiwan, Thailand, Turkey), and 3 frontier markets (Argentina, Morocco, Jordan).

¹⁰Modelling the time dependence in volatility is an important issue, although not the focus of this work, there is no uniform agreement on whether there is dependence once taking into account the heteroskedasticity of market returns or shifts in standard errors, see [96].

¹¹This is to control for the fact that markets in different countries are not open during the same hours. Results are robust to whether using two-day returns or using daily returns as in older versions of our work. Daily returns are also adjusted for weekends and holidays.

¹²We calculate returns based on U.S. dollars since these were most frequently used in past work on contagion.

¹³This has historically been used in the literature. While we do recognize that there are many different methods calculating volatility measures, volatility measuring itself is a large research study area and is outside the scope of the current work.




FIGURE 2. International Financial Contagion. Note: Lack of edges between nodes indicates that neither node's equity index volatility help predict the other node's volatility at τ -th quantile. Arrows (or directed edges) mean that the source node's volatility helps predict the target node's volatility. These graphs show that the volatility transmission mechanism are asymmetric at different volatility quantiles (or tails).

6.1. Contagion Networks. Figure 2 provides a full-sample analysis of global volatility network spillovers at different tails. The networks are estimated via Algorithm 4.2.

We denote 10% quantile as Low Tail, 50% quantile as Median, 90% quantile as Up Tail. Results learnt from both PQGMs and GGM are presented. GGM or "Gaussian Graph" in Figure 2 means the graph is estimated via graphical lasso (e.g., [53]), and the final graph is chosen by Extended Bayesian Information Criterion (ebic), see [52]. Our purpose is to show the usefulness of PQGM in representing nonlinear tail interdependence allowing for heteroscedasticity and to show that PQGM can measure correlation asymmetry through looking at the tails of the distribution (not specific to any model).

There are significant differences in the network structure in terms of volatility spillovers when using PQGM and GGM. PQGM permits asymmetries in correlation dynamics, suited to investigate the presence of asymmetric responses. We find significant increase interdependence at the up tail between the volatility series, that is we find downside correlations (high volatility) are much larger than upside correlations (low volatility). This confirms findings in the finance literature that financial markets become more interdependent during high volatility periods.

We also find if two countries locate in the same geographic region, with many similarities in terms of market structure and history, they tend to be more closely connected (homophily effect as stated in network terminology), while two economies locate in separate geographic regions are less likely directly connected. In addition, we find among European Union member countries, Germany appears to play a major role in the transmission of shocks to others; while in Asia, Hong Kong, Thailand, and Singapore appear to play major roles; and among all the north and south American countries, Canada and US play major roles.


6.2. Systemic Risk. With the estimated network, we can use different network statistics to measure the systemic risk contributions. Below we focus on the modified $\Delta CoVaR$ measure mentioned in Section 2.2.

Figure 3 provides German's $\Delta CoVaRs$ with $\tau = 0.9$ and their 90% uniform confidence intervals obtained via Corollary 1. It reconfirms that France, Italy and UK contribute the most to German's $\Delta CoVaR$, means conditional on those countries being under distress relative to their median states, German would be affected the most. It is also interesting to find that other countries such as Netherlands can also have effects on German's $\Delta CoVaR$ although less statistically significant in terms of the magnitude of the effect.

In addition, we present $net-\Delta CoVaR$ discussed in Section 2.2 with $\tau = 0.9$, i.e. the Up Tail, in Figure 4 which shows that: globally, total volatility spillovers from Germany and France to the others are much larger than total volatility spillovers from the others to them, and their $net-\Delta CoVaR$ are positive. Both Greece and Spain have negative $net-\Delta CoVaR$.

REFERENCES

- [1] Alberto Abadie, Guido Imbens, and Fanyin Zheng. Inference for misspecified models with fixed regressors. *Journal of the American Statistical Association*, 109(508):1601–1614, 2014.
- [2] Daron Acemoglu, Asuman Ozdaglar, and Alireza Tahbaz-Salehi. Cascades in networks and aggregate volatility. Technical report, National Bureau of Economic Research, 2010.

FIGURE 3. German's $\Delta CoVaR$ and 90% Confidence Intervals.

- [3] Daron Acemoglu, Asuman Ozdaglar, and Alireza Tahbaz-Salehi. Systemic risk and stability in financial networks. *The American Economic Review*, 105(2):564–608, 2015.
- [4] Tobias Adrian and Markus Brunnermeier. Covar. *The American Economic Review*, 106(7):1705–1741, 2016.
- [5] Torben Andersen, Tim Bollerslev, Peter Christoffersen, and Francis Diebold. Financial risk measurement for financial risk management. *Handbook of the Economics of Finance*, 2:1127–1220, 2013.
- [6] Andrew Ang and Joseph Chen. Asymmetric correlations of equity portfolios. *Journal of Financial Economics*, 63(3):443–494, 2002.
- [7] Andrew Ang, Joseph Chen, and Yuhang Xing. Downside risk. *Review of Financial Studies*, 19(4):1191–1239, 2006.
- [8] Joshua Angrist, Victor Chernozhukov, and Iván Fernández-Val. Quantile regression under misspecification, with an application to the us wage structure. *Econometrica*, 74(2):539–563, 2006.
- [9] Patrick Bajari and Lixin Ye. Deciding between competition and collusion. *Review of Economics and statistics*, 85(4):971–989, 2003.
- [10] Onureena Banerjee, Laurent El Ghaoui, and Alexandre d’Aspremont. Model selection through sparse maximum likelihood estimation for multivariate gaussian or binary data. *The Journal of Machine Learning Research*, 9:485–516, 2008.
- [11] Alexandre Belloni, Daniel Chen, Victor Chernozhukov, and Christian Hansen. Sparse models and methods for optimal instruments with an application to eminent domain. *Econometrica*, 80(6):2369–2429, 2012.
- [12] Alexandre Belloni and Victor Chernozhukov. ℓ_1 -penalized quantile regression for high dimensional sparse models. *Annals of Statistics*, 39(1):82–130, 2011.




FIGURE 4. Volatility Spillovers Net Contribution of Each Country

- [13] Alexandre Belloni and Victor Chernozhukov. Inference methods for high-dimensional sparse econometric models. *Advances in Economics and Econometrics, 10th World Congress of Econometric Society*, III:245–295, 2013.
- [14] Alexandre Belloni and Victor Chernozhukov. Least squares after model selection in high-dimensional sparse models. *Bernoulli*, 19(2):521–547, 2013. ArXiv, 2009.
- [15] Alexandre Belloni, Victor Chernozhukov, Denis Chetverikov, and Ying Wei. Uniformly valid post-regularization confidence regions for many functional parameters in z-estimation framework. *arXiv:1512.07619*, 2015.

- [16] Alexandre Belloni, Victor Chernozhukov, and Iván Fernández-Val. Conditional quantile processes based on series or many regressors. *arXiv*, 2011.
- [17] Alexandre Belloni, Victor Chernozhukov, Ivan Fernández-Val, and Christian Hansen. Program evaluation and causal inference with high-dimensional data. *Econometrica*, 85(1):233–298, 2017.
- [18] Alexandre Belloni, Victor Chernozhukov, and Christian Hansen. Inference on treatment effects after selection among high-dimensional controls. *The Review of Economic Studies*, 81(2):608–650, 2014.
- [19] Alexandre Belloni, Victor Chernozhukov, and Kengo Kato. Robust inference in high-dimensional approximately sparse quantile regression models. *arXiv preprint arXiv:1312.7186*, 2013.
- [20] Alexandre Belloni, Victor Chernozhukov, and Kengo Kato. Uniform post-selection inference for least absolute deviation regression and other z-estimation problems. *Biometrika*, 102(1):77–94, 2015.
- [21] Alexandre Belloni, Victor Chernozhukov, and Lie Wang. Square-root-lasso: Pivotal recovery of sparse signals via conic programming. *Biometrika*, 98(4):791–806, 2011. Arxiv, 2010.
- [22] Peter Bickel, Ya’acov Ritov, and Alexandre Tsybakov. Simultaneous analysis of lasso and dantzig selector. *Annals of Statistics*, 37(4):1705–1732, 2009.
- [23] Monica Billio, Mila Getmansky, Andrew W Lo, and Lorian Pelizzon. Measuring systemic risk in the finance and insurance sectors. 2010.
- [24] Pietro Bonaldi, Ali Hortaçsu, and Jakub Kastl. An empirical analysis of funding costs spillovers in the euro-zone with application to systemic risk. Technical report, National Bureau of Economic Research, 2015.
- [25] Tony Cai, Weidong Liu, and Xi Luo. A constrained ℓ_1 minimization approach to sparse precision matrix estimation. *Journal of the American Statistical Association*, 106(494), 2011.
- [26] Mehmet Caner and Anders Kock. Asymptotically honest confidence regions for high dimensional parameters by the desparsified conservative lasso. *Journal of Econometrics*, 203(1):143–168, 2018.
- [27] Victor Chernozhukov. Extremal quantile regression. *Annals of Statistics*, pages 806–839, 2005.
- [28] Victor Chernozhukov, Denis Chetverikov, and Kengo Kato. Gaussian approximations and multiplier bootstrap for maxima of sums of high-dimensional random vectors. *The Annals of Statistics*, 41(6):2786–2819, 2013.
- [29] Victor Chernozhukov, Denis Chetverikov, and Kengo Kato. Gaussian approximation of suprema of empirical processes. *The Annals of Statistics*, 42(4):1564–1597, 2014.
- [30] Victor Chernozhukov, Denis Chetverikov, and Kengo Kato. Comparison and anti-concentration bounds for maxima of gaussian random vectors. *Probability Theory and Related Fields*, 162:47–70, 2015.
- [31] Victor Chernozhukov, Denis Chetverikov, and Kengo Kato. Empirical and multiplier bootstraps for supreme of empirical processes of increasing complexity, and related gaussian couplings. *Stochastic Processes and their Applications*, 126(12):3632–3651, 2016.
- [32] Victor Chernozhukov, Denis Chetverikov, and Kengo Kato. Central limit theorems and bootstrap in high dimensions. *The Annals of Probability*, 45(4):2309–2352, 2017.
- [33] Victor Chernozhukov, Christian Hansen, and Martin Spindler. Post-selection and post-regularization inference in linear models with very many controls and instruments. *American Economic Review: Papers and Proceedings*, 105(5):486–490, 2015.
- [34] Khai Xiang Chiong and Hyungsik Roger Moon. Estimation of graphical models using the $\ell_1, 2$ norm. *The Econometrics Journal*, 21(3):247–263, 2018.
- [35] Stijn Claessens and Kristin Forbes. *International Financial Contagion*. Springer, 2001.

- [36] David Cox and Nanny Wermuth. *Multivariate dependencies: Models, analysis and interpretation*, volume 67. CRC Press, 1996.
- [37] Philip Dawid. Conditional independence in statistical theory. *Journal of the Royal Statistical Society. Series B (Methodological)*, pages 1–31, 1979.
- [38] Aureo De Paula. Econometrics of network models. In *Advances in Economics and Econometrics: Theory and Applications, Eleventh World Congress*, pages 268–323. Cambridge University Press Cambridge, 2017.
- [39] Arthur Dempster. Covariance selection. *Biometrics*, pages 157–175, 1972.
- [40] Francis Diebold and Kamil Yilmaz. Measuring financial asset return and volatility spillovers, with application to global equity markets. *The Economic Journal*, 119(534):158–171, 2009.
- [41] Francis X Diebold and Kamil Yilmaz. Trans-atlantic equity volatility connectedness: Us and european financial institutions, 2004–2014. *Journal of Financial Econometrics*, 14(1):81–127, 2015.
- [42] Peng Ding. On the conditional distribution of the multivariate t distribution. *The American Statistician*, 70(3):293–295, 2016.
- [43] Mathias Drton and Michael Perlman. Model selection for gaussian concentration graphs. *Biometrika*, 91(3):591–602, 2004.
- [44] Mathias Drton and Michael Perlman. Multiple testing and error control in gaussian graphical model selection. *Statistical Science*, 22(3):430–449, 2007.
- [45] Mathias Drton and Michael Perlman. A sinful approach to gaussian graphical model selection. *Journal of Statistical Planning and Inference*, 138(4):1179–1200, 2008.
- [46] David Edwards. *Introduction to graphical modelling*. Springer, 2000.
- [47] Matthew Elliott, Benjamin Golub, and Matthew O Jackson. Financial networks and contagion. *American Economic Review*, 104(10):3115–53, 2014.
- [48] Robert Engle and Raul Susmel. Common volatility in international equity markets. *Journal of Business & Economic Statistics*, 11(2):167–176, 1993.
- [49] Jianqing Fan, Jinchi Lv, and Lei Qi. Sparse high dimensional models in economics. *Annual review of economics*, 3:291, 2011.
- [50] Max Farrell. Robust inference on average treatment effects with possibly more covariates than observations. *Journal of Econometrics*, 174(2):1–23, 2015.
- [51] Kristin J Forbes and Roberto Rigobon. No contagion, only interdependence: measuring stock market comovements. *The journal of finance*, 57(5):2223–2261, 2002.
- [52] Rina Foygel and Mathias Drton. Extended bayesian information criteria for gaussian graphical models. In *Advances in Neural Information Processing Systems*, pages 604–612, 2010.
- [53] Jerome Friedman, Trevor Hastie, and Robert Tibshirani. Sparse inverse covariance estimation with the graphical lasso. *Biostatistics*, 9(3):432–441, 2008.
- [54] Lars Peter Hansen. Challenges in identifying and measuring systemic risk. *Risk Topography: Systemic Risk and Macro Modeling*, 2014.
- [55] Wolfgang Karl Härdle, Weining Wang, and Lining Yu. Tenet: Tail-event driven network risk. *Journal of Econometrics*, 192(2):499–513, 2016.
- [56] Joseph E Harrington. Detecting cartels, handbook of antitrust economics, 2008.
- [57] Nikolaus Hautsch, Julia Schaumburg, and Melanie Schienle. Forecasting systemic impact in financial networks. *International Journal of Forecasting*, 30(3):781–794, 2014.

- [58] Nikolaus Hautsch, Julia Schaumburg, and Melanie Schienle. Financial network systemic risk contributions. *Review of Finance*, 19(2):685–738, 2015.
- [59] Xuming He, Lan Wang, and Hyokyoung Grace Hong. Quantile-adaptive model-free variable screening for high-dimensional heterogeneous data. *The Annals of Statistics*, 41(1):342–369, 2013.
- [60] James Heckman and Rodrigo Pinto. Causal analysis after haavelmo. *Econometric Theory*, 31(1):115–151, 2015.
- [61] Matthew O Jackson. *Social and economic networks*. Princeton university press, 2010.
- [62] Jana Jankova and Sara van de Geer. Confidence intervals for high-dimensional inverse covariance estimation. *Electronic Journal of Statistics*, 9(1):1205–1229, 2015.
- [63] Harry Joe. *Multivariate models and multivariate dependence concepts*. CRC Press, 1997.
- [64] Jakub Kastl. Recent advances in empirical analysis of financial markets: industrial organization meets finance. *Honoré, B., Pakes A., Piazzesi, M. and L. Samuelson (editors) Advances in Economics and Econometrics*, 2:231–270, 2017.
- [65] Kengo Kato. Group lasso for high dimensional sparse quantile regression models. Preprint, ArXiv, 2011.
- [66] Keith Knight. Limiting distributions for L_1 regression estimators under general conditions. *The Annals of Statistics*, 26:755–770, 1998.
- [67] Keith Knight. Asymptotics of the regression quantile basic solution under misspecification. *Applications of Mathematics*, 53(3):223–234, 2008.
- [68] Roger Koenker. *Quantile regression*. Cambridge University Press, New York, 2005.
- [69] Roger Koenker and Gilbert Bassett. Regression quantiles. *Econometrica*, 46(1):33–50, 1978.
- [70] Eric Kolaczyk. *Statistical analysis of network data*. Springer, 2009.
- [71] Steffen Lauritzen. *Graphical models*. Oxford University Press, 1996.
- [72] Michel Ledoux and Michel Talagrand. *Probability in Banach Spaces (Isoperimetry and processes)*. Ergebnisse der Mathematik und ihrer Grenzgebiete, Springer-Verlag, 1991.
- [73] Ying-Ying Lee. Efficiency bounds for semiparametric estimation of quantile regression under misspecification. *University of Wisconsin-Madison*, 2009.
- [74] Hannes Leeb and Benedikt Pötscher. Can one estimate the unconditional distribution of post-model-selection estimators? *Econometric Theory*, 24(02):338–376, 2008.
- [75] Han Liu, Fang Han, Ming Yuan, John Lafferty, and Larry Wasserman. High-dimensional semiparametric gaussian copula graphical models. *The Annals of Statistics*, 40(4):2293–2326, 2012.
- [76] Han Liu, John Lafferty, and Larry Wasserman. The nonparanormal: Semiparametric estimation of high dimensional undirected graphs. *The Journal of Machine Learning Research*, 10:2295–2328, 2009.
- [77] Han Liu and Lie Wang. Tiger: A tuning-insensitive approach for optimally estimating gaussian graphical models. *Electronic Journal of Statistics*, 11(1):241–294, 2017.
- [78] Weidong Liu and Xi Luo. High-dimensional sparse precision matrix estimation via sparse column inverse operator. *arXiv preprint arXiv:1203.3896*, 2012.
- [79] Po-Ling Loh and Martin J Wainwright. Structure estimation for discrete graphical models: Generalized covariance matrices and their inverses. In *Advances in Neural Information Processing Systems*, pages 2087–2095, 2012.
- [80] Francois Longin and Bruno Solnik. Extreme correlation of international equity markets. *The Journal of Finance*, 56(2):649–676, 2001.

- [81] Marloes Maathuis, Mathias Drton, Steffen Lauritzen, and Martin Wainwright. *Handbook of Graphical Models*. CRC Press, 2018.
- [82] Nicolai Meinshausen and Peter Bühlmann. High-dimensional graphs and variable selection with the lasso. *The Annals of Statistics*, pages 1436–1462, 2006.
- [83] Sahand Negahban, Pradeep Ravikumar, Martin Wainwright, and Bin Yu. A unified framework for high-dimensional analysis of m-estimators with decomposable regularizers. *Statistical Science*, 27(4):538–557, 2012.
- [84] Mark Newman. *Networks: an introduction*. Oxford university press, 2010.
- [85] Yang Ning and Han Liu. A general theory of hypothesis tests and confidence regions for sparse high dimensional models. *The Annals of Statistics*, 45(1):158–195, 2017.
- [86] Roberto Imbuzeiro Oliveira. The lower tail of random quadratic forms, with applications to ordinary least squares and restricted eigenvalue properties. *Probability Theory and Related Fields*, 166(3-4):1175–1194, 2016.
- [87] Andrew Patton. On the out-of-sample importance of skewness and asymmetric dependence for asset allocation. *Journal of Financial Econometrics*, 2(1):130–168, 2004.
- [88] Andrew Patton. Modelling asymmetric exchange rate dependence. *International economic review*, 47(2):527–556, 2006.
- [89] Judea Pearl. *Causality*. Cambridge university press, 2009.
- [90] Jie Peng, Pei Wang, Nengfeng Zhou, and Ji Zhu. Partial correlation estimation by joint sparse regression models. *Journal of the American Statistical Association*, 104(486):735–746, 2009.
- [91] Robert H Porter. Detecting collusion. *Review of Industrial Organization*, 26(2):147–167, 2005.
- [92] Pradeep Ravikumar, Martin J Wainwright, and John D Lafferty. High-dimensional ising model selection using ℓ_1 -regularized logistic regression. *The Annals of Statistics*, 38(3):1287–1319, 2010.
- [93] Pradeep Ravikumar, Martin J Wainwright, Garvesh Raskutti, and Bin Yu. High-dimensional covariance estimation by minimizing ℓ_1 -penalized log-determinant divergence. *Electronic Journal of Statistics*, 5:935–980, 2011.
- [94] James Robins. A new approach to causal inference in mortality studies with a sustained exposure period: application to control of the healthy worker survivor effect. *Mathematical modelling*, 7(9-12):1393–1512, 1986.
- [95] Joseph Romano and Michael Wolf. Exact and approximate stepdown methods for multiple hypothesis testing. *Journal of the American Statistical Association*, 100(469):94–108, 2005.
- [96] Catalin Starica and Clive Granger. Nonstationarities in stock returns. *Review of economics and statistics*, 87(3):503–522, 2005.
- [97] Tingni Sun and Cun-Hui Zhang. Sparse matrix inversion with scaled lasso. *The Journal of Machine Learning Research*, 14(1):3385–3418, 2013.
- [98] Sara van de Geer. High-dimensional generalized linear models and the lasso. *Annals of Statistics*, 36(2):614–645, 2008.
- [99] Sara Van de Geer, Peter Bühlmann, Ya’acov Ritov, and Ruben Dezeure. On asymptotically optimal confidence regions and tests for high-dimensional models. *The Annals of Statistics*, 42(3):1166–1202, 2014.
- [100] Aad van der Vaart and Jon Wellner. *Weak Convergence and Empirical Processes*. Springer Series in Statistics, 1996.

- [101] Aad van der Vaart and Jon Wellner. Empirical process indexed by estimated functions. *IMS Lecture Notes-Monograph Series*, 55:234–252, 2007.
- [102] Jialei Wang and Mladen Kolar. Inference for high-dimensional exponential family graphical models. In *Proc. of AISTATS*, volume 51, pages 751–760, 2016.
- [103] Yuanshan Wu and Guosheng Yin. Conditional quantile screening in ultrahigh-dimensional heterogeneous data. *Biometrika*, 102(1):65–76, 2015.
- [104] Lingzhou Xue and Hui Zou. Regularized rank-based estimation of high-dimensional nonparanormal graphical models. *The Annals of Statistics*, 40(5):2541–2571, 2012.
- [105] Lingzhou Xue, Hui Zou, and Tianxi Cai. Nonconcave penalized composite conditional likelihood estimation of sparse ising models. *The Annals of Statistics*, 40(3):1403–1429, 2012.
- [106] Eunho Yang, Pradeep Ravikumar, Genevera I Allen, and Zhandong Liu. Graphical models via univariate exponential family distributions. *The Journal of Machine Learning Research*, 16(1):3813–3847, 2015.
- [107] Ming Yuan. High dimensional inverse covariance matrix estimation via linear programming. *The Journal of Machine Learning Research*, 99:2261–2286, 2010.
- [108] Ming Yuan and Yi Lin. Model selection and estimation in the gaussian graphical model. *Biometrika*, 94(1):19–35, 2007.
- [109] Cun-Hui Zhang and Stephanie Zhang. Confidence intervals for low dimensional parameters in high dimensional linear models. *Journal of the Royal Statistical Society: Series B (Statistical Methodology)*, 76(1):217–242, 2014.
- [110] Qi Zheng, Limin Peng, and Xuming He. Globally adaptive quantile regression with ultra-high dimensional data. *Annals of statistics*, 43(5):2225, 2015.

APPENDIX A. IMPLEMENTATION DETAILS OF ALGORITHMS

This section provides details of the algorithms mentioned in Section 4. Note, for the weighted-Lasso estimator, the choice of penalty level $\lambda := 1.1n^{-1/2}2\Phi^{-1}(1 - \xi/N_n)$ and penalty loading $\hat{\Gamma}_\tau = \text{diag}[\hat{\Gamma}_{\tau kk}, k \in [p] \setminus \{j\}]$ is a diagonal matrix defined by the following procedure: (1) Compute the post-Lasso estimator $\tilde{\gamma}_{a\tau}^j$ based on λ and initial values $\hat{\Gamma}_{\tau kk} = \max_{i \leq n} \|f_{ia\tau} Z_i^a\|_\infty \{\mathbb{E}_n[\|f_{a\tau} Z_k^a\|^2]\}^{1/2}$. (2) Compute the residuals $\hat{v}_{ia\tau j} = f_{ia\tau}(Z_{ij}^a - Z_{i,-j}^a \tilde{\gamma}_{a\tau}^j)$ and update the loadings

$$\hat{\Gamma}_{\tau kk} = \sqrt{\mathbb{E}_n[f_{a\tau}^2 | Z_k^a \hat{v}_{a\tau j}|^2]}, \quad k \in [p] \setminus \{j\} \quad (\text{A.33})$$

and use them to recompute the post-Lasso estimator $\tilde{\gamma}_{a\tau}^j$. In the case of Algorithm 4.1 we can take $N_n = |V|p^3n^3$, in the case of Algorithm 4.3 we take $N_n = |V|p^2\{pn^3\}^{1+dw}$. Denote $\hat{\sigma}_{aj}^Z = \{\mathbb{E}_n[(Z_j^a)^2]\}^{1/2}$, $\hat{\sigma}_{aj}^X = \{\mathbb{E}_n[X_{-a,j}^2]\}^{1/2}$, $\hat{\sigma}_{a\varpi j}^Z = \{\mathbb{E}_n[K_\varpi(W)(Z_j^a)^2]\}^{1/2}$, and $\hat{\sigma}_{a\varpi j}^X = \{\mathbb{E}_n[K_\varpi(W)X_{-a,j}^2]\}^{1/2}$.

Detailed version of Algorithm 4.1 (CIQGM)

For each $a \in V$, $\tau \in \mathcal{T}$, and $j \in [p]$, perform the following:

- (1) Run Post- ℓ_1 -quantile regression of X_a on Z^a ; keep fitted value $Z_{-j}^a \tilde{\beta}_{a\tau, -j}$,

$$\begin{aligned} \hat{\beta}_{a\tau} &\in \arg \min_\beta \mathbb{E}_n[\rho_\tau(X_a - Z^a \beta)] + \lambda_{V\mathcal{T}} \sqrt{\tau(1-\tau)} \sum_{j=1}^p \hat{\sigma}_{aj}^Z |\beta_j| \\ \tilde{\beta}_{a\tau} &\in \arg \min_\beta \mathbb{E}_n[\rho_\tau(X_a - Z^a \beta)] : \beta_j = 0 \text{ if } |\hat{\beta}_{a\tau j}| \leq \lambda_{V\mathcal{T}} \sqrt{\tau(1-\tau)} / \hat{\sigma}_{aj}^Z. \end{aligned}$$

- (2) Run Post-Lasso of $f_{a\tau} Z_j^a$ on $f_{a\tau} Z_{-j}^a$; keep the residual $\tilde{v}_i := f_{ia\tau} \{Z_{ij}^a - Z_{i,-j}^a \tilde{\gamma}_{a\tau}^j\}$,

$$\begin{aligned} \hat{\gamma}_{a\tau}^j &\in \arg \min_\gamma \mathbb{E}_n[f_{a\tau}^2 (Z_j^a - Z_{-j}^a \gamma)^2] + \lambda \|\hat{\Gamma}_\tau \gamma\|_1 \\ \tilde{\gamma}_{a\tau}^j &\in \arg \min_\gamma \mathbb{E}_n[f_{a\tau}^2 (Z_j^a - Z_{-j}^a \gamma)^2] : \text{support}(\gamma) \subseteq \text{support}(\hat{\gamma}_{a\tau}^j). \end{aligned}$$

- (3) Run Instrumental Quantile Regression of $X_a - Z_{-j}^a \tilde{\beta}_{a\tau, -j}$ on Z_j^a using \tilde{v} as the instrument for Z_j^a ,

$$\check{\beta}_{a\tau, j} \in \arg \min_{\alpha \in \mathcal{A}_{a\tau j}} \frac{\{\mathbb{E}_n[(1\{X_a \leq Z_j^a \alpha + Z_{-j}^a \tilde{\beta}_{a\tau, -j}\} - \tau) \tilde{v}]\}^2}{\mathbb{E}_n[(1\{X_a \leq Z_j^a \alpha + Z_{-j}^a \tilde{\beta}_{a\tau, -j}\} - \tau)^2 \tilde{v}^2]},$$

with $\mathcal{A}_{a\tau j} = \{\alpha \in \mathbb{R} : |\alpha - \tilde{\beta}_{a\tau j}| \leq 10/\{\hat{\sigma}_{aj}^Z \log n\}\}$.

Detailed version of Algorithm 4.2 (PQGM)

For each $a \in V$, and $\tau \in \mathcal{T}$, perform the following:

- (1) Run Post- ℓ_1 -quantile regression of X_a on X_{-a} ,

$$\begin{aligned} \hat{\beta}_{a\tau} &\in \arg \min_\beta \mathbb{E}_n[\rho_\tau(X_a - X'_{-a} \beta)] + \lambda_0 \sum_{j \in [d]} \hat{\sigma}_{aj}^X |\beta_j| \\ \tilde{\beta}_{a\tau} &\in \arg \min_\beta \mathbb{E}_n[\rho_\tau(X_a - X'_{-a} \beta)] : \beta_j = 0 \text{ if } |\hat{\beta}_{a\tau j}| \leq \lambda_0 / \hat{\sigma}_{aj}^X. \end{aligned}$$

- (2) Set $\hat{\varepsilon}_{ia\tau} = 1\{X_{ia} \leq X'_{i,-a} \tilde{\beta}_{a\tau}\} - \tau$ for $i \in [n]$. Compute the penalty level $\bar{\lambda}_{V\mathcal{T}}$ via (4.25).

- (3) Run Post- ℓ_1 -quantile regression of X_a on X_{-a} ,

$$\begin{aligned} \hat{\beta}_{a\tau} &\in \arg \min_\beta \mathbb{E}_n[\rho_\tau(X_a - X'_{-a} \beta)] + \bar{\lambda}_{V\mathcal{T}} \sum_{j \in [d]} \{\mathbb{E}_n[\hat{\varepsilon}_{a\tau}^2 X_{-a,j}^2]\}^{1/2} |\beta_j| \\ \tilde{\beta}_{a\tau} &\in \arg \min_\beta \mathbb{E}_n[\rho_\tau(X_a - X'_{-a} \beta)] : \beta_j = 0 \text{ if } |\hat{\beta}_{a\tau j}| \leq \bar{\lambda}_{V\mathcal{T}} / \{\mathbb{E}_n[\hat{\varepsilon}_{a\tau}^2 X_{-a,j}^2]\}^{1/2}. \end{aligned}$$

Detailed version of Algorithm 4.3 (\mathcal{W} -Conditional CIQGM)

For each $u = (a, \tau, \varpi) \in \mathcal{U} = V \times \mathcal{T} \times \mathcal{W}$, and $j \in [p]$, perform the following:

- (1) Run Post- ℓ_1 -quantile regression of X_a on Z^a ; keep fitted value $Z_{-j}^a \tilde{\beta}_{u,-j}$,

$$\begin{aligned} \hat{\beta}_u &\in \arg \min_{\beta} \mathbb{E}_n[K_{\varpi}(W)\rho_{\tau}(X_a - Z^a\beta)] + \lambda_u \sum_{j=1}^p \hat{\sigma}_{a\varpi j}^Z |\beta_j| \\ \tilde{\beta}_u &\in \arg \min_{\beta} \mathbb{E}_n[K_{\varpi}(W)\rho_{\tau}(X_a - Z^a\beta)] : \beta_j = 0 \text{ if } |\hat{\beta}_{uj}| \leq \lambda_u / \hat{\sigma}_{a\varpi j}^Z. \end{aligned}$$

- (2) Run Post-Lasso of $f_u Z_j^a$ on $f_u Z_{-j}^a$; keep the residual $\tilde{v} := f_u(Z_j^a - Z_{-j}^a \tilde{\gamma}_u^j)$,

$$\begin{aligned} \hat{\gamma}_u^j &\in \arg \min_{\gamma} \mathbb{E}_n[K_{\varpi}(W)f_u^2(Z_j^a - Z_{-j}^a\gamma)^2] + \lambda \|\hat{\Gamma}_u \gamma\|_1 \\ \tilde{\gamma}_u^j &\in \arg \min_{\gamma} \mathbb{E}_n[K_{\varpi}(W)f_u^2(Z_j^a - Z_{-j}^a\gamma)^2] : \text{support}(\gamma) \subseteq \text{support}(\hat{\gamma}_u^j). \end{aligned}$$

- (3) Run Instrumental Quantile Regression of $X_a - Z_{-j}^a \tilde{\beta}_{u,-j}$ on Z_j^a using \tilde{v} as the instrument,

$$\check{\beta}_{uj} \in \arg \min_{\alpha \in \mathcal{A}_{uj}} \frac{\{\mathbb{E}_n[K_{\varpi}(W)(1\{X_a \leq Z_j^a \alpha + Z_{-j}^a \tilde{\beta}_{u,-j}\} - \tau)\tilde{v}\}^2}{\mathbb{E}_n[K_{\varpi}(W)(1\{X_a \leq Z_j^a \alpha + Z_{-j}^a \tilde{\beta}_{u,-j}\} - \tau)^2 \tilde{v}^2]}$$

where $\mathcal{A}_{uj} := \{\alpha \in \mathbb{R} : |\alpha - \tilde{\beta}_{uj}| \leq 10 / \{\hat{\sigma}_{a\varpi j}^Z \log n\}\}$.

Detailed version of Algorithm 4.4 (\mathcal{W} -Conditional PQGM)

For each $u = (a, \tau, \varpi) \in \mathcal{U} = V \times \mathcal{T} \times \mathcal{W}$ perform the following:

- (1) Run Post- ℓ_1 -quantile regression of X_a on X_{-a} ,

$$\begin{aligned} \hat{\beta}_u &\in \arg \min_{\beta} \mathbb{E}_n[K_{\varpi}(W)\rho_{\tau}(X_a - X'_{-a}\beta)] + \lambda_{0\mathcal{W}} \sum_{j \in [d]} \hat{\sigma}_{a\varpi j}^X |\beta_j| \\ \tilde{\beta}_u &\in \arg \min_{\beta} \mathbb{E}_n[K_{\varpi}(W)\rho_{\tau}(X_a - X'_{-a}\beta)] : \beta_j = 0 \text{ if } |\hat{\beta}_{uj}| \leq \lambda_{0\mathcal{W}} / \hat{\sigma}_{a\varpi j}^X. \end{aligned}$$

- (2) Set $\hat{\varepsilon}_{iu} = 1\{X_{ia} \leq X'_{i,-a} \tilde{\beta}_u\} - \tau$ for $i \in [n]$, compute $\bar{\lambda}_{V\mathcal{T}\mathcal{W}}$ via (4.27).

- (3) Run Post- ℓ_1 -quantile regression of X_a on X_{-a} ,

$$\begin{aligned} \hat{\beta}_u &\in \arg \min_{\beta} \mathbb{E}_n[K_{\varpi}(W)\rho_{\tau}(X_a - X'_{-a}\beta)] + \bar{\lambda}_{V\mathcal{T}\mathcal{W}} \sum_{j \in [d]} \{\mathbb{E}_n[K_{\varpi}(W)\hat{\varepsilon}_u^2 X_{-a,j}^2]\}^{1/2} |\beta_j|. \\ \tilde{\beta}_u &\in \arg \min_{\beta} \mathbb{E}_n[K_{\varpi}(W)\rho_{\tau}(X_a - X'_{-a}\beta)] : \beta_j = 0 \text{ if } |\hat{\beta}_{uj}| \leq \bar{\lambda}_{V\mathcal{T}\mathcal{W}} / \{\mathbb{E}_n[K_{\varpi}(W)\hat{\varepsilon}_u^2 X_{-a,j}^2]\}^{1/2}. \end{aligned}$$

APPENDIX B. SIMULATIONS OF QUANTILE GRAPHICAL MODELS

In this section, we perform numerical examples to illustrate the performance of the estimators proposed for QGMs. We will consider several different designs. In order to compare with other proposals we will consider both Gaussian and non-Gaussian examples.

B.1. Isotropic Non-Gaussian Example. In general, the equivalence between a zero in the inverse covariance matrix and a pair of conditional independent variables will break down for non-gaussian distributions. The nonparanormal graphical models extends Gaussian Graphical Models to Semiparametric Gaussian Copula models by transforming the variables with smooth functions. We illustrate the applicability of CIQGM in representing the conditional independence structure of a set of variables when the random variables are not even jointly nonparanormal.

Consider i.i.d. copies of an d -dimensional random vector $\tilde{X}_V = (\tilde{X}_1, \dots, \tilde{X}_{d-1}, \tilde{X}_d)$ from the following multivariate normal distribution, $\tilde{X}_V \sim N(0, I_{d \times d})$, where $I_{d \times d}$ is the identity matrix. Further, we generate

$$X_d = -\sqrt{\frac{2}{3\pi-2}} + \sqrt{\frac{\pi}{3\pi-2}} \tilde{X}_{d-1}^2 |\tilde{X}_d|. \quad (\text{B.34})$$

It follows that $E[X_d] = \sqrt{\frac{\pi}{3\pi-2}}(E[|\tilde{X}_d|] - \sqrt{2/\pi}) = 0$ and $\text{Var}(X_d) = \frac{\pi}{3\pi-2}(E[\tilde{X}_d^2 \cdot \tilde{X}_{d-1}^4] - \frac{2}{\pi}) = 1$. In addition, equation (B.34) is a location-scale-shift model in which the conditional median of the response is zero while quantile functions other than the median are nonzero. We define vector X_V as

$$X_V = (X_d, \tilde{X}_1, \dots, \tilde{X}_{d-1})'.$$

In this new set of variables, only X_d and \tilde{X}_{d-1} (i.e., node 1 and 15, when $d = 15$) are not conditionally independent. Nonetheless, the covariance matrix of X_V is still $I_{d \times d}$.

Next we consider an example with $n = 300$ and $d = 15$. We show graphs, in Figure 5 and 6, estimated by both CIQGM(s) and GGMs in this non-Gaussian setting.




FIGURE 5. QGM and GGM

In Figure 5, Gaussian means the graph is estimated by using graphical lasso without any transformation of X_V , and the final graph is chosen by Extended Bayesian Information Criterion (ebic), see [52]. Nonparanormal means the graph is estimated using graphical lasso (likelihood based approach) with nonparanormal

transformation of X_V , see [76], and again the final graph is chosen by ebic. Both graphs are estimated using R-package **huge**.

In Figure 6, as a robustness check, we also compare results produced by CIQGM with those produced by neighborhood selection methods (pseudo-likelihood approach), e.g. TIGER of [77] in R-package **flare** the left graph is when choosing the turning parameter to be $\sqrt{\frac{\log d}{n}}$ while the right graph is when choosing the tuning parameter to be $2\sqrt{\frac{\log d}{n}}$. Throughout, we use Tiger2 represent TIGER with penalty level $2\sqrt{\frac{\log d}{n}}$. As expected, GGM cannot detect the correct dependence structure when the joint distribution is non-Gaussian while CIQGM can still represent the right conditional independence structure.




FIGURE 6. TIGER

B.2. Gaussian Examples.

B.2.1. Graph Recovery. In this subsection, we start with comparing the numerical performance of QGM and other methods, e.g. TIGER of [77] and graphical lasso algorithm (Glasso) of [53], in graph recovery using simulated datasets with different pairs of (n, d) . We start with one simulation for illustration purpose (the results are summarized in Figure 7), and then we show the performance of QGM through estimated degree distribution with 100 simulations (the results are summarized in Figure 8).

We mainly consider the Hub graph, as mentioned in [77], which also corresponds to the star network mentioned in [2, 3]. In line with [77], we generate a d -dimensional sparse graph $G^I = (V, E^I)$ represents the conditional independence structure between the variables. In our simulations, we consider 12 settings to compare these methods: (A) $n = 200, d = 10$; (B) $n = 200, d = 20$; (C) $n = 200, d = 40$; (D) $n = 400, d = 10$; (E) $n = 400, d = 20$; (F) $n = 400, d = 40$; (G) $n = 200, d = 100$; (H) $n = 200, d = 200$; (I) $n = 200, d = 400$; (J) $n = 400, d = 100$; (K) $n = 400, d = 200$; (L) $n = 400, d = 400$. We adopt the following model for generating undirected graphs and precision matrices.

Hub graph. The d nodes are evenly partitioned into $d/20$ (or $d/10$ when $d < 20$) disjoint groups with each group contains 20 (or 10) nodes. Within each group, one node is selected as the hub and we add edges between the hub and the other 19 (or 9) nodes in that group. For example, the resulting graph has 190 edges when $d = 200$ and 380 edges when $d = 400$. Once the graph is obtained, we generate an adjacency matrix E^I by setting the nonzero off-diagonal elements to be 0.3 and the diagonal elements to be 0. We calculate its smallest eigenvalue $\Lambda_{\min}(E^I)$. The precision matrix is constructed as

$$\Theta = \mathbf{D}[E^I + (|\Lambda_{\min}(E^I)| + 0.2) \cdot I_{d \times d}] \mathbf{D} \quad (\text{B.35})$$

where $\mathbf{D} \in \mathbb{R}^{d \times d}$ is a diagonal matrix with $\mathbf{D}_{jj} = 1$ for $j = 1, \dots, d/2$ and $\mathbf{D}_{jj} = 1.5$ for $j = d/2 + 1, \dots, d$. The covariance matrix $\Sigma := \Theta^{-1}$ is then computed to generate the multivariate normal data: $X_1, \dots, X_d \sim N(0, \Sigma)$. Below we provide simulation results using different estimators: PQGM¹⁴, TIGER and Glasso. We start with one simulation as an illustration:




FIGURE 7. One simulation

¹⁴Given the graphs are generated from multivariate Gaussian distribution we can use PQGM to simplify the computation.




FIGURE 7. One simulation (Cont.)

Figure 7 shows that: for the low dimensional cases, $d = 10, 20, 40$, n is large compared to d , CIQGM is comparable to TIGER and both are better than Glasso in terms of false positives; for the high dimensional cases, $d = \{100, 200, 300\}$, we can compare the performance of different graph estimators through looking at the denseness of the estimated graph (e.g., whether it is even or not), and again, both CIQGM and TIGER perform well in terms of graph recovery as compared to Glasso, and their performance are getting better when n is increasing.

In what follows, Figure 8 shows the degree distribution of true graph, the estimated ones, and the standard deviations of the degree difference (between the true graph and the estimated ones). It is based on simulations of Hub graph with $n = 500$ and $d = 40$. Simulated 100 times.




FIGURE 8. Upper panel shows the degree distribution of the true graphs. Middle panel shows the degree distribution of the estimated graphs. Bottom panel shows the standard deviations of the degree difference (between the true graph and the estimated ones). Hub graph with $n = 500$ and $d = 40$. Simulated 100 times.

B.2.2. Inference. In this subsection Table 1 shows the numerical performance of CIQGM, based on Algorithm 4.1, on estimating Erdős-Rényi random graphs. More precisely, we construct approximate 90% confidence intervals for β_{ab} with $\tau = 0.5$, and we report the coverage probabilities. Note, in the jointly Gaussian distributed case, we have closed form solution of β_{ab} as shown in Example 1.

Erdős-Rényi random graph. We add an edge between each pair of nodes with probability $\sqrt{\log d/n}/d$ independently.

Once the graph is obtained, we construct the adjacency matrix E^I and generate the precision matrix Θ using (B.35) but setting $\mathbf{D}_{jj} = 1$ for $j = 1, \dots, d/2$ and $\mathbf{D}_{jj} = 1.5$ for $j = d/2 + 1, \dots, d$. We then invert Θ to get the covariance matrices $\Sigma := \Theta^{-1}$ and generate the multivariate Gaussian data: $X_1, \dots, X_d \sim N(0, \Sigma)$.

TABLE 1. Erdős-Rényi Random Graph

	(a, b)	$n = 200$	$n = 500$	$n = 1000$
d=20	(1,20)	84.0	87.5	91.5
	(10,11)	84.0	88.0	92.5
	(19, 20)	86.5	86.0	90.0
	ACP	86.3	89.4	89.8
d=50	(1,50)	86.5	93.0	90.5
	(25,26)	88.0	87.0	91.0
	(49, 50)	87.5	90.5	87.5
	ACP	86.7	89.4	90.1
d=100	(1,100)	82.5	81	89.0
	(50,51)	85.0	86	92.0
	(99, 100)	78.5	84	87.0
	ACP	86.5	86.8	90

(a, b) , coverage probability for β_{ab} ; ACP, average coverage probability for β_{ab} , with $a \in V$, $b \in V \setminus \{a\}$. Simulated 200 times.

APPENDIX C. PROOFS OF SECTION 4

Proof of Theorem 1. By Lemma 6, under Condition CI, for any θ such that $\|\theta\|_0 \leq C s \ell_n$, $\ell_n \rightarrow \infty$ slowly, we have that

$$\|\sqrt{f_u} Z^a \theta\|_{n, \varpi} / \{E[K_\varpi(W) f_u (Z^a \theta)^2]\}^{1/2} = 1 + o_P(1).$$

Moreover, $E[K_\varpi(W) f_u (Z^a \theta)^2] \geq \underline{f}_u E[K_\varpi(W) (Z^a \theta)^2]$, $E[K_\varpi(W) (Z^a \theta)^2] = E[(Z^a \theta)^2 | \varpi] P(\varpi)$, and $E[(Z^a \theta)^2 | \varpi] \geq c \|\theta\|^2$ by Condition CI. Lemma 6 further implies that the ratio of the minimal and maximal eigenvalues of order $s \ell_n$ are bounded away from zero and from above uniformly over $\varpi \in \mathcal{W}$ and $a \in V$ with probability $1 - o(1)$. Therefore, since $c \{P(\varpi)\}^{1/2} \|\delta\|_1 \leq \|\delta\|_{1, \varpi} \leq C \{P(\varpi)\}^{1/2} \|\delta\|_1$, we have $\kappa_{u, 2c} \geq c$ uniformly over $u \in \mathcal{U}$ with the same probability for n large enough, see for instance [22].

To establish rates of convergence of the estimator obtained in Step 1 we will apply Lemma 1. Consider the events Ω_1, Ω_2 , and Ω_3 as defined in (D.48), (D.49) and (D.50). By the choice of λ_u we have $P(\Omega_1) \geq 1 - o(1)$. By Condition CI with $\bar{R}_{u\xi} \leq C s \log(p|V|n)/n$ and Lemma 2 we have $P(\Omega_2) \geq 1 - o(1)$. Moreover, $P(\Omega_3) \geq 1 - o(1)$ by Lemma 3 with $t_3 \leq C n^{-1/2} \sqrt{(1 + d_W) \log(p|V|n L_f)}$.

Using the same argument (with Z^a replacing X_{-a}) as in (C.45), (C.46), and (C.47), for

$$\delta \in A_u := \Delta_{\varpi, 2c} \cup \{v : \|v\|_{1, \varpi} \leq 2c\bar{R}_{u\xi}/\lambda_u, \|\sqrt{f_u}Z^a v\|_{n, \varpi} \geq C\sqrt{s(1+d_W)\log(p|V|n)/n/\kappa_{u, 2c}}\},$$

with the restricted set defined as $\Delta_{u, 2\tilde{c}} = \{\delta : \|\delta_{T_u^c}\|_1 \leq 2\tilde{c}\|\delta_{T_u}\|_1\}$ for $u \in \mathcal{U}$. we have $\bar{q}_{A_u} \geq c(\underline{f}_{\mathcal{U}}^{3/2}/\bar{f}')\mu_{\mathcal{W}}^{1/2}/\{\sqrt{s}\max_{a \in V, i \leq n} \|Z_i^a\|_\infty\}$ where $\max_{a \in V, i \leq n} \|Z_i^a\|_\infty \lesssim_P M_n$. Thus the conditions on \bar{q}_{A_u} are satisfied since Condition CI assumes $M_n^2 s^2 \log(p|V|n) \leq n\mu_{\mathcal{W}} \underline{f}_{\mathcal{U}}^3$. The conditions on the approximation error are assumed in Condition CI.

Therefore, setting $\xi = 1/\log n$, by Lemma 1 we have uniformly over $u = (a, \tau, \varpi) \in \mathcal{U}$

$$\begin{aligned} \|\sqrt{f_u}Z^a(\hat{\beta}_u - \beta_u)\|_{n, \varpi} &\lesssim \sqrt{(1 + (t_3/\lambda_u)\bar{R}_{u\xi})} + (\lambda_u + t_3)\sqrt{s} \lesssim \sqrt{\frac{s(1+d_W)\log(p|V|n)}{n\tau(1-\tau)}} \\ \|\hat{\beta}_u - \beta_u\|_{1, \varpi} &\lesssim s\sqrt{\frac{(1+d_W)\log(p|V|n)}{n}} \end{aligned} \quad (\text{C.36})$$

here we used that $\lambda_u \leq C\sqrt{\frac{(1+d_W)\log(p|V|n)}{n}}$. Indeed by Lemma 15 with $\tilde{x}_{ij} = K_\varpi(W_i)Z_{ij}^a$ and $\hat{\sigma}_j \geq cP(\varpi)^{1/2}$, we can bound $\Lambda_{a\tau\varpi}(1 - \xi/\{|V|n^{1+2d_W}\}|X_{-a}, W)$ under $M_n^2 \log(p|V|n/\{\tau(1-\tau)\}) = o(n\tau(1-\tau)\mu_{\mathcal{W}})$ for all $\tau \in \mathcal{T}$ and $\varpi \in \mathcal{W}$, and the bound on λ_u follows from the union bound.

Let $\delta_u = \hat{\beta}_u - \beta_u$. By triangle inequality it follows that

$$\{E[K_\varpi(W)f_u(Z^a\delta_u)^2]\}^{1/2} \leq \|\sqrt{f_u}Z^a\delta_u\|_{n, \varpi} + \|\delta_u\|_1\{(|\mathbb{E}_n - \mathbb{E}|[K_\varpi(W)f_u(Z^a\delta_u)^2]/\|\delta_u\|_1^2)\}^{1/2} \quad (\text{C.37})$$

and the last term can be bounded by

$$\begin{aligned} \sup_{\|\delta\|_1 \leq 1} |(\mathbb{E}_n - \mathbb{E})[K_\varpi(W)f_u(Z^a\delta)^2]/\|\delta\|_1^2| &\leq \max_{k,j} |(\mathbb{E}_n - \mathbb{E})[K_\varpi(W)f_u Z_k^a Z_j^a]| \\ &\lesssim \sqrt{\frac{(1+d_W)\log(p|V|n)}{n}} \end{aligned}$$

with probability $1 - o(1)$ by Lemma 19 under our conditions.

Combining the relations above with (C.37), under $(1 + d_W)s^2 \log(p|V|n) = o(n)$ we have uniformly over $u \in \mathcal{U}$

$$\begin{aligned} \|\delta_u\| &\lesssim \{E[(Z^a\delta_u)^2 | \varpi]\}^{1/2} \lesssim \{P(\varpi)\}^{-1/2} \{E[K_\varpi(W)(Z^a\delta_u)^2]\}^{1/2} \\ &\lesssim \{P(\varpi)\underline{f}_u\}^{-1/2} \{E[K_\varpi(W)f_u(Z^a\delta_u)^2]\}^{1/2} \\ &\leq \{P(\varpi)\underline{f}_u\}^{-1/2} \|\sqrt{f_u}Z^a\delta_u\|_{n, \varpi} + \{P(\varpi)\underline{f}_u\}^{-1/2} \sqrt{\frac{(1+d_W)\log(p|V|n)}{n}} \|\delta_u\|_1 \\ &\leq C\sqrt{\frac{s(1+d_W)\log(p|V|n)}{n\underline{f}_u P(\varpi)}}. \end{aligned}$$

given $\|\delta_u\|_1 \leq \|\delta_u\|_{1, \varpi}/P(\varpi)^{1/2}$, (C.36), and $s^2(1 + d_W)\log(p|V|n) = o(n\mu_{\mathcal{W}}^4 \underline{f}_{\mathcal{U}}^2)$ assumed in Condition CI.

Finally, let $\hat{\beta}_u^\lambda$ obtained by thresholding the estimator $\hat{\beta}_u$ with $\bar{\lambda} := \sqrt{(1 + d_W)\log(p|V|n)/n}$ (note that each component is weighted by $\mathbb{E}_n[K_\varpi(W)(Z_j^a)^2]^{1/2}$). By Lemma 17, we have with probability $1 - o(1)$

$$\begin{aligned} \|Z^a(\hat{\beta}_u^\lambda - \beta_u)\|_{n, \varpi} &\lesssim \sqrt{s(1 + d_W)\log(p|V|n)/n} \\ \|\hat{\beta}_u^\lambda - \beta_u\|_{1, \varpi} &\lesssim s\sqrt{(1 + d_W)\log(p|V|n)/n} \\ &\quad |\text{support}(\hat{\beta}_u^\lambda)| \lesssim s \end{aligned}$$

by the choice of $\bar{\lambda}$ and the rates in (C.36)

■

Proof of Theorem 2. We verify Assumption C4 and Condition WL for the weighted Lasso model with index set $\mathcal{U} \times [p]$ where $Y_u = K_\varpi(W)Z_j^a$, $X_u = K_\varpi(W)Z_{-j}^a$, $\theta_u = \bar{\gamma}_u^j$, $a_u = (f_u, \bar{r}_{uj})$, $\bar{r}_{uj} = K_\varpi(W)Z_{-j}^a(\gamma_u^j - \bar{\gamma}_u^j)$, $S_{uj} = K_\varpi(W)f_u^2(Z_j^a - Z_{-j}^a\gamma_u^j)Z_{-j}^a = K_\varpi(W)f_u v_{uj}Z_{-j}^a$, and $w_u = K_\varpi(W)f_u^2$. We will take $N_n = |V|p^2\{pn^3\}^{1+d_W}$ in the definition of λ .

We first verify Condition WL. We have $E[S_{ujk}^2] \leq \bar{f}^2 E[|v_{uj}Z_{-j,k}^a|^2] \leq \bar{f}^2 \{E[|v_{uj}|^4 | Z_{-j,k}^a|^4]\}^{1/2} \leq C$ by the bounded fourth moment condition. We have that

$$\frac{E[|S_{ujk}|^3]^{1/3}}{E[|S_{ujk}|^2]^{1/2}} = \frac{E[|S_{ujk}|^3 | \varpi]^{1/3}}{E[|S_{ujk}|^2 | \varpi]^{1/2}} \{P(\varpi)\}^{-1/6} = \frac{E[|f_u v_{uj} Z_{-j,k}^a|^3 | \varpi]^{1/3}}{E[|f_u v_{uj} Z_{-j,k}^a|^2 | \varpi]^{1/2}} \{P(\varpi)\}^{-1/6} =: M_{uk}$$

By the choice of N_n and $\Phi^{-1}(1-t) \leq C\sqrt{\log(1/t)}$, we have $M_{uk}\Phi^{-1}(1-\xi/\{2pN_n\}) \leq M_{uk}C(1+d_W)\log^{1/2}(pn|V|) \leq C\delta_n n^{1/6}$ where the last inequality holds by Condition CI so Condition WL(i) holds.

To verify Condition WL(ii) we will establish the validity of the choice of N_n . We will consider $u = (a, \tau, \varpi) \in \mathcal{U}$ and $u' = (a, \tau', \varpi') \in \mathcal{U}$. By Condition CI we have that

$$|f_u - f_{u'}| \leq L_f \|u - u'\| \quad \text{and} \quad E[|K_\varpi(W) - K_{\varpi'}(W)|] \leq L_K \|\varpi - \varpi'\|. \quad (\text{C.38})$$

Further, by Lemma 5 we have

$$\|\gamma_u^j - \gamma_{u'}^j\| \leq L_\gamma \{\|u - u'\| + \|\varpi - \varpi'\|^{1/2}\}. \quad (\text{C.39})$$

By definition we have

$$S_{ujk} - S_{u'jk} = \{K_\varpi(W)f_u^2 - K_{\varpi'}(W)f_{u'}^2\} \{Z_j^a - Z_{-j}^a\gamma_u^j\} Z_k^a - K_{\varpi'}(W)f_{u'}^2 \{Z_j^a - Z_{-j}^a\gamma_{u'}^j\} Z_k^a$$

and note that $f_u + f_{u'} \leq 2f_u + L\|u - u'\|$, $|Z_j - Z_{-j}^a\gamma_u^j| \cdot |Z_k^a| \leq |Z_j|^2 + 2|Z_k^a|^2 + |Z_{-j}^a\gamma_u^j|^2$. Moreover,

$$\begin{aligned} |K_\varpi(W)f_u^2 - K_{\varpi'}(W)f_{u'}^2| &\leq K_\varpi(W)K_{\varpi'}(W)|f_u^2 - f_{u'}^2| + (f_u + f_{u'})^2 |K_\varpi(W) - K_{\varpi'}(W)| \\ &\leq 2\bar{f}K_\varpi(W)K_{\varpi'}(W)|f_u - f_{u'}| + 4\bar{f}^2 |K_\varpi(W) - K_{\varpi'}(W)| \end{aligned}$$

Using these relations we have $|\mathbb{E}_n[S_{ujk} - S_{u'jk}]| \leq (I) + (II)$ where

$$\begin{aligned} (I) &= \mathbb{E}_n[|K_\varpi(W)f_u^2 - K_{\varpi'}(W)f_{u'}^2| \cdot |\{Z_j^a - Z_{-j}^a\gamma_u^j\} Z_k^a|] \\ &\leq \max_{i \leq n} \|Z_i\|_\infty^2 (1 + \|\gamma_u^j\|_1) \mathbb{E}_n[2\bar{f}K_\varpi(W)K_{\varpi'}(W)|f_u - f_{u'}| + 4\bar{f}^2 |K_\varpi(W) - K_{\varpi'}(W)|] \\ &\leq (\bar{f} + \bar{f}^2)C\sqrt{s} \max_{i \leq n} \|Z_i\|_\infty^2 \{L_f\|u - u'\| + \mathbb{E}_n[|K_\varpi(W) - K_{\varpi'}(W)|]\} \\ (II) &= \mathbb{E}_n[K_{\varpi'}(W)f_{u'}^2 |Z_{-j}^a(\gamma_u^j - \gamma_{u'}^j)Z_k^a|] \\ &\leq \bar{f}^2 \mathbb{E}_n[\|Z^a\|_\infty^2] \|\gamma_u^j - \gamma_{u'}^j\|_1 \\ &\leq \bar{f}^2 \mathbb{E}_n[\|Z^a\|_\infty^2] \sqrt{p} L_\gamma \{\|u - u'\| + \|\varpi - \varpi'\|^{1/2}\} \end{aligned}$$

Moreover, we have that $\max_{i \leq n} \|Z_i^a\|_\infty^2 \lesssim_P M_n^2$. For $d_{\mathcal{U}} = \|\cdot\|$, an uniform ϵ -cover of \mathcal{U} satisfies $(6\text{diam}(\mathcal{U})/\epsilon)^{1+d_W} \geq N(\epsilon, \mathcal{U}, \|\cdot\|)$. We will set $1/\epsilon = (1 + \bar{f}^2)\{L_\gamma + L_f\}^2 pn M_n^2 \log^2(p|V|n)/\{\mu_{\mathcal{W}} f_{\mathcal{U}}^2\} \leq pn^3$ so that with probability $1 - o(1)$, for any pair $u, u' \in \mathcal{U}$, $\|u - u'\| \leq \epsilon$, we have

$$\begin{aligned} |\mathbb{E}_n[S_{ujk} - S_{u'jk}]| &\lesssim (\bar{f} + \bar{f}^2)\sqrt{s} \max_{i \leq n} \|Z_i\|_\infty^2 \{L_f\epsilon + \mathbb{E}_n[|K_\varpi(W) - K_{\varpi'}(W)|]\} \\ &\quad + \bar{f}^2 \mathbb{E}_n[\|Z_i^a\|_\infty^2] \sqrt{p} L_\gamma \{\epsilon + \epsilon^{1/2}\} \\ &\lesssim \delta_n n^{-1/2} \{\mu_{\mathcal{W}} f_{\mathcal{U}}^2\}^{1/2} + \sqrt{s} M_n \log(n) \mathbb{E}_n[|K_\varpi(W) - K_{\varpi'}(W)|] \end{aligned}$$

by the choice of ϵ . To control the last term, note that \mathcal{W} is a VC-class of events with VC dimension d_W . Thus by Lemma 19, with probability $1 - o(1)$

$$\begin{aligned} \mathbb{E}_n[|K_\varpi(W) - K_{\varpi'}(W)|] &\leq |(\mathbb{E}_n - \mathbb{E})[|K_\varpi(W) - K_{\varpi'}(W)|]| + \mathbb{E}[|K_\varpi(W) - K_{\varpi'}(W)|] \\ &\leq \sup_{\varpi, \varpi' \in \mathcal{W}, \|\varpi - \varpi'\| \leq \epsilon} |(\mathbb{E}_n - \mathbb{E})[|K_\varpi(W) - K_{\varpi'}(W)|]| + L_K \epsilon \\ &\lesssim \sqrt{\frac{d_W \log(n/\epsilon)}{n}} \epsilon^{1/2} + \frac{d_W \log(n/\epsilon)}{n} + L_K \epsilon \end{aligned}$$

which yields uniformly over $u \in \mathcal{U}$ and $j \in [p]$

$$|\mathbb{E}_n[S_{ujk} - S_{u'jk}]| \lesssim \delta_n n^{-1/2} \mu_{\mathcal{W}}^{1/2} \underline{f}_{\mathcal{U}} \quad (\text{C.40})$$

under $\sqrt{\epsilon d_W \log(n/\epsilon)} M_n \log n = o(\mu_{\mathcal{W}}^{1/2} \underline{f}_{\mathcal{U}})$ and $d_W \log(n/\epsilon) M_n \log n = o(n^{1/2} \mu_{\mathcal{W}}^{1/2} \underline{f}_{\mathcal{U}})$ assumed in Condition CI. In turn this implies

$$\sup_{|u-u'| \leq \epsilon} \max_{j, k \in [p], j \neq k} \frac{|\mathbb{E}_n[S_{ujk} - S_{u'jk}]|}{\mathbb{E}[S_{ujk}^2]^{1/2}} \leq \delta_n n^{-1/2}$$

since $\mathbb{E}[S_{ujk}^2] \geq c \mu_{\mathcal{W}} \underline{f}_{\mathcal{U}}^2$. Using the same choice of ϵ , similar arguments also imply

$$\sup_{|u-u'| \leq \epsilon} \max_{j, k \in [p], j \neq k} \frac{|\mathbb{E}[S_{ujk}^2 - S_{u'jk}^2]|}{\mathbb{E}[S_{ujk}^2]} \leq \delta_n \quad (\text{C.41})$$

To establish the last requirement of Condition WL(ii), note that

$$\sup_{u \in \mathcal{U}} \max_{j, k \in [p], j \neq k} |(\mathbb{E}_n - \mathbb{E})[S_{ujk}^2]| \leq \sup_{u \in \mathcal{U}^\epsilon} \max_{j, k \in [p], j \neq k} |(\mathbb{E}_n - \mathbb{E})[S_{ujk}^2]| + \Delta_n \quad (\text{C.42})$$

where $\Delta_n := \sup_{u, u' \in \mathcal{U}, \|u-u'\| \leq \epsilon} \max_{j, k \in [p], j \neq k} |(\mathbb{E}_n - \mathbb{E})[S_{ujk}^2] - (\mathbb{E}_n - \mathbb{E})[S_{u'jk}^2]|$.

To bound the first term, we will apply Corollary 2 with $k = 1$, $\hat{\mathcal{U}} := \mathcal{U}^\epsilon \times [p]$ and the vector $\{(\bar{X})_{uj} = S_{uj}, (u, j) \in \hat{\mathcal{U}}\}$. In this case note that

$$K^2 = \mathbb{E}[\max_{i \leq n} \sup_{u \in \mathcal{U}} \max_{j, k \in [p], j \neq k} S_{ujk}^2] \leq \mathbb{E}[\max_{i \leq n} \sup_{u \in \mathcal{U}, j \in [p]} |v_{iu}|^2 \|f_{iu} Z_i^a\|_\infty^2] \leq \bar{f}^2 M_n^2 L_n^2.$$

Therefore, by Corollary 2 and Markov inequality, we have with probability $1 - o(1)$ that

$$\sup_{u \in \mathcal{U}} \max_{j, k \in [p], j \neq k} |(\mathbb{E}_n - \mathbb{E})[S_{ujk}^2]| \leq C n^{-1/2} M_n L_n \log^{1/2}(p|V|n) \leq C \delta_n \mu_{\mathcal{W}} \underline{f}_{\mathcal{U}} + \Delta_n$$

under $M_n^2 L_n^2 \log(p|V|n) \leq \delta_n n \mu_{\mathcal{W}}^2 \underline{f}_{\mathcal{U}}^2$.

To control Δ_n , note that

$$\begin{aligned} |(\mathbb{E}_n - \mathbb{E})[S_{ujk}^2] - (\mathbb{E}_n - \mathbb{E})[S_{u'jk}^2]| &\leq |\mathbb{E}_n[S_{ujk}^2 - S_{u'jk}^2]| + |\mathbb{E}[S_{ujk}^2 - S_{u'jk}^2]| \\ &\leq \mathbb{E}_n[|S_{ujk} - S_{u'jk}|] \sup_{u \in \mathcal{U}} \max_{i \leq n} |2S_{iu'jk}| + |\mathbb{E}[S_{ujk}^2 - S_{u'jk}^2]| \\ &\lesssim \delta_n n^{-1/2} \mu_{\mathcal{W}}^{1/2} \underline{f}_{\mathcal{U}} \bar{f} \sup_{u \in \mathcal{U}} \max_{i \leq n} |v_{iu}| \|Z_i^a\|_\infty + \delta_n \mu_{\mathcal{W}} \underline{f}_{\mathcal{U}}^2 \\ &\lesssim \delta_n n^{-1/2} \mu_{\mathcal{W}}^{1/2} \underline{f}_{\mathcal{U}} M_n L_n \log n + \delta_n \mu_{\mathcal{W}} \underline{f}_{\mathcal{U}}^2 \end{aligned}$$

with probability $1 - o(1)$ where we used (C.40) and (C.41). Therefore $\Delta_n \lesssim \delta_n \mu_{\mathcal{W}} \underline{f}_{\mathcal{U}}^2$ with probability $1 - o(1)$ as required.

To verify Assumption C4(a), note that $[\partial_\theta M_u(Y_u, X, \theta_u) - \partial_\theta M_u(Y_u, X, \theta_u, a_u)]' \delta = -f_u^2 K_\varpi(W) \bar{r}_{uj} Z_{-j}^a \delta$, so that by Cauchy-Schwartz, we have

$$\mathbb{E}_n[\partial_\theta M_u(Y_u, X, \theta_u) - \partial_\theta M_u(Y_u, X, \theta_u, a_u)]' \delta \leq \|f_u \bar{r}_{uj}\|_{n, \varpi} \|f_u Z_{-j}^a \delta\|_{n, \varpi} \leq C_{un} \|f_u Z_{-j}^a \delta\|_{n, \varpi}$$

where we choose C_{un} so that $\{C_{un} \geq \max_{j \in [p]} \|f_u \bar{r}_{uj}\|_{n,\varpi} : u \in \mathcal{U}\}$ with probability $1 - o(1)$. To bound C_{un} , by Lemma 4, uniformly over $u \in \mathcal{U}, j \in [p]$ we have with probability $1 - o(1)$

$$\|f_u \bar{r}_{uj}\|_{n,\varpi} = \|f_u Z_{-j}^a (\gamma_u^j - \hat{\gamma}_u^j)\|_{n,\varpi} \lesssim \underline{f}_u \{P(\varpi)\}^{1/2} \{n^{-1} s \log(p|V|n)\}^{1/2}$$

so that setting $C_{un} = \underline{f}_u \{P(\varpi)\}^{1/2} \{n^{-1} s \log(p|V|n)\}^{1/2}$ suffices.

Next we show that Assumption C4(b) holds. First, by (C.42) and the corresponding bounds, note the uniform convergence of the loadings

$$\sup_{u \in \mathcal{U}, j, k \in [p], j \neq k} (|\mathbb{E}_n[S_{ujk}^2] - \mathbb{E}[S_{ujk}^2]| + |(\mathbb{E}_n - \mathbb{E})[K_\varpi(W) f_u^2 |Z_j^a Z_{-jk}^a|^2]|) \leq \delta_n \mu_{\mathcal{W}} \underline{f}_{\mathcal{U}}^2$$

so that $\mathbb{E}_n[S_{ujk}^2]/\mathbb{E}[S_{ujk}^2] = 1 + o_P(1)$. It follows that \tilde{c} is bounded above by a constant for n large enough. Indeed, uniformly over $u \in \mathcal{U}, j \in [p]$, since $c \underline{f}_u \leq \mathbb{E}[|f_u v_{uj} Z_k^a|^2 | \varpi]^{1/2} \leq C \underline{f}_u$, with probability $1 - o(1)$ we have $c \underline{f}_u P(\varpi)^{1/2} \leq \hat{\Psi}_{u0jj} \leq C \underline{f}_u P(\varpi)^{1/2}$ so that $c/C \leq \|\hat{\Psi}_{u0}\|_\infty \|\hat{\Psi}_{u0}^{-1}\|_\infty \leq C/c$.

Assumption C4(c) follows directly from the choice of $M_u(Y_u, X_u, \theta) = K_\varpi(W) f_u^2 (Z_j^a - Z_{-j}^a \theta)^2$ with $\bar{q}_{A_u} = \infty$.

The result for the rate of convergence then follows from Lemma 22, namely

$$\|f_u X'_u (\hat{\gamma}_u^j - \gamma_u^j)\|_{n,\varpi} \lesssim \frac{\|\hat{\Psi}_{u0}\|_\infty}{\bar{\kappa}_{u,2\mathbf{c}}} \sqrt{\frac{s \log(p|V|n)}{n}} + C_{un} \lesssim \frac{\underline{f}_u P(\varpi)^{1/2}}{\bar{\kappa}_{u,2\mathbf{c}}} \sqrt{\frac{s \log(p|V|n)}{n}} \quad (\text{C.43})$$

By Lemma 6 we have that for sparse vectors, $\|\theta\|_0 \leq \ell_n s$ satisfies

$$\|f_u Z_{-j}^a \theta\|_{n,\varpi}^2 / \mathbb{E}[K_\varpi(W) f_u^2 (Z_{-j}^a \theta)^2] = 1 + o_P(1)$$

so that $\phi_{\max}(\ell_n s, u_j) \leq C \underline{f}_u^2 P(\varpi)$ and $\hat{s}_{uj} \leq \min_{m \in \mathcal{M}_u} \phi_{\max}(m, u_j) L_u^2 \leq C s$ provided $L_u^2 \lesssim s \{ \underline{f}_u^2 P(\varpi) \}^{-1}$. Indeed, with probability $1 - o(1)$, we have $\|\hat{\Psi}_{u0}^{-1}\|_\infty \leq C \underline{f}_u^{-1} P(\varpi)^{-1/2}$, so that $L_u \lesssim \underline{f}_u^{-1} P(\varpi)^{-1/2} \frac{n}{\lambda} \{C_{un} + L_{un}\}$. Moreover, we can take $C_{un} \lesssim \underline{f}_u \{P(\varpi) n^{-1} s \log(p|V|n)\}^{1/2}$, and $L_{un} \lesssim \{n^{-1} s \log(p|V|n)\}^{1/2}$ in Assumption C4 because

$$\begin{aligned} & |\{\mathbb{E}_n[\partial_\gamma M_u(Y_u, X_u, \hat{\gamma}_u^j) - \partial_\gamma M_u(Y_u, X_u, \gamma_u^j)]\}' \delta| \\ &= 2 |\mathbb{E}_n[K_\varpi(W) f_u^2 \{X'_u (\hat{\gamma}_u^j - \gamma_u^j)\} X'_u \delta]| \\ &\leq 2 \|f_u X'_u (\hat{\gamma}_u^j - \gamma_u^j)\|_{n,\varpi} \|f_u X'_u \delta\|_{n,\varpi} =: L_{un} \|f_u X'_u \delta\|_{n,\varpi}, \end{aligned}$$

where the last inequality hold by (C.43) since $\bar{\kappa}_{u,2\mathbf{c}} \geq c \underline{f}_u \{P(\varpi)\}^{1/2}$. The bound on the restricted eigenvalue $\bar{\kappa}_{u,2\mathbf{c}}$ holds¹⁵ by arguments similar to (C.45) and using that $\|\delta\|_1 \leq C \sqrt{s} \|\delta\|$ for any $\delta \in \Delta_{u,2\mathbf{c}}$, and since for any $\|\delta\| = 1$, we have

$$\begin{aligned} c \underline{f}_u P(\varpi) &\leq \mathbb{E}[K_\varpi(W) f_u (Z^a \delta)^2] \\ &\leq \{\mathbb{E}[K_\varpi(W) f_u^2 (Z^a \delta)^2]\}^{1/2} \{\mathbb{E}[K_\varpi(W) (Z^a \delta)^2]\}^{1/2} \\ &\leq \{\mathbb{E}[K_\varpi(W) f_u^2 (Z^a \delta)^2]\}^{1/2} C \{P(\varpi)\}^{1/2} \end{aligned}$$

where the first inequality follows from the definition of \underline{f}_u , $\|\delta\| = 1$, and Condition CI, so that we have $\{\mathbb{E}[K_\varpi(W) f_u^2 (Z^a \delta)^2]\}^{1/2} \geq c' \underline{f}_u \{P(\varpi)\}^{1/2}$.

¹⁵Note that there are two restricted eigenvalues definitions, one used for the quantile regression $(\kappa_{u,2\mathbf{c}})$, and another used here for the weighted lasso $(\bar{\kappa}_{u,2\mathbf{c}})$. It is a consequence of the use of different norms.

Return to the rate of convergence we have by (C.43) and $\bar{\kappa}_{u,2c} \geq c \underline{f}_u \{P(\varpi)\}^{1/2}$ that

$$\|f_u X'_u (\hat{\gamma}_u^j - \gamma_u^j)\|_{n,\varpi} \lesssim \frac{\underline{f}_u P(\varpi)^{1/2}}{\bar{\kappa}_{u,2c}} \sqrt{\frac{s \log(p|V|n)}{n}} \lesssim \sqrt{\frac{s \log(p|V|n)}{n}} \quad (\text{C.44})$$

and the result follows by noting that $\|f_u X'_u (\hat{\gamma}_u^j - \gamma_u^j)\|_{n,\varpi} \geq c \underline{f}_u P(\varpi)^{1/2} \|\hat{\gamma}_u^j - \gamma_u^j\|$ with probability $1 - o(1)$ by arguments similar to (C.45) under Condition CI.

The sparsity result follows from Lemma 21. The result for Post Lasso follows from Lemma 20 under the growth requirements in Condition CI. ■

Proof of Theorem 3. We will verify Assumptions C1 and C2, and the result follows from Theorem 5. The estimate of the nuisance parameter is constructed from the estimators in Steps 1 and 2 of the Algorithm.

For each $u = (a, \tau, \varpi) \in \mathcal{U}$ and $j \in [p]$, let $W_{uj} = (W, X_a, Z^a, v_{uj}, r_u)$, where $v_{uj} = f_u(Z_j^a - Z_{-j}^a \gamma_u^j)$ and let $\theta_{uj} \in \Theta_{uj} = \{\theta \in \mathbb{R} : |\theta - \beta_{uj}| \leq c/\log n\}$ (Assumption C1(i) holds). The score function is

$$\psi_{uj}(W_{uj}, \theta, \eta_{uj}) = K_\varpi(W) \{\tau - 1\{X_a \leq Z_j^a \theta + Z_{-j}^a \beta_{u,-j} + r_u\}\} f_u(Z_j^a - Z_{-j}^a \gamma_u^j)$$

where the nuisance parameter is $\eta_{uj} = (\beta_{u,-j}, \gamma_u^j, r_u)$ and the last component is a function $r_u = r_u(X)$. Recall that $K_\varpi(W) \in \{0, 1\}$ and let $a_n = \max(n, p, |V|)$. Define the nuisance parameter set $\mathcal{H}_{uj} = \{\eta = (\eta^{(1)}, \eta^{(2)}, \eta^{(3)}) : \|\eta - \eta_{uj}\|_e \leq \tau_n\}$ where $\|\eta - \eta_{uj}\|_e = \|(\delta_\eta^{(1)}, \delta_\eta^{(2)}, \delta_\eta^{(3)})\|_e = \max\{\|\delta_\eta^{(1)}\|, \|\delta_\eta^{(2)}\|, E[|\delta_\eta^{(3)}|^2]^{1/2}\}$, and

$$\tau_n := C \sup_{u \in \mathcal{U}} \frac{1}{1 \wedge \underline{f}_u} \sqrt{\frac{s \log a_n}{n \mu_W}}$$

The differentiability of the mapping $(\theta, \eta) \in \Theta_{uj} \times \mathcal{H}_{uj} \mapsto E\psi_{uj}(W_{uj}, \theta, \eta)$ follows from the differentiability of the conditional probability distribution of X_a given $X_{V \setminus \{a\}}$ and ϖ . Let $\eta = (\eta^{(1)}, \eta^{(2)}, \eta^{(3)})$, $\delta_\eta = (\delta_\eta^{(1)}, \delta_\eta^{(2)}, \delta_\eta^{(3)})$, and $\theta_{\bar{r}} = \theta + \bar{r} \delta_\theta$, $\eta_{\bar{r}} = \eta + \bar{r} \delta_\eta$.

To verify Assumption C1(v)(a) with $\alpha = 2$, for any $(\theta, \eta), (\bar{\theta}, \bar{\eta}) \in \Theta_{uj} \times \mathcal{H}_{uj}$ note that $f_{X_a|X_{-a}, \varpi}$ is uniformly bounded from above by \bar{f} , therefore

$$\begin{aligned} & E[\{\psi_{uj}(W_{uj}, \theta, \eta) - \psi_{uj}(W_{uj}, \bar{\theta}, \bar{\eta})\}^2]^{1/2} \\ & \leq \bar{f} E[|Z_{-j}^a (\eta^{(2)} - \bar{\eta}^{(2)})|^2]^{1/2} + \bar{f}^2 E[(Z_j^a - Z_{-j}^a \bar{\eta}^{(2)})^2 \{|\eta^{(3)} - \bar{\eta}^{(3)}| + |Z_{-j}^a (\eta^{(1)} - \bar{\eta}^{(1)})| + |Z_j^a (\theta - \bar{\theta})|\}]^{1/2} \\ & \leq C \|\eta^{(2)} - \bar{\eta}^{(2)}\| + \bar{f} E[(Z_j^a - Z_{-j}^a \bar{\eta}^{(2)})^4]^{1/4} \{E[|\eta^{(3)} - \bar{\eta}^{(3)}|^2]^{1/4} + C \|\eta^{(1)} - \bar{\eta}^{(1)}\| + |\theta - \bar{\theta}|\}^{1/2} \\ & \leq C' |\theta - \bar{\theta}|^{1/2} \vee \|\eta - \bar{\eta}\|_e^{1/2} \end{aligned}$$

for some constance $C' < \infty$ since by Condition CI we have $E[|Z^a \bar{\xi}|^2]^{1/2} \leq C \|\bar{\xi}\|$ for all vectors $\bar{\xi}$, and the conditions $\sup_{u \in \mathcal{U}, j \in [p]} \|\gamma_u^j\| \leq C$, $\sup_{\theta \in \Theta_{uj}} |\theta| \leq C$, and $\sqrt{s \log(a_n)} \leq \delta_n \sqrt{n}$. This implies that $\|\eta^{(2)} - \bar{\eta}^{(2)}\| \leq \|\eta^{(2)} - \eta_{uj}^{(2)}\| + \|\eta_{uj}^{(2)} - \bar{\eta}^{(2)}\| \leq 1$ so that $\|\eta^{(2)} - \bar{\eta}^{(2)}\| \leq \|\eta^{(2)} - \bar{\eta}^{(2)}\|^{1/2}$.

To verify Assumption C1(v)(b), let $t_{\bar{r}} = Z_j^a \theta_{\bar{r}} + Z_{-j}^a \eta_{\bar{r}}^{(1)} + \eta_{\bar{r}}^{(3)}$. We have

$$\begin{aligned} & \partial_r E(\psi_{uj}(W_{uj}, \theta + r \delta_\theta, \eta + r \delta_\eta))|_{r=\bar{r}} = \\ & -E[K_\varpi(W) f_{X_a|X_{-a}, \varpi}(t_{\bar{r}}) (Z_j^a - Z_{-j}^a \eta_{\bar{r}}^{(2)}) \{Z_j^a \delta_\theta + Z_{-j}^a \delta_\eta^{(1)} + \delta_\eta^{(3)}\}] \\ & -E[K_\varpi(W) \{\tau - F_{X_a|X_{-a}, \varpi}(t_{\bar{r}})\} Z_{-j}^a \delta_\eta^{(2)}] \end{aligned}$$

Applying Cauchy-Schwartz we have that

$$\begin{aligned} & \left| \partial_r \mathbb{E}(\psi_{uj}(W_{uj}, \theta + r\delta_\theta, \eta + r\delta_\eta)) \Big|_{r=\bar{r}} \right| \\ & \leq \bar{f} \mathbb{E}[(Z_j^a - Z_{-j}^a \eta_{\bar{r}}^{(2)})^2]^{1/2} \{ \mathbb{E}[(Z_j^a)^2]^{1/2} |\delta_\theta| + \mathbb{E}[(Z_{-j}^a \delta_\eta^{(1)})^2]^{1/2} + \mathbb{E}[|\delta_\eta^{(3)}|^2]^{1/2} \} + \bar{f} \mathbb{E}[(Z_{-j}^a \delta_\eta^{(2)})^2]^{1/2} \\ & \leq \bar{B}_{1n} (|\delta_\theta| \vee \|\eta - \eta_{uj}\|_e) \end{aligned}$$

where $\bar{B}_{1n} \leq C$ by the same arguments of bounded (second) moments of linear combinations.

Assumption C1(v)(c) follows similarly as

$$\begin{aligned} & \partial_r^2 \mathbb{E}(\psi_{uj}(W_{uj}, \theta + r\delta_\theta, \eta + r\delta_\eta)) \Big|_{r=\bar{r}} = \\ & -\mathbb{E}[K_\varpi(W) f'_{X_a|X_{-a}, \varpi}(t_{\bar{r}})(Z_j^a - Z_{-j}^a \eta_{\bar{r}}^{(2)}) \{Z_j^a \delta_\theta + Z_{-j}^a \delta_\eta^{(1)} + \delta_\eta^{(3)}\}^2] \\ & + 2\mathbb{E}[K_\varpi(W) f_{X_a|X_{-a}, \varpi}(t_{\bar{r}})(Z_{-j}^a \delta_\eta^{(2)}) \{Z_j^a \delta_\theta + Z_{-j}^a \delta_\eta^{(1)} + \delta_\eta^{(3)}\}] \end{aligned}$$

and under $|f'_{X_a|X_{-a}, \varpi}| \leq \bar{f}'$, from Cauchy-Schwartz inequality we have

$$\begin{aligned} & \left| \partial_r^2 \mathbb{E}(\psi_{uj}(W_{uj}, \theta + r\delta_\theta, \eta + r\delta_\eta)) \Big|_{r=\bar{r}} \right| \\ & \leq |\bar{f}'_n \mathbb{E}[(Z_j^a - Z_{-j}^a \eta_{\bar{r}}^{(2)})^2]^{1/2} \{ \mathbb{E}[(Z_j^a)^4] |\delta_\theta|^2 + \mathbb{E}[(Z_{-j}^a \delta_\eta^{(1)})^4]^{1/2} \} + C \mathbb{E}[\{\delta_\eta^{(3)}\}^2] \\ & + 2\bar{f}' \mathbb{E}[(Z_{-j}^a \delta_\eta^{(2)})^2]^{1/2} \{ \mathbb{E}[(Z_j^a)^2]^{1/2} |\delta_\theta| + \mathbb{E}[(Z_{-j}^a \delta_\eta^{(1)})^2]^{1/2} + \mathbb{E}[\{\delta_\eta^{(3)}\}^2]^{1/2} \} \\ & \leq \bar{B}_{2n} (\delta_\theta^2 \vee \|\eta - \eta_{uj}\|_e^2) \end{aligned}$$

where $\bar{B}_{2n} \leq C(1 + \bar{f}'_n)$ by the same arguments of bounded (fourth) moments as before and using that $|\mathbb{E}[(Z_j^a - Z_{-j}^a \eta_{\bar{r}}^{(2)})(\delta_\eta^{(3)})^2]| \leq \{ \mathbb{E}[(Z_j^a - Z_{-j}^a \eta_{\bar{r}}^{(2)})^2 (\delta_\eta^{(3)})^2] \}^{1/2} \mathbb{E}[(\delta_\eta^{(3)})^2]^{1/2} \leq C \mathbb{E}[(\delta_\eta^{(3)})^2]$.

To verify the near orthogonality condition, note that for all $u \in \mathcal{U}$ and $j \in [p]$, since by definition $f_u = f_{X_a|X_{-a}, \varpi}(Z^a \beta_u + r_u)$ we have

$$|\mathbb{D}_{u,j,0}[\tilde{\eta}_{uj} - \eta_{uj}]| = |-\mathbb{E}[K_\varpi(W) f_u \{Z_{-j}^a (\tilde{\eta}^{(2)} - \eta_{uj}^{(2)}) + r_u\} v_{uj}]| \leq \delta_n n^{-1/2}$$

by the relations $\mathbb{E}[K_\varpi(W)(\tau - F_{X_a|X_{-a}, \varpi}(Z^a \beta_u + r_u)) Z_{-j}^a] = 0$ and $\mathbb{E}[K_\varpi(W) f_u Z_{-j}^a v_{uj}] = 0$ implied by the model, and $|\mathbb{E}[K_\varpi(W) f_u r_u v_{uj}]| \leq \delta_n n^{-1/2}$ by Condition CI. Thus, condition (H.78) holds.

Furthermore, since $\Theta_{uj} \subset \theta_{uj} \pm C/\log n$, for $J_{uj} = \partial_\theta \mathbb{E}[\psi_{uj}(W_{uj}, \theta_{uj}, \eta_{uj})] = \mathbb{E}[K_\varpi(W) f_u Z_j^a v_{uj}] = \mathbb{E}[K_\varpi(W) v_{uj}^2] = \mathbb{E}[v_{uj}^2 | \varpi] \mathbb{P}(\varpi)$ as $\mathbb{E}[K_\varpi(W) f_u Z_{-j}^a v_{uj}] = 0$, we have that for all $\theta \in \Theta_{uj}$

$$\mathbb{E}[\psi_{uj}(W_{uj}, \theta, \eta_{uj})] = J_{uj}(\theta - \theta_{uj}) + \frac{1}{2} \partial_\theta^2 \mathbb{E}[\psi_{uj}(W_{uj}, \bar{\theta}, \eta_{uj})](\theta - \theta_{uj})^2$$

where $|\partial_\theta^2 \mathbb{E}[\psi_{uj}(W_{uj}, \bar{\theta}, \eta_{uj})]| \leq \bar{f}' \mathbb{E}[|Z_j^a|^2 | v_{uj}| | \varpi] \mathbb{P}(\varpi) \leq \bar{f}' \mathbb{E}[|Z_j^a|^4 | \varpi]^{1/2} \mathbb{E}[|v_{uj}|^2 | \varpi]^{1/2} \mathbb{P}(\varpi) \leq C \bar{f}' \mathbb{P}(\varpi)$ so that for all $\theta \in \Theta_{uj}$

$$|\mathbb{E}[\psi_{uj}(W_{uj}, \theta, \eta_{uj})]| \geq \{ \mathbb{E}[v_{uj}^2 | \varpi] - (C^2 \bar{f}') / \log n \} \mathbb{P}(\varpi) |\theta - \theta_{uj}|$$

and we can take $j_n \geq c \inf_{\varpi \in \mathcal{W}} \mathbb{P}(\varpi) = c \mu_{\mathcal{W}}$.

Next we verify Assumption C2 with $\mathcal{H}_{ujn} = \{\eta = (\beta, \gamma, 0) : \|\beta\|_0 \leq Cs, \|\gamma\|_0 \leq Cs, \|\beta - \beta_{u,-j}\| \leq C\tau_n, \|\gamma - \gamma_u^j\| \leq C\tau_n, \|\gamma - \gamma_u^j\|_1 \leq C\sqrt{s}\tau_n\}$. We will show that $\hat{\eta}_{uj} = (\tilde{\beta}_{u,-j}, \tilde{\gamma}_u^j, 0) \in \mathcal{H}_{ujn}$ with probability $1 - o(1)$, uniformly over $u \in \mathcal{U}$ and $j \in [p]$.

Under Condition CI and the choice of penalty parameters, by Theorems 1 and 2, with probability $1 - o(1)$, uniformly over $u \in \mathcal{U}$ we have

$$\|\tilde{\beta}_u - \beta_u\| \leq C\tau_n, \quad \max_{j \in [p]} \sup_{u \in \mathcal{U}} \|\tilde{\gamma}_u^j - \gamma_u^j\| \leq C\tau_n, \quad \text{and} \quad \max_{j \in [p]} \sup_{u \in \mathcal{U}} \|\tilde{\gamma}_u^j\|_0 \leq \bar{C}s,$$

further by thresholding we can achieve $\sup_{u \in \mathcal{U}} \|\tilde{\beta}_u\|_0 \leq \bar{C}s$ using Lemma 17.

Next we establish the entropy bounds. For $\eta \in \mathcal{H}_{ujn}$ we have that

$$\psi_{uj}(W_{uj}, \theta, \eta) = K_{\varpi}(W)(\tau - 1\{X_a \leq Z_j^a \theta + Z_{-j}^a \beta_{-j}\})f_u\{Z_j^a - Z_{-j}^a \gamma\}$$

It follows that $\mathcal{F}_1 \subset \mathcal{W}\mathcal{G}_1\mathcal{G}_2\mathcal{G}_3 \cup \bar{\mathcal{F}}_0$ where $\bar{\mathcal{F}}_0 = \{\psi_{uj}(W_{uj}, \theta, \eta_{uj}) : u \in \mathcal{U}, j \in [p], \theta \in \Theta_{uj}\}$, $\mathcal{G}_1 = \{\tau - 1\{X_a \leq Z^a \beta\} : \|\beta\|_0 \leq Cs, \tau \in \mathcal{T}, a \in V\}$, $\mathcal{G}_2 = \{Z^a \rightarrow Z^a(1, -\gamma), \|\gamma\|_0 \leq Cs, \|\gamma\| \leq C, a \in V\}$, $\mathcal{G}_3 = \{f_u : u \in \mathcal{U}\}$. Under Condition CI, \mathcal{W} is a VC class of sets with VC index d_W (fixed). It follows that \mathcal{G}_1 and \mathcal{G}_2 are p choose $O(s)$ VC-subgraph classes with VC indices at most $O(s)$. Therefore, $\text{ent}(\mathcal{G}_1) \vee \text{ent}(\mathcal{G}_2) \vee \text{ent}(\mathcal{W}) \leq Cs \log(a_n/\epsilon) + Cd_W \log(e/\epsilon)$ by Theorem 2.6.7 in [100] and by standard arguments. Also, since f_u is Lipschitz in u by Condition CI, we have $\text{ent}(\mathcal{G}_3) \leq (1 + d_W) \log(a_n L_f/\epsilon)$. Moreover, an envelope F_G for \mathcal{F}_1 satisfies

$$\begin{aligned} \mathbb{E}[F_G^q] &= \mathbb{E}[\sup_{u \in \mathcal{U}, j \in [p], \|\gamma - \gamma_u^j\|_1 \leq C\sqrt{s}\tau_n} |v_{uj} - f_u Z_{-j}^a(\gamma - \gamma_u^j)|^q] \\ &\leq 2^{q-1} \mathbb{E}[\sup_{u \in \mathcal{U}, j \in [p]} |v_{uj}|^q] + 2^{q-1} \bar{f} \mathbb{E}[\max_{a \in V} \|Z^a\|_{\infty}^q] \{C\sqrt{s}\tau_n\}^q \\ &\leq 2^{q-1} L_n^q + 2^{q-1} \bar{f} \{M_n C \sqrt{s}\tau_n\}^q \leq 2^q L_n^q \end{aligned}$$

since $M_n C \sqrt{s}\tau_n \leq \delta_n L_n / \bar{f}$ and $\delta_n \leq 1$ for n large.

Next we bound the entropy in $\bar{\mathcal{F}}_0$. Note that for any $\psi_{uj}(W_{uj}, \theta, \eta_{uj}) \in \bar{\mathcal{F}}_0$, there is some $\delta \in [-C, C]$ such that

$$\psi_{uj}(W_{uj}, \theta, \eta_{uj}) = K_{\varpi}(W)\{\tau - 1\{X_a \leq Z_j^a \delta + Q_{X_a}(\tau | X_{-a}, \varpi)\}\}v_{uj}$$

and therefore $\bar{\mathcal{F}}_0 \subset \mathcal{W}\{\mathcal{T} - \phi(\mathcal{V})\}\mathcal{L}$ where $\phi(t) = 1\{t \leq 0\}$, $\mathcal{V} = \cup_{a \in V, j \in [p]} \mathcal{V}_{aj}$ with

$$\mathcal{V}_{aj} := \{X_a - Z_j^a \delta - Q_{X_a}(\tau | X_{-a}, \varpi) : \tau \in \mathcal{T}, \varpi \in \mathcal{W}, |\delta| \leq C\},$$

and $\mathcal{L} = \cup_{a \in V, j \in [p]} (\mathcal{L}_{aj} + \{v_{\bar{u}j}\})$ where $\mathcal{L}_{aj} = \{(X, W) \mapsto v_{uj} - v_{\bar{u}j} = f_u Z_{-j}^a(\gamma_u^j - \gamma_{\bar{u}}^j) : u \in \mathcal{U}\}$. Note that each \mathcal{V}_{aj} is a VC subgraph class of functions with index $1 + Cd_W$ as $\{Q_{X_a}(\tau | X_{-a}, \varpi) : (\tau, \varpi) \in \mathcal{W} \times \mathcal{T}\}$ is a VC-subgraph with VC-dimension Cd_W for every $a \in V$. Since ϕ is monotone, $\phi(\mathcal{V})$ is also the union of VC-dimension of order $1 + Cd_W$.

Letting $F_1 = 1$ be an envelope for \mathcal{W} and $\mathcal{T} - \phi(\mathcal{V})$. By Lemma 5, it follows that $\|\gamma_u^j - \gamma_{u'}^j\| \leq L_{\gamma} \{\|u - u'\| + \|\varpi - \varpi'\|^{1/2}\}$ for some L_{γ} satisfying $\log(L_{\gamma}) \leq C \log(p|V|n)$ under Condition CI. Therefore, $|v_{uj} - v_{\bar{u}j}| = |Z_{-j}^a(\gamma_u^j - \gamma_{\bar{u}}^j)| \leq \|Z^a\|_{\infty} \sqrt{p} \|\gamma_u^j - \gamma_{\bar{u}}^j\|$. For a choice of envelope $F_a = M_n^{-1} \|Z^a\|_{\infty} + 2 \sup_{u \in \mathcal{U}} |v_{uj}|$ which satisfies $\|F_a\|_{P, q} \lesssim L_n$, we have

$$\begin{aligned} \log N(\epsilon \|F_a\|_{Q, 2}, \mathcal{L}_{aj}, \|\cdot\|_{Q, 2}) &\leq \log N(\frac{\epsilon}{M_n} \| \|Z^a\|_{\infty} \|_{Q, 2}, \mathcal{L}_{aj}, \|\cdot\|_{Q, 2}) \\ &\leq \log N(\epsilon / \{M_n \sqrt{p} L_{\gamma}\}, \mathcal{U}, \|\cdot\|) \leq Cd_u \log(M_n p L_{\gamma} / \epsilon) \end{aligned}$$

Since $\mathcal{L} = \cup_{a \in V, j \in [p]} (\mathcal{L}_{aj} + \{v_{\bar{u}j}\})$, taking $F_L = \max_{a \in V} F_a$, we have that

$$\begin{aligned} \log N(\epsilon \|F_L F_1\|_{Q, 2}, \bar{\mathcal{F}}_0, \|\cdot\|_{Q, 2}) &\leq \log N(\frac{\epsilon}{4} \|F_1\|_{Q, 2}, \mathcal{W}, \|\cdot\|_{Q, 2}) + \log N(\frac{\epsilon}{4} \|F_1\|_{Q, 2}, \mathcal{T} - \phi(\mathcal{V}), \|\cdot\|_{Q, 2}) \\ &\quad + \log \sum_{a \in V, j \in [p]} N(\frac{\epsilon}{2} \|F_a\|_{Q, 2}, \mathcal{L}_{aj}, \|\cdot\|_{Q, 2}) \\ &\leq \log(p|V|) + 1 + C' \{d_W + d_u\} \log(4e M_n |V| p L_{\gamma} / \epsilon) \end{aligned}$$

where the last line follows from the previous bounds.

Next we verify the growth conditions in Assumption C2 with the proposed \mathcal{F}_1 and $K_n \lesssim CL_n$. We take $s_{n(\mathcal{U}, p)} = (1 + d_W)s$ and $a_n = \max\{n, p, |V|\}$. Recall that $\bar{B}_{1n} \leq C$, $\bar{B}_{2n} \leq C$, $j_n \geq c\mu_{\mathcal{W}}$. Thus, we have $\sqrt{n}(\tau_n/j_n)^2 \lesssim \sqrt{n} \frac{s \log(p|V|n)}{n(1 \wedge \underline{f}_{\mathcal{U}}^2) \mu_{\mathcal{W}}^3} \leq \delta_n$ under $s^2 \log^2(p|V|n) \leq n(1 \wedge \underline{f}_{\mathcal{U}}^4) \mu_{\mathcal{W}}^6$. Moreover,

$(\tau_n/j_n)^{\alpha/2} \sqrt{s_{n(\mathcal{U},p)} \log(a_n)} \lesssim \sqrt[4]{\frac{(1+d_W)^3 s^3 \log^3(p|V|n)}{n(1+\underline{f}_{\mathcal{U}}^2) \mu_W^3}} \lesssim \delta_n$ under d_W fixed and $s^3 \log^3(p|V|n) \leq \delta_n^4 n(1 \wedge \underline{f}_{\mathcal{U}}^2) \mu_W^3$ and $s_{n(\mathcal{U},p)} n^{-\frac{1}{2}} K_n \log(a_n) \log n \lesssim (1+d_W) s n^{\frac{1}{q}-\frac{1}{2}} M_n \log(p|V|n) \log n \leq \delta_n$ under our conditions. Finally, the conditions of Corollary 4 hold with $\rho_n = (1+d_W)$ since the score is the product of VC-subgraph classes of function with VC index bounded by $C(1+d_W)$. ■

Proof of Theorem 4. We will invoke Lemma 7 with $\bar{\beta}_u$ as the estimand and $r_{iu} = X'_{i,-a}(\beta_u - \bar{\beta}_u)$, therefore $E[K_{\varpi}(W)(\tau - 1\{X_a \leq X'_{-a}\bar{\beta}_u + r_u\})X_{-a}] = 0$. To invoke the lemma we verify that the events Ω_1 , Ω_2 , Ω_3 and Ω_4 hold with probability $1 - o(1)$

$$\begin{aligned} \Omega_1 &:= \{\lambda_u \geq c|S_{uj}|/\hat{\sigma}_{a\varpi j}^X, \text{ for all } u \in \mathcal{U}, j \in V\}, \\ \Omega_2 &:= \{\hat{R}_u(\bar{\beta}_u) \leq \bar{R}_{u\xi} : u \in \mathcal{U}\} \\ \Omega_3 &:= \left\{ \sup_{u \in \mathcal{U}, 1/\sqrt{n} \leq \|\delta\|_{1,\varpi} \leq \sqrt{n}} |\mathbb{E}_n[g_u(\delta, X, W) - E[g_u(\delta, X, W) | X_{-a}, W]]|/\|\delta\|_{1,\varpi} \leq t_3 \right\} \\ \Omega_4 &:= \{K_u \hat{\sigma}_{a\varpi j}^X \geq |\mathbb{E}_n[h_{uj}(X_{-a}, W)]|, \text{ for all } u \in \mathcal{U}, j \in V \setminus \{a\}\} \end{aligned}$$

where $g_u(\delta, X, W) = K_{\varpi}(W)\{\rho_{\tau}(X_a - X'_{-a}(\bar{\beta}_u + \delta)) - \rho_{\tau}(X_a - X'_{-a}\bar{\beta}_u)\}$, $h_{uj}(X_{-a}, W) = E[K_{\varpi}(W)\{\tau - F_{X_a|X_{-a},W}(X'_{-a}\bar{\beta}_u + r_u)\}X_j | X_{-a}, W]$.

By Lemma 8 with $\xi = 1/n$, by setting $\lambda_u = \lambda_0 = c2(1+1/16)\sqrt{2\log(8|V|^2\{ne/d_W\}^{2d_W}n)/n}$, we have $P(\Omega_1) = 1 - o(1)$. By Lemma 9, setting $\bar{R}_{u\xi} = Cs(1+d_W)\log(|V|n)/n$ we have $P(\Omega_2) = 1 - o(1)$ for some $\xi = o(1)$. By Lemma 10 we have $P(\Omega_3) = 1 - o(1)$ by setting $t_3 := C\sqrt{(1+d_W)\log(|V|nM_n/\xi)}$. Finally, by Lemma 11 with $K_u = C\sqrt{\frac{(1+d_W)\log(|V|n)}{n}}$ we have $P(\Omega_4) = 1 - o(1)$.

Moreover, we have that $\|\bar{\beta}_u\|_{1,\varpi} \leq \sqrt{s}\|\bar{\beta}_u\|_{2,\varpi} \leq C\sqrt{s} = o(\sqrt{n})$ and $\frac{1}{\lambda_u(1-1/c)}\bar{R}_{u\xi} = o(\sqrt{n})$ for all $u \in \mathcal{U}$. Finally, we verify condition (F.63) holds for all

$$\delta \in A_u := \Delta_{u,2c} \cup \{v : \|v\|_{1,\varpi} \leq 2c\bar{R}_{u\xi}/\lambda_u, \|\sqrt{f_u}X'_{-a}v\|_{n,\varpi} \geq C\sqrt{s(1+d_W)\log(n|V|)/n/\kappa_{u,2c}}\},$$

$$\bar{q}_{A_u}/4 \geq (\sqrt{\bar{f}} + 1)\|r_u\|_{n,\varpi} + [\lambda_u + t_3 + K_u] \frac{3c\sqrt{s}}{\kappa_{u,2c}} \text{ and } \bar{q}_{A_u} \geq \{2c\left(1 + \frac{t_3+K_u}{\lambda_u}\right) \bar{R}_{u\gamma}\}^{1/2}.$$

Consider the matrices $\mathbb{E}_n[K_{\varpi}(W)f_uX_{-a}X'_{-a}]$ and $E[K_{\varpi}(W)f_uX_{-a}X'_{-a}]$. By Lemma 6, with probability $1 - o(1)$, it follows that we can take $\eta = \eta_n = CM_n\sqrt{s(1+d_W)\log(|V|n)\log(1+s)\{\log n\}/\sqrt{n}}$ and $D_{kk} = 2\eta$ in Lemma 16. (Note that we increase δ_n by a factor of $\sqrt{\log n}$.) Therefore, with at least the same probability we have (taking $s \geq 2$)

$$\delta' \mathbb{E}_n[K_{\varpi}(W)f_uX_{-a}X'_{-a}]\delta \geq \delta' E[K_{\varpi}(W)f_uX_{-a}X'_{-a}]\delta - 4\eta\|\delta\|_1^2/s \quad (\text{C.45})$$

and by definition of \underline{f}_u we have

$$\mathbb{E}_n[K_{\varpi}(W)f_u|X'_{-a}\delta|^2] \geq \underline{f}_u E[K_{\varpi}(W)|X'_{-a}\delta|^2] - 4\eta\|\delta\|_1^2/s \geq c\underline{f}_u P(\varpi)\|\delta\|^2 - 4\eta\|\delta\|_1^2/s.$$

For $\delta \in \Delta_{u,2c}$ we have $\|\delta\|_1 \leq C\|\delta\|_{1,\varpi}/\{P(\varpi)\}^{1/2} \leq C'\|\delta_{T_u}\|_{1,\varpi}/\{P(\varpi)\}^{1/2} \leq C'\sqrt{s}\|\delta_{T_u}\|_2$. Note that we can assume $\|\delta\| \geq c\sqrt{s(1+d_W)\log(n|V|)/n}$ otherwise we are done. So that for $\delta \in A_u \setminus \Delta_{u,2c}$ we have that $\|\delta\|_1/\|\delta\|_2 \leq Cs\sqrt{\log(|V|n)/n}/\sqrt{s(1+d_W)\log(n|V|)/n} \leq C'\sqrt{s}$.

Similarly we have

$$\mathbb{E}_n[K_{\varpi}(W)|X'_{-a}\delta|^2] \leq E[K_{\varpi}(W)|X'_{-a}\delta|^2] + 4\eta\|\delta\|_1^2/s \leq CP(\varpi)\|\delta\|^2 - 4\eta\|\delta\|_1^2/s. \quad (\text{C.46})$$

Under the condition that $\eta = o(\underline{f}_{\mathcal{U}}\mu_{\mathcal{W}})$, which holds by Condition P, for n sufficiently large we have with probability $1 - o(1)$ that

$$\begin{aligned} \bar{q}_{A_u} &\geq \frac{c}{\bar{f}'} \inf_{\delta \in A_u} \frac{\mathbb{E}_n[K_{\varpi}(W)f_u|X'_{-a}\delta|^2]^{3/2}}{\mathbb{E}_n[K_{\varpi}(W)|X'_{-a}\delta|^3]} \geq \frac{c}{\bar{f}'} \inf_{\delta \in A_u} \frac{\mathbb{E}_n[K_{\varpi}(W)f_u|X'_{-a}\delta|^2]^{3/2}}{\mathbb{E}_n[K_{\varpi}(W)|X'_{-a}\delta|^2] \max_{i \leq n} \|X_i\|_{\infty} \|\delta\|_1} \\ &\geq \frac{c}{\bar{f}'} \inf_{\delta \in A_u} \frac{\{c' \underline{f}_u \mathbb{P}(\varpi) \|\delta\|^2\}^{3/2}}{C' \mathbb{P}(\varpi) \|\delta\|^2 \max_{i \leq n} \|X_i\|_{\infty} \|\delta\|_1} \geq \frac{c}{\bar{f}'} \inf_{\delta \in A_u} \frac{c' \underline{f}_u^{3/2} \mathbb{P}(\varpi)^{1/2} \|\delta\|}{C' \max_{i \leq n} \|X_i\|_{\infty} \|\delta\|_1} \\ &\geq C'' \frac{\underline{f}_{\mathcal{U}}^{3/2}}{\bar{f}'} \frac{\mu_{\mathcal{W}}^{1/2}}{\sqrt{s} \max_{i \leq n} \|X_i\|_{\infty}} \end{aligned} \quad (\text{C.47})$$

where $\max_{i \leq n} \|X_i\|_{\infty} \leq \ell_n M_n$ with probability $1 - o(1)$ for any $\ell_n \rightarrow \infty$. Therefore, under the condition $M_n s \sqrt{\log(p|V|n)} = o(\sqrt{n\mu_{\mathcal{W}}})$ assumed in Condition P, the conditions on \bar{q}_{A_u} are satisfied.

By Lemma 7, we have uniformly over all $u = (a, \tau, \varpi) \in \mathcal{U} := V \times \mathcal{T} \times \mathcal{W}$

$$\|\sqrt{f_u} X'_{-a}(\hat{\beta}_u - \beta_u)\|_{n, \varpi} \leq C \sqrt{\frac{(1 + d_W) \log(n|V|)}{n}} \frac{\sqrt{s}}{\kappa_{u, 2c}} \quad \text{and} \quad \|\hat{\beta}_u - \beta_u\|_{1, \varpi} \leq C \sqrt{\frac{(1 + d_W) \log(n|V|)}{n}} \frac{s}{\kappa_{u, 2c}}$$

where $\kappa_{u, 2c}$ is bounded away from zero with probability $1 - o(1)$ for n sufficiently large. Consider the thresholded estimators $\hat{\beta}_u^{\bar{\lambda}}$ for $\bar{\lambda} = \{(1 + d_W) \log(n|V|)/n\}^{1/2}$. By Lemma 17 we have $\|\hat{\beta}_u^{\bar{\lambda}}\|_0 \leq Cs$ and the same rates of convergence as $\hat{\beta}_u$. Therefore, by refitting over the support of $\hat{\beta}_u^{\bar{\lambda}}$ we have by Lemma 14, the estimator $\tilde{\beta}_u$ has the same rate of convergence where we used that $\hat{Q}_u \leq \lambda_u \|\hat{\beta}_u^{\bar{\lambda}} - \beta_u\|_{1, \varpi} \lesssim Cs(1 + d_W) \log(|V|n)/n$ (the other conditions of Lemma 14 hold as for the conditions in Lemma 7).

Next we will invoke Lemma 7 for the new penalty choice and penalty loadings. (We note that minor modifications cover the new penalty loadings.)

$$\begin{aligned} \Omega_1 &:= \{\lambda_u \geq c|S_{uj}|/\{\mathbb{E}_n[K_w(W)\varepsilon_u^2 X_{-a,j}^2]\}^{1/2}, \text{ for all } u \in \mathcal{U}, j \in V\}, \\ \Omega_2 &:= \{\hat{R}_u(\tilde{\beta}_u) \leq \hat{R}_{u\xi} : u \in \mathcal{U}\} \\ \Omega_3 &:= \left\{ \sup_{u \in \mathcal{U}, 1/\sqrt{n} \leq \|\delta\|_{1, \varpi} \leq \sqrt{n}} |\mathbb{E}_n[g_u(\delta, X, W) - \mathbb{E}[g_u(\delta, X, W)|X_{-a}, W]]|/\{\theta_u \|\delta\|_{1, \varpi}\} \leq t_3 \right\} \\ \Omega_4 &:= \{K_u \theta_u \hat{\sigma}_{a\varpi j}^X \geq |\mathbb{E}_n[h_{uj}(X_{-a}, W)]|, \text{ for all } u \in \mathcal{U}, j \in V \setminus \{a\}\} \\ \Omega_5 &:= \{\theta_u \geq \max_{j \in V} \hat{\sigma}_{a\varpi j}^X / \{\mathbb{E}_n[K_{\varpi}(W)\varepsilon_u^2 X_j^2]\}^{1/2}\} \end{aligned}$$

where event Ω_5 simply makes the relevant norms equivalent, $\|\cdot\|_{1, u} \leq \|\cdot\|_{1, \varpi} \leq \theta_u \|\cdot\|_{1, u}$. Note that we can always take $\theta_u \leq 1/\{\tau(1 - \tau)\} \leq C$ since \mathcal{T} is a fixed compact set.

Next we show that the bootstrap approximation of the score provides a valid choice of penalty parameter. Let $\hat{\varepsilon}_u := 1\{X_a \leq X'_{-a}\tilde{\beta}_u\} - \tau$. For notational convenience for $u \in \mathcal{U}$, $j \in V \setminus \{a\}$ define

$$\bar{\psi}_{iuj} = \frac{K_{\varpi}(W_i)\varepsilon_{iu}X_{ij}}{\mathbb{E}[K_{\varpi}(W)\varepsilon_u^2 X_j^2]^{1/2}}, \quad \psi_{iuj} = \frac{K_{\varpi}(W_i)\varepsilon_{iu}X_{ij}}{\mathbb{E}_n[K_{\varpi}(W)\varepsilon_u^2 X_j^2]^{1/2}}, \quad \hat{\psi}_{iuj} = \frac{K_{\varpi}(W_i)\hat{\varepsilon}_{iu}X_{ij}}{\mathbb{E}_n[K_{\varpi}(W)\hat{\varepsilon}_u^2 X_j^2]^{1/2}}.$$

We will consider the following processes:

$$\bar{S}_{uj} = \frac{1}{\sqrt{n}} \sum_{i=1}^n \bar{\psi}_{iuj}, \quad S_{uj} = \frac{1}{\sqrt{n}} \sum_{i=1}^n \psi_{iuj}, \quad \bar{\mathcal{G}}_{uj} = \frac{1}{\sqrt{n}} \sum_{i=1}^n g_i \bar{\psi}_{iuj}, \quad \hat{\mathcal{G}}_{uj} = \frac{1}{\sqrt{n}} \sum_{i=1}^n g_i \hat{\psi}_{iuj},$$

and \mathcal{N} is a tight zero-mean Gaussian process with covariance operator given by $\mathbb{E}[\bar{\psi}_{uj}\bar{\psi}_{u'j}]$. Their supremum are denoted by $\bar{Z}_S := \sup_{u \in \mathcal{U}, j \in V \setminus \{a\}} |\bar{S}_{uj}|$, $Z_S := \sup_{u \in \mathcal{U}, j \in V \setminus \{a\}} |S_{uj}|$, $\bar{Z}_G^* := \sup_{u \in \mathcal{U}, j \in V \setminus \{a\}} |\bar{\mathcal{G}}_{uj}|$, $\hat{Z}_G^* := \sup_{u \in \mathcal{U}, j \in V \setminus \{a\}} |\hat{\mathcal{G}}_{uj}|$, and $Z_N := \sup_{u \in \mathcal{U}, j \in V \setminus \{a\}} |\mathcal{N}_{uj}|$.

The penalty choice should majorate Z_S and we simulate via \widehat{Z}_G^* . We have that

$$\begin{aligned} |\mathbb{P}(Z_S \leq t) - \mathbb{P}(\widehat{Z}_G^* \leq t)| &\leq |\mathbb{P}(Z_S \leq t) - \mathbb{P}(\bar{Z}_S \leq t)| + |\mathbb{P}(\bar{Z}_S \leq t) - \mathbb{P}(Z_N \leq t)| \\ &\quad + |\mathbb{P}(Z_N \leq t) - \mathbb{P}(\bar{Z}_G^* \leq t)| + |\mathbb{P}(\bar{Z}_G^* \leq t) - \mathbb{P}(\widehat{Z}_G^* \leq t)| \end{aligned}$$

We proceed to bound each term. We have that

$$\begin{aligned} |Z_S - \bar{Z}_S| &\leq \bar{Z}_S \sup_{u \in \mathcal{U}, j \in V \setminus \{a\}} \left| \frac{\mathbb{E}[K_\varpi(W) \varepsilon_u^2 X_j^2]^{1/2}}{\mathbb{E}_n[K_\varpi(W) \varepsilon_u^2 X_j^2]^{1/2}} - 1 \right| \\ &\leq \bar{Z}_S \sup_{u \in \mathcal{U}, j \in V \setminus \{a\}} \left| \frac{(\mathbb{E}_n - \mathbb{E})[K_\varpi(W) \varepsilon_u^2 X_j^2]}{\mathbb{E}_n[K_\varpi(W) \varepsilon_u^2 X_j^2]^{1/2} \{ \mathbb{E}_n[K_\varpi(W) \varepsilon_u^2 X_j^2]^{1/2} + \mathbb{E}[K_\varpi(W) \varepsilon_u^2 X_j^2]^{1/2} \}} \right| \end{aligned}$$

Therefore, since $\{1\{X_a \leq X'_a \beta_u\} : u \in \mathcal{U}\}$ is a VC-subgraph of VC dimension $1 + d_W$, and \mathcal{W} is a VC class of sets of dimension d_W , we apply Lemma 19 with envelope $F = \|X\|_\infty^2$ and $\sigma^2 \leq \max_{j \in V} \mathbb{E}[X_j^4] \leq C$ to obtain with probability $1 - o(1)$

$$\sup_{u \in \mathcal{U}, j \in V \setminus \{a\}} |(\mathbb{E}_n - \mathbb{E})[K_\varpi(W) \varepsilon_u^2 X_j^2]| \lesssim \delta'_{1n} := \sqrt{\frac{(1 + d_W) \log(|V|n)}{n}} + \frac{M_n^2(1 + d_W) \log(|V|n)}{n}$$

where $\delta_{1n} = o(\mu_{\mathcal{W}}^2)$ under Condition P. Note that this implies that the denominator above is bounded away from zero by $c\mu_{\mathcal{W}}$. Therefore,

$$|Z_S - \bar{Z}_S| \lesssim_P \delta_{1n} := \bar{Z}_S \delta'_{1n} / \mu_{\mathcal{W}}.$$

where $\bar{Z}_S \lesssim_P \{(1 + d_W) \log(n|V|)\}^{1/2}$. By Theorem 2.1 in [31], since $\mathbb{E}[\bar{\psi}_{uj}^4] \leq C$, there is a version of Z_N such that

$$|\bar{Z}_S - Z_N| \lesssim_P \delta_{2n} := \left(\frac{M_n(1 + d_W) \log(n|V|)}{n^{1/2}} + \frac{M_n^{1/3}((1 + d_W) \log(n|V|))^{2/3}}{n^{1/6}} \right)$$

and by Theorem 2.2 in [31], there is also a version of

$$|Z_N - \bar{Z}_G^*| \lesssim_P \left(\frac{M_n(1 + d_W) \log(n|V|)}{n^{1/2}} + \frac{M_n^{1/2}((1 + d_W) \log(n|V|))^{3/4}}{n^{1/4}} \right)$$

Finally, we have that

$$|\bar{Z}_G^* - \widehat{Z}_G^*| \leq \sup_{u \in \mathcal{U}, j} \left| \frac{1}{\sqrt{n}} \sum_{i=1}^n g_i(\widehat{\psi}_{iuj} - \bar{\psi}_{iuj}) \right|$$

where conditional on $(X_i, W_i), i = 1, \dots, n$, $\frac{1}{\sqrt{n}} \sum_{i=1}^n g_i(\widehat{\psi}_{iuj} - \bar{\psi}_{iuj})$ is a zero-mean Gaussian with variance $\mathbb{E}_n[(\widehat{\psi}_{iuj} - \bar{\psi}_{iuj})^2] \leq \bar{\delta}_n^2$. Next we bound $\bar{\delta}_n$. We have

$$\bar{\delta}_n \leq \mathbb{E}_n[(\widehat{\psi}_{uj} - \psi_{uj})^2]^{1/2} + \mathbb{E}_n[(\psi_{uj} - \bar{\psi}_{uj})^2]^{1/2} \leq \mathbb{E}_n[(\widehat{\psi}_{uj} - \psi_{uj})^2]^{1/2} + \delta_{1n} / \mu_{\mathcal{W}},$$

and

$$\begin{aligned} \mathbb{E}_n[(\widehat{\psi}_{uj} - \psi_{uj})^2]^{1/2} &\leq \frac{\mathbb{E}_n[(K_\varpi(W) X_{ij} |\widehat{\varepsilon}_u - \varepsilon_u|)^2]^{1/2}}{\mathbb{E}_n[K_\varpi(W) X_{ij}^2 \widehat{\varepsilon}_u^2]^{1/2}} \\ &\quad + \frac{\mathbb{E}_n[K_\varpi(W) X_{ij}^2 \varepsilon_u^2]^{1/2}}{cP(\varpi)} |\mathbb{E}_n[K_\varpi(W) X_{ij}^2 \widehat{\varepsilon}_u^2]^{1/2} - \mathbb{E}_n[K_\varpi(W) X_{ij}^2 \varepsilon_u^2]^{1/2}| \\ &\leq \mathbb{E}_n[K_\varpi(W) X_{ij}^2 |\widehat{\varepsilon}_u - \varepsilon_u|^2]^{1/2} \left\{ \frac{1}{\mathbb{E}_n[K_\varpi(W) X_{ij}^2 \widehat{\varepsilon}_u^2]^{1/2}} + \frac{\mathbb{E}_n[K_\varpi(W) X_{ij}^2 \varepsilon_u^2]^{1/2}}{cP(\varpi)} \right\}, \end{aligned}$$

note that the term in the curly brackets is bounded by $C/P(\varpi)^{1/2}$ with probability $1 - o(1)$. To bound the other term note that $|\widehat{\varepsilon}_u - \varepsilon_u|^2 = |1\{X_a \leq X'_a \widetilde{\beta}_u\} - 1\{X_a \leq X'_a \beta_u\}|$. Note that $\widehat{\varepsilon}_u = 1\{X_a \leq X'_a \widetilde{\beta}_u\} - \tau$

where $\|\tilde{\beta}_u\|_0 \leq Cs$. Therefore, we have $\{1\{X_a \leq X'_{-a}\tilde{\beta}_u\} : u \in \mathcal{U}\} \subset \{1\{X_a \leq X'_{-a}\beta\} : \|\beta\|_0 \leq Cs\}$ which is the union of $\binom{|V|}{Cs}$ VC subgraph classes of functions with VC dimension $C's$. Moreover, we have

$$\begin{aligned} \mathbb{E}[K_\varpi(W)X_{ij}^2|\hat{\varepsilon}_u - \varepsilon_u|^2] &= \mathbb{E}[K_\varpi(W)X_{ij}^2|1\{X_a \leq X'_{-a}\tilde{\beta}_u\} - 1\{X_a \leq X'_{-a}\beta_u\}|] \\ &\leq \bar{f}\mathbb{E}[K_\varpi(W)X_{ij}^2|X'_{-a}(\tilde{\beta}_u - \beta_u)|] \\ &\leq \bar{f}\mathbb{E}[K_\varpi(W)X_{ij}^4]^{1/2}\mathbb{E}[K_\varpi(W)|X'_{-a}(\tilde{\beta}_u - \beta_u)|^2]^{1/2} \\ &\leq C(\bar{f}/\underline{f}_{\mathcal{U}}^{1/2})P(\varpi)^{1/2}\sqrt{s(1+d_W)\log(n|V|)/n} \end{aligned}$$

Therefore, by Lemma 19, with probability $1 - o(1)$ we have

$$|(\mathbb{E}_n - \mathbb{E})[K_\varpi(W)X_{ij}^2|\hat{\varepsilon}_u - \varepsilon_u|^2]| \lesssim \sqrt{\frac{s(1+d_W)\log(n|V|)}{n}} C(\bar{f}/\underline{f}_{\mathcal{U}}^{1/2})\sqrt{s(1+d_W)\log(n|V|)/n}$$

Under $\sqrt{s(1+d_W)\log(n|V|)/n} = o(\underline{f}_{\mathcal{U}}\mu_W)$ we have that with probability $1 - o(1)$ that

$$\bar{\delta}_n \leq C\{s(1+d_W)\log(n|V|)/n\}^{1/4}.$$

Therefore, using again the sparsity of $\tilde{\beta}_u$ in the definition of $\hat{\psi}_{iuj}$

$$\begin{aligned} \sup_{u \in \mathcal{U}, j \in V \setminus \{a\}} \left| \frac{1}{\sqrt{n}} \sum_{i=1}^n g_i(\hat{\psi}_{iuj} - \bar{\psi}_{iuj}) \right| &\lesssim_P \bar{\delta}_n \sqrt{s(1+d_W)\log(|V|n)} \\ &\lesssim_P \delta_{3n} := \{s\log(|V|n)/n\}^{1/4} \sqrt{s(1+d_W)\log(|V|n)} \end{aligned}$$

The rest of the proof follows similarly to Corollary 2.2 in [15] since under Condition P (and the bounds above) we have that $r_n := \delta_{1n} + \delta_{2n} + \delta_{3n} = o(\{\mathbb{E}[Z_N]\}^{-1})$ where $\mathbb{E}[Z_N] \lesssim \{(1+d_W)\log(|V|n)\}^{1/2}$. Then we have $\sup_t |P(Z_S \leq t) - P(\hat{Z}_G^* \leq t)| = o_P(1)$ which in turn implies that

$$\begin{aligned} P(\Omega_1) &= P(Z_S \leq \hat{c}_G^*(\xi)) \\ &\geq P(\hat{Z}_G^* \leq \hat{c}_G^*(\xi)) - |P(Z_S \leq \hat{c}_G^*(\xi)) - P(\hat{Z}_G^* \leq \hat{c}_G^*(\xi))| \\ &\geq 1 - \xi + o_P(1) \end{aligned}$$

Note that the occurrence of the events Ω_2 , Ω_3 and Ω_4 follows by similar arguments. The result follows by Lemma 7, thresholding and applying Lemma 17 and Lemma 14 similarly to before. ■

APPENDIX D. TECHNICAL LEMMAS FOR CONDITIONAL INDEPENDENCE QUANTILE GRAPHICAL MODEL

Let $u = (a, \tau, \varpi) \in \mathcal{U} := V \times \mathcal{T} \times \mathcal{W}$, and $T_u = \text{support}(\beta_u)$ where $|T_u| \leq s$ for all $u \in \mathcal{U}$.

Define the pseudo-norms

$$\|v\|_{n,\varpi}^2 := \frac{1}{n} \sum_{i=1}^n K_\varpi(W_i)(v_i)^2, \quad \|\delta\|_{2,\varpi} := \left\{ \sum_{j=1}^p \{\hat{\sigma}_{a\varpi j}^Z\}^2 |\delta_j|^2 \right\}^{1/2}, \quad \text{and} \quad \|\delta\|_{1,\varpi} := \sum_{j=1}^p \hat{\sigma}_{a\varpi j}^Z |\delta_j|,$$

where $\hat{\sigma}_{a\varpi j}^Z = \{\mathbb{E}_n[\{K_\varpi(W)Z_j^a\}^2]\}^{1/2}$. These pseudo-norms induce the following restricted eigenvalue as

$$\kappa_{u,\mathbf{c}} = \min_{\|\delta_{T_u^c}\|_{1,\varpi} \leq \mathbf{c}\|\delta_{T_u}\|_{1,\varpi}} \frac{\|\sqrt{f_u}Z^a\delta\|_{n,\varpi}}{\|\delta\|_{1,\varpi}/\sqrt{s}}.$$

The restricted eigenvalue $\kappa_{u,\mathbf{c}}$ is an counterpart of the restricted eigenvalue proposed in [22] for our setting. We note that $\kappa_{u,\mathbf{c}}$ typically will vary with the events $\varpi \in \mathcal{W}$.

We will consider three key events in our analysis. Let

$$\Omega_1 := \{\lambda_u \geq c|S_{uj}|/\hat{\sigma}_{a\varpi j}^Z, \text{ for all } u \in \mathcal{U}, j \in [p]\} \quad (\text{D.48})$$

which occurs with probability at least $1 - \xi$ by the choice of λ_u . For CIQGMs, we have $S_{uj} := \mathbb{E}_n[K_\varpi(W)(\tau - 1\{X_a \leq Z^a\beta_u + r_u\})Z_j^a]$, and $\lambda_u = \lambda_{V\tau W}\sqrt{\tau(1-\tau)}$. (In the case of PQGMs, we have $S_{uj} := \mathbb{E}_n[K_\varpi(W)(\tau - 1\{X_a \leq X'_{-a}\beta_u\})X_{-a}]$, $\hat{\sigma}_{a\varpi j}^X = \{\mathbb{E}_n[K_\varpi(W)X_{-a,j}^2]\}^{1/2}$ and $\lambda_u = \lambda_0$.)

To define the next event, for each $u \in \mathcal{U}$, consider the function defined as

$$\hat{R}_u(\beta_u) = \mathbb{E}_n[K_\varpi(W)\{\rho_u(X_a - Z^a\beta) - \rho_u(X_a - Z^a\beta_u - r_u) - (\tau - 1\{X_a \leq Z^a\beta_u + r_u\})(Z^a\beta - Z^a\beta_u - r_u)\}]$$

in the case of CIQGMs. (In the case of PQGMs, we replace Z^a with X_{-a} .) By convexity we have $\hat{R}_u(\beta_u) \geq 0$. The event

$$\Omega_2 := \{\hat{R}_u(\beta_u) \leq \bar{R}_{u\xi} : u \in \mathcal{U}\} \quad (\text{D.49})$$

where $\bar{R}_{u\xi}$ are chosen so that Ω_2 occurs with probability at least $1 - \xi$. Note that by Lemma 2, we have $\mathbb{E}_n[\mathbb{E}[\hat{R}_u(\beta_u)|X_{-a}, W]] \leq \bar{f}\|r_u\|_{n,\varpi}^2/2$ and with probability at least $1 - \xi$, $\hat{R}_u(\beta_u) \leq \bar{R}_{u\xi} := 4\max\{\bar{f}\|r_u\|_{n,\varpi}^2, \|r_u\|_{n,\varpi}C\sqrt{\log(n^{1+d_W}p/\xi)/n}\} \leq C's\log(n^{1+d_W}p/\xi)/n$.

Define $g_u(\delta, X, W) = K_\varpi(W)\{\rho_\tau(X_a - Z^a(\beta_u + \delta)) - \rho_\tau(X_a - Z^a\beta_u)\}$ so that event Ω_3 is defined as

$$\Omega_3 := \left\{ \sup_{u \in \mathcal{U}, 1/\sqrt{n} \leq \|\delta\|_{1,\varpi} \leq \sqrt{n}} \frac{|\mathbb{E}_n[g_u(\delta, X, W)] - \mathbb{E}[g_u(\delta, X, W)|X_{-a}, W]|}{\|\delta\|_{1,\varpi}} \leq t_3 \right\} \quad (\text{D.50})$$

where t_3 is given in Lemma 3 so that Ω_3 holds with probability at least $1 - \xi$.

Lemma 1. Suppose that Ω_1 , Ω_2 and Ω_3 holds. Further assume $2\frac{1+1/c}{1-1/c}\|\beta_u\|_{1,\varpi} + \frac{1}{\lambda_u(1-1/c)}\bar{R}_{u\xi} \leq \sqrt{n}$ for all $u \in \mathcal{U}$, and (F.63) holds for all $\delta \in A_u := \Delta_{u,2c} \cup \{v : \|v\|_{1,\varpi} \leq 2c\bar{R}_{u\xi}/\lambda_u\}$, $\bar{q}_{A_u}/4 \geq (\sqrt{f} + 1)\|r_u\|_{n,\varpi} + [\lambda_u + t_3]\frac{3c\sqrt{s}}{\kappa_{u,2c}}$ and $\bar{q}_{A_u} \geq \{2c(1 + \frac{t_3}{\lambda_u})\bar{R}_{u\xi}\}^{1/2}$. Then uniformly over all $u \in \mathcal{U}$ we have

$$\begin{aligned} \|\sqrt{f_u}Z^a(\hat{\beta}_u - \beta_u)\|_{n,\varpi} &\leq \sqrt{8c\left(1 + \frac{t_3}{\lambda_u}\right)\bar{R}_{u\xi} + (f^{1/2} + 1)\|r_u\|_{n,\varpi} + \frac{3c\lambda_u\sqrt{s}}{\kappa_{u,2c}} + t_3\frac{(1+c)\sqrt{s}}{\kappa_{u,2c}}} \\ \|\hat{\beta}_u - \beta_u\|_{1,\varpi} &\leq (1 + 2c)\sqrt{s}\|\sqrt{f_u}Z^a\delta_u\|_{n,\varpi}/\kappa_{u,2c} + \frac{2c}{\lambda_u}\bar{R}_{u\xi} \end{aligned}$$

Proof of Lemma 1. Let $u = (a, \tau, \varpi) \in \mathcal{U}$ and $\delta_u = \hat{\beta}_u - \beta_u$. By convexity and definition of $\hat{\beta}_u$ we have

$$\begin{aligned} &\hat{R}_u(\hat{\beta}_u) - \hat{R}_u(\beta_u) + S'_u\delta_u \\ &= \mathbb{E}_n[K_\varpi(W)\rho_u(X_a - Z^a\hat{\beta}_u)] - \mathbb{E}_n[K_\varpi(W)\rho_u(X_a - Z^a\beta_u)] \\ &\leq \lambda_u\|\beta_u\|_{1,\varpi} - \lambda_u\|\hat{\beta}_u\|_{1,\varpi} \end{aligned} \quad (\text{D.51})$$

where S_u is defined as in (D.48) so that under Ω_1 we have $\lambda_u \geq c|S_{uj}|/\hat{\sigma}_{a\varpi j}^Z$.

Under $\Omega_1 \cap \Omega_2$, and since $\hat{R}_u(\hat{\beta}_u) \geq 0$, we have

$$\begin{aligned} -\hat{R}_u(\beta_u) - \frac{\lambda_u}{c}\|\delta_u\|_{1,\varpi} &\leq \hat{R}_u(\beta_u + \delta_u) - \hat{R}_u(\beta_u) + \mathbb{E}_n[K_\varpi(W)(\tau - 1\{X_a \leq Z^a\beta_u + r_u\})Z^a\delta_u] \\ &= \mathbb{E}_n[K_\varpi(W)\rho_u(X_a - Z^a(\beta_u + \delta_u))] - \mathbb{E}_n[K_\varpi(W)\rho_u(X_a - Z^a\beta_u)] \\ &\leq \lambda_u\|\beta_u\|_{1,\varpi} - \lambda_u\|\delta_u + \beta_u\|_{1,\varpi} \end{aligned} \quad (\text{D.52})$$

so that for $\mathbf{c} = (c+1)/(c-1)$

$$\|\delta_{T_u^c}\|_{1,\varpi} \leq \mathbf{c}\|\delta_{T_u}\|_{1,\varpi} + \frac{c}{\lambda_u(c-1)}\hat{R}_u(\beta_u).$$

To establish that $\delta_u \in A_u := \Delta_{u,2c} \cup \{v : \|v\|_{1,\varpi} \leq 2c\bar{R}_{u\xi}/\lambda_u\}$ we consider two cases. If $\|\delta_{T_u^c}\|_{1,\varpi} \geq 2c\|\delta_{T_u}\|_{1,\varpi}$ we have

$$\frac{1}{2}\|\delta_{T_u^c}\|_{1,\varpi} \leq \frac{c}{\lambda_u(c-1)}\hat{R}_u(\beta_u)$$

and consequentially

$$\|\delta_u\|_{1,\varpi} \leq \{1 + 1/(2c)\}\|\delta_{T_u^c}\|_{1,\varpi} \leq \frac{2c}{\lambda_u}\hat{R}_u(\beta_u).$$

Otherwise, we have $\|\delta_{T_u^c}\|_{1,\varpi} \leq 2c\|\delta_{T_u}\|_{1,\varpi}$ which implies

$$\|\delta_u\|_{1,\varpi} \leq (1 + 2c)\|\delta_{T_u}\|_{1,\varpi} \leq (1 + 2c)\sqrt{s}\|\sqrt{f_u}Z^a\delta_u\|_{n,\varpi}/\kappa_{u,2c}$$

by definition of $\kappa_{u,2c}$. Thus we have $\delta_u \in A_u$ under $\Omega_1 \cap \Omega_2$.

Furthermore, (D.52) also implies that

$$\begin{aligned} \|\delta_u + \beta_u\|_{1,\varpi} &\leq \|\beta_u\|_{1,\varpi} + \frac{1}{c}\|\delta_u\|_{1,\varpi} + \hat{R}_u(\beta_u)/\lambda_u \\ &\leq (1 + 1/c)\|\beta_u\|_{1,\varpi} + (1/c)\|\delta_u + \beta_u\|_{1,\varpi} + \hat{R}_u(\beta_u)/\lambda_u. \end{aligned}$$

which in turn establishes

$$\|\delta_u\|_{1,\varpi} \leq 2\frac{1+1/c}{1-1/c}\|\beta_u\|_{1,\varpi} + \frac{1}{\lambda_u(1-1/c)}\hat{R}_u(\beta_u) \leq 2\frac{1+1/c}{1-1/c}\|\beta_u\|_{1,\varpi} + \frac{1}{\lambda_u(1-1/c)}\bar{R}_{u\xi}$$

where the last inequality holds under Ω_2 . Thus, $\|\delta_u\|_{1,\varpi} \leq \sqrt{n}$ under our condition. In turn, δ_u is considered in the supremum that defines Ω_3 .

Under $\Omega_1 \cap \Omega_2 \cap \Omega_3$ we have

$$\begin{aligned} &\mathbb{E}_n[\mathbb{E}[K_\varpi(W)\{\rho_u(X_a - Z^a(\beta_u + \delta_u)) - \rho_u(X_a - Z^a\beta_u)\} \mid X_{-a}, W]] \\ &\leq \mathbb{E}_n[K_\varpi(W)\{\rho_u(X_a - Z^a(\beta_u + \delta_u)) - \rho_u(X_a - Z^a\beta_u)\}] + t_3\|\delta_u\|_{1,\varpi} \\ &\leq \lambda_u\|\delta_u\|_{1,\varpi} + t_3\|\delta_u\|_{1,\varpi} \\ &\leq 2c\left(1 + \frac{1}{\lambda_u}t_3\right)\bar{R}_{u\xi} + \|\sqrt{f_u}Z^a\delta_u\|_{n,\varpi}[\lambda_u + t_3]\frac{3c\sqrt{s}}{\kappa_{u,2c}} \end{aligned} \tag{D.53}$$

here we used the bound $\|\delta_u\|_{1,\varpi} \leq (1 + 2c)\sqrt{s}\|\sqrt{f_u}Z^a\delta_u\|_{n,\varpi}/\kappa_{u,2c} + \frac{2c}{\lambda_u}\bar{R}_{u\xi}$ under $\Omega_1 \cap \Omega_2$.

Using Lemma 12, since (F.63) holds, we have for each $u \in \mathcal{U}$

$$\begin{aligned} &\mathbb{E}_n[\mathbb{E}[K_\varpi(W)\{\rho_u(X_a - Z^a(\beta_u + \delta_u)) - \rho_u(X_a - Z^a\beta_u)\} \mid X_{-a}, W]] \\ &\geq -(\sqrt{f} + 1)\|r_u\|_{n,\varpi}\|\sqrt{f_u}Z^a\delta_u\|_{n,\varpi} - \sup_{u \in \mathcal{U}, j \in [p]} |\mathbb{E}_n[\mathbb{E}[S_{uj} \mid X_{-a}, W]/\hat{\sigma}_{a\varpi j}^Z]| \|\delta_u\|_{1,\varpi} \\ &\quad + \frac{\|\sqrt{f_u}Z^a\delta_u\|_{n,\varpi}^2}{4} \wedge \{\bar{q}_{A_u}\|\sqrt{f_u}Z^a\delta_u\|_{n,\varpi}\} \end{aligned}$$

here we have $\mathbb{E}[S_{iuj} \mid X_{i,-a}, W_i] = 0$ since $\tau = P(X_a \leq Z^a\beta_u + r_u \mid X_{-a}, W)$ by the definition of conditional quantile.

Note that for positive numbers $(t^2/4) \wedge qt \leq A + Bt$ implies $t^2/4 \leq A + Bt$ provided $q/2 > B$ and $2q^2 > A$. (Indeed, otherwise $(t^2/4) \geq qt$ so that $t \geq 4q$ which in turn implies that $2q^2 + qt/2 \leq (t^2/4) \wedge qt \leq A + Bt$.) Since $\bar{q}_{A_u}/4 \geq (\sqrt{f} + 1)\|r_u\|_{n,\varpi} + \left[\{\lambda_u + t_3\}\frac{3c\sqrt{s}}{\kappa_{u,2c}}\right]$ and $\bar{q}_{A_u} \geq \{2c\left(1 + \frac{t_3}{\lambda_u}\right)\bar{R}_{u\xi}\}^{1/2}$, the minimum on the right hand side is achieved by the quadratic part for all $u \in \mathcal{U}$. Therefore we have uniformly over $u \in \mathcal{U}$

$$\frac{\|\sqrt{f_u}Z^a\delta_u\|_{n,\varpi}^2}{4} \leq 2c\left(1 + \frac{t_3}{\lambda_u}\right)\bar{R}_{u\xi} + \|\sqrt{f_u}Z^a\delta_u\|_{n,\varpi}\left[(\sqrt{f} + 1)\|r_u\|_{n,\varpi} + \{\lambda_u + t_3\}\frac{3c\sqrt{s}}{\kappa_{u,2c}}\right]$$

which implies that

$$\|\sqrt{f_u}Z^a\delta_u\|_{n,\varpi} \leq \sqrt{8c\left(1 + \frac{t_3}{\lambda_u}\right)\bar{R}_{u\xi}} + \left[(\sqrt{f} + 1)\|r_u\|_{n,\varpi} + \{\lambda_u + t_3\}\frac{3c\sqrt{s}}{\kappa_{u,2c}}\right].$$

■

Lemma 2 (CIQGM, Event Ω_2). *Under Condition CI we have $\mathbb{E}_n[\mathbb{E}[\widehat{R}_u(\beta_u)|X_{-a}, \varpi]] \leq \bar{f}\|r_u\|_{n, \varpi}^2/2$, $\widehat{R}_u(\beta_u) \geq 0$ and*

$$\mathbb{P}\left(\sup_{u \in \mathcal{U}} \widehat{R}_u(\beta_u) \leq C\{1 + \bar{f}\}\{n^{-1}s(1 + d_W)\log(p|V|n)\}\right) = 1 - o(1).$$

Proof of Lemma 2. We have that $\widehat{R}_u(\beta_u) \geq 0$ by convexity of ρ_τ . Let $\varepsilon_{iu} = X_{ia} - Z_i^a \beta_u - r_{iu}$ where $\|\beta_u\|_0 \leq s$ and $r_{iu} = Q_{X_a}(\tau|X_{-a}, \varpi) - Z^a \beta_u$.

By Knight's identity (F.64), $\widehat{R}_u(\beta_u) = -\mathbb{E}_n[K_\varpi(W)r_u \int_0^1 1\{\varepsilon_u \leq -tr_u\} - 1\{\varepsilon_u \leq 0\} dt] \geq 0$.

$$\begin{aligned} \mathbb{E}_n[\mathbb{E}[\widehat{R}_u(\beta_u)|X_{-a}, \varpi]] &= \mathbb{E}_n[K_\varpi(W)r_u \int_0^1 F_{X_a|X_{-a}, \varpi}(Z^a \beta_u + (1-t)r_u) - F_{X_a|X_{-a}, \varpi}(Z^a \beta_u + r_u) dt] \\ &\leq \mathbb{E}_n[K_\varpi(W)r_u \int_0^1 \bar{f} tr_u dt] \leq \bar{f}\|r_u\|_{n, \varpi}^2/2 \leq C\bar{f}s/n. \end{aligned}$$

Since Condition CI assumes $\mathbb{E}[\|r_u\|_{n, \varpi}^2] \leq \mathbb{P}(\varpi)s/n$, by Markov's inequality we have $\mathbb{P}(\widehat{R}_u(\beta_u) \leq C\bar{f}s/n) \geq 1/2$. Define $z_{iu} := -\int_0^1 1\{\varepsilon_{iu} \leq -tr_{iu}\} - 1\{\varepsilon_{iu} \leq 0\} dt$, so that $\widehat{R}_u(\beta_u) = \mathbb{E}_n[K_\varpi(W)r_u z_u]$ where $|z_{iu}| \leq 1$. By Lemma 2.3.7 in [101] (note that the Lemma does not require zero mean stochastic processes), for $t \geq 2C\bar{f}s/n$ we have

$$\frac{1}{2}\mathbb{P}\left(\sup_{u \in \mathcal{U}} |\mathbb{E}_n[K_\varpi(W)r_u z_u]| \geq t\right) \leq 2\mathbb{P}\left(\sup_{u \in \mathcal{U}} |\mathbb{E}_n[\varepsilon K_\varpi(W)r_u z_u]| > t/4\right)$$

where $\varepsilon_i, i = 1, \dots, n$ are Rademacher random variables independent of the data.

Next consider the class of functions $\mathcal{F} = \{-K_\varpi(W)r_u(1\{\varepsilon_{iu} \leq -B_i r_{iu}\} - 1\{\varepsilon_{iu} \leq 0\}) : u \in \mathcal{U}\}$ where $B_i \sim \text{Uniform}(0, 1)$ independent of $(X_i, W_i)_{i=1}^n$. It follows that $K_\varpi(W)r_u z_u = \mathbb{E}[-K_\varpi(W)r_u(1\{\varepsilon_{iu} \leq -B_i r_{iu}\} - 1\{\varepsilon_{iu} \leq 0\})|X_i, W_i]$ where the expectation is taken over B_i only. Thus we will bound the entropy of $\bar{\mathcal{F}} = \{\mathbb{E}[f|X, W] : f \in \mathcal{F}\}$ via Lemma 25. Note that $\mathcal{R} := \{r_u = Q_{X_a}(\tau|X_{-a}, \varpi) - Z^a \beta_u : u \in \mathcal{U}\}$ where $\mathcal{G} := \{Z^a \beta_u : u \in \mathcal{U}\}$ is contained in the union of at most $|V|\binom{p}{s}$ VC-classes of dimension Cs and $\mathcal{H} := \{Q_{X_a}(\tau|X_{-a}, \varpi) : u \in \mathcal{U}\}$ is the union of $|V|$ VC-class of functions of dimension $(1 + d_W)$ by Condition CI. Finally note that $\mathcal{E} := \{\varepsilon_{iu} : u \in \mathcal{U}\} \subset \{X_{ia} : a \in V\} - \mathcal{G} - \mathcal{R}$.

Therefore, we have

$$\begin{aligned} \sup_Q \log N(\epsilon\|\bar{F}\|_{Q, 2}, \bar{\mathcal{F}}, \|\cdot\|_{Q, 2}) &\leq \sup_Q \log N((\epsilon/4)^2\|F\|_{Q, 2}, \mathcal{F}, \|\cdot\|_{Q, 2}) \\ &\leq \sup_Q \log N(\frac{1}{8}(\epsilon^2/16), \mathcal{W}, \|\cdot\|_{Q, 2}) \\ &\quad + \sup_Q \log N(\frac{1}{8}(\epsilon^2/16)\|F\|_{Q, 2}, \mathcal{R}, \|\cdot\|_{Q, 2}) \\ &\quad + \sup_Q \log N(\frac{1}{8}(\epsilon^2/16), 1\{\mathcal{E} + \{B\}\mathcal{R} \leq 0\} - 1\{\mathcal{E} \leq 0\}, \|\cdot\|_{Q, 2}) \end{aligned}$$

We will apply Lemma 19 with envelope $\bar{F} = \sup_{u \in \mathcal{U}} |K_\varpi(W)r_u|$, so that $\mathbb{E}[\max_{i \leq n} \bar{F}_i^2] \leq C$, and $\sup_{u \in \mathcal{U}} \mathbb{E}[K_\varpi(W)r_u^2] \leq Cs/n =: \sigma^2$ by Condition CI. Thus, we have that with probability $1 - o(1)$

$$\sup_{u \in \mathcal{U}} |\mathbb{E}_n[\varepsilon K_\varpi(W)r_u z_u]| \lesssim \sqrt{\frac{s(1 + d_W)\log(p|V|n)}{n}} \sqrt{\frac{s}{n}} + \frac{s(1 + d_W)\log(p|V|n)}{n} \lesssim \frac{s(1 + d_W)\log(p|V|n)}{n}$$

under $M_n \sqrt{s^2/n} \leq C$. ■

Lemma 3 (CIQGM, Event Ω_3). *For $u = (a, \tau, \varpi) \in \mathcal{U} := V \times \mathcal{T} \times \mathcal{W}$, define the function $g_u(\delta, X, W) = K_\varpi(W)\{\rho_\tau(X_a - Z^a(\beta_u + \delta)) - \rho_\tau(X_a - Z^a \beta_u)\}$, and the event*

$$\Omega_3 := \left\{ \sup_{u \in \mathcal{U}, 1/\sqrt{n} \leq \|\delta\|_{1, \varpi} \leq \sqrt{n}} \frac{|\mathbb{E}_n[g_u(\delta, X, W)] - \mathbb{E}[g_u(\delta, X, W) | X_{-a}, W]|}{\|\delta\|_{1, \varpi}} < t_3 \right\}.$$

Then, under Condition CI we have $P(\Omega_3) \geq 1 - \xi$ for any t_3 satisfying

$$t_3\sqrt{n} \geq 12 + 16\sqrt{2\log(64|V|p^2n^{3+2d_W}\log(n)L_\beta^{1+d_W/\kappa}M_n/\xi)}$$

Proof. We have that $\Omega_3^c := \{\max_{a \in V} A_a \geq t_3\sqrt{n}\}$ for

$$A_a := \sup_{(\tau, \varpi) \in \mathcal{T} \times \mathcal{W}, \underline{N} \leq \|\delta\|_1, \varpi \leq \bar{N}} \sqrt{n} \left| \frac{\mathbb{E}_n[g_u(\delta, X, W) - \mathbb{E}[g_u(\delta, X, W) \mid X_{-a}, W]]}{\|\delta\|_1, \varpi} \right|.$$

Therefore, for $\underline{N} = 1/\sqrt{n}$ and $\bar{N} = \sqrt{n}$ we have by Lemma 13 with $\rho = \kappa$, $L_\eta = L_\beta$, $\tilde{x} = Z^a$

$$\begin{aligned} P(\Omega_3^c) &= P(\max_{a \in V} A_a \geq t_3\sqrt{n}) \\ &\leq |V| \max_{a \in V} P(A_a \geq t_3\sqrt{n}) \\ &= |V| \max_{a \in V} \mathbb{E}_{X_{-a}, W} \{P(A_a \geq t_3\sqrt{n} \mid X_{-a}, W)\} \\ &\leq |V| \max_{a \in V} \mathbb{E}_{X_{-a}, W} \left\{ 8p|\hat{N}| \cdot |\hat{W}| \cdot |\hat{T}| \exp(-(t_3\sqrt{n}/4 - 3)^2/32) \right\} \\ &\leq \exp(-(t_3\sqrt{n}/4 - 3)^2/32) |V| 64pn^{1+d_W} \log(n) L_\beta \mathbb{E}_{X_{-a}} \left\{ \frac{\max_{i \leq n} \|Z_i^a\|_\infty^{1+d_W/\kappa}}{\underline{N}^{1+d_W}} \right\} \\ &\leq \xi \end{aligned}$$

by the choice of t_3 and noting that $M_n^{(1+d_W/\kappa)/q} \geq \mathbb{E}_{X_{-a}}[\max_{i \leq n} \|Z_i^a\|_\infty^{1+d_W/\kappa}]$, $1 + d_W/\kappa \leq q$ and $M_n \geq 1$. \blacksquare

Lemma 4 (CIQGM, Uniform Control of Approximation Error in Auxiliary Equation). *Under Condition CI, with probability $1 - o(1)$ uniformly over $u \in \mathcal{U}$ and $j \in [p]$ we have*

$$\mathbb{E}_n[K_\varpi(W)f_u^2\{Z_{-j}^a(\gamma_u^j - \bar{\gamma}_u^j)\}^2] \lesssim \underline{f}_u^2 P(\varpi) \{n^{-1}s \log(p|V|n)\}^{1/2}.$$

Proof. Define the class of functions $\mathcal{G} = \cup_{a \in V, j \in [p]} \mathcal{G}_{aj}$ with $\mathcal{G}_{aj} := \{Z_{-j}^a(\gamma_u^j - \bar{\gamma}_u^j) : \tau \in \mathcal{T}, \varpi \in \mathcal{W}\}$. Under Condition CI we have $\sup_{u \in \mathcal{U}} \|\bar{\gamma}_u^j\|_0 \leq Cs$, $\sup_{u \in \mathcal{U}, j \in [p]} \|\bar{\gamma}_u^j - \gamma_u^j\| \vee \frac{\|\bar{\gamma}_u^j - \gamma_u^j\|_1}{\sqrt{s}} \leq \{n^{-1}s \log(p|V|n)\}^{1/2}$. Without loss of generality we can set $\bar{\gamma}_{uk}^j = \gamma_{uk}^j$ for $k \in \text{support}(\bar{\gamma}_u^j)$. Letting $\mathcal{G}_{aj,T} := \{Z_{-j}^a(\gamma_u^j - \gamma_{u_T}^j) : \tau \in \mathcal{T}, \varpi \in \mathcal{W}\}$ for $T \subset \{1, \dots, p\}$, it follows that $\mathcal{G} \subset \cup_{a \in V, j \in [p]} \cup_{|T| \leq Cs} \mathcal{G}_{aj,T}$.

By Lemma 5, we have $\|\gamma_u^j - \gamma_{u'}^j\| \leq L_\gamma(\|u - u'\| + \|u - u'\|^{1/2})$ for each $a \in V$, $j \in [p]$. (Note that although $\bar{\gamma}_u^j$ might not be Lipschitz in u , however, for each T , $\gamma_{u_T}^j$ satisfies the same Lipschitz relation as γ_u^j , in fact $\|\bar{\gamma}_{u_T}^j - \gamma_{u_T}^j\| \leq \|\gamma_u^j - \gamma_{u'}^j\|$ by construction.) Therefore, for each T we have

$$\begin{aligned} &\|\{Z_{-j}^a(\bar{\gamma}_{u_T}^j - \gamma_u^j)\}^2 - \{Z_{-j}^a(\bar{\gamma}_{u_T}^j - \gamma_{u'}^j)\}^2\|_{Q,2} \\ &\leq \|Z_{-j}^a(\bar{\gamma}_{u_T}^j - \bar{\gamma}_{u_T}^j + \gamma_{u'}^j - \gamma_u^j)Z_{-j}^a(\bar{\gamma}_{u_T}^j - \gamma_u^j + \bar{\gamma}_{u_T}^j - \gamma_{u'}^j)\|_{Q,2} \\ &\leq \| \|Z_{-j}^a\|_\infty^2 \|_{Q,2} \|\bar{\gamma}_{u_T}^j - \bar{\gamma}_{u_T}^j + \gamma_{u'}^j - \gamma_u^j\|_1 \|\bar{\gamma}_{u_T}^j - \gamma_u^j + \bar{\gamma}_{u_T}^j - \gamma_{u'}^j\|_1 \\ &\leq 4 \| \|Z_{-j}^a\|_\infty^2 \|_{Q,2} \sup_{u \in \mathcal{U}} \|\bar{\gamma}_{u_T}^j - \gamma_u^j\|_1 \sqrt{2p} \|\gamma_u^j - \gamma_{u'}^j\| \\ &\leq \| \|Z_{-j}^a\|_\infty^2 \|_{Q,2} L'_\gamma (\|u - u'\| + \|u - u'\|^{1/2}). \end{aligned}$$

where $L'_\gamma = 4\{n^{-1}s^2 \log(p|V|n)\}^{1/2} \sqrt{2p} L_\gamma$. Thus, for the envelope $G = \max_{a \in V} \|Z^a\|_\infty^2 \sup_{u \in \mathcal{U}} \|\bar{\gamma}_u^j - \gamma_u^j\|_1^2$ that

$$\log N(\epsilon \|G\|_{Q,2}, \mathcal{G}, \|\cdot\|_{Q,2}) \leq Cs \log(|V|p) + \log N\left(\epsilon \frac{\sup_{u \in \mathcal{U}} \|\bar{\gamma}_u^j - \gamma_u^j\|_1^2}{L'_\gamma}, \mathcal{U}, d_\mathcal{U}\right) \leq Cs(1 + d_W)^2 \log(L'_\gamma n/\epsilon).$$

Next define the functions $\mathcal{W}_0 = \{K_\varpi(W)f_u^2 : u \in \mathcal{U}\}$, $\mathcal{W}_1 = \{P(\varpi)^{-1} : \varpi \in \mathcal{W}\}$ and $\mathcal{W}_2 = \{K_\varpi(W) : \varpi \in \mathcal{W}\}$. We have that \mathcal{W}_2 is VC class with VC index Cd_W and \mathcal{W}_1 is bounded by $\mu_{\mathcal{W}}^{-1}$ and covering number bounded by $(Cd_W/\{\mu_{\mathcal{W}}\epsilon\})^{1+d_W}$. Finally, since $|K_\varpi(W)f_u^2 - K_{\varpi'}(W)f_{u'}^2| \leq K_\varpi(W)K_{\varpi'}(W)|f_u^2 - f_{u'}^2| +$

$\bar{f}^2 |K_\varpi(W) - K_{\varpi'}(W)| \leq 2\bar{f}L_f \|u - u'\| + \bar{f}^2 |K_\varpi(W) - K_{\varpi'}(W)|$, we have $N(\epsilon, \mathcal{U}, |\cdot|) \leq (C(1+d_W)/\epsilon)^{1+d_W}$. Therefore, using standard bounds we have

$$\log N(\epsilon \|\mu_{\mathcal{W}}^{-1} G \bar{f}\|_{Q,2}, \mathcal{W}_0 \mathcal{W}_1 \mathcal{W}_2 \mathcal{G}, \|\cdot\|_{Q,2}) \lesssim s(1+d_W)^2 \log(L'_\gamma L_f n / \epsilon)$$

By Lemma 19 we have that with probability $1 - o(1)$ that

$$\begin{aligned} & \sup_{u \in \mathcal{U}, j \in [p]} |(\mathbb{E}_n - \mathbb{E})[f_u^2 \{Z_{-j}^a(\gamma_u^j - \bar{\gamma}_u^j)\}^2 / P(\varpi)]| \\ & \lesssim \sqrt{\frac{s(1+d_W)^2 \log(p|V|n) \sup_{u \in \mathcal{U}} \mathbb{E}[K_\varpi(W) f_u^4 \{Z_{-j}^a(\gamma_u^j - \bar{\gamma}_u^j)\}^4] / P(\varpi)^2}{n}} + \frac{s(1+d_W)^2 M_n^2 \mu_{\mathcal{W}}^{-1} \sup_{u \in \mathcal{U}} \|\bar{\gamma}_u^j - \gamma_u^j\|_1^2 \log(p|V|n)}{n} \\ & \lesssim \sqrt{\frac{s(1+d_W)^2 \log(p|V|n)}{\mu_{\mathcal{W}} n}} \frac{s \log(p|V|n)}{n} + \frac{(1+d_W)^2 M_n^2 s^2 \log(p|V|n)}{n \mu_{\mathcal{W}}} \frac{s \log(p|V|n)}{n} \\ & \lesssim \frac{s \log(p|V|n)}{n} \mu_{\mathcal{W}} \underline{f}_{\mathcal{U}} \left\{ \sqrt{\frac{s(1+d_W)^2 \log(p|V|n)}{\mu_{\mathcal{W}}^3 \underline{f}_{\mathcal{U}}^2 n}} + \frac{(1+d_W)^2 M_n^2 s^2 \log(p|V|n)}{n \mu_{\mathcal{W}} \underline{f}_{\mathcal{U}}} \right\} \\ & \lesssim \frac{s \log(p|V|n)}{n} \mu_{\mathcal{W}} \underline{f}_{\mathcal{U}} \{\delta_n^{1/2} + \delta_n^2\} \end{aligned}$$

here we used that $\mathbb{E}[f_u^4 \{Z^a \delta\}^4 | \varpi] \leq \bar{f}^4 \mathbb{E}[\{Z^a \delta\}^4 | \varpi] \leq C \|\delta\|^4$, $\|\bar{\gamma}_u^j - \gamma_u^j\| + s^{-1/2} \|\bar{\gamma}_u^j - \gamma_u^j\|_1 \leq \{n^{-1} s \log(p|V|n)\}^{1/2}$, $s(1+d_W)^2 \log(p|V|n) \leq \delta_n n f_{\mathcal{U}}^2 \mu_{\mathcal{W}}^3$ and $(1+d_W) M_n s \log^{1/2}(p|V|n) \leq \delta_n n^{1/2} \mu_{\mathcal{W}} \underline{f}_{\mathcal{U}}$ by Condition CI. Furthermore, by Condition CI, the result follows from $\mathbb{E}[f_u^2 \{Z_{-j}^a(\bar{\gamma}_u^j - \gamma_u^j)\}^2 | \varpi] \leq C \underline{f}_{\mathcal{U}}^2 \|\bar{\gamma}_u^j - \gamma_u^j\|^2 \leq C \underline{f}_{\mathcal{U}}^2 n^{-1} s \log(p|V|n)$. \blacksquare

Lemma 5. Under Condition CI, for $u = (a, \tau, \varpi) \in \mathcal{U}$ and $u' = (a, \tau', \varpi') \in \mathcal{U}$ we have that

$$\|\gamma_u^j - \gamma_{u'}^j\| \leq \frac{C'}{\underline{f}_{u'}^2 P(\varpi')} \{\bar{f}^2 \mathbb{E}[\{K_{\varpi'}(W) - K_\varpi(W)\}^2]^{1/2} + \mathbb{E}[K_\varpi(W) K_{\varpi'}(W) \{f_{u'}^2 - f_u^2\}^2]^{1/2}\}.$$

In particular, we have $\|\gamma_u^j - \gamma_{u'}^j\| \leq L_\gamma \{\|\varpi - \varpi'\|^{1/2} + \|u - u'\|\}$ for $L_\gamma = C\{L_f + L_K\} / \{\underline{f}_{\mathcal{U}}^2 \mu_{\mathcal{W}}\}$ under $\mathbb{E}[|K_\varpi(W) - K_{\varpi'}(W)|] \leq L_K \|\varpi - \varpi'\|$, $K_\varpi(W) K_{\varpi'}(W) |f_{u'} - f_u| \leq L_f \|u' - u\|$, and $f_u \leq \bar{f} \leq C$.

Proof. Let $u = (a, \tau, \varpi)$ and $u' = (a, \tau', \varpi')$. By Condition CI we have

$$\|\gamma_u^j - \gamma_{u'}^j\|^2 \leq C \mathbb{E}[\{Z_{-j}^a(\gamma_u^j - \gamma_{u'}^j)\}^2 | \varpi'] \leq \{C/P(\varpi')\} \mathbb{E}[K_{\varpi'}(W) \{Z_{-j}^a(\gamma_u^j - \gamma_{u'}^j)\}^2]$$

To bound the last term of the right hand side above, by the definition of $\underline{f}_{u'}$ and Cauchy-Schwarz's inequality we have

$$\begin{aligned} \underline{f}_{u'} \mathbb{E}[K_{\varpi'}(W) \{Z_{-j}^a(\gamma_u^j - \gamma_{u'}^j)\}^2] & \leq \mathbb{E}[K_{\varpi'}(W) f_{u'} \{Z_{-j}^a(\gamma_u^j - \gamma_{u'}^j)\}^2] \\ & \leq \{\mathbb{E}[K_{\varpi'}(W) f_{u'}^2 \{Z_{-j}^a(\gamma_u^j - \gamma_{u'}^j)\}^2] \mathbb{E}[K_{\varpi'}(W) \{Z_{-j}^a(\gamma_u^j - \gamma_{u'}^j)\}^2]\}^{1/2} \end{aligned}$$

so that $\mathbb{E}[K_{\varpi'}(W) \{Z_{-j}^a(\gamma_u^j - \gamma_{u'}^j)\}^2]^{1/2} \leq \{\mathbb{E}[K_{\varpi'}(W) f_{u'}^2 \{Z_{-j}^a(\gamma_u^j - \gamma_{u'}^j)\}^2]\}^{1/2} / \underline{f}_{u'}$. Therefore

$$\|\gamma_u^j - \gamma_{u'}^j\|^2 \leq \{1/\underline{f}_{u'}\}^2 \{C/P(\varpi')\} \mathbb{E}[K_{\varpi'}(W) f_{u'}^2 \{Z_{-j}^a(\gamma_u^j - \gamma_{u'}^j)\}^2]. \quad (\text{D.54})$$

We proceed to bound the last term. The optimality of γ_u^j and $\gamma_{u'}^j$ yields

$$\mathbb{E}[K_\varpi(W) f_u^2 Z_{-j}^a (Z_j^a - Z_{-j}^a \gamma_u^j)] = 0 \quad \text{and} \quad \mathbb{E}[K_{\varpi'}(W) f_{u'}^2 Z_{-j}^a (Z_j^a - Z_{-j}^a \gamma_{u'}^j)] = 0$$

Therefore, we have

$$\begin{aligned} \mathbb{E}[K_{\varpi'}(W) f_{u'}^2 \{Z_{-j}^a(\gamma_u^j - \gamma_{u'}^j)\} Z_{-j}^a] & = -\mathbb{E}[K_{\varpi'}(W) f_{u'}^2 \{Z_j^a - Z_{-j}^a \gamma_u^j\} Z_{-j}^a] \\ & = -\mathbb{E}[\{K_{\varpi'}(W) f_{u'}^2 - K_\varpi(W) f_u^2\} \{Z_j^a - Z_{-j}^a \gamma_u^j\} Z_{-j}^a] \end{aligned} \quad (\text{D.55})$$

Multiplying by $(\gamma_u^j - \gamma_{u'}^j)$ both sides of (D.55), we have

$$\begin{aligned} & \mathbb{E}[K_{\varpi'}(W)f_{u'}^2\{Z_{-j}^a(\gamma_u^j - \gamma_{u'}^j)\}^2] \\ & \leq \mathbb{E}[\{K_{\varpi'}(W)f_{u'}^2 - K_{\varpi}(W)f_u^2\}^2]^{1/2}\{\mathbb{E}[\{Z_j^a - Z_{-j}^a\gamma_u^j\}^2\{Z_{-j}^a(\gamma_u^j - \gamma_{u'}^j)\}^2]\}^{1/2} \\ & \leq \mathbb{E}[\{K_{\varpi'}(W)f_{u'}^2 - K_{\varpi}(W)f_u^2\}^2]^{1/2}C\|\gamma_u^j - \gamma_{u'}^j\| \end{aligned}$$

by the fourth moment assumption in Condition CI. By Condition CI, $f_u, f_{u'} \leq \bar{f}$, and it follows that

$$|K_{\varpi}(W)f_u^2 - K_{\varpi'}(W)f_{u'}^2| \leq K_{\varpi}(W)K_{\varpi'}(W)|f_u^2 - f_{u'}^2| + \bar{f}^2|K_{\varpi}(W) - K_{\varpi'}(W)| \quad (\text{D.56})$$

From (D.54) we obtain

$$\|\gamma_u^j - \gamma_{u'}^j\| \leq \frac{C'}{\underline{f}_{u'}^2 P(\varpi')} \{\bar{f}^2 \mathbb{E}[\{K_{\varpi'}(W) - K_{\varpi}(W)\}^2]^{1/2} + \mathbb{E}[K_{\varpi}(W)K_{\varpi'}(W)\{f_{u'}^2 - f_u^2\}^2]^{1/2}\}.$$

■

Lemma 6. *Let $\mathcal{U} = V \times \mathcal{T} \times \mathcal{W}$. Under Condition CI, for $m = 1, 2$, we have*

$$\mathbb{E} \left[\sup_{u \in \mathcal{U}, \|\theta\|_0 \leq k, \|\theta\|=1} |(\mathbb{E}_n - \mathbb{E})[K_{\varpi}(W)f_u^m(Z^a\theta)^2]| \right] \lesssim C\delta_n \sup_{u \in \mathcal{U}, \|\theta\|_0 \leq k, \|\theta\|=1} \{\mathbb{E}[K_{\varpi}(W)f_u^m(Z^a\theta)^2]\}^{1/2}$$

where $\delta_n = M_n \sqrt{k(1+d_W)C \log(p|V|n) \log(1+k) \sqrt{\log n/n}}$. Moreover, under Condition CI, $\delta_n = o(\mu_{\mathcal{W}})$.

Proof. By symmetrization we have

$$\mathbb{E} \left[\sup_{u \in \mathcal{U}, \|\theta\|_0 \leq k, \|\theta\|=1} |(\mathbb{E}_n - \mathbb{E})[K_{\varpi}(W)f_u^m(Z^a\theta)^2]| \right] \leq 2\mathbb{E} \left[\sup_{u \in \mathcal{U}, \|\theta\|_0 \leq k, \|\theta\|=1} |\mathbb{E}_n[\varepsilon K_{\varpi}(W)f_u^m(Z^a\theta)^2]| \right]$$

where ε_i are i.i.d. Rademacher random variables. We have that $|K_{\varpi}(W)f_u^m - K_{\varpi'}(W)f_{u'}^m| \leq K_{\varpi}(W)K_{\varpi'}(W)|f_u - f_{u'}|(1+2\bar{f}) + \bar{f}^m|K_{\varpi}(W) - K_{\varpi'}(W)|$ for $m = 1, 2$ where u and u' have the same $a \in V$. However, conditional on $\{(W_i, X_i), i = 1, \dots, n\}$, $\{K_{\varpi}(W_i) : i = 1, \dots, n, \varpi \in \mathcal{W}\}$ induces at most n^{d_W} different sequences by Corollary 2.6.3 in [100]. This induces (at most) n^{d_W} partitions of \mathcal{W} such that $K_{\varpi}(W) = K_{\varpi'}(W)$ for any ϖ, ϖ' in the same partition given the conditioning. Thus, for such suitable ϖ' we have $|K_{\varpi}(W)f_u^m - K_{\varpi'}(W)f_{u'}^m| \leq K_{\varpi}(W)|f_u - f_{u'}|(1+2\bar{f})$ for $m = 1, 2$. (Thus it suffices to create a net for each partition.) We can take a cover $\hat{\mathcal{U}}$ of $V \times \mathcal{T} \times \mathcal{W}$ such that $\|u - u'\| \leq \{L_f(1+2\bar{f})nk \max_{i \leq n} \|Z_i^a\|_{\infty}^2\}^{-1}$ so that $|f_u - f_{u'}|(Z^a\theta)^2 \leq |f_u - f_{u'}|\|Z^a\|_{\infty}^2\|\theta\|_1^2 \leq |f_u - f_{u'}|\|Z^a\|_{\infty}^2 k \|\theta\|^2$ which implies

$$\left| \sup_{u \in \mathcal{U}, \|\theta\|_0 \leq k, \|\theta\|=1} |\mathbb{E}_n[\varepsilon K_{\varpi}(W)f_u^m(Z^a\theta)^2]| - \sup_{u \in \hat{\mathcal{U}}, \|\theta\|_0 \leq k, \|\theta\|=1} |\mathbb{E}_n[\varepsilon K_{\varpi}(W)f_u^m(Z^a\theta)^2]| \right| \leq n^{-1}$$

Consequently

$$\mathbb{E} \left[\sup_{u \in \mathcal{U}, \|\theta\|_0 \leq k, \|\theta\|=1} |\mathbb{E}_n[\varepsilon K_{\varpi}(W)f_u^m(Z^a\theta)^2]| \right] \leq \mathbb{E} \left[\sup_{u \in \hat{\mathcal{U}}, \|\theta\|_0 \leq k, \|\theta\|=1} |\mathbb{E}_n[\varepsilon K_{\varpi}(W)f_u^m(Z^a\theta)^2]| \right] + \frac{1}{n}$$

where $|\hat{\mathcal{U}}| \leq |V|n^{d_W}\{L_f(1+2\bar{f})nk \max_{i \leq n} \|Z_i^a\|_{\infty}^2\}^{(1+d_W)}$.

By Lemma 18 with $K = K(W, X) = (1 + \bar{f}^2) \sup_{a \in V} \max_{i \leq n} \|Z_i^a\|_{\infty}$ and

$$\begin{aligned} \delta_n(W, X) &:= \bar{C}K(W, X)\sqrt{k} \left(\sqrt{\log |\hat{\mathcal{U}}|} + \sqrt{1 + \log p} + \log k \sqrt{\log(p \vee n)} \sqrt{\log n} \right) / \sqrt{n} \\ &\lesssim K(W, X) \sqrt{k(1+d_W)C \log(p|V|nK(W, X)) \log(1+k) \sqrt{\log n}/\sqrt{n}} \end{aligned}$$

so that conditional on (W, X) we have

$$\mathbb{E} \left[\sup_{u \in \hat{\mathcal{U}}, \|\theta\|_0 \leq k, \|\theta\|=1} |\mathbb{E}_n[\varepsilon K_\varpi(W) f_u^m(Z^a \theta)^2]| \right] \lesssim \delta_n(W, X) \sup_{u \in \hat{\mathcal{U}}, \|\theta\|_0 \leq k, \|\theta\|=1} \sqrt{\mathbb{E}_n[K_\varpi(W) f_u^m(Z^a \theta)^2]}$$

Therefore,

$$\begin{aligned} & \mathbb{E} \left[\sup_{u \in \mathcal{U}, \|\theta\|_0 \leq k, \|\theta\|=1} |\mathbb{E}_n[\varepsilon K_\varpi(W) f_u^m(Z^a \theta)^2]| \right] \\ & \leq \mathbb{E}_{W, X} \left[\delta_n(W, X) \sup_{u \in \hat{\mathcal{U}}, \|\theta\|_0 \leq k, \|\theta\|=1} \sqrt{\mathbb{E}_n[K_\varpi(W) f_u^m(Z^a \theta)^2]} \right] + \frac{1}{n} \\ & \leq \mathbb{E}_{W, X} [\delta_n^2(W, X)] + \mathbb{E}_{W, X} [\delta_n^2(W, X)]^{1/2} \sup_{u \in \hat{\mathcal{U}}, \|\theta\|_0 \leq k, \|\theta\|=1} \mathbb{E}[K_\varpi(W) f_u^m(Z^a \theta)^2]^{1/2} + \frac{1}{n} \end{aligned}$$

Note that for a random variable $A \geq 1$, we have that $\mathbb{E}[A^2 \sqrt{\log(CA)}] \leq \mathbb{E}[A^2] \sqrt{\log(C)} + \mathbb{E}[A^2 \sqrt{\log(A)}] \leq \mathbb{E}[A^2] \sqrt{\log(C)} + \mathbb{E}[A^{2+1/4}]$. Therefore, under Condition CI, since $q \geq 2 + 1/4$ in the definition of M_n , we have

$$\mathbb{E}_{W, X} [\delta_n^2(W, X)]^{1/2} \lesssim M_n \sqrt{k(1 + d_W) C \log(p|V|n) \log(1 + k) \sqrt{\log n} / \sqrt{n}}.$$

The results follows by setting $\delta_n = \mathbb{E}_{W, X} [\delta_n^2(W, X)]^{1/2}$. \blacksquare

APPENDIX E. RESULTS FOR PREDICTION QUANTILE GRAPHICAL MODELS

In the analysis of PQGM we also use the following event for some sequence $(K_u)_{u \in \mathcal{U}}$

$$\Omega_4 = \{K_u \hat{\sigma}_{a\varpi j}^X \geq \mathbb{E}_n[\mathbb{E}[K_\varpi(W)(\tau - 1\{X_a \leq X'_{-a}\beta_u + r_u\})X_{-a}|X_j, W]], u \in \mathcal{U}, j \in V \setminus \{a\}\}. \quad (\text{E.57})$$

Lemma 7 (Rate for PQGM). *Suppose that $\Omega_1, \Omega_2, \Omega_3$ and Ω_4 hold. Further assume $2^{\frac{1+1/\varepsilon}{1-1/c}} \|\beta_u\|_{1, \varpi} + \frac{1}{\lambda_u(1-1/c)} \bar{R}_{u\xi} \leq \sqrt{n}$ for all $u \in \mathcal{U}$, and (F.63) holds for all $\delta \in A_u := \Delta_{\varpi, 2c} \cup \{v : \|v\|_{1, \varpi} \leq 2c\bar{R}_{u\xi}/\lambda_u\}$, $\bar{q}_{A_u}/4 \geq (\sqrt{f} + 1)\|r_u\|_{n, \varpi} + [\lambda_u + t_3 + K_u] \frac{3c\sqrt{s}}{\kappa_{u, 2c}}$ and $\bar{q}_{A_u} \geq \{2c(1 + \frac{t_3 + K_u}{\lambda_u})\} \bar{R}_{u\xi}^{1/2}$. Then uniformly over all $u = (a, \tau, \varpi) \in \mathcal{U} := V \times \mathcal{T} \times \mathcal{W}$, the $\|\cdot\|_{1, \varpi}$ -penalized estimator $\hat{\beta}_u$ satisfies*

$$\begin{aligned} \|\sqrt{f_u} X'_{-a}(\hat{\beta}_u - \beta_u)\|_{n, \varpi} & \leq \sqrt{8c \left(1 + \frac{t_3}{\lambda_u}\right) \bar{R}_{u\xi} + (\bar{f}^{1/2} + 1)\|r_u\|_{n, \varpi} + [\lambda_u + t_3 + K_u] \frac{3c\sqrt{s}}{\kappa_{u, 2c}}} \\ \|\hat{\beta}_u - \beta_u\|_{1, \varpi} & \leq (1 + 2c)\sqrt{s} \|\sqrt{f_u} X'_{-a} \delta_u\|_{n, \varpi} / \kappa_{u, 2c} + \frac{2c}{\lambda_u} \bar{R}_{u\xi} \end{aligned}$$

Proof of Lemma 7. The proof proceeds similarly to the proof of Lemma 1 by defining

$$\begin{aligned} \hat{R}_u(\beta) & = \mathbb{E}_n[K_\varpi(W) \{\rho_u(X_a - X'_{-a}\beta) - \rho_u(X_a - X'_{-a}\beta_u - r_u)\}] \\ & \quad - \mathbb{E}_n[K_\varpi(W) \{(\tau - 1\{X_a \leq X'_{-a}\beta_u + r_u\})(X'_{-a}\beta - X'_{-a}\beta_u - r_u)\}]. \end{aligned}$$

The same argument yields $\delta_u = \hat{\beta}_u - \beta_u \in A_u := \Delta_{\varpi, 2c} \cup \{v : \|v\|_{1, \varpi} \leq 2c\bar{R}_{u\xi}/\lambda_u\}$ under $\Omega_1 \cap \Omega_2$. (Similarly we also have $\|\delta_u\|_{1, \varpi} \leq \sqrt{n}$.) Furthermore, under $\Omega_1 \cap \Omega_2 \cap \Omega_3$ we have that (D.53) also holds which implies

$$\mathbb{E}_n[\mathbb{E}[K_\varpi(W) \{\rho_u(X_a - X'_{-a}(\beta_u + \delta_u)) - \rho_u(X_a - X'_{-a}\beta_u)\} | X_{-a}, W] \leq (\lambda_u + t_3) \|\delta_u\|_{1, \varpi}$$

Since the conditions of Lemma 12 hold we have

$$\begin{aligned} \mathbb{E}_n[\mathbb{E}[K_\varpi(W) \{\rho_\tau(X_a - X'_{-a}(\beta_u + \delta_u)) - \rho_\tau(X_a - X'_{-a}\beta_u)\} | X_{-a}, W] \\ \geq -(\sqrt{f} + 1)\|r_u\|_{n, \varpi} \|\sqrt{f_u} X'_{-a} \delta_u\|_{n, \varpi} - K_u \|\delta_u\|_{1, \varpi} \\ + \frac{\|\sqrt{f_u} X'_{-a} \delta_u\|_{n, \varpi}^2}{4} \wedge \bar{q}_{A_u} \|\sqrt{f_u} X'_{-a} \delta_u\|_{n, \varpi} \end{aligned}$$

where K_u is given in Ω_4 which accounts for the misspecification the conditional quantile condition. Therefore, we have

$$\begin{aligned} \frac{\|\sqrt{f_u}X'_{-a}\delta_u\|_{n,\varpi}^2}{4} \wedge \bar{q}_{A_u}\|\sqrt{f_u}X'_{-a}\delta_u\|_{n,\varpi} &\leq (\bar{f}^{1/2} + 1)\|r_u\|_{n,\varpi}\|\sqrt{f_u}X'_{-a}\delta_u\|_{n,\varpi} + (\lambda_u + t_3 + K_u)\|\delta_u\|_{1,\varpi} \\ &\leq \{(\bar{f}^{1/2} + 1)\|r_u\|_{n,\varpi} + \frac{3\mathbf{c}\sqrt{s}}{\kappa_{u,2\mathbf{c}}}(\lambda_u + t_3 + K_u)\}\|\sqrt{f_u}X'_{-a}\delta_u\|_{n,\varpi} \\ &\quad + (\lambda_u + t_3 + K_u)\frac{2\mathbf{c}}{\lambda_u}\bar{R}_{u\xi} \end{aligned}$$

The result then follows with the same argument under the current assumptions that account for K_u . \blacksquare

Lemma 8 (PQGM, Event Ω_1). *Under Condition P, we have*

$$\mathbb{P}\left(\sup_{u \in \mathcal{U}, j \in [d]} \frac{|\mathbb{E}_n[K_\varpi(W)(\tau - 1\{X_a \leq X'_{-a}\beta_u + r_u\})X_{-a,j}]|}{\hat{\sigma}_{a\varpi j}^X} > t\right) \leq 8|V|(\frac{ne}{d_W})^{2d_W} \exp\left(-\left\{\frac{t/(1 + \bar{\delta}_n)}{2(1 + 1/16)}\right\}^2\right)$$

where $t \geq 4 \sup_{u \in \mathcal{U}} \{\mathbb{E}[K_\varpi(W)(\tau - 1\{X_a \leq X'_{-a}\beta_u + r_u\})^2 X_{-a,j}^2]\}^{1/2}$ and $\bar{\delta}_n = o(1)$. In particular, the RHS is less than ξ if $t \geq 2(1 + \bar{\delta}_n)(1 + 1/16)\sqrt{\log(8|V|\{ne/d_W\}^{2d_W}/\xi)}$.

Proof. Set $\sigma_{a\varpi j}^X := \mathbb{E}[K_\varpi(W)X_{-a,j}^2]^{1/2}$. We have that for any $\bar{\delta}_n \rightarrow 0$

$$\begin{aligned} &\mathbb{P}(\lambda_0 \leq \sup_{u \in \mathcal{U}, j \in [d]} |\mathbb{E}_n[K_\varpi(W)(\tau - 1\{X_a \leq X'_{-a}\beta_u\})X_{-a,j}]|/\hat{\sigma}_{a\varpi j}^X) \\ &\leq \mathbb{P}(\lambda_0 \leq (1 + \bar{\delta}_n) \sup_{u \in \mathcal{U}, j \in [d]} |\mathbb{E}_n[K_\varpi(W)(\tau - 1\{X_a \leq X'_{-a}\beta_u\})X_{-a,j}]|/\sigma_{a\varpi j}^X) \\ &\quad + \mathbb{P}(\sup_{u \in \mathcal{U}, j \in [d]} \sigma_{a\varpi j}^X/\hat{\sigma}_{a\varpi j}^X \geq (1 + \bar{\delta}_n)) \end{aligned} \tag{E.58}$$

To bound the last term in (E.58), note that under Condition P, $c\mu_{\mathcal{W}} \leq (\sigma_{a\varpi j}^X)^2 \leq C$ and \mathcal{W} is a VC class of set with VC dimension d_W . Therefore, by Lemma 19 we have that with probability $1 - o(1)$

$$\sup_{u \in \mathcal{U}, j \in [d]} (\mathbb{E}_n - \mathbb{E})[K_\varpi(W)X_{-a,j}^2] \lesssim \sqrt{\frac{(1 + d_W)\log(|V|M_n/\sigma_1)}{n}} + \frac{(1 + d_W)M_n^2 \log(|V|M_n/\sigma_1)}{n} \tag{E.59}$$

for $\sigma_1^2 = \max_{u \in \mathcal{U}, j \in [d]} \mathbb{E}[K_\varpi(W)X_{-a,j}^2] \leq \max_{j \in V} \mathbb{E}[X_{-a,j}^2] \leq C$ and envelope $F = \|X\|_\infty^2$ so that $\|F\|_{P,2} \leq \|\max_{i \leq n} F_i\|_{P,2} \leq M_n^2$. Thus for $\bar{\delta}_n \rightarrow 0$, provided $(1 + d_W)M_n^2 \log(|V|n) = o(n^{1/2})$ and $(1 + d_W)\log(|V|n) = o(n\bar{\delta}_n^2\mu_{\mathcal{W}}^2)$, so that the RHS of (E.59) is $o(\bar{\delta}_n\mu_{\mathcal{W}})$, we have

$$\mathbb{P}\left(\frac{1}{1 + \bar{\delta}_n} \leq \frac{\sigma_{a\varpi j}^X}{\hat{\sigma}_{a\varpi j}^X} \leq (1 + \bar{\delta}_n), \text{ for all } u \in \mathcal{U}, j \in [d]\right) = 1 - o(1). \tag{E.60}$$

Now we bound the first term of the RHS of (E.58). and let $\sigma_2^2 = \sup_{u \in \mathcal{U}, j \in [d]} \text{Var}(K_\varpi(W)(\tau - 1\{X_a \leq X'_{-a}\beta_u\})X_{-a,j}/\sigma_{a\varpi j}^X) \leq 1$. By symmetrization (adapting Lemma 2.3.7 in [101] to replace the ‘‘arbitrary’’ factor 2 with $1 + \bar{\delta}_n$), for $\delta := 1/(2(1 + n\sigma_2^2/t^2)) < 1/2$ we have

$$\begin{aligned} (*) &:= \mathbb{P}(\sup_{u \in \mathcal{U}, j \in [d]} |\sum_{i=1}^n K_\varpi(W_i)(\tau - 1\{X_{ia} \leq X'_{i,-a}\beta_u\})X_{i,-aj}/\sigma_{a\varpi j}^X| \geq t) \\ &\leq 2\mathbb{P}(\sup_{u \in \mathcal{U}, j \in [d]} |\sum_{i=1}^n \varepsilon_i K_\varpi(W_i)(\tau - 1\{X_{ia} \leq X'_{i,-a}\beta_u\})X_{i,-aj}/\sigma_{a\varpi j}^X| \geq t\delta) \\ &\leq 2\mathbb{P}\left(\sup_{u \in \mathcal{U}, j \in [d]} \frac{|\sum_{i=1}^n \varepsilon_i K_\varpi(W_i)(\tau - 1\{X_{ia} \leq X'_{i,-a}\beta_u\})X_{i,-aj}|}{\hat{\sigma}_{a\varpi j}^X} \geq t\delta/(1 + \bar{\delta}_n)\right) + o(1) \end{aligned}$$

where $\varepsilon_i, i = 1, \dots, n$, are Rademacher random variables independent of the data, and the last inequality follows from (E.60).

Therefore, by the union bound and symmetry, and iterated expectations we have

$$(*) \leq 4|V| \max_{j \in [d]} \mathbb{E}_{W, X} \left[\mathbb{P}_\varepsilon \left(\sup_{u \in \mathcal{U}} \frac{|\sum_{i=1}^n \varepsilon_i K_\varpi(W_i)(\tau - 1\{X_{ia} \leq X'_{i,-a}\beta_u\})X_{i,-aj}|}{\hat{\sigma}_{a\varpi j}^X} \geq t\delta/(1 + \bar{\delta}_n) \mid W, X \right) \right]$$

Next we use that $\{\varpi \in \mathcal{W}\}$ is a VC class of sets with VC dimension bounded by d_W and $\{1\{X_a \leq X'_{-a}\beta_u\} : (\tau, \varpi) \in \mathcal{T} \times \mathcal{W}\}$ is a VC class of sets with VC dimension bounded by $1 + d_W$. By Corollary 2.6.3 in [100], we have that conditionally on $(W_i, X_i)_{i=1}^n$, the set of (binary) sequences $\{(K_\varpi(W_i))_{i=1, \dots, n} : \varpi \in \mathcal{W}\}$ has at most $\sum_{j=0}^{d_W-1} \binom{n}{j}$ different values. Similarly, $\{1\{X_{ia} \leq X'_{i,-a}\beta_u\}\}_{i=1, \dots, n} : u \in \mathcal{U}\}$ assumes at most $\sum_{j=0}^{d_W} \binom{n}{j}$ different values. Assuming that $n \geq d_W$, we have $\sum_{j=0}^k \binom{n}{j} \leq \{ne/k\}^k$ and

$$\begin{aligned} & \mathbb{P}_\varepsilon \left(\sup_{u \in \mathcal{U}} \frac{|\sum_{i=1}^n \varepsilon_i K_\varpi(W_i)(\tau - 1\{X_{ia} \leq X'_{i,-a}\beta_u\})X_{i,-aj}|}{\hat{\sigma}_{a\varpi j}^X} \geq t\delta/(1 + \bar{\delta}_n) \mid W, X \right) \\ & \leq \left\{ \frac{ne}{d_W - 1} \right\}^{d_W - 1} \left\{ \frac{ne}{d_W} \right\}^{d_W} \sup_{u \in \mathcal{U}} \mathbb{P}_\varepsilon \left(\sup_{\bar{\tau} \in \mathcal{T}} \frac{|\sum_{i=1}^n \varepsilon_i K_\varpi(W_i)(\bar{\tau} - 1\{X_{ia} \leq X'_{i,-a}\beta_u\})X_{i,-aj}|}{\hat{\sigma}_{a\varpi j}^X} \geq t\delta/(1 + \bar{\delta}_n) \mid W, X \right) \\ & \leq \{ne/d_W\}^{2d_W} \sup_{u \in \mathcal{U}, \bar{\tau} \in [\underline{\tau}, \bar{\tau}]} \mathbb{P}_\varepsilon \left(\frac{|\sum_{i=1}^n \varepsilon_i K_\varpi(W_i)(\bar{\tau} - 1\{X_{ia} \leq X'_{i,-a}\beta_u\})X_{i,-aj}|}{\hat{\sigma}_{a\varpi j}^X} \geq t\delta/(1 + \bar{\delta}_n) \mid W, X \right) \\ & \leq 2\{ne/d_W\}^{2d_W} \exp(-\{t\delta/[1 + \bar{\delta}_n]\}^2) \end{aligned}$$

here we used that the expression is linear in τ and so it is maximized at the extremes. Combining the bounds in the last two displayed equations we have

$$(*) \leq 8|V|\{ne/d_W\}^{2d_W} \exp(-\{t\delta/[1 + \bar{\delta}_n]\}^2).$$

Therefore, setting $\lambda_0 = ct/n$ where $t \geq 4\sqrt{n}\sigma_2$ and $t \geq 2(1 + \bar{\delta}_n)(1 + 1/16)\sqrt{2 \log(8p|V|\{ne/d_W\}^{2d_W}/\xi)}$. (Note that $t \geq 4\sqrt{n}\sigma_2$ implies that $\delta \geq 1/\{2(1 + 1/16)\}$.) \blacksquare

Lemma 9 (PQGM, Event Ω_2). *Under Condition P we have*

$$\mathbb{P} \left(\sup_{u \in \mathcal{U}} \hat{R}_u(\bar{\beta}_u) \leq C\{1 + \bar{f}\}\{n^{-1}s(1 + d_W) \log(|V|n)\} \right) = 1 - o(1).$$

Proof of Lemma 9. We have that $\hat{R}_u(\bar{\beta}_u) \geq 0$ by convexity of ρ_τ . Let $\varepsilon_{iu} = X_{ia} - X_{i,-a}\bar{\beta}_u - r_{iu}$ where $\|\bar{\beta}_u\|_0 \leq s$ and $r_{iu} = X'_{-a}(\beta_u - \bar{\beta}_u)$. By Knight's identity (F.64), $\hat{R}_u(\bar{\beta}_u) = -\mathbb{E}_n[K_\varpi(W)r_u \int_0^1 1\{\varepsilon_{iu} \leq -tr_u\} - 1\{\varepsilon_{iu} \leq 0\} dt] \geq 0$.

$$\begin{aligned} \mathbb{E}_n[\mathbb{E}[\hat{R}_u(\bar{\beta}_u)]] &= \mathbb{E}_n[K_\varpi(W)r_u \int_0^1 F_{X_a|X_{-a}, \varpi}(X'_{-a}\bar{\beta}_u + (1-t)r_u) - F_{X_a|X_{-a}, \varpi}(X'_{-a}\bar{\beta}_u + r_u) dt] \\ &\leq \mathbb{E}_n[K_\varpi(W)r_u \int_0^1 \bar{f}tr_u dt] \leq \bar{f}[\|r_u\|_{n, \varpi}^2]/2 \leq C\bar{f}s/n. \end{aligned}$$

Thus, by Markov's inequality we have $\inf_{u \in \mathcal{U}} \mathbb{P}(\hat{R}_u(\bar{\beta}_u) \leq C\bar{f}s/n) \geq 1/2$.

Define $z_{iu} := -\int_0^1 1\{\varepsilon_{iu} \leq -tr_{iu}\} - 1\{\varepsilon_{iu} \leq 0\} dt$, so that $\hat{R}_u(\bar{\beta}_u) = \mathbb{E}_n[K_\varpi(W)r_u z_u]$ with $|z_{iu}| \leq 1$. By Lemma 2.3.7 in [101] (note that the Lemma does not require zero mean stochastic processes), for $t \geq 2C\bar{f}s/n$ we have

$$\frac{1}{2} \mathbb{P} \left(\sup_{u \in \mathcal{U}} |\mathbb{E}_n[K_\varpi(W)r_u z_u]| \geq t \right) \leq 2 \mathbb{P} \left(\sup_{u \in \mathcal{U}} |\mathbb{E}_n[\varepsilon K_\varpi(W)r_u z_u]| > t/4 \right)$$

where $\varepsilon_i, i = 1, \dots, n$, are Rademacher random variables independent of the data.

Consider the class of functions $\mathcal{F} = \{-K_\varpi(W)r_u(1\{\varepsilon_{iu} \leq -B_i r_{iu}\} - 1\{\varepsilon_{iu} \leq 0\}) : u \in \mathcal{U}\}$ where $B_i \sim \text{Uniform}(0, 1)$ independent of $(X_i, W_i)_{i=1}^n$. It follows that $K_\varpi(W)r_u z_u = \mathbb{E}[-K_\varpi(W)r_u(1\{\varepsilon_{iu} \leq -B_i r_{iu}\} - 1\{\varepsilon_{iu} \leq 0\})|X_i, W_i]$ where the expectation is taken over B_i only. Thus we will bound the

entropy of $\bar{\mathcal{F}} = \{E[f|X, W] : f \in \mathcal{F}\}$ via Lemma 25. Note that $\mathcal{R} := \{r_u = X'_{-a}\beta_u - X'_{-a}\bar{\beta}_u : u \in \mathcal{U}\}$ where $\mathcal{G} := \{X'_{-a}\bar{\beta}_u : u \in \mathcal{U}\}$ is contained in the union of at most $|V|\binom{p}{s}$ VC-classes of dimension Cs and $\mathcal{H} := \{X'_{-a}\beta_u : u \in \mathcal{U}\}$ is a VC-class of functions of dimension $(1 + d_W)$ by Condition P. Finally note that $\mathcal{E} := \{\varepsilon_{iu} : u \in \mathcal{U}\} \subset \{X_{ia} : a \in V\} - \mathcal{G} - \mathcal{R}$.

Therefore, we have

$$\begin{aligned} \sup_Q \log N(\epsilon \|\bar{F}\|_{Q,2}, \bar{\mathcal{F}}, \|\cdot\|_{Q,2}) &\leq \sup_Q \log N((\epsilon/4)^2 \|F\|_{Q,2}, \mathcal{F}, \|\cdot\|_{Q,2}) \\ &\leq \sup_Q \log N(\tfrac{1}{8}(\epsilon^2/16), \mathcal{W}, \|\cdot\|_{Q,2}) \\ &\quad + \sup_Q \log N(\tfrac{1}{8}(\epsilon^2/16) \|F\|_{Q,2}, \mathcal{R}, \|\cdot\|_{Q,2}) \\ &\quad + \sup_Q \log N(\tfrac{1}{8}(\epsilon^2/16), 1\{\mathcal{E} + \{B\}\mathcal{R} \leq 0\} - 1\{\mathcal{E} \leq 0\}, \|\cdot\|_{Q,2}) \end{aligned}$$

By Lemma 19 with envelope $\bar{F} = \|X\|_\infty \sup_{u \in \mathcal{U}} \|\beta_u - \bar{\beta}_u\|_1$, and $(\sigma_{\max}^r)^2 = \sup_{u \in \mathcal{U}} E[K_\varpi(W)r_u^2] \lesssim s/n$ by Condition P, we have that with probability $1 - o(1)$

$$\sup_{u \in \mathcal{U}} |\mathbb{E}_n[\varepsilon K_\varpi(W)r_u z_u]| \lesssim \sqrt{\frac{s(1+d_W)\log(|V|n)}{n}} \sqrt{\frac{s}{n}} + \frac{M_n \sqrt{s^2/n} \log(|V|n)}{n} \lesssim \frac{s(1+d_W)\log(|V|n)}{n}$$

under $M_n \sqrt{s^2/n} \leq C$. ■

Lemma 10 (PQGM, Event Ω_3). *Under Condition P, for $u = (a, \tau, \varpi) \in V \times \mathcal{T} \times \mathcal{W}$, define $g_u(\delta, X, W) = K_\varpi(W)\{\rho_\tau(X_a - X'_{-a}(\beta_u + \delta)) - \rho_\tau(X_a - X'_{-a}\beta_u)\}$, and*

$$\Omega_3 := \left\{ \sup_{u \in \mathcal{U}, 1/\sqrt{n} \leq \|\delta\|_{1,\varpi} \leq \sqrt{n}} \frac{|\mathbb{E}_n[g_u(\delta, X, W)] - E[g_u(\delta, X, W)|X_{-a}, W]|}{\|\delta\|_{1,\varpi}} < t_3 \right\}.$$

Then, under Condition CI we have $P(\Omega_3) \geq 1 - \xi$ for any

$$t_3 \sqrt{n} \geq 12 + 16 \sqrt{2 \log(64|V|^2 n^{1+d_W} \log(n)(L_\beta M_n \sqrt{n})^{1+d_W/\kappa}/\xi)}$$

Proof. We have that $\Omega_3^c := \{\max_{a \in V} A_a \geq t_3 \sqrt{n}\}$ for

$$A_a := \sup_{(\tau, \varpi) \in \mathcal{T} \times \mathcal{W}, \underline{N} \leq \|\delta\|_{1,\varpi} \leq \bar{N}} \sqrt{n} \left| \frac{\mathbb{E}_n[g_u(\delta, X, W)] - E[g_u(\delta, X, W)|X_{-a}, W]}{\|\delta\|_{1,\varpi}} \right|.$$

We will apply Lemma 13 with $\rho = \kappa$, $L_\eta = L_\beta$, $\tilde{x} = X_{-a}$ (so we take $p = |V|$), $\underline{N} = 1/\sqrt{n}$ and $\bar{N} = \sqrt{n}$. Therefore, we have by Lemma 13 and the union bound

$$\begin{aligned} P(\Omega_3^c) &= P(\max_{a \in V} A_a \geq t_3 \sqrt{n}) \\ &\leq |V| \max_{a \in V} P(A_a \geq t_3 \sqrt{n}) \\ &= |V| \max_{a \in V} E_{X_{-a}, W} \{P(A_a \geq t_3 \sqrt{n} \mid X_{-a}, W)\} \\ &\leq |V| \max_{a \in V} E_{X_{-a}, W} \left\{ 8|V| |\hat{\mathcal{N}}| \cdot |\hat{\mathcal{W}}| \cdot |\hat{\mathcal{T}}| \exp(-(t_3 \sqrt{n}/4 - 3)^2/32) \right\} \\ &\leq C|V|^2 n^{1+d_W} \log(n) L_\beta^{1+d_W/\kappa} E_{X_{-a}} \left\{ \frac{\max_{i \leq n} \|X_{i,-a}\|_\infty^{1+d_W/\kappa}}{\underline{N}^{1+d_W/\kappa}} \right\} \exp(-(t_3 \sqrt{n}/4 - 3)^2/32) \\ &\leq \xi \end{aligned}$$

where the last step follows by the choice of t_3 . ■

Lemma 11 (PQGM, Event Ω_4). *Under Condition P, and setting $K_u = C \sqrt{\frac{(1+d_W)\log(|V|n)}{n}}$, we have that $P(\Omega_4) = 1 - o(1)$.*

Proof. First note that by Lemma 19 we have that with probability $1 - o(1)$

$$\sup_{\varpi \in \mathcal{W}, j \in V} (\mathbb{E}_n - \mathbb{E})[K_\varpi(W)X_{-a,j}^2] \lesssim \sqrt{\frac{(1+d_W)\log(|V|M_n/\sigma)}{n}} + \frac{(1+d_W)M_n^2\log(|V|M_n/\sigma)}{n}$$

and $\sigma_{a\varpi j}^X \geq cP(\varpi)$. Under $(1+d_W)\log(|V|M_n/\sigma) \leq \delta_n^2\mu_{\mathcal{W}}^2$ and $(1+d_W)M_n^2\log(|V|M_n/\sigma) \leq \delta_n n\mu_{\mathcal{W}}$, we have that $|(\mathbb{E}_n - \mathbb{E})[K_\varpi(W)X_{-a,j}^2]| = o(\sigma_{a\varpi j}^X)$ for all $u \in \mathcal{U}$. Therefore, we have

$$\begin{aligned} & P(\sup_{u \in \mathcal{U}} |\mathbb{E}_n[h_{uj}(X_{-a}, W)]| / \{K_u \hat{\sigma}_{a\varpi j}^X\} > 1) \\ & \leq P(\sup_{u \in \mathcal{U}} |\mathbb{E}_n[h_{uj}(X_{-a}, W)]| / \{K_u \sigma_{a\varpi j}^X\} > 1 + O(\delta_n)) + o(1) \end{aligned}$$

Applying Lemma 19 to $\mathcal{F} = \{h_{uj}(X_{-a}, W)/\sigma_{a\varpi j}^X : u \in \mathcal{U}\}$. For convenience define $\bar{\mathcal{H}}_j = \{h_{uj}(X_{-a}, W) : u \in \mathcal{U}\}$ and $\bar{\mathcal{K}}_j := \{\mathbb{E}[K_\varpi(W)X_j^2] : \varpi \in \mathcal{W}\}$. Note that $\bar{\mathcal{K}}_j$ has covering numbers bounded by the covering number of $\mathcal{K}_j := \{K_\varpi(W)X_j^2 : \varpi \in \mathcal{W}\}$ hence $\sup_Q \log N(\epsilon \|\bar{\mathcal{K}}_j\|_{Q,2}, \bar{\mathcal{K}}_j, \|\cdot\|_{Q,2}) \leq \log \sup_{\tilde{Q}} N((\epsilon/4)^2 \|F\|_{\tilde{Q},2}, \mathcal{K}_j, \|\cdot\|_{\tilde{Q},2})$ by Lemma 25. Similarly, Lemma 25 also allows us to bound covering numbers of $\bar{\mathcal{H}}_j$ via covering numbers of $\mathcal{H}_j = \{K_\varpi(W)(\tau - 1\{X_a \leq X'_a\beta_u\})X_j : u \in \mathcal{U}\}$.

$$\begin{aligned} & \sup_Q \log N(\epsilon \|F\|_{Q,2}, \mathcal{F}, \|\cdot\|_{Q,2}) \leq p \max_{j \in [p]} \sup_Q \log N(\epsilon \|F_j\|_{Q,2}, \mathcal{F}_j, \|\cdot\|_{Q,2}) \\ & \leq p \max_{j \in [p]} \sup_Q \{\log N((1/2)\epsilon \|\bar{\mathcal{H}}_j\|_{Q,2}, \bar{\mathcal{H}}_j, \|\cdot\|_{Q,2}) + \sup_Q \log N((1/2)\epsilon c\mu_{\mathcal{W}}^{1/2}, 1/\bar{\mathcal{K}}_j^{1/2}, \|\cdot\|_{Q,2})\} \\ & \leq p \max_{j \in [p]} \sup_Q \{\log N((1/2)\epsilon \|\bar{\mathcal{H}}_j\|_{Q,2}, \bar{\mathcal{H}}_j, \|\cdot\|_{Q,2}) + \sup_Q \log N((1/2)\epsilon c\mu_{\mathcal{W}}^{3/2}, \bar{\mathcal{K}}_j^{1/2}, \|\cdot\|_{Q,2})\} \\ & \leq p \max_{j \in [p]} \sup_Q \{\log N((1/2)\epsilon \|\bar{\mathcal{H}}_j\|_{Q,2}, \bar{\mathcal{H}}_j, \|\cdot\|_{Q,2}) + \sup_Q \log N(C(1/2)\epsilon c\mu_{\mathcal{W}}^{3/2}, \bar{\mathcal{K}}_j, \|\cdot\|_{Q,2})\} \\ & \leq p \max_{j \in [p]} \sup_Q \{\log N((1/2)\epsilon \|\bar{\mathcal{H}}_j\|_{Q,2}, \bar{\mathcal{H}}_j, \|\cdot\|_{Q,2}) + \sup_Q \log N(1/(2C)\epsilon c\mu_{\mathcal{W}}^{3/2}, \bar{\mathcal{K}}_j, \|\cdot\|_{Q,2})\} \\ & \leq p \max_{j \in [p]} \sup_{\tilde{Q}} \log N((1/4)\epsilon^2 \|H_j\|_{Q,2}, \mathcal{H}_j, \|\cdot\|_{Q,2}) \\ & + p \max_{j \in [p]} \sup_{\tilde{Q}} \log N((1/(4C^2))\epsilon^2 c^2 \mu_{\mathcal{W}}^3, \mathcal{K}_j, \|\cdot\|_{\tilde{Q},2}) \end{aligned}$$

where $F_j = c\|X\|_\infty/\mu_{\mathcal{W}}^{1/2}$, $H_j = \|X\|_\infty$. Since \mathcal{K}_j is the product of a VC subgraph of dimension d_W with a single function, and \mathcal{H}_j is the product of two VC subgraph of dimension $1 + d_W$ and a single function, by Lemma 19 with $\sigma^2 = 1$, we have with probability $1 - o(1)$

$$\sup_{u \in \mathcal{U}} \left| \frac{(\mathbb{E}_n - \mathbb{E})[h_{uj}(X_{-a}, W)]}{\mathbb{E}[K_\varpi(W)X_j^2]^{1/2}} \right| \leq C \sqrt{\frac{(1+d_W)\log(|V|n)}{n}} + C \frac{M_n(1+d_W)\log(|V|n)}{n\mu_{\mathcal{W}}^{1/2}}.$$

Thus, under $M_n(1+d_W)\log(|V|n) \leq n^{1/2}\mu_{\mathcal{W}}^{1/2}$ we have that we can take $K_u = C\sqrt{\frac{(1+d_W)\log(|V|n)}{n}}$. \blacksquare

APPENDIX F. TECHNICAL RESULTS FOR HIGH-DIMENSIONAL QUANTILE REGRESSION

In this section we provide technical results for high-dimensional quantile regression. It is based on a sample $(\tilde{y}_i, \tilde{x}_i, W_i)_{i=1}^n$, independent across i , $\rho_\tau(t) = (\tau - 1\{t \leq 0\})t$, $\tau \in \mathcal{T} \subset (0, 1)$ a compact interval, and a family of indicator functions $K_w(W) = 1$ if $W \in \Omega_w$, $K_w(W) = 0$ otherwise, here $\Omega_w \in \mathcal{W}$. For convenience we index the sets Ω_w by $\varpi \in B_W \subset \mathbb{R}^{d_W}$ where we normalize the diameter of B_W to be less or equal than $1/6$. Let $f_{\tilde{y}|\tilde{x}, r_u, \varpi}(\cdot)$ denote the conditional density function, $f_{\tilde{y}|\tilde{x}, r_u, \varpi}(\cdot) \leq \bar{f}$, $|f'_{\tilde{y}|\tilde{x}, r_u, \varpi}(\cdot)| \leq \bar{f}'$ and $f_u := f_{\tilde{y}|\tilde{x}, r_u, \varpi}(\tilde{x}'\eta_u)$. Moreover, we assume that

$$\|\eta_u - \eta_{\tilde{u}}\|_1 \leq L_\eta \{|\tau - \tilde{\tau}| + \|\varpi - \tilde{\varpi}\|^\rho\}. \quad (\text{F.61})$$

Although the results can be applied more generally, these results will be used for (η_u, r_u) , $u = (\tilde{y}, \tau, \varpi) \in \mathcal{U} := \{\tilde{y}\} \times \mathcal{T} \times \mathcal{W}$ satisfying

$$\mathbb{E}[K_\varpi(W)(\tau - 1\{\tilde{y} \leq \tilde{x}'\eta_u + r_u\})\tilde{x}] = 0.$$

Note that this generality is flexible enough to allow us to cover the case that the τ -conditional quantile function $Q_{\tilde{y}}(\tau|\tilde{x}, \varpi) = \tilde{x}'\tilde{\eta}_u + \tilde{r}_u$ by setting $\eta_u = \tilde{\eta}_u$ and $r_u = \tilde{r}_u$ in which case $E[(\tau - 1\{\tilde{y} \leq \tilde{x}'\eta_u + r_u\})|\tilde{x}, \varpi] = 0$. It also covers the case that

$$\tilde{\eta}_u \in \arg \min_{\beta} E[K_{\varpi}(W)\rho_{\tau}(\tilde{y} - \tilde{x}'\beta)]$$

so that $E[K_{\varpi}(W)(\tau - 1\{\tilde{y} \leq \tilde{x}'\tilde{\eta}_u\})\tilde{x}] = 0$ holds by the first order condition by setting $\eta_u = \tilde{\eta}_u$ and $r_u = 0$. Moreover, it also covers the case that we work with a sparse approximation $\bar{\eta}_u$ of $\tilde{\eta}_u$ by setting $\eta_u = \bar{\eta}_u$ and $r_u = \tilde{x}'(\tilde{\eta}_u - \bar{\eta}_u)$.

Lemma 12 (Identification Lemma). *For $u = (a, \tau, \varpi) \in \mathcal{U}$, and a subset $A_u \subset \mathbb{R}^p$ let*

$$\bar{q}_{A_u} = 1/(2\bar{f}') \cdot \inf_{\delta \in A_u} E_n [K_{\varpi}(W)f_u|\tilde{x}'\delta|^2]^{3/2} / E_n [K_{\varpi}(W)|\tilde{x}'\delta|^3] \quad (\text{F.62})$$

and assume that for all $\delta \in A_u$

$$E_n [K_{\varpi}(W)|r_u| \cdot |\tilde{x}'\delta|^2] + E_n [K_{\varpi}(W)r_u^2 \cdot |\tilde{x}'\delta|^2] \leq 1/(4\bar{f}')E_n [K_{\varpi}(W)f_u|\tilde{x}'\delta|^2]. \quad (\text{F.63})$$

Then we have

$$\begin{aligned} & E_n[E[K_{\varpi}(W)\rho_{\tau}(\tilde{y} - \tilde{x}'(\eta_u + \delta)) | \tilde{x}, r_u, W]] - E_n[E[K_{\varpi}(W)\rho_{\tau}(\tilde{y} - \tilde{x}'\eta_u) | \tilde{x}, r_u, W]] \\ & \geq \frac{\|\sqrt{f_u}\tilde{x}'\delta\|_{n,\varpi}^2}{4} \wedge \left\{ \bar{q}_{A_u} \|\sqrt{f_u}\tilde{x}'\delta\|_{n,\varpi} - K_{n2} \|\sqrt{f_u}\tilde{x}'\delta\|_{n,\varpi} - K_{n1} \|\delta\|_{1,\varpi} \right\}. \end{aligned}$$

where $K_{n2} := (\bar{f}'^{1/2} + 1)\|r_u\|_{n,\varpi}$ and $K_{n1} := \sup_{u \in \mathcal{U}, j \in [p]} \frac{E_n[E[K_{\varpi}(W)(\tau - 1\{\tilde{y} \leq \tilde{x}'\eta_u + r_u\})\tilde{x}_j|\tilde{x}, W]]}{\{E_n[K_{\varpi}(W)\tilde{x}_j^2]\}^{1/2}}$.

Proof of Lemma 12. Let $T_u = \text{support}(\eta_u)$, and $Q_u(\eta) := E_n E[K_{\varpi}(W)\rho_{\tau}(\tilde{y} - \tilde{x}'\eta) | \tilde{x}, r_u, W]$. The proof proceeds in steps.

Step 1. (Minoration) Define the maximal radius over which the criterion function can be minorated by a quadratic function

$$r_{A_u} = \sup_r \left\{ r : \begin{aligned} & Q_u(\eta_u + \delta) - Q_u(\eta_u) + K_{n2} \|\sqrt{f_u}\tilde{x}'\delta\|_{n,\varpi} + K_{n1} \|\delta\|_{1,\varpi} \geq \frac{1}{4} \|\sqrt{f_u}\tilde{x}'\delta\|_{n,\varpi}^2, \\ & \forall \delta \in A_u, \|\sqrt{f_u}\tilde{x}'\delta\|_{n,\varpi} \leq r \end{aligned} \right\}.$$

Step 2 below shows that $r_{A_u} \geq \bar{q}_{A_u}$. By construction of r_{A_u} and the convexity of $Q_u(\cdot)$, $\|\cdot\|_{1,\varpi}$ and $\|\cdot\|_{n,\varpi}$,

$$\begin{aligned} & Q_u(\eta_u + \delta) - Q_u(\eta_u) + K_{n2} \|\sqrt{f_u}\tilde{x}'\delta\|_{n,\varpi} + K_{n1} \|\delta\|_{1,\varpi} \geq \\ & \geq \frac{\|\sqrt{f_u}\tilde{x}'\delta\|_{n,\varpi}^2}{4} \wedge \left\{ \frac{\|\sqrt{f_u}\tilde{x}'\delta\|_{n,\varpi}}{r_{A_u}} \cdot \inf_{\tilde{\delta} \in A_u, \|\sqrt{f_u}\tilde{x}'\tilde{\delta}\|_{n,\varpi} \geq r_{A_u}} Q_u(\eta_u + \tilde{\delta}) - Q_u(\eta_u) + K_{n2} \|\sqrt{f_u}\tilde{x}'\tilde{\delta}\|_{n,\varpi} + K_{n1} \|\tilde{\delta}\|_{1,\varpi} \right\} \\ & \geq \frac{\|\sqrt{f_u}\tilde{x}'\delta\|_{n,\varpi}^2}{4} \wedge \left\{ \frac{\|\sqrt{f_u}\tilde{x}'\delta\|_{n,\varpi}}{r_{A_u}} \cdot \frac{r_{A_u}^2}{4} \right\} \\ & \geq \frac{\|\sqrt{f_u}\tilde{x}'\delta\|_{n,\varpi}^2}{4} \wedge \left\{ \bar{q}_{A_u} \|\sqrt{f_u}\tilde{x}'\delta\|_{n,\varpi} \right\}. \end{aligned}$$

Step 2. ($r_{A_u} \geq \bar{q}_{A_u}$) Let $F_{\tilde{y}|\tilde{x}, r_u, \varpi}$ denote the conditional distribution of \tilde{y} given \tilde{x}, r_u, ϖ . From [66], for any two scalars w and v the Knight's identity is

$$\rho_{\tau}(w - v) - \rho_{\tau}(w) = -v(\tau - 1\{w \leq 0\}) + \int_0^v (1\{w \leq z\} - 1\{w \leq 0\})dz. \quad (\text{F.64})$$

Using (F.64) with $w = \tilde{y}_i - \tilde{x}'_i\eta_u$ and $v = \tilde{x}'_i\delta$ and taking expectations with respect to \tilde{y} , we have

$$\begin{aligned} Q_u(\eta_u + \delta) - Q_u(\eta_u) = & -E_n[E[K_{\varpi}(W)(\tau - 1\{\tilde{y} \leq \tilde{x}'\eta_u\})\tilde{x}'_i\delta | \tilde{x}, r_u, W]] \\ & + E_n \left[\int_0^{K_{\varpi}(W)\tilde{x}'\delta} F_{\tilde{y}|\tilde{x}, r_u, \varpi}(\tilde{x}'\eta_u + t) - F_{\tilde{y}|\tilde{x}, r_u, \varpi}(\tilde{x}'\eta_u) dt \right]. \end{aligned}$$

Using the law of iterated expectations and mean value expansion, the relation

$$\begin{aligned}
& |\mathbb{E}_n[\mathbb{E}[K_\varpi(W)(\tau - 1\{\tilde{y} \leq \tilde{x}'\eta_u\})\tilde{x}'\delta \mid \tilde{x}, r_u, W]]| \\
&= |\mathbb{E}_n[K_\varpi(W)\{F_{\tilde{y}|\tilde{x}, r_u, \varpi}(\tilde{x}'\eta_u + r_u) - F_{\tilde{y}|\tilde{x}, r_u, \varpi}(\tilde{x}'\eta_u)\}\tilde{x}'\delta]| \\
&+ \mathbb{E}_n[K_\varpi(W)\{\tau - F_{\tilde{y}|\tilde{x}, r_u, \varpi}(\tilde{x}'\eta_u + r_u)\}\tilde{x}'\delta \mid \tilde{x}, r_u, W]| \\
&\leq \mathbb{E}_n[K_\varpi(W)f_u|r_u| |\tilde{x}'\delta|] + \bar{f}'\mathbb{E}_n[K_\varpi(W)|r_u|^2|\tilde{x}'\delta|] + K_{n1}\|\delta\|_{1, \varpi} \\
&\leq \|\sqrt{f_u}r_u\|_{n, \varpi}\|\sqrt{f_u}\tilde{x}'\delta\|_{n, \varpi} + \bar{f}'\|r_u\|_{n, \varpi}\|r_u\tilde{x}'\delta\|_{n, \varpi} + K_{n1}\|\delta\|_{1, \varpi} \\
&\leq (\bar{f}^{1/2} + 1)\|r_u\|_{n, \varpi}\|\sqrt{f_u}\tilde{x}'\delta\|_{n, \varpi} + K_{n1}\|\delta\|_{1, \varpi}
\end{aligned}$$

where we used our assumption on the approximation error and we have $K_{n2} = (\bar{f}^{1/2} + 1)\|r_u\|_{n, \varpi}$. With that and similar arguments we obtain for $\tilde{t}_{\tilde{x}_i, t} \in [0, t]$

$$\begin{aligned}
& Q_u(\eta_u + \delta) - Q_u(\eta_u) + K_{n2}\|\sqrt{f_u}\tilde{x}'\delta\|_{n, \varpi} + K_{n1}\|\delta\|_{1, \varpi} \geq \\
& Q_u(\eta_u + \delta) - Q_u(\eta_u) + \mathbb{E}_n[\mathbb{E}[K_\varpi(W)(\tau - 1\{\tilde{y} \leq \tilde{x}'\eta_u\})\tilde{x}'\delta \mid \tilde{x}, r_u, W]] = \\
&= \mathbb{E}_n\left[\int_0^{K_\varpi(W)\tilde{x}'\delta} F_{\tilde{y}|\tilde{x}, r_u, \varpi}(\tilde{x}'\eta_u + t) - F_{\tilde{y}|\tilde{x}, r_u, \varpi}(\tilde{x}'\eta_u) dt\right] \\
&= \mathbb{E}_n\left[\int_0^{K_\varpi(W)\tilde{x}'\delta} t f_{\tilde{y}|\tilde{x}, r_u, \varpi}(\tilde{x}'\eta_u) + \frac{t^2}{2} f'_{\tilde{y}|\tilde{x}, r_u, \varpi}(\tilde{x}'\eta_u + \tilde{t}_{\tilde{x}, t}) dt\right] \\
&\geq \frac{1}{2}\|\sqrt{f_u}\tilde{x}'\delta\|_{n, \varpi}^2 - \frac{1}{6}\bar{f}'\mathbb{E}_n[K_\varpi(W)|\tilde{x}'\delta|^3] - \mathbb{E}_n\left[\int_0^{K_\varpi(W)\tilde{x}'\delta} t[f_{\tilde{y}|\tilde{x}, r_u, \varpi}(\tilde{x}'\eta_u) - f_{\tilde{y}|\tilde{x}, r_u, \varpi}(\tilde{x}'\eta_u + r_u)] dt\right] \\
&\geq \frac{1}{4}\|\sqrt{f_u}\tilde{x}'\delta\|_{n, \varpi}^2 + \frac{1}{4}\|\sqrt{f_u}\tilde{x}'\delta\|_{n, \varpi}^2 - \frac{1}{6}\bar{f}'\mathbb{E}_n[K_\varpi(W)|\tilde{x}'\delta|^3] - (\bar{f}'/2)\mathbb{E}_n[K_\varpi(W)|\tilde{r}_u| \cdot |\tilde{x}'\delta|^2].
\end{aligned} \tag{F.65}$$

Moreover, by assumption we have

$$\mathbb{E}_n[K_\varpi(W)|r_u| \cdot |\tilde{x}'\delta|^2] \leq \frac{1}{4\bar{f}'}\mathbb{E}_n[K_\varpi(W)f_u|\tilde{x}'\delta|^2] \tag{F.66}$$

Note that for any δ such that $\|\sqrt{f_u}\tilde{x}'\delta\|_{n, \varpi} \leq \bar{q}_{A_u}$ we have

$$\|\sqrt{f_u}\tilde{x}'\delta\|_{n, \varpi} \leq \bar{q}_{A_u} \leq 1/(2\bar{f}') \cdot \mathbb{E}_n[K_\varpi(W)f_u|\tilde{x}'\delta|^2]^{3/2} / \mathbb{E}_n[K_\varpi(W)|\tilde{x}'\delta|^3].$$

It follows that $(1/6)\bar{f}'\mathbb{E}_n[K_\varpi(W)|\tilde{x}'\delta|^3] \leq (1/8)\mathbb{E}_n[K_\varpi(W)f_u|\tilde{x}'\delta|^2]$. Combining this with (F.66) we have

$$\frac{1}{4}\mathbb{E}_n[K_\varpi(W)f_u|\tilde{x}'\delta|^2] - \frac{\bar{f}'}{6}\mathbb{E}_n[K_\varpi(W)|\tilde{x}'\delta|^3] - \frac{\bar{f}'}{2}\mathbb{E}_n[K_\varpi(W)|r_u| \cdot |\tilde{x}'\delta|^2] \geq 0. \tag{F.67}$$

Combining (F.65) and (F.67) we have $r_{A_u} \geq \bar{q}_{A_u}$. ■

Lemma 13. *Let \mathcal{W} be a VC-class of sets with VC-index d_W . Conditional on $\{(W_i, \tilde{x}_i), i = 1, \dots, n\}$ we have*

$$P_{\tilde{y}} \left(\sup_{\substack{\tau \in \mathcal{T}, \varpi \in \mathcal{W}, \\ \underline{N} \leq \|\delta\|_{1, \varpi} \leq \bar{N}}} \left| \mathbb{E}_n \left(K_\varpi(W) \frac{\rho_\tau(\tilde{y} - \tilde{x}'(\eta_u + \delta)) - \rho_\tau(\tilde{y} - \tilde{x}'\eta_u)}{\|\delta\|_{1, \varpi}} \right) \right| \geq M \mid (W_i, \tilde{x}_i)_{i=1}^n \right) \leq S_n \exp(-(M/4 - 3)^2/32)$$

where $S_n \leq 8p|\hat{\mathcal{N}}| \cdot |\widehat{\mathcal{W}}| \cdot |\widehat{\mathcal{T}}|$, with

$$|\hat{\mathcal{N}}| \leq 1 + \lfloor 3\sqrt{n} \log(\bar{N}/\underline{N}) \rfloor, \quad |\widehat{\mathcal{T}}| \leq 2\sqrt{n} \frac{\max_{i \leq n} \|\tilde{x}_i\|_\infty}{\underline{N}} L_\eta, \quad |\widehat{\mathcal{W}}| \leq n^{d_W} + \left\{ 2\sqrt{n} \frac{\max_{i \leq n} \|\tilde{x}_i\|_\infty}{\underline{N}} L_\eta \right\}^{d_W/\rho}.$$

Proof of Lemma 13. Let $g_{i\tau\varpi}(b) = K_\varpi(W_i)\{\rho_\tau(\tilde{y}_i - \tilde{x}'_i\eta_{\tau\varpi} + b) - \rho_\tau(\tilde{y}_i - \tilde{x}'_i\eta_{\tau\varpi})\} \leq K_\varpi(W_i)|b|$ since $K_\varpi(W_i) \in \{0, 1\}$. Note that $|g_{i\tau\varpi}(b) - g_{i\tau\varpi}(a)| \leq K_\varpi(W_i)|b - a|$. To easy the notation we omit the conditioning on (\tilde{x}_i, W_i) from the probabilities.

For any $\delta \in \mathbb{R}^p$, since ρ_τ is 1-Lipschitz, we have

$$\text{Var} \left(\mathbb{G}_n \left(\frac{g_{\tau\varpi}(\tilde{x}'\delta)}{\|\delta\|_{1,\varpi}} \right) \right) \leq \frac{\mathbb{E}_n[\{g_{\tau\varpi}(\tilde{x}'\delta)\}^2]}{\|\tilde{x}'\delta\|_{n,\varpi}^2} \leq \frac{\mathbb{E}_n[K_\varpi(W)\tilde{x}'\delta]^2}{\|\tilde{x}'\delta\|_{n,\varpi}^2} = 1$$

since by definition $\|\delta\|_{1,\varpi} = \sum_j \|\delta_j\|_{1,\varpi} = \sum_j \|\tilde{x}'_j \delta_j\|_{n,\varpi} \geq \|\tilde{x}'\delta\|_{n,\varpi}$.

Since we are conditioning on $(W_i, \tilde{x}_i)_{i=1}^n$ the process is independent across i . Then, by Lemma 2.3.7 in [100] (Symmetrization for Probabilities) we have for any $M > 1$

$$\begin{aligned} & \mathbb{P} \left(\sup_{\tau \in \mathcal{T}, \varpi \in \mathcal{W}, \underline{N} \leq \|\delta\|_{1,\varpi} \leq \bar{N}} \left| \mathbb{G}_n \left(\frac{g_{\tau\varpi}(\tilde{x}'\delta)}{\|\delta\|_{1,\varpi}} \right) \right| \geq M \right) \\ & \leq \frac{2}{1-M^{-2}} \mathbb{P} \left(\sup_{\tau \in \mathcal{T}, \varpi \in \mathcal{W}, \underline{N} \leq \|\delta\|_{1,\varpi} \leq \bar{N}} \left| \mathbb{G}_n^o \left(\frac{g_{\tau\varpi}(\tilde{x}'\delta)}{\|\delta\|_{1,\varpi}} \right) \right| \geq M/4 \right) \end{aligned}$$

where \mathbb{G}_n^o is the symmetrized process.

Consider $\mathcal{F}_{t,\tau,\varpi} = \{\delta : \|\delta\|_{1,\varpi} = t\}$. We will consider the families of $\mathcal{F}_{t,\tau,\varpi}$ for $t \in [\underline{N}, \bar{N}]$, $\tau \in \mathcal{T}$ and $\varpi \in \mathcal{W}$.

We will construct a finite net $\hat{\mathcal{T}} \times \hat{\mathcal{W}} \times \hat{\mathcal{N}}$ of $\mathcal{T} \times \mathcal{W} \times [\underline{N}, \bar{N}]$ such that

$$\sup_{\tau \in \mathcal{T}, \varpi \in \mathcal{W}, t \in [\underline{N}, \bar{N}], \delta \in \mathcal{F}_{t,\tau,\varpi}} \left| \mathbb{G}_n^o \left(\frac{g_{\tau\varpi}(\tilde{x}'\delta)}{\|\delta\|_{1,\varpi}} \right) \right| \leq 3 + \sup_{\tau \in \hat{\mathcal{T}}, \varpi \in \hat{\mathcal{W}}, t \in \hat{\mathcal{N}}} \sup_{\delta \in \mathcal{F}_{t,\tau,\varpi}} \left| \mathbb{G}_n^o \left(\frac{g_{\tau\varpi}(\tilde{x}'\delta)}{t} \right) \right| =: 3 + \mathcal{A}^o.$$

By triangle inequality we have

$$\begin{aligned} \left| \mathbb{G}_n^o \left(\frac{g_{\tau\varpi}(\tilde{x}'\delta)}{t} - \frac{g_{\tilde{\tau}\tilde{\varpi}}(\tilde{x}'\delta)}{t} \right) \right| & \leq \left| \mathbb{G}_n^o \left(\frac{g_{\tau\varpi}(\tilde{x}'\delta)}{t} - \frac{g_{\tau\tilde{\varpi}}(\tilde{x}'\delta)}{t} \right) \right| + \left| \mathbb{G}_n^o \left(\frac{g_{\tau\tilde{\varpi}}(\tilde{x}'\delta)}{t} - \frac{g_{\tilde{\tau}\tilde{\varpi}}(\tilde{x}'\delta)}{t} \right) \right| \\ & + \left| \mathbb{G}_n^o \left(\frac{g_{\tilde{\tau}\tilde{\varpi}}(\tilde{x}'\delta)}{t} - \frac{g_{\tilde{\tau}\tilde{\varpi}}(\tilde{x}'\delta)}{t} \right) \right| \end{aligned} \quad (\text{F.68})$$

The first term in (F.68) is such that

$$\begin{aligned} \left| \mathbb{G}_n^o \left(\frac{g_{\tau\varpi}(\tilde{x}'\delta)}{t} - \frac{g_{\tau\tilde{\varpi}}(\tilde{x}'\delta)}{t} \right) \right| & \leq \frac{2\sqrt{n}}{t} \mathbb{E}_n[K_\varpi(W)|\tilde{x}'(\eta_{\tau\varpi} - \eta_{\tau\tilde{\varpi}})|] \\ & \leq \frac{2\sqrt{n}}{\underline{N}} \max_{i \leq n} \|\tilde{x}_i\|_\infty \mathbb{E}_n[K_\varpi(W)] \|\eta_{\tau\varpi} - \eta_{\tau\tilde{\varpi}}\|_1 \\ & \leq \frac{2\sqrt{n}}{\underline{N}} \max_{i \leq n} \|\tilde{x}_i\|_\infty \mathbb{E}_n[K_\varpi(W)] L_\eta |\tau - \tau'|. \end{aligned} \quad (\text{F.69})$$

Define a net $\hat{\mathcal{T}} = \{\tau_1, \dots, \tau_T\}$ such that

$$|\tau_{k+1} - \tau_k| \leq \left\{ 2\sqrt{n} \frac{\max_{i \leq n} \|\tilde{x}_i\|_\infty}{\underline{N}} L_\eta \right\}^{-1}.$$

To bound the second term in (F.68), note that \mathcal{W} is a VC-class. Therefore, by Corollary 2.6.3 in [100] we have that conditional on $(W_i)_{i=1}^n$, there are at most n^{dw} different sets $\varpi \in \mathcal{W}$ that induce a different sequence $\{K_\varpi(W_1), \dots, K_\varpi(W_n)\}$. Thus we can choose a (data-dependent) cover $\hat{\mathcal{W}}$ with at most n^{dw} values of ϖ . Further, similarly to (F.70) we have $\|\eta_{\tilde{\tau}\varpi} - \eta_{\tilde{\tau}\tilde{\varpi}}\|_1 \leq L_\eta \|\varpi - \tilde{\varpi}\|^\rho$ and

$$\begin{aligned} \left| \mathbb{G}_n^o \left(\frac{g_{\tilde{\tau}\varpi}(\tilde{x}'\delta)}{t} - \frac{g_{\tilde{\tau}\tilde{\varpi}}(\tilde{x}'\delta)}{t} \right) \right| & \leq \frac{2\sqrt{n}}{t} \mathbb{E}_n[K_\varpi(W)|\tilde{x}'(\eta_{\tilde{\tau}\varpi} - \eta_{\tilde{\tau}\tilde{\varpi}})|] + \frac{2\sqrt{n}}{t} \mathbb{E}_n[|K_\varpi(W) - K_{\tilde{\varpi}}(W)|\|\tilde{x}'\delta\|] \\ & \leq \frac{2\sqrt{n}}{\underline{N}} \max_{i \leq n} \|\tilde{x}_i\|_\infty \mathbb{E}_n[K_\varpi(W)] \|\eta_{\tilde{\tau}\varpi} - \eta_{\tilde{\tau}\tilde{\varpi}}\|_1 \\ & \leq \frac{2\sqrt{n}}{\underline{N}} \max_{i \leq n} \|\tilde{x}_i\|_\infty \mathbb{E}_n[K_\varpi(W)] L_\eta \|\varpi - \tilde{\varpi}\|^\rho. \end{aligned} \quad (\text{F.70})$$

We define a net $\hat{\mathcal{W}}$ such that $|\hat{\mathcal{W}}| \leq n^{dw} + \left\{ 2\sqrt{n} \frac{\max_{i \leq n} \|\tilde{x}_i\|_\infty}{\underline{N}} L_\eta \right\}^{dw/\rho}$

To bound the third term in (F.68), note that for any $\delta \in \mathcal{F}_{t,\tau,\varpi}$, $t \leq \tilde{t}$, by considering $\tilde{\delta} := \delta(\tilde{t}/t) \in \mathcal{F}_{\tilde{t},\tau,\varpi}$ we have

$$\begin{aligned} \left| \mathbb{G}_n^o \left(\frac{g_{\tau\varpi}(\tilde{x}'\delta)}{t} - \frac{g_{\tau\varpi}(\tilde{x}'\delta(\tilde{t}/t))}{\tilde{t}} \right) \right| &\leq \left| \mathbb{G}_n^o \left(\frac{g_{\tau\varpi}(\tilde{x}'\delta)}{t} - \frac{g_{\tau\varpi}(\tilde{x}'\delta(\tilde{t}/t))}{t} \right) \right| + \left| \mathbb{G}_n^o \left(\frac{g_{\tau\varpi}(\tilde{x}'\delta(\tilde{t}/t))}{t} - \frac{g_{\tau\varpi}(\tilde{x}'\delta(\tilde{t}/t))}{\tilde{t}} \right) \right| \\ &= \frac{1}{t} \left| \mathbb{G}_n^o (g_{\tau\varpi}(\tilde{x}'\delta) - g_{\tau\varpi}(\tilde{x}'\delta[\tilde{t}/t])) \right| + \left| \mathbb{G}_n^o (g_{\tau\varpi}(\tilde{x}'\delta(\tilde{t}/t))) \right| \cdot \left| \frac{1}{t} - \frac{1}{\tilde{t}} \right| \\ &\leq \sqrt{n} \mathbb{E}_n \left(\frac{|K_{\varpi}(W)\tilde{x}'\delta|}{t} \right) \frac{|t-\tilde{t}|}{t} + \sqrt{n} \mathbb{E}_n (|K_{\varpi}(W)\tilde{x}'\delta|) \frac{\tilde{t}}{t} \left| \frac{1}{t} - \frac{1}{\tilde{t}} \right| \\ &= 2\sqrt{n} \mathbb{E}_n \left(\frac{|K_{\varpi}(W)\tilde{x}'\delta|}{t} \right) \left| \frac{t-\tilde{t}}{t} \right| \leq 2\sqrt{n} \left| \frac{t-\tilde{t}}{t} \right|. \end{aligned}$$

We let $\hat{\mathcal{N}}$ be a ε -net $\{\underline{N} =: t_1, t_2, \dots, t_K := \bar{N}\}$ of $[\underline{N}, \bar{N}]$ such that $|t_k - t_{k+1}|/t_k \leq 1/(2\sqrt{n})$. Note that we can achieve that with $|\hat{\mathcal{N}}| \leq 1 + \lfloor 3\sqrt{n} \log(\bar{N}/\underline{N}) \rfloor$.

By Markov bound, we have

$$\begin{aligned} P(\mathcal{A}^o \geq K) &\leq \min_{\psi \geq 0} \exp(-\psi K) \mathbb{E}[\exp(\psi \mathcal{A}^o)] \\ &\leq 8p|\hat{\mathcal{T}}| \cdot |\hat{\mathcal{W}}| \cdot |\hat{\mathcal{N}}| \min_{\psi \geq 0} \exp(-\psi K) \exp(8\psi^2) \\ &\leq 8p|\hat{\mathcal{T}}| \cdot |\hat{\mathcal{W}}| \cdot |\hat{\mathcal{N}}| \exp(-K^2/32) \end{aligned}$$

here we set $\psi = K/16$ and bound $\mathbb{E}[\exp(\psi \mathcal{A}^o)]$ as follows

$$\begin{aligned} \mathbb{E}[\exp(\psi \mathcal{A}^o)] &\leq_{(1)} 2|\hat{\mathcal{T}}| \cdot |\hat{\mathcal{W}}| \cdot |\hat{\mathcal{N}}| \sup_{(\tau,\varpi,t) \in \hat{\mathcal{T}} \times \hat{\mathcal{W}} \times \hat{\mathcal{N}}} \mathbb{E} \left[\exp \left(\psi \sup_{\|\delta\|_{1,\varpi}=t} \mathbb{G}_n^o \left(\frac{g_{\tau\varpi}(\tilde{x}'\delta)}{t} \right) \right) \right] \\ &\leq_{(2)} 2|\hat{\mathcal{T}}| \cdot |\hat{\mathcal{W}}| \cdot |\hat{\mathcal{N}}| \sup_{(\tau,\varpi,t) \in \hat{\mathcal{T}} \times \hat{\mathcal{W}} \times \hat{\mathcal{N}}} \mathbb{E} \left[\exp \left(2\psi \sup_{\|\delta\|_{1,\varpi}=t} \mathbb{G}_n^o \left(\frac{K_{\varpi}(W)\tilde{x}'\delta}{t} \right) \right) \right] \\ &\leq_{(3)} 2|\hat{\mathcal{T}}| \cdot |\hat{\mathcal{W}}| \cdot |\hat{\mathcal{N}}| \sup_{(\tau,\varpi,t) \in \hat{\mathcal{T}} \times \hat{\mathcal{W}} \times \hat{\mathcal{N}}} \mathbb{E} \left[\exp \left(2\psi \left[\sup_{\|\delta\|_{1,\varpi}=t} \frac{\|\delta\|_{1,\varpi}}{t} \max_{j \leq p} \frac{|\mathbb{G}_n^o(K_{\varpi}(W)\tilde{x}_j)|}{\{\mathbb{E}_n[K_{\varpi}(W)\tilde{x}_j^2]\}^{1/2}} \right] \right) \right] \\ &=_{(4)} 2|\hat{\mathcal{T}}| \cdot |\hat{\mathcal{W}}| \cdot |\hat{\mathcal{N}}| \sup_{(\tau,\varpi,t) \in \hat{\mathcal{T}} \times \hat{\mathcal{W}} \times \hat{\mathcal{N}}} \mathbb{E} \left[\exp \left(2\psi \left[\max_{j \leq p} \frac{|\mathbb{G}_n^o(K_{\varpi}(W)\tilde{x}_j)|}{\{\mathbb{E}_n[K_{\varpi}(W)\tilde{x}_j^2]\}^{1/2}} \right] \right) \right] \\ &\leq_{(5)} 4p|\hat{\mathcal{T}}| \cdot |\hat{\mathcal{W}}| \cdot |\hat{\mathcal{N}}| \max_{j \leq p} \sup_{\varpi \in \hat{\mathcal{W}}} \mathbb{E} \left[\exp \left(4\psi \frac{\mathbb{G}_n^o(K_{\varpi}(W)\tilde{x}_j)}{\{\mathbb{E}_n[K_{\varpi}(W)\tilde{x}_j^2]\}^{1/2}} \right) \right] \\ &\leq_{(6)} 8p|\hat{\mathcal{T}}| \cdot |\hat{\mathcal{W}}| \cdot |\hat{\mathcal{N}}| \exp(8\psi^2) \end{aligned}$$

here (1) follows by $\exp(\max_{i \in I} |z_i|) \leq 2|I| \max_{i \in I} \exp(z_i)$, (2) by contraction principle (apply Theorem 4.12 [72] with $t_i = K_{\varpi}(W_i)\tilde{x}'_i\delta$, and $\phi_i(t_i) = \rho_{\tau}(K_{\varpi}(W_i)\tilde{y}_i - K_{\varpi}(W_i)\tilde{x}'_i\eta_{\tau} + t_i) - \rho_{\tau}(K_{\varpi}(W_i)\tilde{y}_i - K_{\varpi}(W_i)\tilde{x}'_i\eta_{\tau})$ so that $|\phi_i(s) - \phi_i(t)| \leq |s - t|$ and $\phi_i(0) = 0$, (3) follows by

$$|\mathbb{G}_n^o(K_{\varpi}(W)\tilde{x}'\delta)| \leq \|\delta\|_{1,\varpi} \max_{j \leq p} |\mathbb{G}_n^o(K_{\varpi}(W)\tilde{x}_j)| / \{\mathbb{E}_n[K_{\varpi}(W)\tilde{x}_j^2]\}^{1/2},$$

(4) by the definition of suprema, (5) we again use $\exp(\max_{i \in I} |z_i|) \leq 2|I| \max_{i \in I} \exp(z_i)$, and (6) $\exp(z) + \exp(-z) \leq 2\exp(z^2/2)$. ■

Lemma 14 (Estimation Error of Refitted Quantile Regression). *Consider an arbitrary vector $\hat{\eta}_u$ and suppose $\|\eta_u\|_0 \leq s$. Let $\|r_{iu} \leq \bar{r}_u\|_{n,\varpi}$, $|\text{support}(\hat{\eta}_u)| \leq \hat{s}_u$ and $\mathbb{E}_n[K_{\varpi}(W)\{\rho_{\tau}(\tilde{y}_i - \tilde{x}'_i\hat{\eta}_u) - \rho_{\tau}(\tilde{y}_i - \tilde{x}'_i\eta_u)\}] \leq \hat{Q}_u$ for all $u \in \mathcal{U}$ hold. Furthermore, suppose that*

$$\sup_{u=(\tau,\varpi) \in \mathcal{U}} \left| \mathbb{E}_n \left(K_{\varpi}(W) \frac{\rho_{\tau}(\tilde{y} - \tilde{x}'\tilde{\eta}_u) - \rho_{\tau}(\tilde{y} - \tilde{x}'\eta_u)}{\|\tilde{\eta}_u - \eta_u\|_{1,\varpi}} \right) - \mathbb{E} \left[K_{\varpi}(W) \frac{\rho_{\tau}(\tilde{y} - \tilde{x}'\tilde{\eta}_u) - \rho_{\tau}(\tilde{y} - \tilde{x}'\eta_u)}{\|\tilde{\eta}_u - \eta_u\|_{1,\varpi}} \mid W, \tilde{x} \right] \right| \leq \frac{t_3}{\sqrt{n}}.$$

Under these events, we have for n large enough,

$$\|\sqrt{f_u}\tilde{x}'_i(\tilde{\eta}_u - \eta_u)\|_{n,\varpi} \lesssim \tilde{N}_u := \sqrt{\frac{(\hat{s}_u + s)}{\phi_{\min}(u, \hat{s}_u + s)}} (K_{n1} + t_3/\sqrt{n}) + K_{n2} + \bar{f}\bar{r}_u + \hat{Q}_u^{1/2}$$

where $\phi_{\min}(u, k) = \inf_{\|\delta\|_0=k} \|\sqrt{f_u} \tilde{x}' \delta\|_{n,\varpi}^2 / \|\delta\|^2$, provided that

$$\sup_{u \in \mathcal{U}, \|\tilde{\delta}\|_0 \leq \hat{s}_u + s} \frac{\tilde{f}' \mathbb{E}_n[K_{\varpi}(W)(|r_u| + |r_u|^2)|\tilde{x}' \tilde{\delta}|^2]}{\mathbb{E}_n[K_{\varpi}(W)f_u|\tilde{x}' \tilde{\delta}|^2]} + \tilde{N}_u / \bar{q}_{A_u} \rightarrow 0. \quad (\text{F.71})$$

where $A_u = \{\delta \in \mathbb{R}^p : \|\delta\|_0 \leq \hat{s}_u + s\}$.

Proof of Lemma 14. Let $\hat{\delta}_u = \hat{\eta}_u - \eta_u$ which satisfies $\|\hat{\delta}_u\|_0 \leq \hat{s}_u + s$. By optimality of $\tilde{\eta}_u$ in the refitted quantile regression we have

$$\begin{aligned} & \mathbb{E}_n[K_{\varpi}(W)\rho_{\tau}(\tilde{y}_i - \tilde{x}'_i \tilde{\eta}_u)] - \mathbb{E}_n[K_{\varpi}(W)\rho_{\tau}(\tilde{y}_i - \tilde{x}'_i \eta_u)] \\ & \leq \mathbb{E}_n[K_{\varpi}(W)\rho_{\tau}(\tilde{y}_i - \tilde{x}'_i \hat{\eta}_u)] - \mathbb{E}_n[K_{\varpi}(W)\rho_{\tau}(\tilde{y}_i - \tilde{x}'_i \eta_u)] \leq \hat{Q}_u \end{aligned} \quad (\text{F.72})$$

where the second inequality holds by assumption.

Moreover, by assumption, uniformly over $u \in \mathcal{U}$, we have conditional on $(W_i, \tilde{x}_i, r_{iu})_{i=1}^n$ that

$$\left| \mathbb{G}_n \left(K_{\varpi}(W) \frac{\rho_{\tau}(\tilde{y} - \tilde{x}'(\eta_u + \tilde{\delta}_u)) - \rho_{\tau}(\tilde{y} - \tilde{x}'\eta_u)}{\|\tilde{\delta}_u\|_{1,\varpi}} \right) \right| \leq t_3. \quad (\text{F.73})$$

Thus combining relations (F.72) and (F.73), we have

$$\mathbb{E}_n[\mathbb{E}[K_{\varpi}(W)\{\rho_u(\tilde{y} - \tilde{x}'(\eta_u + \tilde{\delta}_u)) - \rho_u(\tilde{y} - \tilde{x}'\eta_u)\}|\tilde{x}, \tilde{r}, \varpi]] \leq \|\tilde{\delta}_u\|_{1,\varpi} t_3 / \sqrt{n} + \hat{Q}_u.$$

Invoking the sparse identifiability relation of Lemma 12, since the required condition on the approximation errors r_u 's holds by assumption (F.71), for n large enough

$$\frac{\|\sqrt{f_u} \tilde{x}' \tilde{\delta}_u\|_{n,\varpi}^2}{4} \wedge \left\{ \bar{q}_{A_u} \|\sqrt{f_u} \tilde{x}' \tilde{\delta}_u\|_{n,\varpi} \right\} \leq K_{n2} \|\sqrt{f_u} \tilde{x}' \tilde{\delta}_u\|_{n,\varpi} + \|\tilde{\delta}_u\|_{1,\varpi} (K_{n1} + t_3 / \sqrt{n}) + \hat{Q}_u,$$

where \bar{q}_{A_u} is defined with $A_u := \{\delta : \|\delta\|_0 \leq \hat{s}_u + s\}$. Moreover, by the sparsity of $\tilde{\delta}_u$ we have $\|\tilde{\delta}_u\|_{1,\varpi} \leq \sqrt{(\hat{s}_u + s) / \phi_{\min}(u, \hat{s}_u + s)} \|\sqrt{f_u} \tilde{x}' \tilde{\delta}_u\|_{n,\varpi}$ so that we have for $t = \|\sqrt{f_u} \tilde{x}' \tilde{\delta}_u\|_{n,\varpi}$,

$$\frac{t^2}{4} \wedge \{\bar{q}_{A_u} t\} \leq t(K_{n2} + \sqrt{(\hat{s}_u + s) / \phi_{\min}(u, \hat{s}_u + s)} \{K_{n1} + t_3 / \sqrt{n}\}) + \hat{Q}_u.$$

Note that for positive numbers $(t^2/4) \wedge (\bar{q}_{A_u} t) \leq A + Bt$ implies $t^2/4 \leq A + Bt$ provided $\bar{q}_{A_u}/2 > B$ and $2\bar{q}_{A_u}^2 > A$. (Indeed, otherwise $(t^2/4) \geq qt$ so that $t \geq 4q$, which in turn implies that $2\bar{q}_{A_u}^2 + \bar{q}_{A_u} t/2 \leq (t^2/4) \wedge \bar{q}_{A_u} t \leq A + Bt$.) Note that $\bar{q}_{A_u}/2 > B$ and $2\bar{q}_{A_u}^2 > A$ is implied by condition (F.71) when we set $A = \hat{Q}_u$ and $B = (K_{n2} + \sqrt{(\hat{s}_u + s) / \phi_{\min}(u, \hat{s}_u + s)}) \{K_{n1} + t_3 / \sqrt{n}\}$. Thus the minimum is achieved in the quadratic part. Therefore, for n sufficiently large, we have

$$\|\sqrt{f_u} \tilde{x}' \tilde{\delta}_u\|_{n,\varpi} \leq \hat{Q}_u^{1/2} + K_{n2} + (K_{n1} + t_3 / \sqrt{n}) \sqrt{(\hat{s}_u + s) / \phi_{\min}(u, \hat{s}_u + s)}.$$

■

Under the condition $\max_{i \leq n} \|\tilde{x}_i\|_{\infty}^2 \log(n \vee p) = o(n \min_{\tau \in \mathcal{T}} \tau(1 - \tau))$, the next result provides new bounds for the data driven penalty choice parameter when the quantile indices in \mathcal{T} can approach the extremes.

Lemma 15 (Pivotal Penalty Parameter Bound). *Let $\underline{\tau} = \min_{\tau \in \mathcal{T}} \tau(1 - \tau)$ and $K_n = \max_{i \leq n, j \in [p]} |\tilde{x}_{ij} / \hat{\sigma}_j|$, $\hat{\sigma}_j = \mathbb{E}_n[\tilde{x}_j^2]^{1/2}$. Under $K_n^2 \log(p / \underline{\tau}) = o(n \underline{\tau})$, for n large enough we have that for some constant \bar{C}*

$$\Lambda(1 - \xi | \tilde{x}_1, \dots, \tilde{x}_n) \leq \bar{C} \sqrt{\frac{\log(16p / (\underline{\tau} \xi))}{n}}$$

where $\Lambda(1 - \xi | \tilde{x}_1, \dots, \tilde{x}_n)$ is the $1 - \xi$ quantile of $\max_{j \in [p]} \sup_{\tau \in \mathcal{T}} \left| \frac{\sum_{i=1}^n \tilde{x}_{ij}(\tau - 1\{U_i \leq \tau\})}{\hat{\sigma}_j \sqrt{\tau(1-\tau)}} \right|$ conditional on $\tilde{x}_1, \dots, \tilde{x}_n$, and U_i are independent $\text{uniform}(0, 1)$ random variables.

Proof. Conditional on $\tilde{x}_1, \dots, \tilde{x}_n$, letting $\hat{\sigma}_j^2 = \mathbb{E}_n[x_j^2]$, we have that

$$n\Lambda = \max_{j \in [p]} \sup_{\tau \in \mathcal{T}} \left| \frac{\sum_{i=1}^n \tilde{x}_{ij}(\tau - 1\{U_i \leq \tau\})}{\hat{\sigma}_j \sqrt{\tau(1-\tau)}} \right|.$$

Step 1. (Entropy Calculation) Let $\mathcal{F} = \{\tilde{x}_{ij}(\tau - 1\{U_i \leq \tau\})/\hat{\sigma}_j : \tau \in \mathcal{T}, j \in [p]\}$, $h_\tau = \sqrt{\tau(1-\tau)}$, and $\mathcal{G} = \{f_\tau/h_\tau : \tau \in \mathcal{T}\}$. We have that

$$\begin{aligned} d(f_\tau/h_\tau, f_{\bar{\tau}}/h_{\bar{\tau}}) &\leq d(f_\tau, f_{\bar{\tau}})/h_\tau + d(f_{\bar{\tau}}/h_\tau, f_{\bar{\tau}}/h_{\bar{\tau}}) \\ &\leq d(f_\tau, f_{\bar{\tau}})/h_\tau + d(0, f_{\bar{\tau}}/h_{\bar{\tau}})|h_\tau - h_{\bar{\tau}}|/h_\tau \end{aligned}$$

Therefore, since $\|F\|_Q \leq \|G\|_Q$ by $h_\tau \leq 1$, and $d(0, f_{\bar{\tau}}/h_{\bar{\tau}}) \leq 1/h_{\bar{\tau}}$ we have

$$N(\epsilon \|G\|_Q, \mathcal{G}, Q) \leq N(\epsilon \|F\|_Q / \{2 \min_{\tau \in \mathcal{T}} h_\tau\}, \mathcal{F}, Q) N(\epsilon / \{2 \min_{\tau \in \mathcal{T}} h_\tau^2\}, \mathcal{T}, |\cdot|).$$

Thus we have for some constants K and v that

$$N(\epsilon \|G\|_Q, \mathcal{G}, Q) \leq p(K / \{\epsilon \min_{\tau \in \mathcal{T}} h_\tau^2\})^v.$$

Step 2. (Symmetrization) Since we have $\mathbb{E}[g^2] = 1$ for all $g \in \mathcal{G}$, by Lemma 2.3.7 in [100] we have

$$\mathbb{P}(\Lambda \geq t\sqrt{n}) \leq 4\mathbb{P}(\max_{j \leq p} \sup_{\tau \in \mathcal{T}} |\mathbb{G}_n^o(g)| \geq t/4)$$

here $\mathbb{G}_n^o : \mathcal{G} \rightarrow \mathbb{R}$ is the symmetrized process generated by Rademacher variables. Conditional on $(x_1, u_1), \dots, (x_n, u_n)$, we have that $\{\mathbb{G}_n^o(g) : g \in \mathcal{G}\}$ is sub-Gaussian with respect to the $L_2(\mathbb{P}_n)$ -norm by the Hoeffding inequality. Thus, by Lemma 16 in [12], for $\delta_n^2 = \sup_{g \in \mathcal{G}} \mathbb{E}_n[g^2]$ and $\bar{\delta}_n = \delta_n / \|G\|_{\mathbb{P}_n}$, we have

$$\mathbb{P}(\sup_{g \in \mathcal{G}} |\mathbb{G}_n^o(g)| > CK\delta_n \sqrt{\log(pK/\underline{\tau})} \mid \{\tilde{x}_i, U_i\}_{i=1}^n) \leq \int_0^{\bar{\delta}_n/2} \epsilon^{-1} \{p(K/\{\epsilon \min_{\tau \in \mathcal{T}} h_\tau^2\})^v\}^{-C^2+1} d\epsilon$$

for some universal constant K .

In order to control δ_n , note that $\delta_n^2 = \sup_{g \in \mathcal{G}} \frac{1}{\sqrt{n}} \mathbb{G}_n(g^2) + \mathbb{E}[g^2]$. In turn, since $\sup_{g \in \mathcal{G}} \mathbb{E}_n[g^4] \leq \delta_n^2 \max_{i \leq n} G_i^2$, we have

$$\mathbb{P}(\sup_{g \in \mathcal{G}} |\mathbb{G}_n^o(g^2)| > C\bar{K}\delta_n \max_{i \leq n} G_i \sqrt{\log(pK/\underline{\tau})} \mid \{\tilde{x}_i, U_i\}_{i=1}^n) \leq \int_0^{\bar{\delta}_n/2} \epsilon^{-1} \{p(K/\{\epsilon \underline{\tau}\})^v\}^{-C^2+1} d\epsilon.$$

Thus with probability $1 - \int_0^{1/2} \epsilon^{-1} \{p(K/\{\epsilon \underline{\tau}\})^v\}^{-C^2+1} d\epsilon$, since $\mathbb{E}[g^2] = 1$ and $\max_{i \leq n} G_i \leq K_n/\sqrt{\underline{\tau}}$, we have

$$\delta_n \leq 1 + \frac{C'K_n \sqrt{\log(pK/\underline{\tau})}}{\sqrt{n}\sqrt{\underline{\tau}}}.$$

Therefore, under $K_n \sqrt{\log(pK/\underline{\tau})} = o(\sqrt{n}\sqrt{\underline{\tau}})$, conditionally on $\{\tilde{x}_i\}_{i=1}^n$ and n sufficiently large, with probability $1 - 2 \int_0^{1/2} \epsilon^{-1} \{p(K/\{\epsilon \underline{\tau}\})^v\}^{-C^2+1} d\epsilon$ we have that

$$\sup_{g \in \mathcal{G}} |\mathbb{G}_n^o(g)| \leq 2CK \sqrt{\log(pK/\underline{\tau})}$$

The stated bound follows since for $C > 2$

$$2 \int_0^{1/2} \epsilon^{-1} \{p(K/\{\epsilon \mathcal{I}\})^v\}^{-C^2+1} d\epsilon \leq \{p/\mathcal{I}\}^{-C^2+1} 2 \int_0^{1/2} \epsilon^{-2+C^2} d\epsilon \leq \{p/\mathcal{I}\}^{-C^2+1}.$$

■

APPENDIX G. INEQUALITIES

Lemma 16 (Transfer principle, [86]). *Let $\widehat{\Sigma}$ and Σ be $p \times p$ matrices with non-negative diagonal entries, and assume that for some $\eta \in (0, 1)$ and $s \leq p$ we have*

$$\forall v \in \mathbb{R}^p, \|v\|_0 \leq s, v' \widehat{\Sigma} v \geq (1 - \eta) v' \Sigma v$$

Let D be a diagonal matrix such that $D_{kk} \geq \widehat{\Sigma}_{kk} - (1 - \eta) \Sigma_{kk}$. Then for all $\delta \in \mathbb{R}^p$ we have

$$\delta' \widehat{\Sigma} \delta \geq (1 - \eta) \delta' \Sigma \delta - \|D^{1/2} \delta\|_1^2 / (s - 1).$$

Lemma 17. *Consider $\widehat{\beta}_u$ and β_u with $\|\beta_u\|_0 \leq s$. Denote by $\widehat{\beta}_u^\lambda$ the vector with $\widehat{\beta}_{uj}^\lambda = \widehat{\beta}_{uj} 1\{\widehat{\sigma}_{a\varpi j}^Z |\widehat{\beta}_{uj}| \geq \lambda\}$ where $\widehat{\sigma}_{a\varpi j}^Z = \{\mathbb{E}_n[K_\varpi(W)(Z_j^a)^2]\}^{1/2}$. We have that*

$$\begin{aligned} \|\widehat{\beta}_u^\lambda - \beta_u\|_{1,\varpi} &\leq \|\widehat{\beta}_u - \beta_u\|_{1,\varpi} + \lambda s \\ |\text{support}(\widehat{\beta}_u^\lambda)| &\leq s + \|\widehat{\beta}_u - \beta_u\|_{1,\varpi} / \lambda \\ \|Z^a(\widehat{\beta}_u^\lambda - \beta_u)\|_{n,\varpi} &\leq \|Z^a(\widehat{\beta}_u - \beta_u)\|_{n,\varpi} + \sqrt{\tilde{\phi}_{\max}(s, \varpi)} \{2\sqrt{s}\lambda + \|\widehat{\beta}_u - \beta_u\|_{1,\varpi} / \sqrt{s}\} \end{aligned}$$

here $\tilde{\phi}_{\max}(m, \varpi) = \sup_{1 \leq \|\theta\|_0 \leq m} \|\tilde{Z}^a \theta\|_{n,\varpi} / \|\theta\|$ and $\tilde{Z}_{ij}^a = Z_{ij}^a / \{\mathbb{E}_n[K_\varpi(W)(Z_j^a)^2]\}^{1/2}$.

Proof. Let $T_u = \text{support}(\beta_u)$. The first relation follows from the triangle inequality

$$\begin{aligned} \|\widehat{\beta}_u^\lambda - \beta_u\|_{1,\varpi} &= \|(\widehat{\beta}_u^\lambda - \beta_u)_{T_u}\|_{1,\varpi} + \|(\widehat{\beta}_u^\lambda)_{T_u^c}\|_{1,\varpi} \\ &\leq \|(\widehat{\beta}_u^\lambda - \widehat{\beta}_u)_{T_u}\|_{1,\varpi} + \|(\widehat{\beta}_u - \beta_u)_{T_u}\|_{1,\varpi} + \|(\widehat{\beta}_u^\lambda)_{T_u^c}\|_{1,\varpi} \\ &\leq \lambda s + \|(\widehat{\beta}_u - \beta_u)_{T_u}\|_{1,\varpi} + \|(\widehat{\beta}_u)_{T_u^c}\|_{1,\varpi} \\ &= \lambda s + \|\widehat{\beta}_u - \beta_u\|_{1,\varpi} \end{aligned}$$

To show the second result note that $\|\widehat{\beta}_u - \beta_u\|_{1,\varpi} \geq \{|\text{support}(\widehat{\beta}_u^\lambda)| - s\}\lambda$. Therefore,

$$|\text{support}(\widehat{\beta}_u^\lambda)| \leq s + \|\widehat{\beta}_u - \beta_u\|_{1,\varpi} / \lambda$$

which yields the result.

To show the third bound, we start using the triangle inequality

$$\|Z^a(\widehat{\beta}_u^\lambda - \beta_u)\|_{n,\varpi} \leq \|Z^a(\widehat{\beta}_u^\lambda - \widehat{\beta}_u)\|_{n,\varpi} + \|Z^a(\widehat{\beta}_u - \beta_u)\|_{n,\varpi}.$$

Without loss of generality, assume that the components are ordered so that $|(\widehat{\beta}_u^\lambda - \widehat{\beta}_u)_j| \widehat{\sigma}_{uj}$ is decreasing. Let T_1 be the set of s indices corresponding to the largest values of $|(\widehat{\beta}_u^\lambda - \widehat{\beta}_u)_j| \widehat{\sigma}_{uj}$. Similarly define T_k as the set of s indices corresponding to the largest values of $|(\widehat{\beta}_u^\lambda - \widehat{\beta}_u)_j| \widehat{\sigma}_{uj}$ outside $\cup_{m=1}^{k-1} T_m$. Therefore,

$\hat{\beta}_u^\lambda - \hat{\beta}_u = \sum_{k=1}^{\lceil p/s \rceil} (\hat{\beta}_u^\lambda - \hat{\beta}_u)_{T_k}$. Moreover, given the monotonicity of the components, $\|(\hat{\beta}_u^\lambda - \hat{\beta}_u)_{T_k}\|_{2,\varpi} \leq \|(\hat{\beta}_u^\lambda - \hat{\beta}_u)_{T_{k-1}}\|_{1,\varpi}/\sqrt{s}$. Then, we have

$$\begin{aligned} \|Z^a(\hat{\beta}_u^\lambda - \hat{\beta}_u)\|_{n,\varpi} &= \|Z^a \sum_{k=1}^{\lceil p/s \rceil} (\hat{\beta}_u^\lambda - \hat{\beta}_u)_{T_k}\|_{n,\varpi} \\ &\leq \|Z^a(\hat{\beta}_u^\lambda - \hat{\beta}_u)_{T_1}\|_{n,\varpi} + \sum_{k \geq 2} \|Z^a(\hat{\beta}_u^\lambda - \hat{\beta}_u)_{T_k}\|_{n,\varpi} \\ &\leq \sqrt{\tilde{\phi}_{\max}(s, \varpi)} \|(\hat{\beta}_u^\lambda - \hat{\beta}_u)_{T_1}\|_{2,\varpi} + \sqrt{\tilde{\phi}_{\max}(s, \varpi)} \sum_{k \geq 2} \|(\hat{\beta}_u^\lambda - \hat{\beta}_u)_{T_k}\|_{2,\varpi} \\ &\leq \sqrt{\tilde{\phi}_{\max}(s, \varpi)} \lambda \sqrt{s} + \sqrt{\tilde{\phi}_{\max}(s, \varpi)} \sum_{k \geq 1} \|(\hat{\beta}_u^\lambda - \hat{\beta}_u)_{T_k}\|_{1,\varpi} / \sqrt{s} \\ &= \sqrt{\tilde{\phi}_{\max}(s, \varpi)} \lambda \sqrt{s} + \sqrt{\tilde{\phi}_{\max}(s, \varpi)} \|\hat{\beta}_u^\lambda - \hat{\beta}_u\|_{1,\varpi} / \sqrt{s} \\ &\leq \sqrt{\tilde{\phi}_{\max}(s, \varpi)} \{2\lambda \sqrt{s} + \|\hat{\beta}_u - \beta_u\|_{1,\varpi} / \sqrt{s}\} \end{aligned}$$

here the last inequality follows from the first result and the triangle inequality. ■

Lemma 18 (Supremum of Sparse Vectors on Symmetrized Random Matrices). *Let $\hat{\mathcal{U}}$ denote a finite set and $(X_{iu})_{u \in \hat{\mathcal{U}}}$, $i = 1, \dots, n$, be fixed vectors such that $X_{iu} \in \mathbb{R}^p$ and $\max_{1 \leq i \leq n} \max_{u \in \hat{\mathcal{U}}} \|X_{iu}\|_\infty \leq K$. Furthermore define*

$$\delta_n := \bar{C} K \sqrt{k} \left(\sqrt{\log |\hat{\mathcal{U}}|} + \sqrt{1 + \log p} + \log k \sqrt{\log(p \vee n)} \sqrt{\log n} \right) / \sqrt{n},$$

where \bar{C} is a universal constant. Then,

$$\mathbb{E} \left[\sup_{\|\theta\|_0 \leq k, \|\theta\|=1} \max_{u \in \hat{\mathcal{U}}} |\mathbb{E}_n[\varepsilon(\theta' X_u)^2]| \right] \leq \delta_n \sup_{\|\theta\|_0 \leq k, \|\theta\|=1, u \in \hat{\mathcal{U}}} \sqrt{\mathbb{E}_n[(\theta' X_u)^2]}.$$

Proof. See [15] for the proof. ■

Corollary 2 (Supremum of Sparse Vectors on Many Random Matrices). *Let $\hat{\mathcal{U}}$ denote a finite set and $(X_{iu})_{u \in \hat{\mathcal{U}}}$, $i = 1, \dots, n$, be independent (across i) random vectors such that $X_{iu} \in \mathbb{R}^p$ and*

$$\sqrt{\mathbb{E}[\max_{1 \leq i \leq n} \max_{u \in \hat{\mathcal{U}}} \|X_{iu}\|_\infty^2]} \leq K.$$

Furthermore define

$$\delta_n := \bar{C} K \sqrt{k} \left(\sqrt{\log |\hat{\mathcal{U}}|} + \sqrt{1 + \log p} + \log k \sqrt{\log(p \vee n)} \sqrt{\log n} \right) / \sqrt{n},$$

here \bar{C} is a universal constant. Then,

$$\mathbb{E} \left[\sup_{\|\theta\|_0 \leq k, \|\theta\|=1} \max_{u \in \hat{\mathcal{U}}} |\mathbb{E}_n[(\theta' X_u)^2] - \mathbb{E}[(\theta' X_u)^2]| \right] \leq \delta_n^2 + \delta_n \sup_{\|\theta\|_0 \leq k, \|\theta\|=1, u \in \hat{\mathcal{U}}} \sqrt{\mathbb{E}_n[\mathbb{E}[(\theta' X_u)^2]]}.$$

We will also use the following result of [29].

Lemma 19 (Maximal Inequality). *Work with the setup above. Suppose that $F \geq \sup_{f \in \mathcal{F}} |f|$ is a measurable envelope for \mathcal{F} with $\|F\|_{P,q} < \infty$ for some $q \geq 2$. Let $M = \max_{i \leq n} F(W_i)$ and $\sigma^2 > 0$ be any positive constant such that $\sup_{f \in \mathcal{F}} \|f\|_{P,2}^2 \leq \sigma^2 \leq \|F\|_{P,2}^2$. Suppose that there exist constants $a \geq e$ and $v \geq 1$ such that*

$$\log \sup_Q N(\epsilon \|F\|_{Q,2}, \mathcal{F}, \|\cdot\|_{Q,2}) \leq v \log(a/\epsilon), \quad 0 < \epsilon \leq 1.$$

Then

$$\mathbb{E}_P[\sup_{f \in \mathcal{F}} |\mathbb{G}_n(f)|] \leq K \left(\sqrt{v\sigma^2 \log \left(\frac{a\|F\|_{P,2}}{\sigma} \right)} + \frac{v\|M\|_{P,2}}{\sqrt{n}} \log \left(\frac{a\|F\|_{P,2}}{\sigma} \right) \right),$$

here K is an absolute constant. Moreover, for every $t \geq 1$, with probability $> 1 - t^{-q/2}$,

$$\sup_{f \in \mathcal{F}} |\mathbb{G}_n(f)| \leq (1 + \alpha) \mathbb{E}_P[\sup_{f \in \mathcal{F}} |\mathbb{G}_n(f)|] + K(q) \left[(\sigma + n^{-1/2}\|M\|_{P,q})\sqrt{t} + \alpha^{-1}n^{-1/2}\|M\|_{P,2}t \right],$$

$\forall \alpha > 0$ where $K(q) > 0$ is a constant depends only on q . In particular, setting $a \geq n$ and $t = \log n$, with probability $> 1 - c(\log n)^{-1}$,

$$\sup_{f \in \mathcal{F}} |\mathbb{G}_n(f)| \leq K(q, c) \left(\sigma \sqrt{v \log \left(\frac{a\|F\|_{P,2}}{\sigma} \right)} + \frac{v\|M\|_{P,q}}{\sqrt{n}} \log \left(\frac{a\|F\|_{P,2}}{\sigma} \right) \right), \quad (\text{G.74})$$

here $\|M\|_{P,q} \leq n^{1/q}\|F\|_{P,q}$ and $K(q, c) > 0$ is a constant depending only on q and c .

APPENDIX H. CONFIDENCE REGIONS FOR FUNCTION-VALUED PARAMETERS BASED ON MOMENT CONDITIONS

For completeness, in this section we collect an adaptation of the results of [15] that are invoked in our proofs. The main difference is the weakening of the identification condition (which is allowed to decrease to zero, see the parameter j_n in Condition C1 below). We are interested in function-valued target parameters indexed by $u \in \mathcal{U} \subset \mathbb{R}^{d_u}$. The true value of the target parameter is denoted by

$$\theta^0 = (\theta_{uj})_{u \in \mathcal{U}, j \in [\tilde{p}]}, \text{ where } \theta_{uj} \in \Theta_{uj} \text{ for each } u \in \mathcal{U} \text{ and } j \in [\tilde{p}].$$

For each $u \in \mathcal{U}$ and $j \in [\tilde{p}]$, the parameter θ_{uj} is characterized as the solution to the following moment condition:

$$\mathbb{E}[\psi_{uj}(W_{uj}, \theta_{uj}, \eta_{uj})] = 0, \quad (\text{H.75})$$

where W_{uj} is a random vector that takes values in a Borel set $\mathcal{W}_{uj} \subset \mathbb{R}^{d_w}$, $\eta^0 = (\eta_{uj})_{u \in \mathcal{U}, j \in [\tilde{p}]}$ is a nuisance parameter where $\eta_{uj} \in T_{uj}$ a convex set, and the moment function

$$\psi_{uj} : \mathcal{W}_{uj} \times \Theta_{uj} \times T_{uj} \mapsto \mathbb{R}, \quad (w, \theta, t) \mapsto \psi_{uj}(w, \theta, t) \quad (\text{H.76})$$

is a Borel measurable map.

We assume that the (continuum) nuisance parameter η^0 can be modelled and estimated by $\hat{\eta} = (\hat{\eta}_{uj})_{u \in \mathcal{U}, j \in [\tilde{p}]}$. We will discuss examples where the corresponding η^0 can be estimated using modern regularization and post-selection methods such as Lasso and Post-Lasso (although other procedures can be applied). The estimator $\check{\theta}_{uj}$ of θ_{uj} is constructed as any approximate ϵ_n -solution in Θ_{uj} to a sample analog of the moment condition (H.75), i.e.,

$$\max_{j \in [\tilde{p}]} \sup_{u \in \mathcal{U}} \left\{ |\mathbb{E}_n[\psi_{uj}(W_{uj}, \check{\theta}_{uj}, \hat{\eta}_{uj})]| - \inf_{\theta_j \in \Theta_{uj}} |\mathbb{E}_n[\psi_{uj}(W_{uj}, \theta, \hat{\eta}_{uj})]| \right\} \leq \epsilon_n = o_P(n^{-1/2}\delta_n). \quad (\text{H.77})$$

As discussed before, we rely on an orthogonality condition for regular estimation of θ_{uj} , which we will state next.

Definition 1 (Near Orthogonality Condition). For each $u \in \mathcal{U}$ and $j \in [\tilde{p}]$, we say that ψ_{uj} obeys a general form of orthogonality with respect to \mathcal{H}_{uj} uniformly in $u \in \mathcal{U}$, if the following conditions hold: the Gâteaux derivative map

$$D_{u,j,\bar{r}}[\tilde{\eta}_{uj} - \eta_{uj}] := \partial_r E \left(\psi_{uj} \left\{ W_{uj}, \theta_{uj}, \eta_{uj} + r [\tilde{\eta}_{uj} - \eta_{uj}] \right\} \right) \Big|_{r=\bar{r}}$$

exists for all $r \in [0, 1)$, $\tilde{\eta} \in \mathcal{H}_{uj}$, $j \in \tilde{p}$, and $u \in \mathcal{U}$ and vanishes at $r = 0$, namely,

$$|D_{u,j,0}[\tilde{\eta}_{uj} - \eta_{uj}]| \leq \delta_n n^{-1/2} \quad \text{for all } \tilde{\eta}_{uj} \in \mathcal{H}_{uj}. \quad (\text{H.78})$$

In what follows, we shall denote by c_0 , c , and C some positive constants.

Assumption C1 (Moment Condition). Consider a random element W , taking values in a measure space $(\mathcal{W}, \mathcal{A}_W)$, with law determined by a probability measure $P \in \mathcal{P}_n$. The observed data $((W_{iu})_{u \in \mathcal{U}})_{i=1}^n$ consist of n i.i.d. copies of a random element $(W_u)_{u \in \mathcal{U}}$ which is generated as a suitably measurable transformation with respect to W and u . Uniformly for all $n \geq n_0$ and $P \in \mathcal{P}_n$, the following conditions hold: (i) The true parameter value θ_{uj} obeys (H.75) and is interior relative to Θ_{uj} , namely there is a ball of radius $Cn^{-1/2}u_n \log n$ centered at θ_{uj} contained in Θ_{uj} for all $u \in \mathcal{U}$, $j \in [\tilde{p}]$ with $u_n := E[\sup_{u \in \mathcal{U}, j \in [\tilde{p}]} |\sqrt{n} E_n[\psi_{uj}(W_{uj}, \theta_{uj}, \eta_{uj})]|]$; (ii) For each $u \in \mathcal{U}$ and $j \in [\tilde{p}]$, the map $(\theta, \eta) \in \Theta_{uj} \times \mathcal{H}_{uj} \mapsto E[\psi_{uj}(W_{uj}, \theta, \eta)]$ is twice continuously differentiable; (iii) For all $u \in \mathcal{U}$ and $j \in [\tilde{p}]$, the moment function ψ_{uj} obeys the orthogonality condition given in Definition 1 for the set $\mathcal{H}_{uj} = \mathcal{H}_{ujn}$ specified in Assumption C2; (iv) The following identifiability condition holds: $|E[\psi_{uj}(W_{uj}, \theta, \eta_{uj})]| \geq \frac{1}{2} |J_{uj}(\theta - \theta_{uj})| \wedge c_0$ for all $\theta \in \Theta_{uj}$, with $J_{uj} := \partial_\theta E[\psi_{uj}(W_{uj}, \theta, \eta_{uj})]|_{\theta=\theta_{uj}}$ satisfies $0 < j_n < |J_{uj}| < C < \infty$ for all $u \in \mathcal{U}$ and $j \in [\tilde{p}]$; (v) The following smoothness conditions holds

$$\begin{aligned} \text{(a)} \quad & \sup_{u \in \mathcal{U}, j \in [\tilde{p}], (\theta, \bar{\theta}) \in \Theta_{uj}^2, (\eta, \bar{\eta}) \in \mathcal{H}_{ujn}^2} \frac{E[\{\psi_{uj}(W_{uj}, \theta, \eta) - \psi_{uj}(W_{uj}, \bar{\theta}, \bar{\eta})\}^2]}{\{|\theta - \bar{\theta}| \vee \|\eta - \bar{\eta}\|_e\}^\alpha} \leq C, \\ \text{(b)} \quad & \sup_{u \in \mathcal{U}, (\theta, \eta) \in \Theta_{uj} \times \mathcal{H}_{ujn}, r \in [0, 1]} |\partial_r E[\psi_{uj}(W_{uj}, \theta, \eta_{uj} + r\{\eta - \eta_{uj}\})]| / \|\eta - \eta_{uj}\|_e \leq \bar{B}_{1n}, \\ \text{(c)} \quad & \sup_{u \in \mathcal{U}, j \in [\tilde{p}], (\theta, \eta) \in \Theta_{uj} \times \mathcal{H}_{ujn}, r \in [0, 1]} \frac{|\partial_r^2 E[\psi_{uj}(W_{uj}, \theta_{uj} + r\{\theta - \theta_{uj}\}, \eta_{uj} + r\{\eta - \eta_{uj}\})]|}{\{|\theta - \theta_{uj}|^2 \vee \|\eta - \eta_{uj}\|_e^2\}} \leq \bar{B}_{2n}. \end{aligned}$$

Next we state assumptions on the nuisance functions. In what follows, let $\Delta_n \searrow 0$, $\delta_n \searrow 0$, and $\tau_n \searrow 0$ be sequences of constants approaching zero from above at a speed at most polynomial in n (for example, $\delta_n \geq 1/n^c$ for some $c > 0$).

Assumption C2 (Estimation of Nuisance Functions). The following conditions hold for each $n \geq n_0$ and all $P \in \mathcal{P}_n$. The estimated functions $\hat{\eta}_{uj} \in \mathcal{H}_{ujn}$ with probability at least $1 - \Delta_n$, \mathcal{H}_{ujn} is the set of measurable maps $\tilde{\eta}_{uj}$ such that

$$\sup_{u \in \mathcal{U}} \max_{j \in [\tilde{p}]} \|\tilde{\eta}_{uj} - \eta_{uj}\|_e \leq \tau_n,$$

here the e -norm is the same as in Assumption C1, and whose complexity does not grow too quickly in the sense that $\mathcal{F}_1 = \{\psi_{uj}(W_{uj}, \theta, \eta) : u \in \mathcal{U}, j \in [\tilde{p}], \theta \in \Theta_{uj}, \eta \in \mathcal{H}_{ujn} \cup \{\eta_{uj}\}\}$ is suitably measurable and its uniform covering entropy obeys:

$$\sup_Q \log N(\epsilon \|F_1\|_{Q,2}, \mathcal{F}_1, \|\cdot\|_{Q,2}) \leq s_{n(\mathcal{U}, \tilde{p})}(\log(a_n/\epsilon)) \vee 0,$$

where $F_1(W)$ is an envelope for \mathcal{F}_1 which is measurable with respect to W and satisfies $F_1(W) \geq \sup_{u \in \mathcal{U}, j \in [\tilde{p}], \theta \in \Theta_{uj}, \eta \in \mathcal{H}_{ujn}} |\psi_{uj}(W_{uj}, \theta, \eta)|$ and $\|F_1\|_{P,q} \leq K_n$ for $q \geq 2$. The complexity characteristics

$a_n \geq \max(n, K_n, e)$ and $s_{n(\mathcal{U}, \bar{p})} \geq 1$ obey the growth conditions:

$$\begin{aligned} n^{-1/2} \sqrt{s_{n(\mathcal{U}, \bar{p})} \log(a_n)} + n^{-1} s_{n(\mathcal{U}, \bar{p})} n^{\frac{1}{q}} K_n \log(a_n) &\leq \tau_n \\ \{(1 \vee \bar{B}_{1n})(\tau_n/j_n)\}^{\alpha/2} \sqrt{s_{n(\mathcal{U}, \bar{p})} \log(a_n)} + s_{n(\mathcal{U}, \bar{p})} n^{\frac{1}{q} - \frac{1}{2}} K_n \log(a_n) \log n &\leq \delta_n, \\ \text{and } \sqrt{n} \bar{B}_{2n} (1 \vee \bar{B}_{1n})(\tau_n/j_n)^2 &\leq \delta_n \end{aligned}$$

here \bar{B}_{1n} , \bar{B}_{2n} , j_n , q and α are defined in Assumption C1.

Theorem 5 (Uniform Bahadur representation for a Continuum of Target Parameters). *Under Assumptions C1 and C2, for an estimator $(\check{\theta}_{uj})_{u \in \mathcal{U}, j \in [\bar{p}]}$ that obeys equation (H.77),*

$$\sqrt{n} \sigma_{uj}^{-1} (\check{\theta}_{uj} - \theta_{uj}) = \mathbb{G}_n \bar{\psi}_{uj} + O_P(\delta_n) \text{ in } \ell^\infty(\mathcal{U} \times [\bar{p}]), \text{ uniformly in } P \in \mathcal{P}_n,$$

here $\bar{\psi}_{uj}(W) := -\sigma_{uj}^{-1} J_{uj}^{-1} \psi_{uj}(W_{uj}, \theta_{uj}, \eta_{uj})$ and $\sigma_{uj}^2 = E[J_{uj}^{-2} \psi_{uj}^2(W_{uj}, \theta_{uj}, \eta_{uj})]$.

The uniform Bahadur representation derived in Theorem 5 is useful in the construction of simultaneous confidence bands for $(\theta_{uj})_{u \in \mathcal{U}, j \in [\bar{p}]}$. This is achieved by new high-dimensional central limit theorems that have recently been developed in [28] and [29]. We will make use of the following regularity condition. In what follows $\bar{\delta}_n$ and Δ_n are fixed sequences going to zero, and we denote $\widehat{\psi}_{uj}(W) := -\widehat{\sigma}_{uj}^{-1} \widehat{J}_{uj}^{-1} \psi_{uj}(W_{uj}, \check{\theta}_{uj}, \widehat{\eta}_{uj})$ be the estimators of $\bar{\psi}_{uj}(W)$, with \widehat{J}_{uj} and $\widehat{\sigma}_{uj}$ being suitable estimators of J_{uj} and σ_{uj} . In what follows, $\|\cdot\|_{\mathbb{P}_n, 2}$ denotes the empirical $L_2(\mathbb{P}_n)$ -norm with \mathbb{P}_n as the empirical measure of the data.

Assumption C3 (Score Regularity). *The following conditions hold for each $n \geq n_0$ and all $P \in \mathcal{P}_n$. (i) The class of function induced by the score $\mathcal{F}_0 = \{\bar{\psi}_{uj}(W) : u \in \mathcal{U}, j \in [\bar{p}]\}$ is suitably measurable and its uniform covering entropy obeys:*

$$\sup_Q \log N(\epsilon \|F_0\|_{Q, 2}, \mathcal{F}_0, \|\cdot\|_{Q, 2}) \leq \varrho_n(\log(A_n/\epsilon)) \vee 0,$$

here $F_0(W)$ is an envelope for \mathcal{F}_0 which is measurable with respect to W and satisfies $F_0(W) \geq \sup_{u \in \mathcal{U}, j \in [\bar{p}]} |\bar{\psi}_{uj}(W)|$ and $\|F_0\|_{P, q} \leq L_n$ for $q \geq 4$. Furthermore, $c \leq \sup_{u \in \mathcal{U}, j \in [\bar{p}]} E[|\bar{\psi}_{uj}(W)|^k] \leq CL_n^{k-2}$ for $k = 2, 3, 4$. (ii) The set $\widehat{\mathcal{F}}_0 = \{\bar{\psi}_{uj}(W) - \widehat{\psi}_{uj}(W) : u \in \mathcal{U}, j \in [\bar{p}]\}$ satisfies the conditions $\log N(\epsilon, \widehat{\mathcal{F}}_0, \|\cdot\|_{\mathbb{P}_n, 2}) \leq \bar{\varrho}_n(\log(\bar{A}_n/\epsilon)) \vee 0$, and $\sup_{u \in \mathcal{U}, j \in [\bar{p}]} E_n[\{\bar{\psi}_{uj}(W) - \widehat{\psi}_{uj}(W)\}^2] \leq \bar{\delta}_n \{\rho_n \bar{\rho}_n \log(A_n \vee n) \log(\bar{A}_n \vee n)\}^{-1}$ with probability $1 - \Delta_n$.

Assumption C3 imposes condition on the class of functions induced by $\bar{\psi}_{uj}$ and on its estimators $\widehat{\psi}_{uj}$. Typically the bound L_n on the moment of the envelope is smaller than K_n , and in many settings $\bar{\rho}_n = \rho_n \lesssim d_{\mathcal{U}}$ the dimension of \mathcal{U} .

Next let \mathcal{N} denote a mean zero Gaussian process indexed by $\mathcal{U} \times [\bar{p}]$ with covariance operator given by $E[\bar{\psi}_{uj}(W) \bar{\psi}_{u'j'}(W)]$ for $j, j' \in [\bar{p}]$ and $u, u' \in \mathcal{U}$. Because of the high-dimensionality, indeed \bar{p} can be larger than the sample size n , the central limit theorem will be uniformly valid over “rectangles”. This class of sets are rich enough to construct many confidence regions of interest in applications accounting for multiple testing. Let \mathcal{R} denote the set of rectangles $R = \{z \in \mathbb{R}^{\bar{p}} : \max_{j \in A} z_j \leq t, \max_{j \in B} (-z_j) \leq t\}$ for all $A, B \subset [\bar{p}]$ and $t \in \mathbb{R}$. The following result is a consequence of Theorem 5 above and Corollary 2.2 of [30].

Corollary 3. *Under Assumptions C1, C2 and Assumption C3(i), with $\delta_n = o(\{\rho_n \log(A_n \vee n)\}^{-1/2})$, and $\rho_n \log(A_n \vee n) = o(\{(n/L_n^2)^{1/7} \wedge (n^{1-2/q}/L_n^2)^{1/3}\})$, we have that*

$$\sup_{P \in \mathcal{P}_n} \sup_{R \in \mathcal{R}} \left| P_P \left(\left\{ \sup_{u \in \mathcal{U}} n^{1/2} \sigma_{uj}^{-1} (\check{\theta}_{uj} - \theta_{uj}) \right\}_{j=1}^{\bar{p}} \in R \right) - P_P(\mathcal{N} \in R) \right| = o(1).$$

In order to derive a method to build confidence regions we approximate the process \mathcal{N} by the Gaussian multiplier bootstrap based on estimates $\hat{\psi}_{uj}$ of $\bar{\psi}_{uj}$, namely

$$\hat{\mathcal{G}} = (\hat{\mathcal{G}}_{uj})_{u \in \mathcal{U}, j \in [\bar{p}]} = \left\{ \frac{1}{\sqrt{n}} \sum_{i=1}^n g_i \hat{\psi}_{uj}(W_i) \right\}_{u \in \mathcal{U}, j \in [\bar{p}]}$$

here $(g_i)_{i=1}^n$ are independent standard normal random variables which are independent from the data $(W_i)_{i=1}^n$. Based on Theorem 5.2 of [28], the following result shows that the multiplier bootstrap provides a valid approximation to the large sample probability law of $\sqrt{n}(\check{\theta}_{uj} - \theta_{uj})_{u \in \mathcal{U}, j \in [\bar{p}]}$ over rectangles.

Corollary 4 (Uniform Validity of Gaussian Multiplier Bootstrap). *Under Assumptions C1, C2 and Assumption C3, with $\delta_n = o(\{(1 + d_W)\rho_n \log(A_n \vee n)\}^{-1/2})$ and $\rho_n \log(A_n \vee n) = o(\{(n/L_n^2)^{1/7} \wedge (n^{1-2/q}/L_n^2)^{1/3}\})$, we have that*

$$\sup_{P \in \mathcal{P}_n} \sup_{R \in \mathcal{R}} \left| \mathbb{P}_P \left(\left\{ \sup_{u \in \mathcal{U}} n^{1/2} \sigma_{uj}^{-1} (\check{\theta}_{uj} - \theta_{uj}) \right\}_{j=1}^{\bar{p}} \in R \right) - \mathbb{P}_P(\hat{\mathcal{G}} \in R \mid (W_i)_{i=1}^n) \right| = o(1)$$

APPENDIX I. CONTINUUM OF ℓ_1 -PENALIZED M-ESTIMATORS

For the reader's convenience, this section collects results on the estimation of a continuum of estimation of high-dimensional models via ℓ_1 -penalized estimators. We refer to [15] for the proofs.

Consider a data generating process with a response variable $(Y_u)_{u \in \mathcal{U}}$ and observable covariates $(X_u)_{u \in \mathcal{U}}$ satisfies for each $u \in \mathcal{U}$,

$$\theta_u \in \arg \min_{\theta \in \mathbb{R}^p} \mathbb{E}[M_u(Y_u, X_u, \theta, a_u)], \quad (\text{I.79})$$

here θ_u is a p -dimensional vector, a_u is a nuisance function that capture the misspecification of the model, M_u is a pre-specified function, and the p_u -dimensional ($p_u \leq p$) covariate X_u could have been constructed based on transformations of other variables. This implies that

$$\partial_\theta \mathbb{E}[M_u(Y_u, X_u, \theta_u, a_u)] = 0 \quad \text{for all } u \in \mathcal{U}.$$

The solution θ_u is assumed to be sparse in the sense that for some process $(\theta_u)_{u \in \mathcal{U}}$ satisfies

$$\|\theta_u\|_0 \leq s \quad \text{for all } u \in \mathcal{U}.$$

Because of the nuisance function, such sparsity assumption is very mild and formulation (I.79) encompasses several cases of interest including approximate sparse models. We focus on the estimation of $(\theta_u)_{u \in \mathcal{U}}$ and we assume that an estimate \hat{a}_u of the nuisance function a_u is available and the criterion $M_u(Y_u, X_u, \theta_u) := M_u(Y_u, X_u, \theta_u, \hat{a}_u)$ is used as a proxy for $M_u(Y_u, X_u, \theta_u, a_u)$.

In the case of linear regression we have $M_u(y, x, \theta) = \frac{1}{2}(y - x'\theta)^2$. In the logistic regression case, we have $M_u(y, x, \theta) = -\{1(y = 1) \log G(x'\theta) + 1(y = 0) \log(1 - G(x'\theta))\}$ with G is the logistic link function $G(t) = \exp(t)/(1 + \exp(t))$. Additional examples include quantile regression models for $u \in (0, 1)$.

Example 8 (Quantile Regression Model). Consider a data generating process $Y = F_{Y|X}^{-1}(U) = X'\theta_U + r_U(X)$, with $U \sim \text{Unif}(0, 1)$, and X is a p -dimensional vector of covariates. The criterion $M_u(y, x, \theta) = (u - 1\{y \leq x'\theta\})(y - x'\theta)$ with the (trivial) estimate $\hat{a}_u = 0$ for the nuisance parameter $a_u = r_u$.

Example 9 (Lasso with Estimated Weights). We consider a linear model defined as $f_u Y = f_u X' \theta_u + \bar{r}_u + \zeta_u$, $\mathbb{E}[f_u X \zeta_u] = 0$, here X are \bar{p} -dimensional covariates, θ_u is a s -sparse vector, and \bar{r}_u is an approximation error satisfies $\sup_{u \in \mathcal{U}} \mathbb{E}_n[\bar{r}_u^2] \lesssim_P s \log \bar{p}/n$. In this setting, (Y, X) are observed and only an estimator \hat{f}_u of f_u is available. This corresponds to nuisance parameter $a_u = (f_u, \bar{r}_u)$ and $\hat{a}_u = (\hat{f}_u, 0)$ so that $\mathbb{E}_n[M_u(Y, X, \theta, a_u)] = \mathbb{E}_n[f_u^2(Y - X'\theta - \bar{r}_u)^2]$ and $\mathbb{E}_n[M_u(Y, X, \theta)] = \mathbb{E}_n[\hat{f}_u^2(Y - X'\theta)^2]$.

We assume that n i.i.d. observations from dgps with (I.79) holds, $\{(Y_{iu}, X_{iu})_{u \in \mathcal{U}}\}_{i=1}^n$, are observed to estimate $(\theta_u)_{u \in \mathcal{U}}$. For each $u \in \mathcal{U}$, a penalty level λ , and a diagonal matrix of penalty loadings $\hat{\Psi}_u$, we define the ℓ_1 -penalized M_u -estimator (Weighed-Lasso) as

$$\hat{\theta}_u \in \arg \min_{\theta} \mathbb{E}_n[M_u(Y_u, X_u, \theta)] + \frac{\lambda}{n} \|\hat{\Psi}_u \theta\|_1. \quad (\text{I.80})$$

Furthermore, for each $u \in \mathcal{U}$, the post-penalized estimator (Post-Lasso) based on a set of covariates \tilde{T}_u is then defined as

$$\tilde{\theta}_u \in \arg \min_{\theta} \mathbb{E}_n[M_u(Y_u, X_u, \theta)] \quad : \quad \text{support}(\theta) \subseteq \tilde{T}_u. \quad (\text{I.81})$$

Potentially, the set \tilde{T}_u contains $\text{support}(\hat{\theta}_u)$ and possibly additional variables deemed as important (although in that case the total number of additional variables should also obey the same growth conditions that s obeys). We will set $\tilde{T}_u = \text{support}(\hat{\theta}_u)$ unless otherwise noted.

In order to handle the functional response data, the penalty level λ and penalty loading $\hat{\Psi}_u = \text{diag}(\{\hat{l}_{uk}, k = 1, \dots, p\})$ need to be set to control selection errors uniformly over $u \in \mathcal{U}$. The choice of loading matrix is problem specific and we suggest to mimic the following “ideal” choice $\hat{\Psi}_{u0} = \text{diag}(\{l_{uk}, k = 1, \dots, p\})$ with

$$l_{uk} = \{\mathbb{E}_n[\{\partial_{\theta_k} M_u(Y_u, X_u, \theta_u, a_u)\}^2]\}^{1/2} \quad (\text{I.82})$$

which is motivated by the use of self-normalized moderate deviation theory. In that case, it is suitable to set λ so that with high probability

$$\frac{\lambda}{n} \geq c \sup_{u \in \mathcal{U}} \left\| \hat{\Psi}_{u0}^{-1} \mathbb{E}_n[\partial_{\theta} M_u(Y_u, X_u, \theta_u, a_u)] \right\|_{\infty}, \quad (\text{I.83})$$

here $c > 1$ is a fixed constant. Indeed, in the case that \mathcal{U} is a singleton the choice above is similar to [22], [14], and [21]. This approach was first employed for a continuum of indices \mathcal{U} in the context of ℓ_1 -penalized quantile regression processes by [12].

To implement (I.83), we propose setting the penalty level as

$$\lambda = c\sqrt{n}\Phi^{-1}(1 - \xi/\{2pN_n\}), \quad (\text{I.84})$$

here N_n is a measure of the class of functions indexed by \mathcal{U} , $1 - \xi$ (with $\xi = o(1)$) is a confidence level associated with the probability of event (I.83), and $c > 1$ is a slack constant. In many settings we can take $N_n = n^{d_{\mathcal{U}}}$. If the set \mathcal{U} is a singleton, $N_n = 1$ suffices which corresponds to what is used in [18].

I.1. Generic Finite Sample Bounds. In this subsection we derive finite sample bounds based on Assumption C4 below. This assumption provides sufficient conditions that are implied by a variety of settings including generalized linear models.

Assumption C4 (M-Estimation Conditions). *Let $\{(Y_{iu}, X_{iu}, u \in \mathcal{U}), i = 1, \dots, n\}$ be n i.i.d. observations of the model (I.79) and let $T_u = \text{support}(\theta_u)$, here $\|T_u\|_0 \leq s$, $u \in \mathcal{U}$. With probability $1 - \Delta_n$ we have that for all $u \in \mathcal{U}$ there are weights $w_u = w_u(Y_u, X_u)$ and C_{un} such that:*

- (a) $|\mathbb{E}_n[\partial_\theta M_u(Y_u, X_u, \theta_u) - \partial_\theta M_u(Y_u, X_u, \theta_u, a_u)]'\delta| \leq C_{un} \|\sqrt{w_u} X'_u \delta\|_{\mathbb{P}_{n,2}};$
- (b) $\ell \widehat{\Psi}_{u0} \leq \widehat{\Psi}_u \leq L \widehat{\Psi}_{u0}$ for $\ell > 1/c$, and let $\tilde{c} = \frac{Lc+1}{\ell c-1} \sup_{u \in \mathcal{U}} \|\widehat{\Psi}_{u0}\|_\infty \|\widehat{\Psi}_{u0}^{-1}\|_\infty;$
- (c) for all $\delta \in A_u$ there is $\bar{q}_{A_u} > 0$ such that

$$\begin{aligned} \mathbb{E}_n[M_u(Y_u, X_u, \theta_u + \delta)] - \mathbb{E}_n[M_u(Y_u, X_u, \theta_u)] - \mathbb{E}_n[\partial_\theta M_u(Y_u, X_u, \theta_u)]'\delta + 2C_{un} \|\sqrt{w_u} X'_u \delta\|_{\mathbb{P}_{n,2}} \\ \geq \left\{ \|\sqrt{w_u} X'_u \delta\|_{\mathbb{P}_{n,2}}^2 \right\} \wedge \left\{ \bar{q}_{A_u} \|\sqrt{w_u} X'_u \delta\|_{\mathbb{P}_{n,2}} \right\}. \end{aligned}$$

In many applications we take the weights to be $w_u = w_u(X_u) = 1$ but we allow for more general weights. Assumption C4(a) bounds the impact of estimating the nuisance functions uniformly over $u \in \mathcal{U}$. In the setting with s -sparse estimands, we typically have $C_{un} \lesssim \{n^{-1}s \log(pn)\}^{1/2}$. The loadings $\widehat{\Psi}_u$ are assumed larger (but not too much larger) than the ideal choice $\widehat{\Psi}_{u0}$ defined in (I.82). This is formalized in Assumption C4(b). Assumption C4(c) is an identification condition that will be imposed for specific choices of A_u and q_{A_u} . It relates to conditions in the literature derived for the case of a singleton \mathcal{U} and no nuisance functions, see the restricted strong convexity¹⁶ used in [83] and the non-linear impact coefficients used in [12] and [19].

The following results establish rates of convergence for the ℓ_1 -penalized solution with estimated nuisance functions (I.80), sparsity bounds and rates of convergence for the post-selection refitted estimator (I.81). They are based on restricted eigenvalue type conditions and sparse eigenvalue conditions. With the restricted eigenvalue is defined as $\bar{\kappa}_{u,2\tilde{c}} = \inf_{\delta \in \Delta_{u,2\tilde{c}}} \|\sqrt{w_u} X'_u \delta\|_{\mathbb{P}_{n,2}} / \|\delta\|_{T_u}$. In the results for sparsity and post-selection refitted models, the minimum and maximum sparse eigenvalues,

$$\phi_{\min}(m, u) = \min_{1 \leq \|\delta\|_0 \leq m} \frac{\|\sqrt{w_u} X'_u \delta\|_{\mathbb{P}_{n,2}}^2}{\|\delta\|^2} \quad \text{and} \quad \phi_{\max}(m, u) = \max_{1 \leq \|\delta\|_0 \leq m} \frac{\|\sqrt{w_u} X'_u \delta\|_{\mathbb{P}_{n,2}}^2}{\|\delta\|^2},$$

are also relevant quantities to characterize the behavior of the estimators.

Lemma 20. *Suppose that Assumption C4 holds with $\delta \in A_u = \{\delta : \|\delta_{T_u^c}\|_1 \leq 2\tilde{c}\|\delta_{T_u}\|_1\} \cup \{\delta : \|\delta\|_1 \leq \frac{6c\|\widehat{\Psi}_{u0}^{-1}\|_\infty}{\ell c-1} \frac{n}{\lambda} C_{un} \|\sqrt{w_u} X'_u \delta\|_{\mathbb{P}_{n,2}}\}$ and $\bar{q}_{A_u} > 3 \left\{ (L + \frac{1}{c}) \|\widehat{\Psi}_{u0}\|_\infty \frac{\lambda\sqrt{s}}{n\bar{\kappa}_{u,2\tilde{c}}} + 9\tilde{c}C_{un} \right\}$. Suppose that λ satisfies condition (I.83) with probability $1 - \Delta_n$. Then, with probability $1 - 2\Delta_n$ we have uniformly over $u \in \mathcal{U}$*

$$\begin{aligned} \|\sqrt{w_u} X'_u (\widehat{\theta}_u - \theta_u)\|_{\mathbb{P}_{n,2}} &\leq 3 \left\{ (L + \frac{1}{c}) \|\widehat{\Psi}_{u0}\|_\infty \frac{\lambda\sqrt{s}}{n\bar{\kappa}_{u,2\tilde{c}}} + 9\tilde{c}C_{un} \right\} \\ \|\widehat{\theta}_u - \theta_u\|_1 &\leq 3 \left\{ \frac{(1+2\tilde{c})\sqrt{s}}{\bar{\kappa}_{u,2\tilde{c}}} + \frac{6c\|\widehat{\Psi}_{u0}^{-1}\|_\infty}{\ell c-1} \frac{n}{\lambda} C_{un} \right\} \left\{ (L + \frac{1}{c}) \|\widehat{\Psi}_{u0}\|_\infty \frac{\lambda\sqrt{s}}{n\bar{\kappa}_{u,2\tilde{c}}} + 9\tilde{c}C_{un} \right\} \end{aligned}$$

Lemma 21 (M-Estimation Sparsity). *In addition to conditions of Lemma 20, assume that with probability $1 - \Delta_n$ for all $u \in \mathcal{U}$ and $\delta \in \mathbb{R}^p$ we have*

$$|\{\mathbb{E}_n[\partial_\theta M_u(Y_u, X_u, \widehat{\theta}_u) - \partial_\theta M_u(Y_u, X_u, \theta_u)]'\delta| \leq L_{un} \|\sqrt{w_u} X'_u \delta\|_{\mathbb{P}_{n,2}}.$$

Let $\mathcal{M}_u = \{m \in \mathbb{N} : m \geq 2\phi_{\max}(m, u)L_u^2\}$ with $L_u = \frac{c\|\widehat{\Psi}_{u0}^{-1}\|_\infty}{\ell c-1} \frac{n}{\lambda} \{C_{un} + L_{un}\}$, then with probability $1 - 3\Delta_n$ we have that

$$\widehat{s}_u \leq \min_{m \in \mathcal{M}_u} \phi_{\max}(m, u)L_u^2 \quad \text{for all } u \in \mathcal{U}.$$

¹⁶Assumption C4 (a) and (c) could have been stated with $\{C_{un}/\sqrt{s}\}\|\delta\|_1$ instead of $C_{un}\|\sqrt{w_u} X'_u \delta\|_{\mathbb{P}_{n,2}}$.

Lemma 22. Let $\tilde{T}_u, u \in \mathcal{U}$, be the support used for post penalized estimator (I.81) and $\tilde{s}_u = \|\tilde{T}_u\|_0$ its cardinality. In addition to conditions of Lemma 20, suppose that Assumption C4(c) holds also for $A_u = \{\delta : \|\delta\|_0 \leq \tilde{s}_u + s\}$ with probability $1 - \Delta_n$, $\bar{q}_{A_u} > 2 \left\{ \frac{\sqrt{\tilde{s}_u + s_u} \|\mathbb{E}_n[S_u]\|_\infty}{\sqrt{\phi_{\min}(\tilde{s}_u + s_u, u)}} + 3C_{un} \right\}$ and $\bar{q}_{A_u} > 2\{\mathbb{E}_n[M_u(Y_u, X_u, \tilde{\theta}_u)] - \mathbb{E}_n[M_u(Y_u, X_u, \theta_u)]\}_+^{1/2}$. Then, we have uniformly over $u \in \mathcal{U}$

$$\|\sqrt{w_u} X'_u(\tilde{\theta}_u - \theta_u)\|_{\mathbb{P}_{n,2}} \leq \{\mathbb{E}_n[M_u(Y_u, X_u, \tilde{\theta}_u)] - \mathbb{E}_n[M_u(Y_u, X_u, \theta_u)]\}_+^{1/2} + \frac{\sqrt{\tilde{s}_u + s_u} \|\mathbb{E}_n[S_u]\|_\infty}{\sqrt{\phi_{\min}(\tilde{s}_u + s_u, u)}} + 3C_{un}.$$

In Lemma 22, if $\tilde{T}_u = \text{support}(\hat{\theta}_u)$, we have that

$$\mathbb{E}_n[M_u(Y_u, X_u, \tilde{\theta}_u)] - \mathbb{E}_n[M_u(Y_u, X_u, \theta_u)] \leq \mathbb{E}_n[M_u(Y_u, X_u, \hat{\theta}_u)] - \mathbb{E}_n[M_u(Y_u, X_u, \theta_u)] \leq \lambda C' \|\hat{\theta}_u - \theta_u\|_1$$

and $\sup_{u \in \mathcal{U}} \|\mathbb{E}_n[S_u]\|_\infty \leq C' \lambda$ with high probability, $C' \leq L \sup_{u \in \mathcal{U}} \|\hat{\Psi}_{u0}\|_\infty$.

These results generalize important results of the ℓ_1 -penalized estimators to the case of functional response data and estimated of nuisance functions. A key assumption in Lemmas 20-22 is that the choice of λ satisfies (I.83). We next provide a set of simple generic conditions that will imply the validity of the proposed choice. These generic conditions can be verified in many applications of interest.

Condition WL. For each $u \in \mathcal{U}$, let $S_u = \partial_\theta M_u(Y_u, X_u, \theta_u, a_u)$, suppose that:

- (i) $\sup_{u \in \mathcal{U}} \max_{k \leq p} \{\mathbb{E}[|S_{uk}|^3]\}^{1/3} / \{\mathbb{E}[|S_{uk}|^2]\}^{1/2} \Phi^{-1}(1 - \xi / \{2pN_n\}) \leq \delta_n n^{1/6}$, for all $u \in \mathcal{U}$, $k \in [p]$;
- (ii) $N_n \geq N(\epsilon, \mathcal{U}, d_{\mathcal{U}})$, here ϵ is such that with probability $1 - \Delta_n$:
 $\sup_{d_{\mathcal{U}}(u, u') \leq \epsilon} \max_{k \leq p} \frac{\|\mathbb{E}_n[S_u - S_{u'}]\|_\infty}{\mathbb{E}[|S_{uk}|^2]^{1/2}} \leq \delta_n n^{-\frac{1}{2}}$, and $\sup_{d_{\mathcal{U}}(u, u') \leq \epsilon} \max_{k \leq p} \frac{|\mathbb{E}[S_{uk}^2 - S_{u'k}^2]| + |(\mathbb{E}_n - \mathbb{E})[S_{uk}^2]|}{\mathbb{E}[|S_{uk}|^2]} \leq \delta_n$.

The following technical lemma justifies the choice of penalty level λ . It is based on self-normalized moderate deviation theory.

Lemma 23 (Choice of λ). Suppose Condition WL holds, let $c' > c > 1$ be constants, $\xi \in [1/n, 1/\log n]$, and $\lambda = c' \sqrt{n} \Phi^{-1}(1 - \xi / \{2pN_n\})$. Then for $n \geq n_0$ large enough depends only on Condition WL,

$$\mathbb{P} \left(\lambda/n \geq c \sup_{u \in \mathcal{U}} \|\hat{\Psi}_{u0}^{-1} \mathbb{E}_n[\partial_\theta M_u(Y_u, X_u, \theta_u, a_u)]\|_\infty \right) \geq 1 - \xi - o(\xi) - \Delta_n.$$

We note that Condition WL(ii) contains high level conditions. See [17] for examples that satisfy these conditions. The following corollary summarizes these results for many applications of interest in well behaved designs.

Corollary 5 (Rates under Simple Conditions). Suppose that with probability $1 - o(1)$ we have that $C_{un} \vee L_{un} \leq C \{n^{-1} s \log(pn)\}^{1/2}$, $(Lc+1)/(\ell c-1) \leq C$, $w_u = 1$, and Condition WL holds with $\log N_n \leq C \log(pn)$. Further suppose that with probability $1 - o(1)$ the sparse minimal and maximal eigenvalues are well behaved, $c \leq \phi_{\min}(s\ell_n, u) \leq \phi_{\max}(s\ell_n, u) \leq C$ for some $\ell_n \rightarrow \infty$ uniformly over $u \in \mathcal{U}$. Then with probability $1 - o(1)$ we have

$$\sup_{u \in \mathcal{U}} \|X'_u(\hat{\theta}_u - \theta_u)\|_{\mathbb{P}_{n,2}} \lesssim \sqrt{\frac{s \log(pn)}{n}}, \quad \sup_{u \in \mathcal{U}} \|\hat{\theta}_u - \theta_u\|_1 \lesssim \sqrt{\frac{s^2 \log(pn)}{n}}, \quad \text{and} \quad \sup_{u \in \mathcal{U}} \|\hat{\theta}_u\|_0 \lesssim s.$$

Moreover, if $\tilde{T}_u = \text{support}(\hat{\theta}_u)$, we have that

$$\sup_{u \in \mathcal{U}} \|X'_u(\tilde{\theta}_u - \theta_u)\|_{\mathbb{P}_{n,2}} \lesssim \sqrt{\frac{s \log(pn)}{n}}$$

APPENDIX J. BOUNDS ON COVERING ENTROPY

Let $(W_i)_{i=1}^n$ be a sequence of independent copies of a random element W taking values in a measurable space $(\mathcal{W}, \mathcal{A}_{\mathcal{W}})$ according to a probability law P . Let \mathcal{F} be a set of suitably measurable functions $f: \mathcal{W} \rightarrow \mathbb{R}$, equipped with a measurable envelope $F: \mathcal{W} \rightarrow \mathbb{R}$. The proofs for the following lemmas can be found in [17].

Lemma 24 (Algebra for Covering Entropies). *Work with the setup above.*

(1) *Let \mathcal{F} be a VC subgraph class with a finite VC index k or any other class whose entropy is bounded above by that of such a VC subgraph class, then the uniform entropy numbers of \mathcal{F} obey*

$$\sup_Q \log N(\epsilon \|F\|_{Q,2}, \mathcal{F}, \|\cdot\|_{Q,2}) \lesssim \{1 + k \log(1/\epsilon)\} \vee 0$$

(2) *For any measurable classes of functions \mathcal{F} and \mathcal{F}' mapping \mathcal{W} to \mathbb{R} ,*

$$\log N(\epsilon \|F + F'\|_{Q,2}, \mathcal{F} + \mathcal{F}', \|\cdot\|_{Q,2}) \leq \log N\left(\frac{\epsilon}{2} \|F\|_{Q,2}, \mathcal{F}, \|\cdot\|_{Q,2}\right) + \log N\left(\frac{\epsilon}{2} \|F'\|_{Q,2}, \mathcal{F}', \|\cdot\|_{Q,2}\right),$$

$$\log N(\epsilon \|F \cdot F'\|_{Q,2}, \mathcal{F} \cdot \mathcal{F}', \|\cdot\|_{Q,2}) \leq \log N\left(\frac{\epsilon}{2} \|F\|_{Q,2}, \mathcal{F}, \|\cdot\|_{Q,2}\right) + \log N\left(\frac{\epsilon}{2} \|F'\|_{Q,2}, \mathcal{F}', \|\cdot\|_{Q,2}\right),$$

$$N(\epsilon \|F \vee F'\|_{Q,2}, \mathcal{F} \cup \mathcal{F}', \|\cdot\|_{Q,2}) \leq N(\epsilon \|F\|_{Q,2}, \mathcal{F}, \|\cdot\|_{Q,2}) + N(\epsilon \|F'\|_{Q,2}, \mathcal{F}', \|\cdot\|_{Q,2}).$$

(3) *For any measurable class of functions \mathcal{F} and a fixed function f mapping \mathcal{W} to \mathbb{R} ,*

$$\log \sup_Q N(\epsilon \|f| \cdot F\|_{Q,2}, f \cdot \mathcal{F}, \|\cdot\|_{Q,2}) \leq \log \sup_Q N(\epsilon/2 \|F\|_{Q,2}, \mathcal{F}, \|\cdot\|_{Q,2})$$

(4) *Given measurable classes \mathcal{F}_j and envelopes F_j , $j = 1, \dots, k$, mapping \mathcal{W} to \mathbb{R} , a function $\phi: \mathbb{R}^k \rightarrow \mathbb{R}$ such that for $f_j, g_j \in \mathcal{F}_j$, $|\phi(f_1, \dots, f_k) - \phi(g_1, \dots, g_k)| \leq \sum_{j=1}^k L_j(x) |f_j(x) - g_j(x)|$, $L_j(x) \geq 0$, and fixed functions $\bar{f}_j \in \mathcal{F}_j$, the class of functions $\mathcal{L} = \{\phi(f_1, \dots, f_k) - \phi(\bar{f}_1, \dots, \bar{f}_k) : f_j \in \mathcal{F}_j, j = 1, \dots, k\}$ satisfies*

$$\log \sup_Q N\left(\epsilon \left\| \sum_{j=1}^k L_j F_j \right\|_{Q,2}, \mathcal{L}, \|\cdot\|_{Q,2}\right) \leq \sum_{j=1}^k \log \sup_Q N\left(\frac{\epsilon}{k} \|F_j\|_{Q,2}, \mathcal{F}_j, \|\cdot\|_{Q,2}\right).$$

Proof. See Lemma L.1 in [17]. ■

Lemma 25 (Covering Entropy for Classes obtained as Conditional Expectations). *Let \mathcal{F} denote a class of measurable functions $f: \mathcal{W} \times \mathcal{Y} \rightarrow \mathbb{R}$ with a measurable envelope F . For a given $f \in \mathcal{F}$, let $\bar{f}: \mathcal{W} \rightarrow \mathbb{R}$ be the function $\bar{f}(w) := \int f(w, y) d\mu_w(y)$ here μ_w is a regular conditional probability distribution over $y \in \mathcal{Y}$ conditional on $w \in \mathcal{W}$. Set $\bar{\mathcal{F}} = \{\bar{f} : f \in \mathcal{F}\}$ and let $\bar{F}(w) := \int F(w, y) d\mu_w(y)$ be an envelope for $\bar{\mathcal{F}}$. Then, for $r, s \geq 1$,*

$$\log \sup_Q N(\epsilon \|\bar{F}\|_{Q,r}, \bar{\mathcal{F}}, \|\cdot\|_{Q,r}) \leq \log \sup_{\tilde{Q}} N((\epsilon/4)^r \|F\|_{\tilde{Q},s}, \mathcal{F}, \|\cdot\|_{\tilde{Q},s}),$$

here Q belongs to the set of finitely-discrete probability measures over \mathcal{W} such that $0 < \|\bar{F}\|_{Q,r} < \infty$, and \tilde{Q} belongs to the set of finitely-discrete probability measures over $\mathcal{W} \times \mathcal{Y}$ such that $0 < \|F\|_{\tilde{Q},s} < \infty$. In particular, for every $\epsilon > 0$ and any $k \geq 1$,

$$\log \sup_Q N(\epsilon, \bar{\mathcal{F}}, \|\cdot\|_{Q,k}) \leq \log \sup_{\tilde{Q}} N(\epsilon/2, \mathcal{F}, \|\cdot\|_{\tilde{Q},k}).$$

Proof. See Lemma L.2 in [17]. ■