

Differentially Private Model Selection With Penalized and Constrained Likelihood

Jing Lei^{*1}, Anne-Sophie Charest², Aleksandra Slavkovic³, Adam Smith⁴, and
Stephen Fienberg¹

¹Department of Statistics, Carnegie Mellon University

²Département de mathématiques et statistique, Université Laval

³Department of Statistics, Pennsylvania State University

⁴Department of Computer Science and Engineering, Pennsylvania State University

March 5, 2024

Abstract

In statistical disclosure control, the goal of data analysis is twofold: The released information must provide accurate and useful statistics about the underlying population of interest, while minimizing the potential for an individual record to be identified. In recent years, the notion of *differential privacy* has received much attention in theoretical computer science, machine learning, and statistics. It provides a rigorous and strong notion of protection for individuals' sensitive information. A fundamental question is how to incorporate differential privacy into traditional statistical inference procedures. In this paper we study model selection in multivariate linear regression under the constraint of differential privacy. We show that model selection procedures based on penalized least squares or likelihood can be made differentially private by a combination of regularization and randomization, and propose two algorithms to do so. We show that our private procedures are consistent under essentially the same conditions as the corresponding non-private procedures. We also find that under differential privacy, the procedure becomes more sensitive to the tuning parameters. We illustrate and evaluate our method using simulation studies and two real data examples.

^{*}jinglei@andrew.cmu.edu

1 Introduction

In data privacy research, the goal of data analysis is to provide accurate and useful statistical inference while preventing individual records being identified. Such privacy protection is crucial in many statistical applications, namely for the analysis of census and survey data, medical and clinical studies, genetics data, and web user data collected on the internet. In statistics, the treatment of confidential data has a long history under the name of “Statistical Disclosure Control” or “Statistical Disclosure Limitation”; see for example [Dalenius \(1977\)](#), [Rubin \(1993\)](#), [Willenborg & De Waal \(1996\)](#), [Fienberg & Slavković \(2010\)](#) and [Hundepool et al. \(2012\)](#). A long lasting challenge is to rigorously quantify the disclosure protection offered by privacy-preserving data analysis procedures.

The notion of *differential privacy* has been introduced with the same objective in theoretical computer science by [Dwork \(2006\)](#) and [Dwork et al. \(2006\)](#). The general idea of differential privacy is to require for the outcome of a randomized data analysis procedure not to change much for small perturbations of the input data, so that one can not infer from the output the presence or absence of any individual in the input data set, or infer some of the person’s characteristics. This requirement can be rigorously quantified, and does not depend on assumptions regarding the resources of the intruder, including any access to auxiliary information. Thus, differential privacy compares very favorably to measures of disclosure risk commonly used in the statistical disclosure control literature as it is more encompassing and a *worst case definition*, but at the same time it has been criticized as too stringent from the perspective of achieving needed statistical data utility; e.g., see [Fienberg et al. \(2010\)](#); [Karwa & Slavkovic \(2012\)](#).

Over the last decade, there has been a rapid development of differentially private algorithms and procedures in both the computer science and statistics literature. For example, the main focus in the computer science literature was on designing differentially private mechanisms and efficient algorithms for private data release, e.g., the Laplace noise perturbation mechanism ([Dwork et al., 2006](#)), the exponential mechanism ([McSherry & Talwar, 2007](#)), releasing contingency tables ([Barak et al., 2007](#)), and boosting [Hardt et al. \(2010\)](#). On the statistics side, research efforts include designing consistent and efficient differentially private point estimators ([Dwork & Lei, 2009](#); [Smith, 2011](#); [Chaudhuri et al., 2011](#); [Lei, 2011](#); [Bassily et al., 2014](#); [Karwa & Slavković, 2016](#)), non-parametric density estimation ([Wasserman & Zhou, 2010](#)), hypothesis testing ([Fienberg et al., 2011](#); [Johnson & Shmatikov, 2013](#); [Uhler et al., 2013](#); [Yu et al., 2014](#); [Karwa et al., 2014](#); [Solea, 2014](#); [Dwork et al., 2015](#); [Sheffet, 2015](#); [Wang et al., 2015](#); [Gaboardi et al., 2016](#)), and statistical lower bounds ([Chaudhuri & Hsu, 2011](#); [Duchi et al., 2013](#)); there is also a large literature on private PAC learning, which echoes the concerns of statistical estimation in the context of classification ([Kasiviswanathan et al., 2011](#); [Beimel et al., 2010, 2014, 2015](#); [Bun et al., 2015](#); [Karwa et al., 2015](#)).

In this paper we consider statistical model selection under the constraint of differential privacy. Despite the fast development in combining statistical theory and methodology with differential privacy, model selection under privacy constraints has not been well understood. The problem of differentially private model selection is motivated by practical concerns: when private data analysis procedures are needed, it is rarely known which model is most appropriate for the data. When releasing the whole dataset is impractical due to privacy concerns and releasing point estimates for pre-specified models has limited utility, an appropriate compromise is to first identify the best model and then obtain and release consistent point estimates for this model. Both tasks need to be performed under privacy constraints, and thus we need methods for differentially private model selection.

In particular, we focus on the classical linear regression model selection. In particular, we aim to provide insights for the following two questions. (i) Is it theoretically possible to do model selection with differential privacy under the classical conditions? (ii) What new practical concerns arise in model selection when differential privacy is required?

We first show that the answer to the first question is positive by proposing differentially private model selection procedures based on penalized least squares and likelihood which exhibit asymptotic utility guarantee. Here utility means that the procedure selects the correct model with high probability under appropriate regularity conditions. In other words, differentially private model selection is theoretically possible in the classical setting. More specifically, we propose a two-step differentially private model selection procedure: we first obtain a least square or maximum likelihood estimate under an ℓ_1 constraint, then use noisy optimization with regularization to get the best model. Interestingly, the ℓ_1 constraint, usually employed in high dimensional problems, helps achieve privacy even in low dimensions.

Second, from simulations and real data examples we observe that the finite sample behavior of the proposed method depends crucially on the turning parameter required by the procedure. For example, our algorithm imposes an ℓ_1 norm constraint on the estimated regression coefficient. When the tuning parameter is conservatively chosen, i.e., the imposed upper bound of the ℓ_1 norm is large, the utility is rather limited for the sample or moderate sample sizes. When the sample size is in the thousands, the dependence of utility on tuning parameter is less significant, as predicted by the theory. More importantly, if auxiliary information is available, such as a good upper bound on the ℓ_1 norm of the true regression coefficient, then we may be able to choose the tuning parameter more adaptively, with improved utility. This reveals a distinct feature of differentially private data analysis: Some auxiliary information that does not affect classical inference may lead to significant performance improvement in differentially private analysis. Indeed, the ℓ_1 bound on the regression coefficient which improves the utility of our differentially-private procedure is useful in Lasso when the dimensionality is high, but not so much in the classical

regime.

The remainder of the paper is organized as follows: In section 2, we review the definition and interpretation of differential privacy, as well as two general methods to create differentially private algorithms. Section 3 details the two proposed differentially private model selection procedures, with proofs of their privacy guarantee, and notes on the choice of the tuning parameters. Statements and proofs of the utility guarantee of the two algorithms are given in section 4. Empirical results, including a simulation study and two real data examples are reported in section 5. Section 6 provides a brief discussion. All proofs and technical details are collected in section 7.

2 Differential privacy

Differential privacy requires that the output of a procedure is not drastically altered under small perturbations of the input data set, such that an attacker, regardless of his auxiliary information and computing power, can hardly recover the presence/absence of a particular individual in the data set. The notion of differential privacy is a property of that data analysis procedure, rather than of the output obtained.

2.1 Definition

To formalize, consider a data set $D = \{z_1, \dots, z_n\} \in \mathcal{Z}^n$ consisting of n data points in sample space \mathcal{Z} . A data analysis procedure \mathcal{T} , possibly randomized, maps the data set D , together with a random input ω , to $\mathcal{T}(D) \equiv \mathcal{T}(D, \omega) \in S$, an output space. Here we assume that (S, \mathcal{S}) is a measurable space and $\mathcal{T}(D, \cdot) : \Omega \mapsto S$ is a measurable function. Whenever it is not confusing, we will use $\mathcal{T}(D)$ to denote the random variable $\mathcal{T}(D, \omega)$.

For any two data sets D and D' of the same size, we use $\text{Ham}(D, D')$ to denote their *Hamming distance*, the number of entries at which they differ, regardless of the order. For example, if $\mathcal{Z} = \mathbb{R}$ and $D = \{1, 2, 3, 4\}$, $D' = \{2, 3, 4, 5\}$, then $\text{Ham}(D, D') = 1$. In the rest of this paper, we always use D and D' to denote a pair of adjacent data sets that differ at the last entry: $D = \{z_1, \dots, z_{n-1}, z_n\}$, $D' = \{z_1, \dots, z_{n-1}, z'_n\}$.

We can now state formally the property of differential privacy:

Definition 1 (ϵ -differential privacy, [Dwork et al. \(2006\)](#)). *Given a privacy parameter, $\epsilon > 0$, the procedure \mathcal{T} satisfies ϵ -differential privacy if*

$$\sup_{\text{Ham}(D, D')=1, A \in \mathcal{S}} \left| \log \frac{P_\omega(\mathcal{T}(D', \omega) \in A)}{P_\omega(\mathcal{T}(D, \omega) \in A)} \right| \leq \epsilon$$

where we define $\log \frac{0}{0} = 0$ for convenience.

In the above notation, P_ω denotes the probability with respect to ω , which is the source of randomness in the data analysis procedure. Thus, the definition does not impose any conditions on the distribution of D — the privacy is required to hold for *all pairs of adjacent data sets*. This stringent condition means that procedures which satisfy definition 1 provide very strong privacy guarantees, even against adversaries who have considerable partial information about the data set (Ganta et al., 2008). In order to satisfy this definition, any nonconstant procedure must be randomized.

The ϵ -differential privacy is a strong requirement, as it takes supremum over all possible neighboring data sets of size n . A mild relaxation is the (ϵ, δ) -differential privacy.

Definition 2 ((ϵ, δ) -differential privacy). *Given $\epsilon > 0$, $\delta \in (0, 1)$, a procedure satisfies (ϵ, δ) -differential privacy if, for all measurable $A \subseteq \mathcal{S}$ and all neighboring data sets D, D' ,*

$$P_\omega(\mathcal{T}(D) \in A) \leq e^\epsilon P_\omega(\mathcal{T}(D') \in A) + \delta.$$

Here the requirement of the original ϵ -differential privacy is relaxed so that the distribution of $\mathcal{T}(D)$ only needs to be dominated by that of $\mathcal{T}(D')$ outside of a set with probability no more than δ .

In our discussion of statistical applications we will generally focus on data sets consisting of a sequence of n independent random samples from an underlying distribution, and the corresponding probability will be denoted P_D . Note that differential privacy definition has no such assumption. We will use $P_{D,\omega}$ to denote the overall randomness due to both the data and random mechanism in the analysis.

2.2 Statistical interpretation of differential privacy

Privacy protection due to differential privacy can be interpreted from a Bayesian perspective, as in Abowd & Vilhuber (2008); Kasiviswanathan & Smith (2008). Suppose one has a prior distribution of the input data set D and then gets to observe the random output $\mathcal{T}(D, \omega)$. Denote by P_i and Q_i the marginal prior and posterior of X_i , for the i th entry in D . Then if the procedure \mathcal{T} is differentially-private and the prior P is a product measure on the entries of D , we have that $e^{-\epsilon} \leq dP_i/dQ_i \leq e^\epsilon$, thus limiting the information regarding X_i gained from the output $\mathcal{T}(D, \omega)$. A related hypothesis testing interpretation is given in Theorem 2.4 of Wasserman & Zhou (2010).

One may also intuitively interpret differential privacy as a specific notion of robustness. It requires that the distribution of $\mathcal{T}(D)$ is not changed too much if D is perturbed in only

one entry. However, the definition of differential privacy is also a *worst case definition*, where the probability ratio needs to be uniformly bounded over all pairs of adjacent input data sets. A key ingredient to designing differentially private statistical procedures is to bridge the gap between worst case privacy guarantee and the average case statistical utility. It turns out that robustness and regularization are the most relevant structures to explore, as in [Dwork & Lei \(2009\)](#), [Chaudhuri et al. \(2011\)](#) and [Smith & Thakurta \(2013\)](#).

2.3 Designing differentially private algorithms

For any statistical task which we want to carry, a specific randomized procedure \mathcal{T} must be designed to take as input a database $D \in \mathcal{Z}^n$ and return an element of the output space \mathcal{S} while satisfying differential privacy. There exists a few approaches which are generic enough to be adaptable to various tasks, and which are often used as building blocks for more complicated procedures. We present two of these methods, which we use later in the construction of our procedure. The first adds random noise to a non-private output and the second samples randomly from a set of candidate outputs.

Adding Laplace noise The additive noise approach is applicable when the output space is an Euclidean space. For presentation simplicity we consider here $S = \mathbb{R}$. Let $T(D)$ be a non-private mechanism. Define the *global sensitivity* G_T as

$$G_T = \sup_{\text{Ham}(D, D')=1} |T(D) - T(D')|. \quad (1)$$

If $G_T < \infty$, then it is easy to check ([Dwork et al., 2006](#)) that

$$\mathcal{T}(D) \equiv T(D) + \epsilon^{-1} G_T \zeta \quad (2)$$

satisfies ϵ -differential privacy, where ζ is a standard double exponential random variable with density function $0.5 \exp(-|\zeta|)$ (also known as the Laplace distribution).

The exponential mechanism and noisy optimization The *exponential mechanism* ([McSherry & Talwar, 2007](#)) is designed for discrete output spaces. Suppose $S = \{s_\alpha : \alpha \in \mathbb{N}\}$ and let $q : \mathcal{Z}^n \times S \mapsto \mathbb{R}$ be a score function that measures the quality of $s \in S$ in terms of its agreement with the input data set. Usually $q(D, s) = -|s - T(D)|$ for some deterministic procedure T . Denote $G_q = \sup_s G_{q(\cdot, s)}$, where $G_{q(\cdot, s)}$ is the global sensitivity of the mapping $q(\cdot, s)$. Let $\mathcal{T}(D)$ be the procedure that outputs a random sample from S with probability

$$P_\omega(\mathcal{T}(D) = s) \propto \exp\left(\frac{\epsilon q(D, s)}{2G_q}\right).$$

Then \mathcal{T} satisfies ϵ -differential privacy.

When S is finite, one may also use additive noise to approximately maximize $q(D, s)$ over s . Let

$$\tilde{q}(D, s) = q(D, s) + 2\epsilon^{-1}\zeta G_{q(\cdot, s)}$$

be the privatized score, where ζ is an independent draw from the standard double exponential distribution. Then

$$\mathcal{T}(D) = \arg \max_s \tilde{q}(D, s)$$

satisfies ϵ -differential privacy and usually offers similar performance as the exponential mechanism.

A generic scheme for (ϵ, δ) -differential privacy In many statistical problems the sample space is not compact and hence $G_T = \infty$ for many statistics T such as the sample mean. A general strategy is to show that one can add much less noise for most average case data sets with (ϵ, δ) -differential privacy (Dwork & Lei, 2009). These methods often involves the notion of *local sensitivity* of a deterministic procedure T :

$$G_T(D) = \sup_{D': \text{Ham}(D, D')=1} |T(D) - T(D')|. \quad (3)$$

If $G_T(D)$ is finite and public, then one can show that adding noise to $T(D)$ as in (2) with G_T replaced by $G_T(D)$ also gives ϵ -differential privacy. Unfortunately, $G_T(D)$ depends on the data set and may contain sensitive information. However, there is a generic scheme based on this idea with valid privacy guarantee under the following two general conditions on the deterministic procedure T .

1. For all $\epsilon > 0$ there exists a real-valued function $G^*(D)$ and a randomized procedure $\mathcal{T}_\epsilon(D, g)$, which is ϵ -differentially private if $g \geq G^*(D)$ and is assumed to be non-private.
2. Given $\epsilon > 0$ and $\delta \in (0, 1)$, there exists an ϵ -differentially private mapping $G_{\epsilon, \delta}(D)$ satisfying $P_\omega[G_{\epsilon, \delta}(D) \geq G^*(D)] \geq 1 - \delta$ for all D .

An example of $G^*(D)$ is the local sensitivity of some procedure $T(D)$, and \mathcal{T} is the noisy version as in (2) calibrated to a upper bound of the local sensitivity.

Proposition 2.1. *Under the above two assumptions, for any $\epsilon_1 + \epsilon_2 = \epsilon$, and $\delta \in (0, 1)$, $\mathcal{T}_{\epsilon_2}(D, G_{\epsilon_1, \delta}(D))$ satisfies (ϵ, δ) -differential privacy.*

3 Differentially Private Model Selection Procedures

Popular methods of model selection for linear regression include information criteria such as AIC (Akaike, 1974) and BIC (Schwarz et al., 1978), cross-validation (Picard & Cook, 1984), and the more recent penalized least squares (e.g., Tibshirani, 1996; Fan & Li, 2001). In this paper we focus on penalized least squares and penalized profile likelihood estimators, both being variants of the classical approach based on information criteria.

3.1 Background: Linear regression and model selection

In the linear regression model, the data points are independent, each consisting of a response $Y \in \mathbb{R}^1$, and a covariate $X \in \mathbb{R}^d$, which satisfies

$$Y = X^T \beta_0 + W \quad (4)$$

where $\beta_0 \in \mathbb{R}^d$ is the regression coefficient, and W is a Gaussian random variable, independent of X , with mean zero and variance σ^2 . The observed data set is $D = \{(X_i, Y_i) : 1 \leq i \leq n\}$, where $X_i = (X_{i1}, \dots, X_{id})^T$.

The model selection problem is to find the support of β_0 : $M_0 \equiv \{j : \beta_0(j) \neq 0\}$. To give a precise formulation, consider a class of candidate models $\mathcal{M} \subseteq \{0, 1\}^d$ that contains M_0 . For each $M \in \mathcal{M}$, the corresponding hypothesis is $\beta_0 \in \Theta_M \equiv \{\beta \in \mathbb{R}^d : \beta_0(j) = 0, \forall j \notin M\}$.

Given a parameter (β, σ^2) and an observed data set, the log likelihood is, ignoring constant terms,

$$\ell(\beta, \sigma^2; D) = \sum_{i=1}^n \left[-\frac{1}{2} \log \sigma^2 - \frac{1}{2\sigma^2} (Y_i - X_i^T \beta)^2 \right].$$

We consider two cases separately.

Known variance. When σ^2 is known, we aim to find the β that maximizes the log likelihood, which leads to model selection with the least squares. Without loss of generality, we assume $\sigma^2 = 1$. We then define

$$\ell(\beta; D) = -\frac{1}{2} \sum_{i=1}^n (Y_i - X_i^T \beta)^2, \quad (5)$$

$$\hat{\beta}_M = \arg \min_{\beta \in \Theta_M} -2\ell(\beta; D), \quad (6)$$

$$\ell(M; D) = \ell(\hat{\beta}_M; D). \quad (7)$$

Unknown variance. In the more realistic setting that σ^2 is unknown, we can maximize over the nuisance parameter σ^2 to perform model selection with the profile likelihood. Ignoring constant terms, we obtain the profile log-likelihood for β :

$$\ell^*(\beta; D) = -\frac{n}{2} \log \left[\frac{1}{n} \sum_{i=1}^n (Y_i - X_i^T \beta)^2 \right], \quad (8)$$

Maximizing this over all β in Θ_M , we get

$$\ell^*(M; D) \equiv \sup_{\beta \in \Theta_M} \ell^*(\beta; D). \quad (9)$$

Information criteria. One could simply maximize $\ell(M; D)$ or $\ell^*(M; D)$ as given above. However, more complex models tend to yield larger values of log likelihood. The approach of information criteria thus minimizes the sum of negative log likelihood and a measure of model complexity. For example, in the case of unknown σ^2 , one choose the model by

$$\widehat{M} = \arg \min_{M \in \mathcal{M}} -2\ell^*(M; D) + \phi_n |M|, \quad (10)$$

where ϕ_n is the amount of penalty on the model complexity. The best known and most widely used examples are the AIC ($\phi_n = 2$), and BIC ($\phi_n = \log n$).

In the case of known σ^2 , the penalized minimization becomes

$$\widehat{M} = \arg \min_{M \in \mathcal{M}} -2\ell(M; D) + \phi_n |M|. \quad (11)$$

It is worth noting that the penalized least square estimator, with appropriate choice of ϕ_n , can also be used in the case of unknown σ^2 .

3.2 Towards Differentially Private Model Selection

To perform model selection in a differentially-private manner, we propose to apply the exponential mechanism or noisy minimization to the minimization problems (10) and (11), with $q(D, M)$ (here s in the general definition is replaced by M) given by the penalized likelihood

$$L^*(M; D) \equiv -2\ell^*(M; D) + \phi_n |M|$$

in the case of penalized log profile likelihood (10), or

$$L(M; D) \equiv -2\ell(M; D) + \phi_n |M|$$

in the case of penalized least squares (11).

A key step in this approach is to evaluate and control the sensitivities of $L(M; D)$ and $L^*(M; D)$ as functions of D for any $M \in \mathcal{M}$.

To simplify the notation and concentrate on the main idea, we will assume in the sequel that the data entries are bounded or standardized:

$$\max_{1 \leq i \leq n} |Y_i| \leq r, \quad \max_{1 \leq i \leq n, 1 \leq j \leq d} |X_{ij}| \leq 1, \quad (12)$$

where r is a known number that can grow with n . Boundedness is typically required for differentially private data analysis. Standard methods for finding the range of the data set in a privacy-preserving manner include those given in [Dwork & Lei \(2009\)](#) and [Smith \(2011\)](#).

3.2.1 Sensitivity of least squares and profile log-likelihood

As indicated in Section 3.1, we consider procedures based on two score functions for a model: the sum of squared residuals $\ell(M; D)$, and the profile likelihood $\ell^*(M; D)$.

To bound the sensitivity of either of them, we must bound the possible parameter vectors we consider. We thus consider constrained versions of the two score functions. Given $R > 0$, we define

$$-2\ell_R(M; D) = \min_{\beta \in \Theta_M, \|\beta\|_1 \leq R} -2\ell(\beta, D). \quad (13)$$

and

$$-2\ell_R^*(M; D) = \min_{\beta \in \Theta_M, \|\beta\|_1 \leq R} -2\ell^*(\beta; D). \quad (14)$$

The sensitivity of the least squares loss is now easy to bound:

Lemma 3.1 (Sensitivity of constrained least-squares). *When β has ℓ_1 -norm at most R and the data (X_i, Y_i) are restricted to $[-1, 1]^d \times [-r, r]$, the global sensitivity of the squared error loss functions $-2\ell(\beta; \cdot)$ and $-2\ell_R(M; \cdot)$ is at most $(r + R)^2$.*

The proof is both short and elementary, and hence omitted.

In the case of the constrained profile likelihood, we can not bound the global sensitivity, but we can find a bound on the local sensitivity:

Lemma 3.2 (Local sensitivity of constrained the profile likelihood). *Under the same conditions as in Lemma 3.1, the local sensitivity of $-2\ell_R^*(M; \cdot)$ is no larger than*

$$\frac{n(r + R)^2}{-2\ell_R(M; D) - (r + R)^2}.$$

Algorithm 1 Model Selection via Penalized Constrained Least Squares (PCLS)

Input: Data set $D = \{(X_i, Y_i) : i \in \{1, \dots, n\}\}$, collection of models \mathcal{M} , parameters r, R, ϕ_n, ϵ .

Output: Estimated model $\widehat{M} \in \mathcal{M}$

for each model M in \mathcal{M} **do**

$$\ell_R(M; D) \leftarrow \max_{\beta \in \Theta_M, \|\beta\|_1 \leq R} -\frac{1}{2} \sum_{i=1}^n (Y_i - X_i^T \beta)^2$$

$$L_R(M; D) \leftarrow -2\ell_R(M; D) + \phi_n |M|$$

$$\tilde{L}_R(M; D) \leftarrow L_R(M; D) + \frac{2(r+R)^2}{\epsilon} Z_M, \text{ where } Z_M \text{ is a Laplace random variable}$$

end for

Return $\arg \min_{M \in \mathcal{M}} \tilde{L}_R(M; D)$

3.2.2 Algorithms

Lemma 3.1 implies that the penalized constrained least squares

$$L_R(M; D) = -2\ell_R(M; D) + |M|\phi_n,$$

can easily be minimized in an ϵ -differentially private manner. The complete algorithm is given below.

Next, Lemma 3.2 gives an upper bound of the local sensitivity of $L_R^*(M; D)$. Now we apply the generic scheme of designing (ϵ, δ) -differentially private algorithms described in section 2.3.

First let

$$G^*(D) \equiv \frac{n(r+R)^2}{\min_M -2\ell_R(M; D) - (r+R)^2}.$$

Then $G^*(D)$ is an uniform upper bound of the local sensitivity of $L_R(M; D)$ for all M . The only private part in $G^*(D)$ above is $\min_{M \in \mathcal{M}} -2\ell_R(M; D)$, which has global sensitivity $(r+R)^2$ according to Lemma 3.1. Thus following the general procedure in Section 2.3, a valid choice of $G(D)$ for $-2\ell_R^*(M; D)$ is

$$G(D) = \frac{n(r+R)^2}{\min_M -2\ell_R(M; D) - (r+R)^2 + \epsilon^{-1}(r+R)^2 (Z_G - \log \frac{1}{2\delta})}$$

where Z_G is a standard Laplace random variable.

Then we can construct the final estimators using either exponential mechanism or the noisy minimization, both satisfying $(2\epsilon, \delta)$ -differential privacy. The complete algorithm is given in Algorithm 2.

Algorithm 2 Model Selection via Penalized Constrained Profile Likelihood (PCPL)

Input: Data set $D = \{(X_i, Y_i) : i \in \{1, \dots, n\}\}$, collection of models \mathcal{M} , parameters $r, R, \phi_n, \delta, \epsilon$.

Output: Estimated model $\widehat{M} \in \mathcal{M}$

for each model M in \mathcal{M} **do**

$$\ell_R^*(M; D) \leftarrow \max_{\beta \in \Theta_M, \|\beta\|_1 \leq R} -\frac{n}{2} \log \left[\frac{1}{n} \sum_{i=1}^n (Y_i - X_i^T \beta)^2 \right]$$

$$L_R^*(M; D) \leftarrow \ell_R^*(M; D) + \phi_n |M|$$

$$\widetilde{L}_R^*(M; D) \leftarrow L_R^*(M; D) + \frac{2G(D)}{\epsilon} Z_M \text{ where}$$

$$G(D) = \frac{n(r + R)^2}{\min_M -2\ell_R(M; D) - (r + R)^2 + \epsilon^{-1}(r + R)^2 (Z_G - \log \frac{1}{2\delta})};$$

$\ell_R(M; D)$ is as in Algorithm 1, and Z_G, Z_M are Laplace random variables;

end for

Return $\arg \min_{M \in \mathcal{M}} \widetilde{L}_R^*(M; D)$

3.3 Choosing the tuning parameters

Both algorithms introduced above require two tuning parameters: an upper bound R of the ℓ_1 norm of the regression coefficient, and the penalty parameter ϕ_n . In the context of privacy-preserving data analysis, there are two requirements on the quality of any inference procedure: privacy and utility. Both proposed methods satisfy their corresponding ϵ -differential privacy (Algorithm 1) and $(2\epsilon, \delta)$ -differential privacy (Algorithm 2) for any choices of R and ϕ_n . Regarding utility, our theoretical analysis shows that both algorithms achieve consistent model selection for a wide range of R and ϕ_n .

However, our numerical experiments in section 5 show that the performance of our proposed algorithms is sensitive to the choice of these tuning parameters. We acknowledge that fully data-driven and privacy-preserving methods for choosing these tuning parameters remain a challenging and important open problem, and is beyond the scope of this paper. Data-driven choice of penalty parameter is a hard problem even without privacy constraint. Here we provide some heuristics on potential solutions.

Regarding choosing R , the ideal choice would be the ℓ_1 norm of the true regression coefficient. In practice, a good choice of R should be close to $\|\beta_0\|_1$. This can be achieved by finding a differentially private estimate of the maximum ℓ_1 norm of $\widehat{\beta}_M$ over all $M \in \mathcal{M}$, which is feasible if only a single number is released.

Regarding choosing ϕ_n , a good choice of ϕ_n needs to be large enough so that it dominates

any statistical sampling noise in the data and the additive noise due to privacy constraints. On the other hand, ϕ_n cannot be too large, because otherwise it will introduce substantial bias in the selected model. Taking the penalized least squares estimator for example, a good ϕ_n shall roughly be the difference of the largest $\ell(M; D)$ and the second largest $\ell(M; D)$, which can possibly be estimated with differential privacy as a single number.

4 Utility Analysis

In privacy-preserving data analysis, the privacy shall be protected for *any* possible input data set. In other words, the privacy guarantee needs to cover the worst case and must be established with no distributional conditions on the data set. In the previous sections, our differentially private procedure only requires the data to be bounded, which can be verified or enforced easily in practice. On the other hand, the statistical utility (for example, consistency, rate of convergence) is usually based on common statistical assumptions on the data. To facilitate discussion, we first introduce some notation and assumptions.

Notation. Let \mathbf{X} be the $n \times d$ design matrix, and \mathbf{Y} the $n \times 1$ vector of Y 's. For any $M \in \mathcal{M}$, let \mathbf{X}_M be the $n \times |M|$ design matrix consisting of the columns in M . Then $\hat{\beta}_M$ is a $d \times 1$ vector that is $(\mathbf{X}_M^T \mathbf{X}_M)^{-1} \mathbf{X}_M^T \mathbf{Y}$ on the entries in M and zero elsewhere. It is the ordinary least square estimate under model M . The sample covariance is $\hat{\Sigma} = n^{-1} \mathbf{X}^T \mathbf{X}$.

Assumptions. We state below five assumptions required for our utility results, and discuss their practical significance.

Our first assumption is a sparse linear model with Gaussian noise.

A0. (Linear model with Gaussian noise) The data entries $(X_i, Y_i)_{i=1}^n$ are generated from the linear regression model (4) with a regression coefficient vector β_0 with $d_0 = \|\beta_0\|_0$ and $b_0 \equiv \min_{j: \beta_0(j) \neq 0} |\beta_0(j)|$. The noise W_i are iid Gaussian with mean zero and variance σ^2 .

Next we assume that the number of candidate models grows polynomially with sample size n . This is usually the case when we only search over sparse models and the total number of variables grows polynomially in n .

A1. (Candidate models) The set of candidate models \mathcal{M} contains the true model $M_0 =$

$\{j : \beta_0(j) \neq 0\}$, and has cardinality no more than n^α for some positive number α which is allowed to grow with n . The largest candidate model has \bar{d} variables.

The worst-case sensitivity of the log likelihood is hard to control because the design matrix \mathbf{X} may be poorly conditioned. Thus we need to add some singular value condition on the design matrix.

A2. (Design matrix) The design matrix \mathbf{X} is fixed, and the sample covariance satisfies the sparse eigenvalue condition:

$$\kappa_0 \equiv \inf_{1 \leq \|\beta\|_0 \leq \bar{d} + d_0} \frac{\beta^T \hat{\Sigma} \beta}{\|\beta\|_2^2} > 0.$$

Assumption A2 excludes the situation of linear dependence between columns of \mathbf{X}_{M_0} and $\mathbf{X}_{M_0^c}$. A similar condition has been considered in the literature on model selection consistency using information criteria (Nishii, 1984). The condition stated here is for fixed design matrix \mathbf{X} , but it holds with high probability for many random designs. Assumption A2 also implies an upper bound of the range of $\hat{\beta}_M$. Such a boundedness property will help control the sensitivity of the log likelihood.

Moreover, as we did in section 3, we assume boundedness of the data entries, which are typically needed for developing differentially private procedures.

A3. (Boundedness) Each entry of \mathbf{Y} is bounded by r , and each entry of \mathbf{X} is bounded by 1.

Finally, we assume that the sample size is large enough, when compared to other quantities in the analysis such as the noise variance, and the inverse of privacy parameter ϵ .

A4. (Sample size) The sample size n is large enough so that equations (18), (19), and (20) hold.

4.1 Utility of Penalized Constrained Least Squares

We first give a utility result for the noisy penalized constrained least squares estimator (Algorithm 1).

Theorem 4.1. *Assume A0-A4 hold. If ϕ_n satisfies*

$$2A(1 \vee \sigma^2) \log n \leq \phi_n \leq \frac{1}{4 \vee (1 + 2d_0)} \kappa_0 b_0^2 \sigma^2 n,$$

where $A = 2(\alpha + c)$ for some $c > 0$, and

$$R \geq r \sqrt{\frac{\bar{d}}{\kappa_0}},$$

then the PCLS estimator \widehat{M} given by Algorithm 1 with privacy parameter ϵ and penalty parameter ϕ_n satisfies

$$P_{D,\omega}(\widehat{M} \neq M) \leq n^\alpha \exp\left(-\frac{\phi_n \epsilon}{4(R+r)^2}\right) + \frac{1+2\bar{d}}{\sqrt{2\pi A \log n}} n^{-c}.$$

Recall that $P_{D,\omega}$ stands for taking probability over both the randomness in D and in the generation of Laplace random variables (denoted by ω) in the algorithm. Here we are assuming a fixed design matrix, so the randomness in D is equivalent to the randomness of the additive noise W .

4.2 Utility of Penalized Constrained Profile Likelihood

Now we provide utility result for the noisy penalized profile likelihood estimator.

Theorem 4.2 (Utility of penalized profile likelihood). *Assume A0-A4 hold. If ϕ_n satisfies*

$$4A \log n \leq \phi_n \leq \frac{2}{3|M_0|} n \log\left(1 + \frac{\kappa_0 b_0^2}{4\sigma^2}\right),$$

where $A = 2(\alpha + c)$ for some $c > 0$, and

$$R \geq r \sqrt{\frac{\bar{d}}{\kappa_0}}$$

then for any constant $c > 0$ and n large enough as quantified in (18) to (20), the selected model \widehat{M} given by the noisy penalized constrained profile likelihood (Algorithm 2) satisfies

$$P_{D,\omega}(\widehat{M} \neq M_0) \leq n^\alpha \exp\left(-\frac{\epsilon \sigma^2 \phi_n}{64(R+r)^2}\right) + 3n^{-A/2} + 2\bar{d}n^{-c}.$$

5 Empirical Results

We have shown in the last section that the two proposed differentially-private model selection procedures are consistent under similar conditions as the corresponding non-private algorithms. We now provide some empirical results for model selection via penalized

constrained least squares, to illustrate both the utility of the algorithms and impact of tuning parameter selection as discussed in section 3.3.

5.1 Simulation study

We consider two different regression models, both of the form $Y = X^T \beta_0 + W$, where X is an n by 6 matrix with columns sampled independently from the uniform distribution on $[-1, 1]$ and W is standard normal, but the regression coefficients are different. The first model uses $\beta_0 = (1, 1, 1, 0, 0, 0)$ whereas the second model uses $\beta_0 = c(1.5, 1, 0.5, 0, 0, 0)$. Note that the ℓ_1 norm of β_0 is 3 in both cases, and thus the oracle value for R . However, the first model should be easier to recover from data.

In the simulations, we consider a small sample size ($n = 100$) and a moderately large sample size ($n = 1000$), as well as $R \in \{1, 2.5, 3.5, 10\}$, $\epsilon \in \{0.1, 1, 5, 10\}$ and $\phi \in [0, n/2]$. For each set of parameters, we sample 500 data sets from the true model, apply Algorithm 1 to each of them, and note the proportion of times we correctly identify the correct model. We set r to be the maximum observed value of Y in the data set.

Figure 1 shows the results for model 1 where $\beta_0 = (1, 1, 1, 0, 0, 0)$. Except for with the smallest value of R the procedure is very successful at choosing the correct model over a wide range of ϕ values. The poor results for $R = 1$ are explained by the fact that since R is much smaller than the true ℓ_1 norm of β_0 , the parameter can not be estimated properly. When R is too large, the estimation of β_0 is unchanged, but a larger amount of noise is needed to satisfy differential privacy, which explains the small drop in utility. As expected, decreasing ϵ needed to provide stricter privacy also leads to decreased utility. Note however that for $n = 1000$ the method is very accurate for a wide range of ϕ even for ϵ as small as 1. The results also confirm our claim that the choice of ϕ is less crucial for larger sample size, as the procedure works well on an entire interval. Note also that the actual value of ϕ depends on n , with the optimal ϕ increasing roughly linearly with n .

Figure 2 shows the results for case where $\beta_0 = c(1.5, 1, 0.5, 0, 0, 0)$. The effect of n , R , ϕ and ϵ are very similar in this case, but the choice of R and ϕ are more important to achieve good utility. The sensitivity of the procedure to choices of R and ϕ thus depends on the structure of the true parameter β_0 . Note however that for $n = 1000$ a proper choice for the parameters leads to completely accurate model selection with $\epsilon = 5$, and even very accurate for $\epsilon = 1$ with $R = 2.5$.




Figure 1: Proportion of correct model selection from 500 independent replications of Algorithm 1 selecting among the 63 possible models when the true model has $\beta = (1, 1, 1, 0, 0, 0)$. Various values of n , R , ϕ and ϵ are illustrated, and r is set to the maximum value of Y . There is a large range of ϕ for which the private procedure does as well as without privacy, and in fact picks the correct model. As n increases, the task becomes easier. Note that the proper value of ϕ increases with n as well.

5.2 Application to Real Data Sets

We first illustrate the results of model selection via penalized constrained least squares on a small data set of 97 observations, then on a much larger one with hundreds of thousands observations.

Prostate data set The prostate data set contains several clinical measures for 97 men with prostate cancer. In this paper, our goal is to predict the level of a prostate specific antigen using five continuous variables: the volume of the cancer, the weight of the prostate, the age of the patient, the capsular penetration and the benign prostatic hyperplasia amount. Except for age, all variables are taken on the *log* scale. We also rescale all of the variables to take values between -1 and 1 , which could be done in a differentially private way.

We consider all possible main effects models for the model selection procedure, for a total of 63 models to choose from. Model selection based on the penalized maximum likelihood,




Figure 2: Proportion of correct model selection from 500 independent replications of Algorithm 1 selecting among the 63 possible models when the true model has $\beta = (1.5, 1, 0.5, 0, 0, 0)$. Various values of n , R , ϕ and ϵ are illustrated, and r is set to the maximum value of Y . The utility of the procedure depends crucially on the choices for R and ϕ . As n increases, the task becomes easier. Note that the proper value of ϕ increases with n as well.

without the constraint of differential privacy, selects the model with only two variables in addition to the intercept: the volume of the cancer, and the weight of the prostate. Model selection is then performed using Algorithm 1 for various choices of R , ϕ and ϵ . We set $R = 5.68$, obtained with the non-private algorithm, and we set r to the maximum value of Y .

Figure 3 shows the proportion of times that the private and non-private procedures identify the correct model, for 5000 independent replications. With such a small data set, the choice of R and ϕ are crucial: even without differential privacy, using too large a value of ϕ does not identify the correct model. Due to the small sample size, the utility of the private procedure also decreases quite rapidly with decreasing ϵ which should offer more privacy.

Housing dataset Since the constrained optimization can be implemented based only on sufficient statistics, the private procedure scales very well for much larger data sets. The housing data set contains several variables measured on 348,189 houses sold in San




Figure 3: Model selection on the prostate data set for the private and non-private procedures. Results show the proportion of times that the correct model is selected for 5000 independent replications. Various values of R , ϕ and ϵ are illustrated. Because of the small sample size, the choice of R and ϕ as more impact, and the utility is quite small for usual choices for ϵ .

Francisco Bay Area between 2003 and 2006. In addition to the price of the sale, the data include the year of the transaction (an ordinal variable with 4 levels), the latitude and longitude of the house, the county in which it is located (a categorical variable with 9 levels), and a continuous measure of its size. We preprocess the data to remove houses with price outside the range 105 to 905 thousand dollars, and size larger than 3000 sqft. We also combine some of the small counties into a new indicator variable. All predictors are also scaled so that they take values in $[-1, 1]$. The resulting data set contains 235,760 with 13 variables.

We consider again models with main effects only, for a total 8191 models to choose the best model from. Since the optimal model is not as clear as with the previous example, we do not directly compare the private procedure with a gold standard, but rather with its non-private counterpart. An algorithm which offers differential privacy but recovers the results of a non-private algorithm is successful. As above, we apply the model selection procedures with various values of R , ϕ and ϵ , and use the maximum value of Y as r .

Figure 4 shows the agreement between the private and non-private procedure for various tuning parameters. With this large data set, the results are very positive: even for $\epsilon = 1$ we recover the same model with the differentially-private procedure as without the privacy requirement for ϕ chosen large enough. Note also that, as expected, the scale of ϕ is much larger than for the smaller *Housing* example.

6 Discussion

With modern data acquisition and storage techniques allowing the collection and analysis of huge amounts of personal information in a multitude of formats, protecting individual privacy inevitably becomes of crucial concern for modern data analysis. Although differential privacy offers mathematically strong and elegant privacy guarantee, the nature of such a conservative constraint in the context of everyday statistical analysis remains unclear. Previous works on statistical analysis with differential privacy mainly focused on simple statistical queries, such as location and scale statistics, regression coefficients, and simple hypothesis testing, and network analysis. In this paper we considered the more challenging problem of model selection in the classical setting. We showed that standard techniques for differentially private data analysis can be combined with known statistical tools such as penalized least square or information criteria to construct privacy-preserving model selection procedures with strong utility. We proposed two algorithms for this task, and proved privacy and utility results for each of them.

Our procedures feature a double-regularization, as they include both constrained estimation in the fitting step, and penalization in the model comparison step. Thus the method involves




Figure 4: Model selection on the housing data set. The y axis shows the proportion of time that the DP procedure selects the same model as the non-private version. Results represent 1000 independent replications of Algorithm 1. Various values of R , ϕ and ϵ are illustrated. Even with small values of ϵ , the private procedure returns the same model as the non-private one, for a large range of ϕ .

two tuning parameters. A key observation, illustrated in section 5, is that although we have proven good large-sample properties for a wide range of penalty parameter ϕ_n and R , the practical performance is sensitive to the particular choices for these parameters. In other words, these tuning parameters play a very important and unique role in designing differentially private procedure with good practical performance. While in low-dimensional settings, the regularization parameter is not needed for statistical inference without privacy constraints, when privacy is a concern, a carefully chosen amount of regularization can lead to stable, low-sensitivity estimators even in the worst case. Appropriate choice of both tuning parameters thus reflects the need for some additional information in the data that will be useful for differentially private procedures, but not necessary in traditional inference methods. It will be an interesting future topic to give a general characterization of such privacy related quantities, and to develop differentially private methods to estimate these quantities.

7 Appendix: Proof details

7.1 Proofs

We first provide proofs of the theorems.

Proof of Proposition 2.1

Proof.

$$\begin{aligned}
& P_\omega[\mathcal{T}(D, G(D)) \in A] \\
& \leq P_\omega[\mathcal{T}(D, G(D)) \in A, G(D) \geq G^*(D)] + \mathbb{P}[G(D) < G^*(D)] \\
& \leq \int_{u \geq G^*(D)} P_\omega[\mathcal{T}(D, u) \in A] dP_{G(D)}(u) + \delta \\
& \leq \int_{u \geq G^*(D)} e^{\epsilon_2} P_\omega[\mathcal{T}(D', u) \in A] dP_{G(D)}(u) + \delta \\
& \leq \int_{u \geq G^*(D)} e^{\epsilon_2} P_\omega[\mathcal{T}(D', u) \in A] e^{\epsilon_1} dP_{G(D')}(u) + \delta \\
& \leq e^\epsilon P_\omega[\mathcal{T}(D', G(D')) \in A] + \delta.
\end{aligned}$$

■

Proof of Lemma 3.2

Proof. Given a candidate model M , the log likelihood is, ignoring constant terms,

$$-2\ell^*(M; D) = n \log \left(n^{-1} \sum_{i=1}^n (Y_i - X_i^T \hat{\beta}_M)^2 \right).$$

Let $\hat{\beta}'_{R,M}$ be the constrained least square estimator with input data set D' . By boundedness of Y_i , X_i and $\hat{\beta}_{R,M}$ we have $(Y_i - X_i^T \beta)^2 \leq (r + R)^2$ for all β such that $\|\beta\|_1 \leq R$. Then using the fact that $\log(x/y) \leq (x - y)/y$ for $x \geq y$, we have

$$\begin{aligned} |2\ell_R^*(M; D') - 2\ell_R^*(M; D)| &= n \left| \log \left(\frac{\ell_R(M; D)}{\ell_R(M; D')} \right) \right| \\ &\leq \max \left\{ \frac{(Y'_n - X_n'^T \hat{\beta}_{R,M})^2}{n^{-1} \sum_{i=1}^n (Y_i - X_i^T \hat{\beta}_{R,M})^2}, \frac{(Y_n - X_n^T \hat{\beta}'_{R,M})^2}{n^{-1} \sum_{i=1}^n (Y'_i - X_i'^T \hat{\beta}'_{R,M})^2} \right\} \\ &\leq \frac{n(r + R)^2}{-2\ell_R(M; D) - (r + R)^2}. \end{aligned}$$

■

Next we prove our main utility theorems. The proofs of Theorem 4.1 and Theorem 4.2 rely on the following result, which is a consequence of simple linear algebra.

Proposition 7.1. *We have $\|\hat{\beta}_M\|_1 \leq \sqrt{d/\kappa_0} r$ for all $M \in \mathcal{M}$. As a result, if $R \geq \sqrt{d/\kappa_0} r$ then $\hat{\beta}_{R,M} = \hat{\beta}_M$, $\ell(M; D) = \ell_R(M; D)$, and $\ell_R^*(M; D) = \ell^*(M; D)$ for all $M \in \mathcal{M}$.*

Proposition 7.1 allows us to remove the ℓ_1 constraint in our analysis. Although Proposition 7.1 implies that the ℓ_1 constraint is inactive, such a constraint cannot be removed from the algorithms because it is used to bound the worst-case sensitivity of the estimators.

Proof of Theorem 4.1

Proof. We prove for noisy minimization. The argument can be adapted simply to cover the exponential mechanism.

Define

$$\Delta(M) = [-2\ell(M; D) + 2\ell(M_0; D)] .$$

Then according to Proposition 7.1, we can work directly with the unconstrained version.

$$\begin{aligned}
P_{D,\omega}(\widehat{M} \neq M) &\leq \sum_{M' \in \mathcal{M}} P_{D,\omega} \left(-2\ell(M'; D) + \phi_n |M'| + \frac{2(R+r)^2}{\epsilon} Z_{M'} \leq \right. \\
&\quad \left. -2\ell(M_0; D) + \phi_n |M_0| + \frac{2(R+r)^2}{\epsilon} Z_{M_0} \right) \\
&\leq n^\alpha P_{D,\omega} \left\{ \frac{2(R+r)^2}{\epsilon} (Z_1 - Z_2) \geq \sup_{M' \neq M_0} [\Delta_{M'} + (|M'| - |M_0|)\phi_n] \right\}.
\end{aligned}$$

For $M' \supset M_0$, using Claim 2 in Section 7 we have $\sup_{M': M' \supset M_0} \Delta_{M'} \geq -A\sigma^2(|M'| - |M_0|) \log n$ with probability at least $1 - \frac{\bar{d}-d_0}{\sqrt{2\pi A \log n}} n^{-c}$. Thus with same probability we have

$$\sup_{M' \supset M_0} \Delta_{M'} + (|M'| - |M_0|)\phi_n \geq (|M'| - |M_0|)(\phi_n - A\sigma^2 \log n) \geq \phi_n/2. \quad (15)$$

For $M' \not\supset M_0$, we have by Claim 3 in Section 7, with probability at least $1 - \frac{1+\bar{d}}{\sqrt{2\pi A \log n}} n^{-c}$,

$$\sup_{M' \not\supset M} \Delta_{M'} + (|M'| - |M_0|)\phi_n \geq \frac{1}{2} \kappa_0 b_0^2 \sigma^2 n - |M_0| \phi_n \geq \phi_n/2. \quad (16)$$

Thus conditioning on $E_2^c \cap E_3^c$, which has probability at least $1 - \frac{1+2\bar{d}}{\sqrt{2\pi A \log n}} n^{-c}$, we have

$$P_\omega(\widehat{M} \neq M_0) \leq n^\alpha \exp \left(-\frac{\phi_n}{2} \frac{\epsilon}{2(R+r)^2} \right).$$

■

Proof of Theorem 4.2

Proof. Consider events E_1 – E_5 as defined in Section 7. We focus on the event $\left(\bigcup_{k=0}^5 E_k \right)^c$, which has probability at least $1 - 3n^{-A/2} - 2\bar{d}n^{-c}$.

Because $R \geq r\sqrt{\frac{\bar{d}}{\kappa_0}}$ by Proposition 7.1 we have $-2\ell^*(M; D) = -2\ell_R^*(M; D)$ for all M under consideration.

Next we bound the difference between $-2\ell^*(M; D)$ and $-2\ell^*(M_0; D)$. Denote

$$\Delta^*(M) = -2\ell^*(M; D) + 2\ell^*(M_0; D).$$

In the case of $M \supset M_0$, applying the fact that $-\log(1-x) \leq x/(1-x)$ for $x \in (0,1)$ to

$$x = 1 - \frac{\sum_{i=1}^n (Y_i - X_i^T \hat{\beta}_M)^2}{\sum_{i=1}^n (Y_i - X_i^T \hat{\beta}_0)^2}$$

we have

$$\begin{aligned} 0 \leq -\Delta^*(M) &= n \log \left(\frac{\sum_{i=1}^n (Y_i - X_i^T \hat{\beta}_0)^2}{\sum_{i=1}^n (Y_i - X_i^T \hat{\beta}_M)^2} \right) \leq n \frac{-\Delta(M)}{-2\ell(M; D)} \\ &\leq 2A(|M| - |M_0|) \log n, \end{aligned} \quad (17)$$

where the last step follows from the fact that we are not in the event $E_0 \cup E_1 \cup E_2$ defined in Section 7.

In the case of $M \not\supset M_0$,

$$\Delta^*(M) = n \log \left(1 + \frac{\Delta(M)}{-2\ell(M_0; D)} \right) \geq n \log \left(1 + \frac{\kappa_0 b_0^2}{4\sigma^2} \right)$$

because we are in E_0^c (which implies $-2\ell(M_0; D) \leq 2n\sigma^2$ by Claim 1) and E_3^c (which implies $\Delta(M) \geq n\kappa_0 b_0^2 \sigma^2 / 2$ by Claim 3).

Therefore, when

$$4A \log n \leq \phi_n \leq \frac{2}{3|M_0|} n \log \left(1 + \frac{\kappa_0 b_0^2}{4\sigma^2} \right),$$

we have

$$L_R^*(M; D) - L_R^*(M_0; D) \geq \phi_n / 2, \quad \forall M \neq M_0.$$

On the event considered, we also have $G(D) \leq 4(R+r)^2 \sigma^{-2}$ by Claim 5. Then we can bound the error probability by

$$\begin{aligned} P_\omega[\widehat{M} \neq M_0 | D, G(D)] &= \sum_{M \in \mathcal{M}, M \neq M_0} P_\omega[\widehat{M} = M] \\ &\leq \sum_{M \in \mathcal{M}, M \neq M_0} \exp \left(-\frac{\epsilon}{4G(D)} (L_R^*(M; D) - L_R^*(M_0; D)) \right) \\ &\leq n^\alpha \exp \left(-\frac{\epsilon \phi_n \sigma^2}{64(R+r)^2} \right). \end{aligned}$$

■

7.2 Further proof details

Here we give details and summarize the multiple “with high probability” statements in the utility analysis. Recall that

$$\Delta(M) = [-2\ell(M; D) + 2\ell(M_0; D)] .$$

Let $A = 2(\alpha + c)$, assume

$$\frac{n}{\log n} \geq \max \left\{ \frac{64A\sigma^2}{\kappa_0 b_0^2}, \frac{4\bar{d}\sigma^2}{\kappa_0 b_0^2}, 8A\bar{d} \right\} \quad (18)$$

$$d_0 \leq n/8 \quad (19)$$

$$\epsilon \geq \frac{(R+r)^2 \left(\frac{A}{2} \log n + \log \frac{1}{2\delta} \right)}{\frac{n\sigma^2}{4} - (R+r)^2} \quad (20)$$

We define the following events and give the corresponding probability bounds.

1. Define

$$E_0 := \left\{ D : -2\ell(M_0; D)\sigma^{-2} \geq (n - d_0) + \sqrt{2(n - d_0)A \log n} + A \log n \right\} \quad (21)$$

$$E_1 := \left\{ D : -2\ell(M_0; D)\sigma^{-2} \leq (n - d_0) - \sqrt{2(n - d_0)A \log n} \right\} . \quad (22)$$

Claim 1. $P_D(E_0) \leq n^{-A/2}$, $P_D(E_1) \leq n^{-A/2}$. Also using Eq. (18), we have $-2\ell(M; D) \leq 2n\sigma^2$ on E_0^c .

Proof. Without loss of generality, assume that $\sigma^2 = 1$. Then $-2\ell(M_0; D) = \|\mathbf{Y} - P_{\Pi_0} \mathbf{Y}\|_2^2 = \|P_{\Pi_0^\perp} \mathbf{W}\|_2^2$ is a χ^2 random variable with $n - d_0$ degrees of freedom. The first claim follows from Lemma 1 of [Laurent & Massart \(2000\)](#). The second claim can be verified directly. \blacksquare

2. Define

$$E_2 := \left\{ \inf_{M \in \mathcal{M}, M \supset M_0} \frac{\Delta(M)}{\sigma^2(|M| - |M_0|)} < -A \log n \right\} \quad (23)$$

Claim 2. $P_D(E_2) \leq \frac{\bar{d} - d_0}{\sqrt{2\pi A \log n}} n^{-c}$.

Proof. Because $M \supset M_0$,

$$\begin{aligned} 0 \leq -\Delta(M) &= \|\mathbf{Y} - \mathbf{X}\hat{\beta}_{M_0}\|_2^2 - \|\mathbf{Y} - \mathbf{X}\hat{\beta}_M\|_2^2 \\ &= \|\mathbf{X}\beta_0 + \mathbf{W} - P_{\Pi_0}(\mathbf{X}\beta_0 + \mathbf{W})\|_2^2 - \|\mathbf{X}\beta_0 + \mathbf{W} - P_{\Pi_M}(\mathbf{X}\beta_0 + \mathbf{W})\|_2^2 \\ &= \|\mathbf{W} - P_{\Pi_0}(\mathbf{W})\|_2^2 - \|\mathbf{W} - P_{\Pi_M}(\mathbf{W})\|_2^2 = \|(P_{\Pi_M} - P_{\Pi_0})(\mathbf{W})\|_2^2. \end{aligned}$$

Because $M_0 \subset M$, $P_{\Pi_M} - P_{\Pi_0}$ is a projection operator of dimension $|M| - |M_0|$. Using a tail probability bound for Gaussian random variables, we have, for all $A = 2(\alpha + c) > 0$,

$$\mathbb{P}[\Delta(M)\sigma^{-2} \leq -A(|M| - |M_0|)\log n] \leq \frac{(|M| - |M_0|)}{\sqrt{2\pi A \log n}} n^{-A/2} \leq \frac{\bar{d} - d_0}{\sqrt{2\pi A \log n}} n^{-A/2}.$$

The desired result follows from union bound. \blacksquare

3. For $M \not\supseteq M_0$, let $M_1 = M_0 \setminus M$, $M_2 = M_0 \cap M$, $J_M^* = \sqrt{n\kappa_0 b_0^2 |M_1|}$. Define

$$E_3 := \left\{ \inf_{M \in \mathcal{M}, M \not\supseteq M_0} \Delta(M) \leq (J_M^*)^2 - 2\sigma J_M^* \sqrt{A \log n} - (|M| - |M_2|)A \log n \right\}. \quad (24)$$

Claim 3.

$$P_D(E_3) \leq \frac{1 + \bar{d}}{\sqrt{2\pi A \log n}} n^{-c}.$$

and under E_3^c and equation (18)

$$\Delta(M) \geq \frac{1}{2} \kappa_0 b_0^2 \sigma^2 n, \quad \forall M \in \mathcal{M}, M \not\supseteq M_0.$$

Proof. Denote $\beta_{0,M}$ the vector that agrees with β_0 on M and 0 elsewhere.

$$\begin{aligned} \Delta(M) &= \|\mathbf{Y} - P_{\Pi_M}(\mathbf{Y})\|_2^2 - \|P_{\Pi_0^\perp}(\mathbf{W})\|_2^2 \\ &= \|P_{\Pi_M^\perp}(\mathbf{X}\beta_0) + P_{\Pi_M^\perp}(\mathbf{W})\|_2^2 - \|P_{\Pi_0^\perp}(\mathbf{W})\|_2^2 \\ &= \|P_{\Pi_M^\perp}(\mathbf{X}_{M_1}\beta_{0,M_1})\|_2^2 + 2\langle P_{\Pi_M^\perp}(\mathbf{X}_{M_1}\beta_{0,M_1}), P_{\Pi_M^\perp}(\mathbf{W}) \rangle + \|P_{\Pi_M^\perp}(\mathbf{W})\|_2^2 - \|P_{\Pi_0^\perp}(\mathbf{W})\|_2^2 \\ &\geq \|P_{\Pi_M^\perp}(\mathbf{X}_{M_1}\beta_{0,M_1})\|_2^2 + 2\langle P_{\Pi_M^\perp}(\mathbf{X}_{M_1}\beta_{0,M_1}), \mathbf{W} \rangle - \|P_{\Pi_{0,M} \cap \Pi_0^\perp}(\mathbf{W})\|_2^2, \end{aligned}$$

where $\Pi_{0,M}$ is the linear subspace spanned by $X_{M_0 \cup M}$.

Let $J = \|P_{\Pi_M^\perp}(\mathbf{X}_{M_1}\beta_{0,M_1})\|_2$. Then the second term in the above equation is distributed as $N(0, \sigma^2 J^2)$. By the eigenvalue condition, we have $J^2 \geq \kappa_0 n b_0^2 |M_1|$, where κ_0 is the minimum sparse eigenvalue and b_0 is a lower bound of the signal level. To see this, observe that $M \cap M_1 = \emptyset$ and

$$\begin{aligned} J &= \|P_{\Pi_M^\perp}(\mathbf{X}_{M_1}\beta_{0,M_1})\| = \left\| (\mathbf{X}_{M_1}, \mathbf{X}_M) \begin{pmatrix} \beta_{0,M_1} \\ -(\mathbf{X}_M^T \mathbf{X}_M)^{-1} \mathbf{X}_M^T \mathbf{X}_{M_1} \beta_{0,M_1} \end{pmatrix} \right\| \\ &\geq \sqrt{n\kappa_0} \left\| \begin{pmatrix} \beta_{M_1} \\ -(X_M^T X_M)^{-1} X_M^T X_{M_1} \beta_{M_1} \end{pmatrix} \right\| \\ &\geq \sqrt{n\kappa_0} \|\beta_{M_1}\| \geq \sqrt{n\kappa_0 |M_1|} b_0, \end{aligned}$$

where the first inequality uses the sparse eigenvalue condition.

Note that the dimension of $\Pi_M \cap \Pi_0^\perp$ is at most $|M| - |M_2|$. Let $J^* := \sqrt{\kappa_0 n b_0^2}$. Using (18) we know that $J \geq J^* \geq 4\sqrt{A \log n}$. Then with probability at least $1 - (1 + |M| - |M_2|)n^{-A/2}/\sqrt{2\pi A \log n} \leq 1 - \frac{1+\bar{d}}{\sqrt{2\pi A \log n}}n^{A/2}$, we have

$$\Delta(M)\sigma^{-2} \geq J^2 - 2\sigma J\sqrt{A \log n} - (|M| - |M_2|)\sigma^2 A \log n \quad (25)$$

$$\geq (J^*)^2 - 2\sigma J^*\sqrt{A \log n} - \sigma^2 \bar{d} A \log n \quad (26)$$

$$\geq \frac{1}{2}\kappa_0 b_0^2 n \quad (27)$$

where the last inequality uses eq. (18). The claim follows from union bound. \blacksquare

4. Define

$$E_4 := \left\{ D : \inf_{M \in \mathcal{M}} -2\ell(M; D) \leq \sigma^2 n/2 \right\}. \quad (28)$$

Claim 4. $E_4 \subseteq E_1 \cup E_2 \cup E_3$.

To see this, first note that on $(E_1 \cup E_2 \cup E_3)^c$, we have $\min_{M \in \mathcal{M}} -2\ell(M; D) \geq \sigma^2(n - d_0 - \sqrt{2(n - d_0)A \log n} - A(\bar{d} - d_0) \log n) \geq n\sigma^2/2$ using eq. (18)–(19).

5. Define

$$E_5 := \{z_G < -A \log n/2\}. \quad (29)$$

Claim 5. $P_\omega(E_5) = \frac{1}{2}n^{-A/2} \leq n^{-A/2}$. On $E_4^c \cap E_5^c$, we have, using eq. (18)–(20),

$$G(D) \leq \frac{4(R + r)^2}{\sigma^2}.$$

The proof is elementary and omitted.

References

- Abowd, J. M., & Vilhuber, L. (2008). How protective are synthetic data? In *Privacy in Statistical Databases*, (pp. 239–246). Springer.
- Akaike, H. (1974). A new look at the statistical model identification. *Automatic Control, IEEE Transactions on*, 19(6), 716–723.
- Barak, B., Chaudhuri, K., Dwork, C., Kale, S., McSherry, F., & Talwar, K. (2007). Privacy, accuracy, and consistency too: A holistic solution to contingency table release. In *Proceedings of the 26th Symposium on Principles of Database Systems*, (pp. 273–282).
- Bassily, R., Smith, A., & Thakurta, A. (2014). Private empirical risk minimization: Efficient algorithms and tight error bounds. In *FOCS*, (pp. 464–473). IEEE.

- Beimel, A., Brenner, H., Kasiviswanathan, S. P., & Nissim, K. (2014). Bounds on the sample complexity for private learning and private data release. *Machine Learning*, 94(3), 401–437.
URL <http://dx.doi.org/10.1007/s10994-013-5404-1>
- Beimel, A., Kasiviswanathan, S. P., & Nissim, K. (2010). Bounds on the sample complexity for private learning and private data release. In *TCC*, (pp. 437–454).
- Beimel, A., Nissim, K., & Stemmer, U. (2015). Learning privately with labeled and unlabeled examples. In *Proceedings of the Twenty-Sixth Annual ACM-SIAM Symposium on Discrete Algorithms, SODA 2015, San Diego, CA, USA, January 4-6, 2015*, (pp. 461–477).
URL <http://dx.doi.org/10.1137/1.9781611973730.32>
- Bun, M., Nissim, K., Stemmer, U., & Vadhan, S. P. (2015). Differentially private release and learning of threshold functions. In *IEEE 56th Annual Symposium on Foundations of Computer Science, FOCS 2015, Berkeley, CA, USA, 17-20 October, 2015*, (pp. 634–649).
URL <http://dx.doi.org/10.1109/FOCS.2015.45>
- Chaudhuri, K., & Hsu, D. (2011). Sample complexity bounds for differentially private learning. In *COLT*, vol. 19, (pp. 155–186).
- Chaudhuri, K., Monteleoni, C., & Sarwate, A. D. (2011). Differentially private empirical risk minimization. *The Journal of Machine Learning Research*, 12, 1069–1109.
- Dalenius, T. (1977). *Statistik Tidskrift*, 15, 429–444.
- Duchi, J. C., Jordan, M. I., & Wainwright, M. J. (2013). Local privacy and statistical minimax rates. In *Foundations of Computer Science (FOCS), 2013 IEEE 54th Annual Symposium on*, (pp. 429–438). IEEE.
- Dwork, C. (2006). Differential privacy. In *Proceedings of the 33rd International Colloquium on Automata, Languages and Programming (ICALP)(2)*, (pp. 1–12).
- Dwork, C., & Lei, J. (2009). Differential privacy and robust statistics. In *Proceedings of the 41st Annual ACM Symposium on Theory of Computing*.
- Dwork, C., McSherry, F., Nissim, K., & Smith, A. (2006). Calibrating noise to sensitivity in private data analysis. In *Proceedings of the 3rd Theory of Cryptography Conference*, (pp. 265–284).
- Dwork, C., Su, W., & Zhang, L. (2015). Private false discovery rate control. *arXiv preprint arXiv:1511.03803*.
- Fan, J., & Li, R. (2001). Variable selection via nonconcave penalized likelihood and its oracle properties. *Journal of the American Statistical Association*, 96(456), 1348–1360.

- Fienberg, S., & Slavković, A. (2010). *Data Privacy and Confidentiality*, (pp. 342–345). International Encyclopedia of Statistical Science. Springer-Verlag.
- Fienberg, S. E., Rinaldo, A., & Yang, X. (2010). Differential privacy and the risk-utility tradeoff for multi-dimensional contingency tables. In *Proceedings of the 2010 international conference on Privacy in statistical databases*, PSD’10, (pp. 187–199). Springer-Verlag.
- Fienberg, S. E., Slavkovic, A., & Uhler, C. (2011). Privacy preserving gwas data sharing. In *Data Mining Workshops (ICDMW), 2011 IEEE 11th International Conference on*, (pp. 628–635). IEEE.
- Gaboardi, M., Lim, H., Rogers, R., & Vadhan, S. (2016). Differentially private chi-squared hypothesis testing: Goodness of fit and independence testing. *arXiv preprint arXiv:1602.03090*.
- Ganta, S. R., Kasiviswanathan, S. P., & Smith, A. (2008). Composition attacks and auxiliary information in data privacy. In *Proceedings of the 14th ACM SIGKDD international conference on Knowledge discovery and data mining*, (pp. 265–273). ACM.
- Hardt, M., Ligett, K., & McSherry, F. (2010). A simple and practical algorithm for differentially private data release. *arXiv preprint arXiv:1012.4763*.
- Hundepool, A., Domingo-Ferrer, J., Franconi, L., Giessing, S., Nordholt, E. S., Spicer, K., & de Wolf, P.-P. (2012). *Statistical Disclosure Control*. Wiley.
- Johnson, A., & Shmatikov, V. (2013). Privacy-preserving data exploration in genome-wide association studies. In *Proceedings of the 19th ACM SIGKDD International Conference on Knowledge Discovery and Data Mining*, KDD ’13, (pp. 1079–1087). New York, NY, USA: ACM.
- Karwa, V., Kifer, D., & Slavkovic, A. (2015). Private posterior distributions from variational approximations. *arXiv preprint arXiv:1511.07896*.
- Karwa, V., & Slavkovic, A. (2012). Differentially private graphical degree sequences and synthetic graphs. In J. Domingo-Ferrer, & I. Tinnirello (Eds.) *Privacy in Statistical Databases*, vol. 7556 of *Lecture Notes in Computer Science*, (pp. 273–285). Springer Berlin Heidelberg.
- Karwa, V., & Slavković, A. (2016). Inference using noisy degrees: Differentially private beta-model and synthetic graphs. *The Annals of Statistics*, 44(1), 87–112.
- Karwa, V., Slavkovic, A., & Krivitsky, P. (2014). Differentially private exponential random graphs. In J. Domingo-Ferrer (Ed.) *Privacy in Statistical Databases*, vol. 8744 of *Lecture Notes in Computer Science*, (pp. 143–155). Springer International Publishing.
- Kasiviswanathan, S. P., Lee, H. K., Nissim, K., Raskhodnikova, S., & Smith, A. (2011). What can we learn privately? *SIAM Journal on Computing*, 40(3), 793–826.

- Kasiviswanathan, S. P., & Smith, A. (2008). On the semantics of differential privacy: A bayesian formulation. *arXiv preprint arXiv:0803.3946*.
- Laurent, B., & Massart, P. (2000). Adaptive estimation of a quadratic functional by model selection. *The Annals of Statistics*, 28, 1302–1338.
- Lei, J. (2011). Differentially private m-estimators. In *Advances in Neural Information Processing Systems*, (pp. 361–369).
- McSherry, F., & Talwar, K. (2007). Mechanism design via differential privacy. In *Proceedings of the 48th Annual Symposium on Foundations of Computer Science*, (pp. 94–103).
- Nishii, R. (1984). Asymptotic properties of criteria for selection of variables in multiple regression. *The Annals of Statistics*, 12(2), 758–765.
- Picard, R. R., & Cook, R. D. (1984). Cross-validation of regression models. *Journal of the American Statistical Association*, 79(387), 575–583.
- Rubin, D. B. (1993). Statistical disclosure limitation. *Journal of official Statistics*, 9(2), 461–468.
- Schwarz, G., et al. (1978). Estimating the dimension of a model. *The annals of statistics*, 6(2), 461–464.
- Sheffet, O. (2015). Differentially private least squares: Estimation, confidence and rejecting the null hypothesis. *arXiv preprint arXiv:1507.02482*.
- Smith, A. (2011). Privacy-preserving statistical estimation with optimal convergence rates. In *STOC*.
- Smith, A., & Thakurta, A. (2013). Differentially private feature selection via stability arguments, and the robustness of the lasso. In *COLT 2013 - The 26th Annual Conference on Learning Theory, June 12-14, 2013, Princeton University, NJ, USA*, (pp. 819–850). URL <http://jmlr.org/proceedings/papers/v30/Guha13.html>
- Solea, E. (2014). *Differentially Private Hypothesis Testing For Normal Random Variables*. Master’s thesis, The Pennsylvania State University.
- Tibshirani, R. (1996). Regression shrinkage and selection via the lasso. *Journal of the Royal Statistical Society. Series B (Methodological)*, (pp. 267–288).
- Uhler, C., Slavkovic, A., & Fienberg, S. E. (2013). Privacy-preserving data sharing for genome-wide association studies. *Journal of Privacy and Confidentiality*, 5(1).
- Wang, Y., Lee, J., & Kifer, D. (2015). Differentially private hypothesis testing, revisited. *arXiv preprint arXiv:1511.03376*.

- Wasserman, L., & Zhou, S. (2010). A statistical framework for differential privacy. *Journal of the American Statistical Association*, 105(489), 375–389.
- Willenborg, L., & De Waal, T. (1996). *Statistical disclosure control in practice*. 111. Springer Science & Business Media.
- Yu, F., Fienberg, S. E., Slavkovic, A. B., & Uhler, C. (2014). Scalable privacy-preserving data sharing methodology for genome-wide association studies. *Journal of Biomedical Informatics*, 50, 133–141.