

# Complex Langevin in Lattice QCD: dynamic stabilisation and the phase diagram

GERT AARTS, FELIPE ATTANASIO<sup>†</sup>, BENJAMIN JÄGER

Department of Physics, College of Science, Swansea University, United Kingdom

DÉNES SEXTY

Department of Physics, Bergische Universität Wuppertal, Wuppertal, Germany

Complex Langevin simulations provide an alternative to sample path integrals with complex weights and therefore are suited to determine the phase diagram of QCD from first principles. We use our proposed method of Dynamic Stabilisation (DS) to ensure improved convergence to the right limit and present new systematic tests of this technique. We also show results on QCD in the limit of heavy quarks and an analysis of DS compared to known results from reweighting.

PACS numbers: 12.38.-t, 12.38.Gc, 12.38.Aw

## 1. Introduction

The QCD phase diagram is subject of active research both theoretically and experimentally. On the theoretical side the phase structure of QCD has implications for the understanding of the early and modern universe, where the quark-gluon plasma and compact astrophysical objects still pose unanswered questions. For experimentalists, it of course serves as guide for current and future experiments and facilities.

Studies of the phase diagram at non-zero chemical potential  $\mu$  are extremely difficult due to a complex statistical weight, which results in what is known as the *sign problem*. In QCD this behaviour is caused by the quark determinant, after the quark fields have been integrated out

$$[\det M(U, \mu)]^* = \det M(U, -\mu^*), \quad (1)$$

---

<sup>†</sup> Presented at the Workshop “Excited QCD 2016”, Portugal 6-11 March 2016

where  $U$  generically represents the gauge links. This leads to an exponentially small overlap between the full theory and the “phase quenched” version which can be simulated using Monte Carlo techniques. A review of different methods for the QCD phase diagram can be found in [1].

## 2. Complex Langevin equation

The complex Langevin method is based on stochastic quantisation [2]. In this framework the dynamical variables are evolved along a fictitious time direction  $\theta$  according to a Langevin equation. On the lattice for an SU(3) gauge theory the equation for the gauge links  $U$  reads [3]

$$U_{x\mu}(\theta + \varepsilon) = \exp \left[ i\lambda^a X_{x\mu}^a \right] U_{x\mu}(\theta), \quad X_{x\mu}^a = -\varepsilon D_{x\mu}^a S[U] + \sqrt{\varepsilon} \eta_{x\mu}^a, \quad (2)$$

where  $\lambda^a$  are the Gell-Mann matrices normalised to  $\text{Tr}[\lambda^a \lambda^b] = 2\delta^{ab}$ ,  $\varepsilon$  is the Langevin stepsize chosen adaptively [4],  $\eta_{x\mu}^a$  are white noise fields satisfying  $\langle \eta_{x\mu}^a \eta_{y\nu}^b \rangle = 2\delta^{ab} \delta_{xy} \delta_{\mu\nu}$ ,  $S[U]$  is the action, and the gauge group derivative  $D_{x\mu}^a$  is defined as

$$D_{x\mu}^a f(U) = \left. \frac{\partial}{\partial \alpha} f(e^{i\alpha \lambda^a} U_{x\mu}) \right|_{\alpha=0}. \quad (3)$$

To deal with the sign problem we allow the gauge links to be non-unitary [5–7], which amounts to extending the group SU(3) to SL(3,  $\mathbb{C}$ ). The latter is a non-compact manifold and therefore large excursions into the imaginary directions may occur during the simulation.

As a way to keep track of such excursions and prevent them from becoming large – by using gauge transformations between Langevin updates – we measure the distance from SU(3),

$$d = \frac{1}{N_s^3 N_\tau} \sum_{x,\mu} \text{Tr} \left[ U_{x\mu} U_{x\mu}^\dagger - \mathbf{1} \right]^2 \geq 0, \quad (4)$$

where  $N_s$  and  $N_\tau$  are the lattice extents in the spatial and temporal directions, respectively. These transformations, known as gauge cooling [8], are constructed to minimise  $d$  and are defined as

$$U_{x\mu} \rightarrow \Lambda_x U_{x\mu} \Lambda_{x+\mu}^{-1}, \quad \Lambda_x = \exp \left[ -\varepsilon \alpha \lambda^a f_x^a \right], \quad (5)$$

where

$$f_x^a = 2 \text{Tr} \left[ \lambda^a \left( U_{x\mu} U_{x\mu}^\dagger - U_{x-\mu,\mu}^\dagger U_{x-\mu,\mu} \right) \right]. \quad (6)$$

The parameter  $\alpha$  and the number of cooling steps are chosen adaptively in order to maximise the method’s efficiency [9]. A study of this method applied to one- and two-dimensional QCD can be found in [10].

### 3. QCD with heavy quarks

In the heavy quarks approximation [11,12] the quarks evolve only in the Euclidean time and the fermion determinant decouples as a product over spatial points. This corresponds to the leading order in a spatial hopping expansion,

$$\det M(U, \mu) = \prod_{\vec{x}} \left\{ \det [1 + (2\kappa e^\mu)^{N_\tau} \mathcal{P}_{\vec{x}}] \det [1 + (2\kappa e^{-\mu})^{N_\tau} \mathcal{P}_{\vec{x}}^{-1}] \right\}^2, \quad (7)$$

containing the hopping parameter  $\kappa$ , the Polyakov loop and its inverse

$$\mathcal{P}_{\vec{x}} = \prod_{\tau=0}^{N_\tau-1} U_4(\vec{x}), \quad \mathcal{P}_{\vec{x}}^{-1} = \prod_{\tau=N_\tau-1}^0 U_4^{-1}(\vec{x}). \quad (8)$$

This model exhibits a sign problem and a transition to a high density phase. At zero temperature this is expected to happen at  $\mu = \mu_c^0 \equiv -\ln(2\kappa)$ .

Comparisons with the hopping expansion to all orders [13] as well as multi-parameter reweighting [14] have been investigated recently. For a study of a strong coupling expansion ( $\beta \rightarrow 0$ ) combined with hopping parameter expansion see [15, 16]. Effective model studies for heavy quarks QCD include [17, 18]. Complex Langevin investigations can be found in [19–24].

The expectation value of the Polyakov loop is known to be approximately 0 in a confined phase and non-zero otherwise. We looked at its Binder cumulant [25] to map the boundary between these phases. The resulting plot is shown in figure 1, where a lattice spacing of  $a \sim 0.15$  fm, determined using the Wilson flow [26], has been used to convert the temperature to physical units. Our analysis included only data with small unitarity norm ( $d \leq 0.03$ ) to allow for reliable predictions, as will be explained later.

### 4. Dynamic Stabilisation

There are situations where gauge cooling is not sufficient to keep the system from going too far into  $\text{SL}(3, \mathbb{C})$  and when the distance becomes  $\mathcal{O}(0.1)$  observables seem to converge to a wrong limit [9].

We have developed a new technique called dynamic stabilisation, which consists of adding a  $\text{SU}(3)$  – but not  $\text{SL}(3, \mathbb{C})$  – gauge invariant force to the Langevin drift that is trivial in the continuum limit and grows with  $d$ . It is given by

$$M_x^a[U] = ib_x^a (b_x^c b_x^c)^3, \quad b_x^a = \sum_\nu \text{Tr} \left[ \lambda^a U_{x\nu} U_{x\nu}^\dagger \right]. \quad (9)$$




Fig. 1. Phase boundary of the HDQCD model. The phase boundary was defined as the region where the Binder cumulant transitions from zero to non-zero values within statistical precision. At  $\mu/\mu_c^0 > 1$  we see a similar pattern, which is an effect from particle-hole symmetry [27].

To better benefit from this technique we have also extended our Langevin equation to second order in the Langevin stepsize [28]

$$U_{x\mu}(\theta + \varepsilon/2) = \exp[i\lambda^a X_{x\mu}^a] U_{x\mu}(\theta), \quad (10)$$

$$U_{x\mu}(\theta + \varepsilon) = \exp[i\lambda^a \gamma (X_{x\mu}^{a'} + X_{x\mu}^a)] U_{x\mu}(\theta), \quad (11)$$

where the new drifts read

$$X_{x\mu}^a = -\varepsilon D_{x\mu}^a S[U(\theta)] + i\varepsilon \alpha_{DS} M_x^a[U(\theta)] + \sqrt{\varepsilon} \eta_{x\mu}^a(\theta), \quad (12)$$

$$X_{x\mu}^{a'} = -\varepsilon D_{x\mu}^a S[U(\theta + \varepsilon/2)] + i\varepsilon \alpha_{DS} M_x^a[U(\theta + \varepsilon/2)] + \sqrt{\varepsilon} \eta_{x\mu}^a(\theta), \quad (13)$$

with  $\gamma = 1/2(1 + \varepsilon C_A/6)$ ,  $\alpha_{DS}$  a real parameter, the noise is  $\langle \eta_{x\mu}^a \eta_{y\nu}^b \rangle = 2(1 - \varepsilon C_A/2) \delta^{ab} \delta_{xy} \delta_{\mu\nu}$  and  $C_A = 3$  is the Casimir invariant in the adjoint representation.

A comparison between dynamic stabilisation combined with one gauge cooling step with a known result from reweighting,  $\langle P \rangle = 0.202717(66)$ , is shown in figure 2 for different  $\alpha_{DS}$ 's in a lattice of volume  $10^3 \times 4$ ,  $\beta = 5.8$ ,  $\kappa = 0.04$  and  $\mu = 0.7$ . Agreement with reweighting results is seen for  $\alpha_{DS}$  sufficiently large. For this setup gauge cooling gives  $\langle P \rangle = 0.0027(55)$  if the points with unitarity norm larger than 0.03 are included.

## 5. Conclusions and Outlook

We have shown that the complex Langevin equation method allows simulation of theories that exhibit the sign problem, provided that large ex-




Fig. 2. Polyakov loop in the HDQCD model for different values of  $\alpha_{DS}$ . The inset shows a zoom into the region of agreement.

cursions into the complex directions are suppressed. The gauge cooling technique provided the first consistent way of achieving that and allowed us to map the phase boundary in the limit of heavy quarks.

Our method of dynamic stabilisation, combined with gauge cooling, provides greater control on how far from SU(3) the system can go. It has been shown to agree with known results in the heavy quarks approximation. Further tests with other physical parameters are needed to ensure its reliability.

The next step is to apply these techniques to QCD with fully dynamical quarks [29].

## 6. Acknowledgements

We are grateful for the computing resources made available by HPC Wales. We acknowledge the STFC grants ST/L000369/1, ST/K000411/1, ST/H008845/1, ST/K005804/1 and ST/K005790/1, the STFC DiRAC HPC Facility ([www.dirac.ac.uk](http://www.dirac.ac.uk)), STFC grant ST/L000369/1, the Royal Society and the Wolfson Foundation. FA is grateful for the support through the Brazilian government programme “Science without Borders” under scholarship number BEX 9463/13-5.

## REFERENCES

- [1] P. de Forcrand, PoS LAT **2009** (2009) 010 [arXiv:1005.0539 [hep-lat]].
- [2] G. Parisi and Y. s. Wu, Sci. Sin. **24** (1981) 483.
- [3] P. H. Damgaard and H. Huffer, Phys. Rept. **152** (1987) 227. doi:10.1016/0370-1573(87)90144-X
- [4] G. Aarts, F. A. James, E. Seiler and I. O. Stamatescu, Phys. Lett. B **687** (2010) 154 [arXiv:0912.0617 [hep-lat]].
- [5] J. Klauder, Acta Phys. Austriaca Suppl. **XXXV** (1983) 251; J. Phys. A: Math. Gen. **16**, L317-319 (1983); Phys. Rev. A **29**, 2036-2047 (1984).

- [6] G. Parisi, Phys. Lett. B **131** (1983) 393.
- [7] G. Aarts, E. Seiler and I. O. Stamatescu, Phys. Rev. D **81** (2010) 054508 [arXiv:0912.3360 [hep-lat]].
- [8] E. Seiler, D. Sexty and I. O. Stamatescu, Phys. Lett. B **723** (2013) 213 [arXiv:1211.3709 [hep-lat]].
- [9] G. Aarts, L. Bongiovanni, E. Seiler, D. Sexty and I. O. Stamatescu, Eur. Phys. J. A **49** (2013) 89 doi:10.1140/epja/i2013-13089-4 [arXiv:1303.6425 [hep-lat]].
- [10] J. Bloch, J. Mahr and S. Schmalzbauer, arXiv:1508.05252 [hep-lat].
- [11] I. Bender, T. Hashimoto, F. Karsch, V. Linke, A. Nakamura, M. Plewnia, I. O. Stamatescu and W. Wetzel, Nucl. Phys. Proc. Suppl. **26** (1992) 323.
- [12] G. Aarts and I. O. Stamatescu, JHEP **0809** (2008) 018 [arXiv:0807.1597 [hep-lat]].
- [13] G. Aarts, E. Seiler, D. Sexty and I. O. Stamatescu, Phys. Rev. D **90** (2014) 11, 114505 [arXiv:1408.3770 [hep-lat]].
- [14] Z. Fodor, S. D. Katz, D. Sexty and C. Török, arXiv:1508.05260 [hep-lat].
- [15] P. de Forcrand, J. Langelage, O. Philipsen and W. Unger, Phys. Rev. Lett. **113** (2014) 15, 152002 [arXiv:1406.4397 [hep-lat]].
- [16] J. Glesaaen, M. Neuman and O. Philipsen, JHEP **1603** (2016) 100 doi:10.1007/JHEP03(2016)100 [arXiv:1512.05195 [hep-lat]].
- [17] P. Scior and L. von Smekal, arXiv:1602.04614 [hep-lat].
- [18] S. Ejiri and H. Yoneyama, arXiv:1511.04811 [hep-lat].
- [19] G. Aarts, F. Attanasio, B. Jäger, E. Seiler, D. Sexty and I. O. Stamatescu, PoS LATTICE **2014** (2014) 200 [arXiv:1411.2632 [hep-lat]].
- [20] G. Aarts, F. Attanasio, B. Jäger, E. Seiler, D. Sexty and I. O. Stamatescu, AIP Conf. Proc. **1701** (2016) 020001 doi:10.1063/1.4938590 [arXiv:1412.0847 [hep-lat]].
- [21] G. Aarts, F. Attanasio, B. Jäger, E. Seiler, D. Sexty and I. O. Stamatescu, Acta Phys. Polon. Supp. **8** (2015) 2, 405 [arXiv:1506.02547 [hep-lat]].
- [22] G. Aarts, F. Attanasio, B. Jäger, E. Seiler, D. Sexty and I. O. Stamatescu, arXiv:1510.09098 [hep-lat].
- [23] G. Aarts, F. Attanasio, B. Jäger, E. Seiler, D. Sexty and I. O. Stamatescu, arXiv:1510.09100 [hep-lat].
- [24] G. Aarts, F. Attanasio, B. Jäger and D. Sexty, arXiv:1606.05561 [hep-lat].
- [25] K. Binder, Z. Phys. B **43** (1981) 119. doi:10.1007/BF01293604
- [26] S. Borsanyi *et al.*, JHEP **1209** (2012) 010 doi:10.1007/JHEP09(2012)010 [arXiv:1203.4469 [hep-lat]].
- [27] T. Rindlisbacher and P. de Forcrand, JHEP **1602** (2016) 051 doi:10.1007/JHEP02(2016)051 [arXiv:1509.00087 [hep-lat]].
- [28] M. Fukugita, Y. Oyanagi and A. Ukawa, Phys. Rev. D **36** (1987) 824. doi:10.1103/PhysRevD.36.824
- [29] D. Sexty, Phys. Lett. B **729** (2014) 108 [arXiv:1307.7748 [hep-lat]].