

Entangled Harmonic Oscillators and Space-time Entanglement

Sibel Bařkal¹

Department of Physics, Middle East Technical University, 06531 Ankara, Turkey

Young S. Kim²

Center for Fundamental Physics, University of Maryland,
College Park, Maryland 20742, U.S.A.

Marilyn E. Noz³

Department of Radiology,
New York University New York, New York 10016, U.S.A.

Abstract

The mathematical basis for the Gaussian entanglement is discussed in detail, as well as its implications in the internal space-time structure of relativistic extended particles. It is shown that the Gaussian entanglement shares the same set of mathematical formulas with the harmonic oscillator in the Lorentz-covariant world. It is thus possible to transfer the concept of entanglement to the Lorentz-covariant picture of the bound state which requires both space and time separations between two constituent particles. These space and time variables become entangled as the bound state moves with a relativistic speed. It is shown also that our inability to measure the time-separation variable leads to an entanglement entropy together with a rise in the temperature of the bound state. As was noted by Paul A. M. Dirac in 1963, the system of two oscillators contains the symmetries of $O(3,2)$ de Sitter group containing two $O(3,1)$ Lorentz groups as its subgroups. Dirac noted also that the system contains the symmetry of the $Sp(4)$ group which serves as the basic language for two-mode squeezed states. Since the $Sp(4)$ symmetry contains both rotations and squeezes, one interesting case is the combination of rotation and squeeze resulting in a shear. While the current literature is mostly on the entanglement based on squeeze along the normal coordinates, the shear transformation is an interesting future possibility. The mathematical issues on this problem are clarified.

¹electronic address: baskal@newton.physics.metu.edu.tr

²electronic address: yskim@physics.umd.edu

³electronic address: mairlyne.noz@gmail.com

1 Introduction

Entanglement problems deal with fundamental issues in physics. Among them, the Gaussian entanglement is of current interest not only in quantum optics [1, 2, 3, 4] but also in other dynamical systems [3, 5, 6, 7, 8]. The underlying mathematical language for this form of entanglement is that of harmonic oscillators. In this paper, we present first the mathematical tools which are and may be useful in this branch of physics.

The entangled Gaussian state is based on the formula

$$\frac{1}{\cosh \eta} \sum_k (\tanh \eta)^k \chi_k(x) \chi_k(y), \quad (1)$$

where $\chi_n(x)$ is the n^{th} excited-state oscillator wave function.

In Chapter 16 of their book [9], Walls and Milburn discussed in detail the role of this formula in the theory of quantum information. Earlier, this formula played the pivotal role for Yuen to formulate his two-photon coherent states or two-mode squeezed states [10]. The same formula was used by Yurke and Patasek in 1987 [11] and by Ekert and Knight [12] for the two-mode squeezed state where one of the photons is not observed. The effect of entanglement is to be seen from the beam splitter experiments [13, 14].

In this paper, we point out first that the series of Eq.(1) can also be written as a squeezed Gaussian form

$$\frac{1}{\sqrt{\pi}} \exp \left\{ -\frac{1}{4} \left[e^{-2\eta}(x+y)^2 + e^{2\eta}(x-y)^2 \right] \right\}, \quad (2)$$

which becomes

$$\frac{1}{\sqrt{\pi}} \exp \left\{ -\frac{1}{2} (x^2 + y^2) \right\}, \quad (3)$$

when $\eta = 0$.

We can obtain the squeezed form of Eq.(2) by replacing x and y by x' and y' respectively, where

$$\begin{pmatrix} x' \\ y' \end{pmatrix} = \begin{pmatrix} \cosh \eta & -\sinh \eta \\ -\sinh \eta & \cosh \eta \end{pmatrix} \begin{pmatrix} x \\ y \end{pmatrix}. \quad (4)$$

If x and y are replaced by z and t , Eq.(4) becomes the formula for the Lorentz boost along the z direction. Indeed, the Lorentz boost is a squeeze transformation [3, 15].

The squeezed Gaussian form of Eq.(2) plays the key role in studying boosted bound states in the Lorentz-covariant world [16, 17, 18, 19, 20], where z and t are the space and time separations between two constituent particles. Since the mathematics of this physical system is the same as the series given in Eq.(1), the physical concept of entanglement can be transferred to the Lorentz-covariant bound state, as illustrated in Fig. 1.

We can approach this problem from the system of two harmonic oscillators. In 1963, Paul A. M. Dirac studied the symmetry of this two-oscillator system and discussed all possible transformations applicable to this oscillator [21]. He concluded that there are ten possible generators of transformations satisfying a closed set of commutation




Figure 1: One mathematics for two branches of physics. Let us look at Eq.(1) and Eq.(2) applicable to quantum optics and special relativity respectively. They are the same formula from the Lorentz group with different variables as in the case of the LCR circuit and the mechanical oscillator sharing the same second-order differential equation.

relations. He then noted that this closed set corresponds to the Lie algebra of the $O(3, 2)$ de Sitter group which is the Lorentz group applicable to three space-like and two time-like dimensions. This $O(3, 2)$ group has two $O(3, 1)$ Lorentz groups as its subgroups.

We note that the Lorentz group is the language of special relativity, while the harmonic oscillator is one of the major tools for interpreting bound states. Therefore, Dirac's two-oscillator system can serve as a mathematical framework for understanding quantum bound systems in the Lorentz-covariant world.

Within this formalism, the series given in Eq.(1) can be produced from the ten-generator Dirac system. In discussing the oscillator system, the standard procedure is to use the normal coordinates defined as

$$u = \frac{x + y}{\sqrt{2}}, \quad \text{and} \quad v = \frac{x - y}{\sqrt{2}}. \quad (5)$$

In terms of these variables, the transformation given in Eq.(4) takes the form

$$\begin{pmatrix} u' \\ v' \end{pmatrix} = \begin{pmatrix} e^{-\eta} & 0 \\ 0 & e^{\eta} \end{pmatrix} \begin{pmatrix} u \\ v \end{pmatrix}, \quad (6)$$

where this is a squeeze transformation along the normal coordinates. While the normal-coordinate transformation is a standard procedure, it is interesting to note that it also serves as a Lorentz boost [18].

With these preparations, we shall study in Sec. 2, the system of two oscillators and coordinate transformations of current interest. It is pointed out in Sec. 3 that there are ten different generators for transformations, including those discussed in Sec. 2. It is noted that Dirac derived ten generators of transformations applicable to these

oscillators, and they satisfy the closed set of commutation relations which is the same as the Lie algebra of the $O(3, 2)$ de Sitter group containing two Lorentz groups among its subgroups. In Sec. 4, Dirac's ten-generator symmetry is studied in the Wigner phase-space picture, and it is shown that Dirac's symmetry contains both canonical and Lorentz transformations.

While the Gaussian entanglement starts from the oscillator wave function in its ground state, we study in Sec. 5 the entanglements of excited oscillator states. We give a detailed explanation of how the series of Eq.(1) can be derived from the squeezed Gaussian function of Eq.(2).

In Sec. 6, we study in detail how the sheared state can be derived from a squeezed state. It appears to be a rotated squeezed state, but this is not the case. In Sec. 7, we study what happens when one of the two entangled variables is not observed within the framework of Feynman's rest of the universe [22, 23].

In Sec. 8, we note that most of the mathematical formulas in this paper have been used earlier for understanding relativistic extended particles in the Lorentz-covariant harmonic oscillator formalism [24, 20, 25, 26, 27, 28]. These formulas allow us to transport the concept of entanglement from the current problem of physics to quantum bound states in the Lorentz-covariant world. The time separation between the constituent particles is not observable, and is not known in the present form of quantum mechanics. However, this variable gives its effect in the real world by entangling itself with the longitudinal variable.

2 Two-dimensional Harmonic Oscillators

The Gaussian form

$$\left[\frac{1}{\sqrt{\pi}} \right]^{1/4} \exp\left(-\frac{x^2}{2}\right) \quad (7)$$

is used for many branches of science. For instance, we can construct this function by throwing dice.

In physics, this is the wave function for the one-dimensional harmonic oscillator in the ground state. This function is also used for the vacuum state in quantum field theory, as well as the zero-photon state in quantum optics. For excited oscillator states, the wave function takes the form

$$\chi_n(x) = \left[\frac{1}{\sqrt{\pi} 2^n n!} \right]^{1/2} H_n(x) \exp\left(\frac{-x^2}{2}\right), \quad (8)$$

where $H_n(x)$ is the Hermite polynomial of the n^{th} degree. The properties of this wave function are well known, and it becomes the Gaussian form of Eq.(7) when $n = 0$.

We can now consider the two-dimensional space with the orthogonal coordinate variables x and y , and the same wave function with the y variable:

$$\chi_m(y) = \left[\frac{1}{\sqrt{\pi} 2^m m!} \right]^{1/2} H_m(y) \exp\left(\frac{-y^2}{2}\right), \quad (9)$$

and construct the function

$$\psi^{n,m}(x, y) = [\chi_n(x)] [\chi_m(y)]. \quad (10)$$

This form is clearly separable in the x and y variables. If n and m are zero, the wave function becomes

$$\psi^{0,0}(x, y) = \frac{1}{\sqrt{\pi}} \exp \left\{ -\frac{1}{2} (x^2 + y^2) \right\}. \quad (11)$$

Under the coordinate rotation

$$\begin{pmatrix} x' \\ y' \end{pmatrix} = \begin{pmatrix} \cos \theta & -\sin \theta \\ \sin \theta & \cos \theta \end{pmatrix} \begin{pmatrix} x \\ y \end{pmatrix} \quad (12)$$

this function remains separable. This rotation is illustrated in Fig. 2. This is a transformation very familiar to us.

We can next consider the scale transformation of the form

$$\begin{pmatrix} x' \\ y' \end{pmatrix} = \begin{pmatrix} e^\eta & 0 \\ 0 & e^{-\eta} \end{pmatrix} \begin{pmatrix} x \\ y \end{pmatrix}. \quad (13)$$

This scale transformation is also illustrated in Fig. 2. This area-preserving transformation is known as the squeeze. Under this transformation, the Gaussian function is still separable.

If the direction of the squeeze is rotated by 45° , the transformation becomes the diagonal transformation of Eq.(6). Indeed, this is a squeeze in the normal coordinate system. This form of squeeze is most commonly used for squeezed states of light as well as the subject of entanglements. It is important to note that, in terms of the x and y , variables, this transformation can be written as Eq.(4) [18]. In 1905, Einstein used this form of squeeze transformation for the longitudinal and time-like variables. This is known as the Lorentz boost.

In addition, we can consider the transformation of the form

$$\begin{pmatrix} x' \\ y' \end{pmatrix} = \begin{pmatrix} 1 & 2\alpha \\ 0 & 1 \end{pmatrix} \begin{pmatrix} x \\ y \end{pmatrix}. \quad (14)$$

This transformation shears the system as is shown in Fig. 2.

After the squeeze or shear transformation, the wave function of Eq.(10) becomes non-separable, but it can still be written as a series expansion in terms of the oscillator wave functions. It can take the form

$$\psi(x, y) = \sum_{n,m} A_{n,m} \chi_n(x) \chi_m(y), \quad (15)$$

with

$$\sum_{n,m} |A_{n,m}|^2 = 1,$$

if $\psi(x, y)$ is normalized, as was the case for the Gaussian function of Eq.(11).




Figure 2: Transformations in the two-dimensional space. The object can be rotated, squeezed, or sheared. In all three cases, the area remains invariant.

2.1 Squeezed Gaussian Function

Under the squeeze along the normal coordinate, the Gaussian form of Eq.(11) becomes

$$\psi_\eta(x, y) = \frac{1}{\sqrt{\pi}} \exp \left\{ -\frac{1}{4} \left[e^{-2\eta}(x+y)^2 + e^{2\eta}(x-y)^2 \right] \right\}, \quad (16)$$

which was given in Eq.(2). This function is not separable in the x and y variables. These variables are now entangled. We obtain this form by replacing, in the Gaussian function of Eq.(11), the x and y variables by x' and y' respectively, where

$$x' = (\cosh \eta)x - (\sinh \eta)y, \quad \text{and} \quad y' = (\cosh \eta)y - (\sinh \eta)x. \quad (17)$$

This form of squeeze is illustrated in Fig. 3, and the expansion of this squeezed Gaussian function becomes the series given in Eq.(1) [20, 26]. This aspect will be discussed in detail in Sec. 5.

In 1976 [10], Yuen discussed two-photon coherent states, often called squeezed states of light. This series expansion served as the starting point for two-mode squeezed states. More recently, in 2003, Giedke *et al.* [1] used this formula to formulate the concept of the Gaussian entanglement.

There is another way to derive the series. For the harmonic oscillator wave functions, there are step-down and step-up operators [17]. These are defined as

$$a = \frac{1}{\sqrt{2}} \left(x + \frac{\partial}{\partial x} \right), \quad \text{and} \quad a^\dagger = \frac{1}{\sqrt{2}} \left(x - \frac{\partial}{\partial x} \right). \quad (18)$$

If they are applied to the oscillator wave function, we have

$$a \chi_n(x) = \sqrt{n} \chi_{n-1}(x), \quad \text{and} \quad a^\dagger \chi_n(x) = \sqrt{n+1} \chi_{n+1}(x). \quad (19)$$




Figure 3: Squeeze along the 45° direction, discussed most frequently in the literature.

Likewise, we can introduce b and b^\dagger operators applicable to $\chi_n(y)$:

$$b = \frac{1}{\sqrt{2}} \left(y + \frac{\partial}{\partial y} \right), \quad \text{and} \quad b^\dagger = \frac{1}{\sqrt{2}} \left(y - \frac{\partial}{\partial y} \right). \quad (20)$$

Thus

$$\begin{aligned} (a^\dagger)^n \chi^0(x) &= \sqrt{n!} \chi_n(x), \\ (b^\dagger)^n \chi^0(y) &= \sqrt{n!} \chi_n(y), \end{aligned} \quad (21)$$

and

$$a \chi_0(x) = b \chi_0(y) = 0. \quad (22)$$

In terms of these variables, the transformation leading the Gaussian function of Eq.(11) to its squeezed form of Eq.(16) can be written as

$$\exp \left\{ \frac{\eta}{2} (a^\dagger b^\dagger - a b) \right\}, \quad (23)$$

which can also be written as

$$\exp \left\{ -\eta \left(x \frac{\partial}{\partial y} + y \frac{\partial}{\partial x} \right) \right\}. \quad (24)$$

Next, we can consider the exponential form

$$\exp \left\{ (\tanh \eta) a^\dagger b^\dagger \right\}, \quad (25)$$

which can be expanded as

$$\sum_n \frac{1}{n!} (\tanh \eta)^n (a^\dagger b^\dagger)^n. \quad (26)$$

If this operator is applied to the ground state of Eq.(11), the result is

$$\sum_n (\tanh \eta)^n \chi_n(x) \chi_n(y). \quad (27)$$

This form is not normalized, while the series of Eq.(1) is. What is the origin of this difference?

There is a similar problem with the one-photon coherent state [29, 30]. There, the series comes from the expansion of the exponential form

$$\exp \{ \alpha a^\dagger \}, \quad (28)$$

which can be expanded to

$$\sum_n \frac{1}{n!} \alpha^n (a^\dagger)^n. \quad (29)$$

However, this operator is not unitary. In order to make this series unitary, we consider the exponential form

$$\exp (\alpha a^\dagger - \alpha^* a) \quad (30)$$

which is unitary. This expression can then be written as

$$e^{-\alpha\alpha^*/2} [\exp (\alpha a^\dagger)] [\exp (\alpha^* a)], \quad (31)$$

according to the Baker-Campbell-Hausdorff (BCH) relation [31, 32]. If this is applied to the ground state, the last bracket can be dropped, and the result is

$$e^{-\alpha\alpha^*/2} \exp [\alpha a^\dagger], \quad (32)$$

which is the unitary operator with the normalization constant

$$e^{-\alpha\alpha^*/2}.$$

Likewise, we can conclude that the series of Eq.(27) is different from that of Eq.(1) due to the difference between the unitary operator of Eq.(23) and the non-unitary operator of Eq.(25). It may be possible to derive the normalization factor using the BCH formula, but it seems to be intractable at this time. The best way to resolve this problem is to present the exact calculation of the unitary operator leading to the normalized series of Eq.(11). We shall return to this problem in Sec. 5, where squeezed excited states are studied.

2.2 Sheared Gaussian Function

In addition, there is a transformation called “shear,” where only one of the two coordinates is translated as shown in Fig. 2. This transformation takes the form

$$\begin{pmatrix} x' \\ y' \end{pmatrix} = \begin{pmatrix} 1 & 2\alpha \\ 0 & 1 \end{pmatrix} \begin{pmatrix} x \\ y \end{pmatrix}, \quad (33)$$

which leads to

$$\begin{pmatrix} x' \\ y' \end{pmatrix} = \begin{pmatrix} x + 2\alpha y \\ y \end{pmatrix}. \quad (34)$$

This shear is one of the basic transformations in engineering sciences. In physics, this transformation plays the key role in understanding the internal space-time symmetry of

massless particles [33, 34, 35]. This matrix plays the pivotal role during the transition from the oscillator mode to the damping mode in classical damped harmonic oscillators [36, 37].

Under this transformation, the Gaussian form becomes

$$\psi_{shr}(x, y) = \frac{1}{\sqrt{\pi}} \exp \left\{ -\frac{1}{2} [(x - 2\alpha y)^2 + y^2] \right\}. \quad (35)$$

It is possible to expand this into a series of the form of Eq.(15) [38].

The transformation applicable to the Gaussian form of Eq.(11) is

$$\exp \left(-2\alpha y \frac{\partial}{\partial x} \right), \quad (36)$$

and the generator is

$$-iy \frac{\partial}{\partial x}. \quad (37)$$

It is of interest to see where this generator stands among the ten generators of Dirac.

However, the most pressing problem is whether the sheared Gaussian form can be regarded as a rotated squeezed state. The basic mathematical issue is that the shear matrix of Eq.(33) is triangular and cannot be diagonalized. Therefore, it cannot be a squeezed state. Yet, the Gaussian form of Eq.(35) appears to be a rotated squeezed state, while not along the normal coordinates. We shall look at this problem in detail in Sec. 6.

3 Dirac's Entangled Oscillators

Paul A. M. Dirac devoted much of his life-long efforts to the task of making quantum mechanics compatible with special relativity. Harmonic oscillators serve as an instrument for illustrating quantum mechanics, while special relativity is the physics of the Lorentz group. Thus, Dirac attempted to construct a representation of the Lorentz group using harmonic oscillator wave functions [17, 21].

In his 1963 paper [21], Dirac started from the two-dimensional oscillator whose wave function takes the Gaussian form given in Eq.(11). He then considered unitary transformations applicable to this ground-state wave function. He noted that they can be generated by the following ten Hermitian operators

$$\begin{aligned} L_1 &= \frac{1}{2} (a^\dagger b + b^\dagger a), & L_2 &= \frac{1}{2i} (a^\dagger b - b^\dagger a), \\ L_3 &= \frac{1}{2} (a^\dagger a - b^\dagger b), & S_3 &= \frac{1}{2} (a^\dagger a + b b^\dagger), \\ K_1 &= -\frac{1}{4} (a^\dagger a^\dagger + a a - b^\dagger b^\dagger - b b), \\ K_2 &= \frac{i}{4} (a^\dagger a^\dagger - a a + b^\dagger b^\dagger - b b), \end{aligned}$$

$$\begin{aligned}
K_3 &= \frac{1}{2} (a^\dagger b^\dagger + ab), \\
Q_1 &= -\frac{i}{4} (a^\dagger a^\dagger - aa - b^\dagger b^\dagger + bb), \\
Q_2 &= -\frac{1}{4} (a^\dagger a^\dagger + aa + b^\dagger b^\dagger + bb), \\
Q_3 &= \frac{i}{2} (a^\dagger b^\dagger - ab).
\end{aligned} \tag{38}$$

He then noted that these operators satisfy the following set of commutation relations.

$$\begin{aligned}
[L_i, L_j] &= i\epsilon_{ijk}L_k, & [L_i, K_j] &= i\epsilon_{ijk}K_k, & [L_i, Q_j] &= i\epsilon_{ijk}Q_k, \\
[K_i, K_j] &= [Q_i, Q_j] = -i\epsilon_{ijk}L_k, & [L_i, S_3] &= 0, \\
[K_i, Q_j] &= -i\delta_{ij}S_3, & [K_i, S_3] &= -iQ_i, & [Q_i, S_3] &= iK_i.
\end{aligned} \tag{39}$$

Dirac then determined that these commutation relations constitute the Lie algebra for the $O(3, 2)$ de Sitter group with ten generators. This de Sitter group is the Lorentz group applicable to three space coordinates and two time coordinates. Let us use the notation (x, y, z, t, s) , with (x, y, z) as space coordinates and (t, s) as two time coordinates. Then the rotation around the z axis is generated by

$$L_3 = \begin{pmatrix} 0 & -i & 0 & 0 & 0 \\ i & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 \end{pmatrix}. \tag{40}$$

The generators L_1 and L_2 can be also be constructed. The K_3 and Q_3 generators will take the form

$$K_3 = \begin{pmatrix} 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & i & 0 \\ 0 & 0 & i & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 \end{pmatrix}, \quad Q_3 = \begin{pmatrix} 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & i \\ 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & i & 0 & 0 \end{pmatrix}. \tag{41}$$

From these two matrices, the generators K_1, K_2, Q_1, Q_2 can be constructed. The generator S_3 can be written as

$$S_3 = \begin{pmatrix} 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & -i \\ 0 & 0 & 0 & i & 0 \end{pmatrix}. \tag{42}$$

The last five-by-five matrix generates rotations in the two-dimensional space of (t, s) . If we introduce these two time variables, the $O(3, 2)$ group leads to two coupled Lorentz

groups. The particle mass is invariant under Lorentz transformations. Thus, one Lorentz group cannot change the particle mass. However, with two coupled Lorentz groups we can describe the world with variable masses such as the neutrino oscillations.

In Sec. 2, we used the operators Q_3 and K_3 as the generators for the squeezed Gaussian function. For the unitary transformation of Eq.(23), we used

$$\exp(-i\eta Q_3). \quad (43)$$

However, the exponential form of Eq.(25) can be written as

$$\exp\{-i(\tanh \eta)(Q_3 + iK_3)\}, \quad (44)$$

which is not unitary, as was seen before.

From the space-time point of view, both K_3 and Q_3 generate Lorentz boosts along the z direction, with the time variables t and s respectively. The fact that the squeeze and Lorentz transformations share the same mathematical formula is well known. However, the non-unitary operator iK_3 does not seem to have a space-time interpretation.

As for the sheared state, the generator can be written as

$$Q_3 - L_2, \quad (45)$$

leading to the expression given in Eq.(37). This is a Hermitian operator leading to the unitary transformation of Eq.(36).

4 Entangled Oscillators in the Phase-Space Picture

Also in his 1963 paper, Dirac states that the Lie algebra of Eq. (39) can serve as the four-dimensional symplectic group $Sp(4)$. This group allows us to study squeezed or entangled states in terms of the four-dimensional phase space consisting of two position and two momentum variables [15, 40, 41].

In order to study the $Sp(4)$ contents of the coupled oscillator system, let us introduce the Wigner function defined as [42]

$$W(x, y; p, q) = \left(\frac{1}{\pi}\right)^2 \int \exp\{-2i(px' + qy')\} \\ \times \psi^*(x + x', y + y')\psi(x - x', y - y')dx'dy'. \quad (46)$$

If the wave function $\psi(x, y)$ is the Gaussian form of Eq.(11), the Wigner function becomes

$$W(x, y : p, q) = \left(\frac{1}{\pi}\right)^2 \exp\left\{-\left(x^2 + p^2 + y^2 + q^2\right)\right\}. \quad (47)$$

The Wigner function is defined over the four-dimensional phase space of (x, p, y, q) just as in the case of classical mechanics. The unitary transformations generated by the operators of Eq.(38) are translated into Wigner transformations [40, 39, 41]. As in




Figure 4: Transformations generated by Q_3 and K_3 . As the parameter η becomes larger, both the space and momentum distribution becomes larger.

the case of Dirac's oscillators, there are ten corresponding generators applicable to the Wigner function. They are

$$\begin{aligned}
L_1 &= +\frac{i}{2} \left\{ \left(x \frac{\partial}{\partial q} - q \frac{\partial}{\partial x} \right) + \left(y \frac{\partial}{\partial p} - p \frac{\partial}{\partial y} \right) \right\}, \\
L_2 &= -\frac{i}{2} \left\{ \left(x \frac{\partial}{\partial y} - y \frac{\partial}{\partial x} \right) + \left(p \frac{\partial}{\partial q} - q \frac{\partial}{\partial p} \right) \right\}, \\
L_3 &= +\frac{i}{2} \left\{ \left(x \frac{\partial}{\partial p} - p \frac{\partial}{\partial x} \right) - \left(y \frac{\partial}{\partial q} - q \frac{\partial}{\partial y} \right) \right\}, \\
S_3 &= -\frac{i}{2} \left\{ \left(x \frac{\partial}{\partial p} - p \frac{\partial}{\partial x} \right) + \left(y \frac{\partial}{\partial q} - q \frac{\partial}{\partial y} \right) \right\}, \tag{48}
\end{aligned}$$

and

$$\begin{aligned}
K_1 &= -\frac{i}{2} \left\{ \left(x \frac{\partial}{\partial p} + p \frac{\partial}{\partial x} \right) - \left(y \frac{\partial}{\partial q} + q \frac{\partial}{\partial y} \right) \right\}, \\
K_2 &= -\frac{i}{2} \left\{ \left(x \frac{\partial}{\partial x} + y \frac{\partial}{\partial y} \right) - \left(p \frac{\partial}{\partial p} + q \frac{\partial}{\partial q} \right) \right\}, \\
K_3 &= +\frac{i}{2} \left\{ \left(x \frac{\partial}{\partial q} + q \frac{\partial}{\partial x} \right) + \left(y \frac{\partial}{\partial p} + p \frac{\partial}{\partial y} \right) \right\}, \\
Q_1 &= +\frac{i}{2} \left\{ \left(x \frac{\partial}{\partial x} + q \frac{\partial}{\partial q} \right) - \left(y \frac{\partial}{\partial y} + p \frac{\partial}{\partial p} \right) \right\},
\end{aligned}$$

$$\begin{aligned}
Q_2 &= -\frac{i}{2} \left\{ \left(x \frac{\partial}{\partial p} + p \frac{\partial}{\partial x} \right) + \left(y \frac{\partial}{\partial q} + q \frac{\partial}{\partial y} \right) \right\}, \\
Q_3 &= -\frac{i}{2} \left\{ \left(y \frac{\partial}{\partial x} + x \frac{\partial}{\partial y} \right) - \left(q \frac{\partial}{\partial p} + p \frac{\partial}{\partial q} \right) \right\}.
\end{aligned} \tag{49}$$

These generators also satisfy the Lie algebra given in Eq.(38) and Eq.(39). Transformations generated by these generators have been discussed in the literature [15, 39, 41].

As in the case of Sec. 3, we are interested in the generators Q_3 and K_3 . The transformation generated by Q_3 takes the form

$$\left[\exp \left\{ \eta \left(x \frac{\partial}{\partial y} + y \frac{\partial}{\partial x} \right) \right\} \right] \left[\exp \left\{ -\eta \left(p \frac{\partial}{\partial q} + q \frac{\partial}{\partial p} \right) \right\} \right]. \tag{50}$$

This exponential form squeezes the Wigner function of Eq.(47) in the $x y$ space as well as in their corresponding momentum space. However, in the momentum space, the squeeze is in the opposite direction as illustrated in Fig. 4. This is what we expect from canonical transformation in classical mechanics. Indeed, this corresponds to the unitary transformation which played the major role in Sec. 2.

Even though shown insignificant in Sec. 2, K_3 had a definite physical interpretation in Sec. 3. The transformation generated by K_3 takes the form

$$\left[\exp \left\{ \eta \left(x \frac{\partial}{\partial q} + q \frac{\partial}{\partial x} \right) \right\} \right] \left[\exp \left\{ \eta \left(y \frac{\partial}{\partial p} + p \frac{\partial}{\partial y} \right) \right\} \right]. \tag{51}$$

This performs the squeeze in the $x q$ and $y p$ spaces. In this case, the squeezes have the same sign, and the rate of increase is the same in all directions. We can thus have the same picture of squeeze for both $x y$ and $p q$ spaces as illustrated in Fig. 4. This parallel transformation corresponds to the Lorentz squeeze [20, 25].

As for the sheared state, the combination

$$Q_3 - L_2 = -i \left(y \frac{\partial}{\partial x} + q \frac{\partial}{\partial p} \right), \tag{52}$$

generates the same shear in the $p q$ space.

5 Entangled Excited States

In Sec. 2, we discussed the entangled ground state, and noted that the entangled state of Eq.(1) is a series expansion of the squeezed Gaussian function. In this section, we are interested in what happens when we squeeze an excited oscillator state starting from

$$\chi_n(x) \chi_m(y). \tag{53}$$

In order to entangle this state, we should replace x and y respectively by x' and y' given in Eq.(17).

The question is how the oscillator wave function is squeezed after this operation. Let us note first that the wave function of Eq.(53) satisfies the equation

$$\frac{1}{2} \left\{ \left(x^2 - \frac{\partial^2}{\partial x^2} \right) - \left(y^2 - \frac{\partial^2}{\partial y^2} \right) \right\} \chi_n(x) \chi_m(y) = (n - m) \chi_n(x) \chi_m(y). \quad (54)$$

This equation is invariant under the squeeze transformation of Eq.(17), and thus the eigenvalue $(n - m)$ remains invariant. Unlike the usual two-oscillator system, the x component and the y component have opposite signs. This is the reason why the overall equation is squeeze-invariant [25, 3, 43].

We then have to write this squeezed oscillator in the series form of Eq.(15). The most interesting case is of course for $m = n = 0$, which leads to the Gaussian entangled state given in Eq.(16). Another interesting case is for $m = 0$ while n is allowed to take all integer values. This single-excitation system has applications in the covariant oscillator formalism where no time-like excitations are allowed. The Gaussian entangled state is a special case of this single-excited oscillator system.

The most general case is for nonzero integers for both n and m . The calculation for this case is available in the literature [20, 45]. Seeing no immediate physical applications of this case, we shall not reproduce this calculation in this section.

For the single-excitation system, we write the starting wave function as

$$\chi_n(x) \chi_0(y) = \left[\frac{1}{\pi 2^n n!} \right]^{1/2} H_n(x) \exp \left\{ - \left(\frac{x^2 + y^2}{2} \right) \right\}. \quad (55)$$

There are no excitations along the y coordinate. In order to squeeze this function, our plan is to replace x and y by x' and y' respectively and write $\chi_n(x') \chi_0(y')$ as a series in the form

$$\chi_n(x') \chi_0(y') = \sum_{k',k} A_{k',k}(n) \chi_{k'}(x) \chi_k(y). \quad (56)$$

Since $k' - k = n$ or $k' = n + k$, according to the eigenvalue of the differential equation given in Eq.(54), we write this series as

$$\chi_n(x') \chi_0(y') = \sum_{k',k} A_k(n) \chi_{(k+n)}(x) \chi_k(y), \quad (57)$$

with

$$\sum_k |A_k(n)|^2 = 1. \quad (58)$$

This coefficient is

$$A_k(n) = \int \chi_{k+n}(x) \chi_k(y) \chi_n(x') \chi_0(y') dx dy. \quad (59)$$

This calculation was given in the literature in a fragmentary way in connection with a Lorentz-covariant description of extended particles starting from Ruiz's 1974 paper [44], subsequently by Kim *et al.* in 1979 [26] and by Rotbart in 1981 [45]. In view of the recent developments of physics, it seems necessary to give one coherent calculation of the coefficient of Eq.(59).

We are now interested in the squeezed oscillator function

$$A_k(n) = \left[\frac{1}{\pi^2 2^n n! (k+n)^2 (n+k)! k^2 k!} \right]^{1/2} \times \int H_{n+k}(x) H_k(y) H_n(x') \exp \left\{ - \left(\frac{x^2 + y^2 + x'^2 + y'^2}{2} \right) \right\} dx dy. \quad (60)$$

As was noted by Ruiz [44], the key to the evaluation of this integral is to introduce the generating function for the Hermite polynomials [46, 47]:

$$G(r, z) = \exp(-r^2 + 2rz) = \sum_m \frac{r^m}{m!} H_m(z), \quad (61)$$

and evaluate the integral

$$I = \int G(r, x) G(s, y) G(r', x') \exp \left\{ - \left(\frac{x^2 + y^2 + x'^2 + y'^2}{2} \right) \right\} dx dy. \quad (62)$$

The integrand becomes one exponential function, and its exponent is quadratic in x and y . This quadratic form can be diagonalized, and the integral can be evaluated [20, 26]. The result is

$$I = \left[\frac{\pi}{\cosh \eta} \right] \exp(2rs \tanh \eta) \exp \left(\frac{2rr'}{\cosh \eta} \right). \quad (63)$$

We can now expand this expression and choose the coefficients of r^{n+k} , s^k , r'^n for $H_{(n+k)}(x)$, $H_n(y)$, and $H_n(z')$ respectively. The result is

$$A_{n,k} = \left(\frac{1}{\cosh \eta} \right)^{(n+1)} \left[\frac{(n+k)!}{n!k!} \right]^{1/2} (\tanh \eta)^k. \quad (64)$$

Thus, the series becomes

$$\chi_n(x') \chi_0(y') = \left(\frac{1}{\cosh \eta} \right)^{(n+1)} \sum_k \left[\frac{(n+k)!}{n!k!} \right]^{1/2} (\tanh \eta)^k \chi_{k+n}(x) \chi_k(y). \quad (65)$$

If $n = 0$, it is the squeezed ground state, and this expression becomes the entangled state of Eq.(16).

6 E(2)-sheared States

Let us next consider the effect of shear on the Gaussian form. From Fig. 5 and Fig. 3, it is clear that the sheared state is a rotated squeezed state.

In order to understand this transformation, let us note that the squeeze and rotation are generated by the two-by-two matrices

$$K = \begin{pmatrix} 0 & i \\ i & 0 \end{pmatrix}, \quad J = \begin{pmatrix} 0 & -i \\ i & 0 \end{pmatrix}, \quad (66)$$




Figure 5: Shear transformation of the Gaussian form given in Eq.(11).

which generate the squeeze and rotation matrices of the form

$$\begin{aligned}\exp(-i\eta K) &= \begin{pmatrix} \cosh \eta & \sinh \eta \\ \sinh \eta & \cosh \eta \end{pmatrix}, \\ \exp(-i\theta J) &= \begin{pmatrix} \cos \theta & -\sin \theta \\ \sin \theta & \cos \theta \end{pmatrix},\end{aligned}\quad (67)$$

respectively. We can then consider

$$S = K - J = \begin{pmatrix} 0 & 2i \\ 0 & 0 \end{pmatrix}.\quad (68)$$

This matrix has the property that $S^2 = 0$. Thus the transformation matrix becomes

$$\exp(-i\alpha S) = \begin{pmatrix} 1 & 2\alpha \\ 0 & 1 \end{pmatrix}.\quad (69)$$

Since $S^2 = 0$, the Taylor expansion truncates, and the transformation matrix becomes the triangular matrix of Eq.(34), leading to the transformation

$$\begin{pmatrix} x \\ y \end{pmatrix} \rightarrow \begin{pmatrix} x + 2\alpha y \\ y \end{pmatrix}.\quad (70)$$

The shear generator S of Eq.(68) indicates that the infinitesimal transformation is a rotation followed by a squeeze. Since both rotation and squeeze are area-preserving transformations, the shear should also be an area-preserving transformations.

In view of Fig. 5, we should ask whether the triangular matrix of Eq.(69) can be obtained from one squeeze matrix followed by one rotation matrix. This is not possible mathematically. It can however, be written as a squeezed rotation matrix of the form

$$\begin{pmatrix} e^{\lambda/2} & 0 \\ 0 & e^{-\lambda/2} \end{pmatrix} \begin{pmatrix} \cos \omega & \sin \omega \\ -\sin \omega & \cos \omega \end{pmatrix} \begin{pmatrix} e^{-\lambda/2} & 0 \\ 0 & e^{\lambda/2} \end{pmatrix},\quad (71)$$

resulting in

$$\begin{pmatrix} \cos \omega & e^\lambda \sin \omega \\ -e^{-\lambda} \sin \omega & \cos \omega \end{pmatrix}. \quad (72)$$

If we let

$$(\sin \omega) = 2\alpha e^{-\lambda} \quad (73)$$

Then

$$\begin{pmatrix} \cos \omega & 2\alpha \\ -2\alpha e^{-2\lambda} & \cos \omega \end{pmatrix}. \quad (74)$$

If λ becomes infinite, the angle ω becomes zero, and this matrix becomes the triangular matrix of Eq.(69). This is a singular process where the parameter λ goes to infinity.

If this transformation is applied to the Gaussian form of Eq.(11), it becomes

$$\psi(x, y) = \frac{1}{\sqrt{\pi}} \exp \left\{ -\frac{1}{2} [(x - 2\alpha y)^2 + y^2] \right\}. \quad (75)$$

The question is whether the exponential portion of this expression can be written as

$$\exp \left\{ -\frac{1}{2} [e^{-2\eta}(x \cos \theta + y \sin \theta)^2 + e^{2\eta}(x \sin \theta - y \cos \theta)^2] \right\}. \quad (76)$$

The answer is Yes. This is possible if

$$\begin{aligned} \tan(2\theta) &= \frac{1}{\alpha}, \\ e^{2\eta} &= 1 + 2\alpha^2 + 2\alpha\sqrt{\alpha^2 + 1}, \\ e^{-2\eta} &= 1 + 2\alpha^2 - 2\alpha\sqrt{\alpha^2 + 1}. \end{aligned} \quad (77)$$

In Eq.(74), we needed a limiting case of λ becoming infinite. This is necessarily a singular transformation. On the other hand, the derivation of the Gaussian form of Eq.(75) appears to be analytic. How is it possible? In order to achieve the transformation from the Gaussian form of Eq.(11) to Eq.(75), we need the linear transformation

$$\begin{pmatrix} \cos \theta & -\sin \theta \\ \sin \theta & \cos \theta \end{pmatrix} \begin{pmatrix} e^\eta & 0 \\ 0 & e^{-\eta} \end{pmatrix}. \quad (78)$$

If the initial form is invariant under rotations as in the case of the Gaussian function of Eq.(11), we can add another rotation matrix on the right hand side. We choose that rotation matrix to be

$$\begin{pmatrix} \cos(\theta - \pi/2) & -\sin(\theta - \pi/2) \\ \sin(\theta - \pi/2) & \cos(\theta - \pi/2) \end{pmatrix}, \quad (79)$$

write the three matrices as

$$\begin{pmatrix} \cos \theta' & -\sin \theta' \\ \sin \theta' & \cos \theta' \end{pmatrix} \begin{pmatrix} \cosh \eta & \sinh \eta \\ \sinh \eta & \cosh \eta \end{pmatrix} \begin{pmatrix} \cos \theta' & -\sin \theta' \\ \sin \theta' & \cos \theta' \end{pmatrix}, \quad (80)$$

with

$$\theta' = \theta - \frac{\pi}{4}.$$

The multiplication of these three matrices leads to

$$\begin{pmatrix} (\cosh \eta) \sin(2\theta) & \sinh \eta + (\cosh \eta) \cos(2\theta) \\ \sinh \eta - (\cosh \eta) \cos(2\theta) & (\cosh \eta) \sin(2\theta) \end{pmatrix}. \quad (81)$$

The lower-left element can become zero when $\sinh \eta = \cosh(\eta) \cos(2\theta)$ and consequently this matrix becomes

$$\begin{pmatrix} 1 & 2 \sinh \eta \\ 0 & 1 \end{pmatrix}. \quad (82)$$

Furthermore, this matrix can be written in the form of a squeezed rotation matrix given in Eq.(72), with

$$\begin{aligned} \cos \omega &= (\cosh \eta) \sin(2\theta), \\ e^{-2\lambda} &= \frac{\cos(2\theta) - \tanh \eta}{\cos(2\theta) + \tanh \eta}. \end{aligned} \quad (83)$$

The matrices of the form of Eq.(72) and Eq.(81) are known as the Wigner and Bargmann decompositions respectively [33, 48, 49, 50, 36].

7 Feynman's Rest of the Universe

We need the concept of entanglement in quantum systems of two variables. The issue is how the measurement of one variable affects the other variable. The simplest case is what happens to the first variable while no measurements are taken on the second variable. This problem has a long history since von Neumann introduced the concept of density matrix in 1932 [51]. While there are many books and review articles on this subject, Feynman stated this problem in his own colorful way. In his book on statistical mechanics [22], Feynman makes the following statement about the density matrix.

When we solve a quantum-mechanical problem, what we really do is divide the universe into two parts - the system in which we are interested and the rest of the universe. We then usually act as if the system in which we are interested comprised the entire universe. To motivate the use of density matrices, let us see what happens when we include the part of the universe outside the system.

Indeed, Yurke and Potasek [11] and also Ekert and Knight [12] studied this problem in the two-mode squeezed state using the entanglement formula given in Eq.(16). Later in 1999, Han *et al.* studied this problem with two coupled oscillators where one oscillator is observed while the other is not and thus is in the rest of the universe as defined by Feynman [23].

Somewhat earlier in 1990 [27], Kim and Wigner observed that there is a time separation wherever there is a space separation in the Lorentz-covariant world. The Bohr radius is a space separation. If the system is Lorentz-boosted, the time-separation becomes entangled with the space separation. But, in the present form of quantum mechanics, this time-separation variable is not measured and not understood.

This variable was mentioned in the paper of Feynman *et al.* in 1971 [43], but the authors say they would drop this variable because they do not know what to do with




Figure 6: Feynman's rest of the universe. As the Gaussian function is squeezed, the x and y variables become entangled. If the y variable is not measured, it affects the quantum mechanics of the x variable.

it. While what Feynman *et al.* did was not quite respectable from the scientific point of view, they made a contribution by pointing out the existence of the problem. In 1990, Kim and Wigner [27] noted that the time-separation variable belongs to Feynman's rest of the universe and studied its consequences in the observable world.

In this section, we first reproduce the work of Kim and Wigner using the x and y variables and then study its consequences. Let us introduce the notation $\psi_\eta^n(x, y)$ for the squeezed oscillator wave function given in Eq.(65):

$$\psi_\eta^n(x, y) = \chi_n(x')\chi_0(y'), \quad (84)$$

with no excitations along the y direction. For $\eta = 0$, this expression becomes $\chi_n(x)\chi_0(y)$.

From this wave function, we can construct the pure-state density matrix as

$$\rho_\eta^n(x, y; r, s) = \psi_\eta^n(x, y)\psi_\eta^n(r, s), \quad (85)$$

which satisfies the condition $\rho^2 = \rho$, which means

$$\rho_\eta^n(x, y; r, s) = \int \rho_\eta^n(x, y; u, v)\rho_\eta^n(u, v; r, s)dudv. \quad (86)$$

As illustrated in Fig. 6, it is not possible make measurements on the variable y . We thus have to take the trace of this density matrix along the y axis, resulting in

$$\begin{aligned} \rho_\eta^n(x, r) &= \int \psi_\eta^n(x, y)\psi_\eta^n(r, y)dy \\ &= \left(\frac{1}{\cosh \eta}\right)^{2(n+1)} \sum_k \frac{(n+k)!}{n!k!} (\tanh \eta)^{2k} \chi_{n+k}(x)\chi_{k+n}(r). \end{aligned} \quad (87)$$

The trace of this density matrix is one, but the trace of ρ^2 is

$$\begin{aligned} \text{Tr}(\rho^2) &= \int \rho_\eta^n(x, r) \rho_\eta^n(r, x) dr dx \\ &= \left(\frac{1}{\cosh \eta} \right)^{4(n+1)} \sum_k \left[\frac{(n+k)!}{n!k!} \right]^2 (\tanh \eta)^{4k}, \end{aligned} \quad (88)$$

which is less than one. This is due to the fact that we are not observing the y variable. Our knowledge is less than complete.

The standard way to measure this incompleteness is to calculate the entropy defined as [51, 52, 53]

$$S = -\text{Tr}(\rho(x, r) \ln[\rho(x, r)]), \quad (89)$$

which leads to

$$\begin{aligned} S &= 2(n+1)[(\cosh \eta)^2 \ln(\cosh \eta) - (\sinh \eta)^2 \ln(\sinh \eta)] \\ &\quad - \left(\frac{1}{\cosh \eta} \right)^{2(n+1)} \sum_k \frac{(n+k)!}{n!k!} \ln \left[\frac{(n+k)!}{n!k!} \right] (\tanh \eta)^{2k}. \end{aligned} \quad (90)$$

Let us go back to the wave function given in Eq.(84). As is illustrated in Fig. 6, its localization property is dictated by its Gaussian factor which corresponds to the ground-state wave function. For this reason, we expect that much of the behavior of the density matrix or the entropy for the n^{th} excited state will be the same as that for the ground state with $n = 0$. For this state, the density matrix is

$$\rho_\eta(x, r) = \left(\frac{1}{\pi \cosh(2\eta)} \right)^{1/2} \exp \left\{ -\frac{1}{4} \left[\frac{(x+r)^2}{\cosh(2\eta)} + (x-r)^2 \cosh(2\eta) \right] \right\}, \quad (91)$$

and the entropy is

$$S_\eta = 2 \left[(\cosh \eta)^2 \ln(\cosh \eta) - (\sinh \eta)^2 \ln(\sinh \eta) \right]. \quad (92)$$

The density distribution $\rho_\eta(x, x)$ becomes

$$\rho_\eta(x, x) = \left(\frac{1}{\pi \cosh(2\eta)} \right)^{1/2} \exp \left(\frac{-x^2}{\cosh(2\eta)} \right). \quad (93)$$

The width of the distribution becomes $\sqrt{\cosh(2\eta)}$, and the distribution becomes wide-spread as η becomes larger. Likewise, the momentum distribution becomes wide-spread as can be seen in Fig. 4. This simultaneous increase in the momentum and position distribution widths is due to our inability to measure the y variable hidden in Feynman's rest of the universe [22].

In their paper of 1990 [27], Kim and Wigner used the x and y variables as the longitudinal and time-like variables respectively in the Lorentz-covariant world. In the quantum world, it is a widely accepted view that there are no time-like excitations. Thus, it is fully justified to restrict the y component to its ground state as we did in Sec. 5.

8 Space-time Entanglement

The series given in Eq.(1) plays the central role in the concept of the Gaussian or continuous-variable entanglement, where the measurement on one variable affects the quantum mechanics of the other variable. If one of the variables is not observed, it belongs to Feynman’s rest of the universe.

The series of the form of Eq.(1) was developed earlier for studying harmonic oscillators in moving frames [24, 20, 25, 26, 27, 28]. Here z and t are the space-like and time-like separations between the two constituent particles bound together by a harmonic oscillator potential. There are excitations along the longitudinal direction. However, no excitations are allowed along the time-like direction. Dirac described this as “c-number” time-energy uncertainty relation [16]. Dirac in 1927 was talking about the system without special relativity. In 1945 [17], Dirac attempted to construct space-time wave functions using harmonic oscillators. In 1949 [18], Dirac introduced his light-cone coordinate system for Lorentz boosts, telling that the boost is a squeeze transformation. It is now possible to combine Dirac’s three observations to construct the Lorentz covariant picture of quantum bound states, as illustrated in Fig. 7.

If the system is at rest, we use the wave function

$$\psi_0^n(z, t) = \chi_n(z)\chi_0(t), \quad (94)$$

which allows excitations along the z axis, but no excitations along the t axis, according to Dirac’s c-number time-energy uncertainty relation.

If the system is boosted, the z and t variables are replaced by z' and t' where

$$z' = (\cosh \eta)z - (\sinh \eta)t, \quad \text{and} \quad t' = -(\sinh \eta)z + (\cosh \eta)t. \quad (95)$$

This is a squeeze transformation as in the case of Eq.(17). In terms of these space-time variables, the wave function of Eq.(84), can be written as

$$\psi_\eta^n(z, t) = \chi_n(z')\chi_0(t'), \quad (96)$$

and the series of Eq.(65) then becomes

$$\psi_\eta^n(z, t) = \left(\frac{1}{\cosh \eta} \right)^{(n+1)} \sum_k \left[\frac{(n+k)!}{n!k!} \right]^{1/2} (\tanh \eta)^k \chi_{k+n}(z)\chi_k(t). \quad (97)$$

Since the Lorentz-covariant oscillator formalism shares the same set of formulas with the Gaussian entangled states, it is possible to explain some aspects of space-time physics using the concepts and terminologies developed in quantum optics as illustrated in Fig. 1.

The time-separation variable is a case in point. The Bohr radius is a well-defined spatial separation between the proton and electron in the hydrogen atom. However, if the atom is boosted, this radius picks up its time-like separation. This time-separation variable does not exist in the Schrödinger picture of quantum mechanics. However, this variable plays the pivotal role in the covariant harmonic oscillator formalism. It is gratifying to note that this “hidden or forgotten” variable plays its role in the real




Figure 7: Dirac's form of Lorentz-covariant quantum mechanics. In addition to Heisenberg's uncertainty relation which allows excitations along the spatial direction, there is the "c-number" time-energy uncertainty without excitations. This form of quantum mechanics can be combined with Dirac's light-cone picture of Lorentz boost, resulting in the Lorentz-covariant picture of quantum mechanics. The elliptic squeeze shown in this figure can be called the space-time entanglement.

world while being entangled with the observable longitudinal variable. With this point in mind, let us study some of the consequences of this space-time entanglement.

First of all, does the wave function of Eq.(96) carry a probability interpretation in the Lorentz-covariant world? Since $dzdt = dz'dt'$, the normalization

$$\int |\psi_\eta^n(z, t)|^2 dt dz = 1. \quad (98)$$

This is a Lorentz-invariant normalization. If the system is at rest, the z and t variables are completely dis-entangled, and the spatial component of the wave function satisfies the Schrödinger equation without the time-separation variable.

However, in the Lorentz-covariant world, we have to consider the inner product

$$\left(\psi_\eta^n(z, t), \psi_{\eta'}^m(z, t)\right) = \int \left[\psi_\eta^n(z, t)\right]^* \psi_{\eta'}^m(z, t) dz dt. \quad (99)$$

The evaluation of this integral was carried out by Michael Ruiz in 1974 [44], and the result was

$$\left(\frac{1}{|\cosh(\eta - \eta')|}\right)^{n+1} \delta_{nm}. \quad (100)$$

In order to see the physical implications of this result, let us assume that one of the oscillators is at rest with $\eta' = 0$ and the other is moving with the velocity $\beta = \tanh(\eta)$. Then the result is

$$\left(\psi_\eta^n(z, t), \psi_0^m(z, t)\right) = \left(\sqrt{1 - \beta^2}\right)^{n+1} \delta_{nm}. \quad (101)$$

Indeed, the wave functions are orthonormal if they are in the same Lorentz frame. If one of them is boosted, the inner product shows the effect of Lorentz contraction. We are familiar with the contraction $\sqrt{1 - \beta^2}$ for the rigid rod. The ground state of the oscillator wave function is contracted like a rigid rod.

The probability density $|\psi_\eta^0(z)|^2$ is for the oscillator in the ground state, and it has one hump. For the n^{th} excited state, there are $(n + 1)$ humps. If each hump is contracted like $\sqrt{1 - \beta^2}$, the net contraction factor is $\left(\sqrt{1 - \beta^2}\right)^{n+1}$ for the n^{th} excited state. This result is illustrated in Fig. 8.

With this understanding, let us go back to the entanglement problem. The ground state wave function takes the Gaussian form given in Eq.(11)

$$\psi_0(z, t) = \frac{1}{\sqrt{\pi}} \exp\left\{-\frac{1}{2}(z^2 + t^2)\right\}, \quad (102)$$

where the x and y variables are replaced by z and t respectively. If Lorentz-boosted, this Gaussian function becomes squeezed to [24, 20, 25]

$$\psi_\eta^0(z, t) = \frac{1}{\sqrt{\pi}} \exp\left\{-\frac{1}{4}\left[e^{-2\eta}(z + t)^2 + e^{2\eta}(z - t)^2\right]\right\}, \quad (103)$$

leading to the series

$$\frac{1}{\cosh \eta} \sum_k (\tanh \eta)^k \chi_k(z) \chi_k(t). \quad (104)$$




Figure 8: Orthogonality relations for two covariant oscillator wave functions. The orthogonality relation is preserved for different frames. However, they show the Lorentz contraction effect for two different frames.

According to this formula, the z and t variables are entangled in the same way as the x and y variables are entangled.

Here the z and t variables are space and time separations between two particles bound together by the oscillator force. The concept of the space separation is well defined, as in the case of the Bohr radius. On the other hand, the time separation is still hidden or forgotten in the present form of quantum mechanics. In the Lorentz-covariant world, this variable affects what we observe in the real world by entangling itself with the longitudinal spatial separation.

In Chapter 16 of their book [9], Walls and Milburn wrote down the series of Eq.(1) and discussed what would happen when the η parameter becomes infinitely large. We note that the series given in Eq.(104) shares the same expression as the form given by Walls and Milburn, as well as other papers dealing with the Gaussian entanglement. As in the case of Wall and Milburn, we are interested in what happens when η becomes very large.

As we emphasized throughout the present paper, it is possible to study the entanglement series using the squeezed Gaussian function given in Eq.(103). It is then possible to study this problem using the ellipse. Indeed, we can carry out the mathematics of entanglement using the ellipse shown Fig. 9. This figure is the same as that of Fig. 6, but it tells about the entanglement of the space and time separations, instead of the x and y variables. If the particle is at rest with $\eta = 0$, the Gaussian form corresponds to the circle in Fig. 9. When the particle gains speed, this Gaussian function becomes squeezed into an ellipse. This ellipse becomes concentrated along the light cone with




Figure 9: Feynman’s rest of the universe. This figure is the same as that of Fig. 6. Here, the space variable z and the time variable t are entangled.

$t = z$, as η becomes very large.

The point is that we are able to observe this effect in the real world. These days, the velocity of protons from high-energy accelerators is very close to that of light. According to Gell-Mann [54], the proton is a bound state of three quarks. Since quarks are confined in the proton, they have never been observed, and the binding force must be like that of the harmonic oscillator. Furthermore, the observed mass spectra of the hadrons exhibit the degeneracy of the three-dimensional harmonic oscillator [43]. We use the word “hadron” for the bound state of the quarks. The simplest hadron is thus the bound state of two quarks.

In 1969 [55], Feynman observed that the same proton, when moving with its velocity close to that of light, can be regarded as a collection of partons, with the following peculiar properties.

1. The parton picture is valid only for protons moving with velocity close to that of light.
2. The interaction time between the quarks become dilated, and partons are like free particles.
3. The momentum distribution becomes wide-spread as the proton moves faster. Its width is proportional to the proton momentum.
4. The number of partons is not conserved, while the proton starts with a finite number of quarks.




Figure 10: The transition from the quark to the parton model through space-time entanglement. When $\eta = 0$, the system is called the quark model where the space separation and the time separation are dis-entangled. Their entanglement becomes maximum when $\eta = \infty$. The quark model is transformed continuously to the parton model as the η parameter increases from 0 to ∞ . The mathematics of this transformation is given in terms of circles and ellipses.




Figure 11: Entropy and temperature as functions of $[\tanh(\eta)]^2 = \beta^2$. They are both zero when the hadron is at rest, but they become infinitely large when the hadronic speed becomes close to that of light. The curvature for the temperature plot changes suddenly around $[\tanh(\eta)]^2 = 0.8$ indicating a phase transition.

Indeed, Fig. 10 tells why the quark and parton models are two limiting cases of one Lorentz-covariant entity. In the oscillator regime, the three-particle system can be reduced to two independent two-particle systems [43]. Also in the oscillator regime, the momentum-energy wave function takes the same form as the space-time wave function, thus with the same squeeze or entanglement property as illustrated in this figure. This leads to the wide-spread momentum distribution [20, 56, 57].

Also in Fig. 10, the time-separation between the quarks becomes large as η becomes large, leading to a weaker spring constant. This is why the partons behave like free particles [20, 56, 57].

As η becomes very large, all the particles are confined into a narrow strip around the light cone. The number of particles is not constant for massless particles as in the case of black-body radiation [20, 56, 57].

Indeed, the oscillator model explains the basic features of the hadronic spectra [43]. Does the oscillator model tell the basic feature of the parton distribution observed in high-energy laboratories? The answer is YES. In his 1982 paper [58], Paul Hussar compared the parton distribution observed in high-energy laboratory with the Lorentz-boosted Gaussian distribution. They are close enough to justify that the quark and parton models are two limiting cases of one Lorentz-covariant entity.

To summarize, the proton makes a phase transition from the bound state into a plasma state as it moves faster, as illustrated in Fig. 10. The un-observed time-separation variable becomes more prominent as η becomes larger. We can now go back to the form of this entropy given in Eq.(92) and calculate it numerically. It is plotted against $(\tanh \eta)^2 = \beta^2$ in Fig. 11. The entropy is zero when the hadron is at rest, and it becomes infinite as the hadronic speed reaches the speed of light.

Let us go back to the expression given in Eq.(87). For this ground state, the density matrix becomes

$$\rho_\eta(z, z') = \left(\frac{1}{\cosh \eta} \right)^2 \sum_k (\tanh \eta)^{2k} \chi_k(z) \chi_k(z'). \quad (105)$$

We can now compare this expression with the density matrix for thermally excited oscillator state [22]:

$$\rho_\eta(z, z') = (1 - e^{-1/T}) \sum_k [e^{-1/T}]^k \chi_k(z) \chi_k(z'). \quad (106)$$

By comparing these two expressions, we arrive at

$$[\tanh(\eta)]^2 = e^{-1/T}, \quad (107)$$

and thus

$$T = \frac{-1}{\ln [(\tanh \eta)^2]}. \quad (108)$$

This temperature is also plotted against $(\tanh \eta)^2$ in Fig. 11. The temperature is zero if the hadron is at rest, but it becomes infinite when the hadronic speed becomes close to that of light. The slope of the curvature changes suddenly around $(\tanh \eta)^2 = 0.8$, indicating a phase transition from the bound state to the plasma state.

In this section, we have shown how useful the concept of entanglement is in understanding the role of the time-separation in high energy hadronic physics including Gell-Mann's quark model and Feynman's parton model as two-limiting cases of one Lorentz-covariant entity.

Concluding Remarks

The main point of this paper is the mathematical identity

$$\frac{1}{\sqrt{\pi}} \exp \left\{ -\frac{1}{4} \left[e^{-2\eta}(x+y)^2 + e^{2\eta}(x-y)^2 \right] \right\} = \frac{1}{\cosh \eta} \sum_k (\tanh \eta)^k \chi_k(x) \chi_k(y), \quad (109)$$

which says that the series of Eq.(1) is an expansion of the Gaussian form given in Eq.(2).

The first derivation of this series was published in 1979 [26] as a formula from the Lorentz group. Since this identity is not well known, we explained in Sec. 5 how this formula can be derived from the generating function of the Hermite polynomials.

While the series serves useful purposes in understanding the physics of entanglement, the Gaussian form can be used to transfer this idea to high-energy hadronic physics. The hadron, such as the proton, is a quantum bound state. As was pointed in Sec. 8, the squeezed Gaussian function of Eq.(109) plays the pivotal role for hadrons moving with relativistic speeds.

The Bohr radius is a very important quantity in physics. It is a spatial separation between the proton and electron in the the hydrogen atom. Likewise, there is a space-like separation between constituent particles in a bound state at rest. When the bound state moves, it picks up a time-like component. However, in the present form of quantum mechanics, this time-like separation is not recognized. Indeed, this variable is hidden in Feynman's rest of the universe. When the system is Lorentz-boosted, this variable entangles itself with the measurable longitudinal variable. Our failure to measure this entangled variable appears in the form of entropy and temperature in the real world.

While harmonic oscillators are applicable to many aspects of quantum mechanics, Paul A. M. Dirac observed in 1963 [21] that the system of two oscillators contains also the symmetries of the Lorentz group. We discussed in this paper one concrete case of Dirac's symmetry. There are different languages for harmonic oscillators, such as the Schrödinger wave function, step-up and step-down operators, the Wigner phase-space distribution function. In this paper, we used extensively a pictorial language with circles and ellipses.

Let us go back to Eq.(109), this mathematical identity was published in 1979 as a textbook material in the American Journal of Physics [26], and the same formula was later included in a textbook on the Lorentz group [20]. It is gratifying to note that the same formula serves as a useful tool for the current literature in quantum information theory [59, 60].

References

- [1] Giedke, G.; Wolf, M. M.; Krueger, O; Werner, R. F.; Cirac, J. J. Entanglement of formation for symmetric Gaussian states *Phys. Rev. Lett.* **2003** *91*, 10790.1 - 10790.4.
- [2] Braunstein, S. L.; van Loock, P. Quantum information with continuous variables *Rev. Mod. Phys.* **2005** *28* 513 - 676.
- [3] Kim, Y. S.; Noz, M. E. Coupled oscillators, entangled oscillators, and Lorentz-covariant Oscillators *J. of Optics B: Quantum and Semiclassical* **2003** *7*, s459 - s467.
- [4] Ge, W; Tasgin, M. E.; Suhail Zubairy, S. Conservation relation of nonclassicality and entanglement for Gaussian states in a beam splitter *Phys. Rev. A* **2015** *92* 052328:1 - 9.
- [5] Gingrigh, R. M.; Adami, C. Quantum Engtangement of Moving Bodies *Phys. Rev. Lett.* **2002** *89*, 270402.1 - 4.
- [6] Dodd, P. J.; Halliwell, J. J. Disentanglement and decoherence by open system dynamics *Phys. Rev. A* **2004** *69* 052105.1 - 6.
- [7] Ferraro, A; Olivares, S; Paris, M. G. A. Gaussian states in continuous variable quantum information *EDIZIONI DI FILOSOFIA E SCIENZE (2005)* <http://arxiv.org/abs/quant-ph/0503237>. 1 - 100.
- [8] Adesso, G.; Illuminati, F. Entanglement in continuous-variable systems: recent advances and current perspectives *J. Phys. A* **2007** *40* 7821 - 7880
- [9] Walls, D. F.; Milburn, G. J. *Quantum Optics*. 2nd Ed.; Springer, Berlin, Germany, 2008.
- [10] Yuen, H. P. Two-photon coherent states of the radiation field *Phys. Rev. A* **1976** *13* 2226 - 2243.
- [11] Yurke, B; Potasek, M. Obtainment of Thermal Noise from a Pure State *Phys. Rev. A* **1987** *36* 3464 - 3466.
- [12] Ekert, A. K.; Knight, P. L. Correlations and squeezing of two-mode oscillations *Am. J. Phys.* **1989** *57* 692 - 697.
- [13] Paris, M. G. A. Entanglement and visibility at the output of a Mach-Zehnder interferometer *Phys. Rev. A* **1999** *59* 1615: 1 -7.
- [14] Kim, M. S.; Son, W; Buzek, V; Knight, P. L. Entanglement by a beam splitter: Nonclassicality as a prerequisite for entanglement *Phys. Rev. A* **2002** *65* 02323: 1 - 7.

- [15] Han, D.; Kim, Y. S.; Noz, M. E. Linear Canonical Transformations of Coherent and Squeezed States in the Wigner phase Space III. Two-mode States *Phys. Rev. A* **1990** *41* 6233 - 6244.
- [16] Dirac, P. A. M. The Quantum Theory of the Emission and Absorption of Radiation *Proc. Roy. Soc. (London)* **1927** *A114* 243 - 265.
- [17] Dirac, P. A. M. Unitary Representations of the Lorentz Group. *Proc. Roy. Soc. (London)* **1945** *A183* 284 - 295.
- [18] Dirac, P. A. M. Forms of relativistic dynamics *Rev. Mod. Phys.* **1949** *21* 392 - 399.
- [19] Yukawa, H. Structure and Mass Spectrum of Elementary Particles. I. General Considerations *Phys. Rev.* **1953** *91* 415 - 416.
- [20] Kim, Y. S.; Noz, M. E. *Theory and Applications of the Poincaré Group*; Reidel: Dordrecht, Holland, 1986.
- [21] Dirac, P. A. M. A Remarkable Representation of the 3 + 2 de Sitter Group *J. Math. Phys.* **1963** *4* 901-909.
- [22] Feynman, R. P. *Statistical Mechanics*; Benjamin Cummings: Reading, Massachusetts, U.S.A., 1972.
- [23] Han, D.; Kim, Y. S.; Noz, M. E. Illustrative Example of Feynman's Rest of the Universe *Am. J. Phys.* **1999** *67*, 61 - 66.
- [24] Kim, Y. S.; Noz, M. E. Covariant harmonic oscillators and the quark model *Phys. Rev. D* **1973** *8*, 3521 - 3627.
- [25] Kim, Y. S.; Noz, M. E.; Oh, S. H. Representations of the Poincaré group for relativistic extended hadrons *J. Math. Phys.* **1979** *20* 1341 - 1344.
- [26] Kim, Y. S.; Noz, M. E.; Oh, S. H. A simple method for illustrating the difference between the homogeneous and inhomogeneous Lorentz groups *Am. J. Phys.* **1979** *47* 892 - 897.
- [27] Kim, Y. S.; Wigner, E. P. *Entropy and Lorentz Transformations*. *Phys. Lett. A* **1990** *147* 343 - 347.
- [28] Kim, Y. S.; Noz, M. E. Lorentz Harmonics, Squeeze Harmonics and Their Physical Applications *Symmetry* **2011** *3* 16 - 36.
- [29] Klauder, J. R.; E. C. G. Sudarshan, E. C. G. *Fundamentals of Quantum Optics*; Benjamin: New York, U.S.A., 1968.
- [30] Saleh, B. E. A.; Teich, M. C. *Fundamentals of Photonics*. 2nd Ed.; John Wiley and Sons: Hoboken, New Jersey, U.S.A., 2007.

- [31] Miller, W. *Symmetry Groups and Their Applications*; Academic Press: New York, New York, U.S.A., 1972.
- [32] Hall, B. C. *Lie Groups, Lie Algebras, and Representations. An Elementary Introduction* Second Edition, Springer International, Cham, Switzerland, 2015
- [33] Wigner, E. On Unitary Representations of the Inhomogeneous Lorentz Group *Ann. Math.* **1939** *40* 149-204.
- [34] Weinberg, S. Photons and gravitons in S-Matrix theory: Derivation of charge conservation and equality of gravitational and inertial mass *Phys. Rev.* **1964** *135* B1049 - B1056.
- [35] Kim, Y. S.; Wigner, E. P. Space-time geometry of relativistic-particles *J. Math. Phys.* **1990** *31* 55 - 60.
- [36] Bařkal, S; Kim, Y. S.; Noz, M. E. Wigner's Space-Time Symmetries based on the Two-by-Two Matrices of the Damped Harmonic Oscillators and the Poincar Sphere *Symmetry* **2014** *6* 473 - 515.
- [37] Bařkal, S; Kim, Y. S.; Noz, M. E. *Physics of the Lorentz Group*. IOP Science ; Morgan & Claypool Publishers: San Rafael, California, U.S.A., 2015.
- [38] Kim, Y. S.; Yeh, Y. E(2)-symmetric two-mode sheared states *J. Math. Phys.* **1992** *33*, 1237 - 1246
- [39] Han, D.; Kim, Y. S.; Noz, M. E. O(3, 3)-like Symmetries of Coupled Harmonic Oscillators *J. Math. Phys.* **1995** *36* 3940 - 3954.
- [40] Kim, Y. S.; Noz, M. E. *Phase Space Picture of Quantum Mechanics*; World Scientific Publishing Company: Singapore, 1991.
- [41] Kim, Y. S.; Noz, M. E. *Dirac Matrices and Feynman's Rest of the Universe* Symmetry **2012** *4*, 626 - 643.
- [42] Wigner, E. On the Quantum Corrections for Thermodynamic Equilibrium *Phys. Rev.* **1932** *40* 749 - 759.
- [43] Feynman, R. P.; Kislinger, M; Ravndal, F. Current Matrix Elements from a Relativistic Quark Model *Phys. Rev. D* **1971** *3* 2706 - 2732.
- [44] Ruiz, M. J. Orthogonality relations for covariant harmonic oscillator wave functions *Phys. Rev. D* **1974** *10* 4306 - 4307.
- [45] Rotbart, F. C. Complete orthogonality relations for the covariant harmonic oscillator *Phys.Rev.D* **1981** *12* 3078 - 3090.
- [46] Magnus, W.; Oberhettinger, F.; Soni, R. P. *Formulas and theorems for the special functions of mathematical physics*; Springer-Verlag: Heidelberg, Germany 1966.

- [47] Doman, B. G. S. *The Classical Orthogonal Polynomials* World Scientific: Singapore 2016
- [48] Bargmann, V. Irreducible unitary representations of the Lorentz group *Ann. Math.* **1947** 48 568 - 640.
- [49] Han, D.; Kim, Y. S. Special relativity and interferometers *Phys. Rev. A.* **1988** 37 4494 - 4496.
- [50] Han, D.; Kim, Y. S.; Noz, M.E. Wigner rotations and Iwasawa decompositions in polarization optics *Phys. Rev. E.* **1999** 1 1036-1041.
- [51] von Neumann, J. *Die mathematische Grundlagen der Quanten-mechanik*; Springer: Berlin, Germany, 1932. See also von Neumann, I. *Mathematical Foundation of Quantum Mechanics*; Princeton University, Princeton, New Jersey, U.S.A., 1955.
- [52] Fano, U. Description of States in Quantum Mechanics by Density Matrix and Operator Techniques *Rev. Mod. Phys.* **1957** 29 74 - 93.
- [53] Wigner E. P.; Yanase, M. M. Information Contents of Distributions *Proc. National Academy of Sciences (U.S.A.)* **1963** 49 910 - 918.
- [54] Gell-Mann, M. A Schematic Model of Baryons and Mesons *Phys. Lett.* **1964** 8, 214 - 215.
- [55] Feynman, R. P. Very High-Energy Collisions of Hadrons *Phys. Rev. Lett.* **1969** 23 1415 - 1417.
- [56] Kim, Y. S.; Noz, M. E. Covariant harmonic oscillators and the parton picture *Phys. Rev. D* **1977** 15 335 - 338.
- [57] Kim, Y. S. Observable gauge transformations in the parton picture *Phys. Rev. Lett.* **1989** 63 348 - 351.
- [58] Hussar, P. E. Valons and harmonic oscillators *Phys. Rev. D* **1981** 23 2781 - 2783.
- [59] Leonhardt, U. *Essential Quantum Optics* Cambridge University Press, London, 2010.
- [60] Furusawa, A; Loock, P. V. *Quantum Teleportation and Entanglement: A Hybrid Approach to Optical Quantum Information Processing* Wiley-VCH, Weinheim, 2010.