

Effect of dissipative environment on collapses and revivals of a non-linear quantum oscillator

Maninder Kaur* and Bindiya Arora†

Department of Physics, Guru Nanak Dev University, Amritsar, Punjab-143005, India

Arvind‡

*Department of Physical Sciences, Indian Institute of Science Education
and Research (IISER) Mohali, PO Manauli Punjab 140306, India*

We study the dissipative dynamics of a wave packet passing through two different non-linear media. The effect of dissipation on the phenomenon of collapses and revivals of a wave packet as it evolves in a Kerr-type non-linear medium (represented by the Hamiltonian $(a^\dagger a)^2$) is investigated. We find that partial revivals do take place when dissipation values are moderate. For a certain regime of parameters we find a solution where revivals do not die even in the presence of dissipation and the non-linearity appears to compensate for the energy and coherence loss. We consider the next order non-linearity, represented by the Hamiltonian $(a^\dagger a)^3$, where we observe the phenomena of super revivals. The effect of dissipation in this case has an additional feature of number dependence for the displaced number states. While our simulations explore the degree to which the phenomena of collapses and revivals degrades in a dissipative environment, we also discovered the presence of a situation where degradation is minimal.

PACS numbers: 42.50.-p, 42.50.Lc

I. INTRODUCTION

Dynamics of quantum systems in complex environments plays an important role in diverse fields of science [1–6]. A laser beam that is quantum mechanically represented by a coherent state, while passing through non-linear media, can undergo a variety of complex transformations including collapses and revivals of the quantum state. Over and above any systematic evolution (linear or non-linear), dissipation is always present, and therefore a realistic study must include the effect of a dissipative environment [6–8]. Classically the dissipative effects lead to diminishing of the amplitude, however, since the interactions occur at atomic scales, quantum effects are important. An important set of quantum states is obtained by a phase space displacement of the number states (states with a fixed number of quanta n) [9–11]. A reasonable model for dissipation on the other hand is to model the dissipative medium as a set of oscillators that interact with the system [7]. For linear as well non-linear media the initial motion of the wave packet is periodic with the period of motion termed as the ‘classical period’ T_{cl} [12], corresponding to the natural frequency of the underlying harmonic oscillator. For a linear medium the periodic character of the motion is preserved; however, for a non-linear medium, after a few cycles quantum interference takes over leading to a significant spread of the wave packet. The wave packet is no longer recognizable as a packet and is said to have ‘collapsed’. As time

advances, it resurrects itself leading to its ‘revival’. The time at which this revival takes place is called the ‘revival time’ T_{rev} [13–15]. At times that are rational fractions of T_{rev} , the wave packet turns into a series of subsidiary packets and the phenomenon is called fractional revival. For the case of third order non-linearity, for times beyond revival times, a new sequence of full and fractional revivals start, which are characterized by a longer time scale called the ‘super revival time’ T_{sr} . The phenomenon of collapses and revivals of the wave packets was first discussed in [16] and thereafter, has been studied by a number of authors [17–24]. In most of these studies the medium is assumed to be non-dissipative. However, for situations of practical interest the medium can not be assumed to be non-dissipative or ideal. How to incorporate the effects of dissipation on the phenomena of collapses and revivals of wave packets is the central question that we address in the present paper.

Unlike in classical mechanics, the dissipative terms cannot be directly incorporated in quantum equations of motion as this leads to the decay of the Heisenberg uncertainty relation which is absurd. To overcome this difficulty various approaches have been proposed including those involving complex Hamiltonians [25–28]. We consider the model for dissipation where the system interacts with the environment via an interaction Hamiltonian. The composite system consisting of the system of interest plus the environment evolves unitarily and the environment degrees of freedom are traced over to arrive at the dynamics of the system alone [6, 7]. Concrete physical models involving nonclassical light and dissipative media in cavity QED, quantum wells and plasma systems have been proposed [29–31]. Rydberg atom in a micocavity weakly excited in the strong coupling limit has been investigated from this point of view [32]. Quan-

* mannu.711@yahoo.co.in

† bindiya.phy@gndu.ac.in

‡ arvind@iisermohali.ac.in

tum correlations reflected in quantum discord for open system has also been studied [33].

Our work embarks upon a study of the dynamics of a quantum wave packet through a non-linear medium in the presence of dissipation. In particular we aim to study the effects of dissipation on the phenomenon of wave packet revival and super-revival. We consider a single-mode radiation field as our system and take coherent states and displaced number states as our wave packets. Such packets can arise in many experimental situations [34–36]. For non-linear media we first consider a Kerr medium represented by the Hamiltonian $(a^\dagger a)^2$. The dynamics of the coherent wave packet in this case gives rise to revivals and collapses of the wave packet. To include the effects of dissipation we write down a master equation corresponding to a situation where the single mode radiation field interacts with a set of harmonic oscillators representing a thermal bath. Specifically we have assumed that we have single mode radiation in the visible region (e.g. $\lambda = 600$ nm red light). From the master equation, we numerically calculate the time evolution of the expectation value of the amplitude of the radiation field. Next, we study the one order higher non-linearity represented by a term proportional to $(a^\dagger a)^3$ in the Hamiltonian. This medium is a favorable system to observe the super revivals of the wave packet and we study these revivals under dissipation. Moreover, the dynamics of coherent wave packet in this medium provides the opportunity to analyze the effect of dissipation caused by the medium on the displaced number states. An important result we obtain is that the revivals as well as super-revivals can tolerate a certain amount of dissipation and thus can be observed in experiments where dissipation is inevitable. The tolerance to dissipation, however, depends upon the type and strength of the non-linearity.

A noteworthy result is that for a certain parameter regime for the Kerr-type non-linearity, revivals survive the effects of dissipation over a very long period of time. The simulations indicate that the loss of energy is compensated by the non-linearity. This is a situation which resembles that of solitons in classical systems. Furthermore, along with energy loss being compensated by non-linearity, the coherence is also preserved and the revivals are almost complete. To the best of our knowledge this is the first observation of this phenomenon. In the classical domain solitons are wave forms which travel in a lossy non-linear medium without losing shape and amplitude. Solitons have been extensively discussed for optical situations [37] and also for Kerr-type non-linearity [38, 39]. Solitons in physical system play an important role [40] Interaction of solitons with atomic systems has also been studied [41].

Strictly speaking solitons are associated with situations where dispersion is compensated by nonlinearity, however, the control of dissipation is also sometime called soliton and that is why we call our situation soliton like [42].

There have also been quantum generalizations of solitons for optical situations [43]. However in most of these cases the starting point is classical solitons which are then quantized. In our case the situation is very different, where we are working with an open quantum system with non-linearity being present and the “soliton-like” solutions automatically emerge for certain parameter regimes. These results are valid within the approximations and the assumptions of the dissipation model that we have used and are related to solitons only in the restricted sense as used in [42].

The paper is organized as follows: Section II provides the framework for our study with the model of dissipation discussed in Section II A, the kind of wave packets studied in Section II B and various time scales pertaining to the time evolution of wave packets in non-linear media taken up in Section II C. In Section III, we present the results for the time evolution of the expectation value of the amplitude operator in a dissipative environment. Specifically, Section III A contains results for the effect of dissipation for the case Kerr non-linearity and the “soliton-like” situation is discussed in III B, while Section III C contains results for the higher order non-linearity. In section IV we present some concluding remarks.

II. DISSIPATION MODEL AND REVIVAL TIMES FOR THE NON-LINEAR OSCILLATOR

A. System bath model

We study system-environment interaction in the Born-Markov approximation which yields the quantum master equation for the system. Consider a quantum system S interacting with its environment E , which can be treated as a heat bath or reservoir. The total (system + environment) Hamiltonian for this system can be written in the form [6]

$$H = H_S + H_E + H_{\text{int}} \quad (1)$$

where H_S , H_E , and H_{int} describe the Hamiltonian for the system, environment and interaction between them, respectively. For one mode radiation field represented by bosonic annihilation operator a with $[a, a^\dagger] = 1$, traversing through a non-linear medium the above Hamiltonians are given by the following expressions:

$$H_S = \hbar\omega_0 a^\dagger a + H_{\text{NL}} \quad (2)$$

$$H_E = \sum_j \hbar\omega_j e_j^\dagger e_j$$

$$H_{\text{int}} = \sum_j \hbar k_j [a e_j^\dagger + a^\dagger e_j]. \quad (3)$$

Here ω_0 is the frequency corresponding to the harmonic oscillator representing the mode and H_{NL} is the non-linear part of the Hamiltonian being considered. The dissipation is modeled as a set of harmonic oscillators with

frequencies ω_j , while e_j^\dagger and e_j are the creation and annihilation operators of the environment. The interaction Hamiltonian is in the rotating wave approximation [6] and k_j represents the coupling strength of the j th bath mode with the system.

In the Schrödinger picture, reduced density matrix ρ_S for the system is described by the master equation [3, 6, 7, 44–46]

$$\begin{aligned} \frac{d\rho_S}{dt} = & -\frac{\iota}{\hbar}[H_S, \rho_S] + \gamma(N+1)[a\rho a^\dagger - \frac{1}{2}(a^\dagger a\rho + \rho a^\dagger a)] \\ & + \gamma N[a^\dagger \rho a - \frac{1}{2}(aa^\dagger \rho + \rho aa^\dagger)]. \end{aligned} \quad (4)$$

Here the parameter $N = \left[\exp\left(\frac{\hbar\omega_0}{k_B T}\right) - 1\right]^{-1}$ represents the number of bath quanta at frequency ω_0 for a bath in thermal equilibrium at temperature T and γ is the damping rate. We refer the reader to reference [47] for the detailed derivation. For cases where $\hbar\omega_0 \gg k_B T$ (for instance, a photon of red light traversing an environment at room temperature), the energy will only flow from the system to the environment. This leads to further simplification of the master equation where we drop the last term in Equation (4) [8] to obtain

$$\frac{d\rho_S}{dt} = -\frac{\iota}{\hbar}[H_S, \rho_S] + \gamma[a\rho a^\dagger - \frac{1}{2}(a^\dagger a\rho + \rho a^\dagger a)]. \quad (5)$$

The system Hamiltonian H_S has two parts, the first being the linear part represented by $H_{S_0} = \hbar\omega_0 a^\dagger a$ and the second being the non-linear part H_{NL} leading to

$$\begin{aligned} \frac{d\rho_S}{dt} = & -\frac{\iota}{\hbar}[H_{S_0}, \rho_S] - \frac{\iota}{\hbar}[H_{NL}, \rho_S] + \\ & \gamma[a\rho a^\dagger - \frac{1}{2}(a^\dagger a\rho + \rho a^\dagger a)]. \end{aligned} \quad (6)$$

We consider two cases of non-linear terms H_{NL} as given below

$$\begin{aligned} H_{NL1} &= \hbar b_1 (a^\dagger a)^2, \\ H_{NL2} &= \hbar b_2 (a^\dagger a)^3, \end{aligned} \quad (7)$$

where b_1 and b_2 define the strength of nonlinearity and are in units of frequency.

Equation (6) can be written in the super-operator form as

$$\frac{d\rho_S}{dt} = \mathcal{L}\rho_S, \quad (8)$$

where \mathcal{L} is the Liouvillian given by

$$\mathcal{L} = -\frac{\iota}{\hbar}[H_{S_0} + H_{NL}, \rho_S] + \gamma[a\rho a^\dagger - \frac{1}{2}(a^\dagger a\rho + \rho a^\dagger a)] \quad (9)$$

For the case of a time independent Hamiltonian the solution of Equation (8) can be formally written as

$$\rho_S(t) = \exp[\mathcal{L}(t)]\rho_S(t_0). \quad (10)$$

From the above equation one can compute the time evolution of the expectation value of any physical quantity of interest. In particular, when we have a coherent wave packet, we would like to calculate the time varying expectation value of the wave packet amplitude $\langle a(t) \rangle$.

B. The wave packets

Coherent wave packets that we consider as initial states are obtained by phase space displacements of the vacuum state of the harmonic oscillator [9–11, 48],

$$\begin{aligned} |\alpha\rangle &= D(\alpha)|0\rangle \\ D(\alpha) &= \exp(\alpha a^\dagger - \alpha^* a) \end{aligned} \quad (11)$$

These are well known coherent states parameterized by the complex parameter α and typically represent laser light in quantum optics. For the corresponding harmonic oscillator, wave function for a coherent state is a Gaussian and the time evolution corresponds to a Gaussian wave packet oscillating in space with amplitude governed by $|\alpha|$.

The action of the displacement operator $D(\alpha)$ on number states $|n\rangle$ (states with fixed number of photons) gives rise to displaced number states given by

$$|\alpha, n\rangle = D(\alpha)|n\rangle \quad (12)$$

For the case of higher order non-linearity, we will also consider these states as initial states, in addition to coherent states. Starting with the initial density operator corresponding to the system state we will numerically compute the solution of Equation (10) and then compute the relevant expectation values.

C. Revival time scales

For non-linear oscillators when the Hamiltonian commutes with the number operator, the eigen vectors are same as those of the underlying linear oscillator. The energy of the n^{th} level on the other hand can be expanded as a Taylor series [20] around the central energy n_0 when the energy spread is not too much as follows

$$\begin{aligned} E_n \approx & E_{n_0} + (n - n_0)E'_{n_0} + \frac{1}{2}(n - n_0)^2 E''_{n_0} \\ & + \frac{1}{6}(n - n_0)^3 E'''_{n_0} + \dots \end{aligned} \quad (13)$$

where the primes over the energy terms represent derivatives with respect to n which define the various time scales (although n is discrete, it is useful to treat it as a continuous variable in this analysis). The first time scale is given by

$$T_{cl} = \frac{2\pi\hbar}{E'_{n_0}} \quad (14)$$

and is the ‘classical’ time period for the shortest closed orbit [12]. It controls the initial behavior of the wave packet. The second time scale

$$T_{rev} = \frac{2\pi\hbar}{\frac{1}{2}E''_{n_0}} \quad (15)$$

is the revival time [13, 14]. It governs the appearance of the fractional revivals and the full revivals. The third time scale

$$T_{sr} = \frac{2\pi\hbar}{\frac{1}{6}E_{n_0}'''} \quad (16)$$

is the super revival time. It is a larger time scale as compared to the classical time period and the revival time. These time scales become relevant depending upon the kind of non-linearity under consideration.

For the Kerr-type non-linearity the Hamiltonian is given by

$$H_{S_1} = \hbar\omega_0 a^\dagger a + \hbar b_1 (a^\dagger a)^2 \quad (17)$$

This Hamiltonian has the same eigen states as the original (linear) oscillator namely, $|n\rangle$, and the corresponding energy eigen values are given by

$$E_{n_1} = \hbar\omega_0 n_1 + \hbar b_1 n_1^2 \quad (18)$$

It is clear that the revival time is an important parameter here and is given by

$$T_{rev} = \frac{2\pi\hbar}{\frac{1}{2}(2b_1\hbar)} = \frac{2\pi}{b_1}. \quad (19)$$

Similarly, for a non-linear medium with non-linearity proportional to $(a^\dagger a)^3$ the Hamiltonian and energy are given as

$$H_{S_2} = \hbar\omega_0 a^\dagger a + \hbar b_2 (a^\dagger a)^3 \quad (20)$$

and corresponding energy eigen states are given by

$$E_{n_2} = \hbar\omega_0 n_2 + \hbar b_2 n_2^3. \quad (21)$$

We can compute the revival time as

$$T_{rev} = \frac{2\pi\hbar}{\frac{1}{2}(6b_2 n_2 \hbar)} = \frac{2\pi}{3b_2 n_2} \quad (22)$$

and the super-revival time as

$$T_{sr} = \frac{2\pi\hbar}{\frac{1}{6}(6b_2 \hbar)} = \frac{2\pi}{b_2}. \quad (23)$$

It is evident that the super-revival time becomes relevant in this case because of the presence of higher order non-linear terms in the Hamiltonian.

III. RESULTS AND DISCUSSION

We carry out the numerical solution of Equation (10) to compute the time evolved state of the system. We have evaluated all the physical quantities in atomic units (a.u.) by substituting \hbar and k_B equal to 1. These are convenient units and their use is a standard practice in atomic physics community. The simulations are done for




FIG. 1. The real part of expectation value $\langle a \rangle$ plotted as a function of time for coherent wave packet of red light in the presence of dissipation ($\gamma = 0.01$ a.u.) and without any non-linearity. Time axis is normalized with respect to revival time T_{rev} .




FIG. 2. Collapse and revival of a coherent wave packet of red photon passing through medium with Kerr type non-linearity i.e. proportional to $(a^\dagger a)^2$ for different values of γ . (a) $\gamma = 0.0$ a.u., (b) $\gamma = 0.00001$ a.u., (c) $\gamma = 0.0001$ a.u. and (d) $\gamma = 0.001$ a.u.

a variety of parameters to explore the effects of dissipation on the process of collapses and revivals for both types of non-linearities. We finally calculate the expectation value of oscillator amplitude $\langle a \rangle$ as a function of time. Throughout this paper when plot $\langle a \rangle$ in different circumstances, we mean its real part.




FIG. 3. Effect of weak and strong non-linearity represented by parameter b_1 , on the revival behavior for a coherent wave packet of red photon passing through medium without dissipation. The values of non-linearity are (a) $b_1 = 0.0000005$ a.u., (b) $b_1 = 0.000005$ a.u., (c) $b_1 = 0.05$ a.u. and (d) $b_1 = 1.0$ a.u.

The time step for evolution in the simulations is chosen to be much smaller than the classical time period which in turn is smaller than the revival time and the super revival time. The parameters in the numerical calculations are chosen to match a physically viable situation corresponding to red light with angular frequency $\omega_0 = 0.15\pi$ a.u. ($\lambda = 600$ nm), passing through a non-linear medium. The initial amplitude of the coherent state pertaining to red light is taken to be $\alpha = 1.9$. The damping parameter γ represents the overall coupling strength between the system and environment and $\gamma = 0.00001$ a.u., 0.0001 a.u. and 0.001 a.u. correspond to the weak, intermediate and strong damping respectively (γ value of 0.0001 a.u. corresponds to 6.58×10^{11} Hz). The strength of the non-linearity in the medium is defined by parameters b_1 and b_2 appearing in Equation (7). We study the effect of system-environment coupling on collapses and revivals of the wave packet by varying the dissipation parameter γ . In addition to this, we have also studied the behavior of revival and collapses as a function of non-linearity of the medium.

In the rest of this section, we first present the results obtained for dynamics of a coherent wave packet traversing a dissipative medium in the absence of any non-linearity. Thereafter, we discuss the findings for the media with non-linearity proportional to $(a^\dagger a)^2$ and $(a^\dagger a)^3$. In case of the non-linear term proportional to $(a^\dagger a)^2$, the medium behaves as an optical Kerr medium. The initial coherent state wave packet is constructed by displacing the vacuum state $|0\rangle$ as described in Equation (11). In case of non-linearity proportional to $(a^\dagger a)^3$, the dynamics of coherent wave packet reveals the existence of re-

vivals as well as super-revivals of the expectation values of oscillator amplitude operator $\langle a \rangle$. In this case, the revival time depends upon the principle quantum number n . Therefore we also study the effect of dissipative medium on the revivals of coherent wave packet corresponding to displaced number states for different values of n as defined in Equation (12). The analysis of displaced number states is important only in the case of higher-order non-linearity.

As a first computation, we consider the effect of dissipation on the linear oscillator without any non-linear terms. The results are displayed in Fig. 1 where we have plotted the amplitude $\langle a \rangle$ as a function of time. The simulation has been done for 59 classical time periods which is equivalent to one revival time T_{rev} corresponding to $b_1 = 0.008$ a.u., where we have taken the dissipation parameter $\gamma = 0.01$ a.u. In Fig. 1 and in all subsequent figures we have divided the real time by T_{rev} to have a uniform scale. The energy spectrum is equidistant and there is only one time associated with motion namely the classical time T_{cl} . Dissipation caused by environment is evident through the damping and ultimate decay of the expectation value of coherent wave packet amplitude $\langle a \rangle$ as time evolves. The decay of amplitude is exponential as expected.

A. Results for non-linearity proportional to $(a^\dagger a)^2$

For this case the results are presented in Fig. 2 where we display the expectation value of the amplitude $\langle a \rangle$ as a function of time for different values of dissipation. The value of the non-linearity parameter b_1 is taken to be 0.0005 a.u. It is evident from Fig. 2(a) that in the absence of coupling between system and environment (marked by zero value of γ) full revival of amplitudes takes place with a periodicity of $kT_{\text{rev}}/2$ with $k = 1, 2, 3 \dots$. At these revival times all eigenstates accumulate a phase of $2\pi k$. This behavior has already been studied theoretically [49] as well as experimentally [23]. For finite but small damping parameter $\gamma = 0.00001$ a.u. as shown in Fig. 2(b), due to damping a reduction in revived amplitude is noticeable although the revivals do take place. As the value of γ increases the revivals become weaker and weaker as seen in (Fig. 2(c) & (d)). For a certain value of γ , approximately equal to 0.001 a.u., the revivals disappear altogether. Our results on the one hand show robustness of the revival phenomena in a dissipative environment and on the other hand point towards a threshold level of dissipation after which no revivals occur whatsoever.

In Fig. 3, we study the effect of weak and strong non-linearity on the revival behavior. For zero non-linearity there are perfect revivals with a time period of T_{cl} . Inclusion of non-linear term in the Hamiltonian results in an erratic energy spectrum leading to the collapse and subsequent revival of the wave packet. For very small values of the non-linear parameter there is no perceptible effect on the pattern as seen in fig. 3(a) where we have cho-




FIG. 4. Plots of correlation functions to ascertain the self-similarity of revival patterns. (a) The correlation function is calculated with a time gap which is a multiple of T_{cl} (we have taken $n = 1500$). The sudden drop in this correlation function indicates the onset of collapse at b_{onset} . (b) The correlation function between the the revival patterns is plotted as a function of non-linearity parameter b_1 to ascertain the self-similarity of revival pattern. It turns out that the correlation function decays smoothly and we have to decide on a b_{offset} by visually ascertaining the lack of similarity of the pattern.

sen a value $b_1 = 0.0000005$ a.u. However, as seen from fig. 3(b) the packet undergoes a revival for small values of non-linearity at time T_{rev} . As the value of the non-linearity parameter is increased the revivals survive up to some value (Fig. 3(c)) and then disappear (Fig. 3(d)). For Fig. 3(d), the value of the non-linearity parameter is chosen to be large ($b_1 = 1.0$ a.u.) and the pattern is irregular. As we scan different values of the non-linearity parameter b_1 , we find that there is a threshold value of b_1 for which the state actually collapses and then revives at T_{rev} . We call this value of non-linearity parameter as b_{onset} and in our case $b_{onset} = 0.0000045$ a.u. This value is ascertained from the correlation function plot given in Fig. 4(a). Here it is very clear that the correlation function between the values of the $\langle a \rangle$ separated by a time which is a multiple of T_{cl} (We have taken $1500T_{cl}$) falls suddenly at this value of the non-linear parameter b_1 . As the value of the non-linearity parameter is further increased, the revivals continue. The correlation func-

tion between the values $\langle a \rangle$ separated by $0.5T_{rev}$ drops somewhat smoothly and linearly with the nonlinear parameter b_1 as shown in Fig. 4(b). Therefore, there is no clear threshold after which we can say that the revivals are not taking place. However, a visual inspection of the graphs shows that they survive up to some value of $b_1 \approx 0.05$ (Fig. 3(c) and Fig. 4(b)) and then the pattern looks different from the original (Fig. 3(d)). We thus define non-linearity parameter above which the revival pattern disappears as b_{offset} and assign it a value 0.05 when the correlation function drops to 0.7.

B. Presence of non decaying solutions

Is there a possibility of having non-linearity compensate dissipation leading to a solution where we have the excitation live for a long time? It turns out that such a parameter regime does exist as we have presented in Fig. 5. For a given value of the dissipation we slowly increase the value of non-linearity. We observe that for a given value of dissipation constant we need to increase the strength of non-linearity upto a certain value to achieve a situation where revivals begin to occur with full strength. This is very similar to solitons of classical physics where we can have non-linearity and dissipation acting in such a way that a persistent waveform is generated. It is interesting and surprising that such a situation arises in the case of this fully quantum treatment with dissipation modeled via the master equation. To observe the actual dynamics in this case we choose the non-linearity parameter in such a way that the soliton-like solution is observed and then plot the expectation value of $\langle a \rangle$ as a function of time on a time scale much smaller than T_{rev} . We observe the revivals with full strength upto several periods. This is depicted in Fig. 6. Furthermore, if nonlinearity is increased further the compensation situation disappears again. We would like to caution the reader that these solitons are not of the variety where dispersion is compensated by nonlinearity and it is dissipation that is being compensated and in the literature such situations have been called solitons and therefore, following [42] we call our situation soliton like. Secondly, the presence of these features is linked to the model of dissipation that we have used and the actual physical relevance and explanation of these results require further exploration.

C. Results for non-linearity proportional to $(a^\dagger a)^3$

To begin with we start with a coherent state wave packet and study its behavior in the presence of non-linearity and for different values of the dissipation parameter as was done for the case of Kerr non-linearity. Figures 7 (a),(b), (c) and (d) show the super-revival pattern of coherent wave packet amplitude $\langle a \rangle$ for $\gamma = 0, 0.00001, 0.0001$, and 0.001 a.u., respectively. The




FIG. 5. The plot $\langle a \rangle$ (real part) as a function of time period for coherent wave packet passing through a non-linear medium for (a) $\gamma = 0.00001$ a.u., (b) $\gamma = 0.0001$ a.u. and (c) $\gamma = 0.001$ a.u. at various values of non-linearity parameter. The uppermost curve in all the cases pertains to non-linearity compensating for dissipation leading to persistent revival pattern.

super-revival pattern is evident in these figures. Here the super-revival time T_{sr} is independent of the principle quantum number n (Equation (23)) similar to the case studied in the previous section where the revival time was independent of n . Within one super revival time some full revivals and some partial revivals come into the picture. The revival of the state occurs at the super revival time which is a longer time scale compared to the revival time. For the sake of comparison with the previous section we further study the effect of strength of non-linearity in the system given by the parameter value b_2 . From fig. 8, one can conclude that for a medium with non-linear term proportional to $(a^\dagger a)^3$ revivals are present up to a certain value of non-linearity beyond which the pattern becomes irregular. For values of b_2 (figs. 8(b) & (c)) the revivals are present. For b_2 values greater than 0.01 the revivals are irregular. For strong non-linearity $b_2 = 2.0$ a.u., the revivals are not only irregular but also almost stop appearing, which is depicted in Fig. 8(d). As we discussed for the earlier case we have the notions of b_{onset} and b_{offset} here too. Figure 8(a) ($b_2 = 0.0000002$) a.u. pertains to




FIG. 6. The plot of $\langle a \rangle$ (real part) as a function of time on a fine grained time scale (time step much smaller than T_{rev}) for a coherent wave packet passing through a non-linear medium in the regime where dissipation is compensated by non-linearity. Value of non-linearity parameter in the soliton-like behavior is $b_S = 0.01, 0.47, 0.484$ a.u. in (a), (b) and (c) respectively.

a value of non-linearity below b_{onset} . In this case too, from a scan through different values of b_2 we find that $b_{onset} = 0.000002$ a.u. and $b_{offset} = 0.01$ a.u. approximately.

D. The effect of n

The fact that revival time in this case shows dependence on principle quantum number n (Equation (22)), provides an opportunity to explore the revival pattern for various values n for displaced number states. As defined in Equation (12) we considered displaced number states $|\alpha, n\rangle$.

The results of this study are shown in Figs. 9 and 10. It is seen that the revival patterns for displaced number states are dependent upon n . We have taken the values $n = 1, 2, 3, 4$, with non-linearity $b_2 = 0.0005$ a.u., in the absence and presence of the system-environment coupling, respectively. Fig. 9 represents the results without




FIG. 7. Super-revival pattern of coherent state wave packet. Amplitude $\langle a \rangle$ (real part) is plotted as function of time in a medium with non-linearity proportional to $(a^\dagger a)^3$ for different values of dissipation values γ . (a) $\gamma = 0$, (b) $\gamma = 0.00001$ a.u., (c) $\gamma = 0.0001$ a.u. and (d) $\gamma = 0.001$ a.u.




FIG. 8. Effect of weak and strong non-linearity represented by parameter b_2 , on the revival behavior for a coherent wave packet of red photon passing through medium with non-linearity proportional to $(a^\dagger a)^3$ for various values of the non-linearity parameter b_2 . (a) $b_2 = 0.0000002$ a.u., (b) $b_2 = 0.000002$ a.u., (c) $b_2 = 0.01$ a.u., and (d) $b_2 = 2.0$ a.u.




FIG. 9. Collapse and revival pattern of displaced number states with amplitude $\langle a \rangle$ (real part) being plotted as a function of time with non-linearity proportional to $(a^\dagger a)^3$ and no dissipation. The pattern is shown for states constructed around $n = 1, 2, 3, 4$ at $\gamma = 0$ a.u. The x-axis is normalized with respect to theoretical T_{rev} which is around 4000, 2000, 1330 and 1000 a.u. for $n = 1, 2, 3$ and 4 respectively.




FIG. 10. Collapse and revival pattern of displaced number states with amplitude $\langle a \rangle$ (real part) being plotted as a function of time with non-linearity proportional to $(a^\dagger a)^3$ and in the presence of dissipation. The pattern is shown for states constructed around $n = 1, 2, 3, 4$ at $\gamma = 0.00001$ a.u. The x-axis is normalized with respect to theoretical T_{rev} which is around 4000, 2000, 1330 and 1000 a.u. for $n = 1, 2, 3$ and 4 respectively.

dissipation and Fig. 10 represents the results in the pres-

ence of dissipation. The revival pattern is clearly affected by the dissipative effects.

Theoretically calculated revival times for $n = 1, 2, 3$ and 4 are around 4000, 2000, 1330 and 1000 a.u. respectively. However, from Fig. 9(a), one can notice that the revival of the wave packet occurs at half of the theoretical revival time. This behavior for the $n = 1$ case was discussed in [49] and [23], where it was shown that revivals appear only when all eigenstates have accumulated a phase of $2\pi k$ and in the process some extra revivals may appear. However, from our Figs. 9(c)& (d), we also observe that there are cases where some revivals are missing for higher n values. It is evident from these sub-figures that revivals of wave packets occur at 2000 a.u. independent of the n value for the state $|\alpha, n\rangle$.

A comparison of Fig. 9 with Fig. 10 also indicates that the wave packet constructed around the higher displaced states are dissipated more prominently. In order to elucidate this finding, we extend the value of n for the state $|\alpha, n\rangle$ upto $n = 10$ and study the effect of dissipative medium at the first revival time.

The results are shown in Fig. 11, where the displaced number states constructed for higher n witness more dissipation. This is an interesting observation which is related to the fact that the displaced number states with higher n are more nonclassical and hence the effect of dissipation is rapid. Similar results were found for a related phenomenon of decoherence in [8].




FIG. 11. Effect of dissipative medium on the displaced number states constructed around various n in a medium with non-linearity proportional to $(a^\dagger a)^3$. The amplitude of revived wave packets at their first revival time instant are recorded for various values of n . The dissipation of amplitude is more vigorous for higher values of n .

IV. CONCLUDING REMARKS

In this work the quantum master equation approach has been used to analyze the dissipative dynamics of a coherent wave packet passing through non-linear media of two different types. The effect of a dissipative bath on

the phenomena of collapses and revivals is evident. However, it is observed that the basic features of the revival process survives some amount of dissipation. Although the revival is never complete in the presence of dissipation, its signatures are present. This is relevant for any attempt to observe these revivals experimentally.

The other prominent role played by the dissipative medium is that of an environment which soaks the energy of the wave packet. Dissipation is observed in the damping of successive revived coherent wave packet amplitudes $\langle a \rangle$. The controlling element for dissipation is the damping factor γ which further depends upon the coupling between the coherent wave packet and the medium. With increase in the strength of damping factor γ , the amplitude of the successively revived wave packet decreases and ultimately no revivals occur for a particular value of γ .

A particularly interesting situation arises for a certain range of parameters where the non-linearity and dissipation are poised in such a way that the revivals occur even in the presence of dissipation for a long time. In this regime of parameters, the amplitude of the revivals remains constant and there is no loss of energy. This is very similar to the phenomena of solitons in classical non-linear waves in the presence of dissipation. We believe this observation may have useful consequences for quantum communication protocols where one is interested in fabricating low dissipation communication channels.

For a medium with non-linearity proportional to $(a^\dagger a)^3$, signatures of super revivals are evident. Theoretically, revivals are predicted at different times for wave packets constructed around different displaced quantum numbers n . However, we observe that the revival of the wave packet occurs at a fixed time independent of the n . This behavior sheds light on a very fundamental aspect that revivals appear only when all eigenstates have accumulated a phase of $2\pi k$ and in the process some extra revivals may appear and some predicted ones may be missing.

We also observe that the wave packet build around the higher values of principle quantum number are more prone to the dissipative effects of the medium as compared to the low lying states. The extent of non-linearity of the medium also has an effect on the dynamics of the wave packet. We come across a value of the non-linearity parameter b_{onset} below which no collapses and hence no revivals occur. Similarly, there is an upper limit of b that we call b_{offset} , above which the pattern of revivals is irregular.

We have demonstrated that revivals can tolerate certain amount of dissipation, this certainly is useful in observing this phenomena in the lab. The soliton like solutions that we have found if experimentally realized may turn out to be useful in many practical situations. However, more work is needed to evaluate the relevance and possible applications of these results.

ACKNOWLEDGMENTS

B.A. acknowledges DST, INDIA for funding under Grant No. EMR/2016/001228. Arvind ac-

knowledges DST India for funding under Grant No. EMR/2014/000297.

-
- [1] R. Graham and F. Haake, *Quantum Statistics in Optics and Solid-State Physics (Springer Tracts in Modern Physics)*, Vol. 66 (Springer-Verlag Berlin Heidelberg, 1973).
- [2] W. H. Louisell, *Quantum Statistical Properties of Radiation (Wiley Classics Library)* (Wiley-VCH, 1990).
- [3] A. Caldeira and A. Leggett, *Physica A: Statistical Mechanics and its Applications* **121**, 587 (1983).
- [4] H. Grabert, P. Schramm, and G.-L. Ingold, *Physics Reports* **168**, 115 (1988).
- [5] H. Grabert, U. Weiss, and P. Talkner, *Zeitschrift für Physik B Condensed Matter* **55**, 87 (1984).
- [6] C. W. Gardiner and P. Zoller, *Quantum noise: A handbook of Markovian and non-Markovian quantum stochastic methods with applications to quantum optics (Springer series in synergetics) 3rd edition* (Springer-Verlag Berlin Heidelberg Germany, 2004).
- [7] D. F. Walls and G. J. Milburn, *Quantum Optics 2nd edition* (Springer-Verlag Berlin Heidelberg Germany, 2008).
- [8] P. Földi, M. G. Benedict, A. Cziráj, and B. Molnár, *Phys. Rev. A* **67**, 032104 (2003).
- [9] R. J. Glauber, *Phys. Rev.* **130**, 2529 (1963).
- [10] E. C. G. Sudarshan, *Phys. Rev. Lett.* **10**, 277 (1963).
- [11] J. R. Klauder and E. C. G. Sudarshan, *Fundamentals of Quantum Optics* (W. A. Benjamin Inc., New York, 1968).
- [12] A. Peres, *Phys. Rev. A* **47**, 5196 (1993).
- [13] O. Knospe and R. Schmidt, *Phys. Rev. A* **54**, 1154 (1996).
- [14] R. Arvieu and P. Rozmej, *Phys. Rev. A* **51**, 104 (1995).
- [15] I. S. Averbukh, *Phys. Rev. A* **46**, R2205 (1992).
- [16] C. U. Segre and J. D. Sullivan, *American Journal of Physics* **44**, 729 (1976).
- [17] D. L. Aronstein and C. R. Stroud, *Phys. Rev. A* **55**, 4526 (1997).
- [18] R. W. Robinett, *American Journal of Physics* **68**, 410 (2000).
- [19] D. F. Styer, *American Journal of Physics* **69**, 56 (2001).
- [20] R. Robinett, *Physics Reports* **392**, 1 (2004).
- [21] C. Sudheesh, S. Lakshmibala, and V. Balakrishnan, *Journal of Optics B: Quantum and Semiclassical Optics* **7**, S728 (2005).
- [22] V. Y. Demikhovskii, A. V. Telezhnikov, E. V. Frolova, and N. A. Kravets, *Low Temperature Physics* **39**, 18 (2013).
- [23] B. Gruner, M. Schlesinger, P. Heister, W. T. Strunz, F. Stienkemeier, and M. Mudrich, *Phys. Chem. Chem. Phys.* **13**, 6816 (2011).
- [24] M. Kaur, B. Arora, and M. Mian, *Pramana J. of Phys.* **86**, 31 (2016).
- [25] A. G. Redfield, *IBM Journal of Research and Development* **1**, 19 (1957).
- [26] R. Feynman and F. Vernon, *Annals of Physics* **24**, 118 (1963).
- [27] S. Nakajima, *Progress of Theoretical Physics* **20**, 948 (1958).
- [28] H. Eleuch and I. Rotter, *Phys. Rev. A* **95**, 022117 (2017).
- [29] H. Eleuch, J. M. Courty, G. Messin, C. Fabre, and E. Giacobino, *Journal of Optics B: Quantum and Semiclassical Optics* **1**, 1 (1999).
- [30] H. Eleuch, *The European Physical Journal D* **48**, 139 (2008).
- [31] H. Eleuch, N. Ben Nessib, and R. Bennaceur, *The European Physical Journal D - Atomic, Molecular, Optical and Plasma Physics* **29**, 391 (2004).
- [32] H. Jabri, H. Eleuch, and T. Djerad, *Laser Physics Letters* **2**, 253 (2005).
- [33] K. Berrada, H. Eleuch, and Y. Hassouni, *Journal of Physics B: Atomic, Molecular and Optical Physics* **44**, 145503 (2011).
- [34] S. A. Podoshvedov, *J. Opt. Soc. Am. B* **31**, 2491 (2014).
- [35] N. Pop, D. Popov, and M. Davidovic, *International Journal of Theoretical Physics* **52**, 2275 (2013).
- [36] A. S. Coelho, L. S. Costanzo, A. Zavatta, C. Hughes, M. S. Kim, and M. Bellini, *Phys. Rev. Lett.* **116**, 110501 (2016).
- [37] S. K. Turitsyn, B. G. Bale, and M. P. Fedoruk, *Physics Reports* **521**, 135 (2012), dispersion-Managed Solitons in Fibre Systems and Lasers.
- [38] G. Mehmetallah and P. P. Banerjee, *J. Opt. Soc. Am. B* **23**, 203 (2006).
- [39] S. K. Adhikari, *Phys. Rev. E* **71**, 016611 (2005).
- [40] R. Camassa, J. M. Hyman, and B. P. Luce, *Physica D: Nonlinear Phenomena* **123**, 1 (1998), annual International Conference of the Center for Nonlinear Studies.
- [41] H. Eleuch and R. Bennaceur, *Journal of Optics A: Pure and Applied Optics* **5**, 528 (2003).
- [42] M. Remoissenet, *Waves Called Solitons: Concepts and Experiments (Advanced Texts in Physics) 3rd edition* (Springer-Verlag Berlin Heidelberg, 2003).
- [43] P. Drummond, R. Shelby, S. Friberg, and Y. Yamamoto, *Nature* **365**, 307 (1993).
- [44] W. H. Zurek, *Phys. Rev. D* **26**, 1862 (1982).
- [45] W. G. Unruh and W. H. Zurek, *Phys. Rev. D* **40**, 1071 (1989).
- [46] B. L. Hu, J. P. Paz, and Y. Zhang, *Phys. Rev. D* **45**, 2843 (1992).
- [47] H. J. Carmichael, *Statistical Methods in Quantum Optics 1* (Springer-Verlag Berlin Heidelberg Germany, 2013).
- [48] J. R. Klauder, *Annals of Physics* **11**, 123 (1960).
- [49] M. Kaur, M. Mian, and M. kaur, *i-manager's Journal of Material Science(JMS)* **3**, 31 (2015).